

The Gravitational Ringing of a Spherically Symmetric Black Hole Surrounded by Dark Matter Spike

Yuqian Zhao,^{1,2} Bing Sun,³ Kai Lin,⁴ and Zhoujian Cao^{1,2,5,*}

¹*Institute for Frontiers in Astronomy and Astrophysics,
Beijing Normal University, Beijing 102206, China*

²*Department of Astronomy, Beijing Normal University, Beijing 100875, China*

³*CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,
Chinese Academy of Sciences, Beijing 100190, China*

⁴*Hubei Subsurface Multi-scale Imaging Key Laboratory, School of Geophysics and Geomatics,
China University of Geosciences, Wuhan 430074, Hubei, China*

⁵*School of Fundamental Physics and Mathematical Sciences,
Hangzhou Institute for Advanced Study, UCAS, Hangzhou 310024, China*

Supermassive black holes at the center of each galaxy may be surrounded by dark matter. Such dark matter admits a spike structure and vanishes at a certain distance from the black hole. This dark matter will impact the spacetime near the black hole and the related ringing gravitational waves can show distinguished features to the black hole without dark matter. In the present work, we calculate the quasi-normal modes who dominate the ringdown process of the perturbed black holes surrounded by dark matter spike. We find that the isospectrality for black holes with dark matter spike will break down. And more the relative ringing frequency difference between the black holes with and without dark matter can be as large as 10^{-2} . These features can be used in future gravitational wave detection about extremal mass ratio inspiral systems to probe the existence of dark matter around supermassive black holes.

I. INTRODUCTION

Black holes (BHs) are important predictions of general relativity (GR) [1]. In the past century, many scientists have endeavored to explore the existing evidences and characteristics of black holes [2]. Since the first detection of gravitational wave (GW) event in 2015 [3], more than 90 directly observed black hole binary events have been detected [4–8]. The observations of the shadow from M87 and our galaxy also provided additional evidence for supermassive black holes (SMBHs) [9–16].

The event horizon is the distinguishable feature of BHs. Event horizon, as a one-way causal boundary, prevents us from detecting the inside of a black hole [17]. Black holes in our universe may be affected by astronomical environments, such as the dark matter (DM) near the BHs [18–20]. It is believed that 90% of the host galaxies of SMBHs are composed of DM [21]. It is interesting to ask how the DM around the BHs impacts the spacetime and GWs related to these BHs.

The exploration of DM has a long history. The numerical results from N -body cosmological simulations suggest the density distribution of DM is peaked near the center of galaxies and decreases as a power of $1/r$ with r the distance from the halo's center, as given by [22]:

$$\rho(r) = \rho_0 (r/r_0)^{-\gamma_0} (1 + (r/r_0)^{\alpha_0})^{(\gamma_0 - \beta_0)/\alpha_0} \quad (1)$$

where r_0 and ρ_0 are the scale factors determined by the numerical fitting such as [23], while α_0 , β_0 and γ_0

are the parameters determined by the choice of models with $(\alpha_0, \beta_0, \gamma_0) = (1, 4, 1)$ for Hernquist profile and $(\alpha_0, \beta_0, \gamma_0) = (1, 3, 1)$ for Navarro-Frenk-White (NFW) profile [24, 25].

With the presence of a BH, the density distribution of DM changes. In an early pioneering work [26], a Newtonian method is used to calculate the redistribution of cold DM near the center of galaxies. The BH accretion makes the DM form a cusp (“spike”) structure. For spherically symmetric BHs, the density peaks near $r \sim 4R_s$ with R_s the Schwarzschild radius and has a steep cutoff at $r = 4R_s$, below which the density of DM vanishes due to annihilation or dropping into the BHs. With the relativistic modification [27], the density profile has similar characteristics, but the cutoff radius will be changed to $r = 2R_s$.

In [22], both Hernquist and NFW models with or without relativistic modifications are introduced to explore the impacts of the DM spike on the extreme mass-ratio inspirals (EMRIs) GW waveform. And the relativistic modifications are found to have a positive impact on the detectability of DM for both models. In the current work, we use the same profile to explore the impacts of the ringdown waveform, specifically, the quasi-normal modes (QNMs) of the perturbed BHs [28–32]. This DM profile takes the BHs Eddington accretion effect into consideration instead of Bondi accretion effect [33]. Not like Refs. [34, 35] this DM profile does not consider the self-interactions of DM. Different DM profile may change the QNMs properties.

The detection of ringdown signals with ground-based GW detectors is not satisfactory due to the low signal-to-noise (SNR) [36]. Taking the advantage of high SNR, the impacts of DM spike on ringdown of SMBHs may

* Zhoujian Cao: zjcao@amt.ac.cn

be detected by the future space-based GW detectors including LISA, Taiji, Tianqin and DECIGO [37{39]. As an example, we will discuss the detectability of such signals for the SgrA BH at the center of Milky Way [23], whose ringdown frequency exactly falls into the sensitive frequency bands of space-based GW detectors [40].

In [41{45], the impacts of the cold DM halo on the ringdown process are explored. And the impacts of the DM spike with profile (1) and $\rho_0 = \rho_0$ on the QNMs of scalar perturbations can be found in [46]. Here we extend the investigation to gravitational perturbations with Hernquist profile $(\rho_0; \rho_0; \rho_0) = (1; 4; 1)$ and Navarro-Frenk-White (NFW) profile $(\rho_0; \rho_0; \rho_0) = (1; 3; 1)$. Some new features will be added in the current work, for example, the breaking isospectrality [28, 31], which means the spectra from axial and polar perturbations are not identical.

The structure of this paper is as follows: based on the profile of DM spike and using the TOV equations, we derive the modified gauge metric of spherically symmetric BHs in Sec. II. We then derive the QNMs equation for both axial and polar parities in Sec. III. After that, we solve the QNMs equations with numerical methods and present the results in Sec. IV. We will use the SgrA with the DM spike as an example to explore the impacts of different DM spike on QNMs and discuss the detectability of such impacts. Throughout the paper, we use the geometric unit system so that $c = G = 1$, where c is the speed of light and G is the gravitational constant.

II. BLACK HOLES SURROUNDED BY DARK MATTER SPIKE

In this section, we consider the spherically symmetric BHs surrounded by the DM spike and derive the modified metric. The most well-known solution for the spherically symmetric static BHs spacetime is the Schwarzschild BHs, which characterize the vacuum scenario. In general, such an isolated BH does not exist, particularly for SMBHs in the center of galaxies, which are more or less surrounded by some matters with complex distribution, such as DM. Then, a Schwarzschild BH in DM spike can be well described by the most general metric:

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{g(r)} + r^2(d\theta^2 + \sin^2\theta d\phi^2): \quad (2)$$

and the contribution of the DM can be thought of as the energy-momentum tensor:

$$T = \text{diag} \{ f(r); p(r); p(r); p(r) \} g \quad (3)$$

where $f(r)$ depends on the density distribution of DM. Based on [22], we introduce the density profile of the DM spike as:

$$\rho(r) = \rho_0 \left(1 - \frac{4M}{r} \right)^w \frac{M}{r}^q \quad (4)$$

with $\rho_0 = 4:17 \cdot 10^{11}$ and ρ_0 is the effective density of the DM spike given by:

$$\rho_0 = A \cdot 10^{-10} \frac{\rho_0}{0:3\text{GeV}=\text{cm}^3} \frac{M}{10^6 M_\odot} \frac{r_0}{20\text{kpc}} \quad (5)$$

where the parameter $A = 1(2)$ for relativistic (Newtonian) DM profiles and limits the location of a steep cutoff at $r = 4M$ (corresponding to $2R_s$ and $4R_s$ respectively with R_s the Schwarzschild radius), below which the DM density vanishes due to annihilation or dropping into the BH as shown in Fig. 1. The parameters $(; ; ;)$ depend on whether the Hernquist model or the NFW model is chosen. The parameters $A; w; q$ are the fitting factors. ρ_0 and r_0 are the scale factors in Eq.(1) and we choose the range the same as [22], while M is the mass of the BHs and the range is $10^6 - 10^9$ for SMBHs [32]. Table I lists all the parameter values with the range of ρ_0 .

Inspired by the work of [46], we make the same assumption that the DM distributes like dust and the pressure $p(r)$ can be neglected with respect to $f(r)$ and $(1 - g(r)) = 8r^2$ as shown in Fig. 2. Einstein equations $G = 8T$ or the TOV equations [46{48] now read:

$$rg'(r) + g(r) - 1 + 8r^2 = 0; \quad (6)$$

$$f(r)g(r) - f(r) + rg(r)f'(r) = 0; \quad (7)$$

$$p'(r) = -r \frac{1}{rg(r)} \frac{g(r)}{r}; \quad (8)$$

By using Eq. (4), we then integrate Eq. (6) from $4M$ to the spatial infinity and obtain:

$$g(r) = 1 - \frac{2M}{r} + \frac{8\rho_0(M)^q}{q-3} r^{2-q} {}_2F_1(q-3; w; q-2; \frac{4M}{r}) \quad (9)$$

where ${}_2F_1$ is the hypergeometric function [49, 50]. And we have chosen the constant of integration so that $g(4M) = 1 - \frac{2M}{r}$.

By substituting Eq. (9) and Eq. (4) into Eq. (7), $f(r)$ can be obtained as:

$$f(r) = 1 - \frac{2M}{r} \exp \left[\frac{1}{(q-3)(q-3)} \sum_{n=0}^{N_{\text{max}}} 8(q-2) \rho_0 r^2 \frac{2M^n}{r} \frac{M}{r}^q (q-3) {}_2F_1(q-3; w; q-2; \frac{4M}{r}) \right. \\ \left. - (n+q-2) {}_2F_1(n+q-2; w; n+q-1; \frac{4M}{r}) \right] \\ \cdot M^2 \cdot 2^{7-n-2q} \cdot 2^{n-q} \cdot (q-2) \cdot (w+1) \frac{(q-3)}{(q+w-2)} \frac{(n+q-2)}{(n+q+w-1)} \quad (10)$$




FIG. 1: The DM profiles for Hernquist Newtonian model, Hernquist relativistic model, NFW Newtonian model and NFW relativistic model. The parameters are $M = 10^6 M_\odot$, $\rho_0 = 0.3 \text{ GeV}/\text{cm}^3$, $r_0 = 20 \text{ kpc}$. The two vertical lines correspond $r = 2R_s$ and $r = 4R_s$, below which the density vanishes for relativity and Newtonian DM respectively due to annihilation or dropping into the BH.

TABLE I: The parameters for DM density spike profiles. We choose the same values as [22] except the mass of BHs and we also calculate the realistic range of ρ_R . And we use “rel.” to denote the models with relativistic modification.

	Hernquist (Newton)	Hernquist (rel.)	NFW (Newton)	NFW (rel.)
η	2	1	2	1
α	0.335	0.335	0.331	0.331
β	-1.67	-1.67	-1.66	-1.66
γ	0.31	0.31	0.32	0.32
δ	-0.025	-0.025	-0.000282	-0.000282
$A(10^{-43} M_\odot^{-2})$	4.87	7.90	1.60	6.42
w	2.22	1.83	2.18	1.82
q	1.93	1.90	1.98	1.91
ρ_0 (GeV/cm ³)	0.1 ~ 0.5	0.1 ~ 0.5	0.1 ~ 0.5	0.1 ~ 0.5
M (M_\odot)	$10^5 \sim 10^9$	$10^5 \sim 10^9$	$10^5 \sim 10^9$	$10^5 \sim 10^9$
r_0 (kpc)	0.005 ~ 50	0.005 ~ 50	0.005 ~ 50	0.005 ~ 50
$\rho_R M^2$	1.14×10^{-32} $\sim 7.08 \times 10^{-30}$	1.85×10^{-32} $\sim 1.15 \times 10^{-29}$	3.57×10^{-33} $\sim 2.66 \times 10^{-30}$	1.43×10^{-32} $\sim 1.07 \times 10^{-29}$

where Γ is the Gamma function. Besides choosing suitable integration constant to satisfy $f(4\eta M) = 1 - \frac{2M}{r}$, we use two tricks to simplify the integration process: (1) Noticing that $\rho_R M^2 \ll 1$ as shown in Table I, we apply the Taylor expansion for the function about ρ_R and re-

tain only the constant and first-order terms of ρ_R . (2) Considering the integral region $r \geq 4\eta M$, we expand the function about r to spatial infinity and ignore the higher order terms than $(1/r)^{N_{\max}}$.

It is also important to note that Eq. (4), Eq. (9) and

Eq. (10) only describe the scenarios in the spike region with $r \in (4\eta M, \infty)$. When $r \in (2M, 4\eta M)$ and $r \rightarrow \infty$ or we have $\rho_R = 0$, the density vanishes with $\rho(r) = 0$ and the spacetime returns to Schwarzschild BHs with $f(r) = g(r) = 1 - \frac{2M}{r}$. Table II contains the different scenarios in both regions.

We then discuss the rationality of neglecting the pressure $p(r)$ in the spike region. Following the work of [46], Eq. (8) needs to be considered. By substituting Eq. (4) and Eq. (9), the pressure $p(r)$ can be numerically solved as shown in Fig. 2. In this way, we find it valid for $p(r) \ll \rho(r)$ and $8\pi r^2 p(r) \ll 1 - g(r)$ in the spike region.

III. GRAVITATIONAL PERTURBATIONS OF SCHWARZSCHILD-LIKE BLACK HOLES

In this section, we consider the linear gravitational perturbation for the modified Schwarzschild-like background metric given by Eq. (2). We shall deal with the axial perturbation in Sec. III A and polar case in Sec. III B. The master equations in different regions are contained in Table II and the numerical method for the equations will be discussed in Sec. III C.

In general, the perturbation can be caused by the injection of gravitational waves or the fall of a particle into the BHs [51, 52], in addition to describing the remnants of a binary BH merger event. Such perturbed metric can be described by:

$$g_{\mu\nu} = \dot{g}_{\mu\nu} + h_{\mu\nu} \quad (11)$$

where $\dot{g}_{\mu\nu}$ is the metric of background spacetime given by Eq. (2) and Table II, while $h_{\mu\nu}$ is the linear perturbation term. And higher-order perturbation terms are neglected.

Because of the static spherical symmetry of the background spacetime $\mathcal{M}^4(t, r, \theta, \phi)$, it can be regarded as the product of a Lorentzian 2-dimension manifold $M^2(t, r)$ and a 2-dimension unit sphere surface manifold $S^2(\theta, \phi)$. By taking advantage of this, the metric perturbation $h_{\mu\nu}$ can be decomposed into multipoles known as axial (odd) parity and polar (even) parity as [53, 54]:

$$h_{\mu\nu} = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{m=\ell} \left[(h_{\mu\nu}^{\ell m})^{(\text{axial})} + (h_{\mu\nu}^{\ell m})^{(\text{polar})} \right] \quad (12)$$

where, ℓ and m are the integers from the separation of θ and ϕ respectively. By adopting the Regge-Wheeler (RW) gauge, the axial perturbations can be parameterized as [31, 55]:

$$(h_{\mu\nu}^{\ell m})^{(\text{axial})} = \left(\sin\theta \frac{\partial Y_{\ell 0}(\theta)}{\partial\theta} \right) e^{i\omega t} \epsilon. \quad (13)$$

$$\begin{pmatrix} 0 & 0 & 0 & h_0(r) \\ 0 & 0 & 0 & h_1(r) \\ 0 & 0 & 0 & 0 \\ h_0(r) & h_1(r) & 0 & 0 \end{pmatrix}$$

TABLE II: Summary of the functions or equations in different regions. $\rho(r)$ is the DM profile, $f(r)$ and $g(r)$ are modified gauge functions in Eq.(2) and r_* is the tortoise coordinate.

	$r \in (2M, 4\eta M)$	$r \in (4\eta M, \infty)$
$\rho(r)$	0	Eq. (4)
$f(r)$	$1 - \frac{2M}{r}$	Eq. (10)
$g(r)$	$1 - \frac{2M}{r}$	Eq. (9)
r_*	Eq. (19)	Eq. (16)
Axial equations	Eq. (18)	Eq. (15)
Polar equations	Eq. (34)	Eq. (25)

and that of the polar perturbations is:

$$(h_{\mu\nu}^{\ell m})^{(\text{polar})} = Y_{\ell 0}(\theta) e^{i\omega t} \epsilon. \quad (14)$$

$$\begin{pmatrix} H_0(r)(1 - \frac{2M}{r}) & H_1(r) & 0 & 0 \\ H_1(r) & \frac{H_2(r)}{1 - \frac{2M}{r}} & 0 & 0 \\ 0 & 0 & r^2 K(r) & 0 \\ 0 & 0 & 0 & r^2 K(r) \sin^2\theta \end{pmatrix}$$

where $|\epsilon| \ll 1$ is the real number to track the order of perturbation proposed by [43]. ω is the eigen frequency from the separation of t known as the quasi-normal modes (QNMs) of the BHs. And $Y_{\ell m}$ stands for the spherical harmonics with $m = 0$ for the spherical symmetry case.

A. QNM Equations for Axial Perturbation

Based on [43, 56], we substitute Eq. (13) into the Einstein equations $G_{\mu\nu} = 8\pi T_{\mu\nu}$ and neglect the perturbation of DM. In the DM spike region, the QNM equations of the axial gravitational perturbation in the frequency domain can be obtained:

$$\left[\frac{\partial^2}{\partial r_*^2} + \omega^2 - V_{\text{axial}}(r) \right] \Psi(r) = 0 \quad (15)$$

where r_* is the tortoise coordinate defined by:

$$dr_* = \frac{dr}{\sqrt{f(r)g(r)}} \quad (16)$$

and the effective potential is:

$$V_{\text{axial}}(r) = \frac{r f'(r) g'(r) + g(r) [f'(r) + 2r f''(r)]}{2r} - \frac{g(r) f'(r)^2}{2f(r)} + \frac{f(r) [r g'(r) + 4g(r) + 2(\ell^2 + \ell - 2)]}{2r^2} \quad (17)$$

where $g(r)$ and $f(r)$ are given by Eq. (9) and Eq. (10) respectively. When $r \leq 4\eta M$ or $\rho R = 0$, it returns to




FIG. 2: Self consistency check of the assumption that the pressure $p(r)$ is ignorable in the spike region. The parameters $M = 10^6 M_\odot$, $\rho_0 = 0.3 \text{ GeV/cm}^3$, $r_0 = 20 \text{ kpc}$ are used here to numerically solve Eq. (8). The assumption that $p(r) \ll \rho(r)$ and $8\pi r^2 p(r) \ll 1 - g(r)$ is valid in this region as shown in the figure. The two vertical lines correspond $r = 2R_s$ and $r = 4R_s$, blow which the density vanishes for relativity and Newtonian DM profile respectively.

Schwarzschild case (see Table II) and the QNM equations become:

$$\left\{ \frac{\partial^2}{\partial r_*^2} + \omega^2 - \left(1 - \frac{2M}{r}\right) \left[\frac{\ell(\ell+1)}{r^2} - \frac{6M}{r^3} \right] \right\} \Psi = 0 \quad (18)$$

with r_* given by:

$$dr_* = \left(1 - \frac{2M}{r}\right)^{-1} dr \quad (19)$$

which is known as RW equation [57].

B. QNM Equations for Polar Perturbation

Following [56] and like the discussion in Sec. III A, we neglect the perturbation of DM and substitute Eq. (14) into the Einstein equations. The QNM equations of the polar gravitational perturbation in the frequency domain can be got as:

$$\left[\zeta_2 \frac{\partial^2}{\partial r^2} + \zeta_1 \frac{\partial}{\partial r} + \zeta_0 \right] Z^{\text{II}}(r) = 0 \quad (20)$$

with

$$\zeta_2 \equiv -\frac{N^{\text{II}} (\eta^{\text{II}})^2 \sigma^{\text{II}} (\tau^{\text{II}})^2}{e^{2(\Lambda-\Phi)} N_T^{\text{II}}} \quad (21)$$

$$\zeta_1 \equiv \omega^2 \frac{4r^2 \eta^{\text{II}} \sigma^{\text{II}} \tau^{\text{II}} N^{\text{II}} (\Lambda' + \Phi')}{N_T^{\text{II}}} + \frac{N_R^{\text{II}} (\eta^{\text{II}})^2 \tau^{\text{II}}}{2r e^{2(\Lambda-\Phi)} N_T^{\text{II}}} \quad (22)$$

$$\zeta_0 \equiv \omega^2 + \frac{N_Z^{\text{II}} (\eta^{\text{II}})^2}{2r^2 e^{2(\Lambda-\Phi)} N_T^{\text{II}}} \quad (23)$$

where N^{II} , N_T^{II} , N_R^{II} , N_Z^{II} , σ^{II} , η^{II} and τ^{II} are the functions of Λ , Φ , r as well as ℓ are defined in [56]. In our case,

we have $\Lambda = -\frac{1}{2} \ln(g)$ and $\Phi = \frac{1}{2} \ln(f)$ with $f(r)$ and $g(r)$ given by Eq. (10) and Eq. (9) respectively. Eq. (20) is the same as [42] when $\Lambda = -\Phi$. In order to ensure that the perturbation equation returns to the well-known Zerilli equation when $f = g = 1 - \frac{2M}{r}$, we introduce the following variable substitution

$$\Psi \equiv \frac{(r - 2M)^2}{6M + r(\ell + 2)(\ell - 1)} Z^{\text{II}}. \quad (24)$$

Eq. (20) now reads:

$$\left[\eta_2 \frac{\partial^2}{\partial r^2} + \eta_1 \frac{\partial}{\partial r} + \eta_0 \right] \Psi(r) = 0 \quad (25)$$

with

$$\eta_0 \equiv \left(1 - \frac{2M}{r}\right)^2 \quad (26)$$

$$\eta_1 \equiv \frac{2M - r}{2} (\omega^2 \xi_1 + \xi_2) \quad (27)$$

$$\eta_0 \equiv -\frac{\omega^2 \xi_3 + \xi_4}{2r^3 N^{\text{II}} \sigma^{\text{II}} \tau^{\text{II}} [(\ell^2 + \ell - 2)r + 6M]} \quad (28)$$

and

$$\xi_1 = \frac{4(2M - r)(fg' - gf')}{f^2 g^2 \eta^{\text{II}} \tau^{\text{II}}} \quad (29)$$

$$\xi_2 = \frac{8(\ell^2 + \ell + 4)M + 4(\ell^2 + \ell - 2)r}{r^2 [(\ell^2 + \ell - 2)r + 6M]} \quad (30)$$

$$+ \frac{(r - 2M)N_R^{\text{II}}}{r^3 N^{\text{II}} \sigma^{\text{II}} \tau^{\text{II}}} \quad (31)$$

$$\xi_3 = 2r(r - 2M)^2 [(\ell^2 + \ell - 2)r + 6M] \frac{N_T^{\text{II}}}{fg(\eta^{\text{II}})^2 \tau^{\text{II}}} + 4r^3(2M - r) [2(\ell^2 + \ell + 4)M + (\ell^2 + \ell - 2)r] \cdot \frac{N^{\text{II}} \sigma^{\text{II}} (gf' - fg')}{f^2 g^2 \eta^{\text{II}}} \quad (32)$$

$$\begin{aligned} \xi_4 = & (2M - r)N_R^{\text{II}} [2(\ell^2 + \ell + 4)M + (\ell^2 + \ell - 2)r] \\ & + \frac{(r - 2M)^2 N_Z^{\text{II}} [(\ell^2 + \ell - 2)r + 6M]}{r\tau^{\text{II}}} \\ & - 4rN^{\text{II}}\sigma^{\text{II}}\tau^{\text{II}} [(\ell^2 + \ell - 2)r + 2(2\ell^2 + 2\ell + 5)M] \end{aligned} \quad (33)$$

when we choose the Schwarzschild case with $f(r) = g(r) = 1 - \frac{2M}{r}$, Eq. (25) returns to:

$$\left[\frac{\partial^2}{\partial r_*^2} + \omega^2 - V_{\text{Zerilli}}(r) \right] \Psi(r) = 0 \quad (34)$$

with the effective potential given by:

$$\begin{aligned} V_{\text{Zerilli}}(r) = & \left(\frac{r - 2M}{r} \right) \cdot \\ & \left[\frac{72M^3 + 36M^2 r \tilde{\lambda} + 6Mr^2 \tilde{\lambda}^2 + r^3 \tilde{\lambda}(\tilde{\lambda} + 2)}{r^3(6M + r\tilde{\lambda})^2} \right] \end{aligned} \quad (35)$$

where $\tilde{\lambda} = (\ell + 2)(\ell - 1)$, r_* is defined by Eq. (19) and Eq. (34) is the well-known Zerilli equation [58]. The QNM equation Eq. (25) seems to become more complicated under the rough substitution Eq. (24). However, it is easier to be solved numerically which can be seen in Sec. III C.

C. Matrix Method for QNMs

So far, the physical problem of perturbation has been written in the master equations as summarized in Table II. Obviously, there is a discontinuity point at $r = 4\eta M$ for $\rho_R \neq 0$ as the axial effective potential is shown in Fig. 3 for example. And several standard methods for QNMs cannot be used directly, like WKB approximation methods [59–61]. We then use the modified matrix method to deal with it [62–65]¹.

We begin with the boundary conditions for QNMs of the BHs. As mentioned before, the equations return to the Schwarzschild case when $r \sim 2M$ or $r \sim \infty$. Thus, boundary conditions for QNMs are $e^{-i\omega r_*}$ and $e^{i\omega r_*}$, denoting the incoming waves near the horizon and outgoing waves at spatial infinity respectively [31, 66]. The tortoise coordinate r_* is defined by Eq. (19) or can be explicitly written as $r_* = r + 2M \ln(r - 2M)$. And the boundary conditions now read:

$$\Psi \rightarrow \begin{cases} e^{-i\omega r} (r - 2M)^{-2iM\omega} & r \rightarrow 2M \\ e^{i\omega r} (r - 2M)^{2iM\omega} & r \rightarrow \infty \end{cases} \quad (36)$$

We then introduce the substitution for $r \in (2M, 4\eta M)$:

$$\Psi(r) \equiv e^{-i\omega r} (r - 2M)^{-2iM\omega} L(r) \quad (37)$$




FIG. 3: Comparison of the axial effective potentials for BH with and without DM. In order to make the difference obvious we use parameter $\rho_R M^2 = 10^{-27}$ which is larger than realistic values as listed in Table. I. When $r < 4\eta M$, the DM vanishes and the spacetime returns to Schwarzschild, causing a discontinuous point at $r = 4\eta M$.

$$y = \frac{r - 2M}{4\eta M - 2M} \quad (38)$$

and for $r \in (4\eta M, \infty)$:

$$\Psi(r) \equiv e^{i\omega r} (r - 2M)^{2iM\omega} R(r) \quad (39)$$

$$z = 1 - \frac{4\eta M}{r} \quad (40)$$

where Eq. (39) and Eq. (37) remove the singularity at the boundaries, while Eq. (38) and Eq. (40) map $r \in (2M, 4\eta M)$ and $r \in (4\eta M, \infty)$ to $[0, 1]$ respectively. The QNMs equations in Table II can be rewritten as:

$$A_2(y)L''(y) + A_1(y)L'(y) + A_0(y)L(y) = 0 \quad (41)$$

$$B_2(z)R''(z) + B_1(z)R'(z) + B_0(z)R(z) = 0 \quad (42)$$

where A_2 , A_1 , A_0 , B_2 , B_1 and B_0 are the coefficient functions about ω , y (or z) and the DM parameters in Table I, depending on the specific equations (axial or polar equations). And the boundary conditions Eq. (36) now become:

$$L(z = 0) = \text{const}, \quad R(z = 1) = \text{const} \quad (43)$$

By further introducing:

$$\tilde{L}(y) \equiv yL(y) \quad (44)$$

$$\tilde{R}(z) \equiv (1 - z)R(z) \quad (45)$$

the boundary conditions can be further simplified as:

$$\tilde{L}(z = 0) = \tilde{R}(z = 1) = 0 \quad (46)$$

¹ K. Lin, “Quasinormal Modes and Echo Effect of Cylindrical Anti-de Sitter Black Hole Spacetime with a Thin Shell” (in print).

The corresponding equations now become:

$$\tilde{A}_2(y)\tilde{L}''(y) + \tilde{A}_1(y)\tilde{L}'(y) + \tilde{A}_0(y)\tilde{L}(y) = 0 \quad (47)$$

$$\tilde{B}_2(z)\tilde{R}''(z) + \tilde{B}_1(z)\tilde{R}'(z) + \tilde{B}_0(z)\tilde{R}(z) = 0 \quad (48)$$

with

$$\begin{aligned} \tilde{A}_0(y) &= y^2 A_0(y) - y A_1(y) + 2A_2(y) \\ \tilde{A}_1(y) &= y[y A_1(y) - 2A_2(y)] \\ \tilde{A}_2(y) &= y^2 A_2(y) \\ \tilde{B}_0(z) &= (z-1)^2 B_0(z) - (z-1) B_1(z) + 2B_2(z) \\ \tilde{B}_1(z) &= (z-1)[(z-1) B_1(z) - 2B_2(z)] \\ \tilde{B}_2(z) &= (z-1)^2 B_2(z) \end{aligned} \quad (49)$$

Then, we deal with the discontinuity at $r = r_c = 4\eta M$. Such discontinuity provides the limitation governed by Israel's junction condition [67] and the wave functions on both sides are related to vanishing Wronskian [64, 65, 68]:

$$\Psi'(r = r_c^-)\Psi(r = r_c^+) - \Psi(r = r_c^-)\Psi'(r = r_c^+) = 0 \quad (50)$$

where $r = r_c^-$ and $r = r_c^+$ denote the asymptotic values from left or right to the discontinuity point respectively. We then define the ratio coefficient κ as:

$$\kappa = \frac{\Psi'(r = r_c^-)}{\Psi(r = r_c^-)} = \frac{\Psi'(r = r_c^+)}{\Psi(r = r_c^+)} \quad (51)$$

By substituting Eq. (37)-Eq. (40), it can be rewritten as:

$$\begin{aligned} 0 &= \tilde{L}'(y) \\ &+ y\{2(1-2\eta)\kappa My - 2i\omega[(2\eta-1)My + M] - 1\}\tilde{L}(y) \end{aligned} \quad (52)$$

$$\begin{aligned} 0 &= (z-1)^2(2\eta+z-1)\tilde{R}'(z) \\ &+ [-(2\eta+z-1)(4\eta\kappa M + z-1) + 8i\eta^2 M\omega]\tilde{R}(z) \end{aligned} \quad (53)$$

And Eq. (52) and Eq. (53) are the connection conditions.

Then, the matrix method algorithm for our case is as follows:

- (1) Following [62], any coordinate $x \in [0, 1]$ can be discretized into N points with x_1 to x_N . With the application of Taylor expansion, the value of a function with its 1st order derivatives to its N th order derivatives at each point can be written in the $N \times N$ matrices respectively (details in [62]).
- (2) By substituting the matrices of the function, 1st order derivatives and 2nd order derivatives from (1), Eq. (47) and Eq. (48) can be rewritten as two matrix equations $\overline{\mathcal{M}}_L \mathcal{L} = \overline{\mathcal{M}}_R \mathcal{R} = 0$ respectively. Here, $\overline{\mathcal{M}}_L$ and $\overline{\mathcal{M}}_R$ are the $N_L \times N_L$ and $N_R \times N_R$ matrices only about ω respectively, while $\mathcal{L} = (L(x_1), \dots, L(x_{N_L}))^T$ and $\mathcal{R} = (R(x_1), \dots, R(x_{N_R}))^T$ are the values of functions at each points.

- (3) Then, we use Eq. (46) to replace the 1st line of $\overline{\mathcal{M}}_L$ and N th line of $\overline{\mathcal{M}}_R$, while we use Eq.(52) and Eq. (53) to replace the N th line of $\overline{\mathcal{M}}_L$ and 1st line of $\overline{\mathcal{M}}_R$ respectively. We now obtain $\mathcal{M}_L \mathcal{L} = \mathcal{M}_R \mathcal{R} = 0$, with \mathcal{M}_L and \mathcal{M}_R the modified matrices about ω and κ .

- (4) By using the equations $\det(\mathcal{M}_L) = \det(\mathcal{M}_R) = 0$, the QNMs ω can be numerically solved with the corresponding ratio coefficients κ .

IV. RESULTS

In this section, we shall discuss the numerical results of QNMs and their characteristics affected by the DM spike. With Eq. (5), we have unified the parameters of the DM models r_0 and ρ_0 and the parameter of the BHs M into a single dimensionless parameter $\rho_R M^2$. By analyzing the parameter ρ_R , we can cover the analysis of the parameters r_0 , ρ_0 and M . We then consider the QNMs of the Sgr A* BH at the center of Milky Way in Sec. IV A and explore the QNMs vary with ρ_R to simulate the exploration of various BHs and galaxies in the universe in Sec. IV B. And the detectability of the deviation effected by DM spike will be discussed in Sec. IV C.

Due to the limited computational capacity, it is impossible to use arbitrarily large matrices and unlimited precision in our numerical algorithm. Based on the studies of the Mashhoon method and WKB approximate method for solving QNMs, the peak of the effective potential plays a significant role in QNMs [59, 69]. And since the peak of the effective potential falls exactly in $r \leq 4\eta M$ as shown in Fig. 3, in order to speed up the calculation, we choose $N_L = 24$, $N_R = 12$ and $N_{\max} = 20$.

In order to understand the computational errors that result from this choice, we first calculate the differences between the results of our algorithm with $\rho_R = 0$ and the standard Schwarzschild values as listed in Table III. And the QNMs of the Schwarzschild case from 61 order matrix method are used as standard values, which have been proven to be accurate within 12 significant digits [64]. Notice that the choice of $\rho_R = 0$ returns the equations in Table II back to the Schwarzschild case with corresponding QNMs, the differences in Table II imply that our results can be credible within 6 significant digits.

A. The QNMs of the BH in Milky Way

We certainly want to locate a suitable observational source and target it with our GW detectors, in order to detect the ringing of a BH when it is perturbed. Due to the unsatisfactory detection of ringdown signals with ground-based GW detectors [36], we look to space-based GW detectors in the future.

Based on [40], the corresponding frequency of the ringdown signal from the SMBHs with $M \sim 10^6 M_\odot$ is

TABLE III: The differences between the results of our algorithm with $\rho_R = 0$ and the standard Schwarzschild values validate our numerical algorithm. And we use ω_s , ω_a , and ω_p to denote the standard Schwarzschild values, the results of the axial case from our algorithm, and those of the polar case respectively.

ℓ	n	$2M(\omega_a - \omega_s)$	$2M(\omega_p - \omega_s)$	$2M(\omega_a - \omega_p)$
2	0	$-3.9 \times 10^{-9} + 1.2 \times 10^{-8}i$	$3.2 \times 10^{-10} + 7.9 \times 10^{-9}i$	$-4.2 \times 10^{-9} + 3.6 \times 10^{-9}i$
	1	$-1.6 \times 10^{-7} + 5.6 \times 10^{-7}i$	$-1.4 \times 10^{-7} + 5.2 \times 10^{-7}i$	$-1.7 \times 10^{-8} + 3.9 \times 10^{-8}i$
3	0	$-8.3 \times 10^{-11} + 1.0 \times 10^{-9}i$	$-2.7 \times 10^{-10} + 7.2 \times 10^{-10}i$	$1.8 \times 10^{-10} + 2.8 \times 10^{-10}i$
	1	$-4.0 \times 10^{-8} + 7.2 \times 10^{-8}i$	$-4.2 \times 10^{-8} + 7.3 \times 10^{-8}i$	$2.1 \times 10^{-9} - 4.3 \times 10^{-10}i$

$\sim 10^{-2}$ Hz, which falls exactly in the sensitive frequency bands of LISA. And the Sgr A* BH at the center of the Milky Way galaxy with $M = 4.3 \times 10^6 M_\odot$ can be such a good source. The best-fit values for the parameters of the NFW model are $\rho_0 = 0.51 \text{GeV/cm}^3$ and $r_0 = 8.1 \text{kpc}$, corresponding to $\rho_R M^2 = 1.27 \times 10^{-32}$ for NFW (Newton) and $\rho_R M^2 = 5.09 \times 10^{-32}$ for relativistic DM profile respectively [23]. We then calculate the QNMs of the Sgr A* BH in DM spike and list them in Table IV for the axial case and Table V for the polar case.

As shown in Table IV and Table V, the effects of the DM spike on the QNMs of the Sgr A* BH are ignorably small. The deviations of the fundamental mode from the one without DM are about $\sim 10^{-9}$ Hz and $\sim 0.001s$. The detection of such deviations is not possible in the near future.

B. The Fundamental QNM with Varying ρ_R

We then explore a wider range of observational sources with varying parameters. For given parameters r_0 , ρ_0 and M , the parameter ρ_R is uniquely determined by Eq. (5). Through the exploration of the trend of QNMs with varying ρ_R , the corresponding trend for parameters r_0 , ρ_0 and M can also be obtained. Specifically, the dimensionless parameter $\rho_R M^2$ is closely related to r_0 , ρ_0 and M . This exploration is interesting for three reasons [42, 43]:

- (1) The DM parameters r_0 and ρ_0 are generally derived by fitting the corresponding density profiles with the data of rotation curves in various galaxies [23]. They can accurately reflect the distribution of DM throughout the entire galaxy, but are essentially effected by BHs and basically free near BHs.
- (2) Around a BH, the baryonic component also makes significant contributions to the DM parameters r_0 and ρ_0 .
- (3) The parameters r_0 and ρ_0 change from galaxy to galaxy, while M changes along with BHs. A suitable BH in the DM spike is easier for detection.

As shown in Fig. 4 and Fig. 5, the QNMs of axial and polar cases differ to a certain degree and such difference increases along with $\rho_R M^2$. More specifically, the isospectrality of BHs surrounded by DM spike breaks down [28, 31].

With relativistic modification, both parities' QNMs show a consistent trend. The real parts of the QNMs increase while the absolute values of imaginary parts decrease with the increasing $\rho_R M^2$.

Without relativistic modification, the absolute values of imaginary parts increase as $\rho_R M^2$ increases. However, the trends in the real parts of QNMs for both parities are different. Generally speaking, as $\rho_R M^2$ increases, the real parts for the axial case decrease, and those for the polar case increase. The only exception is the real part of the polar case of the NFW model, where it decreases as $\rho_R M^2$ increases and then increases for $M^2 \rho \gtrsim 4 \times 10^{-30}$. Such a scenario with inflection points for real parts is similar to the trend of QNMs as the Kerr BHs' rotation parameter a varies [70]. These trends can be seen in Fig. 4 and Fig. 5.

C. Hints of detectability by space-based detectors

With the previous exploration, we have found that the DM spike can result in deviations of QNMs from that of BH without DM. We then explore the possibility of detection of such deviations by future space-based detectors. The GW waveform during the ringdown process can be written as [40]:

$$h_+ + ih_\times = \frac{M_z}{D_L} \sum_{\ell mn} \mathcal{A}_{\ell mn} e^{i(f_{\ell mn} t + \phi_{\ell mn})} e^{-t/\tau_{\ell mn}} S_{\ell mn} \quad (54)$$

where M_z is the red-shifted BH mass, D_L is the luminosity distance to the source, $\mathcal{A}_{\ell mn}$ is the amplitude of the corresponding QNM, $\phi_{\ell mn}$ is the phase coefficient and S_ℓ is the 2-spin-weighted spheroidal harmonics depending on the polar and azimuthal angles. And the real waveform is the superposition of the axial and polar parity components. The two parameters associated to QNMs are the GW frequency $f_{\ell mn}$ and the damping time $\tau_{\ell mn}$ defined as:

$$2\pi f_{\ell mn} = \text{Re}(\omega_{\ell mn}), \quad (55)$$




FIG. 4: The behavior of the QNMs with respect to the dimensionless parameter $\rho_R M^2$ for Hernquist models of DM profile. In figures, “Re” in the left column denotes the real part, and “Im” in the right column denotes the imaginary part. The marked points correspond to $\rho_R M^2 = 10^{-35}, 10^{-34}, \dots, 10^{-29}$.




FIG. 5: The behavior of the QNMs with respect to the dimensionless parameter $\rho_R M^2$ for NFW models. The notations here are the same as those of Fig. 4.

TABLE IV: We list the axial QNMs $2M\omega$ of the Sgr A* BH at the center of Milky Way galaxy surrounded by the NFW DM spike.

ℓ	n	Newton	Rel.	Schwarzschild
2	0	$0.747342 - 0.177926i$	$0.747343 - 0.177924i$	$0.747343 - 0.177925i$
	1	$0.693541 - 0.547762i$	$0.693422 - 0.547827i$	$0.693422 - 0.547829i$
3	0	$1.198890 - 0.185406i$	$1.198886 - 0.185406i$	$1.198887 - 0.185406i$
	1	$1.165002 - 0.562909i$	$1.165286 - 0.562594i$	$1.165288 - 0.562596i$

TABLE V: We list the polar QNMs $2M\omega$ of the Sgr A* BH at the center of Milky Way galaxy surrounded by the NFW DM spike.

ℓ	n	Newton	Rel.	Schwarzschild
2	0	$0.747343 - 0.177927i$	$0.747343 - 0.177924i$	$0.747343 - 0.177925i$
	1	$0.693526 - 0.547677i$	$0.693422 - 0.547828i$	$0.693422 - 0.547829i$
3	0	$1.198891 - 0.185405i$	$1.198886 - 0.185406i$	$1.198887 - 0.185406i$
	1	$1.165035 - 0.562981i$	$1.165286 - 0.562594i$	$1.165288 - 0.562596i$

$$\tau_{\ell mn} = -\frac{1}{\text{Im}(\omega_{\ell mn})}, \quad (56)$$

where $\omega_{\ell mn}$ is the QNMs for given (ℓ, m, n) . Here we consider only the fundamental mode with $(\ell, m, n) = (2, 0, 0)$ because it decays slowest. Following [42], the frequency and the damping time can be expanded as:

$$f_{\ell mn} = f_{\ell mn}^{\text{Sch}} (1 + \delta f_{\ell mn}), \quad (57)$$

$$\tau_{\ell mn} = \tau_{\ell mn}^{\text{Sch}} (1 + \delta \tau_{\ell mn}), \quad (58)$$

where $f_{\ell mn}^{\text{Sch}}$ and $\tau_{\ell mn}^{\text{Sch}}$ are the QNM frequency and damping time for Schwarzschild case, while $\delta f_{\ell mn}$ and $\delta \tau_{\ell mn}$ are the corresponding relative deviations. In addition, the possibility of detecting the isospectrality violation is also studied. We investigate the isospectrality violation through

$$\delta f_{\ell mn}^{\text{Iso}} = \frac{f_{\ell mn}^{\text{Polar}} - f_{\ell mn}^{\text{Axial}}}{f_{\ell mn}^{\text{Sch}}}, \quad (59)$$

$$\delta \tau_{\ell mn}^{\text{Iso}} = \frac{\tau_{\ell mn}^{\text{Polar}} - \tau_{\ell mn}^{\text{Axial}}}{\tau_{\ell mn}^{\text{Sch}}}. \quad (60)$$

From Fig.6, we can see that the deviations of axial, polar, and isospectrality violation increase along with the increasing $\rho_R M^2$. In general, the deviation for the Hernquist DM profile is larger than that of NFW profile. The deviation for the Newtonian profile is larger than that of the relativistic profile. Corresponding to the DM profiles listed in Table. I the largest deviation of $\delta f_{\ell mn}$ corresponding to $\rho_R M^2 \sim 10^{-28}$ and with Hernquist DM profile may be as high as $\sim 10^{-2}$ without relativistic modification and about $\sim 10^{-4}$ with relativistic modification. In the meantime, $\delta \tau_{\ell mn}$ can reach 0.1 and 0.001 respectively.

V. CONCLUSION

In this paper, we explore the behavior of the GWs waveform emitted by the perturbed Schwarzschild-like BHs in the DM spike. Accounting for the profile of the DM spike we use the TOV equation to derive the modified metrics. After that, we derive the QNMs equations for both parities under the modified metrics and use modified matrix methods to calculate the QNMs. Finally, we discuss the impacts of the DM spike and analyze the detectability.

The key points of the exploration are as follows:

- (1) We unified the parameters r_0 , ρ_0 and M into a single dimensionless parameter $\rho_R M^2$ by Eq. (5). And the analysis of $\rho_R M^2$ can cover all three parameters.
- (2) We neglect the pressure $p(r)$ in the DM spike and show the rationality in Fig. 2.
- (3) The DM spike profile vanishes for $r \leq 4\eta M$ which divides the entire domain into two different regions as shown in Fig. 1. And the behaviors of the spacetime and QNMs equations differ in both regions as shown in Table II.
- (4) Such differences in both regions lead to the discontinuity of QNMs equations at $4\eta M$ as shown in Fig. 3. And the modified matrix method for such a scenario is used.
- (5) We choose $N_L = 24$, $N_R = 12$ and $N_{\text{max}} = 20$, and the numerical results are as accurate as 6 significant digits as shown in Table III.
- (6) The QNMs of the Sgr A* BH at the center of the Milky Way galaxy in the NFW DM spike are calculated and listed in Table IV and Table V.




FIG. 6: The behavior of the deviations with respect to the dimensionless parameter $\rho_R M^2$ for different models. The points in each plot from the left to the right correspond to $\rho_R M^2 = 10^{-32}, 10^{-31}, \dots, 10^{-28}$ respectively.

- (7) We explore the trend of fundamental QNM with varying ρ_R to simulate the varying BHs and galaxies. The isospectrality breaks down. The trends with relativistic modification for all the cases are monotonic: the real parts increase and the absolute values of the imaginary parts decrease as $\rho_R M^2$ increases. The trends without relativistic modification are not monotonic. And details are shown in Fig. 4 and Fig. 5.
- (8) The detectability of the deviations effected by the DM spike is explored. For the DM profiles listed in Table. I, the relative deviation of the QNM frequency can be as high as $\sim 10^{-2}$.

In summary, the presence of the DM spike does impact the ringing GW waveform of BHs. Only spinless BH is investigated in the current paper. The effect of the DM

spike for spinning BH may be even stronger because the spin of a BH can enhance the spike effect [18] and enlarge the decay time of QNMs. And this will be the focus of our next exploration.

ACKNOWLEDGMENTS

The authors would like to thank Dr. Zhan-Feng Mai and Prof. Run-Qiu Yang for discussing and providing useful ideas to calculate QNMs in this work. The authors thank Wentao Liu and Xiongjun Fang for providing the *Mathematica* code of the results of the polar QNMs equations in their work [56].

This work was supported in part by the National Key Research and Development Program of China Grant No. 2021YFC2203001 and in part by the NSFC (No. 11920101003, No. 12021003 and No. 12005016). Z. Cao was supported by “the Interdiscipline Research Funds of Beijing Normal University” and CAS Project for Young Scientists in Basic Research YSBR-006.

[1] A. Einstein, Die feldgleichungen der gravitation, Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften (Berlin) , 844 (1915).

[2] A. Askar, C. Belczynski, G. Bertone, E. Bon, D. Blas, R. Brito, T. Bulik, C. Burrage, C. T. Byrnes, C. Caprini, *et al.*, Black holes, gravitational waves and fundamental

- physics: a roadmap, *Classical and Quantum Gravity* **36**, 143001 (2019).
- [3] B. P. Abbott, R. Abbott, T. D. Abbott, M. R. Abernathy, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. X. Adhikari, *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), Observation of gravitational waves from a binary black hole merger, *Phys. Rev. Lett.* **116**, 061102 (2016).
- [4] Gravitational wave open science center, <https://www.gw-openscience.org/eventapi/html/allevents/> (2022).
- [5] B. P. Abbott, R. Abbott, T. D. Abbott, S. Abraham, F. Acernese, K. Ackley, C. Adams, R. X. Adhikari, V. B. Adya, C. Affeldt, *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), Gwtc-1: A gravitational-wave transient catalog of compact binary mergers observed by ligo and virgo during the first and second observing runs, *Phys. Rev. X* **9**, 031040 (2019).
- [6] R. Abbott, T. D. Abbott, S. Abraham, F. Acernese, K. Ackley, A. Adams, C. Adams, R. X. Adhikari, V. B. Adya, C. Affeldt, *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), Gwtc-2: Compact binary coalescences observed by ligo and virgo during the first half of the third observing run, *Phys. Rev. X* **11**, 021053 (2021).
- [7] R. Abbott, T. Abbott, F. Acernese, K. Ackley, C. Adams, N. Adhikari, R. Adhikari, V. Adya, C. Affeldt, D. Agarwal, *et al.*, Gwtc-2.1: Deep extended catalog of compact binary coalescences observed by ligo and virgo during the first half of the third observing run, arXiv preprint arXiv:2108.01045 10.48550/ARXIV.2108.01045 (2021).
- [8] R. Abbott, T. Abbott, F. Acernese, K. Ackley, C. Adams, N. Adhikari, R. Adhikari, V. Adya, C. Affeldt, D. Agarwal, *et al.*, Gwtc-3: compact binary coalescences observed by ligo and virgo during the second part of the third observing run, arXiv preprint arXiv:2111.03606 10.48550/ARXIV.2111.03606 (2021).
- [9] T. E. H. T. Collaboration, K. Akiyama, A. Alberdi, W. Alef, K. Asada, R. Azulay, A.-K. Baczkó, D. Ball, M. Baloković, J. Barrett, D. Bintley, *et al.*, First m87 event horizon telescope results. i. the shadow of the supermassive black hole, *The Astrophysical Journal Letters* **875**, L1 (2019).
- [10] T. E. H. T. Collaboration, K. Akiyama, A. Alberdi, W. Alef, K. Asada, R. Azulay, A.-K. Baczkó, D. Ball, M. Baloković, J. Barrett, D. Bintley, *et al.*, First m87 event horizon telescope results. ii. array and instrumentation, *The Astrophysical Journal Letters* **875**, L2 (2019).
- [11] T. E. H. T. Collaboration, K. Akiyama, A. Alberdi, W. Alef, K. Asada, R. Azulay, A.-K. Baczkó, D. Ball, M. Baloković, J. Barrett, D. Bintley, *et al.*, First m87 event horizon telescope results. iii. data processing and calibration, *The Astrophysical Journal Letters* **875**, L3 (2019).
- [12] T. E. H. T. Collaboration, K. Akiyama, A. Alberdi, W. Alef, K. Asada, R. Azulay, A.-K. Baczkó, D. Ball, M. Baloković, J. Barrett, D. Bintley, *et al.*, First m87 event horizon telescope results. iv. imaging the central supermassive black hole, *The Astrophysical Journal Letters* **875**, L4 (2019).
- [13] T. E. H. T. Collaboration, K. Akiyama, A. Alberdi, W. Alef, K. Asada, R. Azulay, A.-K. Baczkó, D. Ball, M. Baloković, J. Barrett, D. Bintley, *et al.*, First m87 event horizon telescope results. v. physical origin of the asymmetric ring, *The Astrophysical Journal Letters* **875**, L5 (2019).
- [14] T. E. H. T. Collaboration, K. Akiyama, A. Alberdi, W. Alef, K. Asada, R. Azulay, A.-K. Baczkó, D. Ball, M. Baloković, J. Barrett, D. Bintley, *et al.*, First m87 event horizon telescope results. vi. the shadow and mass of the central black hole, *The Astrophysical Journal Letters* **875**, L6 (2019).
- [15] T. E. H. T. Collaboration, K. Akiyama, A. Alberdi, W. Alef, K. Asada, R. Azulay, A.-K. Baczkó, D. Ball, M. Baloković, J. Barrett, D. Bintley, *et al.*, First m87 event horizon telescope results. vii. polarization of the ring, *The Astrophysical Journal Letters* **910**, L12 (2021).
- [16] T. E. H. T. Collaboration, K. Akiyama, A. Alberdi, W. Alef, K. Asada, R. Azulay, A.-K. Baczkó, D. Ball, M. Baloković, J. Barrett, D. Bintley, *et al.*, First m87 event horizon telescope results. viii. magnetic field structure near the event horizon, *The Astrophysical Journal Letters* **910**, L13 (2021).
- [17] K. Schwarzschild, Über das Gravitationsfeld einer Kugel aus inkompressibler Flüssigkeit nach der Einsteinschen Theorie, in *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin* (1916) pp. 424–434.
- [18] F. Ferrer, A. Medeiros da Rosa, and C. M. Will, Dark matter spikes in the vicinity of kerr black holes, *Phys. Rev. D* **96**, 083014 (2017).
- [19] S. Nampalliwar, S. Kumar, K. Jusufi, Q. Wu, M. Jamil, and P. Salucci, Modeling the sgr a* black hole immersed in a dark matter spike, *The Astrophysical Journal* **916**, 116 (2021).
- [20] Z. Xu, J. Wang, and M. Tang, Deformed black hole immersed in dark matter spike, *Journal of Cosmology and Astroparticle Physics* **2021** (09), 007.
- [21] K. Jusufi, M. Jamil, and T. Zhu, Shadows of sgr a* black hole surrounded by superfluid dark matter halo, *The European Physical Journal C* **80**, 354 (2020).
- [22] N. Speeney, A. Antonelli, V. Baibhav, and E. Berti, Impact of relativistic corrections on the detectability of dark-matter spikes with gravitational waves, *Phys. Rev. D* **106**, 044027 (2022).
- [23] H.-N. Lin and X. Li, The dark matter profiles in the Milky Way, *Monthly Notices of the Royal Astronomical Society* **487**, 5679 (2019), https://academic.oup.com/mnras/article-pdf/487/4/5679/28897928/stz1698_supplemental_file.pdf.
- [24] L. Hernquist, An analytical model for spherical galaxies and bulges, *Astrophysical Journal*, Part 1 (ISSN 0004-637X), vol. 356, June 20, 1990, p. 359-364. **356**, 359 (1990).
- [25] J. F. Navarro, C. S. Frenk, and S. D. M. White, A universal density profile from hierarchical clustering, *The Astrophysical Journal* **490**, 493 (1997).
- [26] P. Gondolo and J. Silk, Dark matter annihilation at the galactic center, *Phys. Rev. Lett.* **83**, 1719 (1999).
- [27] L. Sadeghian, F. Ferrer, and C. M. Will, Dark-matter distributions around massive black holes: A general relativistic analysis, *Phys. Rev. D* **88**, 063522 (2013).
- [28] S. Chandrasekhar and K. S. Thorne, The mathematical theory of black holes (1985).
- [29] K. D. Kokkotas and B. G. Schmidt, Quasi-normal modes of stars and black holes, *Living Reviews in Relativity* **2**, 1 (1999).

- [30] H.-P. Nollert, Quasinormal modes: the characteristic ‘sound’ of black holes and neutron stars, *Classical and Quantum Gravity* **16**, R159 (1999).
- [31] E. Berti, V. Cardoso, and A. O. Starinets, Quasinormal modes of black holes and black branes, *Classical and Quantum Gravity* **26**, 163001 (2009).
- [32] R. A. Konoplya and A. Zhidenko, Quasinormal modes of black holes: From astrophysics to string theory, *Rev. Mod. Phys.* **83**, 793 (2011).
- [33] W.-X. Feng, A. Parisi, C.-S. Chen, and F.-L. Lin, Self-interacting dark scalar spikes around black holes via relativistic Bondi accretion, *Journal of Cosmology and Astroparticle Physics* **2022** (8), 032, arXiv:2112.05160 [astro-ph.HE].
- [34] S. L. Shapiro and V. Paschalidis, Self-interacting dark matter cusps around massive black holes, *Phys. Rev. D* **89**, 023506 (2014).
- [35] V. D. Luca and J. Khoury, Superfluid dark matter around black holes (2023), arXiv:2302.10286 [astro-ph.CO].
- [36] R. Cotesta, G. Carullo, E. Berti, and V. Cardoso, Analysis of ringdown overtones in gw150914, *Phys. Rev. Lett.* **129**, 111102 (2022).
- [37] W.-H. Ruan, Z.-K. Guo, R.-G. Cai, and Y.-Z. Zhang, Taiji program: Gravitational-wave sources, *International Journal of Modern Physics A* **35**, 2050075 (2020).
- [38] X.-Y. Lu, Y.-J. Tan, and C.-G. Shao, Sensitivity functions for space-borne gravitational wave detectors, *Phys. Rev. D* **100**, 044042 (2019).
- [39] C. J. Moore, R. H. Cole, and C. P. L. Berry, Gravitational-wave sensitivity curves, *Classical and Quantum Gravity* **32**, 015014 (2014).
- [40] E. Berti, V. Cardoso, and C. M. Will, Gravitational-wave spectroscopy of massive black holes with the space interferometer lisa, *Phys. Rev. D* **73**, 064030 (2006).
- [41] D. Liu, Y. Yang, S. Wu, Y. Xing, Z. Xu, and Z.-W. Long, Ringing of a black hole in a dark matter halo, *Phys. Rev. D* **104**, 104042 (2021).
- [42] C. Zhang, T. Zhu, X. Fang, and A. Wang, Imprints of dark matter on gravitational ringing of supermassive black holes, *Physics of the Dark Universe* **37**, 101078 (2022).
- [43] C. Zhang, T. Zhu, and A. Wang, Gravitational axial perturbations of schwarzschild-like black holes in dark matter halos, *Phys. Rev. D* **104**, 124082 (2021).
- [44] V. Cardoso, K. Destounis, F. Duque, R. P. Macedo, and A. Maselli, Black holes in galaxies: Environmental impact on gravitational-wave generation and propagation, *Phys. Rev. D* **105**, L061501 (2022).
- [45] R. Konoplya, Black holes in galactic centers: Quasinormal ringing, grey-body factors and unruh temperature, *Physics Letters B* **823**, 136734 (2021).
- [46] R. G. Daghigh and G. Kunstatter, Spacetime metrics and ringdown waveforms for galactic black holes surrounded by a dark matter spike, *The Astrophysical Journal* **940**, 33 (2022).
- [47] R. A. Konoplya and A. Zhidenko, Solutions of the einstein equations for a black hole surrounded by a galactic halo, *The Astrophysical Journal* **933**, 166 (2022).
- [48] S. M. Carroll, *Spacetime and geometry* (Cambridge University Press, 2019).
- [49] M. Abramowitz and I. Stegun, *Handbook of mathematical functions* (dover publications inc., new york). (1970).
- [50] F. W. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, *NIST handbook of mathematical functions hardback and CD-ROM* (Cambridge university press, 2010).
- [51] J. A. H. Futterman, F. A. Handler, and R. A. Matzner, *Scattering from black holes* (Cambridge ; New York : Cambridge University Press, 1988).
- [52] M. Davis, R. Ruffini, and J. Tiomno, Pulses of gravitational radiation of a particle falling radially into a schwarzschild black hole, *Phys. Rev. D* **5**, 2932 (1972).
- [53] J. E. Thompson, H. Chen, and B. F. Whiting, Gauge invariant perturbations of the schwarzschild spacetime, *Classical and Quantum Gravity* **34**, 174001 (2017).
- [54] A. Nagar and L. Rezzolla, Gauge-invariant non-spherical metric perturbations of schwarzschild black-hole spacetimes, *Classical and Quantum Gravity* **22**, R167 (2005).
- [55] C. V. Vishveshwara, Stability of the schwarzschild metric, *Phys. Rev. D* **1**, 2870 (1970).
- [56] W. Liu, X. Fang, J. Jing, and A. Wang, Gauge invariant perturbations of general spherically symmetric spacetimes, *Science China Physics, Mechanics & Astronomy* **66**, 210411 (2023).
- [57] T. Regge and J. A. Wheeler, Stability of a schwarzschild singularity, *Phys. Rev.* **108**, 1063 (1957).
- [58] F. J. Zerilli, Effective potential for even-parity regge-wheeler gravitational perturbation equations, *Phys. Rev. Lett.* **24**, 737 (1970).
- [59] B. F. Schutz and C. M. Will, Black hole normal modes: a semianalytic approach, *The Astrophysical Journal* **291**, L33 (1985).
- [60] S. Iyer and C. M. Will, Black-hole normal modes: A wkb approach. i. foundations and application of a higher-order wkb analysis of potential-barrier scattering, *Phys. Rev. D* **35**, 3621 (1987).
- [61] R. A. Konoplya, Quasinormal behavior of the d -dimensional schwarzschild black hole and the higher order wkb approach, *Phys. Rev. D* **68**, 024018 (2003).
- [62] K. Lin and W.-L. Qian, A matrix method for quasinormal modes: Schwarzschild black holes in asymptotically flat and (anti-) de sitter spacetimes, *Classical and Quantum Gravity* **34**, 095004 (2017).
- [63] K. Lin, W.-L. Qian, A. B. Pavan, and E. Abdalla, A matrix method for quasinormal modes: Kerr and kerr–sen black holes, *Modern Physics Letters A* **32**, 1750134 (2017).
- [64] K. Lin and W.-L. Qian, High-order matrix method with delimited expansion domain (2022).
- [65] S.-F. Shen, W.-L. Qian, K. Lin, C.-G. Shao, and Y. Pan, Matrix method for perturbed black hole metric with discontinuity, *Classical and Quantum Gravity* **39**, 225004 (2022).
- [66] Y. Zhao, B. Sun, Z.-F. Mai, and Z. Cao, Quasi normal modes of black holes and detection in ringdown process (2022).
- [67] W. Israel, *Nuovo cim b44s10 1*, Erratum: *ibid Nuovo Cim B* **48**, 463 (1966).
- [68] P. T. Leung, Y. T. Liu, W. M. Suen, C. Y. Tam, and K. Young, Perturbative approach to the quasinormal modes of dirty black holes, *Phys. Rev. D* **59**, 044034 (1999).
- [69] V. Ferrari and B. Mashhoon, Oscillations of a black hole, *Phys. Rev. Lett.* **52**, 1361 (1984).
- [70] E. W. Leaver, An analytic representation for the quasinormal modes of kerr black holes, *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences* **402**, 285 (1985).