On Production of Excited Kaluza-Klein States in Large Radius Compactification Scenario

Jnanadeva Maharana¹

Institute of Physics Bhubaneswar-751005, India NISER, Bhubaneswar, India

Abstract

Production of exotic states at LHC is considered in the large radius compactification scenario. We envisage a five dimensional theory for a scalar field in five dimensional flat spacetime. It is compactified on a circle, S^1 , with radius, R. The radius is assumed to be in TeV scale appealing to LRC hypothesis. The production of Kaluza-Klein states whose masses lie in the vicinity of TeV range is considered. Instead of appealing to any specific model, bounds on inelastic cross sections and near forward differental cross section are derived from the Lehmann-Symanzik-Zimmermann (LSZ) formulation. We consider decompactified theory should compactification radius be large enough to unravel the fifth spacial dimension in LHC energy scale. Bounds on cross sections are also derived for this scenario. We present bounds on inclusive cross sections for reactions like $a+b \rightarrow c+X$, X being unobserved states. We plot the bounds as a function of energy and propose that these bounds might be useful for search of exotic states in LHC experiments like ATLAS and CMS.

¹Indian National Science Academy Honorary Scientist at Institute of Physics, Bhubaneswar. e-mail: maharana@iopb.res.in

1 Introduction

The standard model, $SU(3)_C \otimes SU(2)_L \otimes U(1)$, has been tested to a great degree of precision. However, there are reasons to believe that the standard model is not complete and there are several important issues which cannot be resolved within the frameworks of the model. The construction of grand unified theory (GUT) is an attempt to unify strong, weak and electromagnetic interaction with gauge invariance and spontaneously broken symmetry mechanism as underlying cardinal principles. There are hints from experimental observations that an underlying GUT might be present; however in order to test some of the predictions of GUT we do not have accelerators of such high energies. Moreover, other nonaccelerator based tests like instability of proton have not yielded any definite results so far. Therefore, there are no clinching evidences in favor of GUT paradigm. Furthermore, gravitational interaction is not incorporated in GUT. Therefore, a unified theory of four fundamental interaction must incorporate gravitational interaction. Superstring theory offers the prospect of unifying fundamental forces of Nature. Consequences of quantum theory of gravity are expected to be addressed from string theory perspective. A consistent quantum description of superstring string requires that it must live in ten spacetime dimensions. We have no experimental evidence of ten dimensional spacetime manifold so far. In this context, the ideas of Kaluza [1] and Klein [2] (KK) have been revived. A comprehensive review and historical developments of the idea of theories in higher spacetime dimensions and their influence on developments of diverse proposals in supersymmetry, supergravity theories as well as in string theories are presented in the book edited by Appelquist, Chodos and Freund [3]. This book contains collection of large number of important articles on the subject of higher dimensional theories and various schemes of compactifications. Kaluza and Klein had set out to provide a unified description of electromagnetism and gravitation following Einstein's theory of gravitation whose foundation lies in general theory of relativity. It is worthwhile to note that the idea of describing physics in higher dimensions predates works of Kaluza and Klein [4, 5] which was envisaged before Einstein expounded the general theory of relativity. Their endeavors were to unify Maxwell's theory with four dimensional continuum. Kaluza and Klein considered a theory of gravitation in D = 5described by Einstein-Hilbert action. They compactified one spatial dimension on a circle, S^1 , of radius R and argued that $R \ll 1$ so that the experiments of that era could not probe that length scale. The resulting four dimensional action described Einstein's theory of relativity, with a massless spin two graviton, an Abelian gauge field and a massless scalar field appearing as zero modes. These fields interacted through exchange of massless particles (so called zero modes. important for long range forces) which arise as a consequence of KK compactification. Moreover, the resulting 4-D action also contained tower of massive spin two, spin one and spin zero field. Besides these long range forces, there are short range forces due to exchange of massive KK fields. Moreover, the resulting 4-D action also contained tower of massive spin two, spin one and spin zero field. However, the proposal of KK was not successful to describe physical phenomena of that era. The Kaluza-Klein paradigm was examined by several authors; notably among them are Schrödinger [6], Jordan [7], Bergmann and Einstein [8], Pauli [9] and Bergmann [10]. The idea was revived when consistent supersymmetric and supergravity theories were constructed in higher dimensions, D > 4. In an important paper, Scherk and Schwarz [11] presented the procedure for compactification of higher dimensional theories through a systematic prescription of dimensional reduction. This technique has been widely used in present times. Furthermore, string theories offered the prospect of unification of fundamental forces since it was possible to construct theories which had nonabelian massless gauge bosons, chiral fermions and admitted those gauge groups which render the theory to be free from anomalies. The construction of heterotic string in D = 10 had several desirable attributes and led to vigorous activities to obtain four dimensional string effective action though compactification schemes. Thus there were attempt to derive realistic string effective actions in order to investigate phenomenological implications of string theories. It was generally accepted that the compactification scale will be in the vicinity of Planck scale or scale of grand unified theories (GUT). We recall that the string tension is of the order of Planck scale and therefore, massive excited stringy states will have mass spacings of the order of that scale. If the compactification radius corresponds to the scale mentioned above then the KK zero modes are to correspond with the known particle spectrum and KK excitations will have masses proportional to inverse of compactification radius as will be discussed in the sequel. In this optics, there seems to have a slim chance of experimentally observing the excited states of strings in foreseeable future. Thus if the compactification scale is at Planck regime or in the GUT regime, the chances of experimentally observing stringy states, in D = 4, even in distant future is almost hopeless. We do not have accelerator technology to create Planck scale collision energies.

An alternative scenario, for low scale compactification, envisages a low scale (i.e. large radius) compactification of higher dimensional (D > 4) theories [12, 13]. Subsequently, an attractive proposal for large radius compactification was introduced which incorporated several interesting novel features [14, 15, 16]. There is strong basis to pursue phenomenological analysis of this proposal. We mention in passing that the large radius compactification proposal is not specifically intended for string theory effective actions. A theory which is proposed in higher dimensions, D > 4, when compactified to 'physical' lower dimension, D = 4, is amenable to this paradigm. Moreover, there are hopes that the predictions could be subjected to experimental tests. It is proposed, in this scenario, that scale of compactification could be in the vicinity of a few TeV and there are hopes that predictions of this proposal could be tested at LHC energies. Indeed, ATLAS and CMS are searching for such particles which might be produced at LHC. These two experiments have obtained lower limits on mass of such 'exotic' particles which range from 2 TeV to 6 TeV [23, 24]. Of course,

inputs from various phenomenological models.

Instead of proposing another model for production of spectrum of particles arising from large-radius-compactification scheme, we derive bounds on production cross section of the so called exotic KK states. We work in the framework of general field theories with minimum inputs from models. These bounds are derived in a nonperturbative framework; just as the Froissart bound on hadronic total cross section, σ_t , does not appeal to perturbation theory. We shall dwell on this aspect, in detail, in subsequent sections. We mention, in passing, that we are handicapped by the fact that there is no experimental data to be used as inputs. Therefore, these bounds are derived from first principles of a theory. Furthermore, there are a host of models of large-extra-dimensions which are built with the original paradigm of [14, 15, 16]. The topic has been extensively reviewed in the forms of lectures at various schools and workshops [19, 20, 21, 22, 25, 26, 27, 28, 29, 30, 31, 32, 33].

We envisage a five dimensional theory for the sake of simplicity. The arguments presented here can be extended to a theory in D > 4 dimensions since the analyticity properties have been investigated in detail by us [39]. We are also aware that higher dimensional theories, beyond five dimensions might face difficulties while compactified in large radius compactification scenario from phenomenological considerations. However, we present our investigation from a rigorous and new perspective so that insights gained from this work might be useful. We add the following remarks to clarify some statement made in this article from time to time. We shall use terms such as strong bound (result) and weak bound (result) on and off. These two terms are to be understood in the following sense. A strong bound (result) means that given a set of basic assumptions or axioms if the energy dependence of a bound (generally power of *logs* is less compared to another bound (which is derived under same set of assumptions (axioms) has a higher power of *logs* then we designate former to be a stronger bound. In our case, we work in the frameworks of LSZ axioms; therefore, with above definition of stronger and weaker bounds we shall be able to distinguish quality of bounds (i.e. strong or weak).

We consider production of Kaluza-Klein states in high energy collision of protons at LHC. The process under considerations is: $p + p \rightarrow K_1 + K_2$ where K_1 and K_2 are excited KK states. They carry baryon number and also KK charges. The global KK charges are conserved as will discussed later. Therefore, if K_1 is assigned baryon number B = 1 and KK charge q_n then K_2 carries baryon number B = 1 and KK charge $-q_n$. Moreover, KK charges are quantized and are proportional to an integer. If R is the radius of compactification of the circle then the mass formula a KK state is $m_n^2 = m_0^2 + (\frac{n}{R})^2, n \in \mathbb{Z}$; here m_0 is mass of the zero mode (see more discussions later). All these aspects will be dealt with in detail in the next section. Obviously, lowest mass KK 'baryon' is heavier than proton. Correspondingly, the s-channel threshold is $s = (m_p + m_1)^2$ where m_1 is the mass of first excited KK state (carries B = 1 and n = 1). Note that m_p is the proton mass which corresponds to mass of zero mode in KK compactification. The t-channel threshold is also important in that its location

plays an important role to write down fixed-t dispersion relations. The threshold is $(m_p + m_{K_1})^2$. We note that the production process, $p + p \rightarrow K_1 + K_2$ is an inelastic reaction where masses of incoming particles are not same as outgoing particles. Our attentions will be primarily focused on inelastic reactions. Let us note a few simple points. Notice that the lowest mass KK excitation corresponds to $n = \pm 1$ i.e. one of the states, say K_1 carries quantum numbers $B = 1, q_n = 1$ whereas K_2 has quantum numbers $B = 1, q_n = -1$. This reaction is consistent with all conservation laws (all the particles are strongly interacting). We shall discuss the analyticity properties of inelastic reactions, generically denoted as $a + b \rightarrow c + d$ in subsequent sections. The purpose of this articles is twofold. (i) We envisage a situation where one spatial dimension is compactified on a circle; it is the so called S^1 compactification. As we shall describe in detail, in order to study analyticity properties of such a theory it is necessary to construct the Hilbert space where not only the zero modes define a state vector but state vectors associate with KK excitations are to be included. It is essential to derive spectral representation of the amplitude where the complete set of states consists of the states associated with the zero mode sector and the set of states associated with the KK towers. We shall present arguments on the spectral representation in some details. Moreover, the case of inelastic scattering requires some extra care due to additional technical problem. (ii) We also consider the situation when the theory has no compact dimension i.e. noncompact spatial coordinates. It is essentially the problem of investigating a D = 5 field theory and deriving analyticity properties of scattering amplitude. This aspect has been comprehensively investigated by us [39]. For the case at hand, we have to deal with four point amplitude. Here basis functions for partial wave expansion are the Geggenbauer polynomials.

We consider a simple model where it has only one type of field in D = 5 to start with. When we compactify this theory on S^1 and look at the spectrum in manifold $R^{3,1} \otimes S^1$, it consists of a field which has the same mass as the D = 5 theory. This corresponds to the zero mode. Moreover, there are KK towers of massive fields. This argument can be extended to the case where D = 5 theory has several species of fields. In a realistic scenario, indeed, all the fields of the standard model (of D = 4) have their counterparts in the five dimensional world. Therefore, from the perspective of D = 5 theory, the standard model, D = 4 theory, fields are zero modes of compactified D = 5 theory. Naturally, there will be KK modes associated with species present in D = 5 theory. Let us consider a single field in the D = 5 theory. While deriving spectral representation for four point amplitude, in D = 5 theory, we have to sum over complete set of physical states. Moreover, since there is only one field, the four point amplitude describes elastic process. If we have several species of particles in D = 5 then there will be elastic amplitude as well as inelastic amplitude. Therefore, it is important to study analyticity properties of inelastic reactions as well. We shall return to discussions of this subject in later sections.

We assume that the momenta of the produced exotic particles lie in three spatial, physical, dimensions i.e. $k_i = (k_i^0, \mathbf{k}_i), i = 1, 2$, where $\mathbf{k_1}$ and $\mathbf{k_2}$ are three momenta

of two outgoing particles. The energies and momenta of the exotic particle can be measured in LHC experiments. This is our choice of kinematical configurations. We study production of states in LHC. There are two possibilities which we have alluded to above. The excited particles produced in case of S^1 compactification are henceforth referred to as Kaluza-Klein (KK) states, which appear in large-radiuscompactification models, in collisions at LHC. We derive bounds on total production cross section, bounds on differential cross sections and bounds on inclusive cross sections based on principles of general local quantum field theories (QFT). Therefore, our efforts are to minimize the inputs from specific models as much as possible. There are certain differences between existing rigorous bounds derived from axiomatic field theories, in D = 4, and the problem under consideration. There are various scenarios, adopted from different phenomenological perspectives, within LRC paradigm. The references cited here, especially several of the TASI lectures, deal in detail, attributes of LRC models and present predictions from diverse perspectives. We have made the following choice and it is a simple extension of the standard model as far as the underlying assumptions are concerned. It is called universal extra dimension (UED) paradigm (see [31] for a review of this idea). It is proposed that the four dimensional standard model (SM) is promoted to a D dimensional theory where D = 4 + n, nis the number of extra spatial dimensions. Therefore, the fundamental constituents of the standard model such as gauge bosons, fermions and scalars are promoted to fields in 4 + n dimensions. The basic building blocks of the 4-dimensional world appear as zero modes of the compactified theory. Therefore, additional KK states appear after the 4 + n dimensional theory is compactified. It is assumed that extra dimensions are flat and so is our four dimensional world where SM is experimentally tested. This approach has several advantages from phenomenological point of view and it has also shortcoming as is the case with any model in the LCR scenario. We have chosen this from certain advantages it offers from our perspective as will be elaborated soon. In the context of model buildings, the practitioners incorporate extra assumptions so that perturbative corrections coming from states arising from compactifications do not upset results of D = 4 theories at tree level. These aspects (for that matter those assumptions) are not needed here. In this scheme some of their inbuilt results are of interests to us. Parity is an exact symmetry. A single, first excited KK state (n = 1) cannot be produced in isolation. The lightest state, n = 1, is stable. However, it is not very obvious that n = 1 KK quantum number be an absolutely conserved charge since it is not result of an exact local symmetry (see discussions in [37, 38]). We remark that the toroidal compactification of a higher dimensional theory (where D dimensional theory is flat) satisfies LSZ axioms [39]. Moreover, the Hilbert space structure of flat space D = 4 theory derived from a D = 5 flat space theory has been discussed in [37, 38] in detail. Furthermore, toroidal compactification of a higher dimensional theory to D = 4 theory poses no additional difficulties. In view of preceding remarks we consider the essential aspects of UED. We focus on the scalar sector. We intend to explore another scenario where the extra dimension is decompactified, already, at LHC energy. Thus states of the five dimensional (decompactified) model will be produced. A systematic analysis is quite desirable for following reasons. First the bound on total cross sections, the generalized Froissart bound, is different for D > 4. This case has been rigorously studied by us [39]. Furthermore, when we derive spectral representations for absorptive part of four point amplitude in the compactfied, D = 4, theory, the KK states are introduced as physical intermediate states. This aspect is discussed in the next section. On the other hand, in a decompactified theory (here D = 5 theory), the intermediate states appearing in the spectral representation are physical states. We work in the LSZ formulation. Therefore, all the parameters such as mass, charge etc are the physical observables. In this framework, all quantum effects are accounted for. For example let us consider πN scattering. The proof of Froissart bound does not rely on any perturbation theoretic frame work. Similarly, derivation on properties of diffraction peak such as shrinkage of width of diffraction peak and location of minimum of differential cross section as a function of t are derived in this framework. Therefore, LSZ formulation, when applied to decompactified theory will capture effectively the quantum effects without resorting to perturbation theoretic arguments². The presence of spinning particles do not pose any serious problem [45, 46]

The paper is organized as follows. We shall present the general structure of the compactified theory in the next section, Section 2. We briefly summarize the contents of [37, 38] which will be utilized in this work. We refer the interested reader to go through the details in the above mentioned papers. The starting point is to consider a massive scalar field theory in $\hat{D} = 5$ dimensional flat space with Lorentzian signature. It was shown in [39] that LSZ procedure is applicable to study analyticity properties for higher dimensional theories, D > 4. Notice that the asymptotic states i.e. 'in' and 'out' states are defined in LSZ formulation and this concept is not special to four dimensions. Another important point is to be kept in mind is that in LSZ approach the only the physical parameters such as mass of particles and physically measured coupling constants are used. In essence, as is evident from LSZ axioms, all results are derived without addressing to perturbation theory.

We derive results on analyticity properties of scattering amplitude for inelastic two body reactions in Section 3. There have been previous attempts to study analyticity of inelastic reactions in the past [40, 41]. However, these authors were (probably) unaware of the intricacies associated with unequal mass scattering in deriving analyticity domains in the *t*-plane. As we shall present, in the sequel, identification of

²I thank A. P. Balachandran for discussions on this point. He emphasised that Froissart bound and its subsequent refinement by Martin were accomplished long before QCD was proposed (and established) as the theory of strong interactions. Indeed, models to study strong interactions in the field theoretic framework were not proved to be renormalizable, like QED, on various grounds. Therefore, establishing bounds for theories in higher dimensions (with compact or noncompact coordinates) might play an useful role in future.

the holomorphic domain in *t*-plane is to be treated with some care since the physical energy threshold is not identical to elastic threshold. We have invoked rigorous theorems from axiomatic field theory to argue that the amplitude of s-channel reaction can be analytically continued to the *u*-channel reaction. Some of these delicate points were not addressed in the past [40, 41]. Next we discuss the existences of Lehmann ellipses since these results were derived earlier and what is the analog of Martin'e ellipse. Subsequently, a fixed-t dispersion relation is written down. We devote Section 4 to deriving bounds on various cross sections alluded to above. We have employed refined technique of Singh and Roy [43] as well as Einhorn and Blankenbecler [44] to improve results of [40, 41] where the prefactors in their bounds appear as arbitrary constants. This arbitrariness could be removed by invoking more refined procedures which are outlined in this section. Section 5 deals with case when the 5^{th} dimension is decompactified. There are simplifications in kinematical configurations for four point function. It is relatively simple to derive known results of D = 4 theory for the D = 5theory while considering the four point functions. The proof of edge-of-the wedge theorem of D = 4 theory is practically carried over for 4-point function of D = 5 theory. This leads, de facto, to proof of crossing which is crucial to write fixed t dispersion relation. We do not repeat the proof of our derivation [39] here. However, we present the essential arguments. Once we are able to write fixed t dispersion relation for inelastic amplitude we have proved the analyticity properties in this case. We discuss the bounds on inclusive reaction which are obtained by adopting techniques used to derive bounds on inelastic processes.

The last section, Section 6, contains summary of our results and discussions. It is important to note that if the exotic particles are produced in LHC energy regime then new thresholds will open up. If such is the case then the effects of the presence of new threshold will affect the analyticity properties of scattering amplitude. Indeed, a precise measurement of real part of the scattering amplitude (in other words measurement ratio of real to imaginary part) at energies below the threshold will signal the possibility of production of exotic particles. We shall discuss this aspect in discussion section.

2 Properties of inelastic scattering amplitude for S^1 compactified theory

We consider a scalar field, $\hat{\phi}(\hat{x}), \hat{x}^{\hat{\mu}} = 0, 1, 2, 3, 4$ in a five dimensional flat space. It is convenient to decompose the five dimensional spacetime coordinates, $\hat{x}^{\hat{\mu}}$ as follows for later conveniences:

$$\hat{x}^{\mu} = (x^{\mu}, y) \tag{1}$$

where x^{μ} are flat space coordinates, $\mu = 0, 1, 2, 3; y$ is the compact coordinate on S^1 with periodicity $y + 2\pi R = y$, R being the radius of S^1 . As a consequence of

 S^1 compactification the resulting spacetime manifold is $R^{3,1} \otimes S^1$. We remind that asymptotic 'in' and 'out' fields satisfy free field equation in 5-dimensions before we discuss consequences of compactification. The equation of motion, for D = 5 theory, is $[\Box_5 + m_0^2]\hat{\phi}^{in,out}(\hat{x}) = 0$, m_0 being the mass of the scalar field. We expand the field, $\hat{\phi}(x, y)$, in a Fourier series in y-variable after implementing S^1 compactification

$$\hat{\phi}^{in,out}(\hat{x}) = \hat{\phi}^{in,out}(x,y) = \phi_0^{in,out}(x) + \sum_{n=-\infty,n\neq 0}^{+n=\infty} \phi_n^{in,out}(x)e^{\frac{iny}{R}}$$
(2)

where $\phi_0^{in,out}(x)$, the zero mode, has no *y*-dependence. The terms in rest of the series (2) are periodic in *y*. The five dimensional Laplacian, \Box_5 , is decomposed as sum two operators: \Box_4 and $\frac{\partial}{\partial u^2}$. The equation of motion is

$$\left[\Box_4 - \frac{\partial}{\partial y^2} + m_n^2\right]\phi_n^{in,out}(x,y) = 0 \tag{3}$$

where $\phi_n^{in,out}(x,y) = \phi_n^{in,out} e^{\frac{iny}{R}}$ and n = 0 term has no y-dependence being denoted as $\phi_0(x)$; from now on $\Box_4 = \Box$. It follows from equations of motion that $m_n^2 = m_o^2 + \frac{n^2}{R^2}$. Therefore, there is a tower of massive states. The momentum along 5th direction is quantized, $p_4 = q_n = \frac{n}{R}$. q_n is an additive conserved quantum number. We continue to call it Kaluza-Klein (KK) charge and note that we have considered a flat space theory; there is no gravitational interaction in the five dimensional theory we have adopted. For the interacting field $\hat{\phi}(\hat{x})$

$$\hat{\phi}(\hat{x}) = \hat{\phi}(x,y) = \phi_0(x) + \sum_{n=-\infty, n\neq 0}^{n=+\infty} \phi_n(x) e^{\frac{iny}{R}}$$
 (4)

The equation of motion for the interacting fields has a source terms on the *r.h.s.* Naturally, the source current is expanded in Fourier series as is the expansion (4). Each field $\phi_n(x)$ will have a current, $J_n(x)$ associated with it as source current and the expansion is

$$\hat{j}(x,y) = j_0(x) + \sum_{n=-\infty, n\neq 0}^{n=+\infty} J_n(x) e^{iny/R}$$
(5)

The set of currents, $\{J_n(x)\}$, are the source currents associated with the tower of interacting fields $\{\phi_n(x)\}$. These fields carry the discrete KK charge, q_n . Therefore, $J_n(x)$ also carries the same KK charge. This is to be kept in mind while going through LSZ reductions to write down the amplitudes and Greens functions. The zero mode, $\phi_0^{in,out}$, creates its own Fock space. Similarly, each of the KK fields, $\phi_n^{in,out}(x)$, creates its own Fock space. We present some illustrative examples below. A state with spatial momentum, \mathbf{p} , energy, p_0 and discrete momentum q_n (KK state) is created by

$$A^{\dagger}(\mathbf{p}, q_n)|0\rangle = |p, q_n\rangle, \quad p_0 > 0 \quad i.e. \ p_{\mu} \in V^+$$
 (6)

Our starting point was a 5-dimensional theory of a scalar field. After S^1 compactification, the geometry is $R^{3,1} \otimes S^1$ and we have discussed the spectrum of the compactied theory. A massive field of mass m_0 , the mass of zero mode, and a the KK tower characterized by masses and 'charges' $\{(m_n, q_n)\}$ are in the spectrum.

The Hilbert space associated with the five dimensional theory is \mathcal{H} . On compactification, the resulting theory, $R^{3,1} \otimes S^1$, leads to decomposition of the original Hilbert space into direct sum of Hilbert spaces each of which is characterized by its KK charge q_n .

$$\hat{\mathcal{H}} = \sum \oplus \mathcal{H}_n \tag{7}$$

Thus \mathcal{H}_0 is the Hilbert space constructed from $\phi_0^{in,out}$ with charge $q_n = 0$ and it is built by the actions of the creation operators $\{a^{\dagger}(\mathbf{k})\}$ acting on the vacuum. The resulting states span \mathcal{H}_0 . A single particle state is $a^{\dagger}(\mathbf{k})|_0 \ge |\mathbf{k}\rangle$ and multiparticle states are created using the procedure outlined above. Note the orthogonality relation between two states of different values of q_n belonging to two different vector bases,

$$\langle \mathbf{p}, q_n | \mathbf{p}', q_{n'} \rangle = \delta^3(\mathbf{p} - \mathbf{p}')\delta_{n,n'}$$
(8)

It is assumed that there are no bound states in the theory.

The LSZ formalism can be adopted for the compactified theory. If we keep in mind the steps introduced above, it is possible to envisage field operators $\phi_n^{in}(x)$ and $\phi_n^{out}(x)$ for each of the fields for a given n. Therefore, each Hilbert space, \mathcal{H}_n will be spanned by the state vectors created by operators $a^{\dagger}(\mathbf{k})$, for n = 0 and $A^{\dagger}(\mathbf{p}, q_n)$, for $n \neq 0$. *Remark*: Note that in (2) sum over $\{n\}$ runs over positive and negative integers. If there is a parity symmetry $y \to -y$ under which the field is invariant we can reduce the sum to positive n only leading to $\cos \frac{n\theta}{R}$. On the other hand if the field is odd under $y \to -y$ we have sine-function. We make no such assumptions here.

We considered three types of scattering process: (a) scattering of zero modes, $\phi_0 + \phi_0 \rightarrow \phi_0 + \phi_0$. It is the conventional scattering of two massive scalars. (b) A zero mode, ϕ_0 scattering with a KK excitation with charge q_n . This is an elastic scattering of zero KK charge particle with nonzero KK charge particle. The processes (a) and (b) are treated without complications adopting the well known precedures. (c) The reaction $\phi_m + \phi_n \rightarrow \phi_{m'} + \phi_{n'}$ deserves attention as KK charge conservation is required i.e. $q_m + q_n = q_{m'} + q_{n'}$. We had considered elastic reaction since our goal was to prove dispersion relation in order to resolve problems posed by Khuri's [36] analysis in potential scattering. The study of analyticity of elastic process in (c) i.e $\phi_+\phi_n \rightarrow \phi_m + \phi_n$ required careful investigations since the intermediates appear from the entire Hilbert space, $\oplus \mathcal{H}_n$, as long as all conservation laws are satisfied. These issues have been discussed in detail by in references [37, 38]. We are not going to repeat those computations in this article. We focus on inelastic reactions and analyze the analyticity properties of corresponding scattering amplitude. It will be evident that additional care should be exercised when we identify domains of holomorphy in s and in t planes. Moreover, the results presented here on bounds are improvements over previous works where due attention was not carefully paid in study of certain properties of the domains of holomorphy.

We investigate, in this section, two body inelastic scattering amplitude for S^1 compactified theory. We consider inelastic four point amplitude, $F(s,t)^{ab\to cd}$, where c and d are 'KK' states of nonzero KK charge. We shall elaborate on this aspect in the sequel. Consequently, we obtain bounds on $\sigma_t^{ab\to cd}$, on $\frac{d\sigma^{ab\to cd}}{d\Omega}$ in the forward direction as well as in nonforward direction: $\theta_{c,m} \neq 0, \pi$. The bound on $\sigma_t^{ab\to cd}(s)$ is not as stringent as the Froissart bound [34], $\sigma_t(s) \leq \frac{4\pi}{t_0} [\log \frac{s}{s_o}]^2$, and we shall dwell upon this aspect when we arrive at the bound. We also consider inclusive reaction, $a+b\to c+X$ where c is the observed KK state and X could be a single particle or collection of multiple particle. Feynman had introduced the notion of inclusive reactions [35] in the context of hadron collisions and had proposed scaling phenomena based on his parton model. We present bounds on differential cross sections of inclusive reaction in the context of production of KK states.

It is important to mention the difficulties, at this juncture, what one has to surmount is deriving rigorous bounds on inelastic processes in general. We recall that Froissart bound is derived from following ingredients, which are proved from axiomatic field theory. (i) Analyticity of scattering amplitude in the complex s-plane which is also analytic in complex t-plane within the Lehmann ellipses. We need to prove existence of Martin ellipse. (ii) Crossing symmetry. (iii) Convergence of partial wave expension inside Lehmann-Martin ellipse. Whereas the results (i) and (ii) can be proved for inelastic two body reaction, $a + b \rightarrow c + d$; the existence of Martin's ellipse is proved from positivity properties of absorptive elastic scattering amplitude and positivity properties of elastic partial wave amplitudes $0 \leq |f_l(s)^{el}|^2 \leq \text{Im } f_l(s)^{el} \leq 1$ (follows from unitarity). We refer to books and review articles on the study of analyticity properties of scattering amplitude from the perspectives of axiomatic field theory [45, 46, 47, 48, 49, 50, 51, 53, 54, 55, 56, 57].

In fact the partial wave amplitude of inelastic process do not enjoy positivity properties as those of the elastic case. Therefore, proof of existence of Martin's ellipse and proof holomorphy of scattering amplitude in a complex t domain requires some careful analysis. We analyze these issues to identify the domains of analyticity of scattering amplitude. Therefore, a fixed-t dispersion relation is written down only after these issues are addressed and accounted for. Recently, we addressed the case of analyticity of scattering amplitude in field theory where one spatial dimension is compactified on a circle of radius R, S^1 compactification. In fact Khuri [36] had shown, in a nonrelativistic potential model with S^1 compactification, that the forward scattering amplitude does not satisfy dispersion relation. If so were the case in relativistic QFT it would have disastrous consequences. We proved, starting from LSZ axioms, that the scattering amplitude not only satisfies forward dispersion relation but also nonforward dispersion relations are satisfied [37, 38]. The proof was presented for elastic scattering. The proof of dispersion relation for inelastic scattering, for a theory with S^1 compactification, is not so straightforward as we shall discover.

3 Production of Particles in Large-Radius-Compactification Proposal

We discuss production of particles which belong to spectrum of states of large-radiuscompactifaction (LRC) scenario in the UED paradigm. Our purpose is to derive bounds on cross sections without appealing to a specific model. The starting point is to consider an inelastic process $a + b \rightarrow c + d$ where a and b are particles accelerated in an accelerator (they are two colliding protons in LHC experiments) and c and dare two particles which belong to the spectrum of particles (charged KK states) in the LRC models with the hypothesis of UED. Existence of such exotic particles is yet to be confirmed experimentally. We refer to them, generically, as Kaluza-Klein (KK) states since some of the particle of UED scenario belong to KK excitations. We noted that KK, exotic states carry additive quantum numbers (charges), denoted as $\{q_n\}$ which are conserved. The KK charges are quantized since they are associated with momenta in the compact directions. These additive charges may be viewed to be analogous to baryon number or lepton number. In view of preceding remarks, these particles will be produced in pairs and carry equal and opposite KK charges besides other quantum number. For example, if at LHC, in pp collision a pair of these exotic particles carrying two units of KK charge each (equal and opposite charges) one will have quantum number of baryon number and two units of KK charge (this is like strangeness quantum number which is additive). The other one will carry baryon number but opposite KK charge for total KK charge conservation. Thus one of them might decay to a particle carrying baryon number and one unit of KK charge and to a boson with one unit of opposite KK charge (if kinetics permit such decays). We are not proposing a phenomenological analysis of such a production mechanism. We present it, rather as an illustrative reaction. We have noted earlier that there are lower bounds on masses of particle of LRC theories set by ATLAS and CMS collaborations of LHC.

Our goal is to examine analyticity properties of the aforementioned scattering amplitude from principles of general field theory which will be presented shortly. Notice that we envisage inelastic scattering amplitude which does not satisfy all the properties of elastic scattering amplitude. In particular, the elastic partial wave amplitudes, $f_l^{ab\to ab}(s)$, satisfy positivity constraints: $0 \leq |f_l^{ab\to ab}(s)|^2 \leq Im f_l^{ab\to ab}(s) \leq 1$, whereas inelastic amplitudes do not. Moreover, the domain of analyticity of inelastic amplitudes, in the complex t-plane, especially the proof from the existence of Martin's ellipse depends on ingredients which are proved from the properties enjoyed by elastic amplitude. Therefore, our first step is to recapitulate some of the important results on analyticity properties of elastic amplitudes which have been proved in the axiomatic field theoretic framework. These will be utilized to derive analyticity of inelastic scattering amplitudes. We shall adopt the convention that a scattering amplitude will have a superscript denoting the reaction and it will be evident from this notational convention whether we are considering an elastic process or an inelastic process.

Notations and Kinematical variables

We consider inelastic reaction

$$a + b \to c + d \tag{9}$$

For the case at hand masses of particles a and b are equal, $m_a = m_b$ and the masses of two outgoing particles, c and d, are also equal, $m_c = m_d$. The initial momenta of a and b are denoted as p_1 and p_2 whereas final state particles, c and d carry momenta k_1 and k_2 . The Mandelstam variables are

$$s = (p_1 + p_2)^2 = (k_1 + k_2)^2$$
, $t = (p_1 - k_2)^2$, $u = (p_1 - k_1)^2$, $s + t + u = 2(m_a^2 + m_c^2)(10)$

We define $\mathbf{P}_1(s)$ and $\mathbf{K}_1(s)$ to be c.m. momenta of particles (a, b) and (c, d) respectively and

$$P_1^2(s) = \frac{1}{4}(s - 4m_a^2)$$
 and $K_1(s)^2 = \frac{1}{4}(s - 4m_c^2)$ (11)

Let θ be the scattering angle in the c.m. frame then

$$\cos\theta = \frac{\mathbf{P}_1(s)^2 + \mathbf{K}_1(s)^2 + t}{2|\mathbf{P}_1(s)||\mathbf{K}_1(s)|}$$
(12)

We also define following kinematical variables of interest

$$(\mathbf{M}_a^2, \mathbf{M}_b^2, \mathbf{M}_c^2, \mathbf{M}_d^2)$$
 and $(\mathbf{M}_{ab}^2, \mathbf{M}_{ac}^2, \mathbf{M}_{ad}^2, \mathbf{M}_{bc}^2, \mathbf{M}_{bd}^2, \mathbf{M}_{cd}^2)$ (13)

Where \mathbf{M}_{i}^{2} , i = a, b, c, d corresponds to $(mass)^{2}$ of two or more particle states which carry quantum numbers of particle 'i'. Similarly, \mathbf{M}_{ij}^{2} , $i, j = a, b, c, d, i \neq j$ corresponds to two or more particle states which carry quantum numbers of both the particles, 'i' and 'j'. This this the case when we consider various scattering channels. We shall use these definitions in the sequel.

Next, we consider LSZ reduction procedure for four point function: $a + b \rightarrow c + d$. We denote the fields associated with outgoing particles as ϕ_n and ϕ_{-n} so that they carry opposite KK charges; *n* takes positive or negative integer values. Note that the fields might carry other quantum numbers. We adopt LSZ [42] procedure to derive expression for the 4-point inelastic amplitude. The axioms of LSZ are

A1. The states of the system are represented in a Hilbert space, \mathcal{H} . All the physical observables are self-adjoint operators in the Hilbert space, \mathcal{H} .

A2. The theory is invariant under inhomogeneous Lorentz transformations.

A3. The energy-momentum of the states are defined. It follows from the requirements of Lorentz and translation invariance that we can construct a representation of the orthochronous Lorentz group. The representation corresponds to unitary operators,

 $U(a, \Lambda)$, and the theory is invariant under these transformations. Thus there are Hermitian operators corresponding to spacetime translations, denoted as P_{μ} , with $\mu = 0, 1, 2, 3$, which have following properties:

$$\left[P_{\mu}, P_{\nu}\right] = 0 \tag{14}$$

If $\mathcal{F}(x)$ is any Heisenberg operator then its commutator with P_{μ} is

$$\left[P_{\mu}, \mathcal{F}(x)\right] = i\partial_{\mu}\mathcal{F}(x) \tag{15}$$

It is assumed that the operator does not explicitly depend on spacetime coordinates. If we choose a representation where the translation operators, P_{μ} , are diagonal and the basis vectors $|p, \alpha \rangle$ span the Hilbert space, \mathcal{H} ,

$$P_{\mu}|p,\alpha\rangle = p_{\mu}|p,\alpha\rangle \tag{16}$$

then we are in a position to make more precise statements:

• Existence of the vacuum: there is a unique invariant vacuum state $|0\rangle$ which has the property

$$U(a,\Lambda)|0\rangle = |0\rangle \tag{17}$$

The vacuum is unique and is Poincaré invariant.

• The eigenvalue of P_{μ} , p_{μ} , is light-like, with $p_0 > 0$. We are concerned only with massive stated in this discussion. If we implement infinitesimal Poincaré transformation on the vacuum state then

$$P_{\mu}|0>=0, \text{ and } M_{\mu\nu}|0>=0$$
 (18)

from above postulates and note that $M_{\mu\nu}$ are the generators of Lorentz transformations.

A4. The locality of theory implies that a (bosonic) local operator at spacetime point x^{μ} commutes with another (bosonic) local operator at x'^{μ} when their separation is spacelike i.e. if $(x - x')^2 < 0$. Our Minkowski metric convention is as follows: the inner product of two 4-vectors is given by $x \cdot y = x^0 y^0 - x^1 y^1 - \ldots - x^3 y^3$. Note that Hermitian conjugate of ϕ_n , $\phi_n^{\dagger} = \phi_{-n}$. By definition it transforms as a scalar under inhomogeneous Lorentz transformations

$$U(a,\Lambda)\phi_n(x)U(a,\Lambda)^{-1} = \phi_n(\Lambda x + a)$$
(19)

The micro causality, for any two local field operators, $\Phi(x)$ and $\Phi(x')$ is stated to be

$$\left[\Phi(x), \Phi(x')\right] = 0, \quad for \ (x - x')^2 < 0 \tag{20}$$

Remark: Note that, for the compactified theory, the full Hilbert space is direct sum of Hilbert spaces, \mathcal{H}_n , designated by KK charges n. Therefore, while deriving certain spectral representations, we have to sum over all states of the full Hilbert space subject to conservation laws.

We define the amplitude as follows. The incoming particle state is $|p_1, p_2 in \rangle$ and outgoing state, also of two particles is $|k_2, k_1 out \rangle$. Note that we are considering the inelastic amplitude. We mention in passing that the LSZ reduction procedure, spectral representation for retarded, advanced and causal commutators and their spectral representations hold good for elastic amplitudes as well as inelastic counterparts. We shall point out when there are distinctions between their analyticity properties at appropriate juncture. Let us consider the case when the two outgoing particle states are reduced.

$$< k_{2}k_{1}, out|p_{2}p_{1}, in > - < k_{2}k_{1}, in|p_{2}p_{1}, in > = -\frac{i}{2\pi^{4}} \int d^{4}x d^{4}y e^{+ik_{1}.x+ik_{2}.y} (\Box_{x} + m_{c}^{2})(\Box_{y} + m_{c}^{2}) < 0|R\phi_{n}(x)\phi_{-n}(x)|p_{2}p_{1} >$$

$$(21)$$

the retarded product is defined to be $R\phi_n(x)\phi_{-n}(y) = -\theta(x_0-y_0)[\phi_n(x), \phi_{-n}(y)]$. The matrix element of R-products are Lorentz and translation invariant. Consequently, the latter invariance implies that it depend on difference of coordinates. We shall use this property extensively. We define the scattering amplitude as

$$c < k_2 k_1, out | p_2 p_1, in > - < k_1 k_2, in | p_2 p_1, in > = 2\pi i \delta^4 (p_1 + p_2 - k_1 - k_2) F(p_1, p_2, k_1, k_2) (22)$$

Therefore,

$$F(p_1, p_2, k_1, k_2) = -\int d^4x d^4y e^{i(k_1 \cdot x + k_2 \cdot y)} (\Box_x + m_c^2) (\Box_y + m_c^2) < 0 |R\phi_n(x)\phi_{-n}(y)| p_2 p_1, in >$$

$$= -\int d^4x d^4y e^{i(k_1 \cdot x + k_2 \cdot y)} < 0 |RJ_n(x)J_{-n}(y)| p_2 p_1 >$$
(23)

We have used equations of motion i.e. $(\Box_x + m_c^2)\phi_n(x) = J_n(x)$ where $J_n(x)$ is the source current. Moreover, we used a relation $(\Box_x + m_c^2)(\Box_y + m_c^2)R\phi_n(x)\phi_{-n}(y) =$ $RJ_n(x)J_{-n}(y)$. This relation, deserves following qualifying remarks since additional terms are present on the right hand side which have not been displayed explicitly. Notice that the two Klein-Gordon operators $(\Box_y + m_c^2)(\Box_y + m_c^2)$ act on the R-product; $R\phi_n(x)\phi_{-n}(y)$. It is evident that the term $RJ_n(x)J_{-n}(y)$ would appear. In addition, the operation of KG operators on *R*-product of two field operators will give derivatives of δ -functions. Moreover, there will be equal time commutators of operators. Notice that derivatives of delta function when Fourier transformed will give products of momenta. However, these products have to be Lorentz invariant since *S*-matrix elements are Lorentz invariant. Therefore, such terms can be expressed in terms of Mandelstam variables, *s*, *t* and *u*. As far as study of analyticity properties of the amplitude is concerned, presence of such terms (polynomials in s,t and u) do not affect us; moreover, for local field theories, only finite number of such δ -function derivative terms will appear. The third type of terms are equal time commutator of operators. These terms vanish from microcausality arguments: $[\mathcal{O}(x_0, \mathbf{x}), \mathcal{O}(y_0, \mathbf{y})]\delta(x_0 - y_0) = 0$.

We have denoted the fields arising from reduction of out states as $\phi_n(x)$ and $\phi_{-n}(x)$; one carries KK charge of n units and other carries charge of -n units; recall the momentum along compact direction, 'charge' is quantized as $\frac{n}{R}$ where R is radius of the circle. We may reduce one particle from 'out' state and another from 'in' state. In order to study analyticity property of amplitudes with requirements of Lorentz invariance and microcausality, it suffices to consider reductions of two states in whatever combinations we desire. We shall focus attention on the LSZ reduced amplitude (21) in what follows. Moreover, we could reduce all the four particle states i.e. $|k_2k_1, out >$ and $|p_2p_1, in >$; however, as is well known, in order to study analyticity properties and to write down dispersion relations, it is most convenient to reduce two particles from 'in' or 'out' states. Let us define three distributions as follows. We shall recognize shortly the importance of the three distributions.

$$F_{R}(q) = \int d^{4}\xi e^{iq.\xi} \theta(\xi_{0}) < P_{f} | [J_{n}(\xi/2), J_{-n}(-\xi/2)] | P_{i} >$$

$$F_{A}(q) = -\int d^{4}\xi e^{iq.\xi} \theta(-\xi_{0}) < P_{f} | J_{n}(\xi/2), J_{-n}(-\xi/2)] | P_{i} >$$

$$F_{C}(q) = \int d^{4}\xi e^{iq.\xi} < P_{f} | [J_{n}(\xi/2), J_{-n}(-\xi/2)] | P_{i} >$$
(24)

Here $|P_i\rangle$ and $|P_f\rangle$ are arbitrary momentum states. We may treat P_i and P_f as parameters for our purpose since we do not intend to reduce them. There are several pertinent points which need mentioning. First thing to note is that each of the distributions when Fourier transformed respectively to $\tilde{F}_R(\xi)$, $\tilde{F}_A(\xi)$ and $\tilde{F}_C(\xi)$ vanish for $\xi^2 < 0$.

We briefly discuss a few aspects of domain of analyticity of the distributions, defined above, as a function of complex q. It will suffice to focus attention on $F_R(q)$ since it is related to the amplitude we introduced earlier. Moreover, these arguments are easily extended to study of analyticity of $F_A(q)$ and $F_C(q)$. Notice that the Fourier transform $F_R(q)$, $\tilde{F}_R(\xi)$, is defined in the forward light cone, V^+ , i.e. $\xi_0 > 0$ and $\xi^2 > 0$ and it vanishes for $\xi^2 < 0$ due to microcausality. Let us turn attention to properties of the function, $F_R(q)$ for complex q. The function will be analytic in the complex plane if following properties are satisfied. (i) The exponential must converge in all directions in complex plane for $\xi \in V^+$. Thus we desire that Im q > 0 so that exponential is damped in the upper half complex q-plane; however, there is no restriction on real part of q. Moreover, for $\xi \in V^+$, it is necessary that $Im q \in V^+$ also i.e. $(Im q)^2 > 0$ and $Im q_0 > 0$. This defines a forward tube T^+ . Similar arguments is used to define the domain of holomorphy of $F_A(q)$ whose domain of holomorphicity, in the complex q-plane, is defined in the backward light cone, V^- , and the corresponding tube is T^- . We shall argue in what follows that both $F_R(q)$ and $F_A(q)$ vanish for unphysical real values of q-variables and, here $F_R(q) = F_A(q)$ since $F_C(q) = 0$. Let us open up the commutators of the two currents in the expression for $F_C(q)$. Let the first term of the two be $F_{1C}(q)$ which is expressed as

$$F_{1C}(q) = \int d^4 \xi e^{iq.\xi} < P_f |J_n(\xi/2)J_{-n}(-\xi/2)| P_i >$$
(25)

Now we insert a complete set of physical states, $\sum_{p_N,q_N} |p_N,q_N| > \langle p_N,q_N| = 1$, where q_N refer to all other quantum numbers designating the state

$$F_{1C}(q) = \int d^4 \xi e^{iq.\xi} \sum_{N,q_N} \langle P_f | J_n(\xi/2) | N, q_N \rangle \langle N, q_N | J_{-n}(-\xi/2) | P_i \rangle$$
(26)

Note that currents carry KK charges. Therefore, the matrix elements such as $P_f|J_n(\xi/2)|N, q_N > \text{and} < N, q_N|J_{-n}(-\xi/2)|P_f > \text{will be nonvanishing only when}$ all conservation laws are respected. Next we go step by step as follows: (i) use translational symmetry transformations on each of the product matrix elements of (26) such that $J_n(\xi/2) \to J_n(0)$ and $J_{-n}(-\xi/2) \to J_{-n}(0)$ to bring the argument of each current to zero. (ii) Recall that translation operator $e^{-iP.a}$ acting on a state of momentum states, $|p > \text{brings out a factor } e^{-ip.a}$. Thus there will be four such factors coming from this operation and an additional factor of $e^{iq.\xi}$. (iii) When we integrate over ξ , there is a δ function which implies that $p_N = 1/2(P_f + P_i) - q$. The same arguments go through for the second matrix element which appears after the commutator is opened up; the only difference is that product of currents are in reverse order: $J_{-n}(-\xi/2)J_n(\xi/2)$. Now the delta function arising out of ξ integration leads to $p_N = 1/2(P_i + P_f) + q$. We express (24) as

$$F_{C}(q) = \sum_{N} \left[\langle P_{f} | J_{n}(0) | p_{N} = \frac{(P_{i} + P_{f})}{2} - q, q_{N} \rangle \langle p_{N} = \frac{(P_{i} + P_{f})}{2} - q, q_{N} | J_{-n}(0) | P_{i} \rangle - \langle P_{f} | J_{-n}(0) | p_{N} = \frac{(P_{i} + P_{f})}{2} + q, q_{N} \rangle \langle p_{N} = \frac{(P_{i} + P_{f})}{2} + q, q_{N} | J_{n}(0) | P_{i} \rangle \right]$$

$$(27)$$

It is important to emphasize that the intermediate complete set of states, $|p_N, q_N \rangle$ are physical and satisfy mass-shell conditions, $\frac{P_i+P_f}{2} \in V^+$ where V^+ is the forward lightcone. Therefore, it follows that there are states for each of the terms on the r.h.s of (27) which satisfy the following conditions

$$\left(\frac{P_i + P_f}{2} \pm q\right)^2 \ge \mathbf{M}_{\pm}^2$$
, and $\left(\frac{P_i + P_f}{2} \pm q\right)_0 \ge 0$ (28)

In other words, for each of the matrix elements there exists a minimum mass state with positive energy. If this condition is not satisfied by one of the matrix elements that is $(\frac{P_i+P_f}{2}+q)^2 < \mathbf{M_+}^2$, and $(\frac{P_i+P_f}{2}+q)_0 < 0$ then it vanishes. Similarly, when

 $(\frac{P_i+P_f}{2}-q)^2 < \mathbf{M}_2^2$, and $(\frac{P_i+P_f}{2}-q)_0 < 0$ then it also vanishes. Therefore, there are two regions for the two matrix elements where each can vanish. There is a third region where the above conditions (28) are not satisfied for each of the matrix elements and consequently, both matrix elements vanish simultaneously. This is unphysical region for both. The range of q_0 is easily determined as follows. We go to a frame where $1/2(P_i + P_f) = (m_0, \mathbf{0})$. Thus $[1/2(P_i + P_f) \pm q]^2 = (m_0 \pm q_0)^2 \pm (\mathbf{q})^2$. Thus the region where both matrix elements vanish we have a relation: $(m_0 - \sqrt{\mathbf{q}^2 + m_1^2}) \leq q_0 \leq (-m_0 + \sqrt{\mathbf{q}^2 + m_2^2})$ for real values of momenta.

The important question to ask is what is the singularity free region in complex q plane? Jost and Lehmann [58] and, independently, Dyson [59] (JLD) obtained a representation for $F_C(q)$ as well as, by extension, for $F_R(q)$ which enables us to find a singularity free domain in complex q-plane³. Lehmann [60] proved existence of two ellipses, known as small Lehman ellipse (SLE) and large Lehmann ellipse (LLE). The scattering amplitude converges for complex t inside SLE. On the other hand absorptive part of the amplitude converges inside larger ellipse (LLE) defined in the complex t-plane. In other words, the partial wave expansion has a larger domain of converges than $-1 \leq \cos\theta \leq +1$. We present below definitions of Lehmann ellipses (i) SLE

$$E_{SLE}^{ab \to cd}: : |t + (P_1 - K_1)^2 + |t + (P_1 + K_1)^2| < 4|\mathbf{P}_1|\mathbf{K}_1|x_0(s)$$
(29)

where $m_a = m_b$ and $m_c = m_d$, for the case at hand. Furthermore,

$$x_0 = \left[1 + \frac{(\mathbf{M}_a^2 - m_a^2)(\mathbf{M}_c^2 - m_c)^2}{K_1(s)^2 s}\right]^{1/2}$$
(30)

We have used $\mathbf{M}_a = \mathbf{M}_b$ and $\mathbf{M}_c = \mathbf{M}_d$ in arriving at the above formula. Note that $x_0 > 1$. (ii) LLE

$$(11)$$
 LLE

$$E_{LLE}^{ab \to cd} : |t + (P_1 - K_1)^2 + |t + (P_1 + K_1)^2| < 4\mathcal{Z}_L(s)$$
(31)

where $\mathcal{Z}_L(s)$ is given by

$$\mathcal{Z}_{L}(s) = \left(\left[\frac{(\mathbf{M}_{a}^{2} - m_{a}^{2})^{2}}{s} \right] \times \left(\left[\frac{(\mathbf{M}_{c}^{2} - m_{c}^{2})^{2}}{s} \right] \right)^{1/2} + \left(\left[\frac{(\mathbf{M}_{a}^{2} - m_{a}^{2})^{2}}{s} + P_{1}(s)^{2} \right]^{1/2} \times \left[\frac{(\mathbf{M}_{c}^{2} - m_{c}^{2})^{2}}{s} + K_{1}(s) \right]^{1/2} \right)$$
(32)

We note that LLE is bigger than SLE. The absorptive part of the scattering amplitude, as noted, converges inside LLE. In fact this proof of the existence of LLE by Lehman was very crucial to write down fixed-t dispersion relations. It is to be noted, as we

³See Schweber [56] for elaborate discussions on JLD theorem.

shall see when we write dispersion relations, that the integral is over values of s from s_{th} to ∞ . Therefore, there is a LLE for every s. Moreover, the ellipse shrinks to a real line segment of t-plane in the limit $s \to \infty$. All the results summarized above follow by demanding Lorentz invariance and microcausality and other LSZ axioms except unitarity. Martin [46, 61] invoked unitarity and utilized the positivity properties of partial wave amplitudes of elastic scattering to derive stronger results. The study of analyticity in the framework, mentioned above to derive results (i) and (ii), is called linear program (procedure) since unitarity of S-matrix is not invoked until now. Martin's work was influenced by a paper of Lehmann [62]. The discussions presented so far, in a concise form, were derived for a scalar field theory which originated as follows. The original theory, to remind the reader, was considered in D = 5 which was compactified to $R^{3,1} \otimes S^1$. We have considered 4-dimensional fields $\{\phi_n\}$ to derive above results.

We need one more ingredient for writing dispersion relations as is established at present. The dispersion relations are written down by closing a contour in the complex *s*-plane. Therefore, it is necessary to know what kind of singularities lie in this plane and their locations. This questions was addresses and answered by Bross, Epstein and Glaser [63]. They considered a two body scattering amplitude, $a + b \rightarrow c + d$. They demonstrated that the (possible) complex singularities are isolated and located in finite region of *s*-plane (usually called a'potato'). They argued that there will exist a path on which scattering amplitude for $a + b \rightarrow c + d$ can be continued analytically to the u-channel physical scattering amlitude i.e. $a + \bar{d} \rightarrow c + \bar{b}$. They also noted that the size of the domain is not fixed; it depends on *t*, however. They estimated the growth property of this domain as *t* grows. The growth is $|t|^{3+\epsilon}$, $\epsilon > 0$ and the parameter is arbitrary. We mention that BEG theorem is valid for elastic as well as for inelastic processes and will be used as an useful ingredient for us. Now we shall state Martin's theorem. The fixed-t dispersion relation, without subtraction, is

$$F(s,t) = \frac{1}{\pi} \int_{s_{thr}}^{\infty} ds' \frac{ImF(s',t)}{s'-s} + \frac{1}{\pi} \int_{u_{thr}}^{\infty} du' \frac{ImF(u',t)}{u'-u}$$
(33)

when $t \in LLE$.

Martin's Theorem [61]

If the elastic amplitude (consider equal mass scattering of mass, m, for simplicity) satisfies following conditions:

1. F(s,t) satisfies fixed-t dispersion relations in s with finite number of subtractions, $N \leq 2$ with $-t_M \leq t \leq 0$ where

$$t_M = \frac{4}{s' > min(\mathbf{M_{ab}^2}, \mathbf{M_{ac}^2})} \left[P_1(s')^2 + \frac{(\mathbf{M}_a^2 - m_a^2)^2}{s'} \right]$$
(34)

Note that $t_M > 0$.

2. F(s,t) is an analytic function of s and t (in both the variables) in the neighborhood

of any \bar{s} in some interval below the physical threshold: $s_{thr} - \eta < 4m^2$ and in some neighborhood $|t| < R(\bar{s})$ of t = 0 [63, 62].

3. $A_s(s',t)$ and $A_u(u',t)$, the absorptive parts of F(s,t) from discontinuities of right hand cut, $s' > 4m^2$ and left hand cut, $u' > 4m^2$, satisfy following positivity properties. They are also holomorphic in LLE.

4. The positivity properties of $A_s(s',t)$ and $A_u(u',t)$ for $s' > 4m^2$ and $u' > 4m^2$ are

$$\left|\left(\frac{\partial}{\partial t}\right)^{n} A_{s}(s',t)\right| \leq \left(\frac{\partial}{\partial t}\right)^{n} A_{s}(s',t)|_{t=0}, \quad -4k^{2} \leq t \leq 0$$
(35)

and

$$\left|\left(\frac{\partial}{\partial t}\right)^{n} A_{u}(u',t)\right| \leq \left(\frac{\partial}{\partial t}\right)^{n} A_{u}(u',t)|_{t=0}, \quad -4k^{2} \leq t \leq 0$$
(36)

Martin proved, based under assumptions (1)-(4) that F(s,t) is analytic in a domain of quasi topological product

$$\left\{s, t \middle| s \in \text{cutplane}, s \neq (4m^2 + \rho, \rho > 0 \text{ and } s \neq 0 - t - \rho', \rho' > 0 \right.$$
(37)

$$|t| < 2R \text{ with some R}$$
 (38)

The consequence of Martin's theorem is that there exists a radius, R_M , such that the dispersion relation is valid for $|t| < R_M$. The domain of analyticity of F(s,t) is the quasi-topological product, s-t cut-plane \otimes a circle in t-plane with center at t = 0 and the radius is $|t| = R_M$.

Determination of R_M : Martin used analyticity arguments on the amplitude to determine R_M and proved that it is independent of s for large s,

$$R_M = \min[t_M, \max \frac{r_{sL}(s')}{s'}]$$
(39)

where $r_{sL}(s') = 2P_1(s')^2(x_0(s') - 1)$. Here $\mathbf{P}_1(s')$ is the c.m. momentum and its magnitude is the same for both incoming pair and outgoing pair for the case at hand. r_{sL} corresponds to the right extremity of the large Lehmann ellipse. We shall discuss the analyticity domain in the *t*-plane for inelastic scattering where Martin's theorem cannot be adopted directly since the absorptive inelastic amplitudes do not satisfy the positivity conditions (35) and (36). Therefore, a chain of arguments will be presented to study analyticity domains in *s* and in *t* planes.

4 Analyticity Properties of Inelastic Scattering Amplitudes

We study the analyticity properties of inelastic process $a+b \rightarrow c+d$ in this section. It is assumed that a and b are equal mass particles, $m_a = m_b$ and so are the two final

state particle, $m_c = m_d$. Moreover, it is also assumed that produced particles are heavier than accelerated initial colliding particles i.e. $m_c > m_a$. A point to note is that $s_{thr} \neq s_{phys}$. Consequently, the analyticity properties of inelastic reactions are to be analyzed carefully and the results derived for elastic scattering do not automatically hold good here. A few additional steps are necessary to derive various characteristics of the inelastic amplitude, $F^{ab \rightarrow cd}$. The first important point in this case, which differs from elastic case, is that the inelastic partial wave amplitudes, $f_l(s)^{ab \rightarrow cd}$ do not enjoy the positivity properties of elastic partial wave amplitude i.e. $0 \leq |f_l(s)^{ab \to ab}|^2 \leq$ Im $f_l(s)^{ab\to ab} \leq 1$. We also know that there is no analog of optical theorem for inelastic amplitudes. We recall that analyticity properties of inelastic amplitudes have been studied, in the past, in a rigorous framework [40, 41]. These authors assumed that Im F(s,t) is analytic inside an ellipse demanding certain inequality, to be satisfied by partial wave amplitudes. They had not determined the properties of their ellipse from basic principles. However, once the existence of Lehmann-Martin ellipse is proved, as $s \to \infty$, in the complex t-plane there is a domain which does not shrink to real line segment as was proved by Martin [61]. Indeed, the proof of existence of Martin's radius, R_M , is crucial in the sense that R_M is independent of s. In view of preceding remarks, the results derived by those authors [40, 41], at that juncture, were based not on strictly rigorous arguments. Subsequently, Sommer [64] carefully studied analyticity property in inelastic reaction for hadronic production processes. He obtained t-plane domain of analyticity with refinement of Martin's work [61]. We shall adopt similar line of arguments to study production of KK states in high energy collisions at LHC. Notice, however, the difference, Martin [61] was able to determine value of the radius of the circle, R_M , from first principles in terms of the experimentally observed mass. For most of the cases, it was determined to be $t_0 = 4m_{\pi}^2$. For the case under study we cannot identify the analog of R_M to an experimentally measured number and we can only present it in terms of the mass parameters of the theory. From our perspective, we determine the domain of analyticity and we write fixed-t dispersion relations for the right hand cut $F(s, t)^{ab \to cd}$. One more step is to be completed. We must determine the analyticity property of the u-channel amplitude, $F(u,t)^{a\bar{d}\to c+\bar{b}}$, to write the dispersion relation for the left hand cut. Therefore, it is necessary to go through the same procedure, as is adopted for the s-channel reaction, in order to determine the domain of holomorphy in t.

We have discussed the existence of Martin's ellipse for elastic amplitude and the domain of convergence of the absorptive amplitude ; the semi-major axis of the ellipse is $\cos\theta_0 = 1 + \frac{t_o}{2P_1(s)^2}$. We have summarized the results in the preceding section. It follows from unitarity arguments that $\sigma_t(s)^{ab} = \sigma_{el}(s)^{ab} + \sigma(s)^{ab \to cd} + all inelastic channels. Also note that <math>\operatorname{Im} f_l(s)^{ab \to ab} = |f_l(s)^{ab \to ab}|^2 + |f_l(s)^{ab \to cd}|^2 + \dots$ where (...) stand for other two body amplitudes. This is due to the fact $\operatorname{Im} f_l(s)^{elastic}$ is related to total cross section which is sum of elastic and all inelastic cross sections (when we write in terms of partial wave amplitudes). Thus we write inequality for

imaginary part of inelastic partial wave amplitude as

$$|f_l(s)^{ab \to cd}| \le \sqrt{\text{Im } f_l(s)^{ab \to ab}} \tag{40}$$

since partial wave amplitudes $f_l(s)^{ab\to cd}$ do not enjoy positivity properties anymore. We also know that, due to polynomial boundedness of scattering amplitude and convergence of partial wave expansion inside Martin ellipse, the elastic partial wave amplitude has following behavior for asymptotic energy as l tends to large values. Note that $P_l(1 + \frac{t_0}{2K_1(s)^2})$ grows exponentially since $\cos\theta > 1$ in this region. Therefore, the imaginary part of partial wave amplitude must be damped for large l when we take s to be large

$$\lim_{l \to \infty} \text{Im } f_l(s)^{ab \to ab} \le \left[\frac{1}{1 + \frac{R_M}{2P_1(s)^2} + \sqrt{\left(1 + \frac{R_M}{2P_1(s)^2} - 1\right)^2}} \right]^l$$
(41)

The inelastic amplitude $F(s,t)^{ab\to cd}$ can be expanded in partial wave expansion, taking into account the inequality satisfied by imaginary part of its partial wave (40). The partial wave expansion is

$$F(s,t)^{ab\to cd}(s,t) = \frac{\sqrt{s}}{|\mathbf{P}_1(s)|} \sum_{l=0}^{\infty} (2l+1) f_l(s)^{ab\to cd} P_l(\cos\theta)$$
(42)

Notice that the prefactor $\frac{\sqrt{s}}{|\mathbf{P}_1(s)|} \to 1$ as $s \to \infty$. We shall not generally include such constant prefactors at various places unless explicit mention of them carries important implications. We recall that for the inelastic scattering $\cos\theta = \frac{t+P_1(s)^2+K_1(s)^2}{2P_1(s)K_1(s)}$. The argument presented above is similar to the arguments of [40, 41]; however, the power of Martin's theorem and the existence of Martin'e ellipse was not recognized by them. Moreover, one of their primary goal was to rigorously study properties of inclusive processes. In fact it is not necessary to assume existence of an ellipse inside which inelastic partial wave converges. Indeed, with preceding arguments, it can be proved that $F(s,t)^{ab\to cd}$ is analytic inside an ellipse which is defined, in the *t*-plane by the condition

$$|t + (\mathbf{P}_{1}(s) - \mathbf{K}_{1}(s))^{2}| + |t + (\mathbf{P}_{1}(s) + \mathbf{K}_{1}(s))^{2}| < 4|\mathbf{P}_{1}(s)||\mathbf{K}_{1}(s)|\sqrt{1 + \frac{R_{M}}{4P_{1}(s)^{2}}}$$
(43)

We mention in passing that, for elastic process, $\mathbf{P}_1(s) = \mathbf{K}_1(s)$ and thus $(P_1(s) - K_1(s))^2 = 0$. Notice that as $s \to \infty$, the right hand side is $\frac{R_M}{4}$ and is independent of s. We have argued in Section 2 that $\text{Im } F(s,t)^{ab\to cd}$ is analytic inside LLE besides being analytic in Martin's ellipse. In the high energy limit, we can see that Martin's ellipse is bigger than LLE; however, in the low s region, the case is opposite. We noted earlier that the domains defines by SLE and LLE depend on s. Thus, while writing

dispersion relations (integral over s'-variable) and identifying domain of holomorphy in t, this aspect should be kept in mind in the sense that the domain depends on s'. Since we are considering unequal mass scattering the integration regions of s' variable is to be taken into considerations. (i) $s_{thr} \leq s' \leq s_{phys}$, (ii) $s' \geq s_{phys}$. (iii) Another region is to be considered due to the complications of unequal mass inelastic scattering i.e. masses of initial colliding (equal mass) particles producing particle (although they are of equal mass) whose masses are different from initial colliding particles (that is why s_{thr} and s_{phys} are not equal). This region is called s-critical region which has been discussed in detail by Sommer [64]. For our problem this value is determine from the relation

$$|\mathbf{K}_{1}(s)|\sqrt{\mathbf{P}_{1}(s)^{2} + \frac{R_{M}}{4}} = \zeta(s_{sc})$$
 (44)

where

$$\zeta(s) = \left[\frac{(\mathbf{M}_a^2 - m_a^2)^2}{s}\right]^{1/2} \times \left[\frac{(\mathbf{M}_c^2 - m_c^2)^2}{s}\right]^{1/2} + \left[\frac{(\mathbf{M}_a^2 - m_a^2)^2}{s} + P_1(s)^2\right]^{1/2} \times \left[\frac{(\mathbf{M}_c^2 - m_c^2)^2}{s} + K_1(s)^2\right]^{1/2}$$
(45)

A remark is in order at this stage. In case of hadronic reactions, $\mathbf{M}_a, \mathbf{M}_c, m_a$ and m_c are obtained from experimental data from data tables. Therefore, one can use numerical factors to determine the critical value. We do not have values of these mass parameters (except we may choose that m_a is mass of proton). However, as we shall discuss in the sequel, the growth of cross sections as a function of s will be bounded. We mention that for $s' > s_{cr}$ LLE is inside Martin's ellipse. Now, for region (i), $s_{thr} \leq s' \leq s_{phys}$, the domain of holomorphy is LLE. We can write fixed-t dispersion relation for $-t_M \leq t \leq -t_m$ (see the definition of t_m below). We denote this domain as \mathcal{D}_1 where

$$\mathcal{D}_1 = \bigcap_{s_{thr} \le s' \le s_{phys}} E_L(s')^{inel} \tag{46}$$

The second domain of holomorphy, \mathcal{D}_2 , is where the Lehmann ellipse is inside the Martin's ellipse. Therefore, in the $s \to \infty$ limit this complex-*t* domain does not degenerate to a real line

$$\mathcal{D}_2 = \bigcap_{s_{cr} \le s' < \infty} E_{Martin}(s')^{inel} \tag{47}$$

This is defined when t lies in the interval $-t_M \leq t \leq t_m$ (see more discussions below). Moreover, in this s' region, each Martin's ellipse, $E_{mart}(s')$ is larger than corresponding LLE, $E_L(s')$. Furthermore, for asymptotic s', $R_M(s) \rightarrow R_M$ where R_M is s-independent. As asserted above, non of the Martin's ellipses collapse to real *t*-axis and we conclude the \mathcal{D}_2 is a complex domain. Let us look at the interval $s_{phys} < s' \leq s_{cr}$ and denote the corresponding domain as \mathcal{D}_3 . It is given by

$$\mathcal{D}_3 = \bigcap_{s_{phys} < s' \le s_{cr}} E_L^{inel} \tag{48}$$

Here $-t_M \leq t \leq -t_m$. Notice that s' is restricted to a finite interval. Consequently, the complex domain in t-plane is not permitted to shrink real t line. We note that it is necessary to find the two extreme values of real t within which interval t lies; we remit t to go out of physically allowed values. However, it is restricted to above interval. The delicate issues arising for unequal mass inelastic scatterings is discussed briefly in what follows.

$$t_{m} = \max\left\{\max_{s \ge \mathbf{M}_{ab}^{2}}\left[2P_{1}(s)^{2} + 2\left(\frac{m_{a}^{2} - m_{c}^{2}}{4s}\right) - 2\mathcal{Z}_{L}(s)\right], \\ \max_{u \ge \mathbf{M}_{ac}^{2}}\left[\bar{P}_{1}(u)^{2} + \bar{K}_{1}(u)^{2} + 2\left(\frac{m_{a}^{2} - m_{c}^{2}}{4u}\right) - 2\mathcal{Z}_{L}(u)\right]\right\}$$
(49)

where $\mathcal{Z}_L(u)$ is an expression similar to $\mathcal{Z}(s)$ (32) and directly related to the *u*-channel process.

$$\mathcal{Z}_{L}(u) = \left[\frac{(\mathbf{M}_{a}^{2} - m_{a}^{2})(\mathbf{M}_{c}^{2} - m_{c}^{2})}{u - (\mathbf{M}_{a}^{2} - \mathbf{M}_{c}^{2})}\right] + \left[\frac{(\mathbf{M}_{a}^{2} - m_{a}^{2})(\mathbf{M}_{c}^{2} - m_{c}^{2})}{u - (\mathbf{M}_{a}^{2} - \mathbf{M}_{c}^{2})} + \tilde{P}_{1}(u)^{2}\right]^{1/2} \times \left[\frac{(\mathbf{M}_{a}^{2} - m_{a}^{2})(\mathbf{M}_{c}^{2} - m_{c}^{2})}{u - (\mathbf{M}_{a}^{2} - \mathbf{M}_{c}^{2})} + \tilde{K}_{1}(u)^{2}\right]^{1/2}$$
(50)

where $\tilde{P}_1(u)$ and $\tilde{K}_1(u)$ are the c.m momenta of initial pair of states and final pair of states in the u-channel reaction. These formulae, (46) and (47) arise in case of unequal mass inelastic scattering and the corresponding formulae for elastic case assume rather simple form. A few important comments deserve mention at this juncture. (i) One has to be careful to check that the c.m. momenta for direct channel and c.m. momenta for crossed channel process do not become complex. In such a situation there will be a gap in s where we are unable to write dispersion relations. Same argument is valid for the cross channel. (ii) A situation might arise when $\mathcal{Z}_L(s)$ might become complex and/or $\mathcal{Z}_L(u)$ might be complex in some region. In such a situation the Lehmann would not be defined (i.e. will not exist) in those intervals of s and/or u. In absence of Lehmann ellipse for some interval of s and/or u we cannot write fixed-t dispersion relations. It is necessary that $t_M > t_m$. We record these comments at this stage to draw attention of the reader to the delicate issues to be addressed, in the context of study of analyticity properties, for inelastic unequal mass scattering in general. We also note that, in the study of hadronic scattering, the masses of particles, involved in scattering, are known. Therefore, the values obtained from experiments are utilized to chart the domains. For the case at hand, it is not possible to actually display the forbidden domains. It is to be noted that, in the high energy

limit, when s and u take asymptotic values, the formulae are considerably simplified. The purpose of this detour, in our discussions, is that in the works of Russian group [40, 41] these delicate arguments were not presented. Therefore, there were, to some extent additional assumptions in their work; moreover, domains of analyticity were not charted out carefully for unequal mass case involving inelastic reactions. Those aspects have been taken into account in analyzing domains of holomorphy with care in the present investigation.

We have identified the domains of analyticity in various values of s' in order to write down dispersion relations. Our focus has been to identify the holomorphic domain for writing a dispersion relation associated with the right hand cut, the *s*channel reaction, $a + b \rightarrow c + d$; it is

$$\mathcal{I}_R = \frac{1}{\pi} \int_{s_{thr}}^{\infty} ds' \frac{\text{Im } F(s', t)^{ab \to cd}}{s' - s}$$
(51)

This is an unsubtracted dispersion relation. The scattering amplitude is polynomially bounded in s for t lying within Lehmann-Matrin ellipse i.e. $|F(s,t)^{ab\to cd}| \leq s^N$ and $N \leq 2$ as follows from Jin-Martin theorem [65]. Therefore, if at all we need to write a subtracted dispersion relation, it will need at most two subtractions. Moreover the fixed-t dispersion relation is valid in the domain of complex t-plane given by

$$\mathcal{D}_{inel}^{R} = \left[\bigcap_{s_{thr} \le s' \le s_{phys}} E_{L}^{inel}(s') \right] \bigcap_{s' \ge s_{phys}} \left[\bigcap \left(E_{M}^{inel}(R, s') \right) \cup E_{L}^{inel}(s') \right]$$
(52)

Here R stands for right hand side of the threshold where we identify domain of holomorphicity in complex t-plane. We have argued that the union of complex domains is such that each of them does not shrink to real t-line. In order to write dispersion relation for $F(s,t)^{ab\to cd}$, we have to consider properties of the *u*-channel amplitude. Therefore, we consider the reaction $a + \bar{c} \rightarrow d + \bar{b}$. Thus scattering of two particles of mass (m_a, m_c) to (m_c, m_a) is to be considered, (note $m_a = m_b$ and $m_c = m_d$). This process is practically, as far as kinematics is concerned, like elastic scattering of unequal mass particles. We recall that masses of anti particles (b, d) respectively are (m_a, m_c) . We can determine the semi-major axis of the u-channel Martin ellipse and SLE and LLE in this case. It is obvious that we do not encounter any technical difficulties in determining the t-plane analyticity domain for the crossed channel reaction. Note that the magnitudes of c.m. momenta of initial pair and final pair are the same. The mathematical expressions are exactly the same as those for elastic process. Consequently, calculations are simplified to identify the domain of holomorphicity of t. The unsubtracted dispersion relation for the inelastic amplitude is (including contributions from right hand cut and left hand cut) is

$$F(s,t)^{ab\to cd} = \frac{1}{\pi} \int_{s_{thr}}^{\infty} ds' \frac{\text{Im } F(s,t)^{ab\to cd}}{s'-s} + \frac{1}{\pi} \int_{u_{thr}}^{\infty} du' \frac{\text{Im } F(u',t)^{a\bar{c}\to d\bar{b}}}{u'-u}$$
(53)

Bounds on Cross Sections for Inelastic Reaction $F(s,t)^{ab \rightarrow cd}$

We proceed to derive bounds on cross sections utilizing the analyticity properties of the amplitudes derived thus far. The first point to note is that we are unable to derive analog of the Froissart bound for inelastic total cross sections $\sigma(s)_t^{inel}$ from basic principles. The reason is that the absorptive amplitude, Im $F(s,t)^{ab\to cd}$ of two body reaction lacks the positivity properties of its elastic counter part. Thus the insight gained in the preceding discussions will be utilized to derive bounds of our interests. Furthermore, we note that $F(s,t)^{ab \to cd}$ is polynomially bounded, s, for fixed t. Thus it is possible to derive upper bounds on differential cross sections. Logunov and collaborators [40, 41] derived such upper bounds; however, an unknown constant appeared in such a bound. They did not employ the full power of the analyticity properties of amplitude in s and t variables. Singh and Roy [43]. used Lagrangian multiplies technique to derive very powerful bound. One of their important results was that the bound on elastic differential cross section was obtained without any unknown constant. Moreover, the bound is expressed in terms of experimentally measured quantities such as σ_t and σ_t^{el} . Indeed the bound on, differential elastic cross section, $\frac{d\sigma^{el}}{d\Omega}$, was tested against experimental data over a wide range of high energy scattering experiments and no violations were found. Furthermore, all ingredients that went into deduce the bound have been proved, in the past, from axiomatic field theory. In case of inelastic scattering not all those results hold good. We shall utilize the rigorous results reported here and present them which will not be as strong as on the elastic cross sections which is not a surprise. Moreover, one of our limitations is that, at present, we are unable to test the bounds against experimental data. We briefly recollect the technique employed by Singh and Roy [43] for obtaining bound $\frac{d\sigma^{inel}}{d\Omega}$. We collect the necessary formulae to be useful in the sequel. (1). The elastic differential cross section is expressed in terms of partial wave expan-

$$\frac{d\sigma^{el}}{d\Omega} = \sum_{l=0}^{l=\infty} (2l+1)|f_l(s)^{el}|^2$$
(54)

An upper bound is to be derived with following conditions which are obtained already from axiomatic field theory. The conditions are (I)

$$\sigma_t(s) = \frac{1}{s} \sum_{l=0}^{l=L} (2l+1) \text{Im } f_l(s)^{el} P_l(\cos\theta = 0)$$
(55)

 $L \sim \sqrt{slog} \ s$

sions

(II) $|F(s,t)|^2$ satisfies following inequality in the physical region of $\cos\theta$

$$|F(s,t)|^{2} \leq \sum_{l=0}^{l=L} (2l+1)|f_{l}(s)^{el}|^{2}|P_{l}(1+\frac{2t}{\mathbf{P}_{1}^{2}})|$$

$$\leq \sum_{l=0}^{l=L} (2l+1) \operatorname{Im} f_{l}(s)^{el}|P_{l}(1+\frac{2t}{\mathbf{P}_{1}^{2}})| < s^{N}, N \leq 2$$
(56)

It is known from convergence of partial wave expansion, inside Lehmann-Martin ellipse, because $cos\theta_0 > 1$, there is some cutoff value of l = L, $L \sim \sqrt{slog} s$, beyond which sum of all partial wave amplitudes, when we obtain expression for $\sigma_t(s)$, beyond $l > L_0$ is subdominant. The problem of deriving the desired upper bound is reduced to using (I) and (II) as constraints. This is accomplished by appealing to Lagrangian multiplier method. Thus, the upper bound on differential cross section is expressed in terms of $\sigma_t^{el}(s)$,

and (III)

$$\frac{\sqrt{s}}{R_M} \sum_{l=0}^{\infty} (2l+1) |f_l(s)^{el}|^2 P_l(1 + \frac{t_0 - \epsilon}{\mathbf{P}_1^2}) \le \text{Im } F(s, t_0 - \epsilon) < s^2$$
(57)

Here t_0 is the right extremity of the LLE. Note that the argument of P_l is greater than 1 hence we did not write $|P_l(x)|$ in the above equation since x > 1. Thus $|f_l(s)|$ was bounded and the bound involved $P_L^2(\cos\theta)$ where $-1 \leq \cos\theta \leq +1$ and square of its derivative of Legendre polynomial.

For the inelastic case, (Singh and Roy studied some hadronic reactions also) we have derived inequalities for inelastic partial wave amplitudes from unitarity considerations. Moreover, the inelastic total cross section, $\sigma_t(s)^{ab\to cd}$ gives an analogous expression as (54). However, there is no optical theorem for inelastic reaction. Our results are, however, not as strong as those of [43]. Note that $t_0 = 4m_{\pi}^2$ for most hadronic processes [43]. The inequality that follows from unitarity condition is

$$|f_l(s)^{ab \to cd}|^2 \le \operatorname{Im} \, f_l(s)^{ab \to ab} \tag{58}$$

and inelastic total cross section is bounded as

$$\sigma_t^{ab \to cd} \le \frac{4\pi}{T_0} (ln \frac{s}{s_0})^2 \tag{59}$$

Here $\sigma_t^{ab\to cd}$ contains undetermined parameters T_0 and s_0 . We have explained why we cannot provide a value of T_0 . Moreover, the polynomial boundedness of imaginary part of inelastic process is indirectly used from polynomial boundedness of absorptive part elastic amplitude (which is the Jin-Martin bound [65]). We get a bound nevertheless for differential cross sections at high energies in forward directions i.e. $\theta = 0$

and in nonforward directions, $\theta \neq 0, \pi$ as given below.

(i) Forward scattering:

$$\frac{d\sigma^{ab\to cd}}{d\Omega}|_{\theta=0} \le (\frac{s}{16\pi T_0^2})[\log\frac{s}{s_0}]^4 \tag{60}$$

Note that T_0 corresponds to threshold mass of *t*-channel reaction (= $4m_{\pi}^2$ usual hadronic reaction): for $a + \bar{d} \rightarrow \bar{b} + c$ in *our hadronic* reactions. Moreover, differential cross section shows a peak in the forward direction.

(ii) Nonforward scattering, $\theta \neq 0$ or π

$$\frac{d\sigma^{ab\to cd}}{d\Omega} \le \left(\frac{1}{4\pi T_0^{3/2}}\right) \frac{1}{\sqrt{ssin\theta}} \left[\log\frac{s}{s_0}\right]^3 \tag{61}$$

For small angles, we may approximate $\sin\theta \approx \theta$. We discuss another type of inelastic reaction which might be of interests. Let us consider a reaction where only a single particle is observed experimentally. Thus the process, known as *inclusive reaction* is $a+b \rightarrow c+X$ where X is a single particle or multiparticle state and is not subject to observation. Feynman [35] coined this word in the context of parton model to analyze attributes of hadronic scattering. He argued that inclusive cross sections will exhibit scaling phenomena in various kinematical regions. Those feature were experimentally observed. The most familiar example of an inclusive reaction is deep inelastic e + pscattering where energy of the scattered electron is measured at different angles. The structure functions exhibit Feynman scaling. Our goal is not to discuss inclusive reaction and scaling of its inclusive cross sections. It is worthwhile to note that inclusive cross section measurements have some advantages over exclusive reactions such as an inelastic process like $a + b \rightarrow c + d$ since inclusive reactions have several final state channels available. Moreover, if there is conservation of a charge, when we search for such a charged particle, c, the unobserved inclusive state, X, must carry equal and opposite charge of the experimentally observed particle, c. Another point deserves mention here. For two body scattering, the total cross section is $a+b \rightarrow X$ is related to the imaginary part of the forward elastic cross section, Im $F(s, t = 0)^{ab \to ab}$. from the optical theorem.

A generalized optical theorem was derived by Mueller [66] for inclusive reaction. The differential cross section for $a + b \rightarrow c + X$ is related to the imaginary part of a forward scattering amplitude, Im $F^{(a+b+\bar{c})\rightarrow(a+b+\bar{c})}(s, M_X^2, t)$, where \bar{c} is the antiparticle of c. The forward six point amplitude depends on three kinematical variables s, M_X^2 and t where t is the momentum transfer squared between particle a and c i.e. $t = (p_a - p_c)^2$. Note that the integration of the differential cross section $(E_c \frac{d\sigma^{ab\rightarrow cX}}{d^3p_c})$ is not just $\sigma_t(s)$ but it is $\sigma_t(s) < n_c >$ where $< n_c >$ is the average multiplicity of outgoing particle c. We may consider the inclusive reaction to be a two body inelastic reaction, $a + b \rightarrow c + X$ where X could be a single particle or collection of many particles where p_X is sum of four momentum of all unobserved outgoing particles. Therefore, $p_a + p_b = p_c + p_X$ is the energy momentum conserving relation and $t = (p_a - p_c)^2$. These kinematical aspects were considered by [40, 41]. Moreover, they extended their arguments for inelastic two body collision case to the case of inclusive reaction. However, as noted in preceding discussions, these authors [40, 41] did not take into account the delicate issues associated with unequal mass inelastic scattering and analyticity in the complex t variable. We have discussed how to identify the domain of holomorphy in complex t-plane for two body unequal mass inelastic scattering earlier. To remind the reader, the real value of t is bounded as $-t_M \leq t \leq -t_m$ and we have given expression for t_m which is not same as that for equal mass case. Second point to recall is the complex-t domain is expressed as union and intersections of various complex domains as noted in (52) - see the expression for $\mathcal{D}_R^{inelastic}$. Those arguments go through for the inclusive reactions as well.

The inclusive forward differential cross section is

$$\left(\frac{d\sigma^{ab\to cX}}{d\Omega_c}\right)_{\theta_c=0} \le \left(\frac{s}{4T_0^2}\right)\left[\log\left(\frac{s}{s_0}\right)\right]^4 \tag{62}$$

Note that the bound is an improvement of [40, 41] result since the prefactor is fixed; whereas there was an arbitrary constant as prefactor in their analysis. Moreover, the prefactor can only be determined after invoking Martin's theorem. Our bound is weaker than that of [43] since we have not utilized the Lagrange multiplier technique as there are no experimental data for our case. If we sum over all states X then the bound for differential cross section is

$$\sum_{X} \frac{d\sigma^{ac \to cX}}{d\Omega_c}(s, \cos\theta) \le \frac{1}{\pi T_0^{3/2}} \frac{\sqrt{s}}{\sin\theta_c} [\log(\frac{s}{s_0})]^3 < n_c(s)(\theta_c) >$$
(63)

where $\langle n_c(s) \rangle$ is average multiplicity of particle c. We mention in passing that derivation of upper bound on average multiplicity in a general setting has been discussed in [67]. We note that inclusive reactions to detect production of 'exotic' states may be a more optimistic scenario.

5 Analyticity Properties of Inelastic Scattering Amplitude for D = 5, Decompactified Theory

This section contains analyticity properties for D = 5 theory where all spatial dimensions are noncompact. In case of a theory where it has only one field, in D = 4, according to UED hypothesis the five dimensional theory also has only one field. Therefore, for a single specie of field in D = 5, the four point amplitude is an elastic amplitude. We shall recall the essential results for this reaction in the first place. In case of multiple species of fields in a D = 4 theory, the higher dimensional theory (as per UED hypothesis) will be endowed with those many fields. In this case inelastic four point inelastic amplitudes will be permitted; moreover, in general, these will be unequal mass scatterings. We allude to this aspect in some detail in this section. We

recall that we discussed analyticity properties of inelastic amplitude rigorously in the previous section. Next, based on those properties we derived bounds on differential cross section and inclusive cross sections. We recapitulate essential results from [39] for D = 5 theory with single type of field. Let the 4-point amplitude be denoted by $F^{5}(k_{1}k_{2}, p_{1}p_{2})$ where (p_{1}, p_{2}) are momenta of incoming particles and the corresponding 'in' state is $|p_1p_2|$ in >. Similarly, (k_1, k_2) are momenta of outgoing particles and 'out' state is $|k_1k_2 out >$. We follow a convention where masses of particles, in D = 5are also denoted by m_a, m_b, m_c etc. These masses becomes zero modes of compactified $R^{3,1} \otimes S^1$ theory. Indeed, if we appeal to UED hypothesis, the masses appearing in D = 5 (noncompact) theory will correspond to zero modes of compactified theory. Whenever, we need to specify amplitudes, distributions and other entities of interests in D = 5 theory we shall explicitly use subscripts/superscripts, '5'. The expression for $F^5(k_1k_2, p_1p_2)$ is derived from LSZ reduction technique. We choose to reduce the state with momenta (k_1, k_2) . We refer the reader to our paper [39] for detail derivation of the amplitude following LSZ technique. The sequence of arguments are that the two fields in 'out' state are reduced. The operation of KG operators on the interacting fields leads to R product of two source currents associated with these two fields besides derivatives of δ -functions. The presence of such delta functions do not affect analyticity properties of scattering amplitudes as is well known. We do not intend to repeat these details here. For D = 5 case, the amplitude assumes the following form.

$$F^{5}(k_{1}k_{2}, p_{1}p_{2}) = -\int d^{5}x d^{5}y e^{i(k_{1}.x+k_{2}.y)} < 0|RJ^{5}(x)_{c}J^{5}(y)_{d}|p_{1}p_{2} >$$
(64)

Here k_1 and k_2 are reduced. J_5 is the source current and we have used subscripts c and d for later conveniences as will be evident soon. In order to study analyticity properties, it is essential to define three distributions. This is recognized from the fact that KG operators act on R-product of interacting fields $R\phi_c(x)\phi_d(y)$. The operation of KG operators on the interacting fields leads to R product of two source currents associated with these two fields: $(\Box_x + m_c^2)(\Box_y + m_d^2)R\phi_c(x)\phi_d(y) = RJ_c^5(x)J_d^5(y)$ and in addition finite number of derivatives of δ -functions are present on the right hand side of this expression as discussed earlier. Here $\Box + m^2$ stands for five dimensional KG operator. In a local field theory only finite number of derivatives of δ -function appear. Their presence does not affection analyticity properties of the amplitude for following reasons. Let us consider derivative of the δ -functions (which is define in terms coordinate space variables. Then polynomials of momenta will appear; however, scattering amplitude is a function of Lorentz invariant variables (Mandelstam variables: s, t and u. Thus polynomials in s, t and u will appear. With this argument, we qualitatively argue that scattering amplitudes are polynomially bounded. The presence of of such delta functions do not affect analyticity properties of scattering amplitudes as is well known. In such cases, we write subtracted dispersion relations. We define following matrix elements VEV of R-products products of operators which are distributions as has been discussed earlier. However, we derive spectral representation to study support properties without going into all details (see [39]). We shall recall the previous results for elastic amplitude and then mention how analyticity properties inelastic four point amplitude are analyzed.

$$F_R^5(q) = \int d^5 z e^{iq.z} \theta(z_0) < P_f R[J_l^5(z/2), J_m^5(-z/2)] |P_i\rangle$$
(65)

where $|P_i\rangle$ and $|P_f\rangle$ are arbitrary momentum states.

$$F_A^5(q) = \int d^5 z e^{iq.z} \theta(-z_0) < P_f R[J_l^5(z/2), J_m^5(-z/2)] |P_i\rangle$$
(66)

and

$$F_C^5(q) = \int d^5 z e^{iq.z} < P_f R[J_l^5(z/2), J_m^5(-z/2)]|P_i >$$
(67)

Moreover, note that

$$F_C^5(q) = F_R^5(q) - F_A^5(q)$$
(68)

Note also that, as we have noted earlier, $F_C^5(q) = 0$ when both $F_R^5(q)$ and $F_A^5(q)$ vanish. We introduce a complete set of states $\sum_n |p_n, \alpha_n| > < p_n, \alpha_n| = 1$ and $\sum_{n'} |p_{n'}\beta_{n'}| > < p_{n'}\beta_{n'}| = 1$. Here α_n and $\beta_{n'}$ stand for all the quantum numbers permitted according to conservation laws of the theory. Note that unlike the case of compactified theory where intermediate states also include KK towers, here we have no such considerations. For the D = 5 theory, the intermediate states, $|p_n, \alpha_n >$ and $|\alpha_{n'}\beta_{n'} >$ are physical states, i.e. $p_n \ge 0$, $p_n^0 \ge 0$ and $p_{n'}^2 \ge 0$, $p_{n'}^0 \ge 0$. The spectral representation for F_C^5 , after opening up of the current commutator consists of two terms, $A_s^5(q)$ and A_u^5 which are absorptive parts of s-channel and u-channel amplitudes respectively. These are

$$2A_s^5 = \Sigma_{n'} < P_f |J_m^5|(0)| p_{n'} = \frac{1}{2}(P_f + P_i) + q, \beta_{n'} > < \beta_{n'}, p_{n'} = \frac{1}{2}(P_f + P_i) + q|J_l^5(0)|P_i > (69)$$

and

$$2A_u^5 = \Sigma_n < P_f |J_l^5(0)| p_n = \frac{1}{2} (P_f + P_i) - q, \alpha_n > <\alpha_n, p_n = \frac{1}{2} (P_f + P_i) - q |J_m^5(0)| P_i > (70)$$

The arguments of Section 3 can be appropriately rephrased so that the conditions $F_R^5(q) = 0$ and $F_A^5(q) = 0$ hold simultaneously in unphysical regions. Thus $F_C^5(q) = 0$ in this domain. The JLD representation for D > 4 has been obtained in [39]. Finally, the existence of SLE and SLE has been proved my us.

Now we summarize our results for determination of Lehmann ellipses in D = 5 dimensional theory for scattering of unequal mass particles. This analysis is relevant for the case where two or more species of fields are present i.e. the case of a UED model which would have field contents of D = 4 and they are promoted to higher

dimensions. A detail discussion of derivation of Lehmann ellipses have been presented by us in Section III of our paper [39]. We consider inelastic process $a + b \rightarrow c + d$ with $m_a = m_b$ and $m_c = m_d$. Furthermore, \mathbf{P}_1 and \mathbf{K}_1 represent, respectively, the c.m. momenta of initial (incoming particles) and outgoing (final) particles. The small Lehmann Ellipse is parametrized as SLE

$$E_{SLE}^{ab \to cd}: : |t + (P_1 - K_1)^2 + |t + (P_1 + K_1)^2| < 4|\mathbf{P}_1|\mathbf{K}_1|x_0(s)$$
(71)

where $m_a = m_b$ and $m_c = m_d$, for the case at hand. Furthermore,

$$x_0 = \left[1 + \frac{(\mathbf{M}_a^2 - m_a^2)(\mathbf{M}_c^2 - m_c)^2}{K_1(s)^2 s}\right]^{1/2}$$
(72)

We are considering a D = 5 theory. Here \mathbf{M}_a corresponds to two or more particle states states which which have same quantum number as particle 'a'. A similar definition holds \mathbf{M}_c . We have used $\mathbf{M}_a = \mathbf{M}_b$ and $\mathbf{M}_c = \mathbf{M}_d$ in arriving at the above formula. Note that $x_0 > 1$.

(ii) LLE

$$E_{LLE}^{ab \to cd} : |t + (P_1 - K_1)^2 + |t + (P_1 + K_1)^2| < 4\mathbf{Z}_L(s)$$
(73)

where $\mathcal{Z}_L(s)$ is given by

$$\mathbf{Z}_{L}(s) = \left(\left[\frac{(\mathbf{M}_{a}^{2} - m_{a}^{2})^{2}}{s} \right] \times \left(\left[\frac{(\mathbf{M}_{c}^{2} - m_{c}^{2})^{2}}{s} \right] \right)^{1/2} + \left(\left[\frac{(\mathbf{M}_{a}^{2} - m_{a}^{2})^{2}}{s} + P_{1}(s)^{2} \right]^{1/2} \times \left[\frac{(\mathbf{M}_{c}^{2} - m_{c}^{2})^{2}}{s} + K_{1}(s) \right]^{1/2} \right)$$
(74)

Here $\mathbf{Z}_L(s)$ is a function of kinematical variables and masses which are associated with D = 5 theory. We repeat that in case of a D = 4 theory, with several field contents, when we adopt UED hypothesis all those fields are defined in D = 5. Note that upon S^1 compactification, when the spacetime geometry is $R^{3,1} \otimes S^1$, the zero modes of the manifold are identified with the field contents of D = 4 theory which was promoted to D = 5 theory. Thus, now the theory with S^1 compactification will have KK towers. In summary, we have identified the spectrum of UED model and we are able to discuss inelastic reactions in this scenario. The purpose of this investigation (in D = 5) is point out that the spectral representation is nonperturbative in character. Thus it accounts for all quantum corrections as is the characteristic of LSZ approach. It is pertinent to cite the example of derivation of Froissart bound and host of rigorous bounds in hadronic scattering. Historically, the rigorous proof of Froissart does not rely on any perturbation theoretic approach. Of all the parameters; there is one undermined parameter, s_0 , in Froissart bound. All other parameters are physically measured quantities. This bound has been tested against experiment (for D = 4

case). The same argument holds for derivation of bound on slope slope of diffraction peak and some of the lower bounds.

It remains to prove crossing property of four point function. We shall present, for D > 4, a general argument which is only valid for four point amplitude. This result is adequate to prove analyticity of four point scattering amplitude. We may assign following momentum configurations, without loss of generality, to the four particles scattering process. Note that the expression for Lehmann ellipses is defined in terms of Lorentz invariant kinematical variables. We choose a frame where the momenta are given following kinematical configurations.

$$p_a = (p_a^0, 0, 0, 0, 0), \ p_b = (p_b^0, p_b^1, 0, 0.0), \ p_c = (p_c^0, p_c^1, p_c^2, 0, 0) \ p_d = (p_d^0, p_d^1, p_d^2, p_d^3, 0)(75)$$

Note that, with the above assignment of five momenta for the external particles, a, b, c, d, we work in a four dimensional submanifold of the five dimensional momentum space. The four point scattering amplitude is Lorentz invariant and it depends on two Mandelstam variables. Therefore, we are in a position to invoke theorem of Bros-Epstein-Glaser [63] to prove crossing for four point function. We cannot prove crossing for N-point functions, N > 4 in D >, by invoking this argument; moreover, it remains a challenge to prove crossing for the general case of N-point functions.

The analyticity properties of four point amplitude, for elastic processes, has been established rigorously for higher dimensional spacetime [39] already. Moreover, analog of Martin's theorem was proved in the above article. Consequently fixed-t dispersion relations have been written down for the amplitude when t lies inside Lehmann-Martin ellipse. We argue that our results on analyticity properties of scattering amplitude derived in Sections 3 and 4 will go through [39]. However, there is a distinction. If we have a single field in D = 5 then the four point amplitude corresponds to elastic process. Thus inessential complication for inelastic reaction do not appear. On the other hand if D = 5 theory is endowed with several fields in accordance with UED hypothesis then four point amplitude describes elastic as well inelastic reaction; indeed allows provisions for unequal mass scattering. Note that for 4-point amplitude, describing inelastic reactions in D = 5, is not so complicated when we study its analyticity. The reason is that, as we have argued, it is possible to reduce the problem to scattering in D = 4 spacetime which is special to four point amplitude. In view of preceding remarks, we can use the results derived in previous sections. However, there are certain differences which we shall point out at the appropriate juncture.

We can write down fixed-t unsubtracted dispersion relations; if the dispersion integrals do not converge we can implement subtraction which will have finite number of terms. We recall that the amplitude is polynomially bounded in s (see [39]). Thus the dispersion relation is

$$F_5(s,t)^{(ab\to cd)} = \frac{1}{\pi} \int_{s_{thr}}^{\infty} \frac{ds' \text{Im } F_5^{(ab\to cd)}(s,t)}{(s'-s)} + \frac{1}{\pi} \int_{u_{thr}}^{\infty} \frac{du' \text{Im } F_5^{(a\bar{c}\to d\bar{b})}(u',t)}{(u'-u)}$$
(76)

Now we proceed to discuss bounds on near forward amplitude and near forward differential cross section at asymptotic energies. We consider a five dimensional theory which accommodates several species of field; thus inelastic reactions are allowed. The case of a single scalar field has been investigated thoroughly in [39]. The, domain of *t*-plain analyticity is the Large Lehmenn Ellipse (LLE). Let us recall that the amplitude is expanded in partial waves with Geggenbauer polynomials, $C_l^{\lambda}(1 + \frac{2t}{s})$, as basis functions [39].

$$F_5^{(\lambda,ab\to cd)}(s,t) = \text{Const} \ \Sigma_{l=0}^{\infty}(l+\lambda)f_l^{(\lambda,ab\to cd)}(s)C_l^{\lambda}(1+\frac{2t}{s})$$
(77)

where $\lambda = \frac{1}{2}(D-3)$, D is number of spacetime dimensions and $\lambda = 1$ for D = 5. Note that, for inelastic reactions, we are unable to utilize positivity properties of partial wave amplitudes, $f_l^{\lambda ab \to cd}$. We have derived a bound [68] on $F^{\lambda}(s,t)$ for $|t| < T_0$ where T_0 is right extremity of LLE.

$$|F^{(\lambda,ab\to cd)}(s,t)| \le A_2 (\frac{1}{|t|})^{1/2(1+\lambda)} (\frac{1}{T_0})^{\lambda/2} s^{(1+(N-1))\sqrt{\frac{|t|}{T_0}}} (\log s/s_0)^{\lambda}$$
(78)

Here A_2 is a constant, independent of $s, t; s_0$, a dimensionful constant is introduced to make argument of log dimensionless in the above equation. s_0 cannot be determined from first principle (as is the case in D = 4 for Froissart bound). The appearance of T_0 and its detrimination deserves some discussion. It is the right extremity of the LLE. Intuitively, T_0 is related to the lowest two particle threshold in the *t*-channel i.e. mass squared of lightest two particle state. For D = 4, in the case of hadronic collision, $T_0 = t_0 = 4m_{\pi}^2$, first determined by Martin. In case of production of exotic states in D = 5 it will be threshold for t-channel cut. Thus value of T_0 cannot be determined. So is the case for S^1 compactified theory in four dimensions. T_0 is an independent free parameter and it determined from the value of threshold for the crossed t-channel reaction: $p + K_1 \rightarrow p + K_2$ from the direct channel process $p + K_1 \rightarrow p + K_2$. Here K_1, K_2 are respectively anti-particles of K_1 and K_2 . Let take $m_p \approx 1$ GeV and $m_{K_1} \approx 1$ TeV equal to mass of m_{K_2} as a ball park range. We note that the *t*-channel cut begings at $t_{th} = 4m_{K^2}$, if we ignore mass of proton compared to m_K which is order of TeV in LRC scheme. We have discussed the important role of t_{th} in derivation of Lehmann ellepses. Therefore, $T_0 = 4m_K^2$ which is the analog of $t_0 = 4m_\pi^2$ is strong interactions, appearing as the prefactor of Froissart bound. For the case at hand T_0 is not determined as a number from experimental data. Then $T_0 = (m_p + m_{K_1})^2$ and is order of 1 TeV. Moreover, $m_{K_2}^2 \approx \frac{n^2}{R}, n \in \mathbb{Z}$, where R is compactification. Therefore, the mass scale is related to compactification radius as is expected. We have introduced s_0 so that argument of log is dimensionless; s_0 cannot be determined from first principle. We have emphasized that due to polynomial boundedness and convergence of partial wave expansion in a larger complex domain the partial wave amplitudes (rather absorptive part) are cut off beyond certain L_0 exponentially by cut off by logs. We have proved L - 0 is independent of D. In order to derive bound on near forward inelastic scattering amplitude for the five dimensional theory, Note that forward scattering amplitude is bounded as

$$|F_5(s,t=0)|^2 \le \text{const} \ (\frac{1}{2\sqrt{R_M}})^3 (\log \frac{s}{s_0})^3 (1 + (R_M/Const)^2)^{-1/2}$$
(79)

The inelastic total cross section is bounded as

$$\sigma_{5t}^{ab \to cd} \le \frac{4\pi}{T_0} (\log \frac{s}{s_0})^3 \tag{80}$$

Here R_M is the radius of the Martin's circle. Note that this bound is weaker one in the sense that it is $(\log s)^3$ compared to compactified theory in D = 4 where cross section is bounded by $(\log \frac{s}{s_0})^2$. Thus, in what follows, from a phenomenological perspective in the accessible LHC energies the bound for D = 5 theory is higher by a factor of $\log \frac{s}{s_0}$. This 'weaker' bound trend continues to appear for other measurable parameters. If the data exhibits violations of D = 4 bounds (for compactified case) there will be reasons to argue that an extra dimension is decompactified.

For D = 5, differential cross section is bounded as (θ small)

$$\frac{d\sigma}{d\Omega} \le \frac{\text{const}}{T_0^{3/2}\sqrt{s\theta}} (\log \frac{s}{s_0})^4 \tag{81}$$

We note that the differential cross section, $\frac{d\sigma}{d\Omega}$, after approximating Gegenbauer polynomial for small scattering angle in near forward direction, has a weaker bound in asymptotic energies. The method adopted in Section 4, for D = 4 (with S^1 compactification) is applicable to derive similar to bounds in case D = 5 theory. Therefore, we shall not repeat those computations here. These bounds will also be weaker compared to those of Section 4 so far as the energy dependence is concerned.

6 Summary and Discussions

We considered production of Kalulza-Klein states, arising from S^1 compactification of a D = 5 theory. In the LRC scenario, the excited KK states might have masses in the range of TeV and might be discovered at CERN-LHC collider. We have discussed some aspects of the production of the exhotic states in pp collision in the introduction section. We have assumed that KK charge is conserved in the collision of high energy protons and consequently, final two body KK states carry equal and opposite KK charge. Moreover, baryon number is conserved. There are several phenomenological models which envisage detail properties of the idea of LRC paradigm and suggest that experimentalists might succeed in observing such states. We derived bounds on cross section starting from general principles of relativistic quantum field theories.




Figure 1: Depicts bound $\sigma_t^{ab\to cd}$ for the S^1 compactified theory. It is plotted against energy.




Figure 2: This shows the bound for inelastic cross section, $\sigma_{t5}^{ab\to cd}$ for decompactified theory.




Figure 3: This shows bound for forward differential cross section for the S^1 compactified theory.




Figure 4: Depicts bounds on near forward differential cross sections for $ab \rightarrow cd$ at angles $\theta = 2^0, 3^0$ and $\theta = 5^0$.

These bounds might be useful to them. We envisaged a scenario where the theory does not have any compact spatial dimension. In other words, all spatial dimensions are noncompact and the geometry is $R^{4,1}$. One of the advantages of our approach is that the results do not depend on detail inputs of a model. We had to overcome several obstacles in deriving the results. First of all, as we have presented in detail, the case of unequal mass inelastic two body scattering process. It was argued that such scatterings have to be treated carefully. We obtained the desired analyticity properties of inelastic amplitude in Section 4. It is worthwhile to note that some of the issues, which were overlooked in previous investigations, have been addressed in this article and have been resolved. We employed refined techniques to this end. We have noted that in the previous works [40, 41] arbitrary constants appeared in the expressions for bounds and some weak assumptions were incorporated. On this occasion, since issues related to analyticity domains of amplitude have been resolved, those arbitrary constants have been determined. Now more refined bounds on forward differential and nonforward differential cross sections have been presented here. We have considered 'inclusive' processes for our purpose and we have obtained various bounds. We utilized an old theorem, derived for average multiplicity, to argue that the resulting bound for integrated inclusive reaction could be improved. It is important to mention that the energy scale, s_0 which invariably appears in the derivation of these bounds in the power of logarithms in the form of $\log \frac{s}{s_0}$, is not fixed. Even, in the derivation of Froissart-Martin bound, for hadronic collisions, the dimensionful constant, s_0 , remains undetermined starting from axioms of field theory whereas t_0 is determined. Prudently, s_0 may be guessed to be order of 1 GeV^2 relating to mass of proton. Therefore, the question arises what value of s_0 is to be adopted for our situation? A plausible answer is that it should be the scale of compactification. However, so far there is no experimentally confirmed evidence for large-radius-compactification hypothesis. The positive aspect is that the energy dependence of bounds have been determined from first principles in the frameworks of local quantum field theories.

Petrov [69] has advanced argument that a D > 4 theory below decompactification scale will exhibit attributes of a D = 4 theory with specific model calculations. We have already shown that a 5-dimensional theory with S^1 compactification is endowed with analyticity properties of a D = 4 theory in a rigorous framework. We confirm conjecture of Petrov that the compactified theory leads to a bound on total cross sections same as the Froissart bound proved for a D = 4 theory. We have proved fixed-t dispersion relations for elastic processes. This article deals with study of analyticity properties of scattering amplitude for inelastic reactions. Therefore, the production of 'exotic' state were dealt with in our new results. Moreover, we have derived bounds on production cross sections in forward direction and in small finite scattering angles. Therefore, our results will be useful for experiments currently going on at LHC. We have also studied asymptotic behaviors of amplitudes for a decompactified, D = 5, theory. We have proposed to test presence of extra spatial dimensions in high energy collisions in the accessible energy range. Let us imagine that one extra spatial dimension is decompactified. We would like to ask is there a way to detect decompactification in high energy collisions? Our proposal [70] is to analyse energy dependence of σ_t . The particle data group has presented the fit to hadron total cross sections, σ_t , over a wide range of energies for several processes where the energy dependendence is assumed to be Froissart-Martin bound saturating i.e. $\sigma_t \approx (\log \frac{s}{s_0})^2$ with the energy scale fitted to be $s_0 = 16 \ GeV^2$. In case an extra spatial dimension is decompactified, there is a possibility that data might exhibit deviation from Froissart-Martin bound. Nayak and Maharana [70] considered total cross section, σ_t data for combined pp and $p\bar{p}$ scatterings and carried out a fit from ISR energy to Tevatron energy, and up to LHC energy. Furthermore, data from Cosmic rays were included in the analysis. We found no evidence for violation of D = 4 Froissart bound. An optimistic argument might be that an extra domension is decompactified at LHC energy which does not saturate D = 5 bound. However, our fit considered data from ISR energy upwards and therefore, we do not believe measurements of total cross sections are going to provide grounds for decompactication at LHC.

It has been suggested by André Martin [71] that if a new threshold opens up in a high energy scattering it will have important implications for analyticity properties of amplitude. One of the implications is that the amplitude might have a large real part. In the context of present investigation, when a pair of KK state is produced, it opens up a new threshold. Consider a scenario when a pair of KK states are produced, consistent with all conservation laws. At the threshold of production of a pair of exotic particles, the cross section for collision of ordinary particles gets affected. According to Martin, dispersion relations survive, furthermore, it is necessary to put an extra term in total cross section. This extra term has to be put at a high energy which might upset the real part at lower energies below the new threshold when we write dispersion relation. In view of above remarks, if there is a threshold for production of KK states in LHC energy regime, measurements of real part of scattering bellow threshold energies will provide indirect evidence for existence of new threshold. Of course, the real part measurements would not provide a clinching evidence for production of KK states; however, it might provide a possible lead to look for KK states.

We have plotted our bounds as a function of energies and gone beyond the existing LHC energies. We recall that T_0 or its fractional power appear as prefactor in all the equations for the bounds. As we have mentioned repeatedly, we cannot extract value of T_0 from experimental data. Therefore, while plotting the bounds, we have factored out T_0 (or its powers from the expressions for bounds). In view of above mentioned remarks, the figures exhibit the energy bounds. Figure 1 shows the bound on $\sigma_t^{ab\to cd}$ for S^1 compactifies case as a function of energy. We have plotted in figure 2, bound for $\sigma_{t5}^{ab\to cd}$, for the five dimensional theory. Figure 3 shows the bound for inelastic differential cross section in forward direction for the compactified d = 5 theory. Figure 4 depicts bounds on differential cross section in near forward direction for the compactified theory, $R^{3,1}$ at angles $\theta = 2^0, 3^0$ and 5^0 and note that differential

cross section falls of rapidly as θ increases. This bound might interest experimentalists who are searching for exhotic states at LHC.

We have refrained from presenting a figure for bound on differential cross sections for the decompactified theory since this bound is weaker compared to the differential cross section for compactified theory.

We end with following closing remarks. Note that the results presented here are not specific to LRC proposal alone. LHC is also searching for other exotic states besides KK states, such as SUSY particles. The bounds presented here do not specifically utilize ingredients of LRC models. Therefore, they hold good for production of other particles in inelastic reactions. For example, there is a high expectation that supersymmetric particle will be produced in LHC. The bounds derived here will play an useful role. If exotic particles are discovered at CERN collider, their properties can be used to study several attributes in a model independent manner. We remind that, in the context of collisions of high energy hadron, the axiomatic results have played important roles to constrain models not only in high energy collisions but have imposed restrictions in low energy physics like $\pi\pi$ and πK reactions. Moreover, those bounds have been experimentally tested and violations have not been reported.

Acknowledgements: I would like to thank Ignatios Antoniadis for very useful discussions and valuable suggestion. I thank Aruna Kumar Nayak and Sanjay Kumar Swain for their critical remarks and providing me some information about search for states arising in LRC models at CERN. I also thank Aruna Kumar Nayak for his comments to improve the manuscript. I thank Matthias Gaberdiel for a very warm and gracious hospitality at the Institute of Theoretical Physics of ETH. It is a pleasure to acknowledge Pauli Center at ETH, Zurich, for very cordial atmosphere and for generous financial supports.

References

- 1. Th. Kaluza, Sitzungsber. Preuss. Akad. Wiss. Phys Math. Klasse 966 (1921).
- 2. O. Klein, Nature, **118**, 516 (1926); Z. F. Physik, **37**, 895 (1926).
- T. Appelquist, A. Chodos and P. G. O. Freund, Modern Kaluza-Klein Theories, Addition-Wesley Publishing Co. Inc. 1987.
- G. Nordström, Phys. Zeitsch. 13, 1126 (1912): Ann. d. Physik 40, 872 (1913);
 42, 533 (1913).
- 5. G. Nordström, Phys. Zeitsch. 15, 504 (1914)
- 6. E. Schrödinger, Ann. der Phys. 82, 257 (1927).
- 7. P. Jordan, Ann. d. Phys. 1, 219 (1947)
- 8. P. Bergmann and A. Einstein, Ann. Math. **39**, 683 (1938).
- 9. W. Pauli, Ann. d. Physik 18, 305 (1933)
- 10. P. Bergmann, Ann. Math. 49, 255 (1948).
- J. Scherk and J. H. Schwarz, Phys. Lett. 57B, 643 (1975); Nucl. Phys. B153 61 (1979)
- 12. I. Antoniadis, Phys. Lett. **B246**, 377 (1990).
- 13. I. Antoniadis, C. Munoz and M. Quiros, Nucl. Phys. B397, 515 (1993).
- N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. B429, 263 (1998).
- N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Rev. D59, 086004 (1999).
- A. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. 436, 257 (1998)
- 17. J. Kretzschmar, Nucl. Part. Phys. Proc. 273-275, 541 (2016).
- 18. S. Rappoccio, Rev. in Phys.4, 100027 (2019).
- 19. I. Antoniadis and K. Benakli, Mod. Phys. Lett. A 30, 1502002 (2015)
- 20. D. Luest and T. R. Taylor, Mod. Phys. Lett. A 30, 15040015 (2015).

- 21. E. Accomando, Mod. Phys. Lett. A30 1504001106 (2015).
- 22. G. Servant, Mod. Phys. Lett. A30, 1540400118 (2015).
- 23. J. Kretzschmar, Nucl. Part. Phys. Proc. 273-275, 541 (2016).
- 24. S. Rappoccio, Rev. in Phys.4, 100027 (2019).
- 25. C. Csaki, TASI Lecture, hep-th/0404096
- 26. T. G. Rizzo, Pedagogical Introduction to Extra Dimensions hep-th/0409309
- 27. G. D. Kribs, Phenomenology of Extra Dimensions, TASI-2004, hep-th/0605325.
- 28. R. Rattazi, Cargese Lecture on Extra Dimensions hep-th/0607055
- 29. H. -C. Cheng, Introduction to Extra Dimensions, TASI-2009, arXiv:1003.1162.
- M. Shifman, Large Extra Dimensions: Becoming Acquainted with an Alternative Paradigm, Int. J. Mod. Phys. A25, 120101, 2012; arXiv: 0909.3074.
- 31. E. P. Panton, Four Lectures on TeV Scale Extra Dimensions, arXiv:1207.3827
- 32. B. Berenji, E. Bloom and J. Cohen-Tanuji, arXiv:1201.2460.
- 33. Yu- Xiao Li, Introduction to Extra Dimensions and Thick Braneworld, arXiv:1707.08541.
- 34. M. Froissart, Phys. Rev. **123**, 1053 (1961).
- R. P. Feynman, Phys. Rev. Lett. 23, 1415(1969); R. P. Feynman, Photon Hadron Interactions, CRC Press, 1972.
- 36. N. N. Khuri, Ann. Phys. **242**, 332(1995)
- 37. J. Maharana, Nucl. Phys. **B943**, 114619 (2019).
- 38. J. Maharana, J. High Energ. Phys. **2020**, 139 (2020).
- 39. J. Maharana, J. Math. Phys. 58, 012302 (1917).
- 40. V. V. Ezhela, A. A. Logunov and M. Mestvirishvili, Th. Math. Phys. 6, 4 (1971).
- A.A. Logunov, N. Van Hieu (helped by M.A. Mestvieishvilli and N.N. Thuan) Proc. Topical Conf. on High energy collisions of hadrons, Vol. II, CERN, Geneva (1968), p. 74.
- 42. H. Lehmann, K. Symanzik and W. Zimmermann, Nuovo Cim. 1, 205 (1955).

- 43. V. Singh and S. M. Roy, Ann. Phys. 57, 461 (1970).
- 44. M. Einhorn and R. Blankenbecler, Ann. Phys. 57, 480 (1971)
- 45. A. Martin, Scattering Theory: unitarity, analyticity and crossing, Springer-Verlag, Berlin-Heidelberg-New York, (1969).
- 46. A. Martin and F. Cheung, Analyticity properties and bounds of the scattering amplitudes, Gordon and Breach, New York (1970).
- 47. C. Itzykson and J.-B. Zubber, Quantum Field Theory; Dover Publications, Mineola, New York, (2008).
- 48. M. Froissart, in Dispersion Relations and their Connection with Causality (Academic, New York); Varrena Summer School Lectures (1964).
- 49. H. Lehmann, Varrena Lecture Notes, Nuovo Cimen. Supplemento, **14** (1959) 153 (1959) series X.
- 50. G. Sommer, Fortschritte. Phys. 18, 577 (1970).
- 51. R. J. Eden, Rev. Mod. Phys. 43, 15 (1971).
- 52. S. M. Roy, Phys. Rep. C5, 125 (1972).
- 53. R. Jost, The General Theory of Quantized Fields, American Mathematical Society, Providence, Rhodes Island (1965).
- 54. J. F. Streater, Rep. Prog. Phys. **38** (1975) 771.
- 55. L. Klein, Dispersion Relations and Abstract Approach to Field Theory Field Theory, Gordon and Breach, Publisher Inc, New York (1961).
- 56. S. S. Schweber, An Introduction to Relativistic Quantum Field Theory, Raw, Peterson and Company, Evaston, Illinois1(961).
- 57. N. N. Bogolibov, A. A. Logunov, A. I. Oksak, I. T. Todorov, General Principles of Quantum Field Theory, Klwer Academic Publisher,
- 58. R. Jost and H. Lehmann, Nuovo Cim. 5, 1598 (1957).
- 59. F. J. Dyson, Phys. Rev, **110**, 1460 (1958).
- 60. H. Lehmann, Nuovo. Cim. **10**, 579 (1958).
- 61. A. Martin, Nuovo Cim. **44A**, 1219 (1966).
- 62. H. Lehmann, Commun. Math. Phys. 2, 375 (1966).

- 63. J. Bros, H. Epstein and V. Glaser, Commun. Math. Phys. 1, 240 (1965).
- 64. G. Sommer, Nuovo Cim. **52A**, 866 (1967)
- 65. Y. S. Jin and A. Martin, Phys. Rev. 135, B1369, (1964).
- 66. A. Mueller, Phys. Rev. **D2**, 2963 (1970)
- 67. T. Kinoshita and J. Maharana, J. Math. Phys. 16, 2294 (1975) (2015).
- 68. J. Maharana, J. Math. Phys. 50, 1510.03008.
- 69. V. A. Petrov, Mod. Phys. Lett. A16, 151 (2001).
- 70. A. Nayak and J. Maharana, Phys. Rev. D 102, 034018, 2020.
- 71. E-mail communication, 27 Jan 2017, to L. Alvarez-Gaume, N. N. Khuri and J. Maharana.