

# THE SOCLE OF THE GROUP ALGEBRA OF A FINITE $p$ -GROUP

DAVID J. BENSON

**ABSTRACT.** Let  $G$  be a finite  $p$ -group, and  $\alpha$  an automorphism of the group algebra  $\mathbb{F}_p G$ . Then  $\alpha$  fixes the socle of  $\mathbb{F}_p G$  pointwise. More generally, if  $k$  is a field of characteristic  $p$ , and  $\alpha$  is a  $k$ -algebra automorphism of  $kG$ , then  $\alpha$  induces a linear action on the dimension subquotients of the group, and the action on the socle is scalar multiplication by the  $(p-1)$ st power of the product of the determinants of this action. The scalar is thus an element of  $(k^\times)^{p-1}$ .

## 1. INTRODUCTION

Throughout this paper, we let  $G$  be a finite  $p$ -group and  $k$  a field of characteristic  $p$ . It is well known that the socle of  $kG$  is one dimensional, spanned by the sum of the group elements. Therefore given any algebra automorphism  $\alpha$  of  $kG$ , there is an element  $\lambda \in k^\times$  such that the action of  $\alpha$  on the socle is given by multiplication by  $\lambda$ . We use the Jennings–Quillen theory, describing the associated graded of the radical filtration to prove the following theorem.

**Theorem 1.1.** *For any algebra automorphism  $\alpha$  of  $kG$ , we have  $\lambda \in (k^\times)^{p-1}$ . In particular if  $k = \mathbb{F}_p$  then  $\lambda = 1$ .*

The exact value of  $\lambda$  can be computed as the  $(p-1)$ st power of the determinant of a matrix given by the action on the dimension subquotients of  $G$ .

**Acknowledgement.** This paper grew out of a question posed by Henrik Rüping on the online forum MathOverflow.

## 2. THE JENNINGS–QUILLEN THEORY

In this section, we prove Theorem 1.1. Recall that  $G$  is a finite  $p$ -group and  $k$  is a field of characteristic  $p$ . Jennings [3] described the radical layers of  $kG$  in terms of a certain descending central series of  $G$ , and Quillen [4] reinterpreted Jennings’ theorem in terms of restricted enveloping algebras. See also Section 3.14 of [1] and Section V.7 of [2] for background.

Let  $J(kG)$  be the Jacobson radical of  $kG$ , spanned by the elements  $g - 1$  for  $g \in G$ . The associated graded algebra is defined by

$$\mathrm{gr}_*(kG) = \bigoplus_{n \geq 0} J^n(kG) / J^{n+1}(kG).$$

For  $r \geq 1$ , the dimension groups  $F_r(G)$  are defined by

$$F_r(G) = \{g \in G \mid g - 1 \in J^r(kG)\}.$$

---

2020 *Mathematics Subject Classification.* Primary: 20C20. Secondary: 20C05, 16S34.

*Key words and phrases.* Finite  $p$ -group, group algebra, socle.

Thus  $F_1(G) = G$ ,  $F_2(G) = \Phi(G)$ ,  $[F_r(G), F_s(G)] \subseteq F_{r+s}(G)$ , and if  $g \in F_r(G)$  then  $g^p \in F_{pr}(G)$ . Furthermore,  $F_r(G)$  is the most rapidly descending central series with these properties. We define

$$\mathbf{Jen}_*(G) = \bigoplus_{r \geq 1} k \otimes_{\mathbb{F}_p} F_r(G)/F_{r+1}(G).$$

Then  $\mathbf{Jen}_*(G)$  is a  $p$ -restricted Lie algebra over  $k$  with Lie bracket induced by taking commutators in  $G$  and with the  $p$ -restriction map  $x \mapsto x^{[p]}$  induced by taking  $p$ th powers in  $G$ . As a  $p$ -restricted Lie algebra,  $\mathbf{Jen}_*(G)$  is generated by its degree one elements. Let  $\mathcal{U}\mathbf{Jen}_*(G)$  be the restricted universal enveloping algebra of  $\mathbf{Jen}_*(G)$  over  $k$ . As an associative algebra,  $\mathcal{U}\mathbf{Jen}_*(G)$  is generated by  $\mathbf{Jen}_*(G)$ , subject to relations given by the commutators and  $p$ th powers. Note that multiplicative commutators in  $G$  become additive commutators in  $\mathcal{U}\mathbf{Jen}_*(G)$ .

The Jennings–Quillen theorem states that there is a  $k$ -algebra isomorphism

$$(2.1) \quad \mathcal{U}\mathbf{Jen}_*(G) \rightarrow \mathbf{gr}_*(kG)$$

which, for any  $r$  and any  $g \in F_r(G)$ , sends the image of  $g$  in  $F_r(G)/F_{r+1}(G)$  to the image of  $g - 1$  in  $\mathbf{gr}_*(kG)$ .

Now let  $|G| = p^m$ . Then  $\dim_k \mathbf{Jen}_*(G) = m$  and  $\dim_k \mathcal{U}\mathbf{Jen}_*(G) = p^m$ . Let  $x_1, \dots, x_m$  be a basis of homogeneous elements in  $\mathbf{Jen}_*(G)$ , with  $x_j$  the image of an element  $g_j$  in the corresponding dimension subgroup of  $G$ , for  $1 \leq j \leq m$ . We write  $\bar{x}_j$  for the corresponding generators of  $\mathcal{U}\mathbf{Jen}_*(G)$ .

Next, we introduce an increasing filtration on  $\mathcal{U}\mathbf{Jen}_*(G)$  as in the proof of the Poincaré–Birkhoff–Witt theorem for Lie algebras. Let  $\mathcal{U}^{(0)}\mathbf{Jen}_*(G) = k \cdot 1$  in degree zero,  $\mathcal{U}^{(1)}\mathbf{Jen}_*(G) = k \cdot 1 \oplus \mathbf{Jen}_*(G)$ , and for  $i \geq 2$ ,  $\mathcal{U}^{(i)}\mathbf{Jen}_*(G) = \mathbf{Jen}_*(G)\mathcal{U}^{(i-1)}\mathbf{Jen}_*(G)$ . Thus  $\mathcal{U}^{(i)}\mathbf{Jen}_*(G)$  is the subspace of  $\mathcal{U}\mathbf{Jen}_*(G)$  spanned by words of length at most  $i$  in the generators  $\bar{x}_j$  of  $\mathcal{U}\mathbf{Jen}_*(G)$ . Then

$$\begin{aligned} \mathcal{U}^{(i)}\mathbf{Jen}_*(G) \cdot \mathcal{U}^{(j)}\mathbf{Jen}_*(G) &\subseteq \mathcal{U}^{i+j}\mathbf{Jen}_*(G), \\ [\mathcal{U}^{(i)}\mathbf{Jen}_*(G), \mathcal{U}^{(j)}\mathbf{Jen}_*(G)] &\subseteq \mathcal{U}^{i+j-1}\mathbf{Jen}_*(G), \end{aligned}$$

and if  $x$  is in  $\mathcal{U}^{(1)}\mathbf{Jen}_*(G)$  then  $x^p$  is also in  $\mathcal{U}^{(1)}\mathbf{Jen}_*(G)$ . Then the associated graded

$$(2.2) \quad \mathcal{S}\mathbf{Jen}_*(G) = \bigoplus_{i \geq 0} \mathcal{U}^{(i)}\mathbf{Jen}_*(G)/\mathcal{U}^{(i-1)}\mathbf{Jen}_*(G)$$

carries a double grading. The first degree keeps track of the dimension subgroups, and the second degree comes from the filtration. It is generated by  $\mathcal{U}^{(1)}\mathbf{Jen}_*(G)$ , is commutative, and  $p$ th powers of positive degree elements are zero. Writing  $\bar{\bar{x}}_j$  for the image of  $\bar{x}_j$  in  $\mathcal{S}\mathbf{Jen}_*(G)$ , the analogue of the Poincaré–Birkhoff–Witt theorem for  $p$ -restricted Lie algebras shows that

$$\mathcal{S}\mathbf{Jen}_*(G) \cong k[\bar{\bar{x}}_1, \dots, \bar{\bar{x}}_m]/(\bar{\bar{x}}_1^p, \dots, \bar{\bar{x}}_m^p),$$

is the truncated symmetric algebra on  $\mathbf{Jen}_*(G)$ . Thus the elements  $\bar{x}_1^{i_1} \dots \bar{x}_m^{i_m}$  form a basis for  $\mathcal{U}\mathbf{Jen}_*(G)$ , where  $0 \leq i_j < p$  for  $1 \leq j \leq m$ , and the elements  $\bar{\bar{x}}_1^{i_1} \dots \bar{\bar{x}}_m^{i_m}$  form a basis for  $\mathcal{S}\mathbf{Jen}_*(G)$ .

### 3. PROOF OF THE THEOREM

The socle of  $kG$  is one dimensional, spanned by the sum of the group elements,

$$n = \sum_{g \in G} g = \prod_{j=1}^m (g_j^{p-1} + g_j^{p-2} + \cdots + g_j + 1) = \prod_{j=1}^m (g_j - 1)^{p-1} \in kG.$$

The image of this in the restricted universal enveloping algebra  $\mathcal{U}\text{Jen}_*(G)$ , using the isomorphisms (2.1), is  $\bar{n} = \prod_{j=1}^m \bar{x}_j^{p-1}$ . The image in  $\mathcal{S}\text{Jen}_*(G)$ , using (2.2), is  $\bar{\bar{n}} = \prod_{j=1}^m \bar{\bar{x}}_j^{p-1}$ .

Now let  $\alpha$  be a  $k$ -algebra automorphism of the group algebra  $kG$ . Then  $\alpha$  induces an automorphism  $\bar{\alpha}$  of  $\mathcal{U}\text{Jen}_*(G)$  and an automorphism  $\bar{\bar{\alpha}}$  of  $\mathcal{S}\text{Jen}_*(G)$ . Since the socle of  $kG$  is one dimensional, the action of  $\alpha$  on  $n$  is multiplication by some scalar  $\lambda$ , so  $\alpha(n) = \lambda n$  with  $\lambda \in k^\times$ . Taking associated graded with respect to the radical filtration and using the isomorphism 2.1, we obtain  $\bar{\alpha}(\bar{n}) = \lambda \bar{n}$ . Then, taking the associated graded with respect to the Poincaré–Birkhoff–Witt filtration using (2.2), we have

$$(3.1) \quad \bar{\bar{\alpha}}(\bar{\bar{n}}) = \lambda \bar{\bar{n}}.$$

To compute the value of  $\lambda$ , we look at the action of  $\alpha$  on  $\mathcal{S}\text{Jen}_*(G)$ . This preserves the double grading, and so it is given by an  $m \times m$  invertible matrix  $A$  on the generators  $\bar{\bar{x}}_1, \dots, \bar{\bar{x}}_m$ . Denote by  $\det(A)$  the determinant of the matrix  $A$ . Then

$$(3.2) \quad \bar{\bar{\alpha}}(\bar{\bar{n}}) = \prod_{j=1}^m \bar{\bar{\alpha}}(\bar{\bar{x}}_j)^{p-1} = \det(A)^{p-1} \prod_{j=1}^m \bar{\bar{x}}_j^{p-1} = \det(A)^{p-1} \bar{\bar{n}}.$$

To check the second equality, it suffices to check it on generators for the general linear group. Taking the generators to be the elementary matrices and the diagonal matrices, the check is easy.

It follows from (3.1) and (3.2) that

$$\lambda = \det(A)^{p-1} \in (k^\times)^{p-1},$$

and this completes the proof of Theorem 1.1.

### REFERENCES

1. D. J. Benson, *Representations and cohomology I: Basic representation theory of finite groups and associative algebras*, Cambridge Studies in Advanced Mathematics, vol. 30, Cambridge University Press, 1991, reprinted in paperback, 1998.
2. N. Jacobson, *Lie algebras*, John Wiley & Sons Inc., New York, 1962.
3. S. A. Jennings, *The structure of the group ring of a  $p$ -group over a modular field*, Trans. Amer. Math. Soc. **50** (1941), 175–185.
4. D. G. Quillen, *On the associated graded ring of a group ring*, J. Algebra **10** (1968), 411–418.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF ABERDEEN, ABERDEEN AB24 3UE, UNITED KINGDOM  
Email address: d.j.benson@abdn.ac.uk