

Structure of Nucleons and Their Interaction in the Concept of Nonperturbative QCD as a Pressing Issue of 21st-Century Physics

V.I. Komarov*

Joint Institute for Nuclear Research, Dubna, 141980 Russia

Abstract

The problem of the structure of nucleons and their interaction in the concept of nonperturbative QCD is discussed as an approach to studying the transformation of current quarks into constituent ones and the search for the mechanism of such a transformation, creating the bulk of the nucleon mass. Attention is drawn to the possibilities offered by studying central nucleon-nucleon collisions in this aspect.

Modern physics operates with hundreds of elementary particles. The first parameters of particle identification are the average value and variance of its mass. In this case, the mass of nucleons determines the mass of objects in the world around us. Naturally, the problem of the origin of mass is the primary fundamental problem of particle physics. The discovery of the mass generation mechanism proposed by Braut, Engler, and Higgs, which was triumphantly confirmed in experiments, turned out to be the greatest achievement of particle physics in recent decades. This mechanism has become a widely known scientific fact. If you ask how the masses of elementary particles are generated, most physicists will probably say “the Higgs mechanism.” And that would be wrong. Wrong because the Higgs boson mechanism generates only masses of leptons and current quarks and has nothing to do with the mass of constituent quarks that form nucleons [1]. This mass is at least 80% of the mass of the Universe of the mass of the visible Universe (hadrons, leptons, and the electromagnetic field), and an even larger share on Earth.

There is no reliable and generally accepted approach to describe the generation of nucleon mass. It is only known that such generation is due to the spontaneous violation of the chirality of the vacuum, but the question of the mechanism of the transition from a “light,” 5-MeV, current quark to a “heavy,” 300-MeV, constituent quark remains completely open. How the interaction with the vacuum of three high-momentum current quarks converts their kinetic energy into the mass of constituent quarks that stick together into a nucleon is unknown. It is obvious that the fundamental nature of such a transition is not inferior to the fundamental nature of the Higgs mechanism.

The discovery of the mechanism of this transition certainly requires knowledge of the structure of the nucleon and the dynamics of the interaction of

*e-mail: komarov@jinr.ru

constituent quarks in the concept nonperturbative Quantum Chromodynamics (NPQCD). This QCD has achieved significant success. Effective chiral Lagrangians made it possible to reproduce all the basic properties of light mesons and their interactions at low energies [2, 3]. The most successful, consistent and universal microscopic description of light baryonic systems, achieved in the chiral model of constituent quarks [4], simultaneously reproduces baryon spectra and baryon-baryon interactions at low energies. Phase shifts of NN scattering, binding energies of the lightest nuclei, wave function and form factors of the deuteron, masses and channels of strong decay of dibaryon resonances, cross sections for hyperon-nucleon interactions, and even certain properties of pentaquarks [5] are available for this model. However, NPQCD has no direct derivation from the basic postulates of QCD, and its achievements do not indicate a mechanism for the transition of current quarks to constituent ones.

Information on the structure of nucleons obtained at high energies in the concept of perturbative QCD is also not constructive in this respect. Enormous efforts have been made to date to elucidate the structure of nucleons in this concept, but the successfully obtained high-energy structure of nucleons is very far from the low-energy one. Suffice it to recall that the bulk of the data is in the form of one-dimensional parton distributions, while the distributions of quarks in the ground state of a nucleon are certainly three-dimensional in space. In recent decades, the theory of generalized quark distributions, which are a function of three variables [6], and attempts are made to relate the “high-energy” and “low-energy” distributions for a unified description of the entire energy region (see, for example [7]).

However, the problem is not only in the dimension of the distributions. The difference is more fundamental: High-energy data deal with current quarks, and low-energy data deal with constituent ones. It would be possible to obtain information on the distributions of constituent quarks directly from the distributions of current ones, only knowing the relationship of current quarks with their gluon and quark-antiquark environment in constituent quarks. However, no such information is available. Therefore, the quark structure of nucleons in the low-energy region remains a completely special and self-sufficient problem. Its solution obviously demands new experimental data, and the most effective direction is to obtain them in a wide range of energies, including both low-energy and high-energy ones.

In this aspect, the NICA complex being created at JINR has a great advantage, energetically covering both areas of interest. The transition of the meson-baryon phase of matter to the quark-gluon phase has traditionally been studied in collisions of heavy nuclei. However, for the study of the nonperturbative QCD structure of nucleons, such collisions create very significant complications: (1) the low quark density of nuclei in comparison with the density of nucleons requires high collision energies to achieve the baryon and energy densities of interest; (2) fluctuations of the nucleon density in the initial state of nuclei lead to a significant scatter of local baryon and energy densities in the intermediate state; (3) a large number of nucleons participating in the collision creates a complex dynamic scenario of collision, in particular, with the presence of a mixed meson-baryon and quark-gluon phase.

Therefore, the study of the elementary process of transition between constituent and current quarks in nucleon-nucleon collisions has radically more def-

inite initial conditions and is more accessible for interpretation than in collisions of heavy nuclei.

The NICA project provides for the creation of a facility for the study of nucleon–nucleon collisions (the Spin Physics Detector, SPD) [8]. One of the priority goals of this facility may be precisely the study of the structure of nucleons in the low-energy region, that is, in the region of nonperturbative QCD, as well as in the region of the transition to the perturbative region. Optimal research conditions are created in the kinematics of collisions, providing overlapping of the central quark regions of nucleons [9]. Collisions with large impact parameters determined by meson-baryon processes are ineffective for studying the quark degrees of freedom of the arising states. Therefore, it is of particular importance to distinguish collisions with impact parameters on the order and less than the radius of the quark region of nucleons, that is, $r_c \approx 0.4$ fm. The corresponding criterion for centrality was recently discussed in [10].

It should be emphasized that inelastic central collisions of nucleons have a significant probability, which does not disappear with increasing energy. R. Feynman in 1969 [11] drew attention to the fact that the constant with the energy part of the total inelastic cross section of hadron collisions is determined by events with a high multiplicity of produced particles. High multiplicity is now widely used as a criterion for collision centrality. Therefore, the part of the total inelastic nucleon collision cross section discussed by Feynman, which does not vanish with energy, can be associated with the collision of their quark core.

The total cross section for central nucleon collision is at the level $\sigma_{cNN}^{\text{inel}} = \pi r_c^2 = 5$ mbn. In the collision of protons, due to the conservation of the baryon number, a nucleon pair appears in the final state: pp , np , nn . Therefore, each collision is accompanied with a probability close to 100%, the almost isotropic emission of at least one proton. As a result, for almost isotropic emission, the differential cross section for the emission of one proton at an angle close to 90° , is about $\sigma_{cNN}^{\text{inel}}/4\pi = 0.4$ mbn/sr.

Numerous experimental data show that, in a wide range of processes, as well as angular and energy measurement conditions, the emission of protons is accompanied by the emission of deuterons with a ratio of about 3×10^{-3} . If we use the registration of a deuteron at an angle close to 90° [10], the process of interest to us occurs with a differential cross section $\approx 1\mu\text{b/sr}$, which is quite sufficient for carrying out correlation and polarization measurements.

Central NN Collisions open up a wide range of studies on the structure and dynamics of nucleon interaction in the nonperturbative QCD regime. The object of research is the characteristics of the reaction

$$p + p \rightarrow d(90^\circ) + M, \quad (1)$$

where M is a system of mesons with an admixture of baryon-antibaryon pairs at sufficiently high energies.

Of primary interest is the search for the effects of the transition of constituent quarks to current ones. It is essential that the region of such a transition should be within the SPD energy range (4–28 GeV). Indeed, according to long-motivated estimates [12], the characteristic impulse of spontaneous chirality breaking has a value close to $P_\chi \approx 1.2$ GeV/ c . If the transition occurs during the destruction of all six colliding constituent quarks, the total energy required for this in the QCD is $\sqrt{s_{NN}} \approx 12$ GeV. This means that, in the central NN

collisions in the energy region from the overlapping energy of the quark content of nucleons, $\sqrt{s_{NN}} \approx 3$, up to ≈ 12 GeV, a phase state is created in which constituent quarks, Goldstone bosons, and gluons are acting degrees of freedom. This state can be called a constituent-quark phase of strongly interacting matter.

The study of the energy dependence of the processes of central collision of nucleons can be an effective means of observing the discussed fundamental transition of constituent quarks to current ones. It can be expected that such a transition will manifest itself in the energy dependence of the momentum and multiple distributions of particles formed in certain channels of the final state and in their correlations and polarization characteristics.

Note that, over the past decade, considerable interest has arisen in the study of central pp collisions at high energies. However, the energies of hundreds of GeV [13] and TeV [14] used are too large to observe the discussed transition of constituent quarks to current ones, while the unique conditions for such an observation can probably be provided by SPD.

The study of dibaryon resonances should also be among the top-priority tasks. QCD spectroscopy arose in the pioneering work [15] in the 1960s. Light dibaryon resonances with masses 2.1–2.2 GeV [16, 17] were originally viewed as loosely coupled $N\Delta(1238)$ couples. The difficulties that arose in the application of the meson-baryon description of dibaryon resonances increased even more after the observation [18] with respect to heavy resonance with a mass of 2.38 GeV, close to the mass $\Delta(1238)\Delta(1238)$ couples. This stimulated the successful consideration of dibaryon resonances in the framework of the chiral model of constituent quarks [19]. The QCD nature of the heavier dibaryon resonances is beyond doubt. The very existence of such resonances with a mass of 3.0 GeV [20] and 2.65 GeV [21] is shown in the experiments that have already been carried out. A study of dibaryon resonances in the central mode NN collisions can become a promising program for the study of reaction mechanisms determined by the degrees of freedom of nonperturbative QCD (NPQCD). The sensitivity of the processes of pion generation in NN collisions in the GeV energy region to such mechanisms was shown in the works of V.I. Kukulin [22]. Central collisions can become a test bench for NPQCD models of the reaction mechanism determined by the interaction of chiral constituent quarks.

Among the tasks of the first stage, one should also mention the study of the expected dependence of the parameters of the produced mesons and meson correlations on the energy and baryon densities in NN collisions [23, 24].

Having limited ourselves to a brief indication of experiments initiated by modern theoretical expectations, we should also mention the possibility that completely new unexpected effects will be manifested, as is often the case when a new experimental field is opened.

When discussing the proposed experiments on central nucleon–nucleon collisions, it should be borne in mind that the SPD program from the very beginning of the project was focused on polarization studies of the nucleon spin structure [25].

On the whole, the creation of the SPD will open up opportunities for a broad program of investigating the structure of nucleons in nonperturbative QCD. It is difficult to overestimate the possibilities of SPD for carrying out polarization studies in this aspect. The accumulation of empirical information and the development of adequate phenomenological models can become the

basis for creating a rigorous nonperturbative QCD theory of baryons. Such a program requires, on the one hand, intensifying the theoretical work in the discussed problems and, on the other, involving experimenters in preparing the corresponding experiments on the SPD. The first step in solving the problem under discussion should be the realization of its fundamental nature.

Acknowledgments

I thank V.A. Bednyakov, A.V. Kulikov, E.A. Strokovsky, and D.A. Tsirkov for their interest in the problem and our helpful discussions. The support provided by V.I. Kukulin, whose recent passing was untimely, was indispensable.

References

- [1] C. D. Roberts, “Insights into the origin of mass,” *J. Phys.: Conf. Ser.* 1643, 012194 (2020).
- [2] M. K. Volkov, “Effective chiral lagrangians and Nambu-Jona-Lasinio model,” *Phys. Part. Nucl. Lett.* 24, 82 (1993).
- [3] E. R. Arriola, “Pion structure at high and low energies in chiral quark models,” *Acta Phys. Polon. B* 33, 4443 (2002).
- [4] A. Valcarce, H. Garcilazo, F. Fernandez, and P. Gonzalez, “Quark-model study of few-baryon systems,” *Rep. Prog. Phys.* 68, 965 (2005).
- [5] Y. Dong, P. Shen, F. Huang, and Z. Zhang, “Selected strong decays of pentaquark State $P_c(4312)$ in a chiral constituent quark model,” *Eur. Phys. J. C* 80, 341 (2020).
- [6] M. Diehl, “Generalized parton distributions,” *Phys. Rep.* 388, 41 (2003).
- [7] S. Boff, B. Pasquini, and M. Traini, “Linking generalized parton distributions to constituent quark models,” *Nucl. Phys. B* 649, 243 (2003).
- [8] V. M. Abazov et al. (SPD NICA Collab.), “Conceptual design of the spin physics detector,” *arXiv: 2102.00442* (2021).
- [9] V. I. Komarov, “On the possibility of revealing the transition of a baryon pair state to a six-quark confinement state,” *Phys. Part. Nucl. Lett.* 15, 69 (2018).
- [10] V. I. Komarov, B. Baimurzinova, A. Kunsafina, and D. Tsirkov, “Centrality criteria of the inelastic nucleon-nucleon collisions,” *Phys. Part. Nucl. Lett.* 17, 290 (2020).
- [11] R. Feynman, “Very high-energy collisions of hadrons,” *Phys. Rev. Lett.* 23, 1415 (1969).
- [12] A. Manohar and H. Georgi, “Chiral quarks and the non-relativistic quark model,” *Nucl. Phys. B* 234, 189 (1984).

- [13] K. Welsh, J. Singer, and U. Heinz, “Initial state fluctuations in collisions between light and heavy ions,” *Phys. Rev. C* 94, 024919 (2016).
- [14] B. Nachman and M. Mangano, “Observables for possible QGP signatures in central pp collisions,” *Eur. Phys. J. C* 78, 343 (2018).
- [15] F. Dyson and N. H. Xuong, “ $Y = 2$ states in $SU(6)$ theory,” *Phys. Rev. Lett.* 13, 815 (1964).
- [16] R. A. Arndt, D. Roper, R. L. Workman, and M. W. McNaughton, “Nucleon-nucleon partial-wave analysis to 1.6 GeV,” *Phys. Rev. D* 45, 3995 (1992).
- [17] V. I. Komarov, D. Tsirkov, T. Azaryan, Z. Bagdasarian, S. Dymov, R. Gebel, B. Gou, A. Kacharava, A. Khoukaz, A. Kulikov, V. Kurbatov, B. Lorentz, G. Macharashvili, D. Mchedlishvili, S. Merzliakov, et al., “Evidence for excitation of two resonance states in the isovector two-baryon system with a mass of 2.2 GeV/c²,” *Phys. Rev. C* 93, 065206 (2016).
- [18] H. Clement, “On the history of dibaryons and their final observation,” *Prog. Part. Nucl. Phys.* 93, 195 (2017).
- [19] Y. Dong, P. Shen, F. Huang, and Z. Zhang, “Theoretical study of the $d^*(2380) \rightarrow \pi\pi$ decay width,” *Phys. Rev. C* 91, 064002 (2015).
- [20] H. L. Andersen, M. S. Dixit, H. J. Evans, K. A. Klare, D. A. Larson, M. V. Sherbrook, R. L. Martin, D. Kessler, D. E. Nagle, H. A. Thiessen, C. K. Hargrove, E. P. Hincks, and S. Fukui, “Forward differential cross sections for the reaction $p + p \rightarrow d + \pi^+$ in the range 3.4–12.3 GeV/c,” *Phys. Rev. D* 3, 1536 (1971).
- [21] D. Tsirkov, V. Komarov, T. Azaryan, S. Dymov, A. Kunsafina, V. Kurbatov, Z. Kurmanaliyev, and Y. Uzikov, “Recent dibaryon studies at ANKE,” *EPJ Web Conf.* 199, 02016 (2019).
- [22] V. I. Kukulin, I. T. Obukhovskiy, V. N. Pomerantsev, and A. Faessler, “Two-component dressed-bag model for NN interaction: deuteron structure and phase shifts up to 1 GeV,” *Int. J. Mod. Phys. E* 11, 1 (2002).
- [23] T. Hatsuda and T. Kunihiro, “Character changes of pion and σ -meson at finite temperatures,” *Phys. Lett. B* 185, 304 (1987).
- [24] D. Blaschke, Y. L. Kalinovsky, A. E. Radzhabov, and M. K. Volkov, “Scalar σ meson at finite temperature in nonlocal quark model,” *Phys. Part. Nucl. Lett.* 3, 92 (2006).
- [25] A. Arbusov et al. (SPD NICA Collab.), “On the physics potential to study the gluon content of proton and deuteron at NICA SPD,” *Prog. Part. Nucl. Phys.*, 103858 (2021).