Non-degenerate Rigid Alignment in a Patch Framework

Dhruv Kohli

Department of Mathematics University of California San Diego CA 92093, USA

Gal Mishne

Halicioğlu Data Science Institute University of California San Diego CA 92093, USA

Alexander Cloninger

Department of Mathematics and Halicioğlu Data Science Institute University of California San Diego CA 92093, USA DHKOHLI@UCSD.EDU

GMISHNE@UCSD.EDU

ACLONINGER@UCSD.EDU

Abstract

Given a set of overlapping local views (patches) of a dataset, we consider the problem of finding a rigid alignment of the views that minimizes a 2-norm based alignment error. In general, the views are noisy and a perfect alignment may not exist. In this work, we characterize the non-degeneracy of an alignment in the noisy setting based on the kernel and positivity of a certain matrix. This leads to a polynomial time algorithm for testing the non-degeneracy of a given alignment. Consequently, we focus on Riemannian gradient descent for minimization of the error and obtain a sufficient condition on an alignment for the algorithm to converge (locally) linearly to it. In the case of noiseless views, a perfect alignment exists, resulting in a realization of the points that respects the geometry of the views. Under a mild condition on the views, we show that the non-degeneracy of a perfect alignment is equivalent to the local rigidity of the resulting realization. By specializing the characterization of a non-degenerate alignment to the noiseless setting, we obtain necessary and sufficient conditions on the overlapping structure of the views for a locally rigid realization. Similar results are also obtained in the context of global rigidity. Keywords: Nondegeneracy, uniqueness, rigid alignment, local rigidity, global rigidity, linear convergence, Riemannian gradient descent, Morse-Bott function.

1. Introduction

There exist a plethora of problems [9; 7; 26; 30] which involve the task of aligning local views (also known as registration of point clouds) so as to obtain a global view of the data that respects the geometry of the local views. Although it is not uncommon for the correspondence between the points and the views to not be known apriori [27], nevertheless, we assume that the overlapping structure of the local views is available. A suitable example of our setup is rendered by the bottom-up manifold learning techniques, LTSA [33] and LDLE [15], which first construct low distortion local views of high dimensional data into lower dimension followed by the alignment of the views to obtain a low dimensional global embedding of the data. Here we focus on the alignment step.

The correspondence between the local views and the points is captured by a bipartite graph, which when combined with the local coordinates of the points due to the views containing them, form a *patch framework* Θ [3; 12]. The alignment of the views amounts to finding a suitable transformation for each view so that the local coordinates of a point in its transformed views are close to each other. While LTSA seeks affine transformations, LDLE pursues the rigid ones for alignment.

In the case where the views undergo rigid transformation, the problem of aligning views can be posed as the minimization of a 2-norm based alignment error, given by a quadratic F over the product of orthogonal groups $\mathbb{O}(d)^m$ (see Section 2). With our definition of the alignment error, we identify an "alignment" of the local views by an element of $\mathbb{O}(d)^m$. Note that rotating/reflecting each view by the same amount does not affect the alignment error. For a given alignment $\mathbf{S} \in \mathbb{O}(d)^m$, this translates to the alignment error being the same due to \mathbf{SQ} for all $\mathbf{Q} \in \mathbb{O}(d)$ i.e. $F(\mathbf{S}) = F(\mathbf{SQ})$. In this sense, every alignment is degenerate and every optimal alignment is non-unique. With a slight abuse of convention, we define a non-degenerate alignment to be a local minimum of F which is non-degenerate up to an orthogonal transformation. Similarly, we say that an optimal alignment (a global minimum of F) is unique if every other optimal alignment can be obtained by an orthogonal transformation of it. This brings us to our first contribution.

Contribution 1. In Section 3, we derive a characterization of non-degenerate alignment of (possibly noisy) local views that can be tested in polynomial time. By specializing it to the case of two views, we obtain a necessary and sufficient condition for an alignment of two views to be non-degenerate.¹

Given an alignment $\mathbf{S} \in \mathbf{O}(d)^m$ of the local views, a consensus representation $\Theta(\mathbf{S})$ of the points can be obtained by averaging the local coordinates of the points due to the (rigidly transformed) views containing them. In the noiseless setting where the local views are clean measurements of the data (obtained by applying an unknown rigid transformation to a subset of data points), a perfect alignment of views is possible. Equivalently, when the views are noiseless, a value of zero for F is attainable, and an \mathbf{S} that achieves it is called a "perfect alignment". Clearly, a perfect alignment is an optimal one, while the converse may not hold. To be consistent with previous works [12; 13], the consensus representation of the points $\Theta(\mathbf{S})$ due to a perfect alignment \mathbf{S} of the views is called a realization of the framework.

An understanding of affine, global (Euclidean), or local (Euclidean) rigidity [12] of a realization $\Theta(\mathbf{S})$ has importance in several areas such as molecular dynamics [18; 4; 8] or sensor network localization [34; 32]. Under a mild assumption on the structure of the local views, [3; 31; 12] characterized the affine rigidity of a realization by the rank of a certain matrix derived from the framework Θ . It was shown in [23] that deriving a similar characterization of global rigidity is NP-Hard. Nevertheless a characterization of the local rigidity is useful from an algorithmic standpoint as we show in this work. Furthermore, necessary and sufficient conditions on the overlapping structure of the local views for affine rigidity were derived in [31]. Similar results in the context of local and global rigidity form our second set of contributions.

^{1.} A similar condition for the uniqueness of an optimal alignment of two views exists (see [25]). Nevertheless, we provide a short proof based on the constructs defined in this work.

Contribution 2.

- 1. In Section 4, under a mild assumption on the structure of the local views, we show that the local and global rigidity of a realization $\Theta(\mathbf{S})$ are equivalent to the non-degeneracy and uniqueness of the perfect alignment \mathbf{S} , respectively. As a corollary, we obtain a characterization of the local rigidity of a realization.
- 2. By specializing the characterization of a non-degenerate alignment to noiseless setting, we obtain necessary and sufficient conditions on the overlapping structure of the local views for a perfect alignment to be non-degenerate (equivalently, for a realization to be locally rigid). Similar conditions are also derived for the uniqueness of a perfect alignment.

Several algorithms exist to obtain an approximate solution of the alignment problem at hand. A few important ones are: semidefinite programming (SDP) [3; 2], spectral relaxation (SPEC) [3; 2], Procrustes analysis (PROC) [15; 6], generalized power method (GPM) [19] and Riemannian gradient descent (RGD) [17]. It is common to obtain stability and convergence guarantees of an algorithm under rigidity constraints over the framework. For example, [3] derived stability guarantees on the SPEC and SDP solutions under affine rigidity constraints. Also, [19; 20] derived stability guarantees and showed global linear convergence of GPM with SPEC initialization under the setting where each view is affinely non-degenerate and contains all the points. It is easy to deduce that this framework exhibits global rigidity when the views have no noise. Our last contribution is along the same lines.

Contribution 3. In Section 5, we show that RGD converges locally linearly to a nondegenerate alignment of views. A corollary of that is: when the local views are affinely nondegenerate and noiseless then RGD converges locally linearly to a non-degenerate perfect alignment and thus to a locally rigid realization.

The notation and proofs are provided in Appendix A.

2. A Quadratic Program in Orthogonal Groups for Aligning Views

In this section we borrow and build upon the patch framework setup described in [3], elucidate the structure of the objects underlying a framework, define an alignment, optimal alignment and perfect alignment of views, and the realization of a framework due to a perfect alignment.

Suppose that a sequence of m point clouds in \mathbb{R}^d is available where each cloud represents a local view of the dataset $(\mathbf{x}_k)_1^n$. Let Γ be a graph of m + n vertices, where the kth vertex represents the kth point for $k \in [1, n]$ and the (n + i)th vertex represents the ith view for $i \in [1, m]$. An edge $(k, i) \in E(\Gamma)$ means that the kth point has a local representation due to ith view, given by $\mathbf{x}_{k,i} \in \mathbb{R}^d$. In particular, Γ is bipartite. The tuple $\Theta = (\Gamma, (\mathbf{x}_{k,i}))$ is called the patch framework.

Given a patch framework Θ , the task is to align the overlapping views, precisely, to find orthogonal matrices $(\mathbf{S}_i)_1^m \subseteq \mathbb{O}(d)$ and translation vectors $(\mathbf{t}_i)_1^m \subseteq \mathbb{R}^d$ so that the local representations of the *k*th point across the rigidly transformed views are close in the 2-norm.

This naturally leads to the following problem,

$$\mathcal{A}_{0} \coloneqq \min_{\substack{(\mathbf{S}_{i})_{1}^{m} \subseteq \mathbb{Q}(d) \\ (\mathbf{t}_{i})_{1}^{m} \subseteq \mathbb{R}^{d} \\ (k,j) \in E(\Gamma)}} \sum_{\substack{(k,i) \in E(\Gamma) \\ (k,j) \in E(\Gamma)}} \left\| (\mathbf{S}_{i}^{T} \mathbf{x}_{k,i} + \mathbf{t}_{i}) - (\mathbf{S}_{j}^{T} \mathbf{x}_{k,j} + \mathbf{t}_{j}) \right\|_{2}^{2}.$$
(1)

An equivalent problem is to find $(\mathbf{x}_k)_1^n$, $(\mathbf{S}_i)_1^m$ and $(\mathbf{t}_i)_1^m$ that minimize

$$\mathcal{A}_{1} \coloneqq \min_{\substack{(\mathbf{S}_{i})_{1}^{m} \subseteq \mathbb{Q}(d) \\ (\mathbf{t}_{i})_{1}^{m} \subseteq \mathbb{R}^{d} \\ (\mathbf{x}_{k})_{1}^{n} \subseteq \mathbb{R}^{d}}} \sum_{\substack{(k,i) \in E(\Gamma) \\ (\mathbf{x}_{k})_{1}^{n} \subseteq \mathbb{R}^{d}}} \left\| \mathbf{x}_{k} - (\mathbf{S}_{i}^{T} \mathbf{x}_{k,i} + \mathbf{t}_{i}) \right\|_{2}^{2}.$$
(2)

Here, \mathcal{A}_0 and \mathcal{A}_1 are the optimal alignment errors. The proof of the following proposition derives the optimal \mathbf{x}_k in Eq. (2) from which it follows that the minimizers $(\mathbf{S}_i)_1^m$ and $(\mathbf{t}_i)_1^m$ of the two problems coincide.

Proposition 1. $A_0 = 2A_1$.

Note that translating the optimal \mathbf{x}_k and \mathbf{t}_i by the same amount does not change the objective and still leads to an optimal solution. To prevent that, we add a centering constraint $\sum_{1}^{n} \mathbf{x}_k = 0$ so that the mean of \mathbf{x}_k is at the origin. A more concise representation of \mathcal{A}_1 with the centering constraint is as follows. Define

$$\mathbf{H} \coloneqq [\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n, \mathbf{t}_1, \mathbf{t}_2, \dots, \mathbf{t}_m] \in \mathbb{R}^{d \times (n+m)}.$$

Define $\mathbf{e}_{ki} \coloneqq \mathbf{e}_{k}^{n+m} - \mathbf{e}_{n+i}^{n+m}$, then $\mathbf{x}_k - \mathbf{t}_i = \mathbf{H}\mathbf{e}_{ki}$. Let $\mathbf{S} = [\mathbf{S}_i]_1^m \in \mathbb{O}(d)^m \subseteq \mathbb{R}^{md \times d}$. Then

$$\mathcal{A}_{1} = \min_{\substack{\mathbf{S} \in \mathbb{O}(d)^{m} \\ \mathbf{H} \in \mathbb{R}^{d \times (n+m)} \\ \mathbf{H}\mathbf{I}_{n}^{n+m} = 0}} \sum_{\substack{(k,i) \in E(\Gamma) \\ \mathbf{H}\mathbf{I}_{n}^{n+m} = 0}} \left\| \mathbf{H}\mathbf{e}_{ki} - \mathbf{S}^{T}(\mathbf{e}_{i}^{m} \otimes \mathbf{I}_{d})\mathbf{x}_{k,i} \right\|_{2}^{2}$$
$$= \min_{\substack{\mathbf{S} \in \mathbb{O}(d)^{m} \\ \mathbf{H} \in \mathbb{R}^{d \times (n+m)} \\ \mathbf{H}\mathbf{I}_{n}^{n+m} = 0}} \operatorname{Tr}\left(\begin{bmatrix} \mathbf{H} & \mathbf{S}^{T} \end{bmatrix} \begin{bmatrix} \mathcal{L}_{\Gamma} & -\mathbf{B}^{T} \\ -\mathbf{B}^{T} & \mathbf{D} \end{bmatrix} \begin{bmatrix} \mathbf{H}^{T} \\ \mathbf{S} \end{bmatrix} \right)$$
(3)

where

$$\mathcal{L}_{\Gamma} = \sum_{(k,i)\in E(\Gamma)} \mathbf{e}_{ki} \mathbf{e}_{ki}^{T}$$
(4)

$$\mathbf{B} = \sum_{(k,i)\in E(\Gamma)} (\mathbf{e}_i^m \otimes \mathbf{I}_d) \mathbf{x}_{k,i} \mathbf{e}_{ki}^T$$
(5)

$$\mathbf{D} = \sum_{(k,i)\in E(\Gamma)} \mathbf{x}_{k,i} \mathbf{x}_{k,i}^T (\mathbf{e}_i^m \otimes \mathbf{I}_d)^T.$$
(6)

Remark 1. Note that \mathcal{L}_{Γ} is the combinatorial Laplacian of the graph Γ . Due to the bipartite structure of Γ , $\mathcal{L}_{\Gamma}^{\dagger}$ can be computed in $O(nm^2)$ [14], against the complexity of $O((m + n)^3)$ for the general case. The matrix \mathbf{D} is a block diagonal matrix where the *i*th block is $\sum_{(k,i)\in E(\Gamma)} \mathbf{x}_{k,i}\mathbf{x}_{k,i}^T$. The matrix $\mathbf{B} = [\mathbf{B}_i]_1^m$ is a vertical stack of m matrices, one for each view, where each $\mathbf{B}_i \in \mathbb{R}^{d \times (n+m)}$. For a fixed i, $\mathbf{B}_i(:, n + i) = -\sum_{(k,i)\in E} \mathbf{x}_{k,i}$, if $(k,i) \in E(\Gamma)$ then $\mathbf{B}_i(:,k) = \mathbf{x}_{k,i}$ otherwise zero. Thus, $\mathbf{1}_{n+m} \in \ker(\mathbf{B}_i)$. Finally, the *j*th row of \mathbf{B}_i contains information about the *j*th coordinates of the points in the *i*th view.

We will use the following result later in the text,

Proposition 2. $\ker(\mathcal{L}_{\Gamma}) \subseteq \ker(\mathbf{B}).$

Assumption 1. The dimension of the kernel of \mathcal{L}_{Γ} equals the number of connected components in Γ . To keep the subsequent calculations simple, we assume that Γ is connected and thus ker(\mathcal{L}_{Γ}) is the span of a single vector $\mathbf{1}_{n+m}$.

It is clear from Eq. (3) that the the optimal **H** satisfies $\mathbf{H}^* \mathcal{L}_{\Gamma} = \mathbf{S}^T \mathbf{B}$. Using the fact that $\mathcal{L}_{\Gamma}^{\dagger} \mathcal{L}_{\Gamma} = \mathcal{L}_{\Gamma} \mathcal{L}_{\Gamma}^{\dagger} = \mathbf{I}_{n+m} - (n+m)^{-1} \mathbf{1}_{n+m} \mathbf{1}_{n+m}^T$, the solution \mathbf{H}^* is of the form $\mathbf{S}^T \mathbf{B} \mathcal{L}_{\Gamma}^{\dagger} - \mathbf{h} \mathbf{1}_{n+m}^T$ for some translation vector $\mathbf{h} \in \mathbb{R}^d$. Since $\mathbf{H} \mathbf{1}_n^{n+m} = 0$, the optimal value of **h** is $\mathbf{S}^T \mathbf{B} \mathcal{L}_{\Gamma}^{\dagger} \mathbf{1}_n^{n+m} / n$. Substituting back,

$$\mathbf{H}^* = \mathbf{S}^T \mathbf{B} \mathcal{L}_{\Gamma}^{\dagger} \left(\mathbf{I}_{n+m} - \frac{\mathbf{1}_n^{n+m} \mathbf{1}_{n+m}^T}{n} \right).$$
(7)

Substituting Eq. (7) back in Eq. (3), the problem reduces to

$$\mathcal{A}_1 = \min_{\mathbf{S} \in \mathbb{O}(d)^m} F(\mathbf{S}) = \min_{\mathbf{S} \in \mathbb{O}(d)^m} \operatorname{Tr}(\mathbf{C}\mathbf{S}\mathbf{S}^T) \text{ where } \mathbf{C} = \mathbf{D} - \mathbf{B}\mathcal{L}_{\Gamma}^{\dagger}\mathbf{B}^T.$$
(8)

Here \mathbf{C} is named the *patch-stress matrix* and is positive semidefinite [3].

Definition 1. Since the objective F depends only on \mathbf{S} , we identify an alignment of local views with an element of $\mathbb{O}(d)^m$. If \mathbf{S} is a global minimum of F then the alignment is called an optimal alignment. If \mathbf{S} is such that $F(\mathbf{S}) = 0$ then the alignment \mathbf{S} is called a perfect alignment.

Since $\mathbf{C} \succeq 0$, every perfect alignment is optimal, while the converse may not hold.

Definition 2. The consensus representation of the framework Θ due to an alignment **S** is given by

$$\Theta(\mathbf{S}) \coloneqq \mathbf{H}^*(:, 1:n) = \mathbf{S}^T \mathbf{B} \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger}(:, 1:n) \left(\mathbf{I}_n - n^{-1} \mathbf{1}_n \mathbf{1}_n^T \right).$$

The representation due to a perfect alignment is called a realization of the framework [12].

3. Non-degeneracy and Uniqueness in the General Setting

The organization of the section is as follows. In Section 3.1 we derive an expression involving the Hessian of \tilde{F} (Eq. (20), (25)), the function induced by F on a certain quotient space $\mathbb{O}(d)^m / \sim$ (along with all the necessary geometrical objects). We refer the reader to [1, Chapter 3 and 5] for the definitions of differential of a mapping, metric, gradient, connection, Hessian etc. in the context of Riemannian manifolds. Then in Section 3.2 we obtain the equations governing the non-singularity and positivity of the Hessian (Eq. (27), (30)). Subsequently, in Section 3.3, we obtain a characterization of the non-degeneracy of an alignment in the general setting (Theorem 13). We also present a necessary and sufficient condition on the overlapping structure of two views under which an alignment of the views is non-degenerate (Theorem 16). Finally, in Section 3.4, we derive a similar condition under which an optimal alignment of two views is unique (Theorem 17).

3.1 Preliminaries

Recall that the problem under consideration is the minimization of $F(\mathbf{S}) = \text{Tr}(\mathbf{CSS}^T)$ over $\mathbf{S} \in \mathbb{O}(d)^m$ where $\mathbf{C} \succeq 0$ is the patch-stress matrix defined in Eq. (8). Note that the objective is invariant to the action of $\mathbb{O}(d)$ i.e. for any $\mathbf{Q} \in \mathbb{O}(d)$, $F(\mathbf{S}) = F(\mathbf{SQ})$.

Assumption 2. If Γ has K connected components, then the objective is invariant to the action of $\mathbb{O}(d)^K$ on $\mathbb{O}(d)^m = \prod_1^K \mathbb{O}(d)^{m_j}$ where $\sum_1^K m_j = m$ and where each $\mathbb{O}(d)$ acts independently on $\mathbb{O}(d)^{m_j}$ for $j \in [1, K]$. To keep the computations clean, we assume that the bipartite graph Γ is connected (as in Assumption 1) throughout the rest of this work.

Subsequently, we define an equivalence relation on $\mathbb{O}(d)^m$; $\mathbf{S}_1 \sim \mathbf{S}_2$ if and only if $\mathbf{S}_1 = \mathbf{S}_2 \mathbf{Q}$ for some $\mathbf{Q} \in \mathbb{O}(d)$. Given $\mathbf{S} \in \mathbb{O}(d)^m$, its equivalence class is $[\mathbf{S}] = {\mathbf{S}\mathbf{Q} : \mathbf{Q} \in \mathbb{O}(d)}$. Clearly, there exists a bijection between $\mathbb{O}(d)^m / \sim$ and $\mathbb{O}(d)^{m-1}$, thus an element of $\mathbb{O}(d)^m / \sim$ will be identified with an element of $\mathbb{O}(d)^{m-1}$. Define the projection,

$$\pi: \mathbb{O}(d)^m \mapsto \mathbb{O}(d)^m / \sim$$
$$\pi(\mathbf{S}_{1:m}) = \mathbf{S}_{2:m} \mathbf{S}_1^T.$$

Let $\widetilde{\mathbf{S}} \in \mathbb{O}(d)^m / \sim$, then

$$\pi^{-1}(\widetilde{\mathbf{S}}) = \left\{ \begin{bmatrix} \mathbf{Q} \\ \widetilde{\mathbf{S}}\mathbf{Q} \end{bmatrix} : \mathbf{Q} \in \mathbb{O}(d) \right\} = \{ \mathbf{S} \in \mathbb{O}(d)^m : \mathbf{S}_{i+1}\mathbf{S}_1^T = \widetilde{\mathbf{S}}_i, i \in [1, m-1] \}.$$
(9)

The Riemannian metric g on $\mathbb{O}(d)^m$ is the canonical one given by

$$g(\mathbf{Z}, \mathbf{W}) \coloneqq \operatorname{Tr}(\mathbf{Z}^T \mathbf{W}) = \sum_{1}^{m} \operatorname{Tr}(\mathbf{Z}_i^T \mathbf{W}_i) \text{ where } \mathbf{Z}, \mathbf{W} \in T_{\mathbf{S}} \mathbb{O}(d)^m \subseteq \mathbb{R}^{md \times d}.$$
(10)

By a simple extension of the m = 1 case [1], it is easy to deduce the following result.

Proposition 3. For $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$, the tangent space to $\mathbb{O}(d)^m$ at \mathbf{S} is

$$T_{\mathbf{S}}\mathbb{O}(d)^m = \{ [\mathbf{S}_i \mathbf{\Omega}_i]_1^m : \mathbf{\Omega}_i \in \text{Skew}(d) \}.$$
(11)

The orthogonal projection of $\boldsymbol{\xi} = [\boldsymbol{\xi}_i]_1^m$, where $\boldsymbol{\xi}_i \in \mathbb{R}^{d \times d}$, onto $T_{\mathbf{S}} \mathbb{O}(d)^m$ is

$$P_{\mathbf{S}}(\boldsymbol{\xi}) = \underset{\substack{[\mathbf{S}_{i} \mathbf{\Omega}_{i}]_{1}^{m} \\ \mathbf{\Omega}_{i} \in \text{Skew}(d)}}{\operatorname{argmin}} \sum_{1}^{m} \|\boldsymbol{\xi}_{i} - \mathbf{S}_{i} \mathbf{\Omega}_{i}\|_{F}^{2} = [\mathbf{S}_{i} \text{Skew}(\mathbf{S}_{i}^{T} \boldsymbol{\xi}_{i})]_{1}^{m}$$

Then, $\pi^{-1}(\widetilde{\mathbf{S}})$ admits a tangent space at $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$ called the vertical space $\mathcal{V}_{\mathbf{S}}$ at \mathbf{S} . The horizontal space $\mathcal{H}_{\mathbf{S}}$ at \mathbf{S} is the subspace of $T_{\mathbf{S}}\mathbb{O}(d)^m$ that is the orthogonal complement to the vertical space $\mathcal{V}_{\mathbf{S}}$.

Proposition 4. The vertical space $\mathcal{V}_{\mathbf{S}}$ at $\mathbf{S} \in \pi^{-1}(\mathbf{\tilde{S}})$ is

$$\mathcal{V}_{\mathbf{S}} = \{\mathbf{S}\mathbf{\Omega} : \mathbf{\Omega} \in \operatorname{Skew}(d)\}$$

The orthogonal projection of $\mathbf{Z} = [\mathbf{S}_i \mathbf{\Omega}_i]_1^m \in T_{\mathbf{S}} \mathbb{O}(d)^m$ onto $\mathcal{V}_{\mathbf{S}}$ is

$$P_{\mathbf{S}}^{v}([\mathbf{S}_{i}\mathbf{\Omega}_{i}]_{1}^{m}) = \left[\mathbf{S}_{i} \operatorname{argmin}_{\mathbf{\Omega}\in\operatorname{Skew}(d)} \sum_{1}^{m} \|\mathbf{S}_{j}(\mathbf{\Omega}_{j}-\mathbf{\Omega})\|_{F}^{2}\right]_{1}^{m} = \left[\mathbf{S}_{i}\left(\frac{\sum_{1}^{m}\mathbf{\Omega}_{i}}{m}\right)\right]_{1}^{m}.$$
 (12)

The horizontal space $\mathcal{H}_{\mathbf{S}}$ at $\mathbf{S} \in \pi^{-1}(\mathbf{\tilde{S}})$ is

$$\mathcal{H}_{\mathbf{S}} = \left\{ [\mathbf{S}_i \mathbf{\Omega}_i]_1^m, \mathbf{\Omega}_i \in \text{Skew}(d), \sum_1^m \mathbf{\Omega}_i = 0 \right\}.$$

The orthogonal projection of $\mathbf{Z} = [\mathbf{S}_i \mathbf{\Omega}_i]_1^m \in T_{\mathbf{S}} \mathbb{O}(d)^m$ to $\mathcal{H}_{\mathbf{S}}$ is

$$P_{\mathbf{S}}^{h}([\mathbf{S}_{i}\boldsymbol{\Omega}_{i}]_{1}^{m}) = [\mathbf{S}_{i}\boldsymbol{\Omega}_{i}]_{1}^{m} - P_{\mathbf{S}}^{v}([\mathbf{S}_{i}\boldsymbol{\Omega}_{i}]_{1}^{m}) = \left[\mathbf{S}_{i}\left(\boldsymbol{\Omega}_{i} - \frac{\sum_{1}^{m}\boldsymbol{\Omega}_{i}}{m}\right)\right]_{1}^{m}$$
(13)

Note that $T_{\mathbf{S}}\mathbb{O}(d)^m$ is a vector space of dimension md(d-1)/2 and $\mathcal{V}_{\mathbf{S}}$ forms a d(d-1)/2dimensional subspace of $T_{\mathbf{S}}\mathbb{O}(d)^m$. The dimension of $\mathcal{H}_{\mathbf{S}}$ and of $T_{\widetilde{\mathbf{S}}}\mathbb{O}(d)^m/\sim$ is (m-1)d(d-1)/2. In particular, $\mathcal{H}_{\mathbf{S}}$ can be identified with $T_{\widetilde{\mathbf{S}}}\mathbb{O}(d)^m/\sim$. Let $\widetilde{\mathbf{S}} \in \mathbb{O}(d)^m/\sim$ and $\widetilde{\mathbf{Z}} \in T_{\widetilde{\mathbf{S}}}\mathbb{O}(d)^m/\sim$. Then the horizontal lift of $\widetilde{\mathbf{Z}}$ at $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$ is defined as $\overline{\widetilde{\mathbf{Z}}} \in \mathcal{H}_{\mathbf{S}}$ such that for each $i \in [1, m-1]$,

$$D\pi[\mathbf{S}]\left(\widetilde{\mathbf{Z}}\right)_{i} = \widetilde{\mathbf{Z}}_{i}.$$
(14)

Proposition 5. Let $\widetilde{\mathbf{S}} \in \mathbb{O}(d)^m / \sim$ and $\widetilde{\mathbf{Z}} \in T_{\widetilde{\mathbf{S}}} \mathbb{O}(d)^m / \sim$. Let \mathbf{Z} be the horizontal lift of $\widetilde{\mathbf{Z}}$ at $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$. If $(\widetilde{\mathbf{\Omega}}_i)_1^{m-1} \subseteq \text{Skew}(d)$ are such that $\widetilde{\mathbf{Z}}_i = \widetilde{\mathbf{S}}_i \widetilde{\mathbf{\Omega}}_i$, and $(\mathbf{\Omega}_i)_1^m \subseteq \text{Skew}(d)$ are such that $\mathbf{Z}_i = \mathbf{S}_i \mathbf{\Omega}_i$ and $\sum_{1}^m \mathbf{\Omega}_i = 0$, then

$$\mathbf{\Omega}_{1} = -\frac{1}{m} \mathbf{S}_{1}^{T} \left(\sum_{1}^{m-1} \widetilde{\mathbf{\Omega}}_{i} \right) \mathbf{S}_{1}$$

$$(15)$$

$$\mathbf{\Omega}_{i+1} = \mathbf{S}_1^T \widetilde{\mathbf{\Omega}}_i \mathbf{S}_1 + \mathbf{\Omega}_1 \text{ for all } i \in [1, m-1].$$
(16)

Moreover, the linear system above has full rank, and thus the horizontal lift \mathbf{Z} of $\widetilde{\mathbf{Z}}$ at $\mathbf{S} \in \mathbb{O}(d)^m$ is a unique element of $\mathcal{H}_{\mathbf{S}}$.

Proposition 6. Let $\widetilde{\mathbf{Z}}, \widetilde{\mathbf{W}} \in T_{\widetilde{\mathbf{S}}} \mathbb{O}(d)^m / \sim$ and $\mathbf{Z}, \mathbf{W} \in T_{\mathbf{S}} \mathbb{O}(d)^m$ be their the horizontal lifts at $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$. Then

$$\widetilde{g}(\widetilde{\mathbf{Z}},\widetilde{\mathbf{W}})\coloneqq g(\mathbf{Z},\mathbf{W})$$

defines a Riemannian metric on $\mathbb{O}(d)^m / \sim$.

We note that $\mathcal{H}_{\mathbf{S}}$ with the canonical metric g, is isometric to $T_{\mathbf{S}}\mathbb{O}(d)^m/\sim$ (equivalently $T_{\mathbf{S}}\mathbb{O}(d)^{m-1}$) when equipped with the above metric \tilde{g} .

Now, coming back to the alignment error $F(\mathbf{S}) = \text{Tr}(\mathbf{C}\mathbf{S}\mathbf{S}^T)$ defined on $\mathbb{O}(d)^m$. It induces the following function on $\mathbb{O}(d)^m / \sim$ (again identified with $\mathbb{O}(d)^{m-1}$),

$$\widetilde{F}(\widetilde{\mathbf{S}}) = \operatorname{Tr}\left(\mathbf{C}\begin{bmatrix}\mathbf{I}_d\\\widetilde{\mathbf{S}}\end{bmatrix}\begin{bmatrix}\mathbf{I}_d\\\widetilde{\mathbf{S}}\end{bmatrix}^T\right).$$

In particular, $F = \widetilde{F} \circ \pi$. It is easy to see that if $\mathbf{S}_1 \sim \mathbf{S}_2$ then $\widetilde{F} \circ \pi(\mathbf{S}_1) = \widetilde{F} \circ \pi(\mathbf{S}_2)$.

Proposition 7. The horizontal lift of $\operatorname{grad} \widetilde{F}(\widetilde{\mathbf{S}})$ at $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$ is

$$\operatorname{grad} \widetilde{F}(\widetilde{\mathbf{S}}) = \operatorname{grad} F(\mathbf{S}) = [\mathbf{S}_i \mathbf{\Omega}_i]_1^m$$
(17)

where $\Omega_i \coloneqq \mathbf{S}_i^T[\mathbf{CS}]_i - [\mathbf{CS}]_i^T \mathbf{S}_i \in \text{Skew}(d) \text{ and } \sum_{1}^m \Omega_i = \mathbf{S}^T \mathbf{CS} - \mathbf{S}^T \mathbf{CS} = 0 \text{ (which validates that } \overline{\text{grad}} \widetilde{F}(\widetilde{\mathbf{S}}) \text{ is indeed in } \mathcal{H}_{\mathbf{S}}, \text{ see Proposition 4).}$

Due to the above proposition, the set of critical points of \widetilde{F} is given by

$$\widetilde{\mathcal{C}} = \{ \widetilde{\mathbf{S}} \in \mathbb{O}(d)^m / \sim : \operatorname{grad} \widetilde{F}(\widetilde{\mathbf{S}}) = 0 \}$$

= $\{ \widetilde{\mathbf{S}} \in \mathbb{O}(d)^m / \sim : \mathbf{S}_i^T [\mathbf{CS}]_i = [\mathbf{CS}]_i^T \mathbf{S}_i, \text{ for all } i \in [1, m], \mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}}) \},$ (18)

and that of F is,

$$\mathcal{C} = \{ \mathbf{S} \in \mathbb{O}(d)^m : \operatorname{grad} F(\mathbf{S}) = 0 \}$$

= $\{ \mathbf{S} \in \mathbb{O}(d)^m : \mathbf{S}_i^T [\mathbf{CS}]_i = [\mathbf{CS}]_i^T \mathbf{S}_i, \text{ for all } i \in [1, m] \}.$ (19)

The following remark follows trivially from Eq. (18) and (19).

Remark 2. If **S** is a critical point of F i.e. $\mathbf{S} \in \mathcal{C}$ then $\pi(\mathbf{S})$ is a critical point of \widetilde{F} i.e. $\pi(\mathbf{S}) \in \widetilde{C}$. Similarly, if $\widetilde{\mathbf{S}} \in \widetilde{C}$ then $\mathbf{S} \in \mathcal{C}$ for all $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$.

Proposition 8. Let $\widetilde{\mathbf{S}} \in \widetilde{\mathcal{C}}$ then for every $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$ and $\mathbf{Z} \in T_{\mathbf{S}} \mathbb{O}(d)^m$,

$$D \operatorname{grad} F(\mathbf{S})[\mathbf{Z}] = \left[\mathbf{S}_i (\mathbf{S}_i^T [\mathbf{C}\mathbf{Z}]_i - [\mathbf{C}\mathbf{Z}]_i^T \mathbf{S}_i - [\mathbf{C}\mathbf{S}]_i^T \mathbf{Z}_i + \mathbf{Z}_i^T [\mathbf{C}\mathbf{S}]_i)\right]_1^m$$

Let ∇ be the Levi-Civita connection (also known as the Riemannian connection) on $\mathbb{O}(d)^m$ and $\widetilde{\nabla}$ be the induced connection on $\mathbb{O}(d)^m / \sim$. Then the following result follows from Proposition 8 and the definition of the Riemannian Hessian operator [1, Section 5.5].

Proposition 9. Let $\widetilde{\mathbf{S}} \in \widetilde{\mathcal{C}}$. Let $\widetilde{\mathbf{Z}} \in T_{\widetilde{\mathbf{S}}} \mathbb{O}(d)^m / \sim$. Let \mathbf{Z} be the horizontal lift of $\widetilde{\mathbf{Z}}$ at $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$. Then the horizontal lift of $\operatorname{Hess}\widetilde{F}(\widetilde{\mathbf{S}})[\widetilde{\mathbf{Z}}]$ at \mathbf{S} is

$$\operatorname{Hess}\widetilde{F}(\widetilde{\mathbf{S}})[\widetilde{\mathbf{Z}}] = [\mathbf{S}_i\widehat{\mathbf{\Omega}}_i]_1^m \tag{20}$$

where

$$\widehat{\mathbf{\Omega}}_i = \mathbf{S}_i^T [\mathbf{C}\mathbf{Z}]_i - [\mathbf{C}\mathbf{Z}]_i^T \mathbf{S}_i - [\mathbf{C}\mathbf{S}]_i^T \mathbf{Z}_i + \mathbf{Z}_i^T [\mathbf{C}\mathbf{S}]_i.$$

In the following, we obtain a compact representation for $\widehat{\Omega}_i$. We first define certain matrices, then use them to obtain an expression for $\widehat{\Omega}_i$ and then describe their structure. Recall that $\mathbf{C} = \mathbf{D} - \mathbf{B} \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}^T$ (see Eq. (8) and Remark 1) and for convenience define

$$\mathbf{B}(\mathbf{S}) \coloneqq \text{block-diag}((\mathbf{S}_i)_1^m)^T \ \mathbf{B} = [\mathbf{S}_i^T \mathbf{B}_i]_1^m$$

$$\mathbf{D}(\mathbf{S}) \coloneqq \text{block-diag}((\mathbf{S}_i)_1^m)^T \ \mathbf{D} \ \text{block-diag}((\mathbf{S}_i)_1^m) = \text{block-diag}((\mathbf{S}_i^T \mathbf{D}_{ii} \mathbf{S}_i)_1^m).$$
(21)

Then define

$$\mathbf{C}(\mathbf{S}) \coloneqq \text{block-diag}((\mathbf{S}_i)_1^m)^T \ \mathbf{C} \ \text{block-diag}((\mathbf{S}_i)_1^m) = \mathbf{D}(\mathbf{S}) - \mathbf{B}(\mathbf{S})\boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger}\mathbf{B}(\mathbf{S})^T$$
(22)

$$\widehat{\mathbf{C}}(\mathbf{S}) \coloneqq \text{block-diag}(([\mathbf{C}(\mathbf{S})\mathbf{I}_d^m]_i)_1^m)$$
(23)

$$\mathbf{L}(\mathbf{S}) \coloneqq \widehat{\mathbf{C}}(\mathbf{S}) - \mathbf{C}(\mathbf{S}).$$
(24)

Proposition 10. Consider the same setup as in Proposition 9. Then

$$\widehat{\mathbf{\Omega}}_{i} = [\mathbf{L}(\mathbf{S})\mathbf{\Omega}]_{i}^{T} - [\mathbf{L}(\mathbf{S})\mathbf{\Omega}]_{i}$$
(25)

where $\mathbf{\Omega} = [\mathbf{\Omega}_i]_1^m$ and $\mathbf{\Omega}_i \in \text{Skew}(d)$ is such that $\mathbf{Z}_i = \mathbf{S}_i \mathbf{\Omega}_i$ and $\sum_{i=1}^m \mathbf{\Omega}_i = 0$.

Combining the above equation with Proposition 9, we obtain a slightly compact representation of the horizontal lift of the Hessian. We end this subsection by giving remarks that reveal the structure of $\mathbf{C}(\mathbf{S})$, $\widehat{\mathbf{C}}(\mathbf{S})$ and $\mathbf{L}(\mathbf{S})$.

Remark 3. Since block-diag $((\mathbf{S}_i)_1^m)$ is an orthogonal matrix, $\mathbf{C}(\mathbf{S})$ is unitarily equivalent to \mathbf{C} . Thus, $\mathbf{C}(\mathbf{S}) \in \text{Sym}(md)$, $\mathbf{C}(\mathbf{S}) \succeq 0$, $\text{rank}(\mathbf{C}(\mathbf{S})) = \text{rank}(\mathbf{C})$ and the (i, j)th block of size d in $\mathbf{C}(\mathbf{S})$ is $\mathbf{C}(\mathbf{S})_{ij} = \delta_{ij}\mathbf{D}(\mathbf{S})_{ii} - \mathbf{B}(\mathbf{S})_i \mathcal{L}_{\Gamma}^{\dagger}\mathbf{B}(\mathbf{S})_j^T$ where $\mathbf{B}(\mathbf{S})_i$ is the ith row block of $\mathbf{B}(\mathbf{S})$ of dimension $d \times (m + n)$.

Remark 4. From the definition of C (see Eq. (19)), $\mathbf{S} \in C$ if and only if $\widehat{\mathbf{C}}(\mathbf{S}) \in \text{Sym}(md)$ i.e. for each $i \in [1, m]$,

$$[\mathbf{C}(\mathbf{S})\mathbf{I}_d^m]_i = \sum_{j=1}^m \mathbf{C}(\mathbf{S})_{ij} = \mathbf{S}_i^T [\mathbf{C}\mathbf{S}]_i = [\mathbf{C}\mathbf{S}]_i^T \mathbf{S}_i = \sum_{j=1}^m \mathbf{C}(\mathbf{S})_{ij}^T = [\mathbf{C}(\mathbf{S})\mathbf{I}_d^m]_i^T.$$
 (26)

Remark 5. If $\widetilde{\mathbf{S}} \in \widetilde{\mathcal{C}}$ then for every $\mathbf{S} \in \pi^{-1}(\mathbf{S})$, the following are easy to deduce.

1. $\mathbf{L}(\mathbf{S}) \in \text{Sym}(md)$.

2.
$$\sum_{j=1}^{m} \mathbf{L}(\mathbf{S})_{ij} = \sum_{j=1}^{m} \mathbf{L}(\mathbf{S})_{ji} = 0$$
 for all $i \in [1, m]$ (see Eq. (26)).

- 3. For each $i \in [1, d]$, the vector $\mathbf{1}_m \otimes \mathbf{e}_i^d$ lies in the kernel of $\mathbf{L}(\mathbf{S})$ and thus the rank of $\mathbf{L}(\mathbf{S})$ is at most (m-1)d.
- 4. If $\Omega = [\Omega_0]_1^m$ for some $\Omega_0 \in \text{Skew}(d)$ then $\mathbf{L}(\mathbf{S})\Omega = 0$.
- 5. The (i, j)th block of size d in $\mathbf{L}(\mathbf{S})$ is

$$\begin{split} \mathbf{L}(\mathbf{S})_{ij} &= -\delta_{ij} \mathbf{B}(\mathbf{S})_i \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})^T \mathbf{I}_d^m + \mathbf{B}(\mathbf{S})_i \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_j^T \\ &= -\delta_{ij} (\mathbf{I}_d^m)^T \mathbf{B}(\mathbf{S}) \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_i^T + \mathbf{B}(\mathbf{S})_i \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_j^T \end{split}$$

and in particular, $\mathbf{L}(\mathbf{S})$ does not depend on $\mathbf{D}(\mathbf{S})$.

6. Let $\mathbf{Q} \in \mathbb{O}(d)$ then $\mathbf{L}(\mathbf{SQ}) = (\mathbf{I}_m \otimes \mathbf{Q})^T \mathbf{L}(\mathbf{S})(\mathbf{I}_m \otimes \mathbf{Q})$ (follows from Eq. (22, 23, 26)) and since $\mathbf{I}_m \otimes \mathbf{Q}$ is an orthogonal matrix, $L(\mathbf{SQ})$ is unitarily equivalent to $\mathbf{L}(\mathbf{S})$.

3.2 Non-singular and Positive Definite Hessian

A non-degenerate local minimum is defined to be the critical point at which the Hessian is positive definite. Similarly, a non-degenerate critical point is the one where the Hessian is non-singular. We proceed to derive the equations to identify the conditions under which the Hessian is non-singular and positive definite, which in turn characterize the non-degenerate critical points and local minima. First, we need the following result. **Proposition 11.** Let $\widetilde{\mathbf{S}} \in \widetilde{\mathcal{C}}$. Let $\widetilde{\mathbf{Z}} \in T_{\widetilde{\mathbf{S}}} \mathbb{O}(d)^m / \sim$. Let \mathbf{Z} be the horizontal lift of $\widetilde{\mathbf{Z}}$ at $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$. Then

$$\widetilde{g}(\text{Hess}\widetilde{F}(\widetilde{\mathbf{S}})[\widetilde{\mathbf{Z}}],\widetilde{\mathbf{Z}}) = -2 \operatorname{Tr}(\mathbf{\Omega}^T \mathbf{L}(\mathbf{S})\mathbf{\Omega})$$
(27)

where $\mathbf{\Omega} = [\mathbf{\Omega}_i]_1^m$ and $\mathbf{\Omega}_i \in \text{Skew}(d)$ is such that $\mathbf{Z}_i = \mathbf{S}_i \mathbf{\Omega}_i$ and $\sum_{i=1}^m \mathbf{\Omega}_i = 0$.

Then the non-singularity and the positive definiteness of the Hessian amounts to the right side of Eq. (27) being non-zero and positive, respectively, for every non-zero Ω . Although $\mathbf{C}(\mathbf{S})$, and thus $\mathbf{L}(\mathbf{S})$, can be calculated from the patch framework Θ and the alignment \mathbf{S} , it is not obvious how to test the above practically. The main reason being that Ω in Eq. (27) is not unconstrained, and in fact has a specific structure.

Therefore, for the above reason, we are going to manipulate Eq. (27), utilizing the structure of Ω . The aim is to obtain an expression of the form $\boldsymbol{\omega}^T \mathbb{L}(\mathbf{S})\boldsymbol{\omega}$ where (i) the vector $\boldsymbol{\omega}$ is essentially unconstrained and (ii) $\boldsymbol{\Omega}$ and $\mathbf{L}(\mathbf{S})$ are related to $\boldsymbol{\omega}$ and $\mathbb{L}(\mathbf{S})$, respectively, through permutation matrices and vectorization operations (and thus the two pairs carry the same information). To achieve that, we first define certain matrices, then rewrite Eq. (27) in terms of those matrices and then describe their structure.

To this end, for $\mathbf{\Omega} = [\mathbf{\Omega}_i]_1^m$ where $\mathbf{\Omega}_i \in \text{Skew}(d)$, let $\{\mathbf{\Omega}_i(r,s) : 1 \leq r < s \leq d\}$ be the elements in the upper triangular region of $\mathbf{\Omega}_i$. For a fixed pair (r,s) such that $1 \leq r < s \leq d$, define the column vector $\boldsymbol{\omega}_{r,s} \coloneqq [\mathbf{\Omega}_i(r,s)]_{i=1}^m \in \mathbb{R}^m$, a vertical stack of the (r,s)th element of each $\mathbf{\Omega}_i$. Then there exists a permutation matrix \mathbf{P} such that

$$\mathbf{P}\mathbf{\Omega} = egin{bmatrix} \mathbf{0}_m & oldsymbol{\omega}_{1,2} & \dots & oldsymbol{\omega}_{1,d-1} & oldsymbol{\omega}_{1,d} \ -oldsymbol{\omega}_{1,2} & \mathbf{0}_m & \dots & oldsymbol{\omega}_{2,d-1} & oldsymbol{\omega}_{2,d} \ dots & dots & dots & dots & dots & dots \ -oldsymbol{\omega}_{1,d-1} & -oldsymbol{\omega}_{2,d-1} & \dots & oldsymbol{0}_m & oldsymbol{\omega}_{d-1,d} \ -oldsymbol{\omega}_{1,d-1} & -oldsymbol{\omega}_{2,d} & \dots & -oldsymbol{\omega}_{d-1,d} & oldsymbol{0}_m \end{bmatrix}.$$

In words, for $1 \leq r < s \leq d$, the (r, s)th block of $\mathbf{P}\Omega$ is a vertical stack of the (r, s)th element of each Ω_i . For r = s, this is just a zero vector and for r > s, this is $-\omega_{s,r}$.

Then, we collect the (strictly) upper triangular elements of $\mathbf{P}\mathbf{\Omega}$ in the column-major order in the vector $\boldsymbol{\omega}$. Note that $\mathbf{P}\mathbf{\Omega}$ can be fully described by $\boldsymbol{\omega}$. In particular, there exist a block matrix $\overline{\mathbf{P}}$ of size $d(d-1)/2 \times d^2$ such that $\operatorname{vec}(\mathbf{P}\mathbf{\Omega}) = \overline{\mathbf{P}}^T \boldsymbol{\omega}$. The blocks of $\overline{\mathbf{P}}$ when indexed using tuples (r, s) and (p, q) where $1 \leq r < s \leq d$ and $p, q \in [1, d]$, are given by,

$$\overline{\mathbf{P}}_{(r,s),(p,q)} = \begin{cases} \mathbf{0}_{m \times m}, & p = q\\ \delta_{pr} \delta_{qs} \mathbf{I}_m, & p < q\\ -\delta_{ps} \delta_{qr} \mathbf{I}_m, & p > q. \end{cases}$$

Finally, we define

$$\mathcal{B}(\mathbf{S}) \coloneqq \mathbf{PB}(\mathbf{S})$$
$$\mathcal{L}(\mathbf{S}) \coloneqq \mathbf{PL}(\mathbf{S})\mathbf{P}^T$$
(28)

$$\mathbb{L}(\mathbf{S}) \coloneqq \overline{\mathbf{P}}(\mathbf{I}_d \otimes \mathcal{L}(\mathbf{S})) \overline{\mathbf{P}}^T$$
(29)

and then note the following,

Proposition 12. Consider the same setup as in Proposition 11. Then

$$Tr(\mathbf{\Omega}^T \mathbf{L}(\mathbf{S})\mathbf{\Omega}) = \boldsymbol{\omega}^T \mathbb{L}(\mathbf{S})\boldsymbol{\omega}$$
(30)

The following remarks reveal the structure of $\mathcal{B}(\mathbf{S})$, $\mathcal{L}(\mathbf{S})$ and $\mathbb{L}(\mathbf{S})$.

Remark 6. For $p \in [1, d]$, the pth row-block of $\mathcal{B}(\mathbf{S})$, $\mathcal{B}(\mathbf{S})_p$, is of size $m \times (n + m)$, and can be viewed as a vertical stack of the pth row of each $\mathbf{B}(\mathbf{S})_i$, $i \in [1, m]$. In particular, $\mathcal{B}(\mathbf{S})_p$ depends only on the pth coordinate of the local views (also see Remark 1).

Remark 7. If $\widetilde{\mathbf{S}} \in \widetilde{\mathcal{C}}$ then for every $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$, the following are easy to deduce.

1. For $p, q \in [1, d]$, it is easy to deduce

$$\mathcal{L}(\mathbf{S})_{p,q} = -\mathrm{diag}(\mathcal{B}(\mathbf{S})_p \mathcal{L}_{\Gamma}^{\dagger} \mathcal{B}(\mathbf{S})_q^T \mathbf{1}_m) + \mathcal{B}(\mathbf{S})_p \mathcal{L}_{\Gamma}^{\dagger} \mathcal{B}(\mathbf{S})_q^T$$

Thus $\mathcal{L}(\mathbf{S})_{p,q}$ depends on Γ through $\mathcal{L}_{\Gamma}^{\dagger}$ and the pth and qth coordinates of the rigidly transformed local views $\mathbf{B}(\mathbf{S})$.

- 2. Since $\mathbf{L}(\mathbf{S})$ is symmetric, $\mathcal{L}(\mathbf{S})_{q,p} = \mathcal{L}(\mathbf{S})_{p,q}^T$.
- 3. $\mathcal{L}(\mathbf{S})_{p,q}$ is a Laplacian-like matrix and the constant vectors are in its kernel.
- 4. For each $p \in [1,d]$, the vector $\mathbf{e}_p^d \otimes \mathbf{1}_m$ lies in the kernel of $\mathcal{L}(\mathbf{S})$, thus the rank of $\mathcal{L}(\mathbf{S})$ is at most (m-1)d.
- 5. If $\mathbf{Q} \in \mathbb{O}(d)$ then, from Remark 5 and Eq. (28), it follows that $\mathcal{L}(\mathbf{SQ})$ is unitarily equivalent to $\mathcal{L}(\mathbf{S})$.

Remark 8. If $\widetilde{\mathbf{S}} \in \widetilde{\mathcal{C}}$ then for every $\mathbf{S} \in \pi^{-1}(\mathbf{S})$, the following are easy to deduce.

1. $\mathbb{L}(\mathbf{S})$ is a block matrix of size d(d-1)/2 where each block is of size m. Indexing the rows and columns of \mathbb{L} by tuples of the form (r, s) where $1 \leq r < s \leq d$ we have,

$$\begin{split} \mathbb{L}(\mathbf{S})_{(r_{1},s_{1}),(r_{2},s_{2})} &= \sum_{\substack{p_{1},q_{1} \in [1,d] \\ p_{2},q_{2} \in [1,d]}} \overline{\mathbf{P}}_{(r_{1},s_{1}),(p_{1},q_{1})} (\mathbf{I}_{d} \otimes \boldsymbol{\mathcal{L}}(\mathbf{S}))_{(p_{1},q_{1}),(p_{2},q_{2})} \overline{\mathbf{P}}_{(r_{2},s_{2}),(p_{2},q_{2})}^{T} \\ &= \sum_{\substack{p,q_{1},q_{2} \in [1,d] \\ p,q_{1},q_{2} \in [1,d]}} \overline{\mathbf{P}}_{(r_{1},s_{1}),(p,q_{1})} \boldsymbol{\mathcal{L}}(\mathbf{S})_{q_{1},q_{2}} \overline{\mathbf{P}}_{(r_{2},s_{2}),(p,q_{2})}^{T} \\ &= \begin{cases} \boldsymbol{\mathcal{L}}(\mathbf{S})_{r,r} + \boldsymbol{\mathcal{L}}(\mathbf{S})_{s,s}, & r_{1} = r_{2} = r, s_{1} = s_{2} = s \\ 0, & \{r_{1},s_{1}\} \cap \{r_{2},s_{2}\} = \emptyset \\ \boldsymbol{\mathcal{L}}(\mathbf{S})_{s_{1},s_{2}}, & r_{1} = r_{2}, s_{1} \neq s_{2} \\ \boldsymbol{\mathcal{L}}(\mathbf{S})_{r_{1},r_{2}}, & s_{1} = s_{2}, r_{1} \neq r_{2} \\ -\boldsymbol{\mathcal{L}}(\mathbf{S})_{r_{1},s_{2}}, & s_{1} = r_{2} \\ -\boldsymbol{\mathcal{L}}(\mathbf{S})_{s_{1},r_{2}}, & s_{2} = r_{1}. \end{cases} \end{split}$$

- 2. Since $\mathcal{L}(\mathbf{S})$ is symmetric, $\mathbb{L}(\mathbf{S}) = \mathbb{L}(\mathbf{S})^T$.
- 3. The set of vectors of the form $\boldsymbol{\omega} = [\boldsymbol{\omega}_{r,s}]_{1 \leq r < s \leq d}$ where each $\boldsymbol{\omega}_{r,s}$ is a constant vector, lie in the kernel of $\mathbb{L}(\mathbf{S})$.

- 4. The rank of $\mathbb{L}(\mathbf{S})$ is at most (m-1)d(d-1)/2.
- 5. If $\mathbf{Q} \in \mathbb{O}(d)$ then $\mathbb{L}(\mathbf{SQ})$ is unitarily equivalent to $\mathbb{L}(\mathbf{S})$.

Remark 9. For d = 2, 3 and 4, $\mathbb{L}(\mathbf{S})$ is given by (for brevity, we use \mathcal{L} in place of $\mathcal{L}(\mathbf{S})$)

 $[\mathcal{L}_{1,1} + \mathcal{L}_{2,2}],$

$$\begin{bmatrix} \mathcal{L}_{1,1} + \mathcal{L}_{2,2} & \mathcal{L}_{2,3} & -\mathcal{L}_{1,3} \\ \mathcal{L}_{3,2} & \mathcal{L}_{1,1} + \mathcal{L}_{3,3} & \mathcal{L}_{1,2} \\ -\mathcal{L}_{3,1} & \mathcal{L}_{2,1} & \mathcal{L}_{2,2} + \mathcal{L}_{3,3} \end{bmatrix}$$

and

$$\begin{bmatrix} \mathcal{L}_{1,1} + \mathcal{L}_{2,2} & \mathcal{L}_{2,3} & \mathcal{L}_{2,4} & -\mathcal{L}_{1,3} & -\mathcal{L}_{1,4} & 0 \\ \mathcal{L}_{3,2} & \mathcal{L}_{1,1} + \mathcal{L}_{3,3} & \mathcal{L}_{3,4} & \mathcal{L}_{1,2} & 0 & -\mathcal{L}_{1,4} \\ \mathcal{L}_{4,2} & \mathcal{L}_{4,3} & \mathcal{L}_{1,1} + \mathcal{L}_{4,4} & 0 & \mathcal{L}_{1,2} & \mathcal{L}_{1,3} \\ -\mathcal{L}_{3,1} & \mathcal{L}_{2,1} & 0 & \mathcal{L}_{2,2} + \mathcal{L}_{3,3} & \mathcal{L}_{3,4} & -\mathcal{L}_{2,4} \\ -\mathcal{L}_{4,1} & 0 & \mathcal{L}_{2,1} & \mathcal{L}_{4,3} & \mathcal{L}_{2,2} + \mathcal{L}_{4,4} & \mathcal{L}_{2,3} \\ 0 & -\mathcal{L}_{4,1} & \mathcal{L}_{3,1} & -\mathcal{L}_{4,2} & \mathcal{L}_{3,2} & \mathcal{L}_{3,3} + \mathcal{L}_{4,4} \end{bmatrix}$$

respectively.

3.3 Non-degenerate Alignment in the General Setting

As argued in Section 2, since $F(\mathbf{S}) = F(\mathbf{SQ})$ for all $\mathbf{Q} \in \mathbb{O}(d)$, every alignment \mathbf{S} is degenerate in this sense. With a slight abuse of notation, we define a non-degenerate alignment as follows.

Definition 3. An alignment $\mathbf{S} \in \mathbb{O}(d)^m$ is non-degenerate if $\pi(\mathbf{S})$ is a non-degenerate local minimum of \widetilde{F} .

With the above definition, to characterize the non-degenerate alignments, it suffices to characterize the non-degenerate local minima of \tilde{F} . We accomplish the same in the following theorem. Note that we have not made any assumption about the affine non-degeneracy of the points and the noise in the local views.

Theorem 13. (Condition for $\widetilde{\mathbf{S}}$ to be a non-degenerate local minimum of \widetilde{F}). Let $\widetilde{\mathbf{S}} \in \widetilde{\mathcal{C}}$ and $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$. Then the following are equivalent,

- 1. $\widetilde{\mathbf{S}}$ is a non-degenerate local minimum of \widetilde{F} .
- 2. $\widetilde{g}(\operatorname{Hess}\widetilde{F}(\widetilde{\mathbf{S}})[\widetilde{\mathbf{Z}}],\widetilde{\mathbf{Z}}) > 0$ for all $\widetilde{\mathbf{Z}} \in T_{\widetilde{\mathbf{S}}}\mathbb{O}(d)^m / \sim$ such that $\widetilde{\mathbf{Z}} \neq 0$.
- 3. $Tr(\mathbf{\Omega}^T \mathbf{L}(\mathbf{S})\mathbf{\Omega}) < 0$ for all $\mathbf{\Omega} = [\mathbf{\Omega}_i]_1^m$ where $\mathbf{\Omega}_i \in \text{Skew}(d)$, $\sum_1^m \mathbf{\Omega}_i = 0$ and not all $\mathbf{\Omega}_i$ equal zero.
- 4. $Tr(\mathbf{\Omega}^T \mathbf{L}(\mathbf{S})\mathbf{\Omega}) < 0$ for all $\mathbf{\Omega} = [\mathbf{\Omega}_i]_1^m$ where $\mathbf{\Omega}_i \in \text{Skew}(d)$ and not all $\mathbf{\Omega}_i$ are equal.
- 5. $\boldsymbol{\omega}^T \mathbb{L}(\mathbf{S}) \boldsymbol{\omega} < 0$ for all $\boldsymbol{\omega} = [\boldsymbol{\omega}_{r,s}]_{1 \leq r < s \leq d}$ where not all $\boldsymbol{\omega}_{r,s}$ are constant vectors.

6. $\mathbb{L}(\mathbf{S})$ is negative semi-definite and of rank (m-1)d(d-1)/2.

Remark 10. Given the patch framework Θ and the alignment \mathbf{S} , one can compute the matrix $\mathbb{L}(\mathbf{S})$ in polynomial time in m, n and d, and then verify the non-degeneracy of the alignment \mathbf{S} by testing the last condition in the above theorem (which again requires polynomial time in m and d).

Although, for $\widetilde{\mathbf{S}}$ to be a non-degenerate local minimum of \widetilde{F} , the above theorem demands any of the equivalent conditions 3-6 to hold for every $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$, the following result shows that if a conditions hold for one $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$ then it holds for all other elements as well i.e. for all \mathbf{SQ} where $\mathbf{Q} \in \mathbb{O}(d)$ is arbitrary.

Proposition 14. Let $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$ and $\mathbf{Q} \in \mathbb{O}(d)$. Fix $i \in [3, 6]$. Suppose condition i in Theorem 13 holds for \mathbf{S} then it holds for \mathbf{SQ} also. Consequently, an alignment \mathbf{S} is non-degenerate if $\mathbf{S} \in \mathcal{C}$ (see Eq. (19)) and it satisfies any of the (equivalent) conditions 3-6 in Theorem 13.

A sufficient condition for \mathbf{S} to be a non-degenerate alignment is as follows.

Corollary 15. If $\mathbf{L}(\mathbf{S}) \leq 0$ and is of rank (m-1)d, then \mathbf{S} is a non-degenerate alignment. The same holds when $\mathbf{L}(\mathbf{S})$ is replaced by $\mathcal{L}(\mathbf{S})$ as the two are unitarily equivalent.

Note that the rank of $\mathbf{L}(\mathbf{S})$ being (m-1)d is not a necessary condition for non-degeneracy as demonstrated in Figure 1.



Figure 1: The dotted lines represent views and the filled points represent points on the overlaps. Here d = 2. It will be clear from Proposition 18 in Section 4 that $\mathbf{L}(\mathbf{S}) \leq 0$, and thus $\mathbb{L}(\mathbf{S}) \leq 0$. Through simple calculations one can deduce that the rank of $\mathbb{L}(\mathbf{S})$ is 3 (which equals (m-1)d(d-1)/2) while the rank of $\mathbf{L}(\mathbf{S})$ is 3 < 6 = (m-1)d.

We end this subsection by deriving a necessary and sufficient condition for an alignment of two views to be non-degenerate. First we need the following definitions (note that the objects in these definitions are related but not identical to \mathbf{B}_i (see Eq. (5), Remark 1) and $\mathbf{B}(\mathbf{S})_i$ (see Eq. (21), Remark 3)),

Definition 4. Let $i, j \in [1, m]$ be two views. Define $\mathbf{B}_{i,j}$ to be a matrix whose columns are $\mathbf{x}_{k,i}$ (in the increasing order of k) where $(k, i), (k, j) \in E(\Gamma)$. Generally, $\mathbf{B}_{i,j} \neq \mathbf{B}_{j,i}$. Also, define

$$\overline{\mathbf{B}}_{i,j} = \mathbf{B}_{i,j} \left(\mathbf{I}_{n'} - rac{\mathbf{1}_{n'} \mathbf{1}_{n'}^T}{n'}
ight)$$

where $n' = |\{k : (k, i), (k, j) \in E(\Gamma)\}|$ is the number of points on the overlap of the *i*th view and the *j*th view, or equivalently the number of columns in $\mathbf{B}_{i,j}$. **Definition 5.** Let **S** be an alignment. Let $i, j \in [1, m]$. Define $\mathbf{B}(\mathbf{S})_{i,j}$ to be a matrix whose columns are $\mathbf{S}_i^T \mathbf{x}_{k,i} + \mathbf{t}_i$ (in increasing order of k) where $(k, i), (k, j) \in E(\Gamma)$ and where \mathbf{t}_i is obtained using Eq. (7). Also, define

$$\overline{\mathbf{B}(\mathbf{S})}_{i,j} = \mathbf{B}(\mathbf{S})_{i,j} \left(\mathbf{I}_{n'} - \frac{\mathbf{1}_{n'} \mathbf{1}_{n'}^T}{n'} \right).$$

Remark 11. Let $i, j \in [1, m]$ and **S** be an alignment. Let $n' = |\{k : (k, i), (k, j) \in E(\Gamma)\}|$ be the number of points on the overlap of the two views. Then

$$\mathbf{B}(\mathbf{S})_{i,j} = \mathbf{S}_i^T \mathbf{B}_{i,j} + \mathbf{t}_i \mathbf{1}_{n'}^T$$

where \mathbf{t}_i is obtained using Eq. (7). Thus, we have

$$\operatorname{rank}(\overline{\mathbf{B}}_{i,j}) = \operatorname{rank}(\overline{\mathbf{B}(\mathbf{S})}_{i,j})$$

Moreover,

$$\mathbf{B}(\mathbf{S})_{i,j}\left(\mathbf{I}_{n'} - \frac{\mathbf{1}_{n'}\mathbf{1}_{n'}^T}{n'}\right)\mathbf{B}(\mathbf{S})_{j,i}^T = \mathbf{S}_i^T\mathbf{B}_{i,j}\left(\mathbf{I}_{n'} - \frac{\mathbf{1}_{n'}\mathbf{1}_{n'}^T}{n'}\right)\mathbf{B}_{j,i}^T\mathbf{S}_j = \mathbf{S}_1^T\overline{\mathbf{B}}_{i,j}\overline{\mathbf{B}}_{j,i}^T\mathbf{S}_2$$

and in particular rank $(\overline{\mathbf{B}(\mathbf{S})}_{i,j}\overline{\mathbf{B}(\mathbf{S})}_{j,i}^T) = \operatorname{rank}(\overline{\mathbf{B}}_{i,j}\overline{\mathbf{B}}_{j,i}^T).$

Theorem 16. Consider m = 2 and let $\mathbf{S} \in \mathbb{O}(d)^2$. Then \mathbf{S} is a non-degenerate alignment if and only if all of the following hold: (see Figures 2a and 2b for intuition when d = 2)

1. $\overline{\mathbf{B}(\mathbf{S})}_{1,2}\overline{\mathbf{B}(\mathbf{S})}_{2,1}^{T}$ is symmetric. 2. $\operatorname{Tr}(\mathbf{\Omega}^{T}\overline{\mathbf{B}(\mathbf{S})}_{1,2}\overline{\mathbf{B}(\mathbf{S})}_{2,1}^{T}\mathbf{\Omega}) \geq 0$ for all $\mathbf{\Omega} \in \operatorname{Skew}(d)$. 3. $\operatorname{rank}\left(\overline{\mathbf{B}(\mathbf{S})}_{1,2}\overline{\mathbf{B}(\mathbf{S})}_{2,1}^{T}\right) \geq d-1$ (equivalently $\operatorname{rank}(\overline{\mathbf{B}}_{1,2}\overline{\mathbf{B}}_{2,1}^{T}) \geq d-1$).

3.4 Unique Optimal Alignment in the General Setting

Since $F(\mathbf{S}) = F(\mathbf{S}\mathbf{Q})$ for all $\mathbf{Q} \in \mathbb{O}(d)$, if **S** is an optimal alignment i.e. a global minimum then so is **SQ**. In this sense, no optimal alignment is unique. With a slight abuse of convention we define a unique optimal alignment as follows.

Definition 6. An alignment $\mathbf{S} \in \mathbb{O}(d)^m$ is a unique optimal alignment if $\pi(\mathbf{S})$ is the unique global minimum of \widetilde{F} or equivalently, \mathbf{S} is an optimal alignment that is unique up to the action of $\mathbb{O}(d)$: for each optimal alignment \mathbf{S}' , there exist $\mathbf{Q} \in \mathbb{O}(d)$ such that $\mathbf{S}' = \mathbf{S}\mathbf{Q}$.

Now we provide a necessary and sufficient condition for an optimal alignment of two views to be unique. Although the proof can be found in [25], for completeness, we provide a proof using the constructs derived so far.

Theorem 17. Consider m = 2 and let **S** be an optimal alignment. Then **S** is unique if and only if rank $(\overline{\mathbf{B}}_{1,2}\overline{\mathbf{B}}_{2,1}^T) = d$ (see Figures 2b and 2c for intuition when d = 2).




Figure 2: The dotted lines represent views and the filled points represent points on the overlaps. All the pair of views are perfectly aligned. In (2a), $\operatorname{rank}(\overline{\mathbf{B}}_{1,2}\overline{\mathbf{B}}_{2,1}^T) = 0$ and clearly the two views can be rotated by a different amount while still being perfectly aligned. In (2b), $\operatorname{rank}(\overline{\mathbf{B}}_{1,2}\overline{\mathbf{B}}_{2,1}^T) = 1$ and in order for the views to be perfectly aligned, every infinitesimal rotation of the two views must be identical. However the perfect alignment of the views is not unique because the second view can be flipped (a non-infinitesimal rotation) to obtain another perfect alignment of the views. In (2c), $\operatorname{rank}(\overline{\mathbf{B}}_{1,2}\overline{\mathbf{B}}_{2,1}^T) = 2$ and the perfect alignment is unique.

4. Non-degeneracy and Uniqueness in the Noiseless Regime

As discussed in [3], in the noiseless case, the patch-stress matrix \mathbf{C} is constructed from Γ and clean measurements. In particular, there exists a global minimum \mathbf{S} of F such $F(\mathbf{S}) = 0$ or in our terminology, there exists a perfect alignment \mathbf{S} .

The organization of the section is as follows. We start by deriving some important consequences of the noiseless setting in Section 4.1 which are used in the subsequent sections. In Section 4.2, under a mild assumption on the structure of the local views, we show that the non-degeneracy and uniqueness of a perfect alignment is equivalent to the local and global rigidity [12] of the resulting realization (Theorem 21 and 23), respectively. Combined with the results in the previous section, we obtain a characterization of the local rigidity of a realization. Then, in Section 4.3 and Section 4.4, we provide necessary and sufficient conditions on the overlapping structure of the views for their realization to be locally/globally rigid. These conditions should be contrasted with those for the affine rigidity of a realization, as presented in [31].

4.1 Consequences of Noiseless Setting

The following consequences of the noiseless setting will play a crucial role in the subsequent sections.

Proposition 18. Let **S** be a perfect alignment. Then $\widehat{\mathbf{C}}(\mathbf{S}) = 0$ and consequently $\mathbf{L}(\mathbf{S}) = -\mathbf{C}(\mathbf{S})$ (see Eq. (22, 23, 24)) and $\mathbb{L}(\mathbf{S}) = -\overline{\mathbf{P}}(\mathbf{I}_d \otimes (\mathbf{PC}(\mathbf{S})\mathbf{P}^T))\overline{\mathbf{P}}^T$ (see Eq. (28, 29)).

Remark 12. Due to the above proposition and Remark 3, it is easy to deduce that for a perfect alignment \mathbf{S} , $\mathbf{L}(\mathbf{S}) \leq 0$ and $\mathbb{L}(\mathbf{S}) \leq 0$.

Using the above proposition, we provide a simplified characterization of a non-degenerate perfect alignment that will be useful in proving the subsequent results. First, along the similar lines as in [31], we define a certificate of $\mathbf{L}(\mathbf{S})$ as follows.

Definition 7. An $\Omega \in \text{Skew}(d)^m$ is said to be a certificate of $\mathbf{L}(\mathbf{S})$ if $\mathbf{L}(\mathbf{S})\Omega = 0$. It is a trivial certificate if $\Omega_i = \Omega_0$ for all $i \in [1, m]$ and for some $\Omega_0 \in \text{Skew}(d)$.

Then the characterization of a non-degenerate perfect alignment (obtained trivially from Proposition 14 and 18) is as follows.

Proposition 19. If \mathbf{S} is a perfect alignment then \mathbf{S} is non-degenerate if and only if every certificate of $\mathbf{L}(\mathbf{S})$ is trivial.

The following corollary follows trivially from Corollary 15, Proposition 18 and Remark 3.

Corollary 20. If **C** is of rank (m-1)d then every perfect alignment of F is non-degenerate.

4.2 Locally and Globally Rigid Realization

In the following, we reveal the relation between non-degenerate perfect alignment and local rigidity of the realization due to it, and unique perfect alignment and the global rigidity of the resulting realization. First note that due to Definition 2, for any two perfect alignments $\mathbf{S}, \mathbf{O} \in \mathbb{O}(d)^m$, we have

$$\min_{\mathbf{Q}\in\mathbb{O}(d),\mathbf{t}\in\mathbb{R}^{d}} \left\| \Theta(\mathbf{O}) - \mathbf{Q}^{T}\Theta(\mathbf{S}) - \mathbf{t}\mathbf{1}_{n}^{T} \right\|_{F} = \min_{\mathbf{Q}\in\mathbb{O}(d)} \left\| \Theta(\mathbf{O}) - \mathbf{Q}^{T}\Theta(\mathbf{S}) \right\|_{F} \\
= \min_{\mathbf{Q}\in\mathbb{O}(d)} \left\| \Theta(\mathbf{O}) - \Theta(\mathbf{S}\mathbf{Q}) \right\|_{F}.$$
(31)

Now we are ready to define the local and global rigidity of a realization $\Theta(\mathbf{S})$. Although phrased differently, the definitions are the same as those in [12].

Definition 8. Let **S** be a perfect alignment. Then $\Theta(\mathbf{S})$ is said to be locally rigid if there exists $\epsilon > 0$ such that for any other perfect alignment $\mathbf{O} \in \mathbb{O}(d)^m$ with $\|\Theta(\mathbf{O}) - \Theta(\mathbf{S})\|_F < \epsilon$, we have $\Theta(\mathbf{O})$ to be a rigid transformation of $\Theta(\mathbf{S})$ or equivalently (due to Eq. 31) $\Theta(\mathbf{O}) = \mathbf{Q}^T \Theta(\mathbf{S}) = \Theta(\mathbf{S}\mathbf{Q})$ for some $\mathbf{Q} \in \mathbb{O}(d)$.

Definition 9. Let **S** be a perfect alignment. Then $\Theta(\mathbf{S})$ is said to be globally rigid if for any other perfect alignment $\mathbf{O} \in \mathbb{O}(d)^m$ we have $\Theta(\mathbf{O}) = \mathbf{Q}^T \Theta(\mathbf{S}) = \Theta(\mathbf{S}\mathbf{Q})$ for some $\mathbf{Q} \in \mathbb{O}(d)$.

Examples of realizations that are not locally rigid, locally rigid but not globally rigid and globally rigid are provided in Figure 3a, Figure 1 and Figure 3b, respectively.

Assumption 3. Let S and O be perfect alignments then $\Theta(S) = \Theta(O)$ if and only if S = O.

Remark 13. Clearly, Assumption 3 holds when each view is affinely non-degenerate i.e. has at least d+1 points whose affine span has a rank of d. In this case, the perfect alignment of the local views can be uniquely determined by their realization.

In the following, we show that a non-degenerate perfect alignment gives rise to a locally rigid realization and vice versa.

Theorem 21. Let **S** be a perfect alignment and suppose Assumption 3 holds. Then **S** is non-degenerate if and only if $\Theta(\mathbf{S})$ is locally rigid.

Remark 14. Due to Corollary 20 and Theorem 21, under Assumption 3, if C is of rank (m-1)d then every realization is locally rigid.

The above remark is consistent with the results in [3; 12] in that, if each view has at least d+1 affinely non-degenerate points and the rank of **C** is (m-1)d then the underlying patch-stress framework is affinely rigid and thus locally (as well as globally) rigid too.

Finally, following Theorem 21, Remark 12, and Theorem 13, we immediately obtain a characterization of the local rigidity, as follows.

Corollary 22. Let **S** be a perfect alignment and suppose Assumption 3 holds. Then $\Theta(\mathbf{S})$ is locally rigid if and only if $\mathbb{L}(\mathbf{S})$ is of rank (m-1)d(d-1)/2.

In contrast to the general case, here, $\mathbb{L}(\mathbf{S})$ is already negative semidefinite as a consequence of the noiseless views (see Remark 12). Finally, from Definition 6 and Definition 9, it trivially follows that a unique perfect alignment gives rise to a globally rigid realization and vice versa.

Theorem 23. Let **S** be a perfect alignment and suppose Assumption 3 holds. Then **S** is unique if and only if $\Theta(\mathbf{S})$ is globally rigid.

4.3 Conditions on Overlapping Views for a Locally Rigid Realization

Under Assumption 3, the local rigidity of a realization is equivalent to the non-degeneracy of the corresponding perfect alignment (Theorem 21). We therefore focus on deriving necessary and sufficient conditions on the overlapping structure of the views for a perfect alignment to be non-degenerate. First, we need the following definition.

Definition 10. Let **S** be a perfect alignment. Let A and B be non-empty disjoint subsets of [1,m]. Define $\mathbf{B}(\mathbf{S})_{A,B}$ to be a matrix whose columns are $\mathbf{S}_i^T \mathbf{x}_{k,i} + \mathbf{t}_i$ (in the increasing order of k) where $(k,i), (k,j) \in E(\Gamma)$ for some $i \in A$ and $j \in B$, and where \mathbf{t}_i is obtained using Eq. (7). Also define

$$\overline{\mathbf{B}(\mathbf{S})}_{A,B} = \mathbf{B}(\mathbf{S})_{A,B} \left(\mathbf{I}_{n'} - \frac{\mathbf{1}_{n'} \mathbf{1}_{n'}^T}{n'} \right)$$

where $n' = |\{k : (k,i), (k,j) \in E(\Gamma) \text{ for some } (i,j) \in A \times B\}|$. For brevity, we denote $\mathbf{B}(\mathbf{S})_{\{i\},\{j\}}$ and $\overline{\mathbf{B}(\mathbf{S})}_{\{i\},\{j\}}$ by $\mathbf{B}(\mathbf{S})_{i,j}$ and $\overline{\mathbf{B}(\mathbf{S})}_{i,j}$ respectively, where $i \neq j$. Note that the notation is consistent with that of Definition 5.

Remark 15. Since **S** is a perfect alignment $\mathbf{S}_i^T \mathbf{x}_{k,i} + \mathbf{t}_i = \mathbf{S}_j^T \mathbf{x}_{k,j} + \mathbf{t}_j$ for all $(k,i), (k,j) \in E(\Gamma)$ and thus $\mathbf{B}(\mathbf{S})_{A,B}$ is well defined. Also note that $\mathbf{B}(\mathbf{S})_{A,B} = \mathbf{B}(\mathbf{S})_{B,A}$.

Remark 16. Let $i, j \in [1, m]$ and **S** be a perfect alignment. Since $\mathbf{B}(\mathbf{S})_{i,j} = \mathbf{B}(\mathbf{S})_{j,i}$ we conclude the following from Remark 11,

$$\operatorname{rank}(\overline{\mathbf{B}}_{i,j}) = \operatorname{rank}(\overline{\mathbf{B}(\mathbf{S})}_{i,j}) = \operatorname{rank}(\overline{\mathbf{B}(\mathbf{S})}_{j,i}) = \operatorname{rank}(\overline{\mathbf{B}}_{j,i})$$
$$= \operatorname{rank}(\overline{\mathbf{B}(\mathbf{S})}_{i,j}\overline{\mathbf{B}(\mathbf{S})}_{j,i}^{T}) = \operatorname{rank}(\overline{\mathbf{B}}_{i,j}\overline{\mathbf{B}}_{j,i}^{T}).$$

Due to the above two remarks and Theorem 16, a necessary and sufficient condition for a perfect alignment of two views to be non-degenerate is easily obtained and is as follows. **Theorem 24.** Consider m = 2 and let **S** be a perfect alignment. Then **S** is non-degenerate if and only if rank $(\overline{\mathbf{B}}_{1,2}) \ge d-1$.

A necessary condition for a perfect alignment of $m \ge 3$ views to be non-degenerate is as follows. The converse of the theorem does not hold, as demonstrated in Figure 3a.

Theorem 25. Let **S** be a perfect alignment. If **S** is non-degenerate then rank($\mathbf{B}(\mathbf{S})_{A,B}$) is at least d-1 for all non-empty partitions A and B of [1,m] i.e. for all $A, B \subseteq [1,m]$, $A, B \neq \emptyset, A \cap B = \emptyset$ and $A \cup B = [1,m]$.

Now we derive a sufficient condition for a perfect alignment of $m \geq 3$ views to be non-degenerate. As in [31], we construct a graph \mathbb{G} with m vertices where each vertex corresponds to a view and an edge exists between the *i*th and *j*th vertices if and only if rank $(\overline{\mathbf{B}}_{i,j}) \geq d - 1$. The Theorem 24 and the following propositions will play a crucial role in our next set of results,

Lemma 26. Let **S** be a perfect alignment and Ω be a certificate of **L**(**S**). Consider removing the ith view and the points that lie exclusively in it. Then $\mathbf{S}_{-i} = [\mathbf{S}_j]_{j \in [1,m] \setminus \{i\}}$ is a perfect alignment of the remaining set of views and $[\Omega_j]_{j \in [1,m] \setminus \{i\}}$ is a certificate of $\mathbf{L}_{-i}(\mathbf{S}_{-i})$, the matrix in Eq. (24) associated with the remaining set of views.

Proposition 27. Let **S** be a perfect alignment. Let Ω be a certificate of **L**(**S**). If ith and *j*th view lie in the same connected component of \mathbb{G} then $\Omega_i = \Omega_j$.

Similar to the one in [31], consider the following coarsening procedure on \mathbb{G} given a perfect alignment **S**: (i) transform all the views using **S** (and **t** computed using Eq. 7) (ii) merge the views that lie in the same connected component of \mathbb{G} and replace them with a single view (iii) then construct the graph (in the same manner as \mathbb{G}) associated with the new set of views (iv) repeat the procedure from (ii). Let the final graph over the remaining views be $\mathbb{G}^*(\mathbf{S})$, then the following holds.

Theorem 28. Let **S** be a perfect alignment. If $|\mathbb{G}^*(\mathbf{S})| = 1$ then **S** is non-degenerate.

The converse of the above theorem may not hold, as demonstrated in Figure 3b. The following corollary follows trivially.

Corollary 29. If \mathbb{G} is connected then every perfect alignment is non-degenerate.




Figure 3: Counterexamples for the converse of various Theorems. The dotted lines represent views and the filled points represent points on the overlaps. (3a) Clearly for every pair of nonempty partitions A and B of [1, 4], rank $(\overline{\mathbf{B}(\mathbf{S})}_{A,B}) \ge 1$ but **S** is degenerate. (3b) Clearly **S** is non-degenerate but $|\mathbb{G}^*(\mathbf{S})| = 3$.

4.4 Conditions on Overlapping Views for a Globally Rigid Realization

Under Assumption 3, the global rigidity of a realization is equivalent to the uniqueness of the corresponding perfect alignment (Theorem 23). We therefore focus on deriving necessary and sufficient conditions on the overlapping structure of the views for a perfect alignment to be unique. From Remark 16 and Theorem 17, it is easy to deduce the following.

Theorem 30. Consider m = 2 and let **S** be a perfect alignment. Then **S** is unique (see Definition 6) if and only if $\operatorname{rank}(\overline{\mathbf{B}}_{1,2}) = d$.

A necessary condition for a perfect alignment of $m \geq 3$ views to be unique is as follows.

Theorem 31. Let **S** be a perfect alignment. If **S** is unique then $\operatorname{rank}(\mathbf{B}(\mathbf{S})_{A,B}) = d$ for all non-empty partitions A and B of [1, m].

Now we derive a sufficient condition for a perfect alignment of $m \geq 3$ to be unique. As in the previous section, we construct a graph $\overline{\mathbb{G}}$ with m vertices where each vertex corresponds to a view and an edge exists between the *i*th and *j*th vertices if and only if rank $(\overline{\mathbf{B}}_{i,j}) = d$. We need Lemma 26, Theorem 30 and the following proposition for our next result,

Proposition 32. Let **S** and **S'** be perfect alignments. If ith and jth view lie in the same connected component of $\overline{\mathbb{G}}$ then there exist $\mathbf{Q} \in \mathbb{O}(d)$ such that $\mathbf{S}'_i = \mathbf{S}_i \mathbf{Q}$ and $\mathbf{S}'_j = \mathbf{S}_j \mathbf{Q}$.

Now consider the same coarsening procedure as used for Theorem 28, except that \mathbb{G} and $\mathbb{G}^*(\mathbf{S})$ are replaced by $\overline{\mathbb{G}}$ and $\overline{\mathbb{G}}^*$, respectively. Then the following holds.

Theorem 33. Let **S** be a perfect alignment. If $|\overline{\mathbb{G}}^*(\mathbf{S})| = 1$ then **S** is unique.

The converse of the above theorem may not hold, as demonstrated in Figure 4. The following corollary is obtained trivially from the above theorem.

Corollary 34. Let **S** be a perfect alignment. If $\overline{\mathbb{G}}$ is connected, then **S** is unique.



Figure 4: Counterexample for the converse of Theorem 33. The dotted lines represent views and the filled points represent points on the overlaps. Clearly **S** is unique but $|\overline{\mathbb{G}}^*(\mathbf{S})| = 3$.

5. Linear Convergence of RGD

In Section 5.1, we describe the RGD algorithm for solving the alignment problem in Eq. (8). Then in Section 5.2 we prove the local sublinear convergence of RGD to a critical point of F. In Section 5.3, using the theory of Morse-Bott functions, we extend the result to the local linear convergence of RGD to a non-degenerate alignment (see Section 3.3).

5.1 RGD Algorithm

A standard way to find a local minimum of Eq. (8) is to use RGD with a suitable initial point, step size and retraction strategy. In this work, our choice of retraction is based on QR decomposition,

$$R_{\text{QR}} : \bigcup_{\mathbf{S} \in \mathbb{O}(d)^m} (\{\mathbf{S}\} \times T_{\mathbf{S}} \mathbb{O}(d)^m) \mapsto \mathbb{O}(d)^m$$
$$R_{\text{QR}} \left(\begin{bmatrix} \mathbf{S}_1 \\ \vdots \\ \mathbf{S}_m \end{bmatrix}, \begin{bmatrix} \boldsymbol{\xi}_1 \\ \vdots \\ \boldsymbol{\xi}_m \end{bmatrix} \right) = \begin{bmatrix} \operatorname{qf}(\mathbf{S}_1 + \boldsymbol{\xi}_1) \\ \vdots \\ \operatorname{qf}(\mathbf{S}_m + \boldsymbol{\xi}_m) \end{bmatrix}.$$
(32)

where qf(**A**) denotes the **Q** factor in the thin QR decomposition of **A** [29; 1]. The step direction will always be $\boldsymbol{\xi} = -\text{grad}F(\mathbf{S})$ which is the projection of the antigradient $-\nabla F(\mathbf{S})$ onto $T_{\mathbf{S}}\mathbb{O}(d)^m$. Recall (from the proof of Proposition 7) that $\text{grad}F(\mathbf{S}) = [[\mathbf{CS}]_i - \mathbf{S}_i[\mathbf{CS}]_i^T\mathbf{S}_i]_1^m$. Then the step size α is calculated using the Armijo-type rule with parameters $\beta, \gamma \in (0, 1)$ (here g is the canonical metric on $\mathbb{O}(d)^m$ as defined in Eq. (10)),

$$\alpha = \max_{l \ge 0} \{ \beta^l \mid F(R_{\text{QR}}(\mathbf{S}, -\beta^l \text{grad}F(\mathbf{S}))) - F(\mathbf{S}) \le -\gamma \beta^l g(\nabla F(\mathbf{S}), \text{grad}F(\mathbf{S})) \}.$$
(33)

Algorithm 1 Riemannian gradient descent for solving GPOP

Require: $\mathbf{S}^0 \in \mathbb{O}(d)^m$, Γ , $\{\mathbf{x}_{k,i} : (k,i) \in E(\Gamma)\}, \beta, \gamma \in (0,1)$

- 1: Construct \mathbf{C} as in Eq. (8).
- 2: repeat
- 3: calculate the descent direction $-\operatorname{grad} F(\mathbf{S}^k)$ at \mathbf{S}^k using Eq. (17).
- 4: calculate the step size α_k according to the Armijo-type rule (see Eq. (33))
- 5: set $\mathbf{S}^{k+1} = R_{\text{QR}}(\mathbf{S}^k, -\alpha_k \text{grad}F(\mathbf{S}^k))$ using Eq. (32).
- $6: \quad k \leftarrow k+1.$
- 7: **until** convergence.

5.2 Local Sublinear Convergence of RGD

We proceed to show the local sublinear convergence of Algorithm 1 to a non-degenerate alignment (see Definition 3). Our main tool will be the convergence analysis framework presented in [24, Section 2.3] as used in [21]. To this end, we first note that F is a realanalytic function bounded from below by zero and $\mathbb{O}(d)^m$ is a compact submanifold of $\mathbb{R}^{md\times d}$. Thus, the Lojasiewicz gradient inequality [22], [24, Section 2.2] holds at every $\mathbf{S}^* \in \mathbb{O}(d)^m$ and in particular for every $\mathbf{S}^* \in \mathcal{C}$ (see Eq. (19)) i.e. there exist $\delta, \eta > 0$ and $\theta \in (0, 1/2]$ (generally dependent on \mathbf{S}^*) such that

$$|F(\mathbf{S}) - F(\mathbf{S}^*)|^{1-\theta} \le \eta \, \|\operatorname{grad} F(\mathbf{S})\|_F.$$
(34)

holds for every $\mathbf{S} \in \mathbb{O}(d)^m$ satisfying $\|\mathbf{S} - \mathbf{S}^*\|_F < \delta$.

Then using Theorem 2.3 in [24] and the fact that $\mathbb{O}(d)^m$ is compact (thus every sequence on it has a cluster point), for the Algorithm 1 to converge atleast sublinearly to a nondegenerate alignment \mathbf{S}^* , it suffices to show that the iterates $\{\mathbf{S}^k\}_{k\geq 0}$ generated by the algorithm satisfy the following: (A1). (Sufficient Descent) There exist constants $\kappa_0 > 0$ and $k_1 \in \mathbb{N}$ such that for $k \geq k_1$,

$$F(\mathbf{S}^{k+1}) - F(\mathbf{S}^{k}) \le -\kappa_0 \left\| \operatorname{grad} F(\mathbf{S}^{k}) \right\|_F \cdot \left\| \mathbf{S}^{k+1} - \mathbf{S}^{k} \right\|_F$$

(A2). (Stationarity) There exist $k_2 \in \mathbb{N}$ such that for $k \geq k_2$,

$$\left\| \operatorname{grad} F(\mathbf{S}^k) \right\|_F = 0 \implies \mathbf{S}^{k+1} = \mathbf{S}^k$$

(A3). (Safeguard) There exist a constant $\mu > 0$ and $k_3 \in \mathbb{N}$ such that for $k \ge k_3$,

$$\left\| \operatorname{grad} F(\mathbf{S}^k) \right\|_F \le \mu \left\| \mathbf{S}^{k+1} - \mathbf{S}^k \right\|_F.$$

The sequence $\{\mathbf{S}^k\}_{k\geq 0}$ satisfies (A2) trivially. We need the following propositions to prove (A1) and (A3).

Proposition 35. [21, Appendix E.2] There exist $\phi, M > 0$ such that for all $\mathbf{S}_i \in \mathbb{O}(d)$ and $\boldsymbol{\xi}_i \in T_{\mathbf{S}_i} \mathbb{O}(d)$ satisfying $\|\boldsymbol{\xi}_i\|_F \leq \phi$,

$$\| qf(\mathbf{S}_{i} + \boldsymbol{\xi}_{i}) - (\mathbf{S}_{i} + \boldsymbol{\xi}_{i}) \|_{F} \le M \| \boldsymbol{\xi}_{i} \|_{F}^{2}.$$

In particular, $M = \sqrt{10}/4$ and $\phi = 1/2$ satisfy the above inequality.

Proposition 36. There exist $\phi, M > 0$ such that for all $\mathbf{S} \in \mathbb{O}(d)^m$ and $\boldsymbol{\xi} \in T_{\mathbf{S}}\mathbb{O}(d)^m$ satisfying $\|\boldsymbol{\xi}\|_F \leq \phi$,

$$||R_{\text{QR}}(\mathbf{S}, \boldsymbol{\xi}) - (S + \boldsymbol{\xi})||_{F} \le M ||\boldsymbol{\xi}||_{F}^{2}$$

Proposition 37. The sequence $(\alpha_k)_{k\geq 0}$ and $(\mathbf{S}^k)_{k\geq 0}$ satisfies $\lim \alpha_k \|\operatorname{grad} F(\mathbf{S}^k)\|_F = 0$.

5.3 Local Linear Convegence of RGD

Now we extend the above result to the local linear convergence of Algorithm 1 to a nondegenerate alignment \mathbf{S}^* . It suffices to show that $\theta = 1/2$ in Eq. (34) [24, Theorem 2.3]. In turn, it suffices to show that F is a Morse-Bott function at \mathbf{S}^* [28, Section 6.2] [10, Definition 1.5].

Remark 17. Since $F(\mathbf{SQ}) = F(\mathbf{S})$ for all $\mathbf{Q} \in \mathbb{O}(d)$, therefore no critical point of F is non-degenerate and in particular F is not a Morse-function [5].

Proposition 38. Let \mathbf{S}^* be a non-degenerate alignment. Then \widetilde{F} is Morse-Bott at $\pi(\mathbf{S}^*)$ and consequently F is Morse-Bott at \mathbf{S}^* .

As a consequence of the above proposition, we have the following result

Theorem 39. If \mathbf{S}^* is a non-degenerate alignment, then there exist $\delta > 0$ such that RGD (Algorithm 1) converges to \mathbf{S}^* linearly when initialized with \mathbf{S}^0 such that $\|\mathbf{S}^0 - \mathbf{S}^*\|_F < \delta$.

Combining the above theorem with Theorem 13, Corollary 29 and Theorem 21, we obtain the following corollaries.

Corollary 40. If \mathbf{S}^* is an alignment such that $\mathbf{S}^* \in C$, and $\mathbb{L}(\mathbf{S}^*)$ is negative semi-definite and of rank (m-1)d(d-1)/2, then RGD converges locally linearly to \mathbf{S}^* .

Corollary 41. In the local views are noiseless and \mathbb{G} is connected then RGD converges locally linearly to a perfect alignment.

Corollary 42. If each local view is affinely non-degenerate (see Remark 13) and \mathbf{S}^* is a perfect alignment such that the realization $\Theta(\mathbf{S}^*)$ is locally rigid then RGD converges locally linearly to \mathbf{S}^* .

6. Conclusion and Open Problems

In this work we derived a characterization of a non-degenerate alignment of possibly noisy local views (Theorem 13) that can be tested in polynomial time. Thereafter, we worked in the noiseless setting, and under a mild assumption on the structure of the local views (which is satisfied when the views are affinely non-degenerate), we established equivalence of non-degeneracy and uniqueness of a perfect alignment with the local and global rigidity of the resulting perfect realization (Theorems 21 and 23).

Then, by specializing the characterization of non-degenerate alignment to the noiseless setting, we derived certain necessary and sufficient conditions on the overlapping structure of the views for a locally rigid realization (equivalently a non-degenerate perfect alignment) (Theorems 25 and 28). Similar conditions for a globally rigid realization (equivalently a unique perfect alignment) were also presented (Theorems 31 and 33).

Finally, we showed that RGD to solve the alignment problem in Eq. (8) converges linearly to a non-degenerate alignment when initialized close to it (Theorem 39). We ended with several corollaries connecting the overlapping structure of the views and the local rigidity of the realization with the convergence speed of RGD.

The following questions remain unanswered and we aim to address them in future work.

- 1. A non-degenerate alignment can be characterized using both $\mathbf{L}(\mathbf{S})$ and $\mathbb{L}(\mathbf{S})$ (see Theorem 13). The physically-interpretable consequences of the algebraic structure of $\mathbf{L}(\mathbf{S})$ (see Remark 5) are founded in the results that connect the overlapping structure of the views with the non-degeneracy of a perfect alignment (Section 4.3). Although, Remark 8 and 9 reveal the algebraic structure of $\mathbb{L}(\mathbf{S})$, the physical ramifications are still unclear.
- 2. Analogous to those presented in Section 4 but in the case when the views are noisy, necessary and sufficient conditions on the overlapping structure of m > 2 views for a non-degenerate alignment and for a unique optimal alignment are still unknown.
- 3. The proof/counterexample of the converse of Theorem 31 is to be investigated.
- 4. Requiring an alignment to be non-degenerate for the local linear convergence of RGD to it, is in a sense a strong ask. The authors of [21] presented a much stronger result where they showed the local linear convergence of RGD to any critical point (not necessarily non-degenerate) in a similar setting as ours. An extension of their result to our problem in Eq. (8) seems unclear. We aim to investigate the local linear convergence of RGD to arbitrary critical points using a geometric approach based on [28, Section 6.2].

Acknowledgments

DK was partly supported by a grant from Kavli Institute for Brain and Mind (UCSD). GM was partially funded by NSF CCF-2217058. AC was partially funded by NSF DMS-2012266 and a gift from Intel Research.

References

- P.-A. Absil, R. Mahony, and R. Sepulchre. Optimization algorithms on matrix manifolds. In *Optimization Algorithms on Matrix Manifolds*. Princeton University Press, 2009.
- [2] A. S. Bandeira, M. Charikar, A. Singer, and A. Zhu. Multireference alignment using semidefinite programming. In *Proceedings of the 5th conference on Innovations in* theoretical computer science, pages 459–470, 2014.
- [3] K. N. Chaudhury, Y. Khoo, and A. Singer. Global registration of multiple point clouds using semidefinite programming. *SIAM Journal on Optimization*, 25(1):468–501, 2015.
- [4] C. Clementi, H. Nymeyer, and J. N. Onuchic. Topological and energetic factors: what determines the structural details of the transition state ensemble and "en-route" intermediates for protein folding? an investigation for small globular proteins. *Journal of* molecular biology, 298(5):937–953, 2000.
- [5] R. L. Cohen, K. Iga, and P. Norbury. Topics in morse theory: Lecture notes, 2006. URL http://math.stanford.edu/~ralph/morsecourse/biglectures.pdf.
- [6] F. Crosilla and A. Beinat. Use of generalised Procrustes analysis for the photogrammetric block adjustment by independent models. *ISPRS Journal of Photogrammetry* and Remote Sensing, 56:195–209, 04 2002.
- [7] M. Cucuringu, Y. Lipman, and A. Singer. Sensor network localization by eigenvector synchronization over the euclidean group. ACM Transactions on Sensor Networks (TOSN), 8(3):1–42, 2012.
- [8] M. Cucuringu, A. Singer, and D. Cowburn. Eigenvector synchronization, graph rigidity and the molecule problem. *Information and Inference: A Journal of the IMA*, 1(1): 21–67, 2012.
- [9] X. Fang and K.-C. Toh. Using a distributed sdp approach to solve simulated protein molecular conformation problems. In *Distance Geometry: Theory, Methods, and Applications*, pages 351–376. Springer, 2012.
- [10] P. M. Feehan. Optimal łojasiewicz-simon inequalities and morse-bott yang-mills energy functions. Advances in Calculus of Variations, 2021.
- [11] J. E. Gentle. Matrix algebra. Springer texts in statistics, Springer, New York, NY, doi, 10:978–0, 2007.

- [12] S. J. Gortler, C. Gotsman, L. Liu, and D. P. Thurston. On affine rigidity. arXiv preprint arXiv:1011.5553, 2010.
- B. Hendrickson. Conditions for unique graph realizations. SIAM journal on computing, 21(1):65–84, 1992.
- [14] N.-D. Ho and P. Van Dooren. On the pseudo-inverse of the laplacian of a bipartite graph. Applied Mathematics Letters, 18(8):917–922, 2005.
- [15] D. Kohli, A. Cloninger, and G. Mishne. Ldle: low distortion local eigenmaps. Journal of Machine Learning Research, 22(282):1–64, 2021.
- [16] P. Kovanic. On the pseudoinverse of a sum of symmetric matrices with applications to estimation. *Kybernetika*, 15(5):341–348, 1979.
- [17] S. Krishnan, P. Y. Lee, J. B. Moore, S. Venkatasubramanian, et al. Global registration of multiple 3d point sets via optimization-on-a-manifold. In *Symposium on Geometry Processing*, pages 187–196, 2005.
- [18] A. Lee. Geometric constraint systems with applications in CAD and biology. University of Massachusetts Amherst, 2008.
- [19] S. Ling. Generalized power method for generalized orthogonal procrustes problem: global convergence and optimization landscape analysis. *arXiv preprint arXiv:2106.15493*, 2021.
- [20] S. Ling. Near-optimal bounds for generalized orthogonal procrustes problem via generalized power method. arXiv preprint arXiv:2112.13725, 2021.
- [21] H. Liu, A. M.-C. So, and W. Wu. Quadratic optimization with orthogonality constraint: explicit łojasiewicz exponent and linear convergence of retraction-based line-search and stochastic variance-reduced gradient methods. *Mathematical Programming*, 178(1): 215–262, 2019.
- [22] S. Łojasiewicz. Ensembles semi-analytiques. Institut des Hautes Etudes Scientifiques, 1965.
- [23] J. B. Saxe. Embeddability of weighted graphs in k-space is strongly np-hard. In Proc. of 17th Allerton Conference in Communications, Control and Computing, Monticello, IL, pages 480–489, 1979.
- [24] R. Schneider and A. Uschmajew. Convergence results for projected line-search methods on varieties of low-rank matrices via lojasiewicz inequality. SIAM Journal on Optimization, 25(1):622–646, 2015.
- [25] P. H. Schönemann. A generalized solution of the orthogonal procrustes problem. Psychometrika, 31(1):1–10, 1966.
- [26] G. C. Sharp, S. W. Lee, and D. K. Wehe. Multiview registration of 3d scenes by minimizing error between coordinate frames. *IEEE Transactions on Pattern Analysis* and Machine Intelligence, 26(8):1037–1050, 2004.

- [27] R. Szeliski et al. Image alignment and stitching: A tutorial. Foundations and Trends® in Computer Graphics and Vision, 2(1):1–104, 2007.
- [28] K. Usevich, J. Li, and P. Comon. Approximate matrix and tensor diagonalization by unitary transformations: convergence of jacobi-type algorithms. SIAM Journal on Optimization, 30(4):2998–3028, 2020.
- [29] C. F. Van Loan and G. Golub. Matrix computations (johns hopkins studies in mathematical sciences). *Matrix Computations*, 1996.
- [30] J. A. Williams and M. Bennamoun. Simultaneous registration of multiple point sets using orthonormal matrices. In 2000 IEEE International Conference on Acoustics, Speech, and Signal Processing. Proceedings (Cat. No. 00CH37100), volume 4, pages 2199–2202. IEEE, 2000.
- [31] H. Zha and Z. Zhang. Spectral properties of the alignment matrices in manifold learning. SIAM review, 51(3):545–566, 2009.
- [32] L. Zhang, L. Liu, C. Gotsman, and S. J. Gortler. An as-rigid-as-possible approach to sensor network localization. ACM Transactions on Sensor Networks (TOSN), 6(4): 1–21, 2010.
- [33] Z. Zhang and H. Zha. Principal manifolds and nonlinear dimensionality reduction via tangent space alignment. *SIAM journal on scientific computing*, 26(1):313–338, 2004.
- [34] Z. Zhu, A. M.-C. So, and Y. Ye. Universal rigidity: Towards accurate and efficient localization of wireless networks. In 2010 Proceedings IEEE INFOCOM, pages 1–9. IEEE, 2010.

A. Notation and Proofs

| $\left[a, b \right]$ | the set $\{a, \ldots, b\}$ where $a, b \in \mathbb{Z}$ |
|---|---|
| $(a_i)_1^k$ | the sequence a_1, \ldots, a_k where a_i is a scalar/vector/matrix |
| \mathbf{e}_{q}^{p} | a vector of zeros of length p with 1 at the q th location |
| $\mid 1_{q}^{\hat{p}}$ | a vector of zeros of length p whose first q elements are 1s |
| $ 1_{p}^{\dagger} $ | 1_p^p |
| 0_{p} | a vector of zeros of length p |
| $0_{m 	imes n}$ | a matrix of zeros with m rows and n columns |
| \mathbf{I}_d | the identity matrix of size d |
| $[\mathbf{A}_i]_1^n$ | a matrix obtained by vertically stacking the matrices $(\mathbf{A}_i)_1^m$ |
| $[\mathbf{A}]_1^n$ | $[\mathbf{A}_i]_1^n$ where $\mathbf{A}_i = \mathbf{A}$ for all $i \in [1, n]$ |
| \mathbf{I}_{d}^{m} | $[\mathbf{I}_d]_1^m$ |
| $ $ \mathbf{A}_i | i th row block of \mathbf{A}^{\sharp} |
| \mathbf{A}_{ij} | (i, j) th block of the block matrix \mathbf{A}^{\sharp} |
| $\operatorname{vec}(\mathbf{A})$ | the column-major vectorization of the matrix \mathbf{A} |
| block-diag $((\mathbf{A}_i)_1^m)$ | a block diagonal matrix with \mathbf{A}_i as the <i>i</i> th block |
| $\operatorname{diag}(\mathbf{v})$ | a diagonal matrix with $\mathbf{v}(i)$ as the <i>i</i> th diagonal element |
| $\mathbf{A}(k,:) \ (\mathbf{A}(:,k))$ | k th row (column) of \mathbf{A} |
| $ \mathbf{A}(i:j,:) (\mathbf{A}(:,i:j)) $ | a matrix obtained by stacking <i>i</i> th to <i>j</i> th rows (columns) of \mathbf{A} |
| $\mathbb{O}(n)$ | the set of orthogonal matrices of size n |
| Sym (n) , Skew (n) , | the set of symmetric and skew-symmetric matrices of size n |
| Sym (\mathbf{A}) , Skew (\mathbf{A}) | $(\mathbf{A} + \mathbf{A}^T)/2, (\mathbf{A} - \mathbf{A}^T)/2$ |
| $\mathbf{A} \succeq 0 \ (\mathbf{A} \preceq 0)$ | \mathbf{A} is symmetric and positive (negative) semidefinite |
| $\mathbf{A} \succ 0 \ (\mathbf{A} \prec 0)$ | \mathbf{A} is symmetric and positive (negative) definite |
| δ_{ij} | equals 1 if and only if i equals j otherwise it is zero |
| \mathbf{A}^{\dagger} | the pseudoinverse i.e. the Moore-Penrose inverse of \mathbf{A} . |
| $\mathbf{A}\otimes\mathbf{B}$ | the Kronecker product of \mathbf{A} and \mathbf{B} |
| WLOG | without loss of generality |

[‡]Dimensions are contextual.

Table 1: Notation.

Proof of Proposition 1. By differentiating the objective in Eq. (2) with respect to \mathbf{x}_k , the optimal value is

$$\mathbf{x}_k^* \coloneqq \mathbf{x}_k((\mathbf{S}_i)_1^m, (\mathbf{t}_i)_1^m) = \frac{\sum_{(k,i) \in E(\Gamma)} (\mathbf{S}_i^T \mathbf{x}_{k,i} + \mathbf{t}_i)}{|\{i : (k,i) \in E(\Gamma)\}|}$$

which is the consensus of all the views for the kth point. Since

$$\sum_{(k,i)\in E(\Gamma)} (\mathbf{S}_i^T \mathbf{x}_{k,i} + \mathbf{t}_i - \mathbf{x}_k^*) = 0,$$

by adding and subtracting \mathbf{x}_k^* , we obtain

$$\sum_{\substack{(k,i)\in E(\Gamma)\\(k,j)\in E(\Gamma)}} \left\| (\mathbf{S}_i^T \mathbf{x}_{k,i} + \mathbf{t}_i) - (\mathbf{S}_j^T \mathbf{x}_{k,j} + \mathbf{t}_j) \right\|_2^2 = 2 \sum_{(k,i)\in E(\Gamma)} \left\| (\mathbf{S}_i^T \mathbf{x}_{k,i} + \mathbf{t}_i) - \mathbf{x}_k^* \right\|_2^2$$

and the result follows. The above equation also shows that the minimizers $(\mathbf{S}_i)_1^m$ and $(\mathbf{t}_i)_1^m$ of Eq. (1) and Eq. (2) are the same.

Proof of Proposition 2. Let $\mathbf{u} \in \mathbb{R}^{n+m}$. Since $\mathcal{L}_{\Gamma} \succeq 0$, $\mathcal{L}_{\Gamma}\mathbf{u} = 0 \iff \mathbf{u}^{T}\mathcal{L}_{\Gamma}\mathbf{u} = 0$. From Eq. (4), the latter holds if and only if $\mathbf{e}_{ki}^{T}\mathbf{u} = 0$ for all $(k,i) \in E(\Gamma)$. Thus, using Eq. (5), $\mathcal{L}_{\Gamma}\mathbf{u} = 0$ implies $\mathbf{B}\mathbf{u} = 0$.

Proof of Proposition 4. Using the constraints characterizing $\pi^{-1}(\widetilde{\mathbf{S}})$ as in Eq. (9),

$$\mathcal{V}_{\mathbf{S}} = T_{\mathbf{S}} \pi^{-1}(\mathbf{\tilde{S}})$$

= { $\mathbf{Z} \in \mathbb{R}^{md \times d} : \mathbf{Z}_i \mathbf{S}_1^T + \mathbf{S}_i \mathbf{Z}_1^T = 0, \mathbf{Z}_j \mathbf{S}_j^T + \mathbf{S}_j \mathbf{Z}_j^T = 0, i \in [2, m], j \in [1, m]$ }
= { [$\mathbf{S}_i \mathbf{\Omega}_i$]^m : $\mathbf{\Omega}_i \in \text{Skew}(d), \mathbf{\Omega}_i + \mathbf{\Omega}_1^T = 0, i \in [1, m]$ }
= { $\mathbf{S}\mathbf{\Omega} : \mathbf{\Omega} \in \text{Skew}(d)$ }.

Then note that the objective in Eq. (12) is strictly convex in Ω . Therefore, it suffices to solve for the critical point and the Eq. (12) follows immediately.

Then, using Eq. (10),

$$\begin{aligned} \mathcal{H}_{\mathbf{S}} &= \mathcal{V}_{\mathbf{S}}^{\perp} = \{ \mathbf{W} \in T_{\mathbf{S}} \mathbb{O}(d)^{m} : \operatorname{Tr}(\mathbf{Z}^{T} \mathbf{W}) = 0 \text{ for all } \mathbf{Z} \in \mathcal{V}_{\mathbf{S}} \} \\ &= \left\{ [\mathbf{S}_{i} \mathbf{\Omega}_{i}]_{1}^{m} : \mathbf{\Omega}_{i} \in \operatorname{Skew}(d) \text{ and } \operatorname{Tr}\left(\mathbf{\Omega}^{T}\left(\sum_{1}^{m} \mathbf{\Omega}_{i}\right)\right) = 0 \text{ for all } \mathbf{\Omega} \in \operatorname{Skew}(d) \right\} \\ &= \left\{ [\mathbf{S}_{i} \mathbf{\Omega}_{i}]_{1}^{m}, \mathbf{\Omega}_{i} \in \operatorname{Skew}(d), \sum_{1}^{m} \mathbf{\Omega}_{i} \in \operatorname{Sym}(d) \right\} \\ &= \left\{ [\mathbf{S}_{i} \mathbf{\Omega}_{i}]_{1}^{m}, \mathbf{\Omega}_{i} \in \operatorname{Skew}(d), \sum_{1}^{m} \mathbf{\Omega}_{i} = 0 \right\}. \end{aligned}$$

Finally, since the horizontal space $\mathcal{H}_{\mathbf{S}}$ is the orthogonal complement to the vertical space $\mathcal{V}_{\mathbf{S}}$ in $T_{\mathbf{S}}\mathbb{O}(d)^m$, Eq. (13) follows trivially.

Proof of Proposition 5. By Eq. (14), for each $i \in [1, m - 1]$ we obtain,

$$\lim_{t \to 0} \frac{\pi (\mathbf{S} + t\mathbf{Z})_i - \pi(\mathbf{S})_i}{t} = \widetilde{\mathbf{Z}}_i \implies \mathbf{S}_{i+1}\mathbf{Z}_1^T + \mathbf{Z}_{i+1}\mathbf{S}_1^T = \widetilde{\mathbf{Z}}_i.$$

Since $T_{\widetilde{\mathbf{S}}} \mathbb{O}(d)^m / \sim$ is identified with $T_{\widetilde{\mathbf{S}}} \mathbb{O}(d)^{m-1}$, therefore there exist $(\widetilde{\mathbf{\Omega}}_i)_1^{m-1} \subseteq \operatorname{Skew}(d)$, such that $\widetilde{\mathbf{Z}}_i = \widetilde{\mathbf{S}}_i \widetilde{\mathbf{\Omega}}_i$. Also, since $\mathbf{Z} \in \mathcal{H}_{\mathbf{S}}$, there exist $(\mathbf{\Omega}_i)_1^m \subseteq \operatorname{Skew}(d)$ such that $\sum_1^m \mathbf{\Omega}_i = 0$ and $\mathbf{Z}_i = \mathbf{S}_i \mathbf{\Omega}_i$. Substituting $\widetilde{\mathbf{Z}}_i = \widetilde{\mathbf{S}}_i \widetilde{\mathbf{\Omega}}_i$ and $\mathbf{Z}_i = \mathbf{S}_i \mathbf{\Omega}_i$ in the above equation, we obtain

$$\mathbf{S}_{i+1}\mathbf{\Omega}_{1}^{T}\mathbf{S}_{1}^{T} + \mathbf{S}_{i+1}\mathbf{\Omega}_{i+1}\mathbf{S}_{1}^{T} = \widetilde{\mathbf{S}}_{i}\widetilde{\mathbf{\Omega}}_{i} \implies \mathbf{\Omega}_{i+1} - \mathbf{\Omega}_{1} = \mathbf{S}_{1}^{T}\widetilde{\mathbf{\Omega}}_{i}\mathbf{S}_{1}, i \in [1, m-1], \quad (35)$$

where we used the fact that $\widetilde{\mathbf{S}}_i = \mathbf{S}_{i+1}\mathbf{S}_1^T$. It is easy to see that the linear system of equations in $[\mathbf{\Omega}_i]_1^m$ is of full rank. Summation over *i* from 1 to m-1 and using $\sum_{1}^{m} \mathbf{\Omega}_i = 0$ gives Eq. (15) and (16).

Proof of Proposition 6. It suffices to show that $g(\mathbf{Z}, \mathbf{W})$ does not depend on the choice of $\mathbf{S} \in \pi^{-1}(\widetilde{\mathbf{S}})$. Let $\{\widetilde{\mathbf{U}}_i, \widetilde{\mathbf{V}}_i\}_1^{m-1}, \{\mathbf{U}_i, \mathbf{V}_i\}_1^m$ be elements of Skew(d) such that $\widetilde{\mathbf{Z}}_i = \widetilde{\mathbf{S}}_i \widetilde{\mathbf{U}}_i$, $\widetilde{\mathbf{W}}_i = \widetilde{\mathbf{S}}_i \widetilde{\mathbf{V}}_i, \mathbf{Z}_i = \mathbf{S}_i \mathbf{U}_i$ and $\mathbf{W}_i = \mathbf{S}_i \mathbf{V}_i$. By the definition of $\mathcal{H}_{\mathbf{S}}, \sum_1^m \mathbf{U}_i = \sum_1^m \mathbf{V}_i = 0$ and the relation between \mathbf{U}_i and $\widetilde{\mathbf{U}}_i$, and \mathbf{V}_i and $\widetilde{\mathbf{V}}_i$ is given by Eq. (15, 16) (through Eq. (35)). Then we have

$$g(\mathbf{Z}, \mathbf{W})$$

$$= \sum_{1}^{m} \operatorname{Tr}(\mathbf{Z}_{i}^{T} \mathbf{W}_{i}) = \sum_{1}^{m} \operatorname{Tr}(\mathbf{U}_{i}^{T} \mathbf{V}_{i}) = \sum_{1}^{m} \operatorname{Tr}((\mathbf{U}_{i} - \mathbf{U}_{1})^{T} \mathbf{V}_{i}) + \operatorname{Tr}(\mathbf{U}_{1}^{T} \mathbf{V}_{i})$$

$$= \sum_{1}^{m} \operatorname{Tr}((\mathbf{U}_{i} - \mathbf{U}_{1})^{T} \mathbf{V}_{i}) = \sum_{1}^{m} \operatorname{Tr}((\mathbf{U}_{i} - \mathbf{U}_{1})^{T} (\mathbf{V}_{i} - \mathbf{V}_{1})) + \operatorname{Tr}((\mathbf{U}_{i} - \mathbf{U}_{1})^{T} \mathbf{V}_{1})$$

$$= \sum_{1}^{m-1} \operatorname{Tr}((\mathbf{U}_{i+1} - \mathbf{U}_{1})^{T} (\mathbf{V}_{i+1} - \mathbf{V}_{1})) - \operatorname{Tr}(\mathbf{U}_{1}^{T} \mathbf{V}_{1}) = \sum_{1}^{m-1} \operatorname{Tr}(\widetilde{\mathbf{U}}_{i}^{T} \widetilde{\mathbf{V}}_{i}) - \operatorname{Tr}(\mathbf{U}_{1}^{T} \mathbf{V}_{1})$$

$$= \sum_{1}^{m-1} \operatorname{Tr}(\widetilde{\mathbf{U}}_{i}^{T} \widetilde{\mathbf{V}}_{i}) - \frac{1}{m^{2}} \sum_{i,j=1}^{m-1} \operatorname{Tr}(\widetilde{\mathbf{U}}_{i}^{T} \widetilde{\mathbf{V}}_{j}).$$

We observe that the last equation is independent of the choice of \mathbf{S} , therefore, the result follows.

Proof of Proposition 7. Using Proposition 3 and [1, Section 3.6.2],

$$\operatorname{grad} F(\mathbf{S}) = \operatorname{grad} F(\mathbf{S}) = \mathbf{P}_{\mathbf{S}}(\nabla F(\mathbf{S})) = \mathbf{P}_{\mathbf{S}}(2\mathbf{CS})$$
$$= [2\mathbf{S}_{i}\operatorname{skew}(\mathbf{S}_{i}^{T}[\mathbf{CS}]_{i}])]_{1}^{m}$$
$$= [\mathbf{CS}]_{i} - \mathbf{S}_{i}[\mathbf{CS}]_{i}^{T}\mathbf{S}_{i}$$
$$= \mathbf{S}_{i}(\mathbf{S}_{i}^{T}[\mathbf{CS}]_{i} - [\mathbf{CS}]_{i}^{T}\mathbf{S}_{i}).$$

Using the fact that C is symmetric, the result follows.

Proof of Proposition 8. For each $i \in [1, m]$,

$$D \operatorname{grad} F(\mathbf{S})[\mathbf{Z}]_{i}$$

$$= \lim_{t \to 0} \frac{\operatorname{grad} F(\mathbf{S} + t\mathbf{Z})_{i} - \operatorname{grad} F(\mathbf{S})_{i}}{t}$$

$$= \lim_{t \to 0} \frac{([\mathbf{C}(\mathbf{S} + t\mathbf{Z})]_{i} - (\mathbf{S}_{i} + t\mathbf{Z}_{i})[\mathbf{C}(\mathbf{S} + t\mathbf{Z})]_{i}^{T}(\mathbf{S}_{i} + t\mathbf{Z}_{i})) - ([\mathbf{C}\mathbf{S}]_{i} - \mathbf{S}_{i}[\mathbf{C}\mathbf{S}]_{i}^{T}\mathbf{S}_{i})}{t}$$

$$= [\mathbf{C}\mathbf{Z}]_{i} - \mathbf{S}_{i}[\mathbf{C}\mathbf{Z}]_{i}^{T}\mathbf{S}_{i} - \mathbf{S}_{i}[\mathbf{C}\mathbf{S}]_{i}^{T}\mathbf{Z}_{i} - \mathbf{Z}_{i}[\mathbf{C}\mathbf{S}]_{i}^{T}\mathbf{S}_{i}$$

$$= \mathbf{S}_{i}(\mathbf{S}_{i}^{T}[\mathbf{C}\mathbf{Z}]_{i} - [\mathbf{C}\mathbf{Z}]_{i}^{T}\mathbf{S}_{i} - [\mathbf{C}\mathbf{S}]_{i}^{T}\mathbf{Z}_{i} - \mathbf{S}_{i}^{T}\mathbf{Z}_{i}[\mathbf{C}\mathbf{S}]_{i}^{T}\mathbf{S}_{i}).$$

Note that we have not yet used the fact that $\widetilde{\mathbf{S}} \in \widetilde{\mathcal{C}}$ or equivalently, $[\mathbf{CS}]_i = \mathbf{S}_i [\mathbf{CS}]_i^T \mathbf{S}_i$ (see the proof of Proposition 7) and $\mathbf{Z} \in T_{\mathbf{S}} \mathbb{O}(d)^m$ or equivalently, $\mathbf{S}_i^T \mathbf{Z}_i + \mathbf{Z}_i^T \mathbf{S}_i = 0$ (see Eq. (11) in Proposition 3). Combining the two, we get $\mathbf{S}_i^T \mathbf{Z}_i [\mathbf{CS}]_i^T \mathbf{S}_i = -\mathbf{Z}_i^T [\mathbf{CS}]_i$ and the result follows.

Proof of Proposition 9. Using Proposition 3, 4 and 8, and [1, Chapter 5],

$$\begin{aligned} \operatorname{Hess} \widetilde{F}(\widetilde{\mathbf{S}})[\widetilde{\mathbf{Z}}] &= \widetilde{\nabla}_{\widetilde{\mathbf{Z}}} \operatorname{grad} \widetilde{F}(\widetilde{\mathbf{S}}) = P_{\mathbf{S}}^{h}(\nabla_{\overline{\widetilde{\mathbf{Z}}}} \operatorname{grad} \widetilde{F}(\widetilde{\mathbf{S}})) = P_{\mathbf{S}}^{h}(\nabla_{\mathbf{Z}} \operatorname{grad} F(\mathbf{S})) \\ &= P_{\mathbf{S}}^{h}(P_{\mathbf{S}}(D \operatorname{grad} F(\mathbf{S})[\mathbf{Z}])) \\ &= P_{\mathbf{S}}^{h}\left(\left[\mathbf{S}_{i}(\mathbf{S}_{i}^{T}[\mathbf{C}\mathbf{Z}]_{i} - [\mathbf{C}\mathbf{Z}]_{i}^{T}\mathbf{S}_{i} - [\mathbf{C}\mathbf{S}]_{i}^{T}\mathbf{Z}_{i} + \mathbf{Z}_{i}^{T}[\mathbf{C}\mathbf{S}]_{i})\right]_{1}^{m}\right) \\ &= P_{\mathbf{S}}^{h}([\mathbf{S}_{i}\widehat{\Omega}_{i}]_{1}^{m}) = [\mathbf{S}_{i}(\widehat{\Omega}_{i} - \overline{\widehat{\Omega}})]_{1}^{m}\end{aligned}$$

where

$$\overline{\widehat{\mathbf{\Omega}}} = \frac{1}{m} \sum_{1}^{m} \widehat{\mathbf{\Omega}}_{i} = \frac{1}{m} \sum_{1}^{m} (\mathbf{S}_{i}^{T} [\mathbf{C} \mathbf{Z}]_{i} - [\mathbf{C} \mathbf{Z}]_{i}^{T} \mathbf{S}_{i} - [\mathbf{C} \mathbf{S}]_{i}^{T} \mathbf{Z}_{i} + \mathbf{Z}_{i}^{T} [\mathbf{C} \mathbf{S}]_{i})$$
$$= \frac{1}{m} (\mathbf{S}^{T} \mathbf{C} \mathbf{Z} - \mathbf{Z}^{T} \mathbf{C} \mathbf{S} - \mathbf{S}^{T} \mathbf{C} \mathbf{Z} + \mathbf{Z}^{T} \mathbf{C} \mathbf{S}) = 0.$$

This validates that $\overline{\operatorname{Hess}\widetilde{F}(\widetilde{\mathbf{S}})[\widetilde{\mathbf{Z}}]} = [\mathbf{S}_i\widehat{\mathbf{\Omega}}_i]_1^m$ indeed lies in $\mathcal{H}_{\mathbf{S}}$ (see Proposition 4).

Proof of Proposition 10. Since $\mathbf{Z} \in \mathcal{H}_{\mathbf{S}}$, using Proposition 4, there exist $\mathbf{\Omega} = [\mathbf{\Omega}_i]_1^m$ such that $\mathbf{\Omega}_i \in \text{Skew}(d), \sum_{1}^{m} \mathbf{\Omega}_i = 0$ and $\mathbf{Z}_i = \mathbf{S}_i \mathbf{\Omega}_i$. Then

$$\begin{aligned} \widehat{\boldsymbol{\Omega}}_{i} &= (\mathbf{S}_{i}^{T}[\mathbf{C}\mathbf{Z}]_{i} - [\mathbf{C}\mathbf{Z}]_{i}^{T}\mathbf{S}_{i}) - ([\mathbf{C}\mathbf{S}]_{i}^{T}\mathbf{Z}_{i} - \mathbf{Z}_{i}^{T}[\mathbf{C}\mathbf{S}]_{i}) \\ &= ([\mathbf{C}(\mathbf{S})\boldsymbol{\Omega}]_{i} - [\mathbf{C}(\mathbf{S})\boldsymbol{\Omega}]_{i}^{T}) - ([\widehat{\mathbf{C}}(\mathbf{S})\boldsymbol{\Omega}]_{i} - [\widehat{\mathbf{C}}(\mathbf{S})\boldsymbol{\Omega}]_{i}^{T}) \\ &= [\mathbf{L}(\mathbf{S})\boldsymbol{\Omega}]_{i}^{T} - [\mathbf{L}(\mathbf{S})\boldsymbol{\Omega}]_{i}. \end{aligned}$$

Proof of Proposition 11. Using Proposition 6, 9 and 10,

$$\begin{split} \widetilde{g}(\operatorname{Hess}\widetilde{F}(\widetilde{\mathbf{S}})[\widetilde{\mathbf{Z}}],\widetilde{\mathbf{Z}}) &= g(\overline{\operatorname{Hess}\widetilde{F}(\widetilde{\mathbf{S}})[\widetilde{\mathbf{Z}}]},\overline{\widetilde{\mathbf{Z}}}) = g(\overline{\operatorname{Hess}\widetilde{F}(\widetilde{\mathbf{S}})[\widetilde{\mathbf{Z}}]},\mathbf{Z}) \\ &= \sum_{1}^{m} \operatorname{Tr}((\mathbf{S}_{i}\widehat{\mathbf{\Omega}}_{i})^{T}(\mathbf{S}_{i}\mathbf{\Omega}_{i})) = \sum_{1}^{m} \operatorname{Tr}(\widehat{\mathbf{\Omega}}_{i}^{T}\mathbf{\Omega}_{i}) \\ &= \sum_{1}^{m} \operatorname{Tr}([\mathbf{L}(\mathbf{S})\mathbf{\Omega}]_{i}\mathbf{\Omega}_{i} - [\mathbf{L}(\mathbf{S})\mathbf{\Omega}]_{i}^{T}\mathbf{\Omega}_{i}) \\ &= -2\operatorname{Tr}(\mathbf{\Omega}^{T}\mathbf{L}(\mathbf{S})\mathbf{\Omega}) \end{split}$$

Proof of Proposition 12. Using the fact that $vec(\mathbf{AXB}) = (\mathbf{B}^T \otimes \mathbf{A})vec(\mathbf{X})$,

$$Tr(\mathbf{\Omega}^{T}\mathbf{L}(\mathbf{S})\mathbf{\Omega}) = Tr((\mathbf{P}\mathbf{\Omega})^{T}\mathbf{P}\mathbf{L}(\mathbf{S})\mathbf{P}^{T}(\mathbf{P}\mathbf{\Omega}))$$
$$= vec(\mathbf{P}\mathbf{\Omega})^{T}vec(\mathcal{L}(\mathbf{S})(\mathbf{P}\mathbf{\Omega}))$$
$$= vec(\mathbf{P}\mathbf{\Omega})^{T}(\mathbf{I}_{d} \otimes \mathcal{L}(\mathbf{S}))vec(\mathbf{P}\mathbf{\Omega})$$

$$= \boldsymbol{\omega}^T \overline{\mathbf{P}} (\mathbf{I}_d \otimes \mathcal{L}(\mathbf{S})) \overline{\mathbf{P}}^T \boldsymbol{\omega}$$

 $= \boldsymbol{\omega}^T \mathbb{L}(\mathbf{S}) \boldsymbol{\omega}.$

Proof of Theorem 13. 1, 2 and 3 are equivalent by definitions and Proposition 11. For (3 \iff 4), by comparing dimensions, we note that the set of $\Omega = [\Omega_i]_1^m$ where $\Omega_i \in \text{Skew}(d)$ and $\sum_{1}^{m} \Omega_i = 0$, is the same as the set of $\Omega = [\Omega_i - \Omega_0]_1^m$ where $\Omega_i \in \text{Skew}(d)$ and $\Omega_0 = \frac{1}{m} \sum_{1}^{m} \Omega_i$. Using Remark 5, we know that $\mathbf{L}(\mathbf{S}) = \mathbf{L}(\mathbf{S})^T$ and $\mathbf{L}(\mathbf{S})[\Omega_0]_1^m = 0$. Thus $(\Omega - [\Omega_0]_1^m)^T \mathbf{L}(\mathbf{S})(\Omega - [\Omega_0]_1^m) = \Omega^T \mathbf{L}(\mathbf{S})\Omega$ and the result follows.

 $(4 \iff 5)$: follows directly from the definition of ω and Proposition 12.

 $(5 \iff 6)$: follows from Remark 8.

Proof of Proposition 14. It suffices to pick conditions 4 and 6 (one involving L(S)) and one involving L(S)). Suppose L(S) satisfies condition 4 and let Ω be as described in the condition. Then using Remark 5,

$$Tr(\mathbf{\Omega}^{T}\mathbf{L}(\mathbf{S}\mathbf{Q})\mathbf{\Omega}) = Tr(\mathbf{\Omega}^{T}(\mathbf{I}_{m} \otimes \mathbf{Q})^{T}\mathbf{L}(\mathbf{S})(\mathbf{I}_{m} \otimes \mathbf{Q})\mathbf{\Omega})$$
$$= Tr(\mathbf{Q}\mathbf{\Omega}^{T}(\mathbf{I}_{m} \otimes \mathbf{Q})^{T}\mathbf{L}(\mathbf{S})(\mathbf{I}_{m} \otimes \mathbf{Q})\mathbf{\Omega}\mathbf{Q}^{T})$$
$$= Tr(\overline{\mathbf{\Omega}}^{T}\mathbf{L}(\mathbf{S})\overline{\mathbf{\Omega}}) < 0$$

because $\overline{\Omega}_i = \mathbf{Q} \Omega_i \mathbf{Q}^T \in \text{Skew}(d)$ for all $i \in [1, m]$ and not all $\overline{\Omega}_i$ are equal (if $\overline{\Omega}_i = \overline{\Omega}_j$ then $\Omega_i = \Omega_j$, a contradiction). Thus the condition 4 holds for $\mathbf{S}\mathbf{Q}$ too.

Now, suppose $\mathbb{L}(\mathbf{S})$ satisfy condition 6. Then using Remark 8, we conclude that \mathbf{SQ} also satisfies condition 6.

Proof of Corollary 15. Since the rank of $\mathbf{L}(\mathbf{S})$ is (m-1)d, using Remark 5, $\mathbf{L}(\mathbf{S})[\Omega_i]_1^m = 0$ (equivalently, $\operatorname{Tr}(([\Omega_i]_1^m)^T \mathbf{L}(\mathbf{S})[\Omega_i]_1^m) = 0)$ if and only if $\Omega_i = \Omega_0$ for all $i \in [1, m]$ and for some $\Omega_0 \in \operatorname{Skew}(d)$. Thus the condition 4 in Theorem 13 is satisfied.

Proof of Theorem 16. The proof is divided into three parts specialized to the three conditions in the statement.

Part 1. First note that $\mathbf{S} \in \mathcal{C}$ if and only if $\mathbf{S}_i^T[\mathbf{CS}]_i = [\mathbf{CS}]_i^T \mathbf{S}_i$ for i = 1, 2 (see Eq. (19)). Since $\mathbf{S}_1^T[\mathbf{CS}]_1 - [\mathbf{CS}]_1^T \mathbf{S}_1 = [\mathbf{CS}]_2^T \mathbf{S}_2 - \mathbf{S}_2^T[\mathbf{CS}]_2$ and

$$\begin{split} \mathbf{S}_1^T [\mathbf{C}\mathbf{S}]_1 &- [\mathbf{C}\mathbf{S}]_1^T \mathbf{S}_1 = \mathbf{S}_1^T \mathbf{B}_1 \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}_2^T \mathbf{S}_2 - \mathbf{S}_2^T \mathbf{B}_2 \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}_1^T \mathbf{S}_1 \\ &= \mathbf{B}(\mathbf{S})_1 \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_2^T - \mathbf{B}(\mathbf{S})_2 \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_1^T, \end{split}$$

thus $\mathbf{S} \in \mathcal{C}$ if and only if $\mathbf{B}(\mathbf{S})_1 \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_2^T$ is symmetric. At this point, we note that the forms of $\mathbf{B}(\mathbf{S})_1$ and $\mathbf{B}(\mathbf{S})_2$ are

where (see Remark 1) $\mathbf{X}_1 \in \mathbb{R}^{d \times n_1}$ and $\mathbf{X}_3 \in \mathbb{R}^{d \times n_3}$ correspond to the local coordinates, due to the first view, of the n_1 points that lie exclusively in the first view and the n_3 points that lie on the overlap of both views, respectively. Similarly, $\mathbf{Y}_2 \in \mathbb{R}^{d \times n_2}$ and $\mathbf{Y}_3 \in \mathbb{R}^{d \times n_3}$

correspond to the local coordinates, due to the second view, of the n_2 points that lie exclusively in the second view and the n_3 points which lie on the overlap of both views, respectively. In particular, $\mathbf{X}_3 = \mathbf{B}(\mathbf{S})_{1,2}$ and $\mathbf{Y}_3 = \mathbf{B}(\mathbf{S})_{2,1}$ (perhaps after permuting the points). Moreover,

$$\mathcal{L}_{\Gamma} = egin{bmatrix} \mathbf{I}_{n_1} & -\mathbf{1}_{n_1} & \mathbf{0}_{n_1} \ & \mathbf{I}_{n_2} & \mathbf{0}_{n_2} & -\mathbf{1}_{n_2} \ & 2\mathbf{I}_{n_3} & -\mathbf{1}_{n_3} & -\mathbf{1}_{n_3} \ -\mathbf{1}_{n_1}^T & \mathbf{0}_{n_2}^T & -\mathbf{1}_{n_3}^T & n_1+n_3 \ & \mathbf{0}_{n_1}^T & -\mathbf{1}_{n_2}^T & -\mathbf{1}_{n_3}^T & n_2+n_3 \end{bmatrix}.$$

Through simple calculations, we obtain

$$\boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} = \frac{1}{2n_3} \begin{bmatrix} 2n_3\mathbf{I}_{n_1} + \mathbf{1}_{n_1}\mathbf{1}_{n_1}^T & -\mathbf{1}_{n_1}\mathbf{1}_{n_2}^T & \mathbf{1}_{n_1} & -\mathbf{1}_{n_1} \\ -\mathbf{1}_{n_2}\mathbf{1}_{n_1}^T & 2n_3\mathbf{I}_{n_2} + \mathbf{1}_{n_2}\mathbf{1}_{n_2}^T & -\mathbf{1}_{n_2} & \mathbf{1}_{n_2} \\ & & & n_3\mathbf{I}_{n_3} & & \\ \mathbf{1}_{n_1}^T & -\mathbf{1}_{n_2}^T & & \mathbf{1} & -\mathbf{1} \\ -\mathbf{1}_{n_1}^T & \mathbf{1}_{n_2}^T & & -\mathbf{1} & \mathbf{1} \end{bmatrix}$$

Thus,

$$\mathbf{B}(\mathbf{S})_{1} \mathcal{L}_{\Gamma}^{\dagger} = \begin{bmatrix} \mathbf{X}_{1} - \frac{\mathbf{X}_{3} \mathbf{1}_{n_{3}} \mathbf{1}_{n_{1}}^{T}}{2n_{3}}, & \frac{\mathbf{X}_{3} \mathbf{1}_{n_{3}} \mathbf{1}_{n_{2}}^{T}}{2n_{3}}, & \frac{1}{2} \mathbf{X}_{3}, & -\frac{\mathbf{X}_{3} \mathbf{1}_{n_{3}}}{2n_{3}}, & \frac{\mathbf{X}_{3} \mathbf{1}_{n_{3}}}{2n_{3}} \end{bmatrix}$$

and (since $\mathbf{S} \in \mathcal{C}$ and part (i) of the proof),

$$\begin{split} \mathbf{B}(\mathbf{S})_{2} \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_{1}^{T} &= \mathbf{B}(\mathbf{S})_{1} \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_{2}^{T} \\ &= \frac{1}{2} \mathbf{X}_{3} \left(\mathbf{I}_{n_{3}} - \frac{1}{n_{3}} \mathbf{1}_{n_{3}} \mathbf{1}_{n_{3}}^{T} \right) \mathbf{Y}_{3}^{T} \\ &= \frac{1}{2} \mathbf{B}(\mathbf{S})_{1,2} \left(\mathbf{I}_{n_{3}} - \frac{1}{n_{3}} \mathbf{1}_{n_{3}} \mathbf{1}_{n_{3}}^{T} \right) \left(\mathbf{I}_{n_{3}} - \frac{1}{n_{3}} \mathbf{1}_{n_{3}} \mathbf{1}_{n_{3}}^{T} \right)^{T} \mathbf{B}(\mathbf{S})_{2,1}^{T} \\ &= \frac{1}{2} \overline{\mathbf{B}(\mathbf{S})}_{1,2} \overline{\mathbf{B}(\mathbf{S})}_{2,1}^{T} \end{split}$$

Combining the above equations, we conclude that $\mathbf{S} \in \mathcal{C}$ if and only if $\overline{\mathbf{B}(\mathbf{S})}_{1,2}\overline{\mathbf{B}(\mathbf{S})}_{2,1}^T$ is symmetric.

Part 2. For $\mathbf{S} \in \mathcal{C}$, from the Remark 5 and Part 1, we have,

$$\begin{split} \mathbf{L}(\mathbf{S}) &= \begin{bmatrix} -\mathbf{B}(\mathbf{S})_1 \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_2^T & \mathbf{B}(\mathbf{S})_1 \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_2^T \\ \mathbf{B}(\mathbf{S})_2 \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_1^T & -\mathbf{B}(\mathbf{S})_2 \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_1^T \end{bmatrix} \\ &= \begin{bmatrix} -\mathbf{B}(\mathbf{S})_1 \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_2^T & \mathbf{B}(\mathbf{S})_1 \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_2^T \\ \mathbf{B}(\mathbf{S})_1 \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_2^T & -\mathbf{B}(\mathbf{S})_1 \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_2^T \end{bmatrix} \end{split}$$

Let $\Omega_1, \Omega_2 \in \text{Skew}(d)$ such that $\Omega_1 + \Omega_2 = 0$. Then, using the above equations,

$$\operatorname{Tr}\left(\begin{bmatrix}\boldsymbol{\Omega}_1^T & \boldsymbol{\Omega}_2^T\end{bmatrix}\mathbf{L}(\mathbf{S})\begin{bmatrix}\boldsymbol{\Omega}_1\\\boldsymbol{\Omega}_2\end{bmatrix}\right) = \operatorname{Tr}\left(\begin{bmatrix}-\boldsymbol{\Omega}_1 & \boldsymbol{\Omega}_1\end{bmatrix}\mathbf{L}(\mathbf{S})\begin{bmatrix}\boldsymbol{\Omega}_1\\-\boldsymbol{\Omega}_1\end{bmatrix}\right)$$

$$= -2\mathrm{Tr}(\mathbf{\Omega}_1^T \overline{\mathbf{B}(\mathbf{S})}_{1,2} \overline{\mathbf{B}(\mathbf{S})}_{2,1}^T \mathbf{\Omega}_1).$$

Combining the above equation and Part 1 with Proposition 11, we conclude that $\pi(\mathbf{S})$ is a local minimum of \widetilde{F} if and only if the first two conditions of the statement are met.

Part 3. Here we deal with the non-degeneracy of $\widetilde{\mathbf{S}} = \pi(\mathbf{S})$. For d = 1, $\widetilde{\mathbf{S}}$ is trivially non-degenerate. So we assume that $d \geq 2$. From Part 2, we note that for $\Omega_1, \Omega_2 \in \text{Skew}(d)$ such that $\Omega_1 + \Omega_2 = 0$, $\mathbf{L}(\mathbf{S})[\Omega_i]_1^2 = 0$ if and only if $\overline{\mathbf{B}(\mathbf{S})}_{1,2}\overline{\mathbf{B}(\mathbf{S})}_{2,1}^T\Omega = 0$. Thus $\widetilde{\mathbf{S}}$ is non-degenerate if and only if $\overline{\mathbf{B}(\mathbf{S})}_{1,2}\overline{\mathbf{B}(\mathbf{S})}_{2,1}^T\Omega = 0 \iff \Omega = 0$. It suffices to show that $\overline{\mathbf{B}(\mathbf{S})}_{1,2}\overline{\mathbf{B}(\mathbf{S})}_{2,1}^T\Omega = 0$ if and only if $\operatorname{rank}(\overline{\mathbf{B}(\mathbf{S})}_{1,2}\overline{\mathbf{B}(\mathbf{S})}_{2,1}^T) \geq d-1$.

 (\Leftarrow) Suppose rank $(\overline{\mathbf{B}(\mathbf{S})}_{1,2}\overline{\mathbf{B}(\mathbf{S})}_{2,1}^T) \ge d-1$ then the null space of $\overline{\mathbf{B}(\mathbf{S})}_{1,2}\overline{\mathbf{B}(\mathbf{S})}_{2,1}^T$ is at most one-dimensional. Moreover, rank of a non-zero skew symmetric matrix of size $d \ge 2$, is at least two. Thus $\overline{\mathbf{B}(\mathbf{S})}_{1,2}\overline{\mathbf{B}(\mathbf{S})}_{2,1}^T \mathbf{\Omega} = 0 \iff \mathbf{\Omega} = 0$. We conclude that $\mathbf{L}(\mathbf{S})$ has trivial certificates only, and thus $\widetilde{\mathbf{S}}$ is non-degenerate.

 (\Longrightarrow) Suppose rank $(\overline{\mathbf{B}(\mathbf{S})}_{1,2}\overline{\mathbf{B}(\mathbf{S})}_{2,1}^T) \leq d-2$, then there exist non-zero vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^d$ in the kernel of $\overline{\mathbf{B}(\mathbf{S})}_{1,2}\overline{\mathbf{B}(\mathbf{S})}_{2,1}^T$ such that $\mathbf{u}^T\mathbf{v} = 0$. Let $\mathbf{\Omega} = \mathbf{u}\mathbf{v}^T - \mathbf{v}\mathbf{u}^T$ then clearly $\mathbf{\Omega} \in \text{Skew}(d), \ \mathbf{\Omega} \neq 0$ and $\overline{\mathbf{B}(\mathbf{S})}_{1,2}\overline{\mathbf{B}(\mathbf{S})}_{2,1}^T\mathbf{\Omega} = 0$.

Proof of Theorem 17. Let **S** be an optimal alignment i.e. a global minimum of F, then from the proof of Theorem 16 we have

$$\mathbf{C}(\mathbf{S})_{1,2} = -\mathbf{B}(\mathbf{S})_1 \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_2^T = -\frac{1}{2} \mathbf{S}_1^T \overline{\mathbf{B}}_{1,2} \overline{\mathbf{B}}_{2,1}^T \mathbf{S}_2.$$

Then note that the minimizer of $\operatorname{Tr}(\mathbf{CSS}^T)$ is same as the minimizer of $\operatorname{Tr}(\mathbf{C}(\mathbf{S})_{1,2})$ because $\mathbf{S}_1\mathbf{S}_1^T = \mathbf{S}_2\mathbf{S}_2^T = \mathbf{I}_d$ and $\mathbf{C}_{2,1} = \mathbf{C}_{1,2}^T$ is symmetric. WLOG, take $\mathbf{S}_1 = \mathbf{I}_d$. Then it suffices to show that the following problem has unique solution

$$\min_{\mathbf{S}_2 \in \mathbb{O}(d)} \operatorname{Tr}(\mathbf{C}(\mathbf{S})_{12}) = -\frac{1}{2} \max_{\mathbf{S}_2 \in \mathbb{O}(d)} \operatorname{Tr}(\overline{\mathbf{B}}_{1,2}\overline{\mathbf{B}}_{2,1}^T \mathbf{S}_2)$$

if and only if rank $(\overline{\mathbf{B}}_{1,2}\overline{\mathbf{B}}_{2,1}^T) = d$.

Let $\overline{\mathbf{B}}_{1,2}\overline{\mathbf{B}}_{2,1}^T = \mathbf{U}\Sigma\mathbf{V}^T$ be its singular value decomposition. Note that $\Sigma \succeq 0$. If $\operatorname{rank}(\overline{\mathbf{B}}_{1,2}\overline{\mathbf{B}}_{2,1}^T) = d$ then $\Sigma \succ 0$ and clearly, the unique optimizer is $\mathbf{S}_2 = \mathbf{V}\mathbf{U}^T$. If $\operatorname{rank}(\overline{\mathbf{B}}_{1,2}\overline{\mathbf{B}}_{2,1}^T) \leq d-1$ then there exist orthogonal matrix $\mathbf{U}' \neq \mathbf{I}_d$ such that $\Sigma = \mathbf{U}'\Sigma$ and thus $\mathbf{V}\mathbf{U}^T \neq \mathbf{V}(\mathbf{U}'\mathbf{U})^T$ are both optimizers of the above objective.

Proof of Proposition 18. Since **S** is a perfect alignment $F(\mathbf{S}) = \text{Tr}(\mathbf{CSS}^T) = 0$. Since $\mathbf{C} \succeq 0$, the columns of **S** lie in the kernel of **C**. In particular $\mathbf{CS} = \mathbf{0}$. It follows that $\widehat{\mathbf{C}}(\mathbf{S}) = \mathbf{0}$ (see Eq. (23)). We conclude that $\mathbf{L}(\mathbf{S}) = -\mathbf{C}(\mathbf{S})$.

Proof of Theorem 21. First note that under Assumption 3, the equation $\Theta(\mathbf{O}) = \Theta(\mathbf{SQ})$ in Definition 8 is equivalent to $\pi(\mathbf{S}) = \pi(\mathbf{O})$.

(\Leftarrow) Suppose **S** is a degenerate perfect alignment. Thus $\pi(\mathbf{S})$ is degenerate (see Definition 3). Let $\epsilon > 0$ be arbitrary and define

$$\eta \coloneqq \left\| \mathbf{B} \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger}(:, 1:n) \left(\mathbf{I}_{n} - n^{-1} \mathbf{1}_{n} \mathbf{1}_{n}^{T} \right) \right\|_{F}$$

Then there exists another perfect alignment $\mathbf{O} \in \mathbb{O}(d)^m$ such that $\|\mathbf{S} - \mathbf{O}\|_F < \epsilon/\eta$ and $\pi(\mathbf{S}) \neq \pi(\mathbf{O})$. Due to Assumption 3, we have $\Theta(\mathbf{O}) \neq \Theta(\mathbf{SQ})$ for any $\mathbf{Q} \in \mathbb{O}(d)$, however, (from Definition 2),

$$\|\Theta(\mathbf{O}) - \Theta(\mathbf{S})\|_{F} \leq \|\mathbf{S} - \mathbf{O}\|_{F} \left\|\mathbf{B}\mathcal{L}_{\Gamma}^{\dagger}(:, 1:n) \left(\mathbf{I}_{n} - n^{-1}\mathbf{1}_{n}\mathbf{1}_{n}^{T}\right)\right\|_{F} < \eta \,\|\mathbf{S} - \mathbf{O}\|_{F} < \epsilon.$$

Since ϵ is arbitrary, we conclude that $\Theta(\mathbf{S})$ is not locally rigid.

 (\Longrightarrow) Let $\epsilon > 0$ be arbitrary. Suppose $\Theta(\mathbf{S})$ is not locally rigid, then there exist another perfect alignment $\mathbf{O}_{\epsilon} \in \mathbb{O}(d)^m$ such that $\|\Theta(\mathbf{O}_{\epsilon}) - \Theta(\mathbf{S})\|_F < \epsilon$ but $\Theta(\mathbf{O}_{\epsilon}) \neq \Theta(\mathbf{SQ})$, equivalently $\|\mathbf{O}_{\epsilon} - \mathbf{S}\|_F < \epsilon$ but $\pi(\mathbf{O}_{\epsilon}) \neq \pi(\mathbf{S})$. Since this true for all $\epsilon > 0$, we conclude that $\pi(\mathbf{S})$ is degenerate.

Proof of Theorem 25. Consider a partition of [1, m] into two non-empty subsets A and B. Suppose rank of $\overline{\mathbf{B}}(\mathbf{S})_{A,B}$ is at most d-2. Let $\Omega_0 \in \text{Skew}(d)$ be such that $\Omega_0 \neq 0$ and $\overline{\mathbf{B}}(\mathbf{S})_{A,B}^T \Omega_0 = 0$ (the existence of Ω_0 follows from the third part of the proof of Theorem 16). WLOG assume that $\mathbf{B}(\mathbf{S})_{A,B}\mathbf{1}_{n'} = 0$ (here n' is as in Definition 10) (perhaps by translating all aligned views by $-\mathbf{B}(\mathbf{S})_{A,B}\mathbf{1}_{n'}$). Then $\mathbf{B}(\mathbf{S})_{A,B}^T\Omega_0 = 0$.

Let $\Omega = [\Omega_i]_1^m$ be such that $\Omega_i = \Omega_0$ for $i \in A$ and $\Omega_i = -\Omega_0$ for $i \in B$. Clearly, $\Omega \in \text{Skew}(d)^m$ such that not all Ω_i are equal. It suffices to show that Ω is a nontrivial certificate of $\mathbf{L}(\mathbf{S})$, equivalently $\text{Tr}(\Omega^T \mathbf{L}(\mathbf{S})\Omega) = 0$.

For $i \in A$,

$$\begin{split} [\mathbf{L}(\mathbf{S})\mathbf{\Omega}]_i &= (-\mathbf{B}(\mathbf{S})_i \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})^T \mathbf{I}_d^m + \sum_{j \in A} \mathbf{B}(\mathbf{S})_i \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_j^T - \sum_{j \in B} \mathbf{B}(\mathbf{S})_i \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_j^T) \mathbf{\Omega}_0 \\ &= -2 \sum_{j \in B} \mathbf{B}(\mathbf{S})_i \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_j^T \mathbf{\Omega}_0 \end{split}$$

and similarly, for $i \in B$

$$[\mathbf{L}(\mathbf{S})\mathbf{\Omega}]_i = 2\sum_{j\in A} \mathbf{B}(\mathbf{S})_i \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}(\mathbf{S})_j^T \mathbf{\Omega}_0.$$

Denote by \mathbf{B}_A and \mathbf{B}_B , the matrices $\sum_{i \in A} \mathbf{B}(\mathbf{S})_i$ and $\sum_{j \in B} \mathbf{B}(\mathbf{S})_j$, respectively. Thus,

$$\operatorname{Tr}(\mathbf{\Omega}^T \mathbf{L}(\mathbf{S})\mathbf{\Omega}) = -4\operatorname{Tr}(\mathbf{\Omega}_0^T \mathbf{B}_A \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}_B^T \mathbf{\Omega}_0).$$
(36)

We are going to show that the above evaluates to zero. Note that the following calculations up to Eq. (41) are general and will be reused in the proof of Theorem 31.

WLOG assume that the first n_1 points lie in the views with indices in $A \setminus B$, next n_2 points lie in the views with indices in $B \setminus A$ and the remaining n_3 points lie in the views with indices in $A \cap B$. Note that $n_1 + n_2 + n_3 = n$ and |A| + |B| = m. Then the matrices \mathbf{B}_A and \mathbf{B}_B (perhaps after permuting the views) have the following structure.

$$egin{array}{rcl} \mathbf{B}_A &=& [& \mathbf{X}_1 & \mathbf{0}_{d imes n_2} & \mathbf{X}_3 & \mathbf{U}_1 + \mathbf{U}_3 & \mathbf{0}_{d imes |B|} &] \ \mathbf{B}_B &=& [& \mathbf{0}_{d imes n_1} & \mathbf{Y}_2 & \mathbf{Y}_3 & \mathbf{0}_{d imes |A|} & \mathbf{V}_2 + \mathbf{V}_3 &] \end{array}$$

where $\mathbf{X}_1 \in \mathbb{R}^{d \times n_1}$ and $\mathbf{Y}_2 \in \mathbb{R}^{d \times n_2}$ contain the sum of the local coordinates of the n_1 and n_2 points, respectively. Also, $\mathbf{X}_3 \in \mathbb{R}^{d \times n_3}$ and $\mathbf{Y}_3 \in \mathbb{R}^{d \times n_3}$ contain the sum of the local coordinates of the remaining n_3 points due to the views with indices in A and B respectively. The matrices $\mathbf{U}_1 \in \mathbb{R}^{d \times |A|}$, $\mathbf{V}_2 \in \mathbb{R}^{d \times |B|}$, $\mathbf{U}_3 \in \mathbb{R}^{d \times |A|}$ and $\mathbf{V}_3 \in \mathbb{R}^{d \times |B|}$ follow from Remark 1.

Further define

then $\mathbf{B}_A = \mathbf{B}_{A \setminus B} + \mathbf{B}_{A,B}$ and $\mathbf{B}_B = \mathbf{B}_{B \setminus A} + \mathbf{B}_{B,A}$. Note that $\mathbf{1}_{n+m}$ lies in the kernel of the four matrices defined above (see Remark 1) and, $\mathbf{B}_{A \setminus B} \mathbf{B}_{B \setminus A}^T = 0$, $\mathbf{B}_{A \setminus B} \mathbf{B}_{B,A}^T = 0$ and $\mathbf{B}_{B \setminus A} \mathbf{B}_{A,B}^T = 0$.

Now, the structure of \mathcal{L}_{Γ} is as follows,

$$\boldsymbol{\mathcal{L}}_{\Gamma} = \begin{bmatrix} \mathcal{D}_{1} & & -\mathcal{K}_{1} \\ & \mathcal{D}_{2} & & -\mathcal{K}_{2} \\ & & \mathcal{D}_{3} + \mathcal{D}_{3} & -\mathcal{K}_{A_{3}} & -\mathcal{K}_{B_{3}} \\ -\mathcal{K}_{1}^{T} & & -\mathcal{K}_{A_{3}}^{T} & \mathcal{D}_{A} & \\ & -\mathcal{K}_{2}^{T} & -\mathcal{K}_{B_{3}}^{T} & & \mathcal{D}_{B} \end{bmatrix}$$

where $\mathcal{K}_1 \in \mathbb{R}^{n_1 \times |A|}$ is the adjacency between the first n_1 points and the views with indices in $A, \mathcal{K}_2 \in \mathbb{R}^{n_2 \times |B|}$ is the adjacency between the next n_2 points and the views with indices in B, and $\mathcal{K}_{A_3} \in \mathbb{R}^{n_3 \times |A|}$ and $\mathcal{K}_{B_3} \in \mathbb{R}^{n_3 \times |B|}$ are the adjacencies between the remaining n_3 points and the views with indices in A and B respectively. As for the remaining matrices, $\mathcal{D}_i = \operatorname{diag}(\mathcal{K}_i \mathbf{1}_{|A|})$ represents the degrees of the points in the bipartite adjacency $\mathcal{K}_i, \mathcal{D}_A = \operatorname{diag}(\mathcal{K}_1^T \mathbf{1}_{n_1} + \mathcal{K}_{A_3}^T \mathbf{1}_{n_3})$ represents the degree of the views i.e. the number of points contained in the views with indices in A, Similarly, $\mathcal{D}_j = \operatorname{diag}(\mathcal{K}_j \mathbf{1}_{|B|})$. and $\mathcal{D}_B = \operatorname{diag}(\mathcal{K}_2^T \mathbf{1}_{n_2} + \mathcal{K}_{B_3}^T \mathbf{1}_{n_3})$.

Since **S** is a perfect alignment, the local coordinates of a point due to all the views containing it are the same. Using the fact that $\mathbf{B}(\mathbf{S})_{A,B}$ represent the local coordinates of the n_3 points contained in views with indices in $A \cap B$ (see Definition 10), we obtain the following equations,

$$\begin{aligned}
\mathbf{X}_3 &= \mathbf{B}(\mathbf{S})_{A,B}\mathcal{D}_3 \\
\mathbf{Y}_3 &= \mathbf{B}(\mathbf{S})_{A,B}\mathcal{D}_3 \\
\mathbf{U}_3 &= -\mathbf{B}(\mathbf{S})_{A,B}\mathcal{K}_{A_3} \\
\mathbf{V}_3 &= -\mathbf{B}(\mathbf{S})_{A,B}\mathcal{K}_{B_3}.
\end{aligned}$$
(38)

Analogous to the above equations, it follows that $\mathbf{U}_1 = \mathbf{X}_1 \mathcal{D}_1^{-1} \mathcal{K}_1$ and $\mathbf{V}_2 = \mathbf{Y}_2 \mathcal{D}_2^{-1} \mathcal{K}_2$. Thus,

$$\mathbf{B}_{A\setminus B} = \begin{bmatrix} \mathbf{X}_1 \mathcal{D}_1^{-1} & \mathbf{0}_{d \times n_2} & \mathbf{0}_{d \times n_3} & \mathbf{0}_{d \times |A|} & \mathbf{0}_{d \times |B|} \end{bmatrix} \mathcal{L}_{\Gamma}
\mathbf{B}_{B\setminus A} = \begin{bmatrix} \mathbf{0}_{d \times n_1} & \mathbf{Y}_2 \mathcal{D}_2^{-1} & \mathbf{0}_{d \times n_3} & \mathbf{0}_{d \times |A|} & \mathbf{0}_{d \times |B|} \end{bmatrix} \mathcal{L}_{\Gamma}$$
(39)

Subsequently,

$$\mathbf{B}_{A\setminus B} \mathcal{L}_{\Gamma}^{\dagger} = \begin{bmatrix} \mathbf{X}_{1} \mathcal{D}_{1}^{-1} & \mathbf{0}_{d \times n_{2}} & \mathbf{0}_{d \times n_{3}} & \mathbf{0}_{d \times |A|} & \mathbf{0}_{d \times |B|} \end{bmatrix} + \mathbf{t}_{A\setminus B} \mathbf{1}_{n+m}^{T} \\
\mathbf{B}_{B\setminus A} \mathcal{L}_{\Gamma}^{\dagger} = \begin{bmatrix} \mathbf{0}_{d \times n_{1}} & \mathbf{Y}_{2} \mathcal{D}_{2}^{-1} & \mathbf{0}_{d \times n_{3}} & \mathbf{0}_{d \times |A|} & \mathbf{0}_{d \times |B|} \end{bmatrix} + \mathbf{t}_{B\setminus A} \mathbf{1}_{n+m}^{T}$$
(40)

for some translation vectors $\mathbf{t}_{A \setminus B}$, $\mathbf{t}_{B \setminus A} \in \mathbb{R}^d$. Since $\mathbf{1}_{n+m}$ lies in ker $(\mathbf{B}_{A \setminus B})$ and ker $(\mathbf{B}_{A \setminus B})$ (see the paragraph after Eq. (37)), thus

$$\mathbf{B}_{A\setminus B}\boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger}\mathbf{B}_{B\setminus A}^{T} = \mathbf{B}_{A\setminus B}\boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger}\mathbf{B}_{B,A}^{T} = \mathbf{B}_{B\setminus A}\boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger}\mathbf{B}_{A,B}^{T} = 0.$$
(41)

Since $\mathbf{B}(\mathbf{S})_{A,B}^T \mathbf{\Omega}_0 = 0$ (by assumption), combining it with Eq. (38) and Eq. (37) yields

$$\mathbf{B}_{A,B}^T \mathbf{\Omega}_0 = 0 \\ \mathbf{B}_{B,A}^T \mathbf{\Omega}_0 = 0.$$

Substituting the above equation and Eq. (41) into Eq. (36), we obtain

$$\operatorname{Tr}(\boldsymbol{\Omega}_{0}^{T}\mathbf{B}_{A}\boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger}\mathbf{B}_{B}^{T}\boldsymbol{\Omega}_{0}) = \operatorname{Tr}(\boldsymbol{\Omega}_{0}^{T}(\mathbf{B}_{A\setminus B} + \mathbf{B}_{A,B})\boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger}(\mathbf{B}_{B\setminus A} + \mathbf{B}_{B,A})^{T}\boldsymbol{\Omega}_{0})$$
$$= \operatorname{Tr}(\boldsymbol{\Omega}_{0}^{T}\mathbf{B}_{A,B}\boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger}\mathbf{B}_{B,A}^{T}\boldsymbol{\Omega}_{0}) = 0.$$

We conclude that Ω is a non-trivial certificate of L(S) and thus $\pi(S)$ is degenerate.

Proof of Lemma 26. WLOG, let the *m*th vertex be removed. The setup is as follows. Let Γ be the bipartite graph representing the correspondence between *m* views and *n* vertices, as described in Section 2. Let Γ_{-} be the bipartite graph obtained after the removal of the vertices representing the *m*th view and the points which lie exclusively in it. Let $\mathbf{D} \in \mathbb{R}^{md \times md}$, $\mathbf{B} \in \mathbb{R}^{md \times (n+m)}$, $\mathbf{D}_{-} \in \mathbb{R}^{(m-1)d \times (m-1)d}$ and $\mathbf{B}_{-} \in \mathbb{R}^{(m-1)d \times (n_{1}+n_{2}+m-1)}$ be the matrices defined in Remark 1 for graphs Γ and Γ_{-} . The structure of the combinatorial Laplacian of Γ and Γ_{-} are

$$\boldsymbol{\mathcal{L}}_{\Gamma} = \begin{bmatrix} \mathcal{D}_{1} & & -\mathcal{K}_{1} & \boldsymbol{0}_{n_{1}} \\ & \mathcal{D}_{2} + \boldsymbol{I}_{n_{2}} & & -\mathcal{K}_{2} & -\boldsymbol{1}_{n_{2}} \\ & & \boldsymbol{I}_{n_{3}} & \boldsymbol{0}_{n_{3} \times (m-1)} & -\boldsymbol{1}_{n_{3}} \\ -\mathcal{K}_{1}^{T} & -\mathcal{K}_{2}^{T} & \boldsymbol{0}_{n_{3}}^{T} & \overline{\mathcal{D}} & \boldsymbol{0}_{m-1} \\ \boldsymbol{0}_{n_{1}}^{T} & -\boldsymbol{1}_{n_{2}}^{T} & -\boldsymbol{1}_{n_{3}}^{T} & \boldsymbol{0}_{m-1}^{T} & n_{2} + n_{3} \end{bmatrix}$$

and

$$\mathcal{L}_{\Gamma_{-}} = egin{bmatrix} \mathcal{D}_1 & -\mathcal{K}_1 \ & \mathcal{D}_2 & -\mathcal{K}_2 \ -\mathcal{K}_1^T & -\mathcal{K}_2^T & \overline{\mathcal{D}} \end{bmatrix},$$

respectively, where

- $\mathcal{K}_1 \in \mathbb{R}^{n_1 \times (m-1)}$ is the bipartite adjacency matrix between the first m-1 views and the n_1 points which lie exclusively in them. Note that the adjacency between such points and the *m*th view is $\mathbf{0}_{n_1}$.
- $\mathcal{K}_2 \in \mathbb{R}^{n_2 \times (m-1)}$ is the bipartite adjacency matrix between the first m-1 views and the n_2 points which lie on the overlap of the *m*th view and the union of the first m-1 views. Note that the adjacency between such points and the *m*th view is $\mathbf{1}_{n_2}$.

- the fifth and the third column in \mathcal{L}_{Γ} correspond to the *m*th view and the n_3 points that lie exclusively in it, respectively. The adjacency between such points and the first m-1 views is $\mathbf{0}_{n_3 \times (m-1)}$, and that with the *m*th view is $\mathbf{1}_{n_3}$.
- $\mathcal{D}_1 = \operatorname{diag}(\mathcal{K}_1 \mathbf{1}_{m-1}), \ \mathcal{D}_2 = \operatorname{diag}(\mathcal{K}_2 \mathbf{1}_{m-1}) \ \text{and} \ \overline{\mathcal{D}} = \operatorname{diag}(\mathcal{K}_1^T \mathbf{1}_{n_1} + \mathcal{K}_2^T \mathbf{1}_{n_2}).$

Note that since Γ is connected by assumption in this work, $n_2 > 0$.

The matrices \mathcal{L}_{Γ} and $\mathcal{L}_{\Gamma_{-}}$ are related as follows. There exist a permutation matrix

$$\mathcal{P}_0 = egin{bmatrix} \mathbf{I}_{n_1} & & & & \ & & \mathbf{I}_{n_2} & & & \ & & & \mathbf{I}_{n_3} & & \ & & & \mathbf{I}_{m-1} & & \ & & & & 1 \end{bmatrix}$$

and a diagonal matrix $\mathcal{D}_0 = \text{diag}((\mathbf{0}_{n_1}, \mathbf{1}_{n_2}, \mathbf{0}_{m-1}))$ such that

$$\mathcal{P}_{0}\mathcal{L}_{\Gamma}\mathcal{P}_{0}^{T} = \begin{bmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{bmatrix} = \begin{bmatrix} \mathcal{L}_{\Gamma_{-}} + \mathcal{D}_{0} & \mathbf{0}_{n_{2} \times n_{3}} & \mathbf{0}_{n_{1}} \\ \mathbf{0}_{n_{1} \times n_{3}} & \mathbf{0}_{n_{2} \times n_{3}} & \mathbf{0}_{(m-1) \times n_{3}} & \mathbf{0}_{m-1} \\ \mathbf{0}_{n_{1} \times n_{3}}^{T} & \mathbf{0}_{n_{2} \times n_{3}}^{T} & \mathbf{0}_{(m-1) \times n_{3}}^{T} & \mathbf{I}_{n_{3}} & -\mathbf{1}_{n_{3}} \\ \mathbf{0}_{n_{1}}^{T} & -\mathbf{1}_{n_{2}}^{T} & \mathbf{0}_{m-1}^{T} & -\mathbf{1}_{n_{3}}^{T} & n_{2} + n_{3} \end{bmatrix}$$
(42)

The rest is divided into three parts. First, we derive the pseudoinverse of the above block matrix using [11, Section 3.6.2]. Then we show that $\mathbf{S}_{-} \coloneqq \mathbf{S}_{-m}$ is a perfect alignment of the m-1 views and finally we show that $[\mathbf{\Omega}_i]_1^{m-1}$ is a certificate of $\mathbf{L}_{-}(\mathbf{S}_{-}) \coloneqq \mathbf{L}_{-m}(\mathbf{S}_{-m})$ when $[\mathbf{\Omega}_i]_1^m$ is a certificate of $\mathbf{L}(\mathbf{S})$.

Part 1. Here we derive the pseudoinverse of the block matrix in Eq. (42). First, we need the following result,

Proposition 43. $\mathcal{L}_{\Gamma_{-}} + \mathcal{D}_0 \succ 0.$

Proof. Since $\mathcal{L}_{\Gamma_{-}} \succeq 0$ and $\mathcal{D}_0 \succeq 0$, it suffices to show that $\ker(\mathcal{L}_{\Gamma_{-}}) \cap \ker(\mathcal{D}_0) = \{0\}$. Recall that the *m*th view contains $n_2 + n_3$ points where $n_2 > 0$ points lie on the overlap of *m*th view and the union of first m - 1 views, and n_3 points lie exclusively in the *m*th view. Removal of the *m*th view and the n_3 points that lie exclusively in it may disconnect Γ to produce Γ_{-} with at most n_2 connected components. The binary vectors \mathbf{u}_i with ones at the indices of the vertices in the *i*th component and zeros elsewhere, form an orthogonal basis of the $\ker(\mathcal{L}_{\Gamma_{-}})$. Note that there exists at least one $k \in [n_1 + 1, n_1 + n_2]$ such that $\mathbf{u}_i(k) = 1$. Thus $\mathbf{u}_i^T \mathcal{D}_0 \mathbf{u}_i > 0$. Moreover, for $i \neq j$, $\mathbf{u}_i^T \mathcal{D}_0 \mathbf{u}_j = 0$. The result follows. \Box

Since $\mathcal{L}_{\Gamma_{-}} + \mathcal{D}_0 \succ 0$, thus $(\mathcal{L}_{\Gamma_{-}} + \mathcal{D}_0)^{\dagger} = (\mathcal{L}_{\Gamma_{-}} + \mathcal{D}_0)^{-1}$ and

$$(\mathcal{L}_{\Gamma_{-}} + \mathcal{D}_{0}) \begin{bmatrix} \mathbf{1}_{n_{1}} \\ \mathbf{1}_{n_{2}} \\ \mathbf{1}_{m-1} \end{bmatrix} = \begin{bmatrix} \mathbf{0}_{n_{1}} \\ \mathbf{1}_{n_{2}} \\ \mathbf{0}_{m-1} \end{bmatrix} \implies (\mathcal{L}_{\Gamma_{-}} + \mathcal{D}_{0})^{\dagger} \begin{bmatrix} \mathbf{0}_{n_{1}} \\ \mathbf{1}_{n_{2}} \\ \mathbf{0}_{m-1} \end{bmatrix} = \begin{bmatrix} \mathbf{1}_{n_{1}} \\ \mathbf{1}_{n_{2}} \\ \mathbf{1}_{m-1} \end{bmatrix}.$$

Using the above equation, the matrix $\mathbf{Z} \coloneqq [(\mathcal{P}_0 \mathcal{L}_{\Gamma} \mathcal{P}_0^T)^{\dagger}]_{22} = \mathbf{A}_{22} - \mathbf{A}_{21} \mathbf{A}_{11}^{\dagger} \mathbf{A}_{12}$ and its pseudoinverse is,

$$\mathbf{Z} = \begin{bmatrix} \mathbf{I}_{n_3} & -\mathbf{1}_{n_3} \\ -\mathbf{1}_{n_3}^T & n_3 \end{bmatrix} \implies \mathbf{Z}^{\dagger} = \begin{bmatrix} \mathbf{I}_{n_3} & \mathbf{0}_{n_3} \\ \mathbf{0}_{n_3}^T & 0 \end{bmatrix}$$

Next,

$$\begin{aligned} &[(\mathcal{P}_{0}\mathcal{L}_{\Gamma}\mathcal{P}_{0}^{T})^{\dagger}]_{11} \\ &= (\mathcal{L}_{\Gamma_{-}} + \mathcal{D}_{0})^{\dagger} + ((\mathcal{L}_{\Gamma_{-}} + \mathcal{D}_{0})^{\dagger}\mathbf{A}_{12})\mathbf{Z}^{\dagger}(\mathbf{A}_{21}(\mathcal{L}_{\Gamma_{-}} + \mathcal{D}_{0})^{\dagger}) \\ &= (\mathcal{L}_{\Gamma_{-}} + \mathcal{D}_{0})^{\dagger} + \begin{bmatrix} \mathbf{0}_{n_{1}\times n_{3}} & \mathbf{1}_{n_{1}} \\ \mathbf{0}_{n_{2}\times n_{3}} & \mathbf{1}_{n_{2}} \\ \mathbf{0}_{(m-1)\times n_{3}} & \mathbf{1}_{m-1} \end{bmatrix} \begin{bmatrix} \mathbf{I}_{n_{3}} & \mathbf{0}_{n_{3}} \\ \mathbf{0}_{n_{3}}^{T} & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{0}_{n_{1}\times n_{3}}^{T} & \mathbf{0}_{n_{2}\times n_{3}}^{T} & \mathbf{0}_{(m-1)\times n_{3}}^{T} \\ \mathbf{1}_{n_{1}}^{T} & \mathbf{1}_{n_{2}}^{T} & \mathbf{1}_{m-1} \end{bmatrix} \\ &= (\mathcal{L}_{\Gamma_{-}} + \mathcal{D}_{0})^{\dagger} \end{aligned} \tag{43}$$

As above, $[(\mathcal{P}_0 \mathcal{L}_{\Gamma} \mathcal{P}_0^T)^{\dagger}]_{12} = -(\mathbf{A}_{11}^{\dagger} \mathbf{A}_{12}) \mathbf{Z}^{\dagger} = 0$ and similarly $[(\mathcal{P}_0 \mathcal{L}_{\Gamma} \mathcal{P}_0^T)^{\dagger}]_{21} = 0$. Thus,

$$(\mathcal{P}_0 \mathcal{L}_{\Gamma} \mathcal{P}_0^T)^{\dagger} = \begin{bmatrix} (\mathcal{L}_{\Gamma_-} + \mathcal{D}_0)^{\dagger} & & \\ & \mathbf{I}_{n_3} & \\ & & 0 \end{bmatrix}.$$

Part 2. Now, let $\mathbf{S} = [\mathbf{S}_i]_1^m$ and $\mathbf{S}_- = [\mathbf{S}_i]_1^{m-1}$. Intuitively, it should be clear that \mathbf{S}_- is a perfect alignment for the m-1 views. Since \mathbf{S} is a perfect alignment, the alignment error $\operatorname{Tr}(\mathbf{S}^T(\mathbf{D} - \mathbf{B}\mathcal{L}_{\Gamma}^{\dagger}\mathbf{B}^T)\mathbf{S}) = 0$ (see Eq. (8)). Here we will show that the alignment error after the removal of the *m*th view is still zero i.e. $\operatorname{Tr}(\mathbf{S}_-^T(\mathbf{D}_- - \mathbf{B}_-\mathcal{L}_{\Gamma_-}\mathbf{B}_-^T)\mathbf{S}_-) = 0$. Let $\mathbf{B}_1^* \in \mathbb{R}^{d \times n_1}, \mathbf{B}_2^* \in \mathbb{R}^{d \times n_2}$ and $\mathbf{B}_3^* \in \mathbb{R}^{d \times n_3}$ contain the coordinates (after alignment

Let $\mathbf{B}_1^* \in \mathbb{R}^{d \times n_1}$, $\mathbf{B}_2^* \in \mathbb{R}^{d \times n_2}$ and $\mathbf{B}_3^* \in \mathbb{R}^{d \times n_3}$ contain the coordinates (after alignment with **S**) of the n_1 points that lie exclusively in the first m-1 views, of the n_2 points that lie on the overlap of the *m*th view with the remaining views, and of the n_3 points that lie exclusively in the *m*th view, respectively. Then, it suffices to show the following (by taking the trace of the difference of the two equations below, the result follows),

Proposition 44.

$$\mathbf{S}^T \mathbf{B} \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}^T \mathbf{S} = \mathbf{S}_{-}^T \mathbf{B}_{-} \boldsymbol{\mathcal{L}}_{\Gamma_{-}}^{\dagger} \mathbf{B}_{-}^T \mathbf{S}_{-} + \mathbf{B}_2^* \mathbf{B}_2^{*^T} + \mathbf{B}_3^* \mathbf{B}_3^{*^T}$$

and

$$\mathbf{S}^T \mathbf{D} \mathbf{S} = \mathbf{S}_{-}^T \mathbf{D}_{-} \mathbf{S}_{-} + \mathbf{B}_2^* \mathbf{B}_2^{*T} + \mathbf{B}_3^* \mathbf{B}_3^{*T}.$$

Proof. The second equation follows trivially from Eq. (6), Remark 1 and the fact that, since the *m* views are perfectly aligned, the local coordinates of the points are the same as those in the matrices \mathbf{B}_1^* , \mathbf{B}_2^* and \mathbf{B}_3^* . We proceed to prove the first equation.

Since \mathcal{P}_0 is a permutation matrix, $\mathbf{S}^T \mathbf{B} \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}^T \mathbf{S} = (\mathbf{S}^T \mathbf{B} \mathcal{L}_{\Gamma}^{\dagger} \mathcal{P}_0^T) (\mathcal{P}_0 \mathbf{B}^T \mathbf{S})$. Then, using the same idea as in Eq. (38), Eq. (39) and Eq. (40) in the proof of Theorem 25, we obtain

$$\mathbf{S}^{T}\mathbf{B} = \begin{bmatrix} \mathbf{B}_{1}^{*}\mathcal{D}_{1} & \mathbf{B}_{2}^{*}(\mathcal{D}_{2} + \mathbf{I}_{n_{2}}) & \mathbf{B}_{3}^{*} & -(\mathbf{B}_{1}^{*}\mathcal{K}_{1} + \mathbf{B}_{2}^{*}\mathcal{K}_{2}) & -(\mathbf{B}_{2}^{*}\mathbf{1}_{n_{2}} + \mathbf{B}_{3}^{*}\mathbf{1}_{n_{3}}) \end{bmatrix} \quad (44)$$
$$= \begin{bmatrix} \mathbf{B}_{1}^{*} & \mathbf{B}_{2}^{*} & \mathbf{B}_{3}^{*} & \mathbf{0}_{d \times (m-1)} & \mathbf{0}_{d} \end{bmatrix} \mathcal{L}_{\Gamma}$$

and thus

$$\mathbf{S}^{T}\mathbf{B}\boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} = \begin{bmatrix} \mathbf{B}_{1}^{*} & \mathbf{B}_{2}^{*} & \mathbf{B}_{3}^{*} & \mathbf{0}_{d \times (m-1)} & \mathbf{0}_{d} \end{bmatrix} + \mathbf{t}\mathbf{1}_{n+m}^{T}$$
(45)

for some translation vector $\mathbf{t} \in \mathbb{R}^d$. Similarly,

$$\mathbf{S}_{-}^{T}\mathbf{B}_{-} = \begin{bmatrix} \mathbf{B}_{1}^{*}\mathcal{D}_{1} & \mathbf{B}_{2}^{*}\mathcal{D}_{2} & -(\mathbf{B}_{1}^{*}\mathcal{K}_{1} + \mathbf{B}_{2}^{*}\mathcal{K}_{2}) \end{bmatrix} = \begin{bmatrix} \mathbf{B}_{1}^{*} & \mathbf{B}_{2}^{*} & \mathbf{0}_{d \times (m-1)} \end{bmatrix} \mathcal{L}_{\Gamma_{-}}$$
(46)

and thus

$$\mathbf{S}_{-}^{T}\mathbf{B}_{-}\boldsymbol{\mathcal{L}}_{\Gamma_{-}}^{\dagger} = \begin{bmatrix} \mathbf{B}_{1}^{*} & \mathbf{B}_{2}^{*} & \mathbf{0}_{d\times(m-1)} \end{bmatrix} + \mathbf{t}_{-}\mathbf{v}_{-}^{T}$$

for some translation vector $\mathbf{t}_{-} \in \mathbb{R}^{d}$ and $\mathbf{v}_{-} \in \ker(\mathcal{L}_{\Gamma_{-}})$. From Proposition 2, we have $\ker(\mathcal{L}_{\Gamma_{-}}) \subseteq \ker(\mathbf{B}_{-})$, thus

$$\mathbf{S}_{-}^{T}\mathbf{B}_{-}\boldsymbol{\mathcal{L}}_{\Gamma_{-}}^{\dagger}\mathbf{B}_{-}^{T}\mathbf{S}_{-} = \begin{bmatrix} \mathbf{B}_{1}^{*} & \mathbf{B}_{2}^{*} & \mathbf{0}_{d\times(m-1)} \end{bmatrix} \mathbf{B}_{-}^{T}\mathbf{S}_{-}.$$
(47)

Now, combining Eq. (44) and Eq. (46) we obtain

$$\begin{split} \mathbf{S}^T \mathbf{B} \mathcal{P}_0^T &= \begin{bmatrix} \mathbf{B}_1^* \mathcal{D}_1 & \mathbf{B}_2^* (\mathcal{D}_2 + \mathbf{I}_{n_2}) & -(\mathbf{B}_1^* \mathcal{K}_1 + \mathbf{B}_2^* \mathcal{K}_2) & \mathbf{B}_3^* & -(\mathbf{B}_2^* \mathbf{1}_{n_2} + \mathbf{B}_3^* \mathbf{1}_{n_3}) \end{bmatrix} \\ &= \begin{bmatrix} \mathbf{S}_-^T \mathbf{B}_- & \mathbf{0}_{d \times n_3} & \mathbf{0}_d \end{bmatrix} + \begin{bmatrix} \mathbf{0}_{d \times n_1} & \mathbf{B}_2^* & \mathbf{0}_{d \times (m-1)} & \mathbf{B}_3^* & -(\mathbf{B}_2^* \mathbf{1}_{n_2} + \mathbf{B}_3^* \mathbf{1}_{n_3}) \end{bmatrix}. \end{split}$$

By observing that $\mathbf{1}_{n+m}$ is in the kernel of $\mathbf{S}^T \mathbf{B} \mathcal{P}_0^T$, combining the above equation with Eq. (45), and using Eq. (47), we obtain

$$\begin{aligned} \mathbf{S}^{T} \mathbf{B} \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \mathbf{B}^{T} \mathbf{S} \\ &= (\mathbf{S}^{T} \mathbf{B} \boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger} \boldsymbol{\mathcal{P}}_{0}^{T}) (\boldsymbol{\mathcal{P}}_{0} \mathbf{B}^{T} \mathbf{S}) \\ &= (\begin{bmatrix} \mathbf{B}_{1}^{*} & \mathbf{B}_{2}^{*} & \mathbf{0}_{d \times (m-1)} & \mathbf{B}_{3}^{*} & \mathbf{0}_{d} \end{bmatrix} + \mathbf{t} \mathbf{1}_{n+m}^{T}) (\boldsymbol{\mathcal{P}}_{0} \mathbf{B}^{T} \mathbf{S}) \\ &= \begin{bmatrix} \mathbf{B}_{1}^{*} & \mathbf{B}_{2}^{*} & \mathbf{0}_{d \times (m-1)} & \mathbf{B}_{3}^{*} & \mathbf{0}_{d} \end{bmatrix} \begin{bmatrix} \mathbf{S}_{-}^{T} \mathbf{B}_{-} & \mathbf{0}_{d \times n_{3}} & \mathbf{0}_{d} \end{bmatrix}^{T} + \mathbf{B}_{2}^{*} \mathbf{B}_{2}^{*T} + \mathbf{B}_{3}^{*} \mathbf{B}_{3}^{*T} \\ &= \mathbf{S}_{-}^{T} \mathbf{B}_{-} \boldsymbol{\mathcal{L}}_{\Gamma-}^{\dagger} \mathbf{B}_{-}^{T} \mathbf{S}_{-} + \mathbf{B}_{2}^{*} \mathbf{B}_{2}^{*T} + \mathbf{B}_{3}^{*} \mathbf{B}_{3}^{*T} \end{aligned}$$

Part 3. Now let $\mathbf{L}(\mathbf{S})$ and $\mathbf{L}_{-}(\mathbf{S}_{-})$ be the matrices, as described in Eq. (24) for the two graphs Γ and Γ_{-} and the corresponding views. Let $\mathbf{\Omega} = [\mathbf{\Omega}_{i}]_{1}^{m}$ be a certificate of $\mathbf{L}(\mathbf{S})$. By Remark 5, $\mathbf{\Omega}' = \mathbf{\Omega} - [\mathbf{\Omega}_{m}]_{1}^{m}$ is also a certificate of $\mathbf{L}(\mathbf{S})$ and in particular $\mathbf{\Omega}'_{m} = 0$. Define $\mathbf{\Omega}_{-} = [\mathbf{\Omega}'_{i}]_{1}^{m-1}$. We are going to show that $\operatorname{Tr}(\mathbf{\Omega}_{-}^{T}\mathbf{L}_{-}(\mathbf{S}_{-})\mathbf{\Omega}_{-}) = 0$ i.e. $\mathbf{\Omega}_{-}$ is a certificate of $\mathbf{L}_{-}(\mathbf{S}_{-})$. Then using Remark 5, it follows that $[\mathbf{\Omega}_{i}]_{1}^{m-1}$ (which equals $\mathbf{\Omega}_{-} + [\mathbf{\Omega}_{m}]_{1}^{m-1}$) is a certificate of $\mathbf{L}_{-}(\mathbf{S}_{-})$.

First, we note that

$$\begin{split} \mathbf{B}(\mathbf{S})\mathcal{P}_{0}^{T} &= \begin{bmatrix} \mathbf{B}_{-}(\mathbf{S}_{-}) & \mathbf{0}_{(m-1)d \times n_{3}} & \mathbf{0}_{(m-1)d} \\ \begin{bmatrix} \mathbf{0}_{d \times n_{1}} & \mathbf{B}_{2}^{*} & \mathbf{0}_{d \times (m-1)} \end{bmatrix} & \mathbf{B}_{3}^{*} & -(\mathbf{B}_{2}^{*}\mathbf{1}_{n_{2}} + \mathbf{B}_{3}^{*}\mathbf{1}_{n_{3}}) \end{bmatrix} \\ \mathbf{D}(\mathbf{S}) &= \begin{bmatrix} \mathbf{D}_{-}(\mathbf{S}_{-}) & \\ & \mathbf{B}_{2}^{*}\mathbf{B}_{2}^{*T} + \mathbf{B}_{3}^{*}\mathbf{B}_{3}^{*T} \end{bmatrix} \end{split}$$

and then using the fact that $\mathbf{\Omega}'$ is a certificate of $\mathbf{L}(\mathbf{S})$, then $\mathbf{\Omega}'_m = 0$, followed by the above equations and Eq. (43), we obtain

$$0 = \operatorname{Tr}(\boldsymbol{\Omega}^{T} \mathbf{L}(\mathbf{S})\boldsymbol{\Omega}^{\prime})$$

= $\operatorname{Tr}(\boldsymbol{\Omega}^{T} (\mathbf{B}(\mathbf{S})\mathcal{P}_{0}^{T}\mathcal{P}_{0}\mathcal{L}_{\Gamma}^{\dagger}\mathcal{P}_{0}^{T}\mathcal{P}_{0}\mathbf{B}(\mathbf{S})^{T} - \mathbf{D}(\mathbf{S}))\boldsymbol{\Omega}^{\prime})$
= $\operatorname{Tr}(\boldsymbol{\Omega}^{T} (\mathbf{B}(\mathbf{S})\mathcal{P}_{0}^{T}(\mathcal{P}_{0}\mathcal{L}_{\Gamma}\mathcal{P}_{0}^{T})^{\dagger}\mathcal{P}_{0}\mathbf{B}(\mathbf{S})^{T} - \mathbf{D}(\mathbf{S}))\boldsymbol{\Omega}^{\prime})$
= $\operatorname{Tr}(\boldsymbol{\Omega}_{-}^{T}(\mathbf{B}_{-}(\mathbf{S}_{-})(\mathcal{P}_{0}\mathcal{L}_{\Gamma}^{\dagger}\mathcal{P}_{0}^{T})_{11}\mathbf{B}_{-}(\mathbf{S}_{-})^{T} - \mathbf{D}_{-}(\mathbf{S}))\boldsymbol{\Omega}_{-})$
= $\operatorname{Tr}(\boldsymbol{\Omega}_{-}^{T}(\mathbf{B}_{-}(\mathbf{S}_{-})(\mathcal{L}_{\Gamma_{-}} + \mathcal{D}_{0})^{\dagger}\mathbf{B}_{-}(\mathbf{S}_{-})^{T} - \mathbf{D}_{-}(\mathbf{S}))\boldsymbol{\Omega}_{-})$

From Proposition 18 and Eq. (22), we have $\mathbf{L}_{-}(\mathbf{S}_{-}) = \mathbf{B}_{-}(\mathbf{S}_{-})\boldsymbol{\mathcal{L}}_{\Gamma_{-}}^{\dagger}\mathbf{B}_{-}(\mathbf{S}_{-})^{T} - \mathbf{D}_{-}(\mathbf{S}_{-})$, thus

$$\operatorname{Tr}(\mathbf{\Omega}_{-}^{T}(\mathbf{B}_{-}(\mathbf{S}_{-})((\mathcal{L}_{\Gamma_{-}}+\mathcal{D}_{0})^{\dagger}-\mathcal{L}_{\Gamma_{-}}^{\dagger})\mathbf{B}_{-}(\mathbf{S}_{-})^{T})\mathbf{\Omega}_{-})+\operatorname{Tr}(\mathbf{\Omega}_{-}^{T}\mathbf{L}_{-}(\mathbf{S}_{-})\mathbf{\Omega}_{-})=0$$

Since $\mathbf{L}_{-}(\mathbf{S}_{-}) \leq 0$, to show that $\operatorname{Tr}(\mathbf{\Omega}_{-}^{T}\mathbf{L}_{-}(\mathbf{S}_{-})\mathbf{\Omega}_{-}) = 0$, it suffices to show the following

Proposition 45.

$$\mathbf{B}_{-}(\mathbf{S}_{-})((\boldsymbol{\mathcal{L}}_{\Gamma_{-}}+\boldsymbol{\mathcal{D}}_{0})^{\dagger}-\boldsymbol{\mathcal{L}}_{\Gamma_{-}}^{\dagger})\mathbf{B}_{-}(\mathbf{S}_{-})^{T} \leq 0.$$

Proof. Since $\mathcal{L}_{\Gamma_{-}} \succeq 0$, consider the decomposition

$$\mathcal{L}_{\Gamma_{-}} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^{T} = \begin{bmatrix} \mathbf{U}_{1} & \mathbf{U}_{2} \end{bmatrix} \begin{bmatrix} \mathbf{\Lambda}_{1} & \ & 0 \end{bmatrix} \begin{bmatrix} \mathbf{U}_{1}^{T} \\ \mathbf{U}_{2}^{T} \end{bmatrix}$$

where the $1 \leq n' \leq n_2$ columns of $\mathbf{U}_2 \in \mathbb{R}^{(n_1+n_2+m-1)\times n'}$ form the orthogonal basis of the $\ker(\mathcal{L}_{\Gamma_-})$ as described in the proof of Proposition 43, and $\Lambda_1 \succ 0$. From Proposition 2 and Eq. (21), $\mathbf{B}_-(\mathbf{S}_-)\mathbf{U}_2 = 0$. Thus,

$$\mathbf{B}_{-}(\mathbf{S}_{-})\mathbf{U} = \begin{bmatrix} \mathbf{B}_{-}(\mathbf{S}_{-})\mathbf{U}_{1} & 0 \end{bmatrix}$$
(48)

Then note that,

$$\left(\left(\mathcal{L}_{\Gamma_{-}}+\mathcal{D}_{0}\right)^{\dagger}-\mathcal{L}_{\Gamma_{-}}^{\dagger}\right)=\mathbf{U}\left\{\left(\begin{bmatrix}\mathbf{\Lambda}_{1}\\0\end{bmatrix}+\mathbf{U}^{T}\mathcal{D}_{0}\mathbf{U}\right)^{\dagger}-\begin{bmatrix}\mathbf{\Lambda}_{1}^{-1}\\0\end{bmatrix}\right\}\mathbf{U}^{T}.$$
(49)

Using [16][Eq. (10,11,17,19)] and straightforward calculations, we obtain

$$\left(\begin{bmatrix} \mathbf{\Lambda}_1 \\ & 0 \end{bmatrix} + \mathbf{U}^T \mathcal{D}_0 \mathbf{U} \right)^{\dagger} = \begin{bmatrix} \mathbf{\Lambda}_1^{-1} \\ & 0 \end{bmatrix} + \begin{bmatrix} \mathbf{W}_1 \\ & \mathbf{W}_2 \end{bmatrix}$$

where $\mathbf{W}_1 = -\mathbf{\Lambda}_1^{-1} \mathbf{U}_1^T \mathcal{D}_0 (\mathbf{I} + \mathcal{D}_0 \mathbf{U}_1 \mathbf{\Lambda}_1^{-1} \mathbf{U}_1^T \mathcal{D}_0)^{-1} \mathcal{D}_0 \mathbf{U}_1 \mathbf{\Lambda}_1^{-1}$ and $\mathbf{W}_2 = (\mathbf{U}_2^T \mathcal{D}_0 \mathbf{U}_2)^{\dagger}$. In particular $\mathbf{W}_1 \leq 0$ and $\mathbf{W}_2 \succeq 0$. Combining above equation with Eq. (48) and Eq. (49),

$$\mathbf{B}_{-}(\mathbf{S}_{-})((\boldsymbol{\mathcal{L}}_{\Gamma_{-}}+\mathcal{D}_{0})^{\dagger}-\boldsymbol{\mathcal{L}}_{\Gamma_{-}}^{\dagger})\mathbf{B}_{-}(\mathbf{S}_{-})^{T}=\mathbf{B}_{-}(\mathbf{S}_{-})\mathbf{U}_{1}\mathbf{W}_{1}\mathbf{U}_{1}^{T}\mathbf{B}_{-}(\mathbf{S}_{-})^{T} \leq 0.$$

We conclude that $\operatorname{Tr}(\mathbf{\Omega}_{-}^{T}\mathbf{L}_{-}(\mathbf{S}_{-})\mathbf{\Omega}_{-}) = 0$ and thus $\mathbf{\Omega}_{-}$ is a certificate of $\mathbf{L}_{-}(\mathbf{S}_{-})$.

Proof of Proposition 27. It suffices to show that if the *i*th and *j*th vertices are adjacent in \mathbb{G} then $\Omega_i = \Omega_j$. For m = 2, the result is a direct consequence of Theorem 24. Suppose the result holds for m - 1 views for some m > 2. We show that the result holds for mviews. If there are no edges in \mathbb{G} , then the result holds trivially. Suppose *i*th and *j*th vertices are adjacent in \mathbb{G} . Let $r \in [1, m] \setminus \{i, j\}$. We remove the *r*th view and the points which lie exclusively in it. Then by Lemma 26, $[\Omega_k]_{k \in [1,m] \setminus r}$ is a certificate of $\mathbf{L}_{-r}(\mathbf{S}_{-r})$. Now construct \mathbb{G}_{-r} (in the same way as \mathbb{G}) and note that the vertices corresponding to the *i*th and *j*th views are still adjacent. Thus, by the induction hypothesis we conclude that $\Omega_i = \Omega_j$.

Proof of Theorem 28. The result holds for two views (see Theorem 24). Suppose the result holds for m - 1 views for some m > 2. We will show that the result holds for m views. Suppose $|\mathbb{G}^*(\mathbf{S})| = 1$. Let $\mathbf{\Omega}$ be a certificate of $\mathbf{L}(\mathbf{S})$. We need to show that $\mathbf{\Omega}$ is trivial. Since $|\mathbb{G}^*(\mathbf{S})| = 1$, \mathbb{G} must have a connected component with at least two views. Pick one such component and note that there exist a view in it such that removing it will not disconnect the component. Let it be the *i*th view. Consider removing the *i*th view and the points which lie exclusively in it. Note that for the new set of views we still have $|\mathbb{G}^*(\mathbf{S}_{-i})| = 1$ (where $\mathbb{G}^*_{-i}(\mathbf{S}_{-i})$ is constructed in the same manner as $\mathbb{G}^*(\mathbf{S})$). By Lemma 26 and the induction hypothesis, we conclude that $[\mathbf{\Omega}_j]_{j\in[1,m]\setminus\{i\}}$ must be trivial. Then, by Proposition 27 we conclude that $\mathbf{\Omega}$ is trivial.

Proof of Theorem 31. Consider a partition of [1, m] into two non-empty subsets A and B. Suppose the rank of $\overline{\mathbf{B}(\mathbf{S})}_{A,B}$ is at most d-1. As in the proof of Theorem 25, WLOG assume that $\mathbf{B}(\mathbf{S})_{A,B}\mathbf{1}_{n'} = 0$ (where n' is as in Definition 10). Then the rank of $\mathbf{B}(\mathbf{S})_{A,B}$ is at most d-1. We are going to construct another perfect alignment \mathbf{S}' such that $\pi(\mathbf{S}) \neq \pi(\mathbf{S}')$, thus concluding that \mathbf{S} is not unique.

Let $\mathbf{V}_1, \mathbf{V}_2 \in \mathbb{O}(d)$ and Σ be the diagonal matrix containing the singular values of $\mathbf{B}(\mathbf{S})_{A,B}$ such that $\mathbf{B}(\mathbf{S})_{A,B} = \mathbf{V}_1 \Sigma \mathbf{V}_2^T$. Since rank of $\mathbf{B}(\mathbf{S})_{A,B} \leq d-1$, there exist $\mathbf{U} \in \mathbb{O}(d)$ such that $\mathbf{U} \neq \mathbf{I}_d$ and $\Sigma = \mathbf{U}\Sigma$. Define $\mathbf{Q} = \mathbf{V}_1 \mathbf{U}^T \mathbf{V}_1^T$. Then $\mathbf{Q} \neq \mathbf{I}_d$ and

$$\mathbf{Q}^T \mathbf{B}(\mathbf{S})_{A,B} = \mathbf{B}(\mathbf{S})_{A,B}.$$
(50)

Define $\mathbf{S}' \in \mathbb{O}(d)^m$ such that $\mathbf{S}'_i = \mathbf{S}_i \mathbf{Q}$ for all $i \in A$ and $\mathbf{S}'_j = \mathbf{S}_j$ for all $j \in B$. Clearly, $\mathbf{S}' \neq \mathbf{S}$. We will show that \mathbf{S}' is another perfect alignment. To this end,

$$\operatorname{Tr}(\mathbf{S}^{T}\mathbf{C}\mathbf{S}^{\prime}) = \sum_{i=1}^{m} \sum_{j=1}^{m} \operatorname{Tr}(\mathbf{S}_{i}^{T}\mathbf{C}_{ij}\mathbf{S}_{j}^{\prime})$$
$$= \sum_{\substack{i \in A, j \in A \\ i \in B, j \in B}} \operatorname{Tr}(\mathbf{S}_{i}^{T}\mathbf{C}_{ij}\mathbf{S}_{j}) + 2\sum_{i \in A} \sum_{j \in B} \operatorname{Tr}(\mathbf{Q}^{T}\mathbf{S}_{i}^{T}\mathbf{C}_{ij}\mathbf{S}_{j})$$

Since, for $i \in A$ and $j \in B$, $\mathbf{C}_{ij} = \mathbf{B}_i \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}_j^T$, it suffices to show that

$$\operatorname{Tr}\left(\left(\sum_{i\in A}\mathbf{B}(\mathbf{S})_i\right)\mathcal{L}_{\Gamma}^{\dagger}\left(\sum_{j\in B}\mathbf{B}(\mathbf{S})_j^T\right)\right) = \operatorname{Tr}\left(\mathbf{Q}^T\left(\sum_{i\in A}\mathbf{B}(\mathbf{S})_i\right)\mathcal{L}_{\Gamma}^{\dagger}\left(\sum_{j\in B}\mathbf{B}(\mathbf{S})_j^T\right)\right)$$

Define \mathbf{B}_A , \mathbf{B}_B , $\mathbf{B}_{A\setminus B}$, $\mathbf{B}_{A,B}$, $\mathbf{B}_{B\setminus A}$ and $\mathbf{B}_{B,A}$ as in the proof of Theorem 25, then it suffices to show that $\operatorname{Tr}(\mathbf{Q}^T \mathbf{B}_A \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}_B^T) = \operatorname{Tr}(\mathbf{B}_A \mathcal{L}_{\Gamma}^{\dagger} \mathbf{B}_B^T)$. Using Eq. (50, 37, 38) we obtain $\mathbf{Q}^T \mathbf{B}_{A,B} = \mathbf{B}_{A,B}$ and this combined with Eq. (41), yields

$$\begin{aligned} \operatorname{Tr}(\mathbf{Q}^{T}\mathbf{B}_{A}\boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger}\mathbf{B}_{B}^{T}) &= \operatorname{Tr}(\mathbf{Q}^{T}(\mathbf{B}_{A\setminus B} + \mathbf{B}_{A,B})\boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger}(\mathbf{B}_{B\setminus A} + \mathbf{B}_{B,A})^{T}) \\ &= \operatorname{Tr}(\mathbf{Q}^{T}\mathbf{B}_{A,B}\boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger}\mathbf{B}_{B,A}^{T}) \\ &= \operatorname{Tr}(\mathbf{B}_{A,B}\boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger}\mathbf{B}_{B,A}^{T}) \\ &= \operatorname{Tr}((\mathbf{B}_{A\setminus B} + \mathbf{B}_{A,B})\boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger}(\mathbf{B}_{B\setminus A} + \mathbf{B}_{B,A})^{T}) \\ &= \operatorname{Tr}(\mathbf{B}_{A}\boldsymbol{\mathcal{L}}_{\Gamma}^{\dagger}\mathbf{B}_{B}^{T}). \end{aligned}$$

Proof of Proposition 32. By replacing \mathbb{G} with $\overline{\mathbb{G}}$ and Theorem 24 with Theorem 30, the inductive proof is essentially same as the proof of Proposition 27.

Proof of Theorem 33. Again, by replacing \mathbb{G} with $\overline{\mathbb{G}}$, Theorem 24 with Theorem 30 and Proposition 27 with Proposition 32, the inductive proof is essentially same as the proof of Theorem 28.

Proof of Proposition 36. Fix $\phi = 1/2$. Then $\|\boldsymbol{\xi}_i\|_F \leq 1/2$. From Proposition 35,

$$\begin{aligned} \|R_{\text{QR}}(\mathbf{S}, \boldsymbol{\xi}) - (\mathbf{S} + \boldsymbol{\xi})\|_{F}^{2} &= \sum_{1}^{m} \|\text{qf}(\mathbf{S}_{i} + \boldsymbol{\xi}_{i}) - (\mathbf{S}_{i} + \boldsymbol{\xi}_{i})\|_{F}^{2} \le M^{2} \sum_{1}^{m} \|\boldsymbol{\xi}_{i}\|_{F}^{4} \\ &\leq M^{2} \left(\sum_{1}^{m} \|\boldsymbol{\xi}_{i}\|_{F}^{2}\right)^{2} = M^{2} \|\boldsymbol{\xi}\|_{F}^{4}. \end{aligned}$$

Thus, the result follows for the same values of ϕ and M as in Proposition 35.

Proof of Proposition 37. The proof is essentially the same as the one in [21]. Due to (A2) in Section 5.2, WLOG we assume that $\operatorname{grad} F(\mathbf{S}^k) \neq 0$ for all $k \geq 0$. Then, from the proof of Proposition 7, since $\operatorname{grad} F(\mathbf{S})_i = \frac{1}{2} \left(\nabla F(\mathbf{S})_i - \mathbf{S}_i \nabla F(\mathbf{S})_i^T \mathbf{S}_i \right)$, thus

$$g(\nabla F(\mathbf{S}^k)_i, \operatorname{grad} F(\mathbf{S}^k)_i) = \frac{1}{2} \left(\left\| \nabla F(\mathbf{S}^k)_i \right\|_F^2 - g(\nabla F(\mathbf{S}^k)_i, \mathbf{S}_i^k \nabla F(\mathbf{S}^k)_i^T \mathbf{S}_i^k) \right)$$
$$= \frac{1}{4} \left\| \nabla F(\mathbf{S}^k)_i - \mathbf{S}_i^k \nabla F(\mathbf{S}^k)_i^T \mathbf{S}_i^k \right\|_F^2$$
$$= \left\| \operatorname{grad} F(\mathbf{S}^k)_i \right\|_F^2.$$

Subsequently,

$$g(\nabla F(\mathbf{S}^k), \operatorname{grad} F(\mathbf{S}^k)) = \sum_{1}^{m} g(\nabla F(\mathbf{S}^k)_i, \operatorname{grad} F(\mathbf{S}^k)_i)$$
$$= \sum_{1}^{m} \left\| \operatorname{grad} F(\mathbf{S}^k)_i \right\|_F^2 = \left\| \operatorname{grad} F(\mathbf{S}^k) \right\|_F^2$$

This, together with lines 3 and 4 of Algorithm 1, and Eq. (33), implies that

$$F(\mathbf{S}^{k+1}) - F(\mathbf{S}^k) \le -\gamma \alpha_k \left\| \operatorname{grad} F(\mathbf{S}^k) \right\|_F^2$$
(51)

Since $\mathbb{O}(d)^m$ is compact and F is analytic, thus F is bounded. Since $\alpha_k \in (0,1]$ for all $k \ge 0$, using the above equation,

$$\sum_{0}^{\infty} \alpha_k^2 \left\| \operatorname{grad} F(\mathbf{S}^k) \right\|_F^2 \le \sum_{0}^{\infty} \alpha_k \left\| \operatorname{grad} F(\mathbf{S}^k) \right\|_F^2 \le \frac{1}{\gamma} (F(\mathbf{S}^0) - \lim F(\mathbf{S}^k)) < \infty.$$

The result follows.

Proof of (A1) in Section 5.2. From Proposition 37, there exists $k_1 \ge 0$ such that $\alpha_k \| \operatorname{grad} F(\mathbf{S}^k) \| \le 1/2$ for all $k \ge k_1$. Then from Proposition 36, for all $k \ge k_1$, we have

$$\begin{split} \left\| \mathbf{S}^{k+1} - \mathbf{S}^{k} \right\|_{F} &= \left\| R_{QR}(\mathbf{S}^{k}, -\alpha_{k} \operatorname{grad} F(\mathbf{S}^{k})) - \mathbf{S}^{k} \right\|_{F} \\ &\leq M \alpha_{k}^{2} \left\| \operatorname{grad} F(\mathbf{S}^{k}) \right\|_{F}^{2} + \alpha_{k} \left\| \operatorname{grad} F(\mathbf{S}^{k}) \right\|_{F} \\ &\leq (M/2 + 1) \alpha_{k} \left\| \operatorname{grad} F(\mathbf{S}^{k}) \right\|_{F}. \end{split}$$

Finally, for all $k \ge k_1$, using Eq (51),

$$F(\mathbf{S}^{k+1}) - F(\mathbf{S}^k) \le -\frac{2\gamma}{M+2} \left\| \operatorname{grad} F(\mathbf{S}^k) \right\|_F \cdot \left\| \mathbf{S}^{k+1} - \mathbf{S}^k \right\|_F.$$

Thus, (A1) holds for $\kappa_0 = 2\gamma/(M+2)$.

Proof of (A3) in Section 5.2. Since $\nabla F(\mathbf{S}) = 2\mathbf{CS}$ is Lipschitz with Lipschitz constant $L_F \leq 2 \|\mathbf{C}\|_F$, the proof of (A3) is exactly same as in [21, Pg. 235] or alternatively [24, Theorem 2.10].

Proof of Proposition 38. Since $\pi(\mathbf{S}^*)$ is non-degenerate, \widetilde{F} being Morse-Bott at $\pi(\mathbf{S}^*)$ follows from [28, Definition 6.5]. Since non-degenerate critical points of a smooth function are isolated from other critical points, it follows again from [28] that \widetilde{F} is Morse-Bott in a sufficiently small neighborhood of $\pi(\mathbf{S}^*)$.

Then, since the action of $\mathbb{O}(d)$ on $\mathbb{O}(d)^m$ is free and the function F is invariant under the action of $\mathbb{O}(d)$ ($\mathbf{SQ} = \mathbf{S} \iff \mathbf{Q} = \mathbf{I}_d$ and $F(\mathbf{SQ}) = F(\mathbf{S})$ for all $\mathbf{S} \in \mathbb{O}(d)^m, \mathbf{Q} \in \mathbb{O}(d)$), the result follows from [5, Section 15.2].