# Random Inverse Problems Over Graphs: Decentralized Online Learning

Tao Li and Xiwei Zhang

## Abstract

We establish a framework of distributed random inverse problems over network graphs with online measurements, and propose a decentralized online learning algorithm. This unifies the distributed parameter estimation in Hilbert spaces and the least mean square problem in reproducing kernel Hilbert spaces (RKHS-LMS). We transform the convergence of the algorithm into the asymptotic stability of a class of inhomogeneous random difference equations in Hilbert spaces with  $L_2$ -bounded martingale difference terms and develop the  $L_2$ -asymptotic stability theory in Hilbert spaces. It is shown that if the network graph is connected and the sequence of forward operators satisfies the *infinite-dimensional spatio-temporal persistence of excitation* condition, then the estimates of all nodes are mean square and almost surely strongly consistent. Moreover, we propose a decentralized online learning algorithm in RKHS based on non-stationary and non-independent online data streams, and prove that the algorithm is mean square and almost surely strongly consistent if the operators induced by the random input data satisfy the *infinite-dimensional spatio-temporal persistence of excitation condition* condition.

## **Index Terms**

Decentralized online learning, random inverse problem, reproducing kernel Hilbert space, randomly time-varying difference equation, persistence of excitation.

This work was supported in part by the National Natural Science Foundation of China under Grant 62261136550, and in part by the Basic Research Project of Shanghai Science and Technology Commission under Grant 20JC1414000. (*Corresponding author: Tao Li.*)

Tao Li is with the Key Laboratory of Pure Mathematical Practice, School of Mathematical Sciences, East China Normal University, Shanghai 200241, China (e-mail: tli@math.ecnu.edu.cn).

Xiwei Zhang is with the School of Mathematical Sciences, East China Normal University, Shanghai 200241, China (e-mail: xwzhangmath@sina.com).

# 1. INTRODUCTION

Inverse problems have wide applications such as medical imaging, geophysics and oil exploration ([1–6]). An inverse problem is to determine the system input (cause) from the measurable system output (result). In reality, measurements are usually affected by external disturbances, and the inverse problems with noisy measurements have been widely studied including the cases with deterministic noises ([7–10]) and those with Gaussian white noises ([11–15]). It is of practical significance to consider inverse problems with both randomly time-varying forward operators and random measurement noises. For example, consider the online learning problem in RKHS. Let  $\mathscr{X} \subseteq \mathbb{R}^n$  be the input space and  $(\mathscr{H}_K, \langle \cdot, \cdot \rangle_K)$  be the Hilbert space with Mercer kernel  $K : \mathscr{X} \times \mathscr{X} \to \mathbb{R}$ . At time instant k, the random (with unknown distribution) input data  $x(k) \in \mathscr{X}$  and the output data  $y(k) \in \mathbb{R}$  satisfy the measurement equation  $y(k) = f_0(x(k)) + v(k), k \ge 0$ , where  $f_0 \in \mathscr{H}_K$  is the unknown function, and  $v(k) \in \mathbb{R}$  is the random measurement noise ([16, 17]). The online learning problem in  $\mathscr{H}_K$  is to reconstruct  $f_0$ based on the online data stream  $\{(x(k), y(k))\}_{k=0}^{\infty}$ . By the reproducing property of RKHS ([18]), the above measurement equation can be written as

$$y(k) = H(k)f_0 + v(k), \ k \ge 0, \tag{1.1}$$

where H(k) is the randomly time-varying forward operator induced by the input data x(k), satisfying  $H(k)f := \langle f, K(x(k), \cdot) \rangle_K$ ,  $\forall f \in \mathscr{H}_K$ . Thus, an online learning problem in RKHS ([19–22]) comes down to an inverse problem associated with the measurement model (1.1). Most of the existing works on statistical inverse problems ([11–15, 23–30]) assumed the forward operators to be deterministic and time-invariant, which can not cover the measurement model (1.1).

In addition to online data streams, many practical problems are required to be solved in a decentralized or distributed information structure. The overall large amounts of data are usually divided into several data sets, and the learning process are performed with multiple parallel processors ([31]). For example, in the distributed multi-area state estimation of the power grid ([32]), all buses of the grid are divided into several non-overlapping regions. The communication structure among regions are represented by a graph  $\mathcal{G} = \{\mathcal{V}, \mathcal{E}_{\mathcal{G}}\}$ , where the node set  $\mathcal{V}$  represents the set of regions and the edge set  $\mathcal{E}_{\mathcal{G}}$  represents the set of inter-regional communication channels. The state of the grid  $f_0$  to be estimated consists of the voltage magnitudes and phase angles of all buses,  $y_i(k)$  is the real-time measurement in the region  $i \in \mathcal{V}$ , containing the active and

reactive currents, the bus injected power and voltage magnitude information, and  $v_i(k)$  is the random measurement noise. The problem of distributed multi-region power grid state estimation is formulated as a random inverse problem over the graph  $\mathcal{G}$  with the measurement equations

$$y_i(k) = H_i(k)f_0 + v_i(k), \ k \ge 0, \ i \in \mathcal{V},$$
(1.2)

where the observation matrix  $H_i(k) = \delta_i(k)H_i$ . Here,  $\{\delta_i(k), k \ge 0\}$  is a Bernoulli sequence representing the possible random data losses and  $H_i$  is the observation matrix without data loss.

With the development of online learning in recent years ([33]), decentralized online learning algorithms for finite-dimensional parameter estimation have been widely studied. Pioneering works on the decentralized online parameter estimation in finite-dimensional spaces were achieved in [34, 35] and fruitful results were obtained in [34-45]. Specifically, the decentralized online learning algorithms with randomly spatio-temporal independent observation matrices were proposed in [34, 36–39] via the collaborative strategy of diffusion. Kar and Moura [35] established a distributed observability condition (global observability plus mean connectedness) with temporally independent observation sequences, under which the distributed estimates are consistent and asymptotically normal. Kar and Moura [40] considered decentralized estimation with i.i.d. graph sequences and showed that the algorithm achieves weak consensus under a weak distributed detectability condition. Kar et al. [41, 42] proposed the decentralized parameter estimation algorithms based on consensus plus innovation with measurement and communication noises. It is required that the expectations of the regression matrices be known, and the graphs, regression matrices, measurement and communication noises be spatially i.i.d. Piggott et al. [43, 44] proposed decentralized algorithms over fixed communication graphs with time-dependent observation matrices. The convergence of decentralized online learning algorithms were analyzed by Ishihara and Alghunaim [45] for the case with spatial independent observation matrices.

Besides the finite dimensional parameter estimation, infinite-dimensional supervised online learning in RKHS is another important topic of random inverse problems ([46]). Based on the systematic study of batch learning in [18], rich results of online learning algorithms based on i.i.d. online data streams were obtained in [19–21, 47, 48]. Ying and Pontil [19] considered the least-square online gradient descent algorithm in RKHS and presented a novel capacity independent approach to derive error bounds and convergence results for this algorithm. Tarrès and Yao [20] showed that the online regularized algorithm can achieve the strong convergence rate of batch learning, and the weak convergence rate is optimal in the sense that it reaches the minimax and individual lower rates. Dieuleveut and Bach [21] considered the random-design least-squares regression problem within the RKHS framework, and showed that the averaged unregularized LMS algorithm with a given sufficient large step-size can attain optimal rates of convergence for a variety of regimes for the smoothnesses of the optimal prediction function in RKHS. Shin *et al.* [47] proposed decentralized adaptive learning algorithms over graphs in RKHS in a deterministic framework. Deng *et al.* [48] used the multiplicative operator in the saddle point problem to carve out the communication structure of decentralized networks and proposed a distributed consensus-based online learning algorithm with i.i.d. measurements. The nonlinear online learning problems in RKHS with spatio-temporal independent measurements were studied in [49–51].

Up to now, the existing theory of inverse problems in statistical and stochastic frameworks are far from mature and the existing related results may be divided into three categories: (i) statistical inverse problems based on deterministic time-invariant compact forward operators in Hilbert spaces ([25, 26, 52]); (ii) distributed parameter estimation in finite-dimensional spaces ([32, 34, 36–39, 43–45]); (iii) decentralized learning based on stationary e.g., i.i.d. measurements in RKHS ([47–51]). Some basic problems are still open, such as

- inverse problems with randomly time-varying forward operators;
- to establish a unified framework for random inverse problems in infinite-dimensional Hilbert spaces, distributed parameter estimation problems in finite-dimensional spaces and online learning problems in reproducing kernel Hilbert spaces;
- to develop a decentralized RKHS learning theory based on non-stationary and non-independent data streams.

Motivated by the above problems, we consider a class of random inverse problems over graphs, establish a unified framework to deal with the above three types of problems, and propose a decentralized online learning algorithm in Hilbert spaces. The learning algorithm of each node in the network consists of an innovation term and a consensus term. The innovation term is to update the node' estimate by using the node's own measurement data, and the consensus term is a weighted sum of its own estimate and the estimates of its neighboring nodes. The forward operator of the measurement of each node is randomly time-varying and is not required to satisfy special statistical properties, such as temporal independence (the forward operator of each node over the graph is independent with respect to time), spatial independence (the forward

operators of different nodes are independent of each other at each moment) and stationarity, and the random measurement noise is no longer restricted to Gaussian white noises. These general settings bring essential difficulties to the convergence analysis of the algorithm. Tarrès and Yao [20], Smale and Yao [53] transformed the online learning problem with i.i.d. data streams in RKHS into the inverse problem with the deterministic time-invariant Hilbert-Schmidt forward operator, and then obtained the convergence result by using the singular value decomposition (SVD) of the linear compact operators in the Hilbert space. By using the i.i.d. properties of the data, Dieuleveut and Bach [21] transformed the estimation error equation into the random difference equation equivalently, where the homogeneous part is deterministically time-invariant and the inhomogeneous part is the martingale difference sequence in the Hilbert space, from which the mean-square convergence of the algorithm were obtained by means of the spectral decomposition property of the compact operator. Note that the SVD of linear compact operators in Hilbert spaces are only applicable for the inverse problems with deterministic and time-invariant linear compact forward operators. The existing methodologies in [20, 21, 23, 25, 26, 28, 52–54] are no longer applicable for our problems.

Note that for decentralized online learning in finite-dimensional spaces over random graphs based on non-stationary data, we have established the stochastic spatio-temporal persistence of excitation (SSTPE) condition to ensure convergence of the algorithm [32, 55]. However, for this kind of finite-dimensional excitation conditions, the information matrices are all required to be positive definite, i.e., the eigenvalues of the matrix have strictly positive lower bounds. Obviously, the SSTPE condition is not applicable for inverse problems in infinite-dimensional Hilbert spaces. It is known that even for a strictly positive compact operator, the infimum of its eigenvalues is zero, and then the SSTPE condition can not hold.

To this end, firstly, we explore the properties concerning the measurability, conditional expectation and conditional independence of random elements with values in topological spaces by means of measurability and integration theory of mappings with values in Banach spaces, spectral decomposition theory of bounded self-adjoint operators, and martingale convergence methods. Then, we propose the  $L_p^q$ -stability condition on the sequence of operator-valued random elements. We establish the theory of a class of  $L_2$ -asymptotic stability of infinite-dimensional inhomogeneous random difference equations with  $L_2$ -bounded martingale difference terms. Under the above theoretical framework, we give a sufficient condition for a class of operator-valued random element sequences, which are composed of random forward operators and Laplacian matrices of graphs, to be  $L_2^2$ -stable, i.e., the *infinite-dimensional spatio-temporal persistence of excitation* condition, and establish the convergence condition for decentralized algorithms of random inverse problems over network graphs. Finally, we develop a theory of decentralized learning in RKHS with non-stationary and non-independent data streams. Compared with the existing literature, our contributions are summarized as follows.

- The proposed algorithm involves the sequences of random forward operators induced by random input data. To develop a completely self-contained theoretical framework, we regard both vector-valued mappings and operator-valued mappings as random elements with values in topological spaces. Based on the analytical theory of Banach spaces, we further explore the probability structure of random elements with values in different topological spaces, including (i) the measurable structure of random elements taking values in a strong operator topological space; (ii) the properties of expectations and conditional expectations of random elements with values in a uniform operator topological space or a strong operator topological space; (iii) the properties of independence and conditional independence of random elements with values in a topological space.
- We weaken the constraints on the forward operators compared with the most of the existing studies on inverse problems ([23, 24, 26, 27]). We consider the cases with general bounded linear forward operators instead of compact ones. Besides, we allow the forward operators to be randomly time-varying, which are not restricted to be deterministic and time-invariant.
- We transform the convergence analysis of the algorithm into the  $L_2$ -asymptotic stability of time-varying random difference equations in Hilbert spaces. Since the forward operators of inverse problems in infinite-dimensional Hilbert spaces usually do not have bounded inverses, the existing asymptotic stability theory on infinite-dimensional random difference equations with compressive operators in Hilbert spaces ([56–61]) cannot be applied to inverse problems. To this end, we propose the  $L_p^q$ -stability condition on the sequence of the products of operator-valued random elements, and establish the  $L_2$ -asymptotic stability theory for a class of inhomogeneous random difference equations in Hilbert spaces with  $L_2$ -bounded martingale difference terms. We give sufficient conditions on the stability of a class of operator-valued random sequences composed of forward operators and Laplacian matrices of graphs. We prove that if the graph is connected, and the sequence of forward operators satisfy the *infinite-dimensional spatio-temporal persistence of excitation* condition,

then all nodes' estimates are all mean square and almost surely strongly consistent with the unknown function of the inverse problem.

• We develop a theory of decentralized online learning in RKHS. Almost the existing literature on online supervised learning in RKHS (e.g., [47, 51, 62–64]) are based on i.i.d. data, while we propose a decentralized online learning algorithm based on non-stationary and non-independent data streams in RKHS. We establish the convergence condition by equivalently transforming the distributed learning problem in RKHS into the random inverse problems over graphs. Especially, if the graph has only one node and the random input data is i.i.d., then our algorithm degenerates to the centralized online learning algorithm without regularization parameters in [19, 21].

The following notations will be used throughout the paper. For any given sub- $\sigma$ -algebra  $\mathcal{F}_0 \subseteq \mathcal{F}$ , denote  $f \in \mathcal{F}_0$  if f is measurable with respect to  $\mathcal{F}_0$ . For any given sets  $\{A_i, i \in \mathscr{I}\}$ , where  $\mathscr{I}$  is a set of indices, denote  $\sigma \left(\bigcup_{i \in \mathscr{I}} A_i\right)$  by  $\bigvee_{i \in \mathscr{I}} A_i$ . Let  $(\mathscr{X}_i, \langle \cdot, \cdot \rangle_{\mathscr{X}_i})$  be a Hilbert space, where the norm induced by the inner product is defined by  $||x_i||_{\mathscr{X}_i} := \sqrt{\langle x_i, x_i \rangle_{\mathscr{X}_i}}, x_i \in \mathscr{X}_i, i = 1, \dots, n$ . The Hilbert direct sum space is denoted by

$$\bigoplus_{i=1} \mathscr{X}_i = \left\{ x = (x_1, \cdots, x_n) : x_i \in \mathscr{X}_i, 1 \le i \le n \right\},\$$

where the inner product is defined by  $\langle x, y \rangle_{\bigoplus_{i=1}^{n} \mathscr{X}_{i}} := \sum_{i=1}^{n} \langle x_{i}, y_{i} \rangle_{\mathscr{X}_{i}}, \forall x = (x_{1}, \cdots, x_{n}), y = (y_{1}, \cdots, y_{n}) \in \bigoplus_{i=1}^{n} \mathscr{X}_{i}$ . Denote  $\bigoplus_{i=1}^{n} \mathscr{X} := \mathscr{X}^{n}$ . Let  $\mathscr{X}$  and  $\mathscr{Y}$  be Hilbert spaces. Denote the Kronecker product of the vector  $\mathbf{1}_{n} \in \mathbb{R}^{n}$  and  $\phi \in \mathscr{X}$  by  $\mathbf{1}_{n} \otimes f := (f, \cdots, f) \in \mathscr{X}^{n}$  and the Kronecker product of the matrix  $A \in \mathbb{R}^{n \times m}$  and  $B \in \mathscr{L}(\mathscr{X}, \mathscr{Y})$  by the matrix of operators

$$A \otimes B := \begin{pmatrix} a_{11}B & \cdots & a_{1m}B \\ \vdots & \ddots & \vdots \\ a_{n1}B & \cdots & a_{nm}B \end{pmatrix} \in \mathscr{L}(\mathscr{X}^m, \mathscr{Y}^n)$$

The operations and properties of operator matrices in Hilbert direct sum spaces can be found in [65]. Let  $\mathscr{X}$  be a Hilbert space and  $T \in \mathscr{L}(\mathscr{X})$ . If  $T^* = T$ , i.e.,  $\langle Tx, y \rangle_{\mathscr{X}} = \langle x, Ty \rangle_{\mathscr{X}}$ ,  $\forall x, y \in \mathscr{X}$ , then T is called the self-adjoint operator. If the self-adjoint operator  $T \in \mathscr{L}(\mathscr{X})$ satisfies  $\langle Tx, x \rangle_{\mathscr{X}} \ge 0$ ,  $\forall x \in \mathscr{X}$ , then T is said to be positive self-adjoint, and is denoted by  $T \ge 0$ , and in particular, if  $\langle Tx, x \rangle_{\mathscr{X}} > 0$  for any given non-zero element x in  $\mathscr{X}$ , then T is said to be strictly positive self-adjoint and is denoted by T > 0.

Let  $\mathcal{G} = \{\mathcal{V}, \mathcal{E}_{\mathcal{G}}, \mathcal{A}_{\mathcal{G}}\}$  denote a weighted graph, where  $\mathcal{V} = \{1, ..., N\}$  is the set of nodes and  $\mathcal{E}_{\mathcal{G}}$  is the set of edges. The unordered pair  $(j, i) \in \mathcal{E}_{\mathcal{G}}$  if and only if there exists an edge between

nodes j and i. Denote the set of neighboring nodes of node i by  $\mathcal{N}_i = \{j \in \mathcal{V} : (j,i) \in \mathcal{E}_{\mathcal{G}}\}$ . The matrix  $\mathcal{A}_{\mathcal{G}} = [a_{ij}] \in \mathbb{R}^{N \times N}$  is called the weighted adjacency matrix of  $\mathcal{G}$ , and for any given  $i, j \in \mathcal{V}, a_{ii} = 0$  and  $a_{ij} = a_{ji} > 0$  if and only if  $j \in \mathcal{N}_i$ . The Laplacian matrix of  $\mathcal{G}$  is defined by  $\mathcal{L}_{\mathcal{G}} = \mathcal{D}_{\mathcal{G}} - \mathcal{A}_{\mathcal{G}}$ , where the degree matrix  $\mathcal{D}_{\mathcal{G}} = \text{diag}\{\sum_{j=1}^{N} a_{1j}, \sum_{j=1}^{N} a_{2j}, \cdots, \sum_{j=1}^{N} a_{Nj}\}$ .

#### 2. THEORETICAL FRAMEWORK OF RANDOM ELEMENTS WITH VALUES IN A TOPOLOGICAL

#### SPACE

The proposed algorithm involves the sequences of random forward operators induced by random input data. Conventionally, a random element with values in a Banach space is required to be strongly measurable, which is almost separably valued with respect to the topology induced by the norm in the Banach space ([66]). It is known that operator-valued mappings may not be strongly measurable since the Banach space of bounded linear operators is generally non-separable with respect to the uniform operator topology (the topology induced by the operator norm) ([67]). In this section, we develop a self-contained theoretical framework of random elements with values in a topological space.

# 2.1 Random elements with values in a topological space

We first introduce the following definitions.

**Definition 2.1.** Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a complete probability space, and  $(\mathcal{U}, \tau)$  a topological space. Given the mapping  $f : \Omega \to \mathcal{U}$ , if there exists a separable closed subset  $\mathcal{U}_0$  of  $\mathcal{U}$  and a subset  $\Omega_0$  of  $\Omega$  with probability measure 1, such that

$$f(\Omega_0) := \{ f(x) : x \in \Omega_0 \} \subseteq \mathscr{U}_0,$$

then f is called almost separably valued with respect to  $\tau$ .

**Definition 2.2.** Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a complete probability space and  $(\mathcal{U}, \mathcal{B}(\mathcal{U}; \tau))$  be a measurable space, where  $\tau$  is the topology on  $\mathcal{U}$ , and  $\mathcal{B}(\mathcal{U}; \tau)$  is the Borel  $\sigma$ -algebra of the topological space  $(\mathcal{U}, \tau)$ , i.e., the smallest  $\sigma$ -algebra containing all open sets in  $\mathcal{U}$ . A mapping  $f: \Omega \to \mathcal{U}$  is said to be a random element with values in the topological space  $(\mathcal{U}, \tau)$  if it is  $\mathcal{F}/\mathcal{B}(\mathcal{U}; \tau)$ -measurable and almost separably valued with respect to  $\tau$ .

**Definition 2.3.** If f is a random element with values in the topological space  $(\mathcal{U}, \tau)$ , then the distribution of f is defined by the Borel probability measure  $\mu_f(B) := \mathbb{P}(f^{-1}(B))$  on  $(\mathcal{U}, \tau)$ ,  $\forall B \in \mathscr{B}(\mathcal{U}; \tau)$ .

Let  $\tau_{N}(\mathscr{X})$  be the topology induced by the norm  $\|\cdot\|_{\mathscr{X}}$  in a Banach space  $\mathscr{X}$ . It follows from Lemma A.4 that the mapping  $f: \Omega \to \mathscr{X}$  is strongly measurable with respect to  $\tau_{N}(\mathscr{X})$  if and only if f is a random element with values in the Banach space  $(\mathscr{X}, \tau_{N}(\mathscr{X}))$ . Especially, if  $\mathscr{X}$  is separable with respect to  $\tau_{N}(\mathscr{X})$ , then any  $\mathcal{F}/\mathscr{B}(\mathscr{X}; \tau_{N}(\mathscr{X}))$ -measurable mapping  $f: \Omega \to \mathscr{X}$ is a random element with values in  $(\mathscr{X}, \tau_{N}(\mathscr{X}))$ .

Let  $\mathscr{L}(\mathscr{Y}, \mathscr{Z})$  be a linear space consisting of all bounded linear operators mapping from the Banach space  $\mathscr{Y}$  to the Banach space  $\mathscr{Z}$ , in particular,  $\mathscr{L}(\mathscr{Z}) := \mathscr{L}(\mathscr{Z}, \mathscr{Z})$ . Let  $\tau_{\mathsf{S}}(\mathscr{L}(\mathscr{Y}, \mathscr{Z}))$  be the strong operator topology of  $\mathscr{L}(\mathscr{Y}, \mathscr{Z})$ .

For a random element  $f : \Omega \to \mathscr{X}$  with values in  $(\mathscr{X}, \tau(\mathscr{X}))$ , we denote the  $\sigma$ -algebra generated by f in the sense of the topology  $\tau(\mathscr{X})$  by

$$\sigma(f;\tau(\mathscr{X})) := \left\{ f^{-1}(B) : B \in \mathscr{B}(\mathscr{X};\tau(\mathscr{X})) \right\}.$$

Based on the above definitions, we have the following proposition, whose proof is in Appendix B.

**Proposition 2.4.** If  $f : \Omega \to \mathscr{X}$  is a random element with values in the Banach space  $(\mathscr{X}, \tau_N(\mathscr{X}))$ , and  $T : \Omega \to \mathscr{L}(\mathscr{X}, \mathscr{Y})$  is a random element with values in the topological space  $(\mathscr{L}(\mathscr{X}, \mathscr{Y}),$ 

 $\tau_{\mathcal{S}}(\mathscr{L}(\mathscr{X},\mathscr{Y})))$ , then  $Tf:\omega\mapsto T(\omega)f(\omega)$  satisfies

$$Tf \in \left(\bigvee_{x \in \mathscr{X}} \sigma(Tx; \tau_N(\mathscr{Y}))\right) \bigvee \sigma(f; \tau_N(\mathscr{X})).$$

## 2.2 Conditional expectation

As for mathematical expectations and conditional expectations of the random elements with values in the Banach space  $(\mathscr{X}, \tau_{N}(\mathscr{X}))$ , we introduce the following notations. Let  $L^{0}(\Omega; \mathscr{X})$ be a linear space composed of all mappings which take values in  $(\mathscr{X}, \tau_{N}(\mathscr{X}))$  and are strongly measurable with respect to  $\tau_{N}(\mathscr{X})$ . Let  $L^{0}(\Omega, \mathscr{G}; \mathscr{X})$  be the linear space composed of all  $\mathscr{G}/\mathscr{B}(\mathscr{X}; \tau_{N}(\mathscr{X}))$ -measurable functions in  $L^{0}(\Omega; \mathscr{X})$ , where  $\mathscr{G}$  is a sub- $\sigma$ -algebra of  $\mathcal{F}$ . Let  $L^{p}(\Omega; \mathscr{X})$  be the Bochner space composed of all mappings which are strongly measurable with respect to  $\tau_{\mathbf{N}}(\mathscr{X})$ . Define  $\|f\|_{L^{p}(\Omega;\mathscr{X})} := \left(\int_{\Omega} \|f\|_{\mathscr{X}}^{p} d\mathbb{P}\right)^{\frac{1}{p}} < \infty, 1 \leq p < \infty$ . Denote  $L^{p}(\Omega) := L^{p}(\Omega; \mathbb{R})$ .

By Lemma A.9, it follows that there exists a unique conditional expectation  $\mathbb{E}[f|\mathscr{G}] \in L^0(\Omega, \mathscr{G}; \mathscr{X})$  of the Bochner integrable random element f with values in  $(\mathscr{X}, \tau_N(\mathscr{X}))$ , and  $\mathbb{E}[f|\mathscr{G}]$  is a random element with values in  $(\mathscr{X}, \tau_N(\mathscr{X}))$ . We have the following propositions of conditional expectations, whose proofs are in Appendix B.

**Proposition 2.5.** If  $f \in L^1(\Omega; \mathscr{L}(\mathscr{Y}, \mathscr{Z}))$  is a random element with values in Banach space  $(\mathscr{L}(\mathscr{Y}, \mathscr{Z}), \tau_N(\mathscr{L}(\mathscr{Y}, \mathscr{Z})))$ , then  $fy \in L^1(\Omega; \mathscr{Z})$  is the random element with values in Banach space  $(\mathscr{Z}, \tau_N(\mathscr{Z}))$ , and  $\mathbb{E}[fy] = \mathbb{E}[f]y, \forall y \in \mathscr{Y}$ .

**Proposition 2.6.** If  $f \in L^2(\Omega; \mathscr{L}(\mathscr{Y}, \mathscr{Z}))$  is a random element with values in the Banach space  $(\mathscr{L}(\mathscr{Y}, \mathscr{Z}), \tau_N(\mathscr{L}(\mathscr{Y}, \mathscr{Z})))$ , and  $y \in L^2(\Omega, \mathscr{G}; \mathscr{Y})$  is a random element with values in the Banach space  $(\mathscr{Y}, \tau_N(\mathscr{Y}))$ , where  $\mathscr{G}$  is a sub- $\sigma$ -algebra of  $\mathcal{F}$ , then  $fy \in L^1(\Omega; \mathscr{Z})$  is a random element with values in the Banach space  $(\mathscr{Z}, \tau_N(\mathscr{Z}))$  and  $\mathbb{E}[fy|\mathscr{G}] = \mathbb{E}[f|\mathscr{G}]y$  a.s.

## 2.3 Independence and conditional independence

In terms of the independence and conditional independence of random elements with values in a topological space, we have the following definitions.

**Definition 2.7.** Let  $\mathscr{I}$  be a set of indices,  $f_j$ ,  $j \in \mathscr{I}$  be the random elements with values in  $(\mathscr{X}_j, \tau(\mathscr{X}_j))$ . If  $\mathbb{P}(f_{\alpha_1} \in B_1, \dots, f_{\alpha_n} \in B_n) = \prod_{j=1}^n \mathbb{P}(f_{\alpha_j} \in B_j)$ , for arbitrarily different indices  $\alpha_1, \dots, \alpha_n \in \mathscr{I}$  and arbitrary  $B_1 \in \mathscr{B}(\mathscr{X}_1; \tau(\mathscr{X}_{\alpha_1})), \dots, B_n \in \mathscr{B}(\mathscr{X}_n; \tau(\mathscr{X}_{\alpha_n}))$ , then  $f_j$ ,  $j \in \mathscr{I}$  are said to be mutually independent.

**Definition 2.8.** Let  $\mathcal{F}_i$ ,  $i \in \mathscr{I}$  and  $\mathscr{G}$  be sub- $\sigma$ -algebra of  $\mathcal{F}$ . If  $\mathbb{P}\left(\bigcap_{j=1}^n A_j \middle| \mathscr{G}\right) = \prod_{j=1}^n \mathbb{P}(A_j \middle| \mathscr{G})$ a.s., for arbitrarily different indices  $\alpha_1, \dots, \alpha_n \in \mathscr{I}$  and arbitrary  $A_1 \in \mathcal{F}_{\alpha_1}, \dots, A_n \in \mathcal{F}_{\alpha_n}$ , then it is said that  $\mathcal{F}_{\alpha}$  is conditionally independent w.r.t.  $\mathscr{G}$ ,  $\alpha \in \mathscr{I}$ . If  $\sigma(f_i; \tau(\mathscr{X}_i))$  is conditionally independent w.r.t.  $\mathscr{G}$ ,  $i \in \mathscr{I}$ , then we say that the random elements  $f_i$ ,  $i \in \mathscr{I}$  with values in  $(\mathscr{X}_i, \tau(\mathscr{X}_i))$  are conditionally independent w.r.t.  $\mathscr{G}$ .

By Definitions 2.7 and 2.8, we have the following propositions, the proofs of which are in Appendix B.

**Proposition 2.9.** If the family of random elements  $\{f_k, k \ge 1\}$  with values in  $(\mathscr{X}_1, \tau(\mathscr{X}_1))$ and the family of random elements  $\{g_k, k \ge 1\}$  with values in  $(\mathscr{X}_2, \tau(\mathscr{X}_2))$  are independent, then  $\bigvee_{i=k}^{\infty} \sigma(f_i; \tau(\mathscr{X}_1))$  is conditionally independent of  $\bigvee_{i=k}^{\infty} \sigma(g_i; \tau(\mathscr{X}_2))$  with respect to  $\bigvee_{i=1}^{k-1} (\sigma(f_i; \tau(\mathscr{X}_1)) \bigvee \sigma(g_i; \tau(\mathscr{X}_2))), \forall k \ge 2.$ 

**Proposition 2.10.** Let  $\mathscr{G}$  be the sub- $\sigma$ -algebra of  $\mathcal{F}$ . The random element  $T : \Omega \to \mathscr{L}(\mathscr{X}, \mathscr{Y})$ with values in  $(\mathscr{L}(\mathscr{X}, \mathscr{Y}), \tau_{\mathcal{S}}(\mathscr{L}(\mathscr{X}, \mathscr{Y})))$  satisfies  $\mathbb{E}[||T||^{2}_{\mathscr{L}(\mathscr{X}, \mathscr{Y})}] < \infty$  and  $f \in L^{2}(\Omega; \mathscr{X})$  is a random element with values in the Banach space  $(\mathscr{X}, \tau_{N}(\mathscr{X}))$ . If T is conditionally independent of f w.r.t.  $\mathscr{G}$ , then  $\mathbb{E}[Tf|\mathscr{G}] = \mathbb{E}[T\mathbb{E}[f|\mathscr{G}]|\mathscr{G}]$  a.s.

# 3. ONLINE LEARNING FOR RANDOM INVERSE PROBLEMS OVER GRAPHS

#### 3.1 Online random inverse problems over graphs

Consider a distributed communication network modeled by a weighted graph  $\mathcal{G} = \{\mathcal{V}, \mathcal{E}_{\mathcal{G}}, \mathcal{A}_{\mathcal{G}}\}$ consisting of N nodes. The measurement  $y_i(k)$  of node i at instant k is given by

$$y_i(k) = H_i(k)f_0 + v_i(k), \ k = 0, 1, 2, \dots \ i = 1, \dots, N,$$
(3.1)

where  $f_0 \in \mathscr{X}$  to be estimated is an unknown element in the Hilbert space  $\mathscr{X}$ , the random forward operator  $H_i(k) : \Omega \to \mathscr{L}(\mathscr{X}, \mathscr{Y}_i)$  is an operator-valued random element with values in  $(\mathscr{L}(\mathscr{X}, \mathscr{Y}_i), \tau_{\mathsf{S}}(\mathscr{L}(\mathscr{X}, \mathscr{Y}_i)))$ , and the measurement noise  $v_i(k) : \Omega \to \mathscr{Y}_i$  is a random element with values in the Hilbert space  $(\mathscr{Y}_i, \tau_{\mathsf{N}}(\mathscr{Y}_i))$ .

**Remark 3.1.** For the finite-dimensional Euclidean space  $\mathscr{X} = \mathbb{R}^n$ , the measurement model (3.1) has been widely studied in [34–42], where the forward operator  $H_i(k)$  degenerates to the random observation matrix and  $f_0 \in \mathbb{R}^n$  degenerates to the unknown finite-dimensional parameter vector. Kar et al. [41] investigated the decentralized estimation algorithms with nonlinear measurement models, where they introduced separably estimable measurement models that generalize the observability condition in linear centralized estimation to nonlinear decentralized estimation. It is worth pointing out that even for the nonlinear measurement model in [41], the unknown quantity to be estimated is a parameter vector in a finite-dimensional space, different from which, the unknown quantity  $f_0$  in the measurement model (3.1) can be a nonlinear function, which is an element in the infinite-dimensional Hilbert space  $\mathscr{X}$ . The measurement model (3.1) is essentially different from that in [41]. This will be further clarified in Remark 5.1.

Denote  $y(k) = (y_1(k), \dots, y_N(k)) : \Omega \to \bigoplus_{i=1}^N \mathscr{Y}_i, v(k) = (v_1(k), \dots, v_N(k)) : \Omega \to \bigoplus_{i=1}^N \mathscr{Y}_i \text{ and } H(k) = (H_1(k), \dots, H_N(k)) : \Omega \to \mathscr{L}(\mathscr{X}, \bigoplus_{i=1}^N \mathscr{Y}_i).$  We can write (3.1) in the compact form

$$y(k) = H(k)f_0 + v(k), \ k \ge 0.$$
(3.2)

The online random inverse problem means reconstructing  $f_0$  by real-time random measurements  $\{y(k), k \ge 0\}.$ 

The measurement equation (3.2), which covers the existing models of inverse problems, is more general in the sense that the forward operator can be randomly time-varying. Besides, different from the existing literature, which has a centralized information structure, here, the reconstruction of  $f_0$  is constrained by the information structure of the graph, i.e., there is no centralized fusion center collecting the overall measurements y(k), and at each moment k, node i can only use its own observation  $y_i(k)$  and its neighbors' estimates  $f_j(k), j \in \mathcal{N}_i$  to give its next estimate  $f_i(k+1): \Omega \to \mathscr{X}$  for  $f_0$ , i.e.,

$$f_i(k) \in \left(\bigvee_{s=0,1,2,\dots,k-1} \sigma(y_i(s);\tau_{\mathsf{N}}(\mathscr{Y}_i))\right) \bigvee \left(\bigvee_{j\in\mathcal{N}_i\cup\{i\}} \sigma\left(f_j(k-1);\tau_{\mathsf{N}}(\mathscr{X})\right)\right), \ i\in\mathcal{V}.$$

Here, the problem of cooperatively estimating  $f_0$  by the nodes over the graph based on each node's local measurements, its own and neighbors' estimates is called the *random inverse problem over the graph*.

**Remark 3.2.** Most of the existing literature on inverse problems assume the forward operator to be a deterministic and time-invariant linear compact operator H, associated with the measurement equation

$$y = Hf_0 + v. ag{3.3}$$

Here, different from the existing literature, the forward operator in the measurement equation (3.2) is allowed to be random and time-varying. In the classical inverse problem, the noise v in the measurement equation (3.3) is modeled as a deterministic perturbation ([68]). In the statistical inverse problem, it is modeled as Gaussian white noise ([23, 24, 27]). Based on the stochastic gradient descent (SGD) algorithm, Lu and Mathe [28], Jahn and Jin [29] and Jin and Lu [30] obtained the centralized learning strategy by the random discrete sampling of the forward operator and minimizing the loss functional

$$\widetilde{J}(x) = \frac{1}{2} \mathbb{E}\left[ \|y - Hx\|^2 \right], \ \forall \ x \in \mathscr{X}.$$
(3.4)

13

Specifically, Jahn and Jin [29] and Jin and Lu [30] investigated the regularization property of SGD with a priori and a posteriori stopping rules, and Lu and Mathe [28] gave an upper bound on the estimation error of SGD with discrete level. Recently, Iglesias et al. [25] and Lu et al. [26] solved the statistical inverse problem based on real-time measurements  $\{y(k) :$  $y(k) = Hf_0 + v(k), k \ge 0\}$ , where  $\{v(k), k \ge 0\}$  is an i.i.d. Gaussian white noise sequence. The statistical inverse problems in [25, 26] are special cases of the online random inverse problem with random forward operators.

# 3.2 Decentralized online learning algorithm

Denote  $\mathcal{H}(k) = \text{diag}\{H_1(k), \dots, H_N(k)\}$ . Based on the loss functional (3.4), we consider minimizing the loss functional  $J : \mathscr{X}^N \to \mathbb{R}$  with the Laplacian regularization term given by

$$J(f) = \frac{1}{2} \left( \mathbb{E} \left[ \| y(k) - \mathcal{H}(k) f \|_{\bigoplus_{i=1}^{N} \mathscr{Y}_{i}}^{2} \right] + \left\langle \left( \mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}} \right) f, f \right\rangle_{\mathscr{X}^{N}} \right), \ \forall \ f \in \mathscr{X}^{N}$$

where  $I_{\mathscr{X}}$  is the identical operator on the Hilbert space  $\mathscr{X}$ . The loss functional J(f) consists of two terms: the mean-square estimation error term  $\mathbb{E}[||y(k) - \mathcal{H}(k)f||^2_{\bigoplus_{i=1}^N \mathscr{Y}_i}]$  and the Laplacian regularization term  $\langle (\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})f, f \rangle_{\mathscr{X}^N} = \frac{1}{2} \sum_{i=1}^N \sum_{j=1}^N a_{ij} ||f_i - f_j||^2_{\mathscr{X}}$ , where  $f_i \in \mathscr{X}$  is the *i*-th component of f.

Suppose  $\mathcal{H}(k) \in L^2(\Omega; \mathscr{L}(\mathscr{X}^N, \bigoplus_{i=1}^N \mathscr{Y}_i))$  and  $v(k) \in L^2(\Omega; \bigoplus_{i=1}^N \mathscr{Y}_i)$ . If the sequences  $\{\mathcal{H}(k), k \geq 0\}$  and  $\{v(k), k \geq 0\}$  are both i.i.d, then it follows from Definition 2.3 and Proposition A.5.(b) that  $\{y(k) - \mathcal{H}(k)f, k \geq 0\}$  is a sequence of i.i.d. random elements with values in the Hilbert space  $(\bigoplus_{i=1}^N \mathscr{Y}_i, \tau_N(\bigoplus_{i=1}^N \mathscr{Y}_i))$ . Noting that  $\mathcal{H}^*(k)\mathcal{H}(k) \in L^1(\Omega; \mathscr{L}(\mathscr{X}^N))$ , by Proposition 2.5, the gradient operator grad  $J: \mathscr{X}^N \to \mathscr{X}^N$  is given by

$$\begin{aligned} \operatorname{grad} J(f) &= \frac{1}{2} \operatorname{grad} \left( \mathbb{E} \left[ \left\| y(k) - \mathcal{H}(k) f \right\|_{\bigoplus_{i=1}^{N} \mathscr{Y}_{i}}^{2} \right] + \left\langle \left( \mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}} \right) f, f \right\rangle_{\mathscr{X}^{N}} \right) \\ &= \frac{1}{2} \operatorname{grad} \mathbb{E} \left[ \left\langle \mathcal{H}^{*}(k) \mathcal{H}(k) f, f \right\rangle_{\mathscr{X}^{N}} \right] - \operatorname{grad} \mathbb{E} \left[ \left\langle \mathcal{H}(k) f, y(k) \right\rangle_{\bigoplus_{i=1}^{N} \mathscr{Y}_{i}} \right] \\ &+ \frac{1}{2} \operatorname{grad} \left\langle \left( \mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}} \right) f, f \right\rangle_{\mathscr{X}^{N}} \\ &= \frac{1}{2} \operatorname{grad} \left\langle \mathbb{E} \left[ \mathcal{H}^{*}(k) \mathcal{H}(k) \right] f, f \right\rangle_{\mathscr{X}^{N}} - \operatorname{grad} \mathbb{E} \left[ \left\langle \mathcal{H}(k) f, y(k) \right\rangle_{\bigoplus_{i=1}^{N} \mathscr{Y}_{i}} \right] \\ &+ \frac{1}{2} \operatorname{grad} \left\langle \left( \mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}} \right) f, f \right\rangle_{\mathscr{X}^{N}} . \end{aligned}$$

Similarly, we have

$$\langle \mathbb{E}[\mathcal{H}^*(k)\mathcal{H}(k)]x, y \rangle_{\mathscr{X}^N} = \mathbb{E}[\langle \mathcal{H}^*(k)\mathcal{H}(k)x, y \rangle_{\mathscr{X}^N}]$$

$$= \mathbb{E}[\langle x, \mathcal{H}^*(k)\mathcal{H}(k)y\rangle_{\mathscr{X}^N}]$$
$$= \langle x, \mathbb{E}[\mathcal{H}^*(k)\mathcal{H}(k)]y\rangle_{\mathscr{X}^N}, \ \forall \ x, y \in \mathscr{X}^N$$

Therefore,  $\mathbb{E}[\mathcal{H}^*(k)\mathcal{H}(k)]: \mathscr{X}^N \to \mathscr{X}^N$  is a self-adjoint operator. Noting that the Laplacian matrix  $\mathcal{L}_{\mathcal{G}}$  is positive semi-definite, it follows that  $\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}$  is a self-adjoint operator. Then

grad 
$$\langle \mathbb{E}[\mathcal{H}^*(k)\mathcal{H}(k)]f, f \rangle_{\mathscr{X}^N} = 2\mathbb{E}[\mathcal{H}^*(k)\mathcal{H}(k)]f, \ \forall \ f \in \mathscr{X}^N,$$

and

grad 
$$\langle (\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}) f, f \rangle_{\mathscr{X}^N} = 2(\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}) f, \forall f \in \mathscr{X}^N.$$

Noting that  $\mathcal{H}^*(k)y(k) \in L^1(\Omega; \mathscr{X}^N)$ , it follows that

$$\begin{split} &\lim_{t\to 0} \frac{1}{t} \left( \mathbb{E} \left[ \langle \mathcal{H}(k)(f+tg), y(k) \rangle_{\bigoplus_{i=1}^{N} \mathscr{Y}_{i}} \right] - \mathbb{E} \left[ \langle \mathcal{H}(k)f, y(k) \rangle_{\bigoplus_{i=1}^{N} \mathscr{Y}_{i}} \right] \right) \\ &= \mathbb{E} [\langle \mathcal{H}^{*}(k)y(k), g \rangle_{\mathscr{X}^{N}}] \\ &= \langle \mathbb{E} [\mathcal{H}^{*}(k)y(k)], g \rangle_{\mathscr{X}^{N}}, \ \forall \ g \in \mathscr{X}^{N}. \end{split}$$

It follows that grad  $\mathbb{E}[\langle \mathcal{H}(k)f, y(k)\rangle_{\bigoplus_{i=1}^{N} \mathscr{Y}_{i}}] = \mathbb{E}[\mathcal{H}^{*}(k)y(k)]$ . Thus, we have

grad 
$$J(f) = -\mathbb{E}[\mathcal{H}^*(k)(y(k) - \mathcal{H}(k)f)] + (\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})f, \ \forall \ f \in \mathscr{X}^N.$$

Then we have the stochastic gradient descent (SGD) algorithm in the Hilbert space

$$f(k+1) = f(k) + a(k)\mathcal{H}^*(k)(y(k) - \mathcal{H}(k)f(k)) - b(k)(\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})f(k), \ k \ge 0.$$
(3.5)

Let  $f(k) = (f_1(k), \dots, f_N(k))$ . From (3.5), we obtain the decentralized online learning algorithm

$$f_{i}(k+1) = f_{i}(k) + a(k)H_{i}^{*}(k)(y_{i}(k) - H_{i}(k)f_{i}(k)) + b(k)\sum_{j\in\mathcal{N}_{i}}a_{ij}(f_{j}(k) - f_{i}(k)), \ k \ge 0, \ i \in \mathcal{V}.$$
(3.6)

**Remark 3.3.** In contrast to the centralized algorithm with a fusion center, the algorithm (3.6) gives a decentralized online learning strategy for each node, i.e., there is no fusion center collecting measurements and estimations of all nodes, and each node estimates  $f_0$  through its own measurements and its neighbors' estimations. Especially, for the measurement model (3.1) in finite-dimensional space, the corresponding learning strategy (3.6) is exactly the "consensus+innovations" type algorithm, which has been introduced by [41, 42]. Intuitively, node i obtains the innovation term  $y_i(k) - H_i(k)f_i(k)$  and the consensus term  $\sum_{j \in \mathcal{N}_i} a_{ij}(f_j(k) - f_i(k))$ , based on which the estimation  $f_i(k)$  is updated for the next instant.

# 4. CONVERGENCE ANALYSIS

Although the algorithm (3.5) is designed by assuming that  $\{\mathcal{H}(k) \in L^2(\Omega; \mathscr{L}(\mathscr{X}^N, \bigoplus_{i=1}^N \mathscr{Y}_i)), k \ge 0\}$  and  $\{v(k) \in L^2(\Omega; \bigoplus_{i=1}^N \mathscr{Y}_i), k \ge 0\}$  are both i.i.d. In fact, in this section, we will show that even for the non-independence and non-stationarity sequence of operator-valued random elements  $\{\mathcal{H}(k), k \ge 0\}$  and the noise sequence  $\{v(k), k \ge 0\}$ , the algorithm (3.6) still converge under mild conditions. The proofs of lemmas, theorems and corollaries of this section are in Appendix C.

Denote the global estimation error by  $e(k) = f(k) - \mathbf{1}_N \otimes f_0$ . Note that  $(\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})(\mathbf{1}_N \otimes f_0) = 0$ and  $\mathcal{H}(k)(\mathbf{1}_N \otimes f_0) = H(k)f_0$ . Subtracting  $\mathbf{1}_N \otimes f_0$  on both sides of equation (3.5) yields

$$e(k+1) = (I_{\mathscr{X}^{N}} - a(k)\mathcal{H}^{*}(k)\mathcal{H}(k) - b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})f(k) + a(k)\mathcal{H}^{*}(k)y(k) - \mathbf{1}_{N} \otimes f_{0}$$

$$= (I_{\mathscr{X}^{N}} - a(k)\mathcal{H}^{*}(k)\mathcal{H}(k) - b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})(f(k) - \mathbf{1}_{N} \otimes f_{0} + \mathbf{1}_{N} \otimes f_{0})$$

$$+ a(k)\mathcal{H}^{*}(k)y(k) - \mathbf{1}_{N} \otimes f_{0}$$

$$= (I_{\mathscr{X}^{N}} - a(k)\mathcal{H}^{*}(k)\mathcal{H}(k) - b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})e(k) - (a(k)\mathcal{H}^{*}(k)\mathcal{H}(k)$$

$$+ b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})(\mathbf{1}_{N} \otimes f_{0}) + a(k)\mathcal{H}^{*}(k)y(k)$$

$$= (I_{\mathscr{X}^{N}} - a(k)\mathcal{H}^{*}(k)\mathcal{H}(k) - b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})e(k)$$

$$+ a(k)\mathcal{H}^{*}(k)(y(k) - \mathcal{H}(k)(\mathbf{1}_{N} \otimes f_{0}))$$

$$= (I_{\mathscr{X}^{N}} - a(k)\mathcal{H}^{*}(k)\mathcal{H}(k) - b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})e(k) + a(k)\mathcal{H}^{*}(k)v(k).$$
(4.1)

The estimation error equation (4.1) belongs to the following family of randomly time-varying difference equations:

$$x(k+1) = (I_{\mathscr{X}_1} - F(k))x(k) + G(k)u(k), \ k \ge 0,$$
(4.2)

where u(k) is a random element with values in the Hilbert space  $(\mathscr{X}_2, \tau_N(\mathscr{X}_2)), F(k) : \Omega \to \mathscr{L}(\mathscr{X}_1)$  and  $G(k) : \Omega \to \mathscr{L}(\mathscr{X}_2, \mathscr{X}_1)$  are random elements with values in the topological spaces  $(\mathscr{L}(\mathscr{X}_1), \tau_S(\mathscr{L}(\mathscr{X}_1)))$  and  $(\mathscr{L}(\mathscr{X}_2, \mathscr{X}_1), \tau_S(\mathscr{L}(\mathscr{X}_2, \mathscr{X}_1)))$ , respectively. To analyze the convergence of the estimation error equation (4.1), we will first develop an asymptotic stability theory of the randomly time-varying difference equation (4.2).

# 4.1 Asymptotic stability of random difference equations in Hilbert spaces

To rigorously study the asymptotic stability of the random difference equations in the Hilbert space  $(\mathscr{X}, \tau_N(\mathscr{X}))$ , we introduce the following definitions.

**Definition 4.1.** If the sequence of random elements  $\{x(k), k \ge 0\}$  with values in the Hilbert space  $(\mathscr{X}, \tau_N(\mathscr{X}))$  satisfies  $\sup_{k\ge 0} \mathbb{E}[||x(k)||_{\mathscr{X}}^p] < \infty$ , where p > 0, then  $\{x(k), k \ge 0\}$  is said to be  $L_p$ -bounded.

**Definition 4.2.** If the sequence of random elements  $\{x(k), k \ge 0\}$  with values in the Hilbert space  $(\mathscr{X}, \tau_N(\mathscr{X}))$  satisfies  $\lim_{k\to\infty} \mathbb{E}[||x(k)||_{\mathscr{X}}^p] = 0$ , where p > 0, then  $\{x(k), k \ge 0\}$  is said to be  $L_p$ -asymptotically stable.

**Definition 4.3.** Let  $\{A(k), k \ge 0\}$  be a sequence of operator-valued random elements with values in  $(\mathscr{L}(\mathscr{X}), \tau_{S}(\mathscr{L}(\mathscr{X})))$  and  $\{\mathcal{F}(k), k \ge 0\}$  be a filter in  $(\Omega, \mathcal{F}, \mathbb{P})$ . If for any given  $L_{q}$ -bounded adaptive sequence  $\{x(k), \mathcal{F}(k), k \ge 0\}$  with values in the Hilbert space  $\mathscr{X}$ ,

$$\lim_{m \to \infty} \mathbb{E}\left[ \left\| \prod_{k=n+1}^{m} A(k) x(n) \right\|_{\mathscr{X}}^{p} \right] = 0, \ \forall \ n \ge 0, \ \text{where } p, \ q > 0,$$

then  $\{A(k), k \ge 0\}$  is said to be  $L_p^q$ -stable with respect to the filter  $\{\mathcal{F}(k), k \ge 0\}$ .

The strong convergence of the sequence of products of deterministic non-expansive operators has attracted the attentions of many scholars. By assuming strong convergence of operator products, the convergence results on infinite-dimensional deterministic time-varying difference equations in a general metric space were obtained [69–72]. Reich and Zaslavski [69] studied deterministic time-varying compressive operators in general metric spaces and obtained strong convergence results on the sequence of operator products; Pustylnik *et al.* [70–72] studied the strong convergence of the sequence of operator products consisting of finite number of projection operators.

Noting that a sequence of deterministic operator products converging strongly to 0 can be regarded as a  $L_p^q$ -stable sequence of operators w.r.t. the trivial filter  $\{\mathcal{F}(k) = \{\emptyset, \Omega\}, k \in \mathbb{Z}\}$ , the concept of strong convergence for the sequence of operator products in the above literature can be regarded as a special case of Definition 4.3. Besides, for the case in finite-dimensional spaces, Guo [73] proposed the concept of  $L_p$ -exponentially stable random matrix sequence  $\{I - B(k) \in \mathbb{R}^{N \times N}, k \ge 0\}$ , i.e., there exist constants M > 0 and  $\lambda \in (0, 1)$  such that

$$\mathbb{E}\left[\left\|\prod_{k=n+1}^{m} (I-B(k))\right\|_{\mathscr{L}(\mathbb{R}^{N})}^{p}\right] \le M\lambda^{m-n}, \ \forall \ m > n \ge 0.$$

From Definition 4.3, it can be seen that  $\{I - B(k) \in \mathbb{R}^{N \times N}, k \ge 0\}$  is  $L_r^s$ -stable w.r.t. any given filter  $\{\mathcal{F}(k), k \ge 0\}$ , where  $r = pa^{-1}$ ,  $s = pba^{-1}$  and a, b are positive real numbers with  $a^{-1} + b^{-1} = 1$ .

Let  $\mathcal{F}(k) = \bigvee_{i=0}^{k} (\sigma(F(i); \tau_{S}(\mathscr{L}(\mathscr{X}_{1}))) \bigvee \sigma(G(i); \tau_{S}(\mathscr{L}(\mathscr{X}_{2}, \mathscr{X}_{1}))) \bigvee \sigma(u(i); \tau_{N}(\mathscr{X}_{2}))), k \geq 0$  and  $\mathcal{F}(-1) = \{\emptyset, \Omega\}$ . For the  $L_{2}$ -asymptotic stability of the solution sequence of the random difference equation (4.2), we have the following theorem.

**Theorem 4.4.** For the random difference equation (4.2), let  $\{u(k), \mathcal{F}(k), k \geq 0\}$  be a  $L_2$ bounded sequence of martingale differences, and the sequence  $\{u(k), k \geq 0\}$  be independent of  $\{F(k), k \geq 0\}$  and  $\{G(k), k \geq 0\}$ . If (i)  $\{I_{\mathscr{X}_1} - F(k), k \geq 0\}$  is  $L_2^2$ -stable w.r.t.  $\{\mathcal{F}(k), k \geq 0\}$ , (ii) there exists a sequence of nonnegative real numbers  $\{\gamma(k), k \geq 0\}$  such that

$$\mathbb{E}\left[\left\|I_{\mathscr{X}_{1}}-F(k)\right\|_{\mathscr{L}(\mathscr{X}_{1})}^{4}\right|\mathcal{F}(k-1)\right] \leq 1+\gamma(k) \text{ a.s.}, \quad \sum_{k=0}^{\infty}\gamma(k)<\infty,$$
(4.3)

and (iii)

$$\sum_{k=0}^{\infty} \mathbb{E}\left[\|G(k)\|_{\mathscr{L}(\mathscr{X}_{2},\mathscr{X}_{1})}^{2}\right] < \infty, \quad \sup_{k\geq 0} \mathbb{E}\left[\|G(k)\|_{\mathscr{L}(\mathscr{X}_{2},\mathscr{X}_{1})}^{4}\right] < \infty, \tag{4.4}$$

then, the solution  $\{x(k), k \ge 0\}$  of (4.2) is  $L_2$ -asymptotically stable.

**Remark 4.5.** Systematic results on the stability of randomly time-varying difference equations in finite-dimensional spaces were achieved in [73–78], while the results on randomly time-varying difference equations in infinite-dimensional spaces remain fragmented. Kubrusly [57], Vajjha et al. [58] and the references therein transformed the analysis of the mean square convergence of stochastic approximation algorithms in Hilbert spaces into the L<sub>2</sub>-asymptotic stability analysis of randomly time-varying difference equation. Ungureanu et al. [56], Ungureanu [59, 60] and Zhang et al. [61] investigated the L<sub>2</sub>-asymptotic stability of the solution sequence of the randomly time-varying difference equation x(k + 1) = A(k)x(k) + b(k) in Hilbert space. The above literature assumed  $A(k) : \Omega \to \mathcal{L}(\mathcal{X})$  to be mean square exponentially stable, i.e., there exist constants M > 0 and  $\lambda \in (0, 1)$  such that

$$\mathbb{E}\left[\left\|\left(\prod_{k=n+1}^{m} A(k)\right) x\right\|_{\mathscr{X}}^{2}\right] \le M\lambda^{m-n} \|x\|^{2}, \ \forall \ m > n \ge 0, \ \forall \ x \in \mathscr{X}.$$
(4.5)

For the randomly time-varying difference equation (4.2), even if  $F(k) \equiv F$ , where F is a deterministic self-adjoint compact operator with  $||F||_{\mathscr{L}(\mathscr{X})} \leq 1$ , the operator  $A(k) \equiv I_{\mathscr{X}} - F$  does not satisfy (4.5). In fact, if (4.5) holds, then

$$\left\|\prod_{k=n+1}^{n+2^l} A(k)\right\|_{\mathscr{L}(\mathscr{X})} \le \sqrt{M}\lambda^{2^{l-1}}, \ \forall \ l,n \ge 0.$$
(4.6)

Noting that  $||I_{\mathscr{X}} - F||_{\mathscr{L}(\mathscr{X})} = \sup_{||x||_{\mathscr{X}}=1} |\langle (I_{\mathscr{X}} - F)x, x \rangle_{\mathscr{X}}| = 1 - \inf_{||x||_{\mathscr{X}}=1} \langle Fx, x \rangle_{\mathscr{X}} = 1$ , thus we have

$$\left\|\prod_{k=n+1}^{n+2^l} A(k)\right\|_{\mathscr{L}(\mathscr{X})}^2 = \left\|(I_{\mathscr{X}} - F)^{2^l}\right\|_{\mathscr{L}(\mathscr{X})} = \|I_{\mathscr{X}} - F\|_{\mathscr{L}(\mathscr{X})}^{2^l} = 1, \ \forall \ l, n \ge 0.$$

which is in contradiction to (4.6). Thereby, the existing results and methods on the stability of infinite-dimensional random difference equations are not applicable to random inverse problems.

# 4.2 Convergence of the decentralized algorithm

Let

$$\mathcal{F}(k) = \bigvee_{i=0}^{k} \left( \sigma \left( \mathcal{H}(i); \tau_{\mathbf{S}} \left( \mathscr{L} \left( \mathscr{X}^{N}, \bigoplus_{j=1}^{N} \mathscr{Y}_{j} \right) \right) \right) \right) \bigvee \sigma \left( v(i); \tau_{\mathbf{N}} \left( \bigoplus_{j=1}^{N} \mathscr{Y}_{j} \right) \right) \right), \ k \ge 0,$$

and  $\mathcal{F}(-1) = \{\emptyset, \Omega\}$ , where  $\mathcal{H}(i)$  and v(i) are given in Section 3-B. We need the following assumptions.

Assumption 4.1. The noises  $\{v(k), k \geq 0\}$  with values in  $(\bigoplus_{i=1}^{N} \mathscr{Y}_{i}, \tau_{N}(\bigoplus_{i=1}^{N} \mathscr{Y}_{i}))$  and the random forward operator sequence  $\{H(k), k \geq 0\}$  with values in  $(\mathscr{L}(\mathscr{X}, \bigoplus_{i=1}^{N} \mathscr{Y}_{i}), \tau_{S}(\mathscr{L}(\mathscr{X}, \bigoplus_{i=1}^{N} \mathscr{Y}_{i})))$  are mutually independent.

**Assumption 4.2.** The noises  $\{v(k), \mathcal{F}(k), k \ge 0\}$  is a martingale difference sequence and there exists a constant  $\beta_v > 0$  such that

$$\sup_{k\geq 0} \mathbb{E}\left[ \left\| v(k) \right\|_{\bigoplus_{i=1}^{N} \mathscr{Y}_{i}}^{2} \middle| \mathcal{F}(k-1) \right] \leq \beta_{v} \text{ a.s.}$$

For the gains of the algorithm (3.6), we may need the following conditions.

**Condition 4.1.** The algorithm gains  $\{a(k), k \ge 0\}$  and  $\{b(k), k \ge 0\}$  are both monotonically decreasing sequences of positive real numbers.

Condition 4.2.  $\sum_{k=0}^{\infty} a^2(k) < \infty$  and  $\sum_{k=0}^{\infty} b^2(k) < \infty$ .

**Condition 4.3.**  $\sum_{k=0}^{\infty} a(k) = \infty$  and  $\max\{a(k) - a(k+1), b(k) - a(k)\} = \mathcal{O}(a^2(k) + b^2(k)).$ 

We are now in the position to analyze the convergence of the algorithm (3.6). Firstly, based on Theorem 4.4, we have the following key lemma.

**Lemma 4.6.** For the algorithm (3.6), suppose that Assumptions 4.1, 4.2 and Condition 4.2 hold, there exists a sequence of nonnegative real numbers  $\{\gamma(k), k \ge 0\}$  with  $\sum_{k=0}^{\infty} \gamma(k) < \infty$ , such that

$$\mathbb{E}\left[\left\|I_{\mathscr{X}^{N}}-(a(k)\mathcal{H}^{*}(k)\mathcal{H}(k)+b(k)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}\right)\right\|_{\mathscr{L}(\mathscr{X}^{N})}^{4}\left|\mathcal{F}(k-1)\right] \leq 1+\gamma(k) \text{ a.s.},\tag{4.7}$$

and the sequence of operator-valued random elements  $\{I_{\mathscr{X}^N} - (a(k)\mathcal{H}^*(k)\mathcal{H}(k) + b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}), k \geq 0\}$  is  $L_2^2$ -stable w.r.t.  $\{\mathcal{F}(k), k \geq 0\}$ . I. If

$$\sup_{k\geq 0} \mathbb{E}\left[ \|\mathcal{H}(k)\|_{\mathscr{L}\left(\mathscr{X}^{N},\bigoplus_{i=1}^{N}\mathscr{Y}_{i}\right)}^{2} \right] < \infty,$$

then the algorithm (3.6) is mean square consistent, i.e.,  $\lim_{k\to\infty} \mathbb{E}[\|f_i(k) - f_0\|_{\mathscr{X}}^2] = 0, \ i \in \mathcal{V}.$ II. If there exists a constant  $\rho_0 > 0$  such that

$$\mathbb{E}\left[\left\|\mathcal{H}(k)\right\|_{\mathscr{L}\left(\mathscr{X}^{N},\bigoplus_{i=1}^{N}\mathscr{Y}_{i}\right)}^{2}\right|\mathcal{F}(k-1)\right] \leq \rho_{0} \ a.s.,$$

then the algorithm (3.6) is almost surely strongly consistent, i.e.,  $\lim_{k\to\infty} ||f(k) - f_0||_{\mathscr{X}} = 0$  a.s.,  $i \in \mathcal{V}$ .

**Remark 4.7.** The boundedness condition (4.7) in Lemma 4.6 guarantees the existence of the expectation and conditional expectation of the estimate f(k) in the algorithm (3.6), It is not difficult to verify that (4.7) holds if the norm of the random forward operator has a uniform upper bound independent of the sample path.

From Lemma 4.6, we can see that the  $L_2^2$ -stability of the homogeneous part of the error equation (4.1) plays a key role in the convergence of the algorithm (3.6). We will next give a more intuitive convergence condition of the algorithm (3.6) by further studying the  $L_2^2$ -stability of the sequence of random operators  $\{I_{\mathscr{X}^N} - a(k)\mathcal{H}^*(k)\mathcal{H}(k) - b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}, k \ge 0\}$ . First, we have the following fundamental lemmas.

**Lemma 4.8.** For the algorithm (3.6), suppose that Assumptions 4.1, 4.2, Conditions 4.1 and 4.2 hold. If there exists an integer h > 0 and a constant  $\rho_0 > 0$  such that

$$\left\{I_{\mathscr{X}^N} - \sum_{i=kh}^{(k+1)h-1} (a(i)\mathcal{H}^*(i)\mathcal{H}(i) + b(i)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}), k \ge 0\right\}$$

is  $L_2^2$ -stable w.r.t.  $\{\mathcal{F}((k+1)h-1), k \ge 0\};$ (ii)  $\sup_{k\ge 0} \left(\mathbb{E}\left[\|\mathcal{H}^*(k)\mathcal{H}(k)\|_{\mathscr{L}(\mathscr{X}^N)}^{2^{\max\{h,2\}}} \middle| \mathcal{F}(k-1)\right]\right)^{\frac{1}{2^{\max\{h,2\}}}} \le \rho_0 \text{ a.s.,}$ then  $\{I_{\mathscr{X}^N} - a(k)\mathcal{H}^*(k)\mathcal{H}(k) - b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}, k \ge 0\}$  is  $L_2^2$ -stable w.r.t.  $\{\mathcal{F}(k), k \ge 0\},$ 

If the graph  $\mathcal{G}$  is connected, then by Lemma 4.8, we can further obtain a more intuitive sufficient condition, under which the operator-valued random sequence  $\{I_{\mathscr{X}^N} - a(k)\mathcal{H}^*(k)\mathcal{H}(k) - b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}, k \geq 0\}$  is  $L_2^2$ -stable w.r.t.  $\{\mathcal{F}(k), k \geq 0\}$ .

**Lemma 4.9.** For the algorithm (3.6), let  $\mathcal{G}$  be connected and assume that Assumptions 4.1, 4.2 and Conditions 4.1-4.3 hold. If there exist positive self-adjoint operators  $\mathscr{H}_i \in \mathscr{L}(\mathscr{X})$ ,  $i = 1, \dots, N$  satisfying  $\sum_{i=1}^{N} \mathscr{H}_i > 0$ , and an integer h > 0, such that

$$\sum_{j=1}^{N}\sum_{k=0}^{\infty}\mathbb{E}\left[\left\|\mathscr{H}_{j}x(k)-\sum_{i=kh}^{(k+1)h-1}\mathbb{E}\left[H_{j}^{*}(i)H_{j}(i)x(k)\middle|\mathcal{F}(kh-1)\right]\right\|_{\mathscr{X}}^{2}\right]<\infty,$$
(4.8)

for any  $L_2$ -bounded adaptive sequence  $\{x(k), \mathcal{F}(kh-1), k \geq 0\}$  with values in the Hilbert space  $\mathscr{X}$ , and there exists a constant  $\rho_0 > 0$  such that

$$\sup_{k\geq 0} \left( \mathbb{E}\left[ \left\| \mathcal{H}^*(k)\mathcal{H}(k) \right\|_{\mathscr{L}(\mathscr{X}^N)}^{2^{\max\{h,2\}}} \middle| \mathcal{F}(k-1) \right] \right)^{\frac{1}{2^{\max\{h,2\}}}} \leq \rho_0 \ a.s., \tag{4.9}$$
$$-a(k)\mathcal{H}^*(k)\mathcal{H}(k) - b(k)\mathcal{L}_{\mathscr{L}} \otimes L_{\mathscr{X}} \ k\geq 0 \} \text{ is } L^2_{\ast}\text{-stable wrt} \{\mathcal{F}(k) \ k\geq 0 \}$$

then  $\{I_{\mathscr{X}^N} - a(k)\mathcal{H}^*(k)\mathcal{H}(k) - b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}, k \ge 0\}$  is  $L_2^2$ -stable w.r.t.  $\{\mathcal{F}(k), k \ge 0\}$ .

Combing Lemmas 4.6 and 4.9, we will next give intuitive sufficient conditions on the mean square and almost sure strong consistency of the algorithm (3.6).

**Theorem 4.10.** For the algorithm (3.6), suppose that all the conditions in Lemma 4.9 hold. If there exists a sequence of nonnegative real numbers  $\{\Gamma(k), k \ge 0\}$  with  $\sum_{k=0}^{\infty} \Gamma(k) < \infty$ , such that

$$\mathbb{E}\left[\left\|I_{\mathscr{X}^{N}}-4\left(a(k)\mathcal{H}^{*}(k)\mathcal{H}(k)+b(k)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}\right)\right\|_{\mathscr{L}(\mathscr{X}^{N})}\right|\mathcal{F}(k-1)\right]$$

$$\leq 1+\Gamma(k) \ a.s.,$$
(4.10)

then the algorithm (3.6) is both mean square and almost surely strongly consistent.

Especially, if  $\{\mathcal{H}(k), k \geq 0\}$  with values in  $(\mathscr{L}(\mathscr{X}^N, \bigoplus_{i=1}^N \mathscr{Y}_i), \tau_{\mathsf{S}}(\mathscr{L}(\mathscr{X}^N, \bigoplus_{i=1}^N \mathscr{Y}_i)))$  and the random sequence  $\{v(k), k \geq 0\}$  with values in  $(\bigoplus_{i=1}^N \mathscr{Y}_i, \tau_{\mathsf{N}}(\bigoplus_{i=1}^N \mathscr{Y}_i))$  are both i.i.d. and they are mutually independent, then the following corollary follows from Theorem 4.10.

**Corollary 4.11.** For the algorithm (3.6), assume that  $\mathcal{G}$  is connected, Assumptions 4.1, 4.2 and Conditions 4.1-4.3 hold, and  $\{\mathcal{H}(k), k \ge 0\}$  and  $\{v(k), k \ge 0\}$  are both i.i.d. sequences and they are mutually independent. If there exists a constant  $\rho_0 > 0$  such that  $\|\mathcal{H}(0)\| \le \rho_0$  a.s. and

$$\sum_{j=1}^{N} \mathbb{E}\left[ \|H_j(0)x\|_{\mathscr{Y}_j}^2 \right] > 0, \ \forall \ x \in \mathscr{X} \setminus \{0\},$$

$$(4.11)$$

then the algorithm (3.6) is both mean square and almost surely strongly consistent.

The graph  $\mathcal{G}$  describes the communication topology among nodes, and its connectivity ensures the nodes to collaboratively reconstruct the unknown function  $f_0$  successfully. The condition (4.8) in Lemma 4.9 plays an important role in the convergence analysis of the decentralized algorithm, which we call the *infinite-dimensional spatio-temporal persistence of excitation* condition. Note that

$$\begin{aligned} &\left|\sum_{j=1}^{N}\sum_{i=kh}^{(k+1)h-1}\mathbb{E}\left[\left\|H_{j}(i)x\right\|_{\mathscr{Y}_{j}}^{2}\right] - \left\langle\sum_{j=1}^{N}\mathscr{H}_{j}x,x\right\rangle_{\mathscr{X}}\right| \\ &= \left|\sum_{j=1}^{N}\left\langle\mathbb{E}\left[\sum_{i=kh}^{(k+1)h-1}H_{j}^{*}(i)H_{j}(i)x - \mathscr{H}_{j}x\right],x\right\rangle_{\mathscr{X}}\right| \\ &= \left|\sum_{j=1}^{N}\left\langle\mathbb{E}\left[\sum_{i=kh}^{(k+1)h-1}\mathbb{E}\left[H_{j}^{*}(i)H_{j}(i)x\right|\mathcal{F}(kh-1)\right] - \mathscr{H}_{j}x\right],x\right\rangle_{\mathscr{X}}\right| \\ &\leq \sum_{j=1}^{N}\left[\mathbb{E}\left[\left\|\mathscr{H}_{j}x - \sum_{i=kh}^{(k+1)h-1}\mathbb{E}\left[H_{j}^{*}(i)H_{j}(i)x\right|\mathcal{F}(kh-1)\right]\right\|_{\mathscr{X}}^{2}\right]\right]^{\frac{1}{2}}\|x\|_{\mathscr{X}}, x \in \mathscr{X}, k \ge 0. \end{aligned}$$

Therefore, the infinite-dimensional spatio-temporal persistence of excitation condition is equivalent to the combination of the following two conditions.

• For any  $x \in \mathscr{X} \setminus \{0\}$ , there exists an integer K(x) > 0 such that

$$\sum_{j=1}^{N} \sum_{i=kh}^{(k+1)h-1} \mathbb{E}\left[ \|H_j(i)x\|_{\mathscr{Y}_j}^2 \right] > 0, \ k \ge K(x);$$
(4.12)

• there exist deterministic time-invariant operators  $\mathscr{H}_j$ ,  $j = 1, \dots, N$  such that

$$\sum_{j=1}^{N}\sum_{k=0}^{\infty}\mathbb{E}\left[\left\|\mathscr{H}_{j}x(k)-\sum_{i=kh}^{(k+1)h-1}\mathbb{E}\left[H_{j}^{*}(i)H_{j}(i)x(k)\middle|\mathcal{F}(kh-1)\right]\right\|_{\mathscr{X}}^{2}\right]<\infty,$$

for any  $L_2$ -bounded adaptive sequence  $\{x(k), \mathcal{F}(kh-1), k \ge 0\}$  with values in the Hilbert space  $\mathscr{X}$ .

The *spatio-temporal persistence of excitation* implies that the non-zero orbits of the random forward operators of all nodes are non-degenerate in the mean square sense for a fixed length time period, where *spatio-temporal* refers specifically to the temporal and spatial states of the operator orbit  $x \mapsto H_j(i)x$  of the random forward operator. By (4.12) we know that we neither need the temporal orbit of the forward operator of each node over the graph to be non-degenerate, i.e., (k+1)h-1

$$\sum_{i=kh}^{k+1)h-1} \mathbb{E}\left[ \|H_j(i)x\|_{\mathscr{Y}_j}^2 \right] > 0, \ \forall \ j \in \mathcal{V},$$

nor need the spatial orbit of the forward operator of all nodes to be non-degenerate at each instant, i.e.,

$$\sum_{j=1}^{N} \mathbb{E}\left[ \|H_j(i)x\|_{\mathscr{Y}_j}^2 \right] > 0, \ \forall \ i \ge 0$$

Notably, if the sequences of random forward operator and noises are both i.i.d. and they are mutually independent, then the *infinite-dimensional spatio-temporal persistence of excitation* condition degenerates to the case that the spatial orbits of the forward operators of all nodes are non-degenerate at the initial moment k = 0, i.e., the condition (i) in the Corollary 4.11.

In the past decades, to solve the problems of finite-dimensional parameter estimation and signal tracking with non-stationary and non-independent data, many scholars have proposed excitation conditions based on the conditional expectation of the observation/regression matrix. The stochastic persistence of excitation condition was first proposed by Guo [74] in the analysis of centralized Kalman filtering algorithms. Xie and Guo [79] proposed the cooperative information condition based on the conditional expectations of the observation matrices for the distributed adaptive filtering algorithm over connected graphs. Wang *et al.* [32] proposed the stochastic spatio-temporal persistence of excitation condition for the decentralized online estimation algorithm over randomly time-varying graphs. Zhang *et al.* [80] proved that if the graph is connected and the randomly time-varying regression matrix satisfies the uniformly conditionally spatio-temporally joint observability condition, i.e., there exists an integer h > 0 and a constant  $\theta > 0$ , respectively, such that

$$\inf_{k\geq 0} \lambda_{\min}\left(\sum_{j=1}^{N} \sum_{i=kh}^{(k+1)h-1} \mathbb{E}\left[H_j^T(i)H_j(i) \middle| \mathcal{F}(kh-1)\right]\right) \geq \theta \text{ a.s.},$$

then the algorithm achieves mean square and almost sure convergence. For this case, it is not difficult to verify that the random matrix sequence  $\{I_{Nn} - a(k)\mathcal{H}^*(k)\mathcal{H}(k) - b(k)\mathcal{L}_{\mathcal{G}} \otimes I_n, k \geq 0\}$ of satisfies the  $L_2^2$ -stability condition w.r.t.  $\{\mathcal{F}(k), k \geq 0\}$ , and thus the excitation conditions proposed in [32, 80] are all special cases of the  $L_2^2$ -stability condition in Lemma 4.6.

The persistence of excitation conditions proposed for finite-dimensional systems all require that the conditional expectation of the information matrix consisting of the observation (regression) matrices is positive definite, i.e., the information matrix has strictly positive minimum eigenvalues; however, inverse problems in infinite-dimensional Hilbert spaces are usually illposed. Even for a strictly positive linear compact operator, the excitation condition similar to the persistence excitation conditions for finite-dimensional systems can not hold any more since the infimum of the eigenvalues of the compact operator is always 0.

#### 5. DECENTRALIZED ONLINE LEARNING IN REPRODUCING KERNEL HILBERT SPACE

We will discuss in this section a special class of online random inverse problems: decentralized online learning problems in reproducing kernel Hilbert spaces (RKHS). Let  $\mathscr{X}$  be a non-empty subset of  $\mathbb{R}^n$ ,  $K : \mathscr{X} \times \mathscr{X} \to \mathbb{R}$  be a Mercer kernel, and  $(\mathscr{H}_K, \langle \cdot, \cdot \rangle_K)$  be a reproducing kernel Hilbert space with kernel K, which is consisted of functions with domain  $\mathscr{X}$ . Suppose  $f_0 : \mathscr{X} \to \mathbb{R}$  is an unknown function in  $\mathscr{H}_K$ . The nodes cooperatively estimate  $f_0$  by information exchanging among them. The observation data  $y_i(k)$  of the *i*-th node at instant k is given by

$$y_i(k) = f_0(x_i(k)) + v_i(k), \ k \ge 0, \ i \in \mathcal{V},$$
(5.1)

where  $x_i(k) : \Omega \to \mathscr{X}$  is a random vector with values in the Hilbert space  $(\mathscr{X}, \tau_N(\mathscr{X}))$  at instant k, called the random input data, and the observation noise  $v_i(k) : \Omega \to \mathbb{R}$  is a random vector with values in the Hilbert space  $(\mathbb{R}, \tau_N(\mathbb{R}))$ .

**Remark 5.1.** In [41], the measurement of each node is given by  $y_i(k) = f_i(\theta^*) + v_i(k)$ , i = 1, 2, ..., N, in which the nonlinear mappings  $f_i(\cdot)$ , i = 1, 2, ..., N are completely known in prior and it is the parameter vector  $\theta^*$  in a finite-dimensional space, who is unknown and to be estimated. While for the measurement model (5.1), it is the mapping  $f_0$  in the infinite-dimensional space  $\mathscr{H}_K$ , who is unknown and to be estimated.

For any given  $x \in \mathscr{X}$ , the function  $K_x : \mathscr{X} \to \mathscr{H}_K$  induced by the Mercer kernel is given by

$$K_x(y) = K(x, y), \ \forall \ y \in \mathscr{X}.$$

Define the random forward operator  $H_i(k)$ :

$$H_i(k)(f) := f(x_i(k)), \ f \in \mathscr{H}_K, \ k \ge 0, \ i \in \mathcal{V},$$

then the measurement model (5.1) can be represented as the random inverse problem with the measurement equation (3.1). Based on the algorithm (3.6), the decentralized online learning strategy in  $\mathcal{H}_K$  is given by

$$f_i(k+1) = f_i(k) + a(k)(y_i(k) - f_i(k)(x_i(k)))K_{x_i(k)} + b(k)\sum_{j\in\mathcal{N}_i} a_{ij}(f_j(k) - f_i(k)),$$
  
$$k \ge 0, \ i \in \mathcal{V}.$$
(5.2)

Given  $\phi, \psi \in \mathscr{H}_K$ , denote the rank 1 tensor product operator  $\phi \otimes \psi : \mathscr{H}_K \to \mathscr{H}_K$  by

$$(\phi \otimes \psi)(f) := \langle f, \psi \rangle_K \phi, \ f \in \mathscr{H}_K.$$

Let

$$\mathcal{F}(k) = \bigvee_{s=0}^{k} \bigvee_{i=1}^{N} \left( \bigvee_{f \in \mathscr{H}_{K}} \sigma\left(f(x_{i}(s)); \tau_{N}(\mathbb{R})\right) \bigvee \sigma\left(v_{i}(s); \tau_{N}(\mathbb{R})\right) \right), \ k \ge 0$$

and  $\mathcal{F}(-1) = \{\emptyset, \Omega\}$ . For the algorithm (5.2), we need the following assumptions.

Assumption 5.1. The sequence  $\{x_i(k), i \in \mathcal{V}, k \geq 0\}$  of random vectors with values in the Hilbert space  $(\mathscr{X}, \tau_N(\mathscr{X}))$  and the sequence  $\{v_i(k), i \in \mathcal{V}, k \geq 0\}$  of random variables with values in the Hilbert space  $(\mathbb{R}, \tau_N(\mathbb{R}))$  are mutually independent.

**Assumption 5.2.** The noises  $\{v_i(k), \mathcal{F}(k), k \ge 0\}$ ,  $i \in \mathcal{V}$ , are martingale difference sequences and there exists a constant  $\beta > 0$ , such that

$$\max_{i \in \mathcal{V}} \sup_{k \ge 0} \mathbb{E} \left[ \| v_i(k) \|_{\mathbb{R}}^2 \big| \mathcal{F}(k-1) \right] \le \beta \text{ a.s.}$$

Assumption 5.3.  $\sup_{x \in \mathscr{X}} K(x, x) < \infty$ .

**Remark 5.2.** The existing works on RKHS online learning (e.g., [19–22, 53, 81]) all require the random input data to be i.i.d. Notice that in Assumption 5.1, the sequence of random forward operators with values in  $(\mathscr{L}(\mathscr{X}, \bigoplus_{i=1}^{N} \mathscr{Y}_{i}), \tau_{S}(\mathscr{L}(\mathscr{X}, \bigoplus_{i=1}^{N} \mathscr{Y}_{i})))$  is not required to satisfy special statistical properties such as independence, stationarity, etc.

**Remark 5.3.** Assumption 5.3 is often used in the learning theory in RKHS ([20, 21, 53, 81]). There are many kernel functions satisfying Assumption 5.3, such as the Gaussian kernel  $K : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ ,  $K(x, y) = e^{-\frac{\|x-y\|_{\mathbb{R}^n}^2}{c^2}}$  and the homogeneous polynomial kernel  $K : \mathbb{S}^{n-1} \times \mathbb{S}^{n-1} \to \mathbb{R}$  on the unit sphere in  $\mathbb{R}^n$ ,  $K(x, y) = \langle x, y \rangle_{\mathbb{R}^n}^d$ , where the integer d > 0. In addition to these common kernel functions, note that the Mercer kernel is continuous, thus Assumption 5.3 holds for arbitrary compact set  $\mathscr{X} \subseteq \mathbb{R}^n$ .

Noting the continuity of the Mercer kernel function,  $K_x$  is therefore a continuous function. For any  $f \in \mathscr{H}_K$  and random input data  $x_i(k)$ ,  $i \in \mathcal{V}$ ,  $k \ge 0$ , it follows from the reproducing properties of RKHS that  $f(x_i(k)) = \langle f, K_{x_i(k)} \rangle_K$  ([18, 82]), thus  $f(x_i(k)) : \Omega \to \mathbb{R}$  is a random variable with values in  $(\mathbb{R}, \tau_N(\mathbb{R}))$ . It follows from Assumption 5.3 that  $H_i(k)$  is the random element with values in  $(\mathscr{L}(\mathscr{H}_K, \mathbb{R}), \tau_S(\mathscr{L}(\mathscr{H}_K, \mathbb{R})))$ , which is induced by the random input data  $x_i(k)$ .

Based on the convergence results of the algorithm (3.6), we can obtain the following convergence results on the decentralized online learning algorithm (5.2) in RKHS. The proofs of theorems and corollaries of this section are in Appendix C.

**Theorem 5.4.** For the algorithm (5.2), suppose that  $\mathcal{G}$  is connected, Assumptions 5.1-5.3 and Conditions 4.1-4.3 hold. If there exist positive self-adjoint operators  $N_i \in \mathscr{L}(\mathscr{H}_K)$ ,  $i = 1, \dots, N$ satisfying  $\sum_{i=1}^{N} N_i > 0$  and there exists an integer h > 0, such that

$$\sum_{j=1}^{N}\sum_{k=0}^{\infty}\mathbb{E}\left[\left\|\left(N_{j}-\sum_{i=kh}^{(k+1)h-1}\mathbb{E}\left[K_{x_{j}(i)}\otimes K_{x_{j}(i)}\middle|\mathcal{F}(kh-1)\right]\right)g(k)\right\|_{K}^{2}\right]<\infty,$$
(5.3)

for arbitrary  $L_2$ -bounded adaptive sequence  $\{g(k), \mathcal{F}(kh-1), k \ge 0\}$ , then the algorithm (5.2) is both mean square and almost surely strongly consistent.

**Corollary 5.5.** For the algorithm (5.2), suppose that  $\mathcal{G}$  is connected, Assumptions 5.1-5.3 and Conditions 4.1-4.3 hold. If there exist positive self-adjoint operators  $N_i \in \mathscr{L}(\mathscr{H}_K)$ ,  $i = 1, \dots, N$ satisfying  $\sum_{i=1}^{N} N_i > 0$ , and there exists an integer h > 0, a constant  $\mu_0 > 0$  and a nonnegative real sequence  $\{\tau(k), k \ge 0\}$ , respectively, such that

$$\max_{j\in\mathcal{V}} \left\| N_j - \sum_{i=kh}^{(k+1)h-1} \mathbb{E}\left[ K_{x_j(i)} \otimes K_{x_j(i)} \middle| \mathcal{F}(kh-1) \right] \right\|_{\mathscr{L}(\mathscr{H}_K)}^2 \le \mu_0 \tau(k) \ a.s., \tag{5.4}$$

where  $\sum_{k=0}^{\infty} \tau(k) < \infty$ , then the algorithm (5.2) is both mean square and almost surely strongly consistent.

**Remark 5.6.** Noting that Assumption 5.3 implies that  $K_{x_j(i)} \otimes K_{x_j(i)}$  is a Bochner integrable random element with values in the Banach space  $(\mathscr{L}(\mathscr{H}_K), \tau_N(\mathscr{L}(\mathscr{H}_K))))$ , then the conditional expectations  $\mathbb{E}[K_{x_j(i)} \otimes K_{x_j(i)} | \mathcal{F}(kh-1)]$ ,  $i, k \ge 0, j \in \mathcal{V}$  uniquely exist by Lemma A.9.

Especially, if the input data  $\{(x_1(k), \dots, x_N(k)), k \ge 0\}$  are i.i.d, then we have the following corollary.

**Corollary 5.7.** For the algorithm (5.2), suppose that  $\{(x_1(k), \dots, x_N(k)), k \ge 0\}$  and  $\{(v_1(k), \dots, v_N(k)), k \ge 0\}$  be i.i.d. sequences and they are mutually independent, and  $\mathcal{G}$  is connected. If Assumptions 5.1-5.3 and Conditions 4.1-4.3 hold, and

$$\mathbb{E}\left[\sum_{j=1}^{N} K_{x_j(0)} \otimes K_{x_j(0)}\right] > 0,$$
(5.5)

then the algorithm (5.2) is both mean square and almost surely strongly consistent.

**Remark 5.8.** For the centralized online learning with input data drawn independently from the probability measure  $\rho_{\mathscr{X}}$  on  $\mathscr{X}$ , Tarres and Yao [20] defined the covariance operator of the probability measure  $\rho_{\mathscr{X}}$  in  $\mathscr{H}_K$  as  $L_K : \mathscr{H}_K \to \mathscr{H}_K$ ,

$$L_K(f)(y) = \int_{\Omega} K(x, y) f(x) \, d\rho_{\mathscr{X}}, \ \forall \ f \in \mathscr{H}_K.$$

By the reproducing property and Assumption 5.3, it follows that  $L_K = \mathbb{E}[K_x \otimes K_x]$ . This implies that for the centralized online learning problem in RKHS with i.i.d. data, the condition (5.5) in Corollary 5.7 just degenerates to that in [20]: the covariance operator  $L_K > 0$ .

# 6. CONCLUSIONS

We have established a framework of random inverse problems with real-time measurements over graphs, and present a decentralized online learning algorithm with online data streams, which unifies the distributed parameter estimation in Hilbert spaces and the least mean square problem in reproducing kernel Hilbert spaces (RKHS-LMS). It is not required that the random forward operators satisfy special statistical assumptions such as mutual independence, spatio-temporal independence or stationarity. Each node updates its estimate at the next instant by using its new observation and a weighted sum of its own and neighbors' estimates. Firstly, based on the existing analytical results in Banach space, we further exploited the probabilistic properties of random elements with values in different topological spaces in a stochastic framework. Then, we proposed the  $L_p^q$ -stability condition on the sequence of operator-valued random elements, and established the  $L_2$ -asymptotic stability theory of a class of inhomogeneous random difference equations in Hilbert spaces with  $L_2$ -bounded martingale difference terms. Subsequently, we transform the asymptotic stability of these kinds of infinite-dimensional random difference equations into the  $L_2^2$ -stability condition on the operator-valued random elements. We have obtained an intuitive sufficient condition on the convergence of decentralized online learning algorithms for random inverse problems over graphs which is imposed on the random forward operators and the Laplacian matrix of the graph, i.e., the *infinite-dimensional spatio-temporal persistence of excitation* condition. We have proved that if the forward operators over connected graphs satisfy the *infinitedimensional spatio-temporal persistence of excitation* condition, then all nodes' estimates are mean square and almost surely strongly consistent. Finally, by equivalently transforming the distributed learning problem in RKHS to the random inverse problem over graphs, we propose a decentralized online learning algorithm in RKHS with non-stationary and non-independent online data streams, and prove that the algorithm is mean square and almost surely strongly consistent if the operators induced by the random input data satisfy the *infinite-dimensional spatio-temporal persistence of excitation* condition.

#### APPENDIX A

## PRELIMINARY ANALYTICAL THEORY OF BANACH SPACES

**Definition A.1** ([67]). Let  $(S, \mathscr{A})$  and  $(T, \mathscr{B})$  be measurable spaces, if the mapping  $f : S \to T$  satisfies

$$f^{-1}(B) := \{ x \in S : f(x) \in B \} \in \mathscr{A}, \ \forall \ B \in \mathscr{B},$$

then f is said to be  $\mathscr{A}/\mathscr{B}$ -measurable.

**Definition A.2** ([67]). The function  $f = \sum_{i=1}^{n} \mathbf{1}_{A_i} \otimes x_i$  is called a simple function with values in the Banach space  $\mathscr{X}$ , where  $A_i \in \mathcal{F}$ ,  $x_i \in \mathscr{X}$ ,  $i = 1, \dots, n$ ,  $\mathbf{1}_A$  is the indicator function of the set A and  $(\mathbf{1}_A \otimes x)(s) := \mathbf{1}_A(s)x, \forall x \in \mathscr{X}, s \in \Omega$ .

**Definition A.3** ([67]). Given a mapping  $f : \Omega \to \mathscr{X}$  with values in the Banach space  $(\mathscr{X}, \tau_N(\mathscr{X}))$ , if there exists a sequence of simple functions  $\{f_n, n \ge 1\}$  almost everywhere converging to f in the topology  $\tau_N(\mathscr{X})$ , then f is said to be strongly measurable with respect to  $\tau_N(\mathscr{X})$ .

**Lemma A.4** ([83]). The mapping  $f : \Omega \to \mathscr{X}$  is strongly measurable with respect to the topology  $\tau_N(\mathscr{X})$  if and only if f is a random element with values in the Banach space  $(\mathscr{X}, \tau_N(\mathscr{X}))$ .

Combining the Definition 2.2 and Corollary 1.4.7, Proposition 1.1.28 and Corollary 1.1.29 in [67], we directly obtain the following properties of operator-valued random elements.

**Proposition A.5.** Let  $f_0 : \Omega \to \mathscr{L}(\mathscr{Y}, \mathscr{Z}), f_1 : \Omega \to \mathscr{X}, f_2 : \Omega \to \mathscr{L}(\mathscr{X}, \mathscr{Y}), g_1 : \Omega \to \mathscr{L}(\mathscr{X}, \mathscr{Y}) \text{ and } g_2 : \Omega \to \mathscr{L}(\mathscr{Y}, \mathscr{Z}).$ 

(a). The mapping  $f_0$  is a random element with values in  $(\mathscr{L}(\mathscr{Y}, \mathscr{Z}), \tau_s(\mathscr{L}(\mathscr{Y}, \mathscr{Z})))$  if and only if the mapping  $f_0y : \omega \mapsto f_0(\omega)y$  is a random element with values in  $(\mathscr{Z}, \tau_N(\mathscr{Z}))$  for arbitrary  $y \in \mathscr{Y}$ .

(b). If  $f_1$  is strongly measurable with respect to  $\tau_N(\mathscr{X})$  and  $g_1x : \Omega \to \mathscr{Y}$  is strongly measurable with respect to  $\tau_N(\mathscr{Y}), \forall x \in \mathscr{X}$ , then  $g_1f_1 : \Omega \to \mathscr{Y}$  is strongly measurable with respect to  $\tau_N(\mathscr{Y})$ .

(c). For arbitrary  $x \in \mathscr{X}$  and  $y \in \mathscr{Y}$ , if the mapping  $f_2x : \Omega \to \mathscr{Y}$  is strongly measurable with respect to  $\tau_N(\mathscr{Y})$  and the mapping  $g_2y : \Omega \to \mathscr{Z}$  is strongly measurable with respect to  $\tau_N(\mathscr{Z})$ , then the mapping  $(g_2f_2)x' : \Omega \to \mathscr{Z}$  is strongly measurable with respect to  $\tau_N(\mathscr{Z})$  for arbitrary  $x' \in \mathscr{X}$ .

If the operator-valued mapping  $f : \Omega \to \mathscr{L}(\mathscr{Y}, \mathscr{Z})$  is strongly measurable with respect to the uniform operator topology  $\tau_{N}(\mathscr{L}(\mathscr{Y}, \mathscr{Z}))$ , then for arbitrary  $y \in \mathscr{Y}$ ,  $fy : \Omega \to \mathscr{Z}$  is strongly measurable with respect to  $\tau_{N}(\mathscr{Z})$ . By Proposition A.5. (a) and Pettis measurability theorem ([84]), it follows that a random element with values in  $(\mathscr{L}(\mathscr{Y}, \mathscr{Z}), \tau_{N}(\mathscr{L}(\mathscr{Y}, \mathscr{Z})))$  is a random element with values in  $(\mathscr{L}(\mathscr{Y}, \mathscr{Z}), \tau_{S}(\mathscr{L}(\mathscr{Y}, \mathscr{Z})))$ . For the measurability of mappings with values in the space of operators with different topologies, one may refer to [67, 83, 84].

**Definition A.6** ([66]). If  $f \in L^1(\Omega; \mathscr{X})$ , then the mathematical expectation of f is defined as the Bochner integral

$$\mathbb{E}[f] = (\mathbf{B}) \int_{\Omega} f \, \mathrm{d}\mathbb{P}.$$

Without raising ambiguity, the Bochner integral in this paper will omit the capital letter (B) in front of the integral symbol.

**Definition A.7** ([67]). Let  $\mathscr{G} \subseteq \mathcal{F}$  be a sub- $\sigma$ -algebra,  $f \in L^0(\Omega; \mathscr{X})$  and  $g \in L^0(\Omega, \mathscr{G}; \mathscr{X})$ . If

$$\int_{F} g \, \mathrm{d}\mathbb{P} = \int_{F} f \, \mathrm{d}\mathbb{P}, \ \forall \ F \in \mathscr{G}_{f} \cap \mathscr{G}_{g},$$

where  $\mathscr{G}_{\phi} := \{F \in \mathscr{G} : \mathbf{1}_F \phi \in L^1(\Omega; \mathscr{X})\}, \phi \in L^0(\Omega; \mathscr{X}), \text{ then } g \text{ is called the conditional expectation of } f \text{ with respect to } \mathscr{G}.$ 

**Definition A.8** ([67]). Let  $\{\mathcal{F}_k, k \ge 0\}$  be a family of sub- $\sigma$ -algebras of  $\mathcal{F}$  and  $\{f_k, k \ge 0\}$  be a family of random elements with values in the Hilbert space  $(\mathscr{X}, \tau_N(\mathscr{X}))$ .

- 1. If  $\mathcal{F}_n \subseteq \mathcal{F}_m$ ,  $\forall m \ge n \ge 0$ , then  $\{\mathcal{F}_k, k \ge 0\}$  is called a filter in  $(\Omega, \mathcal{F}, \mathbb{P})$ .
- 2. If  $\{\mathcal{F}_k, k \ge 0\}$  is a filter in  $(\Omega, \mathcal{F}, \mathbb{P})$ ,  $f_k \in L^0(\Omega, \mathcal{F}_k; \mathscr{X})$ ,  $\forall k \ge 0$ , then  $\{f_k, \mathcal{F}_k, k \ge 0\}$  is called an adaptive sequence.
- 3. If  $\{f_k, \mathcal{F}_k, k \ge 0\}$  is an adaptive sequence,  $f_k$  is Bochner integrable on  $\mathcal{F}_{k-1}$  and

$$\mathbb{E}[f_k | \mathcal{F}_{k-1}] = 0, \ \forall \ k \ge 0,$$

then  $\{f_k, \mathcal{F}_k, k \ge 0\}$  is called a sequence of martingale differences.

**Lemma A.9** ([67]). If  $\mathscr{G} \subseteq \mathcal{F}$  is a sub- $\sigma$ -algebra and  $f \in L^1(\Omega; \mathscr{X})$ , then there exists a unique conditional expectation  $\mathbb{E}[f|\mathscr{G}] \in L^0(\Omega, \mathscr{G}; \mathscr{X}) \cap L^1(\Omega; \mathscr{X})$  satisfying

$$\int_{A} \mathbb{E}[f|\mathscr{G}] \, d\mathbb{P} = \int_{A} f \, d\mathbb{P}, \,\, \forall \,\, A \in \mathscr{G}.$$

**Lemma A.10** ([67]). Let  $\mathscr{X}_1, \mathscr{X}_2$  and  $\mathscr{Y}$  be Banach spaces,  $\beta : \mathscr{X}_1 \times \mathscr{X}_2 \to \mathscr{Y}$  be a bounded bilinear mapping and  $\mathscr{G}$  be a sub- $\sigma$ -algebra of  $\mathcal{F}$ . If  $g \in L^0(\Omega, \mathscr{G}; \mathscr{X}_1)$  and  $f \in L^0(\Omega; \mathscr{X}_2)$  is Bochner integrable on  $\mathscr{G}$ , then  $\beta(g, f) \in L^0(\Omega; \mathscr{Y})$  is Bochner integrable on  $\mathscr{G}$ and  $\mathbb{E}[\beta(g, f)|\mathscr{G}] = \beta(g, \mathbb{E}[f|\mathscr{G}])$  a.s.

## APPENDIX B

# **PROOFS OF SECTION 2**

*Proof of Proposition 2.4.* This proposition is proved by the **two steps** as follows.

Step 1: If f is a simple function with values in the Banach space  $\mathscr{X}$ , without loss of generality, let  $f = \sum_{i=1}^{n} \mathbf{1}_{A_i} \otimes y_i$ , where  $A_i \in \mathcal{F}$ ,  $y_i \in \mathscr{X}$ ,  $A_i \cap A_j = \emptyset$ ,  $1 \le i \ne j \le n$ . Note that  $\sigma(\mathbf{1}_{A_i} \otimes y_i; \tau_{\mathrm{N}}(\mathscr{X})) = \sigma(\mathbf{1}_{A_i}; \tau_{\mathrm{N}}(\mathscr{X})) = \{\emptyset, \Omega, A_i, A_i^{\complement}\}$ . For any given  $B \in \mathscr{B}(\mathscr{Y}; \tau_{\mathrm{N}}(\mathscr{Y}))$ , we have

$$= \begin{cases} \left(T\left(\sum_{i=1}^{n} \mathbf{1}_{A_{i}} \otimes y_{i}\right)\right)^{-1}(B) \\ \left(\bigcup_{i=1}^{n} (Ty_{i})^{-1}(B)\right) \bigcap \left(\bigcup_{i=1}^{n} A_{i}\right) \bigcup \left(\bigcup_{i=1}^{n} A_{i}\right)^{\complement}, \quad 0 \in B; \\ \left(\bigcup_{i=1}^{n} (Ty_{i})^{-1}(B)\right) \bigcap \left(\bigcup_{i=1}^{n} A_{i}\right), \quad 0 \notin B, \end{cases}$$

which gives  $T(\sum_{i=1}^{n} \mathbf{1}_{A_i} \otimes y_i) \in (\bigvee_{x \in \mathscr{X}} \sigma(Tx; \tau_{\mathbf{N}}(\mathscr{Y}))) \bigvee \sigma(f; \tau_{\mathbf{N}}(\mathscr{X})).$ 

Step 2: For the random element f with values in the Banach space  $(\mathscr{X}, \tau_{N}(\mathscr{X}))$ , by Lemma A.4 and Definition A.3, we know that there exists a sequence of simple functions  $\{f_n, n \ge 1\}$ with values in  $\mathscr{X}$  such that  $\lim_{n\to\infty} ||f_n - f||_{\mathscr{X}} = 0$  a.s. It follows from Proposition A.5.(b) that  $Tf: \Omega \to \mathscr{Y}$  is strongly measurable with respect to the topology  $\tau_{N}(\mathscr{Y})$ , i.e., there exists  $\Omega_0 \subseteq \Omega, \mathbb{P}(\Omega_0) = 1$  and a separable closed subspace  $\mathscr{Y}_0 \subseteq \mathscr{Y}$  such that  $Tf(\Omega_0) \subseteq \mathscr{Y}_0$ . Let  $\mathscr{Y}^*$ be the dual of the Banach space  $\mathscr{Y}$ , and by the Hahn-Banach extension theorem we obtain

$$\|y\| = \sup_{\|y^*\|_{\mathscr{Y}^*} \le 1} |y^*(y)|, \ \forall \ y \in \mathscr{Y}.$$

Thus, for the closed ball  $B_{\mathscr{Y}_0}(y_0, r) := \{y \in \mathscr{Y}_0 : ||y - y_0|| \le r\}$  in the Banach space  $\mathscr{Y}_0$ , it follows from Proposition B.1.10 in [67] that there exists a unit sequence  $\{y_n^*, n \ge 1\}$  such that

$$(Tf)^{-1}(B(y_0, r))$$

$$= \{ \omega \in \Omega : ||T(\omega)f(\omega) - y_0|| \le r \}$$

$$= \bigcap_{\substack{n=1\\\infty}} \{ \omega \in \Omega : |y_n^*(T(\omega)f(\omega) - y_0)| \le r \}$$

$$= \bigcap_{n=1}^{\infty} \{ \omega \in \Omega : \lim_{\substack{m \to \infty}} |y_n^*(T(\omega)f_m(\omega) - y_0)| \le r \}.$$
(B.1)

From the analysis in Step 1, we know that  $Tf_n \in (\bigvee_{x \in \mathscr{X}} \sigma(Tx; \tau_N(\mathscr{Y}))) \bigvee \sigma(f; \tau_N(\mathscr{X}))$ . Noting that  $y_n^*$  is a linear measurable functional mapping the Banach space  $\mathscr{Y}$  to  $\mathbb{R}$ , we have

$$\left\{ \begin{split} & \left\{ \omega \in \Omega : \lim_{m \to \infty} |y_n^*(T(\omega)f_m(\omega) - y_0)| \le r \right\} \\ & \in \left( \bigvee_{x \in \mathscr{X}} \sigma(Tx; \tau_{\mathsf{N}}(\mathscr{Y})) \right) \bigvee \sigma(f; \tau_{\mathsf{N}}(\mathscr{X})). \end{split}$$
(B.2)

Combining (B.1)-(B.2) yields  $(Tf)^{-1}(B(y_0, r)) \in (\bigvee_{x \in \mathscr{X}} \sigma(Tx; \tau_N(\mathscr{Y}))) \bigvee \sigma(f; \tau_N(\mathscr{X}))$ , and since the open set of the Banach space  $(\mathscr{Y}_0, \tau_N(\mathscr{Y}_0))$  is generated by a countable number of closed spheres, we have

$$Tf|_{\mathscr{Y}_0} \in \left(\bigvee_{x \in \mathscr{X}} \sigma(Tx; \tau_{\mathbf{N}}(\mathscr{Y}))\right) \bigvee \sigma(f; \tau_{\mathbf{N}}(\mathscr{X})).$$

Given  $B \in \mathscr{B}(\mathscr{Y}; \tau_{\mathbb{N}}(\mathscr{Y}))$ , noting that  $B_0 = B \cap \mathscr{Y}_0$ , it is known that  $B_0 \in \mathscr{B}(\mathscr{Y}_0; \tau_{\mathbb{N}}(\mathscr{Y}_0))$  and

$$(Tf)^{-1}(B) = \{ \omega \in \Omega : T(\omega)f(\omega) \in B \} = \{ \omega \in \Omega : T(\omega)f(\omega) \in B_0 \} = (Tf)^{-1}(B_0),$$

which leads to  $Tf \in \left(\bigvee_{x \in \mathscr{X}} \sigma(Tx; \tau_{N}(\mathscr{Y}))\right) \bigvee \sigma(f; \tau_{N}(\mathscr{X})).$ 

Proof of Proposition 2.5. Since the random elements with values in the Banach space  $(\mathscr{L}(\mathscr{Y}, \mathscr{Z}), \tau_{N}(\mathscr{L}(\mathscr{Y}, \mathscr{Z})))$  are certainly the random elements with values in  $(\mathscr{L}(\mathscr{Y}, \mathscr{Z}), \tau_{N}(\mathscr{L}(\mathscr{Y}, \mathscr{Z})))$ ,

it follows that  $f \in L^1(\Omega; \mathscr{L}(\mathscr{Y}, \mathscr{Z}))$  implies  $fy \in L^1(\Omega; \mathscr{Z}), \forall y \in \mathscr{Y}$  by Proposition (A.5).(a). Considering the simple function  $\sum_{i=1}^n \mathbf{1}_{A_i} \otimes T_i$  with values in Banach space  $\mathscr{L}(\mathscr{Y}, \mathscr{Z})$ , where  $A_i \in \mathcal{F}, T_i \in \mathscr{L}(\mathscr{Y}, \mathscr{Z}), A_i \cap A_j = \emptyset, 1 \leq i \neq j \leq n$ , we have

$$\mathbb{E}\left[\left(\mathbf{1}_{A_i} \otimes T_i\right)y\right] = \mathbb{P}(A)(T_i y) = (\mathbb{P}(A_i)T_i)y = \mathbb{E}\left[\mathbf{1}_{A_i} \otimes T_i\right]y, \ 1 \le i \le n,$$

which leads to

$$\mathbb{E}\left[\left(\sum_{i=1}^{n} \mathbf{1}_{A_{i}} \otimes T_{i}\right) y\right] = \sum_{i=1}^{n} \mathbb{E}\left[(\mathbf{1}_{A_{i}} \otimes T_{i})y\right]$$
$$= \sum_{i=1}^{n} \mathbb{E}\left[\mathbf{1}_{A_{i}} \otimes T_{i}\right] y$$
$$= \mathbb{E}\left[\sum_{i=1}^{n} \mathbf{1}_{A_{i}} \otimes T_{i}\right] y.$$
(B.3)

Since  $f \in L^1(\Omega; \mathscr{L}(\mathscr{Y}, \mathscr{Z}))$  is the random element with values in the Banach space  $(\mathscr{L}(\mathscr{Y}, \mathscr{Z}), \tau_N(\mathscr{L}(\mathscr{Y}, \mathscr{Z})))$ , it follows from Lemma A.4 and Definition A.3 that there exists a sequence of simple functions  $\{T_n, n \ge 1\}$  such that  $\lim_{n\to\infty} \|f - T_n\| = 0$  a.s. and  $\|T_n\| \le \|f\|$  a.s. Noting that  $\|f - T_n\| \le 2\|f\| \in L^1(\Omega)$  and  $\lim_{n\to\infty} \|fy - T_ny\| = 0$  a.s.,  $\forall y \in \mathscr{Y}$ , by the dominated convergence theorem, it follows that  $\lim_{n\to\infty} \mathbb{E}[(f - T_n)y] = 0, \forall y \in \mathscr{Y}$ , and  $\lim_{n\to\infty} \mathbb{E}[T_n] = \mathbb{E}[f]$ . Thus, it follows from (B.3) that

$$\mathbb{E}[fy] = \lim_{n \to \infty} \mathbb{E}[(f - T_n)y] + \lim_{n \to \infty} \mathbb{E}[T_n y] = \lim_{n \to \infty} \mathbb{E}[T_n]y = \mathbb{E}[f]y, \ \forall \ y \in \mathscr{Y}.$$

Proof of Proposition 2.6. Since the random elements with values in the Banach space  $(\mathscr{L}(\mathscr{Y}, \mathscr{Z}), \tau_{N}(\mathscr{L}(\mathscr{Y}, \mathscr{Z})))$  are certainly the random elements with values in the topological space  $(\mathscr{L}(\mathscr{Y}, \mathscr{Z}), \tau_{S}(\mathscr{L}(\mathscr{Y}, \mathscr{Z})))$ , it follows from Proposition A.5.(b) that fy is the random element with values in the Banach space  $(\mathscr{L}, \tau_{N}(\mathscr{Z})), f \in L^{2}(\Omega; \mathscr{L}(\mathscr{Y}, \mathscr{Z}))$ , and  $y \in L^{2}(\Omega; \mathscr{Y})$  implies  $fy \in L^{1}(\Omega; \mathscr{Z})$ . Consider the simple function  $\sum_{i=1}^{n} \mathbf{1}_{A_{i}} \otimes y_{i}$  with values in the Banach space  $\mathscr{Y}$ , where  $A_{i} \in \mathscr{G}, y_{i} \in \mathscr{Y}, A_{i} \cap A_{j} = \emptyset, 1 \leq i \neq j \leq n$ . For any given  $F \in \mathscr{G}$  and  $1 \leq i \leq n$ , by Lemma A.9, we have

$$\int_{F} \mathbb{E}[f(\mathbf{1}_{A_{i}} \otimes y_{i})|\mathscr{G}] \, \mathrm{d}\mathbb{P} = \int_{F} f(\mathbf{1}_{A_{i}} \otimes y_{i}) \, \mathrm{d}\mathbb{P} = \int_{F \cap A_{i}} fy_{i} \, \mathrm{d}\mathbb{P} = \mathbb{E}[(\mathbf{1}_{F \cap A_{i}} f)y_{i}]. \tag{B.4}$$

Noting that  $F \cap A_i \in \mathcal{G}$ , by Lemma A.9 and Proposition 2.5, we get

$$\mathbb{E}[(\mathbf{1}_{F\cap A_i}f)y_i] = \mathbb{E}[\mathbf{1}_{F\cap A_i}f]y_i = \left(\int_{F\cap A_i}f\,\mathrm{d}\mathbb{P}\right)y_i = \left(\int_{F\cap A_i}\mathbb{E}[f|\mathscr{G}]\,\mathrm{d}\mathbb{P}\right)y_i.$$

31

Noting that  $\mathbf{1}_{F \cap A_i} \mathbb{E}[f|\mathscr{G}] \in L^2(\Omega, \mathscr{G}; \mathscr{L}(\mathscr{G}, \mathscr{Z}))$ , it follows from Lemma A.9 and Proposition 2.5 that

$$\left(\int_{F\cap A_i} \mathbb{E}[f|\mathscr{G}] \, \mathrm{d}\mathbb{P}\right) y_i = \mathbb{E}\left[\mathbf{1}_{F\cap A_i} \mathbb{E}[f|\mathscr{G}] y_i\right] = \int_F \mathbb{E}[f|\mathscr{G}](\mathbf{1}_{A_i} \otimes y_i) \, \mathrm{d}\mathbb{P}. \tag{B.5}$$

Noting that  $\mathbb{E}[f|\mathscr{G}](\mathbf{1}_{A_i} \otimes y_i) \in L^0(\Omega, \mathscr{G}; \mathscr{Z})$ , it follows from (B.4)-(B.5) and Lemma A.9 that

$$\mathbb{E}\left[f\left(\sum_{i=1}^{n} \mathbf{1}_{A_{i}} \otimes y_{i}\right) \middle| \mathscr{G}\right] = \mathbb{E}[f|\mathscr{G}]\left(\sum_{i=1}^{n} \mathbf{1}_{A_{i}} \otimes y_{i}\right) \text{ a.s.}$$
(B.6)

Given the random element  $y \in L^2(\Omega, \mathscr{G}; \mathscr{Y})$  with values in the Banach space  $(\mathscr{Y}, \tau_N(\mathscr{Y}))$ , there exists a sequence of simple functions  $\{y_n \in L^0(\Omega, \mathscr{G}; \mathscr{Y}), n \geq 1\}$  with values in the Banach space  $\mathscr{Y}$  such that  $\lim_{n\to\infty} ||y - y_n|| = 0$  a.s. and  $||y_n|| \leq ||y||$  a.s. Noting that  $||fy_n|| \leq$  $||f||^2 + ||y||^2 \in L^1(\Omega)$  and  $\mathbb{E}[f|\mathcal{G}] \in L^2(\Omega, \mathscr{G}; \mathscr{L}(\mathscr{Y}, \mathscr{Z}))$ , by the dominated convergence theorem of conditional expectation and (B.6), we have

$$\mathbb{E}[fy|\mathscr{G}] = \mathbb{E}\left[f\left(\lim_{n \to \infty} y_n\right) \middle| \mathscr{G}\right] = \lim_{n \to \infty} \mathbb{E}\left[fy_n \middle| \mathscr{G}\right] = \mathbb{E}\left[f \middle| \mathscr{G}\right] \left(\lim_{n \to \infty} y_n\right) = \mathbb{E}\left[f \middle| \mathscr{G}\right] y \text{ a.s.}$$

Proof of Proposition 2.9. Given an integer 
$$k \geq 2$$
, denote  $\mathscr{F}^k = \bigvee_{i=k}^{\infty} \sigma(f_i; \tau(\mathscr{X}_1)), \mathscr{G}^k = \bigvee_{i=k}^{\infty} \sigma(g_i; \tau(\mathscr{X}_2)), \mathscr{F}_k = \bigvee_{i=1}^{k-1} \sigma(f_i; \tau(\mathscr{X}_1)), \mathscr{G}_k = \bigvee_{i=1}^{k-1} \sigma(g_i; \tau(\mathscr{X}_2))$  and  
 $\mathscr{F}(k) = \bigvee_{i=1}^{k-1} (\sigma(f_i; \tau(\mathscr{X}_1)) \bigvee \sigma(g_i; \tau(\mathscr{X}_2))).$ 

Let  $E \in \mathscr{F}^k$ ,  $F \in \mathscr{G}^k$ ,  $A \in \mathscr{F}_k$  and  $B \in \mathscr{G}_k$ . On the one hand, noting that  $\{f_k, k \ge 1\}$  and  $\{g_k, k \ge 1\}$  are mutually independent, we know that  $\bigvee_{i=1}^{\infty} \sigma(f_i; \tau(\mathscr{X}_1))$  and  $\bigvee_{i=1}^{\infty} \sigma(g_i; \tau(\mathscr{X}_2))$  are also mutually independent. Since  $A \cap E \in \bigvee_{i=1}^{\infty} \sigma(f_i; \tau(\mathscr{X}_1))$  and  $B \cap F \in \bigvee_{i=1}^{\infty} \sigma(g_i; \tau(\mathscr{X}_2))$ , we get  $\mathbb{P}(E \cap F \cap A \cap B) = \mathbb{P}(E \cap A)\mathbb{P}(F \cap B)$ , which together with  $A \cap B \in \mathscr{F}(k)$  gives

$$\int_{A\cap B} \mathbb{P}(E\cap F|\mathscr{F}(k)) \, \mathrm{d}\mathbb{P} = \mathbb{P}(E\cap F\cap A\cap B) = \mathbb{P}(E\cap A)\mathbb{P}(F\cap B). \tag{B.7}$$

On the other hand, noting that  $\mathbf{1}_F \in \mathscr{G}^k$ ,  $\mathscr{F}(k) = \mathscr{F}_k \bigvee \mathscr{G}_k$  and  $\mathscr{G}^k \bigvee \mathscr{G}_k = \bigvee_{i=1}^{\infty} \sigma(g_i; \tau(\mathscr{X}_2))$ is independent of  $\mathscr{F}_k$ , by Corollary 7.3.3 in [85], we obtain  $\mathbb{E}[\mathbf{1}_F|\mathscr{F}(k)] = \mathbb{E}[\mathbf{1}_F|\mathscr{G}_k]$ , which further implies

$$\mathbb{E}[\mathbf{1}_E|\mathscr{F}(k)]\mathbb{E}[\mathbf{1}_F|\mathscr{F}(k)] = \mathbb{E}[\mathbf{1}_E\mathbb{E}[\mathbf{1}_F|\mathscr{F}(k)]|\mathscr{F}(k)] = \mathbb{E}[\mathbf{1}_E\mathbb{E}[\mathbf{1}_F|\mathscr{G}_k]|\mathscr{F}(k)],$$

from which we get

$$\int_{A \cap B} \mathbb{P}(E|\mathscr{F}(k)) \mathbb{P}(F|\mathscr{F}(k)) \, \mathrm{d}\mathbb{P} = \int_{A \cap B} \mathbb{E}[\mathbf{1}_E \mathbb{E}[\mathbf{1}_F|\mathscr{G}_k]|\mathscr{F}(k)] \, \mathrm{d}\mathbb{P}$$

$$= \int_{A \cap B} \mathbf{1}_E \mathbb{E}[\mathbf{1}_F | \mathscr{G}_k] \, \mathrm{d}\mathbb{P}. \tag{B.8}$$

Noting that  $\mathbf{1}_B \in \mathscr{G}_k$ ,  $\mathbf{1}_{A \cap E}$  and  $\mathbb{E}[\mathbf{1}_{B \cap F} | \mathscr{G}_k]$  are mutually independent, we have

$$\int_{A \cap B} \mathbf{1}_{E} \mathbb{E}[\mathbf{1}_{F} | \mathscr{G}_{k}] \, \mathrm{d}\mathbb{P} = \mathbb{E}[\mathbf{1}_{A \cap E} \mathbb{E}[\mathbf{1}_{B \cap F} | \mathscr{G}_{k}]]$$
$$= \mathbb{E}[\mathbf{1}_{A \cap E}] \mathbb{E}[\mathbf{1}_{B \cap F}]$$
$$= \mathbb{P}(A \cap E) \mathbb{P}(B \cap F). \tag{B.9}$$

It follows from (B.8)-(B.9) that

$$\int_{A \cap B} \mathbb{P}(E|\mathscr{F}(k))\mathbb{P}(F|\mathscr{F}(k)) \, \mathrm{d}\mathbb{P} = \mathbb{P}(A \cap E)\mathbb{P}(B \cap F). \tag{B.10}$$

Combining (B.7) and (B.10) gives

$$\int_{A\cap B} \mathbb{P}(E\cap F|\mathscr{F}(k)) \, \mathrm{d}\mathbb{P} = \int_{A\cap B} \mathbb{P}(E|\mathscr{F}(k))\mathbb{P}(F|\mathscr{F}(k)) \, \mathrm{d}\mathbb{P}. \tag{B.11}$$

Denote

$$\Pi = \left\{ \bigcup_{i=1}^{n} (A_i \cap B_i) \middle| A_i \in \mathscr{F}_k, B_i \in \mathscr{G}_k, A_i \cap B_i \cap A_j \cap B_j = \emptyset, 1 \le i \ne j \le n \right\}.$$

If  $\bigcup_{i=1}^{n} (A_i \cap B_i) \in \Pi$  and  $\bigcup_{i=1}^{m} (C_i \cap D_i) \in \Pi$ , then we have

$$\begin{pmatrix} \bigcup_{i=1}^{n} (A_i \cap B_i) \end{pmatrix} \bigcap \left( \bigcup_{i=1}^{m} (C_i \cap D_i) \right) \\
= \bigcup_{j=1}^{m} \left( \left( \bigcup_{i=1}^{n} (A_i \cap B_i) \right) \cap C_j \cap D_j \right) \\
= \bigcup_{j=1}^{m} \bigcup_{i=1}^{n} (A_i \cap B_i \cap C_j \cap D_j), \quad (B.12)$$

noting that  $A_i \cap B_i \cap A_j \cap B_j = \emptyset$ ,  $1 \le i \ne j \le n$ , and  $C_s \cap D_s \cap C_t \cap D_t = \emptyset$ ,  $1 \le s \ne t \le m$ , we know that  $A_i \cap B_i \cap A_j \cap B_j \cap C_s \cap D_s \cap C_t \cap D_t = \emptyset$ ,  $(i, j) \ne (s, t)$ , which together with (B.12) gives

$$\left(\bigcup_{i=1}^{n} (A_i \cap B_i)\right) \bigcap \left(\bigcup_{i=1}^{m} (C_i \cap D_i)\right) \in \Pi,$$

thus, we conclude that  $\Pi$  is a  $\pi$ -class, i.e.,  $\Pi$  is closed under the intersection operation of the set. Denote

$$\Lambda = \left\{ M \in \mathscr{F}(k) \middle| \int_M \mathbb{P}(E \cap F | \mathscr{F}(k)) \, \mathrm{d}\mathbb{P} = \int_M \mathbb{P}(E | \mathscr{F}(k)) \mathbb{P}(F | \mathscr{F}(k)) \, \mathrm{d}\mathbb{P} \right\}$$

Noting that  $\Pi$  is composed of finite disjoint union of elements in  $\mathscr{F}_k \cap \mathscr{G}_k$ , it follows from (B.11) that  $\Omega \in \Lambda$ . If  $\bigcup_{i=1}^n (A_i \cap B_i) \in \Pi$ , then we have

$$\int_{\bigcup_{i=1}^{n} (A_i \cap B_i)} \mathbb{P}(E \cap F | \mathscr{F}(k)) \, \mathrm{d}\mathbb{P} = \int_{\bigcup_{i=1}^{n} (A_i \cap B_i)} \mathbb{P}(E | \mathscr{F}(k)) \mathbb{P}(F | \mathscr{F}(k)) \, \mathrm{d}\mathbb{P},$$

which shows that  $\bigcup_{i=1}^{n} (A_i \cap B_i) \in \Lambda$  and further gives  $\Pi \subseteq \Lambda$ . If  $M_i \in \Lambda$  and  $M_i \subseteq M_{i+1}$ , then we obtain

$$\int_{\Omega} \mathbf{1}_{M_i} \mathbb{P}(E \cap F | \mathscr{F}(k)) \, \mathrm{d}\mathbb{P} = \int_{\Omega} \mathbf{1}_{M_i} \mathbb{P}(E | \mathscr{F}(k)) \mathbb{P}(F | \mathscr{F}(k)) \, \mathrm{d}\mathbb{P}, \ \forall i \ge 1.$$

Noting that  $M_i \subseteq M_{i+1}$  implies  $\lim_{i\to\infty} \mathbf{1}_{M_i} = \mathbf{1}_{\bigcup_{i=1}^{\infty} M_i}$ , by Lebesgue dominated convergence theorem, we get

$$\int_{\Omega} \mathbf{1}_{\bigcup_{i=1}^{\infty} M_i} \mathbb{P}(E \cap F | \mathscr{F}(k)) \, \mathrm{d}\mathbb{P} = \int_{\Omega} \mathbf{1}_{\bigcup_{i=1}^{\infty} M_i} \mathbb{P}(E | \mathscr{F}(k)) \mathbb{P}(F | \mathscr{F}(k)) \, \mathrm{d}\mathbb{P}$$

Thus, we have  $\bigcup_{i=1}^{\infty} M_i \in \Lambda$ . If  $M_1, M_2 \in \Lambda$  and  $M_1 \subseteq M_2$ , then  $M_2 = M_1 \cup (M_2 \setminus M_1)$  implies

$$\left(\int_{M_1} + \int_{M_2 \setminus M_1}\right) \mathbb{P}(E \cap F | \mathscr{F}(k)) \, \mathrm{d}\mathbb{P} = \left(\int_{M_1} + \int_{M_2 \setminus M_1}\right) \mathbb{P}(E | \mathscr{F}(k)) \mathbb{P}(F | \mathscr{F}(k)) \, \mathrm{d}\mathbb{P}.$$

It follows from  $M_1 \in \Lambda$  that

$$\int_{M_2 \setminus M_1} \mathbb{P}(E \cap F | \mathscr{F}(k)) \, \mathrm{d}\mathbb{P} = \int_{M_2 \setminus M_1} \mathbb{P}(E | \mathscr{F}(k)) \mathbb{P}(F | \mathscr{F}(k)) \, \mathrm{d}\mathbb{P},$$

which shows that  $M_2 \setminus M_1 \in \Lambda$ . From the above analysis, we know that  $\Lambda$  is a  $\lambda$ -class, by Theorem 1.3.2 in [85], we get  $\sigma(\Pi) \subseteq \Lambda$ . On the one hand, it follows from  $\Pi \subseteq \mathscr{F}(k) = \mathscr{F}_k \bigvee \mathscr{G}_k$  that  $\sigma(\Pi) \subseteq \mathscr{F}(k)$ . On the other hand, noting that  $\mathscr{F}_k \subseteq \Pi$  and  $\mathscr{G}_k \subseteq \Pi$ , we have  $\mathscr{F}_k \cup \mathscr{G}_k \subseteq \sigma(\Pi)$ , which leads to  $\mathscr{F}(k) = \mathscr{F}_k \bigvee \mathscr{G}_k \subseteq \sigma(\Pi)$  and further gives  $\mathscr{F}(k) = \sigma(\Pi) \subseteq \Lambda$ . Noting that  $E \in \mathscr{F}^k$  and  $F \in \mathscr{G}^k$ , we obtain

$$\int_{M} \mathbb{P}(E \cap F | \mathscr{F}(k)) \, \mathrm{d}\mathbb{P} = \int_{M} \mathbb{P}(E | \mathscr{F}(k)) \mathbb{P}(F | \mathscr{F}(k)) \, \mathrm{d}\mathbb{P}, \ \forall M \in \mathscr{F}(k),$$

which together with  $\mathbb{P}(E|\mathscr{F}(k))\mathbb{P}(F|\mathscr{F}(k)) \in \mathscr{F}(k)$  gives

$$\mathbb{P}(E\cap F|\mathscr{F}(k))=\mathbb{P}(E|\mathscr{F}(k))\mathbb{P}(F|\mathscr{F}(k))$$
 a.s.,

thus, we conclude that  $\mathscr{F}^k$  is conditionally independent of  $\mathscr{G}^k$  with respect to  $\mathscr{F}(k)$ .

Before proving Proposition 2.10, we introduce the following lemma.

**Lemma B.1.** Let  $\mathscr{G}$  be a sub- $\sigma$ -algebra of  $\mathcal{F}$ ,  $A \in \mathcal{F}$  and  $f \in L^1(\Omega; \mathscr{X})$ . If f is conditionally independent of  $\mathbf{1}_A$  with respect to  $\mathscr{G}$ , then

$$\mathbb{E}[f\mathbf{1}_A|\mathscr{G}] = \mathbb{E}[f|\mathscr{G}]\mathbb{E}[\mathbf{1}_A|\mathscr{G}] \text{ a.s.}$$

*Proof.* Let  $f = \mathbf{1}_B \otimes x$ , where  $B \in \mathcal{F}$  and  $x \in \mathscr{X}$ . Noting that

$$\sigma(\mathbf{1}_B \otimes x; \tau_{\mathsf{N}}(\mathscr{X})) = \left\{ (\mathbf{1}_B \otimes x)^{-1}(E) : E \in \mathscr{B}(\mathscr{X}; \tau_{\mathsf{N}}(\mathscr{X})) \right\},$$
(B.13)

it follows that

$$(\mathbf{1}_{B} \otimes x)^{-1}(E) = \begin{cases} \Omega, & \text{If } 0 \in E, x \in E; \\ B, & \text{If } 0 \in E, x \notin E; \\ B^{\complement}, & \text{If } 0 \notin E, x \in E; \\ \emptyset, & \text{If } 0 \notin E, x \notin E. \end{cases}, \forall E \in \mathscr{B}(\mathscr{X}; \tau_{N}(\mathscr{X})), \tag{B.14}$$

which leads to  $\sigma(\mathbf{1}_B \otimes x; \tau_N(\mathscr{X})) = \sigma(\mathbf{1}_B; \tau_N(\mathscr{X}))$ . Since  $\mathbf{1}_B \otimes x$  is conditionally independent of  $\mathbf{1}_A$  with respect to  $\mathscr{G}$ , it follows from Definition 2.8 that  $\sigma(\mathbf{1}_B \otimes x; \tau_N(\mathscr{X}))$  is conditionally independent of  $\sigma(\mathbf{1}_A; \tau_N(\mathscr{X}))$  with respect to  $\mathscr{G}$ , from which we can further conclude that  $\sigma(B)$ is conditionally independent of  $\sigma(A)$  with respect to  $\mathscr{G}$ . Given  $F \in \mathscr{G}$ , on the one hand, we have

$$\int_{F} \mathbb{E}[f\mathbf{1}_{A}|\mathscr{G}] \, d\mathbb{P} = x \int_{F} \mathbb{E}[\mathbf{1}_{A\cap B}|\mathscr{G}] \, d\mathbb{P}$$
$$= x \int_{F} \mathbb{P}(A \cap B|\mathscr{G}) \, d\mathbb{P}$$
$$= x \int_{F} \mathbb{P}(A|\mathscr{G})\mathbb{P}(B|\mathscr{G}) \, d\mathbb{P}.$$
(B.15)

On the other hand, noting that  $\mathbb{E}[\mathbf{1}_B \otimes x | \mathscr{G}] = \mathbb{E}[\mathbf{1}_B | \mathscr{G}] \otimes x$  a.s., we get

$$\int_{F} \mathbb{E}[f|\mathscr{G}]\mathbb{E}[\mathbf{1}_{A}|\mathscr{G}] \, \mathrm{d}\mathbb{P} = \int_{F} \mathbb{E}[\mathbf{1}_{B} \otimes x|\mathscr{G}]\mathbb{E}[\mathbf{1}_{A}|\mathscr{G}] \, \mathrm{d}\mathbb{P}$$
$$= x \int_{F} \mathbb{P}(A|\mathscr{G})\mathbb{P}(B|\mathscr{G}) \, \mathrm{d}\mathbb{P}.$$
(B.16)

Then, by (B.15)-(B.16), we obtain

$$\mathbb{E}[(\mathbf{1}_B \otimes x)\mathbf{1}_A|\mathscr{G}] = \mathbb{E}[\mathbf{1}_B \otimes x|\mathscr{G}]\mathbb{E}[\mathbf{1}_A|\mathscr{G}] \text{ a.s.}$$
(B.17)

Let  $f = \sum_{i=1}^{n} \mathbf{1}_{B_i} \otimes x_i$ , where  $B_i \in \mathcal{F}$ ,  $B_i \cap B_j = \emptyset$ ,  $x_i \in \mathscr{X}$ ,  $1 \le i \ne j \le n$ . Following the same way as (B.13)-(B.14), we have

$$\sigma\left(\sum_{i=1}^{n} \mathbf{1}_{B_{i}} \otimes x_{i}; \tau_{\mathsf{N}}(\mathscr{X})\right) = \sigma\left(\bigcup_{i=1}^{n} B_{i}\right).$$

Thus, f is conditionally independent of  $\mathbf{1}_A$  with respect to  $\mathscr{G}$  implies that  $\sigma(\bigcup_{i=1}^n B_i)$  is conditionally independent of  $\sigma(A)$  with respect to  $\mathscr{G}$ , from which we know that  $B_i$  is conditionally independent of A with respect to  $\mathscr{G}$ ,  $1 \le i \le n$ . It follows from (B.17) that

$$\int_{F} \mathbb{E}\left[\sum_{i=1}^{n} (\mathbf{1}_{B_{i}} \otimes x_{i}) \mathbf{1}_{A} \middle| \mathscr{G}\right] d\mathbb{P} = \sum_{i=1}^{n} \int_{F} \mathbb{E}\left[\mathbf{1}_{B_{i}} \otimes x \middle| \mathscr{G}\right] \mathbb{E}[\mathbf{1}_{A} \middle| \mathscr{G}] d\mathbb{P}, \tag{B.18}$$

which leads to

$$\mathbb{E}\left[\left.\sum_{i=1}^{n} (\mathbf{1}_{B_{i}} \otimes x_{i}) \mathbf{1}_{A} \right| \mathscr{G}\right] = \mathbb{E}\left[\left.\sum_{i=1}^{n} \mathbf{1}_{B_{i}} \otimes x \right| \mathscr{G}\right] \mathbb{E}[\mathbf{1}_{A}| \mathscr{G}] \text{ a.s.}$$

For the random element f with values in Banach space  $(\mathscr{X}, \tau_{N}(\mathscr{X}))$ , there exists a sequence of simple functions with values in  $\mathscr{X}$  such that  $\lim_{n\to\infty} f_n = f$  a.s. and  $||f_n|| \le ||f||$  a.s. Noting that  $||f_n \mathbf{1}_A|| \le ||f||$  a.s. together with  $f \in L^1(\Omega; \mathscr{X})$  implies that  $\mathbb{E}[||f||] < \infty$ , by the dominated convergence theorem, we get

$$\int_{F} \mathbb{E} \left[ \lim_{n \to \infty} f_{n} \mathbf{1}_{A} \middle| \mathscr{G} \right] d\mathbb{P} = \int_{F} \lim_{n \to \infty} \mathbb{E} [f_{n} \mathbf{1}_{A} \middle| \mathscr{G}] d\mathbb{P}, \tag{B.19}$$

and

$$\int_{F} \mathbb{E} \left[ \lim_{n \to \infty} f_n \middle| \mathscr{G} \right] \mathbb{E}[\mathbf{1}_A | \mathscr{G}] \, \mathrm{d}\mathbb{P} = \int_{F} \lim_{n \to \infty} \mathbb{E}[f_n | \mathscr{G}] \mathbb{E}[\mathbf{1}_A | \mathscr{G}] \, \mathrm{d}\mathbb{P}.$$

It follows from (B.18) that

$$\int_{F} \mathbb{E}[f_{n}\mathbf{1}_{A}\mathscr{G}] \, \mathrm{d}\mathbb{P} = \int_{F} \mathbb{E}[f_{n}|\mathscr{G}]\mathbb{E}[\mathbf{1}_{A}|\mathscr{G}] \, \mathrm{d}\mathbb{P}.$$
(B.20)

Thus, by (B.19)-(B.20), we have

$$\int_{F} \mathbb{E}[f\mathbf{1}_{A}|\mathscr{G}] \, \mathrm{d}\mathbb{P} = \int_{F} \mathbb{E}[f|\mathscr{G}]\mathbb{E}[\mathbf{1}_{A}|\mathscr{G}] \, \mathrm{d}\mathbb{P}, \,\, \forall F \in \mathscr{G}.$$

Proof of Proposition 2.10. We first consider the case with  $f = \sum_{i=1}^{n} \mathbf{1}_{A_i} \otimes x_i$ , where  $A_i \in \mathcal{F}$ ,  $A_i \cap A_j = \emptyset$ ,  $x_i \in \mathscr{X}$ ,  $1 \le i \ne j \le n$ . Since T is conditionally independent of f with respect to  $\mathscr{G}$ , it follows that  $\sigma(Tx_i; \tau_N(\mathscr{G}))$  is conditionally independent of  $\sigma(\sum_{i=1}^{n} \mathbf{1}_{A_i} \otimes x_i; \tau_N(\mathscr{X}))$  with respect to  $\mathscr{G}$ ,  $1 \le i \le n$ , from which we know that  $\sigma(Tx_i; \tau_N(\mathscr{G}))$  is conditionally independent of  $\mathbf{1}_{A_i}$  with respect to  $\mathscr{G}$ , thus,  $Tx_i$  is conditionally independent of  $\mathbf{1}_{A_i}$  with respect to  $\mathscr{G}$ , then by Lemma B.1, we have

$$\mathbb{E}[T(\mathbf{1}_{A_i} \otimes x_i)|\mathscr{G}] = \mathbb{E}[(Tx_i)\mathbf{1}_{A_i}|\mathscr{G}]$$
  
$$= \mathbb{E}[Tx_i|\mathscr{G}]\mathbb{E}[\mathbf{1}_{A_i}|\mathscr{G}]$$
  
$$= \mathbb{E}[(Tx_i)\mathbb{E}[\mathbf{1}_{A_i}|\mathscr{G}]|\mathscr{G}] \text{ a.s.}$$
(B.21)

Noting that

$$\mathbb{E}[(Tx_i)\mathbb{E}[\mathbf{1}_{A_i}|\mathscr{G}]|\mathscr{G}] = \mathbb{E}[T(\mathbb{E}[\mathbf{1}_{A_i}|\mathscr{G}] \otimes x_i)|\mathscr{G}] = \mathbb{E}[T\mathbb{E}[\mathbf{1}_{A_i} \otimes x_i|\mathscr{G}]|\mathscr{G}] \text{ a.s..}$$
(B.22)

36

By (B.21)-(B.22), we have

$$\mathbb{E}[Tf|\mathscr{G}] = \sum_{\substack{i=1\\n}}^{n} \mathbb{E}[T(\mathbf{1}_{A_i} \otimes x_i)|\mathscr{G}] \\ = \sum_{\substack{i=1\\i\in I}}^{n} \mathbb{E}[T\mathbb{E}[\mathbf{1}_{A_i} \otimes x_i|\mathscr{G}]|\mathscr{G}] \\ = \mathbb{E}[T\mathbb{E}[f|\mathscr{G}]|\mathscr{G}] \text{ a.s.}$$
(B.23)

Given the random element  $f \in L^2(\Omega; \mathscr{X})$  with values in Banach space  $(\mathscr{X}, \tau_N(\mathscr{X}))$ , by Lemma A.4 and Definition A.3, we know that there exists a sequence  $\{f_n, n \ge 1\}$  of simple functions, such that  $\lim_{n\to\infty} f_n = f$  a.s. and  $||f_n|| \le ||f||$  a.s. If  $\mathbb{E}[||T||^2] < \infty$  and  $f \in L^2(\Omega; \mathscr{X})$ , noting that  $T(\omega) \in \mathscr{L}(\mathscr{X}, \mathscr{Y})$ , we get  $||(Tf_n)(\omega)|| = ||T(\omega)f_n(\omega)|| \le ||T(\omega)|| ||f_n(\omega)|| \le$  $||T(\omega)||^2 + ||f(\omega)||^2 \in L^2(\Omega)$ , which together with conditional dominated convergence theorem gives

$$\mathbb{E}[Tf|\mathscr{G}] = \mathbb{E}\left[T\left(\lim_{n \to \infty} f_n\right) \middle| \mathscr{G}\right] = \mathbb{E}\left[\lim_{n \to \infty} Tf_n \middle| \mathscr{G}\right] = \lim_{n \to \infty} \mathbb{E}[Tf_n|\mathscr{G}] \text{ a.s.}$$
(B.24)

By (B.23), we obtain

$$\mathbb{E}[Tf_n|\mathscr{G}] = \mathbb{E}[T\mathbb{E}[f_n|\mathscr{G}]|\mathscr{G}] \text{ a.s.}$$

It follows from  $||T\mathbb{E}[f_n|\mathscr{G}]|| \leq ||T|| ||\mathbb{E}[f_n|\mathscr{G}]|| \leq ||T||^2 + \mathbb{E}[||f_n||^2|\mathscr{G}] \leq ||T||^2 + \mathbb{E}[||f||^2|\mathscr{G}] \in L^2(\Omega)$  and conditional dominated convergence theorem that

$$\lim_{n \to \infty} \mathbb{E}[T\mathbb{E}[f_n|\mathscr{G}]|\mathscr{G}] = \mathbb{E}\left[\lim_{n \to \infty} T\mathbb{E}[f_n|\mathscr{G}]\Big|\mathscr{G}\right] = \mathbb{E}\left[T\mathbb{E}\left[\lim_{n \to \infty} f_n\Big|\mathscr{G}\right]\Big|\mathscr{G}\right] \text{ a.s.}$$
(B.25)

Hence, by (B.24)-(B.25), we get  $\mathbb{E}[Tf|\mathscr{G}] = \mathbb{E}[T\mathbb{E}[f|\mathscr{G}]|\mathscr{G}]$  a.s.

# APPENDIX C

## **PROOFS OF SECTIONS 4-5**

For the convenience of notational writing without giving rise to ambiguity, the subscripts of the parametrization in Banach spaces and the subscripts of the inner product in Hilbert spaces will be omitted in the sequel.

*Proof of Theorem 4.4.* Given the initial value  $x(0) \in \mathscr{X}_1$ , by Proposition A.5.(a)-(c), we know that x(k) is a random element with values in the Hilbert space  $(\mathscr{X}_1, \tau_N(\mathscr{X}_1))$ . It follows from the random difference equation (4.2) that

$$x(k+1) = \left(\prod_{i=0}^{k} (I_{\mathscr{X}_1} - F(i))\right) x(0)$$

$$+\sum_{i=0}^{k} \left(\prod_{j=i+1}^{k} (I_{\mathscr{X}_{1}} - F(j))\right) G(i)u(i), \ k \ge 0,$$
(C.1)

by (C.1) and Cauchy inequality, we have

$$\mathbb{E}[\|x(k+1)\|^2] \leq 2\mathbb{E}\left[\left\|\left(\prod_{i=0}^k (I_{\mathscr{X}_1} - F(i))\right) x(0)\right\|^2\right] + 2\mathbb{E}\left[\left\|\sum_{i=0}^k \left(\prod_{j=i+1}^k (I_{\mathscr{X}_1} - F(j))\right) G(i)u(i)\right\|^2\right].$$
(C.2)

It follows from Definition 2.8 that  $\sigma(u(k); \tau_N(\mathscr{X}_2))$  is independent of

$$\sigma\left(F(k);\tau_{\mathsf{S}}(\mathscr{L}(\mathscr{X}_{1}))\right)\bigvee\sigma\left(G(k);\tau_{\mathsf{S}}(\mathscr{L}(\mathscr{X}_{2},\mathscr{X}_{1}))\right),$$

which leads to

$$\mathbb{E}\left[\|G(t)u(t)\|^{2}\right] \leq \mathbb{E}\left[\|G(t)\|^{2}\right] \mathbb{E}\left[\|u(t)\|^{2}\right]$$
$$\leq \sup_{k\geq 0} \mathbb{E}\left[\|u(k)\|^{2}\right] \mathbb{E}\left[\|G(t)\|^{2}\right], t \geq 0.$$
(C.3)

By the condition (4.4), we get

$$\sup_{k\geq 0} \mathbb{E}[\|G(k)\|^2] < \infty,$$

thus, (C.3) implies  $\sup_{k\geq 0} \mathbb{E}[\|G(k)u(k)\|^2] < \infty$ . For  $0 \leq t < k$ , by the condition (4.3), we obtain

$$\mathbb{E}\left[\prod_{j=t+1}^{k} \|I_{\mathscr{X}_{1}} - F(j)\|^{4} \middle| \mathcal{F}(t)\right] \\
= \mathbb{E}\left[\mathbb{E}\left[\prod_{j=t+1}^{k} \|I_{\mathscr{X}_{1}} - F(j)\|^{4} \middle| \mathcal{F}(k-1)\right] \middle| \mathcal{F}(t)\right] \\
= \mathbb{E}\left[\mathbb{E}\left[\|I_{\mathscr{X}_{1}} - F(k)\|^{4} \middle| \mathcal{F}(k-1)\right] \prod_{j=t+1}^{k-1} \|I_{\mathscr{X}_{1}} - F(j)\|^{4} \middle| \mathcal{F}(t)\right] \\
\leq (1+\gamma(k))\mathbb{E}\left[\prod_{j=t+1}^{k-1} \|I_{\mathscr{X}_{1}} - F(j)\|^{4} \middle| \mathcal{F}(t)\right] \\
\vdots \\
\leq \prod_{j=t+1}^{k} (1+\gamma(j)) \text{ a.s.} \tag{C.4}$$

It follows from  $\sup_{k\geq 0}\mathbb{E}[\|G(k)\|^4]<\infty$  and (C.4) that

$$\mathbb{E}\left[\left\|\left(\prod_{j=s+1}^{k}(I_{\mathscr{X}_{1}}-F(j))\right)^{*}\left(\prod_{j=t+1}^{k}(I_{\mathscr{X}_{1}}-F(j))\right)G(t)\right\|^{2}\right]$$

$$\leq \mathbb{E} \left[ \left( \prod_{j=t+1}^{k} \|I_{\mathscr{X}_{1}} - F(j)\|^{4} \right) \left( \prod_{j=s+1}^{t} \|I_{\mathscr{X}_{1}} - F(j)\|^{2} \right) \|G(t)\|^{2} \right]$$

$$= \mathbb{E} \left[ \mathbb{E} \left[ \prod_{j=t+1}^{k} \|I_{\mathscr{X}_{1}} - F(j)\|^{4} \right| \mathcal{F}(t) \right] \left( \prod_{j=s+1}^{t} \|I_{\mathscr{X}_{1}} - F(j)\|^{2} \right) \|G(t)\|^{2} \right]$$

$$\leq \left( \prod_{j=t+1}^{k} (1+\gamma(j)) \right) \mathbb{E} \left[ \left( \prod_{j=s+1}^{t} \|I_{\mathscr{X}_{1}} - F(j)\|^{2} \right) \|G(t)\|^{2} \right]$$

$$\leq \left( \prod_{j=t+1}^{k} (1+\gamma(j)) \right) \left( \mathbb{E} \left[ \prod_{j=s+1}^{t} \|I_{\mathscr{X}_{1}} - F(j)\|^{4} \right] + \mathbb{E} \left[ \|G(t)\|^{4} \right] \right)$$

$$\leq \left( \prod_{j=t+1}^{k} (1+\gamma(j)) \right) \left( \prod_{j=s+1}^{t} (1+\gamma(j)) + \sup_{k\geq 0} \mathbb{E} \left[ \|G(k)\|^{4} \right] \right)$$

$$\leq \left( \prod_{k=0}^{\infty} (1+\gamma(k)) \right) \left( 1 + \sup_{k\geq 0} \mathbb{E} \left[ \|G(k)\|^{4} \right] \right)$$

$$< \infty, \ 0 \le s < t \le k,$$

$$(C.5)$$

where the last inequality is due to  $\sum_{k=0}^{\infty} \gamma(k) < \infty$ . By  $\sup_{k\geq 0} \mathbb{E}[||u(k)||^2] < \infty$ , Proposition A.5 and (C.5), we have

$$\left(\prod_{j=s+1}^{k} (I_{\mathscr{X}_{1}} - F(j))\right)^{*} \left(\prod_{j=t+1}^{k} (I_{\mathscr{X}_{1}} - F(j))\right) G(t)u(t)$$
  

$$\in L^{1}(\Omega; \mathscr{X}_{1}), \ 0 \le s < t \le k.$$
(C.6)

Noting that  $\mathbb{E}[u(t)|\mathcal{F}(s)] = \mathbb{E}[\mathbb{E}[u(t)|\mathcal{F}(t-1)]|\mathcal{F}(s)] = 0, 0 \le s < t$ , and Proposition 2.4 implies  $G(s)u(s) \in L^0(\Omega, \mathcal{F}(s); \mathscr{X}^N)$ , it follows from (C.5)-(C.6), Proposition A.5.(a)-(c), Lemma A.10 and Propositions 2.9-2.10 that

$$\mathbb{E}\left[\left\langle \left(\prod_{j=s+1}^{k} (I_{\mathscr{X}_{1}} - F(j))\right) G(s)u(s), \left(\prod_{j=t+1}^{k} (I_{\mathscr{X}_{1}} - F(j))\right) G(t)u(t)\right\rangle \right] \\ = \mathbb{E}\left[\left\langle G(s)u(s), \left(\prod_{j=s+1}^{k} (I_{\mathscr{X}_{1}} - F(j))\right)^{*} \left(\prod_{j=t+1}^{k} (I_{\mathscr{X}_{1}} - F(j))\right) G(t)u(t)\right\rangle \right] \\ \times \left(\prod_{j=t+1}^{k} (I_{\mathscr{X}_{1}} - F(j))\right) G(t)u(t) \left| \mathcal{F}(s) \right] \right] \\ = \mathbb{E}\left[\left\langle G(s)u(s), \mathbb{E}\left[\left(\prod_{j=s+1}^{k} (I_{\mathscr{X}_{1}} - F(j))\right)^{*} \\ \times \left(\prod_{j=t+1}^{k} (I_{\mathscr{X}_{1}} - F(j))\right) G(t)u(t) \left| \mathcal{F}(s) \right] \right) \right] \right]$$

$$= \mathbb{E}\left[\left\langle G(s)u(s), \mathbb{E}\left[\left(\prod_{j=s+1}^{k} (I_{\mathscr{X}_{1}} - F(j))\right)^{*} \times \left(\prod_{j=t+1}^{k} (I_{\mathscr{X}_{1}} - F(j))\right) G(t)\mathbb{E}[u(t)|\mathcal{F}(s)] \middle| \mathcal{F}(s)\right]\right\rangle\right]$$
  
= 0. (C.7)

It follows from Markov inequality that  $\mathbb{P}(\mathbb{E}[||G(i)u(i)||^2] = 0) = \mathbb{P}(G(i)u(i) = 0) = 1, \forall i \ge 0.$ Denote  $\Lambda = \{i \in \mathbb{N} : \mathbb{E}[||G(i)u(i)||^2] > 0\}$ . By (C.2), (C.3) and (C.7), we obtain

$$\mathbb{E}\left[\|x(k+1)\|^{2}\right] = \mathbb{E}\left[\left\|\left(\prod_{i=0}^{k}(I_{\mathscr{X}_{1}}-F(i))\right)x(0)\right\|^{2}\right] + \sum_{i=0}^{k}\mathbb{E}\left[\left\|\left(\prod_{j=i+1}^{k}(I_{\mathscr{X}_{1}}-F(j))\right)G(i)u(i)\right\|^{2}\right] + \sum_{i=0}^{\infty}\mathbb{E}\left[\left\|\left(\prod_{j=i+1}^{k}(I_{\mathscr{X}_{1}}-F(j))\right)G(i)u(i)\right\|^{2}\right] + \sum_{i=0}^{\infty}\mathbb{E}\left[\left\|\left(\prod_{j=i+1}^{k}(I_{\mathscr{X}_{1}}-F(j))\right)G(i)u(i)\right\|^{2}\right] + \sum_{i\in\Lambda}\mathbb{E}\left[\left\|G(i)u(i)\|^{2}\right]\mathbb{E}\left[\left\|\left(\prod_{j=i+1}^{k}(I_{\mathscr{X}_{1}}-F(j))\right)\frac{G(i)u(i)}{(\mathbb{E}\left[\left\|G(i)u(i)\|^{2}\right]\right)^{\frac{1}{2}}}\right\|^{2}\right] \\ \leq \mathbb{E}\left[\left\|\left(\prod_{i=0}^{k}(I_{\mathscr{X}_{1}}-F(i))\right)x(0)\right\|^{2}\right] + \sup_{k\geq0}\mathbb{E}\left[\left\|u(k)\|^{2}\right]\sum_{i\in\Lambda}\mathbb{E}\left[\left\|G(i)\|^{2}\right] \\ \times \mathbb{E}\left[\left\|\left(\prod_{j=i+1}^{k}(I_{\mathscr{X}_{1}}-F(j))\right)\frac{G(i)u(i)}{(\mathbb{E}\left[\left\|G(i)u(i)\|^{2}\right]\right)^{\frac{1}{2}}}\right\|^{2}\right].$$
(C.8)

Noting that the operator-valued random sequence  $\{I_{\mathscr{X}_1} - F(k), k \ge 0\}$  is  $L_2^2$ -stable with respect to the filter  $\{\mathcal{F}(k), k \ge 0\}$ , we get

$$\lim_{k \to \infty} \mathbb{E}\left[ \left\| \left( \prod_{i=0}^{k} (I_{\mathscr{X}_{1}} - F(i)) \right) x(0) \right\|^{2} \right] = 0.$$
 (C.9)

By Proposition 2.4, we know that  $G(i)u(i) \in L^0(\Omega, \mathcal{F}(i); \mathscr{X}_1)$  and

$$\mathbb{E}\left[\left\|\frac{G(i)u(i)}{\left(\mathbb{E}\left[\|G(i)u(i)\|^{2}\right]\right)^{\frac{1}{2}}}\right\|^{2}\right] = 1, \ i \in \Lambda,$$

which leads to

$$\lim_{k \to \infty} \mathbb{E}\left[ \left\| \left( \prod_{j=i+1}^{k} (I_{\mathscr{X}_{1}} - F(j)) \right) \frac{G(i)u(i)}{(\mathbb{E}\left[ \|G(i)u(i)\|^{2} \right])^{\frac{1}{2}}} \right\|^{2} \right] = 0, \ i \in \Lambda.$$
(C.10)

It follows from  $G(i)u(i) \in L^0(\Omega, \mathcal{F}(i); \mathscr{X}_1)$  and (C.4) that

$$\begin{split} \sup_{\substack{k\geq 0\\i\in\Lambda}} \mathbb{E} \left[ \left\| \left( \prod_{j=i+1}^{k} (I_{\mathscr{X}_{1}} - F(j)) \right) \frac{G(i)u(i)}{(\mathbb{E} \left[ \|G(i)u(i)\|^{2} \right])^{\frac{1}{2}}} \right\|^{2} \right] \\ &= \sup_{\substack{k\geq 0\\i\in\Lambda}} \mathbb{E} \left[ \mathbb{E} \left[ \left\| \left( \prod_{j=i+1}^{k} (I_{\mathscr{X}_{1}} - F(j)) \right) \frac{G(i)u(i)}{(\mathbb{E} \left[ \|G(i)u(i)\|^{2} \right])^{\frac{1}{2}}} \right\|^{2} \right] \mathcal{F}(i) \right] \right] \\ &\leq \sup_{\substack{k\geq 0\\i\in\Lambda}} \mathbb{E} \left[ \mathbb{E} \left[ \left\| \prod_{j=i+1}^{k} (I_{\mathscr{X}_{1}} - F(j)) \right\|^{2} \right| \mathcal{F}(i) \right] \frac{G(i)u(i)}{(\mathbb{E} \left[ \|G(i)u(i)\|^{2} \right])^{\frac{1}{2}}} \right\|^{2} \right] \\ &\leq \sup_{\substack{k\geq 0\\i\in\Lambda}} \mathbb{E} \left[ \mathbb{E} \left[ \left\| \prod_{j=i+1}^{k} (I_{\mathscr{X}_{1}} - F(j)) \right\|^{4} \right| \mathcal{F}(i) \right]^{\frac{1}{2}} \left\| \frac{G(i)u(i)}{(\mathbb{E} \left[ \|G(i)u(i)\|^{2} \right])^{\frac{1}{2}}} \right\|^{2} \right] \\ &\leq \sup_{\substack{k\geq 0\\i\in\Lambda}} \mathbb{E} \left[ \sqrt{\prod_{j=i+1}^{k} (1 + \gamma(j))} \left\| \frac{G(i)u(i)}{(\mathbb{E} \left[ \|G(i)u(i)\|^{2} \right])^{\frac{1}{2}}} \right\|^{2} \right] \\ &\leq \sqrt{\prod_{\substack{k=0\\i\in\Lambda}} (1 + \gamma(k))} \sup_{i\in\Lambda} \mathbb{E} \left[ \left\| \frac{G(i)u(i)}{(\mathbb{E} \left[ \|G(i)u(i)\|^{2} \right])^{\frac{1}{2}}} \right\|^{2} \right] \\ &= \sqrt{\prod_{k=0}^{\infty} (1 + \gamma(k))} < \infty. \end{split}$$
(C.11)

By the condition (4.4), (C.10)-(C.11) and Lemma D.2, we obtain

$$\lim_{k \to \infty} \sum_{i \in \Lambda} \mathbb{E} \left[ \|G(i)\|^2 \right] \mathbb{E} \left[ \left\| \left( \prod_{j=i+1}^k (I_{\mathscr{X}_1} - F(j)) \right) \frac{G(i)u(i)}{(\mathbb{E} \left[ \|G(i)u(i)\|^2 \right])^{\frac{1}{2}}} \right\|^2 \right] = 0.$$
(C.12)

Thus, substituting (C.9) and (C.12) into (C.8) gives  $\lim_{k\to\infty} \mathbb{E}[||x(k)||^2] = 0.$ 

Proof of Lemma 4.6. Denote  $F(k) = a(k)\mathcal{H}^*(k)\mathcal{H}(k) + b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}$  and  $G(k) = a(k)\mathcal{H}^*(k)$ , respectively. Notice that  $F(k) \ge 0$  and the estimation error equation (4.1) can be rewritten as the following random difference equation

$$e(k+1) = (I_{\mathscr{X}^N} - F(k)) e(k) + G(k)v(k).$$

Given the initial value  $e(0) \in \mathscr{X}^N$ , by Proposition A.5.(a)-(c), we know that  $\{e(k), k \ge 0\}$  is a random sequence with values in the Hilbert space  $(\mathscr{X}^N, \tau_N(\mathscr{X}^N))$ . On the one hand, it follows from Assumptions 4.1 and 4.2 that  $\{v(k), k \ge 0\}$  is independent of  $\{F(k), G(k), k \ge 0\}$ , and  $\sup_{k\ge 0} \mathbb{E}[||v(k)||^2] \le \beta_v$ . On the other hand, by the condition (4.7), we get  $\mathbb{E}[||I_{\mathscr{X}^N} - F(k)||^4|\mathcal{F}(k-1)] \le 1 + \gamma(k)$  a.s., which gives

$$\mathbb{E}\left[\|a(k)\mathcal{H}^*(k)\mathcal{H}(k) + b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}\|^2\right]$$

$$\leq \mathbb{E} \left[ \|I_{\mathscr{X}^{N}} - (I_{\mathscr{X}^{N}} - F(k))\|^{2} \right]$$
  

$$\leq 2 \left( 1 + \mathbb{E} \left[ \|I_{\mathscr{X}^{N}} - F(k)\|^{2} \right] \right)$$
  

$$\leq 2 \left( 1 + \mathbb{E} \left[ \|I_{\mathscr{X}^{N}} - F(k)\|^{4} \right]^{\frac{1}{2}} \right)$$
  

$$\leq 2 \left( 1 + \sqrt{1 + \gamma(k)} \right).$$
(C.13)

Noting that  $\mathcal{L}_{\mathcal{G}}$  is positive semi-definite, we have  $a(k)\mathcal{H}^*(k)\mathcal{H}(k)+b(k)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}} \geq a(k)\mathcal{H}^*(k)\mathcal{H}(k)$  $\geq 0$  a.s., which leads to  $||a(k)\mathcal{H}^*(k)\mathcal{H}(k)||^2 \leq ||a(k)\mathcal{H}^*(k)\mathcal{H}(k)+b(k)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}||^2$ . By (C.13), we obtain

$$\sup_{k\geq 0} \mathbb{E} \left[ \|G(k)\|^4 \right]$$

$$= \sup_{k\geq 0} \left\{ a^2(k) \mathbb{E} \left[ \|a(k)\mathcal{H}^*(k)\mathcal{H}(k)\|^2 \right] \right\}$$

$$\leq \sup_{k\geq 0} \left\{ a^2(k) \mathbb{E} \left[ \|a(k)\mathcal{H}^*(k)\mathcal{H}(k) + b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}} \|^2 \right] \right\}$$

$$\leq 2 \sup_{k\geq 0} \left\{ a^2(k) \left( 1 + \sqrt{1 + \gamma(k)} \right) \right\}$$

$$< \infty, \qquad (C.14)$$

where the last inequality is obtained from Condition 4.2 and  $\sum_{k=0}^{\infty} \gamma(k) < \infty$ . (I) If  $\sup_{k\geq 0} \mathbb{E}[\|\mathcal{H}(k)\|^2] < \infty$ , it follows from Condition 4.2 that

$$\sum_{k=0}^{\infty} \mathbb{E}\left[\|G(k)\|^2\right] \le \left\{\sup_{k\ge 0} \mathbb{E}\left[\|\mathcal{H}(k)\|^2\right]\right\} \sum_{k=0}^{\infty} a^2(k) < \infty.$$

By Theorem 4.4, the sequence of solutions to the estimation error equation (4.1) is  $L_2$ -asymptotically stable, i.e.,  $\lim_{k\to\infty} \mathbb{E}[\|f_i(k) - f_0\|^2] = 0, \ i \in \mathcal{V}.$ 

(II) If  $\mathbb{E}[||\mathcal{H}(k)||^2 |\mathcal{F}(k-1)] \leq \rho_0$  a.s., on the one hand, noting that  $\sup_{k\geq 0} \mathbb{E}[||\mathcal{H}(k)||^2] \leq \rho_0 < \infty$ , by above conclusion (I), we have  $\lim_{k\to\infty} \mathbb{E}[||e(k)||^2] = 0$ . On the other hand, it follows from the condition (4.7) and (C.14) that

$$\sup_{k \ge 0} \mathbb{E} \left[ \| (I_{\mathscr{X}^N} - F(k)) G(k) \|^2 \right] \le \sup_{k \ge 0} \mathbb{E} \left[ \| I_{\mathscr{X}^N} - F(k) \|^4 + \| G(k) \|^4 \right] < \infty.$$

Thus, by the estimation error equation (4.1), Assumptions 4.1 and 4.2, Lemma A.10 and Proposition 2.10, we get

$$\mathbb{E} \left[ \|e(k+1)\|^2 | \mathcal{F}(k-1) \right] \\ = \mathbb{E} \left[ \| (I_{\mathscr{X}^N} - F(k)) e(k) + G(k)v(k)\|^2 | \mathcal{F}(k-1) \right] \\ = \mathbb{E} \left[ \| (I_{\mathscr{X}^N} - F(k))e(k)\|^2 | \mathcal{F}(k-1) \right] + \mathbb{E} \left[ \| G(k)v(k)\|^2 | \mathcal{F}(k-1) \right] \\ + 2\mathbb{E} \left[ \langle e(k), (I_{\mathscr{X}^N} - F(k))G(k)v(k) \rangle | \mathcal{F}(k-1) \right] \\ = \mathbb{E} \left[ \| (I_{\mathscr{X}^N} - F(k))e(k)\|^2 | \mathcal{F}(k-1) \right] + \mathbb{E} \left[ \| G(k)v(k)\|^2 | \mathcal{F}(k-1) \right]$$

$$\begin{split} &+ 2\langle e(k), \mathbb{E}\left[(I_{\mathscr{X}^{N}} - F(k))G(k)v(k)|\mathcal{F}(k-1)\right]\rangle \\ &= \mathbb{E}\left[\left\|(I_{\mathscr{X}^{N}} - F(k))e(k)\right\|^{2}\right|\mathcal{F}(k-1)\right] + \mathbb{E}\left[\left\|G(k)v(k)\right\|^{2}\right|\mathcal{F}(k-1)\right] \\ &+ 2\langle e(k), \mathbb{E}\left[(I_{\mathscr{X}^{N}} - F(k))G(k)\mathbb{E}[v(k)|\mathcal{F}(k-1)]\right]\mathcal{F}(k-1)\right]\rangle \\ &= \mathbb{E}\left[\left\|(I_{\mathscr{X}^{N}} - F(k))e(k)\right\|^{2}\right|\mathcal{F}(k-1)\right] + \mathbb{E}\left[\left\|G(k)v(k)\right\|^{2}\right|\mathcal{F}(k-1)\right] \\ &\leq \mathbb{E}\left[\left\|(I_{\mathscr{X}^{N}} - F(k))\right\|^{2}\right|\mathcal{F}(k-1)\right] \|e(k)\|^{2} + \mathbb{E}\left[\left\|G(k)v(k)\right\|^{2}\right|\mathcal{F}(k-1)\right] \\ &\leq \mathbb{E}\left[\left\|(I_{\mathscr{X}^{N}} - F(k))\right\|^{4}\right|\mathcal{F}(k-1)\right]^{\frac{1}{2}}\|e(k)\|^{2} + \mathbb{E}\left[\left\|G(k)v(k)\right\|^{2}\right|\mathcal{F}(k-1)\right] \\ &\leq (1 + \gamma(k))^{\frac{1}{2}}\|e(k)\|^{2} + \beta_{v}\mathbb{E}\left[\left\|G(k)\right\|^{2}\right|\mathcal{F}(k-1)\right] \\ &\leq \left(1 + \frac{1}{2}\gamma(k)\right)\|e(k)\|^{2} + \rho_{0}\beta_{v}a^{2}(k) \text{ a.s.} \end{split}$$

Noting that  $\sum_{k=0}^{\infty} \gamma(k) < \infty$  and  $\sum_{k=0}^{\infty} a^2(k) < \infty$ , it follows from Lemma D.1 that  $||e(k)||^2$  converges almost surely, which together with  $\lim_{k\to\infty} \mathbb{E}[||e(k)||^2] = 0$  gives  $\lim_{k\to\infty} e(k) = 0$  a.s.

Proof of Lemma 4.8. Given the  $L_2$ -bounded adaptive sequence  $\{x(k), \mathcal{F}(kh-1), k \ge 0\}$  with values in the Hilbert space  $\mathscr{X}^N$  and the nonnegative integer m, we define a new sequence  $\{u(k), k \ge 0\}$  by

$$u(k+1) = \Phi_P((k+1)h - 1, kh)u(k), \ k \ge m,$$
(C.15)

where  $u(m) = x(m), u(i) = 0, i = 0, \dots, m-1$ . It follows from Proposition A.5.(a)-(c) that  $\{u(k), k \ge 0\}$  is a random sequence with values in the Hilbert space  $(\mathscr{X}^N, \tau_N(\mathscr{X}^N))$ . On one hand, from (C.15), by iterative calculations, we get

$$u(k+1) = \left(\prod_{i=m}^{k} \Phi_P((i+1)h - 1, ih)\right) u(m) \\ = \Phi_P((k+1)h - 1, mh)x(m), \ k \ge m.$$
(C.16)

Noting that  $x(m) \in L^0(\Omega, \mathcal{F}(mh-1); \mathscr{X}^N)$ , it is known from Lemma D.7 that

$$\mathbb{E}\left[\|u(k+1)\|^{2}\right] = \mathbb{E}\left[\mathbb{E}\left[\|\Phi_{P}((k+1)h-1,mh)x(m)\|^{2}|\mathcal{F}(mh-1)\right]\right] \leq d_{1}\mathbb{E}\left[\|x(m)\|^{2}\right],$$

thus,  $\sup_{k\geq 0} \mathbb{E}[||x(k)||^2] < \infty$  implies  $\sup_{k\geq 0} \mathbb{E}[||u(k)||^2] < \infty$ . On the other hand, we can rewrite (C.15) as

$$u(i+1) = \left(I_{\mathscr{X}^N} - \sum_{s=ih}^{(i+1)h-1} (a(s)\mathcal{H}^*(s)\mathcal{H}(s) + b(s)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})\right)u(i) + \left(\Phi_P((i+1)h-1,ih) - \left(I_{\mathscr{X}^N} - \sum_{s=ih}^{(i+1)h-1} (a(s)\mathcal{H}^*(s)\mathcal{H}(s) + b(s)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})\right)\right)u(i),$$

which leads to

$$u(k+1) = \left(\prod_{i=m}^{k} \left( I_{\mathscr{X}^{N}} - \sum_{s=ih}^{(i+1)h-1} (a(s)\mathcal{H}^{*}(s)\mathcal{H}(s) + b(s)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}) \right) \right) x(m) + \sum_{i=m}^{k} \left( \prod_{j=i+1}^{k} \left( I_{\mathscr{X}^{N}} - \sum_{s=jh}^{(j+1)h-1} (a(s)\mathcal{H}^{*}(s)\mathcal{H}(s) + b(s)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}) \right) \right) \right) \times \left( \Phi_{P}((i+1)h-1,ih) - \left( I_{\mathscr{X}^{N}} - \sum_{s=ih}^{(i+1)h-1} (a(s)\mathcal{H}^{*}(s)\mathcal{H}(s) + b(s)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}) \right) \right) u(i).$$
(C.17)

Denote the s-th order term in the binomial expansion of  $\Phi_P((i+1)h-1,ih)$  by  $M_s(i)$ ,  $s = 2, \dots, h$ . By (C.16)-(C.17), we have

$$\Phi_P((k+1)h-1,mh)x(m)$$

$$= \left(\prod_{i=m}^k \left(I_{\mathscr{X}^N} - \sum_{s=ih}^{(i+1)h-1} (a(s)\mathcal{H}^*(s)\mathcal{H}(s) + b(s)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})\right)\right)x(m)$$

$$+ \sum_{i=m}^k \left(\prod_{j=i+1}^k \left(I_{\mathscr{X}^N} - \sum_{s=jh}^{(j+1)h-1} (a(s)\mathcal{H}^*(s)\mathcal{H}(s) + b(s)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})\right)\right) \left(\sum_{s=2}^h M_s(i)\right)u(i),$$

from which we get

$$\mathbb{E}\left[\left\|\Phi_{P}((k+1)h-1,mh)x(m)\right\|^{2}\right] \leq 2\mathbb{E}\left[\left\|\left(\prod_{i=m}^{k}\left(I_{\mathscr{X}^{N}}-\sum_{s=ih}^{(i+1)h-1}(a(s)\mathcal{H}^{*}(s)\mathcal{H}(s)+b(s)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}})\right)\right)x(m)\right\|^{2}\right] \\ + 2\mathbb{E}\left[\left\|\sum_{i=m}^{k}\left(\prod_{j=i+1}^{k}\left(I_{\mathscr{X}^{N}}-\sum_{s=jh}^{(j+1)h-1}(a(s)\mathcal{H}^{*}(s)\mathcal{H}(s)+b(s)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}})\right)\right)\right) \\ \times\left(\sum_{s=2}^{h}M_{s}(i)\right)u(i)\right\|^{2}\right].$$
(C.18)

Noting that  $x(m) \in L^0(\Omega, \mathcal{F}(mh-1); \mathscr{X}^N)$  and  $\{I_{\mathscr{X}^N} - \sum_{i=kh}^{(k+1)h-1} (a(i)\mathcal{H}^*(i)\mathcal{H}(i) + b(i)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}), k \ge 0\}$  is  $L^2_2$ -stable with respect to the filter  $\{\mathcal{F}((k+1)h-1), k \ge 0\}$ , we obtain

$$\lim_{k \to \infty} \mathbb{E} \left[ \left\| \left( \prod_{i=m}^{k} \left( I_{\mathscr{X}^{N}} - \sum_{s=ih}^{(i+1)h-1} (a(s)\mathcal{H}^{*}(s)\mathcal{H}(s) + b(s)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}) \right) \right) x(m) \right\|^{2} \right] = 0.$$

Hereafter, we will analyze the second term on the right-hand side of the inequality in (C.18). Denote  $D(s) = a(s)\mathcal{H}^*(s)\mathcal{H}(s) + b(s)(\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})$ . On one hand, for  $2 \leq r \leq 2^h$ ,  $ih \leq s \leq (i+1)h-1$ , by Cr-inequality and the conditional Lyapunov inequality, we have

$$\mathbb{E}\left[\|D(s)\|^{r}|\mathcal{F}(ih-1)\right] \leq \left(\mathbb{E}\left[\|a(s)\mathcal{H}^{*}(s)\mathcal{H}(s)+b(s)(\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}})\|^{2^{h}}\middle|\mathcal{F}(ih-1)\right]\right)^{\frac{r}{2^{h}}} \leq \max\{a(s),b(s)\}^{r}\left(2^{2^{h}-1}\mathbb{E}\left[\|\mathcal{H}^{*}(s)\mathcal{H}(s)\|^{2^{h}}\middle|\mathcal{F}(ih-1)\right] + 2^{2^{h}-1}\|\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}\|^{2^{h}}\right)^{\frac{r}{2^{h}}} \leq (a^{r}(s)+b^{r}(s))\left(\left(\mathbb{E}\left[\|\mathcal{H}^{*}(s)\mathcal{H}(s)\|^{2^{h}}\middle|\mathcal{F}(ih-1)\right]\right)^{\frac{r}{2^{h}}}+\|\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}\|^{r}\right) \leq (a^{r}(s)+b^{r}(s))(\rho_{0}^{r}+\|\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}\|^{r}) \leq (a(s)+b(s))^{r}(\rho_{0}+\|\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}\|)^{r} \text{ a.s.} \tag{C.19}$$

Denote  $\rho_1 = \rho_0 + \|\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}\|$ . For  $ih \leq n_1 < \cdots < n_r \leq (i+1)h-1$ , by the conditional Hölder inequality, the conditional Lyapunov inequality and (C.19), we get

$$\mathbb{E}\left[\left\|\prod_{j=1}^{r} D(n_{j})\right\|^{2} \middle| \mathcal{F}(ih-1)\right] \\ \leq \left(\mathbb{E}\left[\left\|\prod_{j=1}^{r-1} D(n_{j})\right\|^{4} \middle| \mathcal{F}(ih-1)\right]\right)^{\frac{1}{2}} \left(\mathbb{E}\left[\left\|D(n_{r})\right\|^{4} \middle| \mathcal{F}(ih-1)\right]\right)^{\frac{1}{2}} \\ \leq \rho_{1}^{2}(a(n_{r})+b(n_{r}))^{2} \left(\mathbb{E}\left[\left\|\prod_{j=1}^{r-1} D(n_{j})\right\|^{4} \middle| \mathcal{F}(ih-1)\right]\right)^{\frac{1}{2}} \\ \leq \rho_{1}^{2r} \prod_{j=1}^{r} (a(n_{j})+b(n_{j}))^{2} \text{ a.s.}$$
(C.20)

On the other hand, for  $2 \le s \le h$ , it follows from Condition 4.1 and (C.20) that

$$\mathbb{E} \left[ \|M_{s}(i)\|^{2} | \mathcal{F}(ih-1) \right] \\= \mathbb{E} \left[ \left\| \sum_{ih \le n_{1} < \dots < n_{s} \le (i+1)h-1} \prod_{j=1}^{s} D(n_{j}) \right\|^{2} | \mathcal{F}(ih-1) \right] \\\leq \mathbb{C}_{h}^{s} \sum_{ih \le n_{1} < \dots < n_{s} \le (i+1)h-1} \mathbb{E} \left[ \left\| \prod_{j=1}^{s} D(n_{j}) \right\|^{2} | \mathcal{F}(ih-1) \right] \\\leq \mathbb{C}_{h}^{s} \sum_{ih \le n_{1} < \dots < n_{s} \le (i+1)h-1} \rho_{1}^{2s} \prod_{j=1}^{s} (a(n_{j}) + b(n_{j}))^{2} \\\leq \mathbb{C}_{h}^{s} \sum_{ih \le n_{1} < \dots < n_{s} \le (i+1)h-1} \rho_{1}^{2s} \prod_{j=1}^{s} (a(i) + b(i))^{2}$$

$$= (\mathbb{C}_{h}^{s})^{2} \rho_{1}^{2s} (a(i) + b(i))^{2s}$$
  

$$\leq \mathbb{C}_{h}^{s} \rho_{1}^{2s} (a(i) + b(i))^{2s} h! \text{ a.s.}, \qquad (C.21)$$

where the last inequality is obtained from  $\mathbb{C}_{h}^{s} \leq h!$ . Noting that  $\{a(k), k \geq 0\}$  and  $\{b(k), k \geq 0\}$ both monotonically vanish, it follows that there exists a constant  $c_{0} > 0$ , such that  $\sup_{k\geq 0}(a(k) + b(k)) \leq c_{0}$ . Then, by (C.21), we get

$$\sum_{s=2}^{h} \mathbb{E} \left[ \|M_{s}(i)\|^{2} |\mathcal{F}(ih-1) \right] \leq 4h! c_{0}^{-4} \left( a^{2}(i) + b^{2}(i) \right)^{2} \sum_{s=2}^{h} \mathbb{C}_{h}^{s} \rho_{1}^{2s} c_{0}^{2s} \\ = \rho_{2} \left( a^{2}(i) + b^{2}(i) \right)^{2} \text{ a.s.}, \qquad (C.22)$$

where  $\rho_2 = 4h!c_0^{-2}((\rho_1^2c_0^2+1)^h - 1 - h\rho_1^2c_0^2)$ . It follows from (C.22) that

$$\mathbb{E}\left[\left\|\left(\sum_{s=2}^{h} M_{s}(i)\right) u(i)\right\|^{2}\right]$$
  

$$\leq h\mathbb{E}\left[\sum_{s=2}^{h} \mathbb{E}\left[\left\|M_{s}(i)\right\|^{2}\right] \mathcal{F}(ih-1)\right] \|u(i)\|^{2}\right]$$
  

$$\leq h\rho_{2}\left(a^{2}(i)+b^{2}(i)\right)^{2} \mathbb{E}\left[\left\|u(i)\right\|^{2}\right],$$

which together with  $\sup_{i\geq 0}\mathbb{E}[\|u(i)\|^2]<\infty$  leads to

$$\sup_{i\geq 0} \mathbb{E}\left[\left\|R(i)\right\|^2\right] \leq h\rho_2 \sup_{i\geq 0} \mathbb{E}\left[\left\|u(i)\right\|^2\right] < \infty,\tag{C.23}$$

where

$$R(i) = \frac{1}{a^2(i) + b^2(i)} \left(\sum_{s=2}^h M_s(i)\right) u(i), \ i \ge m.$$

By the Minkowski inequality, we obtain

$$\mathbb{E}\left[\left\|\sum_{i=m}^{k} \left(\prod_{j=i+1}^{k} \left(I_{\mathscr{X}^{N}} - \sum_{s=jh}^{(j+1)h-1} D(s)\right)\right) \left(\sum_{s=2}^{h} M_{s}(i)\right) u(i)\right\|^{2}\right] \\ = \mathbb{E}\left[\left\|\sum_{i=m}^{k} \left(a^{2}(i) + b^{2}(i)\right) \left(\prod_{j=i+1}^{k} \left(I_{\mathscr{X}^{N}} - \sum_{s=jh}^{(j+1)h-1} D(s)\right)\right) R(i)\right\|^{2}\right] \\ \le \left(\sum_{i=m}^{k} \left(a^{2}(i) + b^{2}(i)\right) \left(\mathbb{E}\left[\left\|\left(\prod_{j=i+1}^{k} \left(I_{\mathscr{X}^{N}} - \sum_{s=jh}^{(j+1)h-1} D(s)\right)\right) R(i)\right\|^{2}\right]\right)^{\frac{1}{2}}\right)^{2}.$$
 (C.24)

From Proposition 2.4, we know that  $R(i) \in L^0(\Omega, \mathcal{F}((i+1)h-1); \mathscr{X}^N)$ , thus, by (C.23) and Lemma D.8, we know that there exists a constant  $d_3 > 0$  such that

$$\sup_{\substack{k\geq 0\\i\geq 0}} \mathbb{E}\left[ \left\| \left( \prod_{j=i+1}^{k} \left( I_{\mathscr{X}^{N}} - \sum_{s=jh}^{(j+1)h-1} D(s) \right) \right) R(i) \right\|^{2} \right]$$

$$\leq d_3 \sup_{i \geq 0} \mathbb{E}\left[ \|R(i)\|^2 \right]$$
  
< \infty, (C.25)

which together with that the operator-valued random sequence

$$\left\{I_{\mathscr{X}^N} - \sum_{i=kh}^{(k+1)h-1} (a(i)\mathcal{H}^*(i)\mathcal{H}(i) + b(i)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}), k \ge 0\right\}$$

is  $L_2^2$ -stable w.r.t.  $\{\mathcal{F}((k+1)h-1), k \ge 0\}$  gives

$$\lim_{k \to \infty} \mathbb{E} \left[ \left\| \left( \prod_{j=i+1}^{k} \left( I_{\mathscr{X}^N} - \sum_{s=jh}^{(j+1)h-1} D(s) \right) \right) R(i) \right\|^2 \right] = 0, \ \forall \ i \ge 0.$$
(C.26)

By Condition 4.2, (C.25)-(C.26) and Lemma D.2, we have

$$\lim_{k \to \infty} \sum_{i=m}^{k} \left( a^2(i) + b^2(i) \right) \left( \mathbb{E} \left[ \left\| \left( \prod_{j=i+1}^{k} \left( I_{\mathscr{X}^N} - \sum_{s=jh}^{(j+1)h-1} D(s) \right) \right) R(i) \right\|^2 \right] \right)^{\frac{1}{2}} = 0.$$

Given the  $L_2$ -bounded adaptive sequence  $\{x(k), \mathcal{F}(kh-1), k \ge 0\}$  with values in the Hilbert space  $\mathscr{X}^N$ , from (C.18) and (C.24), we know that

$$\lim_{k \to \infty} \mathbb{E}\left[ \|\Phi_P((k+1)h - 1, mh)x(m)\|^2 \right] = 0, \ \forall \ m \ge 0.$$
(C.27)

For  $j \in \mathbb{N}$ , denote  $m_j = \lfloor \frac{j}{h} \rfloor$ ,  $\widetilde{m}_j = \lceil \frac{j}{h} \rceil$ . Let  $\{y(k), \mathcal{F}(k), k \ge 0\}$  be a  $L_2$ -bounded adaptive sequence with values in the Hilbert space  $\mathscr{X}^N$ . For  $0 \le i < k-3h$ , noting that  $0 \le k-m_kh < h$ , from Propositions A.5-2.4, it is known that  $\Phi_P(m_kh - 1, \widetilde{m}_{i+1}h)\Phi_P(\widetilde{m}_{i+1}h - 1, i+1)y(i) \in$  $L^0(\Omega, \mathcal{F}(m_kh - 1); \mathscr{X}^N)$ , by Lemma D.7, we know that there exists a constant  $d_2 > 0$  such that

$$\mathbb{E} \left[ \|\Phi_{P}(k, i+1)y(i)\|^{2} \right] \\
= \mathbb{E} \left[ \|\Phi_{P}(k, m_{k}h)\Phi_{P}(m_{k}h-1, \widetilde{m}_{i+1}h)\Phi_{P}(\widetilde{m}_{i+1}h-1, i+1)y(i)\|^{2} \right] \\
= \mathbb{E} \left[ \mathbb{E} \left[ \left\| \Phi_{P}(k, m_{k}h)\Phi_{P}(m_{k}h-1, \widetilde{m}_{i+1}h) \right\|^{2} |\mathcal{F}(m_{k}h-1) \right] \right] \\
\times \Phi_{P}(\widetilde{m}_{i+1}h-1, i+1)y(i) \|^{2} |\mathcal{F}(m_{k}h-1)] \right] \\
\leq d_{2} \mathbb{E} \left[ \|\Phi_{P}(m_{k}h-1, \widetilde{m}_{i+1}h)\Phi_{P}(\widetilde{m}_{i+1}h-1, i+1)y(i)\|^{2} \right], \ 0 \leq i < k - 3h. \ (C.28)$$

Noting that  $0 \leq \widetilde{m}_{i+1}h - (i+1) < h$  and  $y(i) \in \mathcal{F}(i)$ , it follows from Lemma D.7 that

$$\sup_{i\geq 0} \mathbb{E} \left[ \left\| \Phi_P(\widetilde{m}_{i+1}h - 1, i+1)y(i) \right\|^2 \right]$$
  
= 
$$\sup_{i\geq 0} \mathbb{E} \left[ \mathbb{E} \left[ \left\| \Phi_P(\widetilde{m}_{i+1}h - 1, i+1)y(i) \right\|^2 \right] \mathcal{F}(i) \right] \right]$$
  
$$\leq d_2 \sup_{i\geq 0} \mathbb{E} \left[ \left\| y(i) \right\|^2 \right]$$
  
$$< \infty.$$
(C.29)

By Propositions A.5-2.4, we get  $\Phi_P(\widetilde{m}_{i+1}h - 1, i + 1)y(i) \in L^0(\Omega, \mathcal{F}(\widetilde{m}_{i+1}h - 1); \mathscr{X}^N)$ . Substituting (C.27) and (C.29) into (C.28) gives

$$\lim_{k \to \infty} \mathbb{E}\left[ \|\Phi_P(k, i+1)y(i)\|^2 \right] = 0, \ \forall i \ge 0,$$

which implies that the operator-valued random sequence  $\{I_{\mathscr{X}^N} - a(k)\mathcal{H}^*(k)\mathcal{H}(k) - b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}, k \geq 0\}$  is  $L_2^2$ -stable w.r.t.  $\{\mathcal{F}(k), k \geq 0\}$ .

Proof of Lemma 4.9. It follows from Condition 4.3 that there exists a constant  $C_1 > 0$ , such that  $|b(k) - a(k)| \le C_1(a^2(k) + b^2(k))$ , which gives

$$\sum_{i=0}^{k} a(k) \leq \sum_{i=0}^{k} (|a(k) - b(k)| + b(k))$$
  
$$\leq C_1 \sum_{i=0}^{k} (a^2(k) + b^2(k)) + \sum_{i=0}^{k} b(k), \ k \geq 0.$$
(C.30)

For the integer h > 0, denote

$$c(k) = \sum_{s=kh}^{(k+1)h-1} b(s),$$
(C.31)

by (C.30) and Condition 4.2, we get

$$\sum_{k=0}^{\infty} c(k) = \sum_{k=0}^{\infty} \sum_{s=kh}^{(k+1)h-1} b(s) = \sum_{k=0}^{\infty} b(k) = \infty.$$

Denote

$$\mathscr{H} = \operatorname{diag}\left\{\frac{1}{h}\mathscr{H}_{1}, \cdots, \frac{1}{h}\mathscr{H}_{N}\right\} + \mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}.$$
(C.32)

Noting that  $\mathcal{L}_{\mathcal{G}}$  is positive semi-definite and  $\sum_{i=1}^{N} \mathscr{H}_{i} > 0$ , by Lemma D.5, we know that  $\mathscr{H} \in \mathscr{L}(\mathscr{X}^{N})$  is a strictly positive self-adjoint operator. Let  $\{x(k), \mathcal{F}(kh-1), k \geq 0\}$  be a  $L_{2}$ -bounded adaptive sequence with values in the Hilbert space  $\mathscr{X}^{N}$ , then we can write  $x(k) = (x_{1}(k), \dots, x_{N}(k))$ , where  $x_{i}(k) : \Omega \to \mathscr{X}$ ,  $i = 1, \dots, N$  are the random elements with values in the Hilbert space  $(\mathscr{X}, \tau_{N}(\mathscr{X}))$ . Denote

$$\mu(i) = c(i)\mathscr{H}x(i) - \sum_{\substack{s=ih\\ s=ih}}^{(i+1)h-1} (a(s)\mathbb{E}[\mathcal{H}^*(s)\mathcal{H}(s)x(i)|\mathcal{F}(ih-1)] + b(s)(\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})x(i)).$$
(C.33)

By (C.31)-(C.33), we get

 $\mu(i)$ 

$$= \sum_{s=ih}^{(i+1)h-1} b(s) \operatorname{diag} \left\{ \frac{1}{h} \mathscr{H}_{1}, \cdots, \frac{1}{h} \mathscr{H}_{N} \right\} x(i) - \sum_{s=ih}^{(i+1)h-1} a(s) \mathbb{E}[\mathcal{H}^{*}(s)\mathcal{H}(s)x(i)|\mathcal{F}(ih-1)]$$

$$= \operatorname{diag} \left\{ \frac{1}{h} \sum_{s=ih}^{(i+1)h-1} b(s) \mathscr{H}_{1}x_{1}(i) - \sum_{s=ih}^{(i+1)h-1} a(s) \mathbb{E}[H_{1}^{*}(s)H_{1}(s)x_{1}(i)|\mathcal{F}(ih-1)], \cdots, \frac{1}{h} \sum_{s=ih}^{(i+1)h-1} b(s) \mathscr{H}_{N}x_{N}(i) - \sum_{s=ih}^{(i+1)h-1} a(s) \mathbb{E}[H_{N}^{*}(s)H_{N}(s)x_{N}(i)|\mathcal{F}(ih-1)] \right\}.$$

It follows from Lemma D.6 that there exists a constant  $C_2 > 0$  such that

$$\max_{ih \le s \le (i+1)h-1} \left( \frac{1}{h} \left( \sum_{s=ih}^{(i+1)h-1} b(s) \right) - a(s) \right)^2 \le C_2 \left( a^4(i) + b^4(i) \right).$$
(C.34)

Therefore, by Conditions 4.1-4.3, the condition (4.9) and (C.34), we have

$$\begin{split} & \mathbb{E}\left[\|\mu(i)\|^{2}\right] \\ &= \sum_{j=1}^{N} \mathbb{E}\left[\left\|\frac{1}{h}\sum_{s=ih}^{(i+1)h-1} b(s)\mathscr{H}_{j}x_{j}(i) - \sum_{s=ih}^{(i+1)h-1} a(s)\mathbb{E}[H_{j}^{*}(s)H_{j}(s)x_{j}(i)|\mathcal{F}(ih-1)]\right]\right|^{2}\right] \\ &= \sum_{j=1}^{N} \mathbb{E}\left[\left\|\left(\frac{1}{h}\sum_{s=ih}^{(i+1)h-1} b(s)\right)\left(\mathscr{H}_{j}x_{j}(i) - \sum_{s=ih}^{(i+1)h-1} \mathbb{E}\left[H_{j}^{*}(s)H_{j}(s)x_{j}(i)|\mathcal{F}(ih-1)\right]\right]\right|^{2}\right] \\ &+ \sum_{s=ih}^{N} \left(2b^{2}(i)\mathbb{E}\left[\left\|\mathscr{H}_{j}x_{j}(i) - \sum_{s=ih}^{(i+1)h-1} \mathbb{E}\left[H_{j}^{*}(s)H_{j}(s)x_{j}(i)|\mathcal{F}(ih-1)\right]\right\|^{2}\right] \\ &+ 2h\mathbb{E}\left[\sum_{s=ih}^{(i+1)h-1} \left(\frac{1}{h}\left(\sum_{s=ih}^{(i+1)h-1} b(s)\right) - a(s)\right)^{2} \\ &\times \left\|\mathbb{E}\left[H_{j}^{*}(s)H_{j}(s)x_{j}(i)|\mathcal{F}(ih-1)\right]\right\|^{2}\right]\right) \\ &\leq \sum_{j=1}^{N} \left(2b^{2}(i)\mathbb{E}\left[\left\|\mathscr{H}_{j}x_{j}(i) - \sum_{s=ih}^{(i+1)h-1} \mathbb{E}\left[H_{j}^{*}(s)H_{j}(s)x_{j}(i)|\mathcal{F}(ih-1)\right]\right\|^{2}\right] \\ &+ 2h\mathbb{E}\left[\sum_{s=ih}^{(i+1)h-1} \left(\frac{1}{h}\left(\sum_{s=ih}^{(i+1)h-1} \mathbb{E}\left[H_{j}^{*}(s)H_{j}(s)x_{j}(i)|\mathcal{F}(ih-1)\right]\right\|^{2}\right] \\ &+ 2h\mathbb{E}\left[\left\|\mathscr{H}_{j}^{*}(s)H_{j}(s)\right\|^{2}|\mathcal{F}(ih-1)\right]\|x_{j}(i)\|^{2}\right]\right) \\ &\leq \sum_{j=1}^{N} \left(2b^{2}(i)\mathbb{E}\left[\left\|\mathscr{H}_{j}x_{j}(i) - \sum_{s=ih}^{(i+1)h-1} \mathbb{E}\left[H_{j}^{*}(s)H_{j}(s)x_{j}(i)|\mathcal{F}(ih-1)\right]\right\|^{2}\right]\right) \\ &\leq \sum_{j=1}^{N} \left(2b^{2}(i)\mathbb{E}\left[\left\|\mathscr{H}_{j}x_{j}(i) - \sum_{s=ih}^{(i+1)h-1} \mathbb{E}\left[H_{j}^{*}(s)H_{j}(s)x_{j}(i)|\mathcal{F}(ih-1)\right]\right\|^{2}\right]\right) \end{aligned}$$

$$+ 2h\rho_{0}^{2} \sup_{k\geq 0} \mathbb{E} \left[ \|x(k)\|^{2} \right] \sum_{s=ih}^{(i+1)h-1} \left( \frac{1}{h} \left( \sum_{s=ih}^{(i+1)h-1} b(s) \right) - a(s) \right)^{2} \right)$$

$$\leq \sum_{j=1}^{N} \left( 2b^{2}(i) \mathbb{E} \left[ \left\| \mathscr{H}_{j}x_{j}(i) - \sum_{s=ih}^{(i+1)h-1} \mathbb{E} \left[ H_{j}^{*}(s)H_{j}(s)x_{j}(i) \right| \mathcal{F}(ih-1) \right] \right\|^{2} \right]$$

$$+ 2h^{2}\rho_{0}^{2} \sup_{k\geq 0} \mathbb{E} \left[ \|x(k)\|^{2} \right] \max_{ih\leq s\leq (i+1)h-1} \left( \frac{1}{h} \left( \sum_{s=ih}^{(i+1)h-1} b(s) \right) - a(s) \right)^{2} \right)$$

$$\leq \sum_{j=1}^{N} 2b^{2}(i) \mathbb{E} \left[ \left\| \mathscr{H}_{j}x_{j}(i) - \sum_{s=ih}^{(i+1)h-1} \mathbb{E} \left[ H_{j}^{*}(s)H_{j}(s)x_{j}(i) \right| \mathcal{F}(ih-1) \right] \right\|^{2} \right]$$

$$+ 2C_{2}Nh^{2}\rho_{0}^{2} \sup_{k\geq 0} \mathbb{E} \left[ \|x(k)\|^{2} \right] \left( a^{4}(i) + b^{4}(i) \right).$$

$$(C.35)$$

Noting that  $\{x_j(k), \mathcal{F}(kh-1), k \geq 0\}, j = 1, \dots, N$  are the  $L_2$ -bounded adaptive sequences with values in the Hilbert space  $\mathscr{X}$ , by (4.8), we obtain

$$\sum_{i=0}^{\infty} \mathbb{E} \left[ \left\| \mathscr{H}_{j} x_{j}(i) - \sum_{s=ih}^{(i+1)h-1} \mathbb{E} \left[ H_{j}^{*}(s) H_{j}(s) x_{j}(i) \right| \mathcal{F}(ih-1) \right] \right\|^{2} \right]$$

$$< \infty, \ j = 1, \cdots, N.$$
(C.36)

By Condition 4.2, (C.35)-(C.36) and Cauchy inequality, we get

$$\begin{split} &\sum_{i=0}^{\infty} \mathbb{E} \left[ \|\mu(i)\|^2 \right]^{\frac{1}{2}} \\ &\leq \sqrt{2} \sum_{i=0}^{\infty} b(i) \sum_{j=1}^{N} \left( \mathbb{E} \left[ \left\| \mathscr{H}_j x_j(i) - \sum_{s=ih}^{(i+1)h-1} \mathbb{E} \left[ H_j^*(s) H_j(s) x_j(i) \right| \mathcal{F}(ih-1) \right] \right\|^2 \right] \right)^{\frac{1}{2}} \\ &+ h \rho_0 \sup_{k \ge 0} \mathbb{E} \left[ \|x(k)\|^2 \right]^{\frac{1}{2}} \sqrt{2C_2 N} \sum_{i=0}^{\infty} \left( a^2(i) + b^2(i) \right) \\ &\leq \sqrt{2} N C_3 \left( \sum_{i=0}^{\infty} \sum_{j=1}^{N} \mathbb{E} \left[ \left\| \mathscr{H}_j x_j(i) - \sum_{s=ih}^{(i+1)h-1} \mathbb{E} \left[ H_j^*(s) H_j(s) x_j(i) \right| \mathcal{F}(ih-1) \right] \right\|^2 \right] \right) \\ &+ h \rho_0 \sup_{k \ge 0} \mathbb{E} \left[ \|x(k)\|^2 \right]^{\frac{1}{2}} \sqrt{2C_2 N} \sum_{i=0}^{\infty} \left( a^2(i) + b^2(i) \right) \\ &= \sqrt{2} N C_3 \left( \sum_{j=1}^{N} \sum_{i=0}^{\infty} \mathbb{E} \left[ \left\| \mathscr{H}_j x_j(i) - \sum_{s=ih}^{(i+1)h-1} \mathbb{E} \left[ H_j^*(s) H_j(s) x_j(i) \right| \mathcal{F}(ih-1) \right] \right\|^2 \right] \right) \\ &+ h \rho_0 \sup_{k \ge 0} \mathbb{E} \left[ \|x(k)\|^2 \right]^{\frac{1}{2}} \sqrt{2C_2 N} \sum_{i=0}^{\infty} \left( a^2(i) + b^2(i) \right) \\ &< \infty, \end{split}$$

where  $C_3 = \sum_{i=0}^{\infty} b^2(i)$ . For any given integer m > 0, denote  $\Gamma_m = \{i \ge m : \mathbb{E}[\|\mu(i)\|^2] > 0, i \in \mathbb{N}\}$ . Noting that  $\mathbb{E}[\|\mu(i)\|^2] = 0$  implies that  $\mu(i) = 0$  a.s., we obtain

$$\sum_{i=m}^{k} \left( \mathbb{E} \left[ \left\| \left( \prod_{j=i+1}^{k} (I_{\mathscr{X}^{N}} - c(j)\mathscr{H}) \right) \mu(i) \right\|^{2} \right] \right)^{\frac{1}{2}} \right]$$

$$\leq \sum_{i=m}^{\infty} \left( \mathbb{E} \left[ \left\| \left( \prod_{j=i+1}^{k} (I_{\mathscr{X}^{N}} - c(j)\mathscr{H}) \right) \mu(i) \right\|^{2} \right] \right)^{\frac{1}{2}} \right]$$

$$= \sum_{i\in\Gamma_{m}} \mathbb{E} \left[ \left\| \left( \prod_{j=i+1}^{k} (I_{\mathscr{X}^{N}} - c(j)\mathscr{H}) \right) \mu(i) \right\|^{2} \right] \right)^{\frac{1}{2}} \left( \mathbb{E} \left[ \left\| \left( \prod_{j=i+1}^{k} (I_{\mathscr{X}^{N}} - c(j)\mathscr{H}) \right) \eta(i) \right\|^{2} \right] \right)^{\frac{1}{2}}, \quad (C.37)$$

where  $\eta(i) = \mu(i)\mathbb{E}[\|\mu(i)\|^2]^{-\frac{1}{2}}, i \in \Gamma_m$ . Noting that  $\mathbb{E}[\|\eta(i)\|^2] = 1$ , it follows from Lemma D.3 that there exist constants M, d > 0, such that

$$\sup_{\substack{k\geq 0\\i\in\Gamma_m}} \left( \mathbb{E}\left[ \left\| \left( \prod_{j=i+1}^k (I_{\mathscr{X}^N} - c(j)\mathscr{H}) \right) \eta(i) \right\|^2 \right] \right)^{\frac{1}{2}} \\ \leq M^d \sup_{i\in\Gamma_m} \mathbb{E}\left[ \|\eta(i)\|^2 \right]^{\frac{1}{2}} \\ = Md.$$
(C.38)

By Lemma D.3 and Lebesgue dominated convergence theorem, we get

$$\lim_{k \to \infty} \left( \mathbb{E}\left[ \left\| \left( \prod_{j=i+1}^{k} (I_{\mathscr{X}^{N}} - c(j)\mathscr{H}) \right) \eta(i) \right\|^{2} \right] \right)^{\frac{1}{2}} = 0, \ \forall \ i \ge 0.$$
(C.39)

Therefore, combining (C.37)-(C.39) and Lemma D.2 leads to

$$\lim_{k \to \infty} \sum_{i=m}^{k} \left( \mathbb{E}\left[ \left\| \left( \prod_{j=i+1}^{k} (I_{\mathscr{X}^{N}} - c(j)\mathscr{H}) \right) \mu(i) \right\|^{2} \right] \right)^{\frac{1}{2}} = 0, \ \forall \ m \ge 0.$$

It follows from Lemma D.9 that the operator-valued random sequence

$$\left\{I_{\mathscr{X}^N} - \sum_{i=kh}^{(k+1)h-1} (a(i)\mathcal{H}^*(i)\mathcal{H}(i) + b(i)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}), k \ge 0\right\}$$

is  $L_2^2$ -stable w.r.t. { $\mathcal{F}((k+1)h-1), k \ge 0$ }. Hence, from Lemma 4.8, it is known that the operator-valued random sequence { $I_{\mathscr{X}^N} - a(k)\mathcal{H}^*(k)\mathcal{H}(k) - b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}, k \ge 0$ } is  $L_2^2$ -stable w.r.t. { $\mathcal{F}(k), k \ge 0$ }.

Proof of Theorem 4.10. By the conditions (4.8)-(4.9) and Lemma 4.9, it is known that  $\{I_{\mathscr{X}^N} - a(k)\mathcal{H}^*(k)\mathcal{H}(k) - b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}, k \geq 0\}$  is  $L_2^2$ -stable w.r.t.  $\{\mathcal{F}(k), k \geq 0\}$ . Denote  $D(k) = a(k)\mathcal{H}^*(k)\mathcal{H}(k) + b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}$ . By the condition (4.9) that

$$\mathbb{E}\left[\|D(k)\|^{r}|\mathcal{F}(k-1)\right] \leq \left(\mathbb{E}\left[\|a(k)\mathcal{H}^{*}(k)\mathcal{H}(k)+b(k)(\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}})\|^{2^{h}}\middle|\mathcal{F}(k-1)\right]\right)^{\frac{r}{2^{h}}} \leq \max\{a(k),b(k)\}^{r}\left(2^{2^{h}-1}\mathbb{E}\left[\|\mathcal{H}^{*}(k)\mathcal{H}(k)\|^{2^{h}}\middle|\mathcal{F}(k-1)\right]\right] + 2^{2^{h}-1}\|\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}\|^{2^{h}}\right)^{\frac{r}{2^{h}}} \leq (a^{r}(k)+b^{r}(k))\left(\left(\mathbb{E}\left[\|\mathcal{H}^{*}(k)\mathcal{H}(k)\|^{2^{h}}\middle|\mathcal{F}(k-1)\right]\right)^{\frac{r}{2^{h}}}+\|\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}\|^{r}\right) \leq (a^{r}(k)+b^{r}(k))(\rho_{0}^{r}+\|\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}\|^{r}) \leq (a^{r}(k)+b^{r}(k))\rho_{1}^{r} \text{ a.s.}, \ \forall 1 \leq r \leq 4,$$
(C.40)

where  $\rho_1 = \rho_0 + \|\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}\|$ . By the condition (4.10) and (C.40), we get

$$\mathbb{E} \left[ \|I_{\mathscr{X}^{N}} - (a(k)\mathcal{H}^{*}(k)\mathcal{H}(k) + b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}) \|^{4} | \mathcal{F}(k-1) \right] \\
= \mathbb{E} \left[ \|(I_{\mathscr{X}^{N}} - (a(k)\mathcal{H}^{*}(k)\mathcal{H}(k) + b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}))^{4} \| | \mathcal{F}(k-1) \right] \\
= \mathbb{E} \left[ \|I_{\mathscr{X}^{N}} - 4D(k) + 6D^{2}(k) - 4D^{3}(k) + D^{4}(k) \| | \mathcal{F}(k-1) \right] \\
\leq \mathbb{E} \left[ \|I_{\mathscr{X}^{N}} - 4D(k)\| + 6\|D(k)\|^{2} + 4\|D(k)\|^{3} + \|D(k)\|^{4} | \mathcal{F}(k-1) \right] \\
\leq 1 + \gamma(k) \text{ a.s.},$$
(C.41)

where  $\gamma(k) = \Gamma(k) + 6(a^2(k) + b^2(k))\rho_1^2 + 4(a^3(k) + b^3(k))\rho_1^3 + (a^4(k) + b^4(k))\rho_1^4$ . From Condition 4.2 and the condition (4.10), we know that  $\sum_{k=0}^{\infty} \gamma(k) < \infty$ , which together with Lemma 4.6 implies that algorithm (3.6) is both mean square and almost surely strongly consistent.

Proof of Corollary 4.11. It follows from  $\|\mathcal{H}(0)\| \leq \rho_0$  a.s. and Proposition A.5.(a) that  $H_j^*(0) \times H_j(0)x \in L^1(\Omega; \mathscr{X}), x \in \mathscr{X}$ . For the integer h > 0 and  $j \in \mathcal{V}$ , we define the operator  $\mathscr{H}_j : \mathscr{X} \to \mathscr{X}$  by

$$\mathscr{H}_{j}(x) = h\mathbb{E}\left[H_{j}^{*}(0)H_{j}(0)x\right], \ x \in \mathscr{X}, \ j \in \mathcal{V}.$$
(C.42)

For any given  $x_1, x_2 \in \mathscr{X}$  and  $c_1, c_2 \in \mathbb{R}$ , we have

$$\mathcal{H}_{j}(c_{1}x_{1} + c_{2}x_{2}) = c_{1}h\mathbb{E}\left[H_{j}^{*}(0)H_{j}(0)x_{1}\right] + c_{2}h\mathbb{E}\left[H_{j}^{*}(0)H_{j}(0)x_{2}\right]$$
  
$$= c_{1}\mathcal{H}_{j}(x_{1}) + c_{2}\mathcal{H}_{j}(x_{2}).$$
(C.43)

Noting that  $H_j^*(0)H_j(0)x \in L^1(\Omega; \mathscr{X})$ , by Lemma A.10, we get

$$\langle \mathscr{H}_j(x_1), x_2 \rangle = h \mathbb{E} \left[ \left\langle H_j^*(0) H_j(0) x_1, x_2 \right\rangle \right]$$

$$= h\mathbb{E}\left[\left\langle x_1, H_j^*(0)H_j(0)x_2\right\rangle\right]$$
  
=  $\langle x_1, \mathscr{H}_j(x_2)\rangle, \ j \in \mathcal{V}.$  (C.44)

From (C.43)-(C.44), it is known that  $\mathscr{H}_j$  is a linear self-adjoint operator, which gives

$$\|\mathscr{H}_{j}(x)\| \leq h\mathbb{E}\left[\left\|H_{j}^{*}(0)H_{j}(0)\right\| \|x\|\right] \leq h\rho_{0}\|x\|, \ \forall \ x \in \mathscr{X}, \ j \in \mathcal{V},$$

thus, the self-adjoint operator  $\mathscr{H}_j \in \mathscr{L}(\mathscr{X})$ . Denote  $\mathscr{H}_j(x) := \mathscr{H}_j x, \forall x \in \mathscr{X}$ . Noting that  $H_j^*(0)H_j(0)x \in L^1(\Omega; \mathscr{X})$ , it follows from Lemma A.10 that

$$\langle \mathscr{H}_{j}x, x \rangle = h \left\langle \mathbb{E} \left[ H_{j}^{*}(0)H_{j}(0)x \right], x \right\rangle$$

$$= h \mathbb{E} \left[ \left\langle H_{j}^{*}(0)H_{j}(0)x, x \right\rangle \right]$$

$$= h \mathbb{E} \left[ \|H_{j}(0)x\|^{2} \right] \geq 0,$$
(C.45)

from which we know that the operator  $\mathscr{H}_j$  defined in (C.42) is positive bounded linear selfadjoint,  $j \in \mathcal{V}$ . Noting that  $\{\mathcal{H}(k), k \geq 0\}$  is an i.i.d. sequence with values in the topology space  $(\mathscr{L}(\mathscr{X}^N, \bigoplus_{i=1}^N \mathscr{Y}_i), \tau_{\mathsf{S}}(\mathscr{L}(\mathscr{X}^N, \bigoplus_{i=1}^N \mathscr{Y}_i)))$ , it follows from Definition 2.7 that  $\{\mathcal{H}^*(k)\mathcal{H}(k)x, x \in \mathscr{X}^N, k \geq 0\}$  is an i.i.d. sequence with values in the Banach space  $(\mathscr{X}^N, \tau_{\mathsf{N}}(\mathscr{X}^N))$ . By Proposition E.1.10 in [66], we know that  $\{\|\mathcal{H}^*(k)\mathcal{H}(k)\|, k \geq 0\}$  and  $\{\|I_{\mathscr{X}^N} - (a(k)\mathcal{H}^*(k)\mathcal{H}(k) + b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})\|, k \geq 0\}$  are both independent random sequences. By Condition 4.2, there exists an integer  $s_0 > 0$ , such that  $a(k) + b(k) \leq (4\rho_0^2 + 4\|\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}\|)^{-1}, \forall k \geq s_0$ . We define the nonnegative real sequence  $\{\Gamma(k), k \geq 0\}$  by

$$\Gamma(k) = \begin{cases} 4(a(k) + b(k)) \left(\rho_0^2 + \|\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}\|\right), & 0 \le k < s_0; \\ 0, & k \ge s_0, \end{cases}$$
(C.46)

which shows that  $\sum_{k=0}^{\infty} \Gamma(k) < \infty$ . Noting that  $\|I_{\mathscr{X}^N} - 4(a(k)\mathcal{H}^*(k)\mathcal{H}(k) + b(k)\mathcal{L}_{\mathscr{G}} \otimes I_{\mathscr{X}})\| \le 1 + 4(a(k) + b(k))(\rho_0^2 + \|\mathcal{L}_{\mathscr{G}} \otimes I_{\mathscr{X}}\|)$  a.s.,  $\forall k \ge 0$ , we obtain

$$\mathbb{E}\left[\left\|I_{\mathscr{X}^{N}}-4\left(a(k)\mathcal{H}^{*}(k)\mathcal{H}(k)+b(k)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}\right)\right\|\left|\mathcal{F}(k-1)\right]\right]$$

$$=\mathbb{E}\left[\left\|I_{\mathscr{X}^{N}}-4\left(a(k)\mathcal{H}^{*}(k)\mathcal{H}(k)+b(k)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}\right)\right\|\right]$$

$$=\mathbb{E}\left[\sup_{\|x\|=1}\left|\langle I_{\mathscr{X}^{N}}-4\left(a(k)\mathcal{H}^{*}(k)\mathcal{H}(k)+b(k)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}\right)x,x\rangle\right|\right]$$

$$=\mathbb{E}\left[\sup_{\|x\|=1}\left|1-4\left\langle\left(a(k)\mathcal{H}^{*}(k)\mathcal{H}(k)+b(k)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}\right)x,x\right\rangle\right|\right]$$

$$=\mathbb{E}\left[1-4\inf_{\|x\|=1}\left\langle\left(a(k)\mathcal{H}^{*}(k)\mathcal{H}(k)+b(k)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}\right)x,x\right\rangle\right]$$

$$\leq 1 \text{ a.s., } \forall k \geq s_{0}.$$
(C.47)

Then, by (C.46)-(C.47), we get

$$\mathbb{E}\left[\left\|I_{\mathscr{X}^{N}}-4\left(a(k)\mathcal{H}^{*}(k)\mathcal{H}(k)+b(k)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}\right)\right\|\left|\mathcal{F}(k-1)\right]\leq1+\Gamma(k) \text{ a.s.}, \ \forall \ k\geq0.$$

It can be verified that

$$\sup_{k \ge 0} \left( \mathbb{E} \left[ \|\mathcal{H}^{*}(k)\mathcal{H}(k)\|^{2^{\max\{h,2\}}} \middle| \mathcal{F}(k-1) \right] \right)^{\frac{1}{2^{\max\{h,2\}}}} \\ = \sup_{k \ge 0} \left( \mathbb{E} \left[ \|\mathcal{H}^{*}(k)\mathcal{H}(k)\|^{2^{\max\{h,2\}}} \right] \right)^{\frac{1}{2^{\max\{h,2\}}}} \\ = \sup_{k \ge 0} \left( \mathbb{E} \left[ \|\mathcal{H}^{*}(0)\mathcal{H}(0)\|^{2^{\max\{h,2\}}} \right] \right)^{\frac{1}{2^{\max\{h,2\}}}} \\ \le \rho_{0}^{\frac{2}{9}} \text{ a.s.}$$
(C.48)

For any given integers  $k \ge 0$ , h > 0 and  $x \in \mathscr{X}$ , if  $A \in \mathcal{F}(kh-1)$ , then it follows from Proposition A.5 and (C.48) that  $H_j^*(i)H_j(i)(\mathbf{1}_A \otimes x) \in L^1(\Omega; \mathscr{X})$ ,  $i \ge kh$ , which together with Lemma A.9 implies that  $\mathbb{E}[H_j^*(i)H_j(i)(\mathbf{1}_A \otimes x)|\mathcal{F}(kh-1)]$  uniquely exists. Since  $\{\mathcal{H}(k), k \ge 0\}$ and  $\{v(k), k \ge 0\}$  are both i.i.d sequences and they are mutually independent, it follows from Definition 2.7 that  $H_j^*(i)H_j(i)x$  is independent of  $\mathbf{1}_{F\cap A}$  with  $F \in \mathcal{F}(kh-1)$ . Thus,

$$\begin{split} &\int_{F} \mathbb{E} \left[ \sum_{i=kh}^{(k+1)h-1} H_{j}^{*}(i)H_{j}(i)(\mathbf{1}_{A} \otimes x) \middle| \mathcal{F}(kh-1) \right] \, \mathrm{d}\mathbb{P} \\ &= \int_{F} \sum_{i=kh}^{(k+1)h-1} H_{j}^{*}(i)H_{j}(i)(\mathbf{1}_{A} \otimes x) \, \mathrm{d}\mathbb{P} \\ &= \int_{\Omega} \left( \sum_{i=kh}^{(k+1)h-1} H_{j}^{*}(i)H_{j}(i)x \right) \mathbf{1}_{F \cap A} \, \mathrm{d}\mathbb{P} \\ &= \int_{\Omega} \sum_{i=kh}^{(k+1)h-1} H_{j}^{*}(i)H_{j}(i)x \, \mathrm{d}\mathbb{P} \int_{\Omega} \mathbf{1}_{F \cap A} \, \mathrm{d}\mathbb{P} \\ &= \mathbb{E} \left[ \sum_{i=kh}^{(k+1)h-1} H_{j}^{*}(i)H_{j}(i)x \right] \, \mathbb{P}(F \cap A) \\ &= h\mathbb{E} \left[ H_{j}^{*}(0)H_{j}(0)x \right] \, \mathbb{P}(F \cap A) \\ &= h \int_{F} \mathbb{E} \left[ H_{j}^{*}(0)H_{j}(0)x \right] \, \mathbf{1}_{A} \, \mathrm{d}\mathbb{P}, \ \forall F \in \mathcal{F}(kh-1), \ j \in \mathcal{V}, \end{split}$$

which gives

$$\mathbb{E}\left[\sum_{i=kh}^{(k+1)h-1} H_j^*(i)H_j(i)(\mathbf{1}_A \otimes x) \middle| \mathcal{F}(kh-1)\right] = \mathscr{H}_j(\mathbf{1}_A \otimes x) \text{ a.s., } j \in \mathcal{V},$$

which together with the properties of the conditional expectation, the operator  $\mathscr{H}_j$  and the linearity of Bochner integral leads to

$$\mathscr{H}_{j}y = \mathbb{E}\left[\left|\sum_{i=kh}^{(k+1)h-1} H_{j}^{*}(i)H_{j}(i)y\right| \mathcal{F}(kh-1)\right] \text{ a.s., } j \in \mathcal{V},$$
(C.49)

where  $y \in L^0(\Omega, \mathcal{F}(kh-1); \mathscr{X})$  is a simple function. For  $f \in L^2(\Omega, \mathcal{F}(kh-1); \mathscr{X})$ , by Pettis measurability theorem, we know that there exists a sequence of simple functions  $\{f_n \in L^0(\Omega, \mathcal{F}(kh-1); \mathscr{X}), n \ge 0\}$  satisfying  $||f_n|| \le f$  a.s. and  $\lim_{n\to\infty} f_n = f$  a.s., which together with (C.48) and Cauchy inequality gives  $H_j^*(i)H_j(i)f \in L^1(\Omega; \mathscr{X})$ . Thus, from Lemma A.9, it is known that  $\mathbb{E}[H_j^*(i)H_j(i)f|\mathcal{F}(kh-1)]$  uniquely exists. Noting that  $\mathscr{H}_j \in \mathscr{L}(\mathscr{X})$ , it follows from (C.48)-(C.49) and the dominated convergence theorem of conditional expectation that

$$\mathcal{H}_{j}f$$

$$= \lim_{n \to \infty} \mathcal{H}_{j}f_{n}$$

$$= \lim_{n \to \infty} \mathbb{E} \left[ \sum_{i=kh}^{(k+1)h-1} H_{j}^{*}(i)H_{j}(i)f_{n} \middle| \mathcal{F}(kh-1) \right]$$

$$= \mathbb{E} \left[ \sum_{i=kh}^{(k+1)h-1} H_{j}^{*}(i)H_{j}(i)\lim_{n \to \infty} f_{n} \middle| \mathcal{F}(kh-1) \right]$$

$$= \mathbb{E} \left[ \sum_{i=kh}^{(k+1)h-1} H_{j}^{*}(i)H_{j}(i)f \middle| \mathcal{F}(kh-1) \right] \text{ a.s., } j \in \mathcal{V}. \quad (C.50)$$

Let  $\{x(k), \mathcal{F}(kh-1), k \ge 0\}$  be a  $L_2$ -bounded adaptive sequence with values in the Hilbert space  $\mathscr{X}$ , by (C.50), we get

$$\mathscr{H}_{j}x(k) = \mathbb{E}\left[\left|\sum_{i=kh}^{(k+1)h-1} H_{j}^{*}(i)H_{j}(i)x(k)\right| \mathcal{F}(kh-1)\right] \text{ a.s., } j \in \mathcal{V}.$$

For any non-zero element x in Hilbert space x, from Lemma A.10, the condition (4.11) and (C.42), we have

$$\left\langle \sum_{j=1}^{N} \mathscr{H}_{j} x, x \right\rangle = h \left\langle \mathbb{E} \left[ \sum_{j=1}^{N} H_{j}^{*}(0) H_{j}(0) x \right], x \right\rangle = h \sum_{j=1}^{N} \mathbb{E} \left[ \|H_{j}(0) x\|^{2} \right] > 0.$$
(C.51)

Hence, by (C.45)-(C.48), (C.51) and Theorem 4.10, it is known that the algorithm (3.6) is both mean square and almost surely strongly consistent.

*Proof of Theorem 5.4.* Let  $H_i(k)$  be the mapping induced by random input data  $x_i(k)$ , where

$$H_i(k)(f) = f(x_i(k)), \ f \in \mathscr{H}_K, \ k \ge 0, \ i \in \mathcal{V}.$$

For any given  $f_1, f_2 \in \mathscr{H}_K$  and  $c_1, c_2 \in \mathbb{R}$ , it follows from the reproducing property of  $\mathscr{H}_K$  that

$$H_{i}(k)(c_{1}f_{1}+c_{2}f_{2}) = \langle c_{1}f_{1}+c_{2}f_{2}, x_{i}(k)\rangle_{K}$$
  
=  $c_{1}H_{i}(k)(f_{1})+c_{2}H_{i}(k)(f_{2}), k \ge 0, i \in \mathcal{V},$  (C.52)

thus,  $H_i(k)$  is a linear operator. Noting the continuity of Mercer kernel  $K : \mathscr{X} \times \mathscr{X} \to \mathbb{R}$ , we know that the function  $K_x : \mathscr{X} \to \mathscr{H}_K$  induced by the Mercer kernel K is also continuous. It is known from  $\mathscr{H}_K = \overline{\operatorname{span}\{K_x, x \in \mathscr{X}\}}$  that  $f \in \mathscr{H}_K$  is a Borel measurable function, which implies that  $H_i(k)(f) = f(x_i(k))$  is a random variable with values in the Hilbert space  $(\mathbb{R}, \tau_N(\mathbb{R}))$ . By Assumption 5.3 and the reproducing property of  $\mathscr{H}_K$ , we have

$$||K_x||_K = \sqrt{\langle K_x, K_x \rangle_K} = \sqrt{K(x, x)} \le \sup_{x \in \mathscr{X}} \sqrt{K(x, x)} < \infty.$$
(C.53)

For any given sample path  $\omega \in \Omega$ , by (C.53), we get

$$|H_{i}(k)(\omega)(f)| = |\langle f, K_{x_{i}(k)(\omega)} \rangle_{K}|$$
  

$$\leq \sup_{x \in \mathscr{X}} \sqrt{K(x, x)} ||f||_{K}, \ \forall \ f \in \mathscr{H}_{K}, \ k \ge 0, \ i \in \mathcal{V},$$
(C.54)

then  $||H_i(k)||_{\mathscr{L}(\mathscr{H}_K,\mathbb{R})} \leq \sup_{x\in\mathscr{X}} \sqrt{K(x,x)}$  a.s.. By (C.52), (C.54) and Proposition A.5, we know that  $H_i(k): \Omega \to \mathscr{L}(\mathscr{H}_K,\mathbb{R})$  is a random element with values in the topological space  $(\mathscr{L}(\mathscr{H}_K,\mathbb{R}),\tau_{\mathsf{S}}(\mathscr{L}(\mathscr{H}_K,\mathbb{R})))$ . Denote  $\mathcal{H}(k):=\operatorname{diag}\{H_1(k),\cdots,H_N(k)\}$  and  $v(k):=(v_1(k),$ 

 $\cdots, v_N(k)$ ), it follows from Definition 2.2 that  $\mathcal{H}(k)$  is the random element with values in the topological space  $(\mathscr{L}(\mathscr{H}_K^N, \mathbb{R}^N), \tau_{\mathsf{S}}(\mathscr{L}(\mathscr{H}_K^N, \mathbb{R}^N)))$ , and v(k) is a random vector with values in the Hilbert space  $(\mathbb{R}^N, \tau_{\mathsf{N}}(\mathbb{R}^N))$ . Thus, by Proposition A.5, we have

$$\mathcal{F}(k) = \bigvee_{s=0}^{k} \left( \sigma \left( \mathcal{H}(s); \tau_{\mathbf{S}} \left( \mathscr{L} \left( \mathscr{H}_{K}^{N}, \mathbb{R}^{N} \right) \right) \right) \bigvee \sigma \left( v(s); \tau_{\mathbf{N}} \left( \mathbb{R}^{N} \right) \right) \right), \ k \ge 0.$$
(C.55)

It follows from Assumption 5.1 and (C.55) that Assumption 4.1 holds. By Assumption 5.2 and (C.55), it is known that  $\{v(k), \mathcal{F}(k), k \ge 0\}$  is the martingale difference sequence and there exists a constant  $\beta_v := N\beta > 0$ , such that

$$\sup_{k\geq 0} \mathbb{E}\left[ \left\| v(k) \right\|^2 \middle| \mathcal{F}(k-1) \right] \leq N \max_{i\in\mathcal{V}} \sup_{k\geq 0} \mathbb{E}\left[ \left\| v_i(k) \right\|^2 \middle| \mathcal{F}(k-1) \right] \leq \beta_v \text{ a.s.},$$

which implies that Assumption 4.2 holds. Let  $x \in \mathscr{X}$ , for any given  $f_1, f_2 \in \mathscr{H}_K$  and  $c_1, c_2 \in \mathbb{R}$ , we get

$$(K_x \otimes K_x) (c_1 f_1 + c_2 f_2) = c_1 f_1(x) K_x + c_2 f_2(x) K_x$$
  
=  $c_1 (K_x \otimes K_x) f_1 + c_2 (K_x \otimes K_x) f_2,$  (C.56)

from (C.53), Assumption 5.3 and the reproducing property of  $\mathcal{H}_K$ , it is known that

$$\|(K_x \otimes K_x) f\|_K \le \|K_x\|_K^2 \|f\|_K \le \sup_{x \in \mathscr{X}} K(x, x) \|f\|_K, \ \forall \ f \in \mathscr{H}_K, \ \forall \ x \in \mathscr{X}.$$
(C.57)

Thus, it follows from (C.56)-(C.57) that  $K_x \otimes K_x \in \mathscr{L}(\mathscr{H}_K)$ . Let  $x = \sum_{i=1}^n \mathbf{1}_{A_i} \otimes x_i$  be a random vector with values in the Hilbert space  $(\mathscr{X}, \tau_N(\mathscr{X}))$ , where  $A_i \cap A_j = \emptyset$ ,  $1 \le i \ne j \le n$ . Noting that  $K_x = \sum_{i=1}^n \mathbf{1}_{A_i} \otimes K_{x_i}$ , we have

$$K_x \otimes K_x = \left(\sum_{i=1}^n \mathbf{1}_{A_i} \otimes K_{x_i}\right) \otimes \left(\sum_{i=1}^n \mathbf{1}_{A_i} \otimes K_{x_i}\right) = \sum_{i=1}^n \mathbf{1}_{A_i} \otimes \left(K_{x_i} \otimes K_{x_i}\right),$$

thus,  $K_x \otimes K_x$  is a simple function with values in the Banach space  $\mathscr{L}(\mathscr{H}_K)$ . For any given random vector x with values in the Hilbert space  $(\mathscr{X}, \tau_N(\mathscr{X}))$ , it is known that there exists a simple function sequence  $\{x_n, n \ge 0\}$  with values in  $\mathscr{X}$ , such that  $\lim_{n\to\infty} ||x - x_n|| = 0$  a.s.. This together with the reproducing property of  $\mathscr{H}_K$  and the symmetry of Mercer kernel K gives

$$\|K_{x} - K_{x_{n}}\|_{K}^{2} = \langle K_{x} - K_{x_{n}}, K_{x} - K_{x_{n}} \rangle_{K}$$
  
=  $K(x, x) - 2K(x, x_{n}) + K(x_{n}, x_{n}).$  (C.58)

Noting the continuity of Mercer kernel K and Assumption 5.3, by (C.58), we get

$$\lim_{n \to \infty} \|K_x - K_{x_n}\|_K = 0 \text{ a.s.}$$
(C.59)

It follows from (C.53) and the reproducing property of  $\mathscr{H}_K$  that

$$\|K_{x} \otimes K_{x} - K_{x_{n}} \otimes K_{x_{n}}\|_{\mathscr{L}(\mathscr{H}_{K})}$$

$$\leq \|(K_{x} - K_{x_{n}}) \otimes K_{x}\|_{\mathscr{L}(\mathscr{H}_{K})} + \|K_{x_{n}} \otimes (K_{x} - K_{x_{n}})\|_{\mathscr{L}(\mathscr{H}_{K})}$$

$$= \sup_{\|f\|_{K}=1} \|((K_{x} - K_{x_{n}}) \otimes K_{x}) f\|_{K} + \sup_{\|f\|_{K}=1} \|(K_{x_{n}} \otimes (K_{x} - K_{x_{n}})) f\|_{K}$$

$$= \sup_{\|f\|_{K}=1} \|f(x)(K_{x} - K_{x_{n}})\|_{K} + \sup_{\|f\|_{K}=1} \|(f(x) - f(x_{n}))K_{x_{n}}\|_{K}$$

$$\leq \|K_{x}\|_{K} \|K_{x} - K_{x_{n}}\|_{K} + \|K_{x_{n}}\|_{K} \|K_{x} - K_{x_{n}}\|_{K}$$

$$\leq 2 \sup_{x \in \mathscr{X}} \sqrt{K(x, x)} \|K_{x} - K_{x_{n}}\|_{K}$$

$$(C.60)$$

By Assumption 5.3 and (C.59)-(C.60), we have

$$\lim_{n\to\infty} \|K_x \otimes K_x - K_{x_n} \otimes K_{x_n}\|_{\mathscr{L}(\mathscr{H}_K)} = 0 \text{ a.s.}$$

Noting that  $K_{x_n} \otimes K_{x_n}$  is the simple function with values in Banach space  $\mathscr{L}(\mathscr{H}_K)$ , by Definition A.3, it is known that  $K_x \otimes K_x$  is strongly measurable with respect to the topology  $\tau_N(\mathscr{L}(\mathscr{H}_K))$ , which together with Pettis measurability theorem shows that  $K_x \otimes K_x$  is the random element with values in Banach space  $(\mathscr{L}(\mathscr{H}_K), \tau_N(\mathscr{L}(\mathscr{H}_K)))$ . Thus, by Assumption 5.3, we get  $K_{x_j(i)} \otimes K_{x_j(i)} \in L^1(\Omega; \mathscr{L}(\mathscr{H}_K))$ , which together with Lemma A.9 gives the fact that  $\mathbb{E}[K_{x_j(i)} \otimes K_{x_j(i)} | \mathcal{F}(kh-1)]$  uniquely exists. Let  $\{g(k), \mathcal{F}(kh-1), k \geq 0\}$  be the  $L_2$ -bounded adaptive sequence with values in  $\mathscr{H}_K$ , by Assumption 5.3, Proposition 2.6 and the condition (5.3), we obtain

$$\sum_{j=1}^{N} \sum_{k=0}^{\infty} \mathbb{E} \left[ \left\| N_{j}g(k) - \sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[ H_{j}^{*}(i)H_{j}(i)g(k) \middle| \mathcal{F}(kh-1) \right] \right\|_{K}^{2} \right]$$

$$= \sum_{j=1}^{N} \sum_{k=0}^{\infty} \mathbb{E} \left[ \left\| N_{j}g(k) - \sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[ K_{x_{j}(i)} \otimes K_{x_{j}(i)}g(k) \middle| \mathcal{F}(kh-1) \right] \right\|_{K}^{2} \right]$$

$$= \sum_{j=1}^{N} \sum_{k=0}^{\infty} \mathbb{E} \left[ \left\| \left( N_{j} - \sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[ K_{x_{j}(i)} \otimes K_{x_{j}(i)} \middle| \mathcal{F}(kh-1) \right] \right) g(k) \right\|_{K}^{2} \right]$$

$$< \infty.$$
(C.61)

Denote  $\rho_0 = N \sup_{x \in \mathscr{X}} K(x, x)$ . Given the integer h > 0, by Assumption 5.3 and (C.53), we have

$$\begin{split} \sup_{k\geq 0} \left( \mathbb{E} \left[ \left\| \mathcal{H}^{*}(k) \mathcal{H}(k) \right\|_{\mathscr{L}(\mathscr{H}_{K}^{N})}^{2^{\max\{h,2\}}} \middle| \mathcal{F}(k-1) \right] \right)^{\frac{1}{2^{\max\{h,2\}}}} \\ &\leq N \sup_{k\geq 0} \left( \mathbb{E} \left[ \sup_{i\in\mathcal{V}} \left\| H_{i}^{*}(k) H_{i}(k) \right\|_{\mathscr{L}(\mathscr{H}_{K})}^{2^{\max\{h,2\}}} \middle| \mathcal{F}(k-1) \right] \right)^{\frac{1}{2^{\max\{h,2\}}}} \\ &= N \sup_{k\geq 0} \left( \mathbb{E} \left[ \sup_{i\in\mathcal{V}} \left\| K_{x_{i}(k)} \otimes K_{x_{i}(k)} \right\|_{\mathscr{L}(\mathscr{H}_{K})}^{2^{\max\{h,2\}}} \middle| \mathcal{F}(k-1) \right] \right)^{\frac{1}{2^{\max\{h,2\}}}} \\ &\leq N \sup_{k\geq 0} \left( \mathbb{E} \left[ \sup_{i\in\mathcal{V}} \sup_{\|f\|_{K}=1} \left\| \left( K_{x_{i}(k)} \otimes K_{x_{i}(k)} \right) f \right\|_{K}^{2^{\max\{h,2\}}} \middle| \mathcal{F}(k-1) \right] \right)^{\frac{1}{2^{\max\{h,2\}}}} \\ &= N \sup_{k\geq 0} \left( \mathbb{E} \left[ \sup_{i\in\mathcal{V}} \sup_{\|f\|_{K}=1} \left\| f(x_{i}(k)) K_{x_{i}(k)} \right\|_{K}^{2^{\max\{h,2\}}} \middle| \mathcal{F}(k-1) \right] \right)^{\frac{1}{2^{\max\{h,2\}}}} \\ &\leq N \sup_{k\geq 0} \left( \mathbb{E} \left[ \sup_{i\in\mathcal{V}} \sup_{\|f\|_{K}=1} \left\| f(x_{i}(k)) \right\|^{2^{\max\{h,2\}}} \left\| K_{x_{i}(k)} \right\|_{K}^{2^{\max\{h,2\}}} \middle| \mathcal{F}(k-1) \right] \right)^{\frac{1}{2^{\max\{h,2\}}}} \\ &\leq N \sup_{k\geq 0} \left( \mathbb{E} \left[ \sup_{i\in\mathcal{V}} \sup_{\|f\|_{K}=1} \left\| f(x_{i}(k)) \right\|^{2^{\max\{h,2\}}} \left\| K_{x_{i}(k)} \right\|_{K}^{2^{\max\{h,2\}}} \middle| \mathcal{F}(k-1) \right] \right)^{\frac{1}{2^{\max\{h,2\}}}} \\ &\leq N \sup_{k\geq 0} \left( \mathbb{E} \left[ \sup_{i\in\mathcal{V}} \sup_{\|f\|_{K}=1} \left\| f \right\|_{K}^{2^{\max\{h,2\}}} \left( \sup_{x\in\mathscr{X}} K(x,x) \right)^{2^{\max\{h,2\}}} \middle| \mathcal{F}(k-1) \right] \right)^{\frac{1}{2^{\max\{h,2\}}}} \\ &\leq N \sup_{x\in\mathscr{X}} K(x,x) \\ &\leq N \sup_{x\in\mathscr{X}} K(x,x) \\ &= \rho_{0} \text{ a.s.} \end{split}$$

It follows from Condition 4.2 that there exists a constant  $j_0 > 0$  and an integer  $t_0 > 0$ , such that

 $\sup_{k\geq 0}(4\rho_0 a(k) + 4\|\mathcal{L}_{\mathcal{G}}\|b(k)) \leq j_0$  and  $\sup_{k\geq t_0}(a(k) + b(k))(4\rho_0 + 4\|\mathcal{L}_{\mathcal{G}}\|) \leq 1$ . Noting that

$$\|4a(k)\mathcal{H}^*(k)\mathcal{H}(k) + 4b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{H}_K}\|_{\mathscr{L}(\mathscr{H}_K^N)} \leq \begin{cases} j_0, & k < t_0; \\ 1, & k \ge t_0 \end{cases}$$
a.s. (C.62)

By (C.62), we get

$$\left\|I_{\mathscr{H}_{K}^{N}}-4\left(a(k)\mathcal{H}^{*}(k)\mathcal{H}(k)+b(k)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{H}_{K}}\right)\right\|_{\mathscr{L}\left(\mathscr{H}_{K}^{N}\right)} \leq 1+\Gamma(k) \text{ a.s.},\tag{C.63}$$

where

$$\Gamma(k) = \begin{cases} j_0, & k < t_0, \\ 0, & k \ge t_0, \end{cases}$$

and satisfies  $\sum_{k=0}^{\infty} \Gamma(k) < \infty$ . It follows from (C.63) that

$$\mathbb{E}\left[\left\|I_{\mathscr{H}_{K}^{N}}-4\left(a(k)\mathcal{H}^{*}(k)\mathcal{H}(k)+b(k)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{H}_{K}}\right)\right\|_{\mathscr{L}\left(\mathscr{H}_{K}^{N}\right)}\right|\mathcal{F}(kh-1)\right] \leq 1+\Gamma(k) \text{ a.s.}$$
(C.64)

Hence, combining (C.61)-(C.64) and Theorem 4.10 gives the fact that the algorithm (5.2) is mean square and almost surely strongly consistent.  $\Box$ 

Proof of Corollary 5.5. It follows from Assumption 5.3 and Theorem 5.4 that  $K_{x_j(i)} \otimes K_{x_j(i)} \in L^1(\Omega; \mathscr{L}(\mathscr{H}_K))$ , which together with Lemma A.9 implies that  $\mathbb{E}[K_{x_j(i)} \otimes K_{x_j(i)} | \mathcal{F}(kh-1)]$  uniquely exists. Let  $\{g(k), \mathcal{F}(kh-1), k \geq 0\}$  be a  $L_2$ -bounded adaptive sequence with values in  $\mathscr{H}_K$ , by the condition (5.4), we get

$$\sum_{j=1}^{N} \sum_{k=0}^{\infty} \mathbb{E} \left[ \left\| \left( N_{j} - \sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[ K_{x_{j}(i)} \otimes K_{x_{j}(i)} \middle| \mathcal{F}(kh-1) \right] \right) g(k) \right\|_{K}^{2} \right]$$

$$\leq \sum_{j=1}^{N} \sum_{k=0}^{\infty} \mathbb{E} \left[ \left\| N_{j} - \sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[ K_{x_{j}(i)} \otimes K_{x_{j}(i)} \middle| \mathcal{F}(kh-1) \right] \right\|_{\mathscr{L}(\mathscr{H}_{K})}^{2} \| g(k) \|_{K}^{2} \right]$$

$$\leq \sum_{j=1}^{N} \sum_{k=0}^{\infty} \mathbb{E} \left[ \max_{j \in \mathcal{V}} \left\| N_{j} - \sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[ K_{x_{j}(i)} \otimes K_{x_{j}(i)} \middle| \mathcal{F}(kh-1) \right] \right\|_{\mathscr{L}(\mathscr{H}_{K})}^{2} \| g(k) \|_{K}^{2} \right]$$

$$\leq N \mu_{0} \sup_{k \geq 0} \mathbb{E} \left[ \| g(k) \|_{K}^{2} \right] \sum_{k=0}^{\infty} \tau(k)$$

$$< \infty,$$

where the last inequality is obtained from  $\sum_{k=0}^{\infty} \tau(k) < \infty$ . This together with Theorem 5.4 implies that the algorithm (5.2) is both mean square and almost surely strongly consistent.  $\Box$ 

Proof of Corollary 5.7. Noting that  $\{(x_1(k), \dots, x_N(k)), k \ge 0\}$  and  $\{(v_1(k), \dots, v_N(k)), k \ge 0\}$ 

0} are both i.i.d. sequences and they are mutually independent, we have

$$\sum_{i=kh}^{k+1)h-1} \mathbb{E} \left[ K_{x_j(i)} \otimes K_{x_j(i)} \middle| \mathcal{F}(kh-1) \right] = \sum_{i=kh}^{(k+1)h-1} \mathbb{E} \left[ K_{x_j(i)} \otimes K_{x_j(i)} \right]$$
$$= h \mathbb{E} \left[ K_{x_j(0)} \otimes K_{x_j(0)} \right].$$

Denote  $N_j : \mathscr{H}_K \to \mathscr{H}_K$  by

$$N_j = h\mathbb{E}\left[K_{x_j(0)} \otimes K_{x_j(0)}\right], \ j \in \mathcal{V}$$

It follows from Lemma A.10 that  $N_j \in \mathscr{L}(\mathscr{H}_K)$  is a positive self-adjoint operator. Noting that

$$\left\| N_j - \sum_{i=kh}^{(k+1)h-1} \mathbb{E}\left[ K_{x_j(i)} \otimes K_{x_j(i)} \middle| \mathcal{F}(kh-1) \right] \right\|_{\mathscr{L}(\mathscr{H}_K)}^2 = 0 \text{ a.s.},$$

by the condition (5.5), we get

$$\left\langle \sum_{j=1}^{N} N_j f, f \right\rangle_K = h \left\langle \mathbb{E} \left[ \sum_{j=1}^{N} K_{x_j(0)} \otimes K_{x_j(0)} \right] f, f \right\rangle_K > 0,$$

where f is an arbitrary non-zero function in  $\mathscr{H}_K$ . Hence, it follows from Corollary 5.5 that the algorithm (5.2) is both mean square and almost surely strongly consistent.

#### APPENDIX D

### KEY LEMMAS

**Lemma D.1** ([86]). Let  $\{x(k), \mathcal{F}(k)\}$ ,  $\{\alpha(k), \mathcal{F}(k)\}$ ,  $\{\beta(k), \mathcal{F}(k)\}$  and  $\{\gamma(k), \mathcal{F}(k)\}$  be nonnegative adaptive sequences satisfying

$$\mathbb{E}[x(k+1)|\mathcal{F}(k)] \le (1+\alpha(k))x(k) - \beta(k) + \gamma(k), \ k \ge 0 \ a.s.,$$

and  $\sum_{k=0}^{\infty} (\alpha(k) + \gamma(k)) < \infty$  a.s. Then x(k) converges to a finite random variable a.s., and  $\sum_{k=0}^{\infty} \beta(k) < \infty$  a.s.

**Lemma D.2.** Let  $\{a(i), i \in \Lambda\}$  be a nonnegative real sequence, where  $\Lambda \subseteq \mathbb{N}$  and  $\sum_{i \in \Lambda} a(i) < \infty$ ,  $\{b(k,i), k \in \mathbb{N}, i \in \Lambda\}$  be a double index real sequence. If  $\lim_{k\to\infty} b(k,i) = 0$ ,  $\forall i \in \Lambda$ , and there exists a constant c > 0 such that  $|b(k,i)| \leq c$ ,  $\forall k \in \mathbb{N}$ ,  $\forall i \in \Lambda$ , then

$$\lim_{k \to \infty} \sum_{i \in \Lambda} a(i)b(k,i) = 0.$$
(D.1)

*Proof.* For any given  $\varepsilon > 0$ , it follows from  $\sum_{i \in \Lambda} a(i) < \infty$  that there exists an integer N > 0, such that  $\sum_{i \in \Lambda_N} a(i) < \varepsilon$ , where  $\Lambda_N = \{i \in \Lambda : i \ge N+1\}$ . We obtain

$$\left|\sum_{i\in\Lambda_N}a(i)b(k,i)\right|\leq c\varepsilon, \ \forall k\geq 0.$$

On the other hand, noting that  $\lim_{k\to\infty} b(k,i) = 0$ ,  $\forall i \in \Lambda$ , it follows that there exists a constant  $M_i > 0$ , such that  $|b(k,i)| \le \varepsilon$ ,  $\forall k \ge M_i$ . Denote  $M = \max\{M_i : i \in \{1, \dots, N\} \cap \Lambda\}$ . It follows from  $|b(k,i)| \leq \varepsilon, \ \forall \ k \geq M$  that

$$\left|\sum_{i\in\Lambda}a(i)b(k,i)\right| \le \left|\sum_{i\in\{1,\cdots,N\}\cap\Lambda}a(i)b(k,i)\right| + \left|\sum_{i\in\Lambda_N}a(i)b(k,i)\right| \le \varepsilon \sum_{i\in\Lambda}a(i) + c\varepsilon, \ k \ge M,$$
  
which gives (D.1).

which gives (D.1).

**Lemma D.3.** Let  $\mathscr{X}$  be a Hilbert space,  $H \in \mathscr{L}(\mathscr{X})$  be a strictly positive self-adjoint operator, and  $\{\mu(k), k \ge 0\}$  be a real sequence monotonically decreasing to 0 with  $\sum_{k=0}^{\infty} \mu(k) = \infty$ . Then there exist positive constants M and d, such that

$$\sup_{s\geq 0} \left\| \left( \prod_{j=t}^{s} (I_{\mathscr{X}} - \mu(j)H) \right) x \right\| \le M^{d} \|x\|, \ \forall \ x \in \mathscr{X}, \ \forall \ t \ge 0,$$
(D.2)

and

$$\lim_{k \to \infty} \left( \prod_{j=0}^{k} (I_{\mathscr{X}} - \mu(j)H) \right) x = 0, \ \forall \ x \in \mathscr{X}.$$
 (D.3)

*Proof.* Noting that  $H \in \mathscr{L}(\mathscr{X})$  is the bounded self-adjoint operator, it is shown that H has the following spectral decomposition

$$H = \int_{-\infty}^{+\infty} \lambda \, \mathrm{d}p_{\lambda} = \int_{\sigma(H)} \lambda \, \mathrm{d}p_{\lambda}$$

From the property of the self-adjoint operator, we have

$$\prod_{j=0}^{k} (I_{\mathscr{X}} - \mu(j)H)^2 = \int_{\sigma(H)} \prod_{j=0}^{k} (1 - \mu(j)\lambda)^2 \,\mathrm{d}p_{\lambda}, \tag{D.4}$$

which together with (D.4) gives

$$\left\| \prod_{j=0}^{k} (I_{\mathscr{X}} - \mu(j)H)x \right\|^{2} = \left\langle \prod_{j=0}^{k} (I_{\mathscr{X}} - \mu(j)H)^{2}x, x \right\rangle$$
$$= \int_{\sigma(H)} \prod_{j=0}^{k} (1 - \mu(j)\lambda)^{2} d\langle p_{\lambda}x, x \rangle, \ \forall \ x \in \mathscr{X}.$$
(D.5)

It follows from H > 0 that  $\sigma(H) \subset [0, ||H||]$ , which leads to

$$(1 - \mu(j)\lambda)^2 \le \max\{(1 - \mu(j)||H||)^2, 1\}, \ \forall \ \lambda \in \sigma(H).$$

Noting that  $\mu(k) \to 0, k \to \infty$ , it follows that there exists an integer d > 0 such that  $\mu(j) \leq d$  $||H||^{-1}, \forall j > 2d$ , which further shows that  $(1 - b(j)\lambda)^2 \leq 1, \forall j > 2d$ . Denote  $M = \max\{(1 - b(j)\lambda)^2 \leq 1, \forall j > 2d\}$ .  $\mu(j) \|H\|^2, 1: 0 \le j \le 2d\}$ , we have

$$\prod_{j=t}^{s} (1 - \mu(j)\lambda)^2 \le M^{2d}, \ \forall \lambda \in \sigma(H), \ \forall \ t \ge 0,$$

and

$$\left\|\prod_{j=t}^{s} (I_{\mathscr{X}} - \mu(j)H)x\right\|^{2} \le M^{2d} \int_{\sigma(H)} \mathbf{d}\langle p_{\lambda}x, x\rangle = M^{2d} \|x\|^{2}, \ \forall \ x \in \mathscr{X}, \ \forall \ t \ge 0.$$

This gives (D.2). Noting the inequality  $1 - a \le e^{-a}$ ,  $\forall a \ge 0$ , we obtain

$$\prod_{j=0}^{k} (1 - \mu(j)\lambda)^2 \le M^{2d} \mathbf{e}^{-2\lambda \sum_{j=2d+1}^{k} \mu(j)}.$$

It follows from  $\sum_{j=0}^{\infty} \mu(k) = \infty$  that

$$\lim_{k \to \infty} \prod_{j=0}^{k} (1 - \mu(j)\lambda)^2 = 0, \ \forall \ \lambda \in \sigma(H) \cap \mathbb{R}^+.$$
(D.6)

Since H is the strictly positive self-adjoint operator, it follows that 0 is not in the point spectrum of the operator H, which gives

$$\int_{\sigma(H)} \prod_{j=0}^{k} (1-\mu(j)\lambda)^2 \,\mathrm{d}\langle p_\lambda x, x\rangle = \int_{\sigma(H)\cap\mathbb{R}^+} \prod_{j=0}^{k} (1-\mu(j)\lambda)^2 \,\mathrm{d}\langle p_\lambda x, x\rangle, \ \forall \ x \in \mathscr{X}.$$

By (D.5), (D.2), (D.6) and the dominated convergence theorem, we get

$$\lim_{k \to \infty} \left\| \prod_{j=0}^{k} (I_{\mathscr{X}} - \mu(j)H)x \right\|^{2} = \int_{\sigma(H) \cap \mathbb{R}^{+}} \lim_{k \to \infty} \prod_{j=0}^{k} (1 - \mu(j)\lambda)^{2} \, \mathrm{d}\langle p_{\lambda}x, x \rangle = 0, \ \forall \ x \in \mathscr{X},$$
  
hich gives (D.3).

which gives (D.3).

**Lemma D.4.** Let  $\mathscr{X}$  be a Hilbert space,  $H \in \mathscr{L}(\mathscr{X})$  be a strictly positive self-adjoint operator, and  $\{x(k), k \ge 0\}$  be a  $L_2$ -bounded random sequence with values in Hilbert space. If  $\{\mu(k), k \ge 0\}$ 0 and  $\{\gamma(k), k \ge 0\}$  are both real sequences monotonically decreasing to 0 with  $\sum_{k=0}^{\infty} \mu(k) = 0$  $\infty$  and  $\sum_{k=0}^{\infty} \gamma(k) < \infty$ , then

$$\lim_{k \to \infty} \sum_{i=0}^{k} \gamma(i) \left( \mathbb{E} \left[ \left\| \left( \prod_{j=i+1}^{k} (I_{\mathscr{X}} - \mu(j)H) \right) x(i) \right\|^2 \right] \right)^{\frac{1}{2}} = 0.$$

Proof. Denote

$$b(k,i) = \left( \mathbb{E}\left[ \left\| \prod_{j=i+1}^{k} (I_{\mathscr{X}} - \mu(j)H)x(i) \right\|^2 \right] \right)^{\frac{1}{2}}, \ \forall \ k, i \ge 0.$$

it follows from Lemma D.3 that

$$0 \le b(k,i) \le M^d \sup_{i \ge 0} \left( \mathbb{E} \left[ \|x(i)\|^2 \right] \right)^{\frac{1}{2}} < \infty, \ \forall \ k, i \ge 0,$$
(D.7)

and

$$\left\|\prod_{j=i+1}^{k} (I_{\mathscr{X}} - \mu(j)H)x(i)\right\| \le M^d \|x(i)\| \text{ a.s., } \forall i \ge 0.$$

Thus, by the Lebesgue dominated convergence theorem and Lemma D.3, we get

$$\lim_{k \to \infty} \mathbb{E}\left[ \left\| \prod_{j=i+1}^{k} (I_{\mathscr{X}} - \mu(j)H)x(i) \right\|^2 \right] = 0, \ \forall \ i \ge 0,$$
(D.8)

which further gives  $\lim_{k\to\infty} b(k,i) = 0, \forall i \ge 0$ . Noting that  $\sum_{k=0}^{\infty} \gamma(k) < \infty$ , by (D.7)-(D.8) and Lemma D.2, we obtain

$$\lim_{k \to \infty} \sum_{i=0}^{k} \gamma(i) \left( \mathbb{E} \left[ \left\| \left( \prod_{j=i+1}^{k} (I_{\mathscr{X}} - \mu(j)H) \right) x(i) \right\|^2 \right] \right)^{\frac{1}{2}} = 0.$$

**Lemma D.5.** Let  $\mathcal{A} = [a_{ij}] \in \mathbb{R}^{N \times N}$  be the adjacency matrix of an undirected connected graph  $\mathcal{G}$ ,  $\mathcal{L}_{\mathcal{G}}$  be the Laplacian matrix,  $H_i \in \mathscr{L}(\mathscr{X})$ ,  $i = 1, \dots, N$  are positive self-adjoint operators. If

$$\sum_{i=1}^{N} H_i > 0, \tag{D.9}$$

then diag $\{H_1, \cdots, H_N\} + \mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}} > 0.$ 

*Proof.* Given the non-zero element  $x = (x_1, \dots, x_N)$  with values in Hilbert space  $\mathscr{X}^N$ , where  $x_i \in \mathscr{X}, i = 1, \dots, N$ . Here, we will prove  $\langle (\operatorname{diag}\{H_1, \dots, H_N\} + \mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})x, x \rangle_{\mathscr{X}^N} > 0$  in two steps.

If there exists a non-zero element a ∈ X with x<sub>1</sub> = ··· = x<sub>N</sub> = a, then x = 1<sub>N</sub> ⊗ a. Noting that L<sub>G</sub>1<sub>N</sub> = 0, it follows from (D.9) that

$$\langle (\operatorname{diag}\{H_1,\cdots,H_N\} + \mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}) x, x \rangle_{\mathscr{X}^N}$$

$$= \langle (\operatorname{diag}\{H_{1}, \cdots, H_{N}\} + \mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}) (\mathbf{1}_{N} \otimes a), (\mathbf{1}_{N} \otimes a) \rangle_{\mathscr{X}^{N}} \\ = \langle \operatorname{diag}\{H_{1}, \cdots, H_{N}\} (\mathbf{1}_{N} \otimes a), (\mathbf{1}_{N} \otimes a) \rangle_{\mathscr{X}^{N}} \\ + \langle (\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}) (\mathbf{1}_{N} \otimes a), (\mathbf{1}_{N} \otimes a) \rangle_{\mathscr{X}^{N}} \\ = \left\langle \left(\sum_{i=1}^{N} H_{i}\right) a, a \right\rangle_{\mathscr{X}} + \langle (\mathcal{L}_{\mathcal{G}}\mathbf{1}_{N}) \otimes a, (\mathbf{1}_{N} \otimes a) \rangle_{\mathscr{X}^{N}} \\ = \left\langle \left(\sum_{i=1}^{N} H_{i}\right) a, a \right\rangle_{\mathscr{X}} + \langle (\mathcal{L}_{\mathcal{G}}\mathbf{1}_{N}) \otimes a, (\mathbf{1}_{N} \otimes a) \rangle_{\mathscr{X}^{N}} \\ > 0. \qquad (D.10)$$

If there exist 1 ≤ i<sub>0</sub> ≠ j<sub>0</sub> ≤ N, such that x<sub>i0</sub> ≠ x<sub>j0</sub>. It follows from H<sub>i</sub> ≥ 0, i = 1, · · · , N that diag{H<sub>1</sub>, · · · , H<sub>N</sub>} ≥ 0. Noting that the graph is undirected and connected, it is shown that a<sub>ij</sub> = a<sub>ji</sub> > 0, 1 ≤ i ≠ j ≤ N. Thus, we get

$$\langle (\operatorname{diag}\{H_1, \cdots, H_N\} + \mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}) x, x \rangle_{\mathscr{X}^N}$$

$$\geq \langle (\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}) x, x \rangle_{\mathscr{X}^N}$$

$$= \frac{1}{2} \sum_{i=1}^N \sum_{j=1}^N a_{ij} \|x_i - x_j\|_{\mathscr{X}}^2$$

$$\geq \frac{1}{2} a_{i_0 j_0} \|x_{i_0} - x_{j_0}\|_{\mathscr{X}}^2$$

$$> 0.$$

$$(D.11)$$

Combining (D.10)-(D.11) gives that diag $\{H_1, \cdots, H_N\} + \mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}} \in \mathscr{L}(\mathscr{X}^N)$  is strictly positive.

**Lemma D.6.** Let  $\{a(k), k \ge 0\}$  and  $\{b(k), k \ge 0\}$  be monotonically decreasing sequences of positive real numbers. If

$$\max\{a(k) - a(k+1), b(k) - a(k)\} = \mathcal{O}\left(a^2(k) + b^2(k)\right),$$
(D.12)

then

$$\max_{ih \le s \le (i+1)h-1} \left( \frac{1}{h} \left( \sum_{s=ih}^{(i+1)h-1} b(s) \right) - a(s) \right)^2 = \mathcal{O} \left( a^4(i) + b^4(i) \right), \ \forall \ h = 1, 2, \dots$$
(D.13)

*Proof.* Let h be any given positive integer. Noting that  $\{a(k), k \ge 0\}$  and  $\{b(k), k \ge 0\}$  are both monotonically decreasing sequences, it follows that

$$\begin{cases} \frac{1}{h} \left( \sum_{s=ih}^{(i+1)h-1} b(s) \right) \in [b((i+1)h-1), b(ih)], \\ a(s) \in [a((i+1)h-1), a(ih)], \ ih \le s \le (i+1)h-1, \end{cases}$$

from which we obtain

$$\max_{\substack{ih \le s \le (i+1)h-1 \\ ih \le s \le (i+1)h-1 \\ lim \le (i+1)h-1 \\ lim \le s \le (i+1)h-1 \\ lim \le (i+1$$

It follows from (D.12) that there exists a constant C > 0, such that

$$\max\left\{(a(k) - a(k+1))^2, (b(k) - a(k))^2\right\} \le C\left(a^4(k) + b^4(k)\right),\tag{D.15}$$

which gives

$$\max\left\{(b((i+1)h-1) - a((i+1)h-1))^2, (b(ih) - a(ih))^2\right\} \le C\left(a^4(i) + b^4(i)\right). \quad (D.16)$$

Combining (D.15) and the monotonicity of sequences  $\{a(k), k \ge 0\}$  and  $\{b(k), k \ge 0\}$  leads to

$$(b((i+1)h-1) - a(ih))^{2} = (b((i+1)h-1) - a((i+1)h-1) + a((i+1)h-1) - a(ih))^{2} = \left(b((i+1)h-1) - a((i+1)h-1) + \sum_{s=ih}^{(i+1)h-2} (a(s+1) - a(s))\right)^{2} \le h \left( (b((i+1)h-1) - a((i+1)h-1))^{2} + \sum_{s=ih}^{(i+1)h-2} (a(s) - a(s+1))^{2} \right) \le h \left( C \left( a^{4}((i+1)h-1) + b^{4}((i+1)h-1) \right) + C \sum_{s=ih}^{(i+1)h-2} (a^{4}(s) + b^{4}(s)) \right) \le Ch^{2} \left( a^{4}(i) + b^{4}(i) \right).$$
(D.17)

Similarly, we obtain

$$\begin{aligned} & (a((i+1)h-1) - b(ih))^2 \\ &= (b(ih) - a((i+1)h-1))^2 \\ &= (b(ih) - a(ih) + a(ih) - a((i+1)h-1))^2 \\ &= \left(b(ih) - a(ih) + \sum_{s=ih}^{(i+1)h-2} (a(s) - a(s+1))\right)^2 \\ &\leq h \left( (b(ih) - a(ih))^2 + \sum_{s=ih}^{(i+1)h-2} (a(s) - a(s+1))^2 \right) \end{aligned}$$

$$\leq h\left(C\left(a^{4}(ih) + b^{4}(ih)\right) + C\sum_{s=ih}^{(i+1)h-2} \left(a^{4}(s) + b^{4}(s)\right)\right)$$
  
$$\leq Ch^{2}\left(a^{4}(i) + b^{4}(i)\right).$$
 (D.18)

Combining (D.14) and (D.16)-(D.18) gives (D.13).

Lemma D.7. For the algorithm (3.6), suppose that Conditions 4.1 and 4.2 hold, and there exists an integer h > 0 and a constant  $\rho > 0$ , such that

$$\sup_{k \ge 0} \mathbb{E} \left[ \left\| \mathcal{H}^*(k) \mathcal{H}(k) \right\|^{2^{\max\{h,2\}}} \middle| \mathcal{F}(k-1) \right]^{\frac{1}{2^{\max\{h,2\}}}} \le \rho \text{ a.s.}$$
(D.19)

Then (I) for any given integer  $n \ge 0$  and  $x \in L^0(\Omega, \mathcal{F}(nh-1); \mathscr{X}^N)$ , there exists a constant  $d_1 > 0$  such that

$$\sup_{m \ge 0} \mathbb{E} \left[ \|\Phi_P(mh-1,nh)x\|^2 \, \Big| \, \mathcal{F}(nh-1) \right] \le d_1 \|x\|^2 \, a.s.; \tag{D.20}$$

(II) for any given integer  $n \ge 0$  and  $x \in L^0(\Omega, \mathcal{F}(n-1); \mathscr{X}^N)$ , then there exists a constant  $d_2 > 0$  such that

$$\sup_{0 \le m \le n+h} \mathbb{E}\left[ \|\Phi_P(m,n)x\|^2 |\mathcal{F}(n-1) \right] \le d_2 \|x\|^2 \text{ a.s.}$$
(D.21)

*Proof.* Denote  $D(k) = a(k)\mathcal{H}^*(k)\mathcal{H}(k) + b(k)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}$  and  $P(k) = I_{\mathscr{X}^N} - D(k)$ . Noting that  $\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}} \geq 0$ , we have  $D(k) \geq 0$  a.s. Let x be a random element with values in Hilbert space  $(\mathscr{X}^N, \tau_{\mathbf{N}}(\mathscr{X}^N))$ , we get

$$\begin{split} \|\Phi_{P}(m,n)x\|^{2} \\ &= \langle \Phi_{P}(m,n)x, \Phi_{P}(m,n)x \rangle \\ &= \langle x, \Phi_{P}^{*}(m,n)\Phi_{P}(m,n)x \rangle \\ &= \langle x, (I_{\mathscr{X}^{N}} - D^{*}(n)) \cdots (I_{\mathscr{X}^{N}} - D^{*}(m))(I_{\mathscr{X}^{N}} - D(m)) \cdots (I_{\mathscr{X}^{N}} - D(n))x \rangle \\ &= \langle x, x - 2\sum_{k=n}^{m} D(k)x + \sum_{k=n}^{m} M_{k}x \rangle \\ &= \|x\|^{2} - 2\sum_{k=n}^{m} \langle x, D(k)x \rangle + \langle x, \sum_{k=n}^{m} M_{k}x \rangle \\ &\leq \|x\|^{2} + \langle x, \sum_{k=n}^{m} M_{k}x \rangle \\ &\leq \left(1 + \sum_{k=n}^{m} \|M_{k}\|\right) \|x\|^{2} \text{ a.s., } 0 \leq n \leq m, \end{split}$$
(D.22)

66

where  $M_k, k = 2, \dots, 2(m - n + 1)$  denote the k-th order terms in the binomial expansion of  $\Phi_P^*(m, n)\Phi_P(m, n)$ . For  $0 \le m - n \le h$ , by the conditional Lyapunov inequality and (D.19), we obtain

$$\sup_{k \ge 0} \mathbb{E} \left[ \|D(k)\|^i | \mathcal{F}(k-1) \right] \le \sup_{k \ge 0} \mathbb{E} \left[ \|D(k)\|^{2^h} | \mathcal{F}(k-1) \right]^{\frac{i}{2h}} \le \rho_0^i (a(k) + b(k))^i \text{ a.s.},$$
  
$$2 \le i \le 2^h, \text{ (D.23)}$$

where  $\rho_0 := \rho + \|\mathcal{L}_{\mathcal{G}}\|$ . Note that

$$\mathbb{E}\left[\|D(k)\|^{i} | \mathcal{F}(n-1)\right]$$
  
=  $\mathbb{E}\left[\mathbb{E}\left[\|D(k)\|^{i} | \mathcal{F}(k-1)\right] | \mathcal{F}(n-1)\right], \ 2 \le i \le 2^{h}, \ k \ge n.$  (D.24)

Since the real sequences  $\{a(k), k \ge 0\}$  and  $\{b(k), k \ge 0\}$  are both monotonically decreasing to 0, it follows that there exists a constant  $c_0 > 0$  such that  $\sup_{k\ge 0}(a(k) + b(k)) \le c_0$ . For  $0 \le m - n \le h$ , from the definition of  $M_k$  and (D.23)-(D.24), by termwise multiplication and using the Hölder inequality repeatedly, we have

$$\mathbb{E}[\|M_k\||\mathcal{F}(n-1)] \le \mathbb{C}_{2h}^k \rho_0^k (a(n) + b(n))^k \le 2c_0^{-2} \left(a^2(n) + b^2(n)\right) \mathbb{C}_{2h}^k \rho_0^k c_0^k \text{ a.s.},$$
  
$$k = 2, \cdots, 2(m-n+1). \quad (D.25)$$

Denote  $c_k = 2c_0^{-2}(1 + \rho_0 c_0)^{2k}$ , by (D.25), we get

$$\sum_{k=n}^{m} \mathbb{E}[\|M_k\| | \mathcal{F}(n-1)] \le c_{m-n+1} \left( a^2(n) + b^2(n) \right) \text{ a.s.}$$
(D.26)

If  $x \in L^0(\Omega, \mathcal{F}(n-1); \mathscr{X}^N)$ , substituting (D.26) into (D.22) gives

$$\mathbb{E}\left[\left\|\Phi_{P}(m,n)x\right\|^{2}\right|\mathcal{F}(n-1)\right] \leq \left(1 + c_{h+1}a^{2}(n) + c_{h+1}b^{2}(n)\right)\|x\|^{2} \text{ a.s.}$$
(D.27)

If  $x \in L^0(\Omega, \mathcal{F}(ih-1); \mathscr{X}^N)$ , it follows from (D.27) that

$$\mathbb{E}\left[\left\|\Phi_{P}((i+1)h-1,ih)x\right\|^{2}\right|\mathcal{F}(ih-1)\right] \leq \left(1+c_{h}a^{2}(i)+c_{h}b^{2}(i)\right)\|x\|^{2} \text{ a.s., } i \geq 0.$$
(D.28)

(I) For the given integer n and  $x \in L^0(\Omega, \mathcal{F}(nh-1); \mathscr{X}^N)$ , by (D.28), we obtain

$$\mathbb{E} \left[ \|\Phi_P(mh-1,nh)x\|^2 | \mathcal{F}(nh-1) \right] \\ = \mathbb{E} \left[ \|\Phi_P(mh-1,(m-1)h)\Phi_P((m-1)h-1,nh)x\|^2 | \mathcal{F}(nh-1) \right] \\ = \mathbb{E} \left[ \mathbb{E} \left[ \|\Phi_P(mh-1,(m-1)h) \times \Phi_P((m-1)h-1,nh)x\|^2 | \mathcal{F}((m-1)h-1) \right] | \mathcal{F}(nh-1) \right] \\ \times \Phi_P((m-1)h-1,nh)x\|^2 | \mathcal{F}((m-1)h-1) \right] | \mathcal{F}(nh-1) \right]$$

$$\leq \left(1 + c_h a^2 (m-1) + c_h b^2 (m-1)\right) \mathbb{E} \left[ \|\Phi_P((m-1)h - 1, nh)x\|^2 \right| \mathcal{F}(nh-1) \right]$$
  
$$\leq \prod_{k=n}^{m-1} \left(1 + c_h a^2 (k) + c_h b^2 (k)\right) \|x\|^2 \text{ a.s.}$$
(D.29)

Denote  $d_1 = \prod_{k=0}^{\infty} (1 + c_h a^2(k) + c_h b^2(k))$ , from (D.29) and Condition 4.2, we get (D.20).

(II) For the given integer n and  $x \in L^0(\Omega, \mathcal{F}(n-1); \mathscr{X}^N)$ , it follows from Condition 4.1 that there exists a constant  $d_2 > 0$ , such that  $\sup_{k \ge 0} (1 + c_{h+1}a^2(k) + c_{h+1}b^2(k)) \le d_2$ , which together with (D.27) gives (D.21).

**Lemma D.8.** For the algorithm (3.6), if Conditions 4.1 and 4.2 hold, and there exists an integer h > 0 and a constant  $\rho > 0$ , such that

$$\sup_{k\geq 0} \mathbb{E}\left[\left\|\mathcal{H}^*(k)\mathcal{H}(k)\right\|^{2^{\max\{h,2\}}} \middle| \mathcal{F}(k-1)\right]^{\frac{1}{2^{\max\{h,2\}}}} \leq \rho \text{ a.s.},$$

then there exits a constant  $d_3 > 0$ , such that

$$\sup_{\substack{k\geq 0\\ \leq d_3 \mathbb{E}\left[\|y\|^2\right], \ \forall \ i\geq 0,}} \mathbb{E}\left[\left\|\left(\prod_{j=i+1}^k \left(I_{\mathscr{X}^N} - \sum_{s=jh}^{(j+1)h-1} (a(s)\mathcal{H}^*(s)\mathcal{H}(s) + b(s)(\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}}))\right)\right)y\right\|^2\right]$$

for any given  $y \in L^0(\Omega, \mathcal{F}((i+1)h-1); \mathscr{X}^N)$ .

*Proof.* Denote  $D'(k) = \sum_{s=kh}^{(k+1)h-1} (a(s)\mathcal{H}^*(s)\mathcal{H}(s) + b(s)(\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}))$  and  $P'(k) = I_{\mathscr{X}^N} - D'(k)$ . Noting that  $\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}} \ge 0$ , we have  $D'(k) \ge 0$  a.s. Let  $x \in L^0(\Omega, \mathcal{F}(nh^2-1); \mathscr{X}^N)$  be a random element with values in the Hilbert space  $\mathscr{X}^N$ . Then we get

$$\mathbb{E}\left[\left\|\Phi_{P'}((n+1)h-1,nh)x\right\|^{2}\right|\mathcal{F}(nh^{2}-1)\right] \leq \left(1 + \sum_{k=nh}^{(n+1)h-1} \mathbb{E}\left[\left\|M'_{k}\right\|\right|\mathcal{F}(nh^{2}-1)\right]\right)\|x\|^{2},$$

where  $M'_k, k = 2, \dots, 2h$  denote the k-th order terms in the binomial expansion of  $\Phi^*_{P'}((n + 1)h - 1, nh)\Phi_{P'}((n + 1)h - 1, nh)$ . It follows from the conditional Lyapunov inequality and Condtion 4.1 that

$$\sup_{k\geq 0} \mathbb{E}\left[ \|D'(k)\|^i \left| \mathcal{F}(k-1) \right] \leq h\rho_0^i \sum_{s=kh}^{(k+1)h-1} (a(s)+b(s))^i \leq h^2 \rho_0^i (a(n)+b(n))^i \text{ a.s.}, \\ 2 \leq i \leq 2^h, \quad (D.30)$$

where  $\rho_0 = \rho + \|\mathcal{L}_{\mathcal{G}}\|$ . Since  $\{a(k), k \ge 0\}$  and  $\{b(k), k \ge 0\}$  are both monotonically decreasing to 0, there exists a constant  $c_0 > 0$  such that  $\sup_{k\ge 0}(a(k) + b(k)) \le c_0$ . From the definition of  $M_k'$ , by termwise multiplication and using the Hölder inequality of the conditional expectation, we have

$$\mathbb{E} \left[ \|M_k'\| | \mathcal{F}(nh^2 - 1) \right] \leq h^2 \mathbb{C}_{2h}^k \rho_0^k (a(n) + b(n))^k \\ \leq 2h^2 c_0^{-2} \left( a^2(n) + b^2(n) \right) \mathbb{C}_{2h}^k \rho_0^k c_0^k \text{ a.s., } k = 2, \cdots, 2h.$$
(D.31)

Noting that (D.31) and  $x \in L^0(\Omega, \mathcal{F}(nh^2 - 1); \mathscr{X}^N)$ , we get

$$\mathbb{E}\left[\left\|\Phi_{P'}((n+1)h-1,nh)x\right\|^{2}\right|\mathcal{F}(nh^{2}-1)\right] \leq \left(1+c'a^{2}(n)+c'b^{2}(n)\right)\|x\|^{2} \text{ a.s.,}$$
(D.32)

where  $c' = 2h^2 c_0^{-2} (1 + \rho_0 c_0)^{2h}$ . By (D.32), we obtain

$$\begin{split} & \mathbb{E}\left[ \|\Phi_{P'}(mh-1,nh)x\|^2 \Big| \,\mathcal{F}(nh^2-1) \right] \\ &= \mathbb{E}\left[ \|\Phi_{P'}(mh-1,(m-1)h)\Phi_{P'}((m-1)h-1,nh)x\|^2 \Big| \,\mathcal{F}(nh^2-1) \right] \\ &= \mathbb{E}\Big[ \mathbb{E}\Big[ \|\Phi_{P'}(mh-1,(m-1)h) \\ &\times \Phi_{P'}((m-1)h-1,nh)x\|^2 |\mathcal{F}((m-1)h^2-1)\Big] |\mathcal{F}(nh^2-1) \Big] \\ &\leq \left(1+c'a^2(m-1)+c'b^2(m-1)\right) \mathbb{E}\left[ \|\Phi_{P'}((m-1)h-1,nh)x\|^2 \Big| \,\mathcal{F}(nh^2-1) \right] \\ &\leq \prod_{k=n}^{m-1} \left(1+c'a^2(k)+c'b^2(k)\right) \|x\|^2 \text{ a.s., } m>n \ge 0. \end{split}$$

Denote  $q_1 = \prod_{k=0}^{\infty} (1 + c'a^2(k) + c'b^2(k))$ , it follows from Condition 4.2 that

$$\sup_{m \ge 0} \mathbb{E} \left[ \|\Phi_{P'}(mh-1,nh)x\|^2 \right] \mathcal{F}(nh^2-1) \leq q_1 \|x\|^2 \text{ a.s.}$$

Following the same way as the proof of Lemma D.7, it shows that there exists a constant  $q_2 > 0$ , such that

$$\sup_{0 \le m \le n+h} \mathbb{E} \left[ \|\Phi_{P'}(m,n)x\|^2 \big| \mathcal{F}(nh-1) \right] \le q_2 \|x\|^2 \text{ a.s.}$$
(D.33)

For any given positive integer j, denote  $m_j = \lfloor \frac{j}{h} \rfloor$ ,  $\widetilde{m}_j = \lceil \frac{j}{h} \rceil$ . Firstly, if  $0 \le i < k - 3h$ , then  $m_k h > \widetilde{m}_{i+1}h$ . Let  $y \in L^0(\Omega, \mathcal{F}((i+1)h-1); \mathscr{X}^N)$  be the random element with values in the Hilbert space  $\mathscr{X}^N$ . Noting that  $0 \le k - m_k h < h$ ,  $0 \le \widetilde{m}_{i+1}h - (i+1) < h$ ,  $\Phi_{P'}(m_k h - 1, \widetilde{m}_{i+1}h)\Phi_{P'}(\widetilde{m}_{i+1}h - 1, i+1)y \in \mathcal{F}(m_k h^2 - 1)$  and  $\Phi_{P'}(\widetilde{m}_{i+1}h - 1, i+1)y \in \mathcal{F}(\widetilde{m}_{i+1}h^2 - 1)$ , by (D.32)-(D.33), we get

$$\mathbb{E} \left[ \|\Phi_{P'}(k,i+1)y\|^2 \right] \\ = \mathbb{E} \left[ \|\Phi_{P'}(k,m_kh)\Phi_{P'}(m_kh-1,\widetilde{m}_{i+1}h)\Phi_{P'}(\widetilde{m}_{i+1}h-1,i+1)y\|^2 \right] \\ \le q_2 \mathbb{E} \left[ \|\Phi_{P'}(m_kh-1,\widetilde{m}_{i+1}h)\Phi_{P'}(\widetilde{m}_{i+1}h-1,i+1)y\|^2 \right]$$

$$= q_{2}\mathbb{E}\left[\mathbb{E}\left[\left\|\Phi_{P'}(m_{k}h-1,\widetilde{m}_{i+1}h)\Phi_{P'}(\widetilde{m}_{i+1}h-1,i+1)y\right\|^{2}\right]\mathcal{F}(\widetilde{m}_{i+1}h^{2}-1)\right]\right]$$

$$\leq q_{1}q_{2}\mathbb{E}\left[\left\|\Phi_{P'}(\widetilde{m}_{i+1}h-1,i+1)y\right\|^{2}\right]$$

$$= q_{1}q_{2}\mathbb{E}\left[\mathbb{E}\left[\left\|\Phi_{P'}(\widetilde{m}_{i+1}h-1,i+1)y\right\|^{2}\right]\mathcal{F}((i+1)h-1)\right]\right]$$

$$\leq q_{1}q_{2}^{2}\mathbb{E}\left[\left\|y\right\|^{2}\right], \ 0 \leq i < k-3h.$$
(D.34)

Secondly, it follows from (D.33) that

$$\mathbb{E}\left[\left\|\Phi_{P'}(k, i+1)y\right\|^{2}\right]$$
  
=  $\mathbb{E}\left[\mathbb{E}\left[\left\|\Phi_{P'}(k, i+1)y\right\|^{2}\right]\mathcal{F}((i+1)h-1)\right]\right]$   
 $\leq q_{2}\mathbb{E}\left[\left\|y\right\|^{2}\right], \ k-h \leq i < k.$  (D.35)

From (D.33) and (D.35), it is known that

$$\mathbb{E}\left[ \|\Phi_{P'}(k,i+1)y\|^{2} \right] = \mathbb{E}\left[ \mathbb{E}\left[ \|\Phi_{P'}(k,i+1)y\|^{2} \right] \mathcal{F}((i+1)h-1) \right] \\ = \mathbb{E}\left[ \mathbb{E}\left[ \|\Phi_{P'}(k,k-h+1)\Phi_{P'}(k-h,i+1)y\|^{2} \right] \mathcal{F}((i+1)h-1) \right] \\ \leq q_{2}\mathbb{E}\left[ \mathbb{E}\left[ \|\Phi_{P'}(k-h,i+1)y\|^{2} \right] \mathcal{F}((i+1)h-1) \right] \\ \leq q_{2}^{2}\mathbb{E}\left[ \|y\|^{2} \right], \ k-2h \leq i < k-h.$$
(D.36)

Finally, it follows from (D.33) and (D.36) that

$$\mathbb{E}\left[\|\Phi_{P'}(k, i+1)y\|^{2}\right] = \mathbb{E}\left[\mathbb{E}\left[\|\Phi_{P'}(k, i+1)y\|^{2}\right|\mathcal{F}((i+1)h-1)\right]\right] \\ = \mathbb{E}\left[\mathbb{E}\left[\|\Phi_{P'}(k, k-h+1)\Phi_{P'}(k-h, i+1)y\|^{2}\right|\mathcal{F}((i+1)h-1)\right]\right] \\ \leq q_{2}\mathbb{E}\left[\mathbb{E}\left[\|\Phi_{P'}(k-h, i+1)y\|^{2}\right|\mathcal{F}((i+1)h-1)\right]\right] \\ \leq q_{2}^{3}\mathbb{E}\left[\|y\|^{2}\right], \ k-3h \leq i < k-2h.$$
(D.37)

Combining (D.35)-(D.37), we get

$$\mathbb{E}\left[\|\Phi_{P'}(k, i+1)y\|^2\right] \le \max\left\{q_2, q_2^2, q_2^3\right\} \mathbb{E}\left[\|y\|^2\right], \ 0 < k-i \le 3h.$$
(D.38)

Denote  $d_3 = \max\{q_1q_2^2, q_2, q_2^2, q_2^3\}$ . By (D.34) and (D.38), we have

$$\sup_{k\geq 0} \mathbb{E}\left[\|\Phi_{P'}(k,i+1)y\|^2\right] \leq d_3 \mathbb{E}\left[\|y\|^2\right], \ \forall \ i\geq 0.$$

**Lemma D.9.** For the algorithm (3.6), suppose that Assumptions 4.1, 4.2, Conditions 4.1 and 4.2 hold, there exists an integer h > 0, a constant  $\rho_0 > 0$ , a strictly positive self-adjoint operator

 $\mathscr{H} \in \mathscr{L}(\mathscr{X}^N)$  and a nonnegative real sequence  $\{c(k), k \geq 0\}$ , respectively, satisfying the following conditions:

(i) for any given  $L_2$ -bounded adaptive sequence  $\{x(k), \mathcal{F}(kh-1), k \geq 0\}$  with values in the Hilbert space  $\mathscr{X}^N$ ,

$$\lim_{k \to \infty} \sum_{i=m}^{k} \left( \mathbb{E}\left[ \left\| \left( \prod_{j=i+1}^{k} (I_{\mathscr{X}^{N}} - c(j)\mathscr{H}) \right) \mu(i) \right\|^{2} \right] \right)^{\frac{1}{2}} = 0, \ \forall \ m \in \mathbb{N}, \quad \sum_{k=0}^{\infty} c(k) = \infty,$$

where  $\mu(i) := c(i)\mathscr{H}x(i) - \sum_{s=ih}^{(i+1)h-1} (a(s)\mathbb{E}\left[\mathcal{H}^*(s)\mathcal{H}(s)x(i) \mid \mathcal{F}(ih-1)\right] + b(s)\left(\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}\right)x(i)\right);$ (*ii*)  $\sup_{k \ge 0} \left( \mathbb{E} \left[ \|\mathcal{H}^*(k)\mathcal{H}(k)\|^{2^{\max\{h,2\}}} \middle| \mathcal{F}(k-1) \right] \right)^{\frac{1}{2^{\max\{h,2\}}}} \le \rho_0 \ a.s.$ 

Then the sequence of operator-valued random elements

$$\left\{I_{\mathscr{X}^N} - \sum_{i=kh}^{(k+1)h-1} (a(i)\mathcal{H}^*(i)\mathcal{H}(i) + b(i)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}), k \ge 0\right\}$$

is  $L_2^2$ -stable w.r.t. { $\mathcal{F}((k+1)h-1), k \ge 0$ }.

*Proof.* Let  $\{x(k), \mathcal{F}(kh-1), k \geq 0\}$  be a  $L_2$ -bounded adaptive sequence with values in the Hilbert space  $\mathscr{X}^N$ . For any given integer  $m \geq 0$ , let

$$= \left(I_{\mathscr{X}^N} - \sum_{s=kh}^{(k+1)h-1} (a(s)\mathcal{H}^*(s)\mathcal{H}(s) + b(s)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})\right) w(k), \ k \ge m,$$
(D.39)

where  $w(m) = x(m), w(i) = 0, i = 0, \dots, m-1$ . It follows from Proposition A.5.(a)-(c) that  $\{w(k), k \ge 0\}$  is a random sequence with values in the Hilbert space  $(\mathscr{X}^N, \tau_N(\mathscr{X}^N))$ . On one hand, from the definition of w(k), it follows that

$$= \left(\prod_{i=m}^{k} \left( I_{\mathscr{X}^{N}} - \sum_{s=ih}^{(i+1)h-1} (a(s)\mathcal{H}^{*}(s)\mathcal{H}(s) + b(s)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}) \right) \right) x(m), \ k \ge m. (D.40)$$

Noting that  $x(m) \in L^0(\Omega, \mathcal{F}(mh-1); \mathscr{X}^N)$ , it follows from Lemma D.8 that there exists a constant  $d_3 > 0$ , such that

$$\sup_{k \ge 0} \mathbb{E}\left[ \|w(k+1)\|^2 \right] \le d_3 \mathbb{E}\left[ \|x(m)\|^2 \right] \le d_3 \sup_{k \ge 0} \mathbb{E}\left[ \|x(k)\|^2 \right] < \infty.$$
(D.41)

Moreover, (D.39) can be rewritten as

$$w(i+1) = (I_{\mathscr{X}^N} - c(i)\mathscr{H})w(i)$$

$$+\left(c(i)\mathscr{H}-\sum_{s=ih}^{(i+1)h-1}(a(s)\mathcal{H}^*(s)\mathcal{H}(s)+b(s)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}})\right)w(i),\ i\geq m,$$

which gives

$$= \left(\prod_{i=m}^{k} (I_{\mathscr{X}^{N}} - c(i)\mathscr{H})\right) x(m) + \sum_{i=m}^{k} \left(\prod_{j=i+1}^{k} (I_{\mathscr{X}^{N}} - c(j)\mathscr{H})\right) \times \left(c(i)\mathscr{H} - \sum_{s=ih}^{(i+1)h-1} (a(s)\mathscr{H}^{*}(s)\mathscr{H}(s) + b(s)\mathscr{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})\right) w(i), \ k \ge m.$$
(D.42)

From (D.41) and the condition (ii), it is known that

$$\sup_{\substack{s\geq 0\\i\geq 0}} \mathbb{E}\left[\left\|\mathcal{H}^*(s)\mathcal{H}(s)w(i)\right\|\right] \le \sup_{s\geq 0} \mathbb{E}\left[\left\|\mathcal{H}^*(s)\mathcal{H}(s)\right\|^2\right] + \sup_{i\geq 0} \mathbb{E}\left[\left\|w(i)\right\|^2\right] < \infty,$$

which together Proposition 2.4 leads to  $\mathcal{H}^*(s)\mathcal{H}(s)w(i) \in L^1(\Omega, \mathcal{F}(s); \mathscr{X}^N)$ ,  $s \ge ih$ . Thus, by Lemma A.9, we know that there exists a unique conditional expectation  $\mathbb{E}[\mathcal{H}^*(s)\mathcal{H}(s)w(i)|\mathcal{F}(ih - 1)]$  of  $\mathcal{H}^*(s)\mathcal{H}(s)w(i)$  w.r.t.  $\mathcal{F}(ih - 1)$ . Then we have

$$\left(c(i)\mathscr{H} - \sum_{s=ih}^{(i+1)h-1} (a(s)\mathcal{H}^*(s)\mathcal{H}(s) + b(s)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})\right)w(i) = w_1(i) + w_2(i), \quad (D.43)$$

where

$$\begin{cases} w_{1}(i) = c(i)\mathscr{H}w(i) - \sum_{s=ih}^{(i+1)h-1} (a(s)\mathbb{E}[\mathcal{H}^{*}(s)\mathcal{H}(s)w(i)|\mathcal{F}(ih-1)] \\ + b(s)(\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}})w(i)), \\ w_{2}(i) = \sum_{s=ih}^{(i+1)h-1} a(s)(\mathbb{E}[\mathcal{H}^{*}(s)\mathcal{H}(s)w(i)|\mathcal{F}(ih-1)] - \mathcal{H}^{*}(s)\mathcal{H}(s)w(i)). \end{cases}$$
(D.44)

By (D.40) and (D.42)-(D.43), we get

$$\left(\prod_{i=m}^{k} \left( I_{\mathscr{X}^{N}} - \sum_{s=ih}^{(i+1)h-1} (a(s)\mathcal{H}^{*}(s)\mathcal{H}(s) + b(s)\mathcal{L}_{\mathcal{G}} \otimes I_{\mathscr{X}}) \right) \right) x(m)$$
  
= 
$$\left(\prod_{i=m}^{k} (I_{\mathscr{X}^{N}} - c(i)\mathscr{H}) \right) x(m) + \sum_{i=m}^{k} \left(\prod_{j=i+1}^{k} (I_{\mathscr{X}^{N}} - c(j)\mathscr{H}) \right) (w_{1}(i) + w_{2}(i)),$$

which together with Cauchy inequality leads to

$$\mathbb{E}\left[\left\|\left(\prod_{i=m}^{k}\left(I_{\mathscr{X}^{N}}-\sum_{s=ih}^{(i+1)h-1}(a(s)\mathcal{H}^{*}(s)\mathcal{H}(s)+b(s)\mathcal{L}_{\mathcal{G}}\otimes I_{\mathscr{X}})\right)\right)x(m)\right\|^{2}\right]$$

$$\leq 2\mathbb{E}\left[\left\|\left(\prod_{i=m}^{k}(I_{\mathscr{X}^{N}}-c(i)\mathscr{H})\right)x(m)\right\|^{2}\right] + 2\mathbb{E}\left[\left\|\sum_{i=m}^{k}\left(\prod_{j=i+1}^{k}(I_{\mathscr{X}^{N}}-c(j)\mathscr{H})\right)(w_{1}(i)+w_{2}(i))\right\|^{2}\right], \ k \geq m.$$
(D.45)

By Lemma D.3, it is known that there exist constants M, d > 0, such that

$$\left\| \left( \prod_{i=m}^k (I_{\mathscr{X}^N} - c(i)\mathscr{H}) \right) x(m) \right\|^2 \le M^{2d} \|x(m)\|^2 \text{ a.s.}$$

Noting that  $\sup_{k\geq 0} \mathbb{E}[||x(k)||^2] < \infty$ , it follows from Lebesgue dominated convergence theorem and Lemma D.3 that

$$\lim_{k \to \infty} \mathbb{E}\left[ \left\| \left( \prod_{i=m}^{k} (I_{\mathscr{X}^{N}} - c(i)\mathscr{H}) \right) x(m) \right\|^{2} \right] = 0, \ \forall \ m \ge 0.$$
 (D.46)

By using Cauchy inequality again, we obtain

$$\mathbb{E}\left[\left\|\sum_{i=m}^{k}\left(\prod_{j=i+1}^{k}(I_{\mathscr{X}^{N}}-c(j)\mathscr{H})\right)\left(w_{1}(i)+w_{2}(i)\right)\right\|^{2}\right]$$

$$\leq 2\mathbb{E}\left[\left\|\sum_{i=m}^{k}\left(\prod_{j=i+1}^{k}(I_{\mathscr{X}^{N}}-c(j)\mathscr{H})\right)w_{1}(i)\right\|^{2}\right]$$

$$+2\mathbb{E}\left[\left\|\sum_{i=m}^{k}\left(\prod_{j=i+1}^{k}(I_{\mathscr{X}^{N}}-c(j)\mathscr{H})\right)w_{2}(i)\right\|^{2}\right].$$
(D.47)

We now consider the right-hand side of (D.47) term by term. Firstly, by Minkowski inequality, we get

$$\mathbb{E}\left[\left\|\sum_{i=m}^{k}\left(\prod_{j=i+1}^{k}(I_{\mathscr{X}^{N}}-c(j)\mathscr{H})\right)w_{1}(i)\right\|^{2}\right]$$

$$\leq \left(\sum_{i=m}^{k}\left(\mathbb{E}\left[\left\|\left(\prod_{j=i+1}^{k}(I_{\mathscr{X}^{N}}-c(j)\mathscr{H})\right)w_{1}(i)\right\|^{2}\right]\right)^{\frac{1}{2}}\right)^{2}.$$
(D.48)

It follows from (D.39) and (D.41) that  $\{w(k), \mathcal{F}(kh-1), k \ge 0\}$  is an adaptive sequence with  $\sup_{k\ge 0} \mathbb{E}[\|w(k)\|^2] < \infty$ , which together with (D.44), the condition (i) and (D.48) gives

$$\lim_{k \to \infty} \mathbb{E}\left[ \left\| \sum_{i=m}^{k} \left( \prod_{j=i+1}^{k} (I_{\mathscr{X}^{N}} - c(j)\mathscr{H}) \right) w_{1}(i) \right\|^{2} \right] = 0.$$
 (D.49)

Secondly, noting that  $w_2(i) \in \mathcal{F}((i+1)h-1)$ ,  $w(i) \in \mathcal{F}(ih-1)$ , by Condition 4.1, it follows from the condition (ii) and (D.41) that

$$\begin{split} & \mathbb{E} \left[ \|w_{2}(i)\|^{2} \right] \\ &= \mathbb{E} \left[ \left\| \sum_{s=ih}^{(i+1)h-1} a(s)(\mathbb{E}[\mathcal{H}^{*}(s)\mathcal{H}(s)w(i)|\mathcal{F}(ih-1)] - \mathcal{H}^{*}(s)\mathcal{H}(s)w(i)) \right\|^{2} \right] \\ &\leq ha^{2}(i)\mathbb{E} \left[ \sum_{s=ih}^{(i+1)h-1} \|\mathbb{E}[\mathcal{H}^{*}(s)\mathcal{H}(s)w(i)|\mathcal{F}(ih-1)] - \mathcal{H}^{*}(s)\mathcal{H}(s)w(i)\|^{2} \right] \\ &\leq 2ha^{2}(i)\mathbb{E} \left[ \sum_{s=ih}^{(i+1)h-1} \left( \mathbb{E} \left[ \|\mathcal{H}^{*}(s)\mathcal{H}(s)w(i)\|^{2} |\mathcal{F}(ih-1) \right] \\ &+ \|\mathcal{H}^{*}(s)\mathcal{H}(s)w(i)\|^{2} \right) \right] \\ &\leq 2ha^{2}(i)\mathbb{E} \left[ \sum_{s=ih}^{(i+1)h-1} \left( \rho_{0}^{2} \|w(i)\|^{2} + \|\mathcal{H}^{*}(s)\mathcal{H}(s)w(i)\|^{2} \right) \right] \\ &\leq 2ha^{2}(i)\mathbb{E} \left[ \|w(i)\|^{2} \sum_{s=ih}^{(i+1)h-1} \left( \rho_{0}^{2} + \|\mathcal{H}^{*}(s)\mathcal{H}(s)\|^{2} \right) \right] \\ &= 2ha^{2}(i)\mathbb{E} \left[ \mathbb{E} \left[ \|w(i)\|^{2} \sum_{s=ih}^{(i+1)h-1} \left( \rho_{0}^{2} + \|\mathcal{H}^{*}(s)\mathcal{H}(s)\|^{2} \right) \right| \mathcal{F}(ih-1) \right] \\ &= 2ha^{2}(i)\mathbb{E} \left[ \mathbb{E} \left[ \|w(i)\|^{2} \right], \tag{D.50} \end{split}$$

which together with Condition 4.2 gives

$$\sup_{i\geq 0} \mathbb{E}\left[\|w_2(i)\|^2\right] < \infty. \tag{D.51}$$

Thus, it follows from Lemma A.9 that  $\mathbb{E}[w_2(i)|\mathcal{F}(ih-1)]$  exists and

$$\mathbb{E}[w_{2}(i)|\mathcal{F}(ih-1)] = \mathbb{E}\left[\sum_{s=ih}^{(i+1)h-1} a(s)(\mathbb{E}[\mathcal{H}^{*}(s)\mathcal{H}(s)w(i)|\mathcal{F}(ih-1)] - \mathcal{H}^{*}(s)\mathcal{H}(s)w(i))\right|\mathcal{F}(ih-1)\right] \\
= \sum_{s=ih}^{(i+1)h-1} a(s)\mathbb{E}[\mathbb{E}[\mathcal{H}^{*}(s)\mathcal{H}(s)w(i)|\mathcal{F}(ih-1)] - \mathcal{H}^{*}(s)\mathcal{H}(s)w(i)|\mathcal{F}(ih-1)] \\
= 0.$$
(D.52)

Meanwhile, from Lemma D.3, it is known that there exist constants M, d > 0 such that

$$\sup_{\substack{\|x\|=1\\x\in\mathscr{X}^N}} \inf\left\{r \ge 0: \mathbb{P}\left(\left\|\left(\prod_{j=t+1}^k (I_{\mathscr{X}^N} - c(j)\mathscr{H})\right) x\right\| < r\right) = 1\right\}$$
  
$$\le \sup_{\substack{\|x\|=1\\x\in\mathscr{X}^N}} M^d \|x\| < \infty.$$
(D.53)

For  $m \le s < t \le k$ , it follows from Proposition 2.4, Lemma A.10, Lemma 3.5.2 in [67], Proposition 2.10 and (D.51)-(D.53) that

$$\mathbb{E}\left[\left\langle \left(\prod_{j=s+1}^{k} (I_{\mathscr{X}^{N}} - c(j)\mathscr{H})\right) w_{2}(s), \left(\prod_{j=t+1}^{k} (I_{\mathscr{X}^{N}} - c(j)\mathscr{H})\right) w_{2}(t)\right\rangle \right] = 0. \quad (D.54)$$

On one hand, by Lemma D.3, it is known that there exist positive constants M and d, such that

$$\left\| \left( \prod_{j=i+1}^{k} (I_{\mathscr{X}^N} - c(j)\mathscr{H}) \right) \frac{1}{a(i)} w_2(i) \right\| \le M^d \left\| \frac{1}{a(i)} w_2(i) \right\| \text{ a.s.}$$
(D.55)

Thus, by Lemma D.3 and (D.54)-(D.55), we get

$$\mathbb{E}\left[\left\|\sum_{i=m}^{k}\left(\prod_{j=i+1}^{k}\left(I_{\mathscr{X}^{N}}-c(j)\mathscr{H}\right)\right)w_{2}(i)\right\|^{2}\right] \\
=\sum_{i=m}^{k}\mathbb{E}\left[\left\|\left(\prod_{j=i+1}^{k}\left(I_{\mathscr{X}^{N}}-c(j)\mathscr{H}\right)\right)w_{2}(i)\right\|^{2}\right] \\
+2\sum_{\substack{m\leq s< t\leq k}}\mathbb{E}\left[\left\langle\left(\prod_{j=s+1}^{k}\left(I_{\mathscr{X}^{N}}-c(j)\mathscr{H}\right)\right)w_{2}(s),\left(\prod_{j=t+1}^{k}\left(I_{\mathscr{X}^{N}}-c(j)\mathscr{H}\right)\right)w_{2}(i)\right)\right] \\
=\sum_{i=m}^{k}\mathbb{E}\left[\left\|\left(\prod_{j=i+1}^{k}\left(I_{\mathscr{X}^{N}}-c(j)\mathscr{H}\right)\right)w_{2}(i)\right\|^{2}\right] \\
=\sum_{i=m}^{k}a^{2}(i)\mathbb{E}\left[\left\|\left(\prod_{j=i+1}^{k}\left(I_{\mathscr{X}^{N}}-c(j)\mathscr{H}\right)\right)\frac{1}{a(i)}w_{2}(i)\right\|^{2}\right] \\
\leq M^{d}\sum_{i=m}^{k}a^{2}(i)\left[\mathbb{E}\left[\left\|\left(\prod_{j=i+1}^{k}\left(I_{\mathscr{X}^{N}}-c(j)\mathscr{H}\right)\right)\frac{1}{a(i)}w_{2}(i)\right\|^{2}\right]\right]^{\frac{1}{2}} \\
\times\left[\mathbb{E}\left[\left\|\frac{1}{a(i)}w_{2}(i)\right\|^{2}\right]\right]^{\frac{1}{2}}.$$
(D.56)

On the other hand, by (D.41) and (D.50), we have

$$\sup_{i \ge 0} \mathbb{E}\left[ \left\| \frac{1}{a(i)} w_2(i) \right\|^2 \right] \le 4h^2 \rho_0^2 \sup_{i \ge 0} \mathbb{E}\left[ \|w(i)\|^2 \right] < \infty.$$
(D.57)

Substituting (D.57) into (D.56) gives

$$\mathbb{E}\left[\left\|\sum_{i=m}^{k}\left(\prod_{j=i+1}^{k}\left(I_{\mathscr{X}^{N}}-c(j)\mathscr{H}\right)\right)w_{2}(i)\right\|^{2}\right]\right]^{2}$$

$$\leq M^{d}\sup_{i\geq0}\left[\mathbb{E}\left[\left\|\frac{1}{a(i)}w_{2}(i)\right\|^{2}\right]\right]^{\frac{1}{2}}$$

$$\times\sum_{i=m}^{k}a^{2}(i)\left(\mathbb{E}\left[\left\|\left(\prod_{j=i+1}^{k}\left(I_{\mathscr{X}^{N}}-c(j)\mathscr{H}\right)\right)\frac{1}{a(i)}w_{2}(i)\right\|^{2}\right]\right)^{\frac{1}{2}},$$

which together with Condition 4.2, (D.57) and Lemma D.4 leads to

$$\lim_{k \to \infty} \mathbb{E}\left[ \left\| \sum_{i=m}^{k} \left( \prod_{j=i+1}^{k} (I_{\mathscr{X}^{N}} - c(j)\mathscr{H}) \right) w_{2}(i) \right\|^{2} \right] = 0, \ \forall \ m \ge 0.$$
 (D.58)

Hence, substituting (D.46)-(D.47), (D.49) and (D.58) into (D.45) completes the proof of Lemma D.9.  $\Box$ 

## REFERENCES

- [1] BERTERO, M. and BOCCACCI, P. (1998). Introduction to inverse problems in imaging. CRC press.
- [2] CAKONI, G. and COLTON, D. (2005). Open problems in the qualitative approach to inverse electromagnetic scattering theory. *European Journal of Applied Mathematics*. **16** 411–425.
- [3] COLTON, D. and KRESS, R. (2012). *Inverse acoustic and electromagnetic scattering theory* 93. Springer Science and Business Media.
- [4] ISAKOV, V. (2008). On inverse problems in secondary oil recovery. *European Journal of Applied Mathematics*. **19** 459–478.
- [5] ENGL, H. W., LOUIS, A. K. and Rundell, W. (1996). Inverse problems in medical imaging and nondestructive testing. *Proceeding of the Conference in Oberwolfach*. Federal Republic of Germany, Feb. 4–10.
- [6] KIRSCH, A. (1996). An Introduction to the mathematical theory of inverse problems. New York: Springer-Verlag.
- [7] ENGL, H. W., HANKE, M. and NEUBAUER, A. (1996). *Regularization of inverse problems* 375. Springer Science and Business Media.
- [8] MOROZOV, V. A. (1984). Regular methods for solving linear and nonlinear ill-posed problems," in *Methods for Solving Incorrectly Posed Problems* 65–122, Springer, New York.

- [9] WERSCHULZ, A. G. (1991). *The computational complexity of differential and integral equations: An information-based approach*. Oxford University Press, Inc.
- [10] TIKHONOV, A. N. (1963). Regularization of incorrectly posed problems. Soviet Mathematics Doklady. 4 1624–1627.
- [11] BISSANTZ, N., HOHAGE, T. and MUNK, A. (2004). Consistency and rates of convergence of nonlinear Tikhonov regularization with random noise. *Inverse Problems*, **20** 1773.
- [12] CAVALIER, L. (2008). Nonparametric statistical inverse problems. *Inverse Problems*. 24 034004.
- [13] KEKKONEN, H., LASSAS, M. and SILTANEN, S. (2014). Analysis of regularized inversion of data corrupted by white Gaussian noise. *Inverse Problems*. **30** 045009.
- [14] GINE, E. and NICKL, R. (2015). Mathematical foundations of infinite-dimensional statistical models 40. Cambridge University Press.
- [15] HOHAGE, T. and Werner, F. (2016). Inverse problems with poisson data: statistical regularization theory, applications and algorithms. *Inverse Problems*. **32** 093001.
- [16] JANZ, D., BURT, D. and GONZALEZ, J. (2020). Bandit optimisation of functions in the Matern kernel RKHS. *Proceeding of International Conference on Artificial Intelligence and Statistics.* Jun. 3. 2486–2495.
- [17] TAKEMORI, S. and SATO, M. (2021). Approximation theory based methods for RKHS bandits. *Proceeding of International Conference on Machine Learning*. Jul. 1. 10076-10085.
- [18] SMALE, S. and CUCKER, F. (2002). On the mathematical foundations of learning. American Mathematical Society. 39 1–49.
- [19] YING, Y., and PONTIL, M. (2008). Online gradient descent learning algorithms. Foundations of Computational Mathematics. 8 561–596.
- [20] TARRES, P. and YAO, Y. (2014). Online learning as stochastic approximation of regularization paths: Optimality and almost-sure convergence. *IEEE Transactions on Information Theory.* **60** 5716–5735.
- [21] DIEULEVEUT, A. and BACH, F. (2016). Nonparametric stochastic approximation with large step-sizes. *The Annals of Statistics*. 44 1363–1399.
- [22] LEI, Y., SHI, L. and GUO, Z. C. (2018). Convergence of unregularized online learning algorithms. *Journal of Machine Learning Research*. 18 6269–6301.
- [23] MATHE, P. and PEREVERZEV, S. (2006). Regularization of some linear ill-posed problems with discretized random noisy data. *Mathematics of Computation*. **75** 1913–1929.

- [24] LU, S. and MATHE, P. (2014). Discrepancy based model selection in statistical inverse problems. *Journal of Complexity.* **30** 290–308.
- [25] IGLESIAS, M. A., LIN, K., LU, S. and STUART, A. M. (2017). Filter based methods for statistical linear inverse problems. *Communications in Mathematical Sciences*. 15 1867– 1896.
- [26] LU, S., NIU, P. and WERNER, F. (2021). On the asymptotical regularization for linear inverse problems in presence of white noise. SIAM/ASA Journal on Uncertainty Quantification. 9 1–28.
- [27] LU, S., MATHE, P. and PEREVERZEV, S. V. (2020). Randomized matrix approximation to enhance regularized projection schemes in inverse problems. *Inverse Problems*. **36** 085013.
- [28] LU. S. and MATHE, P. (2017). Stochastic gradient descent for linear inverse problems in Hilbert spaces. *Mathematics of Computation*. **91** 1763–1788.
- [29] JAHN, T. and JIN, B. (2020). On the discrepancy principle for stochastic gradient descent. *Inverse Problems.* 36 095009.
- [30] JIN, B. and LU, X. (2018). On the regularizing property of stochastic gradient descent. *Inverse Problems.* 35 015004.
- [31] ROSENBLATT, J. D. and NADLER, B. (2016). On the optimality of averaging in distributed statistical learning. *Information and Inference: A Journal of the IMA*. **5** 379–404.
- [32] WANG, J., LI, T. and ZHANG, X. (2021). Decentralized Cooperative Online Estimation With Random Observation Matrices, Communication Graphs and Time Delays. *IEEE Transactions on Information Theory*. 67 4035–4059.
- [33] IPPEL, L., KAPTEIN, M. and VERMUNT, J. (2016). Dealing with data streams: An online, row-by-row, estimation tutorial. *Methodology: European Journal of Research Methods for the Behavioral and Social Sciences.* **12** 124.
- [34] LOPES, C. G. and SAYED, A. H. (2008). Diffusion least-mean squares over adaptive networks: Formulation and performance analysis. *IEEE Transactions on Signal Processing*. 56 3122–3136.
- [35] KAR, S. and MOURA, J. M. F. (2011). Convergence rate analysis of distributed gossip (linear parameter) estimation: fundamental limits and tradeoff. *IEEE J. Sel. Topics Signal Processing.* 5 674–690.
- [36] CATTIVELLI, F. S. and SAYED, A. H. (2010). Diffusion strategies for distributed Kalman filtering and smoothing. *IEEE Transactions on Automatic Control.* **55** 2069–2084.

- [37] AL-SAYED, S., ZOUBIR, A. M. and SAYED, A. H. (2017). Robust distributed estimation by networked agents. *IEEE Transactions on Signal Processing*. **65** 3909–3921.
- [38] GHOLAMI, M. R., JANSSON, M., STRÖM, E. G. and SAYED, A. H. (2016). Diffusion estimation over cooperative multi-agent networks with missing data. *IEEE Transactions on Signal and Information Processing over Networks*. 2 276–289.
- [39] ABDOLEE, R., CHAMPAGNE, B. and SAYED, A. H. (2016). Diffusion adaptation over multi-agent networks with wireless link impairments. *IEEE Transactions on Mobile Computing.* 15 1362–1376.
- [40] KAR, S. and MOURA, J. M. F. (2011). Gossip and distributed Kalman filtering: Weak consensus under weak detectability. *IEEE Trans. Signal Processing.* 59 1766–1786.
- [41] KAR, S., MOURA, J. M. F. and RAMANAN, K. (2012). Distributed parameter estimation in sensor networks: Nonlinear observation models and imperfect communication. *IEEE Trans. Information Theory.* 58 3575–3605.
- [42] KAR, S. and MOURA, J. M. F. (2013). Consensus+innovations distributed inference over networks: Cooperation and sensing in networked systems. *IEEE Signal Processing Magazine*. **30** 99–109.
- [43] PIGGOTT, M. J. and SOLO, V. (2016). Diffusion LMS with correlated regressors I: Realization-wise stability. *IEEE Transactions on Signal Processing*. 64 5473–5484.
- [44] PIGGOTT, M. J. and SOLO, V. (2017). Diffusion LMS with correlated regressors II: Performance. *IEEE Transactions on Signal Processing*. 65 3934–3947.
- [45] ISHIHARA, J. Y. and ALGHUNAIM, S. C. (2017). Diffusion LMS filter for distributed estimation of systems with stochastic state transition and observation matrices. *Proceeding* of 2017 American Control Conference (ACC). May. 24, 5199–5204.
- [46] LYAQINI, S., QUAFAFOU, M., NACHAOUI, M. and CHAKIB, A. (2020). Supervised learning as an inverse problem based on non-smooth loss function. *Knowledge and Information Systems.* 62 3039–3058.
- [47] SHIN, B., YUKAWA, M., CAVALCANTE, R. L. G. and DEKORSY, A. (2018). Distributed Adaptive Learning With Multiple Kernels in Diffusion Networks. *IEEE Transactions on Signal Processing.* 66 5505–5519.
- [48] DENG, Z., GREGORY, J. and KURDILA, A. (2012). Learning theory with consensus in reproducing kernel Hilbert spaces. *Proceeding of 2012 American Control Conference* (ACC). Jun. 27, 1400–1405.

- [49] MITRA, R. and BHATIA, V. (2014). The diffusion-KLMS algorithm. Proceeding of 2014 International Conference on Information Technology. Dec. 22, 256–259.
- [50] CHOUVARDAS, R. and DRAIEF, M. (2016). A diffusion kernel LMS algorithm for nonlinear adaptive networks. *Proceeding of 2016 IEEE International Conference on Acoustics, Speech* and Signal Processing (ICASSP). Mar. 20, 4164–4168.
- [51] BOUBOULIS, P., CHOUVARDAS, S. and THEODORIDIS, S. (2018). Online distributed learning over networks in RKH spaces using random Fourier features. *IEEE Transactions* on Signal Processing. 66 1920–1932.
- [52] JONES, F. G. and SIMPSON, G. (2022). Iterate averaging, the Kalman filter, and 3DVAR for linear inverse problems. *Numerical Algorithms*. 1–21.
- [53] SMALE, S. and Yao, Y. (2006). Online learning algorithms. *Foundations of Computational Mathematics*. 6 145–170.
- [54] MATHE, P. and PEREVERZEV, S. V. (2001). Optimal discretization of inverse problems in Hilbert scales. Regularization and self-regularization of projection methods. *SIAM Journal on Numerical Analysis.* **38** 1999–2021.
- [55] ZHANG, X., LI, T. and GU, Y. (2020). Consensus+ innovations distributed estimation with random network graphs, observation matrices and noises. *Proceeding of 2020 59th IEEE Conference on Decision and Control (CDC)*. Dec. 14. 4318–4323.
- [56] UNGUREANU, V. M. and CHENG, S. S. (2008). Mean stability of a stochastic difference equation. *Proceeding of Annales Polonici Mathematici*. 1 33–52. Institute of Mathematics Polish Academy of Sciences.
- [57] KUBRUSLY, S. C. (1978). Applied stochastic approximation algorithms in Hilbert space. International Journal of Control. 28 23–31.
- [58] VAJJHA, K., TRAGER, B., SHINNAR, A. and PESTUN, V. (2022). Formalization of a Stochastic Approximation Theorem. arXiv:2202.05959.
- [59] UNGUREANU, V. M. (2014). Stability, stabilizability and detectability for Markov jump discrete-time linear systems with multiplicative noise in Hilbert spaces. *Optimization*. 63 1689–1712.
- [60] UNGUREANU, V. M. (2009). Optimal control for linear discrete-time systems with Markov perturbations in Hilbert spaces. *IMA Journal of Mathematical Control and Information*. 26 105–127.
- [61] ZHANG, W., ZHENG, W. X. and CHEN, B. S. (2017). Detectability, observability and

Lyapunov-type theorems of linear discrete time-varying stochastic systems with multiplicative noise. *International Journal of Control.* **90** 2490–2507.

- [62] GUO, Z. C., LIN, S. B. and SHI, L. (2019). Distributed learning with multi-penalty regularization. *Applied and Computational Harmonic Analysis*. **46** 478–499.
- [63] GUO, Z. C., LIN, S. B. and ZHOU, D. X. (2017). Learning theory of distributed spectral algorithms. *Inverse Problems*. **33** 074009.
- [64] LIN, J. and CEVHER, V. (2020). Optimal convergence for distributed learning with stochastic gradient methods and spectral algorithms. *Journal of Machine Learning Research*. 21 1–63.
- [65] SCHMITT, U. and LOUIS, A. K. (2002). Efficient algorithms for the regularization of dynamic inverse problems: I. Theory. *Inverse Problems*. 18 645.
- [66] HYTÖNEN. T., VAN NEERVEN, J., VERAAR, M. and WEIS, L. (2018). Analysis in Banach Spaces: Volume II: Probabilistic Methods and Operator Theory. 67 Springer.
- [67] HYTÖNEN. T., VAN NEERVEN, J., VERAAR, M. and WEIS, L. (2016). Analysis in Banach spaces. 12 Berlin: Springer.
- [68] BENNING, M. and BURGER, M. (2018). Modern regularization methods for inverse problems. Acta Numerica. 27 1–111.
- [69] REICH, S. and ZASLAVSKI, A. J. (1999). Convergence of generic infinite products of affine operators. *Abstract and Applied Analysis*. **4** 1–19.
- [70] PUSTYLNIK, E. and REICH, S. (2014). Infinite products of arbitrary operators and intersections of subspaces in Hilbert space. *Journal of Approximation Theory*. 91–102.
- [71] PUSTYLNIK, E., REICH, S. and ZASLAVSKI, A. J. (2011). Convergence of non-cyclic infinite products of operators. *Journal of Mathematical Analysis and Applications*. 380 759–767.
- [72] PUSTYLNIK, E., REICH, S. and ZASLAVSKI, A. J. (2012). Convergence of non-periodic infinite products of orthogonal projections and nonexpansive operators in Hilbert space. *Journal of Approximation Theory.* 164 611–624.
- [73] GUO, L. (1994). Stability of recursive stochastic tracking algorithms. SIAM Journal on Control and Optimization. 32 1195–1225.
- [74] GUO, L. (1990). Estimating time-varying parameters by the Kalman filter based algorithm: stability and convergence. *IEEE Transactions on Automatic Control.* **35** 141–147.
- [75] GUO, L. and LJUNG, L. (1995). Exponential stability of general tracking algorithms. IEEE

August 28, 2023

- [76] GUO, L. and LJUNG, L. (1995). Performance analysis of general tracking algorithms. *IEEE Transactions on Automatic Control.* 40 1388–1402.
- [77] GUO, L., LJUNG, L. and PRIOURET, P. (1993). Performance analysis of forgetting factor RLS algorithms. *International Journal of Adaptive Control and Signal Processing*. 7 525– 537.
- [78] GUO, L., LJUNG, L. and WANG, G. (1997). Necessary and sufficient conditions for stability of LMS. *IEEE Transactions on Automatic Control.* 42 761–770.
- [79] XIE, S. and GUO, L. (2018). Analysis of normalized least mean squares-based consensus adaptive filters under a general information condition. SIAM Journal on Control and Optimization. 56 3404–3431.
- [80] ZHANG, X., LI, T. and FU, X. (2022). Decentralized Online Regularized Learning Over Random Time-Varying Graphs. arXiv: 2206.03861.
- [81] CHEN, X., TANG, B., FAN, J. and GUO, X. (2022). Online gradient descent algorithms for functional data learning. *Journal of Complexity*. 70 101635.
- [82] THEODORIDIS, S. (2015). *Machine learning: a Bayesian and optimization perspective*. Academic press.
- [83] HILLE, E. and PHILLIPS, R. C. (1996). Functional analysis and semi-groups. American Mathematical Society. 31.
- [84] BLASCO, O. and GARCIA-BAYONA, I. (2016). Remarks on measurability of operatorvalued functions. *Mediterranean Journal of Mathematics*. **13** 5147–5162.
- [85] CHOW, Y. S. and TEICHER, H. (2012). *Probability theory: independence, interchangeability, martingales.* Springer Science and Business Media.
- [86] ROBBINS, H. and SIEGMUND, D. (1971). A convergence theorem for non-negative almost supermartingales and some applications. *Optimizing Methods in Statistics*. 233–257, Academic Press.