

REGULAR DESSINS WITH PRIMITIVE AUTOMORPHISM GROUPS

GARETH A. JONES AND MARTIN MAČAJ

ABSTRACT. We classify the dessins \mathcal{D} for which the automorphism group G acts primitively and faithfully on the points over one of the three critical values (without loss of generality the black vertices in the usual bipartite map representation). We show that they are all generalised Paley dessins, in which the black vertices are the elements of a finite field \mathbb{F}_q , and G is a subgroup of the affine group $\text{AGL}_1(q)$. Using earlier results obtained with Streit and Wolfart, we determine the orbits of the absolute Galois group on these dessins, we show that they are all defined over certain cyclotomic fields, and we obtain defining equations in some special cases. Relaxing the condition of a faithful action allows only cyclic regular coverings of these dessins.

1. INTRODUCTION

Every dessin is a quotient of a regular dessin (one with the greatest possible symmetry) by a subgroup of its automorphism group, so much of the study of dessins can be reduced to that of regular dessins and their automorphism groups. Unfortunately, the great proliferation of regular dessins (resulting from that of two-generator finite groups) makes it almost impossible to study them systematically. They all have the property that the automorphism group acts transitively on each of the three sets of critical points, the fibres over the critical values of the Belyĭ function. However, if we strengthen this transitivity property by requiring that the action on at least one of these sets should be primitive (preserving no nontrivial equivalence relations), then only a very specific and easily-understood family of dessins arises, namely the generalised Paley dessins (when the action is also faithful) and their cyclic regular covers (when it is not faithful).

We will say that a dessin \mathcal{D} is *primitive* or *faithful* if its automorphism group $G = \text{Aut } \mathcal{D}$ acts primitively or faithfully on the set of black vertices. (Throughout this paper we will represent dessins as bipartite graphs on surfaces, with vertices coloured black or white; by applying suitable dualities of dessins one can replace statements about black vertices with similar statements about white vertices or face centres.) Our main result is as follows:

Theorem 1.1. *A regular dessin is primitive and faithful if and only if it is a generalised Paley dessin.*

In a generalised Paley dessin \mathcal{D} , the black vertices are identified with the elements of a finite field \mathbb{F}_q of order $q = p^d$ for some prime p , and the automorphism group $G = \text{Aut } \mathcal{D}$ is a subgroup of the 1-dimensional affine group $\text{AGL}_1(q)$ acting primitively on the field. Equivalently, G is a

2010 *Mathematics Subject Classification.* Primary 20B25; secondary 05C10, 14H57, 20B15.

Key words and phrases. Regular dessin, automorphism group, primitive, generalised Paley dessin.

semidirect product $T \rtimes S$, where T is the translation subgroup (an elementary abelian group of order q) and S is a subgroup of the multiplicative group $\mathbb{F}_q^* = \mathbb{F}_q \setminus \{0\}$ of \mathbb{F}_q acting on the field as an irreducible subgroup of $\text{GL}_d(p)$, so that d is the multiplicative order of $p \bmod (n)$ where $n = |S|$. The standard generators of G are x, y and z , where x is a generator of S , y is an arbitrary element of $G \setminus S$, and $z = (xy)^{-1}$. Thus the black vertices have valency n , while the valencies of the white vertices and the faces, and hence the type and genus of the dessin, depend on which coset Tx^c of T contains y . In particular, if $q \equiv 1 \bmod (4)$ with $n = (q - 1)/2$ and $c = n/2$, then S is the group of quadratic residues in \mathbb{F}_q , and the white vertices have valency 2, so that if they are ignored the embedded graph is the Paley graph P_q , in which vertices are adjacent if and only if they differ by a quadratic residue.

For the origins of the Paley graphs, see [6]. Generalised Paley graphs were introduced by Lim and Praeger in [12], and generalised Paley maps were studied by Jones in [5]. These maps played an important role in the classification by Jones and Mačaj in [8] of regular maps with primitive automorphism groups. Generalised Paley dessins can be regarded as their natural generalisation to orientably regular hypermaps.

For a given n and prime p not dividing n , the number of non-isomorphic generalised Paley dessins \mathcal{D} is $n\phi(n)/d$ where ϕ is Euler's function, with $\phi(n)/d$ dessins for each choice of $c \in \mathbb{Z}_n$. These $\phi(n)/d$ dessins are permuted transitively by the group of hole operations H_j for j coprime to n . It follows from earlier results of Jones, Streit and Wolfart [9] that they form orbits under the absolute Galois group $\text{Gal } \overline{\mathbb{Q}}/\mathbb{Q}$, and are defined over a specific subfield of the cyclotomic field of the n th roots of unity, namely the splitting field of p . This allows us to give defining equations for a few special cases of these dessins.

After two sections giving background on dessins and some simple examples, the basic properties of generalised Paley dessins are explained and determined in Sections 4–7, and further examples are given in Section 8, with some of the techniques illustrated for $n = 7$ and $p = 29$ in Section 9. The main theorem is proved in Section 10. Galois conjugacy and defining equations are considered in Sections 11 and 12, and there is a brief discussion of non-faithful primitive dessins in Section 13. A technical problem concerning number theory is dealt with in Section 14.

2. DESSINS D'ENFANTS

This is just a brief outline of the theory of dessins; for more details see [10]. By Belyi's Theorem, a compact Riemann surface S is defined, as an algebraic curve, over an algebraic number field if and only if it admits a Belyi function, that is, a non-constant meromorphic function $\beta : S \rightarrow \Sigma = \mathbb{P}^1(\mathbb{C}) = \mathbb{C} \cup \{\infty\}$ with at most three critical values. Applying a Möbius transformation to Σ if necessary, one can assume these are 0, 1 and ∞ . Then β can be represented combinatorially by a *dessin d'enfant* \mathcal{D} , a bipartite map on S with black and white vertices at the points over 0 and 1, and edges over the unit interval $[0, 1] \subset \mathbb{R}$; the edges correspond bijectively to the sheets of the covering β , and the faces correspond to the points over ∞ , called the face-centres. (It is sometimes useful to extend \mathcal{D} to a tripartite map, with red vertices over ∞ and edges over \mathbb{R} .)

The automorphism group $G = \text{Aut } \mathcal{D}$ of \mathcal{D} is the group of covering transformations of β , acting semi-regularly on the edges. We say that \mathcal{D} is *regular* if β is a regular covering, or equivalently G

acts regularly on the edges. In this case G acts transitively on the black and the white vertices and the face-centres, so all black and all white vertices have the same valencies n and m , and all faces are $2l$ -gons for some l . The triple (n, m, l) is called the *type* of \mathcal{D} . Then G has generators x, y, z (rotations around an incident black and white vertex and face-centre) satisfying

$$x^n = y^m = z^l = xyz = 1.$$

If $m \leq 2$ one can ignore the white vertices of \mathcal{D} (or more precisely, regard them as ordinary points on their incident edge or edges), giving an oriented map $\mathcal{M} = \mathcal{M}(\mathcal{D})$ in which the vertices are the black vertices of \mathcal{D} , the faces are the same, and the edges are the unions of the edges incident with each white vertex of \mathcal{D} . A second way of converting a dessin into a map, applicable to every dessin \mathcal{D} , is to ignore vertex colours, regarding \mathcal{D} as an uncoloured map \mathcal{D}° on the same surface.

3. SIMPLE EXAMPLES

First we consider some simple examples of primitive and faithful actions.

Example 1 Let \mathcal{D} be the regular planar dessin of type $(1, m, m)$ embedding a complete bipartite graph $K_{m,1}$ for some integer $m \geq 1$. This has one white vertex, joined by a single edge to each of the m black vertices. We will call \mathcal{D} an *m -star dessin* and denote it by $\mathcal{S}t_m$. The corresponding Belyĭ function $\Sigma \rightarrow \Sigma$ is $z \mapsto 1 - z^m$. The automorphism group G is a cyclic group C_m of order m , generated by $z \mapsto e^{2\pi i/m} z$. This group acts transitively and faithfully on the black vertices (the m th roots of 1 in \mathbb{C}), and acts primitively on them if and only if m is prime. Figure 1 shows $\mathcal{S}t_5$.





FIGURE 1. The star dessin $\mathcal{S}t_5$

Lemma 3.1. *Let \mathcal{D} be a regular dessin with automorphism group G . Then G acts regularly (and thus faithfully) on the black vertices of \mathcal{D} if and only if $\mathcal{D} \cong \mathcal{S}t_m$ for some $m \geq 1$.*

Proof. The star dessins clearly have this property. For the converse, if G acts regularly on the black vertices of \mathcal{D} , then since G also acts regularly on the edges, there must be the same number of edges as black vertices, and hence the latter must all have valency 1; since the embedded graph is connected there can therefore be only one white vertex, so this graph is a complete bipartite graph $K_{m,1}$ for some m . Up to isomorphism, this bicoloured graph has a unique embedding as a dessin, namely as $\mathcal{S}t_m$. \square

FIGURE 2. The Fermat dessins \mathcal{F}_3 and \mathcal{F}_4 .

Example 2 The Fermat dessin \mathcal{F}_n of degree n is regular, of type (n, n, n) and genus $(n-1)(n-2)/2$, with

$$G = \langle x, y \mid x^n = y^n = [x, y] = 1 \rangle \cong C_n \times C_n.$$

The embedded graph is the complete bipartite graph $K_{n,n}$. The dessins \mathcal{F}_3 and \mathcal{F}_4 are shown in Figure 2: in the case of \mathcal{F}_3 opposite sides of the outer hexagon are identified to form a torus, while in the case of \mathcal{F}_4 sides of the outer 16-gon are identified, A with A and the others using the obvious D_4 symmetry, to give a surface of genus 3.

The automorphism group G acts transitively but not faithfully on the n black vertices, on the n white vertices and on the n faces. The kernels of these actions of G are the cyclic subgroups generated by the canonical generators x, y and $z := (xy)^{-1}$, so in each case G induces a permutation group $\overline{G} \cong C_n$ acting regularly. Each of these three actions is primitive if and only if n is prime. The kernels of these three actions of G have trivial intersection, so the intransitive action of G on the set of all $3n$ critical points is faithful.

Example 3 The quaternion group Q_8 of order 8 has, up to automorphisms, a unique generating triple of elements x, y, z of order 4 satisfying $xyz = 1$. They determine a regular dessin Q of type $(4, 4, 4)$ and genus 2 with automorphism group $G \cong Q_8$. This is shown in Figure 3 where opposite sides of the outer hexagon are identified with each other. There are two black vertices, two white vertices, and two faces, each pair transposed by G . In each case the action is primitive, and its kernel is the commutator subgroup G' (also the centre) of G , of order 2, generated by a half turn about the centre of the diagram. The corresponding hypermap is RPH2.4 in Conder's census of regular hypermaps [2]. The quotient Q/G' is the Fermat dessin \mathcal{F}_2 , an embedding of $K_{2,2}$ in the sphere.

Example 4 The preceding example can be extended to all odd primes, though the structure of the automorphism group is slightly different. For any prime $p > 2$, let

$$G = \langle a, b \mid a^{p^2} = b^p = 1, a^b = a^{p+1} \rangle \cong C_{p^2} \rtimes C_p$$

be the unique nonabelian group of order p^3 and exponent p^2 . Each element has the unique form $a^i b^j$ for $i \in \mathbb{Z}_{p^2}$ and $j \in \mathbb{Z}_p$, and it has order p^2 if and only if i is coprime to p . The commutator




FIGURE 3. The dessin Q (identify opposite sides of the outer octagon).

subgroup G' (also the centre) is $\langle a^p \rangle \cong C_p$, with $G/G' \cong C_p \times C_p$. By using standard generators $x = a$, $y = ab$ and $z = (xy)^{-1} = a^{-2(p+1)}b^{-1}$ of order p^2 we obtain a regular dessin \mathcal{D}_p of type (p^2, p^2, p^2) and genus $(p-1)^2(p+2)/2$, with automorphism group G acting primitively on the p black vertices, on the p white vertices, and on the p faces. The kernels of these actions, namely the normal subgroups $\langle x \rangle$, $\langle y \rangle$ and $\langle z \rangle$ of index p in G , all contain G' , so the action of G on the $3p$ special points of \mathcal{D}_p is not faithful, with kernel G' . This dessin \mathcal{D}_p is a regular p -sheeted covering of the Fermat dessin $\mathcal{D}_p/A' \cong \mathcal{F}_p$ of degree p , branched over all $3p$ of its critical points; for $p = 3$ or 5 the corresponding hypermap is RPH10.32 or RPH56.70 in [2].

4. GENERALISED PALEY DESSINS

For further examples, and an important definition, we need the following lemma:

Lemma 4.1. *Let G be a subgroup of the affine group $\text{AGL}_1(q)$ for some prime power $q = p^d$. Then G acts primitively on the field \mathbb{F}_q if and only if*

$$G = G_S := \{t \mapsto at + b \mid a \in S, b \in \mathbb{F}_q\}$$

for some subgroup S of order n in the multiplicative group $\mathbb{F}_q^* = \mathbb{F}_q \setminus \{0\}$, where either $n = d = 1$, or $n > 1$ and S satisfies the following equivalent conditions:

- (a) S acts by multiplication on \mathbb{F}_q as an irreducible subgroup of $\text{GL}_d(p)$;
- (b) S generates the additive group of \mathbb{F}_q ;
- (c) p has multiplicative order $d \bmod (n)$.

The proof is elementary, the only point requiring comment being that if G is primitive then it is transitive on \mathbb{F}_q , so it has order divisible by q and hence contains the unique Sylow p -subgroup of $\text{AGL}_1(q)$, namely the translation subgroup $T = \{t \mapsto t + b \mid b \in \mathbb{F}_q\}$ of order q . Note that G_S is a semidirect product $T \rtimes S$ of T , acting regularly on \mathbb{F}_q by translations, and the stabiliser $S = G_0$ of 0 , acting on \mathbb{F}_q by multiplication. In the case when $n = d = 1$ we have $G_S = T \cong C_p$.

We can use this result to construct a wider class of primitive dessins. Let $G = G_S$ where S satisfies the conditions of Lemma 4.1. Since \mathbb{F}_q^* is cyclic, so is its subgroup S , so let x be any generator of S . If y is any element of $G \setminus S$, then $G = \langle x, y \rangle$ since S , as a point stabiliser in the primitive group G , is a maximal subgroup of G . We can therefore define a regular dessin \mathcal{D} with $\text{Aut } \mathcal{D} \cong G$ by taking the generating triple (x, y, z) for G where $z := (xy)^{-1}$. The black vertices of

\mathcal{D} , of valency n , are identified with the cosets of $\langle x \rangle = S$ in G , and hence with the elements of \mathbb{F}_q , in each case admitting the natural action of G , so G acts primitively and faithfully on them. We will call such a dessin \mathcal{D} a *generalised Paley dessin*, since this construction is the natural extension to dessins of the concept of a generalised Paley map (see [5] or [10, Section 9.1.1]).

We will call n , p and d the *valency*, the *field characteristic* and the *dimension* of \mathcal{D} . (Of course, the field characteristic p should not be confused with the Euler characteristic χ of \mathcal{D} .) Let $GP(n)$ and $GP(n, p)$ denote the set of all generalised Paley dessins which have valency n , and those which also have field characteristic p . Thus $GP(1)$ consists of the primitive star dessins $\mathcal{S}t_m$, those with $m = p$ prime. As another simple special case we have the following:

Example 5 Let $\mathcal{D} \in GP(2)$, so $n = 2$, $d = 1$, $S = \{\pm 1\} \subseteq \mathbb{F}_p^*$ for some odd prime $q = p$, and G is the dihedral group D_p . If we choose $y \in T = C_p$ then \mathcal{D} is the regular dessin \mathcal{B}_p of type $(2, p, 2)$ embedding a complete bipartite graph $K_{p,2}$ in Σ , with p black vertices at the p th roots of unity and two white vertices at 0 and ∞ , each black and white pair joined by a single radial edge; the face-centres are obtained by multiplying the black vertices by $e^{\pi i/p}$. (This dessin is the union of $\mathcal{S}t_p$ and its reflection in the equatorial plane of the sphere Σ .) If we choose $y \in G \setminus T = D_p \setminus C_p$ then \mathcal{D} is the dual dessin \mathcal{B}_p^{12} of type $(2, 2, p)$, with white vertices and face-centres transposed, and with edges around the unit circle. In either case the automorphism group G , generated by $z \mapsto e^{2\pi i/p}z$ and $z \mapsto \bar{z}$, acts primitively and faithfully on the black vertices. Figure 4 shows planar drawings of \mathcal{B}_5 , with a 5-valent white vertex at infinity, and of \mathcal{B}_5^{12} .




FIGURE 4. The dessins \mathcal{B}_5 (with a white vertex at infinity) and \mathcal{B}_5^{12} .

The next example is more instructive.

Example 6 Let $q = 13$ and $n = 6$, so that S is the group of quadratic residues (non-zero squares) in \mathbb{F}_{13} . This is generated by the element 4, so we can define $x : t \mapsto 4t$. If we take $y : t \mapsto -t + b$ for any $b \neq 0$ in \mathbb{F}_{13} then x and y generate the unique subgroup $G = \text{AHL}_1(13) \cong C_{13} \rtimes C_6$ of index 2 in $\text{AGL}_1(13)$. Since y and $z := (xy)^{-1}$ have orders 2 and 3 we obtain an orientably regular dessin

\mathcal{D} of type $(6, 2, 3)$ with automorphism group G . Since the white vertices all have valency 2 we can ignore them and regard \mathcal{D} as a 6-valent triangular map \mathcal{M} . This map has 13 vertices, 39 edges and 26 faces, so it has genus 1; it is shown in Figure 5, where opposite sides of the outer hexagon are identified to form a torus. The vertices of \mathcal{M} are labelled with the elements of \mathbb{F}_{13} , and the neighbours of 0 are successive powers of the generator 4 of S . This map $(\{3, 6\}_{3,1})$ in the notation of Coxeter and Moser [3, Section 8.4]) is chiral, with its mirror image formed similarly using the generator $4^{-1} = 10$ of S ; in each case, the embedded graph is the Paley graph P_{13} .




FIGURE 5. A torus embedding of the Paley graph P_{13} .

The importance of generalised Paley dessins lies in Theorem 1.1, which asserts that a regular dessin \mathcal{D} is primitive and faithful if and only if it is a generalised Paley dessin. Before proving this in Section 10 we consider the basic properties of these dessins.

Let us define \mathcal{D} to have *colour constant* $c = c(x, y) = c(\mathcal{D}) \in \mathbb{Z}_n$ if $y \in Tx^c$, and for each $c \in \mathbb{Z}_n$ let $GP(n, p, c)$ denote the set of dessins $\mathcal{D} \in GP(n, p)$ with $c(\mathcal{D}) = c$. If we take $c = 0$, that is, if we choose y to be a non-identity element of T (as in \mathcal{B}_p , see Example 4), then y has order p and $z := (xy)^{-1}$ has order n , so \mathcal{D} has type (n, p, n) . In this case there are $q = p^d$ black vertices and faces, and $nq/p = np^{d-1}$ white vertices, so \mathcal{D} has Euler characteristic

$$\chi = nq \left(\frac{2}{n} + \frac{1}{p} - 1 \right) = p^{d-1}(2p + n - np)$$

and genus $g = 1 - \frac{\chi}{2}$. If we take $c = -1$ then y has order n while z , an element of T , has order p , so we obtain a dessin of type (n, n, p) , with χ and g the same as for $c = 0$ (see \mathcal{B}_p^{12} in Example 4). However, if we take some $c \neq 0, -1$ then y has the same order as x^c , namely $m = n/\gcd(n, c)$, while z has order $l = n/\gcd(n, c + 1)$, so \mathcal{D} has type (n, m, l) with

$$\chi = nq \left(\frac{1}{n} + \frac{1}{m} + \frac{1}{l} - 1 \right) = q(1 + \gcd(n, c) + \gcd(n, c + 1) - n)$$

and $g = 1 - \frac{x}{2}$.

If $m = 2$ then \mathcal{D} can be regarded as a map \mathcal{M} by ignoring its white vertices. This happens if and only if either $c = 0$ and $p = 2$, so that \mathcal{D} has type $(n, 2, n)$ and genus $1 + 2^{d-2}(n-4)$ with n odd, or $c = n/2$ with n even, so that p is odd and \mathcal{D} has type $(n, 2, l)$ with $l = n$ or $n/2$ as $n \equiv 0$ or $2 \pmod{4}$. In this last case the genus is $1 + q(n-4)/4$ or $1 + q(n-6)/4$ respectively.

Two generalised Paley dessins \mathcal{D} and \mathcal{D}' in $GP(n, p)$ are isomorphic if and only if their standard generating pairs x, y and x', y' are equivalent under $\text{Aut } G$. This group is isomorphic to $\text{AGL}_1(q) = \text{AGL}_1(q) \rtimes \text{Gal } \mathbb{F}_q$, acting by conjugation on its normal subgroup G , regarded as a subgroup of $\text{AGL}_1(q)$. It follows that $\mathcal{D} \cong \mathcal{D}'$ if and only if

- $x' = x^\gamma$ for some $\gamma \in \text{Gal } \mathbb{F}_q \cong \mathbb{C}_d$;
- $c(x', y') = c(x, y)$, i.e. $y' \in T(x')^c$ where $y \in Tx^c$.

If we denote \mathcal{D} by $\mathcal{P}(x, c)$ then we have $\mathcal{P}(x, c) \cong \mathcal{P}(x', c)$ if and only if x and x' are Galois conjugate and $c = c'$. It follows from condition (b) of Lemma 4.1 that $\text{Gal } \mathbb{F}_q$ acts semiregularly on generators x for S , so it has $\phi(n)/d$ orbits on them; independently there are n choices for c , so we have:

Proposition 4.2. *For each integer $n \geq 1$ and prime p coprime to n , we have $|GP(n, p)| = n\phi(n)/d$, with $|GP(n, p, c)| = \phi(n)/d$ for each $c \in \mathbb{Z}_n$, where d is the multiplicative order of $p \pmod{n}$.*

The mirror image of $\mathcal{D} = \mathcal{P}(x, c)$ is the generalised Paley dessin $\overline{\mathcal{D}} = \mathcal{P}(x^{-1}, c)$, in the same set $GP(n, p, c)$ as \mathcal{D} . Then \mathcal{D} is real if $\overline{\mathcal{D}} \cong \mathcal{D}$, that is, the corresponding hypermap is fully regular; this happens if and only if x is inverted by some $\gamma \in \text{Gal } \mathbb{F}_q$, and since $\text{Gal } \mathbb{F}_q$ is generated by the Frobenius automorphism $t \mapsto t^p$ we have:

Proposition 4.3. *The dessins $\mathcal{D} \in GP(n, p)$ are all real or all form chiral pairs as -1 is or is not congruent to a power of $p \pmod{n}$.*

Note that if $p^i \equiv -1 \pmod{n}$, then $p^{2i} \equiv 1 \pmod{n}$, so either $n = 2$ and p is odd, or $n > 2$, d is even and $i \equiv d/2 \pmod{d}$. Let \mathbb{Z}_n^* denote the multiplicative group of units mod (n) , and let $\mathbb{Z}_{n,p}^*$ denote the subgroup of \mathbb{Z}_n^* of order d generated by (the congruence class of) p . Then Proposition 4.3 states that the dessins in $GP(n, p)$ are real or chiral as $-1 \in \mathbb{Z}_{n,p}^*$ or not.

In order to characterise the pairs n, p with $-1 \in \mathbb{Z}_{n,p}^*$, let $n_r = r^{e_r}$ (r prime, $e_r \geq 1$) be the prime powers appearing in the factorisation of n , and let d_r be the multiplicative order of $p \pmod{n_r}$. Then we have the following:

Proposition 4.4. *There is some i such that $p^i \equiv -1 \pmod{n}$ if and only if the orders d_r for primes $r > 2$ dividing n all have the same 2-part $2^k \geq 2$, where $p \equiv -1 \pmod{n_2}$ if $n_2 \geq 2$, and $k = 1$ if $n_2 \geq 4$. In these circumstances the congruence $p^i \equiv -1 \pmod{n}$ is satisfied by $i = \text{lcm}\{d_r/2 \mid r > 2\}$ if some prime $r > 2$ divides n , and otherwise by $i = 1$.*

For instance, if $p = 2$ then these conditions are satisfied for $n = 3$ and $n = 5$, but not for $n = 15$: the orders $d_3 = 2$ and $d_5 = 4$ are both even but have different 2-parts. More explicitly, $2^i \equiv -1 \pmod{3}$ if and only if i is odd, whereas $2^i \equiv -1 \pmod{5}$ if and only if $i \equiv 2 \pmod{4}$, and these two conditions are incompatible. However, if $p = 2$ and $n = 99$ then $d_3 = 6$ and $d_{11} = 10$ are both even and have the same 2-part, so the conditions are satisfied with $i = 15$.

The primality of p is irrelevant here: as for many other properties of generalised Paley dessins, the conditions depend only on the congruence class of $p \bmod (n)$. Of course, Dirichlet's Theorem shows that every congruence class in \mathbb{Z}_n^* contains infinitely many primes.

To avoid a longer digression, the proof of Proposition 4.4 is postponed to Section 14.

Warning. If any dessin is real then it admits a reflection, so the underlying Riemann surface is symmetric. However, the converse is not true: for example the dessins in Example 6 (one of them shown in Figure 5) are chiral, whereas their Riemann surface, the hexagonal torus, is symmetric.

5. OPERATIONS ON DESSINS

For each permutation π of $\{0, 1, 2\}$ of order 2 or 3 there is a duality or triality operation D^π on dessins, permuting the sets V_0, V_1 and V_2 of black and white vertices and face-centres by renaming them. These form a group $\Omega \cong S_3$ of operations $D^\pi : \mathcal{D} \mapsto \mathcal{D}^\pi$ on dessins, preserving their underlying surface and automorphism group, and permuting the three periods in their type. If the subgroup of Ω leaving \mathcal{D} invariant has order k , then \mathcal{D} lies in an orbit of Ω consisting of $6/k$ mutually non-isomorphic dessins.

The duality operation $D^{12} : \mathcal{D} \mapsto \mathcal{D}^{12}$, transposing white vertices and faces, preserves each set $GP(n, p)$. Since $\mathcal{D}^{12} = \mathcal{P}(x, c')$ where $1 + c + c' = 0$ in \mathbb{Z}_n , it follows that $\mathcal{D} \cong \mathcal{D}^{12}$ if and only if n is odd and $c = (n - 1)/2$. The other two duality operations, transposing black vertices with white vertices or faces, do not preserve the set of all generalised Paley dessins. For instance, if $\mathcal{D} \in GL(n, p, c)$ then the black/white dual \mathcal{D}^{01} of \mathcal{D} is a generalised Paley dessin if and only if $c \in \mathbb{Z}_n^*$, in which case $\mathcal{D}^{01} \in GP(n, p, c^{-1})$ where c^{-1} denotes the multiplicative inverse of $c \bmod (n)$. In particular, $\mathcal{D} \cong \mathcal{D}^{01}$ if and only if some $\alpha \in \text{Aut } G = \text{AGL}_1(q)$ transposes the standard generators x and y of G . There are now two cases to consider. If $c = 1$ then x and y have the form $t \mapsto at$ and $t \mapsto at + b$ for some $a \neq 1$ and $b \neq 0$, in which case conjugation by $t \mapsto -t + b(1 - a)^{-1}$ transposes them. If $c \neq 1$ then x and y are transposed if and only if x and x^c are transposed by some element of $\text{Gal } \mathbb{F}_q$; this is equivalent to d being even and $p^{d/2} \equiv c \bmod (n)$. Similarly, $\mathcal{D} \cong \mathcal{D}^{02}$ if and only if either $c' = 1$ (equivalently $c = -2$), or d is even and $p^{d/2} \equiv c' (= -1 - c) \bmod (n)$.

The mutually inverse triality operations D^{012} and D^{210} act by cyclically permuting the standard generators x, y, z of G , so \mathcal{D} is invariant under one (and hence both) of these operations if and only if there is some $\alpha \in \text{Aut } G$ inducing this 3-cycle. If $c = 1$ then the existence of α implies that $c' = 1$, so that $n = 3$; thus $d \leq 2$, so $\alpha \in \text{AGL}_1(q)$. Since p is coprime to $n = 3$, α must have the form $t \mapsto at + b$ where the element $a \in S$ has multiplicative order 3. Conversely, given any $y \neq x$ in Tx there exists such an automorphism α inducing the required 3-cycle, so \mathcal{D} is triality-invariant. If $c \neq 1$ then, arguing as for dualities, we find that \mathcal{D} is triality-invariant if and only if d is divisible by 3 and $c \equiv p^{d/3}$ or $p^{2d/3} \bmod (n)$. To summarise, we have:

Proposition 5.1. *Let $\mathcal{D} \in GP(n, p, c)$. Then*

- $\mathcal{D} \cong \mathcal{D}^{12}$ if and only if n is odd and $c = (n - 1)/2$;
- $\mathcal{D} \cong \mathcal{D}^{01}$ if and only if either $c = 1$, or $p^{d/2} \equiv c \bmod (n)$ with d even;
- $\mathcal{D} \cong \mathcal{D}^{02}$ if and only if either $c = -2$, or $p^{d/2} \equiv -1 - c \bmod (n)$ with d even;

- $\mathcal{D} \cong \mathcal{D}^{012}$, or equivalently $\mathcal{D} \cong \mathcal{D}^{210}$, if and only if either $n = 3$ and $c = 1$, or $p^{d/3}$ or $p^{2d/3} \equiv c \pmod{n}$ with d divisible by 3.

Corollary 5.2. *A dessin $\mathcal{D} \in GP(n, p, c)$ is invariant under all six operations in Ω if and only if $n = 3$ and $c = 1$.*

The hole operations H_j ($j \in \mathbb{Z}_n^*$) act on the set of all maps of valency n by preserving the embedded graph and replacing the cyclic rotation of edges around each vertex (in the monodromy group) with its j th power. They preserve the automorphism group, but in general they may change the genus, since they change the faces. (For example, H_2 transposes the icosahedron and the great dodecahedron, which has pentagonal faces and has genus 4.) One can apply these operations to regular dessins of type (n, m, l) , regarded as maps by ignoring their black and white vertex-colours, provided j is coprime to both n and m . This applies to dessins $\mathcal{D} \in GP(n, p, c)$ with $c \neq 0$, since in this case m divides n ; the result is another dessin $H_j(\mathcal{D}) \in GP(n, p, c)$. If $c = 0$, so that \mathcal{D} has type (n, p, n) , one can instead apply $D^{12} \circ H_j \circ D^{12}$, since \mathcal{D}^{12} has type (n, n, p) . If x and x' are both generators of S then $x' = x^j$ for some j coprime to n , so we have:

Proposition 5.3. *The group of operations H_j ($j \in \mathbb{Z}_n^*$) acts transitively on each set $GP(n, p, c)$.*


The stabiliser of each dessin, and hence the kernel of this action, is the subgroup $\{H_j \mid j \in \mathbb{Z}_{n,p}^*\}$ of order d generated by H_p .

The most important hole operation, which applies for all valencies n , is H_{-1} , replacing each map or dessin with its mirror image by reflecting it in an edge. Since this operation has order 2 and commutes with those in Ω , we obtain a group $\Omega^* = \langle \Omega, H_{-1} \rangle \cong S_3 \times C_2 \cong D_6$ of operations on maps and dessins. If the subgroup of Ω^* leaving \mathcal{D} invariant has order k^* , then \mathcal{D} lies in an orbit of Ω^* consisting of $12/k^*$ mutually non-isomorphic dessins. In particular, if \mathcal{D} is real (that is, $H_{-1}(\mathcal{D}) \cong \mathcal{D}$), then so are all other dessins in that particular orbit of Ω , so $k^* = 2k$ and the orbits of Ω and Ω^* are identical; otherwise, $k^* = k$ and the orbit of Ω^* is the disjoint union of that of Ω and its mirror image. Using Proposition 4.3 and Corollary 5.2 we have:


Corollary 5.4. *A dessin $\mathcal{D} \in GP(n, p, c)$ is invariant under all twelve operations in Ω^* if and only if $n = 3$, $p \equiv 2 \pmod{3}$ and $c = 1$.*

Example 7 For each prime $p \equiv 2 \pmod{3}$ there is a unique dessin $\mathcal{D} \in GP(3, p, 1)$. It has type $(3, 3, 3)$ and genus 1. Since $d = 2$ it has automorphism group $G \cong T \rtimes C_3$ with $T \cong C_p^2$. The smallest of these dessins, for $p = 2$, is shown in Figure 6, on the left as a bicoloured map, and on the right, to emphasise its full symmetry under Ω^* , as a tricoloured embedded graph, with red vertices at $\beta^{-1}(\infty)$; in each case, opposite sides of the outer hexagon are identified to form a torus. More generally, the triangle group $\Delta = \Delta(3, 3, 3)$, acting by isometries on \mathbb{C} , has a translation subgroup $\Delta_0 \cong \mathbb{Z}^2$ of index 3, and \mathcal{D} is uniformised by the subgroup of index p^2 in Δ_0 generated by its p th powers. For more on the dessins in $GP(3)$ see Example 9 in Section 8.

Although these dessins $\mathcal{D} \in GP(3, p, 1)$ for $p \equiv 2 \pmod{3}$ are the only generalised Paley dessins which are invariant under the whole group Ω^* , there are other dessins, such as the Fermat dessins, with this property. (Indeed, any characteristic subgroup of finite index in the free group


 FIGURE 6. Two views of an Ω^* -invariant torus dessin.

F_2 uniformises such a dessin; for \mathcal{F}_n this is the subgroup generated by the n th powers and the commutators.) At the other extreme, many generalised Paley dessins are invariant under only the identity operation in Ω^* . These include the chiral torus dessins of type $(6, 2, 3)$ in $GP(6, p, 3)$ for primes $p \equiv 1 \pmod{6}$; see Figure 7 for one of the chiral pair with $p = 7$, drawn as a map.


 FIGURE 7. A chiral dessin of type $(6, 2, 3)$, drawn as a map.

One reason for the importance of understanding the action of this group Ω^* is that in Conder's census of regular and orientably regular maps and hypermaps [2], each entry represents a single orbit of Ω^* . In general, no information is given about the number of non-isomorphic objects in each orbit, although this can be deduced, after some computational labour, from the presentation for the automorphism group given there. The invariance results presented here provide an alternative solution to this problem for the generalised Paley dessins.

We have discussed invariance under operations in Ω , and Proposition 4.3 deals with H_{-1} , so there remain just the compositions of H_{-1} with the duality and triality operations, of orders 2 and 6 respectively. For instance $H_{-1}(\mathcal{D}^{12}) = (H_{-1}(\mathcal{D}))^{12} = \mathcal{P}(x^{-1}, c')$, and this is isomorphic to \mathcal{D} if and only if $-1 \in \mathbb{Z}_{n,p}^*$ and $c = (n-1)/2$, with n odd.

We have $H_{-1}(\mathcal{D}^{01}) \cong \mathcal{D}$ if and only if some $\alpha \in \text{Aut } G$ transposes x and y^{-1} . If $c = -1$, so that $y^{-1} \in Tx$, then conjugation by a suitable involution achieves this, as in the case of \mathcal{D}^{01} with

$c = 1$. If $c \neq -1$ it is necessary and sufficient that $p^{d/2} \equiv c^{-1} \pmod{n}$ with d even. Similarly, $H_{-1}(\mathcal{D}^{02}) \cong \mathcal{D}$ if and only if either $c = 0$, or $p^{d/2} \equiv (-1 - c)^{-1} \pmod{n}$ with d even. Finally, \mathcal{D} is invariant under the mutually inverse operations $H_{-1} \circ D^{012}$ and $H_{-1} \circ D^{210}$ of order 6 if and only if it is invariant under their cubes H_{-1} and their squares D^{012} and D^{210} , so by Propositions 4.3 and 5.1 this happens if and only if (a) $-1 \in \mathbb{Z}_{n,p}^*$, and either (b1) $n = 3$ and $c = 1$, or (b2) $p^{d/3} \equiv c \pmod{n}$ with d divisible by 3. Now (a) and (b1) are equivalent to $\mathcal{D} \in GF(3, p, 1)$ with $p \equiv 2 \pmod{3}$, while (a) and (b2) are equivalent to d being divisible by 6 with $c \equiv p^{d/3}$ or $p^{2d/3} \pmod{n}$ and $p^{d/2} \equiv -1 \pmod{n}$. Of course, we have already met the first condition, that $\mathcal{D} \in GF(3, p, 1)$ with $p \equiv 2 \pmod{3}$ in Corollaries 5.2 and 5.4, while the second condition is met when, for example, $n = 7$, $p \equiv 3$ or $5 \pmod{7}$ and $c = 2$ or 4 .

This proves the following analogue of Proposition 5.1:

Proposition 5.5. *Let $\mathcal{D} \in GP(n, p, c)$. Then*

- $\mathcal{D} \cong H_{-1}(\mathcal{D}^{12})$ if and only if n is odd and $c = (n - 1)/2$;
- $\mathcal{D} \cong H_{-1}(\mathcal{D}^{01})$ if and only if either $c = 1$, or $p^{d/2} \equiv c^{-1} \pmod{n}$ with d even;
- $\mathcal{D} \cong H_{-1}(\mathcal{D}^{02})$ if and only if either $c = 0$, or $p^{d/2} \equiv (-1 - c)^{-1} \pmod{n}$ with d even;
- $\mathcal{D} \cong H_{-1}(\mathcal{D}^{012})$, or equivalently $\mathcal{D} \cong H_{-1}(\mathcal{D}^{210})$, if and only if either $\mathcal{D} \in GF(3, p, 1)$ with $p \equiv 2 \pmod{3}$, or d is divisible by 6 with $c \equiv p^{d/3}$ or $p^{2d/3} \pmod{n}$ and $p^{d/2} \equiv -1 \pmod{n}$.

One can also ask which dessins $\mathcal{D} \in GP(n, p, c)$ are invariant under all the hole operations H_j for $j \in \mathbb{Z}_n^*$. By Proposition 5.3 this is equivalent to $GP(n, p, c)$ containing just one dessin (up to isomorphism), that is, to $\phi(n)/d = 1$. Since d is the multiplicative order of $p \pmod{n}$, this is equivalent to \mathbb{Z}_n^* being a cyclic group with generator p . Now it is known (see [7, Theorem 6.11], for instance) that \mathbb{Z}_n^* is cyclic if and only if $n = 1, 2, 4, r^e$ or $2r^e$ where r is an odd prime, in which case the number of generating elements is $\phi(\phi(n))$. Since we are assuming that $n \geq 2$ we have:

Proposition 5.6. *The following are equivalent:*

- A dessin $\mathcal{D} \in GP(n, p, c)$ is invariant under all the hole operations H_j for $j \in \mathbb{Z}_n^*$;
- $GP(n, p, c)$ contains only one dessin;
- p has multiplicative order $d = \phi(n) \pmod{n}$;
- \mathbb{Z}_n^* is a cyclic group with generator p ;
- $n = 2, 4, r^e$ or $2r^e$ where r is an odd prime, and p generates \mathbb{Z}_n^* .

For instance, $n = 18$ satisfies these conditions, with $\phi(\phi(18)) = \phi(6) = 2$ generating classes, namely $p \equiv 5$ and $p \equiv 11 \pmod{18}$. Note that if these conditions are satisfied, they apply to all n dessins in $GP(n, p)$.

Borrowing a term from the theory of regular maps, let us define a regular dessin to be *kaleidoscopic* if it is invariant under Ω^* and all the relevant hole operations. From Corollary 5.4 and Proposition 5.6 we have:

Corollary 5.7. *A dessin $\mathcal{D} \in GP(n, p, c)$ is kaleidoscopic if and only if $n = 3$, $p \equiv 2 \pmod{3}$ and $c = 1$.*

See Example 7 for more information on these dessins.

6. ACTION ON WHITE VERTICES AND FACE CENTRES

The automorphism group G of any generalised Paley dessin \mathcal{D} acts primitively on the black vertices. It acts primitively on the white vertices if and only if their stabilisers are maximal subgroups of G . If $c \neq 0$ then these stabilisers are the conjugates of $\langle x^c \rangle$, so this action is primitive if and only if c is a unit mod (n) . However, if $c = 0$ then the stabilisers are cyclic subgroups of T , so the action is primitive if and only if n is prime and $d = 1$, that is, $p \equiv 1 \pmod{n}$.

The same argument applies to the face centres, except that we must replace c with $c' = -1 - c$, or equivalently with $c + 1$. We therefore have:

Proposition 6.1. *Let \mathcal{D} be a generalised Paley dessin, with automorphism group G . Then G acts primitively on the white vertices (resp. face-centres) of \mathcal{D} if and only if either*

- (a) c (resp. $c + 1$) is a unit mod (n) , or
- (b) n is prime, $p \equiv 1 \pmod{n}$ and $c = 0$ (resp. $c = -1$).

It follows that G acts primitively on the white vertices and the face centres, in addition to the black vertices, if and only if either

- (1) c and $c + 1$ are both units mod (n) , or
- (2) n is prime, $p \equiv 1 \pmod{n}$ and $c = 0$ or -1 .

Condition (1) implies that n is odd, and conversely if n is odd then one can satisfy this condition by taking $c = 1$.

7. QUOTIENT DESSINS

If \mathcal{D} is a generalised Paley dessin with automorphism group G , then \mathcal{D} has a quotient dessin \mathcal{D}/T where T is the translation subgroup of G , an elementary abelian group of order $q = p^d$. This dessin has a single black vertex since T acts transitively on those of \mathcal{D} . Since T is normal in G , \mathcal{D}/T is a regular dessin, with automorphism group $G/T \cong S \cong C_n$ where n is the valency of \mathcal{D} . If \mathcal{D} has colour constant $c \neq 0$ or -1 then the covering $\mathcal{D} \rightarrow \mathcal{D}/T$ is smooth, so \mathcal{D}/T has the same type as \mathcal{D} and has genus

$$\frac{1}{2} (1 - \gcd(n, c) - \gcd(n, c + 1) + n).$$

However, if $c = 0$ or -1 , so that \mathcal{D} has type (n, p, n) or (n, n, p) , then the covering is branched over the white vertices or face-centres; in these cases \mathcal{D}/T has type $(n, 1, n)$ or $(n, n, 1)$ and is isomorphic to $\mathcal{S}t_n^{01}$ or $\mathcal{S}t_n^{02}$, each of genus 0.

Example 8 If $n = 6$ then \mathcal{D}/T has genus 0, 2, 1, 1, 2, 0 as $c = 0, 1, \dots, 5$ in \mathbb{Z}_6 .

In each case \mathcal{D}/T , as a regular dessin with an abelian automorphism group of exponent n , is a regular quotient of the Fermat dessin \mathcal{F}_n . Now $\text{Aut } \mathcal{F}_n$ is isomorphic to $C_n \times C_n$, with standard generators x, y and z which act on points $[u, v, w]$ of the Fermat curve $u^n + v^n + w^n = 0$ by multiplying u, v or w by $\zeta_n := \exp(2\pi i/n)$ (so that $xyz = 1$ since u, v and w are homogeneous coordinates). Then \mathcal{D}/T is the quotient of \mathcal{F}_n by the subgroup $\langle x^c y^{-1}, x^{c'} z^{-1} \rangle$ where $1 + c + c' = 0$ in \mathbb{Z}_n .

8. FURTHER EXAMPLES

We dealt with the case $n = 2$ rather informally in Example 5. We can now use the above general arguments to deal with the next case.

Example 9 Let $n = 3$. If $p \equiv 1 \pmod{3}$ then $d = 1$, so there are $n\phi(n)/d = 6$ dessins in $GP(3, p)$, forming one chiral pair for each $c \in \mathbb{Z}_3$, whereas if $p \equiv 2 \pmod{3}$ then $d = 2$, so there are 3 real dessins, one for each $c \in \mathbb{Z}_3$. For $c = 0$ or 2 the dessins have type $(3, p, 3)$ or $(3, 3, p)$ respectively, and have genus $(p - 1)/2$ or $(p - 1)(p - 2)/2$ as $p \equiv 1$ or $2 \pmod{3}$, whereas for $c = 1$ they have genus 1 and type $(3, 3, 3)$. These torus dessins \mathcal{T}_p of type $(3, 3, 3)$ are quotients of the universal hypermap of type $(3, 3, 3)$, obtained by factoring out normal subgroups M of the triangle group $\Delta(3, 3, 3)$ which are contained in its translation subgroup with quotient C_p^d ; if $p \equiv 1 \pmod{3}$, so that $d = 1$, there are two such subgroups, conjugate in the extended triangle group, whereas if $p \equiv 2 \pmod{3}$, so that $d = 2$, there is one such subgroup, normal in it. The self-dualities and -trialities are:

- $\mathcal{D} \cong \mathcal{D}^{12}$ if and only if $c = 1$;
- $\mathcal{D} \cong \mathcal{D}^{01}$ if and only if either $c = 1$ or $p \equiv 2 \pmod{3}$ with $c = 2$;
- $\mathcal{D} \cong \mathcal{D}^{02}$ if and only if either $c = 1$ or $p \equiv 2 \pmod{3}$ with $c = 0$;
- $\mathcal{D} \cong \mathcal{D}^{012} \cong \mathcal{D}^{210}$ if and only if $c = 1$.



For instance, if $p = 2$ we have the tetrahedron (with white vertices at the midpoints of its edges) for $c = 0$ and its dual of type $(3, 3, 2)$ for $c = 2$, together with a real torus dessin \mathcal{T}_2 of type $(3, 3, 3)$ for $c = 1$, already shown in Figure 6. If $p = 5$ we have real dessins of type $(3, 5, 3)$ and $(3, 3, 5)$ and genus 6 (see RPH6.1 in [2] for the corresponding hypermaps) and a real torus dessin \mathcal{T}_5 . However, if $p = 7$ we have three chiral pairs, two of genus 3 and types $(3, 7, 3)$ and $(3, 3, 7)$ on Klein's quartic curve (see CH3.1), together with a pair \mathcal{T}_7 and $\overline{\mathcal{T}}_7$ of genus 1; those dessins of type $(3, 3, 7)$ and $(3, 3, 3)$ represent embeddings of the Heawood graph, with black and white vertices corresponding to the points and lines of the Fano plane $\mathbb{P}^2(\mathbb{F}_2)$ (see [15]). The dessin \mathcal{T}_7 is shown in Figure 8; with opposite sides of the outer hexagon identified to form a torus. One of the chiral pair of type $(3, 3, 7)$ is shown in Figure 9; sides of the outer 14-gon are identified, A with A , and the others by C_7 symmetry.

The case $n = 4$ is similar, so we omit it. Higher values of n show a more interesting variety of phenomena.

Example 10 Let $n = 7$, so that $|GP(7, p)|$ is as follows:

- if $p \equiv 1 \pmod{7}$ then $d = 1$, so there are 42 dessins;
- if $p \equiv 6 \pmod{7}$ then $d = 2$, so there are 21 dessins;
- if $p \equiv 2$ or $4 \pmod{7}$ then $d = 3$, so there are 14 dessins in each case;
- if $p \equiv 3$ or $5 \pmod{7}$ then $d = 6$, so there are 7 dessins in each case.

If $p \equiv 3, 5$ or $6 \pmod{7}$ then $p^{d/2} \equiv -1 \pmod{7}$, so by Proposition 4.3 the dessins $\mathcal{D} \in GP(7, p)$ are all real, whereas if $p \equiv 1, 2$ or $4 \pmod{7}$ they form chiral pairs. If $c = 0$ or 6 then \mathcal{D} has type


 FIGURE 8. The dessin \mathcal{T}_7 .

 FIGURE 9. A generalised Paley dessin of type $(3, 3, 7)$ and genus 3.

$(7, p, 7)$ or $(7, 7, p)$ respectively; in either case $\chi = p^{d-1}(7 - 5p)$, so \mathcal{D} has genus

$$g = \frac{1}{2} (5p^d - 7p^{d-1} + 2).$$

However, if $c \neq 0$ or 6 then \mathcal{D} has type $(7, 7, 7)$ and genus

$$g = 2p^d + 1.$$

Applying Proposition 5.1 we have:

- $\mathcal{D} \cong \mathcal{D}^{12}$ if and only if $c = 3$;
- $\mathcal{D} \cong \mathcal{D}^{01}$ if and only if either $c = 1$, or $c = 6$ with $p \equiv 3, 5$ or $6 \pmod{7}$;
- $\mathcal{D} \cong \mathcal{D}^{02}$ if and only if either $c = 5$, or $c = 0$ with $p \equiv 3, 5$ or $6 \pmod{7}$.
- $\mathcal{D} \cong \mathcal{D}^{012} \cong \mathcal{D}^{210}$ if and only if $c = 2$ or 4 with $p \equiv 2, 3, 4$ or $5 \pmod{7}$.

Let us consider the six congruence classes $p \not\equiv 0 \pmod{7}$ in turn. First let $p \equiv 1 \pmod{7}$, so the 42 dessins $\mathcal{D} \in GP(7, p)$ form 21 chiral pairs, three for each $c \in \mathbb{Z}_7$. If $c = 0$ or 6 the dessins have type $(7, p, 7)$ or $(7, 7, p)$ and genus $g = 5(p-1)/2$; these 12 dessins, together with six duals of type $(p, 7, 7)$ which are not generalised Paley maps, form three orbits of length 6 under the action of Ω^* , with each dessin isomorphic to the mirror image its appropriate dual. For instance, if $p = 29$ then $g = 70$, and these three orbits correspond to items CH70.8, CH70.9 and CH70.10 in Conder's census [2], each item representing one chiral pair of each of the above three types. (For details of the calculations supporting these and other similar claims, see the next section.) The remaining 30 dessins in $GP(7, p)$ with $c = 1, 2, 3, 4$ or 5 all have type $(7, 7, 7)$ and genus $2p + 1$; they form three orbits of length 6, each orbit consisting of one chiral pair with each of $c = 1, 3$ and 5, and there is one orbit of length 12 consisting of three chiral pairs with each of $c = 2$ and 4. For instance, if $p = 29$ these four orbits correspond to entries CH59.11–14 in [2], the last having length 12.

If $p \equiv 2$ or 4 $\pmod{7}$ then $d = 3$, and in each case there are 14 dessins, forming 7 chiral pairs, one for each $c \in \mathbb{Z}_7$. If $c = 0$ or 6 the corresponding pair have type $(7, p, 7)$ or $(7, 7, p)$ and genus $g = (5p^3 - 7p^2 + 2)/2$; together with a chiral pair of type $(p, 7, 7)$ not in $GP(7, p)$ they form an orbit of length 6. For instance, if $p = 2$, so that $g = 7$, this orbit corresponds to C7.2 in [2], with the pair for $c = 0$, regarded as maps, giving the Edmonds pair of orientably regular embeddings of the complete graph K_8 . (Historically, this was the first discovery of a chiral pair of maps of genus $g \neq 1$.) If $c \neq 0$ or 6 the corresponding pair have type $(7, 7, 7)$ and genus $g = 2p^3 + 1$; these ten dessins form two orbits, of lengths 6 (one pair each with $c = 1, 3$ and 5) and 4 (with $c = 2$ and 4). For instance, if $p = 2$ these orbits respectively correspond to CH17.7 and CH17.8 in [2].

If $p \equiv 3$ or 5 $\pmod{7}$ then $d = 6$, and in each case there are seven dessins, all real, one for each $c \in \mathbb{Z}_7$. If $c = 0$ or 6 they have type $(7, p, 7)$ or $(7, 7, p)$ and genus $(5p^6 - 7p^5 + 2)/2$; if $c \neq 0$ or 6 they have type $(7, 7, 7)$ and genus $2p^6 + 1$. For each p these seven dessins lie in three orbits of Ω : one of length 2 with $c = 2$ or 4, one of length 3 with $c = 1, 3$ and 5, and another of length 3 with $c = 0$ and 6 and a third dessin of type $(p, 7, 7)$ which is not a generalised Paley dessin. In all cases, even for $p = 3$, the genus is too large for the corresponding hypermaps to appear in [2].

Finally, if $p \equiv 6 \pmod{7}$ then $d = 2$ and there are 21 dessins, all real, three for each $c \in \mathbb{Z}_7$, corresponding to the three mutually inverse pairs $x^{\pm 1}$ of generators of \mathbb{Z}_7^* . If $c = 0$ or 6 they have type $(7, p, 7)$ or $(7, 7, p)$ and genus $(5p^2 - 7p + 2)/2$; if $c \neq 0$ or 6 they have type $(7, 7, 7)$ and genus $2p^2 + 1$. There is one orbit of Ω of length 6, consisting of the dessins with $c = 2$ or 4, and the other 15 dessins, together with three more of type $(p, 7, 7)$, form six orbits of length 3. Again the genus is too large for the corresponding hypermaps to be listed in [2].

For comparison, we briefly consider an example where n is composite, allowing a wider variety of types to appear:

Example 11 Let $n = 8$, so that $|GP(8, p)|$ is as follows:

- if $p \equiv 1 \pmod{8}$ then $d = 1$, with 32 chiral dessins;
- if $p \equiv 3$ or 5 $\pmod{8}$ then $d = 2$, with 16 chiral dessins in each case;
- if $p \equiv 7 \pmod{8}$ then $d = 2$, with 16 real dessins.

If $c = 0$ or 7 then \mathcal{D} has type $(8, p, 8)$ or $(8, 8, p)$ respectively; in either case $\chi = 2p^{d-1}(4 - 3p)$, so \mathcal{D} has genus

$$g = 1 - \frac{\chi}{2} = 3p^d - 4p^{d-1} + 1.$$

Thus if $p \equiv 1 \pmod{8}$ then $d = 1$ and $g = 3(p - 1)$; for instance, if $p = 17$ then $g = 48$ (see CH48.4–6 in [2]). If $p \equiv 3, 5$ or $7 \pmod{8}$ then $d = 2$ and $g = (3p - 1)(p - 1)$; for instance, if $p = 3, 5$ or 7 then $g = 16, 56$ or 120 (see CH16.1 and CH56.5–6 in the first two cases).

If $c = 1, 2, 5$ or 6 then \mathcal{D} has type $(8, 8, 4)$ or $(8, 4, 8)$ as c is odd or even. We have $\chi = -4p^d$, so $g = 2p + 1$ or $2p^2 + 1$ as $p \equiv 1 \pmod{8}$ or otherwise. For example, if $p = 17$ then $g = 35$ (CH35.4–8); if $p = 3$ then $g = 19$ (CH19.8–9); if $p = 5$ then $g = 51$ (CH51.21–24); if $p = 7$ then $g = 99$ (RPH99.21–25).

If $c = 3$ or 4 then \mathcal{D} has type $(8, 8, 2)$ or $(8, 2, 8)$ respectively, with $\chi = -2p^d$ and hence $g = p + 1$ or $p^2 + 1$ as $p \equiv 1 \pmod{8}$ or otherwise. For example, if $p = 17$ then $g = 18$ (the chiral pair C18.1 of Paley maps); if $p = 3$ then $g = 10$ (the chiral pair C10.3 of self-dual Biggs embeddings of K_9); if $p = 5$ then $g = 26$ (C26.1); if $p = 7$ then $g = 50$ (R50.7).

Example 12 The valencies $n = 2, 3, 7$ and 8 which we have considered in the above examples all have the property that $n + 1$ is a prime power. More generally, if $n = p^e - 1$ for some prime p , then for dessins $\mathcal{D} \in GP(n, p)$ we have $d = e$ and $G \cong \text{AGL}_1(q)$ where $q = p^e$. Indeed, since $\text{AGL}_1(q)$ acts doubly transitively on \mathbb{F}_q , but has no proper subgroups with this property, these are the only regular dessins in which G is doubly transitive and faithful on the black vertices. If, in addition, either $c = 0$ and $p = 2$, or n is even and $c = n/2$, then $m = 2$ and \mathcal{D} (with white vertices ignored) is a map \mathcal{M} , specifically one of the orientably regular embeddings of complete graphs K_q constructed by Biggs in [1]. For instance, if $n = 10$ and $c = 5$ we have the two chiral pairs C12.1 and C12.2 of Biggs embeddings \mathcal{M} of K_{11} , corresponding to dessins \mathcal{D} of type $(10, 2, 5)$ and genus 12; if $n = 15$ and $c = 0$ we have the chiral pair C45.2 of Biggs embedding of K_{16} , corresponding to dessins of type $(15, 2, 15)$ and genus 45.

Similarly, if $2n = p^e - 1$ for some prime p (necessarily odd) then again dessins $\mathcal{D} \in GP(n, p)$ have $d = e$, but now G is the subgroup $\text{AHL}_1(q)$ of index 2 in $\text{AGL}_1(q)$. If n is even, so that $q \equiv 1 \pmod{4}$, and $c = n/2$, then \mathcal{D} corresponds to one of the orientably regular Paley maps \mathcal{M} considered in [5]. For instance, if $n = 6$ we obtain a chiral pair of torus embeddings of the Paley graph P_{13} , one of which was shown in Figure 5.

9. TYPICAL CALCULATIONS

One can determine the values of c for generalised Paley dessins appearing in [2] by abelianising the presentations of $G = \text{Aut } \mathcal{D}$ given there, thus determining the quotient dessins \mathcal{D}/T , which share these values of c . With n, p and c known, the invariance results in Section 5 then give the length of the orbit of Ω^* corresponding to each entry in [2], that is, the number of non-isomorphic dessins it represents, information which is not given there.

Example 13 As an illustration, the entries CH70.8, CH70.9 and CH70.10 in [2] represent the orbits of Ω^* containing orientably regular chiral hypermaps of genus 70 and type $(7, 7, 29)$. In our terminology, these include the dessins \mathcal{D} in $GP(7, 29, c)$ with $c = 0$ or 6 (see Example 10 in

Section 8). In the case of CH70.8, the defining relations given for the automorphism group G of \mathcal{D} , in terms of the standard generators x and y (or R and S in the notation of [2]), are:

$$x^{-1}y^{-1}x^2y^2 = x^{-7} = y^{-1}x^{-1}yx^2yx^{-1}y^{-1} = 1.$$

Abelianising these, we see from the first relation that $xy \in G' = T$, so $y \in Tx^{-1}$ and hence $c = -1 = 6$ in \mathbb{Z}_7 . We obtain the same value for c in the cases of CH70.9, where the relations are

$$x^{-7} = xyx^3y^3 = y^{-1}x^{-1}yx^2yx^{-1}y^{-1} = y^{-7} = 1,$$

and CH70.10, with relations

$$x^{-7} = y^{-1}x^{-1}yx^2yx^{-1}y^{-1} = x^{-1}y^2x^2y^{-1}xy = y^{-7} = 1.$$

Since $c = 6$ the invariance results in Section 5 show that the only operations in Ω^* preserving these dessins \mathcal{D} of type $(7, 7, 29)$ are $H_{-1} \circ \mathcal{D}^{01}$ and the identity, so the orbits CH70.8, CH70.9 and CH70.10 of Ω^* containing them have length $12/2 = 6$. Applying triality, we see that each of these orbits contains one chiral pair of generalised Paley dessins of each of the types $(7, 7, 29)$ and $(7, 29, 7)$ (the latter with $c = 0$), together with one pair of type $(29, 7, 7)$ which are not generalised Paley dessins.

In order to determine which generalised Paley dessins \mathcal{D} are in which of these orbits we need to know in each case how x acts on V , or equivalently on T by conjugation. If we define $t := xy$, so that $T = \langle t \rangle$, $y = x^{-1}t$ and $y^{-1} = t^{-1}x$, the second relation for CH70.8 becomes

$$1 = x^{-1}.t^{-1}x.x^2.x^{-1}t.x^{-1}t = x^{-1}t^{-1}x^2tx^{-1}t = (t^{-1})^xt^{x^{-1}}t = t^{-r+r^{-1}+1},$$

where we define $r \in \mathbb{F}_{29}$ by $t^x = t^r$. Thus $-r+r^{-1}+1 = 0$ in \mathbb{F}_{29} , so solving the equivalent quadratic equation $r^2 - r - 1 = 0$ gives $r = -5$ or 6 , and of these only $r = -5$ satisfies $r^7 = 1$, as implied by the second relation. (The third relation gives no useful information.) Thus the dessin \mathcal{D} is defined within $GP(7, 29, 6)$ by the action $x : t \mapsto t^{-5}$ of x on T . Since the multiplicative inverse of -5 in \mathbb{F}_{29} is -6 , in the mirror image $\overline{\mathcal{D}} = H_{-1}(\mathcal{D})$ the action is given by $t \mapsto t^{-6}$. Applying the duality operation D^{12} gives the second chiral pair of generalised Paley dessins in CH70.8, of type $(7, 29, 7)$ with $c = 0$. Similar calculations show that in CH70.9 the corresponding chiral pair \mathcal{D} and $\overline{\mathcal{D}}$ have $r = -4$ and 7 , and in CH70.10 they have $r = -9$ and -13 . This accounts for all the elements r of order 7 in \mathbb{F}_{29}^* , and hence for all the $\phi(7)/d = 6$ dessins in each of $GP(7, 29, 6)$ and $GP(7, 29, 0)$.

One can apply similar arguments to the dessins in $GP(7, 29, c)$ for $c = 1, 2, \dots, 5$; these have type $(7, 7, 7)$ and genus 59, and correspond to entries CH59.11–14 in [2]. (Again, see Example 10 in Section 8.) The dessin \mathcal{D} representing CH59.11 has an automorphism group G with defining relations

$$x^{-7} = x^{-1}y^{-1}x^{-2}yxy^{-1} = xy^{-1}x^{-2}y^{-3} = 1.$$

Abelianising, we see that $y \in Tx^5$ and hence $c = 5$. It follows that the only non-identity operation in Ω^* preserving \mathcal{D} is D^{02} so this orbit has length $12/2 = 6$. If we define $t := yx^2 \in T$, so that $y = tx^{-2}$ and $y^{-1} = x^2t^{-1}$, the second relation gives

$$1 = x^{-1}.x^2t^{-1}.x^{-2}.tx^{-2}.x.x^2t = xt^{-1}x^{-2}txt = (t^{-1})^{x^{-1}}t^xt = t^{-r^{-1}+r+1}$$

where $t^x = t^r$. Thus $-r^{-1} + r + 1 = 0$ in \mathbb{F}_{29} , so $r^2 + r - 1 = 0$ giving $r = 5$ or -6 , and of these only $r = -6$ satisfies $r^7 = 1$. Thus in \mathcal{D} the action of x on T is given by $t \mapsto t^{-6}$, and in $\overline{\mathcal{D}}$ by $t \mapsto t^{-5}$. The other four generalised Paley dessins in the orbit, with $c = 1$ and 3 , are obtained by applying triality operations to this pair.

We obtain the same value for c , and hence for the length and structure of the orbit, in the cases of CH59.12, where the defining relations are

$$x^{-7} = xy^{-1}x^3yxy^{-1} = y^{-7} = x^{-1}yx^{-3}y^{-1}x^2y^{-1} = 1,$$

and CH59.13, where they are

$$x^{-7} = xyx^3y^{-2}x = yxyx^2y^{-1}xy = 1.$$

In these cases x acts on T in \mathcal{D} and $\overline{\mathcal{D}}$ with $r = -4, 7$ and $-9, -13$ respectively. (Note that we have now accounted for all six elements r of order $n = 7$ in \mathbb{F}_{29}^* , and hence for all nontrivial actions of S on T .)

However, in CH59.14 the defining relations

$$x^{-7} = xy^{-1}x^{-2}yx^{-1}y = y^2x^{-2}yxy = 1$$

imply that $c = 2$. In this case no non-identity operations in Ω^* leave \mathcal{D} invariant, so this orbit has length 12, and all the duality and triality operations in Ω must be applied to the pair \mathcal{D} and $\overline{\mathcal{D}}$ to find the other ten dessins, which have $c = 1, 2$ or 4 . In \mathcal{D} the action of x on T is given by $t \mapsto t^{-6}$, and in $\overline{\mathcal{D}}$ by $t \mapsto t^{-5}$.

In this example we have a favourable situation, with small parameters n, p and especially d . In general, if these are larger, and in particular if T and V have dimension $d > 1$ over \mathbb{F}_p , then the calculations will be much more laborious, and the use of computer tools will be required.

10. PROOF OF THEOREM 1.1

In order to prove Theorem 1.1 we need the concept of a *Frobenius group*. This is a transitive permutation group in which stabiliser of any two points is the identity, but the stabiliser of any single point is not (that is, the group does not act regularly). Each Frobenius group G is a semidirect product $N \rtimes G_0$ of a regular normal subgroup N (the *Frobenius kernel*, which is nilpotent by a theorem of Thompson [16]), and a point-stabiliser G_0 (a *Frobenius complement*). For more background on Frobenius groups, see [4, §V.8] or [14].

Proposition 10.1. *Let \mathcal{D} be a primitive regular dessin with automorphism group G and black vertex set V . Then the kernel K of the action of G on V is a cyclic normal subgroup of G , and either*

- (a) *the permutation group $\overline{G} = G/K$ induced by G on V is regular of prime degree, or*
- (b) *\overline{G} is a Frobenius group, and K is contained in the centre of G .*

Proof. The stabilisers G_v of vertices $v \in V$ are cyclic, and K is contained in them, so K is also cyclic. Since \mathcal{D} is primitive, these stabilisers are maximal subgroups of G . Having the same stabiliser is an G -invariant equivalence relation, so either these stabilisers are all equal or they are mutually distinct. If they are all equal, and hence equal to K , then \overline{G} is a regular permutation group

on V , of prime degree by primitivity. We may therefore assume that distinct vertices $v, w \in V$ have distinct stabilisers. These stabilisers are abelian, so the centraliser $C_G(G_v \cap G_w)$ of their intersection contains both G_v and G_w and is therefore equal to G ; thus $G_v \cap G_w$ is in the centre of G , so since the stabilisers G_u ($u \in V$) are mutually conjugate $G_v \cap G_w$ fixes every black vertex and is therefore contained in the kernel $K = \bigcap_{u \in V} G_u$ of the action of G on V . Clearly $G_v \cap G_w \geq K$, so $G_v \cap G_w = K$ and thus K is contained in the centre of G . In the permutation group $\overline{G} = G/K$ induced by G on V , the stabiliser $\overline{G}_v \cap \overline{G}_w = (G_v \cap G_w)/K$ of each pair $v \neq w$ in V is the identity. Since $G_v > K$ we have $\overline{G}_v > 1$, so \overline{G} is a Frobenius group. \square

Example 14 To see how K need not be a central subgroup in case (a), let

$$G = \langle x, y \mid x^n = y^p = 1, x^y = x^r \rangle \cong C_n \rtimes C_p,$$

where r is an element of prime order p in \mathbb{Z}_n^* . Then $G = \text{Aut } \mathcal{D}$ for a regular dessin \mathcal{D} , with standard generators x and y . There are p black vertices of valency n , and n white vertices of valency p . Now G permutes the black vertices primitively, inducing a regular permutation group $\overline{G} = G/K \cong C_p$ on them, with a kernel $K = \langle x \rangle$ which is cyclic but not central in G . For instance, if $p = 2$ and $r = -1$ then $G \cong D_n$ and \mathcal{D} , regarded as a map, is the n -valent hosohedron (or beachball) on the sphere.

Proof of Theorem 1.1. By their construction, generalised Paley dessins are primitive and faithful, as are p -star dessins for each prime p .

Conversely, if a regular dessin \mathcal{D} is primitive and faithful, then $G = \overline{G}$ and by Proposition 10.1 this is a regular or Frobenius group. In the first case Lemma 3.1 shows that \mathcal{D} is a star map, so we may assume that G is a Frobenius group.

As in any Frobenius group, the Frobenius kernel N of G is a nilpotent normal subgroup, acting regularly on V . Since G acts primitively, N is characteristically simple (for otherwise it would contain a non-identity normal and hence transitive subgroup of G as a proper subgroup) and it is therefore a direct product of isomorphic simple groups. Since N is nilpotent these must be cyclic groups of prime order, so N is elementary abelian, of order $q = p^d$ for some prime p , and we can regard N as a d -dimensional vector space over the field \mathbb{F}_p . Since N acts regularly on V we can identify V with N , acting on itself by translations. The Frobenius complement G_0 , the stabiliser of the vector 0, acts by conjugation on this vector space V as a subgroup of $\text{GL}_d(p)$, which is irreducible since G is primitive. Since G_0 is abelian it follows from Schur's Lemma that one can identify V with the additive group of the field \mathbb{F}_q , and G_0 with a subgroup S of the multiplicative group $\mathbb{F}_q^* = \mathbb{F}_q \setminus \{0\}$, which generates V (see, for example, [4, Satz II.3.10] with $f = 1$). Thus $G = V \rtimes G_0$ acts on V as a primitive subgroup of $\text{AGL}_1(q)$, as in Lemma 4.1. The black vertices of \mathcal{D} correspond to the cosets of the subgroup $G_0 = S$, and the cyclic order of edges around the vertex 0 is given by successive powers of a generator x of S , so \mathcal{D} is a generalised Paley dessin, as in Example 3. \square

11. GALOIS ACTION

The absolute Galois group $\Gamma = \text{Gal } \overline{\mathbb{Q}}/\mathbb{Q}$ acts on isomorphism classes of dessins, preserving their automorphism group and type. It therefore acts on the set of generalised Paley dessins \mathcal{D} , preserving their valency n and characteristic p . Recall that for a given n and p (not dividing n)

there are $n\phi(n)/d$ dessins, with $\phi(n)/d$ for each colour constant $c \in \mathbb{Z}_n$, where d is the multiplicative order of $p \bmod (n)$. In fact, Γ also preserves the value of c . One way to see this is to note that there is an induced action of Γ on the set of quotient dessins \mathcal{D}/T , since the translation group T of order $q = p^d$ is a characteristic subgroup of $G := \text{Aut } \mathcal{D}$. These are regular dessins with abelian automorphism groups (isomorphic to C_n), so they are defined over \mathbb{Q} (see [17]) and hence invariant under Γ . Since the values of c are in bijective correspondence with the isomorphism classes of these dessins, they too are preserved by Γ . Thus, for each triple n, p and c there is an action of Γ on the set $GP(n, p, c)$ of generalised Paley dessins with these parameters.

Question: is there a more direct proof that c is invariant under Γ ?

Since the group of hole operations H_j ($j \in \mathbb{Z}_n^*$) acts transitively on each of these sets $GP(n, p, c)$, it follows from [9, Theorem 2] that they are the orbits of Γ on generalised Paley dessins, and that each dessin \mathcal{D} in this orbit is defined over a subfield K of the cyclotomic field $\mathbb{Q}(\zeta_n)$, where $\zeta_n := e^{2\pi i/n}$: specifically, K is the fixed field of the subgroup $H = \{j \in \mathbb{Z}_n^* \mid H_j(\mathcal{D}) \cong \mathcal{D}\}$, where we identify $\text{Gal } \mathbb{Q}(\zeta_n)/\mathbb{Q}$ with \mathbb{Z}_n^* in the usual way. (This generalises a similar result for generalised Paley maps proved in [5]; see also [10, Section 9.1.1].) The length of this orbit is the degree $[K : \mathbb{Q}] = \phi(n)/d$ of this extension. Proposition 5.6 tells us when this degree is 1, so we have:

Proposition 11.1. *A dessin $\mathcal{D} \in GP(n, p, c)$ is defined over \mathbb{Q} if and only if $n = 2, 4, r^e$ or $2r^e$ where r is an odd prime, and p generates \mathbb{Z}_n^* .*

12. EQUATIONS

A significant and difficult problem is that of converting combinatorial data about a dessin into defining equations. Even with the group-theoretic advantage of a regular dessin, this has been accomplished in only a few families of simple cases: see [10, Section 9.2], for instance. One of these families consists of generalised Paley dessins.

Example 15 Suppose that $\mathcal{D} \in GP(n, p)$ where $p \equiv 1 \bmod (n)$, so that $d = 1$ and G is a metacyclic group $C_p \rtimes C_n$. If we take $c = 0$ then \mathcal{D} has type (n, p, n) and genus $(n-2)(p-1)/2$; the embedded graph is the complete bipartite graph $K_{p,n}$, with p black and n white vertices of valencies n and p . There are $\phi(n)$ such dessins \mathcal{D} in $GP(n, p, 0)$, one for each element x of order n in \mathbb{F}_p^* . As explained in Section 11 these $\phi(n)$ dessins are permuted transitively by the hole operations H_j ($j \in \mathbb{Z}_n^*$) and by the absolute Galois group, acting as $\text{Gal } \mathbb{Q}(\zeta_n)/\mathbb{Q} \cong \mathbb{Z}_n^*$. It is therefore sufficient to give defining equations, with coefficients in $\mathbb{Q}(\zeta_n)$, for just one of these dessins, since equations for the others can then be obtained by Galois conjugacy. Here we are in the favourable situation that \mathcal{D} is a cyclic covering of \mathcal{D}/T , which is a dessin on the Riemann sphere Σ . In fact, $\mathcal{D}/T = St_n^{01}$, with a black vertex of valency n at 0, and white vertices of valency 1 at the n th roots of 1, where the p -sheeted covering is branched. In Example 4 of [9] these facts are used to give (with a few minor changes of notation) the affine model

$$w^p = \prod_{j=1}^n (z - \zeta_n^j)^{u_j},$$

in variables $w, z \in \mathbb{C}$, for one of the underlying curves, where x acts on \mathbb{F}_p by $t \mapsto t^x = ut$ for $u \in \mathbb{Z}_n^*$. The corresponding Belyĭ function $(w, z) \mapsto z^n$ is the composition of the covering $\mathcal{D} \rightarrow \mathcal{D}/T = St_n^{01}$, $(w, z) \mapsto z$, with the Belyĭ function $z \mapsto z^n$ of St_n^{01} . Formulae for the standard generators of G are given in [9]; the automorphisms of order p are obvious, but those of order n require careful calculation. Replacing ζ_n with other primitive n th roots of 1 gives the remaining curves and dessins in this Galois orbit. (This example is also considered in [10, Section 9.2.4], where for simplicity n is assumed to be prime.) One can extend this to the Galois orbit $GP(n, p, -1)$ by using the duality D^{12} , replacing the variable z with $z/(z-1)$.

Problem. When $c = 0$ or -1 , so that \mathcal{D}/T has genus 0, can one extend these results for $d = 1$ to cases where $d > 1$? A simple example is the tetrahedron, the unique dessin $\mathcal{D} \in GP(3, 2, 0)$: this is on the sphere Σ , with $d = 2$ and $G \cong A_4 \cong V_4 \rtimes C_3$, and it has Belyĭ function

$$\beta(z) = -64 \frac{(z^3 - 1)^3}{(z^3 - 8)^3 z^3}.$$

(Magot and Zvonkin give this and many other Belyĭ functions on the sphere in [13].) Can one find defining equations for the unique dessin $\mathcal{D} \in GP(3, p, 0)$ for other primes $p \equiv 2 \pmod{3}$, which all give $d = 2$ (and have genus $(p-1)(p-2)/2$ with $G \cong V_{p^2} \rtimes C_3$)?

13. NON-FAITHFUL ACTIONS

In Section 3 we saw some examples of non-faithful actions of G , on black vertices or more generally on all special points. In the case of maps, Li and Širáň [11] have dealt thoroughly with non-faithful actions, and most of their results also apply to dessins, so here we just give a few examples to show how such actions on dessins are related to faithful actions.

If \mathcal{D} is a regular dessin with automorphism group G acting primitively but *not* faithfully on its black vertex set V , then Proposition 10.1 shows that the kernel K of the action of G on V , the intersection of the stabilisers G_v ($v \in V$), is a cyclic normal subgroup of G , contained in the centre of G if G/K is a Frobenius group. Now G/K acts primitively and faithfully on the vertex set $V/K = V$ of \mathcal{D}/K , so Theorem 1.1 shows that \mathcal{D}/K is a generalised Paley dessin. Then \mathcal{D} is a k -sheeted regular cyclic covering of \mathcal{D}/K , branched over its black vertices (and possibly its white vertices and/or face-centres), where $k = |K|$. In Example 2, for instance, where \mathcal{D} is the Fermat dessin \mathcal{F}_n , the kernel K is the subgroup $\langle x \rangle \cong C_n$ of $G = C_n \times C_n$ fixing the n black vertices and cyclically permuting the n white vertices and the n faces, so that \mathcal{D}/K is the star dessin St_n (see Example 1); the n -sheeted regular covering $\mathcal{D} \rightarrow \mathcal{D}/K$ is branched over the black vertices of St_n , but not over its white vertex or face centre. In Examples 3 and 4 the kernel K is the commutator subgroup G' of G , and \mathcal{D}/K is a primitive Fermat dessin \mathcal{F}_p ; in this case the p -sheeted covering is branched over all $3p$ special points of \mathcal{F}_p .

One can reverse this process, and construct non-faithful primitive dessins $\tilde{\mathcal{D}}$ from a faithful primitive dessin \mathcal{D} . Let $\mathcal{D} \in GP(n, p, c)$, with automorphism group $G = T \rtimes S$ as before. Let \tilde{S} be a cyclic group of order kn for some $k \in \mathbb{N}$, so that there is an epimorphism $\theta : \tilde{S} \rightarrow S$, and let $\tilde{G} = T \rtimes \tilde{S}$ where the action of \tilde{S} on T is given by composing that of S with θ . The unique subgroup $K = \ker \theta$ of order k in \tilde{S} is then a central cyclic subgroup of \tilde{G} , and θ extends to an

epimorphism $\tilde{\theta} : \tilde{G} \rightarrow G$ with kernel K , so that $\tilde{G}/K \cong G$. If we choose a generator \tilde{x} of \tilde{S} with $\theta(\tilde{x}) = x$, and any element $\tilde{y} \in \tilde{\theta}^{-1}(y) \setminus \tilde{S}$, they generate \tilde{G} and hence determine a primitive regular dessin $\tilde{\mathcal{D}}$ which is a cyclic covering of \mathcal{D} , with K as the kernel of the action of $\tilde{G} = \text{Aut } \tilde{\mathcal{D}}$ on the black vertices. There are q of these, of valency kn , while the numbers and valencies of the white vertices and faces, and hence the type and genus of $\tilde{\mathcal{D}}$, depend on the choice of \tilde{y} .

Example 16 If k is coprime to the order nq of G then the extension \tilde{G} of G by K splits, by the Schur–Zassenhaus Theorem. Since K is in the centre of \tilde{G} we have $\tilde{G} = G \times K$, so that $\tilde{\mathcal{D}}$ is the direct product (minimal common cover) of \mathcal{D} and a regular dessin \mathcal{K} with one black vertex and with automorphism group $K \cong C_k$. There are k such dessins \mathcal{K} , one for each choice of y as a power of the standard generator x of K ; for instance, taking $y = 1$ gives $\mathcal{K} = St_k^{01}$.

Example 17 For any odd prime p the unique dessin \mathcal{D} in $GP(2, p, 0)$ is the spherical dessin \mathcal{B}_p of type $(2, p, 2)$ in Example 5, with automorphism group $G \cong D_p$ acting primitively and faithfully on the p black vertices (and on the p faces). Here $S \cong C_2$, so we can take \tilde{S} to be a cyclic group of any even order $2k > 2$, with a unique epimorphism $\theta : \tilde{S} \rightarrow S$. To obtain a covering $\tilde{\mathcal{D}} \rightarrow \mathcal{D}$ we need to choose $\tilde{y} \in T\tilde{x}^{\tilde{c}} \setminus \tilde{S}$ for some even $\tilde{c} \in \mathbb{Z}_{2k}$. For instance, if $\tilde{c} = 0$ then $\tilde{y} \in T$, so $\tilde{\mathcal{D}}$ has type $(2k, p, 2k)$ and genus $p - 1$; there is branching over the black vertices and face-centres, but not the white vertices. For $p = 3, 5, 7, 11$ and 13 the corresponding hypermap in [2] is RPH2.2, RPH4.9, RPH6.5, RPH 10.18 and RPH12.4. If $\tilde{c} \neq 0$ then $\tilde{\mathcal{D}}$ has type $(2k, \tilde{m}, \tilde{l})$ where \tilde{m} and \tilde{l} are the orders of \tilde{y} and \tilde{z} ; for instance, if $\tilde{c} = 2$, so that $\tilde{y} = t\tilde{x}^2 = \tilde{x}^2t$ for some non-identity $t \in T$, then $\tilde{m} = \text{lcm}(k, p)$.

14. APPENDIX: THE PROOF OF PROPOSITION 4.4

Clearly $p^i \equiv -1 \pmod{n}$ if and only if $p^i \equiv -1 \pmod{n_r}$ for each prime power $n_r = r^{e_r}$ (r prime) appearing in the factorisation of n .

If $r > 2$ then $\mathbb{Z}_{n_r}^*$ is cyclic, and -1 is its unique involution, so $p^i \equiv -1 \pmod{n_r}$ if and only if $i \equiv d_r/2 \pmod{d_r}$ where p has even order d_r in $\mathbb{Z}_{n_r}^*$. If $r = 2$ then the same applies provided $n_2 = 4$, but if $n_2 > 4$ then $\mathbb{Z}_{n_2}^* = \langle 5 \rangle \times \langle -1 \rangle \cong C_{n_2/4} \times C_2$ (see [7, Theorem 6.10], for instance), a non-cyclic group with three involutions, $\pm 5^{n_2/8} = n_2/2 \pm 1$ and -1 . In this case, and also when $n_2 = 4$, we have $p^i \equiv -1 \pmod{n_2}$ if and only if $p \equiv -1 \pmod{n_2}$ and i is odd; when $n_2 = 2$ one can omit the requirement that i is odd. To summarise, $p^i \equiv -1 \pmod{n}$ if and only if

- a) $i \equiv d_r/2 \pmod{d_r}$ where p has even order d_r in $\mathbb{Z}_{n_r}^*$, for each odd r , and
- b) $p \equiv -1 \pmod{n_2}$ if n is even, with any i or any odd i as $n_2 = 2$ or $n_2 > 2$.

The congruences in (a) require that i is an odd multiple of $d_r/2$ for each odd r , so they are satisfied by the same i if and only if all such d_r have the same 2-part $2^k \geq 2$ in their factorisation: one can then take i to be the least common multiple of the integers $d_r/2$ for odd r . If $n \equiv 0 \pmod{4}$ then (b) implies that i must be odd, so by (a) we have $d_r \equiv 2 \pmod{4}$ for each odd r , that is, $k = 1$. If $n \not\equiv 0 \pmod{4}$ then (b) imposes no restrictions on p , apart from being coprime to n .

Acknowledgments

The authors are grateful to Rubén Hidalgo for some very helpful comments and corrections. The second author acknowledges support from the APVV Research Grant APVV-19-0308 and from the VEGA Research Grants 1/0423/20 and 1/0727/22.

REFERENCES

- [1] N. L. Biggs, Classification of complete maps on orientable surfaces, *Rend. Mat.* (6) 4 (1971), 645–655.
- [2] M. D. E. Conder, Regular maps and hypermaps of Euler characteristic -1 to -200 , *J. Combin. Theory Ser. B* 99 (2009), 455–459, with associated lists of computational data available at <http://www.math.auckland.ac.nz/~conder/hypermaps.html>.
- [3] H. S. M. Coxeter and W. O. J. Moser, *Generators and Relations for Discrete Groups*, Springer.
- [4] B. Huppert, *Endliche Gruppen I*, Springer-Verlag, Berlin / Heidelberg / New York, 1967.
- [5] G. A. Jones, Characterisations and Galois conjugacy of generalised Paley maps, *J. Combin. Theory Ser. B* 103 (2013), 209–219.
- [6] G. A. Jones, Paley and the Paley graphs, in: *Isomorphisms, Symmetries and Computations in Algebraic Graph Theory (Pilsen 2016)*, Springer Proc. Math. Statistics 305 (2020), pp. 155–183. Also available at [arXiv:1702.00285](https://arxiv.org/abs/1702.00285) (math.HO).
- [7] G. A. Jones and J. M. Jones, *Elementary Number Theory*, Springer, 1998.
- [8] G. A. Jones and M. Mačaj, Regular maps with primitive automorphism groups, submitted; for a longer version with additional computational data, see [arXiv:2303.02493](https://arxiv.org/abs/2303.02493).
- [9] G. A. Jones, M. Streit and J. Wolfart, Wilson’s map operations on regular dessins and cyclotomic fields of definition, *Proc. London Math. Soc.* (3) 100 (2010), 510–532.
- [10] G. A. Jones and J. Wolfart, *Dessins d’Enfants on Riemann Surfaces*, Springer, Cham, 2016.
- [11] C. H. Li and J. Širáň, Regular maps whose groups do not act faithfully on vertices, edges, or faces, *Europ. J. Combin.* 26 (2005), 521–541.
- [12] T K. Lim and C. E. Praeger, On generalised Paley graphs and their automorphism groups, *Michigan Math. J.* 58 (2009), 293–308.
- [13] N. Magot and A. Zvonkin, Belyi functions for Archimedean solids, *Discrete Math.* 217 (2000), 249–271.
- [14] D. S. Passman, *Permutation Groups*, W. A. Benjamin, New York – Amsterdam, 1968.
- [15] D. Singerman, Klein’s Riemann surface of genus 3 and regular imbeddings of finite projective planes, *Bull. London Math. Soc.* 18 (1986), 364–370.
- [16] J. G. Thompson, Normal p -complements for finite groups, *Math. Z.* 72 (1960), 332–354.
- [17] J. Wolfart and B. Mühlbauer, Regular dessins with abelian automorphism groups, in: J. Sander, J. Steuding and R. Steuding (eds), *From Arithmetic to Zeta-Functions*, Springer, Cham, 2016, pp. 527–533.

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SOUTHAMPTON, SOUTHAMPTON SO17 1BJ, UK
Email address: G.A.Jones@maths.soton.ac.uk

FACULTY OF MATHEMATICS, PHYSICS AND INFORMATICS, COMENIUS UNIVERSITY, BRATISLAVA, SLOVAKIA
Email address: macaj@dcs.fmph.uniba.sk