

Cosmological studies with VLBI

Cristiana Spingola^{a,*}

^a*INAF – Istituto di Radioastronomia,
Via Gobetti 101, I-40129, Bologna, Italy*

E-mail: spingola@ira.inaf.it

Current cosmological controversies can be solved if a sufficient level of precision is achieved by observations. Future surveys with the next generation of telescopes will offer significantly improved depth and angular resolution with respect to existing observations, opening the so-called “era of precision cosmology”. But, that era can be considered already started at the radio wavelengths with Very Long Baseline Interferometry (VLBI). In this paper, we give an overview on how VLBI is contributing to some open questions in contemporary cosmology by reaching simultaneously the largest distances and the smallest scales.

*15th European VLBI Network Mini-Symposium and Users’ Meeting (EVN2022)
11-15 July 2022
University College Cork, Ireland*

*Speaker

1. Introduction

In the current standard cosmological framework Λ CDM (Dark Energy Cold Dark Matter), simple gas (mostly H and He) evolves into highly complex galaxies, which are gravitationally bound systems made of metal enriched gas, stars, dust, supermassive black holes, that can become active and have a strong effect on the surrounding medium. All of these components are believed to be embedded in a dark matter halo [1]. The cosmological model expects that the growth of these structures is driven by the dark matter component and happens in an expanding Universe, as known for almost a century [2] but still controversial [3]. We are in an era where we can approach cosmology from an observational rather than a theoretical standpoint. The Λ CDM model has been successful at explaining the observations of the large-scale structure of the Universe, but it presents some issues on the small (galactic and sub-galactic) scales [4]. To clearly assess if such small-scale problems affect *all* galaxies, it is mandatory to robustly characterize also high redshift (high- z) galaxies. To do so, a primary requirement is high angular resolution and deep observations.

Current VLBI arrays are ideal to perform such tests, as they can provide milliarcsecond (mas) and sub-mas angular resolution (at high frequency) and $\mu\text{Jy beam}^{-1}$ sensitivity observations. However, if we want to use VLBI arrays, the selected targets must emit radiation at the radio wavelengths. Among the possible sources to investigate the small-scales of high- z galaxies there are active galactic nuclei (AGN). In particular, those AGN that shows collimated jets ("jetted AGN") can reach extremely large luminosities because of the effect of Doppler boosting [5]. Therefore, their radio emission can be detected up to the highest redshifts, making them ideal targets for studying the most distant Universe. Since jetted AGN can be cosmologically distant sources, some of them could be *strongly gravitationally lensed* (see [6] for a review on strong lensing). The magnification and amplification due to the presence of a foreground (lensing) massive and concentrated object can make possible the detection also of the faintest jetted AGN in the form of multiple and distorted images. Thus, strong lensing can provide an unbiased view on the jetted AGN population across the cosmic time. Both lensed and "unlensed" jetted AGN give valuable observational constraints that can directly test the standard model.

In this paper, we describe the contribution of VLBI to some current open issues in cosmology related to AGN and their surrounding medium (Sec. 2), dark matter (Sec. 3) and dark energy (Sec. 4). Finally, we conclude by discussing the role of current and future wide-field VLBI surveys for observational cosmology (Sec. 5).

2. AGN and molecular gas at high redshift

How supermassive black holes (SMBHs) form and evolve is still unclear. The presence of SMBHs of masses of $10^{8-9} M_{\odot}$ when the Universe was less than a Gyr old [7, 8] supports models of high mass BH seeds and super- (or even hyper-) Eddington accretion events. Some models assume an initial phase dominated by BH-BH mergers lasting up to $z \sim 6$. After this epoch, the accretion of the resulting SMBHs is believed to be driven by inflowing of their surrounding medium (e. g., [9]). From an observational point of view, jetted AGN have been found in gas-rich environments, which supports some of these models and at the same time suggests that mergers could be the triggering of radio-loudness in AGN [10, 11]. Therefore, observations of the molecular gas surrounding jetted

AGN can be a powerful method to test both SMBHs formation models and the activation of the radio-loudness across the cosmic time.

Very recently, the James Webb Space Telescope (JWST) has opened a new frontier for the study of high redshift galaxies, with a large number of candidates *during* the reionization epoch (at photometric redshifts $z > 7$, e.g. [12] and references therein). The observations with JWST are suggesting even older and more massive SMBHs than expected, which require complex super-Eddington growth [13, 14]. VLBI observations of jetted AGN are currently reported up to $z = 6.82$ [15], finding jets extended from a few to several hundreds of parsecs [16–34]. As for the molecular gas, at high- z detections of several J transitions of the CO and C⁺ emission lines have been reported in the radio/mm bands [35–41], which revealed the presence of massive molecular gas reservoirs around SMBHs, in agreement with the predictions from hydrodynamical simulations [e.g., 42, 43]. Moreover, detailed analysis of the profile of these emission line showed evidence for on-going mergers in gas-rich radio-loud AGN at $z > 6$ [44]. However, these results have been obtained for the brightest high- z sources and these sources were only partially spatially resolved. In this context, the magnification due to strong lensing is really powerful for detecting the low surface brightness emission of the low-J transitions of CO at high redshift. This boost in spatial resolution and sensitivity revealed both mergers (complex velocity fields) but also molecular gas disks (ordered rotation) in jetted AGN at high redshift [40, 45–49].

As an intermediate stage of the merger process, we expect to observe binary (or multiple) SMBH systems [50–52]. The interesting multiple SMBHs systems for the future pulsar timing arrays and the Laser Interferometer Space Antenna are those at *small separation*, as they will be the emitters of gravitational waves in the μHz – nHz regime [53, 54]. The fraction of the active pairs of SMBHs (binary/dual AGN) is very low in the local Universe, but it increases up to a few percent at high redshift [55]. Hence, ideally, the search for close pairs of SMBHs should be performed at $z > 1$. Only VLBI can spatially resolve such systems on parsec scales even at large distances. Indeed, using VLBI observations several pairs of AGN have been found so far [56–61].

The combination of strong lensing and VLBI has been proven to be a powerful method to unveil dual AGN candidates at high redshift and small separation. Three cases have been reported so far: MG B2016+112 at $z = 3.273$ [62, 63], PS J1721+8842 at $z = 2.37$ [64] and PKS 1830-211 at $z = 2.5$ [65]. As first suggested by [62], the detection of even a single strongly lensed binary AGN implies a larger population of AGN pairs in the early Universe than first thought. However, such estimate is highly limited by the very small number statistics, and these lensed (and unlensed) candidates should also be confirmed. Nevertheless, VLBI provides currently the only way to find the closest AGN pairs, hence probing the bound systems and the final merging stages at the largest distances (as summarized in Fig. 7 of [66]).

3. Dark matter distribution at high- z

Among the current small-scale issues of the ΛCDM model, the most persistent ones are the so-called *missing satellite* and the *cusp-core* problems [4, 67]. Both these observational-theoretical conflicts could be solved by questioning the cold dark matter paradigm and assuming an alternative ("warm") dark matter particle [68]. Gravitational lensing provides a direct way to both find the expected satellites orbiting around massive galaxies and infer the inner mass density profile to the

the cusp-core problem. In particular, the strong lensing systems showing many extended images in radial and tangential directions are ideal for such investigations. One can detect the sub-halos associated with a lensing galaxy or along the line of sight by means of "anomalies" in the flux, position or time delay in the lensed images. Such "anomalies" arise because the presence of a sub-halo that locally changes the magnification of that image, hence its properties (in particular flux and position) will be different from what expected for a smooth mass density distribution (i.e. without sub-halos). For instance, [69] recently detected mas-level positional anomalies in the lensed images positions inferred by VLBI observations in HS 0810+2554. This system shows eight extended lensed images of a background AGN with intrinsic integrated flux density of only 880 nJy. Given the angular scale of these astrometric perturbations, they could be due to a population of low-mass ($10^{5-7} M_{\odot}$) dark matter dominated sub-halos. Milliarcsecond-level astrometric anomalies were found also using global-VLBI observations of the lensing system MG J0751+2716, which shows compact and extended lensed images forming giant gravitational arcs [70]. Therefore, also for this system it is possible that the observed astrometric anomalies are due to low-mass sub-halos (see also [71–76]). The inner density profile of MG J0751+2716 was inferred at high precision, thanks to the many constraints from the VLBI-detected lensed images. The inner mass density profile has been found to be steeper than isothermal: such steepening could be caused by the dissipative process due to *in situ* star formation, which leads to an increased stellar content in the central regions of the lensing galaxy. Therefore, it is possible that the complex processes involving *baryonic* matter might be responsible for the current tensions regarding the small scales of galaxies. Only VLBI observations of extended gravitational arcs can test such complex models, as demonstrated by [77, 78] for testing the origin of the astrometric anomalies in MG J0751+2716.

Despite the advantages of using gravitational lensing and VLBI to study astrometric anomalies at mas and sub-mas level, we are currently limited by the small number of VLBI-detected systems. More strongly lensed jetted AGN observed with VLBI are needed to provide a conclusive interpretation on these dark matter studies.

4. Cosmological parameters from jetted AGN observed with VLBI

The research field of using VLBI observations of AGN to infer cosmological parameters (such as q_0 and H_0) has been pioneered by [79, 80]. They proposed a characteristic angular size θ for compact jetted AGN to build a cosmic distance ladder. For a cosmological rod of intrinsic length l_m , the θ - z relation can be written as $\theta(z) = l_m/D_A(z)$, where D_A is the angular diameter distance, which depends on the cosmological parameters. This method was applied for the first time using ~ 340 VLBI observations of AGN with known redshift by [81]. With these observations they built a $\theta - z$ plane up to $z \simeq 4$ and could test different cosmological models (namely, different values of q_0), despite the large uncertainty given by the limited number of sources. Also, early work of [82] proposed to measure the distance to jetted AGN by combining the observed proper motion, Doppler factor derived from equipartition between the energy density in magnetic and particles fields, and aspect ratio of the (superluminal) jet component. Recently [83] proposed to use VLBI-detected AGN also to measure the cosmic curvature (Ω_k), finding no significant deviations from a flat Universe. The caveat about these methods is whether it is possible to use jetted AGN as

standard rods. For this reason, a lot of research has been done to find the best jetted AGN "category" or some of their components for this cosmological application (see [84] and references therein).

Another methodology to test cosmological models is to measure jet proper motions (μ) across the cosmic time, as μ depends on the cosmological parameters [85–87]. The difficulty of this method is to obtain measurements of proper motions at high- z , as such measurements require long time baselines and high signal-to-noise ratios. Nevertheless, such measurements have been obtained up to $z \approx 5.5$ (e.g., [29, 33, 88, 89] and references therein). Recently, a reverse application of the μ - z plot has been proposed, namely for deriving the redshift of a radio-loud AGN by adopting the standard cosmological model [90]. This method will be particularly important for wide-field surveys, where there are AGN candidates lacking spectroscopic redshift.

A method to estimate H_0 is based on the variability of jetted AGN [91] and has been recently used by [92] on the Boston University blazar monitoring programme [93]. The approach is based on light travel-time ("causality") arguments, and can be extended from low- to high- z AGN. The angular diameter distance D_A can be also expressed as $D_A(z) = c\Delta t/\theta_{\text{VLBI}}(1+z)$, where Δt is the variability timescale and θ_{VLBI} is the size of size of the variable region. By applying this method on the low- z AGN 3C 84, they infer a value for H_0 with a similar precision of other "late Universe methods" applied to samples of sources (summarized in [94]), demonstrating that this approach is a competitive cosmological probe. Recently, [95] revised the most important issue of this methodology, which is the determination of the Doppler factor, which becomes more difficult to obtain for high- z jetted AGN. The authors propose a method to derive the Doppler factor that is independent from the typical equipartition and inverse Compton brightness temperature limits [96].

Finally, gravitational time delays observed in strong lensing systems provide a one-step measurement for H_0 [97], because the time-delay distance is inversely proportional to H_0 [98]. Even if time delays between the multiple images can be estimated also using a spatially unresolved light curve (e.g., [99]), at the radio wavelengths i) VLBI can easily resolve the lensed images (typically separated by arcseconds), ii) AGN show intrinsic time variability on several time scales and iii) there is not the additional challenge of microlensing, as the (background) jetted AGN are typically extended several parsecs in size. Despite the powerful combination of VLBI and gravitational time delays, there has not been any time delay measured using VLBI yet. This is mostly due to the high cadence of the observing epochs required and the relatively short periods for such monitoring programs available (e.g., [100]). Instead, monitoring programs with the Very Large Array have been proven to be quite effective for estimating gravitational time delays also by innovatively using polarization [101–103].

5. Wide-field VLBI for cosmological probes and conclusions

Wide-field VLBI is now possible routinely. In the last ~ 10 years, impressive progress has been done to make this possible, and in particular the development of multi-phase centre correlation (i.e., the entire primary beam of VLBI experiment can be imaged [104, 105]) and multi-source self-calibration [106]. The progress in wide-field VLBI is timely, as there is the need to complement current (and future) surveys with radio observations at sub-arcsec angular resolution. At the same time, the exquisite sensitivity that wide-field VLBI surveys can reach now (e.g., $1.5 \mu\text{Jy beam}^{-1}$ in the *e*MERGE survey [107]) enables us to study at high image fidelity the faintest AGN population,

which only the lensing magnification revealed so far [69, 108, 109]. Hence, wide-field VLBI surveys can detect the different radio sources that are most suitable for each of the cosmological applications described in this paper. For instance, the mJIVE-20 survey at 1.4 GHz [110] has been used to search for radio-loud AGN in gas-rich mergers [111], to find strongly lensed jetted AGN at small angular separation [112] and for the hunt for AGN pairs [113]. The same wide-field VLBI survey has been also used for developing tools to identify and characterize VLBI sources with deep learning [114]. Such tools together with pipelines for calibrating large VLBI datasets (e.g., [115]) are crucial for current and future wide-field VLBI surveys.

In summary, VLBI can provide precise and independent cosmological probes. In this paper, we focused on how VLBI observations of AGN (lensed and unlensed) can test the current cosmological model. Future surveys with the Square Kilometer Array (SKA) and the follow-ups with SKA-VLBI will offer significantly improved sensitivity. In the future, we will not be limited by the small number statistics anymore, but we will also need synergy with multi-wavelength instruments to more robustly provide competitive precision on the estimate of the cosmological parameters.

References

- [1] A. Cimatti, F. Fraternali and C. Nipoti, *Introduction to Galaxy Formation and Evolution: From Primordial Gas to Present-Day Galaxies* (2019).
- [2] E. Hubble, *A Relation between Distance and Radial Velocity among Extra-Galactic Nebulae*, *Proceedings of the National Academy of Science* **15** (1929) 168.
- [3] A.G. Riess, *The expansion of the Universe is faster than expected*, *Nature Reviews Physics* **2** (2020) 10 [2001.03624].
- [4] J.S. Bullock and M. Boylan-Kolchin, *Small-Scale Challenges to the Λ CDM Paradigm*, *ARA&A* **55** (2017) 343 [1707.04256].
- [5] P. Padovani, D.M. Alexander, R.J. Assef, B. De Marco, P. Giommi, R.C. Hickox et al., *Active galactic nuclei: what's in a name?*, *A&A Rev.* **25** (2017) 2 [1707.07134].
- [6] T. Treu, *Strong Lensing by Galaxies*, *ARA&A* **48** (2010) 87 [1003.5567].
- [7] E. Bañados, B.P. Venemans, C. Mazzucchelli, E.P. Farina, F. Walter, F. Wang et al., *An 800-million-solar-mass black hole in a significantly neutral Universe at a redshift of 7.5*, *Nature* **553** (2018) 473 [1712.01860].
- [8] S. Belladitta, A. Caccianiga, A. Diana, A. Moretti, P. Severgnini, M. Pedani et al., *Central engine of the highest redshift blazar*, *A&A* **660** (2022) A74 [2201.08863].
- [9] O. Piana, P. Dayal, M. Volonteri and T.R. Choudhury, *The mass assembly of high-redshift black holes*, *MNRAS* **500** (2021) 2146 [2009.13505].
- [10] M. Chiaberge, R. Gilli, J.M. Lotz and C. Norman, *Radio Loud AGNs are Mergers*, *ApJ* **806** (2015) 147 [1505.07419].

- [11] F. Gao, L. Wang, W.J. Pearson, Y.A. Gordon, B.W. Holwerda, A.M. Hopkins et al., *Mergers trigger active galactic nuclei out to $z \sim 0.6$* , *A&A* **637** (2020) A94 [2004.00680].
- [12] L. Whitler, R. Endsley, D.P. Stark, M. Topping, Z. Chen and S. Charlot, *On the ages of bright galaxies 500 Myr after the big bang: insights into star formation activity at $z \gtrsim 15$ with JWST*, *MNRAS* **519** (2023) 157 [2208.01599].
- [13] F. Pacucci, P. Dayal, Y. Harikane, A.K. Inoue and A. Loeb, *Are the newly-discovered $z \sim 13$ drop-out sources starburst galaxies or quasars?*, *MNRAS* **514** (2022) L6 [2201.00823].
- [14] M. Volonteri, M. Habouzit and M. Colpi, *What if young $z > 9$ JWST galaxies hosted massive black holes?*, *MNRAS* **521** (2023) 241 [2212.04710].
- [15] E. Momjian, E. Bañados, C.L. Carilli, F. Walter and C. Mazzucchelli, *Resolving the Radio Emission from the Quasar P172+18 at $z = 6.82$* , *AJ* **161** (2021) 207 [2103.03481].
- [16] E. Momjian, C.L. Carilli and I.D. McGreer, *Very Large Array and Very Long Baseline Array Observations of the Highest Redshift Radio-Loud QSO J1427+3312 at $Z = 6.12$* , *AJ* **136** (2008) 344 [0805.2897].
- [17] S. Frey, Z. Paragi, L.I. Gurvits, D. Cseh and K.É. Gabányi, *High-resolution images of five radio quasars at early cosmological epochs*, *A&A* **524** (2010) A83 [1009.5023].
- [18] S. Frey, Z. Paragi, L.I. Gurvits, K.É. Gabányi and D. Cseh, *Into the central 10 pc of the most distant known radio quasar. VLBI imaging observations of J1429+5447 at $z = 6.21$* , *A&A* **531** (2011) L5 [1106.0717].
- [19] S. Frey, J.O. Fogasy, Z. Paragi and L.I. Gurvits, *On the Doppler boosting in the compact radio jet of the distant blazar J1026+2542 at $z = 5.3$* , *MNRAS* **431** (2013) 1314 [1302.2209].
- [20] H.M. Cao, S. Frey, L.I. Gurvits, J. Yang, X.Y. Hong, Z. Paragi et al., *VLBI observations of the radio quasar J2228+0110 at $z = 5.95$ and other field sources in multiple-phase-centre mode*, *A&A* **563** (2014) A111 [1402.2353].
- [21] K.E. Gabanyi, D. Cseh, S. Frey, Z. Paragi, L.I. Gurvits, T. An et al., *VLBI observation of the newly discovered $z = 5.18$ quasar SDSS J0131-0321.*, *MNRAS* **450** (2015) L57 [1503.06593].
- [22] Y. Zhang, T. An, S. Frey, K.É. Gabányi, Z. Paragi, L.I. Gurvits et al., *J0906+6930: a radio-loud quasar in the early Universe*, *MNRAS* **468** (2017) 69 [1702.03925].
- [23] H.M. Cao, S. Frey, K.É. Gabányi, Z. Paragi, J. Yang, D. Cseh et al., *VLBI observations of four radio quasars at $z > 4$: blazars or not?*, *MNRAS* **467** (2017) 950 [1701.04760].
- [24] R. Wang, E. Momjian, C.L. Carilli, X.-B. Wu, X. Fan, F. Walter et al., *Milliarcsecond Imaging of the Radio Emission from the Quasar with the Most Massive Black Hole at Reionization*, *ApJL* **835** (2017) L20 [1701.03260].

- [25] E. Momjian, C.L. Carilli, E. Bañados, F. Walter and B.P. Venemans, *Resolving the Powerful Radio-loud Quasar at $z \sim 6$* , *ApJ* **861** (2018) 86 [1807.02560].
- [26] S. Frey, O. Titov, A.E. Melnikov, P. de Vicente and F. Shu, *High-resolution radio imaging of two luminous quasars beyond redshift 4.5*, *A&A* **618** (2018) A68 [1807.06837].
- [27] K. Perger, S. Frey, K.É. Gabányi and L.V. Tóth, *Unveiling the weak radio quasar population at $z \geq 4$* , *MNRAS* **490** (2019) 2542 [1909.11012].
- [28] C. Spingola, D. Dallacasa, S. Belladitta, A. Caccianiga, M. Giroletti, A. Moretti et al., *Parsec-scale properties of the radio brightest jetted AGN at $z > 6$* , *A&A* **643** (2020) L12 [2010.11193].
- [29] T. An, P. Mohan, Y. Zhang, S. Frey, J. Yang, K.É. Gabányi et al., *Evolving parsec-scale radio structure in the most distant blazar known*, *Nature Communications* **11** (2020) 143 [2001.02809].
- [30] Y. Zhang, T. An, S. Frey, X. Yang, M. Krezinger, O. Titov et al., *J2102+6015: a young radio source at $z = 4.575$* , *MNRAS* **507** (2021) 3736 [2108.02142].
- [31] K.É. Gabányi, S. Frey, T. An, H. Cao, Z. Paragi, L.I. Gurvits et al., *A small radio galaxy at $z = 4.026$* , *Astronomische Nachrichten* **342** (2021) 1092 [2110.10964].
- [32] K. Perger, S. Frey, D.A. Schwartz, K.É. Gabányi, L.I. Gurvits and Z. Paragi, *Multi-scale Radio and X-Ray Structure of the High-redshift Quasar PMN J0909+0354*, *ApJ* **915** (2021) 98 [2105.06307].
- [33] Y. Zhang, T. An, A. Wang, S. Frey, L.I. Gurvits, K.É. Gabányi et al., *VLBI observations of VIK J2318–3113, a quasar at $z = 6.44$* , *A&A* **662** (2022) L2 [2205.05859].
- [34] M. Krezinger, K. Perger, K.É. Gabányi, S. Frey, L.I. Gurvits, Z. Paragi et al., *Radio-loud Quasars above Redshift 4: Very Long Baseline Interferometry (VLBI) Imaging of an Extended Sample*, *ApJS* **260** (2022) 49 [2204.02114].
- [35] F. Walter, R. Decarli, M. Sargent, C. Carilli, M. Dickinson, D. Riechers et al., *A Molecular Line Scan in the Hubble Deep Field North: Constraints on the CO Luminosity Function and the Cosmic H_2 Density*, *ApJ* **782** (2014) 79 [1312.6365].
- [36] M. Aravena, J.S. Spilker, M. Bethermin, M. Bothwell, S.C. Chapman, C. de Breuck et al., *A survey of the cold molecular gas in gravitationally lensed star-forming galaxies at $z > 2$* , *MNRAS* **457** (2016) 4406 [1602.00652].
- [37] M. Aravena, R. Decarli, F. Walter, R. Bouwens, P.A. Oesch, C.L. Carilli et al., *The ALMA Spectroscopic Survey in the Hubble Ultra Deep Field: Search for [CII] Line and Dust Emission in 6*, *ApJ* **833** (2016) 71 [1607.06772].
- [38] R. Decarli, F. Walter, M. Aravena, C. Carilli, R. Bouwens, E. da Cunha et al., *ALMA Spectroscopic Survey in the Hubble Ultra Deep Field: CO Luminosity Functions and the Evolution of the Cosmic Density of Molecular Gas*, *ApJ* **833** (2016) 69 [1607.06770].

- [39] D.A. Riechers, R. Pavesi, C.E. Sharon, J.A. Hodge, R. Decarli, F. Walter et al., *COLDz: Shape of the CO Luminosity Function at High Redshift and the Cold Gas History of the Universe*, *ApJ* **872** (2019) 7 [1808.04371].
- [40] C. Spingola, J.P. McKean, S. Vegetti, D. Powell, M.W. Auger, L.V.E. Koopmans et al., *SHARP - VI. Evidence for CO (1-0) molecular gas extended on kpc-scales in AGN star-forming galaxies at high redshift*, *MNRAS* **495** (2020) 2387 [1905.06363].
- [41] M. Bischetti, C. Feruglio, E. Piconcelli, F. Duras, M. Pérez-Torres, R. Herrero et al., *The WISSH quasars project. IX. Cold gas content and environment of luminous QSOs at $z \sim 2.4$ -4.7*, *A&A* **645** (2021) A33 [2009.01112].
- [42] T. Di Matteo, J. Colberg, V. Springel, L. Hernquist and D. Sijacki, *Direct Cosmological Simulations of the Growth of Black Holes and Galaxies*, *ApJ* **676** (2008) 33 [0705.2269].
- [43] R. Davé, R.A. Crain, A.R.H. Stevens, D. Narayanan, A. Saintonge, B. Catinella et al., *Galaxy cold gas contents in modern cosmological hydrodynamic simulations*, *MNRAS* **497** (2020) 146 [2002.07226].
- [44] Y. Khusanova, E. Bañados, C. Mazzucchelli, S. Rojas-Ruiz, E. Momjian, F. Walter et al., *The [CII] and FIR properties of $z > 6$ radio-loud quasars*, *A&A* **664** (2022) A39 [2204.08973].
- [45] D.A. Riechers, F. Walter, B.J. Brewer, C.L. Carilli, G.F. Lewis, F. Bertoldi et al., *A Molecular Einstein Ring at $z = 4.12$: Imaging the Dynamics of a Quasar Host Galaxy Through a Cosmic Lens*, *ApJ* **686** (2008) 851 [0806.4616].
- [46] D.A. Riechers, F. Walter, C.L. Carilli, F. Bertoldi and E. Momjian, *Formation of a Quasar Host Galaxy through a Wet Merger 1.4 Billion Years after the Big Bang*, *ApJL* **686** (2008) L9 [0808.3774].
- [47] D.A. Riechers, C.L. Carilli, R.J. Maddalena, J. Hodge, A.I. Harris, A.J. Baker et al., *CO($J = 1 \rightarrow 0$) in $z > 2$ Quasar Host Galaxies: No Evidence for Extended Molecular Gas Reservoirs*, *ApJL* **739** (2011) L32 [1106.2553].
- [48] R.P. Deane, I. Heywood, S. Rawlings and P.J. Marshall, *The preferentially magnified active nucleus in IRAS F10214+4724 - II. Spatially resolved cold molecular gas*, *MNRAS* **434** (2013) 23 [1307.6565].
- [49] D. Paraficz, M. Rybak, J.P. McKean, S. Vegetti, D. Sluse, F. Courbin et al., *ALMA view of RX J1131-1231: Sub-kpc CO (2-1) mapping of a molecular disk in a lensed star-forming quasar host galaxy*, *A&A* **613** (2018) A34 [1705.09931].
- [50] M.C. Begelman, R.D. Blandford and M.J. Rees, *Massive black hole binaries in active galactic nuclei*, *Nature* **287** (1980) 307.
- [51] T. Di Matteo, V. Springel and L. Hernquist, *Energy input from quasars regulates the growth and activity of black holes and their host galaxies*, *Nature* **433** (2005) 604 [astro-ph/0502199].

- [52] M. Volonteri, M. Habouzit and M. Colpi, *The origins of massive black holes*, *Nature Reviews Physics* **3** (2021) 732 [2110.10175].
- [53] S. Burke-Spolaor, S.R. Taylor, M. Charisi, T. Dolch, J.S. Hazboun, A.M. Holgado et al., *The astrophysics of nanohertz gravitational waves*, *A&A Rev.* **27** (2019) 5 [1811.08826].
- [54] M. Colpi, K. Holley-Bockelmann, T. Bogdanovic, P. Natarajan, J. Bellovary, A. Sesana et al., *Astro2020 science white paper: The gravitational wave view of massive black holes*, *arXiv e-prints* (2019) arXiv:1903.06867 [1903.06867].
- [55] A. De Rosa, C. Vignali, T. Bogdanović, P.R. Capelo, M. Charisi, M. Dotti et al., *The quest for dual and binary supermassive black holes: A multi-messenger view*, *New Astronomy Reviews* **86** (2019) 101525 [2001.06293].
- [56] C. Rodriguez, G.B. Taylor, R.T. Zavala, A.B. Peck, L.K. Pollack and R.W. Romani, *A Compact Supermassive Binary Black Hole System*, *ApJ* **646** (2006) 49 [astro-ph/0604042].
- [57] M. Bondi and M.A. Pérez-Torres, *VLBI Detection of an Active Galactic Nucleus Pair in the Binary Black Hole Candidate SDSS J1536+0441*, *ApJL* **714** (2010) L271 [1003.4849].
- [58] S. Burke-Spolaor, *A radio Census of binary supermassive black holes*, *MNRAS* **410** (2011) 2113 [1008.4382].
- [59] T. An, Z. Paragi, S. Frey, T. Xiao, W.A. Baan, S. Komossa et al., *The radio structure of 3C 316, a galaxy with double-peaked narrow optical emission lines*, *MNRAS* **433** (2013) 1161 [1305.1509].
- [60] R.P. Deane, Z. Paragi, M.J. Jarvis, M. Coriat, G. Bernardi, R.P. Fender et al., *A close-pair binary in a distant triple supermassive black hole system*, *Nature* **511** (2014) 57 [1406.6365].
- [61] K.É. Gabányi, T. An, S. Frey, S. Komossa, Z. Paragi, X.Y. Hong et al., *Four Dual AGN Candidates Observed with the VLBA*, *ApJ* **826** (2016) 106 [1605.09188].
- [62] C. Spingola, J.P. McKean, D. Massari and L.V.E. Koopmans, *Proper motion in lensed radio jets at redshift 3: A possible dual super-massive black hole system in the early Universe*, *A&A* **630** (2019) A108 [1908.11756].
- [63] D. Schwartz, C. Spingola and A. Barnacka, *Resolving Complex Inner X-Ray Structure of the Gravitationally Lensed AGN MG B2016+112*, *ApJ* **917** (2021) 26 [2103.08537].
- [64] C.S. Mangat, J.P. McKean, R. Brilenkov, P. Hartley, H.R. Stacey, S. Vegetti et al., *PS J1721+8842: a gravitationally lensed dual AGN system at redshift 2.37 with two radio components*, *MNRAS* **508** (2021) L64 [2109.03253].
- [65] S. Nair, C. Jin and M.A. Garrett, *Helical jet in the gravitationally lensed blazar PKS1830-211*, *MNRAS* **362** (2005) 1157 [astro-ph/0507015].

- [66] C. Spingola, D. Schwartz and A. Barnacka, *Milliarcsecond X-Ray Astrometry to Resolve Inner Regions of AGN at $z > 1$ Using Gravitational Lensing*, *ApJ* **931** (2022) 68 [2203.04245].
- [67] A. Del Popolo and M. Le Delliou, *Review of Solutions to the Cusp-Core Problem of the Λ CDM Model*, *Galaxies* **9** (2021) 123.
- [68] M.R. Lovell, V. Eke, C.S. Frenk, L. Gao, A. Jenkins, T. Theuns et al., *The haloes of bright satellite galaxies in a warm dark matter universe*, *MNRAS* **420** (2012) 2318 [1104.2929].
- [69] P. Hartley, N. Jackson, D. Sluse, H.R. Stacey and H. Vives-Arias, *Strong lensing reveals jets in a sub-microJy radio-quiet quasar*, *MNRAS* **485** (2019) 3009 [1901.05791].
- [70] C. Spingola, J.P. McKean, M.W. Auger, C.D. Fassnacht, L.V.E. Koopmans, D.J. Lagattuta et al., *SHARP - V. Modelling gravitationally lensed radio arcs imaged with global VLBI observations*, *MNRAS* **478** (2018) 4816 [1807.05566].
- [71] A.D. Biggs, I.W.A. Browne, N.J. Jackson, T. York, M.A. Norbury, J.P. McKean et al., *Radio, optical and infrared observations of CLASS B0128+437*, *MNRAS* **350** (2004) 949 [astro-ph/0402128].
- [72] J.P. McKean, L.V.E. Koopmans, C.E. Flack, C.D. Fassnacht, D. Thompson, K. Matthews et al., *High-resolution imaging of the anomalous flux ratio gravitational lens system CLASS B2045+265: dark or luminous satellites?*, *MNRAS* **378** (2007) 109 [astro-ph/0611215].
- [73] A. More, J.P. McKean, S. More, R.W. Porcas, L.V.E. Koopmans and M.A. Garrett, *The role of luminous substructure in the gravitational lens system MG 2016+112*, *MNRAS* **394** (2009) 174 [0810.5341].
- [74] J.W. Hsueh, C.D. Fassnacht, S. Vegetti, J.P. McKean, C. Spingola, M.W. Auger et al., *SHARP - II. Mass structure in strong lenses is not necessarily dark matter substructure: a flux ratio anomaly from an edge-on disc in B1555+375*, *MNRAS* **463** (2016) L51 [1601.01671].
- [75] J.W. Hsueh, L. Oldham, C. Spingola, S. Vegetti, C.D. Fassnacht, M.W. Auger et al., *SHARP - IV. An apparent flux-ratio anomaly resolved by the edge-on disc in B0712+472*, *MNRAS* **469** (2017) 3713 [1701.06575].
- [76] H.R. Stacey, A. Lafontaine and J.P. McKean, *Smoke on the water: CO and H₂O in a circumnuclear disc around a quasar at redshift 2.64*, *MNRAS* **493** (2020) 5290 [2002.07186].
- [77] D. Powell, S. Vegetti, J.P. McKean, C. Spingola, F. Rizzo and H.R. Stacey, *A novel approach to visibility-space modelling of interferometric gravitational lens observations at high angular resolution*, *MNRAS* **501** (2021) 515 [2005.03609].
- [78] D.M. Powell, S. Vegetti, J.P. McKean, C. Spingola, H.R. Stacey and C.D. Fassnacht, *A lensed radio jet at milliarcsecond resolution I: Bayesian comparison of parametric lens models*, *MNRAS* **516** (2022) 1808 [2207.03375].

- [79] L.I. Gurvits, *Milli-arcsecond structure of AGN at different redshifts (Review)*, in *Sub-arcsecond Radio Astronomy*, R.J. Davis and R.S. Booth, eds., p. 380, Jan., 1993.
- [80] K.I. Kellermann, *The cosmological deceleration parameter estimated from the angular-size/redshift relation for compact radio sources*, *Nature* **361** (1993) 134.
- [81] L.I. Gurvits, *Apparent Milliarcsecond Sizes of Active Galactic Nuclei and the Geometry of the Universe*, *ApJ* **425** (1994) 442.
- [82] D.C. Homan and J.F.C. Wardle, *Direct Distance Measurements to Superluminal Radio Sources*, *ApJ* **535** (2000) 575 [[astro-ph/0001298](#)].
- [83] S. Cao, J. Qi, M. Biesiada, X. Zheng, T. Xu, Y. Pan et al., *Milliarcsecond compact structure of radio quasars and the geometry of the Universe*, *Physics of the Dark Universe* **24** (2019) 100274.
- [84] S. Cao, X. Zheng, M. Biesiada, J. Qi, Y. Chen and Z.-H. Zhu, *Ultra-compact structure in intermediate-luminosity radio quasars: building a sample of standard cosmological rulers and improving the dark energy constraints up to $z \approx 3$* , *A&A* **606** (2017) A15 [[1708.08635](#)].
- [85] M.H. Cohen, P.D. Barthel, T.J. Pearson and J.A. Zensus, *Expanding Quasars and the Expansion of the Universe*, *ApJ* **329** (1988) 1.
- [86] K.I. Kellermann, M.L. Lister, D.C. Homan, R.C. Vermeulen, M.H. Cohen, E. Ros et al., *Sub-Milliarcsecond Imaging of Quasars and Active Galactic Nuclei. III. Kinematics of Parsec-scale Radio Jets*, *ApJ* **609** (2004) 539 [[astro-ph/0403320](#)].
- [87] S.G. Jorstad, A.P. Marscher, D.A. Morozova, I.S. Troitsky, I. Agudo, C. Casadio et al., *Kinematics of Parsec-scale Jets of Gamma-Ray Blazars at 43 GHz within the VLBA-BU-BLAZAR Program*, *ApJ* **846** (2017) 98 [[1711.03983](#)].
- [88] S. Frey, Z. Paragi, J.O. Fogasy and L.I. Gurvits, *The first estimate of radio jet proper motion at $z > 5$* , *MNRAS* **446** (2015) 2921 [[1410.8101](#)].
- [89] K. Perger, S. Frey, K.É. Gabányi, T. An, S. Britzen, H.-M. Cao et al., *Constraining the radio jet proper motion of the high-redshift quasar J2134-0419 at $z = 4.3$* , *MNRAS* **477** (2018) 1065 [[1803.10452](#)].
- [90] T. An, Y. Zhang and S. Frey, *A method for checking high-redshift identification of radio AGNs*, *MNRAS* **497** (2020) 2260 [[2007.09038](#)].
- [91] K. Wiik and E. Valtaoja, *The geometry of the universe from high resolution VLBI data of AGN shocks*, *A&A* **366** (2001) 1061.
- [92] J.A. Hodgson, B. L'Huillier, I. Liodakis, S.-S. Lee and A. Shafieloo, *Using variability and VLBI to measure cosmological distances*, *MNRAS* **495** (2020) L27 [[2003.10278](#)].
- [93] S. Jorstad and A. Marscher, *The VLBA-BU-BLAZAR Multi-Wavelength Monitoring Program*, *Galaxies* **4** (2016) 47.

- [94] L. Verde, T. Treu and A.G. Riess, *Tensions between the early and late Universe*, *Nature Astronomy* **3** (2019) 891 [1907.10625].
- [95] J.A. Hodgson, B. L’Huillier, I. Liodakis, S.-S. Lee and A. Shafieloo, *Estimating the feasibility of ‘standard speed-gun’ distances*, *MNRAS* **521** (2023) L44 [2301.06252].
- [96] A.C.S. Readhead, *Equipartition Brightness Temperature and the Inverse Compton Catastrophe*, *ApJ* **426** (1994) 51.
- [97] S. Refsdal, *On the possibility of determining Hubble’s parameter and the masses of galaxies from the gravitational lens effect*, *MNRAS* **128** (1964) 307.
- [98] T. Treu and P.J. Marshall, *Time delay cosmography*, *A&A Rev.* **24** (2016) 11 [1605.05333].
- [99] A. Barnacka, M.J. Geller, I.P. Dell’Antonio and W. Benbow, *Resolving the High-energy Universe with Strong Gravitational Lensing: The Case of PKS 1830-211*, *ApJ* **809** (2015) 100 [1504.05210].
- [100] C. Spingola, D. Dallacasa, M. Orienti, M. Giroletti, J.P. McKean, C.C. Cheung et al., *Radio follow-up of the γ -ray flaring gravitational lens JVAS B0218+357*, *MNRAS* **457** (2016) 2263 [1601.03591].
- [101] A.D. Biggs and I.W.A. Browne, *A revised lens time delay for JVAS B0218+357 from a reanalysis of VLA monitoring data*, *MNRAS* **476** (2018) 5393 [1802.10088].
- [102] A.D. Biggs, *The time delay of JVAS B1030+074 from VLA polarization monitoring*, *MNRAS* **481** (2018) 1000 [1808.00757].
- [103] A.D. Biggs, *The time delay of CLASS B1600+434 from VLA multi-frequency and polarization monitoring*, *MNRAS* **505** (2021) 2610 [2105.07905].
- [104] A.T. Deller, W.F. Brisken, C.J. Phillips, J. Morgan, W. Alef, R. Cappallo et al., *DiFX-2: A More Flexible, Efficient, Robust, and Powerful Software Correlator*, *PASP* **123** (2011) 275 [1101.0885].
- [105] J.S. Morgan, F. Mantovani, A.T. Deller, W. Brisken, W. Alef, E. Middelberg et al., *VLBI imaging throughout the primary beam using accurate UV shifting*, *A&A* **526** (2011) A140 [1302.6040].
- [106] J.F. Radcliffe, M.A. Garrett, R.J. Beswick, T.W.B. Muxlow, P.D. Barthel, A.T. Deller et al., *Multi-source self-calibration: Unveiling the microJy population of compact radio sources*, *A&A* **587** (2016) A85 [1601.04452].
- [107] T.W.B. Muxlow, A.P. Thomson, J.F. Radcliffe, N.H. Wrigley, R.J. Beswick, I. Smail et al., *The e-MERGE Survey (e-MERLIN Galaxy Evolution Survey): overview and survey description*, *MNRAS* **495** (2020) 1188 [2005.02407].
- [108] N. Jackson, *The Faintest Radio Source Yet: Expanded Very Large Array Observations of the Gravitational Lens SDSS J1004+4112*, *ApJL* **739** (2011) L28 [1105.2141].

- [109] S. Badole, N. Jackson, P. Hartley, D. Sluse, H. Stacey and H. Vives-Arias, *VLA and ALMA observations of the lensed radio-quiet quasar SDSS J0924+0219: a molecular structure in a 3 μ Jy radio source*, *MNRAS* **496** (2020) 138 [2005.13612].
- [110] A.T. Deller and E. Middelberg, *mJIVE-20: A Survey for Compact mJy Radio Objects with the Very Long Baseline Array*, *AJ* **147** (2014) 14 [1310.8191].
- [111] S.S. Shabala, A. Deller, S. Kaviraj, E. Middelberg, R.J. Turner, Y.S. Ting et al., *Delayed triggering of radio active galactic nuclei in gas-rich minor mergers in the local Universe*, *MNRAS* **464** (2017) 4706 [1608.04178].
- [112] C. Spingola, J.P. McKean, M. Lee, A. Deller and J. Moldon, *A novel search for gravitationally lensed radio sources in wide-field VLBI imaging from the mJIVE-20 survey*, *MNRAS* **483** (2019) 2125 [1811.09152].
- [113] S. Burke-Spolaor, “Followup of Dual AGN Candidates from the mJIVE-20 Survey.” Keck Observatory Archive LRIS, id.C271LA, Feb., 2015.
- [114] S. Rezaei, J.P. McKean, M. Biehl and A. Javadpour, *DECORAS: detection and characterization of radio-astronomical sources using deep learning*, *MNRAS* **510** (2022) 5891 [2109.09077].
- [115] M. Janssen, “rPICARD: Radboud Pipeline for the Calibration of high Angular Resolution Data.” Astrophysics Source Code Library, record ascl:1905.015, May, 2019.