

# Kerr stability in external regions

Dawei Shen\*

April 16, 2024

## Abstract

In 2003, Klainerman and Nicolò [14] proved the stability of Minkowski in the case of the exterior of an outgoing null cone. Relying on the method used in [14], Caciotta and Nicolò [2] proved the stability of Kerr spacetime in *external regions*, i.e. outside an outgoing null cone far away from the Kerr *event horizon*. In this paper, we give a new proof of [2]. Compared to [2], we reduce the number of derivatives needed in the proof, simplify the treatment of the last slice, and provide a unified treatment of the decay of initial data which contains in particular the initial data considered by Klainerman and Szeftel in [20]. Also, concerning the treatment of curvature estimates, similar to [25], we replace the vectorfield method used in [2, 14] by  $r^p$ -weighted estimates introduced by Dafermos and Rodnianski in [8].

## Keywords

Double null foliation, Geodesic foliation, Kerr stability, Linearization procedure, Peeling properties,  $r^p$ -weighted estimates

## Contents

|          |                                                      |           |
|----------|------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                  | <b>4</b>  |
| 1.1      | Einstein vacuum equations and the Cauchy problem     | 4         |
| 1.1.1    | Local existence theorem                              | 4         |
| 1.1.2    | Special solutions                                    | 4         |
| 1.2      | Stability of Kerr conjecture                         | 5         |
| 1.3      | Minkowski stability in [14]                          | 6         |
| 1.4      | Previous works of Kerr stability in external regions | 8         |
| 1.5      | Rough version of the main theorem                    | 8         |
| 1.6      | Structure of the paper                               | 9         |
| <b>2</b> | <b>Preliminaries</b>                                 | <b>10</b> |
| 2.1      | Geometric set-up                                     | 10        |
| 2.1.1    | Double null foliation                                | 10        |
| 2.1.2    | Geometry of Kerr spacetime                           | 12        |
| 2.1.3    | Decay of quantities in Kerr spacetime                | 14        |
| 2.1.4    | Initial layer region                                 | 15        |
| 2.1.5    | Bootstrap region                                     | 16        |
| 2.2      | Integral formulae                                    | 18        |
| 2.3      | Hodge systems                                        | 19        |

---

\*Email adress: dawei.shen@polytechnique.edu

Laboratoire Jacques-Louis Lions, Sorbonne Université, 75252 Paris, France

|          |                                                                                                                        |           |
|----------|------------------------------------------------------------------------------------------------------------------------|-----------|
| 2.4      | Elliptic estimates for Hodge systems                                                                                   | 20        |
| 2.5      | Null structure equations                                                                                               | 21        |
| 2.6      | Bianchi equations                                                                                                      | 22        |
| 2.7      | Transport equations for metric components                                                                              | 23        |
| 2.8      | The linearized equations                                                                                               | 24        |
| 2.8.1    | Linearization procedure                                                                                                | 24        |
| 2.8.2    | Schematic notation $\Gamma_g$ and $\Gamma_b$                                                                           | 26        |
| 2.8.3    | Schematic notation $\mathbf{O}_q^p$                                                                                    | 27        |
| 2.8.4    | The linearized null structure equations                                                                                | 29        |
| 2.8.5    | The linearized Bianchi equations                                                                                       | 31        |
| 2.8.6    | Linearized equations for the metric components                                                                         | 33        |
| 2.9      | Commutator identities                                                                                                  | 33        |
| 2.10     | Sobolev inequalities                                                                                                   | 34        |
| <b>3</b> | <b>Main theorem</b>                                                                                                    | <b>35</b> |
| 3.1      | Fundamental norms                                                                                                      | 35        |
| 3.1.1    | $\mathcal{R}$ norms (Curvature components)                                                                             | 35        |
| 3.1.2    | $\mathcal{O}$ norms (Ricci coefficients and metric components)                                                         | 36        |
| 3.1.3    | $\mathring{\mathcal{O}}(\Sigma_0)$ norms and $\mathfrak{R}_0$ norms (Initial data for the foliation of $\mathcal{K}$ ) | 37        |
| 3.1.4    | $\mathcal{O}^*(\underline{C}_*)$ norms (Ricci coefficients and metric components on $\underline{C}_*$ )                | 38        |
| 3.1.5    | $\mathcal{O}_{(0)}$ norms and $\mathfrak{R}_{(0)}$ norms (Initial data for the foliation of $\mathcal{K}_{(0)}$ )      | 38        |
| 3.1.6    | Osc norms (Oscillation control)                                                                                        | 39        |
| 3.2      | Smallness constants                                                                                                    | 39        |
| 3.3      | Main theorem                                                                                                           | 40        |
| 3.4      | Intermediate results                                                                                                   | 41        |
| 3.5      | Proof of the main theorem                                                                                              | 42        |
| 3.6      | Conclusions                                                                                                            | 43        |
| <b>4</b> | <b>Curvature estimates (Theorem M1)</b>                                                                                | <b>44</b> |
| 4.1      | General Bianchi pairs                                                                                                  | 45        |
| 4.2      | Estimates for the Bianchi pair $(\check{\alpha}, \check{\beta})$                                                       | 49        |
| 4.3      | Estimates for the Bianchi pair $(\check{\beta}, (\check{\rho}, \check{\sigma}))$                                       | 50        |
| 4.4      | Estimates for the Bianchi pair $((\check{\rho}, \check{\sigma}), \check{\beta})$                                       | 51        |
| 4.5      | Estimates for the Bianchi pair $(\check{\beta}, \check{\alpha})$                                                       | 53        |
| 4.6      | End of the proof of Theorem M1                                                                                         | 54        |
| <b>5</b> | <b>Ricci coefficients estimates (Theorem M3)</b>                                                                       | <b>54</b> |
| 5.1      | Preliminaries                                                                                                          | 55        |
| 5.1.1    | Null frame transformation                                                                                              | 55        |
| 5.1.2    | Difference of the pull backs of Kerr values                                                                            | 56        |
| 5.1.3    | Evolution lemma                                                                                                        | 57        |
| 5.1.4    | Bootstrap control in the initial data layer                                                                            | 57        |
| 5.2      | Estimates for $\mathcal{O}_{[1]}$ , $\mathcal{O}_\gamma$ and Osc                                                       | 58        |
| 5.2.1    | Estimate for $\mathring{\mathcal{O}}_{0,1}(\widetilde{\Omega\omega})$                                                  | 58        |
| 5.2.2    | Estimates for $\mathcal{O}_0(\widetilde{\Omega\omega})$ and $\mathcal{O}_0(\widetilde{\Omega\omega})$                  | 59        |
| 5.2.3    | Estimates for $\mathcal{O}_1(\widetilde{\Omega\omega})$ and $\mathcal{O}_1(\widetilde{\Omega\omega})$                  | 59        |
| 5.2.4    | Estimate for $\mathcal{O}_{0,1}(\widetilde{\Omega})$                                                                   | 62        |
| 5.2.5    | Estimate for $\text{Osc}(u)$                                                                                           | 62        |
| 5.2.6    | Estimate for $\mathcal{O}_{0,1}(\widetilde{\Omega \text{tr } \chi})$                                                   | 63        |

|          |                                                                                                                     |            |
|----------|---------------------------------------------------------------------------------------------------------------------|------------|
| 5.2.7    | Estimate for $\mathring{\mathcal{O}}_0(\widetilde{\Omega^{-1} \text{tr } \chi})$                                    | 64         |
| 5.2.8    | Estimate for $\mathcal{O}_0(\Omega \text{tr } \underline{\chi})$                                                    | 64         |
| 5.2.9    | Estimate for $\mathcal{O}_{0,1}(\underline{\eta})$                                                                  | 65         |
| 5.2.10   | Estimate for $\text{Osc}(f)$                                                                                        | 66         |
| 5.2.11   | Estimate for $\mathring{\mathcal{O}}_{0,1}(\underline{\eta})$                                                       | 68         |
| 5.2.12   | Estimate for $\mathcal{O}_{0,1}(\underline{\eta})$                                                                  | 69         |
| 5.2.13   | Estimate for $\mathcal{O}_{0,1,2}(\underline{\check{\Omega}})$                                                      | 70         |
| 5.2.14   | Estimate for $\mathring{\mathcal{O}}_1(\widetilde{\Omega^{-1} \text{tr } \chi})$                                    | 71         |
| 5.2.15   | Estimate for $\mathcal{O}_1(\Omega \text{tr } \underline{\chi})$                                                    | 71         |
| 5.2.16   | Estimates for $\mathcal{O}_{0,1}(\underline{\check{\chi}})$ and $\mathcal{O}_{0,1}(\underline{\check{\chi}})$       | 72         |
| 5.2.17   | Estimates for $\mathcal{O}_{0,1}(\underline{\check{b}})$                                                            | 72         |
| 5.2.18   | Estimate for $\text{Osc}(x)$                                                                                        | 73         |
| 5.2.19   | Estimates for $\mathcal{O}_{0,1}(\underline{\check{\gamma}})$ and $\mathcal{O}_{0,1}(\underline{\check{\epsilon}})$ | 74         |
| 5.2.20   | Estimate for $\text{Osc}(r)$                                                                                        | 74         |
| 5.3      | Estimate for $\mathcal{O}_\Gamma$                                                                                   | 75         |
| <b>6</b> | <b>Initialization and extension (Theorems M0, M2 and M4)</b>                                                        | <b>77</b>  |
| 6.1      | Control of the initial data layer (Theorem M0)                                                                      | 77         |
| 6.2      | Control of the last slice (Theorem M2)                                                                              | 78         |
| 6.2.1    | Geodesic type foliation on $\underline{C}_*$                                                                        | 79         |
| 6.2.2    | Estimate for $\mathcal{O}_1^*$                                                                                      | 80         |
| 6.2.3    | Estimate for $\mathcal{O}_\gamma^*$                                                                                 | 85         |
| 6.2.4    | Estimate for $\mathcal{O}_\Gamma^*$                                                                                 | 86         |
| 6.3      | Extension argument (Theorem M4)                                                                                     | 87         |
| <b>A</b> | <b>Proof of results in Section 2.8</b>                                                                              | <b>89</b>  |
| A.1      | Proof of Proposition 2.34                                                                                           | 89         |
| A.2      | Proof of Proposition 2.36                                                                                           | 93         |
| A.3      | Proof of Propositions 2.37 and 2.38                                                                                 | 94         |
| A.4      | Proof of Proposition 2.40                                                                                           | 105        |
| <b>B</b> | <b>Proof of the case <math>s \in (3, 4)</math></b>                                                                  | <b>107</b> |
| B.1      | Fundamental norms                                                                                                   | 107        |
| B.2      | Estimates for Ricci coefficients and curvature components                                                           | 108        |
| B.3      | Conclusion                                                                                                          | 108        |
| <b>C</b> | <b>Proof of the case <math>s &gt; 6</math></b>                                                                      | <b>108</b> |
| C.1      | Fundamental norms                                                                                                   | 108        |
| C.2      | Curvature estimates                                                                                                 | 109        |
| C.3      | Peeling decay for curvature components                                                                              | 117        |
|          | <b>Declarations</b>                                                                                                 | <b>117</b> |
|          | <b>References</b>                                                                                                   | <b>117</b> |

# 1 Introduction

## 1.1 Einstein vacuum equations and the Cauchy problem

A Lorentzian 4-manifold  $(\mathcal{M}, \mathbf{g})$  is called a vacuum spacetime if it solves the Einstein vacuum equations:

$$\mathbf{Ric}(\mathbf{g}) = 0 \quad \text{in } \mathcal{M}, \quad (1.1)$$

where  $\mathbf{Ric}$  denotes the Ricci tensor of the Lorentzian metric  $\mathbf{g}$ . The Einstein vacuum equations are invariant under diffeomorphisms, and therefore one considers equivalence classes of solutions. Expressed in general coordinates, (1.1) is a nonlinear geometric coupled system of second order partial differential equations for  $\mathbf{g}$ . In suitable coordinates, for example so-called wave coordinates, it can be shown that (1.1) is hyperbolic and hence admits an initial value formulation.

The corresponding initial data for the Einstein vacuum equations is given by specifying a triplet  $(\Sigma, g, k)$  where  $(\Sigma, g)$  is a Riemannian 3-dimension manifold and  $k$  is a symmetric 2-tensor on  $\Sigma$  satisfying the constraint equations:

$$\begin{aligned} R &= |k|^2 - (\text{tr } k)^2, \\ D^j k_{ij} &= D_i(\text{tr } k), \end{aligned} \quad (1.2)$$

where  $R$  denotes the scalar curvature of  $g$ ,  $D$  denotes the Levi-Civita connection of  $g$  and

$$|k|^2 := g^{ad}g^{bc}k_{ab}k_{cd}, \quad \text{tr } k := g^{ij}k_{ij}.$$

In the future development  $(\mathcal{M}, \mathbf{g})$  of such initial data  $(\Sigma, g, k)$ ,  $\Sigma \subset \mathcal{M}$  is a spacelike hypersurface with induced metric  $g$  and second fundamental form  $k$ .

### 1.1.1 Local existence theorem

The seminal well-posedness results for the Cauchy problem obtained in [3, 4] ensure that for any smooth Cauchy data, there exists a unique smooth maximal globally hyperbolic development  $(\mathcal{M}, \mathbf{g})$  solution of Einstein equations (1.1) such that  $\Sigma \subset \mathcal{M}$  and  $g, k$  are respectively the first and second fundamental forms of  $\Sigma$  in  $\mathcal{M}$ .

The simplest example of a vacuum spacetime is Minkowski spacetime:

$$\mathcal{M} = \mathbb{R}^4, \quad \mathbf{g} = -dt^2 + (dx^1)^2 + (dx^2)^2 + (dx^3)^2,$$

for which Cauchy data are given by

$$\Sigma = \mathbb{R}^3, \quad g = (dx^1)^2 + (dx^2)^2 + (dx^3)^2, \quad k = 0.$$

### 1.1.2 Special solutions

Minkowski spacetime is the simplest solution of (1.1). We recall two other relevant solutions of (1.1):

- Schwarzschild spacetime  
(1.1) admits a remarkable family of explicit, stationary, solutions given by the one parameter family of Schwarzschild solutions of mass  $M > 0$

$$\mathbf{g}_M = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\sigma_{\mathbb{S}^2}. \quad (1.3)$$




Figure 1: Complete Penrose diagram of Schwarzschild spacetime

Though the metric seems singular at  $r = 2M$ , it turns out that one can glue together two regions  $r > 2M$  and  $r < 2M$  of the Schwarzschild metric to obtain a metric which is smooth along  $\mathcal{H} = \{r = 2M\}$ , called the Schwarzschild *event horizon*. The portion of  $r < 2M$  to the future of the hypersurface  $t = 0$  is a *black hole* whose future boundary  $r = 0$  is singular. The similar region to the past of  $t = 0$  is called a *white hole*. The region  $r > 2M$ , called the *domain of outer communication* (DOC), is free of singularities. The Penrose diagram of Schwarzschild space is recalled in Figure 1.

- Kerr spacetime

The Schwarzschild family is included in a larger two parameter family of solutions  $K(a, M)$  discovered by Kerr in [13]. A given Kerr spacetime, with  $0 \leq |a| \leq M$  has a well defined domain of outer communication  $r > r_+ := M + (M^2 - a^2)^{\frac{1}{2}}$ . In Boyer-Lindquist coordinates, well adapted to  $r > r_+$ , the Kerr metric has the form:

$$g_{a,M} = -\frac{(\Delta - a^2 \sin^2 \theta)}{q^2} dt^2 - \frac{4aMr}{q^2} \sin^2 \theta dt d\varphi + \frac{q^2}{\Delta} dr^2 + q^2 d\theta^2 + \frac{\Sigma^2}{q^2} \sin^2 \theta d\varphi^2, \quad (1.4)$$

with

$$q^2 = r^2 + a^2 \cos 2\theta, \quad \Delta = r^2 + a^2 - 2Mr, \quad \Sigma^2 = (r^2 + a^2)^2 - a^2 (\sin \theta)^2 \Delta.$$

See Figure 2 for the Penrose diagram of Kerr spacetime.

## 1.2 Stability of Kerr conjecture

**Conjecture 1.1** (Stability of Kerr conjecture). *Vacuum initial data sets, sufficiently close to Kerr initial data, have a maximal development with complete future null infinity<sup>1</sup> and with domain of outer communication which approaches (globally) a nearby Kerr solution.*

The stability of Kerr has been proved in the case of small angular momentum,  $|a| \ll M$ , in the sequence of works by Klainerman-Szeftel [18], [19], [20], Giorgi-Klainerman-Szeftel [9]

<sup>1</sup>This means, roughly, that observers which are far away from the black hole may live forever.




Figure 2: Penrose diagram of Kerr for  $0 < |a| < M$ . The surface  $r = r_+$ , the larger root of  $\Delta = 0$ , is the event horizon of the black hole,  $r > r_+$  the domain of outer communication,  $\mathcal{J}^+$  is the future null infinity, corresponding to  $r = +\infty$ .

and the author [24], but the conjecture is open in general. For a complete introduction to this problem, we refer the reader to [21] and to the introduction in [20] and [9].

In order to attack the conjecture, it is convenient to divide the domain of outer communication to the future of an initial spacelike hypersurface  $\Sigma_0$  into three causal regions as in Figure 3:

- Region I lies to the future of  $\Sigma_0$  and to the past of an ingoing null cone  $\underline{C}_0$ . The stability of Kerr in spacetime region I is a direct consequence of the local existence results in [3, 4].
- Region II lies to the future of  $\Sigma_0$  and to the past of an outgoing null cone  $C_0$  sufficiently far away from the event horizon. See Section 1.4 for a discussion of Kerr stability in this region. The goal of this paper is to provide a new proof of the stability of Kerr in Region II.
- Region III lies to the future of the ingoing null cone  $\underline{C}_0$  and of the outgoing null cone  $C_0$ . The stability of Kerr in region III for small angular momentum  $|a| \ll M$  was proved recently in the sequence of works [18], [19], [20], [9] and [24].

As mentioned above, the goal of this paper is to provide a new proof of the stability of Kerr in Region II. In the next section, we review the corresponding problem for Minkowski.

### 1.3 Minkowski stability in [14]

In 1993, Christodoulou and Klainerman [5] proved the stability of Minkowski for the Einstein-vacuum equations, a milestone in the domain of mathematical general relativity. In 2003, Klainerman and Nicolò [14] gave a second proof of this result in the exterior of an outgoing cone. Moreover, Klainerman and Nicolò [15] showed that under stronger asymptotic decay and regularity properties than those used in [5, 14], asymptotically flat initial data sets lead to solutions of the Einstein vacuum equations which have strong peeling properties. For more references




Figure 3: The Penrose diagram of a perturbation of Kerr

on the stability of Minkowski for Einstein vacuum equations or Einstein equations coupled to matter, see for example the introduction of [25].

In this section, we present the main result of [14], which is the motivation for the analog results in Kerr recalled in Section 1.4 and for our main result stated in Section 1.5. First, we recall the definition of a *maximal hypersurface*, which plays an important role in the statements of the main theorems in [14].

**Definition 1.2.** An initial data  $(\Sigma, g, k)$  is posed on a maximal hypersurface if it satisfies

$$\text{tr } k = 0. \quad (1.5)$$

In this case, we say that  $(\Sigma, g, k)$  is a maximal initial data set, and the constraint equations (1.2) reduce to

$$R = |k|^2, \quad \text{div } k = 0, \quad \text{tr } k = 0. \quad (1.6)$$

We introduce the notion of *s-asymptotically flat initial data*.

**Definition 1.3.** Let  $s$  be a real number with  $s > 3$ . We say that a data set  $(\Sigma_0, g, k)$  is *s-asymptotically flat* if there exists a coordinates system  $(x^1, x^2, x^3)$  defined outside a sufficiently large compact set such that<sup>2</sup>

$$\begin{aligned} g &= \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\sigma_{\mathbb{S}^2} + o_4(r^{-\frac{s-1}{2}}), \\ k &= o_3(r^{-\frac{s+1}{2}}). \end{aligned} \quad (1.7)$$

We also introduce the following functional associated to any asymptotically flat initial data set:

$$J_0(\Sigma_0, g, k) := \sup_{\Sigma_0} \left( (d_0^2 + 1)^3 |\text{Ric}|^2 \right) + \int_{\Sigma_0} \sum_{l=0}^3 (d_0^2 + 1)^{l+1} |D^l k|^2 + \int_{\Sigma_0} \sum_{l=0}^1 (d_0^2 + 1)^{l+3} |D^l B|^2, \quad (1.8)$$

where  $d_0$  is the geodesic distance from a fixed point  $O \in \Sigma_0$ , and  $B_{ij} := (\text{curl } \widehat{R})_{ij}$  is the *Bach tensor*,  $\widehat{R}$  is the traceless part of  $\text{Ric}$ . Now, we can state the main theorem of [14].

<sup>2</sup>The notation  $f = o_l(r^{-m})$  means  $\partial^\alpha f = o(r^{-m-|\alpha|})$ ,  $|\alpha| \leq l$ .

**Theorem 1.4** (Klainerman-Nicolò [14]). *Consider an initial data set  $(\Sigma_0, g, k)$ , 4-asymptotically flat in the sense of Definition 1.3 and maximal, and assume that  $J_0(\Sigma_0, g, k)$  is bounded. Then, given a sufficiently large compact set  $K \subset \Sigma_0$  such that  $\Sigma_0 \setminus K$  is diffeomorphic to  $\mathbb{R}^3 \setminus \overline{B}_1$ , and assuming that  $J_0(\Sigma_0, g, k)$  is small enough, there exists a unique development  $(\mathbf{M}, \mathbf{g})$  with the following properties:*

1.  $(\mathbf{M}, \mathbf{g})$  can be foliated by a double null foliation  $\{C_u\}$  and  $\{\underline{C}_{\underline{u}}\}$  whose outgoing leaves  $C_u$  are complete.
2. We have detailed control of all the quantities associated with the double null foliation of the spacetime, see Theorem 3.7.1 of [14].

**Remark 1.5.** [15] and [25] extend Theorem 1.4 to the case of  $s$ -asymptotically flat initial data with  $s > 7$  and  $s > 3$  respectively. See also [10] for a proof in the case  $3 < s < 5$  using wave coordinates, constraint damping, spacetime compactification,  $b$ -Sobolev spaces and Nash-Moser iteration.

## 1.4 Previous works of Kerr stability in external regions

In 2009, Nicolò [23] proved that the asymptotic behaviour of the Riemann components is in agreement with the *Peeling theorem* if the corrections to the Kerr initial data decay sufficiently fast. In 2010, Caciotta and Nicolò [2] proved the nonlinear stability of the Kerr spacetime for any  $|a| \leq M$  and appropriate class of initial data close to Kerr, in an external region where  $r \geq R_0$  and  $\frac{M}{R_0} \ll 1$ . Their main results can be stated as follows.

**Theorem 1.6** (Caciotta-Nicolò [2]). *Assume that the initial data are given on  $\Sigma_0$  outside of a ball centered on the origin of radius  $R_0$ , denoted  $K := B_{R_0}$ . They are perturbations of the initial data of a Kerr spacetime with mass  $M$  and angular momentum  $a$  satisfying*

$$\frac{M}{R_0} \ll 1, \quad |a| \leq M.$$

*Let  $s > 7$ , and assume that the initial data satisfies for  $q \geq 4$ :*

$$\begin{aligned} g_{ij} &= (g_{Kerr})_{ij} + o_{q+1}(r^{-\frac{s-1}{2}}), \\ k_{ij} &= (k_{Kerr})_{ij} + o_q(r^{-\frac{s+1}{2}}). \end{aligned} \tag{1.9}$$

*Then, the initial data set  $(\Sigma_0, g, k)$  has a unique development  $(\mathcal{M}, \mathbf{g})$  defined outside  $I^+(K)$ , the domain of influence of  $K$ . Moreover,  $\mathcal{M}$  can be foliated by a canonical double null foliation  $(u, \underline{u})$  whose outgoing leaves  $C_u$  are complete and suitable norms of Ricci coefficients and curvature components are sufficiently small.*

The goal of this paper is to provide a new proof of Theorem 1.6.

## 1.5 Rough version of the main theorem

In this section, we state a simple version of our main theorem. For the precise statement, see Theorem 3.2.

**Theorem 1.7** (Main Theorem, first version). *Let  $K = B_{R_0} \subset \Sigma_0$  a ball centered at the origin of radius  $R_0$  and let  $\mathbf{g}_{a,M}$  a Kerr metric with mass  $M > 0$  and angular momentum  $a$  satisfying*

$$\frac{M}{R_0} \ll 1, \quad |a| \leq M.$$

Let  $s > 3$  and let an initial data set  $(\Sigma_0, g, k)$  satisfying (1.9) with  $q = 2$ . Let  $\mathcal{K}_{(0)}$ <sup>3</sup> an initial layer region near  $\Sigma_0 \setminus K$  and assume that the initial data in  $\mathcal{K}_{(0)}$  are perturbations from the initial data of  $\mathbf{g}_{a,M}$  of size  $\epsilon_0 \ll \frac{M}{R_0}$ .<sup>4</sup>

Then, the initial data set  $(\Sigma_0, g, k)$  has a unique development  $(\mathcal{M}, \mathbf{g})$  in its future domain of dependence with the following properties:

- $\mathcal{M}$  can be foliated by a double null foliation  $(u, \underline{u})$  whose outgoing leaves  $C_u$  are complete for all  $u \leq u_0$ ;
- we have detailed control of all the quantities associated with the double null foliation of the spacetime, see Theorem 3.2.

**Remark 1.8.** In the particular case of  $s$ -asymptotically flat initial data with  $s > 7$ , we reobtain the results of [2].

**Remark 1.9.** Let  $C_{u_0}$  be the outgoing null cone located near  $\partial K$  and choose  $s = 4 + 2\delta_B$  with  $\delta_B > 0$ . Then, the estimates of Theorem 1.7 on  $C_{u_0}$  are consistent with the initial data considered in [17] and [20].

**Remark 1.10.** Theorem 1.7 is the analog for the Kerr spacetime of the result for Minkowski in [25].

The proof of Theorem 1.7 has the same structure as in [2, 14] and is closely related to the one in [25]. Below, we compare the proof of this paper and that of [2]:

1. To estimate the norms of curvature components, [2] uses the vectorfield method introduced in [5, 14]. Here, we estimate the curvature norms by the  $r^p$ -weighted estimate method introduced in [8]. This allows for a simple treatment of the curvature estimates.
2. [2] uses fourth order derivatives of Ricci coefficients and third order derivatives of curvature components. Thanks to the use of  $r^p$ -weighted estimates, we only need first order derivatives of both Ricci coefficients and curvature components.
3. In order to control one more derivative for Ricci coefficients compared to curvature, [2] relies on the canonical foliation on the last slice  $\underline{C}_*$ . Since we control the same number of derivatives for curvature and Ricci coefficients, instead of introducing the canonical foliation, we use a geodesic type foliation to simplify the estimates on the last slice  $\underline{C}_*$ .

## 1.6 Structure of the paper

- In Section 2, we recall the fundamental notions and basic equations used in this paper. We also introduce the linearization procedure and the linearized equations.
- In Section 3, we state the precise version of the main theorem. We then state intermediate results, and rely on them to prove the main theorem. The rest of the paper focuses on the proof of these intermediary results. In Sections 4–6, we consider the case  $s \in [4, 6]$  and deal with the other cases in Appendices B and C.
- In Section 4, we apply  $r^p$ -weighted estimates to the linearized Bianchi equations to control curvature.

---

<sup>3</sup> $\mathcal{K}_{(0)}$  is defined in Section 2.1.4.

<sup>4</sup>See (3.6) for the explicit initial data assumptions and see Section 3.1.5 for the definitions of the norms on the initial data.

- In Section 5, we estimate the Ricci coefficients using the linearized null structure equations.
- In Section 6, we estimate the curvature in the initial layer region  $\mathcal{K}_{(0)}$ . Then, we estimate Ricci coefficients on the last slice  $\underline{C}_*$ . We also show how to extend the spacetime in the context of a bootstrap argument.
- In Appendix A, we provide the derivation of the linearized equations stated in Section 2.8.
- In Appendix B, we prove Theorem 1.7 in the case  $s \in (3, 4)$ .
- In Appendix C, we prove Theorem 1.7 in the case  $s > 6$  by applying  $r^p$ -weighted estimates to the linearized Teukolsky equation.

## 2 Preliminaries

### 2.1 Geometric set-up

#### 2.1.1 Double null foliation

We first introduce the geometric setup. The material in this section is classical, see for example Chapter 3 in [14].

We denote  $(\mathcal{M}, \mathbf{g})$  a spacetime  $\mathcal{M}$  with the Lorentzian metric  $\mathbf{g}$  and  $\mathbf{D}$  its Levi-Civita connection. Let  $u$  and  $\underline{u}$  be two optical functions on  $\mathcal{M}$ , that is<sup>5</sup>

$$\mathbf{g}(\text{grad } u, \text{grad } u) = \mathbf{g}(\text{grad } \underline{u}, \text{grad } \underline{u}) = 0,$$

The spacetime  $\mathcal{M}$  is foliated by the level sets of  $u$  and  $\underline{u}$  respectively, and the functions  $u, \underline{u}$  are required to increase towards the future. We use  $C_u$  to denote the outgoing null cones which are the level sets of  $u$  and use  $\underline{C}_{\underline{u}}$  to denote the ingoing null cones which are the level sets of  $\underline{u}$ . We denote

$$S(u, \underline{u}) := C_u \cap \underline{C}_{\underline{u}}, \tag{2.1}$$

which is a spacelike 2-sphere. We define the area radius of  $S(u, \underline{u})$ :

$$r(u, \underline{u}) := \sqrt{\frac{|S(u, \underline{u})|}{4\pi}}. \tag{2.2}$$

We introduce the vector fields  $L$  and  $\underline{L}$  by

$$L := -\text{grad } u, \quad \underline{L} := -\text{grad } \underline{u}.$$

We define a positive scalar function  $\Omega$  by:

$$\mathbf{g}(L, \underline{L}) = -\frac{1}{2\Omega^2},$$

called the lapse function. We then define the normalized null pair  $(e_3, e_4)$  by

$$e_3 = 2\Omega \underline{L}, \quad e_4 = 2\Omega L.$$

---

<sup>5</sup>Here,  $\text{grad}$  denotes the spacetime gradient.

We also define the following vector fields:

$$\underline{N} = \Omega e_3, \quad N = \Omega e_4.$$

The flows generated by  $\underline{N}$  and  $N$  preserve the double null foliation. On a given two sphere  $S(u, \underline{u})$ , we choose a local frame  $(e_1, e_2)$ , and we call  $(e_1, e_2, e_3, e_4)$  a null frame. As a convention, throughout the paper, we use capital Latin letters  $A, B, C, \dots$  to denote an index from 1 to 2 and Greek letters  $\alpha, \beta, \gamma, \dots$  to denote an index from 1 to 4, e.g.  $e_A$  denotes either  $e_1$  or  $e_2$ .

The spacetime metric  $\mathbf{g}$  induces a Riemannian metric  $\gamma$  on  $S(u, \underline{u})$ . Using  $(u, \underline{u})$ , we can introduce a local coordinates system  $(u, \underline{u}, x^A)$  on  $\mathcal{M}$  with  $e_4(x^A) = 0$ . The metric  $\mathbf{g}$  takes the form:

$$\mathbf{g} = -2\Omega^2(d\underline{u} \otimes d\underline{u} + du \otimes d\underline{u}) + \gamma_{AB}(dx^A - \underline{b}^A du) \otimes (dx^B - \underline{b}^B du), \quad (2.3)$$

and we have  $\underline{N} = \partial_u + \underline{b}$  and  $N = \partial_{\underline{u}}$  where

$$\underline{b} := \underline{b}^A \partial_{x^A}.$$

We recall the null decomposition of the Ricci coefficients and curvature components of the null frame  $(e_1, e_2, e_3, e_4)$ . We have:

$$\begin{aligned} \underline{\chi}_{AB} &= \mathbf{g}(\mathbf{D}_A e_3, e_B), & \chi_{AB} &= \mathbf{g}(\mathbf{D}_A e_4, e_B), \\ \underline{\xi}_A &= \frac{1}{2} \mathbf{g}(\mathbf{D}_3 e_3, e_A), & \xi_A &= \frac{1}{2} \mathbf{g}(\mathbf{D}_4 e_4, e_A), \\ \underline{\omega} &= \frac{1}{4} \mathbf{g}(\mathbf{D}_3 e_3, e_4), & \omega &= \frac{1}{4} \mathbf{g}(\mathbf{D}_4 e_4, e_3), \\ \underline{\eta}_A &= \frac{1}{2} \mathbf{g}(\mathbf{D}_4 e_3, e_A), & \eta_A &= \frac{1}{2} \mathbf{g}(\mathbf{D}_3 e_4, e_A), \\ \zeta_A &= \frac{1}{2} \mathbf{g}(\mathbf{D}_{e_A} e_4, e_3), \end{aligned} \quad (2.4)$$

and

$$\begin{aligned} \alpha_{AB} &= \mathbf{R}(e_A, e_4, e_B, e_4), & \underline{\alpha}_{AB} &= \mathbf{R}(e_A, e_3, e_B, e_3), \\ \beta_A &= \frac{1}{2} \mathbf{R}(e_A, e_4, e_3, e_4), & \underline{\beta}_A &= \frac{1}{2} \mathbf{R}(e_A, e_3, e_3, e_4), \\ \rho &= \frac{1}{4} \mathbf{R}(e_3, e_4, e_3, e_4), & \sigma &= \frac{1}{4} {}^* \mathbf{R}(e_3, e_4, e_3, e_4), \end{aligned} \quad (2.5)$$

where  $\mathbf{R}$  denotes the curvature tensor of  $\mathbf{g}$  and  ${}^* \mathbf{R}$  denotes the Hodge dual of  $\mathbf{R}$ . The null second fundamental forms  $\chi, \underline{\chi}$  are further decomposed in their traces  $\text{tr } \chi$  and  $\text{tr } \underline{\chi}$ , and traceless parts  $\hat{\chi}$  and  $\hat{\underline{\chi}}$ :

$$\begin{aligned} \text{tr } \chi &:= \delta^{AB} \chi_{AB}, & \hat{\chi}_{AB} &:= \chi_{AB} - \frac{1}{2} \delta_{AB} \text{tr } \chi, \\ \text{tr } \underline{\chi} &:= \delta^{AB} \underline{\chi}_{AB}, & \hat{\underline{\chi}}_{AB} &:= \underline{\chi}_{AB} - \frac{1}{2} \delta_{AB} \text{tr } \underline{\chi}. \end{aligned}$$

We define the horizontal covariant operator  $\nabla$  as follows:

$$\nabla_X Y := \mathbf{D}_X Y - \frac{1}{2} \underline{\chi}(X, Y) e_4 - \frac{1}{2} \chi(X, Y) e_3.$$

We also define  $\nabla_4 X$  and  $\nabla_3 X$  to be the horizontal projections:

$$\begin{aligned}\nabla_4 X &:= \mathbf{D}_4 X - \frac{1}{2} \mathbf{g}(X, \mathbf{D}_4 e_3) e_4 - \frac{1}{2} \mathbf{g}(X, \mathbf{D}_4 e_4) e_3, \\ \nabla_3 X &:= \mathbf{D}_3 X - \frac{1}{2} \mathbf{g}(X, \mathbf{D}_3 e_3) e_3 - \frac{1}{2} \mathbf{g}(X, \mathbf{D}_3 e_4) e_4.\end{aligned}$$

A tensor field  $\psi$  defined on  $\mathcal{M}$  is called tangent to  $S$  if it is a priori defined on the spacetime  $\mathcal{M}$  and all the possible contractions of  $\psi$  with either  $e_3$  or  $e_4$  are zero. We use  $\nabla_3 \psi$  and  $\nabla_4 \psi$  to denote the projection to  $S(u, \underline{u})$  of usual derivatives  $\mathbf{D}_3 \psi$  and  $\mathbf{D}_4 \psi$ . As a direct consequence of (2.4), we have the Ricci formulae:

$$\begin{aligned}\mathbf{D}_A e_B &= \nabla_A e_B + \frac{1}{2} \chi_{AB} e_3 + \frac{1}{2} \underline{\chi}_{AB} e_4, \\ \mathbf{D}_A e_3 &= \underline{\chi}_{AB} e_B + \zeta_A e_3, \\ \mathbf{D}_A e_4 &= \chi_{AB} e_B - \zeta_A e_4, \\ \mathbf{D}_3 e_A &= \nabla_3 e_A + \eta_A e_3 + \underline{\xi}_A e_4, \\ \mathbf{D}_4 e_A &= \nabla_4 e_A + \underline{\eta}_A e_4 + \xi_A e_4, \\ \mathbf{D}_3 e_3 &= -2\underline{\omega} e_3 + 2\underline{\xi}_B e_B, \\ \mathbf{D}_3 e_4 &= 2\underline{\omega} e_4 + 2\underline{\eta}_B e_B, \\ \mathbf{D}_4 e_4 &= -2\omega e_4 + 2\xi_B e_B, \\ \mathbf{D}_4 e_3 &= 2\omega e_3 + 2\underline{\eta}_B e_B.\end{aligned}\tag{2.6}$$

In addition to

$$\xi = \underline{\xi} = 0,\tag{2.7}$$

the following identities hold for a double null foliation:

$$\begin{aligned}\nabla \log \Omega &= \frac{1}{2}(\eta + \underline{\eta}), & \omega &= -\frac{1}{2} \mathbf{D}_4(\log \Omega), & \underline{\omega} &= -\frac{1}{2} \mathbf{D}_3(\log \Omega), \\ \eta &= \zeta + \nabla \log \Omega, & \underline{\eta} &= -\zeta + \nabla \log \Omega,\end{aligned}\tag{2.8}$$

see for example (6) in [16].

### 2.1.2 Geometry of Kerr spacetime

In this section, we recall the basic notions of the geometry of Kerr spacetime used in the sequel. See Section 4 in [12] and Appendix A in [7] for more details.

Let  $\mathcal{M}_{Kerr}$  be a 4-dimensional manifold diffeomorphic to  $\mathbb{R}^2 \times \mathbb{S}^2$  which describe the Kerr black hole. The Kerr metric  $\mathbf{g}_{a,M}$  with mass  $M$  and angular momentum  $a$  is a smooth metric on  $\mathcal{M}_{Kerr}$ , which takes the following form in the Boyer-Lindquist local coordinates  $(t, r, \theta, \phi)$ :

$$\begin{aligned}\mathbf{g}_{a,M} &= -\left(1 - \frac{2Mr}{\Sigma}\right) dt \otimes dt + \frac{\Sigma}{\Delta} dr \otimes dr + \Sigma d\theta \otimes d\theta, \\ &\quad + R^2 \sin^2 \theta d\phi \otimes d\phi - \frac{2Mar \sin^2 \theta}{\Sigma} (d\phi \otimes dt + dt \otimes d\phi),\end{aligned}\tag{2.9}$$

where

$$\Sigma := r^2 + a^2 \cos^2 \theta, \quad R^2 := r^2 + a^2 + \frac{2Ma^2 r \sin^2 \theta}{\Sigma}, \quad \Delta := r^2 + a^2 - 2Mr.$$

We denote

$$r_+ := M + (M^2 - a^2)^{\frac{1}{2}},$$

and recall that  $r = r_+$  defines the event horizons of Kerr black hole. Note that (2.9) is valid in  $r > r_+$ .

**Remark 2.1.** Note that  $(\theta, \phi)$  are spherical coordinates on  $S(t, r)$  and have a coordinate singularity at  $\theta = 0, \pi$ . We thus also introduce

$$x^1 := \sin \theta \cos \phi, \quad x^2 := \sin \theta \sin \phi.$$

Then,  $(x^1, x^2)$  are regular coordinates when  $0 \leq \theta \leq \frac{\pi}{3}$  and  $\frac{2\pi}{3} \leq \theta \leq \pi$ . Hence,  $S(t, r)$  can be covered by three regular coordinates charts. In the sequel, we focus on the spherical coordinates chart  $(\theta, \phi)$ .

**Proposition 2.2.** We can define two new functions  $r_* = r(r, \theta)$  and  $\theta_* = \theta_*(r, \theta)$  such that the Kerr metric takes the following form in  $r > r_+$ :

$$\begin{aligned} \mathbf{g}_{a,M} = & \frac{\Delta}{R^2} (-dt \otimes dt + dr_* \otimes dr_*) + \frac{\ell^2}{R^2} d\theta_* \otimes d\theta_* \\ & + R^2 \sin^2 \theta \left( d\phi - \frac{2Mar}{\Sigma R^2} dt \right) \otimes \left( d\phi - \frac{2Mar}{\Sigma R^2} dt \right), \end{aligned} \quad (2.10)$$

where

$$\begin{aligned} \ell &:= -Ma \sqrt{\sin^2 \theta_* - \sin^2 \theta} \sqrt{(r^2 + a^2)^2 - a^2 \sin^2 \theta_* \Delta}, \\ \Delta &:= r^2 + a^2 - 2Mr, \\ R^2 &:= r^2 + a^2 + \frac{2Ma^2 r \sin^2 \theta}{\Sigma}, \\ \Sigma &:= r^2 + a^2 \cos^2 \theta. \end{aligned}$$

*Proof.* See Appendices A.1–A.3 in [7]. □

We can also write the Kerr metric in double null foliation. We define

$$\underline{u}_{Kerr} := t + r_*, \quad u_{Kerr} := t - r_*, \quad \phi_* := \phi - h(r_*, \theta_*),$$

where

$$\frac{dh}{dr_*}(r_*, \theta_*) = \frac{2Mar}{\Sigma R^2}.$$

We easily obtain in  $r > r_+$ <sup>6</sup>

$$\begin{aligned} \mathbf{g}_{a,M} = & -2\Omega_{Kerr}^2 (d\underline{u}_{Kerr} \otimes du_{Kerr} + du_{Kerr} \otimes d\underline{u}_{Kerr}) \\ & + (\gamma_{Kerr})_{AB} (dx_{Kerr}^A - \underline{b}_{Kerr}^A du_{Kerr}) \otimes (dx_{Kerr}^B - \underline{b}_{Kerr}^B du_{Kerr}), \end{aligned}$$

where  $(x_{Kerr}^1, x_{Kerr}^2) = (\theta_*, \phi_*)$  or  $(x_{Kerr}^1, x_{Kerr}^2) = (\sin \theta_* \cos \phi_*, \sin \theta_* \sin \phi_*)$ .

We also denote

$$L_{Kerr} := -\text{grad } u_{Kerr}, \quad \underline{L}_{Kerr} := -\text{grad } \underline{u}_{Kerr}, \quad \mathbf{g}(L_{Kerr}, \underline{L}_{Kerr}) = -\frac{1}{2\Omega_{Kerr}^2},$$

---

<sup>6</sup>See (A.42) in [7] and notice the exchange of role between  $\underline{u}$  and  $u$ .

and

$$(e_4)_{Kerr} := 2\Omega_{Kerr} L_{Kerr}, \quad (e_3)_{Kerr} := 2\Omega_{Kerr} \underline{L}_{Kerr}.$$

The leaves of  $\mathcal{M}_{Kerr}$  are defined by

$$S_{Kerr}(u_{Kerr}, \underline{u}_{Kerr}) := (C_{Kerr})_{u_{Kerr}} \cap (\underline{C}_{Kerr})_{\underline{u}_{Kerr}},$$

where  $(C_{Kerr})_{u_{Kerr}}$  and  $(\underline{C}_{Kerr})_{\underline{u}_{Kerr}}$  respectively denote the outgoing and ingoing null cones of the  $(u_{Kerr}, \underline{u}_{Kerr})$ -foliation in  $\mathcal{M}_{Kerr}$ . Let also  $r_{Kerr}$  denote the area radius of the  $S(u_{Kerr}, \underline{u}_{Kerr})$ , i.e.

$$r_{Kerr} := \sqrt{\frac{|S(u_{Kerr}, \underline{u}_{Kerr})|}{4\pi}}.$$

Note that  $r_{Kerr}$  is a function of  $r_*(r, \theta)$ .

We define

$$\Sigma_{Kerr} := \{p \in \mathcal{M}_{Kerr} / u_{Kerr} + \underline{u}_{Kerr} = 0\},$$

and

$$w_{Kerr} := \frac{u_{Kerr} - \underline{u}_{Kerr}}{2}.$$

Let  $\delta_0 > 0$  be a fixed small constant. The initial layer region of  $\mathcal{M}_{Kerr}$  is defined by

$$(\mathcal{K}_{(0)})_{Kerr} := \{p \in \mathcal{M}_{Kerr} / 0 \leq u_{Kerr} + \underline{u}_{Kerr} \leq 2\delta_0\} \setminus I^+(K_{Kerr}), \quad (2.11)$$

where

$$K_{Kerr} := \{p \in \Sigma_{Kerr} / w_{Kerr}(p) \leq R_0\}, \quad (2.12)$$

$I^+(K_{Kerr})$  denotes the future domain of influence of  $K_{Kerr}$  and  $R_0 > 0$  is a constant.

### 2.1.3 Decay of quantities in Kerr spacetime

In this section, we provide the decay of quantities associated to the double null foliation in  $\mathcal{M}_{Kerr}$  induced by  $(u_{Kerr}, \underline{u}_{Kerr})$  introduced in Section 2.1.2.

**Definition 2.3.** For any  $S_{Kerr}$ -tangent tensor field  $h_{Kerr}$  defined in  $\mathcal{M}_{Kerr}$ , we denote

$$h_{Kerr} = O_q^p, \quad (2.13)$$

if for any integer  $l \geq 0$

$$|\nabla^{\leq l} h_{Kerr}| \lesssim \frac{M^p}{r_{Kerr}^{q+l}},$$

where

$$\nabla := \left\{ \nabla_{\partial_{\underline{u}_{Kerr}}}, \nabla_{\partial_{u_{Kerr}}}, r_{Kerr}^{-1} \nabla_{\partial_{x_{Kerr}^A}} \right\},$$

where the coordinates  $(u_{Kerr}, \underline{u}_{Kerr}, x_{Kerr}^A)$  and the area radius  $r_{Kerr}$  of  $S(u_{Kerr}, \underline{u}_{Kerr})$  have been introduced in Section 2.1.2.

**Proposition 2.4.** *The Ricci coefficients of the  $(u_{Kerr}, \underline{u}_{Kerr})$  double null foliation satisfy in  $\mathcal{M}_{Kerr}$ :*

$$\begin{aligned}
(\mathrm{tr} \chi)_{Kerr} - \frac{2}{r_{Kerr}} &= O_2^1, & (\mathrm{tr} \underline{\chi})_{Kerr} + \frac{2}{r_{Kerr}} &= O_2^1, \\
\omega_{Kerr} &= O_2^1, & \underline{\omega}_{Kerr} &= O_2^1, \\
\widehat{\chi}_{Kerr} &= O_3^2, & \widehat{\underline{\chi}}_{Kerr} &= O_3^2, \\
(\nabla \mathrm{tr} \chi)_{Kerr} &= O_4^2, & (\nabla \mathrm{tr} \underline{\chi})_{Kerr} &= O_4^2, \\
\eta_{Kerr} &= O_3^2, & \underline{\eta}_{Kerr} &= O_3^2, \\
(\nabla \omega)_{Kerr} &= O_4^2, & (\nabla \underline{\omega})_{Kerr} &= O_4^2, \\
\underline{b}_{Kerr} &= O_2^2, & \Omega_{Kerr} - \frac{1}{2} &= O_1^1.
\end{aligned}$$

*Proof.* See for example Section 2.2.3 in [2] and Appendix A.4 in [7].  $\square$

**Proposition 2.5.** *The curvature components of the  $(u_{Kerr}, \underline{u}_{Kerr})$  double null foliation satisfy in  $\mathcal{M}_{Kerr}$ :*

$$\begin{aligned}
\alpha_{Kerr} &= O_5^3, & \underline{\alpha}_{Kerr} &= O_5^3, \\
\beta_{Kerr} &= O_4^2, & \underline{\beta}_{Kerr} &= O_4^2, \\
\rho_{Kerr} &= O_3^1, & \sigma_{Kerr} &= O_4^2.
\end{aligned}$$

*Proof.* See Section 3.3.5 of [2].  $\square$

#### 2.1.4 Initial layer region

Let  $\Sigma_0$  be a spacelike hypersurface, called the *initial hypersurface*. Let  $K \subset \Sigma_0$  be a compact subset such that  $\Sigma_0 \setminus K$  is diffeomorphic to  $\mathbb{R}^3 \setminus \overline{B_1}$  where  $\overline{B_1}$  is the unit closed ball in  $\mathbb{R}^3$ . We fix a radial foliation on the initial hypersurface  $\Sigma_0 \setminus K$  by the level sets of a scalar function  $w$ , its leaves are denoted by

$$S_{(0)}(w_1) = \{p \in \Sigma_0 / w(p) = w_1\},$$

where  $w_1 \geq 0$ . We assume that

$$\partial K = \{p \in \Sigma_0 / w(p) = R_0\}, \quad K = \{p \in \Sigma_0 / w(p) \leq R_0\},$$

where  $R_0 > 0$  is the constant appearing in (2.12).

We now construct an *initial layer region*  $\mathcal{K}_{(0)}$  near  $\Sigma_0 \setminus K$  endowed with a double null foliation  $(u_{(0)}, \underline{u})$  as in Section 2.1.1 satisfying

$$u_{(0)}(p) = -w(p), \quad \underline{u}(p) = w(p), \quad \text{on } \Sigma_0 \setminus K.$$

We then define

$$\mathcal{K}_{(0)} := \{p / 0 \leq u_{(0)}(p) + \underline{u}(p) \leq 2\delta_0\}, \quad (2.14)$$

where  $\delta_0 > 0$  is the small constant appearing in (2.11). The double null foliation  $(u_{(0)}, \underline{u})$  is called the *initial layer foliation*. Its leaves are denoted by

$$S_{(0)}(u_{(0)}, \underline{u}) := (C_{(0)})_{u_{(0)}} \cap \underline{C}_{\underline{u}}, \quad (2.15)$$

where  $(C_{(0)})_{u_{(0)}}$  is the outgoing null cone emanating from  $S_{(0)}(u_{(0)}, -u_{(0)})$  and  $\underline{C}_{\underline{u}}$  is the ingoing null cone emanating from  $S_{(0)}(-\underline{u}, \underline{u})$ . By construction, we have

$$\Sigma_0 \setminus K = \{p \in \mathcal{K}_{(0)} / u_{(0)}(p) + \underline{u}(p) = 0\}, \quad S_{(0)}(\underline{u}) = S_{(0)}(-\underline{u}, \underline{u}). \quad (2.16)$$

We extend the domain of the function  $w$  to  $\mathcal{K}_{(0)}$  as follows:

$$w(p) := \frac{\underline{u}(p) - u_{(0)}(p)}{2}, \quad \forall p \in \mathcal{K}_{(0)}.$$

Also, as in Section 2.1.1,  $\mathcal{K}_{(0)}$  is covered by coordinates systems  $(u_{(0)}, \underline{u}, x_{(0)}^A)$  with  $L_{(0)}(x_{(0)}^A) = 0$  and the metric  $\mathbf{g}$  in  $\mathcal{K}_{(0)}$  has the form:

$$\mathbf{g} = -2\Omega_{(0)}^2 (d\underline{u} \otimes du_{(0)} + du_{(0)} \otimes d\underline{u}) + (\gamma_{(0)})_{AB} (dx_{(0)}^A - \underline{b}_{(0)}^A du_{(0)}) \otimes (dx_{(0)}^B - \underline{b}_{(0)}^B du_{(0)}). \quad (2.17)$$

Let  $\Phi_{(0)}$  be the map defined on  $\Sigma_0$  such that

$$\begin{aligned} \Phi_{(0)} : \Sigma_0 &\longrightarrow \Sigma_{Kerr}, \\ p &\longrightarrow \Phi_{(0)}(p) / u_{Kerr}(\Phi_{(0)}(p)) = u(p), \quad \underline{u}_{Kerr}(\Phi_{(0)}(p)) = \underline{u}(p), \quad x_{Kerr}^A(\Phi_{(0)}(p)) = x_{(0)}^A(p). \end{aligned}$$

The map  $\Phi_{(0)}$  being defined on  $\Sigma_0$ , we now extend it to  $\mathcal{K}_{(0)}$ . We denote  $L_{(0)}$  and  $L_{Kerr}$  respectively the generator of  $(C_{(0)})_{u_{(0)}}$  and  $C_{Kerr}(u_{Kerr})$ . For any  $p \in \mathcal{K}_{(0)}$ , we denote  $\phi_t^{L_{(0)}}(p)$  the flow of  $L_{(0)}$  emanating from  $p$  to the past. Then, we denote

$$p_{(0)} := \phi_t^{L_{(0)}}(p) \cap \Sigma_0, \quad q_{(0)} := \Phi_{(0)}(p_{(0)}).$$

Next, we denote  $\phi_t^{L_{Kerr}}(q_{(0)})$  the flow of  $L_{Kerr}$  emanating from  $q_{(0)}$  to the future. We define

$$q := \phi_t^{L_{Kerr}}(q_{(0)}) \cap S_{Kerr}(u_{Kerr} = u_{(0)}(p), \underline{u}_{Kerr} = \underline{u}(p)).$$

Finally, we define

$$\Phi_{(0)}(p) = q. \quad (2.18)$$

By construction, we have for any  $p \in \mathcal{K}_{(0)}$ <sup>7</sup>

$$\Phi_{(0)}(p) \in \mathcal{K}_{Kerr}, \quad u_{Kerr}(\Phi_{(0)}(p)) = u(p), \quad \underline{u}_{Kerr}(\Phi_{(0)}(p)) = \underline{u}(p), \quad x_{Kerr}^A(\Phi_{(0)}(p)) = x_{(0)}^A(p).$$

which implies that the map  $\Phi_{(0)}$  is equivalent to an identification of  $\mathcal{K}_{(0)}$  and  $\mathcal{K}_{Kerr}$  via the values of the coordinates functions  $(u_{(0)}, \underline{u}, x_{(0)}^1, x_{(0)}^2)$  and  $(u_{Kerr}, \underline{u}_{Kerr}, x_{Kerr}^1, x_{Kerr}^2)$ .<sup>8</sup>

### 2.1.5 Bootstrap region

For every value  $\underline{u}_* > R_0$ ,

$$S_* := S_{(0)}(-\underline{u}_*, \underline{u}_*)$$

denotes a leaf of the  $(u_{(0)}, \underline{u})$ -foliation on  $\Sigma_0 \setminus K$ . Emanating from this sphere  $S_*$ , we construct an ingoing cone  $\underline{C}_*$  called *the last slice*. For an optical function  $u$  defined on  $\underline{C}_*$  such that  $u = -\underline{u}_*$  on  $S_*$ , the level sets of  $u$  induce a foliation on  $\underline{C}_*$ , and its leaves are denoted by  $S(u, \underline{u}_*)$ . We will take a particular choice of the coordinates systems  $(u, x^A)$  on  $\underline{C}_*$  in the

<sup>7</sup>Note that the identity  $x_{Kerr}^A(\Phi_{(0)}(p)) = x_{(0)}^A(p)$  follows from the fact that it holds on  $\Sigma_0$  and from the identities  $L_{(0)}(x_{(0)}^A) = 0$  and  $L_{Kerr}(x_{Kerr}^A) = 0$ .

<sup>8</sup>See Section 4.3 in [7] for more explanations concerning this identification.




Figure 4: Bootstrap region  $\mathcal{K}$

context of a geodesic type foliation, see Definition 6.1, with  $x^A = x^A_{(0)}$  on  $S_*$ .

We define on  $\underline{C}_*$  a map  $\Phi^9$  by

$$\begin{aligned} \Phi : \underline{C}_* &\longrightarrow (\underline{C}_*)_{Kerr}, \\ p &\longrightarrow \Phi(p) / u_{Kerr}(\Phi(p)) = u(p), \quad \underline{u}_{Kerr}(\Phi(p)) = \underline{u}(p), \quad x^A_{Kerr}(\Phi(p)) = x^A(p), \end{aligned}$$

where

$$(\underline{C}_*)_{Kerr} = \{p \in \mathcal{M}_{Kerr} / \underline{u}_{Kerr} = \underline{u}_*\}.$$

For every  $u \geq -\underline{u}_*$ , we construct an outgoing null cone emanating from  $S(u, \underline{u}_*)$  on  $\underline{C}_*$  to the past, denoted  $C_u$ . Then,  $C_u \cap \Sigma_0$  is a sphere on  $\Sigma_0$ , which in general does not coincide with the leaves of the  $(u_{(0)}, \underline{u})$ -foliation on  $\Sigma_0$ . We also define  $\underline{C}_{\underline{u}}$  the ingoing null cone emanating from  $S_{(0)}(-\underline{u}, \underline{u})$ . Hence, we have constructed a double null foliation  $(u, \underline{u})$  in  $\mathcal{K}$  and its leaves are given by:

$$S(u, \underline{u}) := C_u \cap \underline{C}_{\underline{u}}. \quad (2.19)$$

For every  $(u, \underline{u})$ , we define<sup>10</sup>

$$V(u, \underline{u}) := J^-(S(u, \underline{u})) \cap J^+(\Sigma_0). \quad (2.20)$$

Next, we denote

$$\mathcal{K} := V(u_0, \underline{u}_*), \quad u_0 = -R_0, \quad (2.21)$$

called the *bootstrap region*, see Figure 4. Remark that  $\mathcal{K}$  is covered by the double null foliation  $(u, \underline{u})$ , and that the boundary of  $\mathcal{K}$  consists of:

- a bounded subset of  $\Sigma_0 \setminus K$ ;

<sup>9</sup>Notice that  $\Phi = \Phi_{(0)}$  on  $S_*$  since  $u = u_{(0)}$  and  $x^A = x^A_{(0)}$  on  $S_*$ .

<sup>10</sup>For any set  $A$ ,  $J^\pm(A)$  denotes the causal future (and resp. causal past) of  $A$ .

- a part of the outgoing null cone  $C_{u_0}$ ;
- a part of the last slice  $\underline{C}_*$ .

Recalling that the map  $\Phi$  is defined on  $\underline{C}_*$ , we now extend it to  $\mathcal{K}$ . As in Section 2.1.1, we denote  $L$  the generator of  $C_u$  and extend  $x^A$  to  $\mathcal{K}$  by

$$L(x^A) = 0, \quad A = 1, 2.$$

For any  $p \in \mathcal{K}$ , we denote  $\phi_t^L(p)$  the flow of  $L$  emanating from  $p$  to the future. Then, we denote

$$p_* := \phi_t^L(p) \cap \underline{C}_*, \quad q_* := \Phi(p_*).$$

Next, we denote  $\phi_t^{L_{Kerr}}(q_*)$  the flow of  $L_{Kerr}$  emanating from  $q_*$  to the past. We denote

$$q := \phi_t^{L_{Kerr}}(q_*) \cap S_{Kerr}(u(p), \underline{u}(p)).$$

Finally, we define

$$\Phi(p) = q. \quad (2.22)$$

Thus, we have extended the domain of definition of  $\Phi$  to  $\mathcal{K}$  and we have for any  $p \in \mathcal{K}$ <sup>11</sup>

$$\Phi(p) \in \mathcal{M}_{Kerr}, \quad u_{Kerr}(\Phi(p)) = u(p), \quad \underline{u}_{Kerr}(\Phi(p)) = \underline{u}(p), \quad x_{Kerr}^A(\Phi(p)) = x^A(p). \quad (2.23)$$

In other words, the expression of the application  $\Phi$  in the local coordinates charts  $(u, \underline{u}, x^1, x^2)$  and  $(u_{Kerr}, \underline{u}_{Kerr}, x_{Kerr}^1, x_{Kerr}^2)$  is the identity map, or equivalently

$$u = u_{Kerr} \circ \Phi, \quad \underline{u} = \underline{u}_{Kerr} \circ \Phi, \quad x^A = x_{Kerr}^A \circ \Phi, \quad A = 1, 2.$$

## 2.2 Integral formulae

We define the  $S$ -average of scalar functions.

**Definition 2.6.** *Given any scalar function  $f$ , we denote its average by*

$$\bar{f} := \frac{1}{|S|} \int_S f d\gamma.$$

We recall the following integral formulae, which will be used repeatedly in this paper.

**Lemma 2.7.** *Let  $S(u, \underline{u})$  a leaf of a double null foliation  $(u, \underline{u})$ . For any scalar function  $f$ , the following equations hold:*

$$\begin{aligned} \Omega e_4 \left( \int_{S(u, \underline{u})} f d\gamma \right) &= \int_{S(u, \underline{u})} (\Omega e_4(f) + \Omega \operatorname{tr} \chi f) d\gamma, \\ \Omega e_3 \left( \int_{S(u, \underline{u})} f d\gamma \right) &= \int_{S(u, \underline{u})} (\Omega e_3(f) + \Omega \operatorname{tr} \underline{\chi} f) d\gamma. \end{aligned}$$

Taking  $f = 1$ , we obtain

$$e_4(r) = \frac{\overline{\Omega \operatorname{tr} \chi}}{2\Omega} r, \quad e_3(r) = \frac{\overline{\Omega \operatorname{tr} \underline{\chi}}}{2\Omega} r,$$

where we recall that  $r$  is the area radius defined by

$$r(u, \underline{u}) = \sqrt{\frac{|S(u, \underline{u})|}{4\pi}}.$$

*Proof.* See Lemma 3.1.3 in [14]. □

---

<sup>11</sup>Note that the identity  $x_{Kerr}^A(\Phi(p)) = x^A(p)$  follows from the fact that it holds on  $\underline{C}_*$  and from the identities  $L(x^A) = 0$  and  $L_{Kerr}(x_{Kerr}^A) = 0$ .

### 2.3 Hodge systems

**Definition 2.8.** For tensor fields defined on a 2-sphere  $S$ , we denote by  $\mathfrak{s}_0 := \mathfrak{s}_0(S)$  the set of pairs of scalar functions,  $\mathfrak{s}_1 := \mathfrak{s}_1(S)$  the set of 1-forms and  $\mathfrak{s}_2 := \mathfrak{s}_2(S)$  the set of symmetric traceless 2-tensors.

**Definition 2.9.** Given  $\xi \in \mathfrak{s}_1$ , we define its Hodge dual

$$*\xi_A := \epsilon_{AB} \xi^B.$$

Clearly  $*\xi \in \mathfrak{s}_1$  and

$$*(*\xi) = -\xi.$$

Given  $U \in \mathfrak{s}_2$ , we define its Hodge dual

$$*U_{AB} := \epsilon_{AC} U^C_{\phantom{C}B}.$$

Observe that  $*U \in \mathfrak{s}_2$  and

$$*(*U) = -U.$$

**Definition 2.10.** Given  $\xi, \eta \in \mathfrak{s}_1$ , we denote

$$\begin{aligned} \xi \cdot \eta &:= \delta^{AB} \xi_A \eta_B, \\ \xi \wedge \eta &:= \epsilon^{AB} \xi_A \eta_B, \\ (\widehat{\xi \otimes \eta})_{AB} &:= \xi_A \eta_B + \xi_B \eta_A - \delta_{AB} \xi \cdot \eta, \end{aligned}$$

where  $\widehat{\xi \otimes \eta} \in \mathfrak{s}_2$ . Given  $\xi \in \mathfrak{s}_1$ ,  $U \in \mathfrak{s}_2$ , we denote

$$(\xi \cdot U)_A := \delta^{BC} \xi_B U_{AC}.$$

Given  $U, V \in \mathfrak{s}_2$ , we denote

$$U \wedge V := \epsilon^{AB} U_{AC} V_{CB}.$$

**Definition 2.11.** For a given  $\xi \in \mathfrak{s}_1$ , we define the following differential operators:

$$\begin{aligned} \operatorname{div} \xi &:= \delta^{AB} \nabla_A \xi_B, \\ \operatorname{curl} \xi &:= \epsilon^{AB} \nabla_A \xi_B, \\ (\widehat{\nabla \otimes \xi})_{AB} &:= \nabla_A \xi_B + \nabla_B \xi_A - \delta_{AB} (\operatorname{div} \xi). \end{aligned}$$

**Definition 2.12.** We define the Hodge type operators, as introduced in Section 2.2 in [5]:

- $d_1$  takes  $\mathfrak{s}_1$  into  $\mathfrak{s}_0$  and is given by:

$$d_1 \xi := (\operatorname{div} \xi, \operatorname{curl} \xi),$$

- $d_2$  takes  $\mathfrak{s}_2$  into  $\mathfrak{s}_1$  and is given by:

$$(d_2 U)_A := \nabla^A U_{AB},$$

- $d_1^*$  takes  $\mathfrak{s}_0$  into  $\mathfrak{s}_1$  and is given by:

$$d_1^*(f, f_*) := -\nabla_A f + \epsilon_{AB} \nabla_B f_*,$$

- $d_2^*$  takes  $\mathfrak{s}_1$  into  $\mathfrak{s}_2$  and is given by:

$$d_2^* \xi := -\frac{1}{2} \nabla \widehat{\otimes} \xi.$$

Then, we have the following identities:

$$\begin{aligned} d_1^* d_1 &= -\Delta_1 + \mathbf{K}, & d_1 d_1^* &= -\Delta_0, \\ d_2^* d_2 &= -\frac{1}{2} \Delta_2 + \mathbf{K}, & d_2 d_2^* &= -\frac{1}{2} (\Delta_1 + \mathbf{K}), \end{aligned} \quad (2.24)$$

where  $\mathbf{K}$  denotes the Gauss curvature of  $S$ . See for example (2.2.2) in [5] for a proof of (2.24).

**Definition 2.13.** We define weighted angular derivatives  $\mathfrak{d}$  as follows:

$$\begin{aligned} \mathfrak{d}U &:= r d_2 U, & \forall U &\in \mathfrak{s}_2, \\ \mathfrak{d}\xi &:= r d_1 \xi, & \forall \xi &\in \mathfrak{s}_1, \\ \mathfrak{d}f &:= r d_1^* f, & \forall f &\in \mathfrak{s}_0. \end{aligned}$$

Also, we denote for any tensor  $h \in \mathfrak{s}_k$ ,  $k = 0, 1, 2$ ,

$$h^{(0)} = h, \quad h^{(1)} = (h, r \nabla h).$$

## 2.4 Elliptic estimates for Hodge systems

**Definition 2.14.** For a tensor field  $f$  on a 2-sphere  $S$ , we denote its  $L^p$ -norm by

$$|f|_{L^p} := \left( \int_S |f|^p \right)^{\frac{1}{p}}, \quad (2.25)$$

where  $|f|^2 := f^{A_1 \dots A_k} f_{A_1 \dots A_k}$  for a  $k$ -tensor  $f$  on  $S$ .

**Proposition 2.15** ( $L^p$  estimates for Hodge systems). Assume that  $S$  is an arbitrary compact 2-surface satisfying  $c_1 \leq r^2 K \leq c_2$  where  $K$  denotes its Gauss curvature and  $0 < c_1 \leq c_2 < +\infty$  are two positive constants. Then the following statements hold for all<sup>12</sup>  $p \in (1, +\infty)$ :

1. Let  $\phi \in \mathfrak{s}_0$  be a solution of  $\Delta \phi = f$ . Then we have

$$|\nabla^2 \phi|_{L^p} + r^{-1} |\nabla \phi|_{L^p} + r^{-2} |\phi - \bar{\phi}|_{L^p} \lesssim |f|_{L^p}.$$

2. Let  $\xi \in \mathfrak{s}_1$  be a solution of  $d_1 \xi = (f, f_*)$ . Then we have

$$\int_S |\nabla \xi|^p + r^{-p} |\xi|^p \lesssim \int_S |f|^p + |f_*|^p.$$

3. Let  $U \in \mathfrak{s}_2$  be a solution of  $d_2 U = f$ . Then we have

$$\int_S |\nabla U|^p + r^{-p} |U|^p \lesssim \int_S |f|^p.$$

*Proof.* See Corollary 2.3.1.1 of [5]. □

---

<sup>12</sup>In practice, we only use Proposition 2.15 with  $p = 2$  and  $p = 4$ .

## 2.5 Null structure equations

We recall the standard null structure equations in a double null foliation.

**Proposition 2.16.** *We have the following null structure equations in a double null foliation:*

$$\begin{aligned}
\nabla_4 \eta &= -\chi \cdot (\eta - \underline{\eta}) - \beta, \\
\nabla_3 \underline{\eta} &= -\underline{\chi} \cdot (\underline{\eta} - \eta) + \underline{\beta}, \\
\nabla_4 \widehat{\chi} + (\text{tr } \chi) \widehat{\chi} &= -2\omega \widehat{\chi} - \alpha, \\
\nabla_4 \text{tr } \chi + \frac{1}{2}(\text{tr } \chi)^2 &= -|\widehat{\chi}|^2 - 2\omega \text{tr } \chi, \\
\nabla_3 \widehat{\underline{\chi}} + (\text{tr } \underline{\chi}) \widehat{\underline{\chi}} &= -2\underline{\omega} \widehat{\underline{\chi}} - \underline{\alpha}, \\
\nabla_3 \text{tr } \underline{\chi} + \frac{1}{2}(\text{tr } \underline{\chi})^2 &= -|\widehat{\underline{\chi}}|^2 - 2\underline{\omega} \text{tr } \underline{\chi}, \\
\nabla_4 \widehat{\underline{\chi}} + \frac{1}{2}(\text{tr } \chi) \widehat{\underline{\chi}} &= \nabla \widehat{\otimes} \eta + 2\omega \widehat{\underline{\chi}} - \frac{1}{2}(\text{tr } \underline{\chi}) \widehat{\chi} + \eta \widehat{\otimes} \underline{\eta}, \\
\nabla_3 \widehat{\chi} + \frac{1}{2}(\text{tr } \underline{\chi}) \widehat{\chi} &= \nabla \widehat{\otimes} \underline{\eta} + 2\underline{\omega} \widehat{\chi} - \frac{1}{2}(\text{tr } \chi) \widehat{\underline{\chi}} + \underline{\eta} \widehat{\otimes} \eta, \\
\nabla_4 \text{tr } \underline{\chi} + \frac{1}{2}(\text{tr } \chi) \text{tr } \underline{\chi} &= 2\omega \text{tr } \underline{\chi} + 2\rho - \widehat{\chi} \cdot \widehat{\underline{\chi}} + 2 \text{div } \underline{\eta} + 2|\underline{\eta}|^2, \\
\nabla_3 \text{tr } \chi + \frac{1}{2}(\text{tr } \underline{\chi}) \text{tr } \chi &= 2\underline{\omega} \text{tr } \chi + 2\rho - \widehat{\chi} \cdot \widehat{\underline{\chi}} + 2 \text{div } \eta + 2|\eta|^2.
\end{aligned}$$

We also have the Codazzi equations

$$\begin{aligned}
\text{div } \widehat{\chi} &= \frac{1}{2} \nabla \text{tr } \chi - \zeta \cdot \left( \widehat{\chi} - \frac{1}{2} \text{tr } \chi \right) - \beta, \\
\text{div } \widehat{\underline{\chi}} &= \frac{1}{2} \nabla \text{tr } \underline{\chi} + \zeta \cdot \left( \widehat{\underline{\chi}} - \frac{1}{2} \text{tr } \underline{\chi} \right) + \underline{\beta},
\end{aligned}$$

the torsion equation

$$\text{curl } \eta = -\text{curl } \underline{\eta} = \sigma + \frac{1}{2} \widehat{\underline{\chi}} \wedge \widehat{\chi},$$

and the Gauss equation

$$\mathbf{K} = -\frac{1}{4} \text{tr } \chi \text{tr } \underline{\chi} + \frac{1}{2} \widehat{\underline{\chi}} \cdot \widehat{\chi} - \rho.$$

Moreover, we have

$$\begin{aligned}
\nabla_4 \underline{\omega} &= 2\omega \underline{\omega} + \frac{3}{4} |\eta - \underline{\eta}|^2 - \frac{1}{4} (\eta - \underline{\eta}) \cdot (\eta + \underline{\eta}) - \frac{1}{8} |\eta + \underline{\eta}|^2 + \frac{1}{2} \rho, \\
\nabla_3 \omega &= 2\omega \omega + \frac{3}{4} |\eta - \underline{\eta}|^2 + \frac{1}{4} (\eta - \underline{\eta}) \cdot (\eta + \underline{\eta}) - \frac{1}{8} |\eta + \underline{\eta}|^2 + \frac{1}{2} \rho.
\end{aligned}$$

*Proof.* See (3.1)–(3.5) in [16]. □

**Definition 2.17.** *We define the mass aspect functions as follows:*

$$\mu := -\text{div } \eta + \frac{1}{2} \widehat{\chi} \cdot \widehat{\underline{\chi}} - \rho, \quad \underline{\mu} := -\text{div } \underline{\eta} + \frac{1}{2} \widehat{\underline{\chi}} \cdot \widehat{\chi} - \rho. \quad (2.26)$$

We also define the following renormalized quantities which will be used in Sections 5 and 6:

$$[\mu] := \mu + \frac{1}{4} \text{tr } \chi \text{tr } \underline{\chi}, \quad [\underline{\mu}] := \underline{\mu} + \frac{1}{4} \text{tr } \chi \text{tr } \underline{\chi}. \quad (2.27)$$

**Proposition 2.18.** *We have the following null structure equations:*

$$\begin{aligned}
\Omega \nabla_4(\Omega \operatorname{tr} \chi) + \frac{1}{2}(\Omega \operatorname{tr} \chi)^2 &= -4\Omega \operatorname{tr} \chi(\Omega \omega) - \Omega^2 |\widehat{\chi}|^2, \\
\Omega \nabla_3(\Omega \operatorname{tr} \underline{\chi}) + \frac{1}{2}(\Omega \operatorname{tr} \underline{\chi})^2 &= -4\Omega \operatorname{tr} \underline{\chi}(\Omega \omega) - \Omega^2 |\widehat{\underline{\chi}}|^2, \\
\Omega \nabla_4(\Omega \operatorname{tr} \underline{\chi}) &= -2\Omega^2[\mu] + 2\Omega^2|\eta|^2, \\
\Omega \nabla_3(\Omega \operatorname{tr} \chi) &= -2\Omega^2[\mu] + 2\Omega^2|\eta|^2.
\end{aligned} \tag{2.28}$$

*We have the following equations for the renormalized mass aspect functions:*

$$\begin{aligned}
\Omega \nabla_4[\mu] + \Omega \operatorname{tr} \chi[\mu] &= \Omega[F] + \Omega(\operatorname{tr} \chi \rho - 2\eta \cdot \beta), \\
[F] &:= \widehat{\chi} \cdot (\nabla \widehat{\otimes} \eta) + (\eta - \underline{\eta}) \cdot (\nabla \operatorname{tr} \chi + \operatorname{tr} \chi \zeta) + \frac{1}{2} \operatorname{tr} \chi (|\eta|^2 - |\underline{\eta}|^2) \\
&\quad - \frac{1}{2} (\operatorname{tr} \underline{\chi} |\widehat{\chi}|^2 + \operatorname{tr} \chi (\widehat{\chi} \cdot \widehat{\underline{\chi}})) + 2\eta \cdot \widehat{\chi} \cdot \underline{\eta}, \\
\Omega \nabla_3[\mu] + \Omega \operatorname{tr} \underline{\chi}[\mu] &= \Omega[\underline{F}] + \Omega(\operatorname{tr} \underline{\chi} \rho + 2\underline{\eta} \cdot \beta), \\
[\underline{F}] &:= \widehat{\underline{\chi}} \cdot (\nabla \widehat{\otimes} \underline{\eta}) + (\underline{\eta} - \eta) \cdot (\nabla \operatorname{tr} \underline{\chi} - \operatorname{tr} \underline{\chi} \zeta) + \frac{1}{2} \operatorname{tr} \underline{\chi} (|\underline{\eta}|^2 - |\eta|^2) \\
&\quad - \frac{1}{2} (\operatorname{tr} \underline{\chi} |\widehat{\underline{\chi}}|^2 + \operatorname{tr} \underline{\chi} (\widehat{\underline{\chi}} \cdot \widehat{\chi})) + 2\underline{\eta} \cdot \widehat{\underline{\chi}} \cdot \eta.
\end{aligned} \tag{2.29}$$

*Proof.* Notice that (2.28) follows directly from Proposition 2.16 and (2.8). See Lemma 4.3.2 in [14] for the proof of (2.29).  $\square$

## 2.6 Bianchi equations

We recall the Bianchi equations in a double null foliation.

**Proposition 2.19.** *The Bianchi equations take the following form in a double null foliation:*

$$\begin{aligned}
\nabla_4 \underline{\alpha} + \frac{1}{2} \operatorname{tr} \chi \underline{\alpha} &= -\nabla \widehat{\otimes} \underline{\beta} + 4\omega \underline{\alpha} - 3(\widehat{\chi} \rho - {}^* \widehat{\chi} \sigma) + (\zeta - 4\underline{\eta}) \widehat{\otimes} \underline{\beta}, \\
\nabla_3 \underline{\beta} + 2 \operatorname{tr} \underline{\chi} \underline{\beta} &= -\operatorname{div} \underline{\alpha} - 2\omega \underline{\beta} + (2\zeta - \eta) \cdot \underline{\alpha}, \\
\nabla_4 \underline{\beta} + \operatorname{tr} \chi \underline{\beta} &= -\nabla \rho + {}^* \nabla \sigma + 2\omega \underline{\beta} + 2\widehat{\underline{\chi}} \cdot \beta - 3(\underline{\eta} \rho - {}^* \underline{\eta} \sigma), \\
\nabla_3 \rho + \frac{3}{2} \operatorname{tr} \underline{\chi} \rho &= -\operatorname{div} \underline{\beta} - \frac{1}{2} \widehat{\chi} \cdot \underline{\alpha} + \zeta \cdot \underline{\beta} - 2\eta \cdot \underline{\beta}, \\
\nabla_4 \rho + \frac{3}{2} \operatorname{tr} \chi \rho &= \operatorname{div} \beta - \frac{1}{2} \widehat{\underline{\chi}} \cdot \alpha + \zeta \cdot \beta + 2\underline{\eta} \cdot \beta, \\
\nabla_3 \sigma + \frac{3}{2} \operatorname{tr} \underline{\chi} \sigma &= -\operatorname{curl} \underline{\beta} + \frac{1}{2} \widehat{\chi} \cdot {}^* \underline{\alpha} - \zeta \cdot {}^* \underline{\beta} + 2\eta \cdot {}^* \underline{\beta}, \\
\nabla_4 \sigma + \frac{3}{2} \operatorname{tr} \chi \sigma &= -\operatorname{curl} \beta + \frac{1}{2} \widehat{\underline{\chi}} \cdot {}^* \alpha - \zeta \cdot {}^* \beta - 2\underline{\eta} \cdot {}^* \beta, \\
\nabla_3 \beta + \operatorname{tr} \underline{\chi} \beta &= \nabla \rho + {}^* \nabla \sigma + 2\omega \beta + 2\widehat{\chi} \cdot \underline{\beta} + 3(\eta \rho + {}^* \eta \sigma), \\
\nabla_4 \beta + 2 \operatorname{tr} \chi \beta &= \operatorname{div} \alpha - 2\omega \beta + (2\zeta + \eta) \alpha, \\
\nabla_3 \alpha + \frac{1}{2} \operatorname{tr} \underline{\chi} \alpha &= \nabla \widehat{\otimes} \beta + 4\omega \alpha - 3(\widehat{\chi} \rho + {}^* \widehat{\chi} \sigma) + (\zeta + 4\eta) \widehat{\otimes} \beta.
\end{aligned}$$

*Proof.* See Proposition 3.2.4 in [14].  $\square$

## 2.7 Transport equations for metric components

We introduce the following transport equations for metric components.

**Proposition 2.20.** *The metric components satisfy the following equations:*

$$\begin{aligned}\nabla_4(\gamma_{AB}) &= 2\chi_{AB}, \\ \nabla_4(\underline{b}^A) &= -4\Omega\zeta^A, \\ \nabla_4(\in_{AB}) &= \text{tr } \chi \in_{AB}, \\ \nabla_3(\gamma_{AB}) &= 2\underline{\chi}_{AB} - \Omega^{-1}\partial_A(\underline{b}^C)\gamma_{BC} - \Omega^{-1}\partial_B(\underline{b}^C)\gamma_{AC}, \\ \nabla_3(\in_{AB}) &= \text{tr } \underline{\chi} \in_{AB} - \Omega^{-1}\partial_A(\underline{b}^C) \in_{BC} - \Omega^{-1}\partial_B(\underline{b}^C) \in_{AC}.\end{aligned}$$

*Proof.* We have from  $[\partial_{\underline{u}}, \partial_A] = 0$  that

$$\begin{aligned}0 &= \nabla_{\partial_{\underline{u}}}(\gamma) \left( \frac{\partial}{\partial x^A}, \frac{\partial}{\partial x^B} \right) = \frac{\partial}{\partial \underline{u}} \gamma_{AB} - \gamma(\mathbf{D}_{\partial_{\underline{u}}} \partial_A, \partial_B) - \gamma(\partial_A, \mathbf{D}_{\partial_{\underline{u}}} \partial_B) \\ &= \Omega \nabla_4(\gamma_{AB}) - \Omega \mathbf{g}(\mathbf{D}_{e_4} \partial_A, \partial_B) - \Omega \mathbf{g}(\partial_A, \mathbf{D}_{e_4} \partial_B) \\ &= \Omega \nabla_4(\gamma_{AB}) - 2\Omega \chi_{AB},\end{aligned}$$

which implies

$$\nabla_4(\gamma_{AB}) = 2\chi_{AB}$$

as stated in the first equation. For the proof of the second equation, see (3.1.62) in [14].

We have from  $[\partial_{\underline{u}}, \partial_A] = 0$  that

$$\begin{aligned}0 &= \nabla_{\partial_{\underline{u}}}(\in) \left( \frac{\partial}{\partial x^A}, \frac{\partial}{\partial x^B} \right) = \frac{\partial}{\partial \underline{u}} \in_{AB} - \in(\mathbf{D}_{\partial_{\underline{u}}} \partial_A, \partial_B) - \in(\partial_A, \mathbf{D}_{\partial_{\underline{u}}} \partial_B) \\ &= \frac{\partial}{\partial \underline{u}} \in_{AB} - \in(\mathbf{D}_{\partial_A} \partial_{\underline{u}}, \partial_B) - \in(\partial_A, \mathbf{D}_{\partial_B} \partial_{\underline{u}}) \\ &= \frac{\partial}{\partial \underline{u}} \in_{AB} - \Omega \in(\chi_A^C \partial_C, \partial_B) - \Omega \in(\partial_A, \chi_B^C \partial_C) \\ &= \Omega \nabla_4(\in_{AB}) - \Omega \chi_A^C \in_{CB} - \Omega \chi_B^C \in_{AC},\end{aligned}$$

which implies

$$\nabla_4(\in_{AB}) = \chi_A^C \in_{CB} + \chi_B^C \in_{AC} = \text{tr } \chi \in_{AB}.$$

Next, we compute

$$\begin{aligned}0 &= (\nabla_{\Omega e_3} \gamma) \left( \frac{\partial}{\partial x^A}, \frac{\partial}{\partial x^B} \right) \\ &= \Omega \nabla_3(\gamma_{AB}) - \gamma(\nabla_{\Omega e_3} \partial_A, \partial_B) - \gamma(\partial_A, \nabla_{\Omega e_3} \partial_B) \\ &= \Omega \nabla_3(\gamma_{AB}) - \Omega \mathbf{g}(\mathbf{D}_{\partial_A} e_3, \partial_B) - \Omega \mathbf{g}(\partial_A, \mathbf{D}_{\partial_B} e_3) - \mathbf{g}([\Omega e_3, \partial_A], \partial_B) - \mathbf{g}(\partial_A, [\Omega e_3, \partial_B]) \\ &= \Omega \nabla_3(\gamma_{AB}) - 2\Omega \underline{\chi}_{AB} - \mathbf{g}([\underline{b}^B \partial_C, \partial_A], \partial_B) - \mathbf{g}(\partial_A, [\underline{b}^C \partial_C, \partial_B]) \\ &= \Omega \nabla_3(\gamma_{AB}) - 2\Omega \underline{\chi}_{AB} + \partial_A(\underline{b}^C) \gamma_{BC} + \partial_B(\underline{b}^C) \gamma_{AC},\end{aligned}$$

which implies

$$\nabla_3(\gamma_{AB}) = 2\underline{\chi}_{AB} - \Omega^{-1}\partial_A(\underline{b}^C)\gamma_{BC} - \Omega^{-1}\partial_B(\underline{b}^C)\gamma_{AC}. \quad (2.30)$$

Similarly, we have

$$\nabla_3(\in_{AB}) = \text{tr } \underline{\chi} \in_{AB} - \Omega^{-1}\partial_A(\underline{b}^C) \in_{BC} - \Omega^{-1}\partial_B(\underline{b}^C) \in_{AC}.$$

This concludes the proof of Proposition 2.20.  $\square$

## 2.8 The linearized equations

### 2.8.1 Linearization procedure

Let  $\mathcal{K}$  the bootstrap region defined in Section 2.1.5. By construction, we have

$$L(x^A) = 0.$$

Hence, the metric in  $\mathcal{K}$  takes the following form, see (2.3):

$$\mathbf{g} = -2\Omega^2(d\underline{u} \otimes du + du \otimes d\underline{u}) + \gamma_{AB}(dx^A - \underline{b}^A du) \otimes (dx^B - \underline{b}^B du). \quad (2.31)$$

The inverse metric components are given by:

$$\begin{aligned} \mathbf{g}^{AB} &= \gamma^{AB}, & \mathbf{g}^{uu} &= -\frac{1}{2\Omega^2}, & \mathbf{g}^{uA} &= -\frac{\underline{b}^A}{2\Omega^2}, \\ \mathbf{g}^{uu} &= \mathbf{g}^{\underline{u}\underline{u}} = 0, & \mathbf{g}^{uA} &= 0, & A, B &= 1, 2. \end{aligned} \quad (2.32)$$

Let  $X$  be a tensor defined on  $S := S(u, \underline{u})$ . We denote  $X_{Kerr}$  the corresponding  $S_{Kerr}$ -tangent tensor in  $\mathcal{M}_{Kerr}$ . Then, its pullback  $\Phi^* X_{Kerr}$ <sup>13</sup> is a  $S$ -tangent tensor in  $\mathcal{K}$  denoted by

$$X_K := \Phi^* X_{Kerr}. \quad (2.33)$$

We now define the linearization of  $X$  as follows

$$\check{X} := X - X_K. \quad (2.34)$$

We also denote  $\nabla_K$ , (resp.  $\Gamma_K$ ,  $\text{div}_K$  and  $\text{curl}_K$ ) the Levi-Civita connection, (resp. Christoffel symbols, divergence operator and curl operator) of the metric  $\mathbf{g}_K = \Phi^*(\mathbf{g}_{Kerr})$ .

The following lemma plays an important role in the linearization of the equations.

**Lemma 2.21.** *We have the following formulae:*

$$\begin{aligned} \widetilde{\Omega e_4} &:= \Omega e_4 - \Omega_K(e_4)_K = 0, \\ \widetilde{\Omega e_3} &:= \Omega e_3 - \Omega_K(e_3)_K = \check{\underline{b}}^A \partial_{x^A}. \end{aligned}$$

*Proof.* Notice that we have

$$\begin{aligned} e_4(u) &= 0, & e_4(\underline{u}) &= \mathbf{g}(e_4, \text{grad } \underline{u}) = \mathbf{g}(2\Omega L, -\underline{L}) = \Omega^{-1}, \\ e_3(\underline{u}) &= 0, & e_3(u) &= \mathbf{g}(e_3, \text{grad } u) = \mathbf{g}(2\Omega \underline{L}, -L) = \Omega^{-1}, \\ e_4(x^A) &= 0, & e_3(x^A) &= \Omega^{-1} \underline{b}^A. \end{aligned} \quad (2.35)$$

Hence, we obtain

$$\Omega e_4 = \Omega e_4(u) \partial_u + \Omega e_4(\underline{u}) \partial_{\underline{u}} + \Omega e_4(x^A) \partial_{x^A} = \partial_{\underline{u}}. \quad (2.36)$$

Thus, we infer

$$\Phi_*(\Omega e_4) = \Phi_*(\partial_{\underline{u}}) = \partial_{\underline{u}_{Kerr}} = \Omega_{Kerr}(e_4)_{Kerr},$$

which implies

$$\Omega_K(e_4)_K = \Phi^*(\Omega_{Kerr}(e_4)_{Kerr}) = \Omega e_4.$$

---

<sup>13</sup>Recall that the map  $\Phi : \mathcal{K} \rightarrow \mathcal{M}_{Kerr}$  is defined in (2.22) and satisfies (2.23).

Hence, we deduce

$$\widetilde{\Omega e_4} = \Omega e_4 - \Omega_K(e_4)_K = \partial_{\underline{u}} - \partial_{\underline{u}} = 0. \quad (2.37)$$

Similarly, we have

$$\Omega e_3 = \partial_u + \underline{b}^A \partial_{x^A},$$

which implies

$$\Phi_*(\Omega e_3) = \partial_{u_{Kerr}} + (\Phi^{-1})^*(\underline{b}^A) \partial_{x_{Kerr}^A}.$$

Hence, we obtain

$$\begin{aligned} \widetilde{\Omega e_3} &= \Omega e_3 - \Omega_K(e_3)_K = \Phi_*^{-1}(\Phi_*(\Omega e_3) - \Omega_{Kerr}(e_3)_{Kerr}) \\ &= \Phi_*^{-1}\left((\Phi^{-1})^*\underline{b}^A - \underline{b}_{Kerr}^A\right) \partial_{x_{Kerr}^A} \\ &= (\underline{b}^A - \underline{b}_K^A) \partial_{x^A} = \check{\underline{b}}^A \partial_{x^A}. \end{aligned} \quad (2.38)$$

This concludes the proof of Lemma 2.21.  $\square$

**Proposition 2.22.** Denoting  $\Gamma$  and  $\Gamma_K$  respectively the Christoffel symbols of  $\nabla$  and  $\nabla_K$ . For any  $U \in \mathfrak{s}_2(S)$  and  $\phi \in \mathfrak{s}_1(S)$  where  $S = S(u, \underline{u})$ , we have the following formulae:

$$\begin{aligned} (\nabla - \nabla_K)_A U_{BC} &= -(\Gamma - \Gamma_K)_{AB}^D U_{CD} - (\Gamma - \Gamma_K)_{AC}^D U_{BD}, \\ (\nabla - \nabla_K)_A \phi_B &= -(\Gamma - \Gamma_K)_{AB}^C \phi_C, \\ (\nabla - \nabla_K)_{\partial_{\underline{u}}} U_{AB} &= -(\Gamma - \Gamma_K)_{\underline{u}A}^C U_{CB} - (\Gamma - \Gamma_K)_{\underline{u}B}^C U_{CA}, \\ (\nabla - \nabla_K)_{\partial_{\underline{u}}} \phi_A &= -(\Gamma - \Gamma_K)_{\underline{u}A}^B \phi_B, \\ (\nabla - \nabla_K)_{\partial_u} U_{AB} &= -(\Gamma - \Gamma_K)_{uA}^C U_{CB} - (\Gamma - \Gamma_K)_{uB}^C U_{CA}, \\ (\nabla - \nabla_K)_{\partial_u} \phi_A &= -(\Gamma - \Gamma_K)_{uA}^B \phi_B. \end{aligned} \quad (2.39)$$

We also have:

$$\begin{aligned} (\operatorname{div} - \operatorname{div}_K) U &= (\gamma^{AB} - \gamma_K^{AB}) \nabla_A U_{B\bullet} - \gamma_K^{AB} (\Gamma - \Gamma_K)_{AB}^D U_{\bullet D} \\ &\quad - \gamma_K^{AB} (\Gamma - \Gamma_K)_{A\bullet}^D U_{BD}, \\ (\operatorname{div} - \operatorname{div}_K) \phi &= (\gamma^{AB} - (\gamma_K)^{AB}) \nabla_A \phi_B - (\gamma_K)^{AB} (\Gamma - \Gamma_K)_{AB}^C \phi_C, \\ (\operatorname{curl} - \operatorname{curl}_K) \phi &= (\epsilon^{AB} - (\epsilon_K)^{AB}) \nabla_A \phi_B - (\epsilon_K)^{AB} (\Gamma - \Gamma_K)_{AB}^C \phi_C, \\ (\nabla \widehat{\otimes} \phi - \nabla_K \widehat{\otimes} \phi_K)_{AB} &= (\nabla \widehat{\otimes} \check{\phi})_{AB} - 2(\Gamma - \Gamma_K)_{AB}^C (\phi_K)_C \\ &\quad + \delta_{AB} [(\gamma^{CD} - (\gamma_K)^{CD}) \nabla_C (\phi_K)_D - (\gamma_K)^{CD} (\Gamma - \Gamma_K)_{CD}^E (\phi_K)_E]. \end{aligned} \quad (2.40)$$

*Proof.* The proof follows from a direct computation. For example, we have

$$\begin{aligned} (\nabla - \nabla_K)_A U_{BC} &= \left( \frac{\partial}{\partial x^A} (U_{BC}) - \Gamma_{AB}^D U_{CD} - \Gamma_{AC}^D U_{BD} \right) \\ &\quad - \left( \frac{\partial}{\partial x^A} (U_{BC}) - (\Gamma_K)_{AB}^D U_{CD} - (\Gamma_K)_{AC}^D U_{BD} \right) \\ &= -(\Gamma - \Gamma_K)_{AB}^D U_{CD} - (\Gamma - \Gamma_K)_{AC}^D U_{BD}, \end{aligned}$$

which implies the first equation in (2.39). The proof of the other equations in (2.39) are similar. We also have

$$\begin{aligned} (\operatorname{div} - \operatorname{div}_K) U_C &= \gamma^{AB} \nabla_A U_{BC} - \gamma_K^{AB} (\nabla_K)_A U_{BC} \\ &= (\gamma^{AB} - \gamma_K^{AB}) \nabla_A U_{BC} + \gamma_K^{AB} (\nabla_A U_{BC} - (\nabla_K)_A U_{BC}) \\ &= (\gamma^{AB} - \gamma_K^{AB}) \nabla_A U_{BC} - \gamma_K^{AB} (\Gamma - \Gamma_K)_{AB}^D U_{CD} - \gamma_K^{AB} (\Gamma - \Gamma_K)_{AC}^D U_{BD}, \end{aligned}$$

which implies the first equation in (2.40). The proof of the other equations in (2.40) are similar and left to the reader. This concludes the proof of Proposition 2.22.  $\square$

**Lemma 2.23.** *We have the following identities:*

$$\begin{aligned}\Gamma_{uA}^B &= \frac{1}{2}\gamma^{BC}\partial_u(\gamma_{AC}), \\ \Gamma_{uA}^B &= \frac{1}{2}\gamma^{BC}\left(\partial_u(\gamma_{AC}) - \frac{\partial}{\partial x^A}(\gamma_{CD}\underline{b}^D) + \frac{\partial}{\partial x^C}(\gamma_{AD}\underline{b}^D)\right) + \frac{\underline{b}^B}{4\Omega^2}\left(2\frac{\partial}{\partial x^A}(\Omega^2) - \frac{\partial}{\partial \underline{u}}(\gamma_{AD}\underline{b}^D)\right), \\ \Gamma_{BC}^A &= \frac{1}{2}\gamma^{AD}\left(\frac{\partial\gamma_{BD}}{\partial x^C} + \frac{\partial\gamma_{CD}}{\partial x^B} - \frac{\partial\gamma_{BC}}{\partial x^D}\right) + \frac{\underline{b}^A}{4\Omega^2}\frac{\partial\gamma_{BC}}{\partial \underline{u}}.\end{aligned}$$

*Proof.* We have from (2.31) and (2.32)

$$\Gamma_{uA}^B = \frac{1}{2}\mathbf{g}^{BC}\left(\frac{\partial\mathbf{g}_{uC}}{\partial x^A} + \frac{\partial\mathbf{g}_{AC}}{\partial \underline{u}} - \frac{\partial\mathbf{g}_{uA}}{\partial x^C}\right) + \frac{1}{2}\mathbf{g}^{Bu}\left(\frac{\partial\mathbf{g}_{uu}}{\partial x^A} + \frac{\partial\mathbf{g}_{Au}}{\partial \underline{u}} - \frac{\partial\mathbf{g}_{uA}}{\partial \underline{u}}\right) = \frac{1}{2}\gamma^{BC}\partial_u(\gamma_{AC}),$$

and similarly

$$\begin{aligned}\Gamma_{uA}^B &= \frac{1}{2}\mathbf{g}^{BC}\left(\frac{\partial\mathbf{g}_{uC}}{\partial x^A} + \frac{\partial\mathbf{g}_{AC}}{\partial \underline{u}} - \frac{\partial\mathbf{g}_{uA}}{\partial x^C}\right) + \frac{1}{2}\mathbf{g}^{Bu}\left(\frac{\partial\mathbf{g}_{uu}}{\partial x^A} + \frac{\partial\mathbf{g}_{Au}}{\partial \underline{u}} - \frac{\partial\mathbf{g}_{uA}}{\partial \underline{u}}\right) \\ &= \frac{1}{2}\gamma^{BC}\left(-\frac{\partial}{\partial x^A}(\gamma_{CD}\underline{b}^D) + \frac{\partial\gamma_{AC}}{\partial \underline{u}} + \frac{\partial}{\partial x^C}(\gamma_{AD}\underline{b}^D)\right) + \frac{\underline{b}^B}{4\Omega^2}\left(2\frac{\partial}{\partial x^A}(\Omega^2) - \frac{\partial}{\partial \underline{u}}(\gamma_{AD}\underline{b}^D)\right).\end{aligned}$$

Finally, we compute

$$\begin{aligned}\Gamma_{BC}^A &= \frac{1}{2}\gamma^{AD}\left(\frac{\partial\gamma_{BD}}{\partial x^C} + \frac{\partial\gamma_{CD}}{\partial x^B} - \frac{\partial\gamma_{BC}}{\partial x^D}\right) + \frac{1}{2}\mathbf{g}^{Au}\left(\frac{\partial\mathbf{g}_{Bu}}{\partial x^C} + \frac{\partial\mathbf{g}_{Cu}}{\partial x^B} - \frac{\partial\mathbf{g}_{BC}}{\partial \underline{u}}\right) \\ &= \frac{1}{2}\gamma^{AD}\left(\frac{\partial\gamma_{BD}}{\partial x^C} + \frac{\partial\gamma_{CD}}{\partial x^B} - \frac{\partial\gamma_{BC}}{\partial x^D}\right) + \frac{\underline{b}^A}{4\Omega^2}\frac{\partial\gamma_{BC}}{\partial \underline{u}}.\end{aligned}$$

This concludes the proof of Lemma 2.23.  $\square$

The following tensors play an important role in the linearization of the null structure equations and Bianchi equations.

**Definition 2.24.** *We define the following tensors<sup>14</sup>*

$$\check{J}_A^B := (\Gamma - \Gamma_K)_{uA}^B, \quad \check{J}_A^B := (\Gamma - \Gamma_K)_{\underline{u}A}^B, \quad \check{L}_{AB}^C := (\Gamma - \Gamma_K)_{AB}^C. \quad (2.41)$$

### 2.8.2 Schematic notation $\Gamma_g$ and $\Gamma_b$

We introduce the following schematic notations for metric components, Ricci coefficients and curvature components.

**Definition 2.25.** *We divide the linearized quantities into two parts:*

$$\begin{aligned}\Gamma_g &:= \Gamma_g^{(0)} := \{\check{\eta}, \check{\underline{\eta}}, \check{\zeta}, \check{\chi}, \widetilde{\text{tr}\chi}, \widetilde{\text{tr}\underline{\chi}}, \check{\omega}, r^{-1}\check{\underline{b}}, r^{-1}\check{\Omega}, r^{-1}\check{\gamma}, r^{-1}\check{\epsilon}, r^{-2}\check{\gamma}\}, \\ \Gamma_b &:= \Gamma_b^{(0)} := \{\check{\underline{\chi}}, \check{\underline{\omega}}\}.\end{aligned}$$

We also denote:

$$\begin{aligned}\Gamma_g^{(1)} &:= (r\nabla)^{\leq 1}\Gamma_g^{(0)} \cup \{\check{J}, \check{L}, r\check{\beta}, r\check{\rho}, r\check{\sigma}\}, \\ \Gamma_b^{(1)} &:= (r\nabla)^{\leq 1}\Gamma_b^{(0)} \cup \{\check{\underline{J}}, r\check{\underline{\beta}}\}.\end{aligned}$$

**Remark 2.26.** *Throughout this paper,  $\Gamma_g$  decays better than  $\Gamma_b$ , see Lemma 4.1.*

<sup>14</sup>Remark that the difference of two Christoffel symbols is a tensor.

### 2.8.3 Schematic notation $\mathbf{O}_q^p$

**Definition 2.27.** For any  $S$ -tangent tensor field  $X_K = \Phi^*(X_{Kerr})$ , we denote

$$X_K = \mathbf{O}_q^p,$$

if for any integer  $l \geq 0$  we have

$$|(\partial_K)^l X_K| \lesssim \frac{M^p}{r_K^{q+l}},$$

where  $r_K = \Phi^*(r_{Kerr})$  and

$$\partial_K := \{(\nabla_K)_{(e_3)_K}, \nabla_K, (\nabla_K)_{(e_4)_K}\}.$$

**Proposition 2.28.** We have the following formulae:

$$\begin{aligned} \mathrm{tr}_K \chi_K - \frac{2}{r_K} &= \mathbf{O}_2^1, & \mathrm{tr}_K \underline{\chi}_K + \frac{2}{r_K} &= \mathbf{O}_2^1, \\ \omega_K &= \mathbf{O}_2^1, & \underline{\omega}_K &= \mathbf{O}_2^1, \\ \widehat{\chi}_K &= \mathbf{O}_3^2, & \widehat{\underline{\chi}}_K &= \mathbf{O}_3^2, \\ \nabla_K \mathrm{tr}_K \chi_K &= \mathbf{O}_4^2, & \nabla_K \mathrm{tr}_K \underline{\chi}_K &= \mathbf{O}_4^2, \\ \eta_K &= \mathbf{O}_3^2, & \underline{\eta}_K &= \mathbf{O}_3^2, \\ \underline{b}_K &= \mathbf{O}_2^2, & \Omega_K - \frac{1}{2} &= \mathbf{O}_1^1. \end{aligned}$$

We also have:

$$\begin{aligned} \alpha_K &= \mathbf{O}_5^3, & \underline{\alpha}_K &= \mathbf{O}_5^3, \\ \beta_K &= \mathbf{O}_4^2, & \underline{\beta}_K &= \mathbf{O}_4^2, \\ \rho_K &= \mathbf{O}_3^1, & \sigma_K &= \mathbf{O}_4^2. \end{aligned}$$

*Proof.* It follows directly from Propositions 2.4, 2.5 and Definition 2.27.  $\square$

**Remark 2.29.** In the sequel, we consider the following conventions:

- For a quantity  $h$  satisfying the same or even better decay as  $\Gamma_b$ , (resp.  $\Gamma_g$ ), we write

$$h \in \Gamma_b, \quad (\text{resp. } h \in \Gamma_g).$$

- For a sum of schematic notations, we ignore the terms that decay better. For example, we write

$$\Gamma_g + \Gamma_b = \Gamma_b, \quad \Gamma_g \cdot \mathbf{O}_q^p + \Gamma_b \cdot \mathbf{O}_q^p = \Gamma_b \cdot \mathbf{O}_q^p,$$

since  $\Gamma_g$  decays better than  $\Gamma_b$  throughout this paper.

**Lemma 2.30.** We have the following formulae:

$$\begin{aligned} \mathrm{tr} \chi - \frac{2}{r} &= \mathbf{O}_2^1 + \Gamma_g, & \mathrm{tr} \underline{\chi} + \frac{2}{r} &= \mathbf{O}_2^1 + \Gamma_b, \\ \widehat{\chi} &= \mathbf{O}_3^2 + \Gamma_g, & \widehat{\underline{\chi}} &= \mathbf{O}_3^2 + \Gamma_b, \\ \eta &= \mathbf{O}_3^2 + \Gamma_g, & \underline{\eta} &= \mathbf{O}_3^2 + \Gamma_g, \\ \omega &= \mathbf{O}_2^1 + \Gamma_g, & \underline{\omega} &= \mathbf{O}_2^1 + \Gamma_b, \\ \underline{b} &= \mathbf{O}_2^2 + r\Gamma_g, & \Omega - \frac{1}{2} &= \mathbf{O}_2^1 + r\Gamma_g, \\ \gamma &= \mathbf{O}_0^0 + r\Gamma_g, & \epsilon &= \mathbf{O}_0^0 + r\Gamma_g. \end{aligned}$$

We also have:

$$\begin{aligned}\alpha &= \mathbf{O}_5^3 + \check{\alpha}, & \underline{\alpha} &= \mathbf{O}_5^3 + \check{\underline{\alpha}}, \\ \beta &= \mathbf{O}_4^2 + \check{\beta}, & \underline{\beta} &= \mathbf{O}_4^2 + \check{\underline{\beta}}, \\ \rho &= \mathbf{O}_3^1 + \check{\rho}, & \sigma &= \mathbf{O}_4^2 + \check{\sigma}.\end{aligned}$$

*Proof.* It follows directly from Definitions 2.25, 2.27 and Proposition 2.28.  $\square$

The following lemma allows us to compute the derivatives of  $\mathbf{O}_q^p$ .

**Lemma 2.31.** *Let  $X_K$  be a  $S$ -tangent tensor field satisfying*

$$X_K = \mathbf{O}_q^p.$$

*Then, we have*

$$\begin{aligned}\Omega \nabla_3 X_K &= \mathbf{O}_{q+1}^p + \mathbf{O}_q^p \cdot \Gamma_g + \mathbf{O}_q^p \cdot \check{\underline{J}} + \mathbf{O}_q^p \cdot \check{\underline{L}}, \\ \Omega \nabla_4 X_K &= \mathbf{O}_{q+1}^p + \mathbf{O}_q^p \cdot \check{J}, \\ \nabla X_K &= \mathbf{O}_{q+1}^p + \mathbf{O}_q^p \cdot \check{L}.\end{aligned}$$

*Proof.* We have from Lemma 2.21 and Proposition 2.22

$$\begin{aligned}\Omega \nabla_3 X_K - \Omega_K \nabla_{(e_3)_K} X_K &= (\nabla_{\Omega e_3} - \nabla_{\Omega_K(e_3)_K}) X_K + (\nabla_{\Omega_K(e_3)_K} - (\nabla_K)_{\Omega_K(e_3)_K}) X_K \\ &= \check{\underline{b}}^A \nabla_{\partial_{x^A}} X_K + (\nabla - \nabla_K)_{\partial_u} X_K + \check{\underline{b}}_K^A (\nabla - \nabla_K)_{\partial_{x^A}} X_K \\ &= \check{\underline{b}}^A (\nabla - \nabla_K)_{\partial_{x^A}} X_K + \check{\underline{b}}^A (\nabla_K)_{\partial_{x^A}} X_K + \check{\underline{J}} \cdot \mathbf{O}_q^p + \check{\underline{L}} \cdot \mathbf{O}_q^p \\ &= \Gamma_g \cdot \mathbf{O}_q^p + \check{\underline{J}} \cdot \mathbf{O}_q^p + \check{\underline{L}} \cdot \mathbf{O}_q^p,\end{aligned}$$

where we used Remark 2.29 to ignore the terms having better decay. Combining with Definition 2.27, we obtain

$$\Omega \nabla_3 X_K = \mathbf{O}_{q+1}^p + \Gamma_g \cdot \mathbf{O}_q^p + \check{\underline{J}} \cdot \mathbf{O}_q^p + \check{\underline{L}} \cdot \mathbf{O}_q^p.$$

Similarly, we have from Lemma 2.21 and Proposition 2.22

$$\begin{aligned}\Omega \nabla_4 X_K - \Omega_K \nabla_{(e_4)_K} X_K &= (\nabla_{\Omega e_4} - \nabla_{\Omega_K(e_4)_K}) X_K + (\nabla_{\Omega_K(e_4)_K} - (\nabla_K)_{\Omega_K(e_4)_K}) X_K \\ &= (\nabla - \nabla_K)_{\partial_u} X_K \\ &= \check{J} \cdot \mathbf{O}_q^p.\end{aligned}$$

Combining with Definition 2.27, we deduce

$$\Omega \nabla_4 X_K = \mathbf{O}_{q+1}^p + \check{J} \cdot \mathbf{O}_q^p.$$

Finally, we have from Proposition 2.22 and Definition 2.27

$$\nabla X_K = \nabla_K X_K + (\nabla - \nabla_K) X_K = \mathbf{O}_{q+1}^p + \check{L} \cdot \mathbf{O}_q^p.$$

This concludes the proof of Lemma 2.31.  $\square$

**Corollary 2.32.** *We have the following schematic computation:*

$$\nabla(\mathbf{O}_q^p \cdot \Gamma_g) = \mathbf{O}_{q+1}^p \cdot \Gamma_g^{(1)}, \quad \nabla(\mathbf{O}_q^p \cdot \Gamma_b) = \mathbf{O}_{q+1}^p \cdot \Gamma_b^{(1)}.$$

*Proof.* We have from Lemma 2.31

$$\begin{aligned}
\nabla(\mathbf{O}_q^p \cdot \Gamma_i) &= \mathbf{O}_q^p \cdot \nabla \Gamma_i + \nabla \mathbf{O}_q^p \cdot \Gamma_i \\
&= \mathbf{O}_{q+1}^p \cdot \Gamma_i^{(1)} + (\mathbf{O}_{q+1}^p + \mathbf{O}_q^p \cdot \check{L}) \cdot \Gamma_i \\
&= \mathbf{O}_{q+1}^p \cdot \Gamma_i^{(1)},
\end{aligned}$$

for  $i = g, b$ . This concludes the proof of Corollary 2.32.  $\square$

**Remark 2.33.** In the remainder of this paper, Lemma 2.31 and Corollary 2.32 will be used frequently without explicit mention.

#### 2.8.4 The linearized null structure equations

In this section, we introduce the following linearized null structure equations, which play a fundamental role in Ricci coefficients estimates.

**Proposition 2.34.** We have the following linearized null structure equations:

$$\begin{aligned}
\nabla_4 \widetilde{\Omega \operatorname{tr} \chi} + \operatorname{tr} \chi \widetilde{\Omega \operatorname{tr} \chi} &= \mathbf{O}_1^0 \cdot \widetilde{\Omega \omega} + \mathfrak{L}[\nabla_4 \widetilde{\operatorname{tr} \chi}] + \operatorname{Err}[\nabla_4 \widetilde{\operatorname{tr} \chi}], \\
\mathfrak{L}[\nabla_4 \widetilde{\operatorname{tr} \chi}] &:= \Gamma_g \cdot \mathbf{O}_2^1, \\
\operatorname{Err}[\nabla_4 \widetilde{\operatorname{tr} \chi}] &:= \Gamma_g \cdot \Gamma_g, \\
\nabla_3 \widetilde{\Omega \operatorname{tr} \underline{\chi}} + \operatorname{tr} \underline{\chi} \widetilde{\Omega \operatorname{tr} \underline{\chi}} &= \mathbf{O}_1^0 \cdot \widetilde{\Omega \omega} + \mathfrak{L}[\nabla_3 \widetilde{\operatorname{tr} \underline{\chi}}] + \operatorname{Err}[\nabla_3 \widetilde{\operatorname{tr} \underline{\chi}}], \\
\mathfrak{L}[\nabla_3 \widetilde{\operatorname{tr} \underline{\chi}}] &:= \Gamma_g \cdot \mathbf{O}_2^1, \\
\operatorname{Err}[\nabla_3 \widetilde{\operatorname{tr} \underline{\chi}}] &:= \Gamma_b \cdot \Gamma_b, \\
\nabla_4 (\widetilde{\Omega \operatorname{tr} \underline{\chi}}) &= -2\Omega [\widetilde{\mu}] + \mathfrak{L}[\nabla_4 \widetilde{\operatorname{tr} \underline{\chi}}] + \operatorname{Err}[\nabla_4 \widetilde{\operatorname{tr} \underline{\chi}}], \\
\mathfrak{L}[\nabla_4 \widetilde{\operatorname{tr} \underline{\chi}}] &:= \Gamma_g \cdot \mathbf{O}_3^2, \\
\operatorname{Err}[\nabla_4 \widetilde{\operatorname{tr} \underline{\chi}}] &:= \Gamma_g \cdot \Gamma_g, \\
\nabla_3 (\widetilde{\Omega \operatorname{tr} \chi}) &= -2\Omega [\widetilde{\mu}] + \mathfrak{L}[\nabla_3 \widetilde{\operatorname{tr} \chi}] + \operatorname{Err}[\nabla_3 \widetilde{\operatorname{tr} \chi}], \\
\mathfrak{L}[\nabla_3 \widetilde{\operatorname{tr} \chi}] &:= \Gamma_g \cdot \mathbf{O}_3^2, \\
\operatorname{Err}[\nabla_3 \widetilde{\operatorname{tr} \chi}] &:= \Gamma_g \cdot \Gamma_g.
\end{aligned}$$

We also have the linearized Codazzi equations:

$$\begin{aligned}
\operatorname{div}(\check{\chi}) &= -\check{\beta} + \mathbf{O}_0^0 \cdot \nabla \widetilde{\operatorname{tr} \chi} + \mathbf{O}_1^0 \cdot \check{\zeta} + \mathfrak{L}[\operatorname{div}(\check{\chi})] + \operatorname{Err}[\operatorname{div}(\check{\chi})], \\
\mathfrak{L}[\operatorname{div}(\check{\chi})] &:= \Gamma_g \cdot \mathbf{O}_3^2 + \mathbf{O}_3^2 \cdot \check{L}, \\
\operatorname{Err}[\operatorname{div}(\check{\chi})] &:= \Gamma_g \cdot \Gamma_g, \\
\operatorname{div}(\check{\underline{\chi}}) &= \check{\underline{\beta}} + \mathbf{O}_0^0 \cdot \nabla \widetilde{\operatorname{tr} \underline{\chi}} + \mathbf{O}_1^0 \cdot \check{\zeta} + \mathfrak{L}[\operatorname{div}(\check{\underline{\chi}})] + \operatorname{Err}[\operatorname{div}(\check{\underline{\chi}})], \\
\mathfrak{L}[\operatorname{div}(\check{\underline{\chi}})] &:= \Gamma_b \cdot \mathbf{O}_3^2 + \mathbf{O}_3^2 \cdot \check{L}, \\
\operatorname{Err}[\operatorname{div}(\check{\underline{\chi}})] &:= \Gamma_b \cdot \Gamma_g.
\end{aligned}$$

The linearized mass aspect functions are given by:

$$\begin{aligned}\operatorname{div} \tilde{\eta} &= -\tilde{\mu} - \tilde{\rho} + \mathfrak{L}[\operatorname{div}(\tilde{\eta})] + \operatorname{Err}[\operatorname{div}(\tilde{\eta})], \\ \mathfrak{L}[\operatorname{div}(\tilde{\eta})] &= \Gamma_b \cdot \mathbf{O}_3^2 + \mathbf{O}_3^2 \cdot \check{L}, \\ \operatorname{Err}[\operatorname{div}(\tilde{\eta})] &= \Gamma_g \cdot \Gamma_b,\end{aligned}$$

$$\begin{aligned}\operatorname{div} \underline{\tilde{\eta}} &= -\underline{\tilde{\mu}} - \underline{\tilde{\rho}} + \mathfrak{L}[\operatorname{div}(\underline{\tilde{\eta}})] + \operatorname{Err}[\operatorname{div}(\underline{\tilde{\eta}})], \\ \mathfrak{L}[\operatorname{div}(\underline{\tilde{\eta}})] &= \Gamma_b \cdot \mathbf{O}_3^2 + \mathbf{O}_3^2 \cdot \check{L}, \\ \operatorname{Err}[\operatorname{div}(\underline{\tilde{\eta}})] &= \Gamma_g \cdot \Gamma_b.\end{aligned}$$

The linearized torsion equations are given by:

$$\begin{aligned}\operatorname{curl} \tilde{\eta} &= \check{\sigma} + \mathfrak{L}[\operatorname{curl} \tilde{\eta}] + \operatorname{Err}[\operatorname{curl} \tilde{\eta}], \\ \mathfrak{L}[\operatorname{curl} \tilde{\eta}] &:= \Gamma_b \cdot \mathbf{O}_3^2 + \mathbf{O}_3^2 \cdot \check{L}, \\ \operatorname{Err}[\operatorname{curl} \tilde{\eta}] &:= \Gamma_b \cdot \Gamma_g,\end{aligned}$$

$$\begin{aligned}\operatorname{curl} \underline{\tilde{\eta}} &= -\check{\sigma} + \mathfrak{L}[\operatorname{curl} \underline{\tilde{\eta}}] + \operatorname{Err}[\operatorname{curl} \underline{\tilde{\eta}}], \\ \mathfrak{L}[\operatorname{curl} \underline{\tilde{\eta}}] &:= \Gamma_b \cdot \mathbf{O}_3^2 + \mathbf{O}_3^2 \cdot \check{L}, \\ \operatorname{Err}[\operatorname{curl} \underline{\tilde{\eta}}] &:= \Gamma_b \cdot \Gamma_g.\end{aligned}$$

Moreover, we have

$$\begin{aligned}\Omega \nabla_4(\widetilde{\Omega \omega}) &= \frac{1}{2} \widetilde{\Omega^2 \rho} + \mathfrak{L}[\nabla_4 \check{\omega}] + \operatorname{Err}[\nabla_4 \check{\omega}], \\ \mathfrak{L}[\nabla_4 \check{\omega}] &:= \Gamma_g \cdot \mathbf{O}_2^1, \\ \operatorname{Err}[\nabla_4 \check{\omega}] &:= \Gamma_g \cdot \Gamma_g,\end{aligned}$$

$$\begin{aligned}\Omega \nabla_3(\widetilde{\Omega \omega}) &= \frac{1}{2} \widetilde{\Omega^2 \rho} + \mathfrak{L}[\nabla_3 \check{\omega}] + \operatorname{Err}[\nabla_3 \check{\omega}], \\ \mathfrak{L}[\nabla_3 \check{\omega}] &:= \Gamma_g \cdot \mathbf{O}_2^1, \\ \operatorname{Err}[\nabla_3 \check{\omega}] &:= \Gamma_g \cdot \Gamma_g.\end{aligned}$$

*Proof.* See Appendix A.1. □

**Remark 2.35.** In the linearized equations of Proposition 2.34, as well as in the other linearized equations below,  $\mathfrak{L}[\dots]$  contains linear terms which display more decay in  $\frac{M}{r}$  and  $\operatorname{Err}[\dots]$  contains the nonlinear terms.

**Proposition 2.36.** We have the following linearized equations:

$$\begin{aligned}\Omega \nabla_4[\widetilde{\mu}] + \Omega \operatorname{tr} \chi[\widetilde{\mu}] &:= \Omega \operatorname{tr} \chi \check{\rho} + \mathfrak{L}[\nabla_4[\widetilde{\mu}]] + \operatorname{Err}[\nabla_4[\widetilde{\mu}]], \\ \mathfrak{L}[\nabla_4[\widetilde{\mu}]] &:= \mathbf{O}_3^2 \cdot (d_2^* \tilde{\eta}, \nabla \operatorname{tr} \check{\chi}, \check{\beta}) + \mathbf{O}_3^1 \cdot (\Gamma_g, \check{L}), \\ \operatorname{Err}[\nabla_4[\widetilde{\mu}]] &:= -2\check{\chi} \cdot (d_2^* \tilde{\eta}) + (\tilde{\eta} - \underline{\tilde{\eta}}) \cdot \nabla \operatorname{tr} \check{\chi} - 2\check{\Omega} \tilde{\eta} \cdot \check{\beta} + r^{-1} \Gamma_g \cdot \check{L} + r^{-1} \Gamma_g \cdot \Gamma_b,\end{aligned}$$

$$\begin{aligned}\Omega \nabla_3[\widetilde{\mu}] + \Omega \operatorname{tr} \underline{\chi}[\widetilde{\mu}] &:= \Omega \operatorname{tr} \underline{\chi} \check{\rho} + \mathfrak{L}[\nabla_3[\widetilde{\mu}]] + \operatorname{Err}[\nabla_3[\widetilde{\mu}]], \\ \mathfrak{L}[\nabla_3[\widetilde{\mu}]] &:= \mathbf{O}_3^2 \cdot (d_2^* \tilde{\eta}, \nabla \operatorname{tr} \check{\chi}, \check{\beta}) + \mathbf{O}_3^1 \cdot (\Gamma_g, \check{L}), \\ \operatorname{Err}[\nabla_3[\widetilde{\mu}]] &:= -2\check{\chi} \cdot (d_2^* \tilde{\eta}) + (\tilde{\eta} - \underline{\tilde{\eta}}) \cdot \nabla \operatorname{tr} \check{\chi} + 2\check{\Omega} \underline{\tilde{\eta}} \cdot \check{\beta} + r^{-1} \Gamma_b \cdot \check{L} + r^{-1} \Gamma_b \cdot \Gamma_b.\end{aligned}$$

*Proof.* See Appendix A.2. □

### 2.8.5 The linearized Bianchi equations

In this section, we introduce the linearized Bianchi equations, which play a fundamental role in curvature estimates.

**Proposition 2.37.** *We have the following linearized Bianchi equations:*

$$\begin{aligned}\nabla_4 \check{\alpha} + \frac{1}{2} \operatorname{tr} \chi \check{\alpha} &= \nabla \widehat{\otimes} \check{\beta} + \mathfrak{L}[\nabla_4 \check{\alpha}] + \operatorname{Err}[\nabla_4 \check{\alpha}], \\ \mathfrak{L}[\nabla_4 \check{\alpha}] &:= \Gamma_b^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\alpha} + \mathbf{O}_3^2 \cdot (\check{\beta}, \check{\rho}, \check{\sigma}), \\ \operatorname{Err}[\nabla_4 \check{\alpha}] &:= \Gamma_g \cdot (\check{\alpha}, \check{\beta}) + \Gamma_b \cdot (\check{\rho}, \check{\sigma}),\end{aligned}$$

$$\begin{aligned}\nabla_3 \check{\beta} + 2 \operatorname{tr} \chi \check{\beta} &= -\operatorname{div} \check{\alpha} + \mathfrak{L}[\nabla_3 \check{\beta}] + \operatorname{Err}[\nabla_3 \check{\beta}], \\ \mathfrak{L}[\nabla_3 \check{\beta}] &:= \Gamma_b^{(1)} \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot \check{\beta} + \mathbf{O}_3^2 \cdot \check{\alpha}, \\ \operatorname{Err}[\nabla_3 \check{\beta}] &:= \Gamma_b \cdot \check{\beta} + \Gamma_g \cdot \check{\alpha},\end{aligned}$$

$$\begin{aligned}\nabla_4 \check{\beta} + \operatorname{tr} \chi \check{\beta} &= -\nabla \check{\rho} + {}^* \nabla \check{\sigma} + \mathfrak{L}[\nabla_4 \check{\beta}] + \operatorname{Err}[\nabla_4 \check{\beta}], \\ \mathfrak{L}[\nabla_4 \check{\beta}] &:= \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\beta} + \mathbf{O}_3^2 \cdot (\check{\rho}, \check{\sigma}, \check{\beta}), \\ \operatorname{Err}[\nabla_4 \check{\beta}] &:= \Gamma_g \cdot (\check{\beta}, \check{\rho}, \check{\sigma}) + \Gamma_b \cdot \check{\beta},\end{aligned}$$

$$\begin{aligned}\nabla_3(\check{\rho}, \check{\sigma}) + \frac{3}{2} \operatorname{tr} \chi(\check{\rho}, \check{\sigma}) &= -d_1 \check{\beta} + \mathfrak{L}[\nabla_3(\check{\rho}, \check{\sigma})] + \operatorname{Err}[\nabla_3(\check{\rho}, \check{\sigma})], \\ \mathfrak{L}[\nabla_3(\check{\rho}, \check{\sigma})] &:= \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\alpha}, \check{\beta}), \\ \operatorname{Err}[\nabla_3(\check{\rho}, \check{\sigma})] &:= \Gamma_g \cdot (\check{\alpha}, \check{\beta}),\end{aligned}$$

$$\begin{aligned}\nabla_4(\check{\rho}, -\check{\sigma}) + \frac{3}{2} \operatorname{tr} \chi(\check{\rho}, -\check{\sigma}) &= d_1 \check{\beta} + \mathfrak{L}[\nabla_4(\check{\rho}, -\check{\sigma})] + \operatorname{Err}[\nabla_4(\check{\rho}, -\check{\sigma})], \\ \mathfrak{L}[\nabla_4(\check{\rho}, -\check{\sigma})] &:= \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\alpha}, \check{\beta}), \\ \operatorname{Err}[\nabla_4(\check{\rho}, -\check{\sigma})] &:= \Gamma_b \cdot \check{\alpha} + \Gamma_g \cdot \check{\beta},\end{aligned}$$

$$\begin{aligned}\nabla_3 \check{\beta} + \operatorname{tr} \chi \check{\beta} &= \nabla \check{\rho} + {}^* \nabla \check{\sigma} + \mathfrak{L}[\nabla_3 \check{\beta}] + \operatorname{Err}[\nabla_3 \check{\beta}], \\ \mathfrak{L}[\nabla_3 \check{\beta}] &:= \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\beta} + \mathbf{O}_3^2 \cdot (\check{\beta}, \check{\rho}, \check{\sigma}), \\ \operatorname{Err}[\nabla_3 \check{\beta}] &:= \Gamma_b \cdot \check{\beta} + \Gamma_g \cdot (\check{\beta}, \check{\rho}, \check{\sigma}),\end{aligned}$$

$$\begin{aligned}\nabla_4 \check{\beta} + 2 \operatorname{tr} \chi \check{\beta} &= \operatorname{div} \check{\alpha} + \mathfrak{L}[\nabla_4 \check{\beta}] + \operatorname{Err}[\nabla_4 \check{\beta}], \\ \mathfrak{L}[\nabla_4 \check{\beta}] &:= \Gamma_g^{(1)} \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot (\check{\beta}, \check{\alpha}), \\ \operatorname{Err}[\nabla_4 \check{\beta}] &:= \Gamma_g \cdot (\check{\beta}, \check{\alpha}),\end{aligned}$$

$$\nabla_3 \check{\alpha} + \frac{1}{2} \operatorname{tr} \chi \check{\alpha} = \nabla \widehat{\otimes} \check{\beta} + \mathfrak{L}[\nabla_3 \check{\alpha}] + \operatorname{Err}[\nabla_3 \check{\alpha}],$$

$$\begin{aligned}\mathfrak{L}[\nabla_3 \check{\alpha}] &:= \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\alpha} + \mathbf{O}_3^2 \cdot (\check{\rho}, \check{\sigma}, \check{\beta}), \\ \text{Err}[\nabla_3 \check{\alpha}] &:= \Gamma_b \cdot \check{\alpha} + \Gamma_g \cdot \check{\beta}.\end{aligned}$$

*Proof.* See Appendix A.3. □

**Proposition 2.38.** *We have the following linearized equations:*<sup>15</sup>

$$\begin{aligned}\nabla_4 \check{\alpha} + \frac{1}{2} \text{tr} \chi \check{\alpha} &= -\nabla \widehat{\otimes} \check{\beta} + \mathfrak{L}[\nabla_4 \check{\alpha}] + \text{Err}[\nabla_4 \check{\alpha}], \\ \mathfrak{L}[\nabla_4 \check{\alpha}] &= \Gamma_b^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\alpha}^{(1)} + \mathbf{O}_3^2 \cdot (\check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}), \\ \text{Err}[\nabla_4 \check{\alpha}] &= \Gamma_g^{(1)} \cdot (\check{\alpha}, \check{\beta}) + \Gamma_b^{(1)} \cdot (\check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\alpha}^{(1)}, \check{\beta}^{(1)}) + \Gamma_b \cdot (\check{\rho}^{(1)}, \check{\sigma}^{(1)}), \\ \nabla_3 \check{\beta} + 2 \text{tr} \chi \check{\beta} &= -\text{div} \check{\alpha} + \mathfrak{L}[\nabla_3 \check{\beta}] + \text{Err}[\nabla_3 \check{\beta}], \\ \mathfrak{L}[\nabla_3 \check{\beta}] &:= \Gamma_b^{(1)} \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot \check{\beta}^{(1)} + \mathbf{O}_3^2 \cdot \check{\alpha}^{(1)}, \\ \text{Err}[\nabla_3 \check{\beta}] &:= \Gamma_b^{(1)} \cdot \check{\beta} + \Gamma_b \cdot \check{\beta}^{(1)} + \Gamma_g^{(1)} \cdot \check{\alpha} + \Gamma_g \cdot \check{\alpha}^{(1)}, \\ \nabla_4 \check{\beta} + \text{tr} \chi \check{\beta} &:= -\nabla \check{\rho} + * \nabla \check{\sigma} + \mathfrak{L}[\nabla_4 \check{\beta}] + \text{Err}[\nabla_4 \check{\beta}], \\ \mathfrak{L}[\nabla_4 \check{\beta}] &:= \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\beta}^{(1)} + \mathbf{O}_3^2 \cdot (\check{\rho}^{(1)}, \check{\sigma}^{(1)}, \check{\beta}^{(1)}), \\ \text{Err}[\nabla_4 \check{\beta}] &:= \Gamma_g^{(1)} \cdot (\check{\beta}, \check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}) + \Gamma_b^{(1)} \cdot \check{\beta} + \Gamma_b \cdot \check{\beta}^{(1)}, \\ \nabla_3(\check{\rho}, \check{\sigma}) + \frac{3}{2} \text{tr} \chi(\check{\rho}, \check{\sigma}) &= -d_1 \check{\beta} + \mathfrak{L}[\nabla_3(\check{\rho}, \check{\sigma})] + \text{Err}[\nabla_3(\check{\rho}, \check{\sigma})], \\ \mathfrak{L}[\nabla_3(\check{\rho}, \check{\sigma})] &:= \Gamma_b^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\alpha}^{(1)}, \check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}), \\ \text{Err}[\nabla_3(\check{\rho}, \check{\sigma})] &:= \Gamma_g^{(1)} \cdot (\check{\alpha}, \check{\beta}) + \Gamma_g \cdot (\check{\alpha}^{(1)}, \check{\beta}^{(1)}) + \Gamma_b \cdot (\check{\rho}^{(1)}, \check{\sigma}^{(1)}) + \Gamma_b^{(1)} \cdot (\check{\rho}, \check{\sigma}), \\ \nabla_4(\check{\rho}, -\check{\sigma}) + \frac{3}{2} \text{tr} \chi(\check{\rho}, -\check{\sigma}) &= d_1 \check{\beta} + \mathfrak{L}[\nabla_4(\check{\rho}, -\check{\sigma})] + \text{Err}[\nabla_4(\check{\rho}, -\check{\sigma})], \\ \mathfrak{L}[\nabla_4(\check{\rho}, -\check{\sigma})] &:= \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\alpha}^{(1)}, \check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}), \\ \text{Err}[\nabla_4(\check{\rho}, -\check{\sigma})] &:= \Gamma_b^{(1)} \cdot \check{\alpha} + \Gamma_b \cdot \check{\alpha}^{(1)} + \Gamma_g^{(1)} \cdot (\check{\beta}, \check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}), \\ \nabla_3 \check{\beta} + \text{tr} \chi \check{\beta} &= \nabla \check{\rho} + * \nabla \check{\sigma} + \mathfrak{L}[\nabla_3 \check{\beta}] + \text{Err}[\nabla_3 \check{\beta}], \\ \mathfrak{L}[\nabla_3 \check{\beta}] &:= \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\beta}^{(1)} + \mathbf{O}_3^2 \cdot (\check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}), \\ \text{Err}[\nabla_3 \check{\beta}] &:= \Gamma_b^{(1)} \cdot \check{\beta} + \Gamma_b \cdot \check{\beta}^{(1)} + \Gamma_g^{(1)} \cdot (\check{\beta}, \check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}), \\ \nabla_4 \check{\beta} + 2 \text{tr} \chi \check{\beta} &= \text{div} \check{\alpha} + \mathfrak{L}[\nabla_4 \check{\beta}] + \text{Err}[\nabla_4 \check{\beta}], \\ \mathfrak{L}[\nabla_4 \check{\beta}] &:= \Gamma_g^{(1)} \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot (\check{\beta}^{(1)}, \check{\alpha}^{(1)}), \\ \text{Err}[\nabla_4 \check{\beta}] &:= \Gamma_g^{(1)} \cdot (\check{\beta}, \check{\alpha}) + \Gamma_g \cdot (\check{\beta}^{(1)}, \check{\alpha}^{(1)}),\end{aligned}$$

---

<sup>15</sup>Recall that  $\check{\rho}$  is introduced in Definition 2.13.

$$\begin{aligned}
\nabla_3 \mathfrak{P}\check{\alpha} + \frac{1}{2} \operatorname{tr} \chi \mathfrak{P}\check{\alpha} &= \nabla \widehat{\otimes} \mathfrak{P}\check{\beta} + \mathfrak{L}[\nabla_3 \widetilde{\mathfrak{P}\check{\alpha}}] + \operatorname{Err}[\nabla_3 \widetilde{\mathfrak{P}\check{\alpha}}], \\
\mathfrak{L}[\nabla_3 \widetilde{\mathfrak{P}\check{\alpha}}] &:= \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\alpha}^{(1)} + \mathbf{O}_3^2 \cdot (\check{\rho}^{(1)}, \check{\sigma}^{(1)}, \check{\beta}^{(1)}), \\
\operatorname{Err}[\nabla_3 \widetilde{\mathfrak{P}\check{\alpha}}] &:= \Gamma_b^{(1)} \cdot \check{\alpha} + \Gamma_b \cdot \check{\alpha}^{(1)} + \Gamma_g^{(1)} \cdot \check{\beta} + \Gamma_g \cdot \check{\beta}^{(1)}.
\end{aligned}$$

**Remark 2.39.** Notice that Proposition 2.38 involves the quantities  $\widetilde{\mathfrak{P}\check{\alpha}}$ ,  $\widetilde{\mathfrak{P}\check{\beta}}$ ,  $\widetilde{\mathfrak{P}\check{\rho}}$ ,  $\widetilde{\mathfrak{P}\check{\sigma}}$ ,  $\widetilde{\mathfrak{P}\check{\beta}}$  and  $\widetilde{\mathfrak{P}\check{\alpha}}$  instead of  $\mathfrak{P}\check{\alpha}$ ,  $\mathfrak{P}\check{\beta}$ ,  $\mathfrak{P}\check{\rho}$ ,  $\mathfrak{P}\check{\sigma}$ ,  $\mathfrak{P}\check{\beta}$  and  $\mathfrak{P}\check{\alpha}$ . This allows to avoid a loss of a derivative of  $\check{J}$ ,  $\check{J}$  and  $\check{L}$  in the linearization procedure. Indeed, we have for example

$$\widetilde{\mathfrak{P}\check{\beta}} = \mathfrak{P}\check{\beta} + (\check{J}, \check{J}, \check{L}) \cdot \beta_K,$$

and the norm  $\mathcal{O}_\Gamma$  in Section 3.1.2 only controls  $(\check{J}, \check{J}, \check{L})$  but not its first order derivatives.

*Proof.* See Appendix A.3. □

### 2.8.6 Linearized equations for the metric components

**Proposition 2.40.** We have the following transport equations:

$$\begin{aligned}
\nabla_4(\check{\gamma}_{AB}) - \operatorname{tr} \chi(\check{\gamma}_{AB}) &= \mathbf{O}_{-2}^0 \cdot (\check{\chi}, \widetilde{\operatorname{tr} \chi}) + O(r)\check{\Omega} + r^3 \Gamma_g \cdot \Gamma_g, \\
\nabla_4(\check{\Omega}) &= \mathbf{O}_0^0 \cdot \check{\Omega} \omega + \mathbf{O}_1^1 \cdot \Gamma_g + r \Gamma_g \cdot \Gamma_g, \\
\nabla_4(\check{b}^A) &= \mathbf{O}_1^0 \cdot \check{\zeta} + \mathbf{O}_3^2 \cdot \Gamma_g + \Gamma_g \cdot \Gamma_g, \\
\nabla_4(\check{\epsilon}_{AB}) - \operatorname{tr} \chi(\check{\epsilon}_{AB}) &= \mathbf{O}_{-2}^0 \cdot \widetilde{\operatorname{tr} \chi} + r^3 \Gamma_g \cdot \Gamma_g.
\end{aligned}$$

Moreover, we have the following transport equations in the case  $\check{\Omega} = 0$  and  $\check{b} = 0$ .<sup>16</sup>

$$\begin{aligned}
\nabla_3(\check{\gamma}_{AB}) - \operatorname{tr} \chi(\check{\gamma}_{AB}) &= \mathbf{O}_{-2}^0 \cdot (\widetilde{\operatorname{tr} \chi}, \check{\chi}) + \mathbf{O}_3^2 \cdot \check{\gamma}_{AB}, \\
\nabla_3(\check{\epsilon}_{AB}) - \operatorname{tr} \chi(\check{\epsilon}_{AB}) &= \mathbf{O}_{-2}^0 \cdot \widetilde{\operatorname{tr} \chi} + \mathbf{O}_3^2 \cdot \check{\epsilon}_{AB}.
\end{aligned}$$

*Proof.* See Appendix A.4. □

## 2.9 Commutator identities

We recall the following commutator identities for scalar functions, see Proposition 4.8.1 of [14].

**Proposition 2.41.** For a scalar function  $f$ , we have

$$\begin{aligned}
[\Omega \nabla_4, \nabla] f &= -\Omega \chi \cdot \nabla f, \\
[\Omega \nabla_3, \nabla] f &= -\Omega \underline{\chi} \cdot \nabla f, \\
[\nabla_4, \nabla_3] f &= 2\omega \nabla_3 f - 2\underline{\omega} \nabla_4 f - 4\zeta \cdot \nabla f.
\end{aligned}$$

We also need the commutation identities for more general tensor fields. For this, we record the following commutation lemma.

---

<sup>16</sup>The conditions  $\check{\Omega} = 0$  and  $\check{b} = 0$  hold on  $\underline{C}_*$ , see Definition 6.1.

**Lemma 2.42.** *Let  $U_{a_1 \dots a_k}$  be an  $S$ -tangent  $k$ -covariant tensor on  $(\mathcal{M}, \mathbf{g})$ . Then*

$$\begin{aligned}
[\nabla_4, \nabla_b]U_{a_1 \dots a_k} &= -\chi_{BC} \nabla_c U_{a_1 \dots a_k} + F_{4ba_1 \dots a_k}, \\
F_{4ba_1 \dots a_k} &:= (\zeta_b + \underline{\eta}_b) \nabla_4 U_{a_1 \dots a_k} + \sum_{i=1}^k (\chi_{a_i b} \underline{\eta}_c - \chi_{BC} \underline{\eta}_{a_i} + \epsilon_{a_i c} {}^* \beta_b) U_{a_1 \dots c \dots a_k}, \\
[\nabla_3, \nabla_b]U_{a_1 \dots a_k} &= -\underline{\chi}_{BC} \nabla_c U_{a_1 \dots a_k} + F_{3ba_1 \dots a_k}, \\
F_{3ba_1 \dots a_k} &:= (\eta_b - \zeta_b) \nabla_3 U_{a_1 \dots a_k} + \sum_{i=1}^k (\underline{\chi}_{a_i b} \eta_c - \underline{\chi}_{BC} \eta_{a_i} + \epsilon_{a_i c} {}^* \underline{\beta}_b) U_{a_1 \dots c \dots a_k}, \\
[\nabla_3, \nabla_4]U_{a_1 \dots a_k} &= F_{34a_1 \dots a_k}, \\
F_{34a_1 \dots a_k} &:= -2\omega \nabla_3 U + 2\underline{\omega} \nabla_4 U + 4\zeta_b \nabla_b U_{a_1 \dots a_k} + 2 \sum_{i=1}^k (\underline{\eta}_{a_i} \eta_c - \underline{\eta}_{a_i} \eta_c + \epsilon_{a_i c} \sigma) U_{a_1 \dots c \dots a_k}.
\end{aligned}$$

*Proof.* It is a direct consequence of Lemma 7.3.3 in [5].  $\square$

**Corollary 2.43.** *We have the following schematic commutation identities:*

$$\begin{aligned}
[r\nabla, \Omega \nabla_4] &= \Gamma_g \cdot r\nabla + \mathbf{O}_3^2 \cdot r\nabla + \Gamma_g^{(1)} + \mathbf{O}_3^2, \\
[r\nabla, \Omega \nabla_3] &= \Gamma_b \cdot r\nabla + \mathbf{O}_3^2 \cdot r\nabla + \Gamma_b^{(1)} + \mathbf{O}_3^2, \\
[\nabla, \Omega \nabla_4] &= -\frac{1}{2} \Omega \operatorname{tr} \chi \nabla + \Gamma_g \cdot \nabla + \mathbf{O}_3^2 \cdot \nabla + r^{-1} \Gamma_g^{(1)} + \mathbf{O}_4^2, \\
[\nabla, \Omega \nabla_3] &= -\frac{1}{2} \Omega \operatorname{tr} \underline{\chi} \nabla + \Gamma_b \cdot \nabla + \mathbf{O}_3^2 \cdot \nabla + r^{-1} \Gamma_b^{(1)} + \mathbf{O}_4^2, \\
[\nabla_4, \nabla_3] &= \Gamma_g \cdot \nabla_3 + \Gamma_b \cdot (\nabla_4, \nabla) + \mathbf{O}_2^1 \cdot (\nabla_3, \nabla_4, \nabla) + r^{-1} \Gamma_g^{(1)} + \mathbf{O}_4^2, \\
[\Omega \nabla_4, \Omega \nabla_3] &= \Gamma_b \cdot \nabla + \mathbf{O}_3^2 \cdot \nabla + r^{-1} \Gamma_g^{(1)} + \mathbf{O}_4^2.
\end{aligned}$$

*Proof.* It follows directly from Lemmas 2.42 and 2.30 together with the identities (2.8).  $\square$

## 2.10 Sobolev inequalities

**Definition 2.44.** *For a scalar function  $h$  on a null cone  $C \in \{C_u, \underline{C}_u\}$ , we define its integral on  $C$  as follows:<sup>17</sup>*

$$\int_{C_u} h := \int d\underline{u} \int_{S(u, \underline{u})} \Omega^2 h, \quad \int_{\underline{C}_u} h := \int du \int_{S(u, \underline{u})} \Omega^2 h.$$

*For a tensor field  $h$  on a null cone  $C$ , we define its  $L^2$ -flux:*

$$\|h\|_{2,C} := \left( \int_C |h|^2 \right)^{\frac{1}{2}}. \quad (2.42)$$

We recall the following Sobolev inequalities.

**Proposition 2.45.** *Let  $F$  be a tensor field tangent to  $S := S(u, \underline{u})$  at every point. We have the following estimates:*

$$\begin{aligned}
|rF|_{4,S} &\lesssim |rF|_{4,S(u, \underline{u}_0(u))} + \|F\|_{2,C_u \cap V(u, \underline{u})} + \|r\nabla F\|_{2,C_u \cap V(u, \underline{u})} + \|r\nabla_4 F\|_{2,C_u \cap V(u, \underline{u})}, \\
|r^{\frac{1}{2}}|u|^{\frac{1}{2}}F|_{4,S} &\lesssim |rF|_{4,S(u_0(\underline{u}), \underline{u})} + \|F\|_{2,\underline{C}_u \cap V(u, \underline{u})} + \|r\nabla F\|_{2,\underline{C}_u \cap V(u, \underline{u})} + |u| \|\nabla_3 F\|_{2,\underline{C}_u \cap V(u, \underline{u})},
\end{aligned}$$

<sup>17</sup>The factor  $\Omega^2$  has been added since we have from Section 2.1.1 that  $\partial_{\underline{u}} = 2\Omega^2 L$  and  $\partial_u + \underline{b}^A \partial_A = 2\Omega^2 \underline{L}$ , where  $L$  and  $\underline{L}$  denote the generators of  $C_u$  and  $\underline{C}_u$  respectively.

where  $S(u, \underline{u}_0(u))$  (resp.  $S(u_0(\underline{u}), \underline{u})$ ) denotes the unique leaf of  $C_u$  (resp.  $\underline{C}_{\underline{u}}$ ), which located in the future of  $\Sigma_0$  and touches  $\Sigma_0 \setminus K$ .

*Proof.* See Corollary 3.2.1.1 in [5] and Corollary 4.1.1 in [14].  $\square$

We also need the following standard Sobolev inequalities, see Lemma 4.1.3 in [14].

**Proposition 2.46.** *Let  $F$  be a tensor field tangent to the sphere  $S := S(u, \underline{u})$ . Then, we have*

$$\sup_S r^{\frac{1}{2}} |F| \lesssim \left( \int_S |F|^4 + |r \nabla F|^4 \right)^{\frac{1}{4}}.$$

### 3 Main theorem

#### 3.1 Fundamental norms

Our result holds for initial data sets satisfying (1.9) with  $s > 3$ . We will focus on the case  $s \in [4, 6]$  in Sections 3–6, and we postpone the necessary adaptations to the case  $s \in (3, 4)$  and  $s > 6$  to Appendices B and C. The norms in Sections 3.1.1–3.1.4 are defined with respect to the foliation  $(u, \underline{u})$  in the bootstrap region  $\mathcal{K}$  while the norms in Section 3.1.5 are defined with respect to the foliation  $(u_{(0)}, \underline{u})$  in the initial data layer  $\mathcal{K}_{(0)}$ .

##### 3.1.1 $\mathcal{R}$ norms (Curvature components)

We define

$$\mathcal{R}_0^S := \mathcal{R}_0^S[\check{\beta}], \quad \underline{\mathcal{R}}_0^S := \underline{\mathcal{R}}_0^S[\check{\beta}] + \underline{\mathcal{R}}_0^S[\check{\rho}, \check{\sigma}] + \underline{\mathcal{R}}_0^S[\check{\beta}],$$

where

$$\begin{aligned} \mathcal{R}_0^S[\check{\beta}] &:= \sup_{\mathcal{K}} \sup_{p \in [2, 4]} |r^{\frac{7}{2} - \frac{2}{p}} |u|^{\frac{s-4}{2}} \check{\beta}|_{p, S(u, \underline{u})}, \\ \underline{\mathcal{R}}_0^S[\check{\beta}] &:= \sup_{\mathcal{K}} \sup_{p \in [2, 4]} |r^{\frac{s+2}{2} - \frac{2}{p}} |u|^{\frac{1}{2}} \check{\beta}|_{p, S(u, \underline{u})}, \\ \underline{\mathcal{R}}_0^S[\check{\rho}, \check{\sigma}] &:= \sup_{\mathcal{K}} \sup_{p \in [2, 4]} |r^{3 - \frac{2}{p}} |u|^{\frac{s-3}{2}} (\check{\rho}, \check{\sigma})|_{p, S(u, \underline{u})}, \\ \underline{\mathcal{R}}_0^S[\check{\beta}] &:= \sup_{\mathcal{K}} \sup_{p \in [2, 4]} |r^{2 - \frac{2}{p}} |u|^{\frac{s-1}{2}} \check{\beta}|_{p, S(u, \underline{u})}. \end{aligned}$$

Then we define the norms of flux of curvature components. We denote

$$\mathcal{R} := \sum_{q=0}^1 (\mathcal{R}_q + \underline{\mathcal{R}}_q) + \mathcal{R}_0^S + \underline{\mathcal{R}}_0^S,$$

where

$$\begin{aligned} \mathcal{R}_q &:= \left( \mathcal{R}_q[\check{\alpha}]^2 + \mathcal{R}_q[\check{\beta}]^2 + \mathcal{R}_q[(\check{\rho}, \check{\sigma})]^2 + \mathcal{R}_q[\check{\beta}]^2 \right)^{\frac{1}{2}}, \\ \underline{\mathcal{R}}_q &:= \left( \underline{\mathcal{R}}_q[\check{\beta}]^2 + \underline{\mathcal{R}}_q[(\check{\rho}, \check{\sigma})]^2 + \underline{\mathcal{R}}_q[\check{\beta}]^2 + \underline{\mathcal{R}}_q[\check{\alpha}]^2 \right)^{\frac{1}{2}}, \end{aligned}$$

with

$$\begin{aligned} \mathcal{R}_q[w] &:= \sup_{\mathcal{K}} \mathcal{R}_q[w](u, \underline{u}), \\ \underline{\mathcal{R}}_q[w] &:= \sup_{\mathcal{K}} \underline{\mathcal{R}}_q[w](u, \underline{u}). \end{aligned}$$

We denote

$$V := V(u, \underline{u}), \quad C_u^V := C_u \cap V, \quad \underline{C}_{\underline{u}}^V := \underline{C}_{\underline{u}} \cap V. \quad (3.1)$$

It remains to define  $\mathcal{R}_q[w](u, \underline{u})$  and  $\underline{\mathcal{R}}_q[w](u, \underline{u})$ :

$$\begin{aligned} \mathcal{R}_q[\check{\alpha}](u, \underline{u}) &:= \|r^{\frac{s}{2}}(r\nabla)^q \check{\alpha}\|_{2, C_u^V}, \\ \mathcal{R}_q[\check{\beta}](u, \underline{u}) &:= \|r^2|u|^{\frac{s-4}{2}}(r\nabla)^q \check{\beta}\|_{2, C_u^V}, \\ \mathcal{R}_q[(\check{\rho}, \check{\sigma})](u, \underline{u}) &:= \|r|u|^{\frac{s-2}{2}}(r\nabla)^q (\check{\rho}, \check{\sigma})\|_{2, C_u^V}, \\ \mathcal{R}_q[\check{\beta}](u, \underline{u}) &:= \| |u|^{\frac{s}{2}}(r\nabla)^q \check{\beta}\|_{2, C_u^V}, \\ \underline{\mathcal{R}}_q[\check{\beta}](u, \underline{u}) &:= \|r^{\frac{s}{2}}(r\nabla)^q \check{\beta}\|_{2, \underline{C}_{\underline{u}}^V}, \\ \underline{\mathcal{R}}_q[(\check{\rho}, \check{\sigma})](u, \underline{u}) &:= |u|^{\frac{s-4}{2}} \|r^2(r\nabla)^q (\check{\rho}, \check{\sigma})\|_{2, \underline{C}_{\underline{u}}^V}, \\ \underline{\mathcal{R}}_q[\check{\beta}](u, \underline{u}) &:= |u|^{\frac{s-2}{2}} \|r(r\nabla)^q \check{\beta}\|_{2, \underline{C}_{\underline{u}}^V}, \\ \underline{\mathcal{R}}_q[\check{\alpha}](u, \underline{u}) &:= |u|^{\frac{s}{2}} \|(r\nabla)^q \check{\alpha}\|_{2, \underline{C}_{\underline{u}}^V}. \end{aligned}$$

### 3.1.2 $\mathcal{O}$ norms (Ricci coefficients and metric components)

We denote for any linearized quantity:

$$\mathcal{O}_q(\check{X}) := \sup_{\mathcal{K}} \sup_{p \in [2, 4]} \mathcal{O}_q^{p, S}(\check{X})(u, \underline{u}). \quad (3.2)$$

We first define

$$\mathcal{O}_{[1]} := \mathcal{O}_1 + \mathcal{O}_0 + \underline{\mathcal{O}}_1 + \underline{\mathcal{O}}_0 + \sum_{q=0}^2 \mathcal{O}_q(\check{\Omega}),$$

where

$$\begin{aligned} \mathcal{O}_q &:= \mathcal{O}_q(\widetilde{\Omega \operatorname{tr} \chi}) + \mathcal{O}_q(\check{\chi}) + \mathcal{O}_q(\check{\eta}) + \mathcal{O}_q(\check{\Omega \omega}), \quad q = 0, 1, \\ \underline{\mathcal{O}}_q &:= \mathcal{O}_q(\widetilde{\Omega \operatorname{tr} \underline{\chi}}) + \mathcal{O}_q(\check{\underline{\chi}}) + \mathcal{O}_q(\check{\underline{\eta}}) + \mathcal{O}_q(\check{\Omega \omega}), \quad q = 0, 1. \end{aligned}$$

It remains to define  $\mathcal{O}_q^{p, S}(\check{X})(u, \underline{u})$ :

$$\begin{aligned} \mathcal{O}_q^{p, S}(\widetilde{\Omega \operatorname{tr} \chi})(u, \underline{u}) &:= |r^{2+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q (\widetilde{\Omega \operatorname{tr} \chi})|_{p, S(u, \underline{u})}, \\ \mathcal{O}_q^{p, S}(\widetilde{\Omega \operatorname{tr} \underline{\chi}})(u, \underline{u}) &:= |r^{2+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q (\widetilde{\Omega \operatorname{tr} \underline{\chi}})|_{p, S(u, \underline{u})}, \\ \mathcal{O}_q^{p, S}(\check{\chi})(u, \underline{u}) &:= |r^{2+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \check{\chi}|_{p, S(u, \underline{u})}, \\ \mathcal{O}_q^{p, S}(\check{\underline{\chi}})(u, \underline{u}) &:= |r^{1+q-\frac{2}{p}}|u|^{\frac{s-1}{2}} \nabla^q \check{\underline{\chi}}|_{p, S(u, \underline{u})}, \\ \mathcal{O}_q^{p, S}(\check{\eta})(u, \underline{u}) &:= |r^{2+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \check{\eta}|_{p, S(u, \underline{u})}, \\ \mathcal{O}_q^{p, S}(\check{\underline{\eta}})(u, \underline{u}) &:= |r^{2+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \check{\underline{\eta}}|_{p, S(u, \underline{u})}, \\ \mathcal{O}_q^{p, S}(\check{\Omega \omega})(u, \underline{u}) &:= |r^{2+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \check{\Omega \omega}|_{p, S(u, \underline{u})}, \\ \mathcal{O}_q^{p, S}(\check{\Omega \omega})(u, \underline{u}) &:= |r^{1+q-\frac{2}{p}}|u|^{\frac{s-1}{2}} \nabla^q \check{\Omega \omega}|_{p, S(u, \underline{u})}, \\ \mathcal{O}_q^{p, S}(\check{\Omega})(u, \underline{u}) &:= |r^{1+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \check{\Omega}|_{p, S(u, \underline{u})}. \end{aligned}$$

Next, we define the norms of metric components:

$$\mathcal{O}_\gamma := \sum_{q=0}^1 \left( \mathcal{O}_q(\check{\gamma}) + \mathcal{O}_q(\check{\epsilon}) + \mathcal{O}_q(\check{b}) \right),$$

where

$$\begin{aligned} \mathcal{O}_q^{p,S}(\check{\gamma})(u, \underline{u}) &:= |r^{-1+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \check{\gamma}_{AB}|_{p,S(u,\underline{u})}, \\ \mathcal{O}_q^{p,S}(\check{\epsilon})(u, \underline{u}) &:= |r^{-1+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \check{\epsilon}_{AB}|_{p,S(u,\underline{u})}, \\ \mathcal{O}_q^{p,S}(\check{b})(u, \underline{u}) &:= |r^{2+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \check{b}^A|_{p,S(u,\underline{u})}. \end{aligned}$$

We also define

$$\mathcal{O}_\Gamma := \mathcal{O}_0(\check{J}) + \mathcal{O}_0(\check{\underline{J}}) + \mathcal{O}_0(\check{\underline{L}}),$$

where

$$\begin{aligned} \mathcal{O}_0^{p,S}(\check{\underline{L}})(u, \underline{u}) &:= |r^{2-\frac{2}{p}}|u|^{\frac{s-3}{2}} \check{\underline{L}}|_{p,S(u,\underline{u})}, \\ \mathcal{O}_0^{p,S}(\check{J})(u, \underline{u}) &:= |r^{2-\frac{2}{p}}|u|^{\frac{s-3}{2}} \check{J}|_{p,S(u,\underline{u})}, \\ \mathcal{O}_0^{p,S}(\check{\underline{J}})(u, \underline{u}) &:= |r^{1-\frac{2}{p}}|u|^{\frac{s-1}{2}} \check{\underline{J}}|_{p,S(u,\underline{u})}. \end{aligned}$$

Finally, we denote

$$\mathcal{O} := \mathcal{O}_{[1]} + \mathcal{O}_\gamma + \mathcal{O}_\Gamma. \quad (3.3)$$

### 3.1.3 $\mathring{\mathcal{O}}(\Sigma_0)$ norms and $\mathfrak{R}_0$ norms (Initial data for the foliation of $\mathcal{K}$ )

Notice that for every  $\underline{u}$ , there exists a unique leaf  $S(u_0(\underline{u}), \underline{u})$  of  $\underline{C}_{\underline{u}}$ , which located in the future of  $\Sigma_0$  and touches  $\Sigma_0 \setminus K$ . Moreover, we have

$$S(u_0(\underline{u}), \underline{u}) \subset \mathcal{K}_{(0)}.$$

We define the following norms on the union of spheres  $S(u_0(\underline{u}), \underline{u})$ :

$$\mathring{\mathcal{O}}_q(\check{\Gamma}) := \sup_{p \in [2,4]} \sup_{\Sigma_0 \setminus K} \mathring{\mathcal{O}}_q^{p,S}(\check{\Gamma})(\underline{u}), \quad q = 0, 1.$$

We denote

$$\mathring{\mathcal{O}}(\Sigma_0) := \mathring{\mathcal{O}}_1 + \mathring{\mathcal{O}}_0,$$

where

$$\mathring{\mathcal{O}}_q := \mathring{\mathcal{O}}_q(\widetilde{\Omega \operatorname{tr} \underline{\chi}}) + \mathring{\mathcal{O}}_q(\widetilde{\Omega \omega}) + \mathring{\mathcal{O}}_q(\check{\eta}), \quad q = 0, 1.$$

It remains to define  $\mathring{\mathcal{O}}_q^{p,S}(\check{\Gamma})(\underline{u})$ :

$$\begin{aligned} \mathring{\mathcal{O}}_q^{p,S}(\widetilde{\Omega \operatorname{tr} \underline{\chi}})(\underline{u}) &:= |r^{\frac{s+1}{2}+q-\frac{2}{p}} \nabla^q (\widetilde{\Omega \operatorname{tr} \underline{\chi}})|_{p,S(u_0(\underline{u}), \underline{u})}, \\ \mathring{\mathcal{O}}_q^{p,S}(\check{\eta})(\underline{u}) &:= |r^{\frac{s+1}{2}+q-\frac{2}{p}} \nabla^q \check{\eta}|_{p,S(u_0(\underline{u}), \underline{u})}, \\ \mathring{\mathcal{O}}_q^{p,S}(\widetilde{\Omega \omega})(\underline{u}) &:= |r^{\frac{s+1}{2}+q-\frac{2}{p}} \nabla^q (\widetilde{\Omega \omega})|_{p,S(u_0(\underline{u}), \underline{u})}. \end{aligned}$$

We can extend the foliation  $(u, \underline{u})$  to a neighborhood of  $\Sigma_0 \setminus K$  in  $J^-(\Sigma_0 \setminus K)$  such that it is well defined on  $\Sigma_0 \setminus K$ . The curvature flux on  $\Sigma_0 \setminus K$  is defined by:

$$\mathfrak{R}_0^2 := \int_{\Sigma_0 \setminus K} \sum_{l=0}^1 r^s \left( |\mathfrak{d}^l \check{\alpha}|^2 + |\mathfrak{d}^l \check{\beta}|^2 + |\mathfrak{d}^l(\check{\rho}, \check{\sigma})|^2 + |\mathfrak{d}^l \check{\underline{\beta}}|^2 + |\mathfrak{d}^l \check{\underline{\alpha}}|^2 \right),$$

where

$$\mathfrak{d} := (r \nabla, r \nabla_4, r \nabla_3).$$

### 3.1.4 $\mathcal{O}^*(\underline{C}_*)$ norms (Ricci coefficients and metric components on $\underline{C}_*$ )

For any linearized quantity  $\check{\Gamma}$  on  $\underline{C}_*$ , we define

$$\mathcal{O}_q^*(\check{\Gamma}) := \sup_{p \in [2,4]} \sup_{\underline{C}_*} \mathcal{O}_q^{*p,S}(\check{\Gamma})(u).$$

We denote

$$\mathcal{O}^*(\underline{C}_*) := \mathcal{O}_1^* + \mathcal{O}_0^* + \mathcal{O}_\gamma^* + \mathcal{O}_\Gamma^* + \mathcal{O}_2^*(\widetilde{\text{tr}} \chi) + \mathcal{O}_1^*(\widetilde{[\mu]}),$$

where

$$\begin{aligned} \mathcal{O}_\gamma^* &:= \sum_{q=0}^2 (\mathcal{O}_q^*(\check{\gamma}) + \mathcal{O}_q^*(\check{\epsilon})), \\ \mathcal{O}_\Gamma^* &:= \sum_{q=0}^1 (\mathcal{O}_q^*(\check{L}) + \mathcal{O}_q^*(\check{J})), \\ \mathcal{O}_q^* &:= \mathcal{O}_q^*(\widetilde{\text{tr}} \chi) + \mathcal{O}_q^*(\check{\chi}) + \mathcal{O}_q^*(\check{\zeta}) + \mathcal{O}_q^*(\widetilde{\text{tr}} \underline{\chi}) + \mathcal{O}_q^*(\check{\underline{\chi}}), \quad q = 0, 1. \end{aligned}$$

It remains to define  $\mathcal{O}_q^{*p,S}(\check{\Gamma})(u)$  for  $q = 0, 1, 2$ :

$$\begin{aligned} \mathcal{O}_q^{*p,S}(\check{\chi})(u) &:= |r^{2+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \check{\chi}|_{p,S(u,\underline{u}_*)}, \\ \mathcal{O}_q^{*p,S}(\check{\underline{\chi}})(u) &:= |r^{1+q-\frac{2}{p}}|u|^{\frac{s-1}{2}} \nabla^q \check{\underline{\chi}}|_{p,S(u,\underline{u}_*)}, \\ \mathcal{O}_q^{*p,S}(\widetilde{\text{tr}} \chi)(u) &:= |r^{2+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \widetilde{\text{tr}} \chi|_{p,S(u,\underline{u}_*)}, \\ \mathcal{O}_q^{*p,S}(\widetilde{\text{tr}} \underline{\chi})(u) &:= |r^{2+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \widetilde{\text{tr}} \underline{\chi}|_{p,S(u,\underline{u}_*)}, \\ \mathcal{O}_q^{*p,S}(\check{\zeta})(u) &:= |r^{2+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \check{\eta}|_{p,S(u,\underline{u}_*)}, \\ \mathcal{O}_q^{*p,S}(\check{\gamma})(u) &:= |r^{1+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \check{\gamma}|_{p,S(u,\underline{u}_*)}, \\ \mathcal{O}_q^{*p,S}(\check{\epsilon})(u) &:= |r^{1+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \check{\epsilon}|_{p,S(u,\underline{u}_*)}, \\ \mathcal{O}_q^{*p,S}(\check{L})(u) &:= |r^{2+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \check{L}|_{p,S(u,\underline{u}_*)}, \\ \mathcal{O}_q^{*p,S}(\check{J})(u) &:= |r^{1+q-\frac{2}{p}}|u|^{\frac{s-1}{2}} \nabla^q \check{J}|_{p,S(u,\underline{u}_*)}, \\ \mathcal{O}_q^{*p,S}(\widetilde{[\mu]})(u) &:= |r^{3+q-\frac{2}{p}}|u|^{\frac{s-3}{2}} \nabla^q \widetilde{[\mu]}|_{p,S(u,\underline{u}_*)}. \end{aligned}$$

We also define the flux of curvature components on  $\underline{C}_*$  as follows:

$$\underline{\mathcal{R}}(\underline{C}_*) := \underline{\mathcal{R}}_1|_{\underline{C}_*} + \underline{\mathcal{R}}_0|_{\underline{C}_*},$$

where  $\underline{\mathcal{R}}_q$ ,  $q = 0, 1$  are defined in Section 3.1.1.

### 3.1.5 $\mathcal{O}_{(0)}$ norms and $\mathfrak{R}_{(0)}$ norms (Initial data for the foliation of $\mathcal{K}_{(0)}$ )

In this section, all the norms are defined with respect to the initial layer foliation  $(u_{(0)}, \underline{u})$  of  $\mathcal{K}_{(0)}$ . We define

$$\mathcal{O}_{(0)} := \sum_{q=0}^2 \sum_{\check{\Gamma}_{(0)}} (\mathcal{O}_{(0)})_q(\check{\Gamma}_{(0)}),$$

where

$$(\mathcal{O}_{(0)})_q(\check{\Gamma}_{(0)}) := \sup_{p \in [2,4]} \sup_{\mathcal{K}_{(0)}} (\mathcal{O}_{(0)})_q^{p,S}(\check{\Gamma}_{(0)}),$$

with

$$(\mathcal{O}_{(0)})_q^{p,S}(\check{\Gamma}_{(0)}) := \left| r_{(0)}^{\frac{s+1}{2} - \frac{2}{p}} (r_{(0)} \nabla_{(0)})^q \check{\Gamma}_{(0)} \right|_{p, S_{(0)}(u_{(0)}, \underline{u})},$$

and

$$\check{\Gamma}_{(0)} \in \left\{ \widetilde{\text{tr}} \chi_{(0)}, \widetilde{\text{tr}} \underline{\chi}_{(0)}, \check{\chi}_{(0)}, \check{\underline{\chi}}_{(0)}, \check{\eta}_{(0)}, \check{\underline{\eta}}_{(0)}, \check{\omega}_{(0)}, \check{\underline{\omega}}_{(0)}, r_{(0)}^{-1} \check{\gamma}_{(0)}, r_{(0)}^{-1} \check{\xi}_{(0)}, r_{(0)}^{-1} \check{b}_{(0)}, r_{(0)}^{-1} \check{\Omega}_{(0)} \right\}.$$

The initial curvature flux on  $\Sigma_0 \setminus K$  is defined by:

$$\mathfrak{R}_{(0)}^2 := \int_{\Sigma_0 \setminus K} \sum_{l=0}^2 r_{(0)}^s \left( |\mathfrak{d}_{(0)}^l \check{\alpha}_{(0)}|^2 + |\mathfrak{d}_{(0)}^l \check{\beta}_{(0)}|^2 + |\mathfrak{d}_{(0)}^l (\check{\rho}_{(0)}, \check{\sigma}_{(0)})|^2 + |\mathfrak{d}_{(0)}^l \check{\beta}_{(0)}|^2 + |\mathfrak{d}_{(0)}^l \check{\alpha}_{(0)}|^2 \right),$$

where

$$\mathfrak{d}_{(0)} := (r_{(0)} \nabla_{(0)}, r_{(0)} (\nabla_{(0)})_4, r_{(0)} (\nabla_{(0)})_3).$$

### 3.1.6 Osc norms (Oscillation control)

We have two foliations  $(u, \underline{u})$  and  $(u_{(0)}, \underline{u})$  in the initial layer region  $\mathcal{K}_{(0)}$  which will be compared using Osc norms. We denote  $(f, \lambda)$  the change of frame from  $(u, \underline{u})$  to  $(u_{(0)}, \underline{u})$  introduced in Lemma 5.2. We define

$$\text{Osc} := \text{Osc}(f) + \text{Osc}(u) + \text{Osc}(x) + \text{Osc}(r),$$

where

$$\begin{aligned} \text{Osc}(f) &:= \sup_{\mathcal{K}_{(0)}} |(r_{(0)})^{\frac{s-1}{2}} \mathfrak{d}_{(0)}^{\leq 1} f|, \\ \text{Osc}(u) &:= \sup_{\mathcal{K}_{(0)}} |(r_{(0)})^{\frac{s-3}{2}} (u' - u)|, \\ \text{Osc}(x) &:= \sup_{\mathcal{K}_{(0)}} \sum_{A=1}^2 |(r_{(0)})^{\frac{s-1}{2}} (x_{(0)}^A - x^A)|, \\ \text{Osc}(r) &:= \sup_{\mathcal{K}_{(0)}} |(r_{(0)})^{\frac{s-3}{2}} (r_{(0)} - r)|, \end{aligned}$$

with

$$\mathfrak{d}_{(0)} := \{r_{(0)} (\nabla_{(0)})_3, r_{(0)} \nabla_{(0)}, r_{(0)} (\nabla_{(0)})_4\}.$$

## 3.2 Smallness constants

Before starting our main theorem, we first introduce the following constants that will be used throughout this paper.

- The constant  $M > 0$  and the constant  $a$  are the mass and angular momentum of the initial Kerr spacetime relative to which our initial perturbation is measured.
- $R_0 > 0$  measures the radius of the compact set  $K \subset \Sigma_0$  relative to which the external region is defined by  $\mathcal{M} := D^+(\Sigma_0 \setminus K)$ , the future domain of dependence of  $\Sigma_0 \setminus K$ .

- The size of the initial data layer norm is measured by  $\epsilon_0 > 0$ .
- The size of the bootstrap assumption norms are measured by  $\epsilon > 0$ .
- $\delta_0 > 0$  measures the height of the initial layer region  $\mathcal{K}_{(0)}$ .

In what follows  $M$  and  $a$  are fixed constants such that  $0 \leq |a| \leq M$ . We define

$$\mathcal{E}_0 := \frac{M}{R_0}, \quad \epsilon := \mathcal{E}_0^{-\frac{1}{3}} \epsilon_0. \quad (3.4)$$

We take  $\epsilon_0$  small enough and  $R_0$  large enough such that

$$\epsilon^2 \ll \epsilon \mathcal{E}_0 \ll \epsilon \mathcal{E}_0^{\frac{1}{2}} \ll \epsilon_0 \ll \epsilon \ll \mathcal{E}_0 \ll \delta_0 < 1. \quad (3.5)$$

**Remark 3.1.** In (3.5),  $A \ll B$  means that  $CA < B$  where  $C$  is the largest universal constant among all the constants involved in the proof via  $\lesssim$ .

### 3.3 Main theorem

The goal of this paper is to prove the following theorem, which is similar to Theorem 4.1 in [2].

**Theorem 3.2** (Main theorem, version 2). *Let  $\mathbf{g}_{a,M}$  a Kerr metric in  $\mathcal{M}_{Kerr}$  with  $|a| \leq M$ . Let  $\mathcal{K}_{(0)}$  the initial layer region and  $\Phi_{(0)}$  the map defined in Section 2.1.4. We also assume that<sup>18</sup>*

$$\mathcal{O}_{(0)} \leq \epsilon_0, \quad \mathfrak{R}_{(0)} \leq \epsilon_0. \quad (3.6)$$

*Then,  $\mathcal{K}_{(0)}$  has a unique future development  $(\mathcal{M}, \mathbf{g})$ , defined inside its future domain of dependence, with the following properties:*

1.  *$(\mathcal{M}, \mathbf{g})$  can be foliated by a double null foliation  $(u, \underline{u})$ . Moreover, the outgoing cones  $C_u$  are complete for all  $u \leq u_0$ .*
2. *The norms  $\mathcal{O}, \mathcal{R}$  satisfy*

$$\mathcal{O} \lesssim \epsilon_0, \quad \mathcal{R} \lesssim \epsilon_0. \quad (3.7)$$

**Remark 3.3.** We denote  $(u_{\underline{u}_*}, \underline{u}_{\underline{u}_*})$  the double null foliation associated to the bootstrap region  $V(u_0, \underline{u}_*)$ . The bounds (3.7) allow us to show that  $(u_{\underline{u}_*}, \underline{u}_{\underline{u}_*})$  converges to  $(u_\infty, \underline{u}_\infty)$ <sup>19</sup> as  $\underline{u}_* \rightarrow \infty$  with all the bootstrap bounds valid on  $V(u_0, +\infty)$  for the quantities associated to the double null foliation  $(u_\infty, \underline{u}_\infty)$ .

The proof of Theorem 3.2 is given in Section 3.5. It hinges on a sequence of basic theorems stated in Section 3.4, concerning estimates for  $\mathcal{O}$  and  $\mathcal{R}$  norms.

**Remark 3.4.** Theorem 3.2 is proved in Section 3.5 for  $s \in [4, 6]$ , and extended to  $s \in (3, 4)$  and  $s > 6$  in Appendices B and C.

<sup>18</sup>The norms  $\mathcal{O}_{(0)}$  and  $\mathfrak{R}_{(0)}$  are defined in Section 3.1.5.

<sup>19</sup>Here,  $(u_\infty, \underline{u}_\infty)$  denotes the double null foliation of final spacetime.

### 3.4 Intermediate results

**Theorem M0.** *Assume that*

$$\mathcal{O}_{(0)} \leq \epsilon_0, \quad \mathfrak{R}_{(0)} \leq \epsilon_0, \quad \mathcal{O} \leq \epsilon, \quad \text{Osc} \leq \epsilon. \quad (3.8)$$

*Then, we have*

$$\mathfrak{R}_0 \lesssim \epsilon_0. \quad (3.9)$$

Theorem M0 is proved in Section 6.1. The proof is based on null frame transformation formulae introduced by Klainerman and Szeftel in [18], see Proposition 5.3 in Section 5.1.1.

**Theorem M1.** *Assume that*

$$\mathcal{O}_{(0)} \leq \epsilon_0, \quad \mathfrak{R}_0 \leq \epsilon_0, \quad \mathcal{O} \leq \epsilon, \quad \mathcal{R} \leq \epsilon, \quad \text{Osc} \leq \epsilon. \quad (3.10)$$

*Then, we have*

$$\mathcal{R} \lesssim \epsilon_0. \quad (3.11)$$

Theorem M1 is proved in Section 4. The proof is based on the  $r^p$ -weighted estimate method introduced by Dafermos and Rodnianski in [8].

**Theorem M2.** *Assume that  $\underline{C}_*$  is foliated by the geodesic type foliation of Definition 6.1 and that we have the following assumptions:*

$$\mathcal{O}_{(0)}(\underline{C}_* \cap \Sigma_0) \leq \epsilon_0, \quad \mathcal{R}(\underline{C}_*) \lesssim \epsilon_0, \quad \mathcal{O}^*(\underline{C}_*) \leq \epsilon.$$

*Then, we have*

$$\mathcal{O}^*(\underline{C}_*) \lesssim \epsilon_0.$$

Theorem M2 is proved in Section 6.2. The proof is done by integrating the transport equations along the null generator  $\underline{L}$  of the last slice  $\underline{C}_*$  and applying elliptic estimates on the 2-spheres of the geodesic type foliation of  $\underline{C}_*$ .

**Theorem M3.** *Assume that*

$$\mathcal{O}_{(0)} \leq \epsilon_0, \quad \mathcal{R} \leq \epsilon_0, \quad \mathcal{O}^*(\underline{C}_*) \leq \epsilon_0, \quad \mathcal{O} \leq \epsilon, \quad \text{Osc} \leq \epsilon. \quad (3.12)$$

*Then, we have*

$$\mathcal{O} \lesssim \epsilon_0, \quad \text{Osc} \lesssim \epsilon_0.$$

Theorem M3 is proved in Section 5. The proof is done by integrating the transport equations along the outgoing and ingoing null cones and applying elliptic estimate on the 2-spheres of the double null foliation of the spacetime  $\mathcal{K}$ .

**Theorem M4.** *We consider the spacetime  $\mathcal{K}$  and its double null foliation  $(u, \underline{u})$  which satisfies the assumptions:*

$$\mathcal{O} \lesssim \epsilon_0, \quad \mathcal{R} \lesssim \epsilon_0, \quad \text{Osc} \lesssim \epsilon_0, \quad \mathcal{O}^*(\underline{C}_*) \lesssim \epsilon_0. \quad (3.13)$$

*Then, we can extend the spacetime  $\mathcal{K} = V(u_0, \underline{u}_*)$  and the double null foliation to a new spacetime  $\widehat{\mathcal{K}} = \widehat{V}(u_0, \underline{u}_* + \nu)$ , where  $\nu > 0$  is sufficiently small, and an associated double null foliation  $(\widehat{u}, \widehat{\underline{u}})$ . Moreover, the new foliation  $(\widehat{u}, \widehat{\underline{u}})$  is geodesic on the new last slice  $\underline{C}_{**} := \underline{C}_{\underline{u}_* + \nu}$  and the new corresponding norms satisfy*

$$\widehat{\mathcal{O}} \lesssim \epsilon_0, \quad \widehat{\mathcal{R}} \lesssim \epsilon_0, \quad \widehat{\text{Osc}} \lesssim \epsilon_0, \quad \widehat{\mathcal{O}}^*(\underline{C}_{**}) \lesssim \epsilon_0. \quad (3.14)$$

Theorem M4 is proved in Section 6.3. The proof is based on local existence type arguments.

**Remark 3.5.** *When compared to [2], we have the following similarities and differences:*

1. *We treat the general decay  $s > 3$  in Theorem 3.2 while the main result of [2] corresponds to the particular case  $s > 7$  in Theorem 3.2.*
2. *In [2], fourth order derivatives of Ricci coefficients and third order derivatives of curvature are estimated. In this paper, we only estimate first order derivatives of both Ricci coefficients and curvature respectively in Theorems M2–M3 and M1.*
3. *To estimate the curvature components in Theorem M1, instead of using the classical vectorfield method as in [2], we use  $r^p$ -weighted estimates of [8].*
4. *On the last slice  $\underline{C}_*$ , a canonical foliation is used in [2] while we use a geodesic type foliation in Theorem M2.*
5. *The estimates of Ricci coefficients and the extension argument, i.e. the proof of Theorems M3 and M4, are similar to [2].*

### 3.5 Proof of the main theorem

Now, we use Theorems M0–M4 to prove Theorem 3.2.

**Definition 3.6.** *Let  $\aleph(\underline{u}_*)$  the set of spacetimes  $\mathcal{K}$  associated with a double null foliation  $(u, \underline{u})$  which satisfy the following properties:*

1.  $\mathcal{K} = V(u_0, \underline{u}_*)$ .
2. *The foliation on  $\underline{C}_*$  is the geodesic type foliation of Definition 6.1.*
3. *We have the following bootstrap assumptions:*

$$\mathcal{O} \leq \epsilon, \quad \mathcal{R} \leq \epsilon, \quad \text{Osc} \leq \epsilon, \quad \mathcal{O}^*(\underline{C}_*) \leq \epsilon. \quad (3.15)$$

**Definition 3.7.** *We denote  $\mathcal{U}$  the set of values  $\underline{u}_*$  such that  $\aleph(\underline{u}_*) \neq \emptyset$ .*

Applying the local existence result of Theorem 10.2.1 in [5] and the assumption  $\mathcal{O}_{(0)} \leq \epsilon_0$ , we deduce that (3.15) holds if  $\underline{u}_*$  is sufficiently close to  $R_0$ . So, we have  $\mathcal{U} \neq \emptyset$ .

Define  $\underline{u}_*$  to be the supremum of the set  $\mathcal{U}$ . We want to prove  $\underline{u}_* = \infty$ . We assume by contradiction that  $\underline{u}_*$  is finite. In particular, we may assume that  $\underline{u}_* \in \mathcal{U}$ . We consider the region  $\mathcal{K} = V(u_0, \underline{u}_*)$ . Recall that we have

$$\mathcal{O}_{(0)} \leq \epsilon_0, \quad \mathfrak{R}_{(0)} \leq \epsilon_0.$$

Combining with (3.15) and applying Theorem M0, we obtain

$$\mathfrak{R}_0 \lesssim \epsilon_0. \quad (3.16)$$

Then, we have from Theorem M1

$$\mathcal{R} \lesssim \epsilon_0. \quad (3.17)$$

Applying Theorem M2, we obtain

$$\mathcal{O}^*(\underline{C}_*) \lesssim \epsilon_0. \quad (3.18)$$




Figure 5: External region

Combining (3.15), (3.17), (3.18) and Theorem M3, we deduce

$$\mathcal{O} \lesssim \epsilon_0, \quad \text{Osc} \lesssim \epsilon_0. \quad (3.19)$$

Next, applying Theorem M4, we can extend  $\mathcal{K}$  to  $\widehat{\mathcal{K}} := \widehat{V}(u_0, \underline{u}_* + \nu)$  endowed with a new double null foliation  $(\widehat{u}, \widehat{u})$  and

$$\widehat{\mathcal{O}}(\widehat{\mathcal{K}}) \lesssim \epsilon_0, \quad \widetilde{\mathcal{R}}(\widehat{\mathcal{K}}) \lesssim \epsilon_0, \quad \widehat{\text{Osc}}(\widehat{\mathcal{K}}) \lesssim \epsilon_0, \quad \widehat{\mathcal{O}}^*(\underline{C}_{**} \cap \widehat{\mathcal{K}}) \lesssim \epsilon_0,$$

where  $\underline{C}_{**} = \underline{C}_{\underline{u}_* + \nu}$  is the new last slice. Thus, the region  $\widehat{V}(u_0, \underline{u}_* + \nu)$  with the new double null foliation  $(\widehat{u}, \widehat{u})$  satisfies all the properties in Definition 3.6, and so  $\aleph(\underline{u}_* + \nu) \neq \emptyset$ , a contradiction. So, we have  $\underline{u}_* = \infty$ , which implies property 1 of Theorem 3.2. Moreover, we also have

$$\mathcal{O} \lesssim \epsilon_0, \quad \mathcal{R} \lesssim \epsilon_0, \quad (3.20)$$

in the whole external region, which implies property 2 of Theorem 3.2. This concludes the proof of Theorem 3.2.

### 3.6 Conclusions

The following proposition is a consequence of Theorem 3.2.

**Proposition 3.8.** *Under the assumptions of Theorem 3.2, we have the following estimate in the external region<sup>20</sup>:*

$$|\mathbf{g} - \mathbf{g}_K|_{\infty, S(u, \underline{u})} \lesssim \frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}}. \quad (3.21)$$

*Proof.* We recall that we have

$$\mathbf{g} = -2\Omega^2(d\underline{u} \otimes du + du \otimes d\underline{u}) + \gamma_{AB}(dx^A - \underline{b}^A du) \otimes (dx^B - \underline{b}^B du),$$

<sup>20</sup>See Figure 5 for a description of the external region.

and

$$\mathbf{g}_K = -2\Omega_K^2(d\underline{u} \otimes du + du \otimes d\underline{u}) + (\gamma_K)_{AB}(dx^A - \underline{b}_K^A du) \otimes (dx^B - \underline{b}_K^B du).$$

We compare these two metrics as follows:

$$\begin{aligned} \mathbf{g}_{uu} - (\mathbf{g}_K)_{uu} &= 0, \\ \mathbf{g}_{\underline{u}\underline{u}} - (\mathbf{g}_K)_{\underline{u}\underline{u}} &= -4\Omega^2 + 4\Omega_K^2 = -4\check{\Omega}(\Omega + \Omega_K) = O\left(\frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}}\right), \\ \mathbf{g}_{uA} - (\mathbf{g}_K)_{uA} &= 0, \\ \mathbf{g}_{uu} - (\mathbf{g}_K)_{uu} &= \check{\gamma}_{AB}\underline{b}^A\underline{b}^B + (\gamma_K)_{AB}(\underline{b}^A\underline{b}^B - \underline{b}_K^A\underline{b}_K^B) = O\left(\frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}}\right), \\ \mathbf{g}_{uA} - (\mathbf{g}_K)_{uA} &= -\check{\gamma}_{AB}\underline{b}^B - (\gamma_K)_{AB}\check{\underline{b}}^B = O\left(\frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}\right), \\ \mathbf{g}_{AB} - (\mathbf{g}_K)_{AB} &= \check{\gamma}_{AB} = O\left(\frac{\epsilon_0 r}{|u|^{\frac{s-3}{2}}}\right). \end{aligned}$$

This concludes the proof of Proposition 3.8.  $\square$

**Remark 3.9.** Theorem 3.2 contains also a number of important conclusions following from (3.7): peeling properties, complete future infinity, Bondi mass formula and so on, see Chapter 8 in [14]. See also Section 3.8 in [20] for various conclusions of the Kerr stability in the entire domain of outer communication.

## 4 Curvature estimates (Theorem M1)

In this section, we prove Theorem M1 by the  $r^p$ -weighted estimate method introduced in [8] and applied to Bianchi equations in [11], [17] and [9]. For convenience, we recall the statement below.

**Theorem M1.** Assume that

$$\mathcal{O}_{(0)} \leq \epsilon_0, \quad \mathfrak{R}_0 \lesssim \epsilon_0, \quad \mathcal{O} \leq \epsilon, \quad \mathcal{R} \leq \epsilon, \quad \text{Osc} \leq \epsilon. \quad (4.1)$$

Then, we have

$$\mathcal{R} \lesssim \epsilon_0. \quad (4.2)$$

In the sequel, we denote<sup>21</sup>

$$V := V(u, \underline{u}), \quad C_u^V := C_u \cap V, \quad \underline{C}_{\underline{u}}^V := \underline{C}_{\underline{u}} \cap V. \quad (4.3)$$

The following lemma allows us to estimate the linear terms and error terms in Propositions 2.37 and 2.38.

**Lemma 4.1.** Under the assumptions  $\mathcal{O} \leq \epsilon$  and  $\mathcal{R} \leq \epsilon$ , we have

$$\begin{aligned} r^2|u|^{\frac{s-3}{2}}|\Gamma_g|_{\infty, S} &\lesssim \epsilon, & r|u|^{\frac{s-1}{2}}|\Gamma_b|_{\infty, S} &\lesssim \epsilon \\ |r^{2-\frac{2}{p}}|u|^{\frac{s-3}{2}}\Gamma_g^{(1)}|_{p, S} &\lesssim \epsilon, & |r^{1-\frac{2}{p}}|u|^{\frac{s-1}{2}}\Gamma_b^{(1)}|_{p, S} &\lesssim \epsilon, & p \in [2, 4]. \end{aligned} \quad (4.4)$$

*Proof.* It follows directly from the assumption (4.1), Definition 2.25 and Proposition 2.46.  $\square$

<sup>21</sup>See Figure 6 for a description of  $V(u, \underline{u})$ ,  $C_u^V$  and  $\underline{C}_{\underline{u}}^V$ .




Figure 6: Domain of integration

#### 4.1 General Bianchi pairs

The following lemma provides the general structure of Bianchi pairs. It will be used repeatedly in Section 4. See Lemma 8.24 in [17] for an analog under the assumption of axial polarization and Lemma 15.3.8 in [9] for an analog in the context of non integrable foliations.

**Lemma 4.2.** *Let  $k = 1, 2$  and  $a_{(1)}, a_{(2)}$  real numbers. Then, we have the following properties.*

1. *Let  $\psi_{(1)}, h_{(1)} \in \mathfrak{s}_k$  and  $\psi_{(2)}, h_{(2)} \in \mathfrak{s}_{k-1}$  satisfying*

$$\begin{aligned} \nabla_3(\psi_{(1)}) + a_{(1)} \operatorname{tr} \underline{\chi} \psi_{(1)} &= -k d_k^*(\psi_{(2)}) + h_{(1)}, \\ \nabla_4(\psi_{(2)}) + a_{(2)} \operatorname{tr} \chi \psi_{(2)} &= d_k(\psi_{(1)}) + h_{(2)}. \end{aligned} \quad (4.5)$$

*Then, the pair  $(\psi_{(1)}, \psi_{(2)})$  satisfies for any real number  $p$*

$$\begin{aligned} & \operatorname{Div}(r^p |\psi_{(1)}|^2 e_3) + k \operatorname{Div}(r^p |\psi_{(2)}|^2 e_4) \\ & + \left(2a_{(1)} - 1 - \frac{p}{2}\right) r^p \operatorname{tr} \underline{\chi} |\psi_{(1)}|^2 + k \left(2a_{(2)} - 1 - \frac{p}{2}\right) r^p \operatorname{tr} \chi |\psi_{(2)}|^2 \\ & = 2kr^p \operatorname{div}(\psi_{(1)} \cdot \psi_{(2)}) + 2r^p \psi_{(1)} \cdot h_{(1)} + 2kr^p \psi_{(2)} \cdot h_{(2)} - 2r^p \underline{\omega} |\psi_{(1)}|^2 \\ & - 2kr^p \omega |\psi_{(2)}|^2 + pr^{p-1} \left(e_3(r) - \frac{r}{2} \operatorname{tr} \underline{\chi}\right) |\psi_{(1)}|^2 + kpr^{p-1} \left(e_4(r) - \frac{r}{2} \operatorname{tr} \chi\right) |\psi_{(2)}|^2. \end{aligned} \quad (4.6)$$

2. *Let  $\psi_{(1)}, h_{(1)} \in \mathfrak{s}_{k-1}$  and  $\psi_{(2)}, h_{(2)} \in \mathfrak{s}_k$  satisfying*

$$\begin{aligned} \nabla_3(\psi_{(1)}) + a_{(1)} \operatorname{tr} \underline{\chi} \psi_{(1)} &= d_k(\psi_{(2)}) + h_{(1)}, \\ \nabla_4(\psi_{(2)}) + a_{(2)} \operatorname{tr} \chi \psi_{(2)} &= -k d_k^*(\psi_{(1)}) + h_{(2)}. \end{aligned} \quad (4.7)$$

Then, the pair  $(\psi_{(1)}, \psi_{(2)})$  satisfies for any real number  $p$

$$\begin{aligned}
& k \mathbf{Div}(r^p |\psi_{(1)}|^2 e_3) + \mathbf{Div}(r^p |\psi_{(2)}|^2 e_4) \\
& + k \left( 2a_{(1)} - 1 - \frac{p}{2} \right) r^p \operatorname{tr} \chi |\psi_{(1)}|^2 + \left( 2a_{(2)} - 1 - \frac{p}{2} \right) r^p \operatorname{tr} \chi |\psi_{(2)}|^2 \\
& = 2r^p \operatorname{div}(\psi_{(1)} \cdot \psi_{(2)}) + 2kr^p \psi_{(1)} \cdot h_{(1)} + 2r^p \psi_{(2)} \cdot h_{(2)} - 2kr^p \omega |\psi_{(1)}|^2 \\
& - 2r^p \omega |\psi_{(2)}|^2 + kpr^{p-1} \left( e_3(r) - \frac{r}{2} \operatorname{tr} \chi \right) |\psi_{(1)}|^2 + pr^{p-1} \left( e_4(r) - \frac{r}{2} \operatorname{tr} \chi \right) |\psi_{(2)}|^2.
\end{aligned} \tag{4.8}$$

*Proof.* See Lemma 4.2 in [25].  $\square$

**Remark 4.3.** Note that the linearized Bianchi equations can be written as systems of equations of the type (4.5) and (4.7). In particular

- the Bianchi pair  $(\check{\alpha}, \check{\beta})$  satisfies (4.5) with  $k = 2$ ,  $a_{(1)} = \frac{1}{2}$ ,  $a_{(2)} = 2$ ,
- the Bianchi pair  $(\check{\beta}, (\check{\rho}, -\check{\sigma}))$  satisfies (4.5) with  $k = 1$ ,  $a_{(1)} = 1$ ,  $a_{(2)} = \frac{3}{2}$ ,
- the Bianchi pair  $((\check{\rho}, \check{\sigma}), \check{\beta})$  satisfies (4.7) with  $k = 1$ ,  $a_{(1)} = \frac{3}{2}$ ,  $a_{(2)} = 1$ ,
- the Bianchi pair  $(\check{\beta}, \check{\alpha})$  satisfies (4.7) with  $k = 2$ ,  $a_{(1)} = 2$ ,  $a_{(2)} = \frac{1}{2}$ .

**Proposition 4.4.** For  $\psi_{(1)}$ ,  $\psi_{(2)}$  and  $h_{(1)}$ ,  $h_{(2)}$  satisfying (4.5) or (4.7), we have the following properties for all  $(u, \underline{u}) \in \mathcal{K}$ :

(a) In the case  $2 + p - 4a_{(1)} > 0$  and  $4a_{(2)} - 2 - p > 0$ , we have

$$\begin{aligned}
& \int_{C_u^V} r^p |\psi_{(1)}|^2 + \int_{\underline{C}_{\underline{u}}^V} r^p |\psi_{(2)}|^2 + \int_V r^{p-1} |\psi_{(1)}|^2 + r^{p-1} |\psi_{(2)}|^2 \\
& \lesssim \int_{\Sigma_0 \cap V} r^p |\psi_{(1)}|^2 + r^p |\psi_{(2)}|^2 + \int_V r^p |\psi_{(1)}| |h_{(1)}| + r^p |\psi_{(2)}| |h_{(2)}|.
\end{aligned} \tag{4.9}$$

(b) In the case  $2 + p - 4a_{(1)} > 0$  and  $4a_{(2)} - 2 - p = 0$ , we have

$$\begin{aligned}
& \int_{C_u^V} r^p |\psi_{(1)}|^2 + \int_{\underline{C}_{\underline{u}}^V} r^p |\psi_{(2)}|^2 + \int_V r^{p-1} |\psi_{(1)}|^2 \\
& \lesssim \int_{\Sigma_0 \cap V} r^p |\psi_{(1)}|^2 + r^p |\psi_{(2)}|^2 + \int_V r^p |\psi_{(1)}| |h_{(1)}| + r^p |\psi_{(2)}| |h_{(2)}|.
\end{aligned} \tag{4.10}$$

(c) In the case  $2 + p - 4a_{(1)} \leq 0$  and  $4a_{(2)} - 2 - p \geq 0$ , we have

$$\begin{aligned}
& \int_{C_u^V} r^p |\psi_{(1)}|^2 + \int_{\underline{C}_{\underline{u}}^V} r^p |\psi_{(2)}|^2 \\
& \lesssim \int_{\Sigma_0 \cap V} r^p |\psi_{(1)}|^2 + r^p |\psi_{(2)}|^2 + \int_V r^{p-1} |\psi_{(1)}|^2 + r^p |\psi_{(1)}| |h_{(1)}| + r^p |\psi_{(2)}| |h_{(2)}|.
\end{aligned} \tag{4.11}$$

(d) In the case  $2 + p - 4a_{(1)} > 0$  and  $4a_{(2)} - 2 - p \leq 0$ , we have

$$\begin{aligned}
& \int_{C_u^V} r^p |\psi_{(1)}|^2 + \int_{\underline{C}_{\underline{u}}^V} r^p |\psi_{(2)}|^2 + \int_V r^{p-1} |\psi_{(1)}|^2 \\
& \lesssim \int_{\Sigma_0 \cap V} r^p |\psi_{(1)}|^2 + r^p |\psi_{(2)}|^2 + \int_V r^{p-1} |\psi_{(2)}|^2 + r^p |\psi_{(1)}| |h_{(1)}| + r^p |\psi_{(2)}| |h_{(2)}|.
\end{aligned} \tag{4.12}$$

**Remark 4.5.** In the sequel, For a sum of terms of the form  $\Gamma \cdot R$ , we ignore the terms having the same or even better decay. For example, we write

$$\Gamma_g \cdot \beta + \Gamma_g \cdot \alpha = \Gamma_g \cdot \beta, \quad \Gamma_g \cdot (\check{\rho}, \check{\sigma}) + \Gamma_g \cdot \underline{\beta} = \Gamma_g \cdot \underline{\beta},$$

since  $\alpha$  decays better than  $\beta$  and  $\check{\rho}$  and  $\check{\sigma}$  decay better than  $\underline{\beta}$ .

*Proof.* We obtain from Proposition 2.30 that

$$\omega = \Gamma_g + \mathbf{O}_2^1, \quad \underline{\omega} = \Gamma_b + \mathbf{O}_2^1.$$

Applying Lemma 2.7, we obtain

$$e_3(r) - \frac{r}{2} \operatorname{tr} \underline{\chi} = \frac{\overline{\Omega \operatorname{tr} \underline{\chi}}}{2\Omega} r - \frac{r}{2} \operatorname{tr} \underline{\chi} = \frac{r}{2\Omega} (\overline{\Omega \operatorname{tr} \underline{\chi}} - \Omega \operatorname{tr} \underline{\chi}) = r\Gamma_g + \mathbf{O}_2^1, \quad (4.13)$$

and similarly

$$e_4(r) - \frac{r}{2} \operatorname{tr} \chi = r\Gamma_g + \mathbf{O}_2^1. \quad (4.14)$$

Applying Lemma 4.1, we infer

$$\begin{aligned} & \int_V r^p (|\underline{\omega}| |\psi_{(1)}|^2 + |\omega| |\psi_{(2)}|^2) \\ & + \int_V r^{p-1} \left( e_3(r) - \frac{r}{2} \operatorname{tr} \underline{\chi} \right) |\psi_{(1)}|^2 + r^{p-1} \left( e_4(r) - \frac{r}{2} \operatorname{tr} \chi \right) |\psi_{(2)}|^2 \\ & \lesssim \int_V r^p (\Gamma_b + \mathbf{O}_2^1) |\psi_{(1)}|^2 + \int_V r^p (\Gamma_g + \mathbf{O}_2^1) |\psi_{(2)}|^2 \\ & \lesssim M \int_V r^{p-2} |\psi_{(1)}|^2 + \epsilon \int_V r^{p-1} |u|^{-1} |\psi_{(1)}|^2 + M \int_V r^{p-2} |\psi_{(2)}|^2 \\ & \lesssim M \int_{u_0(\underline{u})}^u \left( |u|^{-2} \int_{C_u^V} r^p |\psi_{(1)}|^2 \right) du + M \int_{|u|}^{\underline{u}} \left( |\underline{u}|^{-2} \int_{\underline{C}_{\underline{u}}^V} r^p |\psi_{(2)}|^2 \right) d\underline{u} \\ & \lesssim M \int_{u_0(\underline{u})}^u |u|^{-2} du \left( \sup_u \int_{C_u^V} r^p |\psi_{(1)}|^2 \right) + M \int_{|u|}^{\underline{u}} |\underline{u}|^{-2} d\underline{u} \left( \sup_{\underline{u}} \int_{\underline{C}_{\underline{u}}^V} r^p |\psi_{(2)}|^2 \right) \\ & \lesssim \mathcal{E}_0 \left( \sup_u \int_{C_u^V} r^p |\psi_{(1)}|^2 \right) + \mathcal{E}_0 \left( \sup_{\underline{u}} \int_{\underline{C}_{\underline{u}}^V} r^p |\psi_{(2)}|^2 \right). \end{aligned} \quad (4.15)$$

Integrating (4.6) or (4.8) and recalling that

$$\operatorname{tr} \chi - \frac{2}{r} = \Gamma_g + \mathbf{O}_2^1, \quad \operatorname{tr} \underline{\chi} + \frac{2}{r} = \Gamma_g + \mathbf{O}_2^1,$$

we obtain

$$\begin{aligned} & \int_{C_u^V} r^p |\psi_{(1)}|^2 + \int_{\underline{C}_{\underline{u}}^V} r^p |\psi_{(2)}|^2 + \int_V (2 + p - 4a_{(1)}) r^{p-1} |\psi_{(1)}|^2 + (4a_{(2)} - 2 - p) r^{p-1} |\psi_{(2)}|^2 \\ & \lesssim \int_{\Sigma_0 \cap V} r^p |\psi_{(1)}|^2 + r^p |\psi_{(2)}|^2 + \int_V r^p |\psi_{(1)}| |h_{(1)}| + r^p |\psi_{(2)}| |h_{(2)}| \\ & + \mathcal{E}_0 \sup_u \int_{C_u^V} r^p |\psi_{(1)}|^2 + \mathcal{E}_0 \sup_{\underline{u}} \int_{\underline{C}_{\underline{u}}^V} r^p |\psi_{(2)}|^2. \end{aligned}$$

Taking the supremum in  $u$  and  $\underline{u}$  and applying (4.15), we obtain for  $\epsilon$  and  $\mathcal{E}_0$  small enough

$$\begin{aligned} & \sup_u \int_{C_u^V} r^p |\psi_{(1)}|^2 + \sup_{\underline{u}} \int_{\underline{C}_{\underline{u}}^V} r^p |\psi_{(2)}|^2 \\ & + \int_V (2 + p - 4a_{(1)}) r^{p-1} |\psi_{(1)}|^2 + (4a_{(2)} - 2 - p) r^{p-1} |\psi_{(2)}|^2 \\ & \lesssim \int_{\Sigma_0 \cap V} r^p |\psi_{(1)}|^2 + r^p |\psi_{(2)}|^2 + \int_V r^p |\psi_{(1)}| |h_{(1)}| + r^p |\psi_{(2)}| |h_{(2)}|. \end{aligned} \quad (4.16)$$

This concludes the proof of Proposition 4.4.  $\square$

The following lemma allows us to compare  $\check{\psi}$  and  $\check{\psi}$ .

**Lemma 4.6.** *We denote for  $\psi \in \{\alpha, \beta, \rho, \sigma, \underline{\beta}, \underline{\alpha}\}$*

$$\widetilde{\psi^{(1)}} := (\check{\psi}, \check{\psi}). \quad (4.17)$$

*Then, we have for  $p < 7$*

$$\begin{aligned} \int_{C_u^V} r^p |\check{\psi}^{(1)}|^2 & \lesssim \int_{C_u^V} r^p |\widetilde{\psi^{(1)}}|^2 + \frac{\epsilon_0^2}{|u|^{s-p}}, \\ \int_{\underline{C}_{\underline{u}}^V} r^p |\check{\psi}^{(1)}|^2 & \lesssim \int_{\underline{C}_{\underline{u}}^V} r^p |\widetilde{\psi^{(1)}}|^2 + \frac{\epsilon_0^2}{|u|^{s-p}}. \end{aligned} \quad (4.18)$$

**Remark 4.7.** *Note that Proposition 2.38 involves the quantities  $\check{\alpha}, \check{\beta}, \check{\rho}, \check{\sigma}, \check{\beta}, \check{\alpha}$  and  $\check{\alpha}$ , while the curvature norms  $\mathcal{R}_1$  and  $\underline{\mathcal{R}}_1$  involve  $\check{\alpha}, \check{\beta}, \check{\rho}, \check{\sigma}, \check{\beta}, \check{\alpha}$ . We thus need to compare  $\check{\psi}$  and  $\check{\psi}$  which is the purpose of Lemma 4.6.*

*Proof.* For  $\psi \in \{\rho, \sigma\}$  which is a scalar function, we have

$$\check{\psi} = \check{\psi}.$$

The estimate (4.18) is trivial in this case. Next, we focus on the case  $\psi \in \{\alpha, \beta, \underline{\beta}, \underline{\alpha}\}$ . We have from Proposition 2.22

$$\check{\psi} = \check{\psi} - \check{\psi}_K \psi_K = \check{\psi} + (\check{\psi} - \check{\psi}_K) \psi_K = \check{\psi} + r \check{L} \cdot \psi_K.$$

Recalling that  $\psi_K = \mathbf{O}_4^2$ , we deduce

$$\begin{aligned} \int_{C_u^V} r^p |\check{\psi}|^2 & \lesssim \int_{C_u^V} r^p |\check{\psi}|^2 + \int_{C_u^V} r^p |\Gamma_g^{(1)} \cdot \mathbf{O}_3^2|^2 \\ & \lesssim \int_{C_u^V} r^p |\check{\psi}|^2 + M^4 \int_{C_u^V} r^{p-6} |\Gamma_g^{(1)}|^2 \\ & \lesssim \int_{C_u^V} r^p |\check{\psi}|^2 + M^4 \int_{|u|}^{\underline{u}} r^{p-8} d\underline{u} \int_S |r \Gamma_g^{(1)}|^2 \\ & \lesssim \int_{C_u^V} r^p |\check{\psi}|^2 + M^4 \epsilon^2 \int_{|u|}^{\underline{u}} r^{p-8} \frac{d\underline{u}}{|u|^{s-3}} \\ & \lesssim \int_{C_u^V} r^p |\check{\psi}|^2 + M^4 \epsilon^2 |u|^{p-s-4} \\ & \lesssim \int_{C_u^V} r^p |\check{\psi}|^2 + \frac{\mathcal{E}_0^4 \epsilon^2}{|u|^{s-p}} \\ & \lesssim \int_{C_u^V} r^p |\check{\psi}|^2 + \frac{\epsilon_0^2}{|u|^{s-p}}, \end{aligned}$$

which implies the first inequality in (4.18). The second inequality in (4.18) is similar. This concludes the proof of Lemma 4.6.  $\square$

**Remark 4.8.** *The proof of Theorem M1 is analogous to Theorem M1 in [25]. In particular, the estimates for the error terms  $\text{Err}[\dots]$  in Propositions 2.37 and 2.38 have already been done in [25]. In the proof of Theorem M1 below, we thus focus on the linear terms  $\mathfrak{L}[\dots]$ .*

## 4.2 Estimates for the Bianchi pair $(\check{\alpha}, \check{\beta})$

**Proposition 4.9.** *We have the following estimate:*

$$\mathcal{R}_0[\check{\alpha}]^2 + \mathcal{R}_0[\check{\beta}]^2 + \mathcal{R}_1[\check{\alpha}]^2 + \mathcal{R}_1[\check{\beta}]^2 \lesssim \epsilon_0^2. \quad (4.19)$$

*Proof.* We recall the following equations from Propositions 2.37 and 2.38:

$$\begin{aligned} \nabla_3 \widetilde{\alpha^{(1)}} + \frac{1}{2} \text{tr} \chi \widetilde{\alpha^{(1)}} &= -2d_2^*(\widetilde{\beta^{(1)}}) + \mathfrak{L}[\nabla_3 \widetilde{\alpha^{(1)}}] + \text{Err}[\nabla_3 \widetilde{\alpha^{(1)}}], \\ \nabla_4 \widetilde{\beta^{(1)}} + 2 \text{tr} \chi \widetilde{\beta^{(1)}} &= d_2(\widetilde{\alpha^{(1)}}) + \mathfrak{L}[\nabla_4 \widetilde{\beta^{(1)}}] + \text{Err}[\nabla_4 \widetilde{\beta^{(1)}}]. \end{aligned} \quad (4.20)$$

Applying (4.9) with  $\psi_{(1)} = \widetilde{\alpha^{(1)}}$ ,  $\psi_{(2)} = \widetilde{\beta^{(1)}}$ ,  $a_{(1)} = \frac{1}{2}$ ,  $a_{(2)} = 2$ ,  $h_{(1)} = \mathfrak{L}[\nabla_3 \widetilde{\alpha^{(1)}}] + \text{Err}[\nabla_3 \widetilde{\alpha^{(1)}}]$ ,  $h_{(2)} = \mathfrak{L}[\nabla_4 \widetilde{\beta^{(1)}}] + \text{Err}[\nabla_4 \widetilde{\beta^{(1)}}]$  and  $p = s$  we obtain

$$\begin{aligned} &\int_{C_u^V} r^s |\widetilde{\alpha^{(1)}}|^2 + \int_{\underline{C}_u^V} r^s |\widetilde{\beta^{(1)}}|^2 + \int_V r^{s-1} |\widetilde{\alpha^{(1)}}|^2 \\ &\lesssim \int_{\Sigma_0 \cap V} r^s |\widetilde{\alpha^{(1)}}|^2 + r^s |\widetilde{\beta^{(1)}}|^2 + \int_V r^s |\widetilde{\alpha^{(1)}}| \left| \mathfrak{L}[\nabla_3 \widetilde{\alpha^{(1)}}] + \text{Err}[\nabla_3 \widetilde{\alpha^{(1)}}] \right| \\ &\quad + \int_V r^s |\widetilde{\beta^{(1)}}| \left| \text{Err}[\nabla_4 \widetilde{\beta^{(1)}}] + \text{Err}[\nabla_4 \widetilde{\beta^{(1)}}] \right|. \end{aligned} \quad (4.21)$$

We recall from Proposition 2.37<sup>22</sup>

$$\begin{aligned} &\int_V r^s |\widetilde{\alpha^{(1)}}| |\mathfrak{L}[\nabla_3 \widetilde{\alpha^{(1)}}]| + r^s |\widetilde{\beta^{(1)}}| |\mathfrak{L}[\nabla_4 \widetilde{\beta^{(1)}}]| \\ &\lesssim \int_V r^s \left( \frac{M}{r^3} |\Gamma_g^{(1)}| |\widetilde{\alpha^{(1)}}| + \frac{M^2}{r^3} |\widetilde{\alpha^{(1)}}| |\check{\rho}^{\leq 1} \check{\rho}| + \frac{M}{r^2} |\check{\rho}^{\leq 1} \check{\beta}|^2 \right). \end{aligned} \quad (4.22)$$

We estimate (4.22) term by term. For the first term:

$$\begin{aligned} \int_V M r^{s-3} |\widetilde{\alpha^{(1)}}| |\Gamma_g^{(1)}| &\lesssim M \int_{u_0(\underline{u})}^u du \left( \int_{C_u^V} r^s |\widetilde{\alpha^{(1)}}|^2 \right)^{\frac{1}{2}} \left( \int_{C_u^V} r^{s-6} |\Gamma_g^{(1)}|^2 \right)^{\frac{1}{2}} \\ &\lesssim M \epsilon \int_{u_0(\underline{u})}^u du \left( \int_{|u|}^u r^{s-8} |r \Gamma_g^{(1)}|_{2,S}^2 d\underline{u} \right)^{\frac{1}{2}} \\ &\lesssim M \epsilon^2 \int_{u_0(\underline{u})}^u du \left( \int_{|u|}^u r^{s-8} |u|^{3-s} d\underline{u} \right)^{\frac{1}{2}} \\ &\lesssim M \epsilon^2 \int_{u_0(\underline{u})}^u |u|^{-2} du \\ &\lesssim \epsilon_0^2. \end{aligned}$$

---

<sup>22</sup>We ignore the terms which decay better.

For the second term:

$$\begin{aligned}
\int_V M^2 r^{s-3} |\widetilde{\alpha^{(1)}}| |\widetilde{\rho^{(1)}}| &\lesssim M^2 \int_{u_0(\underline{u})}^u du \left( \int_{C_u^V} r^s |\widetilde{\alpha^{(1)}}|^2 \right)^{\frac{1}{2}} \left( \int_{C_u^V} r^2 |\widetilde{\rho^{(1)}}|^2 \right)^{\frac{1}{2}} r^{\frac{s}{2}-4} \\
&\lesssim M^2 \int_{u_0(\underline{u})}^u du \frac{\epsilon^2}{|u|^{\frac{s-2}{2}}} |u|^{\frac{s}{2}-4} \\
&\lesssim M^2 \epsilon^2 |u|^{-2} \\
&\lesssim \epsilon_0^2.
\end{aligned}$$

For the last term:

$$\int_V M r^{s-2} |\widetilde{\beta^{(1)}}|^2 \lesssim M \int_{|u|}^u \frac{du}{r^2} \int_{\underline{C}_u^V} r^s |\widetilde{\beta^{(1)}}|^2 \lesssim M \epsilon^2 \int_{|u|}^u \frac{du}{r^2} \lesssim \epsilon_0^2.$$

Injecting all the estimates into (4.22), we deduce

$$\int_V r^s |\widetilde{\alpha^{(1)}}| |\mathfrak{L}[\nabla_3 \widetilde{\alpha^{(1)}}]| + r^s |\widetilde{\beta^{(1)}}| |\mathfrak{L}[\nabla_4 \widetilde{\beta^{(1)}}]| \lesssim \epsilon_0^2. \quad (4.23)$$

Next, we proceed as in Proposition 4.7 in [25] to obtain

$$\int_V r^s \left( |\widetilde{\alpha^{(1)}}| |\text{Err}[\nabla_3 \widetilde{\alpha^{(1)}}]| + |\widetilde{\beta^{(1)}}| |\text{Err}[\nabla_4 \widetilde{\beta^{(1)}}]| \right) \lesssim \epsilon_0^2. \quad (4.24)$$

Combining (4.23) and (4.24), we obtain

$$\int_{C_u^V} r^s |\widetilde{\alpha^{(1)}}|^2 + \int_{\underline{C}_u^V} r^s |\widetilde{\beta^{(1)}}|^2 + \int_V r^{s-1} |\widetilde{\alpha^{(1)}}|^2 \lesssim \epsilon_0^2. \quad (4.25)$$

Applying Lemma 4.6, we deduce (4.19). This concludes the proof of Proposition 4.9.  $\square$

### 4.3 Estimates for the Bianchi pair $(\check{\beta}, (\check{\rho}, \check{\sigma}))$

**Proposition 4.10.** *We have the following estimate:*

$$\mathcal{R}_0[\check{\beta}]^2 + \mathcal{R}_0[(\check{\rho}, \check{\sigma})]^2 + \mathcal{R}_1[\check{\beta}]^2 + \mathcal{R}_1[(\check{\rho}, \check{\sigma})]^2 \lesssim \epsilon_0^2. \quad (4.26)$$

*Proof.* We recall from Propositions 2.37 and 2.38:

$$\begin{aligned}
\nabla_4(\widetilde{\rho^{(1)}}) + \frac{3}{2} \text{tr} \chi(\widetilde{\rho^{(1)}}) - \nabla_4(\widetilde{\sigma^{(1)}}) &= d_1 \check{\beta} + \mathfrak{L}[\nabla_4 \widetilde{\rho^{(1)}}] - \nabla_4 \widetilde{\sigma^{(1)}} + \text{Err}[\nabla_4 \widetilde{\rho^{(1)}}] - \nabla_4 \widetilde{\sigma^{(1)}}], \\
\nabla_3 \widetilde{\beta^{(1)}} + \text{tr} \underline{\chi} \widetilde{\beta^{(1)}} &= -d_1^* (\widetilde{\rho^{(1)}}) - \nabla_3 \widetilde{\sigma^{(1)}} + \mathfrak{L}[\nabla_3 \widetilde{\beta^{(1)}}] + \text{Err}[\nabla_3 \widetilde{\beta^{(1)}}],
\end{aligned}$$

where

$$\mathfrak{L}[\nabla_4 \widetilde{\rho^{(1)}}] - \nabla_4 \widetilde{\sigma^{(1)}} = \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot \check{\rho}^{(1)}, \quad \mathfrak{L}[\nabla_3 \widetilde{\beta^{(1)}}] = \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\beta}^{(1)} + \mathbf{O}_3^2 \cdot \check{\beta}^{(1)}.$$

Applying (4.10) with  $\psi_{(1)} = \widetilde{\beta^{(1)}}$ ,  $\psi_{(2)} = (\widetilde{\rho^{(1)}}) - \nabla_3 \widetilde{\sigma^{(1)}}$ ,  $a_{(1)} = 1$ ,  $a_{(2)} = \frac{3}{2}$ ,  $h_{(1)} = \mathfrak{L}[\nabla_3 \widetilde{\beta^{(1)}}] +$

$\text{Err}[\nabla_3 \widetilde{\beta^{(1)}}]$ ,  $h_{(2)} = \mathfrak{L}[\nabla_4 \widetilde{\rho^{(1)}}, -\nabla_4 \widetilde{\sigma^{(1)}}] + \text{Err}[\nabla_4 \widetilde{\rho^{(1)}}, -\nabla_4 \widetilde{\sigma^{(1)}}]$  and  $p = 4$ , we infer

$$\begin{aligned}
& \int_{C_u^V} r^4 |\widetilde{\beta^{(1)}}|^2 + \int_{\underline{C}_u^V} r^4 |(\widetilde{\rho^{(1)}}, \widetilde{\sigma^{(1)}})|^2 \\
& \lesssim \int_{\Sigma_0 \cap V} \left( r^4 |\widetilde{\beta^{(1)}}|^2 + r^4 |(\widetilde{\rho^{(1)}}, \widetilde{\sigma^{(1)}})|^2 \right) + \int_V r^4 \left| \widetilde{\beta^{(1)}} \cdot \left( \mathfrak{L}[\nabla_3 \widetilde{\beta^{(1)}}] + \text{Err}[\nabla_3 \widetilde{\beta^{(1)}}] \right) \right| \\
& \quad + \int_V r^4 \left| (\widetilde{\rho^{(1)}}, \widetilde{\sigma^{(1)}}) \cdot \left( \mathfrak{L}[\nabla_4 \widetilde{\rho^{(1)}}, \nabla_4 \widetilde{\sigma^{(1)}}] + \text{Err}[\nabla_4 \widetilde{\rho^{(1)}}, \nabla_4 \widetilde{\sigma^{(1)}}] \right) \right| \\
& \lesssim |u|^{4-s} \epsilon_0^2 + \int_V r^4 \left| \widetilde{\beta^{(1)}} \cdot \left( \mathfrak{L}[\nabla_3 \widetilde{\beta^{(1)}}] + \text{Err}[\nabla_3 \widetilde{\beta^{(1)}}] \right) \right| \\
& \quad + \int_V r^4 \left| (\widetilde{\rho^{(1)}}, \widetilde{\sigma^{(1)}}) \cdot \left( \mathfrak{L}[\nabla_4 \widetilde{\rho^{(1)}}, \nabla_4 \widetilde{\sigma^{(1)}}] + \text{Err}[\nabla_4 \widetilde{\rho^{(1)}}, \nabla_4 \widetilde{\sigma^{(1)}}] \right) \right|.
\end{aligned} \tag{4.27}$$

Notice that we have

$$\begin{aligned}
& \int_V r^4 |\widetilde{\beta^{(1)}}| |\mathfrak{L}[\nabla_3 \widetilde{\beta^{(1)}}]| + r^4 |(\widetilde{\rho^{(1)}}, \widetilde{\sigma^{(1)}})| |\mathfrak{L}[\nabla_4 \widetilde{\rho^{(1)}}, -\nabla_4 \widetilde{\sigma^{(1)}}]| \\
& \lesssim \int_V M r^2 |\check{\beta}^{(1)}|^2 + M^2 r |\check{\beta}^{(1)}| |\check{\beta}^{(1)}| + M r |\Gamma_g^{(1)}| |\check{\rho}^{(1)}| \\
& \lesssim M \int_{|u|}^u \frac{du}{r^2} \int_{\underline{C}_u^V} r^4 |\check{\beta}^{(1)}|^2 + M^2 \int_{|u|}^u \frac{du}{r^{\frac{s}{2}}} \int_{\underline{C}_u^V} |r^{\frac{s}{2}} \check{\beta}^{(1)}| |r \check{\beta}^{(1)}| \\
& \quad + M \int_{|u|}^u \frac{du}{r^2} \int_{\underline{C}_u^V} |r \Gamma_g^{(1)}| |r^2 \check{\rho}^{(1)}| \\
& \lesssim M \int_{|u|}^u \frac{du}{r^2} \frac{\epsilon^2}{|u|^{s-4}} + M^2 \int_{|u|}^u \frac{du}{r^{\frac{s}{2}}} \left( \int_{\underline{C}_u^V} |r^{\frac{s}{2}} \check{\beta}^{(1)}| \right)^{\frac{1}{2}} \left( \int_{\underline{C}_u^V} |r \check{\beta}^{(1)}|^2 \right)^{\frac{1}{2}} \\
& \quad + M \int_{|u|}^u \frac{du}{r^2} \left( \int_{u_0(u)}^u du |r \Gamma_g^{(1)}|_{2,S}^2 \right)^{\frac{1}{2}} \left( \int_{\underline{C}_u^V} |r^2 \check{\rho}^{(1)}|^2 \right)^{\frac{1}{2}} \\
& \lesssim \frac{M \epsilon^2}{r |u|^{s-4}} + M^2 \epsilon \int_{|u|}^u \frac{du}{r^{\frac{s}{2}}} \frac{\epsilon}{|u|^{\frac{s-2}{2}}} + M \int_{|u|}^u \frac{du}{r^2} \frac{\epsilon}{|u|^{\frac{s-4}{2}}} \frac{\epsilon}{|u|^{\frac{s-4}{2}}} \\
& \lesssim \frac{M \epsilon^2}{r |u|^{s-4}} + \frac{M^2 \epsilon^2}{|u|^{s-2}} + \frac{M \epsilon^2}{r |u|^{s-4}} \\
& \lesssim \frac{\epsilon_0^2}{|u|^{s-4}}.
\end{aligned} \tag{4.28}$$

Next, we proceed as in Proposition 4.8 in [25] to obtain

$$\int_V r^4 |\widetilde{\beta^{(1)}}| |\text{Err}[\nabla_3 \widetilde{\beta^{(1)}}]| + r^4 |(\widetilde{\rho^{(1)}}, \widetilde{\sigma^{(1)}})| |\text{Err}[\nabla_4 \widetilde{\rho^{(1)}}, \nabla_4 \widetilde{\sigma^{(1)}}]| \lesssim \frac{\epsilon_0^2}{|u|^{s-4}}. \tag{4.29}$$

Combining (4.27), (4.28) and (4.29), we deduce

$$\int_{C_u^V} r^4 |\widetilde{\beta^{(1)}}|^2 + \int_{\underline{C}_u^V} r^4 |(\widetilde{\rho^{(1)}}, \widetilde{\sigma^{(1)}})|^2 \lesssim \frac{\epsilon_0^2}{|u|^{s-4}}. \tag{4.30}$$

Combining with Lemma 4.6, this concludes the proof of Proposition 4.10.  $\square$

#### 4.4 Estimates for the Bianchi pair $((\check{\rho}, \check{\sigma}), \check{\beta})$

**Proposition 4.11.** *We have the following estimates:*

$$\mathcal{R}_0[(\check{\rho}, \check{\sigma})]^2 + \mathcal{R}_0[\check{\beta}]^2 + \mathcal{R}_1[(\check{\rho}, \check{\sigma})]^2 + \mathcal{R}_1[\check{\beta}]^2 \lesssim \epsilon_0^2. \tag{4.31}$$

*Proof.* We recall the following equations from Propositions 2.37 and 2.38:

$$\begin{aligned}\nabla_4 \widetilde{\underline{\beta}}^{(1)} + \text{tr } \chi \widetilde{\underline{\beta}}^{(1)} &= d_1^*(\widetilde{\rho}^{(1)}, \widetilde{\sigma}^{(1)}) + \mathfrak{L}[\nabla_4 \widetilde{\underline{\beta}}^{(1)}] + \text{Err}[\nabla_4 \widetilde{\underline{\beta}}^{(1)}], \\ \nabla_3(\widetilde{\rho}^{(1)}, \widetilde{\sigma}^{(1)}) + \frac{3}{2} \text{tr } \chi(\widetilde{\rho}^{(1)}, \widetilde{\sigma}^{(1)}) &= -d_1 \widetilde{\underline{\beta}}^{(1)} + \mathfrak{L}[\nabla_3 \widetilde{\rho}^{(1)}, \nabla_3 \widetilde{\sigma}^{(1)}] + \text{Err}[\nabla_3 \widetilde{\rho}^{(1)}, \nabla_3 \widetilde{\sigma}^{(1)}],\end{aligned}$$

where

$$\mathfrak{L}[\nabla_3 \widetilde{\rho}^{(1)}, \nabla_3 \widetilde{\sigma}^{(1)}] = \Gamma_b^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot \check{\underline{\alpha}}^{(1)}, \quad \mathfrak{L}[\nabla_4 \widetilde{\underline{\beta}}^{(1)}] = \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\underline{\beta}}^{(1)}.$$

Applying (4.11) with  $\psi_{(1)} = (\widetilde{\rho}^{(1)}, \widetilde{\sigma}^{(1)})$ ,  $\psi_{(2)} = \widetilde{\underline{\beta}}^{(1)}$ ,  $a_{(1)} = 1$ ,  $a_{(2)} = \frac{3}{2}$ ,  $h_{(1)} = \mathfrak{L}[\nabla_3 \widetilde{\rho}^{(1)}, \nabla_3 \widetilde{\sigma}^{(1)}] + \text{Err}[\nabla_3 \widetilde{\rho}^{(1)}, \nabla_3 \widetilde{\sigma}^{(1)}]$ ,  $h_{(2)} = \mathfrak{L}[\nabla_4 \widetilde{\underline{\beta}}^{(1)}] + \text{Err}[\nabla_4 \widetilde{\underline{\beta}}^{(1)}]$  and  $p = 2$ , we obtain

$$\begin{aligned}& \int_{C_u^V} r^2 |(\widetilde{\rho}^{(1)}, \widetilde{\sigma}^{(1)})|^2 + \int_{\underline{C}_u^V} r^2 |\widetilde{\underline{\beta}}^{(1)}|^2 \\ & \lesssim \int_{\Sigma_0 \cap V} r^2 \left( |(\widetilde{\rho}^{(1)}, \widetilde{\sigma}^{(1)})|^2 + |\widetilde{\underline{\beta}}^{(1)}|^2 \right) + \int_V r |(\widetilde{\rho}^{(1)}, \widetilde{\sigma}^{(1)})|^2 \\ & \quad + \int_V r^2 |(\widetilde{\rho}^{(1)}, \widetilde{\sigma}^{(1)})| |\mathfrak{L}[\nabla_3 \widetilde{\rho}^{(1)}, \nabla_3 \widetilde{\sigma}^{(1)}] + \text{Err}[\nabla_3 \widetilde{\rho}^{(1)}, \nabla_3 \widetilde{\sigma}^{(1)}]| \\ & \quad + \int_V r^2 |\widetilde{\underline{\beta}}^{(1)}| |\mathfrak{L}[\nabla_4 \widetilde{\underline{\beta}}^{(1)}] + \text{Err}[\nabla_4 \widetilde{\underline{\beta}}^{(1)}]|.\end{aligned}$$

First, we have

$$\int_{\Sigma_0 \cap V} r^2 \left( |(\widetilde{\rho}^{(1)}, \widetilde{\sigma}^{(1)})|^2 + |\widetilde{\underline{\beta}}^{(1)}|^2 \right) \lesssim |u|^{2-s} \int_{\Sigma_0 \cap V} r^s \left( |(\widetilde{\rho}^{(1)}, \widetilde{\sigma}^{(1)})|^2 + |\widetilde{\underline{\beta}}^{(1)}|^2 \right) \lesssim \frac{\epsilon_0^2}{|u|^{s-2}}. \quad (4.32)$$

Notice that we have from Proposition 4.10 that

$$\int_V r |(\widetilde{\rho}^{(1)}, \widetilde{\sigma}^{(1)})|^2 \lesssim \int_{|u|}^u r^{-3} d\underline{u} \int_{\underline{C}_u^V} r^4 |(\widetilde{\rho}^{(1)}, \widetilde{\sigma}^{(1)})|^2 \lesssim \int_{|u|}^u r^{-3} d\underline{u} \frac{\epsilon_0^2}{|u|^{s-4}} \lesssim \frac{\epsilon_0^2}{|u|^{s-2}}. \quad (4.33)$$

Moreover, we have

$$\begin{aligned}& \int_V r^2 |(\widetilde{\rho}^{(1)}, \widetilde{\sigma}^{(1)})| |\mathfrak{L}[\nabla_3 \widetilde{\rho}^{(1)}, \nabla_3 \widetilde{\sigma}^{(1)}]| + r^2 |\widetilde{\underline{\beta}}^{(1)}| |\mathfrak{L}[\nabla_4 \widetilde{\underline{\beta}}^{(1)}]| \\ & \lesssim \int_V M r^{-1} |\check{\rho}^{(1)}| |\Gamma_b^{(1)}| + M^2 r^{-1} |\check{\rho}^{(1)}| |\check{\underline{\alpha}}^{(1)}| \\ & \lesssim M \int_{|u|}^u \frac{d\underline{u}}{r^3} \int_{\underline{C}_u^V} |r^2 \check{\rho}^{(1)}| |\Gamma_b^{(1)}| + M^2 \int_{|u|}^u \frac{d\underline{u}}{r^3} \int_{\underline{C}_u^V} |r^2 \check{\rho}^{(1)}| |\check{\underline{\alpha}}^{(1)}| \\ & \lesssim M \int_{|u|}^u \frac{d\underline{u}}{r^3} \left( \int_{\underline{C}_u^V} |r^2 \check{\rho}^{(1)}|^2 \right)^{\frac{1}{2}} \left( \int_{u_0(u)}^u du |\Gamma_b^{(1)}|_{2,S}^2 \right)^{\frac{1}{2}} \\ & \quad + M^2 \int_{|u|}^u \frac{d\underline{u}}{r^3} \left( \int_{\underline{C}_u^V} |r^2 \check{\rho}^{(1)}|^2 \right)^{\frac{1}{2}} \left( \int_{\underline{C}_u^V} |\check{\underline{\alpha}}^{(1)}|^2 \right)^{\frac{1}{2}} \\ & \lesssim M \int_{|u|}^u \frac{d\underline{u}}{r^3} \frac{\epsilon}{|u|^{\frac{s-4}{2}}} \frac{\epsilon}{|u|^{\frac{s-2}{2}}} + M^2 \int_{|u|}^u \frac{d\underline{u}}{r^3} \frac{\epsilon}{|u|^{\frac{s-4}{2}}} \frac{\epsilon}{|u|^{\frac{s}{2}}} \\ & \lesssim \frac{M \epsilon^2}{|u|^{s-1}} + \frac{M^2 \epsilon^2}{|u|^s} \\ & \lesssim \frac{\epsilon_0^2}{|u|^{s-2}}.\end{aligned} \quad (4.34)$$

Next, we have from Proposition 4.12 in [25] that

$$\int_V r^2 |(\widetilde{\rho^{(1)}}, \widetilde{\sigma^{(1)}})| |\text{Err}[\nabla_3 \widetilde{\rho^{(1)}}, \nabla_3 \widetilde{\sigma^{(1)}}]| + r^2 |\widetilde{\beta^{(1)}}| |\text{Err}[\nabla_4 \widetilde{\beta^{(1)}}]| \lesssim \frac{\epsilon_0^2}{|u|^{s-2}}. \quad (4.35)$$

Combining (4.32), (4.33), (4.34) and (4.35), we obtain

$$\int_{C_u^V} r^2 |(\widetilde{\rho^{(1)}}, \widetilde{\sigma^{(1)}})|^2 + \int_{\underline{C}_u^V} r^2 |\widetilde{\beta^{(1)}}|^2 \lesssim \frac{\epsilon_0^2}{|u|^{s-2}}.$$

Combining with Lemma 4.6, this concludes the proof of Proposition 4.11.  $\square$

#### 4.5 Estimates for the Bianchi pair $(\check{\beta}, \check{\alpha})$

**Proposition 4.12.** *We have the following estimate:*

$$\mathcal{R}_0[\check{\beta}]^2 + \mathcal{R}_0[\check{\alpha}]^2 + \mathcal{R}_1[\check{\beta}]^2 + \mathcal{R}_1[\check{\alpha}]^2 \lesssim \epsilon_0^2. \quad (4.36)$$

*Proof.* We recall the following Bianchi equations from Propositions 2.37 and 2.38:

$$\begin{aligned} \nabla_4 \widetilde{\alpha^{(1)}} + \frac{1}{2} \text{tr} \chi \widetilde{\alpha^{(1)}} &= -\nabla \widetilde{\beta^{(1)}} + \mathfrak{L}[\nabla_4 \widetilde{\alpha^{(1)}}] + \text{Err}[\nabla_4 \widetilde{\alpha^{(1)}}], \\ \nabla_3 \check{\beta} + 2 \text{tr} \chi \check{\beta} &= -\text{div} \widetilde{\alpha^{(1)}} + \mathfrak{L}[\nabla_3 \widetilde{\beta^{(1)}}] + \text{Err}[\nabla_3 \widetilde{\beta^{(1)}}], \end{aligned} \quad (4.37)$$

where

$$\mathfrak{L}[\nabla_4 \widetilde{\alpha^{(1)}}] = \Gamma_b^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\alpha}^{(1)}, \quad \mathfrak{L}[\nabla_3 \widetilde{\beta^{(1)}}] = \Gamma_b^{(1)} \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot \check{\beta}^{(1)}.$$

Applying (4.11) with  $\psi_{(1)} = \widetilde{\beta^{(1)}}$ ,  $\psi_{(2)} = \widetilde{\alpha^{(1)}}$ ,  $a_{(1)} = 2$ ,  $a_{(2)} = \frac{1}{2}$ ,  $h_{(1)} = \mathfrak{L}[\nabla_3 \widetilde{\beta^{(1)}}] + \text{Err}[\nabla_3 \widetilde{\beta^{(1)}}]$ ,  $h_{(2)} = \mathfrak{L}[\nabla_4 \widetilde{\alpha^{(1)}}] + \text{Err}[\nabla_4 \widetilde{\alpha^{(1)}}]$  and  $p = 0$ , we obtain

$$\begin{aligned} &\int_{C_u^V} |\widetilde{\beta^{(1)}}|^2 + \int_{\underline{C}_u^V} |\widetilde{\alpha^{(1)}}|^2 \\ &\lesssim \int_{\Sigma_0 \cap V} |\widetilde{\beta^{(1)}}|^2 + |\widetilde{\alpha^{(1)}}|^2 + \int_V r^{-1} |\widetilde{\beta^{(1)}}|^2 \\ &\quad + \int_V |\widetilde{\beta^{(1)}}| |\mathfrak{L}[\nabla_3 \widetilde{\beta^{(1)}}] + \text{Err}[\nabla_3 \widetilde{\beta^{(1)}}]| + |\widetilde{\alpha^{(1)}}| |\mathfrak{L}[\nabla_4 \widetilde{\alpha^{(1)}}] + \text{Err}[\nabla_4 \widetilde{\alpha^{(1)}}]|. \end{aligned} \quad (4.38)$$

First, we have

$$\int_{\Sigma_0 \cap V} |\widetilde{\beta^{(1)}}|^2 + |\widetilde{\alpha^{(1)}}|^2 \lesssim |u|^{-s} \int_{\Sigma_0 \cap V} r^s \left( |\widetilde{\beta^{(1)}}|^2 + |\widetilde{\alpha^{(1)}}|^2 \right) \lesssim \frac{\epsilon_0^2}{|u|^s}.$$

We recall from Proposition 4.11 that

$$\int_V r^{-1} |\widetilde{\beta^{(1)}}|^2 \lesssim \int_{|u|}^u \frac{du}{r^3} \int_{\underline{C}_u^V} |r \check{\beta}^{(1)}|^2 \lesssim \int_{|u|}^u \frac{du}{r^3} \frac{\epsilon_0^2}{|u|^{s-2}} \lesssim \frac{\epsilon_0^2}{|u|^s}.$$

Next, we estimate

$$\begin{aligned}
& \int_V |\widetilde{\beta^{(1)}}| |\mathfrak{L}[\nabla_3 \widetilde{\beta^{(1)}}]| + |\widetilde{\alpha^{(1)}}| |\mathfrak{L}[\nabla_4 \widetilde{\alpha^{(1)}}]| \\
& \lesssim \int_V M r^{-3} |\check{\alpha}^{(1)}| |\Gamma_b^{(1)}| + M r^{-2} |\check{\alpha}^{(1)}|^2 \\
& \lesssim M \int_{|u|}^u \frac{du}{r^3} \left( \int_{\underline{C}_u^V} |\check{\alpha}^{(1)}|^2 \right)^{\frac{1}{2}} \left( \int_{u_0(u)}^u |\Gamma_b^{(1)}|_{2,S}^2 \right)^{\frac{1}{2}} + M \int_{|u|}^u \frac{du}{r^2} \int_{\underline{C}_u^V} |\check{\alpha}^{(1)}|^2 \\
& \lesssim M \int_{|u|}^u \frac{du}{r^3} \frac{\epsilon}{|u|^{\frac{s}{2}}} \frac{\epsilon}{|u|^{\frac{s-2}{2}}} + M \int_{|u|}^u \frac{du}{r^2} \frac{\epsilon^2}{|u|^s} \\
& \lesssim \frac{\epsilon_0^2}{|u|^s}.
\end{aligned}$$

We recall from Proposition 4.13 in [25]

$$\int_V |\widetilde{\beta^{(1)}}| |\text{Err}[\nabla_3 \widetilde{\beta^{(1)}}]| + |\widetilde{\alpha^{(1)}}| |\text{Err}[\nabla_4 \widetilde{\alpha^{(1)}}]| \lesssim \frac{\epsilon_0^2}{|u|^s}.$$

Hence, we obtain

$$\int_{C_u^V} |\widetilde{\beta^{(1)}}|^2 + \int_{\underline{C}_u^V} |\widetilde{\alpha^{(1)}}|^2 \lesssim \frac{\epsilon_0^2}{|u|^s}.$$

This concludes the proof of Proposition 4.12.  $\square$

## 4.6 End of the proof of Theorem M1

**Proposition 4.13.** *We have the following estimate:*

$$\mathcal{R}_0^S + \underline{\mathcal{R}}_0^S \lesssim \mathcal{R}_0 + \mathcal{R}_1 + \underline{\mathcal{R}}_0 + \underline{\mathcal{R}}_1. \quad (4.39)$$

*Proof.* It is sufficient to prove that

$$\mathcal{R}_0^S \lesssim \mathcal{R}_0 + \mathcal{R}_1, \quad \underline{\mathcal{R}}_0^S \lesssim \underline{\mathcal{R}}_0 + \underline{\mathcal{R}}_1, \quad (4.40)$$

which is a direct consequence of Propositions 2.37, 2.38 and 2.45.<sup>23</sup>  $\square$

Remark that Propositions 4.9–4.13 imply

$$\mathcal{R}^2 \lesssim \epsilon_0^2.$$

This concludes the proof of Theorem M1.

## 5 Ricci coefficients estimates (Theorem M3)

The goal of this section is to prove Theorem M3, which we recall below for convenience.

**Theorem M3.** *Assume that*

$$\mathcal{O}_{(0)} \leq \epsilon_0, \quad \mathcal{R} \lesssim \epsilon_0, \quad \mathcal{O}^*(\underline{C}_*) \lesssim \epsilon_0, \quad \mathcal{O} \leq \epsilon, \quad \text{Osc} \leq \epsilon. \quad (5.1)$$

*Then, we have*

$$\mathcal{O} \lesssim \epsilon_0, \quad \text{Osc} \lesssim \epsilon_0. \quad (5.2)$$

**Remark 5.1.** *Throughout Section 5, we denote*

$$S_0(\underline{u}) := S(u_0(\underline{u}), \underline{u}), \quad S_*(u) := S(u, \underline{u}_*), \quad S := S(u, \underline{u}).$$

---

<sup>23</sup>Note that the initial conditions in Proposition 2.45 can be controlled as in Lemma 6.6 of [25].

## 5.1 Preliminaries

### 5.1.1 Null frame transformation

In order to control  $\text{Osc}$ , we will need to compare the double null frame of  $\mathcal{K}$  and the one of  $\mathcal{K}_{(0)}$  in  $\mathcal{K}_{(0)}$ . To this end, we will rely on the null frame transformations introduced in [17] and [18].

**Lemma 5.2.** *Given two null frames  $(e_3, e_4, e_1, e_2)$  and  $(e'_3, e'_4, e'_1, e'_2)$  associated to double-null foliations, assume that they have the same generator  $\underline{L}$  for the ingoing direction. Then, a null frame transformation from the null frame  $(e_3, e_4, e_1, e_2)$  to  $(e'_3, e'_4, e'_1, e'_2)$  can be written in the form:*

$$\begin{aligned} e'_4 &= \lambda \left( e_4 + f^B e_B + \frac{1}{4} |f|^2 e_3 \right), \\ e'_3 &= \lambda^{-1} e_3, \\ e'_A &= e_A + \frac{1}{2} f_A e_3, \end{aligned} \tag{5.3}$$

where

$$\lambda = \frac{\Omega}{\Omega'} \tag{5.4}$$

is a scalar function and  $f$  is a 1-form. Moreover, the inverse transform of (5.3) is given by

$$\begin{aligned} e_4 &= \lambda^{-1} e'_4 - f^A e'_A + \frac{\lambda}{4} |f|^2 e'_3, \\ e_3 &= \lambda e'_3, \\ e_A &= e'_A - \frac{\lambda}{2} f_A e'_3. \end{aligned} \tag{5.5}$$

*Proof.* Applying Lemma 3.1 in [18] with  $\underline{f} = 0$ , we obtain (5.3). Notice that (5.5) is a direct consequence of (5.3). Recalling

$$e'_3 = 2\Omega' \underline{L}, \quad e_3 = 2\Omega \underline{L},$$

we deduce immediately (5.4). This concludes the proof of Lemma 5.2.  $\square$

**Proposition 5.3.** *Under the null frame transformation (5.3), some of the Ricci coefficients transform as follows:*

$$\begin{aligned} \lambda^{-1} \text{tr } \chi' &= \text{tr } \chi + \text{div}' f + l.o.t., \\ \text{curl}' f &= l.o.t., \\ \lambda \underline{\chi}' &= \underline{\chi}, \\ \eta' &= \eta + \frac{1}{2} \lambda \nabla'_{e'_3} f + l.o.t., \\ \underline{\eta}' &= \underline{\eta} + \frac{1}{4} \text{tr } \underline{\chi} f + l.o.t., \\ \lambda^{-2} \xi' &= \xi + \frac{1}{2} \nabla'_{\lambda^{-1} e'_4} f + \frac{1}{4} \text{tr } \chi f + l.o.t., \\ \lambda^{-1} \omega' &= \omega + l.o.t., \end{aligned}$$

where the terms denoted by *l.o.t.* have the following schematic form:

$$l.o.t. = \Gamma_b \cdot f + \mathbf{O}_2^1 \cdot f.$$

The curvature components transform as follows:

$$\begin{aligned}\lambda^{-2}\alpha' &= \alpha + (f\widehat{\otimes}\beta - {}^*f\widehat{\otimes}{}^*\beta) + \left(f\widehat{\otimes}f - \frac{1}{2}{}^*f\widehat{\otimes}{}^*f\right)\rho + \frac{3}{2}(f\widehat{\otimes}{}^*f)^*\rho + l.o.t., \\ \lambda^2\underline{\alpha}' &= \underline{\alpha}, \\ \lambda^{-1}\beta' &= \beta + \frac{3}{2}(f\rho + {}^*f\sigma) + l.o.t., \\ \lambda\underline{\beta}' &= \underline{\beta} - \frac{1}{2}\underline{\alpha} \cdot f, \\ \rho' &= \rho - f \cdot \underline{\beta} + l.o.t., \\ \sigma' &= \sigma - f \cdot {}^*\underline{\beta} + l.o.t.,\end{aligned}$$

where the terms denoted by *l.o.t.* have the following schematic form:

$$l.o.t. = O(f^3)(\rho, \sigma) + O(f^2)(\alpha, \beta, \underline{\beta}, \underline{\alpha}). \quad (5.6)$$

*Proof.* All the transformation formulae, with the exception of the one for  $\omega$ , follow immediately from Proposition 3.3 in [18], in view of the fact that  $\underline{f} = 0$ . See Proposition 6.2 in [25] for the proof of the transformation formula for  $\omega$  which uses in addition that  $\lambda = \frac{\Omega}{\Omega'}$ .  $\square$

### 5.1.2 Difference of the pull backs of Kerr values

The goal of this section is to compare the difference of the pull backs of Kerr values by  $\Phi$  and  $\Phi_{(0)}$  in  $\mathcal{K}_{(0)}$  which will appear in the control of  $\text{Osc}$ . In the remainder of this paper, for any quantity  $X$ , we denote the quantity  $X_{(0)}$  associated to the initial data layer region  $\mathcal{K}_{(0)}$  by

$$X' := X_{(0)}, \quad (5.7)$$

to simplify the notations. For example:

$$u' := u_{(0)}, \quad \eta' := \eta_{(0)}, \quad \mathcal{O}' := \mathcal{O}_{(0)}, \quad \mathfrak{R}' := \mathfrak{R}_{(0)}, \quad S' := S_{(0)}, \quad \mathcal{K}' := \mathcal{K}_{(0)}. \quad (5.8)$$

**Lemma 5.4.** *Let  $X$  be a scalar defined in  $\mathcal{K}'$ . We denote*

$$X'_K := \Phi'^*(X_{Kerr}), \quad X_K := \Phi^*(X_{Kerr}),$$

and we assume that

$$X_K = \mathbf{O}_q^k.$$

Then, we have:

$$\sup_S |X'_K - X_K| \lesssim \frac{M^k \epsilon}{r^{q+\frac{s-1}{2}}} \lesssim \frac{\epsilon_0}{r^{q-k+\frac{s-1}{2}}}.$$

*Proof.* Let  $p \in \mathcal{K}$  which has the coordinates  $(u, \underline{u}, x^1, x^2)$  and resp.  $(u', \underline{u}, x'^1, x'^2)$  in two different foliations. Then, we have

$$\begin{aligned}
|X'_K(p) - X_K(p)| &\lesssim |\Phi'^*(X_{Kerr})(p) - \Phi^*(X_{Kerr})(p)| \\
&\lesssim |X_{Kerr}(\Phi'(p)) - X_{Kerr}(\Phi(p))| \\
&\lesssim \left| \frac{\partial X_{Kerr}}{\partial u_{Kerr}}(u' - u) \right| + \left| \frac{\partial X_{Kerr}}{\partial x_{Kerr}^A}(x'^A - x^A) \right| \\
&\lesssim \frac{M^k}{r^{q+1}} |u' - u| + \frac{M^k}{r^q} |x'^A - x^A| \\
&\lesssim \frac{M^k}{r^{q+1}} \frac{\text{Osc}(u)}{r^{\frac{s-3}{2}}} + \frac{M^k}{r^q} \frac{\text{Osc}(x)}{r^{\frac{s-1}{2}}} \\
&\lesssim \frac{M^k}{r^{q+1}} \frac{\epsilon}{r^{\frac{s-3}{2}}} + \frac{M^k}{r^q} \frac{\epsilon}{r^{\frac{s-1}{2}}} \\
&\lesssim \frac{\epsilon_0}{r^{q-k+\frac{s-1}{2}}},
\end{aligned} \tag{5.9}$$

where we used (5.1) to bound  $\text{Osc}(u)$  and  $\text{Osc}(x)$ . This concludes the proof of Lemma 5.4.  $\square$

### 5.1.3 Evolution lemma

**Lemma 5.5** (Evolution lemma). *Assume that the spacetime  $\mathcal{K}$  is foliated by  $(u, \underline{u})$ .*

(1) *Let  $U, F$  be  $k$ -covariant  $S$ -tangent tensor fields satisfying the outgoing evolution equation*

$$\nabla_4 U + \lambda_0 \text{tr} \chi U = F, \tag{5.10}$$

where  $\lambda_0 \geq 0$ . Denoting  $\lambda_1 = 2(\lambda_0 - \frac{1}{p})$ , we have along  $C_u$

$$|r^{\lambda_1} U|_{p,S} \lesssim |r^{\lambda_1} U|_{p,S_*(u)} + \int_{\underline{u}(p)}^{\underline{u}_*(p)} |r^{\lambda_1} F|_{p,S}(u, \underline{u}') d\underline{u}', \tag{5.11}$$

where  $\underline{u}_*(p)$  is such that  $C_{u(p)} \cap \underline{C}_* = S(u(p), \underline{u}_*(p))$  and where  $S_*(u) = S(u(p), \underline{u}_*(p))$ .

(2) *Let  $V, \underline{F}$  be  $k$ -covariant  $S$ -tangent tensor fields satisfying the ingoing evolution equation*

$$\nabla_3 V + \lambda_0 \text{tr} \underline{\chi} V = \underline{F}. \tag{5.12}$$

Denoting  $\lambda_1 = 2(\lambda_0 - \frac{1}{p})$ , we have along  $\underline{C}_u$

$$|r^{\lambda_1} V|_{p,S} \lesssim |r^{\lambda_1} V|_{p,S_0(\underline{u})} + \int_{u_0(\underline{u})}^u |r^{\lambda_1} \underline{F}|_{p,S}(u', \underline{u}) du', \tag{5.13}$$

where  $S(u_0(\underline{u}), \underline{u}) := C_{u_0(\underline{u})} \cap \underline{C}_u \subset \mathcal{K}$  is the unique sphere of  $\underline{C}_u$  which is located in the future of  $\Sigma_0$  and touches  $\Sigma_0$  and where  $S_0(\underline{u}) = S(u_0(\underline{u}), \underline{u})$ .

*Proof.* See Lemma 4.1.5 in [14].  $\square$

### 5.1.4 Bootstrap control in the initial data layer

**Lemma 5.6.** *We denote*

$$\Gamma_w := \Gamma_g \cup \Gamma_b \cup \Gamma'_g \cup \Gamma'_b. \tag{5.14}$$

Then, we have the following estimates in  $\mathcal{K}'$ :

$$\begin{aligned} \sup_{\mathcal{K}'} r^{\frac{s+1}{2}} |\Gamma_w| &\lesssim \epsilon, \\ \sup_{S \subset \mathcal{K}'} r^{\frac{s+1}{2}} |r^{-\frac{2}{p}} \Gamma_w^{(1)}|_{p,S} &\lesssim \epsilon, \quad p \in [2, 4], \end{aligned} \quad (5.15)$$

Moreover, we have

$$f \in r\Gamma_w, \quad \lambda \in r\Gamma_w, \quad u' - u \in r^2\Gamma_w, \quad r' - r \in r^2\Gamma_w.$$

*Proof.* It follows directly from Lemma 4.1 and the assumption (5.1).  $\square$

## 5.2 Estimates for $\mathcal{O}_{[1]}$ , $\mathcal{O}_\gamma$ and $\text{Osc}$

In the remainder of Section 5, we always assume  $p \in [2, 4]$ .

### 5.2.1 Estimate for $\mathring{\mathcal{O}}_{0,1}(\widetilde{\Omega\omega})$

**Proposition 5.7.** *We have the following estimate for  $q = 0, 1$ :*

$$|r^{\frac{s+1}{2} - \frac{2}{p}} (r\nabla)^q \widetilde{\Omega\omega}|_{p, S_0(\underline{u})} \lesssim \epsilon_0, \quad (5.16)$$

where we recall that  $S_0(\underline{u})$  is the unique leaf of  $\underline{C}_{\underline{u}}$  which is located in the future of  $\Sigma_0$  and touches  $\Sigma_0$ .

*Proof.* We recall from Proposition 5.3 and  $\lambda = \frac{\Omega}{\Omega'}$  that

$$\Omega'\omega' = \Omega\omega + f \cdot \Gamma_w + \mathbf{O}_2^1 \cdot f. \quad (5.17)$$

Notice from Proposition 2.28 and Lemma 5.4 that

$$|\Omega_K\omega_K - \Omega'_K\omega'_K| \lesssim \frac{\epsilon_0}{r^{\frac{s+1}{2}}}.$$

Hence, we have

$$\begin{aligned} |\widetilde{\Omega'\omega'} - \widetilde{\Omega\omega}| &\lesssim |\Omega'\omega' - \Omega\omega| + |\Omega'_K\omega'_K - \Omega_K\omega_K| \\ &\lesssim |\Gamma_w \cdot f| + |\mathbf{O}_2^1 \cdot f| + \frac{\epsilon_0}{r^{\frac{s+1}{2}}} \\ &\lesssim \frac{\epsilon_0}{r^{\frac{s+1}{2}}}. \end{aligned}$$

Thus, we infer

$$|r^{\frac{s+1}{2} - \frac{2}{p}} \widetilde{\Omega\omega}|_{p, S_0(\underline{u})} \lesssim |r^{\frac{s+1}{2} - \frac{2}{p}} \widetilde{\Omega'\omega'}|_{p, S_0(\underline{u})} + \epsilon_0 \lesssim \epsilon_0.$$

We have from (5.17) and (5.5)

$$\begin{aligned} e_A(\Omega\omega) &= e'_A(\Omega'\omega') - \frac{\lambda}{2} f_A e'_3(\Omega'\omega') + \Gamma_w^{(1)} \cdot f + \Gamma_w \cdot f^{(1)} + \mathbf{O}_2^1 \cdot f^{(1)} \\ &= e'_A(\Omega'\omega') + \Gamma_w \cdot \Gamma_w^{(1)} + \mathbf{O}_2^1 \cdot \Gamma_w^{(1)}. \end{aligned}$$

Notice that we have from Lemma 5.4

$$|e_A(\Omega_K\omega_K) - e'_A(\Omega'_K\omega'_K)| \lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}}.$$

Hence, we have

$$|r^{-\frac{2}{p}} e_A(\widetilde{\Omega\omega})|_{p,S_0(\underline{u})} \lesssim |e'_A(\widetilde{\Omega'\omega'})|_{\infty,S_0(\underline{u})} + \frac{M}{r^2} \frac{\epsilon}{r^{\frac{s+1}{2}}} + \frac{\epsilon^2}{r^{s+1}} + \frac{\epsilon_0}{r^{\frac{s+3}{2}}} \lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}},$$

which implies

$$|r^{\frac{s+3}{2}-\frac{2}{p}} e_A(\widetilde{\Omega\omega})|_{p,S_0(\underline{u})} \lesssim \epsilon_0.$$

This concludes the proof of Proposition 5.7.  $\square$

### 5.2.2 Estimates for $\mathcal{O}_0(\widetilde{\Omega\omega})$ and $\mathcal{O}_0(\widetilde{\Omega\underline{\omega}})$

**Proposition 5.8.** *We have the following estimates:*

$$|r^{2+\frac{s-3}{6}-\frac{2}{p}} |u|^{\frac{s-3}{3}} \widetilde{\Omega\omega}|_{p,S} \lesssim \epsilon_0, \quad (5.18)$$

$$|r^{2-\frac{2}{p}} |u|^{\frac{s-3}{2}} \widetilde{\Omega\underline{\omega}}|_{p,S} \lesssim \epsilon_0. \quad (5.19)$$

*Proof.* We recall from Proposition 2.34

$$\nabla_4(\widetilde{\Omega\underline{\omega}}) = \mathbf{O}_0^0 \cdot \check{\rho} + \Gamma_g \cdot \mathbf{O}_2^1 + \Gamma_g \cdot \Gamma_g + r\Gamma_g \cdot \check{\rho}.$$

Applying Lemma 5.5, we infer

$$\begin{aligned} |r^{-\frac{2}{p}} \widetilde{\Omega\underline{\omega}}|_{p,S} &\lesssim |r^{-\frac{2}{p}} \widetilde{\Omega\underline{\omega}}|_{p,S_*(u)} + \int_{\underline{u}}^{u_*} \left( |r^{-\frac{2}{p}} \check{\rho}|_{p,S} + \frac{M}{r^2} |r^{-\frac{2}{p}} \Gamma_g|_{p,S} + |r^{-\frac{2}{p}} \Gamma_g \cdot \Gamma_g|_{p,S} \right) d\underline{u} \\ &\lesssim \frac{\epsilon_0}{r^2 |u|^{\frac{s-3}{2}}} + \int_{\underline{u}}^{u_*} \left( \frac{\epsilon_0}{r^3 |u|^{\frac{s-3}{2}}} + \frac{\epsilon M}{r^4 |u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r^4 |u|^{s-3}} \right) d\underline{u} \\ &\lesssim \frac{\epsilon_0}{r^2 |u|^{\frac{s-3}{2}}}, \end{aligned}$$

which implies (5.19). Next, we recall from Proposition 2.34

$$\nabla_3(\widetilde{\Omega\omega}) = \mathbf{O}_0^0 \cdot \check{\rho} + \Gamma_g \cdot \mathbf{O}_2^1 + \Gamma_g \cdot \Gamma_g + r\Gamma_g \cdot \check{\rho}.$$

Applying Lemma 5.5 and Proposition 5.7, we deduce

$$\begin{aligned} |r^{-\frac{2}{p}} \widetilde{\Omega\omega}|_{p,S} &\lesssim |r^{-\frac{2}{p}} \widetilde{\Omega\omega}|_{p,S_0(\underline{u})} + \int_{u_0(\underline{u})}^u \left( |r^{-\frac{2}{p}} \check{\rho}|_{p,S} + \frac{M}{r^2} |r^{-\frac{2}{p}} \Gamma_g|_{p,S} + |r^{-\frac{2}{p}} \Gamma_g \cdot \Gamma_g|_{p,S} \right) du \\ &\lesssim \frac{\epsilon_0}{r^{\frac{s+1}{2}}} + \int_{u_0(\underline{u})}^u \left( \frac{\epsilon_0}{r^3 |u|^{\frac{s-3}{2}}} + \frac{\epsilon M}{r^3 |u|^{\frac{s-1}{2}}} + \frac{\epsilon^2}{r^4 |u|^{s-3}} \right) du \\ &\lesssim \frac{\epsilon_0}{r^{2+\frac{s-3}{6}} |u|^{\frac{s-3}{3}}}, \end{aligned}$$

which implies (5.18). This concludes the proof of Proposition 5.8.  $\square$

### 5.2.3 Estimates for $\mathcal{O}_1(\widetilde{\Omega\omega})$ and $\mathcal{O}_1(\widetilde{\Omega\underline{\omega}})$

**Proposition 5.9.** *We have the following estimates:*

$$|r^{3+\frac{s-3}{6}-\frac{2}{p}} |u|^{\frac{s-3}{3}} \nabla \widetilde{\Omega\omega}|_{p,S(u,\underline{u})} \lesssim \epsilon_0, \quad (5.20)$$

$$|r^{2-\frac{2}{p}} |u|^{\frac{s-1}{2}} \nabla \widetilde{\Omega\underline{\omega}}|_{p,S(u,\underline{u})} \lesssim \epsilon_0. \quad (5.21)$$

*Proof.* We recall from Propositions 2.34

$$\Omega \nabla_4(\widetilde{\Omega \underline{\omega}}) = \frac{1}{2} \widetilde{\Omega^2 \rho} + \Gamma_g \cdot \mathbf{O}_2^1 + \Gamma_g \cdot \Gamma_g.$$

Differentiating it by  $r \nabla$  and applying Corollary 2.43, we infer

$$\nabla_4(r \nabla \widetilde{\Omega \underline{\omega}}) = \frac{\Omega}{2} r \nabla \check{\rho} + \Gamma_g^{(1)} \cdot \mathbf{O}_2^1 + \Gamma_g \cdot \Gamma_g^{(1)}. \quad (5.22)$$

Since we cannot control  $\nabla \check{\rho}$  in  $L^p$  directly, we use a renormalization argument. We recall from Proposition 2.37

$$\nabla_4 \check{\underline{\beta}} + \text{tr } \chi \check{\underline{\beta}} = -\nabla \check{\rho} + {}^* \nabla \check{\sigma} + \Gamma_b^{(1)} \cdot \mathbf{O}_3^1 + r^{-1} \Gamma_g \cdot \Gamma_b^{(1)},$$

which implies from (4.14)

$$\nabla_4(\Omega r \check{\underline{\beta}}) + \frac{1}{2} \text{tr } \chi(\Omega r \check{\underline{\beta}}) = -\Omega r \nabla \check{\rho} + \Omega r {}^* \nabla \check{\sigma} + \Gamma_b^{(1)} \cdot \mathbf{O}_2^1 + \Gamma_g \cdot \Gamma_b^{(1)}. \quad (5.23)$$

We define  $\check{\underline{\omega}}^\dagger$  by

$$\nabla_4 \check{\underline{\omega}}^\dagger = \frac{1}{2} \Omega \check{\sigma}, \quad \check{\underline{\omega}}^\dagger = 0 \quad \text{on } \underline{C}_*.$$

Applying Proposition 2.41, we deduce

$$\Omega \nabla_4(r {}^* \nabla \check{\underline{\omega}}^\dagger) = r {}^* \nabla(\Omega \nabla_4 \check{\underline{\omega}}^\dagger) + [\Omega \nabla_4, r {}^* \nabla] \check{\underline{\omega}}^\dagger = \frac{1}{2} \Omega r {}^* \nabla \check{\sigma} + \Gamma_g \cdot r \nabla \check{\underline{\omega}}^\dagger + \Gamma_g \cdot \Gamma_g^{(1)}. \quad (5.24)$$

We introduce a new quantity

$$\underline{\mathcal{Z}} := -r \nabla \widetilde{\Omega \underline{\omega}} + r {}^* \nabla \check{\underline{\omega}}^\dagger - \frac{1}{2} r \check{\underline{\beta}}.$$

Then, we obtain from (5.22), (5.23) and (5.24)

$$\begin{aligned} \nabla_4 \underline{\mathcal{Z}} &= -\nabla_4(r \nabla \widetilde{\Omega \underline{\omega}}) + \nabla_4(r {}^* \nabla \check{\underline{\omega}}^\dagger) - \frac{1}{2} \nabla_4(r \check{\underline{\beta}}) \\ &= \Gamma_b^{(1)} \cdot O\left(\frac{M}{r^2}\right) + \Gamma_g \cdot \Gamma_b^{(1)} + \Gamma_g \cdot r \nabla \check{\underline{\omega}}^\dagger. \end{aligned} \quad (5.25)$$

Applying Lemma 5.5, we obtain

$$\begin{aligned} |r^{-\frac{2}{p}} \underline{\mathcal{Z}}|_{p,S} &\lesssim |r^{-\frac{2}{p}} \underline{\mathcal{Z}}|_{p,S_*(u)} + \int_{\underline{u}}^{u_*} \left( |r^{-\frac{2}{p}} \Gamma_g \cdot \Gamma_b^{(1)}|_{p,S} + \frac{M}{r^2} |r^{-\frac{2}{p}} \Gamma_b^{(1)}|_{p,S} + |r^{-\frac{2}{p}} \Gamma_g \cdot r \nabla \check{\underline{\omega}}^\dagger|_{p,S} \right) d\underline{u} \\ &\lesssim \frac{\epsilon_0}{r |u|^{\frac{s-1}{2}}} + \int_{\underline{u}}^{u_*} \left( \frac{\epsilon^2}{r^3 |u|^{s-2}} + \frac{M}{r^3} \frac{\epsilon}{|u|^{\frac{s-1}{2}}} + \frac{\epsilon}{r^2 |u|^{\frac{s-3}{2}}} |r^{1-\frac{2}{p}} \nabla \check{\underline{\omega}}^\dagger|_{p,S} \right) d\underline{u} \\ &\lesssim \frac{\epsilon_0}{r |u|^{\frac{s-1}{2}}} + \int_{\underline{u}}^{u_*} \left( \frac{\epsilon}{r^2 |u|^{\frac{s-3}{2}}} |r^{1-\frac{2}{p}} \nabla \check{\underline{\omega}}^\dagger|_{p,S} \right) d\underline{u}. \end{aligned}$$

Denoting  $\langle \underline{\omega} \rangle := (\widetilde{\Omega \underline{\omega}}, \check{\underline{\omega}}^\dagger)$ , then<sup>24</sup>

$$d_1^* \langle \underline{\omega} \rangle = r^{-1} \underline{\mathcal{Z}} + \frac{1}{2} \check{\underline{\beta}}. \quad (5.26)$$

---

<sup>24</sup>Recall that  $d_1^*(f, g) = -\nabla f + {}^* \nabla g$ .

Applying Proposition 2.15 to (5.26), we obtain

$$\begin{aligned} |r^{1-\frac{2}{p}} \nabla \widetilde{\Omega \omega}|_{p,S} + |r^{1-\frac{2}{p}} \nabla \widetilde{\omega}^\dagger|_{p,S} &\lesssim |r^{-\frac{2}{p}} \underline{\omega}|_{p,S} + |r^{1-\frac{2}{p}} \underline{\beta}|_{p,S} \\ &\lesssim \frac{\epsilon_0}{r|u|^{\frac{s-1}{2}}} + \int_{\underline{u}}^{u_*} \left( \frac{\epsilon}{r^2|u|^{\frac{s-3}{2}}} |r^{1-\frac{2}{p}} \nabla \widetilde{\omega}^\dagger|_{p,S} \right) d\underline{u}. \end{aligned}$$

By Gronwall inequality, we deduce

$$|r^{1-\frac{2}{p}} \nabla \widetilde{\Omega \omega}|_{p,S} + |r^{1-\frac{2}{p}} \nabla \widetilde{\omega}^\dagger|_{p,S} \lesssim \frac{\epsilon_0}{r|u|^{\frac{s-1}{2}}},$$

which implies (5.21).

Next, we recall from Proposition 2.34

$$\Omega \nabla_3(\widetilde{\Omega \omega}) = \frac{1}{2} \widetilde{\Omega^2 \rho} + \Gamma_g \cdot \mathbf{O}_2^1 + \Gamma_g \cdot \Gamma_g.$$

Differentiating it by  $r\nabla$  and applying Corollary 2.43, we infer

$$\nabla_3(r\nabla \widetilde{\Omega \omega}) = \frac{\Omega}{2} r\nabla \check{\rho} + \Gamma_g^{(1)} \cdot \mathbf{O}_2^1 + \Gamma_g \cdot \Gamma_g^{(1)}. \quad (5.27)$$

We recall from Proposition 2.37

$$\nabla_3 \check{\beta} + \text{tr} \underline{\chi} \check{\beta} = \nabla \check{\rho} + {}^* \nabla \check{\sigma} + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + r^{-1} \Gamma_g \cdot \Gamma_b^{(1)},$$

which implies from (4.13)

$$\nabla_3(\Omega r \check{\beta}) + \frac{1}{2} \text{tr} \underline{\chi}(\Omega r \check{\beta}) = \Omega r \nabla \check{\rho} + \Omega r {}^* \nabla \check{\sigma} + \Gamma_g^{(1)} \cdot \mathbf{O}_2^1 + \Gamma_g \cdot \Gamma_b^{(1)}. \quad (5.28)$$

We define  $\check{\omega}^\dagger$  by

$$\nabla_3 \check{\omega}^\dagger = \frac{1}{2} \Omega \check{\sigma}, \quad \check{\omega}^\dagger = 0 \quad \text{on } S_0(\underline{u}) \quad \forall \underline{u}.$$

Applying Proposition 2.41, we deduce

$$\nabla_3(r^* \nabla \check{\omega}^\dagger) = \frac{1}{2} r^* \nabla \check{\sigma} + \Gamma_b \cdot r \nabla \check{\omega}^\dagger + \Gamma_g \cdot \Gamma_b^{(1)}. \quad (5.29)$$

Combining (5.27), (5.28) and (5.29), we obtain

$$\nabla_3 \varkappa = \Gamma_b^{(1)} \cdot \mathbf{O}_2^1 + \Gamma_g \cdot \Gamma_b^{(1)} + \Gamma_b \cdot r \nabla \check{\omega}^\dagger, \quad (5.30)$$

where

$$\varkappa := r \nabla \widetilde{\Omega \omega} + r^* \nabla \check{\omega}^\dagger - \frac{1}{2} r \check{\beta}. \quad (5.31)$$

Applying Lemma 5.5, we infer

$$\begin{aligned} |r^{1-\frac{2}{p}} \varkappa|_{p,S} &\lesssim |r^{1-\frac{2}{p}} \varkappa|_{p,S_0(\underline{u})} + \int_{-\underline{u}}^u \left( |r^{-\frac{2}{p}} \Gamma_g \cdot \Gamma_b^{(1)}|_{p,S} + \frac{M}{r^2} |r^{-\frac{2}{p}} \Gamma_b^{(1)}|_{p,S} + |r^{-\frac{2}{p}} \Gamma_b \cdot r \nabla \check{\omega}^\dagger|_{p,S} \right) d\underline{u} \\ &\lesssim \frac{\epsilon_0}{r^{\frac{s-1}{2}}} + \int_{-\underline{u}}^u \left( \frac{\epsilon^2}{r^3|u|^{s-2}} + \frac{M\epsilon}{r^3|u|^{\frac{s-1}{2}}} + \frac{\epsilon}{r|u|^{\frac{s-1}{2}}} |r^{1-\frac{2}{p}} \nabla \check{\omega}^\dagger|_{p,S} \right) d\underline{u} \\ &\lesssim \frac{\epsilon_0}{r^{2+\frac{s-3}{6}}|u|^{\frac{s-3}{3}}} + \int_{-\underline{u}}^u \frac{\epsilon}{r|u|^{\frac{s-1}{2}}} |r^{1-\frac{2}{p}} \nabla \check{\omega}^\dagger|_{p,S} d\underline{u}. \end{aligned}$$

Applying Proposition 2.15, we obtain

$$\begin{aligned} |r^{2-\frac{2}{p}} \nabla \widetilde{\Omega} \omega|_{p,S} + |r^{2-\frac{2}{p}} \nabla \widetilde{\omega}^\dagger|_{p,S} &\lesssim |r^{1-\frac{2}{p}} \mathcal{K}|_{p,S} + |r^{2-\frac{2}{p}} \widetilde{\beta}|_{p,S} + \int_{-\underline{u}}^u \frac{\epsilon}{r|u|^{\frac{s-1}{2}}} |r^{1-\frac{2}{p}} \nabla \widetilde{\omega}^\dagger|_{p,S} d\underline{u} \\ &\lesssim \frac{\epsilon_0}{r^{2+\frac{s-3}{6}} |u|^{\frac{s-3}{3}}} + \int_{-\underline{u}}^u \frac{\epsilon}{r|u|^{\frac{s-1}{2}}} |r^{1-\frac{2}{p}} \nabla \widetilde{\omega}^\dagger|_{p,S} d\underline{u}. \end{aligned}$$

By Gronwall inequality, we deduce

$$|r^{2-\frac{2}{p}} \nabla \widetilde{\Omega} \omega|_{p,S} + |r^{2-\frac{2}{p}} \nabla \widetilde{\omega}^\dagger|_{p,S} \lesssim \frac{\epsilon_0}{r^{2+\frac{s-3}{6}} |u|^{\frac{s-3}{3}}},$$

which implies (5.20). This concludes the proof of Proposition 5.9.  $\square$

#### 5.2.4 Estimate for $\mathcal{O}_{0,1}(\widetilde{\Omega})$

**Proposition 5.10.** *We have the following estimate for  $q = 0, 1$ :*

$$\left| r^{1-\frac{2}{p}} (r \nabla)^q |u|^{\frac{s-3}{2}} \widetilde{\Omega} \right|_{p,S} \lesssim \epsilon_0. \quad (5.32)$$

*Proof.* We recall from Proposition 2.40

$$\nabla_4 \widetilde{\Omega} = \mathbf{O}_0^0 \cdot \widetilde{\Omega} \omega + \mathbf{O}_1^1 \cdot \Gamma_g + r \Gamma_g \cdot \Gamma_g,$$

which implies from Corollary 2.43

$$\nabla_4 (r \nabla \widetilde{\Omega}) = \mathbf{O}_0^0 \cdot r \nabla (\widetilde{\Omega} \omega) + \mathbf{O}_1^1 \cdot \Gamma_g^{(1)} + r \Gamma_g \cdot \Gamma_g^{(1)}.$$

Applying Lemma 5.5, we have

$$\begin{aligned} |r^{-\frac{2}{p}} \widetilde{\Omega}|_{p,S} &\lesssim |r^{-\frac{2}{p}} \widetilde{\Omega}|_{p,S_*(u)} + \int_{\underline{u}}^{u_*} \left( |r^{-\frac{2}{p}} \widetilde{\omega}|_{p,S} + \frac{M}{r} |r^{-\frac{2}{p}} \Gamma_g|_{p,S} + |r^{1-\frac{2}{p}} \Gamma_g \cdot \Gamma_g| \right) d\underline{u} \\ &\lesssim \frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}} + \int_{\underline{u}}^{u_*} \left( \frac{\epsilon_0}{r^2 |u|^{\frac{s-3}{2}}} + \frac{M}{r^2} \frac{\epsilon}{r|u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r^3 |u|^{s-3}} \right) d\underline{u} \\ &\lesssim \frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}}, \end{aligned}$$

and similarly

$$\begin{aligned} |r^{1-\frac{2}{p}} \nabla \widetilde{\Omega}|_{p,S} &\lesssim |r^{1-\frac{2}{p}} \nabla \widetilde{\Omega}|_{p,S_*(u)} + \int_{\underline{u}}^{u_*} \left( |r^{1-\frac{2}{p}} \nabla \widetilde{\Omega} \omega|_{p,S} + \frac{M}{r} |r^{-\frac{2}{p}} \Gamma_g^{(1)}|_{p,S} + |r^{1-\frac{2}{p}} \Gamma_g \cdot \Gamma_g^{(1)}| \right) d\underline{u} \\ &\lesssim \frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}}. \end{aligned}$$

This concludes the proof of Proposition 5.10.  $\square$

#### 5.2.5 Estimate for $\text{Osc}(u)$

**Proposition 5.11.** *We have the following estimate:*

$$\sup_{\mathcal{K}'} r^{\frac{s-3}{2}} |u' - u| \lesssim \epsilon_0. \quad (5.33)$$

*Proof.* We have from (5.3)

$$\begin{aligned}
(\Omega' e'_4 - \Omega' e'_3)(u' - u) &= -1 - (\Omega' e'_4 - \Omega' e'_3) u \\
&= -1 - \Omega \left( e_4 + f^B e_B + \frac{1}{4} |f|^2 e_3 \right) u + \frac{\Omega'^2}{\Omega^2} \\
&= -\frac{1}{4\Omega} |f|^2 + \frac{\Omega'^2 - \Omega^2}{\Omega^2} \\
&= \mathbf{O}_0^0 \cdot (\Omega' - \Omega) + r^2 \Gamma_w \cdot \Gamma_w.
\end{aligned}$$

Applying Propositions 5.10, 2.46 and Lemma 5.4, we obtain

$$|\Omega' - \Omega| \lesssim |\check{\Omega}' - \check{\Omega}| + \left| \left( \Omega'_K - \frac{1}{2} \right) - \left( \Omega_K - \frac{1}{2} \right) \right| \lesssim \frac{\epsilon_0}{r^{\frac{s-1}{2}}}.$$

Integrating it along  $e'_4 - e'_3$ , we deduce

$$|u' - u|_{\infty, S'} \lesssim \int_w^{u_*} \left( \frac{\epsilon_0}{r^{\frac{s-1}{2}}} + \frac{\epsilon^2}{r^{s-1}} \right) dw \lesssim \frac{\epsilon_0}{r^{\frac{s-3}{2}}}.$$

This concludes the proof of Proposition 5.11.  $\square$

### 5.2.6 Estimate for $\mathcal{O}_{0,1}(\widetilde{\Omega \operatorname{tr} \chi})$

**Proposition 5.12.** *We have the following estimate for  $q = 0, 1$ :*

$$\left| r^{2-\frac{2}{p}} |u|^{\frac{s-3}{2}} (r \nabla)^q \widetilde{\Omega \operatorname{tr} \chi} \right|_{p,S} \lesssim \epsilon_0. \quad (5.34)$$

*Proof.* We recall from Proposition 2.34

$$\nabla_4(\widetilde{\Omega \operatorname{tr} \chi}) + \operatorname{tr} \chi \widetilde{\Omega \operatorname{tr} \chi} = \mathbf{O}_1^0 \cdot \widetilde{\Omega \omega} + \Gamma_g \cdot \mathbf{O}_2^1 + \Gamma_g \cdot \Gamma_g. \quad (5.35)$$

Applying Propositions 5.8 and Lemma 5.5, we infer

$$\begin{aligned}
|r^{2-\frac{2}{p}} \widetilde{\Omega \operatorname{tr} \chi}|_{p,S} &\lesssim |r^{2-\frac{2}{p}} \widetilde{\Omega \operatorname{tr} \chi}|_{p,S_*(u)} + \int_{\underline{u}}^{u_*} \left( |r^{1-\frac{2}{p}} \widetilde{\Omega \omega}|_{p,S} + \frac{M}{r^2} |r^{2-\frac{2}{p}} \Gamma_g|_{p,S} + |r^{2-\frac{2}{p}} \Gamma_g \cdot \Gamma_g|_{p,S} \right) d\underline{u} \\
&\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}} + \int_{\underline{u}}^{u_*} \left( \frac{\epsilon_0}{r^{1+\frac{s-3}{6}} |u|^{\frac{s-3}{3}}} + \frac{M}{r^2} \frac{\epsilon}{|u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r^2 |u|^{s-3}} \right) d\underline{u} \\
&\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}.
\end{aligned}$$

Next, differentiating (5.35) by  $r \nabla$  and applying Corollary 2.43, we obtain

$$\nabla_4(r \nabla \widetilde{\Omega \operatorname{tr} \chi}) + \operatorname{tr} \chi (r \nabla \widetilde{\Omega \operatorname{tr} \chi}) = \mathbf{O}_0^0 \cdot \nabla \widetilde{\Omega \omega} + \Gamma_g^{(1)} \cdot \mathbf{O}_2^1 + \Gamma_g \cdot \Gamma_g^{(1)}.$$

Applying Lemma 5.5, we deduce

$$\begin{aligned}
&|r^{3-\frac{2}{p}} \nabla \widetilde{\Omega \operatorname{tr} \chi}|_{p,S} \\
&\lesssim |r^{3-\frac{2}{p}} \nabla \widetilde{\Omega \operatorname{tr} \chi}|_{p,S_*(u)} + \int_{\underline{u}}^{u_*} \left( |r^{2-\frac{2}{p}} \nabla \widetilde{\Omega \omega}|_{p,S} + M |r^{-\frac{2}{p}} \Gamma_g^{(1)}|_{p,S} + |r^{2-\frac{2}{p}} \Gamma_g \cdot \Gamma_g^{(1)}|_{p,S} \right) d\underline{u} \\
&\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}} + \int_{\underline{u}}^{u_*} \left( \frac{\epsilon_0}{r^{1+\frac{s-3}{6}} |u|^{\frac{s-3}{3}}} + \frac{M}{r^2 |u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r^2 |u|^{s-3}} \right) d\underline{u} \\
&\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}.
\end{aligned}$$

This concludes the proof of Proposition 5.12.  $\square$

### 5.2.7 Estimate for $\mathring{\mathcal{O}}_0(\Omega^{-1} \text{tr } \underline{\chi})$

**Proposition 5.13.** *We have the following estimate:*

$$|r^{\frac{s+1}{2}-\frac{2}{p}} \widetilde{\Omega^{-1} \text{tr } \underline{\chi}}|_{p, S_0(\underline{u})} \lesssim \epsilon_0. \quad (5.36)$$

*Proof.* We have from Proposition 5.3

$$\Omega'^{-1} \text{tr } \underline{\chi}' = \Omega^{-1} \text{tr } \underline{\chi},$$

which implies

$$\widetilde{\Omega'^{-1} \text{tr } \underline{\chi}'} = \widetilde{\Omega^{-1} \text{tr } \underline{\chi}} + \Omega_K^{-1} \text{tr}_K \underline{\chi}_K - \Omega_K'^{-1} \text{tr}_K \underline{\chi}'_K. \quad (5.37)$$

Proceeding as in (5.9) and applying Propositions 2.16 and 5.11

$$\begin{aligned} \left| \Omega_K^{-1} \text{tr}_K \underline{\chi}_K - \Omega_K'^{-1} \text{tr}_K \underline{\chi}'_K \right| &\lesssim |\partial_u(\Omega^{-1} \text{tr } \underline{\chi})_{Kerr}| |u' - u| + |\partial_{x^A}(\Omega^{-1} \text{tr } \underline{\chi})_{Kerr}| |x'^A - x^A| \\ &\lesssim \frac{1}{r^2} |u' - u| + \frac{M^2}{r^3} |r \Gamma_w| \\ &\lesssim \frac{1}{r^2} \frac{\epsilon_0}{r^{\frac{s-3}{2}}} + \frac{M^2}{r^3} \frac{\epsilon}{r^{\frac{s-1}{2}}} \\ &\lesssim \frac{\epsilon_0}{r^{\frac{s+1}{2}}}. \end{aligned}$$

Injecting it into (5.37), we deduce

$$\sup_{\mathcal{K}'} r^{\frac{s+1}{2}} |\widetilde{\Omega^{-1} \text{tr } \underline{\chi}}| \lesssim \sup_{\mathcal{K}'} r^{\frac{s+1}{2}} |\widetilde{\Omega'^{-1} \text{tr } \underline{\chi}'}| + \sup_{\mathcal{K}'} r^{\frac{s+1}{2}} |\Omega_K^{-1} \text{tr}_K \underline{\chi}_K - \Omega_K'^{-1} \text{tr}_K \underline{\chi}'_K| \lesssim \epsilon_0.$$

This concludes the proof of Proposition 5.13.  $\square$

### 5.2.8 Estimate for $\mathcal{O}_0(\Omega \text{tr } \underline{\chi})$

**Proposition 5.14.** *We have the following estimate:*

$$|r^{2-\frac{2}{p}} |u|^{\frac{s-3}{2}} \widetilde{\Omega \text{tr } \underline{\chi}}|_{p, S} \lesssim \epsilon_0. \quad (5.38)$$

*Proof.* Next, we recall from Proposition 2.34

$$\nabla_3(\widetilde{\Omega \text{tr } \underline{\chi}}) + \text{tr } \underline{\chi} \widetilde{\Omega \text{tr } \underline{\chi}} = \mathbf{O}_1^0 \cdot \widetilde{\Omega \underline{\omega}} + \Gamma_g \cdot \mathbf{O}_2^1 + \Gamma_g \cdot \Gamma_b.$$

Notice that we have from Propositions 5.10 and 5.13

$$|r^{2-\frac{2}{p}} \widetilde{\Omega \text{tr } \underline{\chi}}|_{p, S_0(\underline{u})} \lesssim |r^{2-\frac{2}{p}} \widetilde{\Omega^{-1} \text{tr } \underline{\chi}}|_{p, S_0(\underline{u})} + |r^{2-\frac{2}{p}} \widetilde{\Omega^2}|_{p, S_0(\underline{u})} + |r^{2-\frac{2}{p}} \widetilde{\Omega \text{tr } \underline{\chi}} \widetilde{\Omega^2}|_{p, S_0(\underline{u})} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}.$$

Applying Lemma 5.5, we infer

$$\begin{aligned} |r^{2-\frac{2}{p}} \widetilde{\Omega \text{tr } \underline{\chi}}|_{p, S} &\lesssim |r^{2-\frac{2}{p}} \widetilde{\Omega \text{tr } \underline{\chi}}|_{p, S_0(\underline{u})} + \int_{u_0(\underline{u})}^u \left( |r^{1-\frac{2}{p}} \widetilde{\Omega \underline{\omega}}|_{p, S} + \frac{M}{r^2} |r^{2-\frac{2}{p}} \Gamma_g|_{p, S} + |r^{2-\frac{2}{p}} \Gamma_b \cdot \Gamma_b|_{p, S} \right) du \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}} + \int_{u_0(\underline{u})}^u \left( \frac{\epsilon_0}{|u|^{\frac{s-1}{2}}} + \frac{M}{r^2} \frac{\epsilon}{|u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{|u|^{s-1}} \right) du \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}. \end{aligned}$$

This concludes the proof of Proposition 5.14.  $\square$

### 5.2.9 Estimate for $\mathcal{O}_{0,1}(\tilde{\eta})$

**Proposition 5.15.** *We have the following estimate for  $q = 0, 1$ :*

$$|r^{2-\frac{2}{p}}|u|^{\frac{s-3}{2}}(r\nabla)^q\tilde{\eta}|_{p,S} \lesssim \epsilon_0. \quad (5.39)$$

*Proof.* We recall from Proposition 2.36

$$\nabla_4[\tilde{\mu}] + \text{tr } \chi[\tilde{\mu}] = \mathbf{O}_1^0 \cdot \check{\rho} + \mathbf{O}_3^1 \cdot \Gamma_g^{(1)} + r^{-1}\Gamma_b \cdot \Gamma_g^{(1)}. \quad (5.40)$$

Applying Lemma 5.5, we obtain

$$\begin{aligned} |r^{2-\frac{2}{p}}[\tilde{\mu}]|_{p,S} &\lesssim |r^{2-\frac{2}{p}}[\tilde{\mu}]|_{p,S_*(u)} + \int_{\underline{u}}^{u_*} \left( |r^{1-\frac{2}{p}}\check{\rho}|_{p,S} + |r^{1-\frac{2}{p}}\Gamma_b \cdot \Gamma_g^{(1)}|_{p,S} + \frac{M}{r^3}|r^{2-\frac{2}{p}}\Gamma_g^{(1)}|_{p,S} \right) d\underline{u} \\ &\lesssim \frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}} + \int_{\underline{u}}^{u_*} \left( \frac{\epsilon_0}{r^2|u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r^2|u|^{s-1}} + \frac{M}{r^3} \frac{\epsilon}{|u|^{\frac{s-3}{2}}} \right) d\underline{u} \\ &\lesssim \frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}}. \end{aligned}$$

We have from Proposition 2.34

$$\text{div } \check{\eta} = -\check{\mu} - \check{\rho} + \mathbf{O}_3^2 \cdot \Gamma_b^{(1)} + \Gamma_g \cdot \Gamma_b.$$

We recall from (2.27) that

$$[\mu] = \mu + \frac{1}{4} \text{tr } \chi \text{ tr } \underline{\chi},$$

which implies

$$[\tilde{\mu}] = \check{\mu} + \mathbf{O}_1^0 \cdot (\widetilde{\text{tr } \chi}, \widetilde{\text{tr } \underline{\chi}}, r^{-1}\check{\Omega}) + \Gamma_g \cdot \Gamma_g. \quad (5.41)$$

Thus, we obtain

$$\text{div } \check{\eta} = -[\tilde{\mu}] - \check{\rho} + \mathbf{O}_3^2 + \mathbf{O}_1^0 \cdot (\widetilde{\Omega \text{tr } \chi}, \widetilde{\Omega \text{tr } \underline{\chi}}, r^{-1}\check{\Omega}) + \Gamma_g \cdot \Gamma_b. \quad (5.42)$$

Recall the linearized torsion equation in Proposition 2.34:

$$\text{curl } \check{\eta} = \check{\sigma} + \mathbf{O}_3^2 \cdot \Gamma_b^{(1)} + \Gamma_b \cdot \Gamma_g. \quad (5.43)$$

Combining (5.42) and (5.43), applying Propositions 2.15, 5.10 and 5.12, we obtain for  $q = 0, 1$ :

$$\begin{aligned} &|r^{2+q-\frac{2}{p}}\nabla^q\check{\eta}|_{p,S} \\ &\lesssim |r^{3-\frac{2}{p}}([\tilde{\mu}], \check{\rho}, \check{\sigma})|_{p,S} + |r^{2-\frac{2}{p}}(\widetilde{\Omega \text{tr } \chi}, \widetilde{\Omega \text{tr } \underline{\chi}}, r^{-1}\check{\Omega})|_{p,S} + \frac{M^2}{r^3}|r^{3-\frac{2}{p}}\Gamma_b^{(1)}|_{p,S} + |r^{3-\frac{2}{p}}\Gamma_g \cdot \Gamma_b|_{p,S} \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}. \end{aligned}$$

This concludes the proof of Proposition 5.15. □

### 5.2.10 Estimate for $\text{Osc}(f)$

**Proposition 5.16.** *We have the following estimate:*

$$\sup_{\mathcal{K}'} |r^{\frac{s-1}{2}} \mathfrak{d}^{\leq 1} f| \lesssim \epsilon_0. \quad (5.44)$$

*Proof.* We have from Proposition 5.3

$$\Omega' \text{tr } \chi' = \Omega \text{tr } \chi + \text{div } f + f \cdot \Gamma_w + \mathbf{O}_2^1 \cdot f.$$

Applying (5.5) and Proposition 2.34, we have

$$\begin{aligned} e_A(\Omega \text{tr } \chi) &= e_A(\Omega' \text{tr } \chi') - e_A(\text{div } f) + r^{-1} f \cdot \Gamma_w^{(1)} + r^{-1} \Gamma_w \cdot f^{(1)} + \mathbf{O}_3^1 \cdot f^{(1)} \\ &= e'_A(\Omega' \text{tr } \chi') - \frac{\lambda}{2} f_A e'_3(\Omega' \text{tr } \chi') - e_A(\text{div } f) + \Gamma_w \cdot \Gamma_w^{(1)} + \mathbf{O}_2^1 \cdot \Gamma_w^{(1)} \\ &= e'_A(\Omega' \text{tr } \chi') + \frac{\Omega'}{4} f_A \text{tr } \chi' \text{tr } \underline{\chi}' - e_A(\text{div } f) + \Gamma_w \cdot \Gamma_w^{(1)} + \mathbf{O}_2^1 \cdot \Gamma_w^{(1)}. \end{aligned}$$

Recall from Lemma 5.4 that

$$|e'_A(\Omega'_K \text{tr}_K \chi'_K) - e_A(\Omega_K \text{tr}_K \chi_K)| \lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}}.$$

Thus, we infer

$$\begin{aligned} &\left| r^{-\frac{2}{p}} \left( e_A(\text{div } f) - \frac{\Omega'}{4} f_A \text{tr } \chi' \text{tr } \underline{\chi}' + e_A(\widetilde{\Omega \text{tr } \chi}) - e'_A(\widetilde{\Omega' \text{tr } \chi'}) \right) \right|_{p,S} \\ &\lesssim |e'_A(\Omega'_K \text{tr}_K \chi'_K) - e_A(\Omega_K \text{tr}_K \chi_K)|_{\infty,S} + \frac{M^2}{r^3} |r^{-\frac{2}{p}} \cdot \Gamma_w^{(1)}| + |r^{-\frac{2}{p}} \Gamma_w \cdot \Gamma_w^{(1)}|_{p,S} \\ &\lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}} + \frac{M^2}{r^3} \frac{\epsilon}{r^{\frac{s+1}{2}}} + \frac{\epsilon^2}{r^{s+1}} \\ &\lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}}. \end{aligned}$$

Recall that

$$-\frac{\Omega'}{4} f \text{tr } \chi' \text{tr } \underline{\chi}' = \frac{f}{2r^2} + \mathbf{O}_3^1 \cdot f + \Gamma_w \cdot \Gamma_w.$$

Hence, we deduce from Proposition 5.12

$$\begin{aligned} &\left| r^{-\frac{2}{p}} \left( e_A(\text{div } f) + \frac{f}{2r^2} \right) \right|_{p,S} \\ &\lesssim |r^{-\frac{2}{p}} e_A(\widetilde{\Omega \text{tr } \chi})|_{p,S} + |e'_A(\widetilde{\Omega' \text{tr } \chi'})|_{\infty,S} + \frac{M}{r^2} |r^{-\frac{2}{p}} \cdot \Gamma_w^{(1)}|_{p,S} + |r^{-\frac{2}{p}} \Gamma_w \cdot \Gamma_w^{(1)}|_{p,S} + \frac{\epsilon_0}{r^{\frac{s+3}{2}}} \\ &\lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}} + \frac{M}{r^2} \frac{\epsilon}{r^{\frac{s+1}{2}}} + \frac{\epsilon^2}{r^{s+1}} \\ &\lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}}. \end{aligned} \quad (5.45)$$

Next, we have from Proposition 5.3

$$\text{curl } f = \mathbf{O}_2^1 \cdot \Gamma_w + r \Gamma_w \cdot \Gamma_w,$$

which implies

$$\begin{aligned}
|r^{-\frac{2}{p}} e_A(\operatorname{curl} f)|_{p,S} &\lesssim \frac{M^2}{r^3} |r^{-\frac{2}{p}} \cdot \Gamma_w^{(1)}|_{p,S} + |r^{-\frac{2}{p}} \Gamma_w \cdot \Gamma_w^{(1)}|_{p,S} \\
&\lesssim \frac{M^2}{r^3} \frac{\epsilon}{r^{\frac{s+1}{2}}} + \frac{\epsilon^2}{r^{s+1}} \\
&\lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}}.
\end{aligned} \tag{5.46}$$

Combining (5.45) and (5.46), we deduce<sup>25</sup>

$$\left| r^{-\frac{2}{p}} \left( d_1^* d_1 f - \frac{f}{2r^2} \right) \right|_{p,S} \lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}}.$$

We have from Proposition 2.16

$$\begin{aligned}
d_1^* d_1 f - \frac{f}{2r^2} &= (-\Delta_1 + \mathbf{K})f - \frac{f}{2r^2} \\
&= -\Delta_1 f - \frac{f}{4} \operatorname{tr} \chi \operatorname{tr} \underline{\chi} - \frac{f}{2r^2} + r^{-1} \Gamma_w^{(1)} \cdot f + \mathbf{O}_3^1 \cdot f \\
&= -\Delta_1 f + \frac{f}{2r^2} + \mathbf{O}_2^1 \cdot \Gamma_w^{(1)} + \Gamma_w \cdot \Gamma_w^{(1)}.
\end{aligned}$$

Thus, we infer

$$\begin{aligned}
\left| r^{-\frac{2}{p}} \left( -\Delta_1 + \frac{1}{2r^2} \right) f \right|_{p,S} &\lesssim \left| r^{-\frac{2}{p}} \left( d_1^* d_1 - \frac{1}{2r^2} \right) f \right|_{p,S} + \frac{M}{r^2} |r^{-\frac{2}{p}} \Gamma_w^{(1)}|_{p,S} + |r^{-\frac{2}{p}} \Gamma_w \cdot \Gamma_w^{(1)}|_{p,S} \\
&\lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}} + \frac{M}{r^2} \frac{\epsilon}{r^{\frac{s+1}{2}}} + \frac{\epsilon^2}{r^{s+1}} \\
&\lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}}.
\end{aligned}$$

Applying Proposition 2.15, we deduce for  $q = 0, 1, 2$ ,

$$|r^{-\frac{2}{p}} (r \nabla)^q f|_{p,S} \lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}}.$$

Then, applying Proposition 2.46, we obtain

$$\sup_{S \subset \mathcal{K}'} |r^{\frac{s+3}{2}} (r \nabla)^{\leq 1} f|_{\infty,S} \lesssim \epsilon_0. \tag{5.47}$$

Next, we recall from Proposition 5.3

$$\lambda \nabla_{e_3} f = 2\eta' - 2\eta + \mathbf{O}_1^1 \cdot \Gamma_w + r \Gamma_w \cdot \Gamma_w. \tag{5.48}$$

Applying Lemma 5.4, we obtain

$$|\eta'_K(e'_A) - \eta_K(e_A)| \lesssim \frac{\epsilon_0}{r^{\frac{s+1}{2}}}.$$

---

<sup>25</sup>Recall that  $d_1^* d_1 f = d_1^*(\operatorname{div} f, \operatorname{curl} f) = -\nabla \operatorname{div} f + * \nabla \operatorname{curl} f$ .

Applying Proposition 5.15 and (5.47), we deduce

$$\begin{aligned}
|r^{-\frac{2}{p}} \nabla_{e_3} f|_{p,S} &\lesssim |r^{-\frac{2}{p}} (\check{\eta}', \check{\eta})|_{p,S} + M |r^{-1-\frac{2}{p}} \Gamma_w|_{p,S} + |r^{1-\frac{2}{p}} \Gamma_w \cdot \Gamma_w|_{p,S} + \frac{\epsilon_0}{r^{\frac{s+1}{2}}} \\
&\lesssim \frac{\epsilon_0}{r^{\frac{s+1}{2}}} + \frac{M\epsilon}{r^{\frac{s+3}{2}}} + \frac{\epsilon^2}{r^s} \\
&\lesssim \frac{\epsilon_0}{r^{\frac{s+1}{2}}}.
\end{aligned} \tag{5.49}$$

Finally, we recall from Proposition 5.3

$$\frac{1}{2} \nabla_{\lambda^{-1} e_4} f = -\frac{1}{4} \operatorname{tr} \chi f + \mathbf{O}_1^1 \cdot \Gamma_w + r \Gamma_w \cdot \Gamma_w,$$

which implies from (5.47)

$$\sup_{\mathcal{K}'} r^{\frac{s+1}{2}} |\nabla_{e_4} f| \lesssim \sup_{\mathcal{K}'} r^{\frac{s-1}{2}} |f| + \sup_{\mathcal{K}'} \left( \frac{M\epsilon}{r} + \frac{\epsilon^2}{r^{\frac{s-3}{2}}} \right) \lesssim \epsilon_0. \tag{5.50}$$

Combining (5.47), (5.49) and (5.50), we deduce

$$\sup_{\mathcal{K}'} r^{\frac{s-1}{2}} |\mathfrak{d}^{\leq 1} f| \lesssim \epsilon_0.$$

This concludes the proof of Proposition 5.16.  $\square$

### 5.2.11 Estimate for $\check{\mathcal{O}}_{0,1}(\check{\eta})$

**Proposition 5.17.** *We have the following estimate for  $q = 0, 1$ :*

$$|r^{2-\frac{2}{p}} |u|^{\frac{s-3}{2}} (r \nabla)^q \check{\eta}|_{p, S_0(\underline{u})} \lesssim \epsilon_0. \tag{5.51}$$

*Proof.* We recall from Proposition 5.3

$$\underline{\eta} = \underline{\eta}' - \frac{1}{4} \operatorname{tr} \underline{\chi} f + \mathbf{O}_1^1 \cdot \Gamma_w + r \Gamma_w \cdot \Gamma_w. \tag{5.52}$$

Recall that we have from Lemma 5.4

$$|\underline{\eta}_K(e_A) - \underline{\eta}'_K(e'_A)| \lesssim \frac{\epsilon_0}{r^{\frac{s+1}{2}}}.$$

Thus, we obtain

$$\begin{aligned}
|\check{\eta}_A| &\lesssim |\check{\eta}'_A| + |\underline{\eta}_K(e_A) - \underline{\eta}'_K(e'_A)| + \frac{M}{r} |\Gamma_w| + r |\Gamma_w \cdot \Gamma_w| \\
&\lesssim \frac{\epsilon_0}{r^{\frac{s+1}{2}}} + \frac{M}{r} \frac{\epsilon}{r^{\frac{s+1}{2}}} + \frac{\epsilon^2}{r^s} \\
&\lesssim \frac{\epsilon_0}{r^{\frac{s+1}{2}}}.
\end{aligned} \tag{5.53}$$

Differentiating (5.52) by  $e_B$ , we obtain

$$e_B(\underline{\eta}_A) = e'_B(\underline{\eta}'_A) + \mathbf{O}_2^0 \cdot f^{(1)} + \mathbf{O}_2^1 \cdot \Gamma_w^{(1)} + \Gamma_w \cdot \Gamma_w^{(1)}.$$

Applying Lemma 5.4, we have

$$|e_B(\underline{\eta}_K)_A - e'_B(\underline{\eta}'_K)_A| \lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}}. \quad (5.54)$$

Hence, we obtain

$$\begin{aligned} & |r^{-\frac{2}{p}} e_B(\underline{\eta}_A)|_{p,S} \\ & \lesssim |r^{-\frac{2}{p}} e'_B(\underline{\eta}'_A)|_{p,S} + |e_B(\underline{\eta}_K)_A - e'_B(\underline{\eta}'_K)_A| + r^{-2} |f^{(1)}|_{\infty,S} + M |r^{-2-\frac{2}{p}} \Gamma_w^{(1)}|_{p,S} + |r^{-\frac{2}{p}} \Gamma_w \cdot \Gamma_w^{(1)}|_{p,S} \\ & \lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}} + \frac{M\epsilon}{r^{\frac{s+5}{2}}} + \frac{\epsilon^2}{r^{s+1}} \\ & \lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}}. \end{aligned}$$

Combining with (5.53), we infer for  $q = 0, 1$

$$|r^{\frac{s+1}{2}-\frac{2}{p}} (r\nabla)^q \underline{\eta}|_{p,S} \lesssim \epsilon_0.$$

This concludes the proof of Proposition 5.17.  $\square$

### 5.2.12 Estimate for $\mathcal{O}_{0,1}(\underline{\eta})$

**Proposition 5.18.** *We have the following estimate for  $q = 0, 1$ :*

$$|r^{2-\frac{2}{p}} |u|^{\frac{s-3}{2}} (r\nabla)^q \underline{\eta}|_{p,S} \lesssim \epsilon_0. \quad (5.55)$$

*Proof.* We have from Propositions 2.34, 5.12, 5.13 and 5.17 that

$$\begin{aligned} |r^{2-\frac{2}{p}} [\underline{\mu}]|_{p,S_0(\underline{u})} & \lesssim |r^{2-\frac{2}{p}} \underline{\mu}|_{p,S_0(\underline{u})} + |r^{1-\frac{2}{p}} (\widetilde{\text{tr}} \chi, \widetilde{\text{tr}} \underline{\chi})|_{p,S_0(\underline{u})} \\ & \lesssim |r^{2-\frac{2}{p}} \text{div} \underline{\eta}|_{p,S_0(\underline{u})} + |r^{2-\frac{2}{p}} \underline{\rho}|_{p,S_0(\underline{u})} + |r^{1-\frac{2}{p}} (\widetilde{\Omega \text{tr}} \chi, \widetilde{\Omega^{-1} \text{tr}} \underline{\chi})|_{p,S_0(\underline{u})} \\ & \quad + M^2 |r^{-1-\frac{2}{p}} \Gamma_b^{(1)}|_{p,S_0(\underline{u})} + |r^{2-\frac{2}{p}} \Gamma_b \cdot \Gamma_g|_{p,S_0(\underline{u})} \\ & \lesssim \frac{\epsilon_0}{r^{\frac{s-1}{2}}}. \end{aligned} \quad (5.56)$$

We recall from Proposition 2.36

$$\nabla_3 [\underline{\mu}] + \text{tr} \underline{\chi} [\underline{\mu}] = O(r^{-1}) \underline{\rho} + \mathbf{O}_3^1 \cdot \Gamma_b^{(1)} + r^{-1} \Gamma_b \cdot \Gamma_b^{(1)}.$$

Applying Lemma 5.5 and (5.56), we obtain

$$\begin{aligned} |r^{2-\frac{2}{p}} [\underline{\mu}]|_{p,S} & \lesssim |r^{2-\frac{2}{p}} [\underline{\mu}]|_{p,S_0(\underline{u})} + \int_{u_0(\underline{u})}^u \left( |r^{1-\frac{2}{p}} \underline{\rho}|_{p,S} + \frac{M}{r^2} |r^{1-\frac{2}{p}} \Gamma_b^{(1)}|_{p,S} + |r^{1-\frac{2}{p}} \Gamma_b \cdot \Gamma_b^{(1)}|_{p,S} \right) du \\ & \lesssim \frac{\epsilon_0}{r^{\frac{s-1}{2}}} + \int_{u_0(\underline{u})}^u \left( \frac{\epsilon_0}{r^2 |u|^{\frac{s-3}{2}}} + \frac{M}{r^2} \frac{\epsilon}{|u|^{\frac{s-1}{2}}} + \frac{\epsilon^2}{r^2 |u|^{s-1}} \right) du \\ & \lesssim \frac{\epsilon_0}{r |u|^{\frac{s-3}{2}}}. \end{aligned}$$

We have from Proposition 2.34

$$\begin{aligned} \text{div} \underline{\eta} & = -[\underline{\mu}] - \underline{\rho} + \mathbf{O}_3^2 \cdot \Gamma_b^{(1)} + \mathbf{O}_1^0 \cdot (\widetilde{\text{tr}} \chi, \widetilde{\text{tr}} \underline{\chi}, r^{-1} \underline{\Omega}) + \Gamma_g \cdot \Gamma_b, \\ \text{curl} \underline{\eta} & = \underline{\sigma} + \mathbf{O}_3^2 \cdot \Gamma_b^{(1)} + \Gamma_g \cdot \Gamma_b. \end{aligned} \quad (5.57)$$

Applying Proposition 2.15, we deduce for  $q = 0, 1$ :

$$\begin{aligned} |r^{2+q-\frac{2}{p}} \nabla^q \underline{\eta}|_{p,S} &\lesssim |r^{3-\frac{2}{p}}([\underline{\mu}], \underline{\rho})|_{p,S} + |r^{2-\frac{2}{p}}(\widetilde{\text{tr}} \chi, \widetilde{\text{tr}} \underline{\chi}, r^{-1} \check{\Omega})|_{p,S} + M^2 |r^{-\frac{2}{p}} \Gamma_b^{(1)}|_{p,S} + |r^{3-\frac{2}{p}} \Gamma_g \cdot \Gamma_b|_{p,S} \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}} + \frac{M^2}{r|u|^{\frac{s-1}{2}}} + \frac{\epsilon^2}{|u|^{s-2}} \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}. \end{aligned}$$

This concludes the proof of Proposition 5.18.  $\square$

### 5.2.13 Estimate for $\mathcal{O}_{0,1,2}(\check{\Omega})$

**Proposition 5.19.** *We have the following estimates:*

$$|r^{1-\frac{2}{p}}|u|^{\frac{s-3}{2}}(r\nabla)^q \check{\Omega}|_{p,S} \lesssim \epsilon_0, \quad q = 0, 1, 2. \quad (5.58)$$

*Proof.* Recall that

$$\eta + \underline{\eta} = 2\nabla \log \Omega,$$

and

$$\eta_K + \underline{\eta}_K = 2\nabla \log \Omega_K.$$

Taking the difference, we obtain

$$\Delta \log \left( \frac{\Omega}{\Omega_K} \right) = \frac{1}{2} \text{div}(\check{\eta} + \underline{\eta}),$$

Applying Propositions 2.15, 5.15 and 5.18, we have for  $q = 1, 2$ :

$$\left| r^{1+q-\frac{2}{p}} \nabla^q \left( \log \left( \frac{\Omega}{\Omega_K} \right) \right) \right|_{p,S} (u, \underline{u}) \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}. \quad (5.59)$$

Applying Proposition 5.10, we deduce that

$$\left| r \log \left( \frac{\Omega}{\Omega_K} \right) \right|_{\infty,S} \leq \left| r \left( \frac{\Omega}{\Omega_K} - 1 \right) \right|_{\infty,S} \lesssim |r \check{\Omega}|_{\infty,S} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}$$

so that (5.59) also holds true for  $q = 0$ . Hence, we have

$$\left\| \log \frac{\Omega}{\Omega_K} \right\|_{W^{2,4}(S)} \lesssim \frac{\epsilon_0}{r^{\frac{1}{2}}|u|^{\frac{s-3}{2}}}.$$

We recall that the Sobolev space  $W^{2,4}(S)$  forms a Banach algebra, i.e.

$$\|uv\|_{W^{2,4}(S)} \lesssim \|u\|_{W^{2,4}(S)} \|v\|_{W^{2,4}(S)}, \quad \forall u, v \in W^{2,4}(S), \quad (5.60)$$

see for example Theorem 4.39 in [1]. Notice that

$$\check{\Omega} = \Omega_K \left( \frac{\Omega}{\Omega_K} - 1 \right) = \Omega_K \left( \exp \left( \log \left( \frac{\Omega}{\Omega_K} \right) \right) - 1 \right) = \Omega_K \sum_{k=1}^{\infty} \frac{1}{k!} \left( \log \frac{\Omega}{\Omega_K} \right)^k. \quad (5.61)$$

Combining (5.59), (5.60) and (5.61), we deduce

$$|(r\nabla)^{\leq 2} \check{\Omega}|_{4,S} \lesssim \sum_{k=1}^{\infty} \left| (r\nabla)^{\leq 2} \log \left( \frac{\Omega}{\Omega_K} \right) \right|_{4,S}^k \lesssim \frac{\epsilon_0}{r^{\frac{1}{2}}|u|^{\frac{s-3}{2}}},$$

which implies for  $p \in [2, 4]$ :

$$|r^{1-\frac{2}{p}}(r\nabla)^q \check{\Omega}|_{p,S} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}, \quad q = 0, 1, 2.$$

This concludes the proof of Proposition 5.19.  $\square$

### 5.2.14 Estimate for $\mathring{\mathcal{O}}_1(\Omega^{-1} \text{tr } \underline{\chi})$

**Proposition 5.20.** *We have the following estimate:*

$$|r^{\frac{s+3}{2}-\frac{2}{p}} \nabla(\Omega^{-1} \text{tr } \underline{\chi})|_{p, S_0(\underline{u})} \lesssim \epsilon_0. \quad (5.62)$$

*Proof.* We have from Proposition 5.3

$$\Omega'^{-1} \text{tr } \underline{\chi}' = \Omega^{-1} \text{tr } \underline{\chi}.$$

Differentiating it by  $e_A$  and applying (5.5), we obtain

$$e_A(\Omega^{-1} \text{tr } \underline{\chi}) = e'_A(\Omega'^{-1} \text{tr } \underline{\chi}') - \frac{\lambda}{2} f_A e'_3(\Omega'^{-1} \text{tr } \underline{\chi}').$$

Applying Propositions 2.16, 5.16 and Lemma 5.4, we infer

$$\begin{aligned} |r^{-\frac{2}{p}} e_A(\Omega^{-1} \text{tr } \underline{\chi})|_{p, S} &\lesssim |r^{-\frac{2}{p}} e'_A(\Omega'^{-1} \text{tr } \underline{\chi}')|_{p, S} + |e_A(\Omega^{-1} \text{tr } \underline{\chi})_K - e'_A(\Omega^{-1} \text{tr } \underline{\chi})'_K| + |r^{-2-\frac{2}{p}} f_A|_{p, S} \\ &\lesssim \frac{\epsilon_0}{r^{\frac{s+3}{2}}}. \end{aligned}$$

This concludes the proof of Proposition 5.20.  $\square$

### 5.2.15 Estimate for $\mathcal{O}_1(\Omega \text{tr } \underline{\chi})$

**Proposition 5.21.** *We have the following estimates:*

$$|r^{3-\frac{2}{p}} |u|^{\frac{s-3}{2}} \nabla \widetilde{\Omega \text{tr } \underline{\chi}}|_{p, S} \lesssim \epsilon_0. \quad (5.63)$$

*Proof.* Notice that we have

$$\widetilde{\Omega \text{tr } \underline{\chi}} = \Omega^2(\Omega^{-1} \text{tr } \underline{\chi}) - \Omega_K^2(\Omega_K^{-1} \text{tr } \underline{\chi}_K) = \widetilde{\Omega}^2(\Omega^{-1} \text{tr } \underline{\chi}) + \Omega_K^2(\Omega^{-1} \text{tr } \underline{\chi}),$$

which implies

$$\nabla \widetilde{\Omega \text{tr } \underline{\chi}} = \mathbf{O}_1^0 \cdot \nabla \widetilde{\Omega} + \mathbf{O}_0^0 \cdot \widetilde{\text{tr } \underline{\chi}} + \Gamma_g \cdot \Gamma_g^{(1)}.$$

Hence, we have from Propositions 5.10 and 5.20

$$\begin{aligned} |r^{3-\frac{2}{p}} \nabla \widetilde{\Omega \text{tr } \underline{\chi}}|_{p, S_0(\underline{u})} &\lesssim |r^{3-\frac{2}{p}} \nabla \widetilde{\Omega^{-1} \text{tr } \underline{\chi}}|_{p, S_0(\underline{u})} + |r^{2-\frac{2}{p}} \nabla \widetilde{\Omega}|_{p, S_0(\underline{u})} + |r^{3-\frac{2}{p}} \Gamma_g \cdot \Gamma_g^{(1)}|_{p, S_0(\underline{u})} \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}. \end{aligned}$$

Next, we recall from Proposition 2.34

$$\nabla_3(\widetilde{\Omega \text{tr } \underline{\chi}}) + \text{tr } \underline{\chi} \widetilde{\Omega \text{tr } \underline{\chi}} = \mathbf{O}_1^0 \cdot \widetilde{\Omega \omega} + \Gamma_g \cdot \mathbf{O}_2^1 + \Gamma_b \cdot \Gamma_b.$$

Differentiating it by  $r \nabla$  and applying Corollary 2.43, we deduce

$$\nabla_3(r \nabla \widetilde{\Omega \text{tr } \underline{\chi}}) + \text{tr } \underline{\chi} r \nabla \widetilde{\Omega \text{tr } \underline{\chi}} = \mathbf{O}_0^0 \cdot \nabla \widetilde{\Omega \omega} + \Gamma_g^{(1)} \cdot \mathbf{O}_2^1 + \Gamma_b \cdot \Gamma_b^{(1)}.$$

Applying Lemma 5.5 and Proposition 5.9, we infer

$$\begin{aligned} &|r^{3-\frac{2}{p}} \nabla \widetilde{\Omega \text{tr } \underline{\chi}}|_{p, S} \\ &\lesssim |r^{3-\frac{2}{p}} \nabla \widetilde{\Omega \text{tr } \underline{\chi}}|_{p, S_0(\underline{u})} + \int_{u_0(\underline{u})}^u \left( |r^{2-\frac{2}{p}} \nabla \widetilde{\Omega \omega}|_{p, S} + \frac{M}{r^2} |r^{2-\frac{2}{p}} \Gamma_g|_{p, S} + |r^{2-\frac{2}{p}} \Gamma_b \cdot \Gamma_b^{(1)}|_{p, S} \right) du \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}} + \int_{u_0(\underline{u})}^u \left( \frac{\epsilon_0}{|u|^{\frac{s-1}{2}}} + \frac{M}{r^2} \frac{\epsilon}{|u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{|u|^{s-1}} \right) du \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}. \end{aligned}$$

This concludes the proof of Proposition 5.21.  $\square$

### 5.2.16 Estimates for $\mathcal{O}_{0,1}(\check{\chi})$ and $\mathcal{O}_{0,1}(\check{\underline{\chi}})$

**Proposition 5.22.** *We have the following estimates for  $q = 0, 1$ :*

$$\begin{aligned} |r^{2-\frac{2}{p}}|u|^{\frac{s-3}{2}}(r\nabla)^q\check{\chi}|_{p,S} &\lesssim \epsilon_0, \\ |r^{1-\frac{2}{p}}|u|^{\frac{s-1}{2}}(r\nabla)^q\check{\underline{\chi}}|_{p,S} &\lesssim \epsilon_0. \end{aligned} \quad (5.64)$$

*Proof.* We recall from Proposition 2.34

$$\operatorname{div} \check{\chi} = -\check{\beta} + \mathbf{O}_0^0 \cdot \nabla \widetilde{\operatorname{tr} \chi} + \mathbf{O}_1^0 \cdot \check{\zeta} + \mathbf{O}_3^2 \cdot \Gamma_g^{(1)} + \Gamma_g \cdot \Gamma_g.$$

Applying Propositions 2.15, 5.12, 5.15 and 5.18, we deduce for  $q = 0, 1$ :

$$\begin{aligned} |r^{2+q-\frac{2}{p}}\nabla^q\check{\chi}|_{p,S} &\lesssim |r^{3-\frac{2}{p}}\nabla\widetilde{\operatorname{tr} \chi}|_{p,S} + |r^{2-\frac{2}{p}}\check{\zeta}|_{p,S} + |r^{3-\frac{2}{p}}\check{\beta}|_{p,S} + \frac{M^2}{r^2}|r^{2-\frac{2}{p}}\Gamma_g^{(1)}|_{p,S} + |r^{3-\frac{2}{p}}\Gamma_g \cdot \Gamma_g|_{p,S} \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}. \end{aligned}$$

Next, we recall from Proposition 2.34

$$\operatorname{div} \check{\underline{\chi}} = \check{\underline{\beta}} + \mathbf{O}_0^0 \cdot \nabla \widetilde{\operatorname{tr} \underline{\chi}} + \mathbf{O}_1^0 \cdot \check{\zeta} + \mathbf{O}_3^2 \cdot \Gamma_b^{(1)} + \Gamma_b \cdot \Gamma_g.$$

Applying Propositions 2.15, 5.21, 5.15 and 5.18, we deduce for  $q = 0, 1$ :

$$\begin{aligned} |r^{1+q-\frac{2}{p}}\nabla^q\check{\underline{\chi}}|_{p,S} &\lesssim |r^{2-\frac{2}{p}}\nabla\widetilde{\operatorname{tr} \underline{\chi}}|_{p,S} + |r^{1-\frac{2}{p}}\check{\zeta}|_{p,S} + |r^{2-\frac{2}{p}}\check{\underline{\beta}}|_{p,S} + \frac{M^2}{r^2}|r^{1-\frac{2}{p}}\Gamma_b^{(1)}|_{p,S} + |r^{2-\frac{2}{p}}\Gamma_b \cdot \Gamma_g|_{p,S} \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-1}{2}}}. \end{aligned}$$

This concludes the proof of Proposition 5.22.  $\square$

### 5.2.17 Estimates for $\mathcal{O}_{0,1}(\check{\underline{b}})$

**Proposition 5.23.** *We have the following estimate for  $q = 0, 1$ :*

$$|r^{2-\frac{2}{p}}|u|^{\frac{s-3}{2}}(r\nabla)^q\check{\underline{b}}^A|_{p,S} \lesssim \epsilon_0. \quad (5.65)$$

*Proof.* We have from Proposition 2.40

$$\nabla_4(\check{\underline{b}}^A) = \mathbf{O}_1^0 \cdot \check{\zeta} + \mathbf{O}_3^2 \cdot \Gamma_g + \Gamma_g \cdot \Gamma_g. \quad (5.66)$$

Applying Lemma 5.5, we deduce from Propositions 5.15 and 5.18<sup>26</sup>

$$\begin{aligned} |r^{-\frac{2}{p}}\check{\underline{b}}^A|_{p,S} &\lesssim \int_{\underline{u}}^{\underline{u}_*} \left( |r^{-1-\frac{2}{p}}\check{\zeta}|_{p,S} + M^2|r^{-3-\frac{2}{p}}\Gamma_g|_{p,S} + |r^{-\frac{2}{p}}\Gamma_g \cdot \Gamma_g|_{p,S} \right) d\underline{u} \\ &\lesssim \int_{\underline{u}}^{\underline{u}_*} \left( \frac{\epsilon_0}{r^3|u|^{\frac{s-3}{2}}} + \frac{M^2\epsilon}{r^5|u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r^4|u|^{s-3}} \right) d\underline{u} \\ &\lesssim \frac{\epsilon_0}{r^2|u|^{\frac{s-3}{2}}} + \frac{M^2\epsilon}{r^4|u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r^3|u|^{s-3}} \\ &\lesssim \frac{\epsilon_0}{r^2|u|^{\frac{s-3}{2}}}, \end{aligned}$$

---

<sup>26</sup>Recall that  $\check{\underline{b}}^A = 0$  on  $\underline{C}_*$  and  $\zeta = \frac{\eta-\eta}{2}$ .

which implies (5.65) in the case  $q = 0$ . Differentiating (5.66) by  $r\nabla$  and applying Corollary 2.43, we obtain

$$\nabla_4(r\nabla\check{b}^A) = \mathbf{O}_1^0 \cdot r\nabla\check{\zeta} + \mathbf{O}_3^2 \cdot \Gamma_g^{(1)} + \Gamma_g \cdot \Gamma_g^{(1)}.$$

Applying Lemma 5.5, we deduce from Propositions 5.15 and 5.18

$$\begin{aligned} |r^{-\frac{2}{p}}(r\nabla)\check{b}^A|_{p,S} &\lesssim \int_{\underline{u}}^{\underline{u}_*} \left( |r^{-\frac{2}{p}}\nabla\check{\zeta}|_{p,S} + M^2|r^{-3-\frac{2}{p}}\Gamma_g^{(1)}|_{p,S} + |r^{-\frac{2}{p}}\Gamma_g \cdot \Gamma_g^{(1)}|_{p,S} \right) d\underline{u} \\ &\lesssim \frac{\epsilon_0}{r^2|u|^{\frac{s-3}{2}}}, \end{aligned}$$

which implies (5.65) in the case  $q = 1$ . This concludes the proof of Proposition 5.23.  $\square$

### 5.2.18 Estimate for $\text{Osc}(x)$

**Proposition 5.24.** *We have the following estimates*

$$\sup_{\mathcal{K}'} r^{\frac{s-1}{2}} |x'^A - x^A| \lesssim \epsilon_0. \quad (5.67)$$

*Proof.* We have from (5.3) and Proposition 2.28

$$\begin{aligned} (\Omega' e'_4 - \Omega' e'_3)(x'^A - x^A) &= -\underline{b}'^A - \Omega' e'_4(x^A) + \Omega' e'_3(x^A) \\ &= -\underline{b}'^A - \Omega \left( e_4 + f^B e_B + \frac{1}{4}|f|^2 e_3 \right) x^A + \frac{\Omega'^2}{\Omega^2} \underline{b}^A \\ &= \underline{b}^A - \underline{b}'^A - \Omega f^B e_B(x^A) - \frac{1}{4}|f|^2 \underline{b}^A + \left( \frac{\Omega'^2}{\Omega^2} - 1 \right) \underline{b}^A \\ &= \underline{b}^A - \underline{b}'^A + \mathbf{O}_1^0 \cdot f + \mathbf{O}_1^1 \cdot \Gamma_w + r\Gamma_w \cdot \Gamma_w. \end{aligned} \quad (5.68)$$

Notice from Lemma 5.4 that

$$|b'^A - b^A| \lesssim |\check{b}'^A| + |\check{b}^A| + |(b'_K)^A - b_K^A| \lesssim \frac{\epsilon_0}{r^{\frac{s+1}{2}}}.$$

Hence, we obtain from Proposition 5.16

$$|(\Omega' e'_4 - \Omega' e'_3)(x'^A - x^A)| \lesssim r^{-1}|f| + \frac{\epsilon_0}{r^{\frac{s+1}{2}}} + \frac{M}{r} |\Gamma_w| + r |\Gamma_w \cdot \Gamma_w| \lesssim \frac{\epsilon_0}{r^{\frac{s+1}{2}}}.$$

Integrating it along  $e'_4 - e'_3$ , we deduce<sup>27</sup>

$$|x'^A - x^A|_{\infty, S'} \lesssim |x'^A - x^A|_{\infty, \underline{C}_* \cap \mathcal{K}'} + \int_w^{\underline{u}_*} \frac{\epsilon_0}{r^{\frac{s+1}{2}}} dw \lesssim \frac{\epsilon_0}{r^{\frac{s-1}{2}}}.$$

This concludes the proof of Proposition 5.24.  $\square$

---

<sup>27</sup>Recall that  $x'^A = x^A$  on  $\underline{C}_* \cap \Sigma_0$  which is chosen in Section 2.1.5. Together with the equation for  $\Omega' e'_3(x'^A - x^A)$ , we can easily deduce the estimate for  $x'^A - x^A$  on  $\underline{C}_* \cap \mathcal{K}'$ .

### 5.2.19 Estimates for $\mathcal{O}_{0,1}(\check{\gamma})$ and $\mathcal{O}_{0,1}(\check{\epsilon})$

**Proposition 5.25.** *We have the following estimates for  $q = 0, 1$ :*

$$|r^{-1+q-\frac{2}{p}}|u|^{\frac{s-3}{2}}\nabla^q\check{\gamma}_{AB}|_{p,S} \lesssim \epsilon_0, \quad (5.69)$$

$$|r^{-1+q-\frac{2}{p}}|u|^{\frac{s-3}{2}}\nabla^q\check{\epsilon}_{AB}|_{p,S} \lesssim \epsilon_0. \quad (5.70)$$

*Proof.* We recall from Proposition 2.40

$$\nabla_4(\check{\gamma}_{AB}) - \text{tr } \chi(\check{\gamma}_{AB}) = \mathbf{O}_{-2}^0 \cdot (\check{\chi}, \widetilde{\text{tr } \chi}) + \mathbf{O}_{-1}^0 \cdot \check{\Omega} + r^3 \Gamma_g \cdot \Gamma_g.$$

Differentiating it by  $r\nabla$  and applying Corollary 2.43, we deduce

$$\nabla_4(r\nabla\check{\gamma}_{AB}) - \text{tr } \chi(r\nabla\check{\gamma}_{AB}) = \mathbf{O}_{-3}^0 \cdot \nabla(\check{\chi}, \widetilde{\text{tr } \chi}) + \mathbf{O}_{-2}^0 \cdot \nabla\check{\Omega} + r^3 \Gamma_g \cdot \Gamma_g^{(1)}.$$

Applying Lemma 5.5 and Propositions 5.12, 5.19 and 5.22, we obtain for  $q = 0, 1$

$$\begin{aligned} & |r^{-2-\frac{2}{p}}(r\nabla)^q\check{\gamma}_{AB}|_{p,S} \\ & \lesssim |r^{-2-\frac{2}{p}}(r\nabla)^q\check{\gamma}_{AB}|_{p,S_*(u)} + \int_{\underline{u}}^{\underline{u}_*} \left( |r^{-\frac{2}{p}}(r\nabla)^q(\check{\chi}, \widetilde{\text{tr } \chi}, r^{-1}\check{\Omega})|_{p,S} + |r^{-1-\frac{2}{p}}\Gamma_g \cdot \Gamma_g^{(1)}|_{p,S} \right) d\underline{u} \\ & \lesssim \frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}} + \int_{\underline{u}}^{\underline{u}_*} \left( \frac{\epsilon_0}{r^2|u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r^3|u|^{s-3}} \right) d\underline{u} \\ & \lesssim \frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}}, \end{aligned}$$

which implies (5.69). The proof of (5.70) is similar and left to the reader. This concludes the proof of Proposition 5.25.  $\square$

### 5.2.20 Estimate for $\text{Osc}(r)$

**Proposition 5.26.** *We have the following estimate:*

$$\sup_{\mathcal{K}'} r^{\frac{s-3}{2}} |r' - r| \lesssim \epsilon_0. \quad (5.71)$$

*Proof.* Notice that we have

$$\begin{aligned} \det \gamma_{AB} - \det(\gamma_K)_{AB} &= (\gamma_{11}\gamma_{22} - \gamma_{12}^2) - ((\gamma_K)_{11}(\gamma_K)_{22} - (\gamma_K)_{12}^2) \\ &= \check{\gamma}_{11}(\gamma_K)_{22} + (\gamma_K)_{11}\check{\gamma}_{22} - 2\check{\gamma}_{12}(\gamma_K)_{12} + \check{\gamma}_{11}\check{\gamma}_{22} - \check{\gamma}_{12}^2 \\ &= \mathbf{O}_{-2}^0 \cdot \check{\gamma}_{AB} + r^6 \Gamma_g \cdot \Gamma_g. \end{aligned}$$

Hence, we obtain

$$\begin{aligned} 4\pi |r^2 - r_K^2| &= \left| \int_{\mathbb{S}^2} \left( \sqrt{\det \gamma_{AB}} - \sqrt{\det(\gamma_K)_{AB}} \right) d\sigma_{\mathbb{S}^2} \right| \\ &\lesssim \int_{\mathbb{S}^2} \left| \frac{\det \gamma_{AB} - \det(\gamma_K)_{AB}}{\sqrt{\det \gamma_{AB}} + \sqrt{\det(\gamma_K)_{AB}}} \right| d\sigma_{\mathbb{S}^2} \\ &\lesssim |\check{\gamma}_{AB}| + r^4 |\Gamma_g \cdot \Gamma_g| \\ &\lesssim \frac{r\epsilon_0}{|u|^{\frac{s-3}{2}}} + r^4 \frac{\epsilon^2}{r^4 |u|^{s-3}} \\ &\lesssim \frac{r\epsilon_0}{|u|^{\frac{s-3}{2}}}, \end{aligned}$$

which implies

$$\sup_{\mathcal{K}} |r - r_K| \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}. \quad (5.72)$$

Similarly, we have

$$\sup_{\mathcal{K}'} |r' - r'_K| \lesssim \frac{\epsilon_0}{r^{\frac{s-3}{2}}}. \quad (5.73)$$

Proceeding as in (5.9), we obtain for any  $p \in \mathcal{K}'$

$$\begin{aligned} |r'_K(p) - r_K(p)| &= |r_{Kerr}(\Phi'(p)) - r_{Kerr}(\Phi(p))| \\ &\lesssim |\partial_u(r_{Kerr})||u' - u| + |\partial_A(r_{Kerr})||x'^A - x^A| \\ &\lesssim |u' - u| \\ &\lesssim \frac{\epsilon_0}{r^{\frac{s-3}{2}}}, \end{aligned}$$

where we used Proposition 5.11. Combining with (5.72) and (5.73), we infer

$$\sup_{\mathcal{K}'} r^{\frac{s-3}{2}} |r' - r| \lesssim \epsilon_0. \quad (5.74)$$

This concludes the proof of Proposition 5.26.  $\square$

### 5.3 Estimate for $\mathcal{O}_\Gamma$

**Proposition 5.27.** *We have the following estimates:*

$$\begin{aligned} |r^{2-\frac{2}{p}}|u|^{\frac{s-3}{2}}(\check{J}, \check{L})|_{p,S} &\lesssim \epsilon_0, \\ |r^{1-\frac{2}{p}}|u|^{\frac{s-1}{2}}\check{\underline{J}}|_{p,S} &\lesssim \epsilon_0. \end{aligned} \quad (5.75)$$

*Proof.* We recall from Lemma 2.23

$$\begin{aligned} \Gamma_{uA}^B &= \frac{1}{2}\gamma^{BC}\partial_u(\gamma_{AC}), \\ \Gamma_{uA}^B &= \frac{1}{2}\gamma^{BC}\left(\partial_u(\gamma_{AC}) - \frac{\partial}{\partial x^A}(\gamma_{CD}b^D) + \frac{\partial}{\partial x^C}(\gamma_{AD}b^D)\right) + \frac{b^B}{4\Omega^2}\left(2\frac{\partial}{\partial x^A}(\Omega^2) - \frac{\partial}{\partial \underline{u}}(\gamma_{AD}b^D)\right), \\ \Gamma_{BC}^A &= \frac{1}{2}\gamma^{AD}\left(\frac{\partial\gamma_{BD}}{\partial x^C} + \frac{\partial\gamma_{CD}}{\partial x^B} - \frac{\partial\gamma_{BC}}{\partial x^D}\right) + \frac{b^A}{4\Omega^2}\frac{\partial\gamma_{BC}}{\partial \underline{u}}. \end{aligned}$$

Hence, we have from Definition 2.24

$$\check{J}_A^B = \mathbf{O}_2^0 \cdot \partial_{\underline{u}}(\check{\gamma}_{AB}) + \mathbf{O}_3^0 \cdot \check{\gamma}_{AB} + r^{-1}\Gamma_g \cdot \partial_{\underline{u}}(\check{\gamma}_{AB}).$$

We have from Proposition 2.40

$$\partial_{\underline{u}}(\check{\gamma}_{AB}) = \text{tr } \chi(\check{\gamma}_{AB}) + \mathbf{O}_{-2}^0 \cdot (\widetilde{\text{tr } \chi}, \check{\chi}, r^{-1}\check{\Omega}) + r^3\Gamma_g \cdot \Gamma_g,$$

which implies from Propositions 5.12, 5.22, 5.19 and 5.25

$$|r^{-\frac{2}{p}}\partial_{\underline{u}}(\check{\gamma}_{AB})|_{p,S} \lesssim |r^{-1-\frac{2}{p}}\check{\gamma}_{AB}|_{p,S} + |r^{2-\frac{2}{p}}(\widetilde{\text{tr } \chi}, \check{\chi}, r^{-1}\check{\Omega})|_{p,S} + |r^{3-\frac{2}{p}}\Gamma_g \cdot \Gamma_g|_{p,S} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}.$$

Thus, we infer from Proposition 5.25

$$|r^{2-\frac{2}{p}}\check{J}|_{p,S} \lesssim |r^{-1-\frac{2}{p}}\check{\gamma}_{AB}|_{p,S} + |r^{-\frac{2}{p}}\partial_{\underline{u}}(\check{\gamma}_{AB})|_{p,S} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}.$$

Next, we have from Definition 2.24

$$\check{L}_{BC}^A = \mathbf{O}_3^2 \cdot \partial_{\underline{u}}(\check{\gamma}_{AB}) + \mathbf{O}_2^0 \cdot \check{\mathfrak{p}}^{\leq 1} \check{\gamma}_{AB} + \mathbf{O}_{-1}^0 \cdot \check{\underline{b}}^A + r^2 \Gamma_g \cdot \Gamma_g^{(1)}.$$

Thus, we deduce from Propositions 5.25 and 5.23

$$\begin{aligned} |r^{1-\frac{2}{p}} \check{L}_{BC}^A|_{p,S} &\lesssim \frac{M^2}{r^2} |r^{-\frac{2}{p}} \partial_{\underline{u}}(\check{\gamma}_{AB})|_{p,S} + |r^{-1-\frac{2}{p}} \check{\mathfrak{p}}^{\leq 1} \check{\gamma}_{AB}|_{p,S} + |r^{2-\frac{2}{p}} \check{\underline{b}}^A|_{p,S} + |r^{3-\frac{2}{p}} \Gamma_g \cdot \Gamma_g^{(1)}|_{p,S} \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r|u|^{s-3}} \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}, \end{aligned}$$

which implies

$$|r^{2-\frac{2}{p}} \check{L}|_{p,S} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}.$$

Finally, we have from Definition 2.24

$$\check{J}_A^B = \mathbf{O}_2^0 \cdot (\partial_u(\check{\gamma}_{AB}), \partial_{\underline{u}}(\check{\gamma}_{AB}), r^{-1} \check{\mathfrak{p}}^{\leq 1} \check{\gamma}_{AB}) + \mathbf{O}_3^2 \cdot \check{\mathfrak{p}}^{\leq 1} \check{\Omega} + \mathbf{O}_0^0 \cdot \check{\mathfrak{p}}^{\leq 1} \check{\underline{b}}^A + \mathbf{O}_1^2 \cdot \partial_{\underline{u}}(\check{\underline{b}}^A) + r \Gamma_g \cdot \Gamma_g^{(1)}.$$

We recall from Propositions 2.40, 5.12, 5.19, 5.22, 5.23 and 5.25

$$\begin{aligned} |r^{-\frac{2}{p}} \partial_u(\check{\gamma}_{AB})|_{p,S} &\lesssim |r^{-1-\frac{2}{p}} \check{\gamma}_{AB}| + |r^{2-\frac{2}{p}} (\check{\chi}, \widetilde{\text{tr}} \check{\chi}, r^{-1} \check{\Omega}, \check{\underline{b}}^A)|_{p,S} + M^2 |r^{-\frac{2}{p}} \Gamma_g|_{p,S} + |r^{3-\frac{2}{p}} \Gamma_g \cdot \Gamma_g|_{p,S} \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}} + \frac{r\epsilon_0}{|u|^{\frac{s-1}{2}}} + \frac{M^2\epsilon}{r^2|u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r|u|^{s-3}} \\ &\lesssim \frac{r\epsilon_0}{|u|^{\frac{s-1}{2}}}, \end{aligned}$$

and

$$\begin{aligned} |r^{-\frac{2}{p}} \partial_{\underline{u}}(\check{\underline{b}}^A)|_{p,S} &\lesssim |r^{-1-\frac{2}{p}} \check{\zeta}|_{p,S} + M^2 |r^{-3-\frac{2}{p}} \Gamma_g|_{p,S} + |r^{-\frac{2}{p}} \Gamma_g \cdot \Gamma_g|_{p,S} \\ &\lesssim \frac{\epsilon_0}{r^3|u|^{\frac{s-3}{2}}} + \frac{M^2\epsilon}{r^5|u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r^4|u|^{s-3}} \\ &\lesssim \frac{\epsilon_0}{r^3|u|^{\frac{s-3}{2}}}. \end{aligned}$$

Combining the above estimates, we deduce from Propositions 5.19, 5.23 and 5.25

$$\begin{aligned} |r^{1-\frac{2}{p}} \check{J}|_{p,S} &\lesssim |r^{-1-\frac{2}{p}} (\partial_u(\check{\gamma}_{AB}), \partial_{\underline{u}}(\check{\gamma}_{AB}), r^{-1} \check{\mathfrak{p}}^{\leq 1} \check{\gamma}_{AB})|_{p,S} + M^2 |r^{-2-\frac{2}{p}} \check{\mathfrak{p}}^{\leq 1} \check{\Omega}|_{p,S} \\ &\quad + |r^{1-\frac{2}{p}} \check{\mathfrak{p}}^{\leq 1} \check{\underline{b}}^A|_{p,S} + M^2 |r^{-\frac{2}{p}} \check{\underline{b}}^A|_{p,S} + |r^2 \Gamma_g \cdot \Gamma_g^{(1)}|_{p,S} \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-1}{2}}} + \frac{M^2\epsilon_0}{r^3|u|^{\frac{s-3}{2}}} + \frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}} + \frac{M^2\epsilon_0}{r^2|u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r^2|u|^{s-3}} \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-1}{2}}}. \end{aligned}$$

This concludes the proof of Proposition 5.27.  $\square$

In view of Propositions 5.7-5.27, we obtain (5.2). This concludes the proof of Theorem M3.

## 6 Initialization and extension (Theorems M0, M2 and M4)

### 6.1 Control of the initial data layer (Theorem M0)

The goal of this section is to prove Theorem M0, which we recall below for convenience.

**Theorem M0.** *Assume that<sup>28</sup>*

$$\mathcal{O}' \leq \epsilon_0, \quad \mathfrak{R}' \leq \epsilon_0, \quad \mathcal{O} \leq \epsilon, \quad \text{Osc} \leq \epsilon. \quad (6.1)$$

Then, we have

$$\mathfrak{R}_0 \lesssim \epsilon_0. \quad (6.2)$$

*Proof.* We denote for  $e_A$  an orthogonal frame on  $S$

$$R := \{\alpha_{AB}, \beta_A, \rho, \sigma, \underline{\beta}_A, \underline{\alpha}_{AB}\},$$

and for  $e'_A$  an orthogonal frame on  $S'$

$$R' := \{\alpha'_{AB}, \beta'_A, \rho', \sigma', \underline{\beta}'_A, \underline{\alpha}'_{AB}\}.$$

The transformation formulae for curvature in Proposition 5.3 can be written as follows:

$$R = R' + f \cdot R' + \text{l.o.t.}, \quad (6.3)$$

which implies

$$\check{R} = \check{R}' + R_K - R'_K + f \cdot (\check{R}' + R'_K) + \text{l.o.t.}$$

We have from Lemma 5.4

$$|R_K - R'_K| \lesssim \frac{M\epsilon}{r^{\frac{s+5}{2}}}.$$

Hence, we obtain

$$|\check{R}| \lesssim |\check{R}'| + |f \cdot \check{R}'| + \frac{M}{r^3} |f| + |R_K - R'_K| \lesssim |\check{R}'| + \frac{M\epsilon}{r^{\frac{s+5}{2}}},$$

where we used  $\text{Osc}(f) \leq \epsilon$ . Then, we have

$$\int_{\Sigma_0 \setminus K} r^s |\check{R}|^2 \lesssim \int_{\Sigma_0 \setminus K} r^s |\check{R}'|^2 + r^s \left( \frac{M^2 \epsilon^2}{r^{s+5}} \right) \lesssim \mathfrak{R}_0'^2 + \frac{M^2 \epsilon^2}{R_0^2} \lesssim \epsilon_0^2.$$

Moreover, we have from (6.3)

$$\partial R = (\partial R)' + \partial f \cdot R' + f \cdot (\partial R)' + \text{l.o.t.}, \quad (6.4)$$

where

$$\partial R := \{(\nabla_\mu \alpha)_{AB}, (\nabla_\mu \beta)_A, \nabla_\mu \rho, \nabla_\mu \sigma, (\nabla_\mu \underline{\beta})_A, (\nabla_\mu \underline{\alpha})_{AB}\}, \quad \mu = 1, 2, 3, 4.$$

We have from Lemma 5.4

$$|(\partial R)_K - (\partial R)'_K| \lesssim \frac{M\epsilon}{r^{\frac{s+7}{2}}}.$$

---

<sup>28</sup>Recall that quantities with prime are associated to the foliation of the initial data layer, see (5.7) and (5.8).

Moreover, we have from Proposition 2.22

$$\widetilde{\partial R} = \partial \check{R} + (\partial - \partial_K) R_K = \partial \check{R} + (\check{J}, \check{L}, \check{J}) \cdot \mathbf{O}_3^1,$$

which implies

$$\begin{aligned} \int_{\Sigma_0 \setminus K} r^{s+2} |\partial \check{R}|^2 &\lesssim \int_{\Sigma_0 \setminus K} r^{s+2} |\widetilde{\partial R}|^2 + r^{s+2} \frac{M^2}{r^6} |\Gamma_w^{(1)}|^2 \\ &\lesssim \int_{\Sigma_0 \setminus K} r^{s+2} |\widetilde{\partial R}|^2 + \int_{R_0}^\infty \frac{M^2}{r^{4-s}} |\Gamma_w^{(1)}|_{2,S}^2 \\ &\lesssim \int_{\Sigma_0 \setminus K} r^{s+2} |\widetilde{\partial R}|^2 + \int_{R_0}^\infty \frac{M^2}{r^{4-s}} \frac{\epsilon^2}{r^{s-1}} \\ &\lesssim \int_{\Sigma_0 \setminus K} r^{s+2} |\widetilde{\partial R}|^2 + \epsilon_0^2. \end{aligned}$$

Similarly, we have

$$\int_{\Sigma_0 \setminus K} r^{s+2} |\widetilde{(\partial R)'}|^2 \lesssim \int_{\Sigma_0 \setminus K} r^{s+2} |\partial' \check{R}|^2 + \epsilon_0^2 \lesssim \epsilon_0^2.$$

Hence, we obtain from (6.4) and  $\text{Osc}(f) \leq \epsilon$  that

$$\begin{aligned} \int_{\Sigma_0 \setminus K} r^{s+2} |\partial \check{R}|^2 &\lesssim \epsilon_0^2 + \int_{\Sigma_0 \setminus K} r^{s+2} |\widetilde{\partial R}|^2 \\ &\lesssim \epsilon_0^2 + \int_{\Sigma_0 \setminus K} r^{s+2} |\widetilde{(\partial R)'}|^2 + r^{s+2} \frac{M^2 \epsilon^2}{r^{s+7}} + |\partial f|^2 r^{s+2} |R'|^2 + |f|^2 r^{s+2} |\partial' R'|^2 \\ &\lesssim \epsilon_0^2 + \int_{\Sigma_0 \setminus K} r^s |\check{R}'|^2 + r^{s+2} |\widetilde{(\partial R)'}|^2 + r^{s+2} |f|^2 |(\partial' R')_K|^2 \\ &\lesssim \epsilon_0^2 + \int_{\Sigma_0 \setminus K} r^{s+2} \frac{\epsilon^2}{r^{s-1}} \frac{M^2}{r^8} \\ &\lesssim \epsilon_0^2. \end{aligned}$$

Thus, we deduce

$$\mathfrak{R}_0^2 = \sum_R \int_{\Sigma_0 \setminus K} r^s |\mathfrak{d}^{\leq 1} \check{R}|^2 \lesssim \sum_R \int_{\Sigma_0 \setminus K} r^s |\check{R}|^2 + r^{s+2} |\partial \check{R}|^2 \lesssim \epsilon_0^2.$$

This concludes the proof of Theorem M0. □

## 6.2 Control of the last slice (Theorem M2)

The goal of this section is to prove Theorem M2 which we recall below for convenience.

**Theorem M2.** *Let  $\underline{C}_*$  endowed with the geodesic type foliation of Definition 6.1. We assume that*

$$\mathcal{O}'(\underline{C}_* \cap \Sigma_0) \leq \epsilon_0, \quad \mathcal{R}(\underline{C}_*) \lesssim \epsilon_0, \quad \mathcal{O}^*(\underline{C}_*) \leq \epsilon.$$

*Then, we have*

$$\mathcal{O}^*(\underline{C}_*) \lesssim \epsilon_0.$$

### 6.2.1 Geodesic type foliation on $\underline{C}_*$

**Definition 6.1.** A foliation  $(u, \underline{u}_*)$  on the last slice  $\underline{C}_*$  is called a geodesic type foliation if it satisfies the following conditions:

$$\check{\Omega} = 0, \quad \check{\underline{b}}^A = 0. \quad (6.5)$$

**Remark 6.2.** Notice that (6.5) is equivalent to

$$e_3(u) = \Omega_K^{-1}, \quad e_3(x^A) = \Omega_K^{-1} \underline{b}_K^A.$$

In particular, the choice of the foliation is prescribed by  $\check{\Omega} = 0$  while  $\check{\underline{b}}^A = 0$  corresponds in fact to the transport of the  $x^A$  coordinates. The local existence of geodesic type foliations as in Definition 6.1 is completely analogous to the one for the standard geodesic foliation, i.e.  $\Omega = \frac{1}{2}$ , with a transport of the  $x^A$  coordinates by  $e_3(x^A) = 0$ .

**Lemma 6.3.** Let  $\underline{C}_*$  endowed with the geodesic type foliation of Definition 6.1. Then, we have

$$\check{\underline{\omega}} = 0, \quad \text{on } \underline{C}_*.$$

*Proof.* We have from Lemma 2.21 and (6.5)

$$\Omega e_3 - \Omega_K(e_3)_K = \check{\underline{b}}^A \partial_{x^A} = 0,$$

which implies from (6.5)

$$e_3 = (e_3)_K. \quad (6.6)$$

Hence, we have from (6.5) and (6.6)

$$\underline{\omega} = -\frac{1}{2} \nabla_3(\log \Omega) = -\frac{1}{2} \nabla_{(e_3)_K}(\log \Omega_K) = \underline{\omega}_K.$$

This concludes the proof of Lemma 6.3.  $\square$

**Remark 6.4.** Under the assumption  $\mathcal{O}^*(\underline{C}_*) \leq \epsilon$ , we have on  $\underline{C}_*$

$$\begin{aligned} \check{\vartheta}^{\leq 1} \widetilde{\text{tr } \underline{\chi}} &\in \Gamma_g, & \check{\vartheta}^{\leq 1} \check{\gamma} &\in r\Gamma_g, & \check{\vartheta}^{\leq 1}(\check{\epsilon}) &\in r\Gamma_g, \\ \check{\underline{L}} &\in \Gamma_g, & \check{\underline{J}} &\in \Gamma_b, & r\check{\underline{\mu}} &\in \Gamma_g. \end{aligned}$$

**Proposition 6.5.** We have the following linearized equations on  $\underline{C}_*$ :

$$\begin{aligned} \Omega \nabla_3(\widetilde{\Omega \text{tr } \underline{\chi}}) + \frac{1}{2} \Omega \text{tr } \underline{\chi} \widetilde{\Omega \text{tr } \underline{\chi}} &= -2\Omega^2 \check{\underline{\mu}} + \Gamma_g \cdot \mathbf{O}_3^2 + \Gamma_g \cdot \Gamma_g, \\ \text{div}(\check{\underline{\chi}}) &= -\check{\underline{\beta}} + \mathbf{O}_0^0 \cdot \nabla \widetilde{\text{tr } \underline{\chi}} + \mathbf{O}_1^0 \cdot \check{\underline{\zeta}} + \mathbf{O}_3^2 \cdot \Gamma_g + \Gamma_g \cdot \Gamma_g, \\ \nabla_3 \widetilde{\Omega \text{tr } \underline{\chi}} + \text{tr } \underline{\chi} \widetilde{\Omega \text{tr } \underline{\chi}} &= \Gamma_g \cdot \mathbf{O}_2^1 + \Gamma_b \cdot \Gamma_b, \\ \text{div}(\check{\underline{\chi}}) &= \check{\underline{\beta}} + \mathbf{O}_0^0 \cdot \nabla \widetilde{\text{tr } \underline{\chi}} + \mathbf{O}_1^0 \cdot \check{\underline{\zeta}} + \mathbf{O}_3^2 \cdot \Gamma_b + \Gamma_b \cdot \Gamma_g, \\ \Omega \nabla_3 \widetilde{\underline{\mu}} + \Omega \text{tr } \underline{\chi} \widetilde{\underline{\mu}} &= \Omega \text{tr } \underline{\chi} \check{\underline{\rho}} + \mathbf{O}_3^2 \cdot (\check{\underline{\rho}}, \check{\underline{\beta}}) + \mathbf{O}_3^1 \cdot \Gamma_g + \mathbf{O}_0^0 \cdot \check{\underline{\eta}} \cdot \check{\underline{\beta}} + r^{-1} \Gamma_b \cdot \Gamma_b. \end{aligned} \quad (6.7)$$

We also have the following equation:

$$\begin{aligned} &\nabla_3(r\Delta \widetilde{\Omega \text{tr } \underline{\chi}}) + \frac{3}{2} \text{tr } \underline{\chi} (r\Delta \widetilde{\Omega \text{tr } \underline{\chi}}) \\ &= \mathbf{O}_2^2 \cdot \nabla^2 \check{\underline{\chi}} + \mathbf{O}_3^1 \cdot \Gamma_g^{(1)} + r\Gamma_b \cdot \nabla^2 \check{\underline{\chi}} + \mathbf{O}_{-1}^0 \cdot |\nabla \check{\underline{\chi}}|^2 + r^{-1} \Gamma_b \cdot \Gamma_g^{(1)}. \end{aligned} \quad (6.8)$$

*Proof.* Notice that (6.7) follows directly from Proposition 2.34 and Remark 6.4. Next, we have from Proposition A.1

$$\begin{aligned}\Omega \nabla_3 \widetilde{\Omega \operatorname{tr} \chi} + (\Omega \operatorname{tr} \chi) \widetilde{\Omega \operatorname{tr} \chi} &= -4\Omega \operatorname{tr} \chi \widetilde{\Omega \omega} + \mathfrak{L}[\nabla_3 \widetilde{\operatorname{tr} \chi}] + \operatorname{Err}[\nabla_3 \widetilde{\operatorname{tr} \chi}], \\ \mathfrak{L}[\nabla_3 \widetilde{\operatorname{tr} \chi}] &:= -2\widetilde{\Omega \chi}(\Omega_K \widehat{\chi}_K) - 4\Omega_K \omega_K \widetilde{\Omega \operatorname{tr} \chi} - \check{b}^A \nabla_{\partial_{x^a}} \operatorname{tr}_K \chi_K, \\ \operatorname{Err}[\nabla_3 \widetilde{\operatorname{tr} \chi}] &:= \frac{1}{2}(\widetilde{\Omega \operatorname{tr} \chi})^2 - |\widetilde{\Omega \chi}|^2.\end{aligned}$$

Recalling from Lemma 6.3 that  $\check{\Omega} = 0$ ,  $\check{\omega} = 0$  and  $\check{b}^A = 0$  on  $\underline{C}_*$ , we obtain

$$\Omega \nabla_3 \widetilde{\Omega \operatorname{tr} \chi} + (\Omega \operatorname{tr} \chi) \widetilde{\Omega \operatorname{tr} \chi} = \mathbf{O}_3^2 \cdot \check{\chi} + \mathbf{O}_2^1 \cdot \widetilde{\operatorname{tr} \chi} + \frac{1}{2}(\widetilde{\Omega \operatorname{tr} \chi})^2 - |\widetilde{\Omega \chi}|^2. \quad (6.9)$$

Differentiating it by  $\nabla$  and applying Proposition 2.41, we deduce

$$\begin{aligned}\Omega \nabla_3 (\nabla \widetilde{\Omega \operatorname{tr} \chi}) + (\Omega \operatorname{tr} \chi) \nabla \widetilde{\Omega \operatorname{tr} \chi} &= \mathbf{O}_3^2 \cdot \nabla \check{\chi} + \mathbf{O}_2^1 \cdot \nabla \widetilde{\operatorname{tr} \chi} + \mathbf{O}_0^0 \cdot \nabla |\check{\chi}|^2 + r^{-1} \Gamma_g \cdot \Gamma_g + [\Omega \nabla_3, \nabla] \widetilde{\Omega \operatorname{tr} \chi} \\ &= \mathbf{O}_3^2 \cdot \nabla \check{\chi} + \mathbf{O}_3^1 \cdot \Gamma_g + \mathbf{O}_0^0 \cdot \nabla |\check{\chi}|^2 - \frac{1}{2} \Omega \operatorname{tr} \chi \nabla \widetilde{\Omega \operatorname{tr} \chi} + r^{-1} \Gamma_b \cdot \Gamma_g,\end{aligned}$$

which implies

$$\Omega \nabla_3 (\nabla \widetilde{\Omega \operatorname{tr} \chi}) + \frac{3}{2} \Omega \operatorname{tr} \chi \nabla \widetilde{\Omega \operatorname{tr} \chi} = \mathbf{O}_3^2 \cdot \nabla \check{\chi} + \mathbf{O}_3^1 \cdot \Gamma_g + \mathbf{O}_0^0 \cdot \nabla |\check{\chi}|^2 + r^{-1} \Gamma_b \cdot \Gamma_g.$$

Differentiating it by  $r \operatorname{div}$  and applying Corollary 2.43, we infer

$$\Omega \nabla_3 (r \Delta \widetilde{\Omega \operatorname{tr} \chi}) + \frac{3}{2} \Omega \operatorname{tr} \chi (r \Delta \widetilde{\Omega \operatorname{tr} \chi}) = \mathbf{O}_2^2 \cdot \nabla^2 \check{\chi} + \mathbf{O}_3^1 \cdot \Gamma_g^{(1)} + r \Gamma_b \cdot \nabla^2 \check{\chi} + \mathbf{O}_{-1}^0 \cdot |\nabla \check{\chi}|^2 + r^{-1} \Gamma_b \cdot \Gamma_g^{(1)}.$$

This concludes the proof of Proposition 6.5.  $\square$

### 6.2.2 Estimate for $\mathcal{O}_1^*$

In the remainder of Section 6.2, we denote

$$S := S(u, \underline{u}_*), \quad S_* := \underline{C} \cap \Sigma_0. \quad (6.10)$$

**Proposition 6.6.** *For a geodesic type foliation on  $\underline{C}_*$  satisfying*

$$\mathcal{O}'(\underline{C}_* \cap \Sigma_0) \leq \epsilon_0, \quad \underline{\mathcal{R}}(\underline{C}_*) \lesssim \epsilon_0, \quad \mathcal{O}^*(\underline{C}_*) \leq \epsilon, \quad (6.11)$$

*we have*

$$\mathcal{O}_1^* \lesssim \epsilon_0. \quad (6.12)$$

*Proof.* Notice that the assumption  $\mathcal{O}'(\underline{C}_* \cap \Sigma_0) \leq \epsilon_0$  controls the initial data on the last sphere  $S_*$ . The idea of the proof is to transport the estimates in the direction of  $\nabla_3$  along  $\underline{C}_*$  using Proposition 6.5 and Lemma 5.5. The proof is similar to Section 5, and in fact easier since  $\check{\Omega} = 0$ ,  $\check{\omega} = 0$  and  $\check{b}^A = 0$  on  $\underline{C}_*$ .

**Step 1.** Estimate for  $\mathcal{O}_{0,1,2}^*(\widetilde{\operatorname{tr} \chi})$ .

The estimates for  $\mathcal{O}_0^*(\widetilde{\operatorname{tr} \chi})$  and  $\mathcal{O}_1^*(\widetilde{\operatorname{tr} \chi})$  follow directly from Propositions 5.13 and 5.21 in the particular case  $\check{\Omega} = 0$  and  $\check{\omega} = 0$ . Then, we focus on  $\mathcal{O}_2^*(\widetilde{\operatorname{tr} \chi})$ .

We recall from (6.8)

$$\nabla_3(r\Delta\widetilde{\Omega\text{tr}\chi}) + \frac{3}{2}\text{tr}\chi(r\Delta\widetilde{\Omega\text{tr}\chi}) = \mathbf{O}_2^2 \cdot \nabla^2\check{\chi} + \mathbf{O}_3^1 \cdot \Gamma_g^{(1)} + r\Gamma_b \cdot \nabla^2\check{\chi} + \mathbf{O}_{-1}^0 \cdot |\nabla\check{\chi}|^2 + r^{-1}\Gamma_b \cdot \Gamma_g^{(1)}.$$

Moreover, we have from Proposition 6.5 and Remark 6.4

$$\nabla^2\check{\chi} = \nabla\check{\beta} + r^{-2}\Gamma_g^{(1)}, \quad \nabla\check{\chi} = \check{\beta} + r^{-1}\Gamma_g.$$

Hence, we obtain

$$\nabla_3(r\Delta\widetilde{\Omega\text{tr}\chi}) + \frac{3}{2}\text{tr}\chi(r\Delta\widetilde{\Omega\text{tr}\chi}) = \mathbf{O}_2^2 \cdot \nabla\check{\beta} + r\Gamma_b \cdot \nabla\check{\beta} + \mathbf{O}_{-1}^0 \cdot |\check{\beta}^{(0)}|^2 + \mathbf{O}_3^1 \cdot \Gamma_g^{(1)} + r^{-1}\Gamma_b \cdot \Gamma_g^{(1)}.$$

We recall from Proposition 2.37

$$\nabla_3(\check{\rho}, \check{\sigma}) + \frac{3}{2}\text{tr}\chi(\check{\rho}, \check{\sigma}) = -d_1\check{\beta} + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot \check{\alpha} + \Gamma_g \cdot \check{\alpha}.$$

We deduce

$$\nabla_3\Xi + \frac{3}{2}\text{tr}\chi\Xi = \mathbf{O}_3^2 \cdot \check{\alpha} + \mathbf{O}_{-1}^0 \cdot |\check{\beta}^{(0)}|^2 + \mathbf{O}_3^1 \cdot \Gamma_g^{(1)} + \Gamma_g \cdot \check{\alpha} + r^{-1}\Gamma_b \cdot \Gamma_g^{(1)},$$

where  $\Xi$  is the following renormalized quantity

$$\Xi = r\Delta\widetilde{\Omega\text{tr}\chi} + \mathbf{O}_2^2 \cdot (\check{\rho}, \check{\sigma}) + r\Gamma_b \cdot (\check{\rho}, \check{\sigma}).$$

Applying Lemma 5.5, we deduce for  $p \in [2, 4]$

$$\begin{aligned} & |r^{3-\frac{2}{p}}\Xi|_{p,S} \\ & \lesssim |r^{3-\frac{2}{p}}\Xi|_{p,S_*} + \int_{u_0(\underline{u}_*)}^u M^2 |r^{-\frac{2}{p}}\check{\alpha}|_{p,S} + |r^{4-\frac{2}{p}}|\check{\beta}^{(0)}|^2|_{p,S} + M|r^{-\frac{2}{p}}\Gamma_g^{(1)}| + |r^{3-\frac{2}{p}}\Gamma_g \cdot \check{\alpha}|_{p,S} + |r^{2-\frac{2}{p}}\Gamma_b \cdot \Gamma_g^{(1)}|_{p,S} \\ & \lesssim \frac{\epsilon_0}{r^{\frac{s-3}{2}}} + \int_{u_0(\underline{u}_*)}^u \frac{M^2\epsilon_0}{r|u|^{\frac{s+1}{2}}} + \frac{M}{r^2|u|^{\frac{s-3}{2}}} + \frac{\epsilon_0}{|u|^{s-1}} + \frac{\epsilon^2}{r|u|^{s-2}} + |r^{4-\frac{2}{p}}|\check{\beta}^{(0)}|^2|_{p,S} \\ & \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}} + \int_{u_0(\underline{u}_*)}^u r^2|r^{\frac{3}{4}}\check{\beta}^{(0)}|_{8,S}^2 \\ & \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}} + \int_{u_0(\underline{u}_*)}^u |r\check{\beta}^{(1)}|_{2,S}^2 \\ & \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{|u|^{s-2}} \\ & \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}, \end{aligned}$$

where we used the Sobolev inequality  $|r^{-\frac{1}{4}}U|_{8,S} \lesssim |r^{-1}U^{(1)}|_{2,S}$  for any  $S$ -tangent tensor  $U$ .

Hence, we infer from Proposition 2.15

$$|r^{4-\frac{2}{p}}\nabla^2\widetilde{\Omega\text{tr}\chi}|_{p,S} \lesssim |r^{3-\frac{2}{p}}\Xi|_{p,S} + \frac{M^2}{r^2}|r^{3-\frac{2}{p}}(\check{\rho}, \check{\sigma})|_{p,S} + |r^{4-\frac{2}{p}}\Gamma_b \cdot (\check{\rho}, \check{\sigma})|_{p,S} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}.$$

Thus, we obtain

$$\mathcal{O}_{0,1,2}^*(\widetilde{\text{tr}\chi}) \lesssim \epsilon_0. \quad (6.13)$$

**Step 2.** Estimate for  $\mathcal{O}_{0,1}^*(\widetilde{[\mu]})$ .

The estimate for  $\mathcal{O}_0^*(\widetilde{[\mu]})$  follows directly from Proposition 5.18 in the particular case  $\check{\Omega} = 0$  and  $\check{\omega} = 0$ . Next, we recall from (6.7)

$$\Omega \nabla_3 \widetilde{[\mu]} + \Omega \operatorname{tr} \chi \widetilde{[\mu]} = \Omega \operatorname{tr} \chi \check{\rho} + \mathbf{O}_3^2 \cdot (\check{\rho}, \check{\beta}) + \mathbf{O}_3^1 \cdot \Gamma_g + \mathbf{O}_0^0 \cdot \check{\eta} \cdot \check{\beta} + r^{-1} \Gamma_b \cdot \Gamma_b.$$

Differentiating it by  $r \nabla$  and applying Corollary 2.43, we infer

$$\nabla_3(r \nabla \widetilde{[\mu]}) + \operatorname{tr} \chi(r \nabla \widetilde{[\mu]}) = \mathbf{O}_0^0 \cdot (\nabla \check{\rho}, \nabla \check{\beta}) + \mathbf{O}_0^0 \cdot \check{\rho} \cdot \check{\beta} + r \Gamma_g \cdot \nabla \check{\beta} + \mathbf{O}_3^1 \cdot \Gamma_g^{(1)} + r^{-1} \Gamma_b \cdot \Gamma_b^{(1)}.$$

We deduce

$$\nabla_3 \underline{U} + \operatorname{tr} \chi \underline{U} = \mathbf{O}_0^0 \cdot \check{\rho} \cdot \check{\beta} + \mathbf{O}_3^1 \cdot \Gamma_g^{(1)} + r^{-1} \Gamma_b \cdot \Gamma_b^{(1)},$$

where the renormalized quantity  $\underline{U}$  is given by

$$\underline{U} = r \nabla \widetilde{[\mu]} + \mathbf{O}_0^0 \cdot (\check{\beta}, \check{\rho}, \check{\sigma}) + r \Gamma_g \cdot (\check{\rho}, \check{\sigma}).$$

Applying Lemma 5.5, we have

$$\begin{aligned} |r^{2-\frac{2}{p}} \underline{U}|_{p,S} &\lesssim |r^{2-\frac{2}{p}} \underline{U}|_{p,S_*} + \int_{u_0(\underline{u}_*)}^u |r^{2-\frac{2}{p}} \check{\rho} \cdot \check{\beta}|_{p,S} + M |r^{-1-\frac{2}{p}} \Gamma_g^{(1)}|_{p,S} + |r^{1-\frac{2}{p}} \Gamma_b \cdot \Gamma_b^{(1)}|_{p,S} \\ &\lesssim \frac{\epsilon_0}{r^{\frac{s-1}{2}}} + \frac{M\epsilon}{r^2 |u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r |u|^{s-2}} + \int_{u_0(\underline{u}_*)}^u |r^{\frac{3}{2}} \check{\rho} \cdot \check{\beta}|_{4,S} \\ &\lesssim \frac{\epsilon_0}{r |u|^{\frac{s-3}{2}}} + \int_{u_0(\underline{u}_*)}^u r^2 |r^{-\frac{1}{4}} \check{\rho}|_{8,S} |r^{-\frac{1}{4}} \check{\beta}|_{8,S} \\ &\lesssim \frac{\epsilon_0}{r |u|^{\frac{s-3}{2}}} + \int_{u_0(\underline{u}_*)}^u |\check{\rho}^{(1)}|_{2,S} |\check{\beta}^{(1)}|_{2,S} \\ &\lesssim \frac{\epsilon_0}{r |u|^{\frac{s-3}{2}}} + r^{-2} \left( \int_{u_0(\underline{u}_*)}^u |r \check{\rho}^{(1)}|_{2,S}^2 \right)^{\frac{1}{2}} \left( \int_{u_0(\underline{u}_*)}^u |r \check{\beta}^{(1)}|_{2,S}^2 \right)^{\frac{1}{2}} \\ &\lesssim \frac{\epsilon_0}{r |u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r^2 |u|^{s-2}} \\ &\lesssim \frac{\epsilon_0}{r |u|^{\frac{s-3}{2}}}, \end{aligned}$$

where we used the Sobolev inequality  $|r^{-\frac{1}{4}} U|_{8,S} \lesssim |r^{-1} U^{(1)}|_{2,S}$  for any  $S$ -tangent tensor  $U$ . Hence, we obtain

$$|r^{3-\frac{2}{p}} \nabla \widetilde{[\mu]}|_{p,S} \lesssim |r^{2-\frac{2}{p}} \underline{U}|_{p,S} + |r^{2-\frac{2}{p}} (\check{\beta}, \check{\rho}, \check{\sigma})|_{p,S} + |r^{3-\frac{2}{p}} \Gamma_g \cdot (\check{\rho}, \check{\sigma})|_{p,S} \lesssim \frac{\epsilon_0}{r |u|^{\frac{s-3}{2}}}.$$

Thus, we have

$$\mathcal{O}_{0,1}^*(\widetilde{[\mu]}) \lesssim \epsilon_0. \quad (6.14)$$

**Step 3.** Estimate for  $\mathcal{O}_{0,1}^*(\operatorname{tr} \chi)$ .

We recall from (6.7) that

$$\Omega \nabla_3 (\widetilde{\Omega \operatorname{tr} \chi}) = -2 \widetilde{\Omega^2 [\mu]} + \Gamma_g \cdot \mathbf{O}_3^2 + \Gamma_g \cdot \Gamma_g. \quad (6.15)$$

We recall from (2.26) and (2.27)

$$[\mu] + [\underline{\mu}] = \frac{1}{2} \operatorname{tr} \chi \operatorname{tr} \underline{\chi} - 2\Delta \log \Omega + \widehat{\chi} \cdot \widehat{\underline{\chi}} - 2\rho,$$

which implies

$$\widetilde{\Omega^2[\mu]} = -\widetilde{\Omega^2[\underline{\mu}]} + \frac{1}{2} \widetilde{\Omega \operatorname{tr} \chi} \Omega \operatorname{tr} \underline{\chi} + \mathbf{O}_1^0 \cdot \widetilde{\operatorname{tr} \chi} - 2\Omega^2 \check{\rho} + \mathbf{O}_3^2 \cdot \Gamma_b + \Gamma_g \cdot \Gamma_b.$$

Injecting it into (6.15), we deduce

$$\Omega \nabla_3 (\widetilde{\Omega \operatorname{tr} \chi}) + \Omega \operatorname{tr} \underline{\chi} \widetilde{\Omega \operatorname{tr} \chi} = 2\widetilde{\Omega^2[\underline{\mu}]} + 4\Omega^2 \check{\rho} + \mathbf{O}_1^0 \cdot \widetilde{\operatorname{tr} \chi} + \mathbf{O}_3^2 \cdot \Gamma_b + \Gamma_b \cdot \Gamma_g. \quad (6.16)$$

Applying Lemma 5.5, we infer

$$\begin{aligned} & |r^{2-\frac{2}{p}} \widetilde{\Omega \operatorname{tr} \chi}|_{p,S} \\ & \lesssim |r^{2-\frac{2}{p}} \widetilde{\Omega \operatorname{tr} \chi}|_{p,S_*} + \int_{u_0(\underline{u}_*)}^u |r^{2-\frac{2}{p}} \widetilde{[\underline{\mu}]}|_{p,S} + |r^{2-\frac{2}{p}} \check{\rho}|_{p,S} + |r^{1-\frac{2}{p}} \widetilde{\operatorname{tr} \chi}|_{p,S} + M^2 |r^{-1-\frac{2}{p}} \Gamma_b|_{p,S} + |r^{2-\frac{2}{p}} \Gamma_b \cdot \Gamma_g|_{p,S} \\ & \lesssim \frac{\epsilon_0}{r^{\frac{s-1}{2}}} + \int_{u_0(\underline{u}_*)}^u \frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}} + \frac{M^2 \epsilon}{r^2 |u|^{\frac{s-1}{2}}} + \frac{\epsilon^2}{r|u|^{s-2}} \\ & \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}. \end{aligned}$$

Next, we differentiate (6.16) by  $\nabla$  and apply Proposition 2.41 to deduce

$$\begin{aligned} & \Omega \nabla_3 (\nabla \widetilde{\Omega \operatorname{tr} \chi}) + \Omega \operatorname{tr} \underline{\chi} (\nabla \widetilde{\Omega \operatorname{tr} \chi}) \\ & = 2\Omega^2 \nabla [\underline{\mu}] + 4\Omega^2 \nabla \check{\rho} - \Omega \underline{\chi} \cdot \nabla \widetilde{\Omega \operatorname{tr} \chi} + \mathbf{O}_1^0 \cdot \nabla \widetilde{\operatorname{tr} \chi} + \Gamma_b^{(1)} \cdot \mathbf{O}_4^2 + r^{-1} \Gamma_b \cdot \Gamma_g^{(1)} \\ & = 2\Omega^2 \nabla [\underline{\mu}] + 4\Omega^2 \nabla \check{\rho} - \frac{1}{2} \Omega \operatorname{tr} \underline{\chi} (\nabla \widetilde{\Omega \operatorname{tr} \chi}) + \mathbf{O}_1^0 \cdot \nabla \widetilde{\operatorname{tr} \chi} + \Gamma_b^{(1)} \cdot \mathbf{O}_4^2 + r^{-1} \Gamma_b \cdot \Gamma_g^{(1)}, \end{aligned}$$

which implies from  $\check{\Omega} = 0$  that

$$\nabla_3 (\nabla \widetilde{\operatorname{tr} \chi}) + \frac{3}{2} \operatorname{tr} \underline{\chi} (\nabla \widetilde{\operatorname{tr} \chi}) = 2\nabla [\underline{\mu}] + 4\nabla \check{\rho} + \mathbf{O}_1^0 \cdot \nabla \widetilde{\operatorname{tr} \chi} + \Gamma_b^{(1)} \cdot \mathbf{O}_4^2 + r^{-1} \Gamma_b \cdot \Gamma_g^{(1)}. \quad (6.17)$$

Next, we define  $\chi^\dagger$  by the following transport equation:

$$\Omega \nabla_3 \chi^\dagger + \frac{1}{2} \Omega \operatorname{tr} \underline{\chi} \chi^\dagger = 4\Omega \check{\sigma} \quad \text{on } \underline{C}_*, \quad \chi^\dagger = 0 \quad \text{on } S_*. \quad (6.18)$$

Differentiating it by  ${}^* \nabla$  and applying Proposition 2.41, we infer

$$\begin{aligned} \Omega \nabla_3 {}^* \nabla \chi^\dagger + \frac{1}{2} \Omega \operatorname{tr} \underline{\chi} ({}^* \nabla \chi^\dagger) + \frac{1}{2} {}^* \nabla (\Omega \operatorname{tr} \underline{\chi}) \chi^\dagger &= 4 {}^* \nabla (\Omega \check{\sigma}) + [\Omega \nabla_3, {}^* \nabla] \chi^\dagger \\ &= 4 {}^* \nabla (\Omega \check{\sigma}) - \frac{1}{2} \Omega \operatorname{tr} \underline{\chi} ({}^* \nabla \chi^\dagger) - \Omega \widehat{\underline{\chi}} \cdot {}^* \nabla \chi^\dagger, \end{aligned}$$

which implies<sup>29</sup>

$$\nabla_3 {}^* \nabla \chi^\dagger + \operatorname{tr} \underline{\chi} {}^* \nabla \chi^\dagger = 4 {}^* \nabla \check{\sigma} + \mathbf{O}_4^2 \cdot \Gamma_g^{(1)} + r^{-1} \Gamma_b \cdot \Gamma_g^{(1)}. \quad (6.19)$$

We recall from Proposition 2.37

$$\nabla_3 \check{\beta} + \frac{3}{2} \operatorname{tr} \underline{\chi} \check{\beta} = \frac{1}{2} \operatorname{tr} \underline{\chi} \check{\beta} + \nabla \check{\rho} + {}^* \nabla \check{\sigma} + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + r^{-1} \Gamma_g \cdot \Gamma_b^{(1)}. \quad (6.20)$$

---

<sup>29</sup>We assume here that  $\chi^\dagger \in \Gamma_g$ . This assumption will be improved at the end of Step 3.

Combining (6.17), (6.19) and (6.20), we deduce

$$\nabla_3 X + \text{tr} \chi X = -2 \text{tr} \chi \check{\beta} + 2 \nabla[\check{\mu}] + \mathbf{O}_1^0 \cdot \nabla \widetilde{\text{tr} \chi} + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + r^{-1} \Gamma_b \cdot \Gamma_g^{(1)},$$

where

$$X := \nabla \widetilde{\text{tr} \chi} + {}^* \nabla \chi^\dagger - 4 \check{\beta}.$$

Applying Lemma 5.5 and (6.14), we obtain

$$\begin{aligned} & |r^{3-\frac{2}{p}} X|_{p,S} \\ & \lesssim |r^{3-\frac{2}{p}} X|_{p,S_*} + \int_{u_0(\underline{u}_*)}^u |r^{2-\frac{2}{p}} \check{\beta}|_{p,S} + |r^{3-\frac{2}{p}} \nabla[\check{\mu}]|_{p,S} + |r^{3-\frac{2}{p}} \nabla \widetilde{\text{tr} \chi}|_{p,S} + M |r^{-\frac{2}{p}} \Gamma_g^{(1)}|_{p,S} + |r^{2-\frac{2}{p}} \Gamma_b \cdot \Gamma_g^{(1)}|_{p,S} \\ & \lesssim \frac{\epsilon_0}{r^{\frac{s-3}{2}}} + \int_{u_0(\underline{u}_*)}^u \frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}} + \frac{M\epsilon}{r^2|u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r|u|^{\frac{s-1}{2}}} \\ & \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}. \end{aligned}$$

Then, we have from Proposition 2.15

$$|r^{3-\frac{2}{p}} \nabla(\widetilde{\text{tr} \chi}, \chi^\dagger)|_{p,S} \lesssim |r^{3-\frac{2}{p}} X|_{p,S} + |r^{3-\frac{2}{p}} \check{\beta}|_{p,S} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}.$$

Next, applying Lemma 5.5 to (6.18), we deduce

$$|r^{1-\frac{2}{p}} \chi^\dagger|_{p,S} \lesssim \int_{u_0(\underline{u}_*)}^u |r^{1-\frac{2}{p}} \check{\sigma}|_{p,S} \lesssim \frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}}.$$

Hence, we improve the assumption  $\chi^\dagger \in \Gamma_g$  and obtain

$$\mathcal{O}_{0,1}(\widetilde{\text{tr} \chi}) \lesssim \epsilon_0. \quad (6.21)$$

**Step 4.** Estimates for  $\mathcal{O}_{0,1}^*(\check{\zeta})$ ,  $\mathcal{O}_{0,1}^*(\check{\chi})$  and  $\mathcal{O}_{0,1}^*(\check{\underline{\chi}})$ .

We have from (6.14)

$$|r^{4-\frac{2}{p}} \nabla[\check{\mu}]|_{p,S} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}},$$

which implies from (2.27) and (6.21)

$$|r^{4-\frac{2}{p}} \nabla \check{\mu}|_{p,S} \lesssim |r^{4-\frac{2}{p}} \nabla[\check{\mu}]|_{p,S} + |r^{3-\frac{2}{p}} \nabla(\widetilde{\text{tr} \chi}, \widetilde{\text{tr} \underline{\chi}})|_{p,S} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}.$$

We proceed as in Proposition 5.18 to deduce

$$|r^{2-\frac{2}{p}} (r \nabla)^q \check{\eta}|_{p,S} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}, \quad q = 0, 1.$$

Recalling that

$$2\check{\zeta} = \check{\eta} - \check{\eta}, \quad \check{\eta} + \check{\eta} = \widetilde{\Delta \log \Omega} = 0,$$

we obtain

$$|r^{2-\frac{2}{p}} (r \nabla)^q \check{\zeta}|_{p,S} = |r^{2-\frac{2}{p}} (r \nabla)^q \check{\eta}|_{p,S} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}, \quad q = 0, 1.$$

Hence, we have

$$\mathcal{O}_{0,1}^*(\check{\zeta}) \lesssim \epsilon_0. \quad (6.22)$$

Finally, we apply Proposition 5.22 in the particular case of  $\check{\Omega} = 0$  to deduce

$$\mathcal{O}_{0,1}^*(\check{\chi}) + \mathcal{O}_{0,1}^*(\check{\underline{\chi}}) \lesssim \epsilon_0. \quad (6.23)$$

Combining the estimates (6.13), (6.14), (6.21), (6.22) and (6.23), this concludes the proof of Proposition 6.6.  $\square$

### 6.2.3 Estimate for $\mathcal{O}_\gamma^*$

**Proposition 6.7.** *For a geodesic type foliation on  $\underline{C}_*$  in the sense of Definition 6.1, we assume that*

$$\mathcal{O}'(\underline{C}_* \cap \Sigma_0) \leq \epsilon_0, \quad \underline{R}(\underline{C}_*) \lesssim \epsilon_0, \quad \mathcal{O}^*(\underline{C}_*) \leq \epsilon. \quad (6.24)$$

Then, we have

$$\mathcal{O}_\gamma^* \lesssim \epsilon_0. \quad (6.25)$$

*Proof.* We recall from Proposition 2.40

$$\nabla_3(\check{\gamma}_{AB}) - \text{tr } \underline{\chi}(\check{\gamma}_{AB}) = \mathbf{O}_{-2}^0 \cdot (\check{\underline{\chi}}, \widetilde{\text{tr } \underline{\chi}}) + \mathbf{O}_3^2 \cdot \check{\gamma}_{AB}. \quad (6.26)$$

Differentiating it by  $\nabla$  and applying Corollary 2.43, we deduce

$$\begin{aligned} \Omega \nabla_3(\nabla \check{\gamma}_{AB}) - \Omega \text{tr } \underline{\chi}(\nabla \check{\gamma}_{AB}) &= \nabla(\Omega \text{tr } \underline{\chi}) \check{\gamma}_{AB} + \mathbf{O}_{-2}^0 \cdot (\nabla \check{\underline{\chi}}, \nabla \widetilde{\text{tr } \underline{\chi}}) + \mathbf{O}_3^2 \cdot \nabla \check{\gamma}_{AB} + [\Omega \nabla_3, \nabla] \check{\gamma}_{AB} \\ &= -\frac{1}{2} \Omega \text{tr } \underline{\chi} \nabla \check{\gamma}_{AB} + \mathbf{O}_{-2}^0 \cdot (\nabla \check{\underline{\chi}}, \nabla \widetilde{\text{tr } \underline{\chi}}) + \mathbf{O}_1^2 \cdot \Gamma_g + r^2 \Gamma_g \cdot \Gamma_g, \end{aligned}$$

which implies<sup>30</sup>

$$\nabla_3(\nabla \check{\gamma}_{AB}) - \frac{1}{2} \text{tr } \underline{\chi}(\nabla \check{\gamma}_{AB}) = \mathbf{O}_{-2}^0 \cdot (\nabla \check{\underline{\chi}}, \nabla \widetilde{\text{tr } \underline{\chi}}) + \mathbf{O}_1^2 \cdot \Gamma_g + r^2 \Gamma_g \cdot \Gamma_g. \quad (6.27)$$

Applying Lemma 5.5, we obtain

$$\begin{aligned} &|r^{-1-\frac{2}{p}} \nabla \check{\gamma}_{AB}|_{p,S} \\ &\lesssim |r^{-1-\frac{2}{p}} \nabla \check{\gamma}_{AB}|_{p,S_*} + \int_{u_0(u_*)}^u \left( |r^{1-\frac{2}{p}} \nabla(\check{\underline{\chi}}, \widetilde{\text{tr } \underline{\chi}})|_{p,S} + M^2 |r^{-2-\frac{2}{p}} \Gamma_g|_{p,S} + |r^{1-\frac{2}{p}} \Gamma_g \cdot \Gamma_g|_{p,S} \right) du \\ &\lesssim \frac{\epsilon_0}{r^{\frac{s-1}{2}}} + \int_{u_0(u_*)}^u \left( \frac{\epsilon_0}{r|u|^{\frac{s-1}{2}}} + \frac{M^2}{r^4|u|^{\frac{s-3}{2}}} + \frac{\epsilon^2}{r^3|u|^{s-3}} \right) du \\ &\lesssim \frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}}. \end{aligned}$$

Differentiating (6.27) by  $r\nabla$  and applying Corollary 2.43, we infer

$$\nabla_3(r \nabla^2 \check{\gamma}_{AB}) - \frac{1}{2} \text{tr } \underline{\chi}(r \nabla^2 \check{\gamma}_{AB}) = \mathbf{O}_{-3}^0 \cdot (\nabla^2 \check{\underline{\chi}}, \nabla^2 \widetilde{\text{tr } \underline{\chi}}) + \mathbf{O}_1^2 \cdot \Gamma_g^{(1)} + r^2 \Gamma_g \cdot \Gamma_g^{(1)}.$$

Next, we have from (6.7)

$$\nabla^2 \check{\underline{\chi}} = \nabla \check{\underline{\beta}} + \mathbf{O}_0^0 \cdot \nabla^2 \widetilde{\text{tr } \underline{\chi}} + \mathbf{O}_1^0 \cdot \nabla \check{\zeta} + \mathbf{O}_4^2 \cdot \Gamma_b^{(1)} + r^{-1} \Gamma_b \cdot \Gamma_g^{(1)}.$$

---

<sup>30</sup>We use the convention introduced in Remark 6.4.

Hence, we obtain

$$\nabla_3(r\nabla^2\check{\gamma}_{AB}) - \frac{1}{2}\text{tr}\check{\chi}(r\nabla^2\check{\gamma}_{AB}) = \mathbf{O}_{-3}^0 \cdot (\nabla\check{\beta}, \nabla^2\widetilde{\text{tr}\check{\chi}}, r^{-1}\nabla\check{\zeta}) + \mathbf{O}_1^2 \cdot \Gamma_b^{(1)} + r^2\Gamma_b \cdot \Gamma_g^{(1)}.$$

We recall from Proposition 2.37

$$\nabla_3(\check{\rho}, \check{\sigma}) + \frac{3}{2}\text{tr}\check{\chi}(\check{\rho}, \check{\sigma}) = -d_1\check{\beta} + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot \check{\alpha} + \Gamma_g \cdot \check{\alpha}.$$

We obtain

$$\nabla_3 G - \frac{1}{2}\text{tr}\check{\chi}G = \mathbf{O}_{-3}^0 \cdot (\nabla^2\widetilde{\text{tr}\check{\chi}}, r^{-1}\nabla\check{\zeta}, r^{-1}\check{\rho}, r^{-1}\check{\sigma}) + \mathbf{O}_0^1 \cdot \Gamma_g^{(1)} + \mathbf{O}_0^2 \cdot \check{\alpha} + r^3\Gamma_g \cdot \check{\alpha} + r^2\Gamma_b \cdot \Gamma_g^{(1)},$$

where the renormalized quantity  $G$  is given by

$$G = r\nabla^2\check{\gamma}_{AB} + \mathbf{O}_{-3}^0 \cdot (\check{\rho}, \check{\sigma}).$$

Applying Lemma 5.5, we infer

$$\begin{aligned} |r^{-1-\frac{2}{p}}G|_{p,S} &\lesssim |r^{-1-\frac{2}{p}}G|_{p,S_*} + \int_{u_0(\underline{u}_*)}^u |r^{2-\frac{2}{p}}\nabla^2\widetilde{\text{tr}\check{\chi}}|_{p,S} + |r^{1-\frac{2}{p}}(\nabla\check{\zeta}, \check{\rho}, \check{\sigma})|_{p,S} \\ &\quad + \int_{u_0(\underline{u}_*)}^u M|r^{-1-\frac{2}{p}}\Gamma_g^{(1)}|_{p,S} + M^2|r^{-1-\frac{2}{p}}\check{\alpha}|_{p,S} + |r^{2-\frac{2}{p}}\Gamma_g \cdot \check{\alpha}|_{p,S} + |r^{1-\frac{2}{p}}\Gamma_b \cdot \Gamma_g^{(1)}|_{p,S} \\ &\lesssim \frac{\epsilon_0}{r^{\frac{s-1}{2}}} + \int_{u_0(\underline{u}_*)}^u \frac{\epsilon_0}{r^2|u|^{\frac{s-3}{2}}} + \frac{M\epsilon}{r^3|u|^{\frac{s-3}{2}}} + \frac{M^2\epsilon}{r^2|u|^{\frac{s+1}{2}}} + \frac{\epsilon^2}{r|u|^{s-1}} + \frac{\epsilon^2}{r^2|u|^{s-2}} \\ &\lesssim \frac{\epsilon_0}{r|u|^{\frac{s-3}{2}}}. \end{aligned}$$

Hence, we deduce

$$|r^{1-\frac{2}{p}}\nabla^2\check{\gamma}_{AB}|_{p,S} \lesssim |r^{-\frac{2}{p}}G|_{p,S} + |r^{3-\frac{2}{p}}(\check{\rho}, \check{\sigma})|_{p,S} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}.$$

Thus, we have

$$\mathcal{O}_{0,1,2}^*(\check{\gamma}) \lesssim \epsilon_0. \quad (6.28)$$

The estimate for  $\mathcal{O}_{0,1,2}(\check{\epsilon})$  is similar. This concludes the proof of Proposition 6.7.  $\square$

### 6.2.4 Estimate for $\mathcal{O}_\Gamma^*$

**Proposition 6.8.** *For a geodesic type foliation on  $\underline{C}_*$  in the sense of Definition 6.1, we assume that*

$$\mathcal{O}'(\underline{C}_* \cap \Sigma_0) \leq \epsilon_0, \quad \underline{\mathcal{R}}(\underline{C}_*) \lesssim \epsilon_0, \quad \mathcal{O}^*(\underline{C}_*) \leq \epsilon. \quad (6.29)$$

Then, we have

$$\mathcal{O}_\Gamma^* \lesssim \epsilon_0. \quad (6.30)$$

*Proof.* We recall from Lemma 2.23

$$\begin{aligned} \Gamma_{uA}^B &= \frac{1}{2}\gamma^{BC} \left( \partial_u(\gamma_{AC}) - \frac{\partial}{\partial x^A}(\gamma_{CD}b^D) + \frac{\partial}{\partial x^C}(\gamma_{AD}b^D) \right) + \frac{b^B}{4\Omega^2} \left( 2\frac{\partial}{\partial x^A}(\Omega^2) - \frac{\partial}{\partial \underline{u}}(\gamma_{AD}b^D) \right), \\ \Gamma_{BC}^A &= \frac{1}{2}\gamma^{AD} \left( \frac{\partial\gamma_{BD}}{\partial x^C} + \frac{\partial\gamma_{CD}}{\partial x^B} - \frac{\partial\gamma_{BC}}{\partial x^D} \right) + \frac{b^A}{4\Omega^2} \frac{\partial\gamma_{BC}}{\partial \underline{u}}. \end{aligned}$$

Applying Definition 2.24, and since  $\check{\Omega} = 0$  and  $\check{b} = 0$ , we infer

$$\begin{aligned}\check{J} &= \mathbf{O}_0^0 \cdot (\partial_u \check{\gamma}, \nabla \check{\gamma}, \partial_{\underline{u}} \check{\gamma}, r^{-1} \check{\gamma}) + r \Gamma_g \cdot \Gamma_b, \\ \check{L} &= \mathbf{O}_0^0 \cdot (\nabla \check{\gamma}, \partial_{\underline{u}} \check{\gamma}, r^{-1} \check{\gamma}) + r \Gamma_g \cdot \Gamma_g,\end{aligned}$$

where we used Remark 6.4. Applying Proposition 2.40, we deduce

$$\begin{aligned}\check{J} &= \mathbf{O}_0^0 \cdot (\widetilde{\text{tr } \chi}, \check{\chi}, \widetilde{\text{tr } \chi}, \check{\chi}, r^{-1} \check{\gamma}, \nabla \check{\gamma}) + r \Gamma_g \cdot \Gamma_b, \\ \check{L} &= \mathbf{O}_0^0 \cdot (\widetilde{\text{tr } \chi}, \check{\chi}, r^{-1} \check{\gamma}, \nabla \check{\gamma}) + r \Gamma_g \cdot \Gamma_g.\end{aligned}$$

Differentiating it by  $r \nabla$ , we obtain

$$\begin{aligned}r \nabla \check{J} &= \mathbf{O}_0^0 \cdot (r \nabla \widetilde{\text{tr } \chi}, r \nabla \check{\chi}, r \nabla \widetilde{\text{tr } \chi}, r \nabla \check{\chi}, \nabla \check{\gamma}, r \nabla^2 \check{\gamma}) + r \Gamma_g \cdot \Gamma_b^{(1)}, \\ r \nabla \check{L} &= \mathbf{O}_0^0 \cdot (r \nabla \widetilde{\text{tr } \chi}, r \nabla \check{\chi}, \nabla \check{\gamma}, r \nabla^2 \check{\gamma}) + r \Gamma_g \cdot \Gamma_g^{(1)}.\end{aligned}$$

Combining with Propositions 6.6 and 6.7, we deduce for  $q = 0, 1$

$$\begin{aligned}|r^{1-\frac{2}{p}}(r \nabla)^q \check{J}|_{p,S} &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-1}{2}}} + |r^{2-\frac{2}{p}} \Gamma_g \cdot \Gamma_b^{(1)}|_{p,S} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-1}{2}}}, \\ |r^{2-\frac{2}{p}}(r \nabla)^q \check{L}|_{p,S} &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}} + |r^{3-\frac{2}{p}} \Gamma_g \cdot \Gamma_g^{(1)}|_{p,S} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-3}{2}}}.\end{aligned}$$

This concludes the proof of Proposition 6.8.  $\square$

Combining Propositions 6.6, 6.7 and 6.8, this concludes the proof of Theorem M2.

### 6.3 Extension argument (Theorem M4)

In this section, we prove Theorem M4 which we recall below for convenience.

**Theorem M4.** *We consider the spacetime  $\mathcal{K}$  and its double null foliation  $(u, \underline{u})$  which satisfies the assumptions:*

$$\mathcal{O} \lesssim \epsilon_0, \quad \mathcal{R} \lesssim \epsilon_0, \quad \text{Osc} \lesssim \epsilon_0, \quad \mathcal{O}^*(\underline{C}_*) \lesssim \epsilon_0. \quad (6.31)$$

*Then, we can extend the spacetime  $\mathcal{K} = V(u_0, \underline{u}_*)$  and the double null foliation to a new spacetime  $\widehat{\mathcal{K}} = \widehat{V}(u_0, \underline{u}_* + \nu)$ , where  $\nu > 0$  is sufficiently small, and an associated double null foliation  $(\widehat{u}, \underline{u})$ . Moreover, the new foliation  $(\widehat{u}, \underline{u})$  is geodesic on the new last slice  $\underline{C}_{**} := \underline{C}_{\underline{u}_* + \nu}$  and the new corresponding norms satisfy*

$$\widehat{\mathcal{O}} \lesssim \epsilon_0, \quad \widehat{\mathcal{R}} \lesssim \epsilon_0, \quad \widehat{\text{Osc}} \lesssim \epsilon_0, \quad \widehat{\mathcal{O}}^*(\underline{C}_{**}) \lesssim \epsilon_0. \quad (6.32)$$

*Proof.* We define the region

$$\mathcal{A}_0 := \{p \in \Sigma_0 / w(p) \in [\underline{u}_*, \underline{u}_* + \nu]\}.$$

For  $\nu > 0$  small enough, we consider the future domain of dependence  $\mathcal{K}_0^\nu$  of  $\mathcal{A}_0$ , with

$$\mathcal{K}_0^\nu := V'(u_*, \underline{u}_* + \nu) \subset \mathcal{K}'.$$

The boundary of  $\mathcal{K}_0^\nu$  contains, see Figure 7:

1. the part  $\mathcal{A}_0$  of  $\Sigma_0$ ;




Figure 7: Extension argument

2. a part of an ingoing cone denoted by

$$\underline{C}_{**}(u_* - \nu) := \underline{C}(\underline{u}_* + \nu; [u_* - \nu, u_*]),$$

which is a part of the new last slice  $\underline{C}_{**} := \underline{C}_{\underline{u}_* + \nu}$ , where

$$u_* - \nu := u|_{\underline{C}_{\underline{u}_* + \nu} \cap \Sigma_0}, \quad u_* = u|_{\underline{C}_{\underline{u}_*} \cap \Sigma_0};$$

3. a part of the outgoing cone emanating from  $S_0(u_*) = \underline{C}_{\underline{u}_*} \cap \Sigma_0$  and in  $V'(u_*, \underline{u}_* + \nu)$ , denoted by  $C_\nu^*$ .

For the double null foliation  $(u', \underline{u})$  in  $\mathcal{K}_0^\nu$ , we have:

$$\mathcal{O}'(\mathcal{K}_0^\nu) \lesssim \epsilon_0, \quad \mathcal{R}'(\mathcal{K}_0^\nu) \lesssim \epsilon_0, \quad (6.33)$$

for  $\nu > 0$  small enough by local existence. By the local existence theorem for the characteristic Cauchy problem in [22], we can extend the spacetime  $\mathcal{K} \cup \mathcal{K}_0^\nu$  to the future domain of dependence of  $C_\nu^* \cup \underline{C}_*$ .<sup>31</sup> Moreover, we extend the outgoing cones of  $\mathcal{K}$  to this new domain, construct the ingoing cones from  $C_\nu^*$  and denote  $(\tilde{u}, \underline{u})$  the global foliation in  $\tilde{\mathcal{K}} := \mathcal{K} \cup \mathcal{K}_u^\nu \cup \mathcal{K}_0^\nu$ . As a consequence of Theorem 1 of [22], we have

$$\tilde{\mathcal{O}}(\mathcal{K}_u^\nu) \lesssim \epsilon_0, \quad \tilde{\mathcal{R}}(\mathcal{K}_u^\nu) \lesssim \epsilon_0. \quad (6.34)$$

Combining (6.31), (6.33) and (6.34), we obtain

$$\tilde{\mathcal{O}}(\tilde{\mathcal{K}}) \lesssim \epsilon_0, \quad \tilde{\mathcal{R}}(\tilde{\mathcal{K}}) \lesssim \epsilon_0.$$

The foliation  $(\tilde{u}, \underline{u})$  on the new last slice  $\underline{C}_{**}$  of  $V(u_0, \underline{u}_* + \nu)$  is in general not of the geodesic type of Definition 6.1. Starting from  $S_{**} := \underline{C}_{**} \cap \Sigma_0$ , we construct a geodesic type foliation  $(\hat{u}, \underline{u}_*)$  on  $\underline{C}_{**}(-\underline{u}_* - \nu < \hat{u} < u_1)$ , where  $u_1$  is sufficiently close to  $-\underline{u}_* - \nu$ . We also denote

$$\widehat{V}(u_1) := \widehat{V}(u_1, \underline{u}_* + \nu) := J^-(\widehat{S}(u_1, \underline{u}_* + \nu)).$$

<sup>31</sup>Even if it means reducing the value of  $\nu$ .

Proceeding as in Section 2.1.5, we can construct a double null foliation  $(\widehat{u}, \underline{u})$  in  $\widehat{V}(u_1)$ . Let  $U$  the set of  $u_1 \leq u_0$  such that we have the following estimates:

$$\widehat{\mathcal{O}}(\widehat{V}(u_1)) \leq \epsilon, \quad \widehat{\mathcal{R}}(\widehat{V}(u_1)) \leq \epsilon, \quad \widehat{\text{Osc}}(\widehat{V}(u_1)) \leq \epsilon, \quad \widehat{\mathcal{O}}^*(\underline{C}_{**} \cap \widehat{V}(u_1)) \leq \epsilon. \quad (6.35)$$

We also denote

$$u_2 := \sup U. \quad (6.36)$$

Assume that  $u_2 < u_0$ . Applying Theorems M0–M3 in  $\widehat{V}(u_2)$ , we obtain

$$\widehat{\mathcal{O}}(\widehat{V}(u_2)) \lesssim \epsilon_0, \quad \widehat{\mathcal{R}}(\widehat{V}(u_2)) \lesssim \epsilon_0, \quad \widehat{\text{Osc}}(\widehat{V}(u_2)) \lesssim \epsilon_0, \quad \widehat{\mathcal{O}}^*(\underline{C}_{**} \cap \widehat{V}(u_2)) \lesssim \epsilon_0.$$

Thus, by the continuity of the norms, we obtain  $u_2 + \delta \in U$  for  $\delta > 0$  small enough, a contradiction with (6.36). Hence, we have  $u_2 = u_0$  which implies

$$\widehat{\mathcal{O}}(\widehat{V}(u_0)) \lesssim \epsilon_0, \quad \widehat{\mathcal{R}}(\widehat{V}(u_0)) \lesssim \epsilon_0, \quad \widehat{\text{Osc}}(\widehat{V}(u_0)) \lesssim \epsilon_0, \quad \widehat{\mathcal{O}}^*(\underline{C}_{**} \cap \widehat{V}(u_0)) \lesssim \epsilon_0.$$

This concludes the proof of Theorem M4.  $\square$

## A Proof of results in Section 2.8

### A.1 Proof of Proposition 2.34

We prove below a more precise version of Proposition 2.34.

**Proposition A.1.** *We have the following linearized null structure equations:*

$$\begin{aligned} \Omega \nabla_4 \widetilde{\Omega \text{tr } \chi} + (\Omega \text{tr } \chi) \widetilde{\Omega \text{tr } \chi} &= -4\Omega \text{tr } \chi \widetilde{\Omega \omega} + \mathfrak{L}[\nabla_4 \widetilde{\text{tr } \chi}] + \text{Err}[\nabla_4 \widetilde{\text{tr } \chi}], \\ \mathfrak{L}[\nabla_4 \widetilde{\text{tr } \chi}] &:= -2\widetilde{\Omega \chi}(\Omega_K \widehat{\chi}_K) - 4\Omega_K \omega_K \widetilde{\Omega \text{tr } \chi}, \\ \text{Err}[\nabla_4 \widetilde{\text{tr } \chi}] &:= \frac{1}{2}(\widetilde{\Omega \text{tr } \chi})^2 - |\widetilde{\Omega \chi}|^2, \\ \\ \Omega \nabla_3 \widetilde{\Omega \text{tr } \underline{\chi}} + (\Omega \text{tr } \underline{\chi}) \widetilde{\Omega \text{tr } \underline{\chi}} &= -4\Omega \text{tr } \underline{\chi} \widetilde{\Omega \omega} + \mathfrak{L}[\nabla_3 \widetilde{\text{tr } \underline{\chi}}] + \text{Err}[\nabla_3 \widetilde{\text{tr } \underline{\chi}}], \\ \mathfrak{L}[\nabla_3 \widetilde{\text{tr } \underline{\chi}}] &:= -2\widetilde{\Omega \underline{\chi}}(\Omega_K \widehat{\underline{\chi}}_K) - 4\Omega_K \omega_K \widetilde{\Omega \text{tr } \underline{\chi}} - \check{b}^A \partial_A (\Omega_K \text{tr}_K \underline{\chi}_K), \\ \text{Err}[\nabla_3 \widetilde{\text{tr } \underline{\chi}}] &:= \frac{1}{2}(\widetilde{\Omega \text{tr } \underline{\chi}})^2 - |\widetilde{\Omega \underline{\chi}}|^2, \\ \\ \Omega \nabla_4 (\widetilde{\Omega \text{tr } \underline{\chi}}) &= -2\widetilde{\Omega^2 [\mu]} + \mathfrak{L}[\nabla_4 (\widetilde{\text{tr } \underline{\chi}})] + \text{Err}[\nabla_4 (\widetilde{\text{tr } \underline{\chi}})], \\ \mathfrak{L}[\nabla_4 (\widetilde{\text{tr } \underline{\chi}})] &:= 4\Omega_K \underline{\eta}_K (\widetilde{\Omega \underline{\eta}}), \\ \text{Err}[\nabla_4 (\widetilde{\text{tr } \underline{\chi}})] &:= 2|\widetilde{\Omega \underline{\eta}}|^2, \\ \\ \Omega \nabla_3 (\widetilde{\Omega \text{tr } \chi}) &= -2\widetilde{\Omega^2 [\mu]} + \mathfrak{L}[\nabla_3 (\widetilde{\text{tr } \chi})] + \text{Err}[\nabla_3 (\widetilde{\text{tr } \chi})], \\ \mathfrak{L}[\nabla_3 (\widetilde{\text{tr } \chi})] &:= 4\Omega_K \eta_K (\widetilde{\Omega \eta}) + \check{b}^A \partial_A (\Omega_K \text{tr}_K \chi_K), \\ \text{Err}[\nabla_3 (\widetilde{\text{tr } \chi})] &:= 2|\widetilde{\Omega \eta}|^2. \end{aligned}$$

We also have the linearized Codazzi equations:

$$\begin{aligned}
\operatorname{div}(\check{\chi}) &= \frac{1}{2} \nabla \widetilde{\operatorname{tr} \chi} + \frac{1}{2} \operatorname{tr} \chi \check{\zeta} - \check{\beta} + \mathfrak{L}[\operatorname{div}(\check{\chi})] + \operatorname{Err}[\operatorname{div}(\check{\chi})], \\
\mathfrak{L}[\operatorname{div}(\check{\chi})] &:= \frac{1}{2} \widetilde{\operatorname{tr} \chi} \zeta_K - \check{\zeta} \cdot \widehat{\chi}_K - \zeta_K \cdot \check{\chi} + (\gamma^{AB} - (\gamma_K)^{AB}) \nabla_A (\widehat{\chi}_K)_{B\bullet} - \gamma_K^{AB} \check{L}_{AB}^C (\widehat{\chi}_K)_{C\bullet}, \\
\operatorname{Err}[\operatorname{div}(\check{\chi})] &:= -\check{\zeta} \cdot \check{\chi}, \\
\\
\operatorname{div}(\check{\underline{\chi}}) &= \frac{1}{2} \nabla \widetilde{\operatorname{tr} \underline{\chi}} - \frac{1}{2} \operatorname{tr} \underline{\chi} \check{\zeta} + \check{\underline{\beta}} + \mathfrak{L}[\operatorname{div}(\check{\underline{\chi}})] + \operatorname{Err}[\operatorname{div}(\check{\underline{\chi}})], \\
\mathfrak{L}[\operatorname{div}(\check{\underline{\chi}})] &:= -\frac{1}{2} \widetilde{\operatorname{tr} \underline{\chi}} \zeta_K + \check{\zeta} \cdot \widehat{\underline{\chi}}_K + \zeta_K \cdot \check{\underline{\chi}} + (\gamma^{AB} - (\gamma_K)^{AB}) \nabla_A (\widehat{\underline{\chi}}_K)_{B\bullet} - \gamma_K^{AB} \check{L}_{AB}^C (\widehat{\underline{\chi}}_K)_{C\bullet}, \\
\operatorname{Err}[\operatorname{div}(\check{\underline{\chi}})] &:= \check{\zeta} \cdot \check{\underline{\chi}}.
\end{aligned}$$

The linearized mass aspect functions are given by:

$$\begin{aligned}
\operatorname{div} \check{\eta} &= -\check{\mu} - \check{\rho} + \mathfrak{L}[\operatorname{div}(\check{\eta})] + \operatorname{Err}[\operatorname{div}(\check{\eta})], \\
\mathfrak{L}[\operatorname{div}(\check{\eta})] &= (\gamma^{AB} - (\gamma_K)^{AB}) \nabla_A (\eta_K)_B - (\gamma_K)^{AB} \check{L}_{AB}^C (\eta_K)_C + \frac{1}{2} \widehat{\chi}_K \cdot \check{\underline{\chi}} + \frac{1}{2} \check{\chi} \cdot \widehat{\underline{\chi}}_K, \\
\operatorname{Err}[\operatorname{div}(\check{\eta})] &= \frac{1}{2} \widehat{\chi} \cdot \check{\underline{\chi}}, \\
\\
\operatorname{div} \check{\underline{\eta}} &= -\check{\underline{\mu}} - \check{\underline{\rho}} + \mathfrak{L}[\operatorname{div}(\check{\underline{\eta}})] + \operatorname{Err}[\operatorname{div}(\check{\underline{\eta}})], \\
\mathfrak{L}[\operatorname{div}(\check{\underline{\eta}})] &= (\gamma^{AB} - (\gamma_K)^{AB}) \nabla_A (\underline{\eta}_K)_B - (\gamma_K)^{AB} \check{L}_{AB}^C (\underline{\eta}_K)_C + \frac{1}{2} \widehat{\chi}_K \cdot \check{\underline{\chi}} + \frac{1}{2} \check{\chi} \cdot \widehat{\underline{\chi}}_K, \\
\operatorname{Err}[\operatorname{div}(\check{\underline{\eta}})] &= \frac{1}{2} \widehat{\chi} \cdot \check{\underline{\chi}}.
\end{aligned}$$

The linearized torsion equations are given by:

$$\begin{aligned}
\operatorname{curl} \check{\eta} &= \check{\sigma} + \mathfrak{L}[\operatorname{curl} \check{\eta}] + \operatorname{Err}[\operatorname{curl} \check{\eta}], \\
\mathfrak{L}[\operatorname{curl} \check{\eta}] &:= \frac{1}{2} \check{\chi} \wedge \widehat{\chi}_K + \frac{1}{2} \widehat{\underline{\chi}}_K \wedge \check{\chi} - (\epsilon^{AB} - \epsilon_K^{AB}) \nabla_A (\eta_K)_B + (\epsilon_K)^{AB} \check{L}_{AB}^C (\eta_K)_C, \\
\operatorname{Err}[\operatorname{curl} \check{\eta}] &:= \frac{1}{2} \check{\chi} \wedge \check{\underline{\chi}}, \\
\\
\operatorname{curl} \check{\underline{\eta}} &= -\check{\sigma} + \mathfrak{L}[\operatorname{curl} \check{\underline{\eta}}] + \operatorname{Err}[\operatorname{curl} \check{\underline{\eta}}], \\
\mathfrak{L}[\operatorname{curl} \check{\underline{\eta}}] &:= -\frac{1}{2} \check{\chi} \wedge \widehat{\chi}_K - \frac{1}{2} \widehat{\underline{\chi}}_K \wedge \check{\chi} - (\epsilon^{AB} - \epsilon_K^{AB}) \nabla_A (\underline{\eta}_K)_B + (\epsilon_K)^{AB} \check{L}_{AB}^C (\underline{\eta}_K)_C, \\
\operatorname{Err}[\operatorname{curl} \check{\underline{\eta}}] &:= -\frac{1}{2} \check{\chi} \wedge \check{\underline{\chi}}.
\end{aligned}$$

Moreover, we have

$$\begin{aligned}
\Omega \nabla_4 (\check{\Omega} \check{\omega}) &= \frac{1}{2} \check{\Omega}^2 \check{\rho} + \mathfrak{L}[\nabla_4 \check{\omega}] + \operatorname{Err}[\nabla_4 \check{\omega}], \\
\mathfrak{L}[\nabla_4 \check{\omega}] &:= \frac{3}{2} (\check{\Omega} \check{\eta} - \check{\Omega} \underline{\eta}) \cdot (\Omega_K \eta_K - \Omega_K \underline{\eta}_K) - \frac{1}{4} (\check{\Omega} \check{\eta} - \check{\Omega} \underline{\eta}) \cdot (\Omega_K \eta_K + \Omega_K \underline{\eta}_K) \\
&\quad - \frac{1}{4} (\Omega_K \eta_K - \Omega_K \underline{\eta}_K) \cdot (\check{\Omega} \check{\eta} + \check{\Omega} \underline{\eta}) - \frac{1}{4} (\check{\Omega} \check{\eta} + \check{\Omega} \underline{\eta}) \cdot (\eta_K + \underline{\eta}_K), \\
\operatorname{Err}[\nabla_4 \check{\omega}] &:= \frac{3}{4} |\check{\Omega} \check{\eta} - \check{\Omega} \underline{\eta}|^2 - \frac{1}{4} (\check{\Omega} \check{\eta} - \check{\Omega} \underline{\eta}) \cdot (\check{\Omega} \check{\eta} + \check{\Omega} \underline{\eta}) - \frac{1}{8} |\check{\Omega} \check{\eta} + \check{\Omega} \underline{\eta}|^2,
\end{aligned}$$

$$\begin{aligned}
\Omega \nabla_3(\widetilde{\Omega \omega}) &= \frac{1}{2} \widetilde{\Omega^2 \rho} + \mathfrak{L}[\nabla_3 \widetilde{\omega}] + \text{Err}[\nabla_3 \widetilde{\omega}], \\
\mathfrak{L}[\nabla_3 \widetilde{\omega}] &:= \frac{3}{2} (\widetilde{\Omega \eta} - \widetilde{\Omega \underline{\eta}}) \cdot (\Omega_K \eta_K - \Omega_K \underline{\eta}_K) + \frac{1}{4} (\widetilde{\Omega \eta} - \widetilde{\Omega \underline{\eta}}) \cdot (\Omega_K \eta_K + \Omega_K \underline{\eta}_K) \\
&\quad + \frac{1}{4} (\Omega_K \eta_K - \Omega_K \underline{\eta}_K) \cdot (\widetilde{\Omega \eta} + \widetilde{\Omega \underline{\eta}}) - \frac{1}{4} (\widetilde{\Omega \eta} + \widetilde{\Omega \underline{\eta}}) \cdot (\eta_K + \underline{\eta}_K) + \check{b}^A \nabla_{\partial_A} (\Omega_K \omega_K), \\
\text{Err}[\nabla_3 \widetilde{\omega}] &:= \frac{3}{4} |\widetilde{\Omega \eta} - \widetilde{\Omega \underline{\eta}}|^2 + \frac{1}{4} (\widetilde{\Omega \eta} - \widetilde{\Omega \underline{\eta}}) \cdot (\widetilde{\Omega \eta} + \widetilde{\Omega \underline{\eta}}) - \frac{1}{8} |\widetilde{\Omega \eta} + \widetilde{\Omega \underline{\eta}}|^2.
\end{aligned}$$

**Remark A.2.** Combining Propositions A.1 and 2.28 and Definitions 2.3 and 2.25, we obtain immediately Proposition 2.34.

*Proof of Proposition 2.34.* We recall from Corollary 2.18

$$\Omega \nabla_4(\Omega \operatorname{tr} \chi) + \frac{1}{2}(\Omega \operatorname{tr} \chi)^2 = -\Omega^2 |\widehat{\chi}|^2 - 4\Omega \omega(\Omega \operatorname{tr} \chi),$$

which implies

$$\Omega_K \nabla_{(e_4)_K}(\Omega_K \operatorname{tr}_K \chi_K) + \frac{1}{2}(\Omega_K \operatorname{tr}_K \chi_K)^2 = -\Omega_K^2 |\widehat{\chi}_K|^2 - 4\Omega_K \omega_K(\Omega_K \operatorname{tr}_K \chi_K).$$

Taking the difference, we infer

$$\begin{aligned}
\Omega \nabla_4(\widetilde{\Omega \operatorname{tr} \chi}) &= -\frac{1}{2}(\Omega \operatorname{tr} \chi)^2 - \Omega^2 |\widehat{\chi}|^2 - 4\Omega \omega(\Omega \operatorname{tr} \chi) + \frac{1}{2}(\Omega_K \operatorname{tr}_K \chi_K)^2 + \Omega_K^2 |\widehat{\chi}_K|^2 + 4\Omega_K \omega_K(\operatorname{tr}_K \chi_K) \\
&= -\Omega \operatorname{tr} \chi \widetilde{\Omega \operatorname{tr} \chi} + \frac{1}{2} \widetilde{\Omega \operatorname{tr} \chi}^2 - 2\widetilde{\Omega \widehat{\chi}}(\Omega_K \widehat{\chi}_K) - |\widetilde{\Omega \widehat{\chi}}|^2 - 4\Omega_K \omega_K \widetilde{\Omega \operatorname{tr} \chi} \\
&\quad - 4\Omega_K \operatorname{tr}_K \chi_K \widetilde{\Omega \omega} - 4\widetilde{\Omega \omega} \widetilde{\Omega \operatorname{tr} \chi}.
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
\Omega \nabla_3(\widetilde{\Omega \operatorname{tr} \underline{\chi}}) &= -\Omega \operatorname{tr} \underline{\chi} \widetilde{\Omega \operatorname{tr} \underline{\chi}} + \frac{1}{2} \widetilde{\Omega \operatorname{tr} \underline{\chi}}^2 - 2(\widetilde{\Omega \underline{\chi}}) \Omega_K \widehat{\underline{\chi}}_K - |\widetilde{\Omega \underline{\chi}}|^2 - 4\Omega_K \omega_K \widetilde{\Omega \operatorname{tr} \underline{\chi}} \\
&\quad - 4\Omega_K \operatorname{tr}_K \underline{\chi}_K \widetilde{\Omega \omega} - 4\widetilde{\Omega \omega} \widetilde{\Omega \operatorname{tr} \underline{\chi}} - \check{b}^A \partial_A (\Omega_K \operatorname{tr}_K \underline{\chi}_K).
\end{aligned}$$

Next, we have from Corollary 2.18

$$\Omega \nabla_4(\Omega \operatorname{tr} \underline{\chi}) = -2\Omega^2 [\underline{\mu}] + 2\Omega^2 |\underline{\eta}|^2,$$

which implies

$$\Omega_K \nabla_{(e_4)_K}(\Omega_K \operatorname{tr}_K \underline{\chi}_K) = -2\Omega_K^2 [\underline{\mu}]_K + 2\Omega_K^2 |\underline{\eta}_K|^2.$$

Taking the difference, we infer

$$\Omega \nabla_4(\widetilde{\Omega \operatorname{tr} \underline{\chi}}) = -2\widetilde{\Omega^2 [\underline{\mu}]} + 4\Omega_K \underline{\eta}_K(\widetilde{\Omega \underline{\eta}}) + 2|\widetilde{\Omega \underline{\eta}}|^2.$$

Similarly, we have

$$\Omega \nabla_3(\widetilde{\Omega \operatorname{tr} \chi}) = -2\widetilde{\Omega^2 [\mu]} + 4\Omega_K \eta_K(\widetilde{\Omega \eta}) + 2|\widetilde{\Omega \eta}|^2 + \check{b}^A \partial_A (\Omega_K \operatorname{tr}_K \chi_K).$$

Now we consider the Codazzi equation

$$\operatorname{div} \widehat{\chi} = \frac{1}{2}(\nabla \operatorname{tr} \chi + \operatorname{tr} \chi \zeta) - \zeta \cdot \widehat{\chi} - \beta,$$

which implies

$$\operatorname{div}_K \widehat{\chi}_K = \frac{1}{2}(\nabla_K \operatorname{tr}_K \chi_K + \operatorname{tr}_K \chi_K \zeta_K) - \zeta_K \cdot \widehat{\chi}_K - \beta_K.$$

Taking the difference, we obtain from Proposition 2.22

$$\begin{aligned} \operatorname{div}(\widetilde{\widehat{\chi}}) &= \operatorname{div} \widehat{\chi} - \operatorname{div} \widehat{\chi}_K \\ &= \operatorname{div} \widehat{\chi} - \operatorname{div}_K \widehat{\chi}_K - (\operatorname{div} - \operatorname{div}_K) \widehat{\chi}_K \\ &= \frac{1}{2} \nabla \widetilde{\operatorname{tr} \chi} + \frac{1}{2} \operatorname{tr} \chi \check{\zeta} + \frac{1}{2} \widetilde{\operatorname{tr} \chi} \zeta_K - \check{\zeta} \cdot \widehat{\chi}_K - \zeta_K \cdot \check{\chi} - \check{\zeta} \cdot \check{\chi} - \check{\beta} - \widetilde{\gamma^{AB}} \nabla_A (\widehat{\chi}_K)_B \\ &\quad - \gamma_K^{AB} \check{L}_{AB}^D (\widehat{\chi}_K)_{\bullet D} - \gamma_K^{AB} \check{L}_{A\bullet}^D (\widehat{\chi}_K)_{BD}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \operatorname{div}(\check{\widehat{\chi}}) &= \frac{1}{2} \nabla \widetilde{\operatorname{tr} \underline{\chi}} - \frac{1}{2} \operatorname{tr} \underline{\chi} \check{\zeta} - \frac{1}{2} \widetilde{\operatorname{tr} \underline{\chi}} \zeta_K + \check{\zeta} \cdot \widehat{\chi}_K + \zeta_K \cdot \check{\chi} + \check{\zeta} \cdot \check{\chi} + \check{\beta} - \widetilde{\gamma^{AB}} \nabla_A (\widehat{\chi}_K)_B \\ &\quad - \gamma_K^{AB} \check{L}_{AB}^D (\widehat{\chi}_K)_{\bullet D} - \gamma_K^{AB} \check{L}_{A\bullet}^D (\widehat{\chi}_K)_{BD}. \end{aligned}$$

Next, we recall from (2.26)

$$\mu = -\operatorname{div} \eta + \frac{1}{2} \widehat{\chi} \cdot \widehat{\chi} - \rho.$$

Hence, we have

$$\mu_K = -\operatorname{div}_K \eta_K + \frac{1}{2} \widehat{\chi}_K \cdot \widehat{\chi}_K - \rho_K.$$

Taking the difference, we obtain

$$\begin{aligned} \check{\mu} &= -\operatorname{div} \check{\eta} + (\operatorname{div} - \operatorname{div}_K) \eta_K + \frac{1}{2} \widehat{\chi}_K \cdot \widehat{\chi} + \frac{1}{2} \check{\chi} \cdot \widehat{\chi}_K + \frac{1}{2} \widehat{\chi} \cdot \check{\chi} - \check{\rho} \\ &= -\operatorname{div} \check{\eta} - \check{\rho} + (\gamma^{AB} - (\gamma_K)^{AB}) \nabla_A (\eta_K)_B - (\gamma_K)^{AB} \check{L}_{AB}^C (\eta_K)_C \\ &\quad + \frac{1}{2} \widehat{\chi}_K \cdot \widehat{\chi} + \frac{1}{2} \check{\chi} \cdot \widehat{\chi}_K + \frac{1}{2} \widehat{\chi} \cdot \check{\chi}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \underline{\check{\mu}} &= -\operatorname{div} \underline{\check{\eta}} - \check{\rho} + (\gamma^{AB} - (\gamma_K)^{AB}) \nabla_A (\underline{\eta}_K)_B - (\gamma_K)^{AB} \check{L}_{AB}^C (\underline{\eta}_K)_C \\ &\quad + \frac{1}{2} \widehat{\chi}_K \cdot \widehat{\chi} + \frac{1}{2} \check{\chi} \cdot \widehat{\chi}_K + \frac{1}{2} \widehat{\chi} \cdot \check{\chi}. \end{aligned}$$

We consider the torsion equation

$$\operatorname{curl} \eta = -\operatorname{curl} \underline{\eta} = \sigma + \frac{1}{2} \widehat{\chi} \wedge \widehat{\chi},$$

which implies

$$\operatorname{curl}_K \eta_K = -\operatorname{curl}_K \underline{\eta}_K = \sigma_K + \frac{1}{2} \widehat{\chi}_K \wedge \widehat{\chi}_K.$$

Taking the difference, we infer from Proposition 2.22 that

$$\begin{aligned} \operatorname{curl} \check{\eta} &= \operatorname{curl} \eta - \operatorname{curl} \eta_K = \operatorname{curl} \eta - \operatorname{curl}_K \eta_K - (\operatorname{curl} - \operatorname{curl}_K) \eta_K \\ &= \check{\sigma} + \frac{1}{2} \check{\chi} \wedge \check{\chi} + \frac{1}{2} \check{\chi} \wedge \widehat{\chi}_K + \frac{1}{2} \widehat{\chi}_K \wedge \check{\chi} - \widetilde{\epsilon^{AB}} \nabla_A (\eta_K)_B + \epsilon_K^{AB} \check{L}_{AB}^C (\eta_K)_C. \end{aligned}$$

Similarly, we have

$$-\operatorname{curl} \check{\eta} = \check{\sigma} + \frac{1}{2} \check{\chi} \wedge \check{\chi} + \frac{1}{2} \check{\chi} \wedge \hat{\chi}_K + \frac{1}{2} \hat{\chi}_K \wedge \check{\chi} + \widetilde{\epsilon^{AB}} \nabla_A (\eta_K)_{B-} \in_K^{AB} \check{L}_{AB}^C (\eta_K)_C.$$

Finally, we recall from Proposition 2.16

$$\Omega \nabla_4 (\Omega \omega) = \frac{1}{2} \Omega^2 \rho + \frac{3}{4} |\Omega \eta - \Omega \underline{\eta}|^2 - \frac{1}{4} (\Omega \eta - \Omega \underline{\eta}) \cdot (\Omega \eta + \Omega \underline{\eta}) - \frac{1}{8} |\Omega \eta + \Omega \underline{\eta}|^2,$$

which implies

$$\begin{aligned} \Omega_K \nabla_{(e_4)_K} \omega_K &= \frac{1}{2} \Omega_K^2 \rho_K + \frac{3}{4} |\Omega_K \eta_K - \Omega_K \underline{\eta}_K|^2 - \frac{1}{4} (\Omega_K \eta_K - \Omega_K \underline{\eta}_K) \cdot (\Omega_K \eta_K + \Omega_K \underline{\eta}_K) \\ &\quad - \frac{1}{8} |\Omega_K \eta_K + \Omega_K \underline{\eta}_K|^2. \end{aligned}$$

Taking the difference, we obtain

$$\begin{aligned} \Omega \nabla_4 (\widetilde{\Omega \omega}) &= \frac{3}{4} \Omega^2 |\eta - \underline{\eta}|^2 - \frac{1}{4} \Omega^2 (\eta - \underline{\eta}) \cdot (\eta + \underline{\eta}) - \frac{1}{8} \Omega^2 |\eta + \underline{\eta}|^2 + \frac{1}{2} \Omega^2 \rho \\ &\quad - \frac{3}{4} \Omega_K^2 |\eta_K - \underline{\eta}_K|^2 + \frac{1}{4} \Omega_K^2 (\eta_K - \underline{\eta}_K) \cdot (\eta_K + \underline{\eta}_K) + \frac{1}{8} \Omega_K^2 |\eta_K + \underline{\eta}_K|^2 - \frac{1}{2} \Omega_K^2 \rho_K \\ &= \frac{3}{4} |\widetilde{\Omega \eta} - \widetilde{\Omega \underline{\eta}}|^2 + \frac{3}{2} (\widetilde{\Omega \eta} - \widetilde{\Omega \underline{\eta}}) \cdot (\Omega_K \eta_K - \Omega_K \underline{\eta}_K) - \frac{1}{4} (\widetilde{\Omega \eta} - \widetilde{\Omega \underline{\eta}}) \cdot (\widetilde{\Omega \eta} + \widetilde{\Omega \underline{\eta}}) \\ &\quad - \frac{1}{4} (\widetilde{\Omega \eta} - \widetilde{\Omega \underline{\eta}}) \cdot (\Omega_K \eta_K + \Omega_K \underline{\eta}_K) - \frac{1}{4} (\Omega_K \eta_K - \Omega_K \underline{\eta}_K) \cdot (\widetilde{\Omega \eta} + \widetilde{\Omega \underline{\eta}}) - \frac{1}{8} |\widetilde{\Omega \eta} + \widetilde{\Omega \underline{\eta}}|^2 \\ &\quad - \frac{1}{4} (\widetilde{\Omega \eta} + \widetilde{\Omega \underline{\eta}}) \cdot (\eta_K + \underline{\eta}_K) + \frac{1}{2} \widetilde{\Omega^2 \rho}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \Omega \nabla_3 (\widetilde{\Omega \omega}) &= \frac{3}{4} |\widetilde{\Omega \eta} - \widetilde{\Omega \underline{\eta}}|^2 + \frac{3}{2} (\widetilde{\Omega \eta} - \widetilde{\Omega \underline{\eta}}) \cdot (\Omega_K \eta_K - \Omega_K \underline{\eta}_K) + \frac{1}{4} (\widetilde{\Omega \eta} - \widetilde{\Omega \underline{\eta}}) \cdot (\widetilde{\Omega \eta} + \widetilde{\Omega \underline{\eta}}) \\ &\quad + \frac{1}{4} (\widetilde{\Omega \eta} - \widetilde{\Omega \underline{\eta}}) \cdot (\Omega_K \eta_K + \Omega_K \underline{\eta}_K) + \frac{1}{4} (\Omega_K \eta_K - \Omega_K \underline{\eta}_K) \cdot (\widetilde{\Omega \eta} + \widetilde{\Omega \underline{\eta}}) - \frac{1}{8} |\widetilde{\Omega \eta} + \widetilde{\Omega \underline{\eta}}|^2 \\ &\quad - \frac{1}{4} (\widetilde{\Omega \eta} + \widetilde{\Omega \underline{\eta}}) \cdot (\eta_K + \underline{\eta}_K) + \frac{1}{2} \widetilde{\Omega^2 \rho} + \check{b}^A \nabla_{\partial_{x^a}} (\Omega_K \omega_K). \end{aligned}$$

This concludes the proof of Proposition A.1.  $\square$

## A.2 Proof of Proposition 2.36

We recall from Corollary 2.18 that

$$\begin{aligned} \Omega \nabla_4 [\mu] + \Omega \operatorname{tr} \chi [\mu] &= \Omega [F] + \Omega (\operatorname{tr} \chi \rho - 2\eta \cdot \beta), \\ [F] &:= \widehat{\chi} \cdot (\nabla \widehat{\otimes} \eta) + (\eta - \underline{\eta}) \cdot (\nabla \operatorname{tr} \chi + \operatorname{tr} \chi \zeta) + \frac{1}{2} \operatorname{tr} \chi (|\eta|^2 - |\underline{\eta}|^2) \\ &\quad - \frac{1}{2} (\operatorname{tr} \chi |\widehat{\chi}|^2 + \operatorname{tr} \chi (\widehat{\chi} \cdot \widehat{\chi})) + 2\eta \cdot \widehat{\chi} \cdot \underline{\eta}, \end{aligned}$$

which implies

$$\Omega_K (\nabla_K)_{(e_4)_K} [\mu]_K + \Omega_K \operatorname{tr}_K \chi_K [\mu]_K = \Omega_K [F]_K + \Omega_K (\operatorname{tr} \chi_K \rho_K - 2\eta_K \cdot \beta_K).$$

Taking the difference, we obtain

$$\begin{aligned}
\Omega \nabla_4 [\widetilde{\mu}] + \Omega \operatorname{tr} \chi [\widetilde{\mu}] &= \widetilde{\Omega[F]} + \Omega(\operatorname{tr} \chi \rho - 2\eta \cdot \beta) - \Omega_K(\operatorname{tr} \chi_K \rho_K - 2\eta_K \cdot \beta_K) \\
&= \widetilde{\Omega[F]} + \widetilde{\Omega \operatorname{tr} \chi \rho_K} + \Omega \operatorname{tr} \chi \check{\rho} - 2\widetilde{\Omega \eta} \cdot \beta_K - 2\Omega_K \eta_K \cdot \check{\beta} - 2\widetilde{\Omega \eta} \cdot \check{\beta} \\
&= \Omega \operatorname{tr} \chi \check{\rho} + \widetilde{\Omega[F]} + \mathbf{O}_3^1 \cdot \Gamma_g + \mathbf{O}_3^2 \cdot \check{\beta} - 2\widetilde{\Omega \eta} \cdot \check{\beta}.
\end{aligned}$$

Next, we compute

$$\begin{aligned}
\Omega[F] &= -2\widehat{\chi} \cdot d_2^* \eta + (\eta - \underline{\eta}) \cdot \nabla \operatorname{tr} \chi + (\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_1^0 + \Gamma_g) \cdot (\mathbf{O}_3^2 + \Gamma_g) \\
&\quad + (\mathbf{O}_1^0 + \Gamma_g) \cdot (\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_3^2 + \Gamma_b) + (\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_3^2 + \Gamma_g) \\
&= -2\widehat{\chi} \cdot d_2^* \eta + (\eta - \underline{\eta}) \cdot \nabla \operatorname{tr} \chi + \mathbf{O}_7^4 + \mathbf{O}_4^2 \cdot \Gamma_b + r^{-1} \Gamma_g \cdot \Gamma_b.
\end{aligned}$$

Hence, we deduce from Proposition 2.22 that

$$\begin{aligned}
\widetilde{\Omega[F]} &= -2\check{\chi} \cdot (d_2^*)_K \eta_K - 2\widehat{\chi}_K \cdot \widetilde{d_2^* \eta} - 2\check{\chi} \cdot \widetilde{d_2^* \eta} + (\check{\eta} - \underline{\eta}) \cdot \nabla_K \operatorname{tr}_K \chi_K + (\eta_K - \underline{\eta}_K) \cdot \nabla \operatorname{tr} \chi \\
&\quad + (\check{\eta} - \underline{\eta}) \cdot \nabla \operatorname{tr} \chi + \mathbf{O}_4^2 \cdot \Gamma_b + r^{-1} \Gamma_g \cdot \Gamma_b \\
&= \mathbf{O}_3^2 \cdot \widetilde{d_2^* \eta} - 2\check{\chi} \cdot \widetilde{d_2^* \eta} + \mathbf{O}_3^2 \cdot \nabla \operatorname{tr} \chi + (\check{\eta} - \underline{\eta}) \cdot \nabla \operatorname{tr} \chi + \mathbf{O}_4^2 \cdot \Gamma_b + r^{-1} \Gamma_g \cdot \Gamma_b \\
&= \mathbf{O}_3^2 \cdot (d_2^* \check{\eta} + \check{L} \cdot \mathbf{O}_3^2) - 2\check{\chi} \cdot (d_2^* \check{\eta} + \check{L} \cdot \mathbf{O}_3^2) + \mathbf{O}_3^2 \cdot \nabla \operatorname{tr} \chi + (\check{\eta} - \underline{\eta}) \cdot \nabla \operatorname{tr} \chi + \mathbf{O}_4^2 \cdot \Gamma_b + r^{-1} \Gamma_g \cdot \Gamma_b \\
&= \mathbf{O}_3^2 \cdot (d_2^* \check{\eta}) - 2\check{\chi} \cdot (d_2^* \check{\eta}) + \mathbf{O}_6^4 \cdot \check{L} + r^{-1} \Gamma_g \cdot \check{L} + \mathbf{O}_3^2 \cdot \nabla \operatorname{tr} \chi + (\check{\eta} - \underline{\eta}) \cdot \nabla \operatorname{tr} \chi + \mathbf{O}_4^2 \cdot \Gamma_b + r^{-1} \Gamma_g \cdot \Gamma_b.
\end{aligned}$$

Thus, we obtain

$$\begin{aligned}
\Omega \nabla_4 [\widetilde{\mu}] + \Omega \operatorname{tr} \chi [\widetilde{\mu}] &= \Omega \operatorname{tr} \chi \check{\rho} + \mathbf{O}_3^2 \cdot (d_2^* \check{\eta}, \nabla \operatorname{tr} \chi, \check{\beta}) + \mathbf{O}_3^1 \cdot (\Gamma_g, \check{L}) - 2\check{\chi} \cdot (d_2^* \check{\eta}) + r^{-1} \Gamma_g \cdot \check{L} \\
&\quad + (\check{\eta} - \underline{\eta}) \cdot \nabla \operatorname{tr} \chi + r^{-1} \Gamma_g \cdot \Gamma_b - 2\widetilde{\Omega \eta} \cdot \check{\beta}.
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
\Omega \nabla_3 [\underline{\mu}] + \Omega \operatorname{tr} \underline{\chi} [\underline{\mu}] &= \Omega \operatorname{tr} \underline{\chi} \check{\rho} + \mathbf{O}_3^2 \cdot (d_2^* \check{\eta}, \nabla \operatorname{tr} \underline{\chi}, \check{\beta}) + \mathbf{O}_3^1 \cdot (\Gamma_g, \check{L}) - 2\check{\underline{\chi}} \cdot (d_2^* \check{\eta}) + r^{-1} \Gamma_b \cdot \check{L} \\
&\quad + (\check{\underline{\eta}} - \underline{\eta}) \cdot \nabla \operatorname{tr} \underline{\chi} + r^{-1} \Gamma_b \cdot \Gamma_b + 2\widetilde{\Omega \underline{\eta}} \cdot \check{\beta}.
\end{aligned}$$

This concludes the proof of Proposition 2.36.

### A.3 Proof of Propositions 2.37 and 2.38

We first prove the following lemma.

**Lemma A.3.** *Let  $k, k' = 0, 1, 2$ ,  $\lambda_0$  a real number and  $\nabla$  a first order elliptic operator. Then, we have the following properties:*

1. Assume that  $\psi, h \in \mathfrak{s}_k$  and  $\psi' \in \mathfrak{s}_{k'}$  satisfy the following equation:

$$\nabla_3 \psi + \lambda_0 \operatorname{tr} \underline{\chi} \psi = \nabla \psi' + h. \quad (\text{A.1})$$

Then, we have

$$\nabla_3 \not\psi + \lambda_0 \operatorname{tr} \underline{\chi} \not\psi = \nabla \not\psi' + h^{[1]}, \quad (\text{A.2})$$

where

$$h^{[1]} = \not\psi h + (\Gamma_g + \mathbf{O}_3^2) h + (\Gamma_g + \mathbf{O}_3^2) \cdot \psi'^{(1)} + (\Gamma_b + \mathbf{O}_3^2) \cdot \psi^{(1)} + \Gamma_b^{(1)} \cdot \psi.$$

2. Assume that  $\psi, h \in \mathfrak{s}_k$  and  $\psi' \in \mathfrak{s}_{k'}$  satisfy the following equation:

$$\nabla_4 \psi + \lambda_0 \operatorname{tr} \chi \psi = \nabla \psi' + h. \quad (\text{A.3})$$

Then, we have

$$\nabla_4 \not\psi + \lambda_0 \operatorname{tr} \chi \not\psi = \nabla \not\psi' + h^{[1]}, \quad (\text{A.4})$$

where

$$h^{[1]} = \not\psi h + (\Gamma_g + \mathbf{O}_3^2)h + (\Gamma_g + \mathbf{O}_3^2) \cdot (\psi'^{(1)}, \psi^{(1)}) + \Gamma_g^{(1)} \cdot \psi.$$

*Proof.* We have from (A.1)

$$\Omega \nabla_3 \psi + \lambda_0 \Omega \operatorname{tr} \underline{\chi} \psi = \Omega \nabla \psi' + \Omega h.$$

Differentiating it by  $\not\psi$ , we obtain

$$\Omega \nabla_3(\not\psi \psi) + \lambda_0 r \nabla(\Omega \operatorname{tr} \underline{\chi}) \psi + \lambda_0 \Omega \operatorname{tr} \underline{\chi}(\not\psi \psi) = r \nabla \Omega \cdot \nabla \psi' + \Omega \nabla(\not\psi \psi') + \not\psi(\Omega h) + [\Omega \nabla_3, \not\psi] \psi,$$

which implies from Corollary 2.43

$$\begin{aligned} & \Omega \nabla_3(\not\psi \psi) + \lambda_0 \Omega \operatorname{tr} \underline{\chi}(\not\psi \psi) \\ &= \Omega \nabla(\not\psi \psi') + (\Gamma_g + \mathbf{O}_3^2) \cdot \nabla \psi' + (\Gamma_g^{(1)} + \mathbf{O}_3^2) \cdot \psi + (\Gamma_g + \mathbf{O}_3^2)h + \Omega \not\psi h \\ &+ (\Gamma_b + \mathbf{O}_3^2) \cdot \not\psi \psi + (\Gamma_b^{(1)} + \mathbf{O}_3^2) \cdot \psi \\ &= \Omega \nabla(\not\psi \psi') + (\Gamma_g + \mathbf{O}_3^2)h + (\Gamma_b + \mathbf{O}_3^2) \cdot \psi'^{(1)} + (\Gamma_g + \mathbf{O}_3^2) \cdot \psi^{(1)} + \Gamma_b^{(1)} \cdot \psi + \Omega \not\psi h. \end{aligned}$$

Hence, we obtain

$$\begin{aligned} & \nabla_3(\not\psi \psi) + \lambda_0 \operatorname{tr} \underline{\chi}(\not\psi \psi) \\ &= \nabla(\not\psi \psi') + (\Gamma_g + \mathbf{O}_3^2)h + (\Gamma_g + \mathbf{O}_3^2) \cdot \psi'^{(1)} + (\Gamma_b + \mathbf{O}_3^2) \cdot \psi^{(1)} + \Gamma_b^{(1)} \cdot \psi + \not\psi h, \end{aligned} \quad (\text{A.5})$$

which implies (A.2). The proof of (A.4) is similar. This concludes the proof of Lemma A.3.  $\square$

We next prove the following lemma concerning the linearization of Bianchi equations.

**Lemma A.4.** *Let  $k, k' = 0, 1, 2$ ,  $\lambda_0$  a real number and  $\nabla$  a first order elliptic operator. Then, we have the following properties:*

1. Assume that  $\psi, h \in \mathfrak{s}_k$  and  $\psi' \in \mathfrak{s}_{k'}$  satisfy the following equation:

$$\nabla_3 \psi + \lambda_0 \operatorname{tr} \underline{\chi} \psi = \nabla \psi' + h. \quad (\text{A.6})$$

Then, we have

$$\nabla_3 \check{\psi} + \lambda_0 \operatorname{tr} \underline{\chi} \check{\psi} = \nabla \psi' + \check{h} + \check{J} \cdot \psi_K + \check{L} \cdot \psi'_K + \Gamma_g \cdot \mathfrak{d}^{\leq 1} \psi_K. \quad (\text{A.7})$$

2. Assume that  $\psi, h \in \mathfrak{s}_k$  and  $\psi' \in \mathfrak{s}_{k'}$  satisfy the following equation:

$$\nabla_4 \psi + \lambda_0 \operatorname{tr} \chi \psi = \nabla \psi' + h. \quad (\text{A.8})$$

Then, we have

$$\nabla_4 \check{\psi} + \lambda_0 \operatorname{tr} \chi \check{\psi} = \nabla \psi' + \check{h} + \check{J} \cdot \psi_K + \check{L} \cdot \psi'_K + \Gamma_g \cdot \mathfrak{d}^{\leq 1} \psi_K. \quad (\text{A.9})$$

*Proof.* Assume that we have

$$\nabla_3 \psi + \lambda_0 \operatorname{tr} \underline{\chi} \psi = \nabla \psi' + h.$$

Then, we obtain

$$(\nabla_K)_{(e_3)_K} \psi_K + \lambda_0 \operatorname{tr}_K \underline{\chi}_K \psi_K = \nabla_K \psi'_K + h_K.$$

Taking the difference, we deduce

$$\nabla_3 \psi - (\nabla_K)_{(e_3)_K} \psi_K + \lambda_0 (\operatorname{tr} \underline{\chi} \psi - \operatorname{tr}_K \underline{\chi}_K \psi_K) = \nabla \psi' - \nabla_K \psi'_K + \check{h}. \quad (\text{A.10})$$

Recall from (2.38) that

$$\begin{aligned} e_3 - (e_3)_K &= \Omega^{-1}(\Omega e_3 - \Omega_K(e_3)_K) + \left( \frac{\Omega_K}{\Omega} - 1 \right) (e_3)_K \\ &= \Omega^{-1} \check{b}^A \partial_{x^A} + \Omega^{-1} \check{\Omega} (e_3)_K. \end{aligned}$$

Thus, we obtain

$$(\nabla_K)_{e_3 - (e_3)_K} \psi_K = \Gamma_g \cdot \partial_{x^A}(\psi_K) + r \Gamma_g \cdot (\nabla_K)_{(e_3)_K} \psi_K = \Gamma_g \cdot \mathfrak{d} \psi_K.$$

Combining with (A.10), we deduce

$$\begin{aligned} \nabla_3 \check{\psi} + \lambda_0 \operatorname{tr} \underline{\chi} \check{\psi} &= \nabla_3(\psi - \psi_K) + \lambda_0 \operatorname{tr} \underline{\chi}(\psi - \psi_K) \\ &= ((\nabla_K)_{(e_3)_K} - \nabla_3) \psi_K + \lambda_0 (\operatorname{tr}_K \underline{\chi}_K - \operatorname{tr} \underline{\chi}) \psi_K + \nabla \check{\psi}' + (\nabla - \nabla_K) \psi'_K + \check{h} \\ &= ((\nabla_K)_{(e_3)_K - e_3} + (\nabla_K - \nabla)) \psi_K - \lambda_0 \widetilde{\operatorname{tr} \underline{\chi}} \psi_K + \nabla \check{\psi}' + (\nabla - \nabla_K) \psi'_K + \check{h} \\ &= \nabla \psi' + \check{h} - (\nabla - \nabla_K)_{e_3} \psi_K + (\nabla - \nabla_K) \psi'_K + \Gamma_g \cdot \mathfrak{d}^{\leq 1} \psi_K. \end{aligned}$$

Recalling Lemma 2.39, we have

$$\begin{aligned} (\nabla - \nabla_K)_{e_3} \psi_K &= \check{J} \cdot \psi_K + \check{L} \cdot \psi_K, \\ (\nabla - \nabla_K) \psi'_K &= \check{L} \cdot \psi_K. \end{aligned}$$

Thus, we obtain

$$\nabla_3 \check{\psi} + \lambda_0 \operatorname{tr} \underline{\chi} \check{\psi} = \nabla \psi' + \check{h} + (\check{J}, \check{L}) \cdot \psi_K + \check{L} \cdot \psi'_K + \Gamma_g \cdot \mathfrak{d}^{\leq 1} \psi_K,$$

which implies (A.7). The proof of (A.9) is similar. This concludes the proof of Lemma A.4.  $\square$

We are now ready to prove the following proposition, which implies directly Propositions 2.37 and 2.38.

**Proposition A.5.** *We have the following linearized Bianchi equations:*

$$\nabla_4 \underline{\check{\alpha}} + \frac{1}{2} \text{tr} \chi \underline{\check{\alpha}} = \nabla \widehat{\otimes} \underline{\check{\beta}} + \mathfrak{L}[\nabla_4 \underline{\check{\alpha}}] + \text{Err}[\nabla_4 \underline{\check{\alpha}}],$$

$$\mathfrak{L}[\nabla_4 \underline{\check{\alpha}}] := \Gamma_b^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \underline{\check{\alpha}} + \mathbf{O}_3^2 \cdot (\underline{\check{\beta}}, \underline{\check{\rho}}, \underline{\check{\sigma}}),$$

$$\text{Err}[\nabla_4 \underline{\check{\alpha}}] := \Gamma_g \cdot (\underline{\check{\alpha}}, \underline{\check{\beta}}) + \Gamma_b \cdot (\underline{\check{\rho}}, \underline{\check{\sigma}}),$$

$$\nabla_4 \underline{\check{\rho}} + \frac{1}{2} \text{tr} \chi \underline{\check{\rho}} = -\nabla \widehat{\otimes} \underline{\check{\beta}} + \mathfrak{L}[\nabla_4 \underline{\check{\rho}}] + \text{Err}[\nabla_4 \underline{\check{\rho}}],$$

$$\mathfrak{L}[\nabla_4 \underline{\check{\rho}}] = \Gamma_b^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \underline{\check{\alpha}}^{(1)} + \mathbf{O}_3^2 \cdot (\underline{\check{\beta}}^{(1)}, \underline{\check{\rho}}^{(1)}, \underline{\check{\sigma}}^{(1)}),$$

$$\text{Err}[\nabla_4 \underline{\check{\rho}}] = \Gamma_g^{(1)} \cdot (\underline{\check{\alpha}}, \underline{\check{\beta}}) + \Gamma_b^{(1)} \cdot (\underline{\check{\rho}}, \underline{\check{\sigma}}) + \Gamma_g \cdot (\underline{\check{\alpha}}^{(1)}, \underline{\check{\beta}}^{(1)}) + \Gamma_b \cdot (\underline{\check{\rho}}^{(1)}, \underline{\check{\sigma}}^{(1)}),$$

$$\nabla_3 \underline{\check{\beta}} + 2 \text{tr} \chi \underline{\check{\beta}} = -\text{div} \underline{\check{\alpha}} + \mathfrak{L}[\nabla_3 \underline{\check{\beta}}] + \text{Err}[\nabla_3 \underline{\check{\beta}}],$$

$$\mathfrak{L}[\nabla_3 \underline{\check{\beta}}] := \Gamma_b^{(1)} \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot \underline{\check{\beta}} + \mathbf{O}_3^2 \cdot \underline{\check{\alpha}},$$

$$\text{Err}[\nabla_3 \underline{\check{\beta}}] := \Gamma_b \cdot \underline{\check{\beta}} + \Gamma_g \cdot \underline{\check{\alpha}},$$

$$\nabla_3 \underline{\check{\rho}} + 2 \text{tr} \chi \underline{\check{\rho}} = -\text{div} \underline{\check{\rho}} + \mathfrak{L}[\nabla_3 \underline{\check{\rho}}] + \text{Err}[\nabla_3 \underline{\check{\rho}}],$$

$$\mathfrak{L}[\nabla_3 \underline{\check{\rho}}] := \Gamma_b^{(1)} \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot \underline{\check{\beta}}^{(1)} + \mathbf{O}_3^2 \cdot \underline{\check{\alpha}}^{(1)},$$

$$\text{Err}[\nabla_3 \underline{\check{\rho}}] := \Gamma_b^{(1)} \cdot \underline{\check{\beta}} + \Gamma_b \cdot \underline{\check{\beta}}^{(1)} + \Gamma_g^{(1)} \cdot \underline{\check{\alpha}} + \Gamma_g \cdot \underline{\check{\alpha}}^{(1)},$$

$$\nabla_4 \underline{\check{\beta}} + \text{tr} \chi \underline{\check{\beta}} = -\nabla \underline{\check{\rho}} + {}^* \nabla \underline{\check{\sigma}} + \mathfrak{L}[\nabla_4 \underline{\check{\beta}}] + \text{Err}[\nabla_4 \underline{\check{\beta}}],$$

$$\mathfrak{L}[\nabla_4 \underline{\check{\beta}}] := \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \underline{\check{\beta}} + \mathbf{O}_3^2 \cdot (\underline{\check{\rho}}, \underline{\check{\sigma}}, \underline{\check{\beta}}),$$

$$\text{Err}[\nabla_4 \underline{\check{\beta}}] := \Gamma_g \cdot (\underline{\check{\beta}}, \underline{\check{\rho}}, \underline{\check{\sigma}}) + \Gamma_b \cdot \underline{\check{\beta}},$$

$$\nabla_4 \underline{\check{\rho}} + \text{tr} \chi \underline{\check{\rho}} := -\nabla \underline{\check{\rho}} + {}^* \nabla \underline{\check{\sigma}} + \mathfrak{L}[\nabla_4 \underline{\check{\rho}}] + \text{Err}[\nabla_4 \underline{\check{\rho}}],$$

$$\mathfrak{L}[\nabla_4 \underline{\check{\rho}}] := \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \underline{\check{\beta}}^{(1)} + \mathbf{O}_3^2 \cdot (\underline{\check{\rho}}^{(1)}, \underline{\check{\sigma}}^{(1)}, \underline{\check{\beta}}^{(1)}),$$

$$\text{Err}[\nabla_4 \underline{\check{\rho}}] := \Gamma_g^{(1)} \cdot (\underline{\check{\beta}}, \underline{\check{\rho}}, \underline{\check{\sigma}}) + \Gamma_g \cdot (\underline{\check{\beta}}^{(1)}, \underline{\check{\rho}}^{(1)}, \underline{\check{\sigma}}^{(1)}) + \Gamma_b^{(1)} \cdot \underline{\check{\beta}} + \Gamma_b \cdot \underline{\check{\beta}}^{(1)},$$

$$\nabla_3(\underline{\check{\rho}}, \underline{\check{\sigma}}) + \frac{3}{2} \text{tr} \chi (\underline{\check{\rho}}, \underline{\check{\sigma}}) = -d_1 \underline{\check{\beta}} + \mathfrak{L}[\nabla_3 \underline{\check{\rho}}] + \text{Err}[\nabla_3 \underline{\check{\rho}}],$$

$$\mathfrak{L}[\nabla_3 \underline{\check{\rho}}] := \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\underline{\check{\alpha}}, \underline{\check{\beta}}),$$

$$\text{Err}[\nabla_3 \underline{\check{\rho}}] := \Gamma_g \cdot (\underline{\check{\alpha}}, \underline{\check{\beta}}),$$

$$\nabla_3(\underline{\check{\rho}}, \underline{\check{\sigma}}) + \frac{3}{2} \text{tr} \chi (\underline{\check{\rho}}, \underline{\check{\sigma}}) = -d_1 \underline{\check{\rho}} + \mathfrak{L}[\nabla_3(\underline{\check{\rho}}, \underline{\check{\sigma}})] + \text{Err}[\nabla_3(\underline{\check{\rho}}, \underline{\check{\sigma}})],$$

$$\mathfrak{L}[\nabla_3(\underline{\check{\rho}}, \underline{\check{\sigma}})] := \Gamma_b^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\underline{\check{\alpha}}^{(1)}, \underline{\check{\beta}}^{(1)}, \underline{\check{\rho}}^{(1)}, \underline{\check{\sigma}}^{(1)}),$$

$$\text{Err}[\nabla_3(\underline{\check{\rho}}, \underline{\check{\sigma}})] := \Gamma_g^{(1)} \cdot (\underline{\check{\alpha}}, \underline{\check{\beta}}) + \Gamma_g \cdot (\underline{\check{\alpha}}^{(1)}, \underline{\check{\beta}}^{(1)}) + \Gamma_b \cdot (\underline{\check{\rho}}^{(1)}, \underline{\check{\sigma}}^{(1)}) + \Gamma_b^{(1)} \cdot (\underline{\check{\rho}}, \underline{\check{\sigma}}),$$

$$\nabla_4(\check{\rho}, -\check{\sigma}) + \frac{3}{2} \operatorname{tr} \chi(\check{\rho}, -\check{\sigma}) = d_1 \check{\beta} + \mathfrak{L}[\nabla_4 \check{\rho}] + \operatorname{Err}[\nabla_4 \check{\rho}],$$

$$\mathfrak{L}[\nabla_4 \check{\rho}] := \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\alpha}, \check{\beta}),$$

$$\operatorname{Err}[\nabla_4 \check{\rho}] := \Gamma_b \cdot \check{\alpha} + \Gamma_g \cdot \check{\beta},$$

$$\nabla_4(\check{\check{\rho}}, -\check{\check{\sigma}}) + \frac{3}{2} \operatorname{tr} \chi(\check{\check{\rho}}, -\check{\check{\sigma}}) = d_1 \check{\check{\beta}} + \mathfrak{L}[\nabla_4(\check{\check{\rho}}, -\check{\check{\sigma}})] + \operatorname{Err}[\nabla_4(\check{\check{\rho}}, \check{\check{\sigma}})],$$

$$\mathfrak{L}[\nabla_4(\check{\check{\rho}}, -\check{\check{\sigma}})] := \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\check{\alpha}}^{(1)}, \check{\check{\beta}}^{(1)}, \check{\check{\rho}}^{(1)}, \check{\check{\sigma}}^{(1)}),$$

$$\operatorname{Err}[\nabla_4(\check{\check{\rho}}, -\check{\check{\sigma}})] := \Gamma_b^{(1)} \cdot \check{\check{\alpha}} + \Gamma_b \cdot \check{\check{\alpha}}^{(1)} + \Gamma_g^{(1)} \cdot (\check{\check{\beta}}, \check{\check{\rho}}, \check{\check{\sigma}}) + \Gamma_g \cdot (\check{\check{\beta}}^{(1)}, \check{\check{\rho}}^{(1)}, \check{\check{\sigma}}^{(1)}),$$

$$\nabla_3 \check{\beta} + \operatorname{tr} \underline{\chi} \check{\beta} = \nabla \check{\rho} + {}^* \nabla \check{\sigma} + \mathfrak{L}[\nabla_3 \check{\beta}] + \operatorname{Err}[\nabla_3 \check{\beta}],$$

$$\mathfrak{L}[\nabla_3 \check{\beta}] := \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\beta} + \mathbf{O}_3^2 \cdot (\check{\beta}, \check{\rho}, \check{\sigma}),$$

$$\operatorname{Err}[\nabla_3 \check{\beta}] := \Gamma_b \cdot \check{\beta} + \Gamma_g \cdot (\check{\beta}, \check{\rho}, \check{\sigma}),$$

$$\nabla_3 \check{\check{\beta}} + \operatorname{tr} \underline{\chi} \check{\check{\beta}} = \nabla \check{\check{\rho}} + {}^* \nabla \check{\check{\sigma}} + \mathfrak{L}[\nabla_3 \check{\check{\beta}}] + \operatorname{Err}[\nabla_3 \check{\check{\beta}}],$$

$$\mathfrak{L}[\nabla_3 \check{\check{\beta}}] := \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\check{\beta}}^{(1)} + \mathbf{O}_3^2 \cdot (\check{\check{\beta}}^{(1)}, \check{\check{\rho}}^{(1)}, \check{\check{\sigma}}^{(1)}),$$

$$\operatorname{Err}[\nabla_3 \check{\check{\beta}}] := \Gamma_b^{(1)} \cdot \check{\check{\beta}} + \Gamma_b \cdot \check{\check{\beta}}^{(1)} + \Gamma_g^{(1)} \cdot (\check{\check{\beta}}, \check{\check{\rho}}, \check{\check{\sigma}}) + \Gamma_g \cdot (\check{\check{\beta}}^{(1)}, \check{\check{\rho}}^{(1)}, \check{\check{\sigma}}^{(1)}),$$

$$\nabla_4 \check{\beta} + 2 \operatorname{tr} \chi \check{\beta} = \operatorname{div} \check{\alpha} + \mathfrak{L}[\nabla_4 \check{\beta}] + \operatorname{Err}[\nabla_4 \check{\beta}],$$

$$\mathfrak{L}[\nabla_4 \check{\beta}] := \Gamma_g^{(1)} \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot (\check{\beta}, \check{\alpha}),$$

$$\operatorname{Err}[\nabla_4 \check{\beta}] := \Gamma_g \cdot (\check{\beta}, \check{\alpha}),$$

$$\nabla_4 \check{\check{\beta}} + 2 \operatorname{tr} \chi \check{\check{\beta}} = \operatorname{div} \check{\check{\alpha}} + \mathfrak{L}[\nabla_4 \check{\check{\beta}}] + \operatorname{Err}[\nabla_4 \check{\check{\beta}}],$$

$$\mathfrak{L}[\nabla_4 \check{\check{\beta}}] := \Gamma_g^{(1)} \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot (\check{\check{\beta}}^{(1)}, \check{\check{\alpha}}^{(1)}),$$

$$\operatorname{Err}[\nabla_4 \check{\check{\beta}}] := \Gamma_g^{(1)} \cdot (\check{\check{\beta}}, \check{\check{\alpha}}) + \Gamma_g \cdot (\check{\check{\beta}}^{(1)}, \check{\check{\alpha}}^{(1)}),$$

$$\nabla_3 \check{\alpha} + \frac{1}{2} \operatorname{tr} \underline{\chi} \check{\alpha} = \nabla \widehat{\otimes} \check{\beta} + \mathfrak{L}[\nabla_3 \check{\alpha}] + \operatorname{Err}[\nabla_3 \check{\alpha}],$$

$$\mathfrak{L}[\nabla_3 \check{\alpha}] := \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\alpha} + \mathbf{O}_3^2 \cdot (\check{\rho}, \check{\sigma}, \check{\beta}),$$

$$\operatorname{Err}[\nabla_3 \check{\alpha}] := \Gamma_b \cdot \check{\alpha} + \Gamma_g \cdot \check{\beta},$$

$$\nabla_3 \check{\check{\alpha}} + \frac{1}{2} \operatorname{tr} \underline{\chi} \check{\check{\alpha}} = \nabla \widehat{\otimes} \check{\check{\beta}} + \mathfrak{L}[\nabla_3 \check{\check{\alpha}}] + \operatorname{Err}[\nabla_3 \check{\check{\alpha}}],$$

$$\mathfrak{L}[\nabla_3 \check{\check{\alpha}}] := \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\check{\alpha}}^{(1)} + \mathbf{O}_3^2 \cdot (\check{\check{\rho}}^{(1)}, \check{\check{\sigma}}^{(1)}, \check{\check{\beta}}^{(1)}),$$

$$\operatorname{Err}[\nabla_3 \check{\check{\alpha}}] := \Gamma_b^{(1)} \cdot \check{\check{\alpha}} + \Gamma_b \cdot \check{\check{\alpha}}^{(1)} + \Gamma_g^{(1)} \cdot \check{\check{\beta}} + \Gamma_g \cdot \check{\check{\beta}}^{(1)}.$$

*Proof.* We recall from Proposition 2.19

$$\nabla_4 \underline{\alpha} + \frac{1}{2} \operatorname{tr} \chi \underline{\alpha} = -\nabla \widehat{\otimes} \underline{\beta} + h[\underline{\alpha}_4],$$

$$h[\underline{\alpha}_4] := 4\omega\underline{\alpha} - 3(\widehat{\chi}\rho - {}^*\widehat{\chi}\sigma) + (\zeta - 4\eta)\widehat{\otimes}\underline{\beta}.$$

Applying Lemma A.3, we deduce

$$\nabla_4 \not\partial \underline{\alpha} + \frac{1}{2} \operatorname{tr} \chi \not\partial \underline{\alpha} = -\nabla \widehat{\otimes} \not\partial \underline{\beta} + h[\underline{\alpha}_4]^{[1]},$$

where

$$\begin{aligned} h[\underline{\alpha}_4]^{[1]} &= \not\partial h[\underline{\alpha}_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\underline{\alpha}_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot (\underline{\alpha}^{(1)}, \underline{\beta}^{(1)}) + \Gamma_g^{(1)} \cdot \underline{\alpha} \\ &= \not\partial h[\underline{\alpha}_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\underline{\alpha}_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot (\check{\underline{\alpha}}^{(1)}, \check{\underline{\beta}}^{(1)}) + \Gamma_g^{(1)} \cdot \check{\underline{\alpha}} \\ &\quad + \Gamma_g \cdot \mathbf{O}_5^3 + \Gamma_g \cdot \mathbf{O}_4^2 + \mathbf{O}_7^4 + \Gamma_g^{(1)} \cdot \mathbf{O}_5^3. \end{aligned}$$

We compute

$$\begin{aligned} h[\underline{\alpha}_4] &= 4\omega\underline{\alpha} - 3(\widehat{\chi}\rho - {}^*\widehat{\chi}\sigma) + (\zeta - 4\eta)\widehat{\otimes}\underline{\beta} \\ &= (\mathbf{O}_2^1 + \Gamma_g)(\mathbf{O}_5^3 + \check{\underline{\alpha}}) + (\mathbf{O}_3^2 + \Gamma_b) \cdot (\mathbf{O}_3^1 + \check{\rho} + \check{\sigma}) + (\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_4^2 + \check{\underline{\beta}}) \\ &= \mathbf{O}_7^4 + \Gamma_g \cdot \mathbf{O}_5^3 + \mathbf{O}_2^1 \cdot \check{\underline{\alpha}} + \Gamma_g \cdot \check{\underline{\alpha}} + \mathbf{O}_6^3 + \Gamma_b \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\rho}, \check{\sigma}) + \Gamma_b \cdot (\check{\rho}, \check{\sigma}) \\ &\quad + \mathbf{O}_7^4 + \Gamma_g \cdot \mathbf{O}_4^2 + \mathbf{O}_3^2 \cdot \check{\underline{\beta}} + \Gamma_g \cdot \check{\underline{\beta}} \\ &= \mathbf{O}_6^3 + \Gamma_b \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\underline{\alpha}} + \mathbf{O}_3^2 \cdot (\check{\underline{\beta}}, \check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\underline{\alpha}}, \check{\underline{\beta}}) + \Gamma_b \cdot (\check{\rho}, \check{\sigma}), \end{aligned} \tag{A.11}$$

which implies

$$\begin{aligned} h[\underline{\alpha}_4]^{[1]} &= \mathbf{O}_6^3 + \Gamma_b^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\underline{\alpha}}^{(1)} + \mathbf{O}_3^2 \cdot (\check{\underline{\beta}}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}) \\ &\quad + \Gamma_g^{(1)} \cdot (\check{\underline{\alpha}}, \check{\underline{\beta}}) + \Gamma_b^{(1)} \cdot (\check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\underline{\alpha}}^{(1)}, \check{\underline{\beta}}^{(1)}) + \Gamma_b \cdot (\check{\rho}^{(1)}, \check{\sigma}^{(1)}). \end{aligned}$$

Applying Lemma A.4, we obtain

$$\begin{aligned} \nabla_4 \check{\underline{\alpha}} + \frac{1}{2} \operatorname{tr} \chi \check{\underline{\alpha}} &= -\nabla \widehat{\otimes} \check{\underline{\beta}} + \widetilde{h[\underline{\alpha}_4]} + \check{J} \cdot \underline{\alpha}_K + \check{L} \cdot \underline{\beta}_K + \Gamma_g \cdot \mathfrak{d}^{\leq 1} \underline{\alpha}_K \\ &= -\nabla \widehat{\otimes} \check{\underline{\beta}} + \widetilde{h[\underline{\alpha}_4]} + \Gamma_g^{(1)} \cdot \mathbf{O}_5^3 + \Gamma_g^{(1)} \cdot \mathbf{O}_4^2 + \Gamma_g \cdot \mathbf{O}_5^3 \\ &= -\nabla \widehat{\otimes} \check{\underline{\beta}} + \widetilde{h[\underline{\alpha}_4]} + \Gamma_g^{(1)} \cdot \mathbf{O}_4^2, \end{aligned}$$

and similarly

$$\nabla_4 \check{\underline{\alpha}} + \frac{1}{2} \operatorname{tr} \chi \check{\underline{\alpha}} = -\nabla \widehat{\otimes} \check{\underline{\beta}} + \widetilde{h[\underline{\alpha}_4]}^{[1]} + \Gamma_g^{(1)} \cdot \mathbf{O}_4^2.$$

Hence, we infer

$$\begin{aligned} \nabla_4 \check{\underline{\alpha}} + \frac{1}{2} \operatorname{tr} \chi \check{\underline{\alpha}} &= -\nabla \widehat{\otimes} \check{\underline{\beta}} + \widetilde{h[\underline{\alpha}_4]} + \Gamma_g^{(1)} \cdot \mathbf{O}_4^2 \\ &= -\nabla \widehat{\otimes} \check{\underline{\beta}} + \Gamma_b^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\underline{\alpha}} + \mathbf{O}_3^2 \cdot (\check{\underline{\beta}}, \check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\underline{\alpha}}, \check{\underline{\beta}}) + \Gamma_b \cdot (\check{\rho}, \check{\sigma}). \end{aligned}$$

Similarly, we have

$$\begin{aligned} \nabla_4 \check{\underline{\alpha}} + \frac{1}{2} \operatorname{tr} \chi \check{\underline{\alpha}} &= -\nabla \widehat{\otimes} \check{\underline{\beta}} + \widetilde{h[\underline{\alpha}_4]}^{[1]} + \Gamma_g^{(1)} \cdot \mathbf{O}_4^2 \\ &= -\nabla \widehat{\otimes} \check{\underline{\beta}} + \Gamma_b^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\underline{\alpha}}^{(1)} + \mathbf{O}_3^2 \cdot (\check{\underline{\beta}}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}) \\ &\quad + \Gamma_g^{(1)} \cdot (\check{\underline{\alpha}}, \check{\underline{\beta}}) + \Gamma_b^{(1)} \cdot (\check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\underline{\alpha}}^{(1)}, \check{\underline{\beta}}^{(1)}) + \Gamma_b \cdot (\check{\rho}^{(1)}, \check{\sigma}^{(1)}). \end{aligned}$$

Next, we recall from Proposition 2.19

$$\begin{aligned}\nabla_3 \underline{\beta} + 2 \operatorname{tr} \underline{\chi} \underline{\beta} &= -\operatorname{div} \underline{\alpha} + h[\underline{\beta}_3], \\ h[\underline{\beta}_3] &:= -2\underline{\omega} \underline{\beta} + (2\zeta - \eta) \cdot \underline{\alpha}.\end{aligned}$$

Applying Lemma A.3, we deduce

$$\nabla_3 \underline{\beta} + 2 \operatorname{tr} \underline{\chi} \underline{\beta} = -\operatorname{div} \underline{\beta} \alpha + h[\underline{\beta}_3]^{[1]},$$

where

$$\begin{aligned}h[\underline{\beta}_3]^{[1]} &= \underline{\beta} h[\underline{\beta}_3] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\underline{\beta}_3] + (\Gamma_g + \mathbf{O}_3^2) \cdot \underline{\alpha}^{(1)} + (\Gamma_b + \mathbf{O}_3^2) \cdot \underline{\beta}^{(1)} + \Gamma_b^{(1)} \cdot \underline{\beta} \\ &= \underline{\beta} h[\underline{\beta}_3] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\underline{\beta}_3] + (\Gamma_g + \mathbf{O}_3^2) \cdot \underline{\alpha}^{(1)} + (\Gamma_b + \mathbf{O}_3^2) \cdot \underline{\beta}^{(1)} + \Gamma_b^{(1)} \cdot \underline{\beta} \\ &\quad + \Gamma_b \cdot \mathbf{O}_5^3 + \Gamma_b \cdot \mathbf{O}_4^2 + \mathbf{O}_7^4 + \Gamma_b^{(1)} \cdot \mathbf{O}_4^2.\end{aligned}$$

We compute

$$\begin{aligned}h[\underline{\beta}_3] &= -2\underline{\omega} \underline{\beta} + (2\zeta - \eta) \cdot \underline{\alpha} \\ &= (\mathbf{O}_2^1 + \Gamma_b) \cdot (\mathbf{O}_4^2 + \underline{\beta}) + (\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_5^3 + \underline{\alpha}) \\ &= \mathbf{O}_6^3 + \Gamma_b \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot \underline{\beta} + \Gamma_b \cdot \underline{\beta} + \mathbf{O}_8^5 + \Gamma_g \cdot \mathbf{O}_5^3 + \mathbf{O}_3^2 \cdot \underline{\alpha} + \Gamma_g \cdot \underline{\alpha} \\ &= \mathbf{O}_6^3 + \Gamma_b \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot \underline{\beta} + \mathbf{O}_3^2 \cdot \underline{\alpha} + \Gamma_b \cdot \underline{\beta} + \Gamma_g \cdot \underline{\alpha},\end{aligned}$$

which implies

$$h[\underline{\beta}_3]^{[1]} = \mathbf{O}_6^3 + \Gamma_b^{(1)} \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot \underline{\beta}^{(1)} + \mathbf{O}_3^2 \cdot \underline{\alpha}^{(1)} + \Gamma_b^{(1)} \cdot \underline{\beta} + \Gamma_b \cdot \underline{\beta}^{(1)} + \Gamma_g^{(1)} \cdot \underline{\alpha} + \Gamma_g \cdot \underline{\alpha}^{(1)}.$$

Applying Lemma A.4, we obtain

$$\begin{aligned}\nabla_3 \underline{\beta} + 2 \operatorname{tr} \underline{\chi} \underline{\beta} &= -\operatorname{div} \underline{\alpha} + \widetilde{h[\underline{\beta}_3]} + \underline{\check{J}} \cdot \underline{\beta}_K + \underline{\check{L}} \cdot \underline{\alpha}_K + \Gamma_g \cdot \underline{\mathfrak{d}}^{\leq 1} \underline{\beta}_K \\ &= -\operatorname{div} \underline{\alpha} + \widetilde{h[\underline{\beta}_3]} + \Gamma_b^{(1)} \cdot \mathbf{O}_4^2 + \Gamma_g^{(1)} \cdot \mathbf{O}_5^3 + \Gamma_g \cdot \mathbf{O}_4^2 \\ &= -\operatorname{div} \underline{\alpha} + \widetilde{h[\underline{\beta}_3]} + \Gamma_b^{(1)} \cdot \mathbf{O}_4^2,\end{aligned}$$

and similarly

$$\nabla_3 \underline{\beta} + 2 \operatorname{tr} \underline{\chi} \underline{\beta} = -\operatorname{div} \underline{\beta} \alpha + \widetilde{h[\underline{\beta}_3]^{[1]}} + \Gamma_b^{(1)} \cdot \mathbf{O}_4^2.$$

Hence, we deduce

$$\nabla_3 \underline{\beta} + 2 \operatorname{tr} \underline{\chi} \underline{\beta} = -\operatorname{div} \underline{\alpha} + \Gamma_b^{(1)} \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot \underline{\beta} + \mathbf{O}_3^2 \cdot \underline{\alpha} + \Gamma_b \cdot \underline{\beta} + \Gamma_g \cdot \underline{\alpha}.$$

Similarly, we have

$$\begin{aligned}\nabla_3 \underline{\beta} + 2 \operatorname{tr} \underline{\chi} \underline{\beta} &= -\operatorname{div} \underline{\beta} \alpha + \Gamma_b^{(1)} \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot \underline{\beta}^{(1)} + \mathbf{O}_3^2 \cdot \underline{\alpha}^{(1)} \\ &\quad + \Gamma_b^{(1)} \cdot \underline{\beta} + \Gamma_b \cdot \underline{\beta}^{(1)} + \Gamma_g^{(1)} \cdot \underline{\alpha} + \Gamma_g \cdot \underline{\alpha}^{(1)}.\end{aligned}$$

Next, we recall from Proposition 2.19

$$\begin{aligned}\nabla_4 \underline{\beta} + \operatorname{tr} \underline{\chi} \underline{\beta} &= -\nabla \rho + {}^* \nabla \sigma + h[\underline{\beta}_4], \\ h[\underline{\beta}_4] &:= 2\underline{\omega} \underline{\beta} + 2\underline{\widehat{\chi}} \cdot \underline{\beta} - 3(\underline{\eta} \rho - {}^* \underline{\eta} \sigma).\end{aligned}$$

Applying Lemma A.3, we infer

$$\nabla_4 \underline{\partial} \underline{\beta} + \text{tr } \chi \underline{\partial} \underline{\beta} = -\nabla \underline{\partial} \rho + {}^* \nabla \underline{\partial} \sigma + h[\underline{\beta}_4]^{[1]},$$

where

$$\begin{aligned} h[\underline{\beta}_4]^{[1]} &= \underline{\partial} h[\underline{\beta}_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\underline{\beta}_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot (\underline{\beta}^{(1)}, \rho^{(1)}, \sigma^{(1)}) + \Gamma_g^{(1)} \cdot \underline{\beta}^{(1)} \\ &= \underline{\partial} h[\underline{\beta}_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\underline{\beta}_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot (\check{\underline{\beta}}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}) + \Gamma_g^{(1)} \cdot \check{\underline{\beta}}^{(1)} \\ &\quad + \Gamma_g \cdot \mathbf{O}_4^2 + \Gamma_g \cdot \mathbf{O}_3^1 + \mathbf{O}_6^3 + \Gamma_g^{(1)} \cdot \mathbf{O}_4^2. \end{aligned}$$

We compute

$$\begin{aligned} h[\underline{\beta}_4] &= (\mathbf{O}_2^1 + \Gamma_g) \cdot (\mathbf{O}_4^2 + \check{\underline{\beta}}) + (\mathbf{O}_3^2 + \Gamma_b) \cdot (\mathbf{O}_4^2 + \check{\underline{\beta}}) + (\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_3^1 + \check{\rho} + \check{\sigma}) \\ &= \mathbf{O}_6^3 + \Gamma_g \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot \check{\underline{\beta}} + \Gamma_g \cdot \check{\underline{\beta}} + \mathbf{O}_7^4 + \Gamma_b \cdot \mathbf{O}_4^2 + \mathbf{O}_3^2 \cdot \check{\underline{\beta}} + \Gamma_b \cdot \check{\underline{\beta}} \\ &\quad + \mathbf{O}_6^3 + \Gamma_g \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\rho}, \check{\sigma}) \\ &= \mathbf{O}_6^3 + \Gamma_g \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\underline{\beta}} + \mathbf{O}_3^2 \cdot (\check{\rho}, \check{\sigma}, \check{\underline{\beta}}) + \Gamma_g \cdot (\check{\underline{\beta}}, \check{\rho}, \check{\sigma}) + \Gamma_b \cdot \check{\underline{\beta}}, \end{aligned}$$

which implies

$$\begin{aligned} h[\underline{\beta}_4]^{[1]} &= \mathbf{O}_6^3 + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\underline{\beta}}^{(1)} + \mathbf{O}_3^2 \cdot (\check{\rho}^{(1)}, \check{\sigma}^{(1)}, \check{\underline{\beta}}^{(1)}) \\ &\quad + \Gamma_g^{(1)} \cdot (\check{\underline{\beta}}, \check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\underline{\beta}}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}) + \Gamma_b^{(1)} \cdot \check{\underline{\beta}} + \Gamma_b \cdot \check{\underline{\beta}}^{(1)}. \end{aligned}$$

Applying Lemma A.4, we obtain

$$\begin{aligned} \nabla_4 \check{\underline{\beta}} + \text{tr } \chi \check{\underline{\beta}} &= -\nabla \check{\rho} + {}^* \nabla \check{\sigma} + \widetilde{h[\underline{\beta}_4]} + \check{J} \cdot \underline{\beta}_K + \check{L} \cdot (\rho_K, \sigma_K) + \Gamma_g \cdot \mathfrak{d}^{\leq 1} \underline{\beta}_K \\ &= -\nabla \check{\rho} + {}^* \nabla \check{\sigma} + \widetilde{h[\underline{\beta}_4]} + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1. \end{aligned}$$

Hence, we infer

$$\nabla_4 \check{\underline{\beta}} + \text{tr } \chi \check{\underline{\beta}} = -\nabla \check{\rho} + {}^* \nabla \check{\sigma} + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\underline{\beta}} + \mathbf{O}_3^2 \cdot (\check{\rho}, \check{\sigma}, \check{\underline{\beta}}) + \Gamma_g \cdot (\check{\underline{\beta}}, \check{\rho}, \check{\sigma}) + \Gamma_b \cdot \check{\underline{\beta}}.$$

Similarly, we have

$$\begin{aligned} \nabla_4 \check{\underline{\partial}} \check{\underline{\beta}} + \text{tr } \chi \check{\underline{\partial}} \check{\underline{\beta}} &= -\nabla \check{\underline{\partial}} \rho + {}^* \nabla \check{\underline{\partial}} \sigma + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\underline{\beta}}^{(1)} + \mathbf{O}_3^2 \cdot (\check{\rho}^{(1)}, \check{\sigma}^{(1)}, \check{\underline{\beta}}^{(1)}) \\ &\quad + \Gamma_g^{(1)} \cdot (\check{\underline{\beta}}, \check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\underline{\beta}}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}) + \Gamma_b^{(1)} \cdot \check{\underline{\beta}} + \Gamma_b \cdot \check{\underline{\beta}}^{(1)}. \end{aligned}$$

Next, we recall from Proposition 2.19

$$\begin{aligned} \nabla_3(\rho, \sigma) + \frac{3}{2} \text{tr } \underline{\chi}(\rho, \sigma) &= -d_1 \underline{\beta} + h[\rho_3, \sigma_3], \\ h[\rho_3, \sigma_3] &= -\frac{1}{2} \widehat{\chi} \cdot (\underline{\alpha}, -{}^* \underline{\alpha}) + \zeta \cdot (\underline{\beta}, -{}^* \underline{\beta}) - 2\eta \cdot (\underline{\beta}, -{}^* \underline{\beta}). \end{aligned}$$

Applying Lemma A.3, we obtain

$$\nabla_3 \underline{\partial}(\rho, \sigma) + \frac{3}{2} \text{tr } \underline{\chi} \underline{\partial}(\rho, \sigma) = -d_1 \underline{\partial} \underline{\beta} + h[\rho_3, \sigma_3]^{[1]},$$

where

$$\begin{aligned} h[\rho_3, \sigma_3]^{[1]} &= \underline{\partial} h[\rho_3, \sigma_3] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\rho_3, \sigma_3] + (\Gamma_b + \mathbf{O}_3^2) \cdot (\rho^{(1)}, \sigma^{(1)}) + (\Gamma_g + \mathbf{O}_3^2) \cdot \underline{\beta}^{(1)} + \Gamma_b^{(1)} \cdot (\rho, \sigma) \\ &= \underline{\partial} h[\rho_3, \sigma_3] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\rho_3, \sigma_3] + (\Gamma_b + \mathbf{O}_3^2) \cdot (\check{\rho}^{(1)}, \check{\sigma}^{(1)}) + (\Gamma_g + \mathbf{O}_3^2) \cdot \check{\underline{\beta}}^{(1)} + \Gamma_b^{(1)} \cdot (\check{\rho}, \check{\sigma}) \\ &\quad + \Gamma_b \cdot \mathbf{O}_3^1 + \mathbf{O}_6^3 + \Gamma_g \cdot \mathbf{O}_4^2 + \mathbf{O}_7^4 + \Gamma_b^{(1)} \cdot \mathbf{O}_3^1. \end{aligned}$$

We compute

$$\begin{aligned}
h[\rho_3, \sigma_3] &= (\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_5^3 + \check{\alpha}) + (\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_4^2 + \check{\beta}) \\
&= \mathbf{O}_8^5 + \Gamma_g \cdot \mathbf{O}_5^3 + \mathbf{O}_3^2 \cdot \check{\alpha} + \Gamma_g \cdot \check{\alpha} + \mathbf{O}_7^4 + \Gamma_g \cdot \mathbf{O}_4^2 + \mathbf{O}_3^2 \cdot \check{\beta} + \Gamma_g \cdot \check{\beta} \\
&= \mathbf{O}_7^4 + \Gamma_g \cdot \mathbf{O}_4^2 + \mathbf{O}_3^2 \cdot (\check{\alpha}, \check{\beta}) + \Gamma_g \cdot (\check{\alpha}, \check{\beta}),
\end{aligned}$$

which implies

$$\begin{aligned}
h[\rho_3, \sigma_3]^{[1]} &= \mathbf{O}_6^3 + \Gamma_b^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\alpha}^{(1)}, \check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}) \\
&\quad + \Gamma_g^{(1)} \cdot (\check{\alpha}, \check{\beta}) + \Gamma_g \cdot (\check{\alpha}^{(1)}, \check{\beta}^{(1)}) + \Gamma_b \cdot (\check{\rho}^{(1)}, \check{\sigma}^{(1)}) + \Gamma_b^{(1)} \cdot (\check{\rho}, \check{\sigma}).
\end{aligned}$$

Applying Lemma A.4, we deduce

$$\begin{aligned}
\nabla_3(\check{\rho}, \check{\sigma}) + \frac{3}{2} \operatorname{tr} \chi(\check{\rho}, \check{\sigma}) &= -d_1 \check{\beta} + h[\rho_3, \sigma_3] + \check{L} \cdot \beta_K + \Gamma_g \cdot (\rho_K, \sigma_K) \\
&= -d_1 \check{\beta} + h[\rho_3, \sigma_3] + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 \\
&= -d_1 \check{\beta} + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\alpha}, \check{\beta}) + \Gamma_g \cdot (\check{\alpha}, \check{\beta}).
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
\nabla_3(\check{\rho}, \check{\sigma}) + \frac{3}{2} \operatorname{tr} \chi(\check{\rho}, \check{\sigma}) &= -d_1 \check{\beta} + \Gamma_b^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\alpha}^{(1)}, \check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}) \\
&\quad + \Gamma_g^{(1)} \cdot (\check{\alpha}, \check{\beta}) + \Gamma_g \cdot (\check{\alpha}^{(1)}, \check{\beta}^{(1)}) + \Gamma_b \cdot (\check{\rho}^{(1)}, \check{\sigma}^{(1)}) + \Gamma_b^{(1)} \cdot (\check{\rho}, \check{\sigma}).
\end{aligned}$$

We have from Proposition 2.19

$$\begin{aligned}
\nabla_4(\rho, -\sigma) + \frac{3}{2} \operatorname{tr} \chi(\rho, -\sigma) &= d_1 \beta + h[\rho_4, \sigma_4], \\
h[\rho_4, \sigma_4] &= -\frac{1}{2} \widehat{\chi} \cdot (\alpha, * \alpha) + (\zeta + 2\eta) \cdot (\beta, * \beta).
\end{aligned}$$

Applying Lemma A.3, we have

$$\nabla_4 \check{\rho}(\rho, -\sigma) + \frac{3}{2} \operatorname{tr} \chi \check{\rho}(\rho, -\sigma) = d_1 \check{\rho} \beta + h[\rho_4, \sigma_4]^{[1]},$$

where

$$\begin{aligned}
h[\rho_4, \sigma_4]^{[1]} &= \check{\rho} h[\rho_4, \sigma_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\rho_4, \sigma_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot (\beta^{(1)}, \rho^{(1)}, \sigma^{(1)}) + \Gamma_g^{(1)} \cdot (\rho, \sigma) \\
&= \check{\rho} h[\rho_4, \sigma_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\rho_4, \sigma_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot (\check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}) + \Gamma_g^{(1)} \cdot (\check{\rho}, \check{\sigma}) \\
&\quad + \Gamma_g \cdot \mathbf{O}_3^1 + \mathbf{O}_6^3 + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1.
\end{aligned}$$

We compute

$$\begin{aligned}
h[\rho_4, \sigma_4] &= (\mathbf{O}_3^2 + \Gamma_b) \cdot (\mathbf{O}_5^3 + \check{\alpha}) + (\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_4^2 + \check{\beta}) \\
&= \mathbf{O}_8^5 + \Gamma_b \cdot \mathbf{O}_5^3 + \mathbf{O}_3^2 \cdot \check{\alpha} + \Gamma_b \cdot \check{\alpha} + \mathbf{O}_7^4 + \Gamma_g \cdot \mathbf{O}_4^2 + \mathbf{O}_3^2 \cdot \check{\beta} + \Gamma_g \cdot \check{\beta} \\
&= \mathbf{O}_7^4 + \Gamma_g \cdot \mathbf{O}_4^2 + \mathbf{O}_3^2 \cdot (\check{\alpha}, \check{\beta}) + \Gamma_b \cdot \check{\alpha} + \Gamma_g \cdot \check{\beta},
\end{aligned}$$

which implies

$$\begin{aligned}
h[\rho_4, \sigma_4]^{[1]} &= \mathbf{O}_6^3 + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\alpha}^{(1)}, \check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}) \\
&\quad + \Gamma_b^{(1)} \cdot \check{\alpha} + \Gamma_b \cdot \check{\alpha}^{(1)} + \Gamma_g^{(1)} \cdot (\check{\beta}, \check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}).
\end{aligned}$$

Applying Lemma A.4, we deduce

$$\begin{aligned}\nabla_4(\check{\rho}, -\check{\sigma}) + \frac{3}{2} \operatorname{tr} \chi(\check{\rho}, -\check{\sigma}) &= d_1 \check{\beta} + h[\check{\rho}_4, \check{\sigma}_4] + \check{J} \cdot (\rho_K, \sigma_K) + \check{L} \cdot \beta_K + \Gamma_g \cdot \mathfrak{d}^{\leq 1}(\rho_K, \sigma_K) \\ &= d_1 \check{\beta} + h[\check{\rho}_4, \check{\sigma}_4] + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 \\ &= d_1 \check{\beta} + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\alpha}, \check{\beta}) + \Gamma_b \cdot \check{\alpha} + \Gamma_g \cdot \check{\beta}.\end{aligned}$$

Similarly, we have

$$\begin{aligned}\nabla_4(\check{\rho}, -\check{\sigma}) + \frac{3}{2} \operatorname{tr} \chi(\check{\rho}, -\check{\sigma}) &= d_1 \check{\beta} + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\alpha}^{(1)}, \check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}) \\ &\quad + \Gamma_b^{(1)} \cdot \check{\alpha} + \Gamma_b \cdot \check{\alpha}^{(1)} + \Gamma_g^{(1)} \cdot (\check{\beta}, \check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}).\end{aligned}$$

Next, we recall from Proposition 2.19

$$\begin{aligned}\nabla_3 \beta + \operatorname{tr} \chi \beta &= \nabla \rho + {}^* \nabla \sigma + h[\beta_3], \\ h[\beta_3] &= 2\omega \beta + 2\hat{\chi} \cdot \underline{\beta} + 3(\eta \rho + {}^* \eta \sigma).\end{aligned}$$

Applying Lemma A.3, we deduce

$$\nabla_3 \check{\rho} \beta + \operatorname{tr} \chi \check{\rho} \beta = \nabla \check{\rho} \rho + {}^* \nabla \check{\rho} \sigma + h[\beta_3]^{[1]},$$

where

$$\begin{aligned}h[\beta_3]^{[1]} &= \check{\rho} h[\beta_3] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\beta_3] + (\Gamma_b + \mathbf{O}_3^2) \cdot \beta^{(1)} + (\Gamma_g + \mathbf{O}_3^2) \cdot (\rho^{(1)}, \sigma^{(1)}) + \Gamma_g^{(1)} \cdot \beta \\ &= \check{\rho} h[\beta_3] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\beta_3] + (\Gamma_b + \mathbf{O}_3^2) \cdot \check{\beta}^{(1)} + (\Gamma_g + \mathbf{O}_3^2) \cdot (\check{\rho}^{(1)}, \check{\sigma}^{(1)}) + \Gamma_g^{(1)} \cdot \check{\beta} \\ &\quad + \Gamma_b \cdot \mathbf{O}_4^2 + \mathbf{O}_7^4 + \Gamma_g \cdot \mathbf{O}_3^1 + \mathbf{O}_6^3 + \Gamma_g^{(1)} \cdot \mathbf{O}_4^2.\end{aligned}$$

We compute

$$\begin{aligned}h[\beta_3] &= (\mathbf{O}_2^1 + \Gamma_b) \cdot (\mathbf{O}_4^2 + \check{\beta}) + (\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_4^2 + \check{\beta}) + (\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_3^1 + \check{\rho} + \check{\sigma}) \\ &= \mathbf{O}_6^3 + \Gamma_b \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot \check{\beta} + \Gamma_b \cdot \check{\beta} + \mathbf{O}_7^4 + \Gamma_g \cdot \mathbf{O}_4^2 + \mathbf{O}_3^2 \cdot \check{\beta} + \Gamma_g \cdot \check{\beta} \\ &\quad + \mathbf{O}_6^3 + \Gamma_g \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\rho}, \check{\sigma}) \\ &= \mathbf{O}_6^3 + \Gamma_g \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\beta} + \mathbf{O}_3^2 \cdot (\check{\beta}, \check{\rho}, \check{\sigma}) + \Gamma_b \cdot \check{\beta} + \Gamma_g \cdot (\check{\beta}, \check{\rho}, \check{\sigma}),\end{aligned}$$

which implies

$$\begin{aligned}h[\beta_3]^{[1]} &= \mathbf{O}_6^3 + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\beta}^{(1)} + \mathbf{O}_3^2 \cdot (\check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}, \check{\beta}^{(1)}) \\ &\quad + \Gamma_b^{(1)} \cdot \check{\beta} + \Gamma_b \cdot \check{\beta}^{(1)} + \Gamma_g^{(1)} \cdot (\check{\beta}, \check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}).\end{aligned}$$

Applying Lemma A.4, we infer

$$\begin{aligned}\nabla_3 \check{\beta} + \operatorname{tr} \chi \check{\beta} &= \nabla \check{\rho} + {}^* \nabla \check{\sigma} + h[\check{\beta}_3] + \check{L} \cdot (\rho_K, \sigma_K) + \check{J} \cdot \beta_K + \Gamma_g \cdot \mathfrak{d}^{\leq 1} \beta_K \\ &= \nabla \check{\rho} + {}^* \nabla \check{\sigma} + h[\check{\beta}_3] + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1.\end{aligned}$$

Hence, we obtain

$$\nabla_3 \check{\beta} + \operatorname{tr} \chi \check{\beta} = \nabla \check{\rho} + {}^* \nabla \check{\sigma} + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\beta} + \mathbf{O}_3^2 \cdot (\check{\beta}, \check{\rho}, \check{\sigma}) + \Gamma_b \cdot \check{\beta} + \Gamma_g \cdot (\check{\beta}, \check{\rho}, \check{\sigma}).$$

Similarly, we have

$$\begin{aligned}\nabla_3 \widetilde{\boldsymbol{\vartheta}} \beta + \operatorname{tr} \underline{\chi} \widetilde{\boldsymbol{\vartheta}} \beta &= \nabla \widetilde{\boldsymbol{\vartheta}} \rho + {}^* \nabla \widetilde{\boldsymbol{\vartheta}} \sigma + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\beta}^{(1)} + \mathbf{O}_3^2 \cdot (\check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}) \\ &\quad + \Gamma_b^{(1)} \cdot \check{\beta} + \Gamma_b \cdot \check{\beta}^{(1)} + \Gamma_g^{(1)} \cdot (\check{\beta}, \check{\rho}, \check{\sigma}) + \Gamma_g \cdot (\check{\beta}^{(1)}, \check{\rho}^{(1)}, \check{\sigma}^{(1)}).\end{aligned}$$

Next, we recall from Proposition 2.19

$$\begin{aligned}\nabla_4 \beta + 2 \operatorname{tr} \chi \beta &= \operatorname{div} \alpha + h[\beta_4] \\ h[\beta_4] &= -2\omega\beta + (2\zeta + \underline{\eta})\alpha.\end{aligned}$$

Applying Lemma A.3, we deduce

$$\nabla_4 \boldsymbol{\vartheta} \beta + 2 \operatorname{tr} \chi \boldsymbol{\vartheta} \beta = \operatorname{div} \boldsymbol{\vartheta} \alpha + h[\beta_4]^{[1]},$$

where

$$\begin{aligned}h[\beta_4]^{[1]} &= \boldsymbol{\vartheta} h[\beta_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\beta_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot (\beta^{(1)}, \alpha^{(1)}) + \Gamma_g^{(1)} \cdot \beta \\ &= \boldsymbol{\vartheta} h[\beta_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\beta_4] + (\Gamma_g + \mathbf{O}_3^2) \cdot (\check{\beta}^{(1)}, \check{\alpha}^{(1)}) + \Gamma_g^{(1)} \cdot \check{\beta} \\ &\quad + \Gamma_g \cdot \mathbf{O}_4^2 + \mathbf{O}_7^4 + \Gamma_g^{(1)} \cdot \mathbf{O}_4^2.\end{aligned}$$

We compute

$$\begin{aligned}h[\beta_4] &= (\mathbf{O}_2^1 + \Gamma_g) \cdot (\mathbf{O}_4^2 + \check{\beta}) + (\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_5^3 + \check{\alpha}) \\ &= \mathbf{O}_6^3 + \Gamma_g \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot \check{\beta} + \Gamma_g \cdot \check{\beta} + \mathbf{O}_8^5 + \Gamma_g \cdot \mathbf{O}_5^3 + \mathbf{O}_3^2 \cdot \check{\alpha} + \Gamma_g \cdot \check{\alpha} \\ &= \mathbf{O}_6^3 + \Gamma_g \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot (\check{\beta}, \check{\alpha}) + \Gamma_g \cdot (\check{\beta}, \check{\alpha}),\end{aligned}$$

which implies

$$h[\beta_4]^{[1]} = \mathbf{O}_6^3 + \Gamma_g^{(1)} \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot (\check{\beta}^{(1)}, \check{\alpha}^{(1)}) + \Gamma_g \cdot (\check{\beta}^{(1)}, \check{\alpha}^{(1)}) + \Gamma_g^{(1)} \cdot (\check{\beta}, \check{\alpha}).$$

Applying Lemma A.4, we obtain

$$\begin{aligned}\nabla_4 \check{\beta} + 2 \operatorname{tr} \chi \check{\beta} &= \operatorname{div} \check{\alpha} + \widetilde{h[\beta_4]} + \check{J} \cdot \beta_K + \check{L} \cdot \alpha_K + \Gamma_g \cdot \beta_K \\ &= \operatorname{div} \check{\alpha} + \widetilde{h[\beta_4]} + \Gamma_g^{(1)} \cdot \mathbf{O}_4^2 \\ &= \operatorname{div} \check{\alpha} + \Gamma_g^{(1)} \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot (\check{\beta}, \check{\alpha}) + \Gamma_g \cdot (\check{\beta}, \check{\alpha}).\end{aligned}$$

Similarly, we have

$$\nabla_4 \widetilde{\boldsymbol{\vartheta}} \beta + 2 \operatorname{tr} \chi \widetilde{\boldsymbol{\vartheta}} \beta = \operatorname{div} \widetilde{\boldsymbol{\vartheta}} \alpha + \Gamma_g^{(1)} \cdot \mathbf{O}_4^2 + \mathbf{O}_2^1 \cdot (\check{\beta}^{(1)}, \check{\alpha}^{(1)}) + \Gamma_g^{(1)} \cdot (\check{\beta}, \check{\alpha}) + \Gamma_g \cdot (\check{\beta}^{(1)}, \check{\alpha}^{(1)}).$$

Finally, we recall from Proposition 2.19

$$\begin{aligned}\nabla_3 \alpha + \frac{1}{2} \operatorname{tr} \underline{\chi} \alpha &= \nabla \widehat{\otimes} \beta + h[\alpha_3], \\ h[\alpha_3] &= 4\underline{\omega}\alpha - 3(\widehat{\chi}\rho + {}^* \widehat{\chi}\sigma) + (\zeta + 4\underline{\eta}) \widehat{\otimes} \beta.\end{aligned}$$

Applying Lemma A.3, we obtain

$$\nabla_3 \boldsymbol{\vartheta} \alpha + \frac{1}{2} \operatorname{tr} \underline{\chi} \boldsymbol{\vartheta} \alpha = \nabla \widehat{\otimes} \boldsymbol{\vartheta} \beta + h[\alpha_3]^{[1]},$$

where

$$\begin{aligned}
h[\alpha_3]^{[1]} &= \not\partial h[\alpha_3] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\alpha_3] + (\Gamma_b + \mathbf{O}_3^2) \cdot \alpha^{(1)} + (\Gamma_g + \mathbf{O}_3^2) \cdot \beta^{(1)} + \Gamma_b^{(1)} \cdot \alpha \\
&= \not\partial h[\alpha_3] + (\Gamma_g + \mathbf{O}_3^2) \cdot h[\alpha_3] + (\Gamma_b + \mathbf{O}_3^2) \cdot \check{\alpha}^{(1)} + (\Gamma_g + \mathbf{O}_3^2) \cdot \check{\beta}^{(1)} + \Gamma_b^{(1)} \cdot \check{\alpha} \\
&\quad + \Gamma_b \cdot \mathbf{O}_5^3 + \mathbf{O}_8^5 + \Gamma_g \cdot \mathbf{O}_4^2 + \mathbf{O}_7^4 + \Gamma_b^{(1)} \cdot \mathbf{O}_5^3.
\end{aligned}$$

We compute

$$\begin{aligned}
h[\alpha_3] &= (\mathbf{O}_2^1 + \Gamma_b) \cdot (\mathbf{O}_5^3 + \check{\alpha}) + (\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_3^1 + \check{\rho} + \check{\sigma}) + (\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_4^2 + \check{\beta}) \\
&= \mathbf{O}_7^4 + \Gamma_b \cdot \mathbf{O}_5^3 + \mathbf{O}_2^1 \cdot \check{\alpha} + \Gamma_b \cdot \check{\alpha} + \mathbf{O}_6^3 + \Gamma_g \cdot \mathbf{O}_3^1 + \mathbf{O}_3^2 \cdot (\check{\rho}, \check{\sigma}) \\
&\quad + \mathbf{O}_7^4 + \Gamma_g \cdot \mathbf{O}_4^2 + \mathbf{O}_3^2 \cdot \check{\beta} + \Gamma_g \cdot \check{\beta} \\
&= \mathbf{O}_6^3 + \Gamma_g \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\alpha} + \mathbf{O}_3^2 \cdot (\check{\rho}, \check{\sigma}, \check{\beta}) + \Gamma_b \cdot \check{\alpha} + \Gamma_g \cdot \check{\beta},
\end{aligned}$$

which implies

$$\begin{aligned}
h[\alpha_3]^{[1]} &= \mathbf{O}_6^3 + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\alpha}^{(1)} + \mathbf{O}_3^2 \cdot (\check{\rho}^{(1)}, \check{\sigma}^{(1)}, \check{\beta}^{(1)}) \\
&\quad + \Gamma_b^{(1)} \cdot \check{\alpha} + \Gamma_b \cdot \check{\alpha}^{(1)} + \Gamma_g^{(1)} \cdot \check{\beta} + \Gamma_g \cdot \check{\beta}^{(1)}.
\end{aligned}$$

Applying Lemma A.4, we infer

$$\begin{aligned}
\nabla_3 \check{\alpha} + \frac{1}{2} \text{tr } \underline{\chi} \check{\alpha} &= \nabla \widehat{\otimes} \check{\beta} + \widetilde{h[\alpha_3]} + \check{J} \cdot \alpha_K + \check{L} \cdot \beta_K + \Gamma_g \cdot \alpha_K \\
&= \nabla \widehat{\otimes} \check{\beta} + \widetilde{h[\alpha_3]} + \Gamma_b^{(1)} \cdot \mathbf{O}_5^3 + \Gamma_g^{(1)} \cdot \mathbf{O}_4^2 + \Gamma_g \cdot \mathbf{O}_5^3 \\
&= \nabla \widehat{\otimes} \check{\beta} + \widetilde{h[\alpha_3]} + \Gamma_g^{(1)} \cdot \mathbf{O}_4^2 \\
&= \nabla \widehat{\otimes} \check{\beta} + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\alpha} + \mathbf{O}_3^2 \cdot (\check{\rho}, \check{\sigma}, \check{\beta}) + \Gamma_b \cdot \check{\alpha} + \Gamma_g \cdot \check{\beta}.
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
\nabla_3 \check{\beta} + \frac{1}{2} \text{tr } \underline{\chi} \check{\beta} &= \nabla \widehat{\otimes} \check{\alpha} + \Gamma_g^{(1)} \cdot \mathbf{O}_3^1 + \mathbf{O}_2^1 \cdot \check{\alpha}^{(1)} + \mathbf{O}_3^2 \cdot (\check{\rho}^{(1)}, \check{\sigma}^{(1)}, \check{\beta}^{(1)}) \\
&\quad + \Gamma_b^{(1)} \cdot \check{\alpha} + \Gamma_b \cdot \check{\alpha}^{(1)} + \Gamma_g^{(1)} \cdot \check{\beta} + \Gamma_g \cdot \check{\beta}^{(1)}.
\end{aligned}$$

This concludes the proof of Proposition A.5. □

#### A.4 Proof of Proposition 2.40

**Proposition A.6.** *We have the following equations:*

$$\begin{aligned}
\Omega \nabla_4(\check{\gamma}_{AB}) &= \Omega \text{tr } \underline{\chi} \check{\gamma}_{AB} + \widetilde{\Omega \text{tr } \underline{\chi}(\gamma_K)_{AB}} + 2\Omega \check{\chi}_{AB} + 2\check{\Omega}(\check{\chi}_K)_{AB}, \\
\Omega \nabla_4(\check{\Omega}) &= -2\Omega(\check{\Omega}\omega) - 2\check{\Omega}(\Omega_K \omega_K), \\
\Omega \nabla_4(\check{b}^A) &= -4 \left( \check{\Omega}^2 \zeta_B \gamma^{AB} + \Omega_K^2 \check{\zeta}_B \gamma^{AB} + \Omega_K^2 (\zeta_K)_B \gamma^{AB} \right), \\
\Omega \nabla_4(\check{\epsilon}_{AB}) &= \Omega \text{tr } \underline{\chi} \check{\epsilon}_{AB} + \widetilde{\Omega \text{tr } \underline{\chi}(\epsilon_K)_{AB}}.
\end{aligned}$$

In the case  $\check{\Omega} = 0$  and  $\check{b} = 0$ , we have the following equations:

$$\begin{aligned}
\Omega \nabla_3(\check{\gamma}_{AB}) - \Omega \text{tr } \underline{\chi} \check{\gamma}_{AB} &= \widetilde{\Omega \text{tr } \underline{\chi}(\gamma_K)_{AB}} + 2\Omega \check{\chi}_{AB} - \partial_A(b_K^C) \check{\gamma}_{BC} - \partial_B(b_K^C) \check{\gamma}_{AC}, \\
\Omega \nabla_3(\check{\epsilon}_{AB}) - \Omega \text{tr } \underline{\chi} \check{\epsilon}_{AB} &= \widetilde{\Omega \text{tr } \underline{\chi}(\epsilon_K)_{AB}} - \partial_A(b_K^C) \check{\epsilon}_{BC} - \partial_B(b_K^C) \check{\epsilon}_{AC}.
\end{aligned}$$

**Remark A.7.** Note that Proposition 2.40 follows directly from Propositions 2.28 and A.6 and Definitions 2.3 and 2.25.

*Proof of Proposition A.6.* Firstly, recall from Proposition 2.20 that

$$\Omega \nabla_4(\gamma_{AB}) = 2\Omega \chi_{AB},$$

which implies

$$\begin{aligned} \Omega \nabla_4(\check{\gamma}_{AB}) &= \Omega \nabla_4(\gamma_{AB} - (\gamma_K)_{AB}) \\ &= \Omega \nabla_4(\gamma_{AB}) - \Omega \nabla_{(e_4)_K}(\gamma_K)_{AB} \\ &= 2\Omega \chi_{AB} - 2\Omega_K(\chi_K)_{AB} \\ &= \Omega \operatorname{tr} \chi \check{\gamma}_{AB} + \widetilde{\Omega \operatorname{tr} \chi}(\gamma_K)_{AB} + 2\Omega \widehat{\chi}_{AB} - 2\Omega_K(\widehat{\chi}_K)_{AB} \\ &= \Omega \operatorname{tr} \chi \check{\gamma}_{AB} + \widetilde{\Omega \operatorname{tr} \chi}(\gamma_K)_{AB} + 2\Omega \check{\chi}_{AB} + 2\check{\Omega}(\widehat{\chi}_K)_{AB}. \end{aligned}$$

Next, we compute

$$\begin{aligned} \Omega \nabla_4(\check{\Omega}) &= \Omega \nabla_4(\Omega - \Omega_K) \\ &= \Omega \nabla_4 \Omega - \Omega_K \nabla_{(e_4)_K} \Omega_K \\ &= -2\omega \Omega^2 + 2\omega_K \Omega_K^2 \\ &= -2\widetilde{\Omega \omega} \Omega - 2\check{\Omega}(\Omega_K \omega_K). \end{aligned}$$

Recall from Proposition 2.20 that

$$\nabla_4(\underline{b}^A) = -4\Omega \zeta^A = -4\Omega \zeta_B \gamma^{AB}.$$

Hence, we have

$$\begin{aligned} \Omega \nabla_4 \check{\underline{b}}^A &= \Omega \nabla_4(\underline{b}^A) - \Omega_K \nabla_{(e_4)_K}(\underline{b}_K^A) \\ &= -4\Omega^2 \zeta_B \gamma^{AB} + 4\Omega_K^2 (\zeta_K)_B (\gamma_K)^{AB} \\ &= -4 \left( \widetilde{\Omega^2 \zeta_B} \gamma^{AB} + \Omega_K^2 \check{\zeta}_B \gamma^{AB} + \Omega_K^2 (\zeta_K)_B \widetilde{\gamma^{AB}} \right). \end{aligned}$$

Next, we recall from Proposition 2.20

$$\Omega \nabla_4(\in_{AB}) = \Omega \operatorname{tr} \chi \in_{AB},$$

which implies

$$\begin{aligned} \Omega \nabla_4(\check{\in}_{AB}) &= \Omega \nabla_4(\in_{AB}) - \Omega_K \nabla_{(e_4)_K}(\in_K)_{AB} \\ &= \Omega \operatorname{tr} \chi \in_{AB} - \Omega_K \operatorname{tr}_K \chi_K (\in_K)_{AB} \\ &= \Omega \operatorname{tr} \chi \check{\in}_{AB} + \widetilde{\Omega \operatorname{tr} \chi}(\in_K)_{AB}. \end{aligned}$$

Then, we recall from Proposition 2.20

$$\Omega e_3(\gamma_{AB}) = 2\Omega \underline{\chi}_{AB} - \partial_A(\underline{b}^C) \gamma_{BC} - \partial_B(\underline{b}^C) \gamma_{AC},$$

which implies

$$\Omega_K(e_3)_K((\gamma_K)_{AB}) = 2\Omega_K(\underline{\chi}_K)_{AB} - \partial_A(\underline{b}_K^C)(\gamma_K)_{BC} - \partial_B(\underline{b}_K^C)(\gamma_K)_{AC}.$$

Thus, we infer

$$\begin{aligned}
\Omega e_3(\check{\gamma}_{AB}) &= \Omega e_3(\gamma_{AB}) - \Omega_K(e_3)_K(\gamma_K)_{AB} - \widetilde{\Omega e_3}(\gamma_K)_{AB} \\
&= 2\Omega \underline{\chi}_{AB} - \partial_A(\underline{b}^C)\gamma_{BC} - \partial_B(\underline{b}^C)\gamma_{AC} - 2\Omega_K(\underline{\chi}_K)_{AB} \\
&\quad + \partial_A(\underline{b}_K^C)(\gamma_K)_{BC} + \partial_B(\underline{b}_K^C)(\gamma_K)_{AC} - \check{\underline{b}}^C \partial_C(\gamma_K)_{AB} \\
&= \Omega \operatorname{tr} \underline{\chi} \gamma_{AB} - \Omega_K \operatorname{tr}_K \underline{\chi}_K (\gamma_K)_{AB} + 2(\Omega \check{\underline{\chi}}_{AB} - \Omega_K(\check{\underline{\chi}}_K)_{AB}) \\
&\quad - \partial_A(\underline{b}^C)\gamma_{BC} - \partial_B(\underline{b}^C)\gamma_{AC} + \partial_A(\underline{b}_K^C)(\gamma_K)_{BC} + \partial_B(\underline{b}_K^C)(\gamma_K)_{AC} - \check{\underline{b}}^C \partial_C(\gamma_K)_{AB} \\
&= \Omega \operatorname{tr} \underline{\chi} \check{\gamma}_{AB} + \widetilde{\Omega \operatorname{tr} \underline{\chi}}(\gamma_K)_{AB} + 2\Omega \check{\underline{\chi}}_{AB} + 2\check{\Omega}(\check{\underline{\chi}}_K)_{AB} \\
&\quad - \partial_A(\check{\underline{b}}^C)\gamma_{BC} - \partial_A(\underline{b}_K^C)\check{\gamma}_{BC} - \partial_B(\check{\underline{b}}^C)\gamma_{AC} - \partial_B(\underline{b}_K^C)\check{\gamma}_{AC} - \check{\underline{b}}^C \partial_C(\gamma_K)_{AB}.
\end{aligned}$$

In the particular case  $\check{\Omega} = 0$  and  $\check{\underline{b}} = 0$ , we deduce

$$\Omega e_3(\check{\gamma}_{AB}) - \Omega \operatorname{tr} \underline{\chi} \check{\gamma}_{AB} = \widetilde{\Omega \operatorname{tr} \underline{\chi}}(\gamma_K)_{AB} + 2\Omega \check{\underline{\chi}}_{AB} - \partial_A(\underline{b}_K^C)\check{\gamma}_{BC} - \partial_B(\underline{b}_K^C)\check{\gamma}_{AC}.$$

Similarly, we have from Proposition 2.20

$$\Omega \nabla_3(\in_{AB}) = \Omega \operatorname{tr} \underline{\chi} \in_{AB} - \partial_A(\underline{b}^C) \in_{BC} - \partial_B(\underline{b}^C) \in_{BC},$$

which yields

$$\Omega_K(e_3)_K(\in_K)_{AB} = \Omega_K \operatorname{tr}_K \underline{\chi}_K (\in_K)_{AB} - \partial_A(\underline{b}_K^C)(\in_K)_{BC} - \partial_B(\underline{b}_K^C)(\in_K)_{BC}.$$

Hence, we have

$$\begin{aligned}
\Omega \nabla_3(\check{\in}_{AB}) &= \Omega e_3(\in_{AB}) - \Omega_K(e_3)_K(\in_K)_{AB} - \widetilde{\Omega e_3}(\in_K)_{AB} \\
&= \Omega \operatorname{tr} \underline{\chi} \in_{AB} - \partial_A(\underline{b}^C) \in_{BC} - \partial_B(\underline{b}^C) \in_{BC} - \Omega_K \operatorname{tr}_K \underline{\chi}_K (\in_K)_{AB} \\
&\quad + \partial_A(\underline{b}_K^C)(\in_K)_{BC} + \partial_B(\underline{b}_K^C)(\in_K)_{BC} - \check{\underline{b}}^C \partial_C(\in_K)_{AB} \\
&= \Omega \operatorname{tr} \underline{\chi} \check{\in}_{AB} + \widetilde{\Omega \operatorname{tr} \underline{\chi}}(\in_K)_{AB} - \partial_A(\check{\underline{b}}^C) \in_{BC} - \partial_A(\underline{b}_K^C)\check{\in}_{BC} \\
&\quad - \partial_B(\check{\underline{b}}^C) \in_{AC} - \partial_B(\underline{b}_K^C)\check{\in}_{AC} - \check{\underline{b}}^C \partial_C(\gamma_K)_{AB},
\end{aligned}$$

which implies from  $\check{\Omega} = 0$  and  $\check{\underline{b}} = 0$

$$\Omega \nabla_3(\check{\in}_{AB}) = \Omega \operatorname{tr} \underline{\chi} \check{\in}_{AB} + \widetilde{\Omega \operatorname{tr} \underline{\chi}}(\in_K)_{AB} - \partial_A(\underline{b}_K^C)\check{\in}_{BC} - \partial_B(\underline{b}_K^C)\check{\in}_{AC}.$$

This concludes the proof of Proposition A.6.  $\square$

## B Proof of the case $s \in (3, 4)$

In Sections 4–6, we have provided the proof of Theorem 3.2 in the case  $s \in [4, 6]$ . In this appendix, we outline the proof in the case  $s \in (3, 4)$ , which is similar to the case of  $s \in [4, 6]$ .

### B.1 Fundamental norms

The definitions of some of the norms are different from Section 3.1. In the sequel, we only mention the norms that differ from the ones in Section 3.1.

We define the  $\mathcal{R}_0^S[\check{\beta}]$  and  $\mathcal{R}_0^S[\check{\rho}, \check{\sigma}]$  as follows:

$$\begin{aligned}\mathcal{R}_0^S[\check{\beta}] &:= \sup_{\mathcal{K}} \sup_{p \in [2,4]} |r^{\frac{s+3}{2} - \frac{2}{p}} \check{\beta}|_{p,S}, \\ \underline{\mathcal{R}}_0^S[\check{\rho}, \check{\sigma}] &:= \sup_{\mathcal{K}} \sup_{p \in [2,4]} |r^{\frac{s+2}{2} - \frac{2}{p}} |u|^{\frac{1}{2}} (\check{\rho}, \check{\sigma})|_{p,S}.\end{aligned}$$

The following flux of curvature are different from the case  $s \in [4, 6]$  for  $q = 0, 1$ :

$$\begin{aligned}\mathcal{R}_q[\check{\beta}](u, \underline{u}) &:= \|r^{\frac{s}{2}} (r\nabla)^q \check{\beta}\|_{2, C_u^V}, \\ \underline{\mathcal{R}}_q[(\check{\rho}, \check{\sigma})](u, \underline{u}) &:= \|r^{\frac{s}{2}} (r\nabla)^q (\check{\rho}, \check{\sigma})\|_{2, \underline{C}_u^V}.\end{aligned}$$

The other norms are the same as in the case  $s \in [4, 6]$ .

## B.2 Estimates for Ricci coefficients and curvature components

We discuss the curvature estimates of Section 4 in this case. Recall that we have four Bianchi pairs:  $(\check{\alpha}, \check{\beta})$ ,  $(\check{\beta}, (\check{\rho}, \check{\sigma}))$ ,  $((\check{\rho}, \check{\sigma}), \check{\beta})$  and  $(\check{\beta}, \check{\alpha})$ . In Section 4, we took respectively  $p = s, 4, 2, 0$ , to estimate the Bianchi pairs. In the case  $s \in (3, 4)$ , we take  $p = s, s, 2, 0$  to estimate  $\mathcal{R}$ . The method is then exactly the same as in Section 4.

Concerning the control of Ricci coefficients, we can proceed by the same method as in Section 5. Recall that Proposition 5.9 used the fact that there exists a constant  $\delta > 0$  such that

$$\sup_{p \in [2,4]} |r^{3+\delta-\frac{2}{p}} \check{\beta}|_{p,S} \lesssim \epsilon_0.$$

This still holds true since we have  $\frac{s+3}{2} > 3$  for  $s > 3$ . The other propositions still hold true since we only used  $s > 3$  in their proofs.

## B.3 Conclusion

The proof of Theorems M0, M2 and M4 remain exactly the same since all the arguments also apply to the case  $s > 3$ . Consequently, we have Theorem 3.2 in this case. Hence, we deduce that Theorem 3.2 holds true for  $s \in (3, 6]$ .

# C Proof of the case $s > 6$

In this appendix, we prove Theorem 3.2 in the case  $s > 6$ . Then, we compare the result to the peeling decay for curvature components obtained in [2, 15].

## C.1 Fundamental norms

The definitions of  $\mathcal{R}$ -norms for  $\check{\alpha}$  and  $\check{\beta}$  are different from Section 3.1.1. We denote

$$s_0 := \min \left\{ s, \frac{29}{4} \right\}. \quad (\text{C.1})$$

We define for  $q = 0, 1$ :

$$\begin{aligned}
\mathcal{R}_q[\check{\beta}] &:= |u|^{\frac{s-4}{2}} \|r^2 (r\nabla)^q \check{\beta}\|_{2, C_u^V}, \\
\underline{\mathcal{R}}_q[\check{\beta}] &:= \|r^3 |u|^{\frac{s-6}{2}} (r\nabla)^q \check{\beta}\|_{2, \underline{C}_u^V}, \\
\mathcal{R}_q[\check{\alpha}] &:= \|r^3 |u|^{\frac{s-6}{2}} (r\nabla)^q \check{\alpha}\|_{2, C_u^V}, \\
\underline{\mathcal{R}}_q[\check{\alpha}] &:= |u|^{\frac{s-s_0}{2}} \|r^{\frac{s_0}{2}} (r\nabla)^q \check{\alpha}\|_{2, \underline{C}_u^V}.
\end{aligned} \tag{C.2}$$

We also define:

$$\underline{\mathcal{R}}_0^S[\check{\beta}] := \sup_{\mathcal{K}} \sup_{p \in [2, 4]} |r^{4-\frac{2}{p}} |u|^{\frac{s-5}{2}} \check{\beta}|_{p, S}. \tag{C.3}$$

The norms  $\mathcal{R}_0^S[\check{\alpha}]$  are defined as follows:

$$\begin{aligned}
\underline{\mathcal{R}}_0^S[\check{\alpha}] &:= \sup_{\mathcal{K}} |r^{\frac{s+1}{2}} \check{\alpha}|_{2, S}, & s \in (6, 7), \\
\mathcal{R}_0^S[\check{\alpha}] &:= \sup_{\mathcal{K}} |r^4 (\log r)^{-\frac{1}{2}} \check{\alpha}|_{2, S}, & s = 7, \\
\mathcal{R}_0^S[\check{\alpha}] &:= \sup_{\mathcal{K}} |r^4 |u|^{\frac{s-7}{2}} \check{\alpha}|_{2, S}, & s > 7.
\end{aligned} \tag{C.4}$$

We also define:

$$\underline{\mathcal{R}}_0^S[\check{\alpha}] := \sup_{\mathcal{K}} \sup_{p \in [2, 4]} |r^{1-\frac{2}{p}} |u|^{\frac{s+1}{2}} \check{\alpha}|_{p, S}. \tag{C.5}$$

All the other norms are defined as in Section 3.1.1.

## C.2 Curvature estimates

In Section 4, we took respectively  $p = s, 4, 2, 0$ , to estimate the Bianchi pairs  $(\check{\alpha}, \check{\beta})$ ,  $(\check{\beta}, (\check{\rho}, -\check{\sigma}))$ ,  $((\check{\rho}, \check{\sigma}), \check{\beta})$  and  $(\check{\beta}, \check{\alpha})$ . In the case  $s > 6$ , we take respectively  $p = 6, 4, 2, 0$  for the Bianchi pairs. Proceeding as in Section 4, we deduce from Propositions 4.9–4.12 that

$$\sum_{q=0}^1 \left( \mathcal{R}_q[\check{\alpha}] + \mathcal{R}_q[\check{\beta}] + \underline{\mathcal{R}}_q[\check{\beta}] + \mathcal{R}_q[\check{\rho}, \check{\sigma}] + \mathcal{R}_q[\check{\beta}] + \underline{\mathcal{R}}_q[\check{\rho}, \check{\sigma}] + \underline{\mathcal{R}}_q[\check{\beta}] + \underline{\mathcal{R}}_q[\check{\alpha}] \right) \lesssim \epsilon_0, \tag{C.6}$$

which implies from Proposition 2.45 that

$$\mathcal{R}_0^S[\check{\beta}] + \mathcal{R}_0^S[\check{\rho}, \check{\sigma}] + \mathcal{R}_0^S[\check{\beta}] \lesssim \epsilon_0. \tag{C.7}$$

In view of (C.7), it remains to estimate  $\check{\alpha}$  and  $\check{\alpha}$ . Next, we state the following *Teukolsky equation* first derived by Teukolsky in the linearized setting in [26].

**Proposition C.1.** *We have the following Teukolsky equations for  $\alpha$  and  $\underline{\alpha}$ :*

$$\begin{aligned}
\Omega \nabla_3 (\Omega \nabla_4 (r \Omega^2 \alpha)) + 2r \Omega^2 d_2^* d_2 (\Omega^2 \alpha) &= -\frac{2\Omega}{r} \nabla_3 (r \Omega^2 \alpha) + \not{\partial}^{\leq 1} ((\mathbf{O}_3^2 + \Gamma_g) \cdot (\mathbf{O}_4^2 + \beta)), \\
\Omega \nabla_4 (\Omega \nabla_3 (r \Omega^2 \underline{\alpha})) + 2r \Omega^2 d_2^* d_2 (\Omega^2 \underline{\alpha}) &= \frac{2\Omega}{r} \nabla_4 (r \Omega^2 \underline{\alpha}) + \not{\partial}^{\leq 1} ((\mathbf{O}_3^2 + \Gamma_b) \cdot (\mathbf{O}_4^2 + \underline{\alpha})).
\end{aligned} \tag{C.8}$$

*Proof.* See for example Propositions 3.4.6 and 3.4.7 in [6]. □

**Lemma C.2.** *We define the following quantities:*

$$\begin{aligned}\dot{\alpha} &:= \frac{1}{r^4} \nabla_4(r^5 \alpha) \in \mathfrak{s}_2, & \phi &:= rd_2 \alpha \in \mathfrak{s}_1, \\ \underline{\dot{\alpha}} &:= \frac{1}{r^4} \nabla_3(r^5 \underline{\alpha}) \in \mathfrak{s}_2, & \underline{\phi} &:= rd_2 \underline{\alpha} \in \mathfrak{s}_1.\end{aligned}\tag{C.9}$$

Then, we have the following equations:

$$\begin{aligned}\nabla_3 \dot{\alpha} &= -2d_2^* \phi + \frac{4\alpha}{r} + \not\leq^1 \left( (\mathbf{O}_2^1 + \Gamma_g) \cdot (\mathbf{O}_4^2 + \check{\beta}) \right), \\ \nabla_4 \phi + \frac{5}{2} \operatorname{tr} \chi \phi &= d_2 \dot{\alpha} + \not\leq^1 \left( (\mathbf{O}_2^1 + \Gamma_g) \cdot (\mathbf{O}_4^2 + \check{\beta}) \right),\end{aligned}\tag{C.10}$$

and

$$\begin{aligned}\nabla_4 \underline{\dot{\alpha}} &= -2d_2^* \underline{\phi} + \frac{4\underline{\alpha}}{r} + \not\leq^1 \left( (\mathbf{O}_2^1 + \Gamma_b) \cdot (\mathbf{O}_4^2 + \check{\alpha}) \right), \\ \nabla_3 \underline{\phi} + \frac{5}{2} \operatorname{tr} \underline{\chi} \underline{\phi} &= d_2 \underline{\dot{\alpha}} + \not\leq^1 \left( (\mathbf{O}_2^1 + \Gamma_b) \cdot (\mathbf{O}_4^2 + \check{\alpha}) \right).\end{aligned}\tag{C.11}$$

*Proof.* It follows directly from Lemma C.3 in [25].<sup>32</sup> □

**Corollary C.3.** *We have the following linearized equations:*

$$\begin{aligned}\nabla_3 \check{\alpha} &= -2d_2^* \check{\phi} + \frac{4\check{\alpha}}{r} + \mathfrak{L}[\alpha] + \operatorname{Err}[\alpha], \\ \nabla_4 \check{\phi} + \frac{5}{2} \operatorname{tr} \chi \check{\phi} &= d_2 \check{\alpha} + \mathfrak{L}[\alpha] + \operatorname{Err}[\alpha], \\ \mathfrak{L}[\alpha] &= \mathbf{O}_4^2 \cdot \Gamma_g^{(1)} + \mathbf{O}_2^1 \cdot \check{\beta}^{(1)}, \\ \operatorname{Err}[\alpha] &= \Gamma_g^{(1)} \cdot \check{\beta} + \Gamma_g \cdot \check{\beta}^{(1)},\end{aligned}$$

and

$$\begin{aligned}\nabla_4 \check{\underline{\alpha}} &= -2d_2^* \check{\underline{\phi}} + \frac{4\check{\underline{\alpha}}}{r} + \mathfrak{L}[\underline{\alpha}] + \operatorname{Err}[\underline{\alpha}], \\ \nabla_3 \check{\underline{\phi}} + \frac{5}{2} \operatorname{tr} \underline{\chi} \check{\underline{\phi}} &= d_2 \check{\underline{\alpha}} + \mathfrak{L}[\underline{\alpha}] + \operatorname{Err}[\underline{\alpha}], \\ \mathfrak{L}[\underline{\alpha}] &= \mathbf{O}_4^2 \cdot \Gamma_b^{(1)} + \mathbf{O}_2^1 \cdot \check{\underline{\alpha}}^{(1)}, \\ \operatorname{Err}[\underline{\alpha}] &= \Gamma_b^{(1)} \cdot \check{\underline{\alpha}} + \Gamma_b \cdot \check{\underline{\alpha}}^{(1)}.\end{aligned}$$

*Proof.* It follows directly from Lemmas C.2 and 2.22. □

**Lemma C.4.** *We have the following identity for any real number  $p$ :*

$$\begin{aligned}& \operatorname{Div}(r^p |\check{\alpha}|^2 e_3) + 2\operatorname{Div}(r^p |\check{\phi}|^2 e_4) + (p+2)r^{p-1} |\check{\alpha}|^2 + 2(8-p)r^{p-1} |\check{\phi}|^2 \\ &= 4r^p d_1(\check{\alpha} \cdot \check{\phi}) + 8r^{p-1} \check{\alpha} \cdot \check{\alpha} + 2r^p (\check{\alpha} + 2\check{\phi}) \cdot (\mathfrak{L}[\alpha] + \operatorname{Err}[\alpha]) + r^p (\mathbf{O}_2^1 + \Gamma_b) \cdot (|\check{\alpha}|^2, |\check{\phi}|^2).\end{aligned}\tag{C.12}$$

*Proof.* Applying Lemma 4.2 with  $\psi_{(1)} = \check{\alpha}$ ,  $\psi_{(2)} = \check{\phi}$ ,  $a_{(1)} = 0$ ,  $a_{(2)} = \frac{5}{2}$ ,  $h_{(1)} = \frac{4\check{\alpha}}{r} + \mathfrak{L}[\alpha] + \operatorname{Err}[\alpha]$ ,  $h_{(2)} = \mathfrak{L}[\alpha] + \operatorname{Err}[\alpha]$  and  $k = 2$ , we infer

$$\begin{aligned}& \operatorname{Div}(r^p |\check{\alpha}|^2 e_3) + 2\operatorname{Div}(r^p |\check{\phi}|^2 e_4) - \left(1 + \frac{p}{2}\right) r^p \operatorname{tr} \chi |\check{\alpha}|^2 + 2 \left(4 - \frac{p}{2}\right) r^p \operatorname{tr} \chi |\check{\phi}|^2 \\ &= 4r^p d_1(\check{\alpha} \cdot \check{\phi}) + 8r^{p-1} \check{\alpha} \cdot \check{\alpha} + 2r^p (\check{\alpha} + 2\check{\phi}) \cdot (\mathfrak{L}[\alpha] + \operatorname{Err}[\alpha]) + r^p (\mathbf{O}_2^1 + \Gamma_b) \cdot (|\check{\alpha}|^2, |\check{\phi}|^2),\end{aligned}$$

which concludes the proof of Lemma C.4. □

---

<sup>32</sup>Note that  $\Gamma_a$  in [25] decays better than  $\mathbf{O}_2^1 + \Gamma_b$ .

We recall the following Poincaré inequality.

**Proposition C.5.** *For any  $\alpha \in \mathfrak{s}_2$ , we have the following Poincaré inequality:*

$$|rd_2\alpha|_{2,S}^2 \geq c_2|\alpha|_{2,S}^2,$$

for sphere  $S = S(u, \underline{u})$ , where  $c_2$  is a constant satisfying:

$$c_2 = 2 - O(\epsilon). \quad (\text{C.13})$$

*Proof.* See for example Proposition 9.3.2 in [6].  $\square$

**Proposition C.6.** *We have the following estimate:*

$$\int_{C_u^V} r^{s_0} |\check{\alpha}|^2 + \int_{\underline{C}_u^V} r^{s_0} |\check{\alpha}|^2 + \int_V r^{s_0-1} (|\check{\alpha}|^2 + |\check{\alpha}|^2) \lesssim \frac{\epsilon_0^2}{|u|^{s-s_0}}.$$

*Proof.* Integrating (C.12) with  $p = s_0$  and proceeding as in Proposition 4.9, we obtain

$$\begin{aligned} & \int_{C_u} r^{s_0} |\check{\alpha}|^2 + \int_{\underline{C}_u} r^{s_0} |\check{\alpha}|^2 + \int_V r^{s_0-1} ((s_0+2)|\check{\alpha}|^2 + 2(8-s_0)|\check{\alpha}|^2 - 8\check{\alpha} \cdot \check{\alpha}) \\ & \lesssim \int_{\Sigma_0 \cap V} r^{s_0} (|\check{\alpha}|^2 + |\check{\alpha}|^2) + \int_V r^{s_0} |(\check{\alpha}, \check{\alpha})| |\mathfrak{L}[\alpha] + \text{Err}[\alpha]| + r^{s_0} (\mathbf{O}_2^1 + \Gamma_b)(|\check{\alpha}|^2 + |\check{\alpha}|^2). \end{aligned}$$

First, we have

$$\int_{\Sigma_0 \cap V} r^{s_0} (|\check{\alpha}|^2 + |\check{\alpha}|^2) \lesssim |u|^{s_0-s} \int_{\Sigma_0 \cap V} r^s (|\check{\alpha}|^2 + |\check{\alpha}|^2) \lesssim \frac{\epsilon_0^2}{|u|^{s-s_0}}.$$

Next, we estimate

$$\begin{aligned} & \int_V r^{s_0} |(\check{\alpha}, \check{\alpha})| |\mathfrak{L}[\alpha]| \\ & \lesssim \int_V r^{s_0} |(\check{\alpha}, \check{\alpha})| \left| \mathbf{O}_4^2 \cdot \Gamma_g^{(1)} + \mathbf{O}_2^1 \cdot \check{\beta}^{(1)} \right| \\ & \lesssim \int_V M^2 r^{s_0-4} |(\check{\alpha}, \check{\alpha})| |\Gamma_g^{(1)}| + M r^{s_0-2} |(\check{\alpha}, \check{\alpha})| |\check{\beta}^{(1)}| \\ & \lesssim M^2 \int_{-\underline{u}}^u \frac{du}{r^{4-\frac{s_0}{2}}} \left( \int_{C_u^V} r^{s_0} |\check{\alpha}|^2 \right)^{\frac{1}{2}} \left( \int_{C_u^V} |\Gamma_g^{(1)}|^2 \right)^{\frac{1}{2}} + M^2 \int_{|u|}^u \frac{d\underline{u}}{r^{5-\frac{s_0}{2}}} \left( \int_{\underline{C}_u^V} r^{s_0} |\check{\alpha}|^2 \right)^{\frac{1}{2}} \left( \int_{\underline{C}_u^V} r^2 |\Gamma_g^{(1)}|^2 \right)^{\frac{1}{2}} \\ & \quad + M \int_{-\underline{u}}^u \frac{du}{r^{4-\frac{s_0}{2}}} \left( \int_{C_u^V} r^{s_0} |\check{\alpha}|^2 \right)^{\frac{1}{2}} \left( \int_{C_u^V} r^4 |\check{\beta}^{(1)}|^2 \right)^{\frac{1}{2}} + M \int_{|u|}^u \frac{d\underline{u}}{r^{5-\frac{s_0}{2}}} \left( \int_{\underline{C}_u^V} r^{s_0} |\check{\alpha}|^2 \right)^{\frac{1}{2}} \left( \int_{\underline{C}_u^V} r^6 |\check{\beta}^{(1)}|^2 \right)^{\frac{1}{2}} \\ & \lesssim M^2 \int_{-\underline{u}}^u \frac{du}{r^{4-\frac{s_0}{2}}} \frac{\epsilon}{|u|^{\frac{s-s_0}{2}}} \left( \int_{|u|}^u d\underline{u} |\Gamma_g^{(1)}|_{2,S}^2 \right)^{\frac{1}{2}} + M^2 \int_{|u|}^u \frac{d\underline{u}}{r^{5-\frac{s_0}{2}}} \frac{\epsilon}{|u|^{\frac{s-s_0}{2}}} \left( \int_{-\underline{u}}^u d\underline{u} |r\Gamma_g^{(1)}|_{2,S}^2 \right)^{\frac{1}{2}} \\ & \quad + M \int_{-\underline{u}}^u \frac{du}{r^{4-\frac{s_0}{2}}} \frac{\epsilon}{|u|^{\frac{s-s_0}{2}}} \frac{\epsilon}{|u|^{\frac{s-4}{2}}} + M \int_{|u|}^u \frac{d\underline{u}}{r^{5-\frac{s_0}{2}}} \frac{\epsilon}{|u|^{\frac{s-s_0}{2}}} \frac{\epsilon}{|u|^{\frac{s-6}{2}}} \\ & \lesssim M^2 \int_{-\underline{u}}^u \frac{du}{|u|^{4-\frac{s_0}{2}}} \frac{\epsilon}{|u|^{\frac{s-s_0}{2}}} \frac{\epsilon}{|u|^{\frac{s-2}{2}}} + M^2 \int_{|u|}^u \frac{d\underline{u}}{r^{5-\frac{s_0}{2}}} \frac{\epsilon}{|u|^{\frac{s-s_0}{2}}} \frac{\epsilon}{|u|^{\frac{s-4}{2}}} + \frac{\epsilon^2 M}{|u|^{s-s_0+1}} \\ & \lesssim \frac{\epsilon^2 M^2}{|u|^{s-s_0+2}} + \frac{\epsilon^2 M}{|u|^{s-s_0+1}} \\ & \lesssim \frac{\epsilon_0^2}{|u|^{s-s_0}}. \end{aligned}$$

Proceeding as in Proposition C.5 in [25], we infer

$$\int_V r^{s_0} |(\check{\alpha}, \check{\phi})| |\text{Err}[\alpha]| \lesssim \frac{\epsilon_0^2}{|u|^{s-s_0}}.$$

We have from Proposition 2.22

$$\check{\phi} = \widetilde{rd_2\alpha} = rd_2\check{\alpha} + (rd_2 - r_K(d_2)_K)\alpha_K = rd_2\check{\alpha} + \Gamma_g^{(1)} \cdot \mathbf{O}_4^3.$$

Combining with Proposition C.5, we obtain

$$\begin{aligned} & \int_V r^{s_0-1} \left( (s_0+2)|\check{\alpha}|^2 + 2(8-s_0)|\check{\phi}|^2 - 8\check{\alpha} \cdot \check{\alpha} \right) \\ & \gtrsim \int_V r^{s_0-1} \left( (s_0+2)|\check{\alpha}|^2 + 2c_2(8-s_0)|\check{\alpha}|^2 - 8\check{\alpha} \cdot \check{\alpha} \right) - \int_V r^{s_0-1} |\Gamma_g^{(1)} \cdot \mathbf{O}_4^3|^2. \end{aligned}$$

Since the discriminant satisfies<sup>33</sup>

$$\Delta = 8^2 - 4(s_0+2) \cdot 2c_2(8-s_0) < 0,$$

we obtain

$$\int_V r^{s_0-1} \left( (s_0+2)|\check{\alpha}|^2 + 2c_2(8-s_0)|\check{\alpha}|^2 - 8\check{\alpha} \cdot \check{\alpha} \right) \gtrsim \int_V r^{s_0-1} (|\check{\alpha}|^2 + |\check{\phi}^{\leq 1}\check{\alpha}|^2).$$

Finally, we also have

$$\int_V r^{s_0} (\mathbf{O}_2^1 + \Gamma_b) (|\check{\alpha}|^2 + |\check{\phi}|^2) \lesssim (\epsilon + \mathcal{E}_0) \int_V r^{s_0-1} (|\check{\alpha}|^2 + |\check{\phi}|^2).$$

Combining the above estimates, we deduce

$$\begin{aligned} & \int_{C_u} r^{s_0} |\check{\alpha}|^2 + \int_{\underline{C}_u} r^{s_0} |\check{\phi}^{\leq 1}\check{\alpha}|^2 + \int_V r^{s_0-1} (|\check{\alpha}|^2 + |\check{\phi}^{\leq 1}\check{\alpha}|^2) \\ & \lesssim \frac{\epsilon_0^2}{|u|^{s-s_0}} + (\epsilon + \mathcal{E}_0) \int_V r^{s_0-1} (|\check{\alpha}|^2 + |\check{\phi}|^2) + \int_{\underline{C}_u^V} r^{s_0} |\Gamma_g^{(1)} \cdot \mathbf{O}_4^3|^2 + \int_V r^{s_0-1} |\Gamma_g^{(1)} \cdot \mathbf{O}_4^3|^2 \\ & \lesssim \frac{\epsilon_0^2}{|u|^{s-s_0}} + (\epsilon + \mathcal{E}_0) \int_V r^{s_0-1} (|\check{\alpha}|^2 + |\check{\phi}|^2) + M^6 \int_{-\underline{u}}^u \frac{du}{r^{8-s_0}} |\Gamma_g^{(1)}|_{2,S}^2 + M^6 \int_{-\underline{u}}^u du \int_{|u|}^{\underline{u}} \frac{d\underline{u}}{r^{9-s_0}} |\Gamma_g^{(1)}|_{2,S}^2 \\ & \lesssim \frac{\epsilon_0^2}{|u|^{s-s_0}} + (\epsilon + \mathcal{E}_0) \int_V r^{s_0-1} (|\check{\alpha}|^2 + |\check{\phi}|^2) + \frac{M^6 \epsilon^2}{|u|^{6+s-s_0}} \\ & \lesssim \frac{\epsilon_0^2}{|u|^{s-s_0}} + (\epsilon + \mathcal{E}_0) \int_V r^{s_0-1} (|\check{\alpha}|^2 + |\check{\phi}|^2), \end{aligned}$$

which implies for  $\epsilon$  and  $\mathcal{E}_0$  small enough

$$\int_{C_u} r^{s_0} |\check{\alpha}|^2 + \int_{\underline{C}_u} r^{s_0} |\check{\phi}^{\leq 1}\check{\alpha}|^2 + \int_V r^{s_0-1} (|\check{\alpha}|^2 + |\check{\phi}^{\leq 1}\check{\alpha}|^2) \lesssim \frac{\epsilon_0^2}{|u|^{s-s_0}}.$$

This concludes the proof of Proposition C.6. □

**Proposition C.7.** *We have the following estimate:*

$$\mathcal{R}_0[\check{\alpha}] + \mathcal{R}_1[\check{\alpha}] + \mathcal{R}_0^S[\check{\alpha}] \lesssim \epsilon_0. \quad (\text{C.14})$$

---

<sup>33</sup>Here, we used  $6 < s_0 \leq \frac{29}{4}$  and  $c_2 = 2 - O(\epsilon)$ .

*Proof.* We have from Propositions 2.15 and C.6

$$\underline{\mathcal{R}}_0[\check{\alpha}]^2 + \underline{\mathcal{R}}_1[\check{\alpha}]^2 \lesssim |u|^{s-s_0} \int_{\underline{C}_u^V} |\check{\alpha}|^2 + |\check{\alpha}|^2 \lesssim \epsilon_0^2. \quad (\text{C.15})$$

Recalling that

$$\check{\alpha} = \frac{1}{r^4} \nabla_4(r^5 \check{\alpha}) + \mathbf{O}_4^3 \cdot \Gamma_g^{(1)},$$

we infer from Lemma 2.7 that

$$\begin{aligned} \Omega e_4 \left( \int_S |r^4 \check{\alpha}|^2 \right) &= \int_S 2\Omega \nabla_4(r^4 \check{\alpha}) \cdot (r^4 \check{\alpha}) + \Omega \operatorname{tr} \chi |r^4 \check{\alpha}|^2 \\ &= \int_S 2\Omega r^4 \check{\alpha} \cdot (\nabla_4(r^4 \check{\alpha}) + r^3 \check{\alpha}) + \Omega \left( \operatorname{tr} \chi - \frac{2}{r} \right) |r^4 \check{\alpha}|^2 \\ &= \int_S 2\Omega r^4 \check{\alpha} \cdot (\nabla_4(r^5 \check{\alpha}) r^{-1} + e_4(r^{-1}) r^5 \check{\alpha} + r^3 \check{\alpha}) + (\mathbf{O}_2^1 + \Gamma_g) |r^4 \check{\alpha}|^2 \\ &= \int_S 2\Omega r^4 \check{\alpha} \cdot (\nabla_4(r^5 \check{\alpha}) r^{-1}) + (\mathbf{O}_2^1 + \Gamma_g) |r^4 \check{\alpha}|^2 \\ &= \int_S 2\Omega r^4 \check{\alpha} \cdot r^3 \check{\alpha} + (\mathbf{O}_2^1 + \Gamma_g) |r^4 \check{\alpha}|^2 + (r^4 \check{\alpha}) \cdot \mathbf{O}_1^3 \cdot \Gamma_g^{(1)}. \end{aligned}$$

Hence, we deduce

$$|e_4(|r^4 \check{\alpha}|_{2,S}^2)| \lesssim |r^4 \check{\alpha}|_{2,S} |r^3 \check{\alpha}|_{2,S} + (M + \epsilon) r^{-2} |r^4 \check{\alpha}|_{2,S}^2 + |r^4 \check{\alpha}|_{2,S} |\mathbf{O}_1^3 \cdot \Gamma_g^{(1)}|_{2,S},$$

which implies

$$|e_4(|r^4 \check{\alpha}|_{2,S})| \lesssim |r^3 \check{\alpha}|_{2,S} + (M + \epsilon) r^{-2} |r^4 \check{\alpha}|_{2,S} + |\mathbf{O}_1^3 \cdot \Gamma_g^{(1)}|_{2,S}. \quad (\text{C.16})$$

Integrating (C.16) along  $C_u$  and applying the initial assumption and Proposition C.6, we infer

$$\begin{aligned} |r^4 \check{\alpha}|_{2,S} &\lesssim \frac{\epsilon_0}{r^{\frac{s-7}{2}}} + \int_{|u|}^u d\underline{u} |r^3 \check{\alpha}|_{2,S} + \int_{|u|}^u \frac{(M + \epsilon) d\underline{u}}{r^2} |r^4 \check{\alpha}|_{2,S} + \int_{|u|}^u d\underline{u} \frac{M^3}{r^2 |u|^{\frac{s-3}{2}}} \\ &\lesssim \frac{\epsilon_0}{|u|^{\frac{s-7}{2}}} + \left( \int_{|u|}^u r^{6-s_0} d\underline{u} \right)^{\frac{1}{2}} \left( \int_{|u|}^u r^{s_0} |\check{\alpha}|^2 \right)^{\frac{1}{2}} + \int_{|u|}^u \frac{(M + \epsilon) d\underline{u}}{r^2} |r^4 \check{\alpha}|_{2,S}. \end{aligned}$$

Applying Gronwall inequality, we obtain

$$|r^4 \check{\alpha}|_{2,S} \lesssim \frac{\epsilon_0}{|u|^{\frac{s-7}{2}}} + \left( \int_{|u|}^u r^{6-s_0} d\underline{u} \right)^{\frac{1}{2}} \left( \int_{|u|}^u r^{s_0} |\check{\alpha}|^2 \right)^{\frac{1}{2}}. \quad (\text{C.17})$$

Notice that we have

$$\int_{|u|}^u r^{6-s_0} d\underline{u} = \begin{cases} \frac{1}{7-s_0} (\underline{u}^{7-s_0} - |u|^{7-s_0}), & s_0 \in (6, 7), \\ \log \left( \frac{\underline{u}}{|u|} \right), & s_0 = 7, \\ \frac{1}{s_0-7} (|u|^{7-s_0} - \underline{u}^{7-s_0}), & s_0 > 7. \end{cases}$$

Hence, we obtain from (C.17)

$$|r^4 \alpha|_{2,S} \lesssim \begin{cases} \epsilon_0 r^{\frac{7-s}{2}}, & s \in (6, 7), \\ \epsilon_0 (\log r)^{\frac{1}{2}}, & s = 7, \\ \epsilon_0 |u|^{\frac{7-s}{2}}, & s > 7, \end{cases} \quad (\text{C.18})$$

which implies

$$\mathcal{R}_0^S[\check{\alpha}] \lesssim \epsilon_0. \quad (\text{C.19})$$

Combining (C.15) and (C.19), this concludes the proof of Proposition C.7.  $\square$

**Proposition C.8.** *We have the following estimate:*

$$\mathcal{R}_0^S[\check{\alpha}] \lesssim \epsilon_0.$$

*Proof.* We recall from Corollary C.3

$$\begin{aligned} \nabla_4 \check{\alpha} &= -2d_2^* \check{\alpha} + \frac{4\check{\alpha}}{r} + \mathfrak{L}[\alpha] + \text{Err}[\alpha], \\ \nabla_3 \check{\alpha} + \frac{5}{2} \text{tr} \chi \check{\alpha} &= d_2 \check{\alpha} + \mathfrak{L}[\alpha] + \text{Err}[\alpha], \\ \mathfrak{L}[\alpha] &= \mathbf{O}_4^2 \cdot \Gamma_b^{(1)} + \mathbf{O}_2^1 \cdot \check{\alpha}^{(1)}, \\ \text{Err}[\alpha] &= \Gamma_b^{(1)} \cdot \check{\alpha} + \Gamma_b \cdot \check{\alpha}^{(1)}. \end{aligned}$$

Applying Lemma 4.2 with  $\psi_{(1)} = \check{\alpha}$ ,  $\psi_{(2)} = \check{\alpha}$ ,  $a_{(1)} = \frac{5}{2}$ ,  $a_{(2)} = 0$ ,  $h_{(1)} = \mathfrak{L}[\alpha] + \text{Err}[\alpha]$ ,  $h_{(2)} = \frac{4\check{\alpha}}{r} + \mathfrak{L}[\alpha] + \text{Err}[\alpha]$ ,  $k = 2$  and  $p = -2$ , we obtain

$$\begin{aligned} &2\mathbf{Div}(r^{-2}|\check{\alpha}|^2 e_3) + \mathbf{Div}(r^{-2}|\check{\alpha}|^2 e_4) + 10r^{-2} \text{tr} \chi |\check{\alpha}|^2 \\ &= 2r^{-2} d_1(\check{\alpha} \cdot \check{\alpha}) + r^{-2}(\check{\alpha}, \check{\alpha}) \cdot (\mathfrak{L}[\alpha] + \text{Err}[\alpha]) + 8r^{-3} \check{\alpha} \cdot \check{\alpha} + r^{-2}(\mathbf{O}_2^1 + \Gamma_b)(|\check{\alpha}|^2, |\check{\alpha}|^2). \end{aligned}$$

We compute for  $c_2$  defined in Proposition C.5

$$\begin{aligned} 2c_2 \mathbf{Div}(r^{-2}|\check{\alpha}|^2 e_3) &= 2c_2 \nabla_3(r^{-2}|\check{\alpha}|^2) + 2c_2 r^{-2} |\check{\alpha}|^2 \mathbf{Div}(e_3) \\ &= 2c_2 \nabla_3(r^{-12}|r^5 \check{\alpha}|^2) + 2c_2 r^{-2} |\check{\alpha}|^2 (\text{tr} \chi - 2\omega) \\ &= 20c_2 r^{-3} |\check{\alpha}|^2 + 4c_2 r^{-3} \check{\alpha} \cdot \check{\alpha} + r^{-2}(\mathbf{O}_2^1 + \Gamma_b) |\check{\alpha}|^2 + \mathbf{O}_7^3 \cdot \Gamma_b^{(1)} \cdot \check{\alpha}, \end{aligned}$$

where we used

$$\begin{aligned} \check{\alpha} &= \frac{1}{r^4} \nabla_3(r^5 \alpha) - \frac{1}{r_K^4} (\nabla_K)_{(e_3)_K} (r_K^5 \alpha_K) \\ &= \frac{1}{r^4} \nabla_3(r^5 \alpha) - \frac{1}{r^4} \nabla_3(r^5 \alpha_K) + \frac{1}{r^4} \nabla_3(r^5 \alpha_K) - \frac{1}{r_K^4} (\nabla_K)_{(e_3)_K} (r_K^5 \alpha_K) \\ &= \frac{1}{r^4} \nabla_3(r^5 \check{\alpha}) + \mathbf{O}_4^3 \cdot \Gamma_b^{(1)}. \end{aligned} \quad (\text{C.20})$$

Combining the above identities, we deduce

$$\begin{aligned} &2\mathbf{Div}(r^{-2}(|\check{\alpha}|^2 - c_2 |\check{\alpha}|^2) e_3) + \mathbf{Div}(r^{-2}|\check{\alpha}|^2) + 10r^{-2} \text{tr} \chi |\check{\alpha}|^2 + 20c_2 r^{-3} |\check{\alpha}|^2 \\ &= 2r^{-2} d_1(\check{\alpha} \cdot \check{\alpha}) + (8 - 4c_2) r^{-3} \check{\alpha} \cdot \check{\alpha} + r^{-2}(\check{\alpha}, \check{\alpha}) \cdot (\mathfrak{L}[\alpha] + \text{Err}[\alpha]) + \mathbf{O}_7^3 \cdot \Gamma_b^{(1)} \cdot \check{\alpha} + r^{-2}(\mathbf{O}_2^1 + \Gamma_b) |\check{\alpha}|^2. \end{aligned}$$

Integrating it in  $V$ , we obtain from Proposition C.5

$$\begin{aligned} \int_{C_u^V} r^{-2}(|\check{\underline{\alpha}}|^2 - c_2|\check{\underline{\alpha}}|^2) + \int_{\underline{C}_u^V} r^{-2}|\check{\underline{\alpha}}|^2 &\lesssim \int_{\Sigma_0 \cap V} r^{-2}(|\check{\underline{\alpha}}|^2 + |\check{\underline{\alpha}}|^2) + \int_V r^{-3}|\check{\underline{\alpha}}|^2 + \epsilon r^{-3}|\check{\underline{\alpha}}||\check{\underline{\alpha}}| \\ &+ \int_V r^{-2}(|\check{\underline{\alpha}}, \check{\underline{\alpha}}||\mathfrak{L}[\underline{\alpha}] + \text{Err}[\underline{\alpha}]| + |\mathbf{O}_7^3 \cdot \Gamma_b^{(1)}||\check{\underline{\alpha}}| + r^{-2}(\mathbf{O}_2^1 + \Gamma_b)|\check{\underline{\alpha}}|^2). \end{aligned}$$

We first have from Proposition 2.22

$$\check{\underline{\alpha}} = \widetilde{rd_2 \underline{\alpha}} = rd_2 \check{\underline{\alpha}} + (rd_2 - r_K(d_2)_K)\underline{\alpha}_K = rd_2 \check{\underline{\alpha}} + \Gamma_g^{(1)} \cdot \mathbf{O}_4^3.$$

Noticing that

$$\int_{C_u^V} r^{-2}|\Gamma_g^{(1)} \cdot \mathbf{O}_4^3|^2 \lesssim \int_{|u|}^u \frac{du}{r^{12}} |r\Gamma_g^{(1)}|_{2,S}^2 \lesssim M^6 \int_{|u|}^u \frac{du}{r^{12}|u|^{s-3}} \lesssim \frac{\epsilon_0^2}{|u|^{s+2}},$$

Hence, we obtain from Proposition C.5

$$\int_{C_u^V} r^{-2}(|\check{\underline{\alpha}}|^2 - c_2|\check{\underline{\alpha}}|^2) \geq \int_{C_u^V} r^{-2}(|rd_2 \check{\underline{\alpha}}|^2 - c_2|\check{\underline{\alpha}}|^2) - \int_{C_u^V} r^{-2}|\Gamma_g^{(1)} \cdot \mathbf{O}_4^3|^2 \gtrsim -\frac{\epsilon_0^2}{|u|^{s+2}}.$$

Next, we have

$$\int_{\Sigma_0 \cap V} r^{-2}(|\check{\underline{\alpha}}|^2 + |\check{\underline{\alpha}}|^2) \lesssim \frac{1}{|u|^{s+2}} \int_{\Sigma_0 \cap V} r^s(|\check{\underline{\alpha}}|^2 + |\check{\underline{\alpha}}|^2) \lesssim \frac{\epsilon_0^2}{|u|^{s+2}}.$$

Then, we compute

$$\begin{aligned} \int_V r^{-3}|\check{\underline{\alpha}}|^2 + \epsilon \int_V r^{-3}|\check{\underline{\alpha}}||\check{\underline{\alpha}}| &\lesssim \int_{|u|}^u \frac{du}{r^3} \int_{\underline{C}_u^V} |\check{\underline{\alpha}}|^2 + \epsilon \int_{|u|}^u r^{-2} \left( \int_{\underline{C}_u^V} r^{-2}|\check{\underline{\alpha}}|^2 \right)^{\frac{1}{2}} \left( \int_{\underline{C}_u^V} |\check{\underline{\alpha}}|^2 \right)^{\frac{1}{2}} du \\ &\lesssim \epsilon_0^2 \int_{|u|}^u \frac{du}{r^3|u|^s} + \epsilon \int_{|u|}^u r^{-2} \left( \frac{\epsilon^2}{|u|^{s+2}} \right)^{\frac{1}{2}} \left( \frac{\epsilon_0^2}{|u|^s} \right)^{\frac{1}{2}} du \\ &\lesssim \frac{\epsilon_0^2}{|u|^{s+2}}. \end{aligned}$$

We also have

$$\begin{aligned} &\int_V r^{-2}|\check{\underline{\alpha}}, \check{\underline{\alpha}}||\mathfrak{L}[\underline{\alpha}]| \\ &\lesssim \int_V r^{-2}|\check{\underline{\alpha}}, \check{\underline{\alpha}}| \left| \mathbf{O}_4^2 \cdot \Gamma_b^{(1)} + \mathbf{O}_2^1 \cdot \check{\underline{\alpha}}^{(1)} \right| \\ &\lesssim \int_V M^2 r^{-6}|\check{\underline{\alpha}}||\Gamma_b^{(1)}| + M^2 r^{-6}|\check{\underline{\alpha}}||\Gamma_b^{(1)}| + Mr^{-4}|\check{\underline{\alpha}}||\check{\underline{\alpha}}^{(1)}| + Mr^{-4}|\check{\underline{\alpha}}||\check{\underline{\alpha}}^{(1)}| \\ &\lesssim M^2 \int_{|u|}^u \frac{du}{r^5} \left( \int_{\underline{C}_u^V} r^{-2}|\check{\underline{\alpha}}|^2 \right)^{\frac{1}{2}} \left( \int_{\underline{C}_u^V} |\Gamma_b^{(1)}|^2 \right)^{\frac{1}{2}} + M^2 \int_{|u|}^u \frac{du}{r^5} \left( \int_{\underline{C}_u^V} |\check{\underline{\alpha}}|^2 \right)^{\frac{1}{2}} \left( \int_{\underline{C}_u^V} |r^{-1}\Gamma_b^{(1)}|^2 \right)^{\frac{1}{2}} \\ &\quad + M \int_{|u|}^u \frac{du}{r^3} \left( \int_{\underline{C}_u^V} r^{-2}|\check{\underline{\alpha}}|^2 \right)^{\frac{1}{2}} \left( \int_{\underline{C}_u^V} |\check{\underline{\alpha}}^{(1)}|^2 \right)^{\frac{1}{2}} + M \int_{|u|}^u \frac{du}{r^4} \left( \int_{\underline{C}_u^V} |\check{\underline{\alpha}}|^2 \right)^{\frac{1}{2}} \left( \int_{\underline{C}_u^V} |\check{\underline{\alpha}}^{(1)}|^2 \right)^{\frac{1}{2}} \\ &\lesssim M^2 \int_{|u|}^u \frac{du}{r^5} \frac{\epsilon}{|u|^{\frac{s+2}{2}}} \left( \int_{-u}^u du |\Gamma_b^{(1)}|_{2,S}^2 \right)^{\frac{1}{2}} + M^2 \int_{|u|}^u \frac{du}{r^5} \frac{\epsilon}{|u|^{\frac{s}{2}}} \left( \int_{-u}^u du |r^{-1}\Gamma_b^{(1)}|_{2,S}^2 \right)^{\frac{1}{2}} \end{aligned}$$

$$\begin{aligned}
& + M \int_{|u|}^u \frac{du}{r^3} \frac{\epsilon}{|u|^{\frac{s+2}{2}}} \frac{\epsilon}{|u|^{\frac{s}{2}}} + M \int_{|u|}^u \frac{du}{r^4} \frac{\epsilon}{|u|^{\frac{s}{2}}} \frac{\epsilon}{|u|^{\frac{s}{2}}} \\
& \lesssim \frac{M^2 \epsilon^2}{r^4 |u|^s} + \frac{M \epsilon^2}{r^2 |u|^{s+1}} + \frac{M \epsilon^2}{r^3 |u|^s} \\
& \lesssim \frac{\epsilon_0^2}{|u|^{s+2}}.
\end{aligned}$$

We proceed as in Proposition C.7 in [25] to deduce

$$\int_V r^{-2} |\check{\underline{\alpha}}, \check{\underline{\alpha}}| |\text{Err}[\underline{\alpha}]| \lesssim \frac{\epsilon_0^2}{|u|^{s+2}}.$$

Moreover, we have

$$\begin{aligned}
\int_V |\check{\underline{\alpha}}| |\mathbf{O}_7^3 \cdot \Gamma_b^{(1)}| & \lesssim M^3 \int_{|u|}^u \frac{du}{r^7} \left( \int_{\underline{C}_u^V} |\check{\underline{\alpha}}|^2 \right)^{\frac{1}{2}} \left( \int_{\underline{C}_u^V} |\Gamma_b^{(1)}|^2 \right)^{\frac{1}{2}} \\
& \lesssim M^3 \int_{|u|}^u \frac{du}{r^7} \frac{\epsilon}{|u|^{\frac{s}{2}}} \left( \int_{-\underline{u}}^u du |\Gamma_b^{(1)}|_{2,S}^2 \right)^{\frac{1}{2}} \\
& \lesssim M^3 \epsilon \int_{|u|}^u \frac{du}{r^7} \frac{\epsilon}{|u|^{\frac{s}{2}}} \frac{\epsilon}{|u|^{\frac{s-2}{2}}} \\
& \lesssim \frac{\epsilon_0^2}{|u|^{s+2}}.
\end{aligned}$$

Finally, we have

$$\int_V r^{-2} (\mathbf{O}_2^1 + \Gamma_b) |\check{\underline{\alpha}}|^2 \lesssim \int_V (\epsilon + \mathcal{E}_0) r^{-3} |\check{\underline{\alpha}}|^2 \lesssim \frac{\epsilon_0^2}{|u|^{s+2}}.$$

Combining the above estimates, we deduce

$$\int_{\underline{C}_u^V} r^{-2} |\check{\underline{\alpha}}|^2 \lesssim \frac{\epsilon_0^2}{|u|^{s+2}},$$

which implies from (C.20) and (4.13)

$$\int_{\underline{C}_u^V} r^{-2} |\nabla_3 \check{\underline{\alpha}}|^2 \lesssim \frac{\epsilon_0^2}{|u|^{s+2}}. \quad (\text{C.21})$$

Combining (C.21) and (C.2) and applying Proposition 2.45, we deduce

$$\underline{\mathcal{R}}_0^S[\check{\underline{\alpha}}] \lesssim \epsilon_0.$$

This concludes the proof of Proposition C.8.  $\square$

Combining Propositions C.7 and C.8 with (C.2), this concludes the proof of Theorem M1 in the case  $s > 6$ . Notice that Theorems M0, M2, M3, M4 also hold true in the case  $s > 6$  since we only used  $s > 3$  in their proofs<sup>34</sup>. Combining with Section B.3, this concludes the proof of Theorem 3.2 for  $s > 3$  as stated.

---

<sup>34</sup>See the discussion in Section B.2.

### C.3 Peeling decay for curvature components

As a consequence of Theorem M1 in the case  $s > 7$ , the curvature components satisfy the following decay:

$$\begin{aligned}
& |r^{5-\frac{2}{p}}|u|^{\frac{s-7}{2}}\check{\alpha}|_{p=2,S} \lesssim \epsilon_0, \\
& \sup_{p \in [2,4]} |r^{4-\frac{2}{p}}|u|^{\frac{s-5}{2}}\check{\beta}|_{p,S} \lesssim \epsilon_0, \\
& \sup_{p \in [2,4]} |r^{3-\frac{2}{p}}|u|^{\frac{s-3}{2}}(\check{\rho}, \check{\sigma})|_{p,S} \lesssim \epsilon_0, \\
& \sup_{p \in [2,4]} |r^{2-\frac{2}{p}}|u|^{\frac{s-1}{2}}\check{\underline{\beta}}|_{p,S} \lesssim \epsilon_0, \\
& \sup_{p \in [2,4]} |r^{1-\frac{2}{p}}|u|^{\frac{s+1}{2}}\check{\underline{\alpha}}|_{p,S} \lesssim \epsilon_0.
\end{aligned} \tag{C.22}$$

**Remark C.9.** Assume  $s > 7$  and that the initial data in  $\mathcal{K}_{(0)}$  has sufficient regularity properties. Then, commuting  $r\nabla$  with the Bianchi equations in Propositions 2.37 and 2.38 and the Teukolsky equation (C.8), proceeding as in Sections 4 and C.2, we deduce the following strong peeling properties:

$$\begin{aligned}
|\check{\alpha}|_{\infty, S(u, \underline{u})} &\lesssim \frac{\epsilon_0}{r^5|u|^{\frac{s-7}{2}}}, & |\check{\beta}|_{\infty, S(u, \underline{u})} &\lesssim \frac{\epsilon_0}{r^4|u|^{\frac{s-5}{2}}}, \\
|\check{\rho}, \check{\sigma}|_{\infty, S(u, \underline{u})} &\lesssim \frac{\epsilon_0}{r^3|u|^{\frac{s-3}{2}}}, & |\check{\underline{\beta}}|_{\infty, S(u, \underline{u})} &\lesssim \frac{\epsilon_0}{r^2|u|^{\frac{s-1}{2}}}, & |\check{\underline{\alpha}}|_{\infty, S(u, \underline{u})} &\lesssim \frac{\epsilon_0}{r|u|^{\frac{s+1}{2}}},
\end{aligned} \tag{C.23}$$

which recovers the results obtained in [2] by the vectorfield method. See also Theorem 1 in [15] and Remark C.8 in [25] for the particular case of Minkowski stability.

## Declarations

**Acknowledgements.** The author is very grateful to Jérémie Szeftel for his support, discussions, encouragements and patient guidance.

**Funding.** No funding was received to assist with the preparation of this manuscript.

**Competing Interest statements.** Conflict of interest does not exist in the manuscript.

## References

- [1] R. Adams and J. Fournier, *Sobolev spaces*, Second edition. Pure and Applied Mathematics (Amsterdam), **140**. Elsevier, Academic Press, 2003.
- [2] G. Caciotta and F. Nicolò, *Non linear perturbations of Kerr spacetime in external regions and the peeling decay*, Ann. Henri Poincaré, **11**, 433–497, 2010.
- [3] Y. Choquet-Bruhat, *Théorème d’existence pour certain systèmes d’équations aux dérivées partielles non linéaires*, Acta Math **88**, 141–225, 1952.
- [4] Y. Choquet-Bruhat and R. Geroch, *Global aspects of the Cauchy problem in general relativity*, Comm. Math. Phys. **14**, 329–335, 1969.

- [5] D. Christodoulou and S. Klainerman, *The Global Nonlinear Stability of Minkowski Space*, Princeton Mathematical Series **41**, 1993.
- [6] M. Dafermos, G. Holzegel, I. Rodnianski and M. Taylor, *The non-linear stability of the Schwarzschild family of black holes*, arXiv:2104.08222.
- [7] M. Dafermos and J. Luk *The interior of dynamical vacuum black holes I: The  $C^0$ -stability of the Kerr Cauchy horizon*, arXiv:1710.01722.
- [8] M. Dafermos and I. Rodnianski, *A new physical-space approach to decay for the wave equation with applications to black hole spacetimes*, XVIth International Congress on Mathematical Physics, World Sci. Publ., Hackensack, NJ, 2010, 421–432.
- [9] E. Giorgi, S. Klainerman and J. Szeftel, *Wave equations estimates and the nonlinear stability of slowly rotating Kerr black holes*, arXiv:2205.14808.
- [10] P. Hintz, *Exterior stability of Minkowski space in generalized harmonic gauge*, Arch. Rational Mech. Anal. **247** (99), 2023.
- [11] G. Holzegel, *Ultimately Schwarzschild spacetimes and the black hole stability problem*, arXiv:1010.3216.
- [12] W. Israel and F. Pretorius, *Quasispherical light cones of the Kerr geometry*, Class. Quantum Grav. **15**, 2289–2301, 1998.
- [13] R. P. Kerr, *Gravitational field of a spinning mass as an example of algebraically special metrics*, Phys. Rev. Lett. **11**, 237, 1963.
- [14] S. Klainerman and F. Nicolò, *The Evolution Problem in General Relativity*. Progress in Mathematical Physics, **25**, Birkhauser, Boston, 2003.
- [15] S. Klainerman and F. Nicolò, *Peeling properties of asymptotically flat solutions to the Einstein vacuum equations*, Classical. Quant. Grav. **20**, 3215–3257, 2003.
- [16] S. Klainerman and I. Rodnianski, *On the formation of trapped surfaces*, Acta Math **208**, 211–333, 2012.
- [17] S. Klainerman and J. Szeftel, *Global nonlinear stability of Schwarzschild spacetime under polarized perturbations*, Annals of Math Studies, **210**. Princeton University Press.
- [18] S. Klainerman and J. Szeftel, *Construction of GCM spheres in perturbations of Kerr*, Ann. PDE **8** (2), Art. 17, 153 pp, 2022.
- [19] S. Klainerman and J. Szeftel, *Effective results in uniformization and intrinsic GCM spheres in perturbations of Kerr*, Ann. PDE **8** (2), Art. 18, 89 pp, 2022.
- [20] S. Klainerman and J. Szeftel, *Kerr stability for small angular momentum*, Pure Appl. Math. Q. **19** (3), 791–1678, 2023.
- [21] S. Klainerman and J. Szeftel, *Brief introduction to the nonlinear stability of Kerr*, arXiv:2210.14400, To appear in Pure Appl. Math. Q..
- [22] J. Luk, *On the Local Existence for the Characteristic Initial Value Problem in General Relativity*, Int. Math. Res. Not. No. **20**, 4625–4678, 2018.

- [23] F. Nicolò, *The peeling in the very external region of non linear perturbations of the Kerr spacetime*, Royal Swedish Academy of Sciences. No. **23**, 2009.
- [24] D. Shen, *Construction of GCM hypersurfaces in perturbations of Kerr*, Ann. PDE **9** (1), Art. 11, 112 pp, 2023.
- [25] D. Shen, *Stability of Minkowski spacetime in exterior regions*, Pure Appl. Math. Q. **20** (2), 757–868, 2024.
- [26] S. A. Teukolsky, *Rotating black holes: Separable wave equations for gravitational and electromagnetic perturbations*, Phys. Rev. Lett. **29**, No. 16, 1114–1118, 1992.