

Electronic-mediated nuclear stopping power in proton irradiated water ice

Daniel Muñoz-Santiburcio,^{1,2,*} Jorge Kohanoff,² and Emilio Artacho^{1,3,4,5}

¹*CIC nanoGUNE BRTA, Tolosa Hiribidea 76, 20018 San Sebastián, Spain*

²*Instituto de Fusión Nuclear “Guillermo Velarde”, Universidad Politécnica de Madrid, C/ José Gutiérrez Abascal 2, 28006 Madrid, Spain*

³*Theory of Condensed Matter, Cavendish Laboratory, University of Cambridge, J. J. Thomson Ave, Cambridge CB3 0HE, United Kingdom*

⁴*Donostia International Physics Center (DIPC), Tolosa Hiribidea 76, 20018 San Sebastián, Spain*

⁵*Ikerbasque, Basque Foundation for Science, 48011 Bilbao, Spain*

(Dated: March 24, 2023)

Traditionally, it has been assumed that the stopping of a swift ion travelling through matter can be understood in terms of two essentially independent components, i.e. electronic *vs.* nuclear. Performing extensive Ehrenfest MD simulations of the process of proton irradiation of water ice that accurately describe not only the non-adiabatic dynamics of the electrons but also of the nuclei, we have found a stopping mechanism involving the interplay of the electronic and nuclear subsystems. This effect, which consists in a kinetic energy transfer from the projectile to the target nuclei thanks to the perturbations of the electronic density caused by the irradiation, is fundamentally different from the atomic displacements and collision cascades characteristic of nuclear stopping. Moreover, it shows a marked isotopic effect depending on the composition of the target, being relevant mostly for light water as opposed to heavy water. This result is consistent with long-standing experimental results which remained unexplained so far.

The interaction of a swift ion with matter has been usually rationalized using the concept of ‘stopping power’, which consists in expressing the capacity of a target material to stop a given ion travelling through it as the energy lost by the projectile per unit of travelled length. It has been usually assumed that this has two components, being the ‘electronic’ and ‘nuclear stopping power’ [1]. The former consists of the energy absorbed by the electronic subsystem of the target, which becomes excited as a consequence of the projectile’ passing, while the latter consists of the energy gained by the target’ nuclei as a consequence of the direct impact of the projectile with them. Both these quantities show a strong dependence on the velocity of the projectile, in such a way that the velocity regimes where each of them has a significant value are basically non-overlapping. While it has been usually acknowledged that there is a certain domain of projectile velocities where both effects coexist, it was assumed that this happened where both components are quite small, and therefore this coincidence was not regarded as important for most purposes. Especially for computer simulations aimed at determining the electronic stopping power of any material by first-principles methods, the motion of the target’ nuclei and even the change of the velocity of the projectile are almost always neglected not only for practical reasons (i.e. alleviating the computational cost of the simulation by skipping the calculation of the atomic forces), but also because nuclear stopping is assumed to be negligible in the domain of projectile velocities that are relevant for the electronic stopping power. In this work, we show that such approximation neglects a channel for the projectile-target energy transfer that has remained hitherto unexplored.

Methods. – We performed Real-Time Time-Dependent DFT simulations within the Ehrenfest MD (EMD) formalism, explicitly considering the forces on all atoms during the irradiation process, employing the CP2K code [2] and using the Perdew-Burke-Ernzerhof (PBE) [3] exchange-correlation functional. All the simulations were carried out at the all-electron level with the 6-311++G2d2p basis set [4, 5], using the GAPW method [6, 7] with a cutoff of 350 Ry for the planewave expansion of the soft part of the density. The model system consists of a 1h ice slab with 144 H₂O molecules in the unit cell constructed from the corresponding bulk system with the experimental density at $T = 145$ K ($\rho = 931.19$ kg/m³, with $a = 15.59$, $b = 13.50$ and $c = 21.98$ Å in the bulk [8], extended to $c = 33.0$ Å in the slab), with a proton projectile initially placed in the vacuum region above the slab and with an initial velocity perpendicular to the slab. We simulated 19 different proton trajectories for each one of 11 different values of the projectile’ velocity, from 0.2 to 8.0 a.u. The timesteps for the EMD simulations were chosen to obtain a projectile displacement of ≈ 0.005 Å per EMD step in each case. Further details will be published elsewhere, including a deeper description of the methods and further analyses of the results.

Stopping power: electronic vs. nuclear. – The electronic, nuclear and total stopping power are readily obtained from each simulation as $S_e = \Delta E_{KS}/d$, $S_n = \Delta E_{kin}^{target}/d$, and $S_t = -\Delta E_{kin}^{H^+}/d$, being respectively the differences between the initial and final EMD steps of the Kohn-Sham energy and of the kinetic energies of the target nuclei and the projectile, in all cases divided by the distance travelled by the projectile within the target.




FIG. 1. Average nuclear and electronic stopping power of protons in water ice computed in this work, compared to that obtained from SRIM 2013 [9, 10] (using a 0.94 compound correction) and ICRU 2014 [11], in addition to the S_e experimental results of Shimizu *et al* [12, 13]. All values are scaled to a water density of 998 kg/m³ for an easier comparison.

These definitions imply that in the following we will employ the terms ‘electronic’ and ‘nuclear’ stopping as mere quantifiers of the energy transferred to the electronic *vs.* nuclear subsystems, with independence of the underlying mechanism of such transfer.

In all the simulations, the relation $S_t = S_e + S_n$ holds, with an excellent energy conservation along the irradiation process. As shown in Fig. 1, the average electronic stopping power agrees remarkably well with that from SRIM 2013 [9, 10] and with the tabulated data in ICRU Report 90 [11], with a perfect agreement with the experimental results of Shimizu *et al.* [12, 13] at high velocities. On the other hand, the average nuclear stopping power differs greatly from the SRIM and ICRU data, occurring the greatest values in the same regime of projectile velocities as the electronic stopping maximum. The small $\langle S_n \rangle$ obtained by us in the low velocity limit is surely a consequence of the small size- and timescales in our simulations, which do not allow to observe the proper collision cascades which are characteristic in the regular nuclear stopping processes. On the other hand, the strikingly high $\langle S_n \rangle$ values in the v_{H+} range from 1 to 2 a.u. require a more complex explanation.

The origin of such unexpected result can be found after careful inspection of all the individual EMD trajectories. As we will expose in detail in further publications, in few cases at the lowest projectile velocity, the dynamics of the system is consistent with a ‘classical’ mechanism where the target nuclei which gain more kinetic energy are those closer to the path of the projectile, existing also a certain deflection of the projectile which supports the usual picture of nuclear stopping in terms of elastic collisions. However, as v_{H+} increases, the behavior of the system is no longer fully consistent with that picture. On one hand, the closest nuclei to the projectile

path (with impact parameters of about 0.15 Å or even less) acquire velocities that are not consistent with a binary collision mechanism (Fig. 2). Moreover, we find several nuclei quite far from the projectile path (belonging to molecules that are first or even second neighbors of the water molecules that are directly affected by the projectile) that gain a significant kinetic energy. In all the cases, the target nuclei that gain such kinetic energy are the hydrogen ones, being the kinetic energy gained by the oxygen nuclei negligible.

Electronic density perturbations and nuclear stopping.— The fact that the highest $\langle S_n \rangle$ and $\langle S_e \rangle$ values are both obtained at the same v_{H+} range suggests that this unexpected energy transfer to the nuclei may be related to the perturbation of the target electronic subsystem during the irradiation process. The analysis of the individual simulations confirms a certain correlation between S_n and S_e , though the S_n/S_e ratio varies significantly among the different trajectories, being as high as 15–20% in some of them. In particular, S_n is higher in the cases where the projectile crosses, or passes very close to, the O–H bonds in the target. Inspection of the electronic density of the system during the irradiation process for such cases reveals that the passing of the projectile causes very severe changes in the density distribution with respect to the ground state (Fig. 3), which can be as high as a three-fold increase of the density in the close vicinity of the closest target atom.

Such perturbations of the electronic density of the water molecules along the projectile path induce further density changes that are propagated to the neighboring waters along the H-bond network, as visualized in Fig. 3. Analysis of the forces in such regions reveals that several H and O atoms therein experience extremely high forces, though only the hydrogen atoms eventually acquire meaningful kinetic energies (note that similar forces on H/O atoms imprint an acceleration on H that is 16-fold that on O, which together with the short timescale of the process implies that only hydrogen atoms can reach a significant velocity). We note in passing that this effect does not involve the oxygen’ core electrons in any way, and actually thorough electronic structure analyses that will be unfolded elsewhere show that the role of oxygen’ core electrons in the proton stopping process is much smaller than previously thought [14].

Isotopic effect in electronic-mediated nuclear stopping.— The findings above disclosed strongly suggest the possibility of an isotopic effect for the stopping of protons in light *vs.* heavy water, as the mass of the target nuclei becomes relevant for this electronic-mediated nuclear stopping to take place. Motivated by this, we performed additional irradiation simulations of protons in heavy ice, employing the same simulation settings previously used for light ice, for the three projectile velocities that showed the highest total stopping in light ice ($v_{H+} = 1.41, 1.72$ and 2.00 a.u.).




FIG. 2. Left/center: Atomic positions and velocities at the end of the simulation with the highest S_n in this work, obtained with a 75 keV proton. The scale of the vectors indicating the velocities is shown above; vectors are only drawn for atoms with velocities greater than 0.001 a.u. The initial/final positions of the projectile are shown as a green circle in the xy projection and its trajectory is shown as a green line in the yz projection; note that the projectile travels in the $-z$ direction. Right: Change of the different energy components during the same simulation with respect to their initial values.




FIG. 3. Analysis of the electronic density for the simulation with the highest S_n at an instant where the projectile (green ball) has just passed close to the target atom with the highest kinetic energy increase. Left/center panels: side/top views of the system, with the blue/yellow isosurfaces (at an isovalue of ± 0.005 e⁻/Bohr³) representing respectively the regions of density depletion/accumulation with respect the ground state, i.e. $\Delta\rho = \rho^{\text{TD}} - \rho^{\text{GS}}$, with $\Delta\rho < 0$ in blue and $\Delta\rho > 0$ in yellow. Right panels: electronic density at clipping planes passing through the said target atom (green “x” symbol) parallel to the xy/yz directions, computed at the ground state *vs.* that obtained during the TD-DFT irradiation simulation, showing also the ratio of the non-adiabatic to the ground state density. The position of the projectile is indicated as a green circle (note that the projectile is slightly offset from the clipping plane)

The results, in Fig. 4, show that the total stopping is indeed smaller in deuterated ice, while the electronic stopping is practically the same. Actually, long-standing results in the literature [15, 16] confirm the existence of an isotopic effect in the irradiation of water by protons, which has remained so far unexplored to the best of our knowledge. Indeed, the precise isotopic composition of the target is usually ignored when analyzing or comparing the electronic stopping power from different experiments or simulations, due to the underlying assumption that S_e is solely determined by the response of the electronic subsystem of the target and thus independent of the nuclear masses.

In Fig. 4 we plot the electronic stopping power for proton projectiles experimentally determined by Wenzel and Whaling using heavy water [15] and the one by Bauer *et al.* using light water [16], together with our simulation results. Remarkably, both experimental datasets basically agree in the points farther from the stopping maximum (at $v_{\text{H}^+} \sim 1$ a.u. and ~ 3.5 a.u.), but differ significantly around $v_{\text{H}^+} \sim 2$ a.u., which is precisely the region where the electronic-mediated nuclear stopping found by us is relevant. At this point, we stress that the electronic-mediated kinetic energy transfer from projectile to target nuclei that we have just described occurs without any noticeable deflection of the projec-

tile. Regarding the experimental determination of the electronic stopping power done via analyses of scattering spectra of the irradiating projectiles, this implies that the stopping mechanism that we just described is experimentally indistinguishable from the usually assumed mechanism purely based on electronic excitations. Therefore, we conclude that the ‘*electronic* stopping power’ experimentally reported for H₂O and D₂O in that regime of projectile velocities in Refs. [15, 16] is actually equivalent to the *total* stopping power determined by us in this work.

Finally, the qualitative differences between our computational results and the experimental results deserve further comments. The experimentally reported isotopic effects are greater than in our simulations, as clearly seen in Fig. 4. On one hand, this could be due to a trivial sampling issue: S_n is much more sensitive to small differences of the projectile trajectory than S_e . This suggests that our current estimation of the average $\langle S_n \rangle$ may be improved with a more intensive sampling of the possible projectile trajectories, maybe including directions not parallel to the hexagonal channel. But on the other hand and more interestingly, the presence of H/D isotopic effects always open the question of the possible role of nuclear quantum effects such as the greater quantum delocalization of H compared to D. At that respect, we note that our simulations employ classical nuclei, and of course the inclusion of nuclear quantum effects in the simulations could greatly affect the dynamical response of the H nuclei to the dramatic changes of the electronic density of the system during the irradiation.




FIG. 4. Electronic stopping power experimentally reported for H₂O [16] and D₂O [15] (respectively filled and empty diamonds) together with the average electronic and total stopping power (resp. circles/squares) found in this work from first-principles simulations for light and heavy ice (resp. red filled vs. blue empty symbols). All values are scaled to correspond to a light water density of 998 kg/m³ for an easier comparison.

Conclusions and outlook.— Our extensive Ehrenfest MD simulations, which strictly consider the atomic forces

during the irradiation process, disclosed a so far unknown channel for the energy transfer from the projectile to the target nuclei. This transfer occurs due to the dramatic changes of the electronic density of the target, which are the origin of great forces on the target nuclei during the passing of the projectile. This phenomenon is fundamentally different from other well-known processes such as the electron-phonon energy transfer that occurs after the irradiation at longer timescales. The fact that this electronic-mediated nuclear stopping occurs at the short timescale of the irradiation itself implies that only light atoms (i.e. hydrogen) experience a meaningful gain of the kinetic energy, and moreover is the reason of an isotopic effect when comparing the stopping of light vs. heavy water, in agreement with experiments that had been so far unexplained. Indeed, despite the fact that this effect occurring with no deflection of the projectile makes it rather elusive when it comes to its experimental determination, the isotopic effect experimentally described for H₂O vs D₂O is a very strong confirmation of our simulation results. Actually, the experimental data show an even stronger isotopic effect than our simulations, which could be due to the missing nuclear quantum effects in our approach. This aspect opens up an interesting –though technically challenging– avenue for further research.

We expect that this effect will be present in aqueous systems, and in particular biological tissues. Regarding the latter, this could represent a further mechanism for the creation of damage in biomolecules, which would add up to other purely electronic damage mechanisms already described [17]. On the other hand, it remains to be seen whether it will also occur in hydrogen-rich materials that lack the precise electronic structure of water and its associated H-bond network (e.g. hydrocarbon-based polymers of industrial relevance). Clearly, these findings open up a new and exciting aspect of the radiation damage processes that deserves to be explored.

This project has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 748673. This work received funding from the Research Executive Agency under the EU’s Horizon 2020 Research and Innovation program ESC2RAD (Grant ID 776410). We acknowledge PRACE for awarding us access to MareNostrum at Barcelona Supercomputing Center (BSC), Spain. The author thankfully acknowledges the computer resources at MareNostrum and the technical support provided by Barcelona Supercomputing Center (RES-FI-2018-1-0024, RES-FI-2018-3-0031, RES-FI-2019-1-0037, RES-FI-2019-2-0017, RES-FI-2019-3-0021, RES-FI-2020-3-0014, RES-FI-2021-1-0035, RES-FI-2022-1-0028, RES-FI-2022-3-0040). This work has been supported by the Madrid Government (Comunidad de Madrid-Spain) under the Multiannual Agreement with Universidad Politécnica de Madrid in the line Support for

R&D projects for Beatriz Galindo researchers, in the context of the V PRICIT (Regional Programme of Research and Technological Innovation).

* E-mail: daniel.munozsan@upm.es

- [1] J. F. Ziegler, J. P. Biersack, and M. D. Ziegler, *SRIM – The stopping and range of ions in matter* (SRIM Co., 2008).
- [2] T. D. Kühne, M. Iannuzzi, M. Del Ben, V. V. Rybkin, P. Seewald, F. Stein, T. Laino, R. Z. Khatullin, O. Schütt, F. Schiffrmann, D. Golze, J. Wilhelm, S. Chulkov, M. H. Bani-Hashemian, V. Weber, U. Borštnik, M. TAILLEFUMIER, A. S. Jakobovits, A. Lazzaro, H. Pabst, T. Müller, R. Schade, M. Guidon, S. Andermatt, N. Holmberg, G. K. Schenter, A. Hehn, A. Bussy, F. Belleflamme, G. Tabacchi, A. Glöb, M. Lass, I. Bethune, C. J. Mundy, C. Plessl, M. Watkins, J. Vandevondele, M. Krack, and J. Hutter, CP2K: An electronic structure and molecular dynamics software package - Quickstep: Efficient and accurate electronic structure calculations, *J. Chem. Phys.* **152**, 194103 (2020).
- [3] J. P. Perdew, K. Burke, and M. Ernzerhof, Generalized gradient approximation made simple, *Phys. Rev. Lett.* **77**, 3865 (1996), Erratum: *Phys. Rev. Lett.* **78**, 1396–1396 (1997).
- [4] R. Krishnan, J. S. Binkley, R. Seeger, and J. A. Pople, Self-consistent molecular orbital methods. XX. A basis set for correlated wave functions, *J. Chem. Phys.* **72**, 650 (1980).
- [5] M. J. Frisch, J. A. Pople, and J. S. Binkley, Self-consistent molecular orbital methods 25. Supplementary functions for Gaussian basis sets, *J. Chem. Phys.* **80**, 3265 (1984).
- [6] G. Lippert, J. Hutter, and M. Parrinello, The Gaussian and augmented-plane-wave density functional method for ab initio molecular dynamics simulations, *Theor. Chem. Acc.* **103**, 124 (1999).
- [7] M. Krack and M. Parrinello, All-electron ab-initio molecular dynamics, *Phys. Chem. Chem. Phys.* **2**, 2105 (2000).
- [8] V. F. Petrenko and R. W. Whitworth, *Physics of ice* (OUP Oxford, 1999).
- [9] J. F. Ziegler, M. D. Ziegler, and J. P. Biersack, SRIM–The stopping and range of ions in matter (2010), *Nucl. Instrum. Methods Phys. Res., Sect. B* **268**, 1818 (2010).
- [10] SRIM 2013 Pro, www.srim.org.
- [11] Appendix. Stopping power and range tables for charged particles, *Journal of the ICRU* **14**, 79 (2014).
- [12] M. Shimizu, M. Kaneda, T. Hayakawa, H. Tsuchida, and A. Itoh, Stopping cross sections of liquid water for MeV energy protons, *Nucl. Instrum. Methods Phys. Res., Sect. B* **267**, 2667 (2009).
- [13] M. Shimizu, T. Hayakawa, M. Kaneda, H. Tsuchida, and A. Itoh, Stopping cross-sections of liquid water for 0.3–2.0 MeV protons, *Vacuum* **84**, 1002 (2010).
- [14] Y. Yao, D. C. Yost, and Y. Kanai, K-shell core-electron excitations in electronic stopping of protons in water from first principles, *Phys. Rev. Lett.* **123**, 066401 (2019).
- [15] W. A. Wenzel and W. Whaling, The stopping cross section of D₂O ice, *Phys. Rev.* **87**, 499 (1952).
- [16] P. Bauer, W. Käferböck, and V. Nečas, Investigation of the electronic energy loss of hydrogen ions in H₂O: influence of the state of aggregation, *Nucl. Instrum. Methods Phys. Res., Sect. B* **93**, 132–136 (1994).
- [17] C. Shepard, D. C. Yost, and Y. Kanai, Electronic excitation response of DNA to high-energy proton radiation in water, *Phys. Rev. Lett.* **130**, 118401 (2023).