

Two fluid cosmological models in $f(R, T)$ theory of gravity

*Y. S. Solanke¹, **Sandhya Mhaske., +D.D. Pawar², ++V. J. Dagwal³

¹ Department of Mathematics, Mungasaji Maharaj Mahavidyalaya, Darwha, Yavatmal 445202, India

²School of Mathematical Sciences, S. R. T. M. University, Nanded, Maharashtra 431606, India

³Department of Mathematics, Government College of Engineering, Nagpur441108, India

Email:- *yadaosolanke@gmail.com, **mhaske.ss924@gmail.com, +dypawar@yahoo.com,

++vdagwal@gmail.com

Abstract:

Present work deals with the two fluid Bianchi Type-V cosmological models consisting of matter and radiating source in the $f(R, T)$ theory of gravity studied by Harko et al. (2011). In this paper, we developed a new idea about $f(R, T)$ gravity with the help of two fluids: one fluid is matter field modeling material content of the Universe and other fluid is radiation field modeling the CMB. We have determined the solution of the two fluid gravitational field equations with the systematic structure in $f(R, T)$ gravity. Here we have deliberated four types of universe such as dust universe, radiation universe, hard universe and Zel'dovich universe and also extended our work to observe the big rip and big bang singularity. We have also tested the cosmological parameters.

Keywords: Two fluid, $f(R, T)$ theory, Bianchi Type V space time.

1. Introduction:

As compare to two sequences of a single-fluid model, the cosmic evolution based on two-fluid big-bang universe appears much better. Two-fluid universe explains the change among radiation-dominated phases to a matter-dominated phase. Cosmologists have described that our present universe is expanding as well as accelerating. Ia supernovae and cosmic microwave background strongly recommend that current model is conquered by the dark energy which arises because of cosmic-accelerated expansion of the universe. Numbers of researchers have studied several aspects of two-fluid cosmological models with matter content of the universe and radiating source in Einstein's general relativity (GR) and also in alternative theory of gravity.

Cosmological models with matter and radiating source in general relativity are examined by Oli (2008). Einstein's field equation and Higher dimension cosmological universe with mixture of fluid in GR are proposed by McIntosh (1972) and Samanta (2013). Physical property of Bianchi type-II universe with

matter and radiating source in general relativity are discussed by Pant and Oli (2002). Singh *et al.* (2013) have obtained the deterministic solution of two-fluid Bianchi type-V cosmological models in GR with the support of negative constant deceleration parameter. Pawar and Dagwal (2014) have explored the physical properties of mixture of fluids with tilted cosmological universe in GR and presented various types of universe such as dust universe, radiation universe, hard universe and Zel'dovich universe. Dagwal and Pawar (2018) have extended their work of tilted cosmological universe with two-fluid sources in general relativity with assistance of a variables G and λ . Samanta (2013) has discussed kinematical properties and behavior of two-fluid sources with the help of variables G and λ in Einstein's field equation. Verma *et al.* (2015) are obtained planar cosmic space time with matter and radiating source in general relativity. Coley (1988) and Dunn (1989) have formulated Einstein's field equation with two-fluid sources of Bianchi type-VI₀ universe. Pawar *et al.* (2015) have explored several types of universe with mesonic two-fluid sources in Lyra Geometry. Harko *et al.* (2011) have discussed of kinetic model for two-fluid in gravitational field equations. Dagwal and pawar (2020) have formulated the new idea about matter and radiating source in $f(T)$ theory of gravity and discussed geometric property with help of MATLAB. Thermodynamics property of two-fluid sources is presented by Coley (1988) and Dunn (1989) has observed two-fluid gödel-type space time cosmological universe. Dagwal (2020) has studied mesonic two-fluid sources cosmic model in $f(T)$ theory of gravity. Solanke *et al.* (2021) have obtained accelerating cosmic model with mixture of fluids, first fluid show the perfect fluid and other show the dark energy.

Current astrophysical remark presented the expansion of the universe in an accelerated era. The observational data of supernovae type Ia1 and cosmic microwave background (CMB) have explored that our universe is expanding at an increasing rate. An amended theory of gravity can be employed to address some problems of current interest and may lead to some main modifications. Many researchers have studied various aspects of modified theory of gravity like as $f(T)$, $f(G)$, $f(R)$, $f(R, T)$ and $f(G, T)$ etc. In this work, we have studied $f(R, T)$ modified theory of gravity, where R is the Ricci scalar and T is the trace of the energy-momentum tensor.

Harko *et al.* (2011); Bertolami *et al.* (2008); Myrzakulov (2012); Sharif and Zubair (2012); Houndjo (2012); Adhav (2012); katore (2012); Alvarenga *et al.* (2013); Samanta (2013a), (2013b); Harko and Lobo (2014); Baffou (2015); Pawar and solanke (2015); Yousaf *et al.* (2017) Moraes and Sahoo (2017); Pradhan *et al.* (2021); Pacif *et al.* (2015), (2020); Dagwal *et al.* (2020) ; Kata's (2019a), (2019b) have developed different aspects of $f(R, T)$ gravity. Yousuf (2020a), (2020b) has discussed self-gravitating system in $f(R, T)$ gravity and electromagnetic field in modified theory of gravity. Dagwal and Pawar (2020a) (2020b) have discussed tilted and non-tilted universe in modified theory of gravity. Nagpalet *al.*

(2018) have discussed geometrical and dynamics properties of cosmological model in $f(R, T)$ modified theory of gravity.

Motivated by above work here we have studied two-fluid Bianchi Type V cosmological models with matter and radiating source in the $f(R, T)$ theory of gravity. We have tested the cosmological parameters.

This paper is organized as - Section 2: Metric and field equation, Section 3: Physical and geometrical Properties, Section 4: Cosmological parameters, Sub section 4.1: Look-back time-redshift, Sub section 4.2: Proper Distance, Sub section 4.3: Luminosity distance, Sub section 4.4: Angular-diameter distance, Sub section 8.5: Distance Modulus, Section 5: Results and discussion, Section 6: Conclusion.

2. Metric and field equation:

We have considered Bianchi type V space time of the form

$$ds^2 = -dt^2 + A^2 dx^2 + e^{2x} (B^2 dy^2 + c^2 dz^2), \quad (1)$$

where A, B, C are functions of cosmic time t alone.

The Einstein's field equation in $f(R, T)$ theory of gravity for the function

$$f(R, T) = R + 2f(T) \quad (2)$$

given as

$$R_{ij} - Rg_{ij} = T_{ij} + 2f'T_{ij} + [2pf'(T) + f(T)]g_{ij}, \quad (3)$$

where T_{ij} is the energy momentum tensor for the perfect fluid and it is given by

$$T_{ij} = T_{ij}^{(m)} + T_{ij}^{(r)}, \quad (4)$$

where $T_{ij}^{(m)}$ is energy momentum tensor for matter field having density ρ_m , pressure p_m and $u_1^{(m)} = (0, 0, 0, 1)$ as four velocities with $g^{ij}u_i^{(m)}u_j^{(m)} = 1$. $T_{ij}^{(r)}$ is energy momentum tensor for radiation field having density ρ_r , pressure p_r and $u_1^{(r)} = (0, 0, 0, 1)$ as four velocity with $g^{ij}u_i^{(r)}u_j^{(r)} = 1$ and

$$p_r = \frac{1}{3}\rho_r.$$

Thus, with above information $T_{ij}^{(m)}$ and $T_{ij}^{(r)}$ are given by

$$T_{ij}^{(m)} = (\rho_m + p_m)u_i^{(m)}u_j^{(m)} - p_m g_{ij}, \quad (5)$$

$$T_{ij}^{(r)} = \frac{4}{3}\rho_r u_i^{(m)}u_j^{(m)} - \frac{1}{3}\rho_r g_{ij}. \quad (6)$$

We have consider

$$f(T) = \lambda T. \quad (7)$$

Equation of state

$$p_m = (\gamma - 1)\rho_m \quad 1 \leq \gamma \leq 2 \quad (8)$$

Now, with the help of (5) and (6) the field equations (2) for metric (1) yield

$$\frac{B_{44}}{B} + \frac{C_{44}}{C} + \frac{B_4 C_4}{BC} - \frac{1}{A^2} = -p_m(1 + 5\lambda) - \frac{\rho_r}{3}(1 + 8\lambda) - \lambda\rho_m, \quad (9)$$

$$\frac{A_{44}}{A} + \frac{C_{44}}{C} + \frac{A_4 C_4}{AC} - \frac{1}{A^2} = -p_m(1 + 5\lambda) - \frac{\rho_r}{3}(1 + 8\lambda) - \lambda\rho_m, \quad (10)$$

$$\frac{A_{44}}{A} + \frac{B_{44}}{B} + \frac{A_4 B_4}{AB} - \frac{1}{A^2} = -p_m(1 + 5\lambda) - \frac{\rho_r}{3}(1 + 8\lambda) - \lambda\rho_m, \quad (11)$$

$$\frac{A_4 B_4}{AB} + \frac{B_4 C_4}{BC} + \frac{A_4 C_4}{AC} - \frac{3}{A^2} = p_m(2 + 7\lambda) + \rho_m(1 + 3\lambda) - \frac{\rho_r}{3}(5 + 17\lambda), \quad (12)$$

$$2\frac{A_4}{A} - \frac{B_4}{B} - \frac{C_4}{C} = 0. \quad (13)$$

Here the suffix 4 to the field variable represents the differentiation with respect to cosmic time t .

From equation (13) we have

$$A^2 = lBC,$$

Here l is constant of integration.

Without any loss of generality we can take this constant as unity, so we have,

$$A^2 = BC \quad (14)$$

Equations (9) to (13) are five independent equations with 6 unknowns. In order to find the determinate solution we use the fact that scalar expansion θ is proportional to shear scalar σ^2 , hence we get

$$B = C^k. \quad (15)$$

Where $k \neq 1$ is constant which preserves the isotropic character of the space time.

Subtracting (10) from (11)

$$\frac{B_{44}}{B} - \frac{C_{44}}{C} + \frac{A_4 B_4}{AB} - \frac{A_4 C_4}{AC} = 0. \quad (16)$$

Using (14) and (15) for solving (16), we get

$$A = b_1 T^{\frac{1}{3}}, \quad B = b_2 T^{\frac{2k}{3k+3}}, \quad C = b_3 T^{\frac{2}{3k+3}}, \quad (17)$$

where b_1, b_2, b_3 are constants and $T = t + d$ where d is constant of integration.

Thus, line element (2.1) reduced to

$$ds^2 = -dT^2 + b_1^2 T^{\frac{2}{3}} dx^2 + e^{2x} \left(b_2^2 T^{\frac{4k}{3k+3}} dy^2 + b_3^2 T^{\frac{4}{3k+3}} dz^2 \right)$$

$$ds^2 = -dT^2 + b_1^2 T^{\frac{2}{3}} dx^2 + e^{2x} T^{\frac{4}{3k+3}} (b_2^2 T^k dy^2 + b_3^2 dz^2) \quad (18)$$

3. Physical and geometrical properties

The directional Hubble parameter in the direction of x , y and z are respectively given by

$$H_1 = \frac{1}{3T}, \quad H_2 = \frac{2k}{3k+3} \frac{1}{T}, \quad H_3 = \frac{2}{3k+3} \frac{1}{T}. \quad (19)$$

The Hubble parameter H is

$$H = \frac{1}{3T}. \quad (20)$$

Average scale factor is given by

$$H = \frac{\dot{R}}{R} = \frac{1}{3T}.$$

Integrating

$$R = c'^2 T^{\frac{1}{3}}, \quad (21)$$

where c' is constant of integration.

The energy conservation equation $T_{j,i}^i = 0$ leads to two equations for matter density and radiation density

$$(7\lambda\gamma - 4\lambda + 2\gamma - 1)\rho_{m_4} + (H_1 + H_2 + H_3)(12\lambda\gamma - 10\lambda + 3\gamma - 2)\rho_m = 0, \quad (22)$$

$$(5 + 17\lambda)\rho_{r_4} + (H_1 + H_2 + H_3)(4 + 9\lambda)\rho_r = 0. \quad (23)$$

From (18), (19) and (21) we get

$$\rho_m = \eta_1 T^{-\left(\frac{12\lambda\gamma - 10\lambda + 3\gamma - 2}{7\lambda\gamma - 4\lambda + 2\gamma + 1}\right)}, \quad (24)$$

$$\rho_r = \eta_2 T^{-\left(\frac{4 + 9\lambda}{5 + 17\lambda}\right)}, \quad (25)$$

where η_1 and η_2 are constants of integration.




Fig.1a Energy density for matter against cosmic time t (Gyr)

with $\eta_1 = 0.1$, $d = 0.5$, $\gamma = 1$ and for the values of $\lambda = 1, 3, 5, 7, 9$.




Fig.1b Energy density for matter against cosmic time t (Gyr)

With $\eta_1 = 0.1$, $d = 0.5$, $\lambda = 3$ and for the values of $\gamma = 1, 4/3, 5/3, 2$.




Fig.1c Pressure for matter against cosmic time t (Gyr)

with $\eta_1 = 0.1$, $d = 0.5$, $\lambda = 3$ and for the values of $\gamma = 1, 4/3, 5/3, 2$.




Fig.2 Energy density for radiation against cosmic time t (Gyr)

with $\eta_2 = 0.1$, $d = 0.5$, $\gamma = 1$ and for the values of $\lambda = 1, 3, 5, 7, 9$.

The density parameters are

$$\Omega_m = 3\eta_1 T^{-\left(\frac{12\lambda\gamma-10\lambda+3\gamma-2}{7\lambda\gamma-4\lambda+2\gamma+1}\right)+2}, \quad (26)$$

$$\Omega_r = 3\eta_2 T^{-\left(\frac{4+9\lambda}{5+17\lambda}\right)+2} \quad (27)$$




Fig.3a Density parameter for matter against cosmic time t (Gyr)

with $\eta_1 = 0.1$, $d = 0.5$, $\gamma = 1$ and for the values of $\lambda = 1, 3, 5, 7, 9$.




Fig.3b Density parameter for matter against cosmic time t (Gyr)

with $\eta_1 = 0.1$, $d = 0.5$, $\lambda = 3$ and different values of $\gamma = 1, 4/3, 5/3, 2$.




Fig.4 Density parameter for radiation against cosmic time t (Gyr)

with $\eta_2 = 0.1$, $d = 0.5$, $\gamma = 1$ and for the values of $\lambda = 1, 3, 5, 7, 9$.

The total density parameter for the derived model is

$$\Omega = 3 \left[\eta_1 T^{-\left(\frac{12\lambda\gamma-10\lambda+3\gamma-2}{7\lambda\gamma-4\lambda+2\gamma+1}\right)+2} + \eta_2 T^{-\left(\frac{4+9\lambda}{5+17\lambda}\right)+2} \right] \quad (28)$$




Fig.5a Total density parameter against the cosmic time t (Gyr)

with $\eta_1 = 0.1$, $\eta_2 = 0.1$, $d = 0.5$, $\gamma = 1$ and for the values of $\lambda = 1, 3, 5, 7, 9$.




Fig.5b Total density parameter for matteruniverses cosmic time t (Gyr)

with $\eta_1 = 0.1$, $\eta_2 = 0.1$, $d = 0.5$, $\lambda = 3$ and for the values of $\gamma = 1, 4/3, 5/3, 2$.

Anisotropic parameter is given by

$$A_m = \frac{2}{3} \left(\frac{k-1}{k+1} \right)^2. \quad (29)$$

The scalar expansion is

$$\theta = \frac{1}{T} \quad (30)$$

and shear scalar is given by

$$\sigma^2 = \frac{(k-1)^2}{9(k+1)^2 T^2} \quad (31)$$




Fig.6 Conduct scalar expansion against time t (Gyr) with $d = 0.5$




Fig.7 Conduct of shear scalar against time t (Gyr)

for $d = 0.5$ and $k = 2.5$

The deceleration parameter is found to be

$$q = 2 \quad (32)$$

The spatial volume is

$$V = b'T, \quad (33)$$

where b' is a integration constant.




Fig.8 Conduct of spatial volume with respect to cosmic time t (Gyr)

with $d = 0.5$ and $b' = 3.1$

The profile of spatial volume verses cosmic time is represented byfigure 8 by taking the values $d = 0.5$ and $b' = 3.1$. The spatial volume is an increasing function of time and it approaches to infinity with the evolution of time. Equation (33) represents that the spatial volume is diverges for $T \rightarrow \infty$. It stops at $T \rightarrow 0$. From figure 8, the spatial volume is increasing with respect to cosmic time.

4. Cosmological parameters

The cosmological parameters are given by

4.1 Look back time-red-shift:

Cosmological red-shift z is directly proportional to size of Universe i. e. scale factor $R(t)$

$$1+z = \frac{R(t_0)}{R(t)} = \frac{R_0}{R} . \quad (34)$$

Suffix ‘0’ refers to present epoch. Here R_0 is the size of universe at the time the light from object is observed and R is the size of Universe at the time it was emitted. i.e. scale factor.

Look back t_L time is given by

$$t_L = t_0 - t(z) = \int_R^{R_0} \frac{dR}{R} \quad (35)$$

From equations (21) and (34)

$$H_0(t_0 - t) = \frac{1}{(q+1)} \left[1 - (1+z)^{-3} \right], \quad (36)$$

$$H_0(t_0 - t) = z - 2z^2 + \dots, \quad (37)$$

where H_0 is the Hubble constant at present and $(q+1)=3$. The value of Hubble constant H_0 is lies between $50-100 \text{ kms}^{-1} \text{ Mpc}^{-1}$.

Taking $z \rightarrow \infty$ in (36) we get

$$H_0(t_0 - t) = \frac{1}{(q+1)},$$

$$t_L = (t_0 - t) = \frac{1}{(q+1)H_0} = \frac{H_0^{-1}}{(q+1)},$$

$$t_L = \frac{H_0^{-1}}{(q+1)}. \quad (38)$$




Fig.9 Conduct of look-back time t_L with respect to redshift z

with $q = -0.38$ and for the value of $H_0 = 61, 62, 63, 64$.

The graph of t_L against z is presented by Figure 9 by taking $q = -0.38$ and different values of H_0 .

4.2 Proper distance:

The proper distance $d(z)$ given by

$$d(z) = r_1 R_0, \quad (39)$$

$$\text{where } r_1 = \int_t^{t_0} \frac{dt}{R}, \quad (40)$$

$$d(z) = \frac{9}{qc'^2} H_0^{-1} \left[1 - (1+z)^{-2} \right]. \quad (41)$$

When $z \rightarrow \infty$, the proper distance $d(z)$ is

$$d(z) = \frac{9}{qc'^2} H_0^{-1} \quad (42)$$




Fig.10 Conduct of proper distance $d(z)$ with respect to z

with $q = -0.38$, $c' = -1$ and for the values of $H_0 = 61, 62, 63, 64$.

The graph of proper distance $d(z)$ verses redshift z is presented by figure 10 by taking the values $q = -0.38$, $c' = -1$ and different values of H_0 .

4.3 Luminosity distance:

Luminosity distance is given by

$$d_L = r_1(z)R_0(1+z) = d(z)(1+z) \quad (43)$$

From equation (41)

$$d_L = \frac{9}{qc'^2} H_0^{-1} \left[1 - (1+z)^{-2} \right] (1+z) \quad (44)$$




Fig.11 Conduct of luminosity distance with respect to redshift z

with $q = -0.38$, $c' = -1$ and for the values of $H_0 = 61, 62, 63, 64$.

The graph of luminosity distance with respect to z is presented by figure 11 by taking the ranges $q = -0.38$, $c' = -1$ and different values of H_0 .

4.4 Angular diameter distance

The angular diameter distance is given by ratio of an object's physical transverse size to its angular size.

Angular diameter distance is given by

$$d_A = d(z)(1+z)^{-1} = d_L(1+z)^{-2} \quad (45)$$

$$d_A = \frac{9}{qc'^2} H_0^{-1} \left[1 - (1+z)^{-2} \right] (1+z)^{-1} \quad (46)$$




Fig.12 Conduct of angular diameter distance d_A with respect to redshift z

with $q = -0.38$, $c' = -1$ and for the values of $H_0 = 61, 62, 63, 64$.

The graph of d_A verses z is presented by figure 12 by taking the values $q = -0.38$, $c' = -1$ and different values of H_0 .

4.5 The distance modulus:

The distance modulus $\mu(z)$ is given by

$$\mu(z) = 5 \log d_L + 25 \quad (47)$$

$$\mu(z) = 5 \log \left\{ \frac{9}{qc'^2} H_0^{-1} \left[1 - (1+z)^{-2} \right] (1+z) \right\} + 25 \quad (48)$$




Fig.13 Conduct of distance modulus with respect to z

with $q = -0.38$, $c' = -1$ and for the values of $H_0 = 61, 62, 63, 64$.

The graph of distance modulus $\mu(z)$ versus redshift z is presented by figure 13 by taking the values $q = -0.38$, $c' = -1$ and different values of H_0 .

The Hubble parameter $H(z)$ is given by

$$H(z) = H_0(1+z)^3 \quad (49)$$




Fig.14 Conduct of Hubble parameter $H(z)$ with respect to redshift z

with various values of $H_0=61, 62, 63, 64$.

The graph of $H(z)$ verses z is presented by figure 14 by taking the different values of H_0 .

The deceleration parameter $q(z)$ is constant.

5. Results and discussion:-

The variation of ρ_m verses t is showed in Figure 1a by choosing the ranges $\eta_1 = 0.1, d = 0.5, \gamma = 1$ and for the values of $\lambda = 1, 3, 5, 7, 9$. Figure 1b shows the profile of energy density for matter ρ_m against cosmic time by choosing the values $\eta_1 = 0.1, d = 0.5, \lambda = 3$ and for the values of $\gamma = 1, 4/3, 5/3, 2$. Figure 1c represented behavior of pressure for matter against cosmic time by choosing the values $\eta_1 = 0.1, d = 0.5, \lambda = 3$ and for the values of $\gamma = 1, 4/3, 5/3, 2$. The graph of energy density for radiation ρ_r verses t is represented in Figure 2 by taking the ranges $\eta_2 = 0.1, d = 0.5, \gamma = 1$ and different values of λ . From the Figures 1a, 1b and 2 we observed that, ρ_m and ρ_r are decreasing

functions of cosmic time and they approaches towards zero with the evolution of time. From equations (24) and (25), one can show that ρ_m and ρ_r approaches infinity when $T \rightarrow 0$. Figures 1a observe that the ρ_m decline as t rises for different interval of λ and figure 1b shows that the ρ_m decline as t rises for different interval of γ . Figure 1b and 1c, respectively shows the energy density and pressure for matter of several universes such as dust ($\gamma=1$), radiation ($\gamma=4/3$), hard ($\gamma=5/3$) and Zel'dovich ($\gamma=2$) models. From equation (8) and (24) we get pressure for matter and it is constant when $\gamma=1$. The ρ_m , p_m and ρ_r are constants for large values of cosmic time. It has big rip singularity at cosmic time equal to $-d$ and big bang singularity at large values of cosmic time. The intermediate phase is between big bang at $T \rightarrow \infty$ and big rip singularity at $T=(t=-d)$.

The graph of density parameter for matter verses cosmic time is given by Figure 3a by choosing the values $\eta_1=0.1$, $d=0.5$, $\gamma=1$ and for the values of $\lambda=1, 3, 5, 7, 9$. Figure 3b presents the behavior of density parameter for matter against cosmic time by choosing the values $\eta_1=0.1$, $d=0.5$, $\lambda=3$ and $\gamma=1, 4/3, 5/3, 2$. The variation of density parameter for radiation against cosmic time is represented by Figure 4 by taking $\eta_2=0.1$, $d=0.5$, $\gamma=1$ and for the values of $\lambda=1, 3, 5, 7, 9$. Figure 3b explores the models such as $\gamma=1, 4/3, 5/3, 2$ of density parameter for radiation. From figure 4 it has been observed that the density parameter for radiation increases with increase in cosmic time increases for the chosen values of λ where as in Figure 3b the density parameter for matter for dust ($\gamma=1$), radiation ($\gamma=4/3$), hard ($\gamma=5/3$) and Zel'dovich ($\gamma=2$) models increases with increase in the cosmic time. From equations (26) and (27) it is seen that the density parameter for matter and density parameter for radiation are infinite when $T \rightarrow 0$ and constant for big range of t . The density parameter for matter and radiation have big rip singularity at cosmic time equal to $-d$ and big bang singularity at large value of cosmic time.

Figure 5a represented the behavior total density parameter against cosmic time by setting the values $\eta_1=0.1$, $\eta_2=0.1$, $d=0.5$, $\gamma=1$ and for the values of $\lambda=1, 3, 5, 7, 9$. The variation of total density parameter verses cosmic time is shows in Figure 5b by taking the values $\eta_1=0.1$, $\eta_2=0.1$, $d=0.5$, $\lambda=3$ and different values of γ . Figure 5b the gives total density parameters for different universes such as $\gamma=1, 4/3, 5/3, 2$. The total density parameter observed for the $\gamma=1, 4/3, 5/3, 2$ increases with increase in cosmic time. The total density parameter increases with increase in cosmic time for different interval of λ in figure 5a and for different values of γ in figure 5b.

The variation of scalar expansion against t is represented in Figure 6 by choosing the value $d = 0.5$. The graph of shear scalar against time by choosing the values $d = 0.5$ and $k = 2.5$ is shown in Figure 7. We observed that Figures 7 and 6 presents that the scalar expansion and shear scalar are decreasing functions of time and they approaches towards zero with the evolution of time. Equations (30) and (31) are displays that the scalar expansion and shear scalar are divergent at the initial stage and vanishes for $T \rightarrow \infty$. From figures 7 and 6, the θ and σ decreases as t increases. When $T \rightarrow 0$, they approaches towards infinity and have singularity at $T \rightarrow \infty$ and $T = (t = -d)$. Equations (30) and (31) shows the universe is expanding and shearing.

The profile of spatial volume verses cosmic time is represented byfigure 8 by taking the values $d = 0.5$ and $b' = 3.1$. The spatial volume is an increasing function of time and it approaches to infinity with the evolution of time. Equation (33) represents that the spatial volume is diverges for $T \rightarrow \infty$. It stops at $T \rightarrow 0$. From figure 8, the spatial volume is increasing with respect to cosmic time.

6. Conclusion:

In present paper, we have explored two-fluid cosmological models in $f(R, T)$ theory of gravity. In this paper, we developed a new idea about $f(R, T)$ gravity with the help of two fluids: one fluid is matter field modeling material content of the Universe and other fluid is radiation field modeling the CMB. Two-fluid universe in theory of $f(R, T)$ gravity is expanding and shearing model.

The energy density and pressure for matter with respective t are shows by Figure 1a, 1b and 1c respectively. The energy density for radiation with respective t is represented by Figure 2 by taking the ranges $\eta_2 = 0.1$, $d = 0.5$, $\gamma = 1$ and different values of λ . We observed from the Figures 1a, 1b and 2, that ρ_m and ρ_r are decreasing function of cosmic time and approaches towards zero with the evolution of time. From equation (8) and (24) we get pressure for matter and it has constant when $\gamma = 1$.

Figure 1b, 1c, 3b and 5b shows that the energy density, pressure, density parameter and total density parameter for matter presented different universe such as $\gamma = 1, 4/3, 5/3, 2$. We noted from Figure 1b and 1c, that energy density and pressure for matter for dust ($\gamma = 1$), radiation ($\gamma = 4/3$), hard ($\gamma = 5/3$) and Zel'dovich ($\gamma = 2$) models are lessening against t . Figure 3b and 5b shows the density parameter and total density parameter for matter of the universes $\gamma = 1, 4/3, 5/3, 2$ are increasing against t .

We observed that Figures 7 and 6 presented the scalar expansion and shear scalar are decreasing functions of time and approaches towards zero with the evolution of time. Equations (30) and (31) are displays that the scalar expansion and shear scalar are divergent at the initial stage and vanishes for $T \rightarrow \infty$. We noted from Figures 7 and 6 that the scalar expansion and shear scalar decreases as cosmic time increases. It has big rip singularity at cosmic time equal to $-d$ and big bang singularity at large value of cosmic time. The transitional stage is between big bang at $T \rightarrow \infty$ and big rip singularity at $T = (t = -d)$. Equation (33) represented that the spatial volume is diverge when $T \rightarrow \infty$. It is stop at $T \rightarrow 0$. Figure 8 shows the spatial volume is increasing with respective cosmic time.

The graph of look-back time t_L verses redshift z is presented by Figure 9 by taking the value $q = -0.38$ and different values of H_0 . The variation of proper distance $d(z)$, luminosity distance, angular diameter distance and distance modulus $\mu(z)$ verses redshift z is presented by Figure 10, 11, 12 and 13 respectively by setting the values $q = -0.38$, $c' = -1$ and different values of H_0 . The deceleration parameter $q(z)$ is constant.

References:

- [1] Adhav KS. LRS Bianchi type-I cosmological model in $f(R, T)$ theory of gravity. Astrophysics and space science. 2012; 339: 365-9.
- [2] Aktaş C. Massive and massless scalar field models for Kaluza–Klein universe in $f(R, T)$ gravity. Modern Physics Letters A. 2019 Apr 10; 34(11): 1950066.
- [3] Aktaş C. Various dark energy models for variables G and Λ in $f(R, T)$ modified theory. Modern Physics Letters A. 2019 Apr 30; 34(13): 1950098.
- [4] Alvarenga FG, De La Cruz-Dombriz A, Houndjo MJ, Rodrigues ME, Sáez-Gómez D. Dynamics of scalar perturbations in $f(R, T)$ gravity. Physical review D. 2013; 87(10):103526.
- [5] Baffou EH, Kpadonou AV, Rodrigues ME, Houndjo MJ, Tossa J. Cosmological viable $f(R, T)$ dark energy model: dynamics and stability. Astrophysics and Space Science. 2015 Mar; 356:173-80.
- [6] Bertolami O, Lobo FS, Páramos J. Nonminimal coupling of perfect fluids to curvature. Physical Review D. 2008 Sep 12; 78(6): 064036.
- [7] Coley A A. Thermodynamics and two-fluid cosmological models. Astrophysics and space science. 1988 ; 140: 175-89.
- [8] Dagwal V J, Pawar D D. Tilted congruences with stiff fluid cosmological models. Indian Journal of Physics. 2020; 94(11):1861-7.

- [9] Dagwal V J, Pawar D D. Tilted Two Fluids Cosmological Models with Variable G and $\{\Lambda\}$ In General Relativity. arXiv preprint arXiv:1704.07107. 2017 Apr 24.
- [10] Dagwal V J, Pawar D D. Two-fluid sources in $f(T)$ theory of gravity. Modern Physics Letters A. 2020; 35(04): 1950357.
- [11] Dagwal V J. Mesonic tilted cosmological model with wet dark fluid in $f(T)$ theory of gravity. Canadian Journal of Physics. 2020; 98(7): 636-42.
- [12] Dagwal VJ, Pawar DD, Solanke YS. Study of cosmic models in $f(R, T)$ gravity with tilted observers. Modern Physics Letters A. 2020; 35(38): 2050316.
- [13] Dagwal VJ, Pawar DD. Tilted cosmological models in $f(T)$ theory of gravity. Indian Journal of Physics. 2021 Sep; 95(9):1923-31.
- [14] Dunn K. Two-fluid cosmological models in Gödel-type space times. General relativity and gravitation. 1989; 21: 137-47.
- [15] Harko T, Lobo FS, Nojiri SI, Odintsov SD. $f(R, T)$ gravity. Physical Review D. 2011; 84(2): 024020.
- [16] Harko T, Lobo FS. Generalized curvature-matter couplings in modified gravity. Galaxies. 2014 Jul 28; 2(3):410-65.
- [17] Harko T, Lobo FS. Two-fluid dark matter models. Physical Review D. 2011 ; 83(12): 124051.
- [18] Houndjo M J. Reconstruction of $f(R, T)$ gravity describing matter dominated and accelerated phases. International Journal of Modern Physics D. 2012; 21(01):1250003.
- [19] Katore S D. Bianchi Type II VIII and IX String Cosmological Models in $f(R)$ Gravity. International Journal of Theoretical Physics. 2015; 54: 2700-11.
- [20] McIntosh C B, Foyster J M. Cosmological Models with Two Fluids. II. Conformal and Conformally Flat Metrics. Australian Journal of Physics. 1972; 25(1): 83-90.
- [21] Moraes P H, Sahoo P K. Modeling wormholes in $f(R, T)$ gravity. Physical Review D. 2017 Aug 28;96(4):044038.
- [22] Myrzakulov R. FRW cosmology in $f(R, T)$ gravity. The European Physical Journal C. 2012; 72(11): 2203.
- [23] Nagpal R, Pacif S K, Singh J K, Bamba K, Beesham A. Analysis with observational constraints in Λ -cosmology in $f(R, T)$ gravity. The European Physical Journal C. 2018; 78:1-7.
- [24] Oli S. Bianchi type I two-fluid cosmological models with a variable G and Λ . Astrophys. Space Sci. 2008; 89: 314.
- [25] Pacif S K, Khan MS, Paikroy LK, Singh S. An accelerating cosmological model from a parametrization of Hubble parameter. Modern Physics Letters A. 2020; 35(05): 2050011.

- [26] Pacif S K, Mishra B. FLRW non-singular cosmological model in general relativity. *Research in Astronomy and Astrophysics*. 2015 Dec 1; 15(12): 2141.
- [27] Pant D N, Oli S. Two-fluid Bianchi type II cosmological models. *Astrophysics and space science*. 2002; 281: 623-31.
- [28] Pawar D D, Dagwal V J, Solanke Y S. Mesonic cosmological model with two-fluid source in Lyra geometry. *International Journal of Theoretical Physics*. 2015; 54: 1926.
- [29] Pawar D D, Dagwal V J. Two fluids tilted cosmological models in general relativity. *International Journal of Theoretical Physics*. 2014; 53: 2441-50.
- [30] Pawar D D, Solanke Y. Cosmological models filled with a perfect fluid source in the $f(R, T)$ theory of gravity. *Turkish Journal of Physics*. 2015; 39(1): 54-9.
- [31] Pradhan A, Garg P, Dixit A. FRW cosmological models with cosmological constant in $f(R, T)$ theory of gravity. *Canadian Journal of Physics*. 2021; 99 (999):741-53.
- [32] Samanta G C. Kantowski-Sachs universe filled with perfect fluid in $f(R, T)$ theory of gravity. *International Journal of Theoretical Physics*. 2013; 52: 2647-56.
- [33] Samanta G C. Universe filled with dark energy (DE) from a wet dark fluid (WDF) in $f(R, T)$ gravity. *International Journal of Theoretical Physics*. 2013; 52: 2303-15.
- [34] Samanta GC. Two-Fluid Anisotropic Cosmological Model with Variable G and Λ . *International Journal of Theoretical Physics*. 2013 ; 52: 4015-24.
- [35] Samanta, G.C., Debata, S. Two-Fluid Cosmological Models in Kaluza-Klein Space Time. *Int J Theor Phys*. 2013; 52: 3999.
- [36] Sharif M, Zubair M. Thermodynamics in $f(R, T)$ theory of gravity. *Journal of Cosmology and Astroparticle Physics*. 2012; 2012(03): 028.
- [37] Singh M K, Verma M K, Ram S. Two-fluid cosmological model of Bianchi type-V with negative constant deceleration parameter. *International Journal of Theoretical Physics*. 2013; 52: 227-32.
- [38] Solanke Y S, Pawar D D, Dagwal V J. Accelerating dark energy universe with LRS Bianchi type-I space-time. *International Journal of Geometric Methods in Modern Physics*. 2021 Mar 22; 18(04): 2150062.
- [39] Verma MK, Chandel S, Ram S. Anisotropic Plane Symmetric Two-Fluid Cosmological Model with Time-Varying G and Λ . *Chinese Physics Letters*. 2015; 32(12):120401.
- [40] Yousaf Z, Zaeem-ul-Haq Bhatti M, Rafaqat A. LTB geometry with tilted and nontilted congruences in $f(R, T)$ gravity. *International Journal of Modern Physics D*. 2017 Aug 9; 26(09): 1750099.
- [41] Yousaf Z. Construction of charged cylindrical gravastar-like structures. *Physics of the Dark Universe*. 2020 May 1; 28:100509.

- [42] Yousaf Z. Definition of complexity factor for self-gravitating systems in Palatini $f(R)$ gravity. *Physica Scripta*. 2020 May 27; 95(7): 075307.

