

The extremal Kerr entropy in higher-derivative gravities

Pablo A. Cano,^a Marina David^a

^a*Instituut voor Theoretische Fysica, KU Leuven. Celestijnenlaan 200D, B-3001 Leuven, Belgium*

E-mail: pabloantonio.cano@kuleuven.be, marina.david@kuleuven.be

ABSTRACT: We investigate higher derivative corrections to the extremal Kerr black hole in the context of heterotic string theory with α' corrections and of a cubic-curvature extension of general relativity. By analyzing the near-horizon extremal geometry of these black holes, we are able to compute the Iyer-Wald entropy as well as the angular momentum via generalized Komar integrals. In the case of the stringy corrections, we obtain the physically relevant relation $S(J)$ at order α'^2 . On the other hand, the cubic theories, which are chosen as Einsteinian cubic gravity plus a new odd-parity density with analogous features, possess special integrability properties that enable us to obtain exact results in the higher-derivative couplings. This allows us to find the relation $S(J)$ at arbitrary orders in the couplings and even to study it in a non-perturbative way. We also extend our analysis to the case of the extremal Kerr-(A)dS black hole.

Contents

1	Introduction	2
1.1	Summary of results	3
2	Conserved charges	5
3	Heterotic string theory	8
3.1	The four-dimensional stringy action	8
3.2	Near-horizon geometry	9
3.3	Entropy and angular momentum	13
3.4	Adding a cosmological constant	16
4	Cubic gravity	19
4.1	Near-horizon geometries	21
4.2	Entropy and angular momentum	24
4.3	The relation $S(J)$	27
5	Discussion	31
A	Near-horizon geometry of the heterotic theory with $\Lambda \neq 0$	33
B	Noether-Komar two-form for the cubic theories	34
B.1	Proof that the ξ^0 -integral vanishes	38

1 Introduction

The investigation of higher-derivative corrections in general relativity (GR) provides us with key insights in our understanding of quantum gravity. In fact, the Einstein-Hilbert Lagrangian is considered to be just the leading term in an effective action which is expected to contain an infinite tower of higher-derivative terms. This is in particular realized in string theory [1–3], although the kind of corrections one can obtain in four dimensions depend greatly on the string type and on the compactification process. For this reason, in many instances it is useful to follow an effective field theory approach, which allows one to capture the most general correction to GR at any given order in the derivative expansion once the massless degrees of freedom have been specified.

From a theoretical standpoint, the study of higher-derivative corrections to the thermodynamic properties of black holes, like the entropy, is especially relevant. This is because black hole entropy allows one to connect the effective gravitational action with its quantum-gravitational origin whenever a microscopic description of the system is known, *e.g.*, [4–11].

However, computing higher-derivative corrections to black hole geometries, even at low orders in the derivative expansion, can be challenging. In particular, there is still a gap in understanding the solution space for rotating backgrounds. One approach that has been taken in the context of corrections to Kerr black holes is to consider an expansion around slow rotation [12–15]. This allows us to perturbatively solve for the fields of the theory at a given order in the angular momentum. However, the solution in the extremal limit cannot be captured by this approach, and one often has to restrict to numeric methods [16, 17] with the corresponding loss of analytic control.

If one is not interested in the full black hole solution but only on its thermodynamic properties, there are several strategies that can be followed. One possibility consists in evaluating the Euclidean action on the uncorrected black hole solution [18–21]. As shown by [18] the leading-order corrections to black hole thermodynamics can be obtained in this way, hence avoiding the intricate task of solving the equations of motion. In the case of extremal black holes, another approach is to investigate the solutions at a specific region of the spacetime, namely, the near-horizon geometry. What makes this favorable is that the near-horizon metric of extremal black holes has additional symmetries, mainly an enhanced $SO(2, 1)$ that does not exist in the full solution. In the case of rotating black holes in four dimensions, this becomes a $U(1)$ fibered over the $SO(2, 1)$. This enhancement of isometries is not altered by higher derivative terms and therefore we can take advantage of the near-horizon extremal geometries (NHEG) to study corrections to the black hole thermodynamic properties. In addition, the near-horizon approach can in principle be applied beyond leading-order corrections, which is one of the main motivations of this paper.

Investigating the NHEGs has shown to be quite fruitful. In fact, it has been well studied that the entropy of extremal black holes can be computed by what is now known as the entropy function formalism [7–9, 22].¹ The strength of this method precisely comes from

¹In the context of AdS/CFT, the near-horizon geometry can also be applied to study the entropy via the

the enhancement of isometries at the horizon in the extremal limit. The first examples of the entropy function were for spherically symmetric black holes [7] and later it was extended to rotating solutions [28–30]. In the static case, the equations of motion for NHEGs are reduced to a system of algebraic equations for the parameters of the solution, hence making the resolution essentially straightforward. Rotating NHEGs, on the other hand, require to solve a system of differential equations, which become ever more involved with the inclusion of higher-derivative corrections. Thus, even though the entropy function formalism provides a general framework to study these near horizon geometries, important work has to be carried out on a case-by-case basis.

On the other hand, in order to obtain a physically meaningful result, the entropy must be expressed in terms of other thermodynamic quantities of the black hole, like its charges. Therefore, the correct identification of the black hole charges from the NHEG is also crucial. In the case of the angular momentum J , a first-principle computation involves the evaluation of the Noether charge associated to Killing vector that generates rotations [31–33]. Such charge is defined as the integral of the Noether charge two-form at infinity and cannot be in principle defined on the near-horizon geometry. Here we make use of generalized Komar integrals [34–38], which are a modification of the Noether charge independent of the surface of integration, in order to provide a first-principle computation of the angular momentum from the NHEG. We note that this approach is in principle different to the one used in [28] to define the angular momentum.²

In this paper, we apply the methods discussed above to obtain the corrections to the relation $S(J)$ for extremal Kerr black holes in two relevant higher-derivative theories. The first is motivated by the effective action of heterotic string theory upon compactification on a six-torus. The other is a general effective field theory extension of GR with six-derivative terms which we choose in a specific way. In the following, we provide an overview of these theories along with our main results.

1.1 Summary of results

Heterotic string theory with α' corrections:

We consider the following string-inspired theory³

$$I_{\text{stringy}} = \frac{1}{16\pi G} \int d^4x \sqrt{|g|} \left[R - \frac{1}{2}(\partial\psi)^2 - \frac{\alpha'}{8}\psi R_{\mu\nu\rho\sigma} \tilde{R}^{\mu\nu\rho\sigma} + \frac{\alpha'}{8}\mathcal{X}_4 \right], \quad (1.1)$$

where $\alpha' = \ell_s^2$ is the square of the string length scale,

dual two-dimensional CFT [23–25], including in the near-extremal limit [26, 27].

²In fact, one must be careful with the identification of charges via the entropy function due to the existence of different notions of charge (*i.e.*, Page, Maxwell or brane-source charges [39]). Interpreting the charges correctly is especially non-trivial in the presence of higher-derivative corrections [40–42].

³Up to the topological Gauss-Bonnet term \mathcal{X}_4 , this theory is known as dynamical Chern-Simons gravity [43, 44].

$$\tilde{R}^{\mu\nu\rho\sigma} = \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} R_{\alpha\beta}{}^{\rho\sigma} \quad (1.2)$$

is the dual Riemann tensor and

$$\mathcal{X}_4 = R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} \quad (1.3)$$

is the Gauss-Bonnet density. We will argue that such theory is obtained from heterotic string theory upon dimensional reduction and stabilization of the dilaton. For this theory we are able to find explicitly the corrections to the near-horizon extremal Kerr (NHEK) metric and we obtain that the entropy of the extremal Kerr black hole is corrected according to⁴

$$S = 2\pi|J| + \alpha' \frac{\pi}{2G} - \alpha'^2 \frac{\pi}{|J|G^2} \left(\frac{493}{3360} + \frac{3\pi}{128} \right) + O(\alpha'^3). \quad (1.4)$$

Cubic gravity

We also consider the following extension of GR,

$$I_{\text{cubic}} = \frac{1}{16\pi G} \int d^4x \sqrt{|g|} \left[R + \lambda \mathcal{P} + \tilde{\lambda} \tilde{\mathcal{P}} \right], \quad (1.5)$$

where \mathcal{P} and $\tilde{\mathcal{P}}$ are the following even- and odd-parity cubic-curvature terms

$$\mathcal{P} = 12R_{\mu}{}^{\rho}{}_{\nu}{}^{\sigma} R_{\rho}{}^{\alpha}{}_{\sigma}{}^{\beta} R_{\alpha}{}^{\mu}{}_{\beta}{}^{\nu} + R_{\mu\nu}{}^{\rho\sigma} R_{\rho\sigma}{}^{\alpha\beta} R_{\alpha\beta}{}^{\mu\nu} - 12R_{\mu\nu\rho\sigma} R^{\mu\rho} R^{\nu\sigma} + 8R_{\mu}{}^{\nu} R_{\nu}{}^{\rho} R_{\rho}{}^{\mu}, \quad (1.6)$$

$$\tilde{\mathcal{P}} = 7R^{\mu\nu}{}_{\alpha\beta} R^{\alpha\beta}{}_{\rho\sigma} \tilde{R}^{\rho\sigma}{}_{\mu\nu} - \frac{9}{2} R R_{\mu\nu\rho\sigma} \tilde{R}^{\mu\nu\rho\sigma} - 20\tilde{R}^{\mu\nu\rho\sigma} R_{\mu\rho} R_{\nu\sigma}. \quad (1.7)$$

On the one hand, up to field redefinitions, this Lagrangian provides a general effective field theory extension of GR to six derivatives [15, 45]. On the other, the cubic operators are chosen in this particular form for a reason: they have special properties that allow us to perform exact (rather than perturbative) computations. In fact, the even-parity Lagrangian (1.6) is nothing but Einsteinian cubic gravity (ECG) [46]. One of the most interesting aspects about this theory is that the thermodynamic properties of its black hole solutions can be obtained analytically. Indeed, the black hole solutions of ECG have been thoroughly studied [47–54]. Generalizations of this theory at higher orders and in other dimensions exist [55–61], but so far only parity-preserving generalizations have been studied. The density (1.7) is a new odd-parity density with similar properties to ECG that we are introducing here for the first time.

For the theory (1.5) we find that the exact relation $S(J)$ can be obtained by solving a system of algebraic equations. In the perturbative limit, this yields

$$S = 2\pi|J| \left[1 - \frac{\lambda}{(GJ)^2} + \frac{20\lambda^2 + 9\tilde{\lambda}^2}{2(GJ)^4} - \frac{184\lambda^3 + 117\lambda\tilde{\lambda}^2}{2(GJ)^6} + O(J^{-8}) \right]. \quad (1.8)$$

⁴Note that the entropy is a function of the absolute value $|J|$, but throughout the paper we will assume $J > 0$ hence avoiding using absolute values.

The leading correction matches precisely with [18], after taking into account that $\lambda = 7\eta_e$. Our approach allows us to go beyond the first-order correction, and in particular our results show for the first time the effect of parity-violating terms on the entropy, which enter at order $\tilde{\lambda}^2$.

For both theories, (1.1) and (1.5), we also extend these results to the case of the extremal Kerr-(A)dS black hole by including a cosmological constant. The resulting expressions are more involved and can be found in (3.61) and (4.50).

This paper is organized as follows. In section 2, we review the procedure for defining conserved charges in gravity along with setting the conventions and notations used throughout the paper. In section 3, we delve into heterotic string theory with higher derivative corrections and compute the corrected near-horizon extremal geometry along with the thermodynamic quantities via the Komar integral and Iyer-Wald entropy. We then move to Einstein gravity with six-derivative corrections in section 4 and investigate how the angular momentum and black hole entropy are affected by the even and odd parity cubic terms. Some final remarks can be found in 5 and additional details of the computation are summarized in appendix A and appendix B.

2 Conserved charges

We review the general formalism of constructing the Noether charges associated to the Killing isometries of the gravitational action. Let us consider a general higher-curvature theory of the form⁵

$$I = \int d^D x \sqrt{|g|} \mathcal{L}(g^{\mu\nu}, R_{\mu\nu\alpha\beta}) . \quad (2.1)$$

This theory has equations of motion [62]

$$\mathcal{E}_{\mu\nu} = \frac{1}{\sqrt{|g|}} \frac{\delta I}{\delta g^{\mu\nu}} = R_{\mu}{}^{\sigma\alpha\beta} P_{\nu\sigma\alpha\beta} - \frac{1}{2} \mathcal{L} g_{\mu\nu} + 2 \nabla^\alpha \nabla^\beta P_{\mu\alpha\nu\beta} , \quad (2.2)$$

where

$$P_{\mu\nu\alpha\beta} = \frac{\partial \mathcal{L}}{\partial R^{\mu\nu\alpha\beta}} . \quad (2.3)$$

Given the existence of a Killing vector ξ^μ , one finds that the $(D-1)$ -form Noether current is given by⁶ [32, 33, 63, 64]

$$\mathbf{J}_\xi = d\mathbf{Q}_\xi + 2\epsilon_\mu \xi^\nu \mathcal{E}^\mu{}_\nu , \quad (2.4)$$

where the second term is proportional to the equations of motion and the Noether charge $(D-2)$ form reads⁷

$$\mathbf{Q}_\xi = -\epsilon_{\mu\nu} \left(\bar{P}^{\mu\nu\alpha\beta} \nabla_\alpha \xi_\beta + 2 \nabla_\beta \bar{P}^{\mu\nu\alpha\beta} \xi_\alpha \right) . \quad (2.5)$$

⁵The same results discussed here apply if the theory also contains scalar fields.

⁶We use the convention of denoting differential forms with boldface.

⁷We define $\epsilon_{\mu_1 \dots \mu_n} = \frac{1}{(D-n)!} \epsilon_{\mu_1 \dots \mu_n \nu_1 \dots \nu_{D-n}} dx^{\nu_1} \wedge \dots \wedge dx^{\nu_{D-n}}$ and $\epsilon_{0123\dots} = \sqrt{|g|} \epsilon^{0123\dots}$.

Here, the bar over the tensor $\bar{P}_{\mu\nu\alpha\beta}$ denotes explicitly that this tensor must satisfy the algebraic Bianchi identity $\bar{P}_{\mu[\nu\alpha\beta]} = 0$, and we can generically write it as

$$\bar{P}_{\mu\nu\alpha\beta} = P_{\mu\nu\alpha\beta} - P_{\mu[\nu\alpha\beta]}. \quad (2.6)$$

Note that, depending on how the derivative with respect to the Riemann tensor is defined in (2.3), the tensor $P_{\mu\nu\alpha\beta}$ may not satisfy the Bianchi identity. It turns out that the anti-symmetric part $P_{\mu[\nu\alpha\beta]}$ is irrelevant for the equations of motion (2.2) and for the Iyer-Wald entropy — see (2.17) below. However, it is crucial that we use $\bar{P}_{\mu\nu\alpha\beta}$ and not $P_{\mu\nu\alpha\beta}$ in the Noether charge.

Now, on-shell $\mathcal{E}_{\mu\nu} = 0$ and the exterior derivative of the current in (2.4) yields $d\mathbf{J}_\xi = 0$. Then, the total charge associated to ξ^μ can be found by integrating \mathbf{J}_ξ on any spatial hypersurface of the spacetime, $V_{(D-1)}$. By applying Stokes' theorem, we can then reduce the integration to one over the boundary of $V_{(D-1)}$, which can be typically considered to be the sphere at infinity $\mathbb{S}_\infty^{(D-2)}$. The total charge reads

$$Q_\xi = \int_{\mathbb{S}_\infty^{(D-2)}} \mathbf{Q}_\xi. \quad (2.7)$$

Note that this charge does not satisfy a Gauss law (*i.e.*, it is not localized), since $d\mathbf{Q}_\xi \neq 0$, so it is important that the integration is carried out at infinity. Integrating over any other surface will yield a different result. To remedy this ambiguity, we note that on-shell the Noether current can be written as

$$\mathbf{J}_\xi = -\frac{1}{2} \star \xi \mathcal{L}, \quad (2.8)$$

where \star is the Hodge dual and ξ is the Killing vector expressed as a one-form. Then, following [36, 38], we can define an improved charge $(D-2)$ -form (the Noether-Komar charge form) by

$$\tilde{\mathbf{Q}}_\xi = -\epsilon_{\mu\nu} \left(\bar{P}^{\mu\nu\alpha\beta} \nabla_\alpha \xi_\beta + 2 \nabla_\beta \bar{P}^{\mu\nu\alpha\beta} \xi_\alpha \right) - \mathbf{\Omega}, \quad (2.9)$$

where $\mathbf{\Omega}$ is a 2-form satisfying

$$d\mathbf{\Omega} = -\frac{1}{2} \star \xi \mathcal{L}. \quad (2.10)$$

Note that the local existence of this 2-form is guaranteed by the fact that $d(\star \xi \mathcal{L}) = 0$ if ξ^ν is a Killing vector. Also, $\mathbf{\Omega}$ is determined up to a closed form,

$$\mathbf{\Omega} \rightarrow \mathbf{\Omega} + \beta, \quad \text{with} \quad d\beta = 0. \quad (2.11)$$

Now, the Noether-Komar form satisfies a Gauss law,

$$d\tilde{\mathbf{Q}}_\xi = 0, \quad (2.12)$$

and hence the total Komar charge

$$\tilde{Q}_\xi = \int_\Sigma \tilde{\mathbf{Q}}_\xi \quad (2.13)$$

is independent of the choice of surface of integration Σ , as long as this is a closed co-dimension two surface homeomorphic to the sphere at infinity. Then, it is not hard to see that this Komar charge is related to the Noether charge of interest by

$$Q_\xi = \tilde{Q}_\xi + \int_{\mathbb{S}_\infty^{(D-2)}} \mathbf{\Omega}. \quad (2.14)$$

Therefore, \tilde{Q}_ξ equals the Noether charge if the integral of $\mathbf{\Omega}$ at infinity vanishes. This can always be achieved by using the ambiguity in $\mathbf{\Omega}$ in (2.11). In particular, one can always add to $\mathbf{\Omega}$ a term proportional to the volume element of the integration surface. On the other hand, the addition of an exact form to $\mathbf{\Omega}$ does not change the charge. Finally, we must remember that Stokes' theorem applies for regular p -forms. Thus, the integral is independent on the surface Σ as long as $\tilde{\mathbf{Q}}_\xi$ is regular. Therefore, when choosing $\mathbf{\Omega}$ we must also make sure that this is a regular $(D-2)$ -form everywhere. Thus, once *any* (regular) $\mathbf{\Omega}$ satisfying

$$\int_{\mathbb{S}_\infty^{(D-2)}} \mathbf{\Omega} = 0 \quad (2.15)$$

has been chosen, the Komar integral computes the Noether charge for any integration surface Σ . For example, this allows us to evaluate the charge with Σ being the horizon surface. What makes this advantageous is that it is now sufficient to find the near-horizon solution of the black hole of interest, sidestepping the difficulties of analyzing the solution in the full space-time. However, there is still the question of how one chooses an $\mathbf{\Omega}$ satisfying (2.15) by using only the near-horizon solution. Fortunately, we will see that this can be achieved unambiguously in the case of the angular momentum for stationary and axisymmetric spacetimes.

In sum, the angular momentum will be given by

$$J = -2 \int_\Sigma \left[\epsilon_{\mu\nu} \left(\bar{P}^{\mu\nu\alpha\beta} \nabla_\alpha \xi_\beta + 2 \nabla_\beta \bar{P}^{\mu\nu\alpha\beta} \xi_\alpha \right) + \mathbf{\Omega} \right], \quad \xi = \partial_\phi, \quad (2.16)$$

where here, Σ is any surface enclosing the black hole horizon, ∂_ϕ is the Killing vector that generates rotations with periodicity 2π and we added the appropriate normalization factor [65].

Besides the angular momentum, we are interested in computing the entropy of black holes. The prescription to compute the entropy over the horizon \mathcal{H} is given by the Iyer-Wald entropy formula [32, 33]

$$S = -2\pi \int_{\mathcal{H}} d^{D-2}x \sqrt{h} P^{\mu\nu\alpha\beta} \epsilon_{\mu\nu} \epsilon_{\alpha\beta}, \quad (2.17)$$

where $P^{\mu\nu\alpha\beta}$ is given by (2.3) (or equivalently in this case (2.6)), h the determinant of the induced $(D-2)$ -dimensional horizon metric and $\epsilon_{\mu\nu}$ is the binormal to the horizon, normalized according to $\epsilon_{\mu\nu} \epsilon^{\mu\nu} = -2$.

3 Heterotic string theory

3.1 The four-dimensional stringy action

As shown in [66], a dimensional reduction of the effective action of heterotic superstring theory on a 6-torus yields the following theory at first order in α'

$$I_{\text{stringy}} = \frac{1}{16\pi G} \int d^4x \sqrt{|g|} \left[R - \frac{1}{2}(\partial\varphi)^2 - \frac{1}{2}e^{2\varphi}(\partial\psi)^2 + \frac{\alpha'}{8} \left(e^{-\varphi} \mathcal{X}_4 - \psi R_{\mu\nu\rho\sigma} \tilde{R}^{\mu\nu\rho\sigma} \right) \right]. \quad (3.1)$$

Here φ is the four-dimensional dilaton, the axion ψ is the dual of the Kalb-Ramond 2-form in four dimensions, \mathcal{X}_4 is the Gauss-Bonnet density (1.3) and $R_{\mu\nu\rho\sigma} \tilde{R}^{\mu\nu\rho\sigma}$ is the Pontryagin density, with the dual Riemann tensor $\tilde{R}^{\mu\nu\rho\sigma}$ given by (1.2). The four-dimensional dilaton φ is related to the ten-dimensional one by $\varphi = 2(\hat{\varphi} - \hat{\varphi}_\infty)$, where $\hat{\varphi}_\infty$ represents its asymptotic value, so that by construction $\varphi \rightarrow 0$ at infinity.

Now, under more general setups, one expects the dilaton to be stabilized with the inclusion of a potential $V(\varphi)$ that may be sourced, for example, from D-terms and nonperturbative corrections to the superpotential [67]. The scalar potential may also be obtained from corrections to the Kahler potential and nonzero flux of p -forms. From a phenomenological point of view, the inclusion of all available effects then leads to more realistic physics [68]. On the other hand, the shift symmetry of the axion, which is associated to its origin as the dual of the Kalb-Ramond 2-form, prevents it from acquiring a mass.

It is indeed interesting to stabilize the dilaton, as the dilaton-Gauss-Bonnet sector of the theory (3.1) is known to introduce singularities in extremal rotating black holes [69, 70]. This is problematic for our purposes, since in that case the entropy and angular momentum become ill-defined, and therefore we must stabilize the dilaton in order to get a sensible result. Thus, in general we can assume that the theory (3.1) can be supplemented with a potential $V(\varphi)$, which takes the form

$$V(\varphi) \sim V_0 + \frac{1}{2}m^2\varphi^2 + \dots \quad (3.2)$$

when expanded around the vacuum expectation value of φ . The effect of the dilaton will then be suppressed by the mass scale m , and as a first approximation we just can assume that the dilaton takes a constant value $\varphi = 0$. However we need to account for the possible presence of a cosmological constant $2\Lambda = V_0$. In this scenario, the theory (3.1) reduces to

$$I_{\text{stabilized}} = \frac{1}{16\pi G} \int d^4x \sqrt{|g|} \left[R - 2\Lambda - \frac{1}{2}(\partial\psi)^2 - \frac{\alpha'}{8} \psi R_{\mu\nu\rho\sigma} \tilde{R}^{\mu\nu\rho\sigma} + \frac{\alpha'}{8} \mathcal{X}_4 \right]. \quad (3.3)$$

This is essentially dynamical Chern-Simons gravity [43, 44], except for the presence of the Gauss-Bonnet term. While this is a topological term that does not affect the equations of motion, it does make a contribution to the black hole entropy.

From the point of view of effective field theory, this action is almost the most general we can have for a shift-symmetric pseudoscalar coupled to gravity. In fact, the only four-derivative operators that could be sourced are of the form $\psi R_{\mu\nu\rho\sigma} \tilde{R}^{\mu\nu\rho\sigma}$, \mathcal{X}_4 , R^2 , $R^{\mu\nu} R_{\mu\nu}$,

$R(\partial\psi)^2$ and $(\partial\psi)^4$. The last one is subleading since the solutions we study have $\psi = \mathcal{O}(\alpha')$, and therefore this term can be neglected. On the other hand, the three terms with Ricci curvature can be reabsorbed in a redefinition of the fields and of the couplings G , Λ and α' . Thus, only the two terms $\psi R_{\mu\nu\rho\sigma} \tilde{R}^{\mu\nu\rho\sigma}$ and \mathcal{X}_4 are relevant. The only possible generalization of (3.3) would be to consider an independent coupling (different from $\alpha'/8$) for the Gauss-Bonnet term. It is straightforward to generalize our results to that case, since the only effect of the Gauss-Bonnet density is to add a constant to the entropy.

The equations of motion of this theory are given by

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + \Lambda g_{\mu\nu} = -\frac{\alpha'}{2}\nabla^\rho\nabla^\sigma\left(\tilde{R}_{\rho(\mu\nu)\sigma}\psi\right) + \frac{1}{2}\left(\partial_\mu\psi\partial_\nu\psi - \frac{1}{2}g_{\mu\nu}(\partial\psi)^2\right). \quad (3.4)$$

$$\nabla^2\psi = \frac{\alpha'}{8}R_{\mu\nu\rho\sigma}\tilde{R}^{\mu\nu\rho\sigma} \quad (3.5)$$

We will study first the near-horizon geometries of (3.3) for $\Lambda = 0$ and consider afterwards the effect of the cosmological constant, which significantly increases the complexity of the solutions.

3.2 Near-horizon geometry

The first step in our construction of the solution is to analyze the near-horizon geometry, which enjoys an enhancement of symmetries, particularly $\text{SO}(2,1) \times \text{U}(1)$ symmetry. The metric ansatz is then given by

$$ds^2 = g(x)ds_{\text{AdS}_2}^2 + \frac{1}{f(x)}dx^2 + f(x)N(x)^2(d\phi - 2ardt)^2, \quad (3.6)$$

where

$$ds_{\text{AdS}_2}^2 = -r^2dt^2 + \frac{dr^2}{r^2}, \quad (3.7)$$

is the metric of the AdS_2 space of unit radius. The parameter a is associated to the angular momentum of the black hole and x can be related to the polar coordinate $\theta \in [0, \pi]$ according to

$$x = x_0 \cos \theta \quad (3.8)$$

for a certain x_0 . In addition, the coordinate ϕ has a periodicity

$$\phi \sim \phi + \frac{2\pi}{\omega}, \quad (3.9)$$

for an undetermined constant ω .

The scalar field $\psi(x)$, the metric functions $f(x)$, $g(x)$ and $N(x)$ and the constants x_0 and ω have to be determined by solving the equations of motion. These quantities can be expressed as a power series in α' and generally we can write

$$\Psi(x) = \sum_{n=0} \alpha'^n \Psi_n(x), \quad (3.10)$$

where $\Psi = \{\psi, f, g, N\}$ are the fields of the theory and the subscript denotes the power of α' . Via the ansatz (3.6), we can find the reduced Lagrangian of the theory

$$L[\psi, f, g, N] = g(x)N(x)\mathcal{L}\Big|_{\text{ansatz}}, \quad (3.11)$$

where $g(x)N(x) = \sqrt{-g}$, from which we can derive the equations of motion for Ψ

$$\mathcal{E}_\Psi \equiv \frac{\delta L}{\delta \Psi} = \frac{\partial \mathcal{L}}{\partial \Psi} - \frac{\partial}{\partial x} \frac{\partial \mathcal{L}}{\partial \Psi'} + \frac{\partial^2}{\partial x^2} \frac{\partial \mathcal{L}}{\partial \Psi''}. \quad (3.12)$$

This is equivalent to solving the Einstein equations (3.4) and the scalar equation (3.5). We note that our ansatz contains gauge freedom, since one of the three metric functions $f(x)$, $g(x)$ and $N(x)$ can be chosen at will (corresponding to a choice of x coordinate). For the solution at hand, we consider $N(x)$ to be a constant function that is exact in all orders of α' , i.e.,

$$N(x) = 1. \quad (3.13)$$

However, $N(x)$ must be included in the Lagrangian and its associated equation \mathcal{E}_N must be computed before setting (3.13). The reason for this is technical. We find that the Euler-Lagrange equations corresponding to $g(x)$ gives the \mathcal{E}_{tt} and \mathcal{E}_{rr} components of the Einstein equations (which are proportional to each other) while $N(x)$ yields $\mathcal{E}_{\phi\phi}$ and $f(x)$, a linear combination of \mathcal{E}_{xx} and $\mathcal{E}_{\phi\phi}$. Therefore, by allowing $N(x)$ to enter into the ansatz, we extract $\mathcal{E}_{\phi\phi}$ and \mathcal{E}_{xx} , which imply the full set of Einstein equations due to the Bianchi identities. However, this is not true for other combinations. For instance solving \mathcal{E}_g and \mathcal{E}_f does not imply the full Einstein's equations, because this is equivalent to solving \mathcal{E}_{tt} and a linear combination of \mathcal{E}_{xx} and $\mathcal{E}_{\phi\phi}$. This does not guarantee that the \mathcal{E}_{xx} component is solved, since this equation always has a reduced order on account of the Bianchi identities. Thus, it is crucial to solve \mathcal{E}_{xx} and this is the reason to include $N(x)$ in our ansatz, even though we impose (3.13) eventually.

The reduced Lagrangian for the near-horizon extremal metric (3.6) is given by

$$\begin{aligned} L = & -2fgN'' + N'(-3gf' - 2fg') \\ & - \frac{1}{2}g^{-1}N(-4a^2fN^2 + 2g^2f'' + 4gf'g' + fg^2(\psi')^2 + 4fgg'' - f(g')^2 + 4g) \\ & + \alpha' \left[\frac{1}{2}ag^{-2}N\psi N'f(-16a^2fN^2 + 2g^2f'' - 2fgg'' + 3f(g')^2 + 4g) \right. \\ & + \frac{1}{2}ag^{-3}N^2\psi(gf' - fg')(-8a^2fN^2 + g^2f'' - gf'g' - fgg'' + f(g')^2 + 2g) \\ & + afg^{-1}\psi(N')^2(3gf' - fg') + N''(2af^2\psi N' + afg^{-1}N\psi(gf' - fg')) \\ & \left. + \frac{1}{2}ag^{-2}N\psi N'(3g^2(f')^2 - 6fgf'g') \right], \end{aligned} \quad (3.14)$$

plus a total derivative term, coming from the Gauss-Bonnet density, which we are discarding. When we insert the expansion (3.10) the equations of motion can be studied order by order in α' and contain subleading terms

$$\mathcal{E}_\Psi = \mathcal{E}_\Psi^{(0)} + \alpha' \mathcal{E}_\Psi^{(1)} + \alpha'^2 \mathcal{E}_\Psi^{(2)} + \dots \quad (3.15)$$

This leads to the following equations of motion $\mathcal{E}_\Psi^{(0)}$ at zeroth order

$$\begin{aligned} \mathcal{E}_\psi^{(0)} &= g_0 f_0' \psi_0' + f_0 g_0' \psi_0' + f_0 g_0 \psi_0'' = 0, \\ \mathcal{E}_g^{(0)} &= \frac{2a^2}{g_0} - g_0'' + \frac{(g_0')^2}{2g_0} - \frac{1}{2} g_0 (\psi_0')^2 = 0, \\ \mathcal{E}_f^{(0)} &= -\frac{2a^2 f_0}{g_0^2} - f_0'' - \frac{f_0' g_0'}{g_0} - \frac{f_0 g_0''}{g_0} + \frac{f_0 (g_0')^2}{2g_0^2} - \frac{1}{2} f_0 (\psi_0')^2 = 0, \\ \mathcal{E}_N^{(0)} &= \frac{6a^2 f_0}{g_0} - f_0' g_0' - 2f_0 g_0'' + \frac{f_0 (g_0')^2}{2g_0} - \frac{1}{2} f_0 g_0 (\psi_0')^2 - 2 = 0, \end{aligned} \quad (3.16)$$

where we have already imposed (3.13). Note that, indeed, the equation $\mathcal{E}_N^{(0)}$ is only of first order in f_0 , and \mathcal{E}_f is actually implied by the rest of the equations.

By setting $\psi_0 = 0$ and demanding that both $f_0(x)$ and $g_0(x)$ be even functions, we find the NHEK solution, which takes the form

$$\psi_0(x) = 0, \quad f_0(x) = \frac{a^2 - x^2}{a^2 + x^2}, \quad g_0(x) = x^2 + a^2. \quad (3.17)$$

Note that this satisfies

$$f_0(a) = f_0(-a) = 0, \quad (3.18)$$

so that $x_0 = a$ in this case. In addition, absence of a conical singularity at the poles $x = \pm x_0$ requires the following relation for the period of ϕ

$$f'(x_0) = -f'(-x_0) = -2\omega, \quad (3.19)$$

from where we get $\omega = 1/(2a)$.

We now look at the α' corrections. The scalar ψ receives corrections at first order and the equation $\mathcal{E}_\psi^{(1)}$ reads

$$0 = (a^2 - x^2) \psi_1'' - 2x \psi_1' + \frac{12a^3 (a^2 - 3x^2) (3a^2 x - x^3)}{(a^2 + x^2)^5}. \quad (3.20)$$

For a solution of ψ that is corrected at order α' , the right hand side of the Einstein equations (3.4) is $\mathcal{O}(\alpha'^2)$. This means that at order α' , only the vacuum solution is to be satisfied for f and g and therefore we can set $f_1 = g_1 = 0$ (a different choice would simply correspond to a redefinition of the integration constants of the zeroth-order solution).

Therefore, the first non-zero higher derivative corrections for $f(x)$ and $g(x)$ are at $\mathcal{O}(\alpha'^2)$. At this order, the equations of motion $\mathcal{E}_g^{(2)}$ and $\mathcal{E}_N^{(2)}$ are⁸

$$0 = \frac{1}{2(a^2 + x^2)^2} \left((a^2 + x^2) \left(- (a^2 + x^2) \left((a^2 + x^2) (\psi_1')^2 + 2g_2'' \right) + 4xg_2' - 4g_2 \right) + 2ax(x^2 - 3a^2)\psi_1'' \right), \quad (3.21)$$

$$0 = \frac{2f_2(a^2 - x^2)}{a^2 + x^2} - 2xf_2' - \frac{2g_2(3a^4 - 2a^2x^2 - x^4)}{(a^2 + x^2)^3} + \frac{2(3a^2x - x^3)g_2'}{(a^2 + x^2)^2} - \frac{2(a^2 - x^2)g_2''}{a^2 + x^2} - \frac{1}{2}(a^2 - x^2)(\psi_1')^2 + \frac{2a(x^3 - 3a^2x)^2\psi_1'}{(a^2 + x^2)^4} - \frac{2ax(3a^4 - 4a^2x^2 + x^4)\psi_1''}{(a^2 + x^2)^3}, \quad (3.22)$$

where we have imposed the zeroth order solution (3.17) and (3.13). For purposes of clarity, from now on, we define the x coordinate such that

$$x = ay, \quad (3.23)$$

and suppress the arguments of the functions, which are now functions of y . Similarly, $x_0 = ay_0$. Then, the solution to (3.20), (3.21), and (3.22) is found to be⁹

$$\psi_1 = -\frac{\arctan y}{2a^2} - \frac{y(y^4 + 2y^2 - 7)}{4a^2(y^2 + 1)^3}, \quad (3.24)$$

$$f_2 = -\frac{y(3y^2 + 43)\arctan y}{32a^4(y^2 + 1)^2} - \frac{(315y^{12} + 5495y^{10} + 21966y^8 + 31030y^6 + 51263y^4 + 963y^2 + 1192)}{3360a^4(y^2 + 1)^7}, \quad (3.25)$$

$$g_2 = \frac{5y\arctan y}{8a^2} + \frac{4095y^{10} + 19285y^8 + 35014y^6 + 35610y^4 - 5861y^2 + 241}{6720a^2(y^2 + 1)^5}, \quad (3.26)$$

where we demanded ψ_1 to be an odd function of y — since it is a pseudoscalar — and f_2 and g_2 to be even functions. In this process there is one integration constant left, but this can be seen to be equivalent to a redefinition of a , and hence we have set it to zero. The full solution is then parametrized by the parameter a that is associated with the angular momentum. The solution is regular at the poles — it does not have divergencies — which are now shifted to $x = \pm x_0 = \pm ay_0$, with

$$y_0 = 1 + \frac{\alpha'^2}{a^4} \delta y. \quad (3.27)$$

⁸As before, we do not need $\mathcal{E}_f^{(2)}$ since it is redundant.

⁹This solution (expressed in different coordinates) was already obtained in [70]. The same reference also computed the black hole entropy and expressed it in terms of a parameter claimed to be the mass. However, such identification is not justified since one cannot obtain the mass of the black hole from the NHEG, as we discuss below.

We can solve for the shift δy from

$$f(x_0) = 0 \quad \rightarrow \quad \delta y = \frac{-334 - 115\pi}{1280}, \quad (3.28)$$

which also allows us to solve for ω from the condition that no conical defects arise at the poles

$$f'(x_0) = -2\omega, \quad \rightarrow \quad \omega = \frac{1}{2a} + \alpha'^2 \left(\frac{1972 + 315\pi}{53760a^5} \right). \quad (3.29)$$

3.3 Entropy and angular momentum

Entropy

With the solution of the near-horizon geometry, we can compute the conserved charges of the theory. We consider the variation of the Lagrangian with respect to the Riemann tensor as defined in (2.6) for the Lagrangian (3.1) and we find

$$\bar{P}^{\mu\nu\rho\sigma} = \frac{1}{16\pi G} \left[g^{\mu[\rho} g^{\sigma]\nu} - \frac{\alpha'}{8} \psi \left(\tilde{R}^{\mu\nu\rho\sigma} + \tilde{R}^{\rho\sigma\mu\nu} - \tilde{R}^{\mu[\nu\rho\sigma]} - \tilde{R}^{[\rho\sigma|\mu|\nu]} \right) - \frac{\alpha'}{4} \tilde{\tilde{R}}^{\mu\nu\rho\sigma} \right], \quad (3.30)$$

where

$$\tilde{\tilde{R}}^{\mu\nu}{}_{\rho\sigma} = \frac{1}{4} \epsilon^{\mu\nu\alpha\beta} \epsilon_{\rho\sigma\lambda\tau} R_{\alpha\beta}{}^{\lambda\tau} = -R^{\mu\nu}{}_{\rho\sigma} + 4R^{\mu[\rho} \delta^{\nu]}_{[\sigma]} - R\delta^{\mu[\rho} \delta^{\nu]}_{\sigma]} \quad (3.31)$$

comes from the variation of the Gauss-Bonnet density. We note that this indeed satisfies the algebraic Bianchi identity $\bar{P}^{\mu[\nu\rho\sigma]} = 0$. For the computation of Iyer-Wald's entropy (2.17) we need the binormal to the horizon, which for the metric (3.6) is given by

$$\epsilon_{tr} = -\epsilon_{rt} = g. \quad (3.32)$$

We then have (setting $N = 1$)

$$S = -8\pi a \int dy d\phi g^2 \bar{P}^{trtr}, \quad (3.33)$$

and we only need the \bar{P}^{trtr} component, which reads

$$\bar{P}^{trtr} = \frac{1}{32\pi G} \left[-\frac{1}{g^2} + \alpha' \frac{\psi}{2a^2 g^3} \frac{d}{dy} \left(\frac{f}{g} \right) + \frac{\alpha'}{4a^2 g^2} \frac{d^2 f}{dy^2} \right]. \quad (3.34)$$

Imposing the near-horizon solution via (3.24), (3.25), and (3.26), the black hole entropy is¹⁰

$$S = \frac{a}{16G} \int dy d\phi \left(4 - \frac{\alpha'}{a^2} \frac{d^2 f}{dy^2} + \frac{2\alpha'^2 y (y^2 - 3) \arctan y}{a^4 (y^2 + 1)^3} + \frac{\alpha'^2 y^2 (y^2 - 3) (y^4 + 2y^2 - 7)}{a^4 (y^2 + 1)^6} \right), \quad (3.35)$$

¹⁰Note that the contribution from the Gauss-Bonnet is a total derivative and in order to evaluate it we only need to use that $\frac{df}{dy}(\pm y_0) = \mp 2a\omega$, imposed from the boundary conditions at the poles.

where we evaluate the integral from $-y_0 \leq y \leq y_0$ and $0 \leq \phi \leq \frac{2\pi}{\omega}$, which gives us

$$\begin{aligned} S &= \frac{\pi a}{G\omega} + \alpha' \frac{\pi}{2G} + \alpha'^2 \frac{\pi (3840\delta y + 45\pi + 64)}{3840a^3 G\omega} + \mathcal{O}(\alpha'^3) \\ &= \frac{2\pi a^2}{G} + \alpha' \frac{\pi}{2G} - \alpha'^2 \frac{\pi (805\pi + 2846)}{4480a^2 G} + \mathcal{O}(\alpha'^3), \end{aligned} \quad (3.36)$$

The values of ω from (3.29) and δy from (3.28) have been imposed in the second equal sign.

Angular Momentum

In order to obtain the physically meaningful relation $S(J)$, we must also identify the angular momentum of the solution. We start with a general Killing vector

$$\xi = \xi^0 \partial_t + \xi^3 \partial_\phi, \quad (3.37)$$

which as a 1-form reads

$$\xi = f(2a\xi^0 r - \xi^3)(2ardt - d\phi) - \xi^0 r^2 gdt. \quad (3.38)$$

The Killing vector that generates 2π rotations corresponds to the choice $\xi^0 = 0$, $\xi^3 = \omega^{-1}$, but for now we will leave these constants to be arbitrary for the sake of generality.

In order to utilize the Komar integral (2.16), we are required to compute a 2-form Ω satisfying (2.10). The Hodge dual of (3.38) reads

$$\star \xi = ag(\xi^3 dr \wedge dt \wedge dy + \xi^0 dr \wedge dy \wedge d\phi), \quad (3.39)$$

and since the Lagrangian depends only on the coordinate y , one possible solution to Ω takes the form

$$\Omega = -\frac{1}{2} agr(\xi^3 dt \wedge dy + \xi^0 dy \wedge d\phi) \mathcal{L}. \quad (3.40)$$

At this point, let us address the issue of the ambiguity in Ω . To this result we can add an exact form — particularly a term of the form $dy \wedge d\phi$ — which could contribute to the total charge. As we discussed in section 2, this ambiguity is fixed by the condition that the integral of Ω at infinity vanishes. Even though we do not know the black hole solution beyond the near-horizon region, the global geometry of a stationary and axisymmetric black hole can always be written in the form

$$ds^2 = g_{tt}dt^2 + 2g_{t\phi}dtd\phi + g_{rr}dr^2 + g_{yy}dy^2 + g_{\phi\phi}d\phi^2, \quad (3.41)$$

where all the components are functions of r and y only. Then, for the rotational Killing vector, $\xi = \xi^3 \partial_\phi$, Ω must satisfy

$$d\Omega = -\frac{1}{2} \star \xi \mathcal{L} = -\frac{1}{2} \sqrt{|g|} \mathcal{L} \xi^3 dr \wedge dt \wedge dy, \quad (3.42)$$

and hence, since $\sqrt{|g|}\mathcal{L}$ is independent of ϕ , we can choose this 2-form as

$$\Omega = -\frac{1}{2}\xi^3 dt \wedge dy \int dr \sqrt{|g|}\mathcal{L}. \quad (3.43)$$

The integral of this Ω on any constant t and r surface — including the sphere at infinity — vanishes because Ω does not have a $dx \wedge d\phi$ component. On the other hand, it is clear that in the near horizon region this choice of Ω corresponds precisely to the ξ^3 component of (3.40). Therefore, (3.40) is the correct choice in order to compute the angular momentum.

Since the Komar integral (2.16) is independent of the chosen Σ , we choose a surface of constant t and r and we find the relevant term in the Noether-Komar charge 2-form (2.9) to be given by

$$-\epsilon_{\mu\nu}\bar{P}^{\mu\nu\alpha\beta}\nabla_\alpha\xi_\beta - 2\nabla_\beta\bar{P}^{\mu\nu\alpha\beta}\xi_\alpha - \Omega = \frac{1}{16\pi G}\left[ady \wedge d\phi (r\xi^0W_0 + \xi^3W_3) + \dots\right], \quad (3.44)$$

where

$$\begin{aligned} W_0 &= \frac{a^2f}{g} + \frac{1}{2a^2}gf'' + \frac{1}{a^2}f'g' + \frac{1}{4a^2}fg(\psi')^2 + \frac{1}{a^2}fg'' - \frac{1}{4a^2g}f(g')^2 \\ &\quad + \frac{\alpha'}{24a^3g^3}\left(\psi'\left(8a^3fg(4a^2f-g) - 4afg^3f'' + 4af^2g^2g'' - 4af^2g(g')^2\right.\right. \\ &\quad \left.\left.+ 4afg^2f'g'\right) + \psi\left(g'\left(4a^3f(8a^2f-g) + 4ag^2(f')^2\right) - 4a^3gf'(8a^2f-g)\right.\right. \\ &\quad \left.\left.+ 4afg^2\left(fg^{(3)} - gf^{(3)}\right) + 8af^2(g')^3 - 12afgf'(g')^2 + g''(12afg^2f' - 12af^2gg')\right.\right. \\ &\quad \left.\left.+ f''(4afg^2g' - 4ag^3f')\right)\right), \\ W_3 &= -\frac{af}{g} + \frac{\alpha'}{24a^3g^3}\left(\psi'\left(-4a^2fg(4a^2f-g) + 2fg^3f'' - 2f^2g^2g'' + 2f^2g(g')^2 - 2fg^2f'g'\right)\right. \\ &\quad \left.+ \psi\left(g'\left(2a^2f(g-20a^2f) + g^2(f')^2\right) + 2a^2gf'(20a^2f-g) + 2fg^2\left(fg^{(3)} - fg^{(3)}\right)\right.\right. \\ &\quad \left.\left.- f^2(g')^3 + g''(3f^2gg' - 3fg^2f') + f''(fg^2g' - g^3f')\right)\right), \end{aligned}$$

where the primes indicate derivatives with respect to y and the ellipses denote the other components of the 2-form. We must then impose the solution of ψ , f and g with their corresponding α' corrections found in (3.24), (3.25) and (3.26).

The integrand (3.44) must be integrated from $-y_0 \leq y \leq y_0$ and $0 \leq \phi \leq \frac{2\pi}{\omega}$. We observe that the term proportional to ξ^0 , coming from the time-like Killing vector is proportional to r . However, by construction the Komar integral should be independent of the integration surface. In fact, one can check that the integration of the ξ^0 term in the interval $-y_0 \leq y \leq y_0$ vanishes at order α'^2 . The charge associated to ξ^0 would be the mass of the black hole, but as this result shows, it is not possible to obtain it from the near-horizon geometry. Ultimately, the reason for this follows from the fact that one cannot identify the asymptotic time coordinate from the near-horizon geometry due to its invariance under rescalings $(t, r) \rightarrow (\gamma t, r/\gamma)$.

The angular momentum is obtained from the integral proportional to ξ^3 with the choice $\xi^3 = 1/\omega$. Once the dust settles, we get the result

$$J = -2 \int_{\Sigma} \left[\epsilon_{\mu\nu} \left(\bar{P}^{\mu\nu\alpha\beta} \nabla_{\alpha} \xi_{\beta} + 2 \nabla_{\beta} \bar{P}^{\mu\nu\alpha\beta} \xi_{\alpha} \right) + \mathbf{\Omega} \right] = \frac{1}{4(G\omega^2)} - \frac{\alpha'^2 (245\pi + 437)}{17920a^4 G \omega^2} + O(\alpha'^3). \quad (3.45)$$

Using the value of ω (3.29), this can be expressed in terms of a alone

$$J = \frac{a^2}{G} - \alpha'^2 \frac{469 + 150\pi}{1920a^2 G} + O(\alpha'^3), \quad (3.46)$$

which in turn allows us to express a as a function of J ,

$$a^2 = GJ + \alpha'^2 \frac{469 + 150\pi}{1920GJ} + O(\alpha'^3), \quad (3.47)$$

The relation $S(J)$

Combining the result for the entropy (3.36) together with (3.47), we obtain the entropy as a function of the angular momentum

$$S = 2\pi J + \alpha' \frac{\pi}{2G} - \alpha'^2 \frac{\pi}{JG^2} \left(\frac{493}{3360} + \frac{3\pi}{128} \right) + O(\alpha'^3). \quad (3.48)$$

Observe that, while the first-order correction coming from the topological Gauss-Bonnet contribution is positive, the $\mathcal{O}(\alpha'^2)$ correction is negative. Thus, the higher derivative corrections from the Pontryagin density in (3.3) lead to a smaller black hole entropy.

3.4 Adding a cosmological constant

Motivated by the stabilization of the dilaton as mentioned in section 3.1, we consider the theory (3.3) with $\Lambda \neq 0$ and study how the near-horizon geometry and the conserved charges are affected.

The solution to the near-horizon geometry can still be found analytically. At leading order in α' , the functions in the metric g_0 and N take their form in (3.17) and (3.13) and the scalar is still zero. However, f_0 is now given by

$$f_0 = \frac{1 - y^2}{1 + y^2} - \frac{a^2 \Lambda (y^4 + 6y^2 - 3)}{3(1 + y^2)}, \quad (3.49)$$

where the second term on the right hand side is proportional to the cosmological constant. Moreover, y_0 — determined by $f_0(y_0) = 0$ — is no longer equal to unity and can be solved via the quartic relation

$$\Lambda = -\frac{3(y_0^2 - 1)}{a^2(y_0^4 + 6y_0^2 - 3)}. \quad (3.50)$$

It is indeed useful to express the solution in terms of y_0 instead of Λ , as we do next. The first nonzero α' corrections for ψ , f , and g all have a similar functional dependence on y and they are given by

$$\psi_1 = \frac{1}{4a^2 (y_0^4 + 6y_0^2 - 3)} \left\{ -2 \arctan y (y_0^4 + 4y_0^2 - 1) - \eta (y_0^2 - 3)^2 \operatorname{arctanh}(y\eta) \right. \\ \left. + \frac{y (7y_0^4 + 2y_0^2 + 19) - y^3 (y^2 + 2) (y_0^4 + 14y_0^2 - 11)}{(y^2 + 1)^3} \right\}, \quad (3.51)$$

$$g_2 = \frac{1}{32a^2 (y_0^4 + 6y_0^2 - 3)^2} \left\{ 2y (10y_0^8 + 29y_0^6 + 25y_0^4 + 99y_0^2 - 3) \arctan y \right. \\ \left. + 3y \frac{(y_0^2 - 3)^2}{(y_0^2 + 3)} \eta (y_0^6 + 3y_0^4 - 5y_0^2 + 9) \operatorname{arctanh}(y\eta) \right. \\ \left. + (y^2 - 1) y_0^2 (y_0^2 - 3)^2 (y_0^2 - 1) \log \left[\frac{y_0^2 + 3 - y^2 (y_0^2 - 1)}{y^2 + 1} \right] + \frac{\sum_{n=0}^5 \gamma_{g,2n} y^{2n}}{105 (y^2 + 1)^5 (y_0^2 + 3)} \right\}, \quad (3.52)$$

$$f_2 = \frac{1}{32a^4 (1 + y^2)^2 (-3 + 6y_0^2 + y_0^4)^3} \\ \times \left\{ y \arctan y \sum_{n=0}^2 \tau_{f,(2n)} y^{2n} + y \frac{(y_0^2 - 3)^2}{(y_0^2 + 3)} \eta \arctan(y\eta) \sum_{n=0}^2 \kappa_{f,2n} y^{2n} \right. \\ \left. + y_0^2 (y_0^2 - 3)^2 (y_0^2 - 1) \log \left[\frac{y_0^2 + 3 - y^2 (y_0^2 - 1)}{y^2 + 1} \right] \sum_{n=0}^2 \rho_{f,2n} y^{2n} - \frac{\sum_{n=0}^7 \gamma_{f,2n} y^{2n}}{150 (1 + y^2)^5 (y_0^2 + 3)} \right\}, \quad (3.53)$$

where we have defined

$$\eta = \sqrt{\frac{1 - y_0^2}{y_0^2 + 3}}. \quad (3.54)$$

The various constants τ , κ , ρ and γ can be found in appendix A. Moreover, the shift in y_0

(such that $f(y_0 + \alpha'^2 \delta y_0 / a^4) = 0$) due to the higher derivative terms is given by

$$\begin{aligned} \delta y_0 = & \frac{1}{(1 + y_0^2)(-3 + 6y_0^2 + y_0^4)^2} \\ & \times \left\{ \frac{1}{64(y_0^2 - 3)} \arctan y_0 \sum_{n=0}^6 \tau_{\delta,2n} y_0^{2n} + \frac{1}{32} \frac{y_0^2 - 3}{y_0^2 + 3} \eta \arctan(y_0 \eta) \sum_{n=0}^5 \kappa_{\delta,2n} y_0^{2n} \right. \\ & \left. - \frac{1}{64} y_0^2 (y_0^2 - 3) (y_0^2 - 1) \log \left[\frac{y_0^2 + 3 - y_0^2 (y_0^2 - 1)}{y_0^2 + 1} \right] \sum_{n=0}^3 \rho_{\delta,2n} y_0^{2n} - \frac{\sum_{n=0}^7 \gamma_{\delta,2n} y_0^{2n}}{6720 y_0 (y_0^2 - 3)(y_0^2 + 3)} \right\}, \end{aligned} \quad (3.55)$$

where

$$\begin{aligned} \tau_{\delta,2n} &= \{-281, 348, -505, -480, -475, -92, 13\}, \\ \kappa_{\delta,2n} &= \{-81, 81, 42, -66, -9, 1\}, \\ \rho_{\delta,2n} &= \{3, -3, 9, -1\}, \\ \gamma_{\delta,2n} &= \{-9180, -71019, -19890, -55693, -234600, -72989, -5250, 1365\}. \end{aligned} \quad (3.56)$$

With the solution at hand, we can now investigate the effect of the cosmological constant on the conserved charges. We start with the same general Killing vector (3.38) and we see that Ω takes the similar form as (3.40) except that the Lagrangian now includes a nonzero cosmological constant. Following the procedure of evaluating the Komar integral (2.16) and the Iyer-Wald entropy (2.17), the conserved charges of the theory are

$$J = \frac{3y_0 (1 - y_0^2)}{G\Lambda (3 - y_0^2)^2} + \mathcal{O}(\alpha'^2), \quad (3.57)$$

$$S = \frac{3\pi (1 - y_0^2)}{G\Lambda (3 - y_0^2)} + \mathcal{O}(\alpha'), \quad (3.58)$$

where the α' corrections take a quite involved form that we are not showing yet. Note that even in Einstein gravity one cannot easily obtain the relation $S(J)$ explicitly and it is more convenient to express the final results in terms of the parameter y_0 . Now, both S and J receive α' corrections which depend on y_0 , but we can simplify these relations by using a different parameter ζ defined as

$$y_0 = \zeta + \alpha'^2 \delta \zeta + \mathcal{O}(\alpha'^3). \quad (3.59)$$

By choosing $\delta \zeta$ appropriately, the angular momentum can be recast in the same form as in Einstein gravity (3.57), *i.e.*,

$$J = \frac{3\zeta (1 - \zeta^2)}{G\Lambda (3 - \zeta^2)^2}. \quad (3.60)$$

Using this parameter ζ , the entropy reads

$$S = \frac{3\pi(1-\zeta^2)}{G\Lambda(3-\zeta^2)} + \alpha' \frac{\pi}{2G} + \alpha'^2 \frac{\pi\Lambda}{G} \left\{ -\frac{(-3+\zeta^2)^4(1+\zeta^2)\arctan\left(\zeta\sqrt{\frac{1-\zeta^2}{3+\zeta^2}}\right)}{192\zeta(-3+6\zeta^2+\zeta^4)\sqrt{3-\zeta^2}(2+\zeta^2)} \right. \\ \left. - \frac{8925+71155\zeta^2-104972\zeta^4+75970\zeta^6+7161\zeta^8+4235\zeta^{10}+630\zeta^{12}}{20160(-3+\zeta^2)(-1+\zeta^2)(1+\zeta^2)(-3+6\zeta^2+\zeta^4)} \right. \\ \left. - \frac{(-166+675\zeta^2-1009\zeta^4+550\zeta^6+112\zeta^8+23\zeta^{10}+7\zeta^{12})\arctan\zeta}{192\zeta(-3+\zeta^2)(-1+\zeta^2)(-3+6\zeta^2+\zeta^4)} \right\} + \mathcal{O}(\alpha'^3). \quad (3.61)$$

The two expressions (3.60) and (3.61) allow us to study the relation $S(J)$ parametrically. We conclude this section by studying two limits in which the entropy can be expressed explicitly in terms of the angular momentum. For small black holes — that is, those with $J \ll 1/(G|\Lambda|)$ — we find that ζ can be expressed in terms of a series expansion in J

$$\zeta = 1 - \frac{2}{3}(JG)\Lambda - \frac{14}{9}(JG)^2\Lambda^2 + \mathcal{O}(J^3), \quad (3.62)$$

and therefore, we can express the entropy as

$$S(J) = 2\pi J + \frac{8\pi}{3}G\Lambda J^2 + \alpha' \frac{\pi}{2G} + \alpha'^2 \frac{\pi}{G} \left[-\frac{(1972+315\pi)}{13440(GJ)} + \frac{(4912+1785\pi)\Lambda}{13440} \right. \\ \left. - \frac{(18586+2625\pi)(JG)\Lambda^2}{30240} - \frac{(118432+2135\pi)(JG)^2\Lambda^3}{60480} \right] + \mathcal{O}(\alpha'^3, J^3) \quad (3.63)$$

Note that the leading terms in the α' expansion reproduces what we have computed previously in (3.48) for asymptotically flat black holes. We also note that there is a term constant in J at order $\mathcal{O}(\alpha'^2)$ that would yield a constant shift to the overall entropy of small black holes.

For large black holes, $J \gg 1/(G|\Lambda|)$, the expansion takes the form

$$\zeta = \sqrt{3} + \frac{\sqrt[4]{3}}{\sqrt{-2JG\Lambda}} - \frac{3}{4\Lambda JG} + \mathcal{O}\left(\frac{1}{J}\right)^{3/2}, \quad (3.64)$$

such that the entropy is given by

$$S(J) = \pi \left(\frac{\sqrt{12}J}{G|\Lambda|} \right)^{1/2} \left[1 - \alpha'^2 \Lambda^2 \left(\frac{983}{6720} + \frac{107\pi}{648\sqrt{3}} \right) \right] + \mathcal{O}(1). \quad (3.65)$$

This limit only exists in the case of a negative cosmological constant, as black holes in de Sitter space cannot be arbitrarily large on account of the cosmological horizon.

4 Cubic gravity

We now push forth in our analysis of higher derivative corrections by considering a different theory, namely the following six-derivative extension of GR

$$I_{\text{cubic}} = \frac{1}{16\pi G} \int d^4x \sqrt{|g|} \left\{ -2\Lambda + R + \lambda\mathcal{P} + \tilde{\lambda}\tilde{\mathcal{P}} \right\}, \quad (4.1)$$

where \mathcal{P} and $\tilde{\mathcal{P}}$ are the even-parity and odd-parity cubic densities given by (1.6) and (1.7). Also, λ and $\tilde{\lambda}$ are couplings with dimensions of length to the fourth. Clearly, this Lagrangian provides a general effective field theory extension of GR to six derivatives. In fact, one can always get rid of higher-derivative terms with Ricci curvature via field redefinitions [15, 45], so the theory can be recast into the more standard form¹¹

$$I_{\text{cubic}} = \frac{1}{16\pi G} \int d^4x \sqrt{|g|} \left\{ R - 2\Lambda + 7\lambda R_{\mu\nu}{}^{\rho\sigma} R_{\rho\sigma}{}^{\alpha\beta} R_{\alpha\beta}{}^{\mu\nu} + 7\tilde{\lambda} R^{\mu\nu}{}_{\alpha\beta} R^{\alpha\beta}{}_{\rho\sigma} \tilde{R}^{\rho\sigma}{}_{\mu\nu} + \mathcal{O}(\lambda^2, \tilde{\lambda}^2) \right\}. \quad (4.2)$$

The corrections to the thermodynamic properties of Kerr black holes in this theory (with $\Lambda = 0$) at linear order in the couplings have been studied [18]. Thus, at linear order in λ and $\tilde{\lambda}$, our results should match those of [18] with the identifications $\eta_e = 7\lambda$, $\eta_o = 7\tilde{\lambda}$. However, the most interesting aspect of the theory (4.1) is that we do not need to restrict to the perturbative regime, as we shall be able to obtain an exact result.

The equations of motion of (4.1) can be written as

$$\mathcal{E}_{\mu\nu} = G_{\mu\nu} + \Lambda g_{\mu\nu} + \lambda \mathcal{E}_{\mu\nu}^{\text{even}} + \tilde{\lambda} \mathcal{E}_{\mu\nu}^{\text{odd}} = 0, \quad (4.3)$$

where

$$\mathcal{E}_{\mu\nu}^{\text{even}} = R_{\mu}{}^{\sigma\alpha\beta} P_{\nu\sigma\alpha\beta}^{\text{even}} - \frac{1}{2} \mathcal{P} g_{\mu\nu} + 2\nabla^\alpha \nabla^\beta P_{\mu\alpha\nu\beta}^{\text{even}}, \quad (4.4)$$

$$\mathcal{E}_{\mu\nu}^{\text{odd}} = R_{\mu}{}^{\sigma\alpha\beta} P_{\nu\sigma\alpha\beta}^{\text{odd}} - \frac{1}{2} \tilde{\mathcal{P}} g_{\mu\nu} + 2\nabla^\alpha \nabla^\beta P_{\mu\alpha\nu\beta}^{\text{odd}}, \quad (4.5)$$

and where

$$P_{\text{even}}^{\mu\nu\alpha\beta} = \frac{\partial \mathcal{P}}{\partial R_{\mu\nu\alpha\beta}}, \quad P_{\text{odd}}^{\mu\nu\alpha\beta} = \frac{\partial \tilde{\mathcal{P}}}{\partial R_{\mu\nu\alpha\beta}}. \quad (4.6)$$

These tensors read explicitly

$$\begin{aligned} P_{\text{even}}^{\beta_1\beta_2\beta_3\beta_4} &= 18R^{\beta_2\alpha\beta_4\beta} R_{\alpha}{}^{\beta_1}{}_{\beta}{}^{\beta_3} - 18R^{\beta_2\alpha\beta_3\beta} R_{\alpha}{}^{\beta_1}{}_{\beta}{}^{\beta_4} + 3R_{\alpha\beta}{}^{\beta_1\beta_2} R^{\beta_3\beta_4\alpha\beta} \\ &\quad - 6R^{\beta_1\beta_3} R^{\beta_2\beta_4} 6R^{\beta_1\beta_4} R^{\beta_2\beta_3} + 12R_{\nu\sigma} \left(R^{\nu\beta_1\sigma[\beta_3} g^{\beta_4]\beta_2} - R^{\nu\beta_2\sigma[\beta_4} g^{\beta_3]\beta_1} \right) \\ &\quad + 12R^{\beta_4\rho} R_{\rho}^{[\beta_2} g^{\beta_1]\beta_3} - 12R^{\beta_3\rho} R_{\rho}^{[\beta_2} g^{\beta_1]\beta_4}, \end{aligned} \quad (4.7)$$

¹¹To arrive to this result one also must take into account that the two different Riemann cube contractions in (1.6) can be related through the identity $R_{\mu\nu}{}^{[\rho\sigma} R_{\rho\sigma}{}^{\alpha\beta} R_{\alpha\beta}{}^{\mu\nu]} = 0$.

$$\begin{aligned}
P_{\text{odd}}^{\beta_1\beta_2\beta_3\beta_4} = & \frac{7}{2} \left(R^{\beta_3\beta_4}{}_{\rho\sigma} \tilde{R}^{\rho\sigma\beta_1\beta_2} + R^{\beta_1\beta_2}{}_{\rho\sigma} \tilde{R}^{\rho\sigma\beta_3\beta_4} \right) + 7R^{\beta_3\beta_4}{}_{\rho\sigma} \tilde{R}^{\beta_1\beta_2\rho\sigma} + 7R^{\beta_1\beta_2}{}_{\rho\sigma} \tilde{R}^{\beta_3\beta_4\rho\sigma} \\
& - \frac{9}{2} g^{\beta_1[\beta_3} g^{\beta_2]\beta_3} R_{\mu\nu\rho\sigma} \tilde{R}^{\mu\nu\rho\sigma} - \frac{9}{2} R \tilde{R}^{\beta_1\beta_2\beta_3\beta_4} - \frac{9}{2} R \tilde{R}^{\beta_3\beta_4\beta_1\beta_2} \\
& - 5g^{-1/2} \epsilon^{\mu\nu\beta_1\beta_2} R_{\mu}^{[\beta_3} R_{\nu}^{\beta_4]} - 5g^{-1/2} \epsilon^{\mu\nu\beta_3\beta_4} R_{\mu}^{[\beta_1} R_{\nu}^{\beta_2]} \\
& - 10R_{\nu\sigma} \left(g^{\beta_3[\beta_1} \tilde{R}^{\beta_2]\nu\beta_4\sigma} - g^{\beta_4[\beta_1} \tilde{R}^{\beta_2]\nu\beta_3\sigma} + g^{\beta_1[\beta_3} \tilde{R}^{\beta_4]\nu\beta_2\sigma} - g^{\beta_2[\beta_3} \tilde{R}^{\beta_4]\nu\beta_1\sigma} \right), \tag{4.8}
\end{aligned}$$

and we must note that, in general, for both of these tensors, $P^{\mu[\nu\alpha\beta]} \neq 0$. As discussed in section 2, the antisymmetric part is irrelevant for the equations of motion and for the entropy. However, in order to construct the Noether charge (2.16) we will need to consider their ‘‘barred’’ versions $\bar{P}^{\mu\nu\alpha\beta}$ as in (2.6), *i.e.*, with the antisymmetric part subtracted.

4.1 Near-horizon geometries

In order to study the near-horizon geometries of the theory (4.1), it is convenient to use the ansatz [52]

$$ds_{N,f}^2 = (a^2 + x^2) \left(-r^2 dt^2 + \frac{dr^2}{r^2} \right) + \frac{dx^2}{f(x)} + N(x)^2 f(x) (d\phi + 2ardt)^2, \tag{4.9}$$

which contains a parameter a and two functions $N(x)$ and $f(x)$. The equations for these functions can be obtained equivalently from the equations of motion (4.3) or from the reduced Lagrangian

$$L[N, f] = N(x)(a^2 + x^2) \mathcal{L}|_{\text{ansatz}}. \tag{4.10}$$

In fact, the two cubic densities (1.6) and (1.7) satisfy the following property (also shared by the Einstein-Hilbert Lagrangian): when evaluated on $N(x) = 1$, the reduced Lagrangian $L[1, f]$ becomes a total derivative. In other words, this means that

$$\left. \frac{\delta L}{\delta f} \right|_{N=1} = 0, \tag{4.11}$$

so the Euler-Lagrange equation for $f(x)$ is solved for $N(x) = 1$. We can refer to this phenomenon by saying that the theories allow for ‘‘single-function’’ solutions, since we only need to determine the function $f(x)$. All of the theories satisfying this property are a subset of the family of Generalized Quasi-topological Gravities (GQTG). Those are defined as the theories allowing for single-function static and spherically symmetric solutions [55, 56] and they satisfy a similar condition to (4.11) in that case. For a subset of GQTGs, this property is extended to solutions with NUT charge [51]. Indeed, one can see that the theories satisfying (4.11) for near-horizon geometries are the same ones as the GQTGs allowing for single-function Taub-NUT solutions, which were originally studied in [51]. This is because one can transform the

near-horizon ansatz into a Taub-NUT ansatz (in this case, with a hyperbolic base space) by performing a complex rotation $t \rightarrow it$, $a \rightarrow ia$, where ψ coordinate would play the role of Euclidean time coordinate.

It was already known that ECG (1.6) belonged to this subset of GQT theories. The density (1.7) is the first example of an odd-parity Lagrangian belonging to this class. While this density is trivial for spherically symmetric solutions, it contributes to the equations of motion for rotating near-horizon geometries (and equivalently for Taub-NUT geometries).

Let us now discuss how to solve the equations of motion. The property (4.11) means that the theory (4.1) allows for solutions with $N(x) = 1$. At the level of the equations of motion, (4.11) implies that

$$\mathcal{E}_{\psi\psi}|_{N=1} = f(x)^2 \mathcal{E}_{xx}|_{N=1}, \quad (4.12)$$

so that, in fact, only one component of the equations of motion remains to be solved.¹² Furthermore, this equation takes the form of a total derivative, namely we have

$$\mathcal{E}_{\psi\psi}|_{N=1} = \frac{x^2 f(x)}{(a^2 + x^2)} \frac{d}{dx} \mathcal{E}_f, \quad (4.13)$$

where

$$\begin{aligned} \mathcal{E}_f = & x - \frac{a^2}{x} + \left(\frac{a^2}{x} + x \right) f + \Lambda \left(-\frac{a^4}{x} + 2a^2 x + \frac{x^3}{3} \right) + \lambda \left[-\frac{24f^3 (a^4 - 9a^2 x^2)}{x (a^2 + x^2)^3} \right. \\ & + \left(\frac{12f^2 (-17a^2 + x^2)}{(a^2 + x^2)^2} + \frac{12f}{a^2 + x^2} \right) f' + \left(\frac{6}{x} + \frac{12a^2 f}{a^2 x + x^3} \right) (f')^2 - 2 (f')^3 \\ & \left. + \left(-\frac{12f}{x} + \frac{f^2 (48a^2 - 12x^2)}{a^2 x + x^3} + 6f f' \right) f'' \right] \\ & + \tilde{\lambda} \left[-\frac{72a f^2}{(a^2 + x^2)^2} + \frac{8f^3 (37a^3 - 9a x^2)}{(a^2 + x^2)^3} + \left(\frac{84f^2 (-a^3 + a x^2)}{x (a^2 + x^2)^2} + \frac{12a f}{a^2 x + x^3} \right) f' \right. \\ & \left. - \frac{12a f (f')^2}{a^2 + x^2} - \frac{6a (f')^3}{x} + \left(-\frac{36a f^2}{a^2 + x^2} + \frac{18a f f'}{x} \right) f'' \right]. \end{aligned} \quad (4.14)$$

Therefore, we have the equation

$$\mathcal{E}_f = n, \quad (4.15)$$

where n is an integration constant. Note that for $\lambda = \tilde{\lambda} = 0$, this equation can be solved right away to obtain the solution in Einstein gravity,

$$N_0(x) = 1, \quad f_0(x) = \frac{a^2 + nx - x^2 + \Lambda (a^4 - 2a^2 x^2 - x^4/3)}{(a^2 + x^2)}. \quad (4.16)$$

¹²By using the Bianchi identities one can show that the rest of the components of the EOMs are proportional to \mathcal{E}_{xx} , $\mathcal{E}_{\psi\psi}$ and their derivatives, so there are no more independent equations.

This is the near-horizon geometry of the extremal Kerr-NUT-(A)dS solution, where n is the NUT charge. Note that n breaks the equatorial symmetry $x \leftrightarrow -x$ and as a consequence the horizon is non-smooth, as it contains a conical defect at least at one of the poles — we will be more precise about this in a moment.

Here we are interested only in solutions without NUT charge and hence with a smooth horizon, so in the case of Einstein gravity we must set $n = 0$. The same applies for the even-parity cubic Lagrangian. However, the parity-breaking Lagrangian necessarily breaks the equatorial symmetry and we will find that a non-vanishing $n = n_{\text{smooth}}$ is necessary in order to obtain a smooth horizon in that case. This should not be interpreted as a NUT charge, though; instead the NUT charge would be measured by the difference $n - n_{\text{smooth}}$.

Boundary conditions

The equation (4.14) cannot be solved analytically when $\lambda, \tilde{\lambda} \neq 0$. One way to go around this consists in solving the equation perturbatively by performing an expansion in the higher-derivative couplings, analogous to what we did in section 3 for the theory (1.1). However, it turns out that we can obtain exact information about the solution by studying the boundary conditions that need to be imposed.

Since the solution must contain a compact horizon, the domain of x must be compact. Thus, $f(x)$ must vanish at two different points

$$f(x_1) = f(x_2) = 0, \quad x_2 > x_1, \quad (4.17)$$

which represent the north and south poles of the horizon. Furthermore, we want these poles to be smooth, *i.e.*, free of conical defects. At this point we must take into account that ϕ is a periodic variable. Its period is a priori undetermined so let us write it as

$$\phi \sim \phi + \frac{2\pi}{\omega}, \quad (4.18)$$

for a certain parameter $\omega > 0$. Then one can check that smoothness at the poles is achieved if

$$f'(x_1) = 2\omega, \quad f'(x_2) = -2\omega. \quad (4.19)$$

We additionally assume that the solution is analytic around the poles, and therefore it is given by series expansions of the form

$$f(x) = \begin{cases} +2\omega(x - x_1) + \sum_{k=2}^{\infty} a_k(x - x_1)^k, & x \sim x_1, \\ -2\omega(x - x_2) + \sum_{k=2}^{\infty} b_k(x - x_2)^k, & x \sim x_2, \end{cases} \quad (4.20)$$

for certain coefficients a_k and b_k . We can now insert these expansions into the equation (4.15) and solve order by order in $(x - x_i)^k$ in each case

$$\mathcal{E}_f = \sum_{k=0}^{\infty} \mathcal{E}_{f,i,k}(x - x_i)^k, \quad i = 1, 2. \quad (4.21)$$

This yields an infinite system of algebraic equations $\mathcal{E}_{f,i,k} = 0$ involving the parameters of the solution and the coefficients a_k and b_k . The first two equations, corresponding to $\mathcal{E}_{f,i,0} = 0$ and $\mathcal{E}_{f,i,1} = 0$, read as follows

$$\begin{aligned}
x_1 - \frac{a^2}{x_1} + \Lambda \left(\frac{x_1^3}{3} + 2a^2 x_1 - \frac{a^4}{x_1} \right) + \frac{8\lambda\omega^2(3 - 2x_1\omega)}{x_1} - \frac{48a\tilde{\lambda}\omega^3}{x_1} &= n, \\
x_2 - \frac{a^2}{x_2} + \Lambda \left(\frac{x_2^3}{3} + 2a^2 x_2 - \frac{a^4}{x_2} \right) + \frac{8\lambda\omega^2(3 + 2x_2\omega)}{x_2} + \frac{48a\tilde{\lambda}\omega^3}{x_2} &= n, \\
\frac{(a^2 + x_1^2)(1 + 2x_1\omega)}{x_1^2} + \frac{\Lambda(a^2 + x_1^2)^2}{x_1^2} &+ \frac{24\lambda\omega^2(x_1^2 + a^2(4x_1\omega - 1))}{x_1^2(a^2 + x_1^2)} + \frac{48a\tilde{\lambda}\omega^2(a^2\omega - x_1^2\omega + x_1)}{x_1^2(a^2 + x_1^2)} = 0, \\
\frac{(a^2 + x_2^2)(1 - 2x_2\omega)}{x_2^2} + \frac{\Lambda(a^2 + x_2^2)^2}{x_2^2} &+ \frac{24\lambda\omega^2(x_2^2 - a^2(4x_2\omega + 1))}{x_2^2(a^2 + x_2^2)} - \frac{48a\tilde{\lambda}\omega^2(a^2\omega - x_2^2\omega - x_2)}{x_2^2(a^2 + x_2^2)} = 0.
\end{aligned} \tag{4.22}$$

These are four equations for the five parameters x_1 , x_2 , n , ω and a . Therefore, they determine four of the parameters in terms of the remaining one, for instance, a . Naturally, the remaining free parameter will be related to the angular momentum. Note that in the case of $\tilde{\lambda} = 0$ we have $n = 0$ and $x_1 = -x_2$, so that the problem is reduced to a system of two equations for three variables.

A very remarkable property about these four equations is that they do not involve the parameters a_k and b_k . These start appearing in $\mathcal{E}_{f,i,k}$ with $k \geq 2$ and when solving these equations one finds that the expansions in (4.20) are entirely determined by the first parameter, a_2 or b_2 (thus, there is only one integration constant in each case). These must be chosen so that the two solutions can be glued producing a complete solution which is regular at both poles. Such solution could in principle be obtained numerically via the shooting method, but however we will not need it for our purposes. As we show next, we can obtain the entropy and the angular momentum even if we do not know the solution for $f(x)$ explicitly.

4.2 Entropy and angular momentum

Entropy

In order to compute the black hole entropy we apply Iyer-Wald's formula (2.17), which for the theory (4.1) reads

$$S = \frac{1}{4G} \int d^2x \sqrt{h} \left[1 - \frac{\lambda}{2} P_{\text{even}}^{\mu\nu\alpha\beta} \epsilon_{\mu\nu} \epsilon_{\alpha\beta} - \frac{\tilde{\lambda}}{2} P_{\text{odd}}^{\mu\nu\alpha\beta} \epsilon_{\mu\nu} \epsilon_{\alpha\beta} \right], \tag{4.23}$$

where the tensors $P_{\text{even}}^{\mu\nu\alpha\beta}$ and $P_{\text{odd}}^{\mu\nu\alpha\beta}$ are those in (4.7) and (4.8). For the near-horizon geometry (4.9) with $N(x) = 1$ the binormal has components

$$\epsilon_{tr} = -\epsilon_{rt} = a^2 + x^2, \quad (4.24)$$

and the integral above becomes

$$S = \frac{\pi}{2\omega G} \int_{x_1}^{x_2} dx \left[1 - 2\lambda(a^2 + x^2)^2 P_{\text{even}}^{trtr} - 2\tilde{\lambda}(a^2 + x^2)^2 P_{\text{odd}}^{trtr} \right], \quad (4.25)$$

where we used the fact that $\sqrt{h} = 1$ and integrated over the ϕ coordinate, which has periodicity $2\pi/\omega$. Remarkably, we find that this integrand takes the form of a total derivative. In particular, we have

$$(a^2 + x^2)^2 P_{\text{even}}^{trtr} = \frac{d}{dx} \left[\frac{3xf'(x)^2}{a^2 + x^2} + \frac{12a^2 f(x)f'(x)}{(a^2 + x^2)^2} - \frac{12a^2 xf(x)^2}{(a^2 + x^2)^3} \right], \quad (4.26)$$

$$(a^2 + x^2)^2 P_{\text{odd}}^{trtr} = \frac{d}{dx} \left[\frac{3af'(x)^2}{(a^2 + x^2)} - \frac{12axf(x)f'(x)}{(a^2 + x^2)^2} - \frac{12a^3 f(x)^2}{(a^2 + x^2)^3} \right], \quad (4.27)$$

and therefore it is possible to carry out the integration explicitly even if we do not know what $f(x)$ is. Furthermore, using the boundary conditions (4.17) and (4.19) we have the following result

$$S = \frac{\pi}{2\omega G} \left[x_2 - x_1 + 24\lambda\omega^2 \left(\frac{x_1}{a^2 + x_1^2} - \frac{x_2}{a^2 + x_2^2} \right) + 24\tilde{\lambda}\omega^2 \left(\frac{a}{a^2 + x_1^2} - \frac{a}{a^2 + x_2^2} \right) \right]. \quad (4.28)$$

In the limit that the $x_2 \rightarrow -x_1$, we find that the contribution coming from the parity-violating term vanishes, as expected. This is an exact expression of the entropy in terms of the various parameters of the solution, but this is still not meaningful as our goal is to obtain the expression of S in terms of the angular momentum J .

Angular momentum

We compute the angular momentum from the Noether-Komar integral (2.16), where in this case

$$\bar{P}^{\mu\nu\rho\sigma} = \frac{1}{16\pi G} \left[g^{\mu[\rho} g^{\sigma]\nu} + \lambda \bar{P}_{\text{even}}^{\mu\nu\rho\sigma} + \tilde{\lambda} \bar{P}_{\text{odd}}^{\mu\nu\rho\sigma} \right], \quad (4.29)$$

where we recall that the bar denotes that we subtract the antisymmetric part in the last three indices. The Killing vector we should consider is $\xi = \omega^{-1}\partial_\phi$, since $\omega\phi$ is the angular coordinate with 2π period. However, it will be useful to compute instead the Komar charge for a general Killing vector

$$\xi = \xi^0 \partial_t + \xi^3 \partial_\phi, \quad (4.30)$$

where ξ^0 and ξ^3 are constants. This will allow us to perform a consistency check of our results and will help us perform the integration for the angular momentum.

The first step is to obtain the two-form Ω . For the metric (4.9) and the Killing vector (4.30), we have

$$d\Omega = \frac{1}{2} (\xi^0 dr \wedge d\phi \wedge dx + \xi^3 dt \wedge dr \wedge dx) (a^2 + x^2) \mathcal{L}. \quad (4.31)$$

Since the on-shell Lagrangian is only a function of x , an obvious solution of this equation is

$$\Omega = -\frac{1}{2} (\xi^0 dx \wedge d\phi + \xi^3 dt \wedge dx) r(a^2 + x^2) \mathcal{L}. \quad (4.32)$$

Now, for the same reasons discussed in section 3 below (3.40), we find that this is the appropriate choice of Ω in order to compute the angular momentum.

Let us also take note that, because the theory (4.1) belongs to the GQT class, the Lagrangian takes explicitly the form of a total derivative for any choice of $f(x)$. In fact, we have

$$(a^2 + x^2) \mathcal{L} = \frac{1}{16\pi G} \frac{d\mathcal{I}}{dx}, \quad (4.33)$$

where

$$\begin{aligned} \mathcal{I}(x) = & -(a^2 + x^2) f' - 2fx - 2x - 2\Lambda \left(a^2 x + \frac{x^3}{3} \right) + \lambda \left[\frac{(10a^2 - 2x^2) (f')^3}{a^2 + x^2} \right. \\ & + (f')^2 \left(\frac{12fx(x^2 - 9a^2)}{(a^2 + x^2)^2} + \frac{12x}{a^2 + x^2} \right) + f' \left(\frac{48a^2 f}{(a^2 + x^2)^2} - \frac{168f^2 (a^4 - a^2 x^2)}{(a^2 + x^2)^3} \right) \\ & + \frac{48f^3 (5a^4 x - a^2 x^3)}{(a^2 + x^2)^4} - \frac{48a^2 f^2 x}{(a^2 + x^2)^3} \left. \right] + \tilde{\lambda} \left[-\frac{12ax (f')^3}{a^2 + x^2} \right. \\ & + (f')^2 \left(\frac{f(48ax^2 - 72a^3)}{(a^2 + x^2)^2} + \frac{12a}{a^2 + x^2} \right) + f' \left(\frac{48f^2 (6a^3 x - ax^3)}{(a^2 + x^2)^3} - \frac{48afx}{(a^2 + x^2)^2} \right) \\ & \left. - \frac{48a^3 f^2}{(a^2 + x^2)^3} + \frac{32f^3 (4a^5 - 5a^3 x^2)}{(a^2 + x^2)^4} \right]. \end{aligned} \quad (4.34)$$

The computation of the rest of the charge 2-form (2.16) — using the tensor (4.29) — is straightforward. The full expression is lengthy and we show it in appendix B. In order to obtain the conserved charge, we integrate it over a surface of constant t and r , and hence we only need the $dx \wedge d\psi$ component. The result takes the form

$$J = \frac{1}{8\pi G} \int dx d\phi (r \xi^0 F(x) + \xi^3 G(x)), \quad (4.35)$$

where the functions $F(x)$ and $G(x)$ are given by (B.7) and (B.8). Note that the term proportional to ξ^0 is also proportional to the radius of integration r . However, by construction, the Komar integral should be independent of the surface of integration. Thus, the only possibility is that this term gives zero contribution to the charge and its integral vanishes. In

appendix B.1, we show that this is indeed the case. The proof only requires using the equations of motion as well as the boundary conditions of the solution. As we already observed in section 3.3, this means that we cannot obtain the mass of the black hole from its near-horizon geometry.

As the integral is independent of r , we are free to fix it to a value that facilitates the integration. In fact, we observe that, for

$$r\xi^0 = \frac{\xi^3}{2a}, \quad (4.36)$$

the integrand becomes an explicit total derivative. We have

$$F(x) + \frac{1}{2a}G(x) = \frac{d\mathcal{J}(x)}{dx}, \quad (4.37)$$

where

$$\begin{aligned} \mathcal{J}(x) = & -\frac{fx}{2a} - \frac{(a^2+x^2)f'}{4a} - \frac{\Lambda x(3a^2+x^2)}{6a} - \lambda \left[\frac{12af^3x(-5a^2+x^2)}{(a^2+x^2)^4} + \frac{42af^2(a^2-x^2)f'}{(a^2+x^2)^3} \right. \\ & + \frac{3f(9a^2x-x^3)(f')^2}{a(a^2+x^2)^2} + \left. \frac{(-5a^2+x^2)(f')^3}{2a(a^2+x^2)} \right] - \tilde{\lambda} \left[-\frac{8a^2f^3(4a^2-5x^2)}{(a^2+x^2)^4} \right. \\ & + \left. \frac{12f^2x(-6a^2+x^2)f'}{(a^2+x^2)^3} + \frac{6f(3a^2-2x^2)(f')^2}{(a^2+x^2)^2} + \frac{3x(f')^3}{a^2+x^2} \right]. \end{aligned} \quad (4.38)$$

Therefore, by performing the integration in (4.35), we find that the angular momentum reads

$$J = \frac{1}{4\omega^2 G} [\mathcal{J}(x_2) - \mathcal{J}(x_1)], \quad (4.39)$$

where we used that $\xi^3 = 1/\omega$ for the properly normalized rotational Killing vector. Finally, we can evaluate $\mathcal{J}(x_{1,2})$ by using the boundary conditions (4.17) and (4.19), and we find

$$\begin{aligned} J = & -\frac{\Lambda(x_2-x_1)(3a^2+x_1^2+x_1x_2+x_2^2)}{24a\omega^2 G} + \frac{2a^2+x_1^2+x_2^2}{8aG\omega} \\ & - \frac{2\lambda\omega(5a^4+2a^2(x_1^2+x_2^2)-x_1^2x_2^2)}{aG(a^2+x_1^2)(a^2+x_2^2)} + \frac{6\tilde{\lambda}\omega(x_1+x_2)(a^2+x_1x_2)}{G(a^2+x_1^2)(a^2+x_2^2)}. \end{aligned} \quad (4.40)$$

4.3 The relation $S(J)$

By combining the result (4.40) together with the constraint equations (4.22), all the constants a , x_1 , x_2 , ω , n , and therefore, the full solution, is determined by the angular momentum — up to the possible existence of several branches of solutions. In turn, using (4.28), this defines implicitly the entropy as a function of the angular momentum $S(J)$. What is remarkable about our approach is that the results so far have involved no approximation whatsoever.




Figure 1: The relation $S(J)$ in the presence of the even-parity cubic correction. Left: $\lambda > 0$, right: $\lambda < 0$. The GR prediction $S = 2\pi|J|$ is shown for comparison.

The six equations (4.40), (4.28) and (4.22) provide the exact result for the relationship $S(J)$ in the theory (4.1). Of course, we still need to solve these equations if we want to obtain an explicit result. Let us therefore study several limits of interest.

Asymptotically flat black holes

Let us study first the case of $\Lambda = 0$. The most relevant solutions correspond to those with large angular momentum, or more precisely, those with $J \gg \lambda^{1/2}/G$. This is the regime where the higher-derivative corrections are perturbatively small and the theory (4.1) makes sense as an effective field theory. The result in this case can be expressed as a power series in $1/J$ (equivalently as a power series in λ and $\tilde{\lambda}$) and the few first terms read

$$S = 2\pi|J| \left[1 - \frac{\lambda}{(GJ)^2} + \frac{20\lambda^2 + 9\tilde{\lambda}^2}{2(GJ)^4} - \frac{184\lambda^3 + 117\lambda\tilde{\lambda}^2}{2(GJ)^6} + O(J^{-8}) \right]. \quad (4.41)$$

Importantly, the first order correction $\mathcal{O}(\lambda)$ agrees with the result by Reall and Santos [18] once we notice that $\lambda = \eta_e/7$. Our method is able to compute the subleading corrections as well, and in particular, we observe that the parity-breaking density does affect the entropy starting at order $\mathcal{O}(\tilde{\lambda}^2)$.

We can go beyond the perturbative result by numerically solving the system of equations (4.22) for finite values of the couplings. For simplicity, we focus on the parity-preserving correction so we set $\tilde{\lambda} = 0$. Also, since there is only one scale in the problem, we can introduce the entropy and angular momentum in “natural units” $\hat{S} = S \times G|\lambda|^{-1/2}$ and $\hat{J} = JG|\lambda|^{-1/2}$, so that the relation $\hat{S}(\hat{J})$ is independent of $|\lambda|$ and we only need to distinguish between $\lambda > 0$ and $\lambda < 0$. The result is shown in Figure 1, where we also include the GR result $S = 2\pi|J|$ for comparison.

In these plots we can observe some genuine non-perturbative effects. For $\lambda > 0$ we see that at $J = 0$ there is a solution with non-vanishing entropy (and also non-vanishing area).

Also for small values of J there is a second branch of solutions that ends at a solution with $S = 0$ and $J \neq 0$. On the other hand, for $\lambda < 0$ we see that there are no solutions below a minimum value of the angular momentum, and we have approximately the bound

$$|J| > 4.1 \frac{\sqrt{|\lambda|}}{G}. \quad (4.42)$$

Above this value there are two solutions: one that asymptotes to Kerr for large J and whose entropy behaves as (4.41), and a second one whose entropy tends to a constant value and which does not exist in the GR limit. It is not clear what credibility one can assign to these non-perturbative phenomena since they lie beyond the limit of validity of effective field theory, but they provide an interesting example of the effects of higher-derivative corrections in a highly non-linear regime.

Adding a cosmological constant

Let us now consider $\Lambda \neq 0$. The cosmological constant introduces a new scale and it gives rise to a non-linear relation $S(J)$ even for Einstein gravity, so let us start by reviewing that case.

When $\lambda = \tilde{\lambda} = 0$ we have $x_1 = -x_2$ and $n = 0$. Then, introducing the dimensionless variable,

$$\zeta = \frac{x_2}{a}, \quad (4.43)$$

we can solve the equations (4.22) as

$$a = \sqrt{\frac{3(1 - \zeta^2)}{\Lambda(\zeta^4 + 6\zeta^2 - 3)}}, \quad \omega = \frac{2\zeta(3 - \zeta^2)}{(\zeta^4 + 6\zeta^2 - 3)a}. \quad (4.44)$$

Observe that demanding that a is real and that $\omega > 0$ leads to the following bounds on ζ depending on the sign of Λ :¹³

$$\begin{aligned} \Lambda > 0 &\Rightarrow \sqrt{2\sqrt{3} - 3} \leq \zeta \leq 1, \\ \Lambda < 0 &\Rightarrow 1 \leq \zeta \leq \sqrt{3}. \end{aligned} \quad (4.45)$$

Now, plugging this into our expressions for the entropy (4.28) and the angular momentum (4.40) we get

$$S_0 = \frac{3\pi(1 - \zeta^2)}{G\Lambda(3 - \zeta^2)}, \quad (4.46)$$

$$J_0 = \frac{3\zeta(1 - \zeta^2)}{G\Lambda(3 - \zeta^2)^2}, \quad (4.47)$$

¹³In the dS case one may also consider the solutions with $0 < \zeta < \sqrt{2\sqrt{3} - 3}$. Interestingly, these do not correspond to near-horizon geometries but to rotating Nariai geometries [71] (consistent of a fibration of dS_2 over S^2) which arise when the black hole horizon merges with the cosmological horizon.

which are precisely the relations for the Kerr-(A)dS black hole that we obtained in (3.57). The explicit expression for $S(J)$ is very involved because it requires solving a fourth degree polynomial, so it is more useful to work with this parametric relation.

Having obtained the result for every quantity a , x_i , ω and n , for Einstein gravity, we can now solve the equations (4.22) in the case of $\lambda \neq 0$, $\tilde{\lambda} \neq 0$ by assuming a series expansion around that result,

$$a = \sum_{p,q} a_{(p,q)} \lambda^p \tilde{\lambda}^q, \quad x_i = \sum_{p,q} x_{i(p,q)} \lambda^p \tilde{\lambda}^q, \quad \omega = \sum_{p,q} \omega_{(p,q)} \lambda^p \tilde{\lambda}^q, \quad n = \sum_{p,q} n_{(p,q)} \lambda^p \tilde{\lambda}^q, \quad (4.48)$$

where $n_{(0,0)} = 0$, $x_{2(0,0)} = -x_{1(0,0)} = a_{(0,0)}\zeta$ and $a_{(0,0)}$ and $\omega_{(0,0)}$ are those given by (4.44). At every order in $\lambda^p \tilde{\lambda}^q$ we obtain a linear system of equations for the coefficients $a_{(p,q)}$, $x_{i(p,q)}$, $\omega_{(p,q)}$ and $n_{(p,q)}$ that we can solve straightforwardly. Plugging back the result in (4.28) and (4.40) we then obtain the corrections to (4.46) and (4.47) expressed as a function of ζ . However, since ζ is an arbitrary parameter, we can consider a redefinition

$$\zeta \rightarrow \zeta + \sum_{p,q} \delta\zeta_{(p,q)} \lambda^p \tilde{\lambda}^q = \zeta + \delta\zeta_{(1,0)}\lambda + \delta\zeta_{(0,1)}\tilde{\lambda} + \dots \quad (4.49)$$

By choosing the coefficients $\delta\zeta_{(p,q)}$ appropriately we can always make the expression of J as a function of ζ identical to that of Einstein gravity (4.47). Thus, with the appropriate choice of ζ , we have $J = J_0$. For that choice, the entropy reads, to quadratic order in λ and $\tilde{\lambda}$

$$S = S_0 \left[1 - \frac{16\lambda\Lambda^2\zeta^2(\zeta^2-3)^2(\zeta^4+13\zeta^2-12)}{9(\zeta^2-1)^2(\zeta^6+7\zeta^4+3\zeta^2-3)} + \frac{64\zeta^2(\zeta^2-3)^4\lambda^2\Lambda^4}{81(\zeta^4-1)^4(\zeta^4+6\zeta^2-3)^3}(\zeta^{18}+70\zeta^{16}-582\zeta^{14}-746\zeta^{12}+13444\zeta^{10}+2946\zeta^8-34650\zeta^6+33282\zeta^4-13797\zeta^2+2592) + \frac{128\tilde{\lambda}^2\Lambda^4\zeta^2(3-5\zeta^2)^2(\zeta^2-3)^4}{9(\zeta^4-1)^4(\zeta^4+6\zeta^2-3)} + \dots \right]. \quad (4.50)$$

An explicit expression $S(J)$ can be obtained in the limits of small and large black holes. For small black holes, $J \ll 1/(G|\Lambda|)$, which are obtained in the limit $\zeta \rightarrow 1$, the effect of the cosmological constant is irrelevant and we reproduce the asymptotically flat result (4.41). Large black holes, *i.e.*, those with $J \gg 1/(G|\Lambda|)$, are only possible in the AdS case $\Lambda < 0$, and in that limit (corresponding to $\zeta \rightarrow \sqrt{3}$) we have

$$S(J) = S_0(J) \left[1 + \lambda \left(\frac{3\sqrt{3}\Lambda}{GJ} - \frac{2^{3/2}3^{1/4}\sqrt{-\Lambda}}{(GJ)^{3/2}} + \frac{1}{4(GJ)^2} \right) + \frac{27(\lambda^2 + \tilde{\lambda}^2)\Lambda^2}{4(GJ)^2} + \mathcal{O}(J^{-5/2}) \right]. \quad (4.51)$$

Unlike the case of the stringy corrections for large AdS black holes (3.65), the correction due to the cubic terms vanishes for $J \rightarrow \infty$.

5 Discussion

In this paper we have computed the corrections to the entropy-angular momentum relation of the extremal Kerr black hole in two different higher-derivative extensions of GR motivated by string theory (1.1) and by general effective field theory arguments (1.5). Our strategy consisted in studying near-horizon extremal geometries, since the existence of an enhanced symmetry in the near-horizon region greatly simplifies the analysis of the solutions. One can directly obtain the black hole entropy from the near-horizon metric thanks to the Iyer-Wald formula. However, one should express the entropy in terms of physically meaningful quantities of the black hole — in this case, the angular momentum — rather than some arbitrary parameters of the solution. Computing the angular momentum from the near-horizon geometry is more challenging, since we lack an asymptotic region in our solutions. In fact, the angular momentum must be identified as the integral of the Noether charge two-form in a sphere at infinity. The integration surface cannot be deformed as, in general, the charge two-form does not satisfy a Gauss law. In order to address this problem, we provided an appropriate Komar-type modification of the charge that is independent of the surface of integration. The Komar generalization of the Noether charge two-form is in general ambiguous, but, for the type of solutions considered, we demonstrated that there is a precise choice of the Noether-Komar two-form whose integral on the black hole horizon computes the angular momentum unambiguously.

Another method to identify the angular momentum from the near-horizon geometry was proposed in [28] in the context of the entropy function approach. The underlying idea of [28] is that the angular momentum can be identified as an electric charge in one dimension less, after compactification of the spacetime in the periodic coordinate. It is not evident that this method is equivalent to the Noether charge approach we employed. However, we checked that for the solutions we studied, both methods yield the same answer.

Besides near-horizon geometries, the only other approach we are aware of that can be used to study the corrections to the entropy of extremal rotating black holes is that of Reall and Santos [18]. This approach has the advantage of being applicable to non-extremal black holes, but has the limitation of only working for first-order corrections. The near-horizon approach on the other hand allows us to go beyond the leading correction. In our case, we were able to obtain the corrections to the entropy at order α'^2 for the stringy action (1.1) while for the cubic theory (1.5) we obtained the exact result — implicitly defined as the solution of a system of algebraic equations.

On the other hand, the near-horizon geometry does not allow us to obtain the mass of the solutions. For first-order corrections, the mass could be obtained using the methods of [18] or [72], but beyond first order the computation of the full global solution seems unavoidable in order to obtain the mass.

Let us now discuss the specific results we obtained. In the case of the heterotic string theory, we were able to find explicitly the corrected NHEK geometry. This is even more remarkable in the Kerr-(A)dS case, where the solution takes a quite involved, yet fully analytic

form. Obtaining the solution was necessary in order to compute both the entropy and the angular momentum, which allowed us to derive the relation (1.4) (and (3.61) for the case of $\Lambda \neq 0$). While the $\mathcal{O}(\alpha')$ correction to the entropy — entirely coming from the topological Gauss-Bonnet term — is positive, we observe that the $\mathcal{O}(\alpha'^2)$ correction is negative. It would be interesting to understand how this is related to the fact that we are assuming a stabilized dilaton. In the case where the dilaton remains massless, the corrections to the entropy of the non-extremal Kerr black hole were shown to be positive [66]. The extremal solution becomes singular in that case — which is the reason why we consider a stabilized dilaton — but studying the near-extremal solutions could shed light on the nature of this singularity.¹⁴ It would be interesting to find out whether there is a well-defined limit for the entropy at extremality when the dilaton is massless.

We also obtained the entropy of extremal rotating black holes for the cubic theories (1.5), that provide a general effective field theory extension of GR to six derivatives. These cubic Lagrangians are chosen in a special way: they correspond to Einsteinian cubic gravity [46] and to a new parity-violating density with analogous properties to ECG that we introduced here for the first time. The most remarkable property of these theories is that certain aspects of the near-horizon geometries, like the entropy and the angular momentum, can be obtained exactly even if we do not know the solution explicitly. Indeed, the full non-perturbative relation $S(J)$ is obtained from the equations (4.28), (4.40) and (4.22). This allows us to easily compute the perturbative expansion of $S(J)$ at virtually any order and even to study it in a non-perturbative fashion, as discussed in section 4.3. The fact that we can obtain the perturbative expansion at higher orders is highly non-trivial and relevant from the point of view of effective field theory. Let us remember that the effective action for gravity contains an infinite tower of higher-derivative corrections. However, each term in the action generates at the same time an infinite number of corrections on the black hole entropy. Here we have been able to provide the complete answer for the corrections to the entropy related exclusively with the six-derivative terms. Higher-derivative terms will induce additional corrections that should be added to this result. For instance, the $\mathcal{O}(\lambda^2, \tilde{\lambda}^2)$ terms in (1.8) contribute to the entropy at the same order as the leading contribution from 10-derivative terms. Our results have also showed for the first time the effect of a parity-breaking correction on the entropy. Interestingly the sign of this correction is fixed since it enters at order $\tilde{\lambda}^2$, and it is always positive.

In sum, the strategy of writing the effective field theory of gravity in terms of these special densities — which belong to the broader family of Generalized Quasitopological gravities [51, 55, 56] — greatly simplifies the study of solutions and allow us to go beyond the leading-order corrections. It would be interesting to explore if this procedure can be generalized to higher-order terms (beyond six derivatives) [45].

¹⁴As recently shown by [73], singularities are in fact very common in extremal rotating black holes with higher-derivative corrections. Even in the cases where the background geometry remains regular, fluctuations around it may render the spacetime singular. In this sense, the singularity due to the dilaton is perhaps not so surprising.

Finally, this work could be extended in several ways. Here we restricted ourselves to the case of pure gravity, but it would be interesting to study charged solutions as well. On the one hand, one may consider a general effective field theory extension of Einstein-Maxwell theory and study the corrections to the extremal Kerr-Newman solution. On the other, one could consider the effective action of heterotic string theory with gauge fields and study the corrections to the Kerr-Sen solution [74]. Other cases with additional fields, higher dimensions and/or supersymmetry — like the BMPV black hole [75, 76] — would also be worth to be explored.

Acknowledgments

We thank Nikolay Bobev, Evan McDonough, Rishi Mouland, David Pereñiguez, Thomas Van Riet and Nicolas Yunes for useful discussions. The work of PAC is supported by a postdoctoral fellowship from the Research Foundation - Flanders (FWO grant 12ZH121N). MD is supported by KU Leuven C1 grant ZKD1118 C16/16/005, and by the Research Programme of The Research Foundation – Flanders (FWO) grant G0F9516N.

A Near-horizon geometry of the heterotic theory with $\Lambda \neq 0$

The first nonzero α' corrections for ψ , f , and g in the presence of a nonzero cosmological constant were found in section 3. The various parameters in (3.51), (3.52), and (3.53) are given by

$$\begin{aligned}\rho_{f,0} &= -y_0^4 - 3, \\ \rho_{f,1} &= -4y^2 y_0^2 (y_0^2 + 3), \\ \rho_{f,2} &= y^4 (y_0^4 - 4y_0^2 + 7),\end{aligned}\tag{A.1}$$

$$\begin{aligned}\kappa_{f,0} &= 7y_0^{10} + 48y_0^8 + 96y_0^6 - 18y_0^4 - 135y_0^2 + 162, \\ \kappa_{f,2} &= y_0^{10} + 15y_0^8 + 78y_0^6 - 6y_0^4 - 255y_0^2 + 135, \\ \kappa_{f,4} &= -3(y_0^8 + 2y_0^6 - 8y_0^4 + 14y_0^2 - 9),\end{aligned}\tag{A.2}$$

$$\begin{aligned}\tau_{f,0} &= -47y_0^{12} - 286y_0^{10} - 683y_0^8 - 1124y_0^6 - 685y_0^4 + 354y_0^2 - 281, \\ \tau_{f,2} &= -7y_0^{12} - 70y_0^{10} - 273y_0^8 - 420y_0^6 + 343y_0^4 + 522y_0^2 - 287, \\ \tau_{f,4} &= 2(10y_0^{10} + 19y_0^8 - 4y_0^6 + 74y_0^4 - 102y_0^2 + 3),\end{aligned}\tag{A.3}$$

$$\begin{aligned}
\gamma_{g,0} &= -4 (88y_0^{10} + 644y_0^8 + 3271y_0^6 + 9237y_0^4 - 14403y_0^2 - 765) , \\
\gamma_{g,2} &= -4873y_0^{10} - 23429y_0^8 - 76486y_0^6 - 224022y_0^4 + 209343y_0^2 - 68085 , \\
\gamma_{g,4} &= 60 (246y_0^{10} + 2093y_0^8 + 4862y_0^6 + 1772y_0^4 + 12452y_0^2 - 2433) , \\
\gamma_{g,6} &= 14 (1093y_0^{10} + 8435y_0^8 + 18310y_0^6 + 7794y_0^4 + 58053y_0^2 - 13653) , \\
\gamma_{g,8} &= 140 (64y_0^{10} + 437y_0^8 + 883y_0^6 + 543y_0^4 + 3273y_0^2 - 792) , \\
\gamma_{g,10} &= 105 (19y_0^{10} + 123y_0^8 + 242y_0^6 + 186y_0^4 + 939y_0^2 - 261) ,
\end{aligned} \tag{A.4}$$

$$\begin{aligned}
\gamma_{f,0} &= 4 (298y_0^{14} + 2324y_0^{12} + 5635y_0^{10} + 13689y_0^8 + 27540y_0^6 + 4266y_0^4 + 24831y_0^2 - 2295) , \\
\gamma_{f,2} &= 3 (636y_0^{14} + 5741y_0^{12} + 14566y_0^{10} + 74875y_0^8 + 96936y_0^6 - 213333y_0^4 + 130550y_0^2 \\
&\quad - 27795) , \\
\gamma_{f,4} &= 2 (27574y_0^{14} + 257531y_0^{12} + 966888y_0^{10} + 2159253y_0^8 + 2287958y_0^6 - 53427y_0^4 \\
&\quad + 966092y_0^2 - 50205) , \\
\gamma_{f,6} &= 37120y_0^{14} + 378777y_0^{12} + 1529990y_0^{10} + 3881051y_0^8 + 3496412y_0^6 - 3369561y_0^4 \\
&\quad + 2335518y_0^2 - 345627 , \\
\gamma_{f,8} &= 12 (2198y_0^{14} + 23291y_0^{12} + 95919y_0^{10} + 254192y_0^8 + 221264y_0^6 - 310209y_0^4 + 217323y_0^2 \\
&\quad - 35370) , \\
\gamma_{f,10} &= 7 (980y_0^{14} + 10879y_0^{12} + 44586y_0^{10} + 142065y_0^8 + 126952y_0^6 - 332583y_0^4 + 256282y_0^2 \\
&\quad - 48201) , \\
\gamma_{f,12} &= 70 (6y_0^{14} + 25y_0^{12} - 56y_0^{10} + 1187y_0^8 + 2006y_0^6 - 10041y_0^4 + 10140y_0^2 - 2115) , \\
\gamma_{f,14} &= -105 (17y_0^{12} + 110y_0^{10} + 139y_0^8 - 116y_0^6 + 735y_0^4 - 1146y_0^2 + 261) .
\end{aligned} \tag{A.5}$$

B Noether-Komar two-form for the cubic theories

The Noether-Komar 2-form (2.9) has the following components

$$\begin{aligned}
\tilde{Q}_\xi &= \frac{1}{16\pi G} \left[dt \wedge dr (Ar\xi^0 + B\xi^3) + dt \wedge dx (Cr^2\xi^0 + Dr\xi^3) + dr \wedge d\phi \xi^0 E \right. \\
&\quad \left. + dx \wedge d\phi (Fr\xi^0 + G\xi^3) \right] ,
\end{aligned} \tag{B.1}$$

where A, B, C, D, E, F and G are the following functions. In these expressions we use the dimensionless variable $y = x/a$ and the derivatives are with respect to y , *e.g.*, $f' = \frac{df}{dy} = a \frac{df}{dx}$.

$$\begin{aligned}
A &= a^2 (-2fy + (1 + y^2) f') + \frac{\lambda}{a^2} \left[\frac{8f^2y(27 - 5y^2)}{(1 + y^2)^3} - \frac{40fy}{(1 + y^2)^2} - \frac{32f^3y(-35 + 58y^2)}{(1 + y^2)^4} \right. \\
&\quad \left. + \left(\frac{20}{1 + y^2} + \frac{24f(-9 + 2y^2)}{(1 + y^2)^2} - \frac{4f^2(29 - 576y^2 + 15y^4)}{(1 + y^2)^3} \right) f' \right]
\end{aligned}$$

$$\begin{aligned}
& + \left(-\frac{68y}{1+y^2} + \frac{8fy(-50+3y^2)}{(1+y^2)^2} \right) (f')^2 + \frac{6(-5+y^2)(f')^3}{1+y^2} + \left(\frac{40fy}{1+y^2} \right. \\
& + \frac{f^2(-676y+60y^3)}{(1+y^2)^2} + \left. \left(10 + \frac{f(206-42y^2)}{1+y^2} \right) f' \right) f'' + 6fy(f'')^2 \\
& + \left(-12f - \frac{12f^2(-4+y^2)}{1+y^2} + 6fyf' \right) f^{(3)} + \frac{\tilde{\lambda}}{a^2} \left[-\frac{24f}{(1+y^2)^2} + \frac{8f^2(23+39y^2)}{(1+y^2)^3} \right. \\
& + \frac{8f^3(-11-432y^2+63y^4)}{(1+y^2)^4} + \left. \left(-\frac{12y}{1+y^2} + \frac{f^2(1952y-720y^3)}{(1+y^2)^3} + \frac{f(76y-12y^3)}{(1+y^2)^2} \right) f' \right. \\
& + \left(\frac{2(-33+5y^2)}{1+y^2} + \frac{4f(-21+53y^2)}{(1+y^2)^2} \right) (f')^2 + \frac{36y(f')^3}{1+y^2} + \left(4 + \frac{4f(5+y^2)}{1+y^2} \right. \\
& \left. + \frac{8f^2(-39+37y^2)}{(1+y^2)^2} + \left(-6y - \frac{216fy}{1+y^2} \right) f' \right) f'' + 18f(f'')^2 + \left(-\frac{36f^2y}{1+y^2} + 18ff' \right) f^{(3)} \Big], \tag{B.2}
\end{aligned}$$

$$\begin{aligned}
B = & \frac{1}{2}a(-1-y^2)f' + \frac{\lambda}{a^3} \left[\frac{8f^2y}{(1+y^2)^3} + \frac{16f^3y(-1+14y^2)}{(1+y^2)^4} + \left(\frac{4f(8+3y^2)}{(1+y^2)^2} \right. \right. \\
& + \left. \left. \frac{4f^2(-31-42y^2+3y^4)}{(1+y^2)^3} \right) f' + \left(\frac{12y}{1+y^2} + \frac{f(-62y+6y^3)}{(1+y^2)^2} \right) (f')^2 - \frac{3(-5+y^2)(f')^3}{1+y^2} \right. \\
& + \left. \left(-\frac{12fy}{1+y^2} + \frac{f^2(88y-12y^3)}{(1+y^2)^2} + \frac{f(-20+6y^2)f'}{1+y^2} \right) f'' \right] + \frac{\tilde{\lambda}}{a^3} \left[-\frac{24f^2(2+3y^2)}{(1+y^2)^3} + \right. \\
& \frac{8f^3(20+45y^2-9y^4)}{(1+y^2)^4} + \left. \left(-\frac{12fy}{(1+y^2)^2} + \frac{4f^2(y+15y^3)}{(1+y^2)^3} \right) f' + \left(\frac{12}{1+y^2} \right. \right. \\
& \left. \left. + \frac{4f(-15+4y^2)}{(1+y^2)^2} \right) (f')^2 - \frac{18y(f')^3}{1+y^2} + \left(\frac{f^2(48-44y^2)}{(1+y^2)^2} - \frac{8f}{1+y^2} + \frac{30fyf'}{1+y^2} \right) f'' \right], \tag{B.3}
\end{aligned}$$

$$\begin{aligned}
C = & a \left(1 - \frac{4f}{1+y^2} \right) + \frac{\lambda}{a^3} \left[\frac{20f(-1+3y^2)}{(1+y^2)^3} + \frac{12f^2(9-50y^2+5y^4)}{(1+y^2)^4} \right. \\
& + \frac{64f^3(5-41y^2+5y^4)}{(1+y^2)^5} + \left. \left(\frac{48f^2y(25-13y^2)}{(1+y^2)^4} - \frac{40y}{(1+y^2)^2} + \frac{f(732y-148y^3)}{(1+y^2)^3} \right) f' \right. \\
& + \left. \left(\frac{8f(19+9y^2)}{(1+y^2)^3} + \frac{2(-83+41y^2)}{(1+y^2)^2} \right) (f')^2 + \frac{88y(f')^3}{(1+y^2)^2} + \left(\frac{10}{1+y^2} \right. \right. \\
& \left. \left. + \frac{192f^2(-1+2y^2)}{(1+y^2)^3} + \frac{f(-76+64y^2)}{(1+y^2)^2} + \left(-\frac{184fy}{(1+y^2)^2} - \frac{66y}{1+y^2} \right) f' - \frac{20(f')^2}{1+y^2} \right) f'' \right]
\end{aligned}$$

$$\begin{aligned}
& + \left(5 + \frac{40f}{1+y^2} \right) (f'')^2 + \left(-\frac{80f^2y}{(1+y^2)^2} - \frac{10fy}{1+y^2} + \left(5 + \frac{40f}{1+y^2} \right) f' \right) f^{(3)} \Big] \\
& + \frac{\tilde{\lambda}}{a^3} \left[\frac{48fy}{(1+y^2)^3} + \frac{96f^2y(-7+4y^2)}{(1+y^2)^4} + \frac{224f^3y(-1+9y^2)}{(1+y^2)^5} + \left(\frac{6(-3+y^2)}{(1+y^2)^2} \right. \right. \\
& \left. \left. - \frac{32f^2(4+55y^2)}{(1+y^2)^4} + \frac{6f(59-108y^2+y^4)}{(1+y^2)^3} \right) f' + \left(-\frac{6y(-39+y^2)}{(1+y^2)^2} \right. \right. \\
& \left. \left. + \frac{16fy(4+y^2)}{(1+y^2)^3} \right) (f')^2 + \frac{(84-20y^2)(f')^3}{(1+y^2)^2} + \left(-\frac{32f^2y(-21+y^2)}{(1+y^2)^3} - \frac{6y}{1+y^2} \right. \right. \\
& \left. \left. - \frac{2fy(-71+3y^2)}{(1+y^2)^2} + \left(\frac{8f(-24+5y^2)}{(1+y^2)^2} + \frac{-77+9y^2}{1+y^2} \right) f' + \frac{12y(f')^2}{1+y^2} \right) f'' \\
& + \left(-3y - \frac{24fy}{1+y^2} \right) (f'')^2 + \left(2 + \frac{16f^2(-6+y^2)}{(1+y^2)^2} + \frac{2f(2+y^2)}{1+y^2} \right. \\
& \left. + \left(-3y - \frac{24fy}{1+y^2} \right) f' \right) f^{(3)} \Big], \tag{B.4}
\end{aligned}$$

$$\begin{aligned}
D = & -1 + \frac{f(1-y^2)}{1+y^2} - 2yf' + \frac{1}{2}(-1-y^2)f'' + (-1-y^2)\Lambda a^2 + \frac{\lambda}{a^4} \left[\frac{f^2(8-40y^2)}{(1+y^2)^4} \right. \\
& - \frac{40f^3(1-10y^2+y^4)}{(1+y^2)^5} + \left(\frac{48f^2y(9-2y^2)}{(1+y^2)^4} - \frac{88fy}{(1+y^2)^3} \right) f' + \left(\frac{50-6y^2}{(1+y^2)^2} \right. \\
& \left. + \frac{f(-298+312y^2-6y^4)}{(1+y^2)^3} \right) (f')^2 + \frac{2y(-55+3y^2)(f')^3}{(1+y^2)^2} + \left(\frac{8f}{(1+y^2)^2} \right. \\
& \left. - \frac{12f^2(-1+9y^2)}{(1+y^2)^3} + \left(\frac{12y}{1+y^2} + \frac{4fy(-4+3y^2)}{(1+y^2)^2} \right) f' + \frac{(25-3y^2)(f')^2}{1+y^2} \right) f'' \\
& - \frac{20f(f'')^2}{1+y^2} + \left(\frac{40f^2y}{(1+y^2)^2} - \frac{20ff'}{1+y^2} \right) f^{(3)} \Big] + \frac{\tilde{\lambda}}{a^4} \left[\frac{144f^2y}{(1+y^2)^4} - \frac{16f^3y(35+33y^2)}{(1+y^2)^5} \right. \\
& + \left(\frac{24f(-4+3y^2)}{(1+y^2)^3} + \frac{8f^2(50-19y^2+9y^4)}{(1+y^2)^4} \right) f' + \left(\frac{8fy(56-13y^2)}{(1+y^2)^3} - \frac{48y}{(1+y^2)^2} \right) (f')^2 \\
& + \frac{(-84+40y^2)(f')^3}{(1+y^2)^2} + \left(-\frac{8fy}{(1+y^2)^2} - \frac{8f^2y(24+y^2)}{(1+y^2)^3} + \left(\frac{16}{1+y^2} + \frac{4f(6+7y^2)}{(1+y^2)^2} \right) f' \right. \\
& \left. - \frac{24y(f')^2}{1+y^2} \right) f'' + \frac{12fy(f'')^2}{1+y^2} + \left(-\frac{8f^2(-6+y^2)}{(1+y^2)^2} - \frac{8f}{1+y^2} + \frac{12fyf'}{1+y^2} \right) f^{(3)} \Big], \tag{B.5}
\end{aligned}$$

$$E = -afy + \frac{\lambda}{a^3} \left[-\frac{32f^2y}{(1+y^2)^3} + \frac{f^3(544y-704y^3)}{(1+y^2)^4} + \left(-\frac{2f(1+9y^2)}{(1+y^2)^2} \right. \right.$$

$$\begin{aligned}
& -\frac{2f^2(91-492y^2+9y^4)}{(1+y^2)^3} \Big) f' + \frac{2fy(-131+9y^2)(f')^2}{(1+y^2)^2} + \left(\frac{18fy}{1+y^2} + \frac{2f^2y(-125+9y^2)}{(1+y^2)^2} \right. \\
& \left. + \frac{f(83-15y^2)f'}{1+y^2} \right) f'' + 3fy(f'')^2 + \left(-6f - \frac{6f^2(-4+y^2)}{1+y^2} + 3fyf' \right) f^{(3)} \Big] \\
& + \frac{\tilde{\lambda}}{a^3} \left[\frac{12f^2(-1+15y^2)}{(1+y^2)^3} + \frac{4f^3(29-342y^2+45y^4)}{(1+y^2)^4} + \left(-\frac{96fy}{(1+y^2)^2} \right. \right. \\
& \left. \left. - \frac{20f^2y(-49+15y^2)}{(1+y^2)^3} \right) f' + \frac{2f(-51+61y^2)(f')^2}{(1+y^2)^2} + \left(\frac{14f}{1+y^2} + \frac{4f^2(-27+26y^2)}{(1+y^2)^2} \right. \right. \\
& \left. \left. - \frac{78fyf'}{1+y^2} \right) f'' + 9f(f'')^2 + \left(-\frac{18f^2y}{1+y^2} + 9ff' \right) f^{(3)} \right], \tag{B.6}
\end{aligned}$$

$$\begin{aligned}
F = & -\frac{f(3+y^2)}{1+y^2} - 2yf' - \frac{1}{2}(1+y^2)f'' - (1+y^2)\Lambda a^2 + \frac{\lambda}{a^4} \left[\frac{32f^2(-1+5y^2)}{(1+y^2)^4} \right. \\
& + \frac{8f^3(35-278y^2+35y^4)}{(1+y^2)^5} + \left(\frac{48f^2y(34-15y^2)}{(1+y^2)^4} - \frac{56fy}{(1+y^2)^3} \right) f' - \left(\frac{20}{(1+y^2)^2} \right. \\
& \left. + \frac{2f(73-192y^2+3y^4)}{(1+y^2)^3} \right) (f')^2 + \frac{2y(-11+3y^2)(f')^3}{(1+y^2)^2} + \left(\frac{16f}{(1+y^2)^2} \right. \\
& \left. + \frac{12f^2(-15+23y^2)}{(1+y^2)^3} + \frac{4fy(-50+3y^2)f'}{(1+y^2)^2} + \frac{(5-3y^2)(f')^2}{1+y^2} \right) f'' + \frac{20f(f'')^2}{1+y^2} \\
& + \left(-\frac{40f^2y}{(1+y^2)^2} + \frac{20ff'}{1+y^2} \right) f^{(3)} \Big] + \frac{\tilde{\lambda}}{a^4} \left[\frac{16f^3y(-49+93y^2)}{(1+y^2)^5} + \left(\frac{24f}{(1+y^2)^3} \right. \right. \\
& \left. \left. + \frac{8f^2(34-239y^2+9y^4)}{(1+y^2)^4} \right) f' + \left(\frac{12y}{(1+y^2)^2} - \frac{8fy(-64+11y^2)}{(1+y^2)^3} \right) (f')^2 + \frac{20y^2(f')^3}{(1+y^2)^2} \right. \\
& \left. + \left(-\frac{40f^2y(-12+y^2)}{(1+y^2)^3} - \frac{16fy}{(1+y^2)^2} + \left(-\frac{4}{1+y^2} + \frac{4f(-42+17y^2)}{(1+y^2)^2} \right) f' - \frac{12y(f')^2}{1+y^2} \right) f'' \right. \\
& \left. - \frac{12fy(f'')^2}{1+y^2} + \left(\frac{8f^2(-6+y^2)}{(1+y^2)^2} + \frac{8f}{1+y^2} - \frac{12fyf'}{1+y^2} \right) f^{(3)} \right], \tag{B.7}
\end{aligned}$$

$$\begin{aligned}
G = & \frac{f}{a(1+y^2)} + \frac{\lambda}{a^5} \left[\frac{f^2(16-80y^2)}{(1+y^2)^4} - \frac{16f^3(5-41y^2+5y^4)}{(1+y^2)^5} + \left(\frac{28fy}{(1+y^2)^3} \right. \right. \\
& \left. \left. + \frac{12f^2y(-25+13y^2)}{(1+y^2)^4} \right) f' + \left(\frac{10}{(1+y^2)^2} - \frac{2f(19+9y^2)}{(1+y^2)^3} \right) (f')^2 - \frac{22y(f')^3}{(1+y^2)^2} \right. \\
& \left. + \left(\frac{f^2(48-96y^2)}{(1+y^2)^3} - \frac{8f}{(1+y^2)^2} + \frac{46fyf'}{(1+y^2)^2} + \frac{5(f')^2}{1+y^2} \right) f'' - \frac{10f(f'')^2}{1+y^2} + \left(\frac{20f^2y}{(1+y^2)^2} \right. \right.
\end{aligned}$$

$$\begin{aligned}
& -\frac{10ff'}{1+y^2} f^{(3)} \Big] + \frac{\tilde{\lambda}}{a^5} \left[\frac{56f^3y(1-9y^2)}{(1+y^2)^5} + \left(-\frac{12f}{(1+y^2)^3} + \frac{8f^2(4+55y^2)}{(1+y^2)^4} \right) f' \right. \\
& - \left(\frac{6y}{(1+y^2)^2} + \frac{4fy(4+y^2)}{(1+y^2)^3} \right) (f')^2 + \frac{(-21+5y^2)(f')^3}{(1+y^2)^2} + \left(\frac{8f^2y(-21+y^2)}{(1+y^2)^3} \right. \\
& + \frac{8fy}{(1+y^2)^2} + \left. \left(\frac{f(48-10y^2)}{(1+y^2)^2} + \frac{2}{1+y^2} \right) f' - \frac{3y(f')^2}{1+y^2} \right) f'' + \frac{6fy(f'')^2}{1+y^2} \\
& \left. + \left(-\frac{4f^2(-6+y^2)}{(1+y^2)^2} - \frac{4f}{1+y^2} + \frac{6fyf'}{1+y^2} \right) f^{(3)} \right]. \tag{B.8}
\end{aligned}$$

By direct computation, one can check that $d\mathbf{Q}_\xi = 0$ once the equation of motion (4.15) is assumed.

B.1 Proof that the ξ^0 -integral vanishes

Let us now show that the charge associated to the timelike Killing vector ∂_t vanishes. To this end, we need to evaluate the integral proportional to ξ^0 in (4.35), this is

$$\int_{x_1}^{x_2} dx F(x). \tag{B.9}$$

It is possible to check that

$$F = -\frac{a^2}{a^2+x^2} \frac{d\mathcal{E}_f}{dx} - \frac{1}{2} x \frac{d^2\mathcal{E}_f}{dx^2} + \frac{d\mathcal{F}}{dx}, \tag{B.10}$$

where \mathcal{E}_f is the equation of motion for f , given by (4.14), and

$$\begin{aligned}
\mathcal{F} = f \Big\{ & -x + \lambda \left[-\frac{32a^2fx}{(a^2+x^2)^3} + \frac{32f^2(17a^4x-22a^2x^3)}{(a^2+x^2)^4} + \left(-\frac{2(a^2+9x^2)}{(a^2+x^2)^2} \right. \right. \\
& \left. \left. - \frac{2f(91a^4-492a^2x^2+9x^4)}{(a^2+x^2)^3} \right) f' + \frac{(-262a^2x+18x^3)(f')^2}{(a^2+x^2)^2} + \left(\frac{18x}{a^2+x^2} \right. \right. \\
& \left. \left. + \frac{f(-250a^2x+18x^3)}{(a^2+x^2)^2} + \frac{(83a^2-15x^2)f'}{a^2+x^2} \right) f'' + 3x(f'')^2 + \left(-6 + \frac{f(24a^2-6x^2)}{a^2+x^2} \right. \right. \\
& \left. \left. + 3xf' \right) f^{(3)} \right] + \tilde{\lambda} \left[-\frac{12f(a^3-15ax^2)}{(a^2+x^2)^3} + \frac{4f^2(29a^5-342a^3x^2+45ax^4)}{(a^2+x^2)^4} + \left(-\frac{96ax}{(a^2+x^2)^2} \right. \right. \\
& \left. \left. + \frac{20f(49a^3x-15ax^3)}{(a^2+x^2)^3} \right) f' - \frac{2(51a^3-61ax^2)(f')^2}{(a^2+x^2)^2} + \left(\frac{14a}{a^2+x^2} \right. \right. \\
& \left. \left. - \frac{4f(27a^3-26ax^2)}{(a^2+x^2)^2} - \frac{78axf'}{a^2+x^2} \right) f'' + 9a(f'')^2 + \left(-\frac{18afx}{a^2+x^2} + 9af' \right) f^{(3)} \right] \Big\}.
\end{aligned}$$

Then, on-shell $\mathcal{E}_f = n$ is constant, and therefore we have

$$\int_{x_1}^{x_2} dx F(x) = \int_{x_1}^{x_2} dx \frac{d\mathcal{F}}{dx} = \mathcal{F}(x) \Big|_{x_1}^{x_2} = 0, \quad (\text{B.11})$$

which vanishes because \mathcal{F} is proportional to f , which vanishes at the poles $x = x_1, x_2$. Thus, the Komar charge is indeed independent of the radius.

References

- [1] D. J. Gross and E. Witten, *Superstring Modifications of Einstein's Equations*, *Nucl. Phys. B* **277** (1986) 1.
- [2] D. J. Gross and J. H. Sloan, *The Quartic Effective Action for the Heterotic String*, *Nucl. Phys. B* **291** (1987) 41–89.
- [3] E. A. Bergshoeff and M. de Roo, *The Quartic Effective Action of the Heterotic String and Supersymmetry*, *Nucl. Phys. B* **328** (1989) 439–468.
- [4] A. Strominger and C. Vafa, *Microscopic origin of the Bekenstein-Hawking entropy*, *Phys. Lett. B* **379** (1996) 99–104, [[hep-th/9601029](#)].
- [5] J. M. Maldacena and A. Strominger, *Statistical entropy of four-dimensional extremal black holes*, *Phys. Rev. Lett.* **77** (1996) 428–429, [[hep-th/9603060](#)].
- [6] C. V. Johnson, R. R. Khuri and R. C. Myers, *Entropy of 4-D extremal black holes*, *Phys. Lett. B* **378** (1996) 78–86, [[hep-th/9603061](#)].
- [7] A. Sen, *Black hole entropy function and the attractor mechanism in higher derivative gravity*, *JHEP* **09** (2005) 038, [[hep-th/0506177](#)].
- [8] A. Sen, *Entropy function for heterotic black holes*, *JHEP* **03** (2006) 008, [[hep-th/0508042](#)].
- [9] A. Sen, *Black Hole Entropy Function, Attractors and Precision Counting of Microstates*, *Gen. Rel. Grav.* **40** (2008) 2249–2431, [[0708.1270](#)].
- [10] F. Benini, K. Hristov and A. Zaffaroni, *Black hole microstates in AdS₄ from supersymmetric localization*, *JHEP* **05** (2016) 054, [[1511.04085](#)].
- [11] A. Zaffaroni, *AdS black holes, holography and localization*, *Living Rev. Rel.* **23** (2020) 2, [[1902.07176](#)].
- [12] N. Yunes and F. Pretorius, *Dynamical Chern-Simons Modified Gravity. I. Spinning Black Holes in the Slow-Rotation Approximation*, *Phys. Rev. D* **79** (2009) 084043, [[0902.4669](#)].
- [13] P. Pani, C. F. B. Macedo, L. C. B. Crispino and V. Cardoso, *Slowly rotating black holes in alternative theories of gravity*, *Phys. Rev. D* **84** (2011) 087501, [[1109.3996](#)].
- [14] V. Cardoso, M. Kimura, A. Maselli and L. Senatore, *Black Holes in an Effective Field Theory Extension of General Relativity*, *Phys. Rev. Lett.* **121** (2018) 251105, [[1808.08962](#)].
- [15] P. A. Cano and A. Ruipérez, *Leading higher-derivative corrections to Kerr geometry*, *JHEP* **05** (2019) 189, [[1901.01315](#)].
- [16] B. Kleihaus, J. Kunz and E. Radu, *Rotating Black Holes in Dilatonic Einstein-Gauss-Bonnet Theory*, *Phys. Rev. Lett.* **106** (2011) 151104, [[1101.2868](#)].

- [17] T. Delsate, C. Herdeiro and E. Radu, *Non-perturbative spinning black holes in dynamical Chern–Simons gravity*, *Phys. Lett. B* **787** (2018) 8–15, [[1806.06700](#)].
- [18] H. S. Reall and J. E. Santos, *Higher derivative corrections to Kerr black hole thermodynamics*, *JHEP* **04** (2019) 021, [[1901.11535](#)].
- [19] J. a. F. Melo and J. E. Santos, *Stringy corrections to the entropy of electrically charged supersymmetric black holes with $AdS_5 \times S^5$ asymptotics*, *Phys. Rev. D* **103** (2021) 066008, [[2007.06582](#)].
- [20] N. Bobev, V. Dimitrov, V. Reys and A. Vekemans, *Higher derivative corrections and AdS5 black holes*, *Phys. Rev. D* **106** (2022) L121903, [[2207.10671](#)].
- [21] D. Cassani, A. Ruipérez and E. Turetta, *Corrections to AdS₅ black hole thermodynamics from higher-derivative supergravity*, *JHEP* **11** (2022) 059, [[2208.01007](#)].
- [22] A. Dabholkar, A. Sen and S. P. Trivedi, *Black hole microstates and attractor without supersymmetry*, *JHEP* **01** (2007) 096, [[hep-th/0611143](#)].
- [23] D. D. K. Chow, M. Cvetič, H. Lu and C. N. Pope, *Extremal Black Hole/CFT Correspondence in (Gauged) Supergravities*, *Phys. Rev. D* **79** (2009) 084018, [[0812.2918](#)].
- [24] M. David, J. Nian and L. A. Pando Zayas, *Gravitational Cardy Limit and AdS Black Hole Entropy*, *JHEP* **11** (2020) 041, [[2005.10251](#)].
- [25] M. David and J. Nian, *Universal entropy and Hawking radiation of near-extremal AdS₄ black holes*, *JHEP* **04** (2021) 256, [[2009.12370](#)].
- [26] F. Larsen and S. Paranjape, *Thermodynamics of near BPS black holes in AdS₄ and AdS₇*, *JHEP* **10** (2021) 198, [[2010.04359](#)].
- [27] M. David, A. Lezcano González, J. Nian and L. A. Pando Zayas, *Logarithmic corrections to the entropy of rotating black holes and black strings in AdS₅*, *JHEP* **04** (2022) 160, [[2106.09730](#)].
- [28] D. Astefanesei, K. Goldstein, R. P. Jena, A. Sen and S. P. Trivedi, *Rotating attractors*, *JHEP* **10** (2006) 058, [[hep-th/0606244](#)].
- [29] J. F. Morales and H. Samtleben, *Entropy function and attractors for AdS black holes*, *JHEP* **10** (2006) 074, [[hep-th/0608044](#)].
- [30] G. Lopes Cardoso, J. M. Oberreuter and J. Perz, *Entropy function for rotating extremal black holes in very special geometry*, *JHEP* **05** (2007) 025, [[hep-th/0701176](#)].
- [31] J. Lee and R. M. Wald, *Local symmetries and constraints*, *J. Math. Phys.* **31** (1990) 725–743.
- [32] R. M. Wald, *Black hole entropy is the Noether charge*, *Phys. Rev. D* **48** (1993) R3427–R3431, [[gr-qc/9307038](#)].
- [33] V. Iyer and R. M. Wald, *Some properties of Noether charge and a proposal for dynamical black hole entropy*, *Phys. Rev. D* **50** (1994) 846–864, [[gr-qc/9403028](#)].
- [34] A. Komar, *Covariant conservation laws in general relativity*, *Phys. Rev.* **113** (1959) 934–936.
- [35] S. L. Bazanski and P. Zyla, *A Gauss type law for gravity with a cosmological constant*, *Gen. Rel. Grav.* **22** (1990) 379–387.
- [36] D. Kastor, *Komar Integrals in Higher (and Lower) Derivative Gravity*, *Class. Quant. Grav.* **25** (2008) 175007, [[0804.1832](#)].

- [37] D. Kastor, S. Ray and J. Traschen, *Enthalpy and the Mechanics of AdS Black Holes*, *Class. Quant. Grav.* **26** (2009) 195011, [[0904.2765](#)].
- [38] T. Ortín, *Komar integrals for theories of higher order in the Riemann curvature and black-hole chemistry*, *JHEP* **08** (2021) 023, [[2104.10717](#)].
- [39] D. Marolf, *Chern-Simons terms and the three notions of charge*, in *International Conference on Quantization, Gauge Theory, and Strings: Conference Dedicated to the Memory of Professor Efim Fradkin*, pp. 312–320, 6, 2000. [hep-th/0006117](#).
- [40] P. A. Cano, P. F. Ramírez and A. Ruipérez, *The small black hole illusion*, *JHEP* **03** (2020) 115, [[1808.10449](#)].
- [41] F. Faedo and P. F. Ramirez, *Exact charges from heterotic black holes*, *JHEP* **10** (2019) 033, [[1906.12287](#)].
- [42] P. A. Cano, A. Murcia, P. F. Ramírez and A. Ruipérez, *On small black holes, KK monopoles and solitonic 5-branes*, *JHEP* **05** (2021) 272, [[2102.04476](#)].
- [43] B. A. Campbell, M. Duncan, N. Kaloper and K. A. Olive, *Gravitational dynamics with lorentz chern-simons terms*, *Nuclear Physics B* **351** (1991) 778–792.
- [44] S. Alexander and N. Yunes, *Chern-Simons Modified General Relativity*, *Phys. Rept.* **480** (2009) 1–55, [[0907.2562](#)].
- [45] P. Bueno, P. A. Cano, J. Moreno and A. Murcia, *All higher-curvature gravities as Generalized quasi-topological gravities*, *JHEP* **11** (2019) 062, [[1906.00987](#)].
- [46] P. Bueno and P. A. Cano, *Einsteinian cubic gravity*, *Phys. Rev. D* **94** (2016) 104005, [[1607.06463](#)].
- [47] R. A. Hennigar and R. B. Mann, *Black holes in Einsteinian cubic gravity*, *Phys. Rev. D* **95** (2017) 064055, [[1610.06675](#)].
- [48] P. Bueno and P. A. Cano, *Four-dimensional black holes in Einsteinian cubic gravity*, *Phys. Rev. D* **94** (2016) 124051, [[1610.08019](#)].
- [49] X.-H. Feng, H. Huang, Z.-F. Mai and H. Lu, *Bounce Universe and Black Holes from Critical Einsteinian Cubic Gravity*, *Phys. Rev. D* **96** (2017) 104034, [[1707.06308](#)].
- [50] P. Bueno, P. A. Cano and A. Ruipérez, *Holographic studies of Einsteinian cubic gravity*, *JHEP* **03** (2018) 150, [[1802.00018](#)].
- [51] P. Bueno, P. A. Cano, R. A. Hennigar and R. B. Mann, *NUTs and bolts beyond Lovelock*, *JHEP* **10** (2018) 095, [[1808.01671](#)].
- [52] P. A. Cano and D. Pereñiguez, *Extremal Rotating Black Holes in Einsteinian Cubic Gravity*, *Phys. Rev. D* **101** (2020) 044016, [[1910.10721](#)].
- [53] A. M. Frassino and J. V. Rocha, *Charged black holes in Einsteinian cubic gravity and nonuniqueness*, *Phys. Rev. D* **102** (2020) 024035, [[2002.04071](#)].
- [54] C. Adair, P. Bueno, P. A. Cano, R. A. Hennigar and R. B. Mann, *Slowly rotating black holes in Einsteinian cubic gravity*, *Phys. Rev. D* **102** (2020) 084001, [[2004.09598](#)].
- [55] R. A. Hennigar, D. Kubizňák and R. B. Mann, *Generalized quasitopological gravity*, *Phys. Rev. D* **95** (2017) 104042, [[1703.01631](#)].

- [56] P. Bueno and P. A. Cano, *On black holes in higher-derivative gravities*, *Class. Quant. Grav.* **34** (2017) 175008, [[1703.04625](#)].
- [57] J. Ahmed, R. A. Hennigar, R. B. Mann and M. Mir, *Quintessential Quartic Quasi-topological Quartet*, *JHEP* **05** (2017) 134, [[1703.11007](#)].
- [58] P. Bueno and P. A. Cano, *Universal black hole stability in four dimensions*, *Phys. Rev. D* **96** (2017) 024034, [[1704.02967](#)].
- [59] P. Bueno, P. A. Cano and R. A. Hennigar, *(Generalized) quasi-topological gravities at all orders*, *Class. Quant. Grav.* **37** (2020) 015002, [[1909.07983](#)].
- [60] P. Bueno, P. A. Cano, R. A. Hennigar, M. Lu and J. Moreno, *Generalized quasi-topological gravities: the whole shebang*, *Class. Quant. Grav.* **40** (2023) 015004, [[2203.05589](#)].
- [61] F. Chen, *Quasi-topological Gravities on General Spherically Symmetric Metric*, [2301.00235](#).
- [62] T. Padmanabhan, *Some aspects of field equations in generalised theories of gravity*, *Phys. Rev. D* **84** (2011) 124041, [[1109.3846](#)].
- [63] T. Azeyanagi, G. Compere, N. Ogawa, Y. Tachikawa and S. Terashima, *Higher-Derivative Corrections to the Asymptotic Virasoro Symmetry of 4d Extremal Black Holes*, *Prog. Theor. Phys.* **122** (2009) 355–384, [[0903.4176](#)].
- [64] P. Bueno, P. A. Cano, V. S. Min and M. R. Visser, *Aspects of general higher-order gravities*, *Phys. Rev. D* **95** (2017) 044010, [[1610.08519](#)].
- [65] J. Katz, *A note on komar’s anomalous factor*, *Classical and Quantum Gravity* **2** (may, 1985) 423.
- [66] P. A. Cano and A. Ruipérez, *String gravity in $D=4$* , *Phys. Rev. D* **105** (2022) 044022, [[2111.04750](#)].
- [67] M. Cicoli, S. de Alwis and A. Westphal, *Heterotic Moduli Stabilisation*, *JHEP* **10** (2013) 199, [[1304.1809](#)].
- [68] F. Quevedo, *Local String Models and Moduli Stabilisation*, *Mod. Phys. Lett. A* **30** (2015) 1530004, [[1404.5151](#)].
- [69] B. Kleihaus, J. Kunz, S. Mojica and E. Radu, *Spinning black holes in Einstein–Gauss–Bonnet–dilaton theory: Nonperturbative solutions*, *Phys. Rev. D* **93** (2016) 044047, [[1511.05513](#)].
- [70] B. Chen and L. C. Stein, *Deformation of extremal black holes from stringy interactions*, *Phys. Rev. D* **97** (2018) 084012, [[1802.02159](#)].
- [71] H. Nariai, *On some static solutions of Einstein’s gravitational field equations in a spherically symmetric case*, *Sci. Rep. Tohoku Univ. Eighth Ser.* **34** (Jan., 1950) 160.
- [72] L. Aalsma, *Corrections to extremal black holes from Iyer–Wald formalism*, *Phys. Rev. D* **105** (2022) 066022, [[2111.04201](#)].
- [73] G. T. Horowitz, M. Kolanowski, G. N. Remmen and J. E. Santos, *Extremal Kerr black holes as amplifiers of new physics*, [2303.07358](#).
- [74] A. Sen, *Rotating charged black hole solution in heterotic string theory*, *Phys. Rev. Lett.* **69** (1992) 1006–1009, [[hep-th/9204046](#)].

- [75] J. C. Breckenridge, R. C. Myers, A. W. Peet and C. Vafa, *D-branes and spinning black holes*, *Phys. Lett. B* **391** (1997) 93–98, [[hep-th/9602065](#)].
- [76] M. Guica, L. Huang, W. Li and A. Strominger, *R^{**2} corrections for 5-D black holes and rings*, *JHEP* **10** (2006) 036, [[hep-th/0505188](#)].