

SLD Fisher information for kinetic uncertainty relations

Satoshi Nakajima* and Yasuhiro Utsumi

Department of Physics Engineering, Faculty of Engineering, Mie University, Tsu, Mie 514-8507, Japan

(Dated: March 28, 2023)

We investigate a symmetric logarithmic derivative (SLD) Fisher information for kinetic uncertainty relations (KURs) of open quantum systems described by the GKSL quantum master equation with and without the detailed balance condition. In a quantum kinetic uncertainty relation derived by Vu and Saito [Phys. Rev. Lett. **128**, 140602 (2022)], the Fisher information of probability of quantum trajectory with a time-rescaling parameter plays an essential role. This Fisher information is upper bounded by the SLD Fisher information. For a finite time and arbitrary initial state, we derive a concise expression of the SLD Fisher information, which is a double time integral and can be calculated by solving coupled first-order differential equations. We also derive a simple lower bound of the Fisher information of quantum trajectory. We point out that the SLD Fisher information also appears in the speed limit based on the Mandelstam-Tamm relation by Hasegawa [arXiv:2203.12421v4]. When the jump operators connect eigenstates of the system Hamiltonian, we show that the Bures angle in the interaction picture is upper bounded by the square root of the dynamical activity at short times, which contrasts with the classical counterpart.

I. INTRODUCTION

In recent years, universal relations that characterize the fluctuations of nonequilibrium systems have been intensively investigated. A primary class of inequalities is the thermodynamic uncertainty relation (TUR) [1–10]. A similar relation, the kinetic uncertainty relation (KUR) imposes another upper bound on the precision of generic counting observables in terms of the dynamical activity [11–13].

Quantum coherence plays an essential role in a broad class of thermodynamics. Concerning the TUR and KUR originally derived for classical stochastic systems, it has been shown through specific examples that these relations can be violated in the quantum realm [14–19]. Several quantum bounds accounting for the quantum coherence have been derived [20–25]. For open quantum systems described by the Gorini-Kossakowski-Sudarshan-Lindblad (GKSL) equation with time-independent Hamiltonian and jump operators, Vu and Saito [26] derived a quantum KUR

$$\frac{\tau^2(\partial_\tau\langle\Phi\rangle)^2}{\text{var}[\Phi]} \leq F \quad (1)$$

using the Cramér-Rao inequality. Here, Φ is a time-integrated counting observable of the system and $\text{var}[\Phi]$ is its variance. F is the Fisher information of probability of quantum trajectory with a time-rescaling parameter (see (9)). The Fisher information is the sum of the dynamical activity and the quantum correction \mathcal{Q} (\mathcal{Q}_2 in Ref.[26]) which vanishes for the classical case. The calculation of F is not light since one has to sum over contributions from a huge number of trajectories. Therefore, upper and lower bounds are useful in practical calculations. Vu and Saito

demonstrated that F is upper bounded by the symmetric logarithmic derivative (SLD) Fisher information:

$$\begin{aligned} \mathcal{I} &:= 4 \left[\partial_{\theta_1} \partial_{\theta_2} \mathcal{C}(\theta_1, \theta_2) \right. \\ &\quad \left. - \partial_{\theta_1} \mathcal{C}(\theta_1, \theta_2) \partial_{\theta_2} \mathcal{C}(\theta_1, \theta_2) \right] \Big|_{\theta_1=0=\theta_2}, \quad (2) \\ \mathcal{C}(\theta_1, \theta_2) &:= \text{Tr}_S \rho^{\theta_1, \theta_2}(\tau). \quad (3) \end{aligned}$$

Here, $\rho^{\theta_1, \theta_2}(t)$ is the solution of the two-sided GKSL equation (see (12)) [27]. Tr_S is the trace of the system. Using this, Vu and Saito estimated an upper bound of \mathcal{Q} in the long-time region. In this paper, we derive a concise expression of the upper bound of \mathcal{Q} for a finite time and arbitrary initial state. We also derive a simple expression of a lower bound of \mathcal{Q} .

Another class of inequalities is the speed limit of state transformation. The speed limits in terms of the entropy production for the open quantum systems described by the GKSL equation have been researched [10, 28–31]. For such open quantum systems, Hasegawa [32] derived a KUR by exploiting the Mandelstam-Tamm speed limit [33, 34]. In this KUR, a dynamical activity-like quantity appeared instead of F . We point out that this quantity equals the SLD Fisher information when Hamiltonian and jump operators are time-independent. Using our expressions, we derive a quantum speed limit described by the dynamical activity when the jump operators connect the eigenstates of the system Hamiltonian.

The structure of the paper is as follows. First, we explain the Fisher information F (§II). In §III, we show that the SLD Fisher information is a sum of the dynamical activity and a quantum correction, which is the upper bound of \mathcal{Q} . In §IV, we study \mathcal{Q} and its upper and lower bounds numerically in a two-level system. Next, we study a speed limit (§V). In §VI, we summarize this paper. In Appendix A, we explain the SLD Fisher information and the two-sided GKSL equation. In Appendix B, we derive (21). In Appendix C, we analyze the upper bound of \mathcal{Q} in the long-time region. In Appendix D, we derive the

* subarusatosi@gmail.com

lower bound of \mathcal{Q} . In Appendix E, we derive (42). In Appendix F, we review Hasegawa's method and results. Appendix G is for the detailed calculations for §V.

II. FISHER INFORMATION FOR QUANTUM TRAJECTORIES

Here, we summarize techniques introduced in Ref.[26]. The GKSL equation is given by

$$\frac{d\rho(t)}{dt} = \mathcal{L}(t)\rho(t), \quad (4)$$

$$\mathcal{L}(t)\bullet := -i[H_S(t), \bullet] + \sum_k \left[L_k(t)\bullet L_k(t)^\dagger - \frac{1}{2}\{L_k(t)^\dagger L_k(t), \bullet\} \right]. \quad (5)$$

Here, $[A, B] := AB - BA$, $\{A, B\} = AB + BA$, and \bullet is an arbitrary linear operator of the system. H_S is the system Hamiltonian and $\{L_k\}$ are jump operators. The following discussion does not require the detailed balance condition. We set $\hbar = 1$.

A quantum trajectory is specified by a list of tuples $\Gamma := \{(t_1, k_1), (t_2, k_2), \dots, (t_N, k_N)\}$. Here, t_α is the time of the α -th jump by a jump operator L_{k_α} . The probability of quantum trajectory is given by [26]

$$\mathcal{P}^\theta(\Gamma) = \text{Tr}_S[M^\theta(\Gamma)\rho(0)M^\theta(\Gamma)^\dagger] \quad (6)$$

where

$$M^\theta(\Gamma) := W^\theta(\tau, t_N) \left(\prod_{\alpha=1}^N L_{k_\alpha}^\theta(t_\alpha) W^\theta(t_\alpha, t_{\alpha-1}) \right) \quad (7)$$

with $t_0 = 0$. $W^\theta(t, s)$ is defined by

$$\frac{\partial W^\theta(t, s)}{\partial t} = \left(-iH_S^\theta - \frac{1}{2} \sum_k (L_k^\theta)^\dagger L_k^\theta \right) W^\theta(t, s) \quad (8)$$

under $W^\theta(s, s) = 1$ with

$$H_S^\theta := (1 + \theta)H_S, \quad L_k^\theta := \sqrt{1 + \theta}L_k. \quad (9)$$

Here, θ is the time-rescaling real parameter. The Fisher information F is defined by

$$F := -\langle \partial_\theta^2 \ln \mathcal{P}^\theta(\Gamma) \rangle_{\theta=0}, \quad (10)$$

where $\langle \bullet \rangle$ denotes the expected value for the probability distribution $\mathcal{P}^\theta(\Gamma)|_{\theta=0}$. F is bounded as (see Appendix A)

$$F \leq \mathcal{I}. \quad (11)$$

\mathcal{I} is given by (2). The two-sided GKSL equation governing $\rho^{\theta_1, \theta_2}$ is given by

$$\frac{d\rho^{\theta_1, \theta_2}(t)}{dt} = \mathcal{L}^{\theta_1, \theta_2}(t)\rho^{\theta_1, \theta_2}(t) \quad (12)$$

under $\rho^{\theta_1, \theta_2}(0) = \rho(0)$ [27]. Here,

$$\mathcal{L}^{\theta_1, \theta_2}(t)\bullet := -iH_S^{\theta_1}\bullet + \bullet iH_S^{\theta_2} + \sum_k \left\{ L_k^{\theta_1}\bullet(L_k^{\theta_2})^\dagger - \frac{1}{2}\left[(L_k^{\theta_1})^\dagger L_k^{\theta_1}\bullet + \bullet(L_k^{\theta_2})^\dagger L_k^{\theta_2} \right] \right\}. \quad (13)$$

F can be calculated only numerically. First, we discretize time and introduce

$$\Omega_0^\theta := 1 + \left(-iH_S^\theta - \frac{1}{2} \sum_{m=1}^M (L_m^\theta)^\dagger L_m^\theta \right) \Delta t, \quad (14)$$

$$\Omega_m^\theta := L_m^\theta \sqrt{\Delta t} \quad (m = 1, \dots, M). \quad (15)$$

Here, $\Delta t := \tau/N$ with a sufficiently large integer number N . M is the number of the jump operator of the GKSL equation. The probability of quantum trajectory $\mathcal{P}^\theta(\Gamma)$ is

$$\begin{aligned} & P^\theta(\{m_i\}) \\ & := \text{Tr}_S[\Omega_{m_{N-1}}^\theta \cdots \Omega_{m_0}^\theta \rho(0) (\Omega_{m_0}^\theta)^\dagger \cdots (\Omega_{m_{N-1}}^\theta)^\dagger] \end{aligned} \quad (16)$$

in $N \rightarrow \infty$. Then by quantum jump method [35], we numerically construct each trajectory and calculate

$$\tilde{F} := - \sum_{m_0, \dots, m_{N-1}=0}^M P^\theta(\{m_i\}) \partial_\theta^2 \ln P^\theta(\{m_i\}) \Big|_{\theta=0} \quad (17)$$

which is F in $N \rightarrow \infty$.

Note that (6) is slightly different from the original definition in Ref.[26], in which the initial state is decomposed as

$$\rho(0) = \sum_\alpha p_\alpha |\alpha\rangle \langle \alpha| \quad (18)$$

where $\{|\alpha\rangle\}$ are normalized but need not be orthogonalized. The probability of quantum trajectory is defined by

$$\begin{aligned} & P^\theta(\alpha, \{m_i\}) \\ & := p_\alpha \text{Tr}_S[\Omega_{m_{N-1}}^\theta \cdots \Omega_{m_0}^\theta |\alpha\rangle \langle \alpha| (\Omega_{m_0}^\theta)^\dagger \cdots (\Omega_{m_{N-1}}^\theta)^\dagger]. \end{aligned} \quad (19)$$

The associated Fisher information is

$$\begin{aligned} \tilde{F}' & := - \sum_\alpha \sum_{m_0, \dots, m_{N-1}=0}^M P^\theta(\alpha, \{m_i\}) \\ & \quad \times \partial_\theta^2 \ln P^\theta(\alpha, \{m_i\}) \Big|_{\theta=0}. \end{aligned} \quad (20)$$

Vu-Saito's original Fisher information F' is $N \rightarrow \infty$ limit of \tilde{F}' . In general, F' does not coincide with F and depends on the decomposition (18). However the upper bound derived from the quantum Cramér-Rao theorem [36, 37] is independent of the definitions and $F' \leq \mathcal{I}$ holds (Appendix A).

III. SLD FISHER INFORMATION

In Ref.[26], the SLD Fisher information \mathcal{I} is calculated in the limit of the long-time. Here, we derive an expression for arbitrary times. The first term of (2) is reduced to the sum of the dynamical activity $B(\tau)$ and two double integrals (Appendix B):

$$\partial_{\theta_1} \partial_{\theta_2} \mathcal{C}(\theta_1, \theta_2) \Big|_{\theta_1=0=\theta_2} = \frac{1}{4} B(\tau) + I_1 + I_2, \quad (21)$$

where

$$B(t) := \int_0^t ds \sum_k \text{Tr}_S [L_k(s) \rho(s) L_k(s)^\dagger], \quad (22)$$

$$I_1 := \int_0^\tau ds \int_0^s du \text{Tr}_S [\mathcal{L}_2(s) \mathcal{U}(s, u) \mathcal{L}_1(u) \rho(u)], \quad (23)$$

$$I_2 := \int_0^\tau ds \int_0^s du \text{Tr}_S [\mathcal{L}_1(s) \mathcal{U}(s, u) \mathcal{L}_2(u) \rho(u)] \quad (24)$$

with

$$\mathcal{L}_1 \bullet := -iH_S \bullet + \frac{1}{2} \sum_k [L_k \bullet L_k^\dagger - L_k^\dagger L_k \bullet], \quad (25)$$

$$\mathcal{L}_2 \bullet := \bullet iH_S + \frac{1}{2} \sum_k [L_k \bullet L_k^\dagger - \bullet L_k^\dagger L_k]. \quad (26)$$

Here, $\mathcal{U}(s, u)$ is defined by

$$\frac{\partial \mathcal{U}(s, u)}{\partial s} = \mathcal{L}(s) \mathcal{U}(s, u) \quad (27)$$

with $\mathcal{U}(u, u) = 1$. The second term of (2) is calculated as

$$\begin{aligned} & -\partial_{\theta_1} \mathcal{C}(\theta_1, \theta_2) \partial_{\theta_2} \mathcal{C}(\theta_1, \theta_2) \Big|_{\theta_1=0=\theta_2} \\ &= -\prod_{i=1}^2 \int_0^\tau ds \text{Tr}_S [\mathcal{L}_i(s) \rho(s)] \\ &= -\left(\int_0^\tau ds \text{Tr}_S [H_S(s) \rho(s)] \right)^2 =: I_3. \end{aligned} \quad (28)$$

Here, we used

$$\text{Tr}_S [\mathcal{L}_1 \bullet] = -i \text{Tr}_S [H_S \bullet] = -\text{Tr}_S [\mathcal{L}_2 \bullet]. \quad (29)$$

Thus, the SLD Fisher information is given by

$$\mathcal{I} = B(\tau) + Q_+(\tau), \quad (30)$$

$$Q_+ := 4(I_1 + I_2 + I_3). \quad (31)$$

Here, Q_+ is the upper bound of the quantum correction $\mathcal{Q} := F - B$.

$Q_+(t)$ can be written as (Appendix B)

$$Q_+ = 2\text{Re}(Q_a) \quad (32)$$

with

$$Q_a(t) := 4 \int_0^t ds \int_0^s du \text{Tr}_S [\mathcal{L}_2(t) \mathcal{U}(s, u) \mathcal{P}(u) \mathcal{L}_1(u) \rho(u)]. \quad (33)$$

Here, $\text{Re}(x)$ is the real part of x and

$$\mathcal{P}(u) \bullet := \bullet - \rho(u) \text{Tr}_S [\bullet]. \quad (34)$$

The double time integral is given by the following coupled differential equations,

$$\frac{dQ_a}{dt} = 4\text{Tr}_S [\mathcal{L}_2(t) q_a(t)], \quad (35)$$

$$\frac{dq_a}{dt} = \mathcal{P}(t) \mathcal{L}_1(t) \rho(t) + \mathcal{L}(t) q_a(t) \quad (36)$$

with $Q_a(0) = 0$, $q_a(0) = 0$, and the GKSL equation (4). (35), (36), and (4) can be solved by standard numerical methods such as the Runge-Kutta method. (32), (35), and (36) are the first main results of this paper. In Appendix C, we check Q_+ in the long-time region reproduces Ref.[26].

The lower bound of \mathcal{Q} is also given by a double time integral (Appendix D):

$$\begin{aligned} \mathcal{Q} &\geq Q_- := B(\tau)^2 \\ &\quad - 2 \int_0^\tau ds \int_0^s du \text{Tr}_S [\hat{\gamma}(s) \mathcal{U}(s, u) \hat{\gamma}(u) \rho(u)] \\ &=: B(\tau)^2 + Q_c(\tau). \end{aligned} \quad (37)$$

Here,

$$\begin{aligned} \hat{\gamma}(t) \bullet &:= \left(-iH_S - \frac{1}{2} \sum_k L_k^\dagger L_k \right) \bullet \\ &\quad + \bullet \left(iH_S - \frac{1}{2} \sum_k L_k^\dagger L_k \right). \end{aligned} \quad (38)$$

$Q_c(t)$ can be calculated in the same way as Q_a .

IV. NUMERICAL ANALYSIS

In the following, we suppose that H_S and L_k are time-independent.

As an example, we consider a system described by

$$H_S = \Delta |1\rangle\langle 1| + \Omega(|0\rangle\langle 1| + |1\rangle\langle 0|), \quad (39)$$

$$L_1 = \sqrt{\gamma n} |1\rangle\langle 0|, \quad (40)$$

$$L_2 = \sqrt{\gamma(n+1)} |0\rangle\langle 1|. \quad (41)$$

Figure 1 shows the time dependence of the dynamical activity B and the quantum correction \mathcal{Q} and its upper and lower bounds (Q_+ and Q_-). F was calculated by using (14), (15), (16) and (17) with the quantum jump method [35]. In Fig.1, we set $\gamma\Delta t = 0.001$ and used 10^6 trajectories. In both panels, we observe $Q_- \leq \mathcal{Q} \leq Q_+$ holds. Figure 1 (a) shows that the quantum correction \mathcal{Q} of this example is comparable to the dynamical activity B . At short times, \mathcal{Q} also takes negative values and Q_- provides a good lower bound as Fig.1 (b) shows. After the state relaxes to the steady state, Q_+ increases with a slope of \dot{Q}_+ . Here, $\dot{Q}_+ := \lim_{\tau \rightarrow \infty} Q_+/\tau$ is given by [26] (Appendix E)

$$\dot{Q}_+ = \frac{8\mathfrak{A}}{\gamma y^3 [4(\Delta^2 + 2\Omega^2) + \gamma^2 y^2]^3}, \quad (42)$$

$$\begin{aligned} \mathfrak{A} := & \Delta^2 x (4\Delta^2 + \gamma^2 y^2)^3 + 8\Omega^2 x (4\Delta^2 + \gamma^2 y^2)^2 (6\Delta^2 + \gamma^2 y^2) \\ & + 16\Omega^4 (\gamma^2 \Delta^2 y^2 (100x + 1) + \gamma^4 y^4 (12x + 1) + 4\Delta^4 (52x + 1)) \\ & + 256\Omega^6 (\gamma^2 (6x + 1) y^2 + 2\Delta^2 (12x + 1)) + 1024\Omega^8 y^2 \end{aligned} \quad (43)$$

with $x := n(n + 1)$ and $y := 2n + 1$.

V. QUANTUM SPEED LIMIT

In this section, we discuss a quantum speed limit derived by Hasegawa [32]:

$$\frac{1}{2} \int_{t_1}^{t_2} dt \sqrt{\mathcal{J}(t)} \geq D(\sigma(t_1), \sigma(t_2)). \quad (44)$$

Here, $\sigma(t) := |\Psi_\tau(\tau; t)\rangle\langle\Psi_\tau(\tau; t)|$. (44) is the Mandelstam-Tamm relation [33, 34] applied to a state $|\Psi_\tau(s; t)\rangle$ defined by

$$|\Psi_\tau(s; t)\rangle := V_\tau(s; t) |\tilde{\psi}(0)\rangle \otimes |0\rangle \quad (45)$$

with

$$\begin{aligned} V_\tau(s; t) := & \text{T exp} \left[\int_0^s du \left\{ -i \frac{t}{\tau} H_S \left(\frac{t}{\tau} u \right) \right. \right. \\ & + \sqrt{\frac{t}{\tau}} \sum_k [L_k \left(\frac{t}{\tau} u \right) \otimes \phi_k^\dagger(u) \\ & \left. \left. - L_k \left(\frac{t}{\tau} u \right)^\dagger \otimes \phi_k(u)] \right\} \right]. \end{aligned} \quad (46)$$

Here, $|\tilde{\psi}(0)\rangle$ is a purification of $\rho(0)$ ($\text{Tr}_A[|\tilde{\psi}(0)\rangle\langle\tilde{\psi}(0)|] = \rho(0)$ where A is the ancilla system). T is the time ordering operator. $\{\phi_k(t)\}$ are field operators having the canonical commutation relation

$$[\phi_k(t), \phi_l^\dagger(s)] = \delta_{kl} \delta(t - s) \quad (47)$$

and $|0\rangle$ is the vacuum state for the fields. $\mathcal{J}(t)$ is the SLD Fisher information for time (see (A16)):

$$\begin{aligned} \mathcal{J}(t) := & 4[\langle \partial_t \Psi_\tau(\tau; t) | \partial_t \Psi_\tau(\tau; t) \rangle \\ & - \langle \partial_t \Psi_\tau(\tau; t) | \Psi_\tau(\tau; t) \rangle \langle \Psi_\tau(\tau; t) | \partial_t \Psi_\tau(\tau; t) \rangle] \end{aligned} \quad (48)$$

where $|\partial_t \Psi_\tau(\tau; t)\rangle := \partial_t |\Psi_\tau(\tau; t)\rangle$. In Mandelstam-Tamm relation, $\mathcal{J}(t)/4$ is reduced to the energy fluctuation. In Hasegawa's theory, the Hamiltonian becomes $i \frac{\partial V_\tau(\tau; t)}{\partial t} V_\tau(\tau; t)^\dagger$, which does not related to the real system dynamics and thus the physical meaning may not be so clear. $D(\rho, \sigma)$ is the Bures angle:

$$D(\rho, \sigma) := \cos^{-1} F(\rho, \sigma), \quad (49)$$

$$F(\rho, \sigma) := \text{Tr} \sqrt{\sqrt{\rho} \sigma \sqrt{\rho}} = F(\sigma, \rho). \quad (50)$$

$F(\rho, \sigma)$ is the fidelity [38]. Because of the contractivity of the Bures angle (p.414 of Ref.[38]) and $\rho(t) = \text{Tr}_{AB} \sigma(t)$ (B is the field system),

$$D(\sigma(t_1), \sigma(t_2)) \geq D(\rho(t_1), \rho(t_2)) \quad (51)$$

holds. Note that the trace distance $T(\rho_1, \rho_2) := \frac{1}{2} \text{Tr}_S \sqrt{(\rho_1 - \rho_2)^2}$ is smaller than the Bures angle $D(\rho_1, \rho_2) \geq T(\rho_1, \rho_2)$ [38]. For time-independent H_S and L_k , we recognize that the two SLD Fisher information (2) and (48) are connected (Appendix F)

$$\mathcal{J}(t) = \frac{\mathcal{I}(t)}{t^2} = \frac{B(t) + Q_+(t)}{t^2}. \quad (52)$$

The quantum correction Q_+ can be eliminated in the interaction picture $\tilde{\rho}(t) := e^{iH_S t} \rho(t) e^{-iH_S t}$ when

$$[L_k, H_S] = \omega_k L_k, \quad (53)$$

where ω_k is a real number. The quantum master equation for $\tilde{\rho}(t)$ is given by

$$\frac{d\tilde{\rho}}{dt} = \sum_k [L_k \tilde{\rho}(t) L_k^\dagger - \frac{1}{2} \{L_k^\dagger L_k, \tilde{\rho}(t)\}]. \quad (54)$$

Repeating the arguments from (44) to (52), we obtain a quantum speed limit of the system expressed with the dynamical activity:

$$l(t_1, t_2) := \frac{1}{2} \int_{t_1}^{t_2} dt \frac{\sqrt{B(t)}}{t} \geq D(\tilde{\rho}(t_1), \tilde{\rho}(t_2)). \quad (55)$$

Here, $0 \leq t_1 \leq t_2$. (55) is the second main result of this paper.

As an instance, we consider a spinless quantum dot coupled to a single lead,

$$\frac{d\rho}{dt} = -i[H_S, \rho] + \gamma[1 - f(\varepsilon)]\mathcal{D}[a](\rho) + \gamma f(\varepsilon)\mathcal{D}[a^\dagger](\rho) \quad (56)$$

where $H_S = \varepsilon a^\dagger a$ and $\mathcal{D}[X](\bullet) := X \bullet X^\dagger - \frac{1}{2} \{X^\dagger X, \bullet\}$. Here, a is the annihilation operator of the electron of the system, ε is the energy level of the system, $f(\varepsilon) = \frac{1}{e^{\beta\varepsilon} + 1}$ is the Fermi distribution, β is the inverse temperature of the lead, and γ is the coupling strength. The jump operators are $L_1 = \sqrt{\gamma[1 - f(\varepsilon)]}a$ and $L_2 = \sqrt{\gamma f(\varepsilon)}a^\dagger$




FIG. 1: Q , Q_+ , Q_- , and B for (a) $\gamma t \leq 10$ and (b) $\gamma t \leq 2$. The horizontal axes are γt . We set $\Omega = \gamma$, $\Delta = 0.25\gamma$, $n = 1$ and $\rho(0) = \frac{1}{2}(1 + 0.2\sigma_x + 0.3\sigma_y - 0.4\sigma_z)$. Here, σ_i is the Pauli matrix ($\sigma_y = -i|0\rangle\langle 1| + i|1\rangle\langle 0|$, etc.).

with $\omega_1 = \varepsilon$ and $\omega_2 = -\varepsilon$. (See Appendix G for detailed calculations). Figure 2(a) shows the Bures angle D , the trace distance T , and geometric length l as functions of final time t_2 . In Fig.2, we set the initial time $\gamma t_1 = 2$. Figure 2(b) shows that the bound achievement ratio $D(\tilde{\rho}(t_1), \tilde{\rho}(t_2))/l(t_1, t_2)$ becomes greater than 0.8 around $\gamma t_2 = 2$. In Fig.3, we showed the results when the initial time $\gamma t_1 = 0$. Around $\gamma t = 0$, because of $B(t) \propto t$ and $l(0, t) \propto \sqrt{t}$, the bound achievement ratio grows in square root of time $D(\tilde{\rho}(0), \tilde{\rho}(t))/l(0, t) \propto \sqrt{t}$. The square root dependence makes the upper bound looser in the short time regime as shown in Fig.3(a).

In the open classical system described by the master equation $\frac{d}{dt}p_n = \sum_m W_{nm}p_m$ with the local detailed balance condition, the following speed limits

$$A(\tau) \geq \frac{1}{2}l(p(0), p(\tau)), \quad (57)$$

$$\sqrt{\frac{A(\tau)\sigma}{2}} \geq \frac{1}{2}l(p(0), p(\tau)) \quad (58)$$

have been known [8, 39]. Here, $l(p(0), p(\tau)) := \sum_n |p_n(0) - p_n(\tau)|$ is L_1 -norm, σ is the total entropy production, and $A(\tau) := \int_0^\tau dt \sum_{n \neq m} W_{nm}p_m$ is the dynamical activity. Naive extensions to the quantum regime may be,

$$B(\tau) \geq T(\tilde{\rho}(0), \tilde{\rho}(\tau)), \quad (59)$$

$$\sqrt{\frac{B(\tau)\sigma}{2}} \geq T(\tilde{\rho}(0), \tilde{\rho}(\tau)). \quad (60)$$

However, we numerically checked that they fail even in the spinless quantum dot due to the quantum effect. Actually the quantum extensions of (58) contain nontrivial quantum corrections [10, 28, 31]. However the quantum

extension of (57) has been less investigated. We speculate

$$\sqrt{B(\tau)} \geq D(\tilde{\rho}(0), \tilde{\rho}(\tau)) \quad (61)$$

from the discussion of Fig. 3. However, from (55), we only could derive a looser bound,

$$\sqrt{b_{\max}(\tau)\tau} \geq D(\tilde{\rho}(0), \tilde{\rho}(\tau)) \quad (62)$$

with

$$b_{\max}(\tau) := \max_{0 \leq t \leq \tau} \frac{1}{t}B(t). \quad (63)$$

Here, we used $\sqrt{b_{\max}(\tau)\tau} \geq l(0, \tau)$. Thus (61) would be correct in the short time limit. Derivation of a simple quantum speed limit at zero initial time is a future work. The extension to the first passage time [40–42] for open quantum systems is also an interesting problem.

VI. SUMMARY

We investigated the symmetric logarithm derivative (SLD) Fisher information, which appears in the context of KUR and is the upper bound of the Fisher information of the quantum trajectory for the time-rescaling parameter. For a finite time and arbitrary initial state, we derived a concise expression of the SLD Fisher information using a double time integral, which can be calculated by numerically solving coupled first-order ordinary differential equations. We also derived a simple lower bound of the Fisher information for the probability of quantum trajectory. Furthermore, we pointed out that for time-independent system, the SLD Fisher information divided by time squared is identical to the SLD Fisher information appeared in the Mandelstam-Tamm speed




FIG. 2: (a) $l(t_1, t_2)$, $D(\tilde{\rho}(t_1), \tilde{\rho}(t_2))$, and $T(\tilde{\rho}(t_1), \tilde{\rho}(t_2))$, (b) $D(\tilde{\rho}(t_1), \tilde{\rho}(t_2))/l(t_1, t_2)$, and $T(\tilde{\rho}(t_1), \tilde{\rho}(t_2))/l(t_1, t_2)$ for $\gamma t_1 = 2$. The horizontal axes are γt_2 . $\beta\varepsilon = 10$, $\rho(0) = \frac{1}{2}(1 - 0.5\sigma_x + 0.3\sigma_y + 0.2\sigma_z)$. σ_i is the Pauli matrix.




FIG. 3: (a) $l(t_1, t_2)$, $D(\tilde{\rho}(t_1), \tilde{\rho}(t_2))$, and $T(\tilde{\rho}(t_1), \tilde{\rho}(t_2))$, (b) $D(\tilde{\rho}(t_1), \tilde{\rho}(t_2))/l(t_1, t_2)$, and $T(\tilde{\rho}(t_1), \tilde{\rho}(t_2))/l(t_1, t_2)$ for $\gamma t_1 = 0$. The horizontal axes are γt_2 . Other parameters are same with Fig.2.

limit by Hasegawa [32]. Based on this observation, we showed that when the jump operators connects energy eigenstates, the upper bound of the Bures angle between the initial and final states in the interaction picture is expressed with the square root of the dynamical activity.

ACKNOWLEDGMENTS

We acknowledge helpful discussions with Tan Van Vu. This work was supported by JSPSKAKENHI Grants No. 18KK0385, and No. 20H01827.

Appendix A: SLD Fisher information and the two-sided GKSL equation

1. Continuous measurement

We introduce $(M + 1)$ -dimensional Hilbert space \mathcal{H} with an orthonormal basis $\{|m\rangle\}_{m=0}^M$ and a fictitious environment system E of which Hilbert space is $\mathcal{H}^{\otimes N}$.

We consider a combined system of S , the ancilla system A , and E . We suppose that the initial state of the combined system is $|\psi(0)\rangle \otimes |0_{N-1}, \dots, 0_1, 0_0\rangle$. Here, $|\tilde{\psi}(0)\rangle$ is the purification of $\rho(0)$ (*i.e.* $\text{Tr}_A[|\tilde{\psi}(0)\rangle\langle\tilde{\psi}(0)|] = \rho(0)$), and $|0_{N-1}, \dots, 0_1, 0_0\rangle = \otimes_{i=0}^{N-1} |0\rangle_i$. For each $i = 0, 1, \dots, N-1$, an environmental subspace i interacts with system S during the time interval $[i\Delta t, (i+1)\Delta t]$ via a unitary operator U_i . Here, $\Delta t := \tau/N$. The state of the combined system at time τ is given by

$$\begin{aligned} |\psi^\theta\rangle &= U_{N-1} \cdots U_1 U_0 |\tilde{\psi}(0)\rangle \otimes |0_{N-1}, \dots, 0_1, 0_0\rangle \\ &= \sum_{m_0, \dots, m_{N-1}=0}^M \Omega_{m_{N-1}}^\theta \cdots \Omega_{m_0}^\theta |\tilde{\psi}(0)\rangle \\ &\quad \otimes |m_{N-1}, \dots, m_1, m_0\rangle \end{aligned} \quad (\text{A1})$$

where $\Omega_{m_i}^\theta$ is defined by

$$\langle k_S | \Omega_{m_i}^\theta | k'_S \rangle = \langle k_S |_i \langle m_i | U_i | k'_S \rangle | 0 \rangle_i. \quad (\text{A2})$$

$|k_S\rangle$ and $|k'_S\rangle$ are bases of the system. We suppose that Ω_m^θ are the same as (14) and (15). The Fisher information associated with POVM (positive operator valued measure) [38] \mathcal{M} is denoted

by $I(\theta, \mathcal{M})$. If we put $\mathcal{M}_0(\{m_i\}) := 1_{SA} \otimes |m_{N-1}, \dots, m_1, m_0\rangle\langle m_{N-1}, \dots, m_1, m_0|$ (1_{SA} is the identity operator of SA), the outcome is given by

$$\text{Tr}[\mathcal{M}_0(\{m_i\})|\psi^\theta\rangle\langle\psi^\theta|] = P^\theta(\{m_i\}). \quad (\text{A3})$$

Thus,

$$\tilde{F} = I(0, \mathcal{M}_0(\{m_i\})) \quad (\text{A4})$$

holds. Because of the quantum Cramér-Rao theorem [36, 37],

$$I(\theta, \mathcal{M}) \leq J^\theta \quad (\text{A5})$$

holds. Here, J^θ is the SLD Fisher information given by (A16). Using (A4), (A5) and (A16), we obtain

$$F \leq \mathcal{I} = 4 \left[\frac{\partial^2}{\partial\theta_1 \partial\theta_2} \langle\psi^{\theta_2}|\psi^{\theta_1}\rangle - \left(\frac{\partial}{\partial\theta_2} \langle\psi^{\theta_2}|\psi^{\theta_1}\rangle \right) \left(\frac{\partial}{\partial\theta_1} \langle\psi^{\theta_2}|\psi^{\theta_1}\rangle \right) \right] \Big|_{\theta_1=0=\theta_2}. \quad (\text{A6})$$

Here,

$$\langle\psi^{\theta_2}|\psi^{\theta_1}\rangle = \text{Tr}_S \rho^{\theta_1, \theta_2}(\tau), \quad (\text{A7})$$

$$\rho^{\theta_1, \theta_2}(\tau) := \text{Tr}_{AE} [|\psi^{\theta_1}\rangle\langle\psi^{\theta_2}|]. \quad (\text{A8})$$

The time evolution equation of $\rho^{\theta_1, \theta_2}(t)$ is given by (12) [27]. Then, we obtain (11).

To obtain (19), we adopt a purification

$$|\tilde{\psi}(0)\rangle = \sum_{\alpha} \sqrt{p_{\alpha}} |\alpha\rangle \otimes |\varphi_{\alpha}\rangle_A \quad (\text{A9})$$

with ${}_A\langle\varphi_{\alpha}|\varphi_{\beta}\rangle_A = \delta_{\alpha\beta}$ and the POVM $\mathcal{M}_0(\alpha, \{m_i\}) := |\varphi_{\alpha}\rangle_{AA}\langle\varphi_{\alpha}| \otimes |m_{N-1}, \dots, m_1, m_0\rangle\langle m_{N-1}, \dots, m_1, m_0|$. Then the outcomes becomes (19):

$$\text{Tr}[\mathcal{M}_0(\alpha, \{m_i\})|\psi^\theta\rangle\langle\psi^\theta|] = P^\theta(\alpha, \{m_i\}). \quad (\text{A10})$$

The associated Fisher information is given by

$$\tilde{F}' = I(0, \mathcal{M}_0(\alpha, \{m_i\})). \quad (\text{A11})$$

Then, $F' \leq \mathcal{I}$ holds.

2. SLD Fisher information

We consider n real parameters $\theta = (\theta^1, \dots, \theta^n)$ and a state ρ^θ . The SLD S_i^θ is defined by

$$\partial_i \rho^\theta = \frac{1}{2} (\rho^\theta S_i^\theta + S_i^\theta \rho^\theta), \quad S_i^\theta = (S_i^\theta)^\dagger. \quad (\text{A12})$$

Here, $\partial_i = \partial/\partial\theta^i$. Although S_i^θ is not unique in general, the SLD Fisher information matrix

$$J_{ij}^\theta := \frac{1}{2} \text{Tr} \left[\rho^\theta \{S_i^\theta, S_j^\theta\} \right] \quad (\text{A13})$$

is unique [37]. J_{ij}^θ can be rewritten as $J_{ij}^\theta = \text{Tr}[\partial_i \rho^\theta S_j^\theta]$.

In the following, we consider a pure state ρ^θ . Differentiating $(\rho^\theta)^2 = \rho^\theta$, we obtain

$$\partial_i \rho^\theta = (\partial_i \rho^\theta) \rho^\theta + \rho^\theta \partial_i \rho^\theta. \quad (\text{A14})$$

Thus, $2\partial_i \rho^\theta$ is an SLD. Using this relation and denoting $\rho^\theta = |\psi^\theta\rangle\langle\psi^\theta|$, we obtain

$$J_{ij}^\theta = 4 \text{Re} \left[\langle\partial_i \psi^\theta|\partial_j \psi^\theta\rangle - \langle\partial_i \psi^\theta|\psi^\theta\rangle\langle\psi^\theta|\partial_j \psi^\theta\rangle \right]. \quad (\text{A15})$$

Here, $|\partial_i \psi^\theta\rangle := \partial_i |\psi^\theta\rangle$. For $n = 1$, $J^\theta := J_{11}^\theta$ becomes

$$J^\theta = 4 \left[\langle\partial_\theta \psi^\theta|\partial_\theta \psi^\theta\rangle - \langle\partial_\theta \psi^\theta|\psi^\theta\rangle\langle\psi^\theta|\partial_\theta \psi^\theta\rangle \right]. \quad (\text{A16})$$

Appendix B: Derivation of (21)

In the following, we use the Liouville space. An arbitrary linear operator X of the system is described by a vector $|X\rangle\rangle$. The inner product is defined by $\langle\langle Y|X\rangle\rangle := \text{Tr}_S(Y^\dagger X)$. In particular, $\langle\langle 1|X\rangle\rangle = \text{Tr}_S(X)$. An arbitrary linear super-operator of the system is described by an operator of Liouville space. The conservation of the probability leads to $\langle\langle 1|\mathcal{L}(t)\rangle\rangle = 0$.

$\mathcal{C}(\theta_1, \theta_2)$ is given by

$$\mathcal{C}(\theta_1, \theta_2) = \langle\langle 1|\mathcal{U}^{\theta_1, \theta_2}(\tau, 0)|\rho(0)\rangle\rangle \quad (\text{B1})$$

where $\mathcal{U}^{\theta_1, \theta_2}(u, s)$ is defined by

$$\frac{\partial \mathcal{U}^{\theta_1, \theta_2}(u, s)}{\partial u} = \mathcal{L}^{\theta_1, \theta_2}(u) \mathcal{U}^{\theta_1, \theta_2}(u, s) \quad (\text{B2})$$

with $\mathcal{U}^{\theta_1, \theta_2}(s, s) = 1$. Note that $\mathcal{U}^{0,0}(u, s) = \mathcal{U}(u, s)$. The first derivative reads to

$$\begin{aligned} \partial_{\theta_i} \mathcal{C}(\theta_1, \theta_2) &= \int_0^\tau du \langle\langle 1|\mathcal{U}^{\theta_1, \theta_2}(\tau, u) \partial_{\theta_i} \mathcal{L}^{\theta_1, \theta_2}(u) \\ &\quad \times \mathcal{U}^{\theta_1, \theta_2}(u, 0)|\rho(0)\rangle\rangle \quad (i = 1, 2). \end{aligned} \quad (\text{B3})$$

Using the above equation, $\mathcal{U}^{0,0}(s, 0)|\rho(0)\rangle\rangle = |\rho(s)\rangle\rangle$, and

$$\langle\langle 1|\mathcal{U}^{0,0}(s, 0)\rangle\rangle = \langle\langle 1|, \quad (\text{B4})$$

we obtain (28). (B4) is derived from $\langle\langle 1|\mathcal{L}(s)\rangle\rangle = 0$. The second derivative (21) consists of

$$\int_0^\tau ds \text{Tr}_S \left[\partial_{\theta_1} \partial_{\theta_2} \mathcal{L}^{\theta_1, \theta_2}(s) \Big|_{\theta_1=0=\theta_2} \rho(s) \right] = \frac{1}{4} B(\tau) \quad (\text{B5})$$

and

$$\begin{aligned} \int_0^\tau ds \int_0^s du \text{Tr}_S \left[\partial_{\theta_i} \mathcal{L}^{\theta_1, \theta_2}(s) \mathcal{U}^{\theta_1, \theta_2}(s, u) \right. \\ \left. \times \partial_{\theta_i} \mathcal{L}^{\theta_1, \theta_2}(u) \Big|_{\theta_1=0=\theta_2} \rho(u) \right] = I_i \quad (i = 1, 2) \end{aligned} \quad (\text{B6})$$

with $1' := 2$ and $2' := 1$. Note that

$$\mathcal{L}_i(s) = \partial_{\theta_i} \mathcal{L}^{\theta_1, \theta_2}(s) \Big|_{\theta_1=0=\theta_2}. \quad (\text{B7})$$

$Q_+(t)$ can be written as

$$Q_+(t) = Q_a(t) + Q_b(t), \quad (\text{B8})$$

where

$$\begin{aligned} Q_b(t) &:= 4 \int_0^t ds \int_0^s du \text{Tr}_S \left[\mathcal{L}_1(t) \mathcal{U}(s, u) \mathcal{P}(u) \mathcal{L}_2(u) \rho(u) \right]. \end{aligned} \quad (\text{B9})$$

q_a in (35) is defined by

$$q_a(s) := \int_0^s du \mathcal{U}(s, u) \mathcal{P}(u) \mathcal{L}_1(u) \rho(u). \quad (\text{B10})$$

Because of

$$(\mathcal{L}\bullet)^\dagger = \mathcal{L}\bullet^\dagger, \quad (\text{B11})$$

$$(\mathcal{L}_i\bullet)^\dagger = \mathcal{L}_{i'}\bullet^\dagger, \quad (\text{B12})$$

$$(\mathcal{P}\mathcal{L}_i\bullet)^\dagger = \mathcal{P}\mathcal{L}_{i'}\bullet^\dagger, \quad (\text{B13})$$

$Q_b = Q_a^*$ holds. Then, we obtain (32).

Appendix C: Quantum correction Q_+ in the long-time region

The left and right eigenvalue equations of the Liouvillian \mathcal{L} are

$$\mathcal{L}|\rho_n\rangle\rangle = \lambda_n|\rho_n\rangle\rangle, \quad (\text{C1})$$

$$\langle\langle l_n|\mathcal{L} = \lambda_n\langle\langle l_n|. \quad (\text{C2})$$

We set $\langle\langle l_m|\rho_n\rangle\rangle = \delta_{mn}$, $\lambda_0 = 0$, and $\langle\langle l_0| = \langle\langle 1|$. Then, $\rho_0(=:\rho^{\text{ss}})$ is the steady state.

If the initial state is the steady state, (B8) becomes

$$\begin{aligned} Q_+(\tau) &= 4 \left(\int_0^\tau ds \int_0^s du \text{Tr}_S \left[\mathcal{L}_2 e^{(s-u)\mathcal{L}} \mathcal{P} \mathcal{L}_1 \rho^{\text{ss}} \right] \right. \\ &\quad \left. + \int_0^\tau ds \int_0^s du \text{Tr}_S \left[\mathcal{L}_1 e^{(s-u)\mathcal{L}} \mathcal{P} \mathcal{L}_2 \rho^{\text{ss}} \right] \right) \end{aligned} \quad (\text{C3})$$

with

$$\mathcal{P} := 1 - |\rho^{\text{ss}}\rangle\rangle\langle\langle 1|. \quad (\text{C4})$$

Using

$$e^{(s-u)\mathcal{L}} = \sum_n e^{(s-u)\lambda_n} |\rho_n\rangle\rangle\langle\langle l_n|, \quad (\text{C5})$$

we obtain

$$\begin{aligned} &\int_0^\tau ds \int_0^s du e^{(s-u)\mathcal{L}} \mathcal{P} \\ &= \sum_{n \neq 0} \int_0^\tau ds \int_0^s du e^{(s-u)\lambda_n} \mathcal{P} |\rho_n\rangle\rangle\langle\langle l_n| \\ &= -\tau \sum_{n \neq 0} \frac{1}{\lambda_n} |\rho_n\rangle\rangle\langle\langle l_n| - \sum_{n \neq 0} \frac{1}{\lambda_n^2} |\rho_n\rangle\rangle\langle\langle l_n| \\ &\quad + \sum_{n \neq 0} \frac{e^{\tau\lambda_n}}{\lambda_n^2} |\rho_n\rangle\rangle\langle\langle l_n| \\ &\approx -\tau \sum_{n \neq 0} \frac{1}{\lambda_n} |\rho_n\rangle\rangle\langle\langle l_n| = -\tau \mathcal{R}, \end{aligned} \quad (\text{C6})$$

where

$$\mathcal{R} := \sum_{n \neq 0} \frac{1}{\lambda_n} |\rho_n\rangle\rangle\langle\langle l_n| = -\int_0^\infty dt e^{t\mathcal{L}} \mathcal{P} \quad (\text{C7})$$

is the pseudo-inverse of the Liouvillian [24, 43–47] (the Drazin inverse [48] of \mathcal{L}). Thus, we obtain

$$Q_+(\tau) \approx -4\tau \left(\text{Tr}_S \left[\mathcal{L}_2 \mathcal{R} \mathcal{L}_1 \rho^{\text{ss}} \right] + \text{Tr}_S \left[\mathcal{L}_1 \mathcal{R} \mathcal{L}_2 \rho^{\text{ss}} \right] \right) \quad (\text{C8})$$

in the long-time limit. The expression of Q_+ in the long-time approximation is consistent with (S63) in Ref.[26].

Appendix D: Lower bound of Q

(17) can be rewritten as

$$\tilde{F} = \sum_{m_0, \dots, m_{N-1}=0}^M P^0(\{m_i\}) \left[\partial_\theta \ln P^\theta(\{m_i\})|_{\theta=0} \right]^2. \quad (\text{D1})$$

The score $\partial_\theta \ln P^\theta(\{m_i\})|_{\theta=0}$ is given by

$$\partial_\theta \ln P^\theta(\{m_i\})|_{\theta=0} = b(\{m_i\}) + d(\{m_i\}) \quad (\text{D2})$$

where

$$b(\{m_i\}) := \sum_{n=0}^{N-1} \delta_{m_n \neq 0}, \quad (\text{D3})$$

$$d(\{m_i\}) := \sum_{n=0}^{N-1} \delta_{m_n, 0} \gamma_n(\{m_i\}) \Delta t, \quad (\text{D4})$$

$$\gamma_n(\{m_i\}) := \frac{\text{Tr}_S [\hat{\Omega}_{m_{N-1}} \cdots \hat{\gamma} \cdots \hat{\Omega}_{m_0} \rho(0)]}{P^0(\{m_i\})}. \quad (\text{D5})$$

Here we introduce the convention that $\delta_{m \neq 0} := 1 - \delta_{m, 0}$ and $\hat{\Omega}_m \bullet := \Omega_m^0 \bullet (\Omega_m^0)^\dagger$. $\hat{\gamma}$ is defined by (38). Then \tilde{F} reads to

$$\tilde{F} = \langle b(\{m_i\})^2 \rangle + 2 \langle b(\{m_i\}) d(\{m_i\}) \rangle + \langle d(\{m_i\})^2 \rangle \quad (\text{D6})$$

where $\langle X(\{m_i\}) \rangle := \sum_{m_0, \dots, m_{N-1}=0}^M P^0(\{m_i\}) X(\{m_i\})$. $b(\{m_i\})^2$ and $b(\{m_i\})d(\{m_i\})$ are calculated as

$$b(\{m_i\})^2 = b(\{m_i\}) + 2 \sum_{n=1}^{N-1} \sum_{l=0}^{n-1} \delta_{m_n \neq 0} \delta_{m_l \neq 0}, \quad (\text{D7})$$

and

$$b(\{m_i\})d(\{m_i\}) = \sum_{n=1}^{N-1} \sum_{l=0}^{n-1} \left(\delta_{m_n \neq 0} \gamma_l(\{m_i\}) \delta_{m_l, 0} \Delta t + \gamma_n(\{m_i\}) \delta_{m_n, 0} \Delta t \delta_{m_l \neq 0} \right). \quad (\text{D8})$$

In (D7), the first and second terms of the right-hand side come from the contributions of the same times and different times respectively. In the limit of $\Delta t \rightarrow 0$, trajectory averages of the above two equations become

$$\langle b(\{m_i\})^2 \rangle = B(\tau) + 2 \int_0^\tau ds \int_0^s du \text{Tr}_S[\hat{\Gamma}(s)\mathcal{U}(s, u)\hat{\Gamma}(u)\rho(u)], \quad (\text{D9})$$

$$\langle b(\{m_i\})d(\{m_i\}) \rangle = \int_0^\tau ds \int_0^s du \left(\text{Tr}_S[\hat{\Gamma}(s)\mathcal{U}(s, u)\hat{\gamma}(u)\rho(u)] + \text{Tr}_S[\hat{\gamma}(s)\mathcal{U}(s, u)\hat{\Gamma}(u)\rho(u)] \right) \quad (\text{D10})$$

where $\hat{\Gamma}(t)\bullet := \sum_k L_k \bullet L_k^\dagger$. Thus, we obtain

$$\langle b(\{m_i\})^2 \rangle + 2\langle b(\{m_i\})d(\{m_i\}) \rangle = B(\tau) - 2 \int_0^\tau ds \int_0^s du \text{Tr}_S[\hat{\gamma}(s)\mathcal{U}(s, u)\hat{\gamma}(u)\rho(u)] \quad (\text{D11})$$

and

$$\mathcal{Q} = \langle d(\{m_i\})^2 \rangle - 2 \int_0^\tau ds \int_0^s du \text{Tr}_S[\hat{\gamma}(s)\mathcal{U}(s, u)\hat{\gamma}(u)\rho(u)]. \quad (\text{D12})$$

Here, $\langle d(\{m_i\})^2 \rangle$ cannot be written as the double time integral because $\gamma_n(\{m_i\})$ in $d(\{m_i\})$ depends on the entire sequence of a trajectory $\{m_i\}$. Using $\langle d(\{m_i\})^2 \rangle \geq \langle d(\{m_i\}) \rangle^2$, $\langle b(\{m_i\}) + d(\{m_i\}) \rangle = 0$, and $\langle b(\{m_i\}) \rangle = B(\tau)$, we obtain (37).

Appendix E: Derivation of (42)

In the long-time region, the SLD Fisher information \mathcal{I} is given by

$$\mathcal{I} = 4\partial_{\theta_1}\partial_{\theta_2} \ln \text{Tr}_S \rho^{\theta_1, \theta_2}(\tau) \Big|_{\theta_1=0=\theta_2} = 4\tau \partial_{\theta_1}\partial_{\theta_2} \lambda(\theta_1, \theta_2) \Big|_{\theta_1=0=\theta_2} + O(1). \quad (\text{E1})$$

Here, $\lambda(\theta_1, \theta_2)$ is the eigenvalue of $\mathcal{L}^{\theta_1, \theta_2}$ which satisfies $\lambda(0, 0) = 0$. We suppose that the matrix representation of $\mathcal{L}^{\theta_1, \theta_2}$ is block diagonalized, with one block $\tilde{\mathcal{L}}^{\theta_1, \theta_2}$ having eigenvalue $\lambda(\theta_1, \theta_2)$. The characteristic polynomial of $\tilde{\mathcal{L}}^{\theta_1, \theta_2}$ is given by

$$\det(\tilde{\mathcal{L}}^{\theta_1, \theta_2} - \lambda 1_d) = \sum_{n=0}^d a^{(n)}(\theta_1, \theta_2) \lambda^n. \quad (\text{E2})$$

Here, d is the dimension of $\tilde{\mathcal{L}}^{\theta_1, \theta_2}$ and 1_d is d -dimensional identity matrix. By differentiating $\sum_{n=0}^d a^{(n)}(\theta_1, \theta_2) \lambda(\theta_1, \theta_2)^n = 0$, we obtain

$$a_i^{(0)} + a^{(1)} \lambda_i = 0 \quad (i = 1, 2), \quad (\text{E3})$$

$$a_{12}^{(0)} + a^{(1)} \lambda_{12} + a_1^{(1)} \lambda_2 + a_2^{(1)} \lambda_1 + 2a^{(2)} \lambda_1 \lambda_2 = 0. \quad (\text{E4})$$

Here, $a^{(n)} := a^{(n)}(0, 0)$, $X_i := \partial_{\theta_i} X|_{\theta_1=0=\theta_2}$, and $X_{12} := \partial_{\theta_1} \partial_{\theta_2} X|_{\theta_1=0=\theta_2}$. λ_{12} is calculated from (E3) and (E4). $\dot{Q}_+ := \lim_{\tau \rightarrow \infty} Q_+/\tau$ is given by

$$\dot{Q}_+ = 4\lambda_{12} - \sum_k \text{Tr}_S[L_k^\dagger L_k \rho^{\text{ss}}]. \quad (\text{E5})$$

For the system described by (39)-(41), $\tilde{\mathcal{L}}^{\theta_1, \theta_2}$ is given by

$$\tilde{\mathcal{L}}^{\theta_1, \theta_2} = \begin{pmatrix} -\frac{1}{2}(\Theta_1 + \Theta_2)\gamma n & i\Theta_2\Omega & -\Theta_1\Omega & \sqrt{\Theta_1\Theta_2}\gamma(n+1) \\ i\Theta_2\Omega & i\Theta_2\Delta - \frac{1}{2}\Theta_2\gamma(n+1) - \frac{1}{2}\Theta_1\gamma n & 0 & -i\Theta_1\Omega \\ -i\Theta_1\Omega & 0 & -i\Theta_1\Delta - \frac{1}{2}\Theta_1\gamma(n+1) - \frac{1}{2}\Theta_2\gamma n & i\Theta_2\Omega \\ \sqrt{\Theta_1\Theta_2}\gamma n & -i\Theta_1\Omega & i\Theta_2\Omega & \mathcal{L}_{44} \end{pmatrix} \quad (\text{E6})$$

with $\mathcal{L}_{44} := -\frac{1}{2}(\Theta_1 + \Theta_2)\gamma(n+1) - i(\Theta_1 - \Theta_2)\Delta$ and $\Theta_i := 1 + \theta_i$. Using (E3), (E4) and (E5), we obtain (42).

Appendix F: Hasegawa's approach

We review Hasegawa's method and results [32]. We introduce a state

$$|\Phi(t)\rangle = U(t)|\Phi(0)\rangle \quad (\text{F1})$$

with

$$|\Phi(0)\rangle := |\tilde{\psi}(0)\rangle \otimes |0\rangle, \quad (\text{F2})$$

$$U(t) = \mathbb{T} \exp \left[\int_0^t ds \left\{ -iH_S(s) + \sum_k [L_k(s) \otimes \phi_k^\dagger(s) - L_k(s)^\dagger \otimes \phi_k(s)] \right\} \right]. \quad (\text{F3})$$

The state $|\Phi(t)\rangle$ provides the solution of the GKSL equation [32]:

$$\rho(t) = \text{Tr}_{AB}[|\Phi(t)\rangle\langle\Phi(t)|]. \quad (\text{F4})$$

Using $d\rho = \text{Tr}_{AB}[d(|\Phi(t)\rangle\langle\Phi(t)|)]$,

$$d(|\Phi(t)\rangle\langle\Phi(t)|) = d|\Phi(t)\rangle\langle\Phi(t)| + |\Phi(t)\rangle d\langle\Phi(t)| + d|\Phi(t)\rangle d\langle\Phi(t)|, \quad (\text{F5})$$

and

$$d|\Phi(t)\rangle = \left((-iH_S - \frac{1}{2} \sum_k L_k^\dagger L_k) dt + \sum_k L_k d\phi_k^\dagger + \frac{1}{2} \sum_{k,l} L_k L_l d\phi_k^\dagger d\phi_l^\dagger \right) |\Phi(t)\rangle, \quad (\text{F6})$$

we obtain the GKSL equation (4). Here, $d\phi_k^\dagger = \int_t^{t+dt} ds \phi_k^\dagger(s)$.
If we put

$$\rho_\tau(s; t_1, t_2) := \text{Tr}_{AB}[|\Psi_\tau(s; t_1)\rangle\langle\Psi_\tau(s; t_2)|], \quad (\text{F7})$$

we obtain $\rho_\tau(\tau; t, t) = \rho(t)$ [32]. The time evolution equation of $\rho_\tau(s; t_1, t_2)$ is given by

$$\frac{\partial \rho_\tau(s; t_1, t_2)}{\partial s} = \mathcal{L}_\tau(s; t_1, t_2) \rho_\tau(s; t_1, t_2) \quad (\text{F8})$$

with

$$\begin{aligned} \mathcal{L}_\tau(s; t_1, t_2) \bullet &= -i \frac{t_1}{\tau} H_S \left(\frac{t_1}{\tau} s \right) \bullet + \bullet i \frac{t_2}{\tau} H_S \left(\frac{t_2}{\tau} s \right) \\ &+ \sqrt{\frac{t_1}{\tau}} \sqrt{\frac{t_2}{\tau}} \sum_k L_k \left(\frac{t_1}{\tau} s \right) \bullet L_k \left(\frac{t_2}{\tau} s \right)^\dagger \\ &- \frac{1}{2} \sum_k \left[\frac{t_1}{\tau} L_k \left(\frac{t_1}{\tau} s \right)^\dagger L_k \left(\frac{t_1}{\tau} s \right) \bullet \right. \\ &\left. + \bullet \frac{t_2}{\tau} L_k \left(\frac{t_2}{\tau} s \right)^\dagger L_k \left(\frac{t_2}{\tau} s \right) \right]. \end{aligned} \quad (\text{F9})$$

(F8) is a two-sided GKSL equation. Because of

$$\mathcal{L}_\tau(s; t, t) = \frac{t}{\tau} \mathcal{L} \left(\frac{t}{\tau} s \right), \quad (\text{F10})$$

we obtain

$$\rho_\tau(s; t, t) = \rho \left(\frac{t}{\tau} s \right). \quad (\text{F11})$$

$\mathcal{J}(t)$ defined by (48) is given by

$$\begin{aligned} \mathcal{J}(t) &= 4[\partial_{t_1} \partial_{t_2} C(t_1, t_2) \\ &- \partial_{t_1} C(t_1, t_2) \partial_{t_2} C(t_1, t_2)] \Big|_{t_1=t=t_2} \end{aligned} \quad (\text{F12})$$

with $C(t_1, t_2) := \text{Tr}_S \rho_\tau(\tau; t_1, t_2)$. $\mathcal{J}(t)$ is independent from τ .

From (44), Hasegawa [32] showed a KUR:

$$\frac{\tau^2 (\partial_\tau \langle \mathcal{C} \rangle_\tau)^2}{\langle \mathcal{C}^2 \rangle_\tau - \langle \mathcal{C} \rangle_\tau^2} \leq \mathcal{B}(\tau). \quad (\text{F13})$$

Here,

$$\mathcal{B}(t) := t^2 \mathcal{J}(t) \quad (\text{F14})$$

is ‘‘the quantum generalization of the dynamical activity’’. $\langle X \rangle_\tau := \langle \Psi_\tau(\tau; \tau) | X | \Psi_\tau(\tau; \tau) \rangle = \langle \Phi(\tau) | X | \Phi(\tau) \rangle$ and \mathcal{C} is an operator of the field system which describes a time-integrated counting observable: \mathcal{C} counts and weights jump events in a quantum trajectory of the system S . In Ref.[32], Hasegawa showed a KUR for more general operators of the field system.

If H_S and L_k are time-independent, (F9) is obtained by replacing $1 + \theta_i$ by t_i/τ ($i = 1, 2$) in (13). In this case, $\mathcal{B}(\tau)$ in Ref.[32] is identical to \mathcal{I} in Ref.[26]:

$$\mathcal{B}(\tau) = \mathcal{I}. \quad (\text{F15})$$

(F13) is consistent with

$$\frac{\tau^2 (\partial_\tau \langle \Phi \rangle)^2}{\text{var}[\Phi]} \leq \mathcal{B}(\tau), \quad (\text{F16})$$

which is derived from (1) and (11).

Appendix G: Quantum dot

We consider the quantum dot (56). The state of the system can be written as $\tilde{\rho}(t) = \frac{1}{2}(1 + \mathbf{r}(t) \cdot \boldsymbol{\sigma})$. Here, $\boldsymbol{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$, σ_i is the Pauli matrix, and $\mathbf{r}(t) = (x, y, z)$ is the Bloch vector. The equation of the motion of the Bloch vector is given by

$$\frac{d}{dt} x = -\frac{1}{2} \gamma x, \quad \frac{d}{dt} y = -\frac{1}{2} \gamma y, \quad (\text{G1})$$

$$\frac{d}{dt} z = -\gamma(z - [1 - 2f(\varepsilon)]). \quad (\text{G2})$$

The dynamical activity is given by

$$B(t) = \int_0^t ds \frac{\gamma}{2} (1 + [2f(\varepsilon) - 1]z(s)). \quad (\text{G3})$$

We put $\rho_i := \tilde{\rho}(t_i)$. If the eigenvalues of $\kappa := \sqrt{\rho_1} \rho_2 \sqrt{\rho_1}$ are λ_1 and λ_2 , the fidelity is given by $F(\rho_1, \rho_2) = \sqrt{\lambda_1} + \sqrt{\lambda_2}$. Then,

$$\begin{aligned} [F(\rho_1, \rho_2)]^2 &= \text{Tr}_S(\kappa) + 2\sqrt{\det(\kappa)} \\ &= \text{Tr}_S(\rho_1 \rho_2) + 2\sqrt{\det(\rho_1) \det(\rho_2)}. \end{aligned} \quad (\text{G4})$$

This leads to

$$F(\tilde{\rho}(t_1), \tilde{\rho}(t_2)) = \sqrt{\frac{1 + \mathbf{r}(t_1) \cdot \mathbf{r}(t_2) + \sqrt{[1 - \mathbf{r}(t_1)^2][1 - \mathbf{r}(t_2)^2]}}{2}}. \quad (\text{G5})$$

The Bures angle is given by $D(\tilde{\rho}(t_1), \tilde{\rho}(t_2)) = \cos^{-1} F(\tilde{\rho}(t_1), \tilde{\rho}(t_2))$. The trace distance is given by

$$T(\tilde{\rho}(t_1), \tilde{\rho}(t_2)) = \frac{1}{2} |\mathbf{r}(t_1) - \mathbf{r}(t_2)|. \quad (\text{G6})$$

Here, $|\mathbf{x}| = \sqrt{\mathbf{x}^2} = \sqrt{\mathbf{x} \cdot \mathbf{x}}$.

-
- [1] A. C. Barato and U. Seifert, “Thermodynamic Uncertainty Relation for Biomolecular Processes”, *Phys. Rev. Lett.* **114**, 158101 (2015).
- [2] T. R. Gingrich, J. M. Horowitz, N. Perunov, and J. L. England, “Dissipation Bounds All Steady-State Current Fluctuations”, *Phys. Rev. Lett.* **116**, 120601 (2016).
- [3] A. Dechant, and S.-i. Sasa, “Current fluctuations and transport efficiency for general Langevin systems”, *J. Stat. Mech.: Theory Exp.* **2018**, 063209.
- [4] K. Liu, Z. Gong, and M. Ueda, “Thermodynamic Uncertainty Relation for Arbitrary Initial States”, *Phys. Rev. Lett.* **125**, 140602 (2020).
- [5] T. Koyuk and U. Seifert, “Thermodynamic Uncertainty Relation for Time-Dependent Driving”, *Phys. Rev. Lett.* **125**, 260604 (2020).
- [6] A. Dechant and S.-i. Sasa, “Continuous time reversal and equality in the thermodynamic uncertainty relation”, *Phys. Rev. Research* **3**, L042012 (2021).
- [7] N. Shiraishi, “Optimal Thermodynamic Uncertainty Relation in Markov Jump Processes”, *J. Stat. Phys.* **185**, 19 (2021).
- [8] V. T. Vo, T. V. Vu, and Y. Hasegawa, “Unified thermodynamic-kinetic uncertainty relation”, *J. Phys. A: Math. Theor.* **55**, 405004 (2022).
- [9] K. Yoshimura, A. Kolchinsky, A. Dechant, and S. Ito, “Housekeeping and excess entropy production for general nonlinear dynamics”, *Phys. Rev. Research* **5**, 013017 (2023).
- [10] T. V. Vu and K. Saito, “Thermodynamic Unification of Optimal Transport: Thermodynamic Uncertainty Relation, Minimum Dissipation, and Thermodynamic Speed Limits”, *Phys. Rev. X* **13**, 011013 (2023).
- [11] J. P. Garrahan, “Simple bounds on fluctuations and uncertainty relations for first-passage times of counting observables”, *Phys. Rev. E* **95**, 032134 (2017).
- [12] I. D. Terlizzi and M. Baiesi, “Kinetic uncertainty relation”, *Journal of Physics A: Mathematical and Theoretical* **52**, 02LT03 (2018).
- [13] K. Hiura and S.-i. Sasa, “Kinetic uncertainty relation on first-passage time for accumulated current”, *Phys. Rev. E* **103**, L050103 (2021).
- [14] K. Ptaszyński, “Coherence-enhanced constancy of a quantum thermoelectric generator”, *Phys. Rev. B* **98**, 085425 (2018).
- [15] L. M. Cangemi, V. Cataudella, G. Benenti, M. Sassetti, and G. De Filippis, “Violation of thermodynamics uncertainty relations in a periodically driven work-to-work converter from weak to strong dissipation”, *Phys. Rev. B* **102**, 165418 (2020).
- [16] A. Rignon-Bret, G. Guarnieri, J. Goold, and M. T. Mitchison, “Thermodynamics of precision in quantum nanomachines”, *Phys. Rev. E* **103**, 012133 (2021).
- [17] A. A. S. Kalae, A. Wacker, and P. P. Potts, “Violating the thermodynamic uncertainty relation in the three-level maser”, *Phys. Rev. E* **104**, L012103 (2021).
- [18] P. Menczel, E. Loisa, K. Brandner, and C. Flindt, “Thermodynamic uncertainty relations for coherently driven open quantum systems”, *J. Phys. A: Math. Theor.* **54**, 314002 (2021).
- [19] P. Bayona-Pena and K. Takahashi, “Thermodynamics of a continuous quantum heat engine: Interplay between population and coherence”, *Phys. Rev. A* **104**, 042203 (2021).
- [20] F. Carollo, R. L. Jack, and J. P. Garrahan, “Unraveling the Large Deviation Statistics of Markovian Open Quantum Systems”, *Phys. Rev. Lett.* **122**, 130605 (2019).
- [21] G. Guarnieri, G. T. Landi, S. R. Clark, and John Goold, “Thermodynamics of precision in quantum nonequilibrium steady states”, *Phys. Rev. Research* **1**, 033021 (2019).
- [22] Y. Hasegawa, “Quantum Thermodynamic Uncertainty Relation for Continuous Measurement”, *Phys. Rev. Lett.* **125**, 050601 (2020).
- [23] Y. Hasegawa, “Thermodynamic Uncertainty Relation for General Open Quantum Systems”, *Phys. Rev. Lett.* **126**, 010602 (2021).
- [24] H. J. D. Miller, M. H. Mohammady. M. Perarnau-Llobet, and G. Guarnieri, “Thermodynamic Uncertainty Relation in Slowly Driven Quantum Heat Engines”, *Phys. Rev. Lett.* **126**, 210603 (2021).
- [25] Y. Hasegawa, “Irreversibility, Loschmidt Echo, and Thermodynamic Uncertainty Relation”, *Phys. Rev. Lett.* **127**, 240602 (2021).
- [26] T. V. Vu and K. Saito, “Thermodynamics of Precision in Markovian Open Quantum Dynamics”, *Phys. Rev. Lett.* **128**, 140602 (2022).
- [27] S. Gammelmark and K. Mølmer, “Fisher Information and the Quantum Cramér-Rao Sensitivity Limit of Continuous Measurements”, *Phys. Rev. Lett.* **112**, 170401 (2014).
- [28] K. Funo, N. Shiraishi, and K. Saito, “Speed limit for open quantum systems”, *New J. Phys.* **21**, 013006 (2019).
- [29] T. V. Vu and Y. Hasegawa, “Geometrical Bounds of the Irreversibility in Markovian Systems”, *Phys. Rev. Lett.* **126**, 010601 (2021).
- [30] T. V. Vu and K. Saito, “Finite-Time Quantum Landauer Principle and Quantum Coherence”, *Phys. Rev. Lett.* **128**, 010602 (2022).
- [31] S. Nakajima and Y. Utsumi, “Speed limits of the trace distance for open quantum system”, *New. J. Phys.* **24**, 095004 (2022).
- [32] Y. Hasegawa, “Thermodynamic Bounds via Bulk-Boundary Correspondence: Speed Limit, Thermodynamic Uncertainty Relation and Heisenberg Principle”,

arXiv:2203.12421v4

- [33] L. Mandelstam and I. Tamm, “The uncertainty relation between energy and time in nonrelativistic quantum mechanics”, *J. Phys. (Moscow)* **9**, 249 (1945).
- [34] S. Deffner and S. Campbell, “Quantum speed limits: from Heisenberg’s uncertainty principle to optimal quantum control”, *J. Phys. A: Math. Theor.* **50**, 453001 (2017).
- [35] A. J. Daley, “Quantum trajectories and open many-body quantum systems”, *Adv. Phys.* **63**, 77 (2014).
- [36] S. L. Braunstein and C. M. Caves, “Statistical distance and the geometry of quantum states”, *Phys. Rev. Lett.* **72**, 3439 (1994).
- [37] A. Fujiwara and H. Nagaoka, “Quantum Fisher metric and estimation for pure state models”, *Phys. Lett. A* **201**, 119 (1995).
- [38] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information: 10th Anniversary Edition*, Cambridge University Press (2010).
- [39] N. Shiraishi, K. Funo, and K. Saito, “Speed Limit for Classical Stochastic Processes”, *Phys. Rev. Lett.* **121**, 070601 (2018).
- [40] A. Pal, S. Reuveni, and S. Rahav, “Thermodynamic uncertainty relation for systems with unidirectional transitions”, *Phys. Rev. Research* **3**, 013273 (2021).
- [41] A. Pal, S. Reuveni, and S. Rahav, “Thermodynamic uncertainty relation for first-passage times on Markov chains”, *Phys. Rev. Research* **3**, L032034 (2021).
- [42] Y. Hasegawa, “Thermodynamic uncertainty relation for quantum first passage process”, *Phys. Rev. E* **105**, 044127 (2022).
- [43] C. Flindt, T. Novotný, A. Braggio, and A.-P. Jauho, “Counting statistics of transport through Coulomb blockade nanostructures: High-order cumulants and non-Markovian effects”, *Phys. Rev. B* **82**, 155407 (2010).
- [44] S. Nakajima, M. Taguchi, T. Kubo, and Y. Tokura, “Interaction effect on adiabatic pump of charge and spin in quantum dot”, *Phys. Rev. B* **92**, 195420 (2015).
- [45] T. Pluecker, M. R. Wegewijs, and J. Splettstoesser, “Gauge freedom in observables and Landsberg’s nonadiabatic geometric phase: Pumping spectroscopy of interacting open quantum systems”, *Phys. Rev. B* **95**, 155431 (2017).
- [46] S. Nakajima and Y. Tokura, “Excess Entropy Production in Quantum System: Quantum Master Equation Approach”, *J. Stat. Phys.* **169**, 902 (2017).
- [47] S. Nakajima and Y. Utsumi, “Asymptotic expansion of the solution of the master equation and its application to the speed limit”, *Phys. Rev. E* **104**, 054139 (2021).
- [48] M. P. Drazin, “Pseudo-Inverses in Associative Rings and Semigroups”, *The American Mathematical Monthly* **65**, 506 (1958).