## Natural explanation of recent results on $e^+e^- \rightarrow \Lambda \bar{\Lambda}$

A.I. Milstein<sup>1,2,\*</sup> and S.G. Salnikov<sup>1,2,†</sup>

<sup>1</sup>Budker Institute of Nuclear Physics of SB RAS, 630090 Novosibirsk, Russia

<sup>2</sup>Novosibirsk State University, 630090 Novosibirsk, Russia

(Dated: May 15, 2023)

We show that the recent experimental data on the cross section of the process  $e^+e^- \to \Lambda\bar{\Lambda}$  near the threshold can be perfectly explained by the final-state interaction of  $\Lambda$  and  $\bar{\Lambda}$ . The enhancement of the cross section is related to the existence of low-energy real or virtual state in the corresponding potential. We present a simple analytical formula that fits the experimental data very well.

Recently, new experimental data have appeared on the cross section of  $e^+e^- \to \Lambda\bar{\Lambda}$  annihilation near the threshold [1]. These data are consistent with the results of previous works [2–4], but have much higher accuracy. All these results demonstrate a strong energy dependence of the cross section near the threshold. A similar phenomenon has been observed in such processes as  $e^+e^- \to p\bar{p}$  [5–12],  $e^+e^- \to n\bar{n}$  [13–15],  $e^+e^- \to \Lambda_c\bar{\Lambda}_c$  [16, 17],  $e^+e^- \to B\bar{B}$  [18], and others. In all these cases the shapes of near-threshold resonances differ significantly from the standard Breit-Wigner parameterization. The origin of the phenomenon is naturally explained by the strong interaction of produced particles near the threshold (the so-called final-state interaction). Since a typical value of the corresponding potential is rather large (hundreds of MeV), existence of either low-energy bound state or virtual state is possible. In the latter case a small deepening of the potential well leads to appearance of a real low-energy bound state. In both cases, the value of the wave function (or its derivative) inside the potential well significantly exceeds the value of the wave function inside the potential well is very strong. Since quarks in  $e^+e^-$  annihilation are produced at small distances of the order of  $1/\sqrt{s}$ , a strong energy dependence of the cross section is determined solely by the energy dependence of the wave function of produced pair of hadrons at small distances. Such a natural approach made it possible to describe well the energy dependence of almost all known near-threshold resonances (see Refs. [19–25] and references therein).

The annihilation  $e^+e^- \to \Lambda\bar{\Lambda}$  near the threshold is the most simple for investigation. This is due to the fact that the  $\Lambda\bar{\Lambda}$  system has a fixed isotopic spin I = 0, and the pair is produced mainly in the state with an angular momentum l = 0 (the contribution of state with l = 2 can be neglected). Moreover, there is no Coulomb interaction between  $\Lambda$  and  $\bar{\Lambda}$ . Our analysis shows that the imaginary part of the optical potential of  $\Lambda\bar{\Lambda}$  interaction, which takes into account the possibility of annihilation of  $\Lambda\bar{\Lambda}$  pair into mesons, has only little effect on the cross section. Therefore, we neglect the imaginary part of the potential. Finally we describe the cross section of the process  $e^+e^- \to \Lambda\bar{\Lambda}$  by a simple analytical formula (see Ref. [25] and references therein for more details):

$$\sigma = \frac{2\pi\beta\alpha^2}{s} g^2 F_D^2(s) |\psi(0)|^2,$$
(1)

where  $\beta = k/M_{\Lambda}$  is the baryon velocity,  $k = \sqrt{M_{\Lambda}E}$ ,  $s = (2M_{\Lambda} + E)^2$ , E is the kinetic energy of the pair,  $F_D(s) = (1 - s/\Lambda^2)^{-2}$  is the dipole form factor, and  $\Lambda$  is some parameter close to 1 GeV. The factor g is related to the probability of pair production at small distance  $\sim 1/\sqrt{s}$  and can be considered as a constant independent of energy. In Eq. (1),  $\psi(0)$  is the wave function of  $\Lambda\bar{\Lambda}$  pair at r = 0.

The cross section (1) is enhanced by the factor  $|\psi(0)|^2 \gg 1$  if there is a loosely bound state or a virtual state of  $\Lambda\bar{\Lambda}$  pair. In both cases the modulus of scattering length a of  $\Lambda$  and  $\bar{\Lambda}$  is large compared to the characteristic radius R of  $\Lambda\bar{\Lambda}$  interaction potential,  $|a| \gg R$ . For a loosely bound state a is positive and the binding energy is  $\varepsilon = -1/M_{\Lambda}a^2$ . For a virtual state a is negative and the energy of virtual state is defined as  $\varepsilon = 1/M_{\Lambda}a^2$ . In both cases  $|\varepsilon|$  is much smaller than the characteristic depth of the potential well. The energy dependence of  $|\psi(0)|^2$  for near-threshold resonances is more or less universal and is determined by the scattering length a and the effective radius of interaction [26]. Therefore, one can use any convenient form of potential U(r) for description of near-threshold resonances.

In the present paper we parametrize the potential as  $U(r) = -U_0 \cdot \theta(R-r)$ . For this potential, the energy dependence of  $|\psi(0)|^2$  is well-known (see, e.g., Ref. [26]):

$$|\psi(0)|^2 = \frac{q^2}{q^2 \cos^2(qR) + k^2 \sin^2(qR)}, \qquad q = \sqrt{M_{\Lambda}(E + U_0)}.$$
(2)

<sup>\*</sup> A.I.Milstein@inp.nsk.su

<sup>&</sup>lt;sup>†</sup> S.G.Salnikov@inp.nsk.su




Figure 1. The cross section of  $e^+e^- \to \Lambda\bar{\Lambda}$  annihilation (left) and the enhancement factor  $|\psi(0)|^2$  (right) as the functions of energy *E*. The parameters of the potential are  $U_0 = 584 \,\text{MeV}$  and  $R = 0.45 \,\text{fm}$ . Experimental data are taken from Ref. [1].

Near the threshold  $k \ll q$  and the cross section (1) is enhanced if

$$q_0 a \approx \pi \left( n + \frac{1}{2} \right) + \delta, \qquad |\delta| \ll 1,$$
(3)

where  $q_0 = \sqrt{M_{\Lambda}U_0}$ , and *n* is an integer. For  $|\delta| \ll 1$ , the scattering length is  $a = 1/q_0\delta$ , where  $\delta > 0$  for the bound state, and  $\delta < 0$  for the virtual state.

By means of Eq. (3) the expression (2) can be simplified:

$$\begin{aligned} |\psi(0)|^2 &\approx \frac{\gamma U_0}{\left(E + \varepsilon_0\right)^2 + \gamma E}, \\ \gamma &= 4\kappa^2 U_0, \qquad \varepsilon_0 = 2\delta \kappa U_0, \qquad \kappa = \frac{1}{\pi (n + 1/2)}. \end{aligned}$$
(4)

The corresponding energy dependence of the cross section (1) is equivalent to the Flatté formula [27], which is expressed in terms of the scattering length and the effective radius  $r_0$  of interaction. Note that for the rectangular potential well  $r_0 = R$ . One can easily verify that the precise and approximate formulas for the cross section are in good agreement with each other for  $|\delta| \ll 1$  and  $E \leq \varepsilon_0 \ll U_0$ . Note that  $|\varepsilon_0| \gg |\varepsilon|$  for both bound and virtual states, namely  $\varepsilon_0 \approx 2 |\varepsilon| a/R$ . However, the position of peak in the cross section, which is proportional to  $\sqrt{E} |\psi(0)|^2$ , is located at energy  $E \approx |\varepsilon|$  for both bound and virtual states.

In Fig. 1, we show our predictions for the cross section compared to experimental data [1], as well as the enhancement factor  $|\psi(0)|^2$ . The parameters of the model are  $U_0 = 584$  MeV, R = 0.45 fm, and g = 0.2. In the energy region under consideration the dependence of our predictions on the parameter  $\Lambda$  is very weak. To be specific, we set  $\Lambda = 1$  GeV. Our model, giving  $\chi^2/N_{\rm df} = 9.8/13$ , provides a good description of experimental data [1]. Note that account for the enhancement factor  $|\psi(0)|^2$  is of great importance for correct description of experimental data.

Within our model, we also predict a bound state with the binding energy  $E_0 \approx -30$  MeV. Observation of this bound state would be very important. The results of Refs. [28–36] indicate the anomalous behavior of the cross sections  $e^+e^- \rightarrow K^+K^-\pi^+\pi^-$ ,  $e^+e^- \rightarrow 2(K^+K^-)$ ,  $e^+e^- \rightarrow \phi K^+K^-$ , and others at  $\sqrt{s} \approx 2.2$  GeV (this value of s corresponds to  $E \approx -30$  MeV). However, a more detailed study of this energy region is required.

In conclusion, the assumption of existence of a low-energy real or virtual state has allowed us to describe perfectly recent and previous experimental data for the cross section of  $e^+e^- \rightarrow \Lambda\bar{\Lambda}$  annihilation near the threshold. Our model indicates possible existence of a bound  $\Lambda\bar{\Lambda}$  state with energy  $E \approx -30$  MeV.

We are grateful to A.E. Bondar for valuable discussions.

- [1] M. Ablikim, et al., Phys. Rev. D 107, 072005 (2023).
- [2] B. Aubert, et al., Phys. Rev. D 76, 092006 (2007).
- [3] M. Ablikim, et al., Phys. Rev. D 97, 032013 (2018).
- [4] M. Ablikim, et al., Phys. Rev. Lett. 123, 122003 (2019).
- [5] B. Aubert, et al., Phys. Rev. D **73**, 012005 (2006).
- [6] J.P. Lees, et al., Phys. Rev. D 87, 092005 (2013).
- [7] R.R. Akhmetshin, et al., Phys. Lett. B 759, 634 (2016).
- [8] M. Ablikim, et al., Phys. Rev. Lett. **124**, 042001 (2020).
- [9] M. Ablikim, et al., Phys. Lett. B 817, 136328 (2021).
- [10] R.R. Akhmetshin, et al., Phys. Lett. B **794**, 64 (2019).
- [11] M. Ablikim, et al., Phys. Rev. D **91**, 112004 (2015).
- [12] M. Ablikim, et al., Phys. Rev. D **99**, 092002 (2019).
- [13] M.N. Achasov, et al., Phys. Rev. D **90**, 112007 (2014).
- [14] M. Ablikim, et al., Nat. Phys. 17, 1200 (2021).
- [15] M.N. Achasov, et al., Eur. Phys. J. C 82, 761 (2022).
- [16] G. Pakhlova, et al., Phys. Rev. Lett. **101**, 172001 (2008).
- [17] M. Ablikim, et al., Phys. Rev. Lett. **120**, 132001 (2018).
- [18] B. Aubert, et al., Phys. Rev. Lett. **102**, 012001 (2009).
- [19] J. Haidenbauer, X.-W. Kang, and U.-G. Meißner, Nucl. Phys. A 929, 102 (2014).
- [20] J. Haidenbauer and U.-G. Meißner, Phys. Lett. B 761, 456 (2016).
- [21] J. Haidenbauer, U.-G. Meißner, and L.-Y. Dai, Phys. Rev. D 103, 014028 (2021).
- [22] A.I. Milstein and S.G. Salnikov, Phys. Rev. D 104, 014007 (2021).
- [23] A.I. Milstein and S.G. Salnikov, Phys. Rev. D 105, 074002 (2022).
- [24] A.I. Milstein and S.G. Salnikov, Phys. Rev. D 105, L031501 (2022).
- [25] A.I. Milstein and S.G. Salnikov, Phys. Rev. D 106, 074012 (2022).
- [26] L. D. Landau and E. M. Lifshitz, Quantum Mechanics Non-relativistic Theory, Pergamon Press (1991).
- [27] S.M. Flatté, Phys. Lett. B 63, 224 (1976).
- [28] B. Aubert, et al., Phys. Rev. D 76, 012008 (2007).
- [29] Y.-W. Chang, et al., Phys. Rev. D 79, 052006 (2009).
- [30] M. Ablikim, et al., Phys. Rev. D **100**, 032009 (2019).
- [31] M. Ablikim, et al., Phys. Rev. D 99, 032001 (2019).
- [32] M. Ablikim, et al., Phys. Rev. Lett. **124**, 112001 (2020).
- [33] M. Ablikim, et al., Phys. Rev. D 102, 012008 (2020).
- [34] M. Ablikim, et al., Phys. Rev. D 104, 032007 (2021).
- [35] M. Ablikim, et al., arXiv:2112.13219 [hep-ex].
- [36] L. Xia, Supl. la Rev. Mex. Física **3**, 1 (2022).