

A tight universal relation between the shape eccentricity and the moment of inertia for rotating neutron stars

YONG GAO,^{1,2} LIJING SHAO,^{2,3} AND JAN STEINHOFF⁴

¹*Department of Astronomy, School of Physics, Peking University, Beijing 100871, China*

²*Kawli Institute for Astronomy and Astrophysics, Peking University, Beijing 100871, China*

³*National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100012, China*

⁴*Max Planck Institute for Gravitational Physics (Albert Einstein Institute), Am Mühlenberg 1, Potsdam 14476, Germany*

ABSTRACT

Universal relations that are insensitive to the equation of state (EoS) are useful in reducing the parameter space when measuring global quantities of neutron stars (NSs). In this paper, we reveal a new universal relation that connects the eccentricity to the radius and moment of inertia of rotating NSs. We demonstrate that the universality of this relation holds for both conventional NSs and bare quark stars (QSs) in the slow rotation approximation, albeit with different relations. The maximum relative deviation is approximately 1% for conventional NSs and 0.1% for QSs. Additionally, we show that the universality still exists for fast-rotating NSs if we use the dimensionless spin to characterize their rotation. The new universal relation will be a valuable tool to reduce the number of parameters used to describe the shape and multipoles of rotating NSs, and it may also be used to infer the eccentricity or moment of inertia of NSs in future X-ray observations.

Keywords: dense matter — methods: numerical — stars: rotation

1. INTRODUCTION

Neutron stars (NSs) are the densest stars in the universe, offering a unique laboratory to study supranuclear matter and gravity in the strong-field regime. Currently, the equation of state (EoS) for the cores of NSs is still poorly understood. Many EoS models with varying compositions and states of dense matter have been developed, leading to significantly different predictions of global properties for NSs (Lattimer & Prakash 2001). Therefore, observed global properties of NSs, such as the mass and the radius, can be used to constrain EoS models.

Despite the fact that the global properties of NSs depend sensitively on the EoS models, there exist EoS-insensitive relations that connect various quantities of NSs. These relations are said to be universal because they are insensitive to EoS models to a high degree of accuracy. For instance, a universal relation connecting the frequency and damping time of the quadrupolar f mode to the mass and moment of inertia of NSs was discovered by Lau et al. (2010). Yagi & Yunes (2013a) found the famous I-Love-Q relation for slowly rotating NSs,

which links the mass, the moment of inertia, the tidal Love number, and the spin-induced quadrupole moment. Universal relations for NSs are of great significance in both astrophysics and fundamental physics. By providing EoS-insensitive connections between different quantities, these relations allow us to extract global properties of NSs with higher accuracy, and help us study the inverse problem of determining the EoS. Moreover, universal relations can break the degeneracy between gravity theories and the uncertainties in EoS, making NSs the ideal laboratories to test gravity (Shao & Yagi 2022). We refer readers to Yagi & Yunes (2013b), Doneva & Pappas (2018), and references therein for reviews.

The exploration of the universal relations for rotating NSs has garnered lots of attention since the discovery of the I-Love-Q relation. The calculations of the I-Q relation were quickly extended to fast rotation by Doneva et al. (2013), it was shown that the universality of the relation is lost and becomes increasingly EoS-dependent as the spin frequency increases. However, Pappas & Apostolatos (2014) and Chakrabarti et al. (2014) demonstrated that the I-Q relation remains universal if dimensionless quantities are used to characterize the spin amplitude instead of the spin frequency f . Later, Pappas & Apostolatos (2014) and Yagi et al. (2014) discovered that the first four multipole moments of rotating NSs are

universal to some extent. This relation allows for a more accurate description of the spacetime geometry around a NS with fewer parameters. Additionally, [Luk & Lin \(2018\)](#) found another universal relation connecting the radius and orbital frequency of the innermost stable circular orbit (ISCO) to the mass and spin frequency of rotating NSs.

Apart from multipoles and ISCO, eccentricity is another important parameter to describe rotating NSs. The oblateness induced by rotation has a large impact on the X-ray emissions from the surface of X-ray pulsars. To study this effect, [Morsink et al. \(2007\)](#) parameterized the oblate shape with the compactness of NSs and discovered that the geometric effect induced by the oblateness can rival the Doppler effect in certain configurations. To reduce the parameter space of X-ray modeling, [Bauböck et al. \(2013\)](#) derived a universal relation between the eccentricity and the compactness of slowly rotating NSs. [AlGendy & Morsink \(2014\)](#) also found an EoS-insensitive fit of the eccentricity, using a slightly different parametrization for the surface other than the one used in the Hartle-Thorne formalism.

In this paper, we discover a new universal relation between the surface eccentricity and the moment of inertia for rotating NSs. The paper is structured as follows. In [Sec. 2](#), we provide a definition of multipoles and eccentricity in the slow rotation approximation and present the universal relation for both conventional NSs and QSs. In [Sec. 3](#), we investigate the universal relation for fast rotating NSs. Discussion of possible applications and connections of the new universal relation to early work is shown in [Sec. 4](#). Throughout the paper, we use geometric units with $G = c = 1$.

2. A NEW UNIVERSAL RELATION IN THE SLOW-ROTATION APPROXIMATION

2.1. Multipole moments and shape parameters

To study the universal relation, we first give an overview of the structures and shape parameters of slowly rotating NSs. Following [Hartle \(1967\)](#) and [Hartle & Thorne \(1968\)](#), we construct these stars by solving the Einstein equations perturbatively in a slow-rotation expansion to quadratic order in the spin. At the zeroth order in spin, we obtain the mass M and the radius R of the non-rotating background. At the first order in spin, we extract the angular momentum J , from which we can define the moment of inertia I and the dimensionless spin χ as $I \equiv J/\Omega$ and $\chi \equiv J/M^2$, where Ω is the angular frequency of the rotating star. Universal relations usually connect dimensionless quantities. The dimensionless moment of inertia \bar{I} is usually defined as $\bar{I} \equiv I/M^3$. At the second order in spin, the star is

deformed into an oblate shape, and we get the spin-induced quadrupole moment $Q \equiv -J^2/M - 8KM^3/5$. The parameter K depends on the EoS of NSs and equals to zero for Kerr black holes according to the no hair theorem. The dimensionless quadrupole moment is defined as $\bar{Q} \equiv -Q/M^3\chi^2$. The I-Q relation connects the dimensionless quantities \bar{I} and \bar{Q} . The exterior spacetime of a slowly rotating NS can be fully described up to the quadratic order in spin by the mass M , the angular momentum J , and the quadrupole moment Q ([Hartle & Thorne 1968](#)).

Observationally, some observation of a rotating NS depends on the geometry of its surface. We use the eccentricity e_s to describe the oblate shape of a NS,

$$e_s \equiv \sqrt{\left(\frac{R_{\text{eq}}}{R_p}\right)^2 - 1}, \quad (1)$$

where R_{eq} and R_p are the equatorial and polar radii in a specific coordinate. In the Hartle-Thorne coordinate, the isodensity surface at radial coordinate r in the non-rotating star is displaced to

$$r \rightarrow r + \xi_0(r) + \xi_2(r)P_2(\cos\theta), \quad (2)$$

in the rotating configuration, where ξ_0 and ξ_2 are spherical and quadrupole displacements respectively, and $P_2(\cos\theta)$ is the Legendre polynomial. Combining [Eqs. \(1–2\)](#), we get the surface eccentricity in the Hartle-Thorne coordinate as

$$e_s^{\text{HT}} = [-3(\xi_2(R)/R)]^{1/2}. \quad (3)$$

Equation (2) describes the isodensity surface in a particular coordinate system. By embedding the isodensity surface into a three-dimensional flat space (denoted by r^*, θ^*, ϕ^*), [Hartle & Thorne \(1968\)](#) found an invariant parametrization of the oblate surface. To the second order of the spin, the desired surface is a spheroid with

$$r^*(\theta^*) = r + \xi_0(r) + \left\{ \xi_2(r) + r[v_2(r) - h_2(r)] \right\} P_2(\cos\theta^*). \quad (4)$$

Here v_2 and h_2 are metric functions at the second order in spin. The surface eccentricity observed in flat space is then given by

$$e_s^* = \left\{ -3[v_2(R) - h_2(R) + \xi_2(R)/R] \right\}^{1/2}, \quad (5)$$

where the superscript “*” denotes the eccentricity observed in the flat space.

2.2. A universal relation for the eccentricity of NSs




Figure 1. The mass-radius relation for selected EoS models of NSs (solid) and Qs (dashed). The $1\text{-}\sigma$ regions of the mass measurements of PSR J0348+0432 (Antoniadis et al. 2013) and PSR J0740+6620 (Fonseca et al. 2021) are illustrated.

The universal relation that we discovered connects the quantity $e_s/R\Omega$ and the dimensionless moment of inertia \bar{I} . Same as the I-Q relation (Yagi & Yunes 2013a) and the three hair relation for the multipole moments (Yagi et al. 2014), the normalization factors M and R are quantities of the non-rotating background in the slow-rotation approximation. For convenience, we define a dimensionless radius, $\hat{R} \equiv R\Omega$. We have verified that the universal relation exists for both the eccentricity in the Hartle-Thorne coordinate, e_s^{HT} , and the eccentricity observed in flat spacetime, e_s^* . In the following, we use e_s^* to illustrate the results.

We first study the universal relation for conventional NSs. Our selection of realistic EoSs includes BSK21 (Goriely et al. 2010), AU (Wiringa et al. 1988), HLPS (Hebeler et al. 2013), PAL1 (Prakash et al. 1988), APR (Akmal et al. 1998), SLy4 (Douchin & Haensel 2001), MS0 (Mueller & Serot 1996), and ENG (Engvik et al. 1994). As shown in Fig. 1, these models cover a wide range in the mass-radius diagram of static NSs, and all of them have maximal NS mass larger than $2M_\odot$. Although the very stiff EoSs MS0 and PAL1 have been ruled out by the tidal deformability from GW170817 (Abbott et al. 2017), we include them to demonstrate that the universality exists for a large family of EoSs.

Our universal relation is described with great accuracy by

$$\frac{e_s^*}{\hat{R}} = \sum_{k=0}^3 a_k (\ln \bar{I})^k, \quad (6)$$

where a_k 's are fitting coefficients with $a_0 = -0.855572$, $a_1 = 2.185502$, $a_2 = -0.428061$, $a_3 = 0.051177$. Note




Figure 2. The $e_s^*/\hat{R}-\bar{I}$ universal relation for slowly-rotating NSs. The upper panel shows the fitted universal relations for both NSs and Qs. The middle (lower) panel presents the relative deviation for NSs (Qs).

that we use the dimensionless moment of inertia $\bar{I} \leq 100$, which corresponds to $M \gtrsim 0.5M_\odot$ for selected models. We define the relative deviation to Eq. (6) as

$$\Delta = \frac{e_s^*/\hat{R} - (e_s^*/\hat{R})_{\text{fit}}}{(e_s^*/\hat{R})_{\text{fit}}}. \quad (7)$$

As shown in Fig. 2, the relative deviation to the universal relation is smaller than 1% for selected models of EoSs. NSs with $M \gtrsim 1M_\odot$ are more relevant for astrophysics, and in this case, the dimensionless moment of inertia $\bar{I} \lesssim 30$ for selected EoS models, and the universal relation takes a simpler form,

$$\frac{e_s^*}{\hat{R}} = 0.11418 + 1.04115 \ln \bar{I}. \quad (8)$$

The relative deviation to this relation is less than $\sim 1\%$.

For Qs, we use the phenomenological MIT bag model to describe the quark matter. This model assumes a nearly equal number of u , d , s quarks and a small fraction of electrons confined within a bag of vacuum energy density B (Farhi & Jaffe 1984; Witten 1984). We account for the mass of the s quark, m_s , and include the quark-gluon interaction to the lowest order in $\alpha_c = g^2/4\pi$. To investigate the universal relation for Qs, we employ six different EoSs with varying combinations of m_s , α_c , and B in Table 1. The resulting mass-radius relation is displayed in Fig. 1.

Table 1. Parameters for QSs in the MIT bag model.

Model	B (MeV fm $^{-3}$)	m_s (MeV)	α_c
SQM1	80	100	0
SQM2	80	50	0.1
SQM3	70	150	0
SQM4	70	50	0.3
SQM5	60	0	0
SQM6	60	100	0.4

The relation between e_s^*/\hat{R} and \bar{I} is different from that of NSs, but it is still universal and can be well fitted by

$$\frac{e_s^*}{\hat{R}} = \sum_{k=0}^4 b_k (\ln \bar{I})^k, \quad (9)$$

with coefficients $b_0 = -1.499749$, $b_1 = 2.911859$, $b_2 = -0.749237$, $b_3 = 0.137057$, and $b_4 = -0.007801$. Interestingly, the deviation from the universal relation for QSs is much smaller than that for NSs, with relative deviation less than $\sim 0.1\%$, as shown in the lower panel of Fig. 2. For QSs in our study, the condition for $M \gtrsim 1 M_\odot$ corresponds to $\bar{I} \lesssim 20$. Within this range, the universal relation can be approximated by a simpler fitting formula,

$$\frac{e_s^*}{\hat{R}} = -0.043372 + 1.210546 \ln(\bar{I} - 0.470579), \quad (10)$$

The relative deviation from this fitting formula is less than $\sim 0.3\%$.

3. UNIVERSAL RELATION FOR FAST ROTATING NSS

Fast rotation is relevant for sub-millisecond pulsars, nascent NSs after supernovae, and NSs formed in binary NS mergers. Rapid rotation causes NSs to develop a more obvious oblate shape. In this section, we explore the universal relation for rapidly rotating NSs using the RNS code developed by Stergioulas & Friedman (1995).

The RNS code uses a quasi-isotropic coordinate system to represent the line element of the stationary axisymmetric spacetime,

$$ds^2 = -e^{2\nu} dt^2 + B^2 r^2 \sin^2 \theta e^{-2\nu} (d\phi - \omega dt)^2 + e^{2(\xi-\nu)} (dr^2 + r^2 d\theta^2), \quad (11)$$

where ν , B , ω , and ξ are metric functions that depend on r and θ . Assuming a perfect fluid and uniform rotation, we obtain the stellar structure and spacetime metric. The conserved angular momentum J can be computed from a volume integration over the matter field. The moment of inertia I and the dimensionless spin have




Figure 3. The $\tilde{e}_s/\hat{R}-\bar{I}$ universal relation for fast rotating NSs. The first panel shows the fitted relation for $\chi = 0.2$, 0.4 , and 0.6 . The second to fourth panels display the relative deviations to the universal relation for $\chi = 0.2$, 0.4 , and 0.6 respectively.

the same definition as before. The quadrupole moment Q can be obtained from the asymptotic expansion of the metric functions. The surface eccentricity is formally given by Eq. (1), with the eccentricity, equatorial and polar radii defined in the quasi-isotropic coordinate. We use the notation \tilde{e}_s to denote the eccentricity in the quasi-isotropic coordinate. Note that, unlike in the slow-rotation approximation, the normalization factor M is the mass for the rotating configuration, and $\hat{R} \equiv R_{\text{eq}}\Omega$.

To study universal relations for rapidly rotating NSs, it is necessary to use a suitable parameter to characterize their spin amplitude. As demonstrated by Doneva et al. (2013), if one uses the spin frequency f as the parameter, the I-Q relation for fast rotating NSs is lost. Similarly, the universal relation that we discovered also breaks down for fixed spin frequencies. However, Pappas & Apostolatos (2014) and Chakrabarti et al. (2014) found that the I-Q relation is still universal for fast rotating NSs if one chooses dimensionless spin parameters such as χ , Mf , and Rf , instead of the dimensionful f . Inspired by their work, we use χ to characterize the spin amplitude and find that the $\tilde{e}_s/\hat{R}-\bar{I}$ relation for both conventional NSs and strange QSs is still universal.

According to Lo & Lin (2011), the maximum value of the dimensionless spin parameter χ for NSs rotating at the Keplerian frequency is about 0.7 for various EoS




Figure 4. The $\tilde{e}_s/\hat{R}-\bar{I}$ universal relation for fast rotating QSS. The first panel shows the fitted relation for $\chi = 0.5$ and 0.8 . The second and third panels display the relative deviations to the universal relation for $\chi = 0.5$ and 0.8 respectively.

models. This limit is nearly independent of the mass of the NS if the mass is larger than $1 M_\odot$. However, for QSS in the MIT bag model, the spin parameter can be larger than unity and does not have a universal upper limit. Its value also depends strongly on the bag constant and the mass of the star. Therefore, in Fig. 3, we display three representative cases with $\chi = 0.2, 0.4,$ and 0.6 for conventional NSs. The relative deviation from the universal relation is less than $\sim 1\%$. The cases for QSS with $\chi = 0.5$ and $\chi = 0.8$ are shown in Fig. 4, and the relative deviation is on the order of 0.1% , which is again much tighter than conventional NSs.

4. DISCUSSION

The relation between e_s/\hat{R} and \bar{I} adds a new tight universal relation to the known ones. Here, e_s is the surface eccentricity formally defined in Eq. (1) without assigning a specific coordinate system. In our paper, it includes e_s^{HT} , e_s^* , and \tilde{e}_s , which are eccentricities defined in different coordinate systems. The effect of gauge choice on the eccentricity is very small. All of these eccentricities satisfy the universal relation very well. Since the I-Q relation (Yagi & Yunes 2013a; Chakrabarti et al. 2014) connects \bar{I} to \bar{Q} , and the three-hair relation (Pappas & Apostolatos 2014; Yagi et al. 2014) connects \bar{Q} to two other higher-order multipoles, the relation be-

tween e_s/\hat{R} and these normalized multipole moments is also universal. Combined with the I-Love relation (Yagi & Yunes 2013a), we have a universal relation between e_s/\hat{R} and the dimensionless tidal Love number. Moreover, the universal relation between the f -mode oscillation and \bar{I} (Lau et al. 2010) helps us connect e_s/\hat{R} to the frequency and damping time of the quadrupolar f mode.

Our work is not the only universal relation for the shape parameter of rotating NSs. To incorporate the effects of the eccentricity of NSs when modelling the X-ray emission, Bauböck et al. (2013) derived an analytic expression connecting the eccentricity of the stellar surface to the compactness C , the spin angular momentum, and the quadrupole moment of the spacetime by using the analytical results in Hartle & Thorne (1968). They also used several other universal relations to express e_s in a single parameter C (see Fig. 3 of Bauböck et al. 2013). Our $e_s/\hat{R}-\bar{I}$ relation is tighter than the e_s-C relation.

Universal relations are a powerful tool to reduce modelling uncertainties and inferring NS parameters. As we discussed before, the eccentricity of rotating NSs is an observable and is an important input for X-ray modelling. The oblateness induced by rotation at frequencies above 300 Hz produces a geometric effect that has imprints in the pulse profile of X-ray pulsars (Morsink et al. 2007). For some emission configurations, the oblateness effect can rival the Doppler effect. Besides, the effects of oblateness need to be taken into account when measuring the radii of NSs from rotationally broadened atomic lines (Bauböck et al. 2013). On one hand, our new universal relation reduces the number of parameters used to describe the shape and multipoles of rotating NSs. On the other hand, the new universal relation can be used to infer the eccentricity of NSs if the mass, radius, spin frequency, and moment of inertia are inferred in future X-ray observations.

We thank Enping Zhou for discussion. This work was supported by the National SKA Program of China (2020SKA0120300), the National Natural Science Foundation of China (11975027, 11991053), the Max Planck Partner Group Program funded by the Max Planck Society, and the High-Performance Computing Platform of Peking University.

Software: RNS (Stergioulas & Friedman 1995)

REFERENCES

Abbott, B. P., et al. 2017, Phys. Rev. Lett., 119, 161101

Akmal, A., Pandharipande, V. R., & Ravenhall, D. G. 1998, Phys. Rev. C, 58, 1804

- AlGendy, M., & Morsink, S. M. 2014, *ApJ*, 791, 78
- Antoniadis, J., Freire, P. C. C., Wex, N., et al. 2013, *Science*, 340, 448
- Bauböck, M., Berti, E., Psaltis, D., & Özel, F. 2013, *ApJ*, 777, 68
- Bauböck, M., Psaltis, D., & Özel, F. 2013, *ApJ*, 766, 87
- Chakrabarti, S., Delsate, T., Gürlebeck, N., & Steinhoff, J. 2014, *Phys. Rev. Lett.*, 112, 201102
- Doneva, D. D., & Pappas, G. 2018, *Astrophys. Space Sci. Libr.*, 457, 737
- Doneva, D. D., Yazadjiev, S. S., Stergioulas, N., & Kokkotas, K. D. 2013, *ApJL*, 781, L6
- Douchin, F., & Haensel, P. 2001, *A&A*, 380, 151
- Engvik, L., Hjorth-Jensen, M., Osnes, E., Bao, G., & Østgaard, E. 1994, *Phys. Rev. Lett.*, 73, 2650
- Farhi, E., & Jaffe, R. L. 1984, *Phys. Rev. D*, 30, 2379
- Fonseca, E., et al. 2021, *ApJL*, 915, L12
- Goriely, S., Chamel, N., & Pearson, J. M. 2010, *Phys. Rev. C*, 82, 035804
- Hartle, J. B. 1967, *ApJ*, 150, 1005
- Hartle, J. B., & Thorne, K. S. 1968, *ApJ*, 153, 807
- Hebeler, K., Lattimer, J. M., Pethick, C. J., & Schwenk, A. 2013, *ApJ*, 773, 11
- Lattimer, J. M., & Prakash, M. 2001, *ApJ*, 550, 426
- Lau, H. K., Leung, P. T., & Lin, L. M. 2010, *ApJ*, 714, 1234
- Lo, K.-W., & Lin, L.-M. 2011, *ApJ*, 728, 12
- Luk, S.-S., & Lin, L.-M. 2018, *ApJ*, 861, 141
- Morsink, S. M., Leahy, D. A., Cadeau, C., & Braga, J. 2007, *ApJ*, 663, 1244
- Mueller, H., & Serot, B. D. 1996, *Nucl. Phys. A*, 606, 508
- Pappas, G., & Apostolatos, T. A. 2014, *Phys. Rev. Lett.*, 112, 121101
- Prakash, M., Ainsworth, T. L., & Lattimer, J. M. 1988, *Phys. Rev. Lett.*, 61, 2518
- Shao, L., & Yagi, K. 2022, *Sci. Bull.*, 67, 1946
- Stergioulas, N., & Friedman, J. L. 1995, *ApJ*, 444, 306
- Wiringa, R. B., Fiks, V., & Fabrocini, A. 1988, *Phys. Rev. C*, 38, 1010
- Witten, E. 1984, *Phys. Rev. D*, 30, 272
- Yagi, K., Kyutoku, K., Pappas, G., Yunes, N., & Apostolatos, T. A. 2014, *Phys. Rev. D*, 89, 124013
- Yagi, K., & Yunes, N. 2013a, *Phys. Rev. D*, 88, 023009
- . 2013b, *Science*, 341, 365