

ON PERTURBATIVE QUANTUM GRAVITY WITH A COSMOLOGICAL CONSTANT

David Prinz*

March 24, 2023

Abstract

We discuss how the incorporation of a cosmological constant affects the perturbative quantization of (effective) Quantum General Relativity. To this end, we derive the gravitational Slavnov–Taylor identities and appropriate renormalization conditions for the cosmological constant. Additionally, we calculate the corresponding Feynman rules for any vertex valence and with general gauge parameter. Furthermore, we provide the BRST setup and generate the Faddeev–Popov ghost and the symmetric ghost via a gauge fixing fermion and a gauge fixing boson, respectively. Finally, we study the transversality of the graviton propagator and the graviton three-valent vertex.

1 Introduction

Perturbative Quantum Gravity is generally believed not to admit a predictive perturbative expansion. This is due to the fact that it is non-renormalizable by power-counting and thus requires higher-derivative counterterms. These have been examined on the level of one and two loops [1, 2] in the framework of the background field method [3]. Surprisingly, the counterterms on one-loop level can be combined, using the Chern–Gauss–Bonnet theorem, to a total derivative [1]. This means that the divergences can be absorbed via a field redefinition. On the other hand, on two-loop level such a relation is missing, leading to the well-accepted conjecture that perturbative Quantum Gravity is not renormalizable at two loops [2]. Notably, it is still possible to find a non-local field redefinition to absorb the divergent contributions [4]. We reapproach this long-standing renormalization problem from a different viewpoint, namely via the framework of Hopf algebraic renormalization [5, 6, 7, 8, 9]: Soon after the development of this formalism, it has been applied to gauge theories [10, 11, 12, 13, 14]. In particular, D. Kreimer applied the same reasoning to perturbative Quantum Gravity and conjectured that its non-renormalizable nature could effectively become harmless, if it satisfies a tower of infinite gravitational Slavnov–Taylor identities [15, 16]. This has been worked out in detail by the author in a series of recent articles [17, 18, 19, 20, 21, 22] and his dissertation [23]. In particular, therein the Connes–Kreimer renormalization theory has been generalized to non-renormalizable (generalized) gauge theories, so that it applies to (effective) Quantum General Relativity. Specifically, we have derived precise identities which are necessary for a well-defined perturbative expansion [18]. These identities need to be checked on the level of the gravitational Feynman rules given in [19], which is a topic of ongoing research. In the present article, we want to address the case of a non-vanishing cosmological constant Λ . In particular, we calculate the graviton vertex Feynman rules for arbitrary vertex valence in Theorem 2.2 and the graviton propagator for the linearized de Donder gauge fixing with general gauge parameter ζ in Theorem 2.3. In addition, we list the three-valent and four-valent graviton vertex Feynman rules explicitly in Example 2.4.

*Riemann fellow at Riemann Center for Geometry and Physics and Institute for Theoretical Physics at Leibniz University Hannover; prinz@itp.uni-hannover.de

To this end, we use the following Lagrange density for (effective) Quantum General Relativity with a cosmological constant (QGR- Λ):

$$\mathcal{L}_{\text{QGR-}\Lambda} := \mathcal{L}_{\text{GR-}\Lambda} + \mathcal{L}_{\text{GR-}\Lambda\text{-GF}} + \mathcal{L}_{\text{GR-}\Lambda\text{-Ghost}} \quad (1a)$$

with the Einstein–Hilbert Lagrange density with a cosmological constant

$$\mathcal{L}_{\text{GR-}\Lambda} := -\frac{1}{2\kappa^2} (R + 2\Lambda) dV_g, \quad (1b)$$

the linearized de Donder gauge fixing Lagrange density

$$\mathcal{L}_{\text{GR-}\Lambda\text{-GF}} := -\frac{1}{4\kappa^2\zeta} b^{\mu\nu} d\mathcal{D}_\mu^{(1)} d\mathcal{D}_\nu^{(1)} \quad (1c)$$

and either the Faddeev–Popov ghost construction, cf. Corollary 3.2,

$$\mathcal{L}_{\text{GR-}\Lambda\text{-FP-Ghost}} := -\frac{1}{2} b^{\rho\sigma} \left(\frac{1}{\zeta} \bar{C}^{\mu} (\partial_\rho \partial_\sigma C_\mu) + \bar{C}^{\mu} \left(\partial_\mu (\Gamma^\nu{}_{\rho\sigma} C_\nu) - 2\partial_\rho (\Gamma^\nu{}_{\mu\sigma} C_\nu) \right) \right) dV_b \quad (1d)$$

or the symmetric ghost construction, cf. Corollary 3.3,

$$\begin{aligned} \mathcal{L}_{\text{GR-}\Lambda\text{-Sym-Ghost}} := & -\frac{1}{2\zeta} b^{\mu\nu} (\partial_\mu \bar{C}^\rho) (\partial_\nu C_\rho) dV_b \\ & -\frac{1}{2} b^{\mu\nu} \bar{C}^\rho \left(\frac{1}{2} \partial_\rho (\Gamma^\sigma{}_{\mu\nu} C_\sigma) - \partial_\mu (\Gamma^\sigma{}_{\rho\nu} C_\sigma) \right) dV_b \\ & +\frac{1}{2} b^{\mu\nu} \left(\frac{1}{2} \partial_\rho (\Gamma^\sigma{}_{\mu\nu} \bar{C}_\sigma) - \partial_\mu (\Gamma^\sigma{}_{\rho\nu} \bar{C}_\sigma) \right) C^\rho dV_b \\ & -\frac{\kappa^2 \zeta}{8} b_{\mu\nu} \left(\bar{C}^\rho (\partial_\rho \bar{C}^\mu) \right) \left(C^\sigma (\partial_\sigma C^\nu) \right) dV_b. \end{aligned} \quad (1e)$$

We remark that we consider its linearization with respect to the metric decomposition¹ $g_{\mu\nu} \equiv b_{\mu\nu} + \varkappa h_{\mu\nu}$, where $h_{\mu\nu}$ is the graviton field, $\varkappa := \sqrt{\kappa}$ the graviton coupling constant (with $\kappa := 8\pi G$ the Einstein gravitational constant) and $b_{\mu\nu}$ a convenient background metric.² In addition, $R := g^{\nu\sigma} R^\mu{}_{\nu\mu\sigma}$ is the Ricci scalar (with $R^\rho{}_{\sigma\mu\nu} := \partial_\mu \Gamma^\rho{}_{\nu\sigma} - \partial_\nu \Gamma^\rho{}_{\mu\sigma} + \Gamma^\rho{}_{\mu\lambda} \Gamma^\lambda{}_{\nu\sigma} - \Gamma^\rho{}_{\nu\lambda} \Gamma^\lambda{}_{\mu\sigma}$ the Riemann tensor) and Λ the cosmological constant. Furthermore, $dV_g := \sqrt{-\text{Det}(g)} dt \wedge dx \wedge dy \wedge dz$ denotes the Riemannian volume form and $dV_b := \sqrt{-\text{Det}(b)} dt \wedge dx \wedge dy \wedge dz$ the background metric volume form. In addition, $d\mathcal{D}_\mu^{(1)} := b^{\rho\sigma} \Gamma_{\mu\rho\sigma} \equiv 0$ denotes the linearized de Donder gauge fixing functional and ζ the gauge fixing parameter. Finally, C_μ and \bar{C}^μ are the graviton-ghost and graviton-antighost, respectively. In addition, we establish the relation between the Faddeev–Popov ghost construction and the symmetric ghost construction in Corollary 3.4, cf. [20, 21].

To study multiplicative renormalization, we decompose $\mathcal{L}_{\text{QGR-}\Lambda}$ into its individual monomials. As has been discussed in [18], the individual monomials can be addressed via the powers in the graviton coupling constant \varkappa , the gauge fixing parameter ζ , the ghost field C and now the cosmological constant Λ as follows³

$$\mathcal{L}_{\text{QGR-}\Lambda} \equiv \sum_{i=0}^{\infty} \sum_{j=-1}^0 \sum_{k=0}^1 \sum_{l=0}^1 \mathcal{L}_{\text{QGR-}\Lambda}^{i,j,k,l}, \quad (2)$$

¹This implies that the indices of the graviton field $h_{\mu\nu}$ and its inverse $h^{\mu\nu}$ are raised and lowered via the background metric $b_{\mu\nu}$ and its inverse $b^{\mu\nu}$, respectively.

²Typically given via a solution of the vacuum Einstein field equations, i.e. either the de Sitter or the anti-de Sitter metric.

³We omit the term $\mathcal{L}_{\text{QGR-}\Lambda}^{-1,0,0,0}$ as it is given by a total derivative.

where we have set $\mathcal{L}_{\text{QGR-}\Lambda}^{i,j,k,l} := (\mathcal{L}_{\text{QGR-}\Lambda})|_{O(\varkappa^i \zeta^j C^k \Lambda^l)}$, cf. [18, Section 1] and [19, Section 3]. Again, to absorb the upcoming divergences in a multiplicative manner, we multiply each monomial with an individual function $Z^r(\varepsilon)$ in the regulator $\varepsilon \in \mathbb{R}$:

$$\mathcal{L}_{\text{QGR-}\Lambda}^{\text{R}}(\varepsilon) := \sum_{i=0}^{\infty} \sum_{j=-1}^0 \sum_{k=0}^1 \sum_{l=0}^1 Z^{i,j,k,l}(\varepsilon) \mathcal{L}_{\text{QGR-}\Lambda}^{i,j,k,l}, \quad (3)$$

where the regulator ε is related to the energy scale through the choice of a regularization scheme. Then, the invariance of $\mathcal{L}_{\text{QGR-}\Lambda}^{\text{R}}(\varepsilon)$ under (residual) diffeomorphisms away from the reference point (where all Z -factors fulfill $Z^r(\varepsilon_0) = 1$) depends on the following identities:⁴

$$\frac{Z^{i,0,0,0}(\varepsilon) Z^{1,0,0,0}(\varepsilon)}{Z^{0,0,0,0}(\varepsilon)} \equiv Z^{(i+1),0,0,0}(\varepsilon), \quad (4a)$$

$$\frac{Z^{i,0,0,1}(\varepsilon) Z^{1,0,0,1}(\varepsilon)}{Z^{0,0,0,1}(\varepsilon)} \equiv Z^{(i+1),0,0,1}(\varepsilon) \quad (4b)$$

and

$$\frac{Z^{i,0,0,0}(\varepsilon)}{Z^{0,-1,0,0}(\varepsilon)} \equiv \frac{Z^{i,0,1,0}(\varepsilon)}{Z^{0,-1,1,0}(\varepsilon)}, \quad (4c)$$

for all $i \in \mathbb{N}_+$ and ε in the domain of the regularization scheme. Here, we have denoted the transversal and longitudinal graviton propagator Z -factors via $Z^{0,0,0,0}(\varepsilon)$ and $Z^{0,-1,0,0}(\varepsilon)$, respectively, as they cannot be distinguished by their degree in the cosmological constant since it behaves like an Euclidian mass term. Rather, we suggest the following renormalization conditions, where $\mathfrak{P}_{\mu\nu\rho\sigma}(p^\sigma, \Lambda; \zeta; \epsilon)$ denotes the dressed graviton propagator in the convenient decomposition of Theorem 4.1:

$$\mathfrak{P}_{\mu\nu\rho\sigma}(p^\sigma, \Lambda; \zeta; \epsilon) = -\frac{2ip^2}{p^2 + \Lambda + i\epsilon + \Sigma_{\mathfrak{P}}(p^2, \Lambda)} \left(\mathbb{G}_{\mu\nu\rho\sigma} - \left(\frac{1 - \zeta}{p^2 + \zeta\Lambda + \Sigma_{\mathfrak{P}}(p^2, \Lambda)} \right) \mathbb{L}_{\mu\nu\rho\sigma} \right) \quad (5)$$

Here, $\Sigma_{\mathfrak{P}}(p^2, \Lambda)$ denotes the renormalized self-energy of the full graviton propagator and $\Sigma_{\mathfrak{P}}(p^2, \Lambda)$ denotes the renormalized self-energy of the longitudinal graviton propagator. Both are subjected to the following two renormalization conditions:

$$\Sigma_{\mathfrak{P}}(-\Lambda, \Lambda) = 0 \quad \Sigma_{\mathfrak{P}}(-\Lambda, \Lambda) = 0 \quad (6a)$$

and

$$\left. \frac{\partial}{\partial p^2} \right|_{p^2=-\Lambda} \Sigma_{\mathfrak{P}}(p^2, \Lambda) = 0 \quad \left. \frac{\partial}{\partial p^2} \right|_{p^2=-\Lambda} \Sigma_{\mathfrak{P}}(p^2, \Lambda) = 0 \quad (6b)$$

This determines all renormalization constants uniquely. We will discuss these relations in future work. In particular, we remark that with the introduction of the Z -factor $Z_{\Lambda}(\varepsilon)$ we obtain an additional energy-curvature relation in addition to the Einstein field equations. However, this is only relevant for intermediate calculations and does not contribute to physical scattering amplitudes.

Finally, we discuss the transversality of (Effective) Quantum General Relativity with a cosmological constant in the sense of [22]: We find that the graviton propagator does not split into the sum of its transversal and longitudinal projection tensors in Theorem 4.1 and that the three-valent vertex Feynman rule does not satisfy the contraction identity in Theorem 4.3. Both these identities are satisfied in the case $\Lambda = 0$. The implication of these results on transversality will be discussed in future work.

⁴These identities are known in the literature as Ward–Takahashi identity in the realm of Quantum Electrodynamics and Slavnov–Taylor identities in the realm of Quantum Chromodynamics [24, 25, 26, 27, 28].

2 Feynman rules

We start this article by calculating the corresponding Feynman rules as a generalization of [19]. In particular, we focus only on the graviton vertex Feynman rules and the graviton propagator: This is due to the fact that the other Feynman rules remain valid with the replacement $\hat{\eta} \rightsquigarrow \hat{b}$. We refer to [17, 19, 22] for further information and references.

Convention 2.1. If the expansion of the gravity-matter Lagrange density is considered with respect to the general background metric $b_{\mu\nu}$, then we obtain the following additional factors: The vertex Feynman rules are multiplied by $\sqrt{-\text{Det}(b)}$ and the propagator Feynman rules by $1/\sqrt{-\text{Det}(b)}$. We omit both factors for a cleaner presentation, as diagrammatic Feynman rules do in fact only depend on them via their loop number. More precisely, a Feynman graph with loop number L obtains an additional factor of $\sqrt{-\text{Det}(b)}^{L-1}$.

Theorem 2.2. *Given the situation of [19, Theorem 4.10] with the metric decomposition $g_{\mu\nu} = b_{\mu\nu} + \varkappa h_{\mu\nu}$, where $b_{\mu\nu}$ is a suitable background metric. Then the graviton 2-point vertex Feynman rule for (effective) Quantum General Relativity with a cosmological constant reads (where ζ denotes the gauge parameter, Λ the cosmological constant and we use momentum conservation on the quadratic term, i.e. set $p_1^\sigma := p^\sigma$ and $p_2^\sigma := -p^\sigma$):*

$$\begin{aligned} \mathfrak{L}_2^{\mu_1\nu_1|\mu_2\nu_2}(p^\sigma, \Lambda; \zeta) &= \frac{i}{4} \left(1 - \frac{1}{\zeta}\right) \left(p^{\mu_1} p^{\nu_1} \hat{b}^{\mu_2\nu_2} + p^{\mu_2} p^{\nu_2} \hat{b}^{\mu_1\nu_1}\right) \\ &\quad - \frac{i}{8} \left(1 - \frac{1}{\zeta}\right) \left(p^{\mu_1} p^{\mu_2} \hat{b}^{\nu_1\nu_2} + p^{\mu_1} p^{\nu_2} \hat{b}^{\nu_1\mu_2} + p^{\nu_1} p^{\mu_2} \hat{b}^{\mu_1\nu_2} + p^{\nu_1} p^{\nu_2} \hat{b}^{\mu_1\mu_2}\right) \\ &\quad - \frac{i}{4} \left(1 - \frac{1}{2\zeta}\right) \left(p^2 \hat{b}^{\mu_1\nu_1} \hat{b}^{\mu_2\nu_2}\right) \\ &\quad + \frac{i}{8} \left(p^2 \hat{b}^{\mu_1\mu_2} \hat{b}^{\nu_1\nu_2} + p^2 \hat{b}^{\mu_1\nu_2} \hat{b}^{\nu_1\mu_2}\right) \\ &\quad + \frac{i\Lambda}{4} \left(\hat{b}^{\mu_1\mu_2} \hat{b}^{\nu_1\nu_2} + \hat{b}^{\mu_1\nu_2} \hat{b}^{\nu_1\mu_2} - \hat{b}^{\mu_1\nu_1} \hat{b}^{\mu_2\nu_2}\right) \end{aligned} \quad (7)$$

Furthermore, the graviton n -point vertex Feynman rules with $n > 2$ for (effective) Quantum General Relativity with a cosmological constant read (where \mathfrak{G}_n denotes the corresponding Feynman rules without a cosmological constant, cf. [19, Theorem 4.10], and \mathfrak{V}_n denotes the corresponding Feynman rules of the Riemannian volume form, cf. [19, Lemma 4.8]):

$$\mathfrak{L}_n^{\mu_1\nu_1|\dots|\mu_n\nu_n}(p_1^\sigma, \dots, p_n^\sigma, \Lambda) = \left(\mathfrak{G}_n^{\mu_1\nu_1|\dots|\mu_n\nu_n}(p_1^\sigma, \dots, p_n^\sigma) + \frac{i\Lambda}{\varkappa^2} \mathfrak{V}_n^{\mu_1\nu_1|\dots|\mu_n\nu_n} \right) \Bigg|_{\hat{\eta} \rightsquigarrow \hat{b}} \quad (8)$$

Proof. This follows directly from the linearity of the Feynman rules together with [19, Theorem 4.10] and [19, Lemma 4.8]. We remark an additional relative minus sign in front of \mathfrak{V}_n coming from the Fourier transform, as it does not depend on momenta. \blacksquare

Theorem 2.3. *Given the situation of Theorem 2.2, the graviton propagator Feynman rule for (effective) Quantum General Relativity with a cosmological constant reads:*

$$\begin{aligned} \mathfrak{P}_{\mu_1\nu_1|\mu_2\nu_2}(p^\sigma, \Lambda; \zeta; \epsilon) &= -\frac{2i}{p^2 + \Lambda + i\epsilon} \left[\left(\hat{b}_{\mu_1\mu_2} \hat{b}_{\nu_1\nu_2} + \hat{b}_{\mu_1\nu_2} \hat{b}_{\nu_1\mu_2} - \hat{b}_{\mu_1\nu_1} \hat{b}_{\mu_2\nu_2} \right) \right. \\ &\quad \left. - \left(\frac{1 - \zeta}{p^2 + \zeta\Lambda} \right) \left(\hat{b}_{\mu_1\mu_2} p_{\nu_1} p_{\nu_2} + \hat{b}_{\mu_1\nu_2} p_{\nu_1} p_{\mu_2} + \hat{b}_{\nu_1\mu_2} p_{\mu_1} p_{\nu_2} + \hat{b}_{\nu_1\nu_2} p_{\mu_1} p_{\mu_2} \right) \right] \end{aligned} \quad (9)$$

Proof. To calculate the graviton propagator, we recall the quadratic Feynman rule from Equation (7) and then invert it to obtain the propagator, i.e. such that⁵

$$\mathfrak{L}_2^{\mu_1\nu_1|\mu_2\nu_2}(p^\sigma, \Lambda; \zeta) \mathfrak{P}_{\mu_2\nu_2|\mu_3\nu_3}(p^\sigma, \Lambda; \zeta; 0) = \frac{1}{2} \left(\hat{\delta}_{\mu_3}^{\mu_1} \hat{\delta}_{\nu_3}^{\nu_1} + \hat{\delta}_{\nu_3}^{\mu_1} \hat{\delta}_{\mu_3}^{\nu_1} \right) \quad (10)$$

holds. ■

Example 2.4. Given the situation of Theorem 2.2, the three- and four-valent graviton vertex Feynman rules read as follows:⁶

$$\mathfrak{L}_3^{\mu_1\nu_1|\mu_2\nu_2|\mu_3\nu_3}(p_1^\sigma, p_2^\sigma, p_3^\sigma, \Lambda) = \frac{i}{8} \sum_{\mu_i \leftrightarrow \nu_i} \sum_{s \in S_3} \mathfrak{I}_3^{\mu_{s(1)}\nu_{s(1)}|\mu_{s(2)}\nu_{s(2)}|\mu_{s(3)}\nu_{s(3)}} \left(p_{s(1)}^\sigma, p_{s(2)}^\sigma \right) \quad (11a)$$

with

$$\begin{aligned} \mathfrak{I}_3^{\mu_1\nu_1|\mu_2\nu_2|\mu_3\nu_3}(p_1^\sigma, p_2^\sigma, \Lambda) = & \frac{\varkappa}{4} \left\{ \frac{1}{2} p_1^{\mu_3} p_2^{\nu_3} \hat{b}^{\mu_1\mu_2} \hat{b}^{\nu_1\nu_2} - p_1^{\mu_3} p_2^{\mu_1} \hat{b}^{\nu_1\mu_2} \hat{b}^{\nu_2\nu_3} \right. \\ & + (p_1 \cdot p_2) \left(-\frac{1}{2} \hat{b}^{\mu_1\nu_1} \hat{b}^{\mu_2\mu_3} \hat{b}^{\nu_2\nu_3} + \hat{b}^{\mu_1\nu_2} \hat{b}^{\mu_2\nu_3} \hat{b}^{\mu_3\nu_1} \right. \\ & \left. \left. - \frac{1}{4} \hat{b}^{\mu_1\mu_2} \hat{b}^{\nu_1\nu_2} \hat{b}^{\mu_3\nu_3} + \frac{1}{8} \hat{b}^{\mu_1\nu_1} \hat{b}^{\mu_2\nu_2} \hat{b}^{\mu_3\nu_3} \right) \right\} \\ & + i\varkappa\Lambda \left\{ \frac{1}{8} \hat{b}^{\mu_1\nu_1} \hat{b}^{\mu_2\nu_2} \hat{b}^{\mu_3\nu_3} - \frac{1}{4} \hat{b}^{\mu_1\nu_1} \hat{b}^{\mu_2\nu_3} \hat{b}^{\mu_3\nu_2} + \hat{b}^{\mu_1\nu_2} \hat{b}^{\mu_2\nu_3} \hat{b}^{\mu_3\nu_1} \right\} \end{aligned} \quad (11b)$$

and

$$\mathfrak{L}_4^{\mu_1\nu_1|\dots|\mu_4\nu_4}(p_1^\sigma, \dots, p_4^\sigma, \Lambda) = \frac{i}{16} \sum_{\mu_i \leftrightarrow \nu_i} \sum_{s \in S_4} \mathfrak{I}_4^{\mu_{s(1)}\nu_{s(1)}|\dots|\mu_{s(4)}\nu_{s(4)}} \left(p_{s(1)}^\sigma, p_{s(2)}^\sigma \right) \quad (12a)$$

⁵Where we treat the tuples of indices $\mu_i\nu_i$ as one index, i.e. exclude the a priori possible term $\hat{b}^{\mu_1\nu_1} \hat{b}_{\mu_3\nu_3}$ on the right hand side.

⁶We have used momentum conservation, i.e. performed a partial integration on the Lagrange density for General Relativity, to obtain a more compact form.

with

$$\begin{aligned}
t_4^{\mu_1\nu_1|\dots|\mu_4\nu_4}(p_1^\sigma, p_2^\sigma, \Lambda) = & \frac{\varkappa}{4} \left\{ -p_1^{\mu_3} p_2^{\nu_3} \hat{b}^{\mu_1\mu_2} \hat{b}^{\nu_1\mu_4} \hat{b}^{\nu_2\nu_4} + 2p_1^{\mu_3} p_2^{\mu_1} \hat{b}^{\nu_1\mu_2} \hat{b}^{\nu_2\mu_4} \hat{b}^{\nu_3\nu_4} \right. \\
& - \frac{1}{2} p_1^{\mu_3} p_2^{\mu_1} \hat{b}^{\nu_1\mu_2} \hat{b}^{\nu_2\nu_3} \hat{b}^{\mu_4\nu_4} + p_1^{\mu_3} p_2^{\nu_3} \hat{b}^{\mu_1\mu_2} \hat{b}^{\nu_1\nu_2} \hat{b}^{\mu_4\nu_4} \\
& - \frac{1}{2} p_1^{\mu_3} p_2^{\mu_4} \hat{b}^{\mu_1\mu_2} \hat{b}^{\nu_1\nu_2} \hat{b}^{\nu_3\nu_4} + p_1^{\mu_3} p_2^{\mu_4} \hat{b}^{\mu_1\mu_2} \hat{b}^{\nu_1\nu_3} \hat{b}^{\nu_2\nu_4} \\
& - \frac{1}{2} p_1^{\mu_2} p_2^{\mu_3} \hat{b}^{\mu_1\nu_1} \hat{b}^{\nu_2\mu_4} \hat{b}^{\nu_3\nu_4} + \frac{1}{4} p_1^{\mu_2} p_2^{\mu_1} \hat{b}^{\nu_1\nu_2} \hat{b}^{\mu_3\mu_4} \hat{b}^{\nu_3\nu_4} \\
& + (p_1 \cdot p_2) \left(-\frac{1}{16} \hat{b}^{\mu_1\mu_2} \hat{b}^{\nu_1\nu_2} \hat{b}^{\mu_3\nu_3} \hat{b}^{\mu_4\nu_4} + \frac{1}{8} \hat{b}^{\mu_1\mu_2} \hat{b}^{\nu_1\nu_2} \hat{b}^{\mu_3\mu_4} \hat{b}^{\nu_3\nu_4} \right. \\
& + \frac{1}{2} \hat{b}^{\mu_1\mu_2} \hat{b}^{\nu_1\mu_3} \hat{b}^{\nu_2\nu_3} \hat{b}^{\mu_4\nu_4} - \hat{b}^{\mu_1\mu_2} \hat{b}^{\nu_1\mu_3} \hat{b}^{\nu_2\mu_4} \hat{b}^{\nu_3\nu_4} \\
& + \frac{1}{2} \hat{b}^{\mu_1\mu_3} \hat{b}^{\nu_1\nu_3} \hat{b}^{\mu_2\mu_4} \hat{b}^{\nu_2\nu_4} - \frac{1}{2} \hat{b}^{\mu_1\mu_3} \hat{b}^{\nu_1\mu_4} \hat{b}^{\mu_2\nu_3} \hat{b}^{\nu_2\nu_4} \\
& - \frac{1}{4} \hat{b}^{\mu_1\nu_1} \hat{b}^{\mu_2\mu_3} \hat{b}^{\nu_2\nu_3} \hat{b}^{\mu_4\nu_4} + \frac{1}{2} \hat{b}^{\mu_1\nu_1} \hat{b}^{\mu_2\mu_3} \hat{b}^{\nu_2\mu_4} \hat{b}^{\nu_3\nu_4} \\
& \left. + \frac{1}{32} \hat{b}^{\mu_1\nu_1} \hat{b}^{\mu_2\nu_2} \hat{b}^{\mu_3\nu_3} \hat{b}^{\mu_4\nu_4} - \frac{1}{8} \hat{b}^{\mu_1\nu_1} \hat{b}^{\mu_2\nu_2} \hat{b}^{\mu_3\mu_4} \hat{b}^{\nu_3\nu_4} \right) \Big\} \\
& + i\varkappa\Lambda \left\{ \frac{57}{16} \hat{b}^{\mu_1\nu_1} \hat{b}^{\mu_2\nu_2} \hat{b}^{\mu_3\nu_3} \hat{b}^{\mu_4\nu_4} + \frac{15}{2} \hat{b}^{\mu_1\nu_1} \hat{b}^{\mu_2\nu_3} \hat{b}^{\mu_3\nu_4} \hat{b}^{\mu_4\nu_2} \right. \\
& - \frac{13}{8} \hat{b}^{\mu_1\nu_1} \hat{b}^{\mu_2\nu_2} \hat{b}^{\mu_3\nu_4} \hat{b}^{\mu_4\nu_3} + \frac{7}{8} \hat{b}^{\mu_1\nu_2} \hat{b}^{\mu_2\nu_1} \hat{b}^{\mu_3\nu_4} \hat{b}^{\mu_4\nu_3} \\
& \left. - 12 \hat{b}^{\mu_1\nu_2} \hat{b}^{\mu_2\nu_3} \hat{b}^{\mu_3\nu_4} \hat{b}^{\mu_4\nu_1} \right\}
\end{aligned} \tag{12b}$$

We remark that the three- and four-valent graviton vertex Feynman rules agree with the cited literature modulo prefactors and minus signs. Additionally, we remark that the three- and four-valent graviton-ghost Feynman rules are given in [19, Example 4.15] and remain valid with the replacement $\hat{\eta} \rightsquigarrow \hat{b}$.

3 BRST setup, Faddeev–Popov ghosts and symmetric ghosts

In this section, we discuss the generalization of the results from [20, 21]: This includes the diffeomorphism BRST and anti-BRST operators, the Faddeev–Popov ghost Lagrange density as well as the symmetric ghost Lagrange density and their relation. More precisely, the diffeomorphism BRST operator P is given as the following odd vector field on the spacetime-matter bundle with graviton-ghost degree 1:

$$P := \frac{1}{\zeta} \left(\nabla_\mu^{TM} C_\nu + \nabla_\nu^{TM} C_\mu \right) \frac{\partial}{\partial h_{\mu\nu}} + \varkappa C^\rho (\partial_\rho C^\sigma) \frac{\partial}{\partial C^\sigma} + \frac{1}{\zeta} B^\sigma \frac{\partial}{\partial C^\sigma} \tag{13}$$

Equivalently, its action on fundamental particle fields is given as follows:

$$Ph_{\mu\nu} := \frac{1}{\zeta} \left(\nabla_\mu^{TM} C_\nu + \nabla_\nu^{TM} C_\mu \right) \tag{14a}$$

$$PC^\rho := \varkappa C^\sigma (\partial_\sigma C^\rho) \tag{14b}$$

$$P\bar{C}^\rho := \frac{1}{\zeta} B^\rho \quad (14c)$$

$$PB^\rho := 0 \quad (14d)$$

$$Pb_{\mu\nu} := 0 \quad (14e)$$

Additionally, we define the diffeomorphism anti-BRST operator \bar{P} as the following odd vector field on the spacetime-matter bundle with graviton-ghost degree -1:

$$\bar{P} := P \Big|_{C \rightsquigarrow \bar{C}} \quad (15a)$$

together with the following additional changes

$$\bar{P}C^\rho := -\frac{1}{\zeta} B^\rho + \varkappa \left(\bar{C}^\sigma (\partial_\sigma C^\rho) - (\partial_\sigma \bar{C}^\rho) C^\sigma \right) \quad (15b)$$

$$\bar{P}\bar{C}^\rho := \varkappa \bar{C}^\sigma (\partial_\sigma \bar{C}_\rho) \quad (15c)$$

$$\bar{P}B^\rho := \varkappa \left(\bar{C}^\sigma (\partial_\sigma B_\rho) - (\partial_\sigma \bar{C}^\rho) B^\sigma \right) \quad (15d)$$

We remark the characteristic identities $[P, P] = [P, \bar{P}] = [\bar{P}, \bar{P}] = 0$,⁷ cf. [20, Proposition 3.2 and Corollary 3.8] and the references therein.

Remark 3.1. The definition of the diffeomorphism BRST operator given above is identical to the case of a flat background metric, cf. [20, Definition 3.1 and Definition 3.7] and [21, Definition 2.2] and the references therein: The crucial point is that the background metric $\eta_{\mu\nu}$ or $b_{\mu\nu}$ is set to be invariant with respect to P , as in Equation (14e). This is achieved by defining its transformation into the graviton field $h_{\mu\nu}$.

Corollary 3.2. *The gauge fixing Lagrange density and its accompanying Faddeev–Popov ghost Lagrange density for (effective) Quantum General Relativity with a cosmological constant read*

$$\begin{aligned} \mathcal{L}_{\text{GR-}\Lambda\text{-GF}} + \mathcal{L}_{\text{GR-}\Lambda\text{-Ghost}} &= -\frac{1}{4\varkappa^2\zeta} b^{\mu\nu} d\mathcal{D}_\mu^{(1)} d\mathcal{D}_\nu^{(1)} dV_b \\ &\quad - \frac{1}{2\zeta} b^{\rho\sigma} \bar{C}^\mu (\partial_\rho \partial_\sigma C_\mu) dV_b \\ &\quad - \frac{1}{2} b^{\rho\sigma} \bar{C}^\mu \left(\partial_\mu (\Gamma^\nu_{\rho\sigma} C_\nu) - 2\partial_\rho (\Gamma^\nu_{\mu\sigma} C_\nu) \right) dV_b \end{aligned} \quad (16)$$

with the linearized de Donder gauge fixing functional $d\mathcal{D}_\mu^{(1)} := b^{\rho\sigma} \Gamma_{\rho\sigma\mu}$. They can be obtained from the gauge fixing fermion

$$\zeta^{(1)} := \frac{1}{2} \bar{C}^\rho \left(\frac{1}{\varkappa} d\mathcal{D}_\rho^{(1)} + \frac{1}{2} B_\rho \right) dV_b \quad (17)$$

via $P\zeta^{(1)}$.

Proof. This can be shown analogously to the proof of [20, Proposition 3.4]. ■

⁷We emphasize that $[\cdot, \cdot]$ denotes the supercommutator, which is equivalent to the anticommutator in the previous equation.

Corollary 3.3. *The gauge fixing Lagrange density and its accompanying symmetric ghost Lagrange density for (effective) Quantum General Relativity with a cosmological constant read*

$$\begin{aligned}
\mathcal{L}_{\text{GR-}\Lambda\text{-GF-Ghost}} = & -\frac{1}{2\zeta} \left(\frac{1}{2\kappa^2} b^{\mu\nu} d\mathcal{D}_\mu^{(1)} d\mathcal{D}_\nu^{(1)} + b^{\mu\nu} (\partial_\mu \bar{C}^\rho) (\partial_\nu C_\rho) \right) dV_b \\
& - \frac{1}{2} b^{\mu\nu} \bar{C}^\rho \left(\frac{1}{2} \partial_\rho (\Gamma^\sigma{}_{\mu\nu} C_\sigma) - \partial_\mu (\Gamma^\sigma{}_{\rho\nu} C_\sigma) \right) dV_b \\
& + \frac{1}{2} b^{\mu\nu} \left(\frac{1}{2} \partial_\rho (\Gamma^\sigma{}_{\mu\nu} \bar{C}_\sigma) - \partial_\mu (\Gamma^\sigma{}_{\rho\nu} \bar{C}_\sigma) \right) C^\rho dV_b \\
& - \frac{\kappa^2 \zeta}{8} b_{\mu\nu} \left(\bar{C}^\rho (\partial_\rho \bar{C}^\mu) \right) \left(C^\sigma (\partial_\sigma C^\nu) \right) dV_b
\end{aligned} \tag{18}$$

with the linearized de Donder gauge fixing functional $d\mathcal{D}_\mu^{(1)} := b^{\rho\sigma} \Gamma_{\rho\sigma\mu}$. It can be obtained from the gauge fixing boson

$$F := \frac{\zeta}{4} \left(\frac{1}{\kappa} b^{\mu\nu} h_{\mu\nu} + \bar{C}^\rho C_\rho \right) dV_b \tag{19}$$

via $\mathcal{P}F$, where $\mathcal{P} := P \circ \bar{P}$ is the diffeomorphism super-BRST operator, cf. [21, Definition 2.5].

Proof. This can be shown analogously to the proof of [21, Proposition 3.1]. ■

Corollary 3.4. *We obtain the following homotopy in $\lambda \in [0, 1]$ between the Faddeev–Popov construction $\lambda = 0$, the symmetric setup $\lambda = 1/2$ and the opposed Faddeev–Popov construction $\lambda = 1$:*

$$\begin{aligned}
\mathcal{L}_{\text{GR-}\Lambda\text{-GF-Ghost}}(\lambda) = & -\frac{1}{2\zeta} \left(\frac{1}{2\kappa^2} b^{\mu\nu} d\mathcal{D}_\mu^{(1)} d\mathcal{D}_\nu^{(1)} + b^{\mu\nu} (\partial_\mu \bar{C}^\rho) (\partial_\nu C_\rho) \right) dV_b \\
& - (1 - \lambda) b^{\mu\nu} \bar{C}^\rho \left(\frac{1}{2} \partial_\rho (\Gamma^\sigma{}_{\mu\nu} C_\sigma) - \partial_\mu (\Gamma^\sigma{}_{\rho\nu} C_\sigma) \right) dV_b \\
& + \lambda b^{\mu\nu} \left(\frac{1}{2} \partial_\rho (\Gamma^\sigma{}_{\mu\nu} \bar{C}_\sigma) - \partial_\mu (\Gamma^\sigma{}_{\rho\nu} \bar{C}_\sigma) \right) C^\rho dV_b \\
& - \frac{\kappa^2 \zeta}{8} \lambda (1 - \lambda) b_{\mu\nu} \left(\bar{C}^\rho (\partial_\rho \bar{C}^\mu) \right) \left(C^\sigma (\partial_\sigma C^\nu) \right) dV_b
\end{aligned} \tag{20}$$

Proof. This can be shown analogously to the proof of [21, Theorem 3.2]. ■

4 Transversality

Finally, we study the transversality of (effective) Quantum General Relativity with a cosmological constant as a generalization of [22]. Therein, in Definition 3.14, we introduced the transversal structure of (effective) Quantum General Relativity with a de Donder gauge fixing (QGR) as the set $\mathcal{T}_{\text{QGR}} := \{\mathbb{L}, \mathbb{I}, \mathbb{T}\}$. These three operators are the longitudinal, identical and transversal projection tensors, given in our generalized setting as follows:⁸

$$\mathbb{L}_{\mu\nu}^{\rho\sigma} := \frac{1}{2p^2} \left(\delta_\mu^\rho p^\sigma p_\nu + \delta_\mu^\sigma p^\rho p_\nu + \delta_\nu^\rho p^\sigma p_\mu + \delta_\nu^\sigma p^\rho p_\mu - 2b^{\rho\sigma} p_\mu p_\nu \right), \tag{21a}$$

$$\mathbb{I}_{\mu\nu}^{\rho\sigma} := \frac{1}{2} \left(\delta_\mu^\rho \delta_\nu^\sigma + \delta_\mu^\sigma \delta_\nu^\rho \right) \tag{21b}$$

⁸From now on, we omit the hat for Fourier transformed quantities to improve readability.

and

$$\mathbb{T}_{\mu\nu}^{\rho\sigma} := \mathbb{I}_{\mu\nu}^{\rho\sigma} - \mathbb{L}_{\mu\nu}^{\rho\sigma}, \quad (21c)$$

where we have set $p^2 := b_{\mu\nu}p^\mu p^\nu$. Lorentz indices on \mathbb{L} , \mathbb{I} and \mathbb{T} are raised and lowered with the metric \mathbb{G} , defined via⁹

$$\mathbb{G}_{\mu\nu\rho\sigma} := \frac{1}{p^2} (b_{\mu\rho}b_{\nu\sigma} + b_{\mu\sigma}b_{\nu\rho} - b_{\mu\nu}b_{\rho\sigma}) \quad (22a)$$

and its inverse

$$\mathbb{G}^{\mu\nu\rho\sigma} := \frac{p^2}{4} (b^{\mu\rho}b^{\nu\sigma} + b^{\mu\sigma}b^{\nu\rho} - b^{\mu\nu}b^{\rho\sigma}). \quad (22b)$$

All these tensors share interesting relations, which are listed in [22, Subsection 3.2].

Theorem 4.1. *The Feynman rule of the graviton propagator with a cosmological constant can be written as follows:*

$$\Phi(\text{wavy}) = -\frac{2ip^2}{p^2 + \Lambda + i\epsilon} \left(\mathbb{G}_{\mu\nu\rho\sigma} - \left(\frac{1 - \zeta}{p^2 + \zeta\Lambda} \right) \mathbb{L}_{\mu\nu\rho\sigma} \right) \quad (23)$$

In particular, we obtain the following two identities:

$$\Phi(\text{wavy}) \Big|_{\zeta=0} = -\frac{2ip^2}{p^2 + \Lambda + i\epsilon} \mathbb{T}_{\mu\nu\rho\sigma} \quad (24a)$$

and

$$\Phi(\text{wavy}) \Big|_{\zeta=1} = -\frac{2ip^2}{p^2 + \Lambda + i\epsilon} \mathbb{G}_{\mu\nu\rho\sigma} \quad (24b)$$

Proof. This follows directly from Theorem 2.3 and [22, Definition 3.14], reproduced in Equations (21) and (22). ■

Remark 4.2. In particular, we see from Theorem 4.1 that the de Donder gauge fixing is not the optimal gauge fixing condition for (effective) Quantum General Relativity with a cosmological constant, cf. [22, Definition 3.2]. This comes from the fact that the cosmological constant behaves in the propagator like an Euclidean mass term and thus alters the longitudinal and transversal modes. Interestingly, the transversal behavior for the two cases $\zeta = 0$ and $\zeta = 1$ agrees with the graviton propagator without a cosmological constant, cf. [22, Theorem 3.21].

Theorem 4.3. *The Feynman rule of the three-valent graviton vertex with a cosmological constant does not satisfy the contraction identity, i.e. we obtain:*

$$\Phi \left(\begin{array}{c} L \\ \text{wavy} \\ L \end{array} \right) \not\stackrel{\text{MC}}{=} 0, \quad (25)$$

where \simeq_{MC} indicates equality modulo momentum conservation.

⁹This metric is known in the literature as the de Witt metric [29], cf. [30]. The reason for the asymmetric definition concerning the factor 1/4 is motivated by [22, Equations (57)].

Proof. This is verified by the following calculation: First, we use the following decomposition of $\mathbb{L}_{\mu\nu}^{\rho\sigma} = \mathcal{L}_\tau^{\rho\sigma} \mathcal{G}_{\mu\nu}^\tau$ into the two tensors $\mathcal{G}_{\mu\nu}^\kappa := \frac{1}{p^2} (p_\mu \delta_\nu^\kappa + p_\nu \delta_\mu^\kappa)$ and $\mathcal{L}_\lambda^{\rho\sigma} := \frac{1}{2} (p^\rho \delta_\lambda^\sigma + p^\sigma \delta_\lambda^\rho - p_\lambda b^{\rho\sigma})$ that has been established in [22, Lemma 3.16]. Then, we recall from [22, Theorem 3.23]

$$\Phi \left(\begin{array}{c} \mathcal{G} \\ \mathcal{G} \text{ --- } \mathcal{G} \\ \mathcal{G} \end{array} \right) \Big|_{O(\Lambda^0)} \simeq_{\text{MC}} 0. \quad (26)$$

Thus, using Equations (11), the claimed result follows from

$$\begin{aligned} \Phi \left(\begin{array}{c} \mathcal{G} \\ \mathcal{G} \text{ --- } \mathcal{G} \\ \mathcal{G} \end{array} \right) \Big|_{O(\Lambda^1)} &\simeq_{\text{MC}} -\frac{i\kappa\Lambda}{p_1^2 p_2^2 (p_1 + p_2)^2} \left(b^{\tau_1 \tau_2} \left(-p_1^2 p_2^{\tau_3} - p_2^2 p_1^{\tau_3} \right) \right. \\ &\quad \left. + b^{\tau_1 \tau_3} \left(p_1^2 (p_1^{\tau_2} + p_2^{\tau_2}) - (p_1 + p_2)^2 p_1^{\tau_2} \right) \right. \\ &\quad \left. + b^{\tau_2 \tau_3} \left(p_2^2 (p_1^{\tau_1} + p_2^{\tau_1}) - (p_1 + p_2)^2 p_2^{\tau_1} \right) \right) \end{aligned} \quad (27)$$

$$\not\simeq_{\text{MC}} 0,$$

which does only vanish on-shell. ■

Remark 4.4. Equation (25) indicates that the question of transversality is more involved, if a cosmological constant is added to (effective) Quantum General Relativity. This will be studied in future work.

5 Conclusion

We have studied (effective) Quantum General Relativity with a cosmological constant from the viewpoint of renormalization. To this end, we started with its Lagrange density in Equations (1) and discussed the corresponding quantum gauge symmetries in Equations (4) together with appropriate renormalization conditions for the graviton propagator in Equations (6), cf. [18] for the situation of a vanishing cosmological constant. Then, we calculated the corresponding graviton vertex Feynman rules for arbitrary valence in Theorem 2.2 and the graviton propagator Feynman rule for a general gauge parameter in Theorem 2.3. Additionally, we displayed the three-valent and four-valent vertex Feynman rules explicitly in Example 2.4, cf. [19] for the situation of a vanishing cosmological constant. Next, we constructed the Faddeev–Popov ghost, symmetric ghost and homotopy ghost Lagrange densities in Corollaries 3.2, 3.3 and 3.4 from the viewpoint of BRST cohomology. We refer to [20, 21] for the situation of a vanishing cosmological constant. In addition, we studied its transversality, in particular the decomposition of the graviton propagator Feynman rule in Theorem 4.1 and the contraction identity of the three-valent graviton vertex Feynman rule in Theorem 4.3, cf. [22] for the situation of a vanishing cosmological constant.

Finally, we want to mention a different view on the cosmological constant: The possibility to use it as a diffeomorphism-invariant regulator for pure gravity. In this case, a convenient choice would be $\Lambda \in \mathbb{C}$ and $b_{\mu\nu} := \eta_{\mu\nu}$, so that the final result (after the limit $\Lambda \mapsto 0$) can be considered on the Minkowski background spacetime. This claim is due to the fact that (effective) Quantum General Relativity with only Λ -vertices is renormalizable by power counting. This will be studied in future work.

Acknowledgments

This article was initiated by a question asked by my supervisor Dirk Kreimer during my disputation. Having said this, it is my pleasure to express my gratitude for his insight and support over the past years! This research is supported by the *Riemann Center for Geometry and Physics* of the Leibniz University Hannover via a *Riemann fellowship* in the group of Domenico Giulini. It is my pleasure to thank Domenico Giulini, Philip Schwartz and Lorenzo Casarin for a welcoming atmosphere and an inspiring intellectual exchange.

References

- [1] G. 't Hooft and M. J. G. Veltman: *One loop divergencies in the theory of gravitation*. Ann. Inst. H. Poincaré Phys. Theor. A, 20:69–94, 1974.
- [2] M. H. Goroff and A. Sagnotti: *Quantum Gravity at Two Loops*. Phys. Lett. B, 160:81–86, 1985.
- [3] L. F. Abbott: *Introduction to the Background Field Method*. Acta Phys. Polon. B, 13:33, 1982.
- [4] K. Krasnov: *Effective metric Lagrangians from an underlying theory with two propagating degrees of freedom*. Phys. Rev. D, 81:084026, 2010. arXiv:0911.4903v1 [hep-th].
- [5] D. Kreimer: *On the Hopf algebra structure of perturbative quantum field theories*. Adv. Theor. Math. Phys., 2:303–334, 1998. arXiv:q-alg/9707029v4.
- [6] A. Connes and D. Kreimer: *Hopf Algebras, Renormalization and Noncommutative Geometry*. Commun. Math. Phys., 199:203–242, 1998. arXiv:hep-th/9808042v1.
- [7] A. Connes and D. Kreimer: *Renormalization in quantum field theory and the Riemann-Hilbert problem*. JHEP, 09:024, 1999. arXiv:hep-th/9909126v3.
- [8] A. Connes and D. Kreimer: *Renormalization in quantum field theory and the Riemann-Hilbert problem I: the Hopf algebra structure of graphs and the main theorem*. Commun. Math. Phys., 210:249–273, 1999. arXiv:hep-th/9912092v1.
- [9] A. Connes and D. Kreimer: *Renormalization in quantum field theory and the Riemann-Hilbert problem II: the β -function, diffeomorphisms and the renormalization group*. Commun. Math. Phys., 216:215–241, 2000. arXiv:hep-th/0003188v1.
- [10] D. Kreimer: *Anatomy of a gauge theory*. Annals Phys., 321:2757–2781, 2006. arXiv:hep-th/0509135v3.
- [11] W. D. van Suijlekom: *The Hopf algebra of Feynman graphs in QED*. Lett. Math. Phys., 77:265–281, 2006. arXiv:hep-th/0602126v2.
- [12] W. D. van Suijlekom: *Renormalization of gauge fields: A Hopf algebra approach*. Commun. Math. Phys., 276:773–798, 2007. arXiv:hep-th/0610137v1.
- [13] W. D. van Suijlekom: *The structure of renormalization Hopf algebras for gauge theories I: Representing Feynman graphs on BV-algebras*. Commun. Math. Phys., 290:291–319, 2009. arXiv:0807.0999v2 [math-ph].
- [14] D. Kreimer and W. D. van Suijlekom: *Recursive relations in the core Hopf algebra*. Nucl. Phys. B, 820:682–693, 2009. arXiv:0903.2849v1 [hep-th].
- [15] D. Kreimer: *A remark on quantum gravity*. Annals Phys., 323:49–60, 2008. arXiv:0705.3897v1 [hep-th].

- [16] D. Kreimer: *Not so non-renormalizable gravity.* "Quantum Field Theory: Competitive Models", B.Fausser, J.Tolksdorf, E.Zeidlers, eds., Birkhaeuser (2009), 2008. arXiv:0805.4545v1 [hep-th].
- [17] D. Prinz: *Algebraic Structures in the Coupling of Gravity to Gauge Theories.* Annals Phys., 426:168395, 2021. arXiv:1812.09919v3 [hep-th].
- [18] D. Prinz: *Gauge Symmetries and Renormalization.* Math. Phys. Anal. Geom., 25(3):20, 2022. arXiv:2001.00104v4 [math-ph].
- [19] D. Prinz: *Gravity-Matter Feynman Rules for any Valence.* Class. Quantum Grav., 38:215003, 2021. arXiv:2004.09543v4 [hep-th].
- [20] D. Prinz: *The BRST Double Complex for the Coupling of Gravity to Gauge Theories,* 2022. arXiv:2206.00780v1 [hep-th].
- [21] D. Prinz: *Symmetric Ghost Lagrange Densities for the Coupling of Gravity to Gauge Theories,* 2022. arXiv:2207.07593v1 [hep-th].
- [22] D. Prinz: *Transversality in the Coupling of Gravity to Gauge Theories,* 2022. arXiv:2208.14166v1 [hep-th].
- [23] D. Prinz: *Renormalization of Gauge Theories and Gravity.* PhD thesis, Humboldt University of Berlin, 2022. Available at <https://doi.org/10.18452/25401> and arXiv:2210.17510v1 [hep-th].
- [24] J. C. Ward: *An Identity in Quantum Electrodynamics.* Phys. Rev., 78:182, 1950.
- [25] Y. Takahashi: *On the Generalized Ward Identity.* Nuovo Cim., 6:371, 1957.
- [26] G. 't Hooft: *Renormalization of Massless Yang-Mills Fields.* Nucl. Phys. B, 33 (1):173–199, 1971.
- [27] J. C. Taylor: *Ward identities and charge renormalization of the Yang-Mills field.* Nucl. Phys. B, 33 (2):436–444, 1971.
- [28] A. A. Slavnov: *Ward identities in gauge theories.* Theor. Math. Phys., 10 (2):99–104, 1972.
- [29] B. S. DeWitt: *Quantum Theory of Gravity. I. The Canonical Theory.* Phys. Rev., 160:1113–1148, 1967.
- [30] D. Giulini and C. Kiefer: *Wheeler–DeWitt metric and the attractivity of gravity.* Phys. Lett. A, 193:21–24, 1994. arXiv:gr-qc/9405040v2.