

Constraint on Early Dark Energy from Isotropic Cosmic Birefringence

Johannes R. Eskilt,^{1,2,*} Laura Herold,³ Eiichiro Komatsu,^{3,4}
Kai Murai,^{5,4} Toshiya Namikawa,⁴ and Fumihiro Naokawa^{6,7}

¹*Institute of Theoretical Astrophysics, University of Oslo, P.O. Box 1029 Blindern, N-0315 Oslo, Norway*

²*Imperial Centre for Inference and Cosmology, Department of Physics, Imperial College London, Blackett Laboratory, Prince Consort Road, London SW7 2AZ, United Kingdom*

³*Max Planck Institute for Astrophysics, Karl-Schwarzschild-Str. 1, D-85748 Garching, Germany*

⁴*Kavli IPMU (WPI), UTIAS, The University of Tokyo, Kashiwa, 277-8583, Japan*

⁵*ICRR, The University of Tokyo, Kashiwa, 277-8582, Japan*

⁶*Department of Physics, Graduate School of Science,*

The University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan

⁷*Research Center for the Early Universe, The University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan*

(Dated: March 28, 2023)

Polarization of the cosmic microwave background (CMB) is sensitive to new physics violating parity symmetry, such as the presence of a pseudoscalar “axionlike” field. Such a field may be responsible for early dark energy (EDE), which is active prior to recombination and provides a solution to the so-called Hubble tension. The EDE field coupled to photons in a parity-violating manner would rotate the plane of linear polarization of the CMB and produce a cross-correlation power spectrum of E - and B -mode polarization fields with opposite parities. In this paper, we fit the EB power spectrum predicted by the photon-axion coupling of the EDE model with a potential $V(\phi) \propto [1 - \cos(\phi/f)]^3$ to polarization data from *Planck*. We find that the unique shape of the predicted EB power spectrum is not favored by the data and obtain a first constraint on the photon-axion coupling constant, $g = (0.04 \pm 0.16)M_{\text{Pl}}^{-1}$ (68% CL), for the EDE model that best fits the CMB and galaxy clustering data. This constraint is independent of the miscalibration of polarization angles of the instrument or the polarized Galactic foreground emission. Our limit on g may have important implications for embedding EDE in fundamental physics, such as string theory.

I. INTRODUCTION

The standard cosmological model, called Λ CDM, includes new physics beyond the standard model of elementary particles and fields, such as dark matter and dark energy [1]. Clues to their physical nature may be found in possible deviations from the Λ CDM model. In recent years, a growing number of such deviations, or “tensions,” have been reported [2], which may point towards new physics. In this paper, we study a fascinating connection between two hints of new physics: early dark energy (EDE) as a solution to the so-called Hubble tension (see Refs. [3, 4] for reviews) and cosmic birefringence, a rotation of the plane of linear polarization of photons (see Ref. [5] for a review).

EDE, which was active prior to the epoch of recombination at a redshift of $z \simeq 1090$ [6–8], can resolve the Hubble tension [9, 10] by modifying the value of the Hubble constant, H_0 , inferred from the cosmic microwave background (CMB) data and brings it into agreement with H_0 inferred from the local distance ladder [11]. But, is EDE *the* solution to the Hubble tension? To make progress, one must look elsewhere for corroborating evidence.

In this paper, we look for a signature of EDE in polarization of the CMB. If the EDE field, ϕ , is a pseudoscalar “axionlike” field, it could couple to electromagnetism in

a parity-violating manner in the Lagrangian density, \mathcal{L} . We write [12, 13]

$$\mathcal{L} = -\frac{1}{2}(\partial\phi)^2 - V(\phi) - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{4}g\phi F_{\mu\nu}\tilde{F}^{\mu\nu}, \quad (1)$$

where g is the photon-axion coupling constant, and $F_{\mu\nu}$ and $\tilde{F}^{\mu\nu}$ are the field strength tensor of the photon field and its dual tensor, respectively.

The last term in Eq. (1) is a Chern-Simons term, which violates parity symmetry in the presence of a spacetime-dependent condensate of ϕ . This term appears naturally for an axionlike field with $g = c_{\phi\gamma}\alpha_{\text{em}}/(2\pi f)$, where $c_{\phi\gamma}$ is an anomaly coefficient, $\alpha_{\text{em}} \simeq 1/137$ the electromagnetic fine-structure constant, and f the axion decay constant (see, e.g., Eq. (24) of Ref. [14]), and has been considered for EDE models in Refs. [15–19].

We assume a “canonical” EDE potential, $V(\phi) = V_0[1 - \cos(\phi/f)]^3$, where V_0 is the normalization. This model can resolve the Hubble tension [10, 20–26]. See Refs. [27–34] for other constraints on this model. For other EDE models that can resolve the Hubble tension, see Ref. [4] and references therein.

To probe violation of parity symmetry in the polarization pattern of the CMB, one can decompose a pixelized map of the observed Stokes parameters into eigenstates of parity called E and B modes [35, 36]:

$$Q(\hat{\mathbf{n}}) \pm iU(\hat{\mathbf{n}}) = - \sum_{\ell=2}^{\ell_{\text{max}}} \sum_{m=-\ell}^{\ell} (E_{\ell m} \pm iB_{\ell m})_{\pm 2} Y_{\ell}^m(\hat{\mathbf{n}}), \quad (2)$$

* j.r.eskilt@astro.uio.no

where $\hat{\mathbf{n}}$ is the direction of an observer’s line of sight, $E_{\ell m}$ and $B_{\ell m}$ are the spherical harmonics coefficients of the E and B modes, respectively, $_{\pm 2}Y_{\ell}^m(\hat{\mathbf{n}})$ are the spin-2 spherical harmonics, and ℓ_{\max} is the maximum multipole used for the analysis. The coefficients transform under inversion of spatial coordinates, $\hat{\mathbf{n}} \rightarrow -\hat{\mathbf{n}}$, as $E_{\ell m} \rightarrow (-1)^{\ell} E_{\ell m}$ and $B_{\ell m} \rightarrow (-1)^{\ell+1} B_{\ell m}$. The cross-power spectrum of E and B modes, $C_{\ell}^{EB} \equiv (2\ell + 1)^{-1} \sum_m \text{Re}(E_{\ell m} B_{\ell m}^*)$, has odd parity and is sensitive to parity violation [37].

In this paper, we use the EB power spectrum presented in Ref. [38] to constrain the Chern-Simons term in Eq. (1). Specifically, this term makes phase velocities of right- and left-handed circular polarization states different, which rotates the plane of linear polarization by an angle $\beta(\hat{\mathbf{n}}) = \frac{1}{2}g[\phi(\eta_o) - \phi(\eta_e, r\hat{\mathbf{n}})]$ as photons have traveled from the conformal time of emission, η_e , to the observation, η_o [39–41]. Here, $r = c(\eta_o - \eta_e)$ is the conformal distance to the emitter. As there is no evidence for anisotropic birefringence [42–46], we assume that β is independent of $\hat{\mathbf{n}}$. See Ref. [15] for a study on anisotropic birefringence from the EDE field.

A distinct feature of the EDE field is that ϕ evolves significantly during the epoch of recombination, which has profound implications for observational tests of EDE models with photon-axion coupling [18, 19, 47]. If ϕ were constant during the epoch of recombination and evolved only later, the observed E and B modes (denoted by the superscript “o”) would be given by $E_{\ell m}^o = E_{\ell m} \cos(2\beta) - B_{\ell m} \sin(2\beta)$ and $B_{\ell m}^o = E_{\ell m} \sin(2\beta) + B_{\ell m} \cos(2\beta)$, respectively, and [48]

$$C_{\ell}^{EB,o} = \frac{\sin(4\beta)}{2} (C_{\ell}^{EE} - C_{\ell}^{BB}) + \cos(4\beta) C_{\ell}^{EB}, \quad (3)$$

where the last term is the intrinsic EB correlation at the time of emission, and C_{ℓ}^{EE} and C_{ℓ}^{BB} are the auto-power spectra of E and B modes at emission, respectively.

Eq. (3) has been assumed in all of the previous constraints on isotropic cosmic birefringence including Refs. [38, 49–51], which reported a tantalizing hint of β with a statistical significance of $> 3\sigma$. If, however, ϕ evolved significantly during the epoch of recombination, $C_{\ell}^{EB,o}$ would no longer be given by Eq. (3) and exhibit complex dependence on ℓ [52, 53], which allows us to distinguish between different origins of cosmic birefringence [18, 19, 47].

In this paper, we present a first constraint on the photon-axion coupling constant, g , from the *shape* of the EB power spectrum. This is a powerful new approach that breaks the degeneracy between cosmic birefringence and an instrumental miscalibration of polarization angles of detectors, α . Specifically, Eq. (3) can be generated not only by cosmic birefringence but also by α [54–57]. Therefore, β in Eq. (3) needs to be replaced by the sum, $\alpha + \beta$, and we cannot distinguish between α and β unless one calibrates α well [58] or uses other information such as the Galactic foreground emission [59] and the so-called “reionization bump” at $\ell \lesssim 10$ [60]. Due to this

complication, the current hint of β remains somewhat inconclusive [61]. Our new approach, based on the shape of the EB power spectrum different from $C_{\ell}^{EE} - C_{\ell}^{BB}$, is free from this complication.

II. METHOD

We use the so-called “NPIPE-processed” full-sky polarization maps from the High-Frequency Instrument (HFI) of the *Planck* mission at frequencies of $\nu = 100, 143, 217,$ and 353 GHz [62, 63]. The NPIPE processing divided the set of detectors for each frequency band into two groups, and produced 8 full-sky maps in total.

Unlike in Ref. [38], we do not include WMAP or *Planck* Low-Frequency Instrument data for simplicity of the data analysis. Including them would improve the constraints modestly by about 10%.

We use two masks [38, 51]. One is a small mask, leaving nearly full-sky data available for the analysis. The other is a large mask, removing 30% of the Galactic plane. Both masks remove pixels containing polarized point sources given in the official *Planck* point-source maps and pixels where the carbon-monoxide (CO) emission is brighter than $45 \text{ K}_{\text{RJ}} \text{ km s}^{-1}$. These masks leave the sky fractions of $f_{\text{sky}} = 0.92$ and 0.62 for small and large masks, respectively, available for the analysis.

We calculate the EB power spectra of 8 masked polarization maps using `PolSpice`¹ [64]. We then beam-deconvolve the EB power spectra by using the official NPIPE beam transfer functions and the pixel window functions from the `HEALPix` library [65].

We calculate the inverse-variance weighted average of the beam-deconvolved EB power spectra from

$$\bar{C}_b^{EB,o} \equiv \text{E} \left(C_b^{EB,o} \right) = \frac{\vec{\mathbf{1}} \cdot \mathbf{M}_b^{-1} \cdot \vec{C}_b^{EB}}{\vec{\mathbf{1}} \cdot \mathbf{M}_b^{-1} \cdot \vec{\mathbf{1}}}, \quad (4)$$

which we call a “stacked EB power spectrum” [38]. The variance is given by

$$\text{Var} \left(C_b^{EB,o} \right) = \frac{1}{\vec{\mathbf{1}} \cdot \mathbf{M}_b^{-1} \cdot \vec{\mathbf{1}}}. \quad (5)$$

Here, $\vec{\mathbf{1}}$ is a unit vector, \vec{C}_b^{EB} is the binned set of all combinations of the observed beam-deconvolved EB power spectra, $C_{\ell}^{E_i B_j, o}$, where i and j denote frequency bands, and \mathbf{M}_b is the binned covariance matrix for \vec{C}_b^{EB} .

We bin $C_{\ell}^{E_i B_j, o}$ and \mathbf{M}_{ℓ} as

$$\vec{C}_b^{EB} = \frac{1}{\Delta\ell} \sum_{\ell \in b} \vec{C}_{\ell}^{EB}, \quad \mathbf{M}_b = \frac{1}{\Delta\ell^2} \sum_{\ell \in b} \mathbf{M}_{\ell}, \quad (6)$$

¹ <http://www2.iap.fr/users/hivon/software/PolSpice/>

where

$$\mathbf{M}_\ell = \text{Cov} \left(C_\ell^{E_i B_j, \circ}, C_\ell^{E_p B_q, \circ} \right) \simeq \frac{C_\ell^{E_i E_p, \circ} C_\ell^{B_j B_q, \circ}}{(2\ell + 1) f_{\text{sky}}}. \quad (7)$$

We neglected the term $C_\ell^{E_i B_q, \circ} C_\ell^{E_p B_j, \circ}$, which is much smaller than the other term. As we use the observed power spectra in \mathbf{M}_ℓ , this term fluctuates around zero and biases \mathbf{M}_ℓ when included; thus, it is best to neglect it [59]. We also excluded the auto-power spectra ($i = j$) due to noise domination at high ℓ for the EE and BB power spectra. This does not affect $C_b^{E_i B_j}$ directly, but it does affect \mathbf{M}_b . Following the previous work [38, 49–51, 66], we bin the power spectra over 20 multipoles, i.e., $\Delta\ell = 20$. Our multipole range is from $\ell_{\text{min}} = 51$ to $\ell_{\text{max}} = 1490$, which gives a total of 72 bins. The stacked EB power spectrum is publicly available².

The aim of this paper is to constrain the coupling, g , to pre-recombination EDE while marginalizing over post-recombination cosmic birefringence and miscalibration angles, $\alpha + \beta$. To this end, we fit the stacked EB power spectrum for g and $\alpha + \beta$ simultaneously. We sample these parameters using a Markov Chain Monte Carlo sampler `emcee` [67]. The log-likelihood function is $-2 \ln \mathcal{L} = \sum_b v_b^2 / \text{Var} \left(C_b^{EB, \circ} \right)$, where

$$v_b \equiv \bar{C}_b^{EB, \circ} - \cos [4(\alpha + \beta)] g M_{\text{Pl}} C_b^{EB, \text{EDE}} - \frac{\sin [4(\alpha + \beta)]}{2} \left(C_b^{EE, \text{CMB}} - C_b^{BB, \text{CMB}} \right). \quad (8)$$

We compute the CMB EE and BB power spectra using `CAMB`³ [68] with the best-fitting Λ CDM parameters of the *Planck* 2018 analysis [69]. The EDE EB power spectrum is computed using a modified version of the `CLASS` code [70] developed in Ref. [18] with $g = M_{\text{Pl}}^{-1}$, as g just gives the amplitude of $C_\ell^{EB, \text{EDE}}$ and can be rescaled later. Here, $M_{\text{Pl}} \simeq 2.4 \times 10^{18}$ GeV is the reduced Planck mass.

The EDE model has 3 parameters in addition to the standard Λ CDM parameters: f_{EDE} , z_c , and θ_i . Here, f_{EDE} is the maximum energy density fraction of the EDE field reached at a redshift z_c , while θ_i is a dimensionless initial value of the EDE field, $\theta_i \equiv \phi_i / f$. The first two are related to the fundamental parameters in the potential, V_0 and f .

We fix all EDE and Λ CDM parameters to the best-fitting parameters from Ref. [26] for two different data sets (see Table I) and sample only g and $\alpha + \beta$. As described in Ref. [26], the baseline data set includes the *Planck* temperature and polarization power spectra [71] and the galaxy power spectra from the Baryon Oscillation Spectroscopic Survey (BOSS) Data Release 12 [72] (second column), while the second data set additionally includes the SH0ES measurement of H_0 [11] from the local distance ladder method (third column).

	Base	Base+SH0ES
f_{EDE}	0.0872	0.1271
$\log_{10} z_c$	3.560	3.563
θ_i	2.749	2.768
$100 \omega_b$	2.265	2.278
ω_{CDM}	0.1282	0.1324
$100 \theta_s$	1.041	1.041
$\ln(10^{10} A_s)$	3.063	3.071
n_s	0.983	0.992
τ	0.0562	0.0568

TABLE I. Best-fitting cosmological parameters under the *Planck* + BOSS (base) and base + SH0ES data sets [26].




FIG. 1. Stacked observed EB power spectrum (black points with error bars), compared to the best-fitting models of $\alpha + \beta$ (red) and g for two EDE models with parameters shown in Table I (blue and green). The χ^2 is 65.8, 77.5, and 103.5 for the red, blue, and green lines, respectively, for 71 degrees of freedom.

III. RESULTS

Before performing a simultaneous inference for $\alpha + \beta$ and g , we first fit the stacked EB power spectrum for each of these parameters alone. The black points with error bars in Fig. 1 show the stacked EB power spectrum for the nearly full-sky data. The red shaded area shows the 1σ band from the $\alpha + \beta$ fit, while the blue and green areas show those from g for the two EDE parameter sets. The shapes of the best-fitting EB power spectra are different for all cases, indicating that we can distinguish them easily. We find that $\alpha + \beta$ fits the data better, especially at the acoustic peak around $\ell \simeq 400$, which is reflected in a lower χ^2 value for this model (see the caption of Fig. 1).

We now present results from jointly sampling $\alpha + \beta$ and g . In Fig. 2, we show the posterior distributions of $\alpha + \beta$ and g for the two EDE parameter sets. To show ro-

² <https://github.com/LilleJohs/Observed-EB-Power-Spectrum/>

³ <https://github.com/cmbant/CAMB>




FIG. 2. Posterior distributions of g/M_{Pl}^{-1} and $\alpha + \beta$ for the best-fitting EDE parameters under the base and base+SH0ES data sets, and two Galactic masks.

bustness of the results against the choice of the Galactic mask, the results are shown for both small ($f_{\text{sky}} = 0.92$) and large ($f_{\text{sky}} = 0.62$) masks. We find that all combinations yield similar results.

As also examined in Ref. [38], we find similar stacked EB power spectra for all sky fractions between $f_{\text{sky}} = 0.92$ and 0.62 , suggesting that the Galactic foreground emission does not contribute significantly to the stacked EB power spectrum. This is because foreground-dominated channels such as 353 GHz have large \mathbf{M}_b and are downweighted in the sum given in Eq. (4). The foreground EB power spectrum plays a role only when used to calibrate α [50]. As we do not separate α and β in this paper but marginalize over $\alpha + \beta$, our results are insensitive to the Galactic foreground emission.

We choose the nearly full-sky result with the base EDE parameters as our baseline result, whose quality of the fit is shown in Fig. 3. We report $\alpha + \beta = 0.27^\circ \pm 0.08^\circ$ and $g/M_{\text{Pl}}^{-1} = 0.04 \pm 0.16$ (68% C.L.). The former agrees well with those reported in the literature [38, 46, 49–51, 66], while the latter is a first constraint on g for the EDE model with $V(\phi) = V_0[1 - \cos(\phi/f)]^3$. The data strongly favor $\alpha + \beta$ over g from EDE.

We also check whether the choice of the EDE parameters affects the results. We repeat the analysis for the best-fitting cosmological parameters for f_{EDE} fixed to a grid of values between 0 to 0.15 reported in Ref. [26]. We find that the minimum χ^2 for the stacked EB power spectrum varies less than $\Delta\chi^2 < 0.1$ when sampling $\alpha + \beta$ and g jointly, and obtain similar constraints on the parameters regardless of the value of f_{EDE} . This is because




FIG. 3. Quality of the fit for the baseline result. The red area shows the 1σ band of the $\alpha + \beta$ term, while the blue area shows that of the EDE term given in Eq. (8).

the shape of the observed EB power spectrum is well described by $\alpha + \beta$. This result confirms robustness of our baseline result.

IV. CONCLUSIONS

We presented a first constraint on the photon-axion coupling constant for the EDE model. Thanks to the unique shape of the predicted EB power spectrum, we were able to derive a constraint on $g/M_{\text{Pl}}^{-1} = 0.04 \pm 0.16$ (68% C.L.) independent of the miscalibration angle or the Galactic foreground emission. We find that the *Planck* data do not favor cosmic birefringence caused by a coupling to canonical pre-recombination EDE, g , but favor cosmic birefringence that occurred after the epoch of recombination, β , or a miscalibration of polarization angles of the *Planck* detectors, α .

The derived value of g is much weaker than the strength of the gravitational interaction, $|g| \ll M_{\text{Pl}}^{-1}$. This may have important implications for embedding EDE in fundamental physics, such as string theory. While the previous attempts [73–75] have not yet studied a Chern-Simons coupling for the EDE field in string theory, it is conceivable that such a term appears.

In particular, Ref. [75] finds that the best-fitting axion decay constant, $f \simeq 0.2 M_{\text{Pl}}$, requires fine-tuning of the microscopic parameters to respect the weak gravity conjecture [76], which states that any gauge force must mediate interactions stronger than gravity for some particles [77]. Similarly, $|g| \ll M_{\text{Pl}}^{-1}$ may also be in tension with a requirement for a consistent theory of quantum gravity including the weak gravity conjecture. Whether such a constraint on g exists remains to be studied. See Refs. [78, 79] for some discussion on the weak gravity conjecture in the presence of a Chern-Simons coupling.

The EDE parameters given in Table I yield $f = 0.15 M_{\text{Pl}}$ and $0.18 M_{\text{Pl}}$ for the base and base+SH0ES parameters, respectively. If we take $g = c_{\phi\gamma}\alpha_{\text{em}}/(2\pi f)$ with $f = 0.15 M_{\text{Pl}}$, our constraint on g yields $c_{\phi\gamma} = 5.2 \pm 21$ (68% C.L.). If the weak gravity conjecture demands $|g| \gtrsim M_{\text{Pl}}^{-1}$, we find $|c_{\phi\gamma}| \gtrsim 130$, which may be too large for an anomaly coefficient, although a natural value for $c_{\phi\gamma}$ depends on the precise mechanism by which the Chern-Simons term arises. In any case such a large anomaly coefficient is ruled out by our measurement.

Our work opens up a new research area in the EB power spectrum analysis. While we have focused on a particular EDE model with $V(\phi) = V_0[1 - \cos(\phi/f)]^n$ for $n = 3$, the same analysis can be repeated for any other model, e.g., $n = 2$ [19]. As our approach is independent of the miscalibration angle or the Galactic foreground emission, it can be applied to both current and future CMB experiments [80–84].

ACKNOWLEDGMENTS

We thank K. Kamada, T. Noumi, I. Obata, M. Scalisi, and M. Shiraishi for useful discussions. This work was supported in part by the European Research Council (ERC) under the Horizon 2020 Research and Innovation Programme (Grant agreement No. 819478), JSPS

KAKENHI Grants No. JP20H05850, No. JP20H05859, No. JP20J20248 and No. JP22K03682, the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - EXC-2094 - 390783311, the Program of Excellence in Photon Science, and the Forefront Physics and Mathematics Program to Drive Transformation (FoPM), a World-leading Innovative Graduate Study (WINGS) Program, the University of Tokyo. This work has also received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 101007633. The Kavli IPMU is supported by World Premier International Research Center Initiative (WPI), MEXT, Japan. We acknowledge the use of the Legacy Archive for Microwave Background Data Analysis (LAMBDA), part of the High Energy Astrophysics Science Archive Center (HEASARC). HEASARC/LAMBDA is a service of the Astrophysics Science Division at the NASA Goddard Space Flight Center. *Planck* is a project of the European Space Agency (ESA) with instruments provided by two scientific consortia funded by ESA member states and led by Principal Investigators from France and Italy, telescope reflectors provided through a collaboration between ESA and a scientific consortium led and funded by Denmark, and additional contributions from NASA (USA). Some of the results in this paper have been derived using the HEALPix package [65] and the GetDist package [85].

-
- [1] S. Weinberg, *Cosmology* (Oxford University Press, New York, 2008).
- [2] E. Abdalla *et al.*, *J. High Energy Astrophys.* **34**, 49 (2022), arXiv:2203.06142 [astro-ph.CO].
- [3] M. Kamionkowski and A. G. Riess, arXiv preprint (2022), arXiv:2211.04492 [astro-ph.CO].
- [4] V. Poulin, T. L. Smith, and T. Karwal, arXiv preprint (2023), arXiv:2302.09032 [astro-ph.CO].
- [5] E. Komatsu, *Nat. Rev. Phys.* **4**, 452 (2022), arXiv:2202.13919 [astro-ph.CO].
- [6] M. Doran, M. J. Lilley, J. Schwindt, and C. Wetterich, *Astrophys. J.* **559**, 501 (2001), arXiv:astro-ph/0012139.
- [7] C. Wetterich, *Phys. Lett. B* **594**, 17 (2004), arXiv:astro-ph/0403289.
- [8] M. Doran and G. Robbers, *J. Cosmol. Astropart. Phys.* **06** (2006), 026, arXiv:astro-ph/0601544.
- [9] T. Karwal and M. Kamionkowski, *Phys. Rev. D* **94**, 103523 (2016), arXiv:1608.01309 [astro-ph.CO].
- [10] V. Poulin, T. L. Smith, T. Karwal, and M. Kamionkowski, *Phys. Rev. Lett.* **122**, 221301 (2019), arXiv:1811.04083 [astro-ph.CO].
- [11] A. G. Riess *et al.*, *Astrophys. J. Lett.* **934**, L7 (2022), arXiv:2112.04510 [astro-ph.CO].
- [12] W.-T. Ni, *Phys. Rev. Lett.* **38**, 301 (1977).
- [13] M. S. Turner and L. M. Widrow, *Phys. Rev. D* **37**, 2743 (1988).
- [14] D. J. E. Marsh, *Phys. Rep.* **643**, 1 (2016), arXiv:1510.07633 [astro-ph.CO].
- [15] L. M. Capparelli, R. R. Caldwell, and A. Melchiorri, *Phys. Rev. D* **101**, 123529 (2020), arXiv:1909.04621 [astro-ph.CO].
- [16] T. Fujita, K. Murai, H. Nakatsuka, and S. Tsujikawa, *Phys. Rev. D* **103**, 043509 (2021), arXiv:2011.11894 [astro-ph.CO].
- [17] S. Nakagawa, F. Takahashi, and W. Yin, *Phys. Rev. D* **107**, 063016 (2023), arXiv:2209.01107 [astro-ph.CO].
- [18] K. Murai, F. Naokawa, T. Namikawa, and E. Komatsu, *Phys. Rev. D* **107**, L041302 (2023), arXiv:2209.07804 [astro-ph.CO].
- [19] M. Galaverni, F. Finelli, and D. Paoletti, arXiv preprint (2023), arXiv:2301.07971 [astro-ph.CO].
- [20] T. L. Smith, V. Poulin, and M. A. Amin, *Phys. Rev. D* **101**, 063523 (2020), arXiv:1908.06995 [astro-ph.CO].
- [21] T. L. Smith, V. Poulin, J. L. Bernal, K. K. Boddy, M. Kamionkowski, and R. Murgia, *Phys. Rev. D* **103**, 123542 (2021), arXiv:2009.10740 [astro-ph.CO].
- [22] R. Murgia, G. F. Abellán, and V. Poulin, *Phys. Rev. D* **103**, 063502 (2021), arXiv:2009.10733 [astro-ph.CO].
- [23] T. L. Smith, M. Lucca, V. Poulin, G. F. Abellan, L. Balkenhol, K. Benabed, S. Galli, and R. Murgia, *Phys. Rev. D* **106**, 043526 (2022), arXiv:2202.09379 [astro-ph.CO].
- [24] L. Herold, E. G. M. Ferreira, and E. Komatsu, *Astrophys. J. Lett.* **929**, L16 (2022), arXiv:2112.12140 [astro-ph.CO].
- [25] T. Simon, P. Zhang, V. Poulin, and T. L. Smith, *Phys. Rev. D* **107**, 063505 (2023), arXiv:2208.05930 [astro-ph.CO].

- [26] L. Herold and E. G. M. Ferreira, arXiv preprint (2022), arXiv:2210.16296 [astro-ph.CO].
- [27] J. C. Hill, E. McDonough, M. W. Toomey, and S. Alexander, Phys. Rev. D **102**, 043507 (2020), arXiv:2003.07355 [astro-ph.CO].
- [28] M. M. Ivanov, E. McDonough, J. C. Hill, M. Simonović, M. W. Toomey, S. Alexander, and M. Zaldarriaga, Phys. Rev. D **102**, 103502 (2020), arXiv:2006.11235 [astro-ph.CO].
- [29] G. D'Amico, L. Senatore, P. Zhang, and H. Zheng, J. Cosmol. Astropart. Phys. **05** (2021), 072, arXiv:2006.12420 [astro-ph.CO].
- [30] J. C. Hill *et al.*, Phys. Rev. D **105**, 123536 (2022), arXiv:2109.04451 [astro-ph.CO].
- [31] A. La Posta, T. Louis, X. Garrido, and J. C. Hill, Phys. Rev. D **105**, 083519 (2022), arXiv:2112.10754 [astro-ph.CO].
- [32] A. Reeves, L. Herold, S. Vagnozzi, B. D. Sherwin, and E. G. M. Ferreira, Mon. Not. Roy. Astron. Soc. **520**, 3688 (2023), arXiv:2207.01501 [astro-ph.CO].
- [33] J. S. Cruz, S. Hannestad, E. B. Holm, F. Niedermann, M. S. Sloth, and T. Tram, arXiv preprint (2023), arXiv:2302.07934 [astro-ph.CO].
- [34] S. Goldstein, J. C. Hill, V. Iršič, and B. D. Sherwin, arXiv preprint (2023), arXiv:2303.00746 [astro-ph.CO].
- [35] M. Zaldarriaga and U. Seljak, Phys. Rev. D **55**, 1830 (1997), arXiv:astro-ph/9609170.
- [36] M. Kamionkowski, A. Kosowsky, and A. Stebbins, Phys. Rev. D **55**, 7368 (1997), arXiv:astro-ph/9611125.
- [37] A. Lue, L.-M. Wang, and M. Kamionkowski, Phys. Rev. Lett. **83**, 1506 (1999), arXiv:astro-ph/9812088.
- [38] J. R. Eskilt and E. Komatsu, Phys. Rev. D **106**, 063503 (2022), arXiv:2205.13962 [astro-ph.CO].
- [39] S. M. Carroll, G. B. Field, and R. Jackiw, Phys. Rev. D **41**, 1231 (1990).
- [40] S. M. Carroll and G. B. Field, Phys. Rev. D **43**, 3789 (1991).
- [41] D. Harari and P. Sikivie, Phys. Lett. B **289**, 67 (1992).
- [42] D. Contreras, P. Boubel, and D. Scott, J. Cosmol. Astropart. Phys. **12** (2017), 046, arXiv:1705.06387 [astro-ph.CO].
- [43] T. Namikawa *et al.* (ACT Collaboration), Phys. Rev. D **101**, 083527 (2020), arXiv:2001.10465 [astro-ph.CO].
- [44] F. Bianchini *et al.* (SPT Collaboration), Phys. Rev. D **102**, 083504 (2020), arXiv:2006.08061 [astro-ph.CO].
- [45] A. Gruppuso, D. Molinari, P. Natoli, and L. Pagano, J. Cosmol. Astropart. Phys. **11** (2020), 066, arXiv:2008.10334 [astro-ph.CO].
- [46] M. Bortolami, M. Billi, A. Gruppuso, P. Natoli, and L. Pagano, J. Cosmol. Astropart. Phys. **09** (2022), 075, arXiv:2206.01635 [astro-ph.CO].
- [47] H. Nakatsuka, T. Namikawa, and E. Komatsu, Phys. Rev. D **105**, 123509 (2022), arXiv:2203.08560 [astro-ph.CO].
- [48] B. Feng, H. Li, M. Li, and X. Zhang, Phys. Lett. B **620**, 27 (2005), arXiv:hep-ph/0406269 [hep-ph].
- [49] Y. Minami and E. Komatsu, Phys. Rev. Lett. **125**, 221301 (2020), arXiv:2011.11254 [astro-ph.CO].
- [50] P. Diego-Palazuelos *et al.*, Phys. Rev. Lett. **128**, 091302 (2022), arXiv:2201.07682 [astro-ph.CO].
- [51] J. R. Eskilt, Astron. Astrophys. **662**, A10 (2022), arXiv:2201.13347 [astro-ph.CO].
- [52] G.-C. Liu, S. Lee, and K.-W. Ng, Phys. Rev. Lett. **97**, 161303 (2006), arXiv:astro-ph/0606248 [astro-ph].
- [53] F. Finelli and M. Galaverni, Phys. Rev. D **79**, 063002 (2009), arXiv:0802.4210 [astro-ph].
- [54] E. Y. S. Wu *et al.* (QUaD Collaboration), Phys. Rev. Lett. **102**, 161302 (2009), arXiv:0811.0618 [astro-ph].
- [55] N. J. Miller, M. Shimon, and B. G. Keating, Phys. Rev. D **79**, 103002 (2009), arXiv:0903.1116 [astro-ph.CO].
- [56] E. Komatsu *et al.* (WMAP Collaboration), Astrophys. J. Suppl. **192**, 18 (2011), arXiv:1001.4538 [astro-ph.CO].
- [57] N. Krachmalnicoff *et al.* (LiteBIRD Collaboration), J. Cosmol. Astropart. Phys. **01** (2022), 039, arXiv:2111.09140 [astro-ph.CO].
- [58] J. Cornelison *et al.*, Proc. SPIE Int. Soc. Opt. Eng. **12190**, 121901X (2022), arXiv:2207.14796 [astro-ph.IM].
- [59] Y. Minami, H. Ochi, K. Ichiki, N. Katayama, E. Komatsu, and T. Matsumura, Prog. Theor. Exp. Phys. **2019**, 083E02 (2019), arXiv:1904.12440 [astro-ph.CO].
- [60] B. D. Sherwin and T. Namikawa, Mon. Not. Roy. Astron. Soc. **520**, 3298 (2023), arXiv:2108.09287 [astro-ph.CO].
- [61] P. Diego-Palazuelos *et al.*, J. Cosmol. Astropart. Phys. **01** (2023), 044, arXiv:2210.07655 [astro-ph.CO].
- [62] Planck Collaboration III, Astron. Astrophys. **641**, A3 (2020), arXiv:1807.06207.
- [63] Planck Collaboration Int. LVII, Astron. Astrophys. **643**, A42 (2020), arXiv:2007.04997.
- [64] G. Chon, A. Challinor, S. Prunet, E. Hivon, and I. Szapudi, Mon. Not. Roy. Astron. Soc. **350**, 914 (2004), arXiv:astro-ph/0303414.
- [65] K. M. Górski, E. Hivon, A. J. Banday, B. D. Wandelt, F. K. Hansen, M. Reinecke, and M. Bartelman, Astrophys. J. **622**, 759 (2005), arXiv:astro-ph/0409513.
- [66] Planck Collaboration Int. XLIX, Astron. Astrophys. **596**, A110 (2016), arXiv:1605.08633.
- [67] D. Foreman-Mackey, D. W. Hogg, D. Lang, and J. Goodman, Publ. Astron. Soc. Pac. **125**, 306 (2013), arXiv:1202.3665 [astro-ph.IM].
- [68] A. Lewis, A. Challinor, and A. Lasenby, Astrophys. J. **538**, 473 (2000), astro-ph/9911177.
- [69] Planck Collaboration VI, Astron. Astrophys. **641**, A6 (2020), arXiv:1807.06209.
- [70] D. Blas, J. Lesgourgues, and T. Tram, J. Cosmol. Astropart. Phys. **07** (2011), 034, arXiv:1104.2933 [astro-ph.CO].
- [71] Planck Collaboration V, Astron. Astrophys. **641**, A5 (2020), arXiv:1907.12875.
- [72] S. Alam *et al.* (BOSS Collaboration), Mon. Not. Roy. Astron. Soc. **470**, 2617 (2017), arXiv:1607.03155 [astro-ph.CO].
- [73] T. Rudelius, J. Cosmol. Astropart. Phys. **01** (2023), 014, arXiv:2203.05575 [hep-th].
- [74] E. McDonough and M. Scalisi, arXiv preprint (2022), arXiv:2209.00011 [hep-th].
- [75] M. Cicoli, M. Licheri, R. Mahanta, E. McDonough, F. G. Pedro, and M. Scalisi, arXiv preprint (2023), arXiv:2303.03414 [hep-th].
- [76] N. Arkani-Hamed, L. Motl, A. Nicolis, and C. Vafa, J. High Energy Phys. **06** (2007), 060, arXiv:hep-th/0601001.
- [77] D. Harlow, B. Heidenreich, M. Reece, and T. Rudelius, arXiv preprint (2022), arXiv:2201.08380 [hep-th].
- [78] B. Heidenreich, M. Reece, and T. Rudelius, J. High Energy Phys. **11** (2021), 004, arXiv:2108.11383 [hep-th].
- [79] S. Kaya and T. Rudelius, J. High Energy Phys. **07** (2022), 040, arXiv:2202.04655 [hep-th].
- [80] P. Ade *et al.* (Simons Observatory Collaboration), J. Cos-

- mol. Astropart. Phys. **02** (2019), 056, arXiv:1808.07445 [astro-ph.CO].
- [81] K. Abazajian *et al.* (CMB-S4 Collaboration), arXiv preprint (2019), arXiv:1907.04473 [astro-ph.IM].
- [82] L. Moncelsi *et al.*, Proc. SPIE Int. Soc. Opt. Eng. **11453**, 1145314 (2020), arXiv:2012.04047 [astro-ph.IM].
- [83] S. Hanany *et al.* (PICO Collaboration), arXiv preprint (2019), arXiv:1902.10541 [astro-ph.IM].
- [84] LiteBIRD Collaboration, arXiv preprint (2022), arXiv:2202.02773 [astro-ph.CO].
- [85] A. Lewis, arXiv preprint (2019), arXiv:1910.13970 [astro-ph.IM].