

Shapes of magnetic monopoles in effective $SU(2)$ models

Petr Beneš^{1,*} and Filip Blaschke^{1,2,†}

¹*Institute of Experimental and Applied Physics,*

Czech Technical University in Prague, Husova 240/5, 110 00 Prague 1, Czech Republic

²*Research Centre for Theoretical Physics and Astrophysics, Institute of Physics,*

Silesian University in Opava, Bezručovo náměstí 1150/13, 746 01 Opava, Czech Republic

We present a systematic exploration of a general family of effective $SU(2)$ models with an adjoint scalar. First, we discuss a redundancy in this class of models and use it to identify seemingly different, yet physically equivalent models. Next, we construct the Bogomol'nyi–Prasad–Sommerfield (BPS) limit and derive analytic monopole solutions. In contrast to the 't Hooft–Polyakov monopole, included here as a special case, these solutions tend to exhibit more complex energy density profiles. Typically, we obtain monopoles with a hollow cavity at their core where virtually no energy is concentrated; instead, most of the monopole's energy is stored in a spherical shell around its core. Moreover, the shell itself can be structured, with several “sub-shells”. A recipe for the construction of these analytic solutions is presented.

For Jiří Hošek on his 80th birthday.

Keywords: Magnetic monopole; electroweak model; exact solutions; BPS limit

I. INTRODUCTION

Magnetic monopoles are amongst the most important hypothetical particles in contemporary physics. Although they still await experimental detection, there are very strong theoretical reasons for their existence. Monopoles play a major rôle in such disparate subjects as cosmology, particle physics, and condensed matter physics. Their detection would constitute a fundamental breakthrough as they would provide a completely new window to high-energy physics.

In [1] we have introduced a generalization of Lee–Weinberg's conceptual scheme [2] and presented a landscape of $U(1)$ gauge theories with a scalar field ϕ and massive, complex vector fields W_μ , namely:

$$\begin{aligned} \mathcal{L}_{U(1)} = & -\frac{f_1^2(\phi/v)}{4g^2} F_{\mu\nu}^2 - \frac{\eta(\phi/v)}{2} d_{\mu\nu} F^{\mu\nu} - \frac{\chi(\phi/v)g^2}{4} d_{\mu\nu}^2 \\ & - \frac{f_2^2(\phi/v)}{2} |D_\mu W_\nu - D_\nu W_\mu|^2 + m^2(\phi/v) |W_\mu|^2 \\ & + \frac{1}{2} (\partial_\mu \phi)^2 - \frac{\lambda}{2} (\phi^2 - v^2)^2, \end{aligned} \quad (1)$$

where $d_{\mu\nu} = i(W_\mu^\dagger W_\nu - W_\nu^\dagger W_\mu)$ is a dipole-moment tensor. These models represent the most general effective field-theoretical descriptions of a Dirac monopole possessing a finite mass. The rôle of form-functions $f_{1,2}(\phi/v)$, $\eta(\phi/v)$, $\chi(\phi/v)$ and $m(\phi/v)$ is to provide control over the properties of the “substrate” in which is the monopole is placed (ϕ/v being the collective coordinate of the coherent state). In particular:

- f_1 is as field-dependent magnetic susceptibility,

- f_2 describes field-dependent non-linear elastic properties of W_μ fields,
- η and χ control the dipole-moment interactions of W_μ ,
- m^2 is a field-dependent mass.

Alternatively, we can think about (1) as an effective action of a more fundamental theory with partially resummed loop diagrams that are embodied in these form-functions. In short, we believe that Eq. (1) is something that one could typically encounter down the stream of the renormalization group river starting from an unknown, more fundamental theory of magnetic monopoles.¹

The utility of considering such a vast landscape is the following: It is perhaps intuitively clear that properties of classical solutions would not be qualitatively different for all choices of form-functions. Instead, they would fall into a small number of universality classes in which a modification of various field-dependent couplings would result in only quantitative difference, but not in a qualitative change in their properties and behavior. By identifying and studying such universality classes, we can understand something about monopoles even though we do not know the ultimate, correct theory, provided, of course, that the “real” monopole in Nature (if ever we found one) falls into one of these universality classes.

In this paper, we make a first step toward this goal and provide a systematic classification of spherically-symmetric solutions of a particular sub-landscape of (1) that is endowed with $SU(2)$ gauge symmetry. The advantages of this restriction are both practical and utilitarian. Practical, as we can focus our efforts more narrowly

* petr.benes@utef.cvut.cz

† filip.blaschke@fpf.slu.cz

¹ This viewpoint is strengthened by appearance of four-derivative terms in the canonical formulation of $SU(2)$ landscape in this paper.

and utilitarian, as we can compare our solutions with the canonical 't Hooft–Polyakov monopole [3, 4], which itself is a particular point of the general $SU(2)$ sub-landscape. Furthermore, we can avoid the troubles with residual singularities that are permanent in $U(1)$ models. However, as we will see, the discussion of whether the solutions must be always singularity-free will be one of the corollaries of this paper.

Magnetic monopoles in extended, exotic, and variously generalized models have been studied before [5–8]. One of the characteristics of spherically symmetric single monopole solutions that seems to be linked to form-functions is the shape of their energy density profiles. We will pay a special attention to them. As already reported by Bazeia et al. [8], monopoles can have unexpected shapes (which they dubbed “small” and “hollow”), in comparison with the 't Hooft–Polyakov’s solution. In particular, the appearance of a hollow cavity at the core of the monopole, where virtually no energy is stored, seems to be more typical than not. In this paper, we provide explicit analytic solutions for these hollow monopoles and we discuss the conditions under which the cavity appears and what controls its extent.

The paper is organized as follows. First, in Sec. II, we introduce a general class of $SU(2)$ models and discuss its form-invariance. Next, in Sec. III, we construct the Bogomol’nyi–Prasad–Sommerfield (BPS) limit and formulate the corresponding BPS equation of motion and energy density. In Sec. IV we specialize to spherical symmetry, i.e., to single-monopole solutions, and present a “recipe” for finding analytic solutions. In the subsequent Sec. V, we show explicit examples. Finally, in Sec. VI, we summarize our results and give an outlook to future directions of research.

II. GENERAL (NON-BPS) MODEL

A. The Lagrangian

We consider the most general $SU(2)$ gauge theory that is quadratic in $\mathbf{F}^{\mu\nu}$ and with scalar ϕ in adjoint (real triplet) representation with at most two derivatives. Accordingly, the Lagrangian must be a linear combination of four algebraically independent terms $(D^\mu\phi)^2$, $(\phi\cdot D^\mu\phi)^2$, $(\mathbf{F}^{\mu\nu})^2$ and $(\phi\cdot\mathbf{F}^{\mu\nu})^2$, with the coefficients being dimensionless and gauge-invariant functions of ϕ . We find most convenient to write these terms as

$$\begin{aligned} \mathcal{L} = & \frac{v^2}{2} \left[f_1^2 \left(\frac{(D^\mu\phi)^2}{\phi^2} - \frac{(\phi\cdot D^\mu\phi)^2}{\phi^4} \right) + f_3^2 \frac{(\phi\cdot D^\mu\phi)^2}{\phi^4} \right] \\ & - \frac{1}{4g^2} \left[f_2^2 \left((\mathbf{F}^{\mu\nu})^2 - \frac{(\phi\cdot\mathbf{F}^{\mu\nu})^2}{\phi^2} \right) + f_4^2 \frac{(\phi\cdot\mathbf{F}^{\mu\nu})^2}{\phi^2} \right] \\ & - V(\phi^2), \end{aligned} \quad (2)$$

where $f_{1,2,3,4}^2$ are the “form-functions” and where

$$D^\mu\phi = \partial^\mu\phi + \mathbf{A}^\mu \times \phi, \quad (3)$$

$$\mathbf{F}^{\mu\nu} = \partial^\mu\mathbf{A}^\nu - \partial^\nu\mathbf{A}^\mu + \mathbf{A}^\mu \times \mathbf{A}^\nu, \quad (4)$$

with the boldface denoting a three-vector in the gauge space. The potential $V(\phi^2) \geq 0$ need not be specified, besides the assumption that for $\phi^2 = v^2$ (where $v^2 > 0$) it vanishes, triggering the spontaneous symmetry breakdown $SU(2) \rightarrow U(1)$.

In the bulk of this paper, we utilize the decomposition

$$\phi = vH\mathbf{n}, \quad (5)$$

where the isovector \mathbf{n} is normalized as $\mathbf{n}^2 = 1$ and H is a dimensionless gauge-invariant scalar function, that serves as an argument of all form-functions:

$$f_i^2 \equiv f_i^2(H). \quad (i = 1, 2, 3, 4) \quad (6)$$

The link of the $SU(2)$ Lagrangian (2) to the $U(1)$ Lagrangian (1) is made by rewriting the former in the unitary gauge and matching the respective form-functions (keeping in mind that $f_{1,2}^2$ in both Lagrangians can be different). At this point, there is no need to make this link explicit and we do not display it here.

Note that the general model (2) includes the usual renormalizable Georgi–Glashow model as the special case $f_1^2 = f_3^2 = H^2$ and $f_2^2 = f_4^2 = 1$.

There are two reasons for writing the Lagrangian (2) with the basis of projector-like structures. The first one is that the condition for the energy density to be bounded from below is very simple: Each of the corresponding functions f_i^2 must be non-negative,

$$f_i^2 \geq 0, \quad (7)$$

as indicated by the square notation.

The second reason is that the form of the Lagrangian (2) is well suited for the formalism to be used later in this paper. E.g., using the decomposition (5) the first two terms in (2) (proportional to f_1^2 and f_3^2) can be respectively rewritten very simply as

$$\frac{(D^\mu\phi)^2}{\phi^2} - \frac{(\phi\cdot D^\mu\phi)^2}{\phi^4} = (D^\mu\mathbf{n})^2, \quad (8a)$$

$$\frac{(\phi\cdot D^\mu\phi)^2}{\phi^4} = \frac{(\partial^\mu H)^2}{H^2}. \quad (8b)$$

After introducing spherically symmetric Ansatz later on, similar simplification will occur also for the gauge part.

Finally, we will for convenience assume, without loss of generality,² the normalization

$$f_1^2(1) = f_2^2(1) = 1. \quad (9)$$

In this way, the kinetic terms are properly normalized in the vacuum $\phi^2 = v^2$.

The non-canonical model (2) has necessarily the same spectrum of fluctuations near the $SU(2)$ -breaking vacuum as does the Georgi–Glashow model. Namely, there

² This normalization can be always enforced by redefining $\phi \rightarrow \phi/f_1(1)$, $v \rightarrow v/|f_1(1)|$ and $g \rightarrow g|f_2(1)|$, which is equivalent to $f_{1,2} \rightarrow f_{1,2}/f_{1,2}(1)$.

is a massless gauge field (photon) corresponding to the unbroken $U(1)$ subgroup, a pair of massive charged vector fields (W^\pm), and a massive real scalar field (Higgs). The latter non-vanishing masses are given, due to the normalization (9), by the standard formulas $M_W = vg$ and $M_{\text{Higgs}} = 2v\sqrt{V''(v^2)}$.

B. Form-invariance and redundancy

Let us consider a transformation (field redefinition) of the scalar triplet ϕ to a new $\tilde{\phi}$ that only changes its length, namely

$$\phi = vH\mathbf{n} \quad \longrightarrow \quad \tilde{\phi} = v\tilde{H}\mathbf{n}, \quad (10a)$$

where \tilde{H} is related to H as

$$H = \alpha(\tilde{H}). \quad (10b)$$

Here $\alpha(\tilde{H})$ is an arbitrary invertible and differentiable function that satisfies $\alpha^2(\pm 1) = 1$ to preserve the vacuum value. Under this transformation, the original Lagrangian \mathcal{L} of Eq. (2) transforms into a new Lagrangian $\tilde{\mathcal{L}}$ of the form

$$\begin{aligned} \tilde{\mathcal{L}} = & \frac{v^2}{2} \left[\tilde{f}_1^2 \left(\frac{(D^\mu \tilde{\phi})^2}{\tilde{\phi}^2} - \frac{(\tilde{\phi} \cdot D^\mu \tilde{\phi})^2}{\tilde{\phi}^4} \right) + \tilde{f}_3^2 \frac{(\tilde{\phi} \cdot D^\mu \tilde{\phi})^2}{\tilde{\phi}^4} \right] \\ & - \frac{1}{4g^2} \left[\tilde{f}_2^2 \left((\mathbf{F}^{\mu\nu})^2 - \frac{(\tilde{\phi} \cdot \mathbf{F}^{\mu\nu})^2}{\tilde{\phi}^2} \right) + \tilde{f}_4^2 \frac{(\tilde{\phi} \cdot \mathbf{F}^{\mu\nu})^2}{\tilde{\phi}^2} \right] \\ & - \tilde{V}(\tilde{\phi}^2), \end{aligned} \quad (11)$$

where $\tilde{V}(\tilde{\phi}^2) = V(\phi^2)$ and where the new functions \tilde{f}_i^2 are given in terms of the old functions f_i^2 as

$$\tilde{f}_i^2(\tilde{H}) = \begin{cases} f_i^2(\alpha(\tilde{H})), & (i = 1, 2, 4) \\ f_i^2(\alpha(\tilde{H})) \left(\frac{\tilde{H} \alpha'(\tilde{H})}{\alpha(\tilde{H})} \right)^2, & (i = 3) \end{cases} \quad (12)$$

The non-trivial transformation law for \tilde{f}_3^2 is due to the fact that only this term contains a derivative of H , as can be seen from (8).

There is a special worth noting case of the transformation (10): $H = 1/\tilde{H}$. Obviously, when this transformation is applied twice, we obtain the original theory, hence we can call it a *duality*. Moreover, since $\tilde{H}\alpha'/\alpha = -1$, all four \tilde{f}_i^2 transform uniformly as $\tilde{f}_i^2(\tilde{H}) = f_i^2(1/\tilde{H})$.

Notably, the transformed Lagrangian $\tilde{\mathcal{L}}$ is of the same form as \mathcal{L} . We can thus say that \mathcal{L} is *form-invariant* under the transformation (10) in the sense that (10) does not introduce a new *kind* of term that was absent beforehand. Moreover, both \mathcal{L} and $\tilde{\mathcal{L}}$ describe the same physics. This means, in particular, that while the corresponding solutions can differ, they must both yield the same energy density. Consequently, of the four seemingly independent functions f_i^2 only three are (physically) independent, since the other one can be fixed by (12).

The form-invariance of the theory implies that a singularity in H is not necessarily an issue, as it may be cured by a suitable transformation (10). We will showcase a particular example of this in Sec. V A.

C. Dressed vs. canonical fields

As mentioned, we can utilize the transformation (10) to make one of the four functions \tilde{f}_i^2 to attain a desired form. Although we shall not adopt any single one, let us point out a few choices of interest.

If, for instance, we want the kinetic term of the scalar triplet ϕ to be canonically normalized, we should demand $\tilde{f}_1^2 = H^2$. Another, slightly less obvious choice, is to demand $\tilde{f}_3^2 = H^2$, which makes the kinetic term for the scalar singlet vH canonical.

To achieve canonical normalization of gauge fields, however, the transformation (10) is not enough, as it cannot arrange $\tilde{f}_2^2 = 1$. In fact, to make $\tilde{f}_2^2 = 1$ we need to perform a *field-redefinition of the gauge fields* as³

$$\mathbf{A}^\mu = \tilde{\mathbf{A}}^\mu + \left(\frac{1}{f_2} - 1 \right) \frac{\phi \times \tilde{D}^\mu \phi}{\phi^2}, \quad (13)$$

where $\tilde{D}^\mu \phi = \partial^\mu \phi + \tilde{\mathbf{A}}^\mu \times \phi$ is a covariant derivative with respect to the transformed gauge field $\tilde{\mathbf{A}}^\mu$. Accordingly, the Lagrangian (2) turns into

$$\begin{aligned} \tilde{\mathcal{L}} = & \frac{v^2}{2} \left[\frac{f_1^2}{f_2^2} \left(\frac{(\tilde{D}^\mu \phi)^2}{\phi^2} - \frac{(\phi \cdot \tilde{D}^\mu \phi)^2}{\phi^4} \right) + f_3^2 \frac{(\phi \cdot \tilde{D}^\mu \phi)^2}{\phi^4} \right] \\ & - \frac{1}{4g^2} \left[(\mathbf{F}^{\mu\nu})^2 + (f_4^2 - 1) \frac{(\phi \cdot \mathbf{F}^{\mu\nu})^2}{\phi^2} \right] - V(\phi^2) \\ & - \frac{1}{2g^2} \mathbf{d}^{\mu\nu} \cdot \left\{ f_2 \left[\tilde{\mathbf{F}}_{\mu\nu} - \frac{(\phi \cdot \tilde{\mathbf{F}}_{\mu\nu})}{\phi^2} \phi \right] + f_4^2 \frac{(\phi \cdot \tilde{\mathbf{F}}_{\mu\nu})}{\phi^2} \phi \right\} \\ & - \frac{1}{4g^2} \left[f_2^2 \left((\mathbf{d}_{\mu\nu})^2 - \frac{(\phi \cdot \mathbf{d}_{\mu\nu})^2}{\phi^2} \right) + f_4^2 \frac{(\phi \cdot \mathbf{d}_{\mu\nu})^2}{\phi^2} \right], \end{aligned} \quad (14)$$

where we denoted (here, the prime represents derivative with respect to H)

$$\begin{aligned} \mathbf{d}_{\mu\nu} \equiv & H \left(\frac{1}{f_2} \right)' \frac{(\phi \cdot \tilde{D}_\mu \phi)(\phi \times \tilde{D}_\nu \phi) - (\mu \leftrightarrow \nu)}{\phi^4} \\ & - 2 \left(\frac{1}{f_2} - 1 \right) \frac{(\phi \cdot \tilde{D}_\mu \phi)(\phi \times \tilde{D}_\nu \phi) - (\mu \leftrightarrow \nu)}{\phi^4} \\ & + 2 \left(\frac{1}{f_2} - 1 \right) \frac{\tilde{D}_\mu \phi \times \tilde{D}_\nu \phi}{\phi^2} \\ & + \left(\frac{1}{f_2} - 1 \right)^2 \frac{\phi \cdot (\tilde{D}_\mu \phi \times \tilde{D}_\nu \phi)}{\phi^4} \phi. \end{aligned} \quad (15)$$

³ We present a generalization of this transformation, also in combination with the transformation (10), in Appendix A.

Indeed, the transformation (13) did the trick: The gauge kinetic term in (14) is canonically normalized. Notice, however, that (13) is not form-invariant, and we pay a price of introducing additional interaction terms (those with $\mathbf{d}_{\mu\nu}$) that are of order $\sim \mathcal{O}((D\phi)^4)$.

However, the appearance of these new terms in (14) has a clear physical interpretation. In the original Lagrangian (2), the form-function f_2^2 models a non-linear response of the substrate (described by the scalar triplet ϕ) to the presence of $SU(2)$ fields. By transforming from the *dressed* fields \mathbf{A}_μ to new *canonical* fields $\tilde{\mathbf{A}}_\mu$, this effect does not disappear, but rather manifests itself through the appearance of the dipole-moment tensor interactions.

These interactions, being higher-order in derivatives and non-renormalizable, are normally omitted from the classical action of Yang–Mill–Higgs theory. However, we can view them as effective, partial resummation of quantum corrections at all orders that are modelled here classically via the function f_2^2 . In other words, the Lagrangian (14) represents a semi-classical description of theory with charged bosons that have non-trivial electric and magnetic moments.

Since the *canonical* Lagrangian (14) is far more complicated than the original *dressed* Lagrangian (2), we will use the “dressed” formalism throughout the paper unless stated otherwise. However, the exact correspondence between both Lagrangians shows that the original theory (2) is nothing but canonical $SU(2)$ theory with explicit dipole-moment interactions.

III. BPS MODEL

A. Derivation

Let us consider the energy density of a static configuration ($\partial^0 = \mathbf{A}^0 = 0$) of (2):

$$\begin{aligned} \mathcal{E} = \frac{v^2}{2} & \left[f_1^2 \frac{(D^i \phi)^2}{\phi^2} + (f_3^2 - f_1^2) \frac{(\phi \cdot D^i \phi)^2}{\phi^2} \right] \\ & + \frac{1}{2g^2} \left[f_2^2 (\mathbf{B}^i)^2 + (f_4^2 - f_2^2) \frac{(\phi \cdot \mathbf{B}^i)^2}{\phi^2} \right] + V(\phi^2), \end{aligned} \quad (16)$$

where we defined $\mathbf{B}^i \equiv -\frac{1}{2}\epsilon^{ijk} \mathbf{F}^{jk}$. This can be rewritten identically as

$$\begin{aligned} \mathcal{E} = \frac{1}{2} & \left[\frac{f_1}{H} D^i \phi - \frac{f_2}{g} \mathbf{B}^i + x \frac{(\phi \cdot \mathbf{B}^i)}{g v^2 H^2} \phi - y \frac{(\phi \cdot D^i \phi)}{v^2 H^3} \phi \right]^2 \\ & + \frac{f_1 f_2}{gH} \partial^i (\phi \cdot \mathbf{B}^i) - \frac{x f_1 + y f_2 - xy}{gH^2} (\partial^i H) (\phi \cdot \mathbf{B}^i) \\ & + \frac{f_4^2 - (x - f_2)^2}{2g^2 v^2 H^2} (\phi \cdot \mathbf{B}^i)^2 + \frac{f_3^2 - (y - f_1)^2}{2v^2 H^4} (\phi \cdot D^i \phi)^2 \\ & + V(\phi^2), \end{aligned} \quad (17)$$

where x, y are arbitrary functions of H . We used $D^i \mathbf{B}^i = 0$ in order to write $(D^i \phi) \cdot \mathbf{B}^i = \partial^i (\phi \cdot \mathbf{B}^i)$.

We can now derive the BPS theory in three steps. First, we discard the potential, setting $V = 0$, but keeping the boundary condition $\phi^2 = v^2$. Next, we demand

$$x = f_2 + f_4, \quad y = f_1 + f_3, \quad (18)$$

so that the terms on the third line in Eq. (17) vanish. Finally, we require

$$\left(\frac{f_1 f_2}{gH} \right)' = -\frac{x f_1 + y f_2 - xy}{gH^2}, \quad (19)$$

as then the terms on the second line in (17) can be combined into a total derivative:

$$\begin{aligned} \frac{f_1 f_2}{gH} \partial^i (\phi \cdot \mathbf{B}^i) - \frac{x f_1 + y f_2 - xy}{gH^2} (\partial^i H) (\phi \cdot \mathbf{B}^i) \\ = \partial^i \left(\frac{f_1 f_2}{gH} \phi \cdot \mathbf{B}^i \right). \end{aligned} \quad (20)$$

Substituting the expressions (18) for x, y into (19), we obtain the equation

$$f_3 f_4 = H (f_1 f_2)', \quad (21)$$

i.e., a condition to be satisfied by the functions f_i to allow for a BPS theory in the limit of a vanishing potential.

B. Formulation

The coveted BPS condition (21) suggests that, for a BPS theory, it may be more convenient to use another parametrization. Namely, let us consider the set of functions $F_i(H)$ defined as

$$F_1 = f_1 f_2, \quad F_2 = f_1 / f_2, \quad (22a)$$

$$H F_3' = f_3 f_4, \quad H F_4' = f_3 / f_4, \quad (22b)$$

with the inverse relations

$$f_1^2 = F_1 F_2, \quad f_2^2 = F_1 / F_2, \quad (23a)$$

$$f_3^2 = H^2 F_3' F_4', \quad f_4^2 = F_3' / F_4'. \quad (23b)$$

The functions F_i are defined in terms of f_i uniquely up to a sign, therefore the conditions $f_i^2 \geq 0$ for the stability of the theory become

$$\text{sgn } F_1 = \text{sgn } F_2, \quad \text{sgn } F_3' = \text{sgn } F_4'. \quad (24)$$

Furthermore, we impose the normalization

$$|F_i(1)| = 1, \quad (i = 1, 2, 3, 4) \quad (25)$$

which, for $F_{1,2}$, is equivalent to $f_{1,2}^2(1) = 1$, while for $F_{3,4}$ we utilized the freedom to choose the constant of integration in $F_{3,4} = \int F_{3,4}' dH$ any way we like. Finally, let us note that in terms of F_i 's the transformation (10) attains a simpler form (compared with (12)):

$$\tilde{F}_i(\tilde{H}) = F_i(\alpha(\tilde{H})). \quad (i = 1, 2, 3, 4) \quad (26)$$

The Lagrangian (2) (with $V = 0$) can be equivalently expressed as

$$\mathcal{L} = \frac{v^2}{2} \left[F_1 F_2 \frac{(D^\mu \phi)^2}{\phi^2} + (H^2 F_3' F_4' - F_1 F_2) \frac{(\phi \cdot D^\mu \phi)^2}{\phi^4} \right] - \frac{1}{4g^2} \left[\frac{F_1}{F_2} (\mathbf{F}^{\mu\nu})^2 + \left(\frac{F_3'}{F_4'} - \frac{F_1}{F_2} \right) \frac{(\phi \cdot \mathbf{F}^{\mu\nu})^2}{\phi^2} \right]. \quad (27)$$

Notice that now the renormalizable Georgi–Glashow model corresponds to a special (and rather symmetric) case $F_1 = F_2 = F_3 = F_4 = H$.

The main benefit of the new parametrization is that the BPS condition (21) acquires a very simple form

$$F_3 = F_1, \quad (28)$$

as well as the BPS equation itself:⁴

$$D^i \phi = \frac{H}{g} \left[\frac{1}{F_2} \left(\mathbf{B}^i - \frac{\phi \cdot \mathbf{B}^i}{\phi^2} \phi \right) + \frac{1}{H F_4'} \frac{\phi \cdot \mathbf{B}^i}{\phi^2} \phi \right]. \quad (29)$$

When it is satisfied, the energy density is just a total derivative, as required:

$$\mathcal{E} = \partial^i \left(\frac{F_1}{gH} \phi \cdot \mathbf{B}^i \right), \quad (30)$$

but it can be also rewritten in two other equivalent ways as

$$\mathcal{E} = v^2 \left[F_1 F_2 \frac{(D^i \phi)^2}{\phi^2} + (H^2 F_1' F_4' - F_1 F_2) \frac{(\phi \cdot D^i \phi)^2}{\phi^4} \right] \quad (31a)$$

$$= \frac{1}{g^2} \left[\frac{F_1}{F_2} (\mathbf{B}^i)^2 + \left(\frac{F_1'}{F_4'} - \frac{F_1}{F_2} \right) \frac{(\phi \cdot \mathbf{B}^i)^2}{\phi^2} \right]. \quad (31b)$$

C. Magnetic field and mass of a static solution

Asymptotically, in the vacuum, the field-strength tensor corresponding to the unbroken “electromagnetic” $U(1)$ subgroup is $F^{\mu\nu} \equiv \frac{1}{v} \phi_\infty \cdot \mathbf{F}^{\mu\nu}$. Therefore, we can define and calculate the asymptotic magnetic field as

$$B^i \equiv -\frac{1}{2} \epsilon^{ijk} F^{jk} \quad (32a)$$

$$= \frac{1}{v} \phi_\infty \cdot \mathbf{B}^i \quad (32b)$$

$$= \frac{1}{2} H(\infty) \epsilon^{ijk} (\partial^j \mathbf{n} \times \partial^k \mathbf{n}) \cdot \mathbf{n}, \quad (32c)$$

where $H(\infty) = \pm 1$ is value of H at spatial infinity. The magnetic charge of a static configuration follows as

$$q_m = \lim_{r \rightarrow \infty} \int_{S^2} dS_i B_i \quad (33a)$$

$$= \frac{1}{2} H(\infty) \underbrace{\lim_{r \rightarrow \infty} \int_{S^2} dS_i \epsilon^{ijk} (\partial^j \mathbf{n} \times \partial^k \mathbf{n}) \cdot \mathbf{n}}_{8\pi N} \quad (33b)$$

$$= H(\infty) 4\pi N, \quad (33c)$$

where $N \in \mathbb{Z}$ is the degree of the mapping \mathbf{n} .

We calculate the mass of a static configuration using the Gauss–Ostrogradsky theorem (valid for regular \mathcal{E}). Using the previous result (33), we obtain

$$M = \int_{\mathbb{R}^3} d^3x \mathcal{E} \quad (34a)$$

$$= \int_{\mathbb{R}^3} d^3x \partial^i \left(\frac{F_1}{gH} \phi \cdot \mathbf{B}^i \right) \quad (34b)$$

$$= \lim_{r \rightarrow \infty} \int_{S^2} dS_i \frac{F_1}{gH} \phi \cdot \mathbf{B}^i \quad (34c)$$

$$= \frac{v F_1(1)}{2g} \underbrace{\lim_{r \rightarrow \infty} \int_{S^2} dS_i \epsilon^{ijk} (\partial^j \mathbf{n} \times \partial^k \mathbf{n}) \cdot \mathbf{n}}_{8\pi N = 2H(\infty) q_m} \quad (34d)$$

$$= F_1(1) \frac{4\pi v}{g} N, \quad (34e)$$

where $F_1(1) = \pm 1$. It follows that for the mass to be positive, the two signs must be related as

$$F_1(1) \operatorname{sgn} N = +1, \quad (35)$$

so that $M = \frac{4\pi v}{g} |N| \geq 0$.

Moreover, the magnetic charge can be rewritten as $q_m = \sigma 4\pi |N|$, where the sign $\sigma \equiv F_1(1)H(\infty) = \pm 1$ informs us whether we have a monopole ($q_m > 0$) or antimonopole ($q_m < 0$).

IV. SPHERICAL SYMMETRY

A. The Ansatz

Let us now focus on a spherically symmetric $N = 1$ solutions for which we adopt the standard “hedgehog” Ansatz:

$$\phi = vH \frac{\mathbf{r}}{r}, \quad \mathbf{A}^i = -\frac{\mathbf{r} \times \partial^i \mathbf{r}}{r^2} (1 - K), \quad (36)$$

or in components,

$$\phi_a = vH \frac{x_a}{r}, \quad A_a^i = -\frac{1}{r^2} \epsilon_{abi} x_b (1 - K). \quad (37)$$

Both form factors H and K are functions of $r = |\mathbf{r}|$ and satisfy the boundary conditions

$$H(\infty) = 1, \quad K(\infty) = 0, \quad (38)$$

⁴ The BPS equation is obtained simply as a condition for the square bracket in (17) to vanish (so that the energy density is just a total derivative). However, as the resulting equation is a bit messy and complicated, some of its consequences have to be used to obtain the elegant form (29).

that follow from the requirement of the finite total energy. Notice that we have deliberately chosen $H(\infty) = +1$ instead of -1 ; this, together with

$$F_1(1) = F_2(1) = +1, \quad (39)$$

that follows from (35) and from $N = 1$, means that we are considering, without loss of generality, only monopoles and not anti-monopoles.

At this point it is customary to impose the conditions $H(0) = 0$ and $K(0) = 1$, so that the fields ϕ , \mathbf{A}^i are regular at the origin. (Provided that $K - 1$ and H go to zero fast enough, i.e., at least linearly and quadratically, respectively.) In case of 't Hooft–Polakov monopole in the Georgi–Glashow theory (i.e., $F_i = H$ for all i), these conditions are critical for having a finite mass. However, in our enlarged settings (i.e., $F_i \neq H$ for some i) we should carefully reconsider their necessity and explore the implications of allowing certain types of singularities in these fields.

Thus, in this paper, we shall adopt a philosophy that a divergence in *gauge-dependent* fields \mathbf{A}^i and *form-dependent* field H (see our comment at the end of Sec. II B) is not – by itself – necessarily a problem. What will bother us, however, will be regularity of the energy density in the origin, which is without any doubts a physical quantity that must not diverge.

B. The BPS equations and the energy density

The BPS equation (29) under the Ansatz (36) decouples into a system of two ordinary differential equations for K and H :

$$\partial_\rho(\log K) = -F_2(H), \quad (40a)$$

$$\partial_\rho H = \frac{1 - K^2}{\rho^2} \frac{1}{F_4(H)}, \quad (40b)$$

where we introduced a dimensionless radius

$$\rho \equiv vgr. \quad (41)$$

From the first equation we can immediately read off (due to (38) and (39)) the asymptotic behavior of K :

$$K \sim \exp(-\rho) \quad \text{as} \quad \rho \rightarrow \infty. \quad (42)$$

The energy density reads

$$\frac{\mathcal{E}}{v^4 g^2} = \frac{\partial_\rho [F_1(1 - K^2)]}{\rho^2} \quad (43)$$

$$= 2F_1 F_2 \frac{K^2}{\rho^2} + \frac{F_1'}{F_4'} \frac{(1 - K^2)^2}{\rho^4}, \quad (44)$$

where, on the second line, we used the BPS equations to eliminate the derivatives with respect to ρ . The first line can be immediately integrated over 3-volume to obtain the mass of the monopole

$$M = \int_{\mathbb{R}^3} d^3x \mathcal{E} = \frac{4\pi v}{g} \int_0^\infty d\rho \rho^2 \frac{\mathcal{E}}{v^4 g^2} = \frac{4\pi v}{g}, \quad (45)$$

in agreement with the general formula (34) (using $N = 1$ and (39)).

C. Solving the BPS equations

We can recast the system of the two first-order equations (40) into a single second-order equation by a straightforward substitution. First, let us for convenience introduce the shorthand

$$\kappa \equiv -\partial_\rho(\log K). \quad (46)$$

Also, from now on we assume F_2 to be invertible.⁵ Then we have

$$H = F_2^{-1}(\kappa). \quad (47)$$

Next, we introduce two functions F , G of κ as

$$F(\kappa) \equiv F_1(F_2^{-1}(\kappa)), \quad (48a)$$

$$G(\kappa) \equiv F_4(F_2^{-1}(\kappa)). \quad (48b)$$

Their derivatives can be expressed as

$$F'(\kappa) = \frac{F_1'}{F_2'}, \quad G'(\kappa) = \frac{F_4'}{F_2'}, \quad (49)$$

where the primes on the right-hand sides are derivatives with respect to H , which itself is understood as a function of κ via (47).

Expressing H from (40a) as (47) and substituting it into (40b), we obtain a single second-order equation for the form factor K :

$$\partial_\rho^2(\log K) = -\frac{1 - K^2}{\rho^2} \frac{1}{G'(\kappa)}, \quad (50)$$

where we used also (49).

Using the definitions above we can rewrite the energy density (44) as

$$\frac{\mathcal{E}}{v^4 g^2} = 2F(\kappa)\kappa \frac{K^2}{\rho^2} + \frac{F'(\kappa)}{G'(\kappa)} \frac{(1 - K^2)^2}{\rho^4}. \quad (51)$$

In this form, the energy density depends only on K (and its derivative via κ) and there is no explicit dependence on H . Therefore the invariance of the energy density under the transformation $H \rightarrow \tilde{H}$, which we showed in Sec. II C, is now manifest. Also note that while the energy density in the form (44) depended on three functions, now it depends only on two. Recall that of the original four functions F_i that define our theory, one is removed by the BPS condition and another is removed by the form-invariance of the Lagrangian. The two functions F and G are exactly the two remaining *physical* functions invariant under the transformation (10): $F \equiv F_1 \circ F_2^{-1} = F_1 \circ \alpha \circ \alpha^{-1} \circ F_2^{-1}$ (and analogously for G).

This suggests to switch from the parameterization of our BPS Lagrangian in terms of three functions F_1 , F_2 ,

⁵ However, see the discussion at the end of the paper in Sec. VI.

F_4 to another parameterization in terms of F and G and one non-physical function that would merely represent the freedom under the form-invariance (10). The most straightforward way to do this is to trade F_1, F_4 for F, G by defining

$$F_1(H) = F(F_2(H)), \quad (52a)$$

$$F_4(H) = G(F_2(H)). \quad (52b)$$

Here, we let F_2 play the rôle of the non-physical function. In this way we can recast the Lagrangian (27) – yet again – as

$$\mathcal{L} = \frac{v^2}{2} \left[FF_2 \frac{(D^\mu \phi)^2}{\phi^2} + [(HF_2')^2 F'G' - FF_2] \frac{(\phi \cdot D^\mu \phi)^2}{\phi^4} \right] - \frac{1}{4g^2} \left[\frac{F}{F_2} (\mathbf{F}^{\mu\nu})^2 + \left(\frac{F'}{G'} - \frac{F}{F_2} \right) \frac{(\phi \cdot \mathbf{F}^{\mu\nu})^2}{\phi^2} \right], \quad (53)$$

where the primes at F, G are derivatives with respect to $\kappa = F_2(H)$. The virtue of this (final!) form is that it separates the functions F, G that are physical from the function F_2 that is arbitrary and merely represents our freedom to perform form-invariant transformation (10).

Lastly, let us comment on a special case of $F(\kappa) = 1$. (It follows that also F_1 must be then a constant function, $F_1(H) = 1$.) There are two consequences. First, the Lagrangian (53) reduces to

$$\mathcal{L} = \frac{v^2}{2} F_2 (D^\mu \mathbf{n})^2 - \frac{1}{4g^2} \frac{1}{F_2} \left[(\mathbf{F}^{\mu\nu})^2 - \frac{(\phi \cdot \mathbf{F}^{\mu\nu})^2}{\phi^2} \right], \quad (54)$$

so that the scalar singlet H , living now only in $F_2(H)$, becomes a non-dynamical field. Second, the energy density cannot be simultaneously regular and positive. This can be seen by assuming power function behavior of K for small ρ as $K = a\rho^n + \dots$. The energy density then reads $\frac{\mathcal{E}}{v^4 g^2} = -\frac{(K^2)'}{\rho^2} = -2na^2 \rho^{2n-3} + \dots$. Thus, either $n \geq 3/2$ (or $n \geq 1/2$, if we allow for integrable singularity), so that \mathcal{E} is in the origin regular (but negative), or $n < 0$, implying positive (but singular) \mathcal{E} . For these reasons, we will not consider the case $F(\kappa) = 1$.

D. Recipe for finding analytic solutions of BPS monopoles

At this point, it is the right time to briefly recapitulate the previous developments and formulate a recipe for finding a theory that allows an analytic monopole solution.

1. The first step is to find a function $G(\kappa)$ such that the equation (50) can be solved in a closed form for $K(\rho)$.

Taking into account the definition (48b), we deduce that $G(\kappa)$ must be defined and continuous on some neighborhood of $\kappa = 1$. This is to allow for the existence of a function $F_2(H)$ that would not only satisfy the condition $F_2(1) = 1$ (recall (39) and (47)), but to be also invertible for all $H \in [0, \infty)$.

2. Next step is to choose a function $F(\kappa)$. There are two requirements that come from the desired physical properties of the energy density. (Notice that in its expression (51) everything except F is already fixed.) First, F must satisfy

$$\text{sgn } F(\kappa) = \text{sgn } \kappa, \quad (55a)$$

$$\text{sgn } F'(\kappa) = \text{sgn } G'(\kappa), \quad (55b)$$

for all κ (i.e., for all $\rho \geq 0$), so that both terms in the energy density (51) are non-negative. Second, F must be such that the energy density is regular.

Having G', F and K , we have all physical ingredients that allow us, e.g., to draw a plot of the energy density (51).

3. The last step is choosing the function $F_2(H)$. This has a twofold meaning. Primarily, it allows us to find the remaining form factor H through the equation (47). More generally, however, with F_2 at hand, we can write down the full Lagrangian (53). Recall that, taking into account the form-invariance (10), all invertible functions F_2 are equally good and lead to physically equivalent Lagrangians.

E. Solvable case

To the best of our knowledge, the only case when an analytic solution to the equation (50) is known⁶ is when

$$G(\kappa) = \kappa. \quad (56)$$

This, by the way, corresponds to

$$F_4 = F_2, \quad (57)$$

which is certainly satisfying from an aesthetic point of view as it sort of symmetrizes the theory – compare it with the BPS condition $F_3 = F_1$.

Upon substituting $G'(\kappa) = 1$ into the equation (50), we obtain the general solution $K = c\rho / \sinh[c(\rho - \rho_0)]$, where the constants of integration c, ρ_0 are in principle arbitrary. After all, this solution satisfies the boundary condition $K(\infty) = 0$ for all values of c, ρ_0 .

To determine the value of c , we have to consider the other form factor, H , that can be calculated from $F_2(H) = \kappa$, where $\kappa = -\partial_\rho(\log K) = c \coth[c(\rho - \rho_0)] -$

⁶ This is not entirely correct. In fact, it *is* possible to find many analytic solutions to (50) by turning the logic upside down and using a kind of “reverse engineering”: Given a prescribed form $K(\rho)$, we can then look for $G'(\kappa)$ that solves (50). However, as simple as it sounds, it is surprisingly difficult to find in this way a G' that meets the conditions for its domain, as discussed at the point 1. of Sec. IV D. In particular, all G' we managed to find using this method happened to be singular at $\kappa = 1$.




Figure 1. Behavior of the function $\kappa = \coth(\rho - \rho_0) - 1/\rho$ for various ρ_0 . For $\rho_0 < 0$ it smoothly interpolates between $-\infty$ and 1, while for $\rho_0 = 0$ between 0 and 1. For $\rho_0 > 0$ there is a singularity at $\rho = \rho_0$.

$1/\rho$. From $\lim_{\rho \rightarrow \infty} \kappa = c$ (together with the normalization condition $F_2(1) = 1$) we learn that only $c = 1$ is consistent with the boundary condition $H(\infty) = 1$.

Next, to determine ρ_0 , we have to resort to the requirement that each of the two terms in the energy density

$$\frac{\mathcal{E}}{v^4 g^2} = F(\kappa) \kappa \frac{2K^2}{\rho^2} + F'(\kappa) \frac{(1 - K^2)^2}{\rho^4} \quad (58)$$

has to be separately both regular and non-negative. However, a simple inspection reveals that these conditions cannot be met for any function $F(\kappa)$ as long as $\rho_0 \neq 0$, due to the behavior of $\kappa(\rho)$ in such a case (see Fig. 1). Accordingly, we conclude that only $\rho_0 = 0$ is viable.

We therefore have

$$\kappa = \coth \rho - \frac{1}{\rho} \quad (59)$$

and the general solution for the form factors read

$$K = \frac{\rho}{\sinh \rho}, \quad H = F_2^{-1}(\kappa). \quad (60)$$

Notice that for F_2 being the identity function this is nothing but the notorious 't Hooft–Polyakov solution in the BPS limit. We also remark that since $K = 1 - \frac{1}{6}\rho^2 + \mathcal{O}(\rho^4)$, the gauge fields are regular at the origin:

$$A_a^i = -\frac{1}{6} \varepsilon_{abix} x_b v^2 g^2 \left[1 + \mathcal{O}(\rho^2) \right], \quad (61)$$

as is customary to require.

For the energy density (51) to be non-negative, the function F must satisfy

$$F(\kappa) \geq 0, \quad F'(\kappa) \geq 0, \quad (62)$$

for all $\kappa \in [0, 1]$, while the requirement of regularity of \mathcal{E} at $\rho = 0$ implies that $F(\kappa \rightarrow 0)$ must go to zero sufficiently fast. For instance, if it behaves near the origin like a power function

$$F(\kappa) \sim \kappa^N, \quad (63)$$

the exponent N must satisfy

$$N \geq 1. \quad (64)$$

We also see that, except in the special case $N = 1$ when $\mathcal{E}(0)/v^4 g^2 = F'(0)/3$, the energy density is always vanishing at $\rho = 0$. In other words, most of the energy of the monopole is concentrated not in its centre, but rather in a spherical shell around it. This is very similar to the *hollow monopoles* discussed in [8]. However, while the authors of [8] found and studied the hollow monopoles only numerically, we are going to present analytic solutions for them.

V. EXAMPLES

By choosing $G(\kappa) = \kappa$ we have fulfilled the first point in our recipe of Sec. IV D, so let us move to the next two points and provide some particular examples of functions $F(\kappa)$ (to be able to draw energy density) and $F_2(H)$ (to be able to write down Lagrangian density.)

A. Power function

As the simplest example let us take

$$F = \kappa^N. \quad (65)$$

In order that the energy density (with $G(\kappa) = \kappa$ and, correspondingly, K and κ given by (59) and (60))

$$\frac{\mathcal{E}}{v^4 g^2} = \kappa^N \left[2\kappa \frac{K^2}{\rho^2} + N \frac{1}{\kappa} \frac{(1 - K^2)^2}{\rho^4} \right] \quad (66a)$$

$$\xrightarrow{\rho \rightarrow 0} \rho^{N-1} \left[\frac{N+2}{3^{N+1}} + \mathcal{O}(\rho^2) \right] \quad (66b)$$

be regular in the origin, we must demand $N \geq 1$ in accordance with (64). In Fig. 2 we plot \mathcal{E} for various values of N . As advertised, for $N > 1$ most of the energy is concentrated away from the centre and the monopoles are hollow.

In order to write down the corresponding Lagrangian, we choose, for simplicity, the function $F_2(H)$ to be a power function, too. Thus, we arrive at

$$\mathcal{L} = \frac{v^2}{2} H^{n+m} \left[\frac{(D^\mu \phi)^2}{\phi^2} + (nm - 1) \frac{(\phi \cdot D^\mu \phi)^2}{\phi^4} \right] - \frac{1}{4g^2} H^{n-m} \left[(\mathbf{F}^{\mu\nu})^2 + \left(\frac{n}{m} - 1 \right) \frac{(\phi \cdot \mathbf{F}^{\mu\nu})^2}{\phi^2} \right], \quad (67)$$




Figure 2. Energy density for a single monopole solution of the power-function theory (67) for various $N = n/m$.

which in the language of F_i 's corresponds to

$$F_1 = F_3 = H^n, \quad (68a)$$

$$F_2 = F_4 = H^m, \quad (68b)$$

provided that

$$N = \frac{n}{m}. \quad (69)$$

The solution follows immediately as

$$K = \frac{\rho}{\sinh \rho}, \quad H = \left(\coth \rho - \frac{1}{\rho} \right)^{\frac{1}{m}}. \quad (70)$$

Interestingly, while the Lagrangian (67) is form-invariant under general transformations $H \rightarrow \tilde{H}$ of Eq. (10), there is also a special subclass of these transformations that affects only the parameters n and m . Namely, upon transforming

$$H \rightarrow H^\sigma, \quad (71)$$

where $\sigma \neq 0$ is arbitrary, we obtain a model of the same form as (67), up to the rescaling of parameters

$$n \rightarrow \sigma n, \quad m \rightarrow \sigma m. \quad (72)$$

Note that the physical (measurable) parameter $N = n/m$ is invariant under (72).

Obviously, for $n = m = 1$ this is just the well known 't Hooft–Polyakov monopole in the renormalizable Georgi–Glashow model (in the BPS limit) [3, 4]. However, any model with $n/m = 1$ is equivalent to the Georgi–Glashow model, even though the Lagrangian and the solution H are different.

The form factor H , Eq. (70), behaves in the origin like $H = (\rho/3 + \mathcal{O}(\rho^3))^{1/m}$ and, correspondingly, $\phi = vH\mathbf{x}/r = v^2 g\mathbf{x}\rho^{1/m-1}(1/3 + \mathcal{O}(\rho^2))^{1/m}$. If $1/m \geq 1$, the field ϕ is regular, as is customary to require. If, on




Figure 3. Energy density for a single monopole solution of the power-exponential-function theory (74) with $N = M = 2$, $a = 100$ and various k .

the other hand, $1/m < 1$, the field ϕ is singular at the origin. However, this singularity is completely unphysical and harmless, as it can be transformed away using (71) with, e.g., $\sigma = 1/m$.

B. Power-exponential function

As a slightly more complicated example, let us consider

$$F = \kappa^N \exp \left[a(\kappa^M - 1)^k \right]. \quad (73)$$

Here we assume $a \neq 0$ and $M \neq 0$, as otherwise we would obtain the previous case.

We take again $G = \kappa$. From the requirements that $F(\kappa = 1) = 1$ and $F' = F[N + akM\kappa^M(\kappa^M - 1)^{k-1}]/\kappa \geq 0$ we find that $N \geq 0$ and $aM > 0$, while k must be a positive, odd integer. Now there are two options with rather different properties:

- $a > 0$ and $M > 0$: In this case the energy density behaves for small ρ like a power function $\mathcal{E} \sim \rho^{N-1}$, so that it must be $N \geq 1$.
- $a < 0$ and $M < 0$: In this case we have $\mathcal{E} \sim \exp(-1/\rho^{k|M|})$ (times a power function) as $\rho \rightarrow 0$ and $N \geq 0$.

As an example of the Lagrangian that corresponds to this F , we can take

$$\begin{aligned} \mathcal{L} = & e^{a(H^\ell - 1)^k} H^{n-m} \\ & \times \left\{ \frac{v^2}{2} H^{2m} \left[\frac{(D^\mu \phi)^2}{\phi^2} + (mns - 1) \frac{(\phi \cdot D^\mu \phi)^2}{\phi^4} \right] \right. \\ & \left. - \frac{1}{4g^2} \left[(\mathbf{F}^{\mu\nu})^2 + \left(\frac{n}{m}s - 1 \right) \frac{(\phi \cdot \mathbf{F}^{\mu\nu})^2}{\phi^2} \right] \right\}, \quad (74) \end{aligned}$$




Figure 4. Energy density for a single monopole solution of the power-exponential-function theory (74) with $N = 2$, $M = -1$, $k = 101$ and various $a \leq 0$.

where we denoted for brevity

$$s \equiv 1 + alk(H^{n\ell} - 1)^{k-1}H^{n\ell}. \quad (75)$$

In the language of F_i 's we would specify the theory by

$$F_1 = F_3 = H^n \exp \left[a(H^{n\ell} - 1)^k \right], \quad (76a)$$

$$F_2 = F_4 = H^m, \quad (76b)$$

which is related to the F by the identification

$$N = \frac{n}{m}, \quad M = \ell \frac{n}{m}. \quad (77)$$

The form factors K and H are given by the very same expressions (70) as in the previous power-function model.

The reason to consider this complicated and rather artificial model is that it showcases more interesting energy density profiles with richer structure (admittedly, for “unnaturally” large values of parameters). While we again obtain the hollow monopoles, this time they have new features. As depicted in Fig. 3, for some range of parameters there can be not only one minimum of the energy density, but two. In Fig. 4, we see that the peak of the energy density can be much sharper. Moreover, since in the latter case the energy density falls off exponentially for small ρ (due to $a < 0$), there is a well-defined finite region in the centre of the monopole with virtually vanishing energy density – accordingly we can dub these solutions the *truly hollow* monopoles.

VI. SUMMARY AND OUTLOOK

In this paper we have investigated spherically symmetric solutions of a family of general $SU(2)$ gauge theories (2) with adjoint scalar in the BPS limit.

First, we discussed the redundancy and physical rôle of the four functions f_i^2 that define our model. There is a form-invariance of our theory (2) that exploits a general field redefinition (10) of the gauge-invariant part of the adjoint scalar ϕ . This redefinition leaves the structure of the model intact and allows to fix one of the four functions (e.g., to have the kinetic term of ϕ canonically normalized). By itself, however, the transformation (10) is not enough to eliminate non-canonical structure of the (dressed) gauge fields. To achieve that, we have introduced a form non-invariant transformation (13) that defines canonically normalized gauge fields and through which the starting Lagrangian (2) is recast into (14). In this way, the presence of field-dependent $SU(2)$ magnetic susceptibility (functions f_2^2 and f_4^2) in Eq. (2) are found in Eq. (14) to control the strength of dipole-moment interactions. This clearly illustrates the physical rôle of f_2^2 and f_4^2 and opens up an avenue for further investigations of their impact on physics that we plan to do in the future. (In App. A, we discuss a generalization of both transformations and showcase its group-like properties and the impact on the hedgehog Ansatz.)

Second, we have formulated a BPS version of the theory with the help of reparametrizing the Lagrangian (2) in terms of new, more “BPS-friendly” functions F_i (rather than f_i^2). This has several advantages: i) the key BPS condition is stated very simply as $F_3 = F_1$, ii) the form-invariance (10) becomes a simple function composition, and iii) the solvable cases that are studied in this paper – all of which can be regarded as certain generalizations of the ’t Hooft–Polyakov monopole – belong to a special subset that is defined simply as $F_4 = F_2$.

Third, in the BPS limit there are only two physically relevant form-functions that we have labelled F and G (see Eq. (52)). We have then constructed an especially convenient form of the Lagrangian, (53), that is written solely in terms of form-functions F , G and F_2 , with the freedom of performing (10) being entirely contained in F_2 . Indeed, this function does not appear in a general formula (51) for energy density of spherically symmetric solutions, which is thus manifestly form-invariant.

Having clarified the rôle of various functions, the rest of the paper presents in Sec. V some concrete examples of analytic BPS solutions that illustrate an application of a general “recipe” described in Sec. IV D. These examples are not meant to be exhaustive, rather they expose a key point of this study: How the shape of a monopole (i.e., distribution of its energy) depends on the form-function F . (The other form-function, G , is being fixed for simplicity, although we suspect its rôle to be qualitatively the same.) While the general dependence might be glimpsed from the formula (51), in particular, we have seen that the position and number of extrema can be controlled by the choice of F (see Fig. 3).

A universal feature seems to be tendency of energy to concentrate in a shell rather than at the monopole’s centre. In other words, a generic monopole of our family of models tends to be *hollow*. The only exception

in the examples presented here are the cases $n = m$ of the simplest power-function model (67) that are actually nothing but the 't Hooft–Polyakov monopole, up to a field redefinition (10). These cases also stand out by the presence of canonically normalized gauge kinetic term in the Lagrangian density.

On the other hand, in all our examples of hollow monopoles the gauge kinetic terms are always modified by some power of the scalar singlet H , manifesting a non-trivial $SU(2)$ magnetic dipole-moment of the adjoint triplet. Intuitively, the presence of hollow monopoles therefore seems to be connected with vanishing (or “freezing”) of the gauge kinetic term at the monopole’s centre, namely that the dipole-moment interactions that *screen* the bare monopole charge become effectively infinite. At this point, we present it as an observation, however, we are planing to examine this issue more thoroughly by investigating the properties of homogeneous “phases” of the theory (2) in a future work.

Lastly, in this paper, we have only expounded the case of *invertible* F_2 that allowed us to condense the system of two first-order BPS equations (40) into a single second-order equation (50). However, there is an entire branch of analytic solutions that correspond to a *non-invertible* choices of F_2 . For instance, taking $F_2 = 1$ yields a particularly simple form of K :

$$K = \xi e^{-\rho}, \quad (78)$$

where ξ is a constant of integration. (The other form factor, H , depends on F_4 and is typically much more complicated.) First of all, near origin $K \sim \xi(1 - \rho + \mathcal{O}(\rho^2))$ and, correspondingly, $A_a^i \sim \varepsilon_{abi} x_b (\xi - 1 + \mathcal{O}(r))/r^2$ as $r \rightarrow 0$. Notice that this singularity only becomes milder if $\xi = 1$, but does not disappear completely. As discussed at the beginning of this paper, by itself a singular behavior might not be problematic, as it can be transformed away by a general field redefinition of the type discussed in Appendix A. Interestingly, however, the physical requirements (like regularity of the energy density) do not constrain ξ to have a single value, but rather $\xi \in [-1, 1]$. In other words, ξ is a completely new moduli of the BPS solution and it is physical in the sense that it controls the shape of the energy density. In particular, it is related to the width of the hollow cavity.

We plan to investigate this class of solutions in a separate paper. There we hope to elucidate the physical

origin of this moduli and whether the corresponding solutions are physically viable. In particular, we have to analyze the stability of these solutions, as presence of ξ might lead to dynamical instabilities. Further, we need to study them in non-BPS cases and identify what are the exact conditions that lead to their presence.

ACKNOWLEDGMENTS

The authors would like to express the gratitude for the institutional support of the Institute of Experimental and Applied Physics, Czech Technical University in Prague (P. B. and F. B.), and of the Research Centre for Theoretical Physics and Astrophysics, Institute of Physics, Silesian University in Opava (F. B.). P. B. is indebted to Aštar Šeran and FSM for invaluable discussions.

Appendix A: General canonical transformation for $SU(2)$ adjoint fields

Let us consider the transformation $\{\mathbf{A}_\mu, \phi\} \rightarrow \{\tilde{\mathbf{A}}_\mu, \tilde{\phi}\}$ generated by the functions α, h, k, ℓ as

$$\phi = \frac{\alpha(\tilde{H})}{\tilde{H}} \tilde{\phi}, \quad (A1a)$$

$$\mathbf{A}_\mu = \tilde{\mathbf{A}}_\mu + h(\tilde{H}) \left[\frac{\tilde{D}_\mu \tilde{\phi}}{v\tilde{H}} - \frac{\tilde{\phi} \cdot \tilde{D}_\mu \tilde{\phi}}{(v\tilde{H})^3} \tilde{\phi} \right] + k(\tilde{H}) \frac{\tilde{\phi} \cdot \tilde{D}_\mu \tilde{\phi}}{(v\tilde{H})^3} \tilde{\phi} + [\ell(\tilde{H}) - 1] \frac{\tilde{\phi} \times \tilde{D}_\mu \tilde{\phi}}{(v\tilde{H})^2}, \quad (A1b)$$

where $\tilde{\phi} = v\tilde{H}\mathbf{n}$ and $\tilde{D}_\mu \tilde{\phi} \equiv \partial_\mu \tilde{\phi} + \tilde{\mathbf{A}}_\mu \times \tilde{\phi}$. Equation (A1a) is equivalent to $H = \alpha(\tilde{H})$.

The transformations of the covariant derivative and field-strength tensor follow as

$$\frac{D_\mu \phi}{vH} = \ell \frac{\tilde{D}_\mu \tilde{\phi}}{v\tilde{H}} - h \frac{\tilde{\phi} \times \tilde{D}_\mu \tilde{\phi}}{(v\tilde{H})^2} - \left(\ell - \tilde{H} \frac{\alpha'}{\alpha} \right) \frac{\tilde{\phi} \cdot \tilde{D}_\mu \tilde{\phi}}{(v\tilde{H})^3} \tilde{\phi}, \quad (A2)$$

$$\mathbf{F}_{\mu\nu} = \ell \tilde{\mathbf{F}}_{\mu\nu} - h \frac{\tilde{\phi} \times \tilde{\mathbf{F}}_{\mu\nu}}{v\tilde{H}} - (\ell - 1) \frac{\tilde{\phi} \cdot \tilde{\mathbf{F}}_{\mu\nu}}{(v\tilde{H})^2} \tilde{\phi} + \mathbf{d}_{\mu\nu}, \quad (A3)$$

where we defined

$$\begin{aligned} \mathbf{d}_{\mu\nu} \equiv & (\tilde{H}h' - k\ell) \frac{(\tilde{\phi} \cdot \tilde{D}_\mu \tilde{\phi})(\tilde{D}_\nu \tilde{\phi}) - (\mu \leftrightarrow \nu)}{(v\tilde{H})^3} + (\tilde{H}\ell' + kh) \frac{(\tilde{\phi} \cdot \tilde{D}_\mu \tilde{\phi})(\tilde{\phi} \times \tilde{D}_\nu \tilde{\phi}) - (\mu \leftrightarrow \nu)}{(v\tilde{H})^4} \\ & - \left[(\ell^2 + h^2 - 1) - (\ell - 1)^2 \right] \left[\frac{(\tilde{\phi} \cdot \tilde{D}_\mu \tilde{\phi})(\tilde{\phi} \times \tilde{D}_\nu \tilde{\phi}) - (\mu \leftrightarrow \nu)}{(v\tilde{H})^4} - \frac{\tilde{D}_\mu \tilde{\phi} \times \tilde{D}_\nu \tilde{\phi}}{(v\tilde{H})^2} \right] + (\ell - 1)^2 \frac{\tilde{\phi} \cdot (\tilde{D}_\mu \tilde{\phi} \times \tilde{D}_\nu \tilde{\phi})}{(v\tilde{H})^4} \tilde{\phi}, \end{aligned} \quad (A4)$$

with the primes being differentiations with respect to \tilde{H} . Under the transformation (A1) the original Lagrangian \mathcal{L} ,

Eq. (2), transforms to another Lagrangian $\tilde{\mathcal{L}}$ of the form

$$\begin{aligned} \tilde{\mathcal{L}} = & \frac{v^2}{2} \left[\tilde{f}_1^2 \left(\frac{(\tilde{D}^\mu \tilde{\phi})^2}{\tilde{\phi}^2} - \frac{(\tilde{\phi} \cdot \tilde{D}^\mu \tilde{\phi})^2}{\tilde{\phi}^4} \right) + \tilde{f}_3^2 \frac{(\tilde{\phi} \cdot \tilde{D}^\mu \tilde{\phi})^2}{\tilde{\phi}^4} \right] - \frac{1}{4g^2} \left[\tilde{f}_2^2 \left((\mathbf{F}^{\mu\nu})^2 - \frac{(\tilde{\phi} \cdot \mathbf{F}^{\mu\nu})^2}{\tilde{\phi}^2} \right) + \tilde{f}_4^2 \frac{(\tilde{\phi} \cdot \mathbf{F}^{\mu\nu})^2}{\tilde{\phi}^2} \right] \\ & - \frac{1}{2g^2} \mathbf{d}^{\mu\nu} \cdot \left\{ f_2^2 \ell \left[\tilde{\mathbf{F}}_{\mu\nu} - \frac{(\tilde{\phi} \cdot \tilde{\mathbf{F}}_{\mu\nu})}{\tilde{\phi}^2} \tilde{\phi} \right] + f_2^2 h \frac{\tilde{\mathbf{F}}_{\mu\nu} \times \tilde{\phi}}{v\tilde{H}} + f_4^2 \frac{(\tilde{\phi} \cdot \tilde{\mathbf{F}}_{\mu\nu})}{\tilde{\phi}^2} \tilde{\phi} \right\} \\ & - \frac{1}{4g^2} \left[f_2^2 \left((\mathbf{d}_{\mu\nu})^2 - \frac{(\tilde{\phi} \cdot \mathbf{d}_{\mu\nu})^2}{\tilde{\phi}^2} \right) + f_4^2 \frac{(\tilde{\phi} \cdot \mathbf{d}_{\mu\nu})^2}{\tilde{\phi}^2} \right] - \tilde{V}(\tilde{\phi}^2), \end{aligned} \quad (\text{A5})$$

where the new functions $\tilde{f}_i^2 = \tilde{f}_i^2(\tilde{H})$ are given as

$$\tilde{f}_i^2 = f_i^2(\ell^2 + h^2), \quad (i = 1, 2) \quad (\text{A6a})$$

$$\tilde{f}_3^2 = f_3^2 \left(\frac{\tilde{H} \alpha'}{\alpha} \right)^2, \quad (\text{A6b})$$

$$\tilde{f}_4^2 = f_4^2, \quad (\text{A6c})$$

with $f_i^2 = f_i^2(\alpha(\tilde{H}))$, and $\tilde{V}(\tilde{\phi}^2) = V(\phi^2)$.

We also obtain

$$\phi \cdot \mathbf{A}_\mu = \frac{\alpha}{\tilde{H}} \left[\tilde{\phi} \cdot \tilde{\mathbf{A}}_\mu + k \frac{\tilde{\phi} \cdot \tilde{D}_\mu \tilde{\phi}}{v\tilde{H}} \right], \quad (\text{A7})$$

$$\phi \cdot \mathbf{F}_{\mu\nu} = \frac{\alpha}{\tilde{H}} \left[\tilde{\phi} \cdot \tilde{\mathbf{F}}_{\mu\nu} + (\ell^2 + h^2 - 1) \frac{\tilde{\phi} \cdot (\tilde{D}_\mu \tilde{\phi} \times \tilde{D}_\nu \tilde{\phi})}{\tilde{\phi}^2} \right]. \quad (\text{A8})$$

By performing two consecutive transformations (A1) we obtain again a transformation of the form (A1). For the scalar it is obvious: If we transform first $H_1 \rightarrow H_2$ and then $H_2 \rightarrow H_3$ with $H_1 = \alpha_2(H_2)$ and $H_2 = \alpha_3(H_3)$, respectively, then the combined transformation $H_1 \rightarrow H_3$ is done via $H_1 = \alpha_2(\alpha_3(H_3))$, i.e., by simple function composition. For the gauge fields the combined transformation is more complicated. If the first transformation is by $h_2(H_2)$, $k_2(H_2)$, $\ell_2(H_2)$ and the second one by

$h_3(H_3)$, $k_3(H_3)$, $\ell_3(H_3)$, then the combination of the two transformation is done via

$$h(h_3) = \ell_2(H_2) h_3(H_3) + h_2(H_2) \ell_3(H_3), \quad (\text{A9a})$$

$$\ell(h_3) = \ell_2(H_2) \ell_3(H_3) - h_2(H_2) h_3(H_3), \quad (\text{A9b})$$

$$k(h_3) = k_3(H_3) + k_2(H_2) H_3 \frac{\alpha'_3(H_3)}{\alpha_3(H_3)}. \quad (\text{A9c})$$

Unless $h = k = 0$, the general transformation (A1) does *not* protect the spherically symmetric Ansatz (37), but rather leads to

$$\tilde{\phi}_a = v \tilde{H} \frac{x_a}{r}, \quad (\text{A10a})$$

$$\tilde{A}_a^i = -\frac{\varepsilon_{abi} x_b}{r^2} (1 - \tilde{K}) + \frac{\delta_{ia} r^2 - x_i x_a}{r^3} \tilde{L} + \frac{x_i x_a}{r^3} \tilde{M}, \quad (\text{A10b})$$

where \tilde{H} is obtained by inverting $\alpha(\tilde{H}) = H$, while

$$\tilde{K} \equiv \frac{\ell}{\ell^2 + h^2} K, \quad \tilde{L} \equiv \frac{-h}{\ell^2 + h^2} K, \quad \tilde{M} \equiv -kr \frac{\tilde{H}'}{\tilde{H}}. \quad (\text{A11})$$

Nevertheless, it is still possible to require the transformation (A1) to protect, at least, the transversality of the gauge and scalar fields: $\phi \cdot \mathbf{A}^i = \tilde{\phi} \cdot \tilde{\mathbf{A}}^i = 0$. In other words, one can demand $\tilde{M} = 0$, implying $k = 0$.

-
- [1] F. Blaschke and P. Beneš, *All finite-mass Dirac monopoles*, *Phys. Rev. D* **106** (2022) 125014, [2210.11854].
- [2] K.-M. Lee and E. J. Weinberg, *Nontopological magnetic monopoles and new magnetically charged black holes*, *Phys. Rev. Lett.* **73** (1994) 1203–1206, [hep-th/9406021].
- [3] G. 't Hooft, *Magnetic Monopoles in Unified Gauge Theories*, *Nucl. Phys. B* **79** (1974) 276–284.
- [4] A. M. Polyakov, *Particle Spectrum in Quantum Field Theory*, *JETP Lett.* **20** (1974) 194–195.
- [5] H. S. Ramadhan, *Some exact BPS solutions for exotic vortices and monopoles*, *Phys. Lett. B* **758** (2016) 140–145, [1512.01640].
- [6] D. Bazeia, M. A. Marques and R. Menezes, *Magnetic monopoles with internal structure*, *Phys. Rev. D* **97** (2018) 105024, [1805.03250].
- [7] R. Casana, M. M. Ferreira, Jr and E. da Hora, *Generalized BPS magnetic monopoles*, *Phys. Rev. D* **86** (2012) 085034, [1210.3382].
- [8] D. Bazeia, M. A. Marques and G. J. Olmo, *Small and hollow magnetic monopoles*, *Phys. Rev. D* **98** (2018) 025017, [1807.01299].