#### SPECTRAL GRID INVARIANTS AND LAGRANGIAN COBORDISMS

MITCHELL JUBEIR, INA PETKOVA, NOAH SCHWARTZ, ZACHARY WINKELER, AND C.-M. MICHAEL WONG

ABSTRACT. We prove that the filtered GRID invariants of Legendrian links in link Floer homology, and consequently their associated invariants in the spectral sequence, obstruct decomposable Lagrangian cobordisms in the symplectization of the standard contact structure on  $\mathbb{R}^3$ , strengthening a result by Baldwin, Lidman, and the fifth author.

#### 1. Introduction

An interesting and difficult problem in contact and symplectic geometry is to decide, given contact 3-manifolds  $(Y_-, \xi_-)$  and  $(Y_+, \xi_+)$ , and two Legendrian links

$$\Lambda_{-} \subset (Y_{-}, \xi_{-}), \qquad \Lambda_{+} \subset (Y_{+}, \xi_{+}),$$

whether  $\Lambda_{-}$  and  $\Lambda_{+}$  are related by an exact Lagrangian cobordism in a Weinstein cobordism from  $(Y_{-}, \xi_{-})$  to  $(Y_{+}, \xi_{+})$ . Even in the simplest case where  $Y_{\pm} = \mathbb{R}^{3}$  and  $\xi_{\pm} = \xi_{\text{std}}$  is the standard contact structure given by the kernel of the 1-form

$$\alpha_{\rm std} = dz - y dx$$

and the Weinstein cobordism is the symplectization

$$(\mathbb{R}_t \times \mathbb{R}^3, \mathrm{d}(e^t \alpha_{\mathrm{std}})),$$

this turns out to be a challenging problem.

Focusing on this case in the present article, we recall that the classical invariants—the Thurston–Bennequin and rotation numbers—do give obstructions: Chantraine [Cha10] shows that if there exists an exact Lagrangian cobordism  $L: \Lambda_- \to \Lambda_+$ , then

$$tb(\Lambda_+) - tb(\Lambda_-) = -\chi(L), \qquad r(\Lambda_+) = r(\Lambda_-).$$

For instance, since the undestabilizable Legendrian unknot  $\Lambda_0$  and the undestabilizable Legendrian representative  $\Lambda_1$  of  $m(6_2)$  both have tb=-1, we know that any exact Lagrangian cobordism from  $\Lambda_0$  to  $\Lambda_1$  (or from  $\Lambda_1$  to  $\Lambda_0$ ) must be a concordance, but since  $m(6_2)$  is not smoothly slice, no such cobordism can exist.<sup>1</sup> However, the classical invariants do not give a complete answer: For example, using the functoriality [EHK16] of Legendrian contact homology from symplectic field theory [EGH00, Che02, BC14], Chantraine [Cha15] shows that for the undestabilizable representative  $\Lambda_2$  of  $m(9_{46})$ , which has tb=-1 and r=0, there exists a Lagrangian concordance  $L:\Lambda_0\to\Lambda_2$  but not a Lagrangian concordance  $L':\Lambda_2\to\Lambda_0$ .<sup>2</sup> Thus, an important goal is to develop effective obstructions to exact Lagrangian cobordisms, which give obstructions beyond the classical invariants, such as Legendrian contact homology as mentioned above. For more examples of obstructions from symplectic field theory, see [ST13, CDGG15, CNS16, Pan17].

<sup>&</sup>lt;sup>1</sup>This contrasts with the situation in smooth topology, where a cobordism exists between any pair of links, but the minimum genus of such a cobordism is difficult to determine.

<sup>&</sup>lt;sup>2</sup>In particular, this also illustrates the important fact that exact Lagrangian cobordisms are directed.

Another source of effective invariants is knot (Heegaard) Floer homology. In earlier work, Baldwin, Lidman, and the fifth author [BLW22] prove that the so-called *GRID invariants* in knot Floer homology can effectively obstruct *decomposable* Lagrangian cobordisms (of any genus), which are cobordisms that can be obtained by concatenating elementary cobordisms associated to Legendrian isotopies, pinches, and births, as depicted in Figure 2. Decomposable cobordisms are exact, and most connected exact Lagrangian cobordisms between non-empty Legendrian links are known to be decomposable; whether all connected exact Lagrangian cobordisms between non-empty Legendrian links are decomposable remains a major open problem. If one restricts to cobordisms of genus zero, there is also a body of work [BS18, BS21, GJ19] that shows that knot Floer homology gives effective obstructions in more general settings.

The goal of the present article is to extend the result in [BLW22] to the context of *filtered* knot Floer chain complexes, which will then imply that certain invariants in the associated spectral sequences also provide effective obstructions to exact Lagrangian cobordisms. The existence of these invariants is known to experts in knot Floer homology, but their definitions have not appeared in the literature thus far; in Section 1.1 below, we first provide the definitions.

1.1. The spectral GRID invariants. We briefly recall the invariants of Legendrian links in  $(\mathbb{R}^3, \xi_{\text{std}})$  defined by Ozsváth, Szabó, and Thurston [OSzT08] using the combinatorial grid-diagram formulation [MOS09, MOSzT07] of knot Floer homology; for more details, see Section 2.2 below. For ease of exposition, we focus on the *tilde* flavor, which computes a stabilized version of knot Floer homology. Let  $\mathbb{G}$  be a grid diagram of size n that represents a Legendrian link  $\Lambda \subset (\mathbb{R}^3, \xi_{\text{std}})$ ; then there are *canonical generators*  $\mathbf{x}^+(\mathbb{G})$  and  $\mathbf{x}^-(\mathbb{G})$ , which are cycles of the *grid chain complex*  $\widetilde{\mathrm{GC}}(\mathbb{G})$ , and thus give rise to homology classes

$$\widetilde{\lambda}^+(\mathbb{G}), \widetilde{\lambda}^-(\mathbb{G}) \in \widetilde{\mathrm{GH}}(\mathbb{G}) \cong \widehat{\mathrm{HFL}}(-S^3, \Lambda) \otimes V^{n-\ell}, {}^3$$

where V is a vector space of dimension 2, and  $\ell$  is the number of components of  $\Lambda$ . (On the chain level,  $\widetilde{\mathrm{GC}}(\mathbb{G})$  corresponds to a stabilized version of  $\widehat{\mathrm{gCFL}}(-S^3,\Lambda)$ , whose homology is  $\widehat{\mathrm{HFL}}(-S^3,\Lambda)$ .) Ozsváth, Szabó, and Thurston prove that  $\widetilde{\lambda}^\pm$  are preserved by the homomorphisms on  $\widetilde{\mathrm{GH}}$  induced by Legendrian isotopy, which implies that they are Legendrian isotopy invariants; they are referred to as the GRID invariants. Similar statements hold for the minus flavor of the theory.

As explained in [MOSzT07, OSSz15], one may in fact define a filtered chain complex  $\widetilde{\mathcal{GC}}(\mathbb{G})$  with the *Alexander filtration*, such that the associated graded object of  $\widetilde{\mathcal{GC}}(\mathbb{G})$  is  $\widetilde{\mathrm{GC}}(\mathbb{G})$ . This corresponds to the filtered chain complex  $\widehat{\mathrm{CFL}}$ , whose associated graded object is  $\widehat{\mathrm{gCFL}}$ . The filtered chain homotopy type of  $\widetilde{\mathcal{GC}}(\mathbb{G})$  is an invariant of the smooth link type of  $\Lambda$  and the grid size of  $\mathbb{G}$  (which controls the amount of stabilization). The associated spectral sequence

$$(1.1) \qquad \widetilde{\mathrm{GH}}(\mathbb{G}) \rightrightarrows \mathrm{H}_*(\widetilde{\mathcal{GC}}(\mathbb{G}))$$

then corresponds to the spectral sequence

$$\widehat{\mathrm{HFL}}(-S^3,\Lambda) \rightrightarrows \widehat{\mathrm{HF}}(-S^3),$$

which is an invariant of the smooth link type of  $\Lambda$ .

The idea of the spectral GRID invariants is to consider the homology classes that  $\mathbf{x}^{\pm}(\mathbb{G})$  represent in the spectral sequence in (1.1). These may be defined recursively. For concreteness, let  $(E^i, \delta_i)$  be the  $i^{\text{th}}$  page in the spectral sequence, such that  $\widetilde{\mathrm{GH}}(\mathbb{G})$  is the  $E^1$ -page, and let  $\widetilde{\lambda}_1^+(\mathbb{G}) = \widetilde{\lambda}^+(\mathbb{G})$ . Suppose that  $\delta_i(\widetilde{\lambda}_i^+(\mathbb{G})) = 0$ ; then we obtain a homology class  $\widetilde{\lambda}_{i+1}^+(\mathbb{G}) = [\widetilde{\lambda}_i^+(\mathbb{G})] \in E^{i+1}$ . If at

<sup>&</sup>lt;sup>3</sup>A Legendrian link in  $(\mathbb{R}^3, \xi_{\text{std}})$  can be naturally viewed as a Legendrian link in the standard contact  $S^3$ . We follow [OSzT08] and view the Legendrian invariants as living in  $\widehat{HFL}(S^3, m(\Lambda)) \cong \widehat{HFL}(-S^3, \Lambda)$ .

any stage  $\delta_i(\widetilde{\lambda}_i^+(\mathbb{G})) \neq 0$ , then we may instead record the integer i, which we will call  $n^+(\mathbb{G}) \in \mathbb{Z}^+$  and the process terminates. Let  $n^+(\mathbb{G}) = \infty$  if  $\delta_i(\widetilde{\lambda}_i^+) = 0$  for all  $i \geq 1$ . Considering the invariant  $\widetilde{\lambda}^-(\mathbb{G}) = [\mathbf{x}^-(\mathbb{G})]$ , we likewise obtain invariants  $n^-(\mathbb{G})$  and  $\widetilde{\lambda}_i^-(\mathbb{G})$ . With some work, that  $n^{\pm}$  and  $\widetilde{\lambda}_i^{\pm}$  are indeed invariants of the Legendrian link  $\Lambda$  can be deduced by extending certain proofs in [OSzT08], which we will carry out later:

**Theorem 1.2.** Suppose that  $\mathbb{G}$  and  $\mathbb{G}'$  are two grid diagrams that represent the same Legendrian link  $\Lambda \subset (S^3, \xi_{\text{std}})$ . Then  $n^+(\mathbb{G}) = n^+(\mathbb{G}')$  and  $n^-(\mathbb{G}) = n^-(\mathbb{G})$ , and there exist filtered chain homomorphisms

$$\Phi\colon \widetilde{\mathcal{GC}}(\mathbb{G}) \to \widetilde{\mathcal{GC}}(\mathbb{G}'), \qquad \Phi'\colon \widetilde{\mathcal{GC}}(\mathbb{G}') \to \widetilde{\mathcal{GC}}(\mathbb{G}),$$

such that the induced maps  $\Phi_i \colon E^i(\widetilde{\mathcal{GC}}(\mathbb{G})) \to E^i(\widetilde{\mathcal{GC}}(\mathbb{G}'))$  and  $\Phi_i' \colon E^i(\widetilde{\mathcal{GC}}(\mathbb{G}')) \to E^i(\widetilde{\mathcal{GC}}(\mathbb{G}))$  on the pages of the associated spectral sequences satisfy

$$\Phi_i(\widetilde{\lambda}_i^{\pm}(\mathbb{G})) = \widetilde{\lambda}_i^{\pm}(\mathbb{G}'), \qquad \Phi_i'(\widetilde{\lambda}_i^{\pm}(\mathbb{G})) = \widetilde{\lambda}_i^{\pm}(\mathbb{G}'),$$

for each  $1 \leq i \leq n^{\pm}(\mathbb{G})$ . The fact that  $n^{\pm}(\mathbb{G}) = n^{\pm}(\mathbb{G}')$  means that we have invariants  $n^{\pm}(\Lambda) \in \mathbb{Z}^+ \cup \{\infty\}$  of the Legendrian link type of  $\Lambda$ . Moreover, since the (non)vanishing of  $\widetilde{\lambda}_i^{\pm}(\mathbb{G})$  depends only on  $\Lambda$ , we may view these also as invariants of the Legendrian link type of  $\Lambda$ , and slightly abuse notation to denote them by  $\widetilde{\lambda}_i^{\pm}(\Lambda) \in E^i(\widetilde{\mathrm{CFL}}(-S^3,\Lambda))$ .

Remark 1.3. Versions of the maps  $\Phi$  and  $\Phi'$  first appear in [OSzT08, Lemmas 6.5 and 6.6], and special cases of Theorem 1.2 are spelled out in [NOT08] and [OSSz15]. Specifically, [NOT08, Theorem 2] distinguishes two Legendrian links  $\Lambda$  and  $\Lambda'$  with the same smooth type and classical invariants by showing that  $n^+(\Lambda) = 1$  while  $n^+(\Lambda') > 1$ . For technical reasons, our formulation of  $\Phi$  and  $\Phi'$  differs slightly from [OSzT08, OSSz15] when grid (de)stabilizations are involved, and our language in homological algebra also differs from [OSSz15, Chapter 14].

Remark 1.4. The idea to obtain refined information about a contact-geometric invariant in knot Floer homology by considering its homology classes in a spectral sequence is also found in [KMVW19]. However, the filtration (and hence the spectral sequence) that we consider in the present article is distinct from that in [KMVW19].

**Remark 1.5.** As shown in [OSzT08], the invariant  $\tilde{\lambda}^+(\Lambda)$  is preserved under negative stabilizations of Legendrian links, implying that it is an invariant of the transverse push-off of  $\Lambda$ . Similarly,  $\tilde{\lambda}_i^+$  can also be shown to be transverse invariants by a minor extension of the proof of Theorem 1.2, even though we do not pursue this further in the present article.

Moreover, there is a version of Theorem 1.2 for the *minus* flavor of knot Floer homology, whose proof is also similar. For simplicity, we will work solely with the *tilde* flavor.

1.2. **Obstructions.** Baldwin, Lidman, and the fifth author [BLW22] prove that, if there exists a decomposable Lagrangian cobordism  $L: \Lambda_- \to \Lambda_+$ , then there exists a homomorphism  $\Phi_L: \widetilde{\mathrm{HFL}}(-S^3, \Lambda_+) \to \widetilde{\mathrm{HFL}}(-S^3, \Lambda_-)$  that sends  $\widetilde{\lambda}^{\pm}(\Lambda_+)$  to  $\widetilde{\lambda}^{\pm}(\Lambda_-)$ . In other words:

**Theorem 1.6** ([BLW22, Theorem 1.2]). Suppose that  $\Lambda_-$  and  $\Lambda_+$  are Legendrian links in  $(\mathbb{R}^3, \xi_{std})$ , such that

- $\tilde{\lambda}^+(\Lambda_+) = 0$  and  $\tilde{\lambda}^+(\Lambda_-) \neq 0$ ; or
- $\widetilde{\lambda}^-(\Lambda_+) = 0$  and  $\widetilde{\lambda}^-(\Lambda_-) \neq 0$ .

Then there does not exist a decomposable Lagrangian cobordism from  $\Lambda_-$  to  $\Lambda_+$ .

As mentioned above, the main goal of this article is to prove the following extension of this result:

**Theorem 1.7.** Suppose that  $\Lambda_-$  and  $\Lambda_+$  are Legendrian links in  $(\mathbb{R}^3, \xi_{\text{std}})$ , such that

- $n^{+}(\Lambda_{+}) > n^{+}(\Lambda_{-}); or$
- $\widetilde{\lambda}_{i}^{+}(\Lambda_{+}) = 0$  and  $\widetilde{\lambda}_{i}^{+}(\Lambda_{-}) \neq 0$  for some  $1 \leq i \leq \min\{n^{+}(\Lambda_{-}), n^{+}(\Lambda_{+})\}$ ; or  $n^{-}(\Lambda_{+}) > n^{-}(\Lambda_{-})$ ; or
- $\widetilde{\lambda}_i^-(\Lambda_+) = 0$  and  $\widetilde{\lambda}_i^-(\Lambda_-) \neq 0$  for some  $1 \leq i \leq \min\{n^+(\Lambda_-), n^+(\Lambda_+)\}$ .

Then there does not exist a decomposable Lagrangian cobordism from  $\Lambda_-$  to  $\Lambda_+$ .

Remark 1.8. Restricting to decomposable Lagrangian concordances (i.e. the case that the genus g=0), the hypotheses involving  $\tilde{\lambda}_1^+$  are covered by [BS18, BS21] and [GJ19, Corollary 1.4]. In addition, for g = 0, [GJ19, Corollary 1.5] covers the hypothesis  $n^+(\Lambda_+) > n^+(\Lambda_-) = 1$ .

Specializing to the case where  $\Lambda_{-}$  is the undestabilizable Legendrian unknot, we obtain the following:

Corollary 1.9. Suppose that  $\Lambda$  is a Legendrian link in  $(\mathbb{R}^3, \xi_{\text{std}})$ , such that  $\widetilde{\lambda}_i^+(\Lambda) = 0$  for some  $1 \le i \le n^+(\Lambda)$ , or  $\widetilde{\lambda}_i^-(\Lambda) = 0$  for some  $1 \le i \le n^-(\Lambda)$ . Then there does not exist a decomposable Lagrangian filling of  $\Lambda$ .

1.3. Functoriality. Analogous to [BLW22], Theorem 1.7 follows from the following theorem, which states that the spectral GRID invariants satisfy a weak functoriality under decomposable Lagrangian cobordisms, in the style of Theorem 1.2.

**Theorem 1.10.** Suppose that  $\mathbb{G}_-$  and  $\mathbb{G}_+$  are two grid diagrams that represent Legendrian links  $\Lambda_-$  and  $\Lambda_+$  in  $(\mathbb{R}^3, \xi_{\mathrm{std}})$  respectively. Suppose that there exists a decomposable Lagrangian cobordism L from  $\Lambda_{-}$  to  $\Lambda_{+}$ . Then there exists a filtered chain homomorphism

$$\Psi \colon \widetilde{\mathcal{GC}}(\mathbb{G}_+) \to \widetilde{\mathcal{GC}}(\mathbb{G}_-) \left[ -\chi(L), \frac{|\Lambda_+| - |\Lambda_-| - \chi(L)}{2} \right],$$

such that the induced maps  $\Psi_i : E^i(\widetilde{\mathcal{GC}}(\mathbb{G}_+)) \to E^i(\widetilde{\mathcal{GC}}(\mathbb{G}_-))$  on the pages of the associated spectral sequences satisfy

$$\Psi_i(\widetilde{\lambda}_i^{\pm}(\mathbb{G}_+)) = \widetilde{\lambda}_i^{\pm}(\mathbb{G}_-),$$

for each  $1 \leq i \leq n^{\pm}(\mathbb{G})$ . Here,  $|\Lambda_{+}|$  is the number of components of  $\Lambda_{+}$ .

Throughout the paper, we denote by  $\mathbb{C}[M,A]$  the filtered chain complex obtained from  $\mathbb{C}$  by shifting the homological (Maslov) grading up by M and shifting the (Alexander) filtration level up by A; in other words,  $\mathcal{C}[M, A]_{(0,0)} = \mathcal{C}_{(-M,-A)}$ .

As in [BLW22], we believe but do not prove  $\Psi$  to be the functorial map of Zemke [Zem19] associated to a certain decorated link cobordism.

1.4. Computation. The invariants  $n^{\pm}$  and  $\tilde{\lambda}_{i}^{\pm}$  can be computed algorithmically and directly from the filtered chain complex  $\widetilde{\mathcal{GC}}(\mathbb{G})$ , without the need to compute each page of the spectral sequence. Using some homological algebra, we prove in Section 5 the following proposition:

**Proposition 1.11.** Let  $\mathbb{G}$  be a grid diagram, and let A be the Alexander filtration level of  $\mathbf{x}^+(\mathbb{G})$ . Then

•  $n^+(\mathbb{G}) = i < \infty$  if and only if i is the smallest number such that

$$(1.12) [\widetilde{\partial} \mathbf{x}^{+}(\mathbb{G})] \neq 0 \in \mathrm{H}_{*}(\mathcal{F}_{A-1}\widetilde{\mathcal{GC}}(\mathbb{G})/\mathcal{F}_{A-i-1}\widetilde{\mathcal{GC}}(\mathbb{G}));$$

and

• Supposing  $i < n^+(\mathbb{G})$ , then  $\widetilde{\lambda}_i^+(\mathbb{G}) = 0$  if and only if

$$[\mathbf{x}^{+}(\mathbb{G})] = 0 \in \mathrm{H}_{*}(\mathcal{F}_{A+i-1}\widetilde{\mathcal{GC}}(\mathbb{G})/\mathcal{F}_{A-1}\widetilde{\mathcal{GC}}(\mathbb{G})).$$

(Note that  $\widetilde{\partial}$  is the total differential of  $\widetilde{\mathcal{GC}}(\mathbb{G})$ .) Analogous statements hold for  $n^-(\mathbb{G})$  and  $\widetilde{\lambda}_i^-(\mathbb{G})$ .

**Remark 1.14.** [NOT08, Proposition 1] and [GJ19, Corollary 1.5] mention homomorphisms  $\widehat{\delta}_i \colon \widehat{\mathrm{HFL}}_d(S^3, L, s) \to \widehat{\mathrm{HFL}}_d(S^3, L, s - i)$ . When i = 1, we have that (1.12) is satisfied if and only if  $\widehat{\delta}_i([\mathbf{x}^+(\mathbb{G})]) = 0$ , since

$$H_*(\mathcal{F}_{A-1}\widetilde{\mathcal{GC}}(\mathbb{G})/\mathcal{F}_{A-2}\widetilde{\mathcal{GC}}(\mathbb{G})) \cong \widetilde{GH}_{A-1}(\mathbb{G}).$$

However, for i > 1, only one of the two implications holds in general.

Given Proposition 1.11, the invariants  $n^{\pm}$  and  $\widetilde{\lambda}_{i}^{\pm}$  can be computed using a modified version of the "zigzag" complex in [NOT08, Section 4]. Note that, since the Alexander filtration on  $\widetilde{\mathcal{GC}}(\mathbb{G})$  is bounded, the spectral sequence collapses in finitely many pages, allowing one to determine when  $n^{\pm}(\mathbb{G}) = \infty$ . The authors [JPS<sup>+</sup>23] have implemented the algorithm described in this subsection in Python.

- 1.5. **Effectiveness.** In [BLW22],  $\tilde{\lambda}^{\pm}$  are shown to be effective in obstructing decomposable Lagrangian cobordisms, meaning that they provide obstructions beyond the classical invariants. Precisely, for every  $g \geq 0$ , there exist Legendrian knots  $\Lambda_{-}$  and  $\Lambda_{+}$ , such that
  - $tb(\Lambda_+) tb(\Lambda_-) = 2g$ ;
  - $r(\Lambda_+) = r(\Lambda_-);$
  - There exists a smooth cobordism of genus g between  $\Lambda_{-}$  and  $\Lambda_{+}$ ; but
  - $\tilde{\lambda}^+(\Lambda_+) = 0$  and  $\tilde{\lambda}^-(\Lambda_-) \neq 0$ , implying that there does not exist a decomposable Lagrangian cobordism from  $\Lambda_-$  to  $\Lambda_+$ .

Since  $\tilde{\lambda}^{\pm} = \tilde{\lambda}_{1}^{\pm}$ , the invariants  $\tilde{\lambda}_{i}^{\pm}$  are indeed effective. In view of Theorem 1.6 and Remark 1.8, it would be interesting to ask:

- For g > 0, do there exist examples where  $n^{\pm}$ , or  $\tilde{\lambda}_i^{\pm}$  with  $i \geq 2$ , obstruct decomposable Lagrangian cobordisms that  $\tilde{\lambda}^{\pm}$  could not?
- For g = 0, do there exist examples where  $n^{\pm}$  with  $n^{+}(\Lambda_{+}) \geq 2$ , or  $\tilde{\lambda}_{i}^{\pm}$  with  $i \geq 2$ , obstruct decomposable Lagrangian cobordisms that  $\tilde{\lambda}^{\pm}$  could not?

While it seems likely that the answer to both questions is in the affirmative, the authors have not yet been able to answer the first question. Below, we provide an example that answers the second question in the affirmative.

**Example 1.15.** The pretzel knot  $P(-4, -3, 3) = m(10_{140})$  has three undestabilizable Legendrian representatives  $\Lambda_1$ ,  $-\Lambda_1$ , and  $\Lambda_2$ , as in the Legendrian knot atlas by Chongchitmate and Ng [CN13]. Using [JPS<sup>+</sup>23], we have computed:

$$\widetilde{\lambda}_{1}^{+}(\Lambda_{1}) \neq 0, \qquad n^{+}(\Lambda_{1}) = 1, \qquad \widetilde{\lambda}_{1}^{-}(\Lambda_{1}) \neq 0, \qquad n^{-}(\Lambda_{1}) = \infty; 
\widetilde{\lambda}_{1}^{+}(-\Lambda_{1}) \neq 0, \qquad n^{+}(-\Lambda_{1}) = \infty, \qquad \widetilde{\lambda}_{1}^{-}(-\Lambda_{1}) \neq 0, \qquad n^{-}(-\Lambda_{1}) = 1; 
\widetilde{\lambda}_{1}^{+}(\Lambda_{2}) \neq 0, \qquad n^{+}(\Lambda_{2}) = 1, \qquad \widetilde{\lambda}_{1}^{-}(\Lambda_{2}) \neq 0, \qquad n^{-}(\Lambda_{2}) = 1.$$

<sup>&</sup>lt;sup>4</sup>In [CN13], they are labeled  $L_1$ ,  $-L_1$ , and  $L_2$ ; we continue to use  $\Lambda$  for Legendrian links, for consistence. Note also that our  $-\Lambda_1$  corresponds to  $\Lambda_1$  in [GJ19].

Golla and Juhász [GJ19, Proposition 1.6] show that there is no decomposable (in fact, regular<sup>5</sup>) Lagrangian concordance  $L: \Lambda_2 \to -\Lambda_1$ , using the fact that  $n^+(-\Lambda_1) > n^+(\Lambda_2) = 1$ . This is based on a computation of  $\hat{\delta}_1(\tilde{\lambda}^+(-\Lambda_1)) = 0$  and  $\hat{\delta}_1(\tilde{\lambda}^+(\Lambda_2)) \neq 0$  in [NOT08] using the computer program [NOT07].<sup>6</sup> While not directly stated in [GJ19], their result on  $n^+$  also implies that there is no decomposable Lagrangian cobordism  $L: \Lambda_1 \to -\Lambda_1$ .

The above is recovered by Theorem 1.7. In addition, by considering  $n^-$ , Theorem 1.7 also implies that there is no decomposable Lagrangian cobordism  $L\colon -\Lambda_1 \to \Lambda_1$  or  $L\colon \Lambda_2 \to \Lambda_1$ , which is previously unknown.

Finally, note that none of the obstructions above can be obtained by considering only linearized Legendrian contact homology (LCH), since  $\Lambda_1$ ,  $-\Lambda_1$ , and  $\Lambda_2$  all have the same linearized LCH. This eliminates the most tractable approach to using the Chekanov–Eliashberg DGA from symplectic field theory to obtain an obstruction.

1.6. **Organization.** In Section 2, we provide the necessary background on Legendrian links, Lagrangian cobordisms, link Floer homology, the GRID invariants, and filtered chain complexes and their associated spectral sequences. Next, in Section 3, we define the spectral invariants in detail, and provide a detailed proof that they are preserved by grid commutation and (de)stabilization, which implies that they are Legendrian invariants. In Section 4, we then show that the spectral invariants are preserved under pinches and (the reverse of) birth moves, proving Theorem 1.7. Finally, we prove Proposition 1.11 in Section 5, establishing an algorithm to compute the spectral invariants directly from the filtered chain complex.

Acknowledgments. This work is the result of the 2022 Summer Hybrid Undergraduate Research (SHUR) program at Dartmouth College, and the authors thank Dartmouth for the support. IP was partially supported by NSF CAREER Grant DMS-2145090. MW was partially supported by NSF Grant DMS-2238131 (previously DMS-2039688). The SHUR program was also partially supported by these NSF grants. Part of the research was conducted while MW was at Louisiana State University, and he thanks LSU for the support.

# 2. Preliminaries

2.1. **Legendrian knots and Lagrangian cobordisms.** In this section, we review the basics of Legendrian knots and Lagrangian cobordisms.

Recall that a smooth link  $\Lambda \in \mathbb{R}^3$  is called *Legendrian* if it is everywhere tangent to the standard contact structure on  $\mathbb{R}^3$ ,

$$\xi_{\rm std} = \ker(\alpha_{\rm std}), \quad \alpha_{\rm std} = dz - y \, dx.$$

Two Legendrian links are Legendrian isotopic if they are isotopic through a family of Legendrian links.

A Legendrian link can be represented by its front diagram, or front projection, the projection of the link onto the xz-plane. In a front diagram, strand crossing information is encoded by the slopes of the strands: strands with lower slope pass over strands with higher slope. See Figure 1 for an example.

Two Legendrian front diagrams represent Legendrian-isotopic links if the diagrams can be related by a sequence of Legendrian planar isotopies (isotopies that preserve left and right cusps) and Legendrian Reidemeister moves. Legendrian Reidemeister moves are the first three diagrams in Figure 2 and their mirror reflections.

<sup>&</sup>lt;sup>5</sup>An exact Lagrangian cobordism L is regular if the Liouville vector field is tangent to L.

<sup>&</sup>lt;sup>6</sup>Note that  $L_1$  and  $L_2$  in [NOT08] correspond to  $-\Lambda_1$  and  $\Lambda_2$  respectively. There is also a newer program [MQR<sup>+</sup>19] that provides bug fixes and improvements in computational speed to [NOT07].




Figure 1. An example of a front projection.




FIGURE 2. The moves on front projections that correspond to elementary cobordisms. The first three moves are Legendrian Reidemeister moves; they preserve the Legendrian knot type and correspond to Lagrangian cylinders. The fourth move is called a pinch, and the fifth move is called a birth. The vertical and horizontal reflecions of the second move are also allowed.

The two classical Legendrian link invariants are the Thurston-Bennequin number  $tb(\Lambda)$  and the rotation number  $r(\Lambda)$ . These can be computed from an oriented front diagram D via the relations

$$tb(\Lambda) = wr(D) - \frac{1}{2}(c_{+}(D) + c_{-}(D)), \qquad r(\Lambda) = \frac{1}{2}(c_{-}(D) - c_{+}(D)),$$

where wr(D) is the writhe of the diagram, and  $c_{-}(D)$  and  $c_{+}(D)$  are the number of downward and upward cusps, respectively.

The symplectization of  $(\mathbb{R}^3, \xi_{\text{std}})$  is the symplectic 4-manifold

$$(\mathbb{R}_t \times \mathbb{R}^3, d(e^t \alpha_{\mathrm{std}})).$$

A Lagrangian cobordism from  $\Lambda_- \subset (\mathbb{R}^3, \xi_{\mathrm{std}})$  to  $\Lambda_+ \subset (\mathbb{R}^3, \xi_{\mathrm{std}})$  is an oriented, embedded surface  $L \subset \mathbb{R}^t \times \mathbb{R}^3$  such that

• L is Lagrangian, i.e.  $d(e^t \alpha_{\rm std})|_L \equiv 0$ ;

• L has cylindrical ends, i.e. for some T > 0,

$$L \cap ((-\infty, -T) \times \mathbb{R}^3) = (-\infty, -T) \times \Lambda_-,$$
  
$$L \cap ((T, \infty) \times \mathbb{R}^3) = (T, \infty) \times \Lambda_+,$$

and 
$$L \cap ([-T,T] \times \mathbb{R}^3)$$
 is compact.

A Lagrangian cobordism is *exact* if there exists a function  $f: L \to \mathbb{R}$  that is constant (and not just locally constant) on each of the two cylindrical ends, and satisfies

$$(e^t \alpha_{\rm std})|_L = df.$$

A Lagrangian concordance is a Lagrangian cobordism of genus zero, which is immediately exact. If there exists a Lagrangian cobordism L from  $\Lambda_-$  to  $\Lambda_+$ , Chantraine [Cha10] proves that the classical invariants of the two links are related by

$$tb(\Lambda_{+}) - tb(\Lambda_{-}) = -\chi(L)$$
 and  $r(\Lambda_{+}) - r(\Lambda_{-}) = 0$ ,

where  $\chi(L)$  is the Euler characteristic of L. This immediately implies that Lagrangian cobordism are not an equivalence relation. In fact, even Lagrangian concordance is not an equivalence relation [Cha15].

One important subclass of exact Lagrangian cobordisms is the class of decomposable Lagrangian cobordisms. Precisely, refer to Figure 2: If  $\Lambda_{-}$  and  $\Lambda_{+}$  are Legendrian links such that

- $\Lambda_{-}$  and  $\Lambda_{+}$  are Legendrian isotopic, as in the first three diagrams;
- $\Lambda_{-}$  is obtained from  $\Lambda_{+}$  by a pinch move, as in the fourth diagram; or
- $\Lambda_+$  is obtained from  $\Lambda_-$  by a Legendrian birth, i.e.  $\Lambda_+$  is the disjoint union of  $\Lambda_-$  with an unlinked component that is the undestabilizable Legendrian unknot, as in the fifth diagram:

then there exists an elementary exact Lagrangian cobordism  $L: \Lambda_- \to \Lambda_+$ , by work of Bourgeois, Sabloff, Traynor [BST15], Chantraine [Cha10], Dimitroglou Rizell [Dim16], and Ekholm, Honda, and Kálmán [EHK16]. Note that, topologically, elementary exact Lagrangian cobordisms are annuli, saddles, and cups, respectively. A Lagrangian cobordism is decomposable if it is isotopic through exact Lagrangian cobordisms to a composition of elementary exact Lagrangian cobordisms. In the smooth category, every link cobordism is decomposable into elementary cobordisms; whether every exact Lagrangian cobordism is decomposable remains a major open question.

2.2. **Knot Floer homology and the GRID invariants.** In this section, we review some basics of grid homology, following the conventions in [OSSz15].

A grid diagram  $\mathbb{G}$  is an  $m \times m$  grid on the plane, along with two sets of markers

$$\mathbb{O} = \{O_1, \dots, O_m\}, \qquad \mathbb{X} = \{X_1, \dots, X_m\},\$$

such that there is exactly one O and exactly one X in each row, as well as in each column, and no square of the grid contains more than one marking. The number m is called the *grid number* of  $\mathbb{G}$ .

A grid diagram  $\mathbb{G}$  specifies a link  $L \subset \mathbb{R}^3$  as follows. Draw oriented segments connecting X's to O's in each column, and O's to X's in each row, and require that vertical segments cross above horizontal ones. We say  $\mathbb{G}$  is a grid diagram for L. Conversely, every link L in  $\mathbb{R}^3$  can be represented by a grid diagram. By a theorem of Cromwell [Cro95], two grid diagrams represent the same link if and only if they are related by a sequence of commutations, in which two eligible adjacent rows or columns commute, stabilizations, in which a  $1 \times 1$  square with an O (resp. X) marker is replaced by a  $2 \times 2$  square with two diagonal O markers and an O marker), creating a new row and a new column, and destabilizations, the inverse operations. Following [OSSz15], we classify (de)stabilizations by the marker type and the location of the empty cell in the  $2 \times 2$  square; for example, a stabilization of type X:SE results in a  $2 \times 2$  square with an

empty southeastern cell, and accordingly an X in the northwestern cell and O's in the northeastern and southwestern cells.

To a grid diagram  $\mathbb{G}$ , we associate a graded, filtered chain complex  $\widetilde{\mathcal{GC}}(\mathbb{G})$  over  $\mathbb{F}_2 = \mathbb{Z}/2$  whose filtered chain homotopy type is an invariant of the isotopy type of L. Before we do this, we introduce a bit more notation. First, we will think of a grid diagram as a diagram on a torus, by identifying the left and right edges, as well as the top and bottom edges of the grid. The horizontal arcs of the grid result in a set of circles  $\boldsymbol{\alpha} = \{\alpha_1, \dots, \alpha_m\}$ , indexed from bottom to top, and the vertical ones result in a set of circles  $\boldsymbol{\beta} = \{\beta_1, \dots, \beta_m\}$ , indexed from left to right.

As an  $\mathbb{F}_2$ -module,  $\widetilde{\mathcal{GC}}(\mathbb{G})$  is generated by *grid states*, i.e. bijections between horizontal and vertical circles. Geometrically, a grid state is an m-tuple of points  $\mathbf{x} = \{x_1, \dots, x_m\}$  with one point on each horizontal circle and one on each vertical circle. The set of grid states for a grid diagram  $\mathbb{G}$  is denoted  $S(\mathbb{G})$ .

Before we define the differential, we first define a partial ordering of points in  $\mathbb{R}^2$  given by  $(x_1, y_1) < (x_2, y_2)$  if  $x_1 < x_2$  and  $y_1 < y_2$ . For any two sets  $P, Q \subset \mathbb{R}^2$ , define

$$\mathcal{I}(P,Q) = \# \{ (p,q) \in P \times Q \mid p < q \}.$$

Next, we define the symmetrized function,

$$\mathcal{J}(P,Q) = \frac{\mathcal{I}(P,Q) + \mathcal{I}(Q,P)}{2}.$$

For any state  $\mathbf{x} \in S(\mathbb{G})$ , we can define it uniquely as integer points on the  $[0, m) \times [0, m) \subset \mathbb{R}^2$ . Then, representing  $\mathbb{X}$  and  $\mathbb{O}$  as half-integer points uniquely on the same subset, we can define the Maslov and Alexander functions  $M(\mathbf{x})$  and  $A(\mathbf{x})$  as follows:

$$M(\mathbf{x}) = M_{\mathbb{O}}(\mathbf{x}) = \mathcal{J}(\mathbf{x}, \mathbf{x}) - 2\mathcal{J}(\mathbf{x}, \mathbb{O}) + \mathcal{J}(\mathbb{O}, \mathbb{O}) + 1,$$
  

$$M_{\mathbb{X}}(\mathbf{x}) = \mathcal{J}(\mathbf{x}, \mathbf{x}) - 2\mathcal{J}(\mathbf{x}, \mathbb{X}) + \mathcal{J}(\mathbb{X}, \mathbb{X}) + 1,$$
  

$$A(\mathbf{x}) = \frac{1}{2} \left( M_{\mathbb{O}}(\mathbf{x}) - M_{\mathbb{X}}(\mathbf{x}) \right) - \frac{m - l}{2},$$

where l is the number of link components in the Legendrian link corresponding to the grid  $\mathbb{G}$ .

Given two grid states  $\mathbf{x}, \mathbf{y} \in S(\mathbb{G})$ , let  $\text{Rect}(\mathbf{x}, \mathbf{y})$  denote the space of rectangles in  $\mathbb{G}$  with the following properties. First,  $\text{Rect}(\mathbf{x}, \mathbf{y})$  is empty if  $\mathbf{x}$  and  $\mathbf{y}$  do not agree at exactly m-2 points. An element  $r \in \text{Rect}(\mathbf{x}, \mathbf{y})$  is an embedded rectangle with right angles, such that:

- $\partial r$  lies on the union of horizontal and vertical circles;
- The vertices of r are exactly the points in  $\mathbf{x} \triangle \mathbf{y}$ , where  $\triangle$  denotes the symmetric difference; and
- $\partial(\partial r \cap \beta) = \mathbf{x} \mathbf{y}$ , in the orientation induced by r.

Given  $r \in \text{Rect}(\mathbf{x}, \mathbf{y})$ , we say that r goes from  $\mathbf{x}$  to  $\mathbf{y}$ . Observe that  $\text{Rect}(\mathbf{x}, \mathbf{y})$  consists of either zero or two rectangles. We say a rectangle  $r \in \text{Rect}(\mathbf{x}, \mathbf{y})$  is empty if  $\mathbf{x} \cap \text{Int}(r) = \mathbf{y} \cap \text{Int}(r) = \emptyset$ . We denote the set of empty rectangles from  $\mathbf{x}$  to  $\mathbf{y}$  by  $\text{Rect}^{\circ}(\mathbf{x}, \mathbf{y})$ .

For any two states  $\mathbf{x}, \mathbf{y}$  with a rectangle  $r \in \text{Rect}(\mathbf{x}, \mathbf{y})$ , one could compute that

(2.1) 
$$M(\mathbf{x}) - M(\mathbf{y}) = 1 - 2\#(r \cap \mathbb{O}) + 2\#(\text{Int}(r) \cap \mathbf{x}),$$

(2.2) 
$$A(\mathbf{x}) - A(\mathbf{y}) = \#(r \cap \mathbb{X}) - \#(r \cap \mathbb{O}).$$

<sup>&</sup>lt;sup>7</sup>In [BLW22], the  $\alpha$ -circles are the vertical ones, and  $\beta$ -circles the horizontal ones. Here, we instead follow the convention of [OSSz15]. However, in later sections, we opt to follow [BLW22] in drawing multidiagrams (e.g. Figure 3 and Figure 8) with multiple horizontal curves; as a result, they have multiple  $\alpha$ -curves rather than multiple  $\beta$ -curves.

The differential on  $\widetilde{\mathcal{GC}}(\mathbb{G})$  is defined on generators by

$$\widetilde{\partial}_{\mathbb{O}}(\mathbf{x}) = \sum_{\mathbf{y} \in S(\mathbb{G})} \sum_{\substack{r \in \operatorname{Rect}^{\circ}(\mathbf{x}, \mathbf{y}) \\ r \cap \mathbb{O} = \emptyset}} \mathbf{y}.$$

Using (2.1) and (2.2), we examine how the differential interacts with the Maslov and Alexander functions. First, note that  $M(\mathbf{x}) - M(\mathbf{y}) = 1$  for each  $\mathbf{y}$  in the summation, which implies that  $\widetilde{\partial}_{\mathbb{Q}}$  drops the Maslov grading by 1. Second, we see that  $A(\mathbf{x}) - A(\mathbf{y}) \geq 0$  for all  $\mathbf{y}$  in the summation. Thus, M induces a  $\mathbb{Z}$ -grading on  $\widetilde{\mathcal{GC}}(\mathbb{G})$ , and A induces a  $\mathbb{Z}$ -filtration. We discuss filtered complexes further in Section 2.3.

The homology  $\widetilde{\mathrm{GH}}(\mathbb{G}) = H_*(\mathrm{gr}(\widetilde{\mathcal{GC}}(\mathbb{G})))$  of the associated graded object is (almost) an invariant of the underlying link, in the following sense. If  $\mathbb{G}$  is a grid of size m for an l-component link L, then we have

$$\widetilde{\mathrm{GH}}(\mathbb{G}) \cong \widehat{\mathrm{HFL}}(L) \otimes W^{\otimes (m-l)}$$
,

where W is a two-dimensional bigraded vector space with one generator in bigrading (0,0) and another in bigrading (-1,-1). Alternatively, a combinatorial proof of the invariance of  $\widetilde{\mathrm{GH}}$  that does not appeal to holomorphic Heegaard Floer theory is given by defining combinatorial filtered chain homotopy equivalences between  $\widetilde{\mathcal{GC}}(\mathbb{G}_1)$  and  $\widetilde{\mathcal{GC}}(\mathbb{G}_2)$  when  $\mathbb{G}_1$  and  $\mathbb{G}_2$  differ by a commutation or (de)stabilization.

Given a planar grid  $\mathbb{G}$ , we construct a Legendrian link in the following manner. First, create the oriented link specified by the grid  $\mathbb{G}$ . The projection of this link onto the grid has corners that can be classified into four types: northeast, northwest, southwest, and southeast. First, we smooth all of the northwest and southeast corners of the projection, and turn the northeast and southwest corners into cusps. Next, we rotate the diagram 45 degrees clockwise. Now, since all the vertical strands cross over the horizontal ones and this convention is opposite to the convention for Legendrian front projections, we now flip all the crossings. This gives a Legendrian link  $\Lambda(\mathbb{G})$  whose smooth type is the mirror of the smooth link associated to  $\mathbb{G}$ . Similar to the smooth case, every Legendrian link in  $(\mathbb{R}^3, \xi_{\text{std}})$  can be represented by a grid diagram. Two grid diagrams represent the same Legendrian link if and only if they are related by a sequence of commutations and (de)stabilizations of type X:SE and X:NW.

Given a grid diagram  $\mathbb{G}$ , the generator  $\mathbf{x}^+(\mathbb{G}) \in S(\mathbb{G})$  is the grid state composed of all the points directly northeast of the X's. Similarly, we define  $\mathbf{x}^-(\mathbb{G}) \in S(\mathbb{G})$  to be the grid state composed of all points directly southwest of the X's. Then, if a grid diagram  $\mathbb{G}$  of a Legendrian link  $\Lambda$  of l components, one can compute the gradings of the generators  $\mathbf{x}^{\pm}$  to be

(2.3) 
$$M(\mathbf{x}^{\pm}(\mathbb{G})) = tb(\Lambda) \mp r(\Lambda) + 1$$
$$A(\mathbf{x}^{\pm}(\mathbb{G})) = \frac{tb(\Lambda) \mp r(\Lambda) + l}{2}.$$

In [OSzT08], both  $[\mathbf{x}^+(\mathbb{G})]$  and  $[\mathbf{x}^-(\mathbb{G})]$  are shown to be cycles in the associated graded object  $gr(\widetilde{\mathcal{GC}}(\mathbb{G})) = \widetilde{GC}(\mathbb{G})$ ; moreover, these cycles are preserved by the filtered chain homotopy equivalences associated to commutations and (de)stabilizations of type X:SE and X:NW, which shows that they are invariants of the Legendrian link  $\Lambda$ .

2.3. Filtered chain complexes. The main algebraic structures that we study in this paper are filtered chain complexes, and the spectral sequences they induce.

A filtration on a chain complex C is a sequence of subcomplexes  $(\mathcal{F}_i C)_{i \in \mathbb{Z}}$  of C such that  $\mathcal{F}_i C \subset \mathcal{F}_j C$  whenever  $i \leq j \in Z$ . To be more precise, this is the definition of an "increasing" filtration. We will assume that our filtrations are bounded, which means that  $\mathcal{F}_s C = 0$  and  $\mathcal{F}_t C = C$ 

for some  $s \leq t \in \mathbb{Z}$ . A filtered complex  $C = (C, \mathcal{F})$  is a complex C equipped with a filtration  $\mathcal{F}$  of C. When the filtration is clear from context, we will often omit it.

A map of filtered complexes is a chain map  $f: C \to C'$  that respects the filtration in the sense that  $f(\mathcal{F}_i C) \subset \mathcal{F}_i C'$ .

Given a filtered complex C, we may construct the associated graded complex gr(C), which is defined to be

$$\operatorname{gr}(\mathcal{C}) = \bigoplus_{p \in \mathbb{Z}} \operatorname{gr}_p(\mathcal{C})$$

where

$$\operatorname{gr}_p(\mathcal{C}) = \mathcal{F}_p C / \mathcal{F}_{p-1} C$$
.

A map  $f: \mathcal{C} \to \mathcal{C}'$  of filtered complexes induces a map  $gr(f): gr(\mathcal{C}) \to gr(\mathcal{C}')$  of associated graded complexes in a natural way.

While the underlying modules of filtered complexes and their associated graded complexes are isomorphic (over a field), the same is not true in general when one considers their differentials. For example, it is not true in general that  $H_*(\operatorname{gr}(\mathcal{C})) \cong H_*(\mathcal{C})$ . Instead, what we can say is that there is a spectral sequence from the former group to the latter. A spectral sequence is a sequence of chain complexes  $(E^r)_{r\in\mathbb{N}\cup\{\infty\}}$  and isomorphisms  $H_*(E^r)\cong E^{r+1}$  for all  $r\in\mathbb{N}\cup\{\infty\}$ . The complex  $E^r$  comes equipped with a bigrading  $E^r=\bigoplus_{p,q}E^r_{p,q}$ .

Given a filtered complex  $\mathcal{C}$ , there is an induced spectral sequence with  $E^1_{p,q}\cong H_{p+q}(\operatorname{gr}_p(\mathcal{C}))$  and

Given a filtered complex  $\mathcal{C}$ , there is an induced spectral sequence with  $E_{p,q}^1 \cong H_{p+q}(\operatorname{gr}_p(\mathcal{C}))$  and  $E_{p,q}^{\infty} \cong \operatorname{gr}_p(H_{p+q}(\mathcal{C}))$ . We will summarize the relevant details, using the conventions from [Sta18, Section 012K] adjusted for homological (vs. cohomological) gradings. The filtration  $\mathcal{F}$  on  $\mathcal{C}$  induces a grading on each page  $E^r$  of the spectral sequence as a module; this is the grading corresponding to p in  $E_{p,q}^r$ . Similarly, q corresponds to the original homological grading on  $\mathcal{C}$ ; we will often suppress this grading for simplicity. As modules, we define

$$E_p^r(\mathcal{C}) \cong \frac{Z_p^r(\mathcal{C})}{B_p^r(\mathcal{C})},$$

where

$$Z_p^r(\mathcal{C}) \cong \frac{\mathcal{F}_pC \cap \partial^{-1}(\mathcal{F}_{p-r}C) + \mathcal{F}_{p-1}C}{\mathcal{F}_{p-1}C} \quad \text{and} \quad B_p^r(\mathcal{C}) \cong \frac{\mathcal{F}_pC \cap \partial(\mathcal{F}_{p+r-1}C) + \mathcal{F}_{p-1}C}{\mathcal{F}_{p-1}C}.$$

The differential  $d^r: E^r_p \to E^r_{p-r}$  is induced by  $\partial$  on the above formula for  $E^r$  as a quotient of sub-quotients of C. The only part of the spectral sequence structure induced by  $\mathcal{F}$  that we have not defined is the isomorphism  $H_*(E^r) \cong E^{r+1}$ ; this is mostly tedious but straightforward algebra. For more details, we refer the reader to [Wei94] or another book on homological algebra.

Since we are interested in the behavior of particular elements under spectral sequences, we want to define what it means to talk about the "class of  $x \in C$  on the r-th page". Given any nonzero  $x \in C$ , let p be the value of p for which  $x \in \mathcal{F}_p C \setminus \mathcal{F}_{p-1} C$  (note that this is the difference as sets and not the quotient). We may think of p the "filtration level" of x; such an integer always exists since  $\mathcal{F}$  is assumed to be bounded. Then, if  $x \in \mathcal{F}_p C \cap \partial^{-1}(\mathcal{F}_{p-r} C) + \mathcal{F}_{p-1} C$ , we define  $[x]^r$  to be the class of x in  $E_p^r$ . Note that, given some  $x \in \mathcal{F}_p C$ , it may not be the case that  $[x]^r$  is defined for all r. However, if  $[x]^r$  is defined, then  $[x]^s$  is also defined for all  $s \leq r$ , since  $\partial^{-1}(\mathcal{F}_{p-s} C) \supseteq \partial^{-1}(\mathcal{F}_{p-r} C)$ . In fact,  $[x]^{r+1} \in E_p^{r+1}$  is defined if and only if  $d^r[x]^r = 0$ , since  $\ker d_p^r = \partial^{-1}(B_{p-r}^r) \cap Z_p^r \cong Z_p^{r+1}$ . In Section 5, we will introduce an alternative way of thinking about the (non)vanishing of the class of an element in  $E^r$  that lends itself nicely to certain computations.

In later sections, our strategy to relate elements in the spectral sequences associated to two filtered complexes will be to relate representatives of those elements in the filtered chain complexes.

Thus, we need the following lemma, which explains how a filtered chain map induces maps on the spectral sequence:

**Lemma 2.4.** Let  $C = (C, \mathcal{F})$  and  $C' = (C', \mathcal{F}')$  be two filtered chain complexes, and let  $f: C \to C'$  be a filtered chain map. Then for each  $r \geq 1$ , the map f induces a chain map

$$E^r(f) \colon E^r(\mathcal{C}) \to E^r(\mathcal{C}').$$

Furthermore, if  $x \in \mathcal{F}_pC$  is an element with a well-defined class  $[x]^r \in E^r(\mathcal{C})$ , such that f(x) can be (non-uniquely) written as

$$f(x) = y_1 + y_2$$

where  $y_1 \in \mathcal{F}_p'C'$  and  $y_2 \in \mathcal{F}_{p-1}'C'$ , then  $[y_1]^r$  is well-defined, and

$$E^{r}(f)([x]^{r}) = [y_{1}]^{r}.$$

*Proof.* First, assuming we are given a map  $f: \mathcal{C} \to \mathcal{C}'$ , we will describe the map  $E^r(f)$ . Since  $E^r(\mathcal{C})$  and  $E^r(\mathcal{C}')$  are sub-quotients of  $\mathcal{C}$  and  $\mathcal{C}'$  respectively, one can check that  $f: \mathcal{C} \to \mathcal{C}'$  induces a map  $E^r(f): E^r(\mathcal{C}) \to E^r(\mathcal{C}')$ . On elements,  $E^r(f)$  is defined such that

$$E^{r}(f)([x]^{r}) = [f(x)]^{r}$$
.

Note that f is a filtered chain map and therefore commutes with the original differentials, i.e.  $f \circ \partial = \partial' \circ f$ . Therefore, we get that  $E^r(f)$  also commutes with the induced differentials, i.e.  $E^r(f) \circ d^r = (d')^r \circ E^r(f)$ . It can be shown further that  $E^r(f)$  is also the map induced by  $E^{r-1}(f)$  on homology (a property shared by all morphisms of spectral sequences).

Next, assume  $x \in \mathcal{F}_pC$  is an element with a well-defined class  $[x]^r \in E^r(\mathcal{C})$ , and that

$$f(x) = y_1 + y_2$$

for some  $y_1 \in \mathcal{F}_pC'$  and  $y_2 \in \mathcal{F}_{p-1}C'$ . Then

$$E^{r}(f)([x]^{r}) = [f(x)]^{r} = [y_1 + y_2]^{r} = [y_1]^{r}$$

by the definition of  $E^r(f)$  and the fact that  $y_2 \in \mathcal{F}_{p-1}C'$  is in the denominator of  $Z^r(C')$ .

## 3. Spectral GRID invariants

3.1. **Definition of the spectral GRID invariants.** Now, we have the necessary background to define our invariants.

**Definition 3.1.** Suppose that  $\mathbb{G}$  is a grid diagram, and let  $A = A(\mathbf{x}^+(\mathbb{G}))$ . We define  $n^+(\mathbb{G})$  to be the smallest integer i for which  $d_A^i[\mathbf{x}^+(\mathbb{G})]^i \neq 0 \in E_A^i$ , or  $\infty$  if  $d_A^i[\mathbf{x}^+(\mathbb{G})]^i = 0$  for all  $i \in \mathbb{Z}_{\geq 1}$ . We define  $n^-(\mathbb{G})$  analogously, replacing  $\mathbf{x}^+(\mathbb{G})$  by  $\mathbf{x}^-(\mathbb{G})$ .

**Definition 3.2.** Suppose that  $\mathbb{G}$  is a grid diagram. For each  $1 \leq i \leq n^+(\mathbb{G})$ , we define

$$\lambda_i^+(\mathbb{G}) = [\mathbf{x}^+(\mathbb{G})]^i \in E^i_{A(\mathbf{x}^+(\mathbb{G}))}, \qquad \lambda_i^-(\mathbb{G}) = [\mathbf{x}^-(\mathbb{G})]^i \in E^i_{A(\mathbf{x}^-(\mathbb{G}))}.$$

For the rest of this section, we will focus on proving Theorem 1.2, which states the invariance of  $n^{\pm}(\mathbb{G})$  and  $\widetilde{\lambda}_{i}(\mathbb{G})$  under the choice of  $\mathbb{G}$ ). This will allow us to denote them by  $n^{\pm}(\Lambda)$  and  $\widetilde{\lambda}_{i}(\Lambda)$ .

In [OSzT08, OSSz15], invariance of  $\tilde{\lambda}^{\pm}$  is proven by considering the isomorphisms, on the homology level, associated to commutation and destabilization, and showing that they carry  $[\mathbf{x}^{\pm}]$  to  $[\mathbf{x}^{\pm}]$ . For filtered invariants, the isomorphisms are shown to be covered by filtered quasi-isomorphisms on the chain level. We take a very similar approach, but with two differences:

• We work directly on the filtered chain level, and show that  $\mathbf{x}^{\pm}$  is carried by the filtered chain map either to  $\mathbf{x}^{\pm}$ , or to  $\mathbf{x}^{\pm} + \mathbf{y}$ , where  $\mathbf{y}$  belongs to a lower filtration; and

• We do not require our destabilization maps to be isomorphisms on homology (of the associated graded object); this is in line with the philosophy of viewing these maps as maps of decorated link cobordisms. (See, for example, [Zem19].) Accordingly, we also separately consider stabilization (and not just destabilization) maps.

#### 3.2. **Commutation.** First, we prove invariance under commutation.

**Lemma 3.3.** Suppose  $\mathbb{G}_1$  and  $\mathbb{G}_2$  differ by a commutation move. Then there exists a filtered chain homomorphism

$$\mathcal{C} \colon \widetilde{\mathcal{GC}}(\mathbb{G}_2) \to \widetilde{\mathcal{GC}}(\mathbb{G}_1)$$

such that

$$\mathcal{C}(\mathbf{x}^{\pm}(\mathbb{G}_2)) = \mathbf{x}^{\pm}(\mathbb{G}_1) + \mathbf{y},$$

where 
$$\mathbf{y} \in \mathcal{F}_{A(\mathbf{x}^{\pm}(\mathbb{G}_2))-1}\widetilde{\mathcal{GC}}(\mathbb{G}_1)$$
.

*Proof.* This is essentially [OSzT08, Lemma 6.6]. We briefly recall here the definition of the map constructed there, as we will also need it in Section 4.1. The map is defined by a count of pentagons, as follows. Suppose the commutation is a row commutation, and combine  $\mathbb{G}_1$  and  $\mathbb{G}_2$  into one diagram as in Figure 3, with  $\alpha$  corresponding to  $\mathbb{G}_1$  and  $\alpha'$  corresponding to  $\mathbb{G}_2$ . For  $\mathbf{x} \in S(\mathbb{G}_2)$  and




FIGURE 3. The combined diagram of a row commutation involving  $\alpha$  and  $\alpha'$ .

 $\mathbf{y} \in S(\mathbb{G}_1)$ , let  $\mathrm{Pent}(\mathbf{x}, \mathbf{y})$  be the space of pentagons in the combined diagram with the following properties. First,  $\mathrm{Pent}(\mathbf{x}, \mathbf{y})$  is empty if  $\mathbf{x}$  and  $\mathbf{y}$  do not agree at exactly m-2 points. An element  $p \in \mathrm{Pent}(\mathbf{x}, \mathbf{y})$  is an embedded pentagon with non-reflex angles whose boundary lies on the horizontal and vertical circles (including  $\alpha$  and  $\alpha'$ ) and whose vertices are points in  $(\mathbf{x} \triangle \mathbf{y}) \cup \{a\}$ , where  $\triangle$  denotes the symmetric difference, such that  $\partial(\partial p \cap \beta) = \mathbf{x} - \mathbf{y}$  in the induced orientation. Let  $\mathrm{Pent}^{\circ}(\mathbf{x}, \mathbf{y})$  be the subset of  $p \in \mathrm{Pent}(\mathbf{x}, \mathbf{y})$  such that  $\mathrm{Int}(p) \cap \mathbf{x} = \emptyset$ , and  $\mathrm{Pent}^{\circ}_{\mathbb{O}}(\mathbf{x}, \mathbf{y})$  the subset of  $p \in \mathrm{Pent}(\mathbf{x}, \mathbf{y})$  such that  $\mathrm{Int}(p) \cap \mathbf{x} = p \cap \mathbb{O} = \emptyset$ . Then, we define a linear map  $\mathcal{C}$  on  $\widetilde{\mathcal{GC}}(\mathbb{G}_2)$  given by

$$\mathcal{C}(\mathbf{x}) = \sum_{\mathbf{y} \in S(\mathbb{G}_1)} \sum_{p \in \mathrm{Pent}_\mathbb{O}^o(\mathbf{x}, \mathbf{y})} \mathbf{y}.$$

This map is known to respect the Maslov grading and the Alexander filtration by [MOSzT07, Lemma 3.1]. In the proof of [OSzT08, Lemma 6.6], only pentagons that do not contain X's are considered, and it is asserted that  $\mathcal{C}(\mathbf{x}^{\pm}(\mathbb{G}_2)) = \mathbf{x}^{\pm}(\mathbb{G}_1)$ ; indeed, there is only one pentagon from  $\mathbf{x}^{\pm}(\mathbb{G}_2)$  that does not contain X's (or intersection points in  $\mathbf{x}$ ), which gives the term  $\mathbf{x}^{\pm}(\mathbb{G}_1)$ . Allowing pentagons that contain X's (but blocking those that contain O's), all other pentagons from  $\mathbf{x}^{\pm}(\mathbb{G}_2)$  contain at least one X, which means that the Alexander filtration of the target generator  $\mathbf{y} \neq \mathbf{x}^{\pm}(\mathbb{G}_1)$  must be lower.

### 3.3. **Stabilization.** Next, we consider stabilization maps.

**Lemma 3.4.** Suppose  $\mathbb{G}_1$  is obtained from  $\mathbb{G}_2$  by a type X:SE or X:NW stabilization. Then there exists a filtered chain homomorphism

$$\mathcal{S} \colon \widetilde{\mathcal{GC}}(\mathbb{G}_2) \to \widetilde{\mathcal{GC}}(\mathbb{G}_1)$$

such that

$$S(\mathbf{x}^{\pm}(\mathbb{G}_2)) = \mathbf{x}^{\pm}(\mathbb{G}_1) + \mathbf{y},$$

where 
$$\mathbf{y} \in \mathcal{F}_{A(\mathbf{x}^{\pm}(\mathbb{G}_2))-1}\widetilde{\mathcal{GC}}(\mathbb{G}_1)$$
.

*Proof.* We focus first on a type X:SE stabilization. In this case, we define the filtered chain homomorphism S to be the map  $S^{oL}$  in [OSSz15, Definition 14.3.6] with all formal variables  $V_i$  set to zero. We now recall  $S^{oL}$  with this modification.

Let c be the intersection point of the two new curves in  $\mathbb{G}_1$ , as shown in Figure 4. Decompose



FIGURE 4. The distingushed 2x2 square of the diagram  $\mathbb{G}_1$  obtained from  $\mathbb{G}_2$  by a type X:SE stabilization.

 $S(\mathbb{G}_1)$  into the disjoint union  $I(\mathbb{G}_1) \sqcup N(\mathbb{G}_1)$ , where  $I(\mathbb{G}_1)$  consists of  $\mathbf{x} \in S(\mathbb{G}_1)$  such that  $c \in \mathbf{x}$ , and  $N(\mathbb{G}_1) = S(\mathbb{G}_1) \setminus I(\mathbb{G}_1)$ . Writing I and N for the corresponding submodules, this allows us to decompose  $\widetilde{\mathcal{GC}}(\mathbb{G}_1)$ , as a module, as  $I \oplus N$ . Note that, as a chain complex,  $\widetilde{\mathcal{GC}}(\mathbb{G}_1)$  is not the direct sum of I and N, or even a mapping cone of them; the differential consists of maps between I and N in both directions.

Now for  $\mathbf{x} \in I(\mathbb{G}_1)$  and  $\mathbf{y} \in S(\mathbb{G}_1)$ , a domain  $p \in \pi(\mathbf{x}, \mathbf{y})$  is said to be of type oL (originally for "out of the left") if it is trivial, in which case it has complexity 1, or it satisfies the following conditions:

- All the local multiplicaties of p are non-negative;
- At each corner in  $\mathbf{x} \cup \mathbf{y} \setminus \{c\}$ , at least three of the four adjacent squares have vanishing local multiplicities;
- The domain p has local multiplicity k-1 at the southeast square adjacent to c, and local multiplicity k at the other three squares adjacent to c; and
- The generator y has 2k + 1 coordinates not in x.

The *complexity* of a non-trivial type oL domain is defined to be 2k + 1. See Figure 5 for examples of type oL domains of complexities 1, 3, 5, and 7. The set of domains of type oL from  $\mathbf{x}$  to  $\mathbf{y}$  is denoted  $\pi^{oL}(\mathbf{x}, \mathbf{y})$ . In the literature, such domains are often called *snail domains*.

We are ready to define

$$\mathcal{S} = \mathcal{S}^{oL} \colon \widetilde{\mathcal{GC}}(\mathbb{G}_2) o \widetilde{\mathcal{GC}}(\mathbb{G}_1)$$

linearly by

$$\mathcal{S}^{oL}(\mathbf{x}) = \sum_{\mathbf{y} \in S(\mathbb{G}_1)} \sum_{p \in \pi^{oL}(e'(\mathbf{x}), \mathbf{y})} \mathbf{y},$$

where  $e': S(\mathbb{G}_2) \to I(\mathbb{G}_1)$  is the bijection  $\mathbf{x} \mapsto \mathbf{x} \cup \{c\}$ . By (reducing from the *minus* to the *tilde* versions of) [OSSz15, Lemmas 13.3.12 and 14.3.8], the map  $\mathcal{S}^{oL}$  is a chain map that respects the




Figure 5. Examples of domains of type oL.

Maslov grading and the Alexander filtration. (Note that, on the level of associated graded objects,  $S^{oL}$  is just the map e', which is the map on the top left of [OSSz15, Figure 5.13].)

We now investigate the image of  $\mathbf{x}^+(\mathbb{G}_2)$  under the map  $\mathcal{S}$ . First of all, the trivial domain in  $\pi^{oL}(e'(\mathbf{x}^+(\mathbb{G}_2)), e'(\mathbf{x}^+(\mathbb{G}_2)))$  obviously contributes a term. Since clearly  $e'(\mathbf{x}^+(\mathbb{G}_2)) = \mathbf{x}^+(\mathbb{G}_1)$ , this term is exactly  $\mathbf{x}^+(\mathbb{G}_1)$ . Now for  $\mathbf{y} \neq e'(\mathbf{x}^+(\mathbb{G}_2))$ , the domain  $p \in \pi^{oL}(e'(\mathbf{x}^+(\mathbb{G}_2)), \mathbf{y})$  contains at least two X's and one O; by an argument analogous to the proof of [OSSz15, Lemma 13.3.12], we must have  $\mathbf{y} \in \mathcal{F}_{A(\mathbf{x})-r}\widetilde{\mathcal{GC}}(\mathbb{G}_1)$ , where  $r \geq 3$ . (Since  $\mathbf{x}^+(\mathbb{G}_1)$  occupies the intersection point to the northeast of  $X_2$ , such a domain p must have multiplicity 1 in the square containing  $X_2$ , which means that p must have complexity 3, i.e. is the second-simplest kind of type oL domain.) This shows that  $\mathbf{x}^+(\mathbb{G}_1)$  is the unique generator that appears in the image of  $\mathbf{x}^+(\mathbb{G}_2)$  that has the same Alexander filtration level.

A similar proof applies for  $\mathbf{x}^-$ . Finally, for a type X:NW stabilization, the proof above can be adapted, with S now counting domains that are obtained from the ones above by rotation in the plane by  $\pi$ .

Remark 3.5. In [OSSz15, Section 14.3], the image of  $\mathbf{x}^{\pm}(\mathbb{G}_2)$  under  $\mathcal{S}$  is not computed; instead, it is observed in [OSSz15, Lemma 14.3.9] that the filtered stabilization map, made up of  $\mathcal{S}^{oL}$  and another map  $\mathcal{S}^{oR}$ , induces a map on the associated graded objects  $\widetilde{GC}$  (or rather,  $GC^-$  there) that sends  $\mathbf{x}^{\pm}$  to  $\mathbf{x}^{\pm}$ . This is sufficient to cover the case i=1 in Theorem 1.2.

## 3.4. **Destabilization.** We now move on to destabilization.

**Lemma 3.6.** Suppose  $\mathbb{G}_1$  is obtained from  $\mathbb{G}_2$  by a type X:SE or X:NW stabilization. Then there exists a filtered chain homomorphism

$$\mathcal{D} \colon \widetilde{\mathcal{GC}}(\mathbb{G}_1) \to \widetilde{\mathcal{GC}}(\mathbb{G}_2)$$

such that

$$\mathcal{D}(\mathbf{x}^{\pm}(\mathbb{G}_1)) = \mathbf{x}^{\pm}(\mathbb{G}_2).$$

*Proof.* This is essentially proved in [OSzT08, Lemma 6.5], but we opt to present a proof here for consistency with the more modern choice of destabilization maps as in [OSSz15].

We again focus on a type X:SE destabilization first. Like S, we also define D using snail domains. While a filtered chain map is not spelled out in [OSSz15] for a type X:SE destabilization, we may draw inspiration from the right half of the commutative diagram in [OSSz15], Figure 5.13] (also alluded to in the proof of Lemma 3.4 above) to figure out which snail domains to use. The chain complexes in [OSSz15], Figure 5.13] are those on the level of associated graded objects in our context. (For example, note that for us, there is also an arrow from I to N.) Our snail domains should include the maps  $I \to L$  and  $N \to L$  in that diagram.

With this insight, we define  $\mathcal{D}$  as follows. We continue our notation of c,  $I(\mathbb{G}_1)$ ,  $N(\mathbb{G}_1)$ , and I and N as in the proof of Lemma 3.4. For  $\mathbf{x} \in S(\mathbb{G}_1)$  and  $\mathbf{y} \in I(\mathbb{G}_1)$ , a domain  $p \in \pi(\mathbf{x}, \mathbf{y})$  is said to be of type iL (originally for "into the left") if it is trivial, in which case it has complexity 1, or it satisfies the following conditions:

- All the local multiplicaties of p are non-negative;
- At each corner in  $\mathbf{x} \cup \mathbf{y} \setminus \{c\}$ , at least three of the four adjacent squares have vanishing local multiplicities;
- The domain p has local multiplicity k-1 at the northeast square adjacent to c, and local multiplicity k at the other three squares adjacent to c; and
- The generator  $\mathbf{y}$  has 2k+1 coordinates not in  $\mathbf{x}$ .

The *complexity* of a non-trivial type iL domain is defined to be 2k + 1. See Figure 6 for examples of type iL domains of complexities 1, 3, 5, and 7. The set of domains of type iL of complexity 1 from  $\mathbf{x}$  to  $\mathbf{y}$  is denoted  $\pi_1^{iL}(\mathbf{x}, \mathbf{y})$ , while the set of domains of type iL of complexity greater than 1 from  $\mathbf{x}$  to  $\mathbf{y}$  is denoted  $\pi_{>1}^{iL}(\mathbf{x}, \mathbf{y})$ .




Figure 6. Examples of domains of type iL.

We now define

$$\mathcal{D} = \mathcal{D}^{iL} \colon \widetilde{\mathcal{GC}}(\mathbb{G}_1) \to \widetilde{\mathcal{GC}}(\mathbb{G}_2)$$


linearly by

(3.7) 
$$\mathcal{D}^{iL}(\mathbf{x}) = \begin{cases} \sum_{\mathbf{y} \in I(\mathbb{G}_1)} \sum_{p \in \pi_{>1}^{iL}(\mathbf{x}, \mathbf{y})} e(\mathbf{y}) & \text{if } \mathbf{x} \in I(\mathbb{G}_1),^{8} \\ \sum_{\mathbf{y} \in I(\mathbb{G}_1)} \sum_{p \in \pi_{>1}^{iL}(\mathbf{x}, \mathbf{y})} e(\mathbf{y}) & \text{if } \mathbf{x} \in N(\mathbb{G}_1), \end{cases}$$

where  $e: I(\mathbb{G}_1) \to S(\mathbb{G}_2)$  is the bijection  $\mathbf{x} \cup \{c\} \mapsto \mathbf{x}$ . In other words, if we separate the two cases in (3.7) into maps

$$\mathcal{D}_1^{iL} \colon I \to \widetilde{\mathcal{GC}}(\mathbb{G}_2), \qquad \mathcal{D}_{>1}^{iL} \colon N \to \widetilde{\mathcal{GC}}(\mathbb{G}_2),$$

then  $\mathcal{D} \colon \widetilde{\mathcal{GC}}(\mathbb{G}_1) \to \widetilde{\mathcal{GC}}(\mathbb{G}_2)$  is given by the horizontal maps in



<sup>&</sup>lt;sup>8</sup>Note that the double sum in this first line is in fact simply  $e(\mathbf{x})$ .

See [OSSz15, (13.6)] for comparison. (The destabilization there is of type X:SW.)

By a case analysis similar to [OSSz15, Lemmas 13.3.12 and 13.3.13], one could see that  $\mathcal{D}$  is a chain map that respects the Maslov grading and the Alexander filtration. For brevity, we omit the details, which are considerably tedious. Alternatively, one could also appeal to the holomorphic Heegaard Floer theory, e.g. via [MOT20, Proposition 3.10 and Lemma 3.11]. (For the astute reader, the map  $\mathcal{D}^{iL}_{>1}$  is needed to ensure that we have a chain map.) We now investigate the image of  $\mathbf{x}^{\pm}(\mathbb{G}_1)$  under the map  $\mathcal{D}$ . Since  $\mathbf{x}^{\pm}(\mathbb{G}_1) \in I(\mathbb{G}_1)$ , we immediately

get that

$$\mathcal{D}(\mathbf{x}^{\pm}(\mathbb{G}_1)) = e(\mathbf{x}^{\pm}(\mathbb{G}_1)) = \mathbf{x}^{\pm}(\mathbb{G}_2),$$

which is what we wanted to prove. The case of type X:NW destabilizations is handled again by rotating domains by  $\pi$ .

*Proof of Theorem 1.2.* The result is obtained by applying Lemma 2.4 to the filtered chain homomorphisms in Lemma 3.3, Lemma 3.4, and Lemma 3.6.

## 4. Obstructions to decomposable Lagrangian cobordisms

With the goal of proving Theorem 1.7, we will define filtered chain maps corresponding to pinch and birth moves such that the induced maps on spectral sequences preserve  $\lambda_i^+$  and  $\lambda_i^-$ .

4.1. Pinches. Suppose  $\Lambda_+$  and  $\Lambda_-$  are Legendrian links such that  $\Lambda_+$  is obtained from  $\Lambda_-$  by a pinch move. Then there exist diagrams  $\mathbb{G}_+$  and  $\mathbb{G}_-$  for  $\Lambda_+$  and  $\Lambda_-$ , respectively, which only differ in the placement of a single pair of X's or O's in adjacent rows, as shown in Figure 7. If the two markings at which the diagrams differ are X's, we will say that  $\mathbb{G}_+$  is obtained from  $\mathbb{G}_-$  by an X swap; if the markings are O's, we will say that  $\mathbb{G}_+$  is obtained from  $\mathbb{G}_-$  by an O swap.





Figure 7. The grid diagrams  $\mathbb{G}_{\pm}$  corresponding to the two types of pinch moves. The top corresponds to an X swap, and the bottom to an O swap.

Through a series of local stabilization and commutation moves, we can ensure that the swap occurs between two markings that are separated by at least two vertical lines. This can be achieved, for example, by performing a stabilization of type X:SE on any X marking to the right of all four markings in the two adjacent rows where the swap is to be done, and then commuting the column containing the new O to the left until it occupies the column between the two markings to be swapped. Alternatively, one could perform a stabilization of type X:NW on any X marking to the left of all four markings in the swap rows, and commute the column with the new O to the right.

We combine the two diagrams  $\mathbb{G}_+$  and  $\mathbb{G}_-$  into a single diagram, as in Figure 8, which we call the combined diagram. On the combined diagram, we can see each of  $\mathbb{G}_+$  and  $\mathbb{G}_-$  by using the same markings, but varying the placement of one horizontal circle ( $\alpha$  for  $\mathbb{G}_+$  and  $\alpha'$  for  $\mathbb{G}_-$ , as seen in Figure 8). Of particular interest are the intersection points of  $\alpha$  and  $\alpha'$  labeled a and b in Figure 8; these points will be used to define maps  $\mathcal{P}_X$  and  $\mathcal{P}_O$ .






FIGURE 8. The combined grid diagrams corresponding to the two types of pinch moves. Left: An X swap. Right: An O swap.

# $4.1.1. \ X \ swap.$

**Lemma 4.1.** Suppose  $\mathbb{G}_+$  is obtained from  $\mathbb{G}_-$  by an X swap. Then there exists a filtered chain homomorphism

$$\mathcal{P}_X \colon \widetilde{\mathcal{GC}}(\mathbb{G}_+) \to \widetilde{\mathcal{GC}}(\mathbb{G}_-) \left[ 1, \frac{|\Lambda_+| - |\Lambda_-| + 1}{2} \right]$$

such that

$$\mathcal{P}_X(\mathbf{x}^{\pm}(\mathbb{G}_+)) = \mathbf{x}^{\pm}(\mathbb{G}_-) + \mathbf{y},$$

where  $\mathbf{y} \in \mathcal{F}_{A(\mathbf{x}^{\pm}(\mathbb{G}_{-}))-1}\widetilde{\mathcal{GC}}(\mathbb{G}_{-})$ .


*Proof.* Given  $\mathbf{x} \in S(\mathbb{G}_+)$  and  $\mathbf{y} \in S(\mathbb{G}_-)$ , define  $\mathrm{Pent}^{\circ}(\mathbf{x}, \mathbf{y})$  and  $\mathrm{Pent}^{\circ}_{\mathbb{O}}(\mathbf{x}, \mathbf{y})$  as in Section 3.2, and let

$$\mathcal{P}_X \colon \widetilde{\mathcal{GC}}(\mathbb{G}_+) \to \widetilde{\mathcal{GC}}(\mathbb{G}_-)$$

be the linear map that counts these pentagons, i.e.

$$\mathcal{P}_X(\mathbf{x}) = \sum_{\mathbf{y} \in S(\mathbb{G}_-)} \sum_{p \in \mathrm{Pent}_\mathbb{O}^\circ(\mathbf{x}, \mathbf{y})} \mathbf{y}.$$

First, we prove that  $\mathcal{P}_X$  is a chain map. The proof is analogous to [MOSzT07, Lemma 3.1]. Every domain that is a juxtaposition of a pentagon p and rectangle r decomposes in exactly two ways, and thus contributes an even number of times to the count of  $\widetilde{\delta}_{\mathbb{G}_-} \circ \mathcal{P}_X(\mathbf{x}) + \mathcal{P}_X \circ \widetilde{\delta}_{\mathbb{G}_+}(\mathbf{x})$ . In other words, the coefficient of any  $\mathbf{y} \in S(\mathbb{G}_-)$  in  $\widetilde{\delta}_{\mathbb{G}_-} \circ \mathcal{P}_X(\mathbf{x}) + \mathcal{P}_X \circ \widetilde{\delta}_{\mathbb{G}_+}(\mathbf{x})$  is zero.






FIGURE 9. Two types of pentagons that contribute to the map  $\mathcal{P}_X$ . Below, we analyze how points in the various marked regions contribute to the combinatorial counts used to compute the gradings of the associated generators.

Next, we investigate how  $\mathcal{P}_X$  interacts with the Maslov grading and the Alexander filtration. Suppose  $p \in \operatorname{Pent}^{\circ}(\mathbf{x}, \mathbf{y})$  is an empty pentagon whose domain includes the small pentagon above the intersection point a, as seen on Figure 8. After performing cyclic commutations if necessary, we may assume the X swap is performed in the first and second rows, and the O in the second row is in the last column; see Figure 9. This allows for a simpler discussion of the gradings computation. Below, we assume that the right edge of p is to the left of the second intersection point of  $\alpha$  and  $\alpha'$ ; see the left of Figure 9. The other case is analogous.

Observe that if q is a point in the interior of the starred regions in Figure 9, then  $\mathcal{J}(\{q\}, \mathbf{x}) = \mathcal{J}(\{q\}, \mathbf{y})$ , where the left hand side is computed on  $\mathbb{G}_+$  and the right hand side on  $\mathbb{G}_-$ . Next, exactly one of the regions labeled with 1 and 2 contains a marking, which is an X. Call this marking  $X_1$  and note that  $\mathcal{J}(\{X_1\}, \{x_1\}) = \mathcal{J}(\{X_1\}, \{y_1\})$ , while  $\mathcal{J}(\{X_1\}, \{x_2\}) = 1/2$  and  $\mathcal{J}(\{X_1\}, \{y_2\}) = 0$ . So  $X_1$  contributes 1/2 to the count  $\mathcal{J}(\mathbf{x} - \mathbf{y}, \mathbb{X})$ . Similarly, let  $X_2$  be the unique marker in regions 3 and 4, and note that  $X_2$  contributes 1/2 to the count  $\mathcal{J}(\mathbf{x} - \mathbf{y}, \mathbb{X})$ . There is no marker in region 5, whereas for any marker q in the top unlabelled region, we have  $\mathcal{J}(\{q\}, \mathbf{x}) = \mathcal{J}(\{q\}, \mathbf{y})$ . (Note that the bottom unlabelled region is in fact empty.) Finally, each point q in the interior of p contributes 1 to  $\mathcal{J}(\mathbf{x} - \mathbf{y}, \{q\})$ .

The above paragraph implies that  $\mathcal{J}(\mathbf{y}, \mathbb{X}) - \mathcal{J}(\mathbf{x}, \mathbb{X}) = -1 - |\mathbb{X} \cap p|$  and  $\mathcal{J}(\mathbf{y}, \mathbb{O}) - \mathcal{J}(\mathbf{x}, \mathbb{O}) = 0$ . Also, observe that  $\mathcal{J}(\mathbb{X}, \mathbb{X})$  decreases by one as we move from  $\mathbb{G}_+$  to  $\mathbb{G}_-$ , whereas  $\mathcal{J}(\mathbb{O}, \mathbb{O})$  is unchanged, and  $\mathcal{J}(\mathbf{y}, \mathbf{y}) - \mathcal{J}(\mathbf{x}, \mathbf{x}) = -1$ . Thus,

$$M(\mathbf{y}) - M(\mathbf{x}) = -1,$$
  
$$A(\mathbf{y}) - A(\mathbf{x}) = |p \cap \mathbb{X}| + \frac{|\Lambda_-| - |\Lambda_+| - 1}{2},$$

where  $\mathbf{x}$  is considered an element of the unshifted complex  $\mathcal{GC}(\mathbb{G}_+)$ . Since  $|p \cap \mathbb{X}| \geq 0$ , shifting the grading and filtration of this complex as in the statement of the lemma, we obtain a filtered chain homomorphism. The case where p is a pentagon below a is analogous.

Finally, we compute the image of  $\mathbf{x}^{\pm}(\mathbb{G}_{+})$  under  $\mathcal{P}_{X}$ . The proof of [BLW22, Lemma 3.3] observes that, if X's and O's are both blocked, there is a unique pentagon that carries  $\mathbf{x}^{+}(\mathbb{G}_{+})$  to  $\mathbf{x}^{+}(\mathbb{G}_{-})$ , and similarly for  $\mathbf{x}^{-}$ . We observe that, allowing X's to be unblocked, we may get other pentagons, but such pentagons always contain at least one X inside, meaning that the target generator necessarily belongs to a lower Alexander filtration level.

4.1.2. O swap.

**Lemma 4.2.** Suppose  $\mathbb{G}_+$  is obtained from  $\mathbb{G}_-$  by an O swap. Then there exists a filtered chain homomorphism

$$\mathcal{P}_O \colon \widetilde{\mathcal{GC}}(\mathbb{G}_+) \to \widetilde{\mathcal{GC}}(\mathbb{G}_-) \left[ 1, \frac{|\Lambda_+| - |\Lambda_-| + 1}{2} \right]$$

such that

$$\mathcal{P}_O(\mathbf{x}^{\pm}(\mathbb{G}_+)) = \mathbf{x}^{\pm}(\mathbb{G}_-).$$

Proof. Given  $\mathbf{x} \in S(\mathbb{G}_+)$  and  $\mathbf{y} \in S(\mathbb{G}_-)$ , let  $\mathrm{Tri}(\mathbf{x}, \mathbf{y})$  be the set of triangles in the combined diagram whose vertices are points in  $(\mathbf{x} \triangle \mathbf{y}) \cup \{b\}$ , with the following conditions. First,  $\mathrm{Tri}(\mathbf{x}, \mathbf{y})$  is empty unless  $\mathbf{x}$  and  $\mathbf{y}$  agree at m-1 points. An element  $p \in \mathrm{Tri}(\mathbf{x}, \mathbf{y})$  is an embedded triangle with non-reflex angles whose boundary lies on the horizontal and vertical circles (including  $\alpha$  and  $\alpha'$ ) and whose vertices are points in  $(\mathbf{x} \triangle \mathbf{y}) \cup \{b\}$ , such that  $\partial(\partial p \cap \beta) = \mathbf{x} - \mathbf{y}$ . Note that a triangle is automatically empty, in the sense that  $\mathrm{Int}(p) \cap \mathbf{x} = \emptyset$ . Let  $\mathrm{Tri}_{\mathbb{Q}}(\mathbf{x}, \mathbf{y})$  be the subset of  $p \in \mathrm{Tri}(\mathbf{x}, \mathbf{y})$  such that  $t \cap \mathbb{Q} = \emptyset$ . Define

$$\mathcal{P}_O \colon \widetilde{\mathcal{GC}}(\mathbb{G}_+) \to \widetilde{\mathcal{GC}}(\mathbb{G}_-)$$

to be the linear map that counts these triangles:

$$\mathcal{P}_O(\mathbf{x}) = \sum_{\mathbf{y} \in S(\mathbb{G}_-)} \sum_{t \in \mathrm{Tri}_{\mathbb{O}}(\mathbf{x}, \mathbf{y})} \mathbf{y}.$$

The proof that  $\mathcal{P}_O$  is a chain map is analogous to the proof for  $\mathcal{P}_X$ . This time, we consider concatenations of rectangles and triangles, rather than rectangles and pentagons; see, for example, [Won17, Lemma 3.4] for details. The proof that  $\mathcal{P}_O$ , with the shifts in the statement of the lemma, respects the Maslov and Alexander filtration, is also a direct computation similar to the proof for  $\mathcal{P}_X$ . Finally, by [BLW22, Theorems 3.7 and 3.8], we know that when X's and O's are both blocked, the image of  $\mathbf{x}^{\pm}(\mathbb{G}_+)$  under  $\mathcal{P}_O$  is  $\mathbf{x}^{\pm}(\mathbb{G}_-)$ . We now allow X's to be unblocked, but in fact, no X's can be in a triangle! Thus our proof is complete.

### 4.2. Birth moves.

**Lemma 4.3.** Suppose  $\mathbb{G}_+$  is obtained from  $\mathbb{G}_-$  by a birth move. Then there exists a filtered chain homomorphism

$$\mathcal{B} \colon \widetilde{\mathcal{GC}}(\mathbb{G}_+) \to \widetilde{\mathcal{GC}}(\mathbb{G}_-) \llbracket -1, 0 \rrbracket$$

such that

$$\mathcal{B}(\mathbf{x}^{\pm}(\mathbb{G}_{+})) = \mathbf{x}^{\pm}(\mathbb{G}_{-}).$$

*Proof.* Our strategy is to extend the birth map in [BLW22, Proposition 3.9] to allow rectangles that contain X's. For completeness, we repeat the set up in [BLW22] below.

Through row and column commutations, there exists a diagram such that the birth occurs directly to the bottom right of an O. Then, if we define the points a and b as shown in Figure 10, we can decompose  $S(\mathbb{G}_+)$  into the disjoint union  $AB \sqcup AN \sqcup NB \sqcup NN$  where

- AB consists of  $\mathbf{x} \in S(\mathbb{G}_+)$  with  $a, b \in \mathbf{x}$ ;
- AN consists of  $\mathbf{x} \in S(\mathbb{G}_+)$  with  $a \in \mathbf{x}$  and  $b \notin \mathbf{x}$ ;
- NB consists of  $\mathbf{x} \in S(\mathbb{G}_+)$  with  $a \notin \mathbf{x}$  and  $b \in \mathbf{x}$ ; and
- NN consists of  $\mathbf{x} \in S(\mathbb{G}_+)$  with  $a, b \notin \mathbf{x}$ .




FIGURE 10. Left: Part of the grid diagram  $\mathbb{G}_{-}$ . Right: The corresponding part of the diagram  $\mathbb{G}_{+}$  obtained from  $\mathbb{G}_{-}$  by a birth move.

This induces a decomposition of the vector space  $\widetilde{\mathcal{GC}}(\mathbb{G}_+)$  as a direct sum,

$$\widetilde{\mathcal{GC}}(\mathbb{G}_+) = \widetilde{AB} \oplus \widetilde{AN} \oplus \widetilde{NB} \oplus \widetilde{NN}$$

where the summands are the subspaces generated by the corresponding subsets. Note that we have a sequence of subcomplexes

$$NN \subset NB \oplus NN \subset NN \oplus NB \oplus AN \subset \widetilde{\mathcal{GC}}(\mathbb{G}_+),$$

since no rectangle arising from the differential can end at a or at b. Let  $(\widetilde{AB}, \widetilde{\delta}_{AB})$  be the quotient complex of  $\widetilde{\mathcal{GC}}(G_+)$  by  $NN \oplus NB \oplus AN$ . There is a natural bijection of the generators in AB and

the generators in  $S(\mathbb{G}_{-})$  given by  $\mathbf{x} \mapsto \mathbf{x}' := \mathbf{x} \setminus \{a, b\}$ . This map extends linearly to an isomorphism  $e: AB \to \mathcal{GC}(\mathbb{G}_{-})$  of (not necessarily filtered) chain complexes.

For any  $\mathbf{x}, \mathbf{y} \in AB$ , there is a bijection of empty rectangles  $r \in \text{Rect}^{\circ}(\mathbf{x}, \mathbf{y})$  not containing O's in  $\widehat{AB}$  and empty rectangles  $r' \in \operatorname{Rect}^{\circ}(\mathbf{x}', \mathbf{y}')$  not containing O's in  $\mathbb{G}_{-}$ , since any rectangle that contains the  $2 \times 2$  square where the new unknot is must necessarily contain an O. By (2.1) and (2.2), this bijection shows that e respects the Alexander filtration and is homogeneous.

Let  $\mathbb{O}$  and  $\mathbb{X}$  denote the sets of O and X markers in  $\mathbb{G}_+$  respectively. Now, for  $\mathbf{x} \in NB$  and  $\mathbf{y} \in AB$ , let

$$\operatorname{Rect}_{AB}(\mathbf{x}, \mathbf{y}) \subset \operatorname{Rect}_{\mathbb{G}_+}(\mathbf{x}, \mathbf{y})$$

be the subset of rectangles p that satisfy

The second bullet item here is the key difference from [BLW22, Proposition 3.9], which requires an equality instead of an inclusion. Let  $\psi$  be the linear map defined on generators by counting such rectangles:

$$\psi(\mathbf{x}) = \sum_{\mathbf{y} \in AB} \sum_{p \in \text{Rect}_{AB}(\mathbf{x}, \mathbf{y})} \mathbf{y}.$$

Let  $\Pi: \mathcal{GC}(\mathbb{G}_+) \to NB$  be the projection onto the summand NB. Finally, let  $\mathcal{B}$  be the linear map defined by the composition:

$$\mathcal{B} = e \circ \psi \circ \Pi.$$

First, we show that  $\mathcal{B}(\mathbf{x}^{\pm}(\mathbb{G}_{+})) = \mathbf{x}^{\pm}(\mathbb{G}_{-})$ . Note that  $\mathbf{x}^{\pm}(\mathbb{G}_{+}) \in NB$ , so  $\Pi(\mathbf{x}^{\pm}(\mathbb{G}_{+})) = \mathbf{x}^{\pm}(\mathbb{G}_{+})$ . Thus,  $\mathcal{B}(\mathbf{x}^{\pm}(\mathbb{G}_{+})) = e \circ \psi(\mathbf{x}^{\pm}(\mathbb{G}_{+}))$ . As shown in Figure 11, there is a unique rectangle of the type that defines  $\psi$  that starts at  $\mathbf{x}^{\pm}(\mathbb{G}_{+})$ ; composing with e, we see that  $\mathcal{B}(\mathbf{x}^{\pm}(\mathbb{G}_{+})) = \mathbf{x}^{\pm}(\mathbb{G}_{-})$ .




FIGURE 11. Left: In  $\mathbb{G}_+$ , the generator  $\mathbf{x}^+(\mathbb{G}_+)$  (solid circles) and the unique generator y (hollow circles) for which  $\operatorname{Rect}_{AB}(\mathbf{x}^+(\mathbb{G}_+),\mathbf{y})$  is non-empty. There is a single rectangle  $p \in \text{Rect}_{AB}(\mathbf{x}^+(\mathbb{G}_+), \mathbf{y})$ , showing that  $\psi(\mathbf{x}^+(\mathbb{G}_+)) = \mathbf{y}$ . The corresponding generator  $\mathcal{B}(\mathbf{x}^+(\mathbb{G}_+)) = \mathbf{x}^+(\mathbb{G}_-)$  is shown in  $\mathbb{G}_-$ . Right: The analogous analysis for  $\mathbf{x}^{-}$ .

The proof that  $\mathcal{B}$  is a chain map is, almost *verbatim*, the same as that of [BLW22, Lemma 3.10]. Turning to the Maslov grading and Alexander filtration, first note that e and  $\Pi$  are both homogeneous with respect to both the Maslov and Alexander functions. From the definition of  $\psi$ , it is clear that  $\psi$  is homogeneous with respect to the Maslov grading; thus, one may use (2.3) together with the fact that  $\Lambda_+$  is the disjoint union of  $\Lambda_-$  with an unknot with tb=-1 and r=0, to compute the Maslov grading shift to be 1. Finally, we can use (2.2) to compute the Alexander filtration shift of

a rectangle  $p \in \text{Rect}_{AB}(\mathbf{x}, \mathbf{y})$  to be  $-\#(r \cap \mathbb{X})$ . Temporarily blocking the X's in  $\psi$  except  $X_2$  and  $X_3$ , [BLW22, Lemma 3.12] shows that  $\mathcal{B}$  is homogeneous with respect to, and in fact preserves, the Alexander function. Unblock the X's now; combining the facts above, we see that if  $\mathbf{y}'$  appears as a term in  $\mathcal{B}(\mathbf{x})$  via a rectangle  $p \in \text{Rect}_{AB}(\mathbf{x}, \mathbf{y})$ , then

$$A(y) - A(x) = -\#(p \cap X) + 2 \le 0,$$

with equality when p does not contain any X's, which is indeed attained. This shows that  $\mathcal{B}$  is a filtered chain map with no filtration level shift. 

## 4.3. Proof of the weak functoriality and obstruction.

*Proof of Theorem 1.10.* The result is obtained by applying Lemma 2.4 on the filtered chain homomorphisms in Lemma 4.1, Lemma 4.2, Lemma 4.3 to obtain the desired maps for pinches and births, and using Theorem 1.2 to obtain maps for Legendrian isotopies. As a decomposable Lagrangian cobordism is made up of these pieces, the associated map can be defined by composing the maps above. For more details, see [BLW22, proof of Theorem 1.5].

Proof of Theorem 1.7. This is immediate from Theorem 1.10 and Lemma 2.4; in particular, the existence of the map in Theorem 1.10 shows that  $n^{\pm}(\Lambda_{-}) > n^{\pm}(\Lambda_{+})$ .

Proof of Corollary 1.9. A decomposable Lagrangian filling of  $\Lambda$  is the concatenation of a Lagrangian birth and a decomposable Lagrangian cobordism from the undestabilizable Legendrian unknot  $\Lambda_U$ to  $\Lambda$ . Noting that  $n^{\pm}(\Lambda_U) = \infty$  and  $\tilde{\lambda}_i^{\pm}(\Lambda_U) \neq 0$  for all  $i \in \mathbb{Z}_{>1}$ , the corollary follows.

# 5. Computations

While our invariants have nice definitions in terms of the classes of  $\mathbf{x}^{\pm}(\mathbb{G})$  in sub-quotients of our complex, for computational purposes we would like to rephrase these definitions in terms of a particular kind of sub-quotient: the homology of a complex. Essentially, given a grid diagram G and an integer  $r \geq 1$ , we would like to be able to answer the questions

- (1) Is  $\widetilde{\lambda}_r^{\pm}(\mathbb{G})$  well-defined?
- (2) Is  $\tilde{\lambda}_r^{\pm}(\mathbb{G}) = 0$  for a given r?

by computing the class of some element in the homology of some complex. This would allow us to use the techniques of [NOT08] to compute the homology class of such an element by doing local searches and reductions to make computations more efficient and feasible for larger knots.

We will tackle the second question first. It turns out that the question of whether  $\lambda_r^{\pm}$  is zero or not is equivalent to the question of whether or not  $\mathbf{x}^{\pm}$  is null-homologous in a particular sub-quotient complex.

**Proposition 5.1.** Given  $x \in \mathcal{F}_pC$  with  $\partial x \in \mathcal{F}_{p-1}$  and  $r \in \mathbb{Z}_{>0}$ , we have that  $[x]^r = 0 \in E^r$  if and only if  $[x] = 0 \in H_*(\mathcal{F}_{p+r-1}C/\mathcal{F}_{p-1}C)$ .

*Proof.* To start, assume  $[x]^r=0\in E^r$ . Therefore,  $[x]^r\in B_p^r$ , so we have that  $x+z=\partial y$ , where  $y \in \mathcal{F}_{p+r-1}C$  and  $z \in \mathcal{F}_{p-1}C$ . Therefore,  $\partial y = x + z \equiv x \pmod{\mathcal{F}_{p-1}C}$ , so  $\partial[y] = [x] \in \mathcal{F}_{p+r-1}C/\mathcal{F}_{p-1}C$ , thus  $[x] = 0 \in H_*(\mathcal{F}_{p+r-1}C/\mathcal{F}_{p-1}C)$ . This proves one direction. To prove the other implication, assume  $[x] = 0 \in H_*(\mathcal{F}_{p+r-1}C/\mathcal{F}_{p-1}C)$ . Then  $x = \partial y + z$ , for  $y \in \mathcal{F}_{p+r-1}C$  and  $z \in \mathcal{F}_{p-1}C$ . Therefore,  $x \in \mathcal{F}_pC \cap \partial(\mathcal{F}_{p+r-1}C) + \mathcal{F}_{p-1}$ , thus  $[x]^r \in B_p^r$  and

 $[x]^r = 0 \in E_p^r$ .

Now, we will attempt to answer the first question in a similar way.

**Proposition 5.2.** Given  $x \in \mathcal{F}_pC$  with  $\partial x \in \mathcal{F}_{p-1}$  and  $r \in \mathbb{Z}_{\geq 0}$ , we have that  $d^r[x]^r = 0 \in E^r$  if and only if  $[\partial x] = 0 \in H_*(\mathcal{F}_{p-1}C/\mathcal{F}_{p-r-1}C)$ .

*Proof.* To start, assume  $d^r[x]^r = 0 \in E^r$ . Therefore,  $[x]^{r+1} \in Z_p^{r+1}$  is defined, so we know that  $\partial(x+z) = y$  for some  $y \in \mathcal{F}_{p-r-1}C$  and  $z \in \mathcal{F}_{p-1}C$ . Rearranging gives us that  $\partial x = \partial z + y$ , so  $\partial x \equiv \partial z \pmod{\mathcal{F}_{p-r-1}C}$ , thus  $[\partial x] = [\partial z] = 0 \in H_*(\mathcal{F}_{p-1}C/\mathcal{F}_{p-r-1}C)$ .

To prove the other direction, assume  $[\partial x] = 0 \in H_*(\mathcal{F}_{p-1}C/\mathcal{F}_{p-r-1}C)$ . This means that  $\partial x = \partial z + y$  for some  $z \in \mathcal{F}_{p-1}C$  and  $y \in \mathcal{F}_{p-r-1}C$ . Rearranging gives us that  $\partial (x+z) = y \in \mathcal{F}_{p-r-1}C$ , so  $[x]^{r+1} \in \mathbb{Z}_p^{r+1}$  is defined, thus  $d^r[x]^r = 0 \in E^r$ .

Given a grid diagram  $\mathbb{G}$ , these two techniques allow us to efficiently compute our invariants  $n^{\pm}(\mathbb{G})$  and  $\widetilde{\lambda}_{i}^{\pm}(\mathbb{G})$  as follows:

- (1) Let r=1. We know that  $\tilde{\lambda}_1^{\pm}(\mathbb{G})$  is always well-defined.
- (2) Use Proposition 5.1 to check if  $\widetilde{\lambda}_r^{\pm}(\mathbb{G}) = 0$ . If so, then we are done, and we can conclude that  $n^{\pm}(\mathbb{G}) = \infty$  and  $\widetilde{\lambda}_i^{\pm}(\mathbb{G}) = 0$  for i > r.
- (3) Otherwise, use Proposition 5.2 to check if  $d^r \widetilde{\lambda}_r^{\pm}(\mathbb{G}) = 0$ . If not, then we are done, and we can conclude that  $n^{\pm}(\mathbb{G}) = r$  and  $\widetilde{\lambda}_i^{\pm}(\mathbb{G})$  is undefined for i > r.
- (4) If  $d^r \widetilde{\lambda}_r^{\pm}(\mathbb{G}) = 0$ , then  $\widetilde{\lambda}_{r+1}^{\pm}(\mathbb{G})$  is well-defined, so we may increment r by 1 and repeat the process from (2).

This algorithm is implemented in [JPS<sup>+</sup>23], which was used to produce the results in Section 1.5.

#### References

- [BC14] Frédéric Bourgeois and Baptiste Chantraine, Bilinearized Legendrian contact homology and the augmentation category, J. Symplectic Geom. 12 (2014), no. 3, 553–583, MR 3248668
- [BLW22] John A. Baldwin, Tye Lidman, and C.-M. Michael Wong, Lagrangian cobordisms and Legendrian invariants in knot Floer homology, Michigan Math. J. **71** (2022), no. 1, 145–175. MR 4389674
- [BS18] John A. Baldwin and Steven Sivek, Invariants of Legendrian and transverse knots in monopole knot homology, J. Symplectic Geom. 16 (2018), no. 4, 959–1000. MR 3917725
- [BS21] \_\_\_\_\_, On the equivalence of contact invariants in sutured Floer homology theories, Geom. Topol. 25 (2021), no. 3, 1087–1164. MR 4268162
- [BST15] Frédéric Bourgeois, Joshua M. Sabloff, and Lisa Traynor, Lagrangian cobordisms via generating families: construction and geography, Algebr. Geom. Topol. 15 (2015), no. 4, 2439–2477. MR 3402346
- [CDGG15] Baptiste Chantraine, Georgios Dimitroglou Rizell, Paolo Ghiggini, and Roman Golovko, *Floer homology* and Lagrangian concordance, Proceedings of the Gökova Geometry-Topology Conference 2014, Gökova Geometry/Topology Conference (GGT), Gökova, 2015, pp. 76–113. MR 3381440
- [Cha10] Baptiste Chantraine, Lagrangian concordance of Legendrian knots, Algebr. Geom. Topol. 10 (2010), no. 1, 63–85. MR 2580429
- [Cha15]  $\underline{\qquad}, Lagrangian\ concordance\ is\ not\ a\ symmetric\ relation,\ Quantum\ Topol.\ {\bf 6}\ (2015),\ no.\ 3,\ 451–474.$  MR 3392961
- [Che02] Yuri Chekanov, Differential algebra of Legendrian links, Invent. Math. 150 (2002), no. 3, 441–483.
  MR 1946550
- [CN13] Wutichai Chongchitmate and Lenhard Ng, An atlas of Legendrian knots, Exp. Math. 22 (2013), no. 1, 26–37. MR 3038780
- [CNS16] Christopher Cornwell, Lenhard Ng, and Steven Sivek, Obstructions to Lagrangian concordance, Algebr. Geom. Topol. 16 (2016), no. 2, 797–824. MR 3493408
- [Cro95] Peter R. Cromwell, Embedding knots and links in an open book. I. Basic properties, Topology Appl. 64 (1995), no. 1, 37–58. MR 1339757
- [Dim16] Georgios Dimitroglou Rizell, Legendrian ambient surgery and Legendrian contact homology, J. Symplectic Geom. 14 (2016), no. 3, 811–901. MR 3548486
- [EGH00] Y. Eliashberg, A. Givental, and H. Hofer, Introduction to symplectic field theory, Geom. Funct. Anal., Special Volume, Part II (2000), 560–673. MR 1826267

[EHK16] Tobias Ekholm, Ko Honda, and Tamás Kálmán, Legendrian knots and exact Lagrangian cobordisms, J. Eur. Math. Soc. (JEMS) 18 (2016), no. 11, 2627–2689. MR 3562353

[GJ19] Marco Golla and András Juhász, Functoriality of the EH class and the LOSS invariant under Lagrangian concordances, Algebr. Geom. Topol. 19 (2019), no. 7, 3683–3699. MR 4045364

[JPS<sup>+</sup>23] Mitchell Jubeir, Ina Petkova, Noah Schwartz, Zachary Winkeler, and C.-M. Michael Wong, FilteredGRID, available at https://github.com/math-SHUR/FilteredGRID, 2023.

[KMVW19] Çağatay Kutluhan, Gordana Matić, Jeremy Van Horn-Morris, and Andy Wand, Filtering the Heegaard Floer contact invariant, preprint, version 5, 2019, arXiv:1603.02673v5.

[MOS09] Ciprian Manolescu, Peter Ozsváth, and Sucharit Sarkar, A combinatorial description of knot Floer homology, Ann. of Math. (2) 169 (2009), no. 2, 633–660. MR 2480614 (2009k:57047)

[MOSzT07] Ciprian Manolescu, Peter Ozsváth, Zoltán Szabó, and Dylan Thurston, On combinatorial link Floer homology, Geom. Topol. 11 (2007), 2339–2412. MR 2372850 (2009c:57053)

[MOT20] Ciprian Manolescu, Peter S. Ozsváth, and Dylan P. Thurston, Grid diagrams and Heegaard Floer invariants, preprint, version 4, 2020, arXiv:0910.0078v4.

[MQR<sup>+</sup>19] Lucas Meyers, Robert Quarles, Brandon Roberts, David Shea Vela-Vick, and C.-M. Michael Wong, transverse-hfk-revision, available at https://github.com/albenzo/transverse-hfk-revision/, 2019, accessed on Jun 7, 2019.

[NOT07] Lenhard Ng, Peter Ozsváth, and Dylan Thurston, TransverseHFK.c, available at https://services.math.duke.edu/~ng/math/TransverseHFK.c, 2007, accessed on Feb 8, 2019.

[NOT08] Lenhard Ng, Peter Ozsváth, and Dylan Thurston, Transverse knots distinguished by knot Floer homology, J. Symplectic Geom. 6 (2008), no. 4, 461–490. MR 2471100

[OSSz15] Peter S. Ozsváth, András I. Stipsicz, and Zoltán Szabó, Grid homology for knots and links, Mathematical Surveys and Monographs, vol. 208, American Mathematical Society, Providence, RI, 2015. MR 3381987

[OSzT08] Peter Ozsváth, Zoltán Szabó, and Dylan Thurston, Legendrian knots, transverse knots and combinatorial Floer homology, Geom. Topol. 12 (2008), no. 2, 941–980. MR 2403802

[Pan17] Yu Pan, The augmentation category map induced by exact Lagrangian cobordisms, Algebr. Geom. Topol. 17 (2017), no. 3, 1813–1870. MR 3677941

[ST13] Joshua M. Sabloff and Lisa Traynor, Obstructions to Lagrangian cobordisms between Legendrians via generating families, Algebr. Geom. Topol. 13 (2013), no. 5, 2733–2797. MR 3116302

[Sta18] The Stacks project contributors, *The Stacks project*, available at https://stacks.math.columbia.edu, 2018, accessed on Jan 17, 2023.

[Wei94] Charles A. Weibel, An introduction to homological algebra, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994. MR 1269324

[Won17] C.-M. Michael Wong, Grid diagrams and Manolescu's unoriented skein exact triangle for knot Floer homology, Algebr. Geom. Topol. 17 (2017), no. 3, 1283–1321. MR 3677929

[Zem19] Ian Zemke, Link cobordisms and functoriality in link Floer homology, J. Topol. 12 (2019), no. 1, 94–220. MR 3905679

Department of Mathematics, Dartmouth College, Hanover, NH 03755

 $Email\ address: {\tt mitchell.jubeir.23@dartmouth.edu}$ 

DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, HANOVER, NH 03755

Email address: ina.petkova@dartmouth.edu URL: https://math.dartmouth.edu/~ina/

DEPARTMENT OF MATHEMATICS, DARTMOUTH COLLEGE, HANOVER, NH 03755

Email address: noah.s.schwartz.23@dartmouth.edu

DEPARTMENT OF MATHEMATICAL SCIENCES, SMITH COLLEGE, NORTHAMPTON, MA 01063

Email address: zwinkeler@smith.edu URL: https://zach-winkeler.github.io/

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTTAWA, OTTAWA, ON K1N 6N5

Email address: Mike.Wong@uOttawa.ca

URL: https://mysite.science.uottawa.ca/cwong/