

Yet another QFT model of neutrino oscillations

W. Grimus*

University of Vienna, Faculty of Physics
Boltzmanngasse 5, A-1090 Vienna, Austria

April 5, 2023

Abstract

We consider a quantum field theory (QFT) model of neutrino oscillations in vacuum that attempts to take into account that the neutrino source particle and the neutrino detection particle both interact with their respective environments by collisions. Our model is minimal in a twofold sense. Firstly we simply assume that neutrino production and detection take place in between collisions in time intervals Δt and τ , respectively. Secondly, we only introduce the two wave packets that are absolutely necessary which are those of the neutrino source and the neutrino detection particle. Within this model we find that, for all practical purposes, there are no decoherence effects in the neutrino oscillation amplitude and oscillations occur in space not in time. Moreover, our model leads to the correct time correlation between neutrino production and detection and to a factorization of the event rate of the compound neutrino production–detection process into decay rate, oscillation probability and detection cross section.

*E-mail: walter.grimus@univie.ac.at

Symbol	Meaning
t_S	Time of creation of the source particle
Δt	Time interval of source particle decay given by $(t_S, t_S + \Delta t)$
T	Time that occurs in $\mathcal{A}_{\nu_\alpha \rightarrow \nu_\beta}$ after integrations
t_D	Time of measurement
τ	Time interval of measurement given by $(t_D - \tau/2, t_D + \tau/2)$
λ	Time parameter that occurs in the integral over the neutrino energy
t_c	Time parameter used to describe the correlation between t_S and t_D
$\mathcal{T}(t_c)$	Temporal function that occurs in the decay rate

Table 1: List of time symbols used in the paper. T is defined in equation (31) and $\mathcal{A}_{\nu_\alpha \rightarrow \nu_\beta}$ denotes the oscillation amplitude referring to neutrino flavours α and β . The parameters λ and t_c are defined in equation (27) and the function $\mathcal{T}(t_c)$ in equation (54).

1 Introduction

QFT is a suitable framework for describing neutrino oscillations—for reviews see for instance [1, 2, 3, 4, 5, 6, 7, 8], for recent literature we refer the reader to [9, 10, 11, 12, 13, 14, 15] and references therein. In particular, QFT makes allowance for the fact that neutrinos are not directly observable. This is accomplished by ascribing a single Feynman diagram to the compound process of neutrino production and detection in which the massive neutrinos are inner lines.

In this paper we introduce a minimal QFT model that tries to take into account that neutrino production and detection do usually not take place in vacuum but in an environment. This is true for both reactor and accelerator neutrinos that we have in mind in this context. In a reactor the fission fragments that produce neutrinos by β -decay are in a thermal environment. We adopt the simple picture that this decay may occur without interruption during a time interval Δt . Thus Δt is the typical time between two collisions. If a fission fragment has not decayed during this interval, it will undergo a collision with a surrounding atom and the process starts anew. Similarly, in the detection process we denote the time interval of uninterrupted measurement by τ . Possibly, τ can also be interpreted as the time resolution of the detector. Note that we assume that the detector particle is at rest apart from thermal motion and is embedded in some kind of gas or liquid. Accelerator neutrinos are generated by the decay of charged pions in a decay tunnel which is the environment in this case. Here Δt is the time between the pion creation and the moment when its trajectory intersects the end of the decay tunnel. The notation concerning the different times used in this paper is explained in table 1.

Our model is also minimal with respect to the number of wave packets. Only wave packets for the neutrino source particle and the detector particle are introduced. We assume plane waves for all particles in the final states, which is the usual practice in QFT computations of decay rates and cross sections. Obviously, there are no neutrino

wave packets since neutrinos correspond to inner lines in the Feynman diagram of the compound process.

The subjects of the paper are

- the computation of the neutrino oscillation amplitude $\mathcal{A}_{\nu_\alpha \rightarrow \nu_\beta}$ in our model,
- potential decoherence effects in $\mathcal{A}_{\nu_\alpha \rightarrow \nu_\beta}$,
- and the factorization of the event rate into (decay rate of the source particle) times (oscillation probability) times (detection cross section).

The paper is organized as follows. In section 2 the oscillation amplitude $\mathcal{A}_{\nu_\alpha \rightarrow \nu_\beta}$ is derived, starting from a preparatory form discussed in appendix A. The integration over the neutrino energy in some suitable approximation is done in section 3. A general integral needed in this context is computed in appendix B. Section 4 treats the correlation between the times t_S , when the source particle is created, and t_D , when the neutrino detection takes place; this correlation results from the integration over the neutrino energy. The connection between this time-correlation function and approximate energy conservation is established in section 5. The above-mentioned factorization is deduced in section 6 and the conclusions are presented in section 7.

2 Oscillation amplitude

The space-time variable at the neutrino source process is denoted by x_1 and that at the detection process by x_2 . We describe the neutrino source and detector particles by the wave functions

$$\int d^3p e^{-i(x_1 - x_S) \cdot p} \psi_S(\vec{p}) e^{-\Gamma m_S(t_1 - t_S)/(2E_S)} \Theta(t_1 - t_S) \Theta(t_S + \Delta t - t_1) \quad (1)$$

and

$$\int d^3k e^{-i(x_2 - x_D) \cdot k} \psi_D(\vec{k}) \Theta_\tau(t_2 - t_D), \quad (2)$$

respectively. For later convenience both wave functions are given as Fourier transforms. The vector \vec{x}_S shifts the peak of the source wave function from $\vec{x}_1 = \vec{0}$ to $\vec{x}_1 = \vec{x}_S$, with the analogue \vec{x}_D for the detection wave function. The 4-momenta p and k are decomposed into time and space components as

$$p = \begin{pmatrix} E_S \\ \vec{p} \end{pmatrix} \quad \text{and} \quad k = \begin{pmatrix} E_D \\ \vec{k} \end{pmatrix}, \quad (3)$$

respectively. We denote the mass of the source particle by m_S and that of the detector particle by m_D .

The Heaviside function $\Theta(t_1 - t_S)$ and the exponential—with the total decay rate Γ —describing the decay of the source particle are incorporated into the source wave function; this means that t_S is the time when the source particle is created. For instance, for reactor neutrinos this is the time when the nucleus that undergoes β -decay is created by fission.

As described in the introduction, we further assume that the decay process of the source particle is stopped at a time $t_S + \Delta t$ which is taken into account by $\Theta(t_S + \Delta t - t_1)$.

The function $\Theta_\tau(t_2 - t_D)$, which we incorporated into the wave function of the detector particle, specifies the uncertainty $t_D \pm \tau/2$ in the detection time. The simplest form of $\Theta_\tau(t)$ is rectangular:

$$\Theta_\tau(t) = \begin{cases} 1 & \text{for } |t| < \tau/2, \\ 0 & \text{otherwise.} \end{cases} \quad (4)$$

Note that in the case of reactor neutrinos measured by the inverse β -decay the time t_D is the time of the prompt signal.

We do not need to specify the dispersion relations

$$p^0 \equiv E_S(\vec{p}), \quad k^0 \equiv E_D(\vec{k}). \quad (5)$$

They could be that of a free particle or something else. The derivations

$$\vec{\beta}_S = \vec{\nabla}_p E_S(\vec{p}), \quad \vec{\beta}_D = \vec{\nabla}_k E_D(\vec{k}) \quad (6)$$

are the group velocities of source and detector particle, respectively.

To keep track of the assumptions we use in the computation of the oscillation amplitude $\mathcal{A}_{\nu_\alpha \rightarrow \nu_\beta}$ we number them. The first one is the following.

Assumption 1: The final states in both the neutrino source and detection process are treated as plane waves.

This is the usual assumption for computing cross sections or decay rates in particle physics. This means that we do not consider possible effects of wave packets of the final states. Note, however, that such wave packets, and also ψ_S and ψ_D , are not to be confused with wave functions of composite particles such as nuclei, nucleons, etc. Such particles could be described by some bound-state wave functions pertaining to their inner structure, however, the inner structure would be accounted for in the weak-current matrix elements. In this sense, ψ_S and ψ_D are associated with distributions of the *total* momenta \vec{p} and \vec{k} of source and detector particle, respectively.

In appendix A, the amplitude $\mathcal{A}_{\nu_\alpha \rightarrow \nu_\beta}$ at a stage before integrations are performed is displayed in equation (A.1).

Since in the following we will not consider the weak matrix elements and all final states are treated on the same footing as plane waves, we are allowed to use a simplified notation where

$$p'_S = \begin{pmatrix} E'_S \\ \vec{p}'_S \end{pmatrix} \quad \text{and} \quad k'_D = \begin{pmatrix} E'_D \\ \vec{k}'_D \end{pmatrix} \quad (7)$$

denote the *sum* over all final momenta in the source and detection process, respectively. We furthermore denote the 4-momentum of the neutrino travelling from \vec{x}_S to \vec{x}_D by q . This is the momentum that appears in the neutrino propagator.

The following integrations have to be performed in order to obtain the oscillation amplitude $\mathcal{A}_{\nu_\alpha \rightarrow \nu_\beta}$ [12]:

1. $\int d^3x_1$ and $\int d^3x_2$,

2. $\int d^3p$ and $\int d^3k$,
3. $\int dt_1$ and $\int dt_2$,
4. $\int d^3q$ in the asymptotic limit $L \equiv |\vec{x}_D - \vec{x}_S| \rightarrow \infty$,
5. $\int dq^0$ in an approximation that will be discussed in section 3.

The first three integrations are done in appendix A.

Implicitly, in the time evolution in equations (1) and (2), we have made another assumption without which it would be difficult to write down an oscillation amplitude.

Assumption 2: The source particle behaves approximately as a free particle in the time interval of length Δt between two collisions. The same is assumed for the detection particle during the measurement time-interval of length τ .

In other words, we assume that for short time intervals the time evolution of the source and detector particle can be written as a superposition of plane waves as done in equations (1) and (2), respectively. This is presumably a good approximation in the case of accelerator neutrinos produced by charged pions in a decay tunnel. It is also plausible that, in the case of reactor neutrinos, the β -decay of a fission product is not influenced by collisions because nuclei are protected by an electron shell; nevertheless nuclei get kicked around by the environment and Δt will be very short in this case. Maybe the environment can be partly taken into account by choosing suitable dispersion relations $E_S(\vec{p})$ and $E_D(\vec{k})$ or by allowing for random changes of $\vec{\beta}_S$ and $\vec{\beta}_D$.

We depart from the amplitude of equation (A.3), taking into account equations (A.5), (A.6) and (A.8), in order to discuss the remaining integrations. Through the integration over \vec{q} in the asymptotic limit $L \rightarrow \infty$, for each neutrino species with mass m_j , the neutrino 4-momentum q becomes on-shell, the denominator of the neutrino propagator is cancelled and the amplitude gets a factor L^{-1} [16].¹ Moreover, the spatial neutrino momenta point from \vec{x}_S to \vec{x}_D . Thus we arrive at the neutrino 4-momenta

$$q_j = \begin{pmatrix} Q \\ \vec{q}_j \end{pmatrix} \quad \text{with} \quad \vec{q}_j = \sqrt{Q^2 - m_j^2} \vec{\ell}, \quad \vec{\ell} = \frac{\vec{x}_D - \vec{x}_S}{L} \quad \text{and} \quad |\vec{\ell}| = 1. \quad (8)$$

Defining [18]

$$E_{Sj} = E_S|_{\vec{p}=\vec{p}'_S+\vec{q}_j}, \quad E_{Dj} = E_D|_{\vec{k}=\vec{k}'_D-\vec{q}_j}, \quad (9)$$

the oscillation amplitude now reads

$$\mathcal{A}_{\nu_\alpha \rightarrow \nu_\beta} \propto \frac{1}{L} \sum_j U_{\beta j} U_{\alpha j}^* \int dQ \quad (10a)$$

$$\times \psi_S(\vec{p}'_S + \vec{q}_j) \frac{1 - \exp \left[\left(-i(E_{Sj} - E'_S - Q) - \frac{\Gamma m_S}{2E_{Sj}} \right) \Delta t \right]}{i(E_{Sj} - E'_S - Q) + \frac{\Gamma m_S}{2E_{Sj}}} \quad (10b)$$

¹The separation between source and detection in the asymptotic limit $L \rightarrow \infty$ is a special case of localization effects in QFT that have recently been discussed in [17].

$$\times \exp \left(-i (t_D - t_S) Q + i L \sqrt{Q^2 - m_j^2} \right) \quad (10c)$$

$$\times \psi_D \left(\vec{k}'_D - \vec{q}_j \right) \tilde{\Theta}_\tau (E_{Dj} - E'_D + Q). \quad (10d)$$

The further discussion is based on this amplitude.

The amplitude in equation (10) has been derived for neutrinos. How is this amplitude modified for antineutrinos? In this case, the neutrino propagator in equation (A.3) has $x_2 - x_1$ instead of $x_1 - x_2$ and the complex conjugate is on the CKM matrix element $U_{\beta j}$ instead of $U_{\alpha j}$. Changing the neutrino momentum q to $q' = -q$, we are back to equation (10), apart from potential effects in the matrix elements which do not concern us here and the shift of the complex conjugation from $U_{\alpha j}$ to $U_{\beta j}$. In this way, the further discussion can be trivially modified for antineutrino oscillations.

Now we turn to the question whether ψ_S and ψ_D can induce decoherence in the oscillation amplitude. Let us for instance consider ψ_S . Suppose the width of this wave packet is σ_S and we have two massive neutrino species with masses m_i and m_j ($m_i \neq m_j$). In order to have oscillations involving these two massive neutrinos, neither should be suppressed by ψ_S . Therefore, we simultaneously require

$$|\vec{p}'_S + \vec{q}_i| \lesssim \sigma_S, \quad |\vec{p}'_S + \vec{q}_j| \lesssim \sigma_S. \quad (11)$$

Consequently, using the triangle inequality and expanding the square root in equation (8), we obtain

$$|\vec{q}_i - \vec{q}_j| \simeq \left| \frac{m_i^2 - m_j^2}{2\bar{Q}} \right| \lesssim 2\sigma_S. \quad (12)$$

In this equation, \bar{Q} is some average neutrino energy. In section 3.1 we will specify \bar{Q} . Denoting by Δm^2 some generic and positive neutrino mass-squared difference and assuming that via the uncertainty relation the width σ_{xS} of ψ_D in coordinate space is approximately given by $\sigma_{xS} \sim 1/(2\sigma_S)$, we reformulate equation (12) as

$$\frac{2\bar{Q}}{\Delta m^2} \gtrsim \sigma_{xS} \quad \text{or} \quad L_{\text{osc}} \equiv 4\pi \frac{\bar{Q}}{\Delta m^2} \gtrsim 2\pi \sigma_{xS}. \quad (13)$$

Since the oscillation length L_{osc} is a macroscopic quantity whereas σ_{xS} is microscopic, we conclude that ψ_S does not cause any decoherence effect. By analogy, the same is true for ψ_D [6, 18].

It remains to investigate potential decoherence caused by the second factor in equation (10b) and by $\tilde{\Theta}_\tau$ in equation (10d). In this context, we use henceforth the specific form of Θ_τ of equation (4). Therefore,

$$\tilde{\Theta}_\tau(E) = 2 \frac{\sin(E\tau/2)}{E}. \quad (14)$$

3 Integration over the neutrino energy Q

The integration over Q cannot be performed exactly. However, since it is essentially limited by $\tilde{\Theta}_\tau$ in equation (10d), we can make reasonable approximations. We read off

from equation (14) that the first zeros of this function are $E = \pm\pi/\tau$. We will take this as the typical energy scale associated with Θ_τ . However, since this is a slowly decreasing function, we might for instance integrate to the 100th zero, *i.e.* over the range of Q defined by $|E_{Dj} - E'_D + Q| \lesssim 100\pi/\tau$, to cover the bulk of the Q -integral.

3.1 Assumptions and approximations

In realistic cases, π/τ is many orders of magnitude below 1 MeV while the typical energy scale occurring in the matrix elements at neutrino source and detection is 1 MeV or larger. Taking into account the approximate energy range of Q discussed above, we are lead to

Assumption 3: The Q -dependence of the weak-current matrix elements in the source and the detection process can be neglected in the Q -integration.

Another limitation of the Q -integration which can be read off from equation (8) is

$$Q > \max_j m_j. \quad (15)$$

However, this limitation is irrelevant because the relevant neutrino energies in the integration fulfill $Q \gg m_j$.

In the following, the strategy will be to define a suitable mean neutrino energy \bar{Q} and a new integration variable δQ via

$$Q = \bar{Q} + \delta Q. \quad (16)$$

We will expand all relevant quantities that depend on Q in the small quantities δQ and $m_j^2/(2\bar{Q})$. In particular, it is important to expand the square root in the neutrino momenta \vec{q}_j , equation (8), for performing the integration over δQ . Making an expansion in $1/\bar{Q}$, we find

$$\sqrt{Q^2 - m_j^2} = \bar{Q} + \delta Q - \frac{m_j^2}{2\bar{Q}} + \frac{m_j^2 \delta Q}{2\bar{Q}^2} + \dots, \quad (17)$$

where the dots indicate terms of order $1/\bar{Q}^3$ and higher. We will content ourselves with the first three terms on the right-hand side of this equation, except in section 4 where we also discuss the fourth term.

We define a mean neutrino energy \bar{Q} via the detection process by requiring exact energy conservation in the approximation of vanishing neutrino masses. This means that \bar{Q} is the solution of

$$E_D|_{\vec{k}=\vec{k}'_D-\bar{Q}\vec{\ell}} = E'_D - \bar{Q}. \quad (18)$$

This allows us to define approximate energies [12]

$$\bar{E}_S = E_S|_{\vec{p}=\vec{p}'_S+\bar{Q}\vec{\ell}} \quad \text{and} \quad \bar{E}_D = E_D|_{\vec{k}=\vec{k}'_D-\bar{Q}\vec{\ell}} \quad (19)$$

of source and detector particle, respectively. Note that because of equation (18) the relation

$$\bar{E}_D + \bar{Q} - E'_D = 0 \quad (20)$$

is valid. The relation corresponding to the source process is, however, non-zero in general since \bar{Q} is defined via the detection process. We rather have

$$\bar{E}_S - \bar{Q} - E'_S = \bar{E}_S + \bar{E}_D - E'_S - E'_D \equiv \Delta\bar{E}, \quad (21)$$

where $\Delta\bar{E}$ is a kind of measure of energy non-conservation in the compound source-detection process. We emphasize that here and in the following all barred quantities refer to the mean energy \bar{Q} defined in equation (18).

Now, based on equation (17), we include δQ and $m_j^2/(2\bar{Q})$ in first order in the further computations. In this way we obtain

$$E_{Sj} = \bar{E}_S + \bar{\beta}_S \left(\delta Q - \frac{m_j^2}{2\bar{Q}} \right) \quad \text{with} \quad \bar{\beta}_S = \vec{\ell} \cdot \left(\vec{\nabla}_p E_S \right) \Big|_{\vec{p}=\vec{p}'_S+\bar{Q}\vec{\ell}}, \quad (22a)$$

$$E_{Dj} = \bar{E}_D - \bar{\beta}_D \left(\delta Q - \frac{m_j^2}{2\bar{Q}} \right) \quad \text{with} \quad \bar{\beta}_D = \vec{\ell} \cdot \left(\vec{\nabla}_k E_D \right) \Big|_{\vec{k}=\vec{k}'_D-\bar{Q}\vec{\ell}}. \quad (22b)$$

In these equations, $\bar{\beta}_S$ and $\bar{\beta}_D$ are mean velocities of the source and detector particle, respectively, projected unto the direction $\vec{\ell}$. The expressions we eventually need are

$$E_{Sj} - Q - E'_S = - (1 - \bar{\beta}_S) \delta Q - \bar{\beta}_S \frac{m_j^2}{2\bar{Q}} + \Delta\bar{E}, \quad (23a)$$

$$E_{Dj} + Q - E'_D = (1 - \bar{\beta}_D) \delta Q + \bar{\beta}_D \frac{m_j^2}{2\bar{Q}}. \quad (23b)$$

In order to integrate over δQ we finally resort to

Assumption 4: We assume that, in the integration over δQ , we are allowed to extend the integration limits to minus and plus infinity. In addition, we neglect the δQ -dependence in the wave functions ψ_S and ψ_D .

The latter implies that the widths σ_S and σ_D corresponding to ψ_S and ψ_D , respectively, are large enough such that this neglect is a good approximation, *i.e.*

$$\sigma_S \gg \frac{\pi}{\tau}, \quad \sigma_D \gg \frac{\pi}{\tau}. \quad (24)$$

At least for the detector particle the inequality is easy to check. Since the detector particle is at rest apart from thermal motion, we estimate [12]

$$\sigma_D \sim \sqrt{3m_D k_B T}. \quad (25)$$

For instance, for a proton as detector particle we find $\sigma_D \sim 0.01$ MeV at room temperature. Let us assume $\sigma_S \sim \sigma_D$, $\tau \sim 10^{-12}$ s, $m_j \lesssim 0.1$ eV and $\bar{Q} \lesssim 0.5$ MeV. Then

$$\frac{\pi}{\tau} \sim 2 \times 10^{-9} \text{ MeV}, \quad \frac{m_j^2}{2\bar{Q}} \lesssim 10^{-14} \text{ MeV} \quad (26)$$

and these quantities are much smaller than the widths σ_S and σ_D .

In summary, we have argued that, in the range of δQ where Θ_τ is unsuppressed, we are allowed to neglect the δQ -dependence of the weak matrix elements and of ψ_S and ψ_D while, in the range of δQ where this dependence becomes relevant, Θ_τ is suppressed. Therefore, it is a reasonable approximation to integrate over δQ from $-\infty$ to $+\infty$ with constant neutrino energy \bar{Q} in the weak matrix elements and in ψ_S and ψ_D .

3.2 Integration over δQ

In order to apply the integral formula of appendix B, we define the times

$$\lambda = (1 - \bar{\beta}_D) \frac{\tau}{2}, \quad t_c = t_D - t_S - L \quad (27)$$

and the quantities

$$z_{1j} = -\frac{1}{1 - \bar{\beta}_S} \left(\frac{\bar{\beta}_S m_j^2}{2\bar{Q}} + i \frac{\Gamma m_S}{2\bar{E}_S} - \Delta \bar{E} \right), \quad (28a)$$

$$z_{2j} = -\frac{1}{1 - \bar{\beta}_D} \frac{\bar{\beta}_D m_j^2}{2\bar{Q}}. \quad (28b)$$

Note that in $\Gamma m_S/E_{Sj}$ we have replaced E_{Sj} by \bar{E}_S since Γ is already pretty small, cf. section 5. Furthermore, we have to consider the exponential function, equation (10c), and expand the exponent according to equation (17):

$$\exp \left(-i(t_D - t_S)Q + iL\sqrt{Q^2 - m_j^2} \right) = \exp \left(-it_c \bar{Q} - i \frac{m_j^2}{2\bar{Q}} L \right) \times \exp(-it_c \delta Q). \quad (29)$$

The first factor on the right-hand side contains the irrelevant phase $t_c \bar{Q}$, but also the phase that gives rise to neutrino oscillations. The second factor has to be taken into account in the integral. With assumption 4 we rewrite equation (10) as

$$\mathcal{A}_{\nu_\alpha \rightarrow \nu_\beta} \propto \frac{1}{L} \psi_S \left(\vec{p}'_S + \bar{Q} \vec{\ell}' \right) \psi_D \left(\vec{k}'_D - \bar{Q} \vec{\ell}' \right) e^{-it_c \bar{Q}} \sum_j U_{\beta j} U_{\alpha j}^* e^{-im_j^2 L/(2\bar{Q})} \quad (30a)$$

$$\times \frac{\tau}{2} \frac{1}{1 - \bar{\beta}_S} [I_j(t_c) - I_j(t_c - T) e^{-iT z_{1j}}] \quad (30b)$$

with

$$T \equiv (1 - \bar{\beta}_S) \Delta t. \quad (31)$$

Note that T is positive. The integrals in equation (30b) have the form

$$I_j(t) = 2i \int d\delta Q \frac{e^{-it\delta Q}}{\delta Q - z_{1j}} \frac{\sin(\lambda(\delta Q - z_{2j}))}{\lambda(\delta Q - z_{2j})} \quad (32)$$

with $t = t_c$ and $t = t_c - T$. They are thus of the type of equation (B.1) treated in appendix B and the results of this appendix can be applied.

It is useful to define

$$M_j(t_c, \Delta \bar{E}) = \frac{\tau}{2} \frac{1}{1 - \bar{\beta}_S} [I_j(t_c) - I_j(t_c - T) e^{-iT z_{1j}}]. \quad (33)$$

This corresponds to equation (30b). Note that the result of the integral $I(c)$, equation (B.4), requires three case distinctions. Since in $M_j(t_c, \Delta \bar{E})$ there are two such integrals, this quantity requires five case distinctions, depending on the relative positions of the intervals

$$\mathcal{I}_c = (t_c - T, t_c) \quad \text{and} \quad \mathcal{I}_\lambda = (-\lambda, \lambda). \quad (34)$$

The two cases with $\mathcal{I}_c \cap \mathcal{I}_\lambda = \emptyset$ simply yield $M_j(t_c, \Delta\bar{E}) = 0$. This is trivial for $t_c < -\lambda$, cf. equation (B.4a), but for $\lambda < t_c - T$ equation (B.4c) has to be invoked. Among the remaining three cases with $\mathcal{I}_c \cap \mathcal{I}_\lambda \neq \emptyset$ there is one case where the intervals completely overlap. In this situation there are two possibilities:

- $T < 2\lambda$ with $\mathcal{I}_c \subset \mathcal{I}_\lambda$ and
- $T > 2\lambda$ with $\mathcal{I}_\lambda \subset \mathcal{I}_c$.

For the sake of clarity we present the result for both possibilities separately in spite of some overlap:

$$\boxed{T < 2\lambda:}$$

$$-\infty < t_c < -\lambda \quad \Rightarrow \quad M_j/f = 0, \quad (35a)$$

$$-\lambda < t_c < -\lambda + T \quad \Rightarrow \quad M_j/f = \frac{e^{-it_c z_{2j}}}{z_{1j} - z_{2j}} \left(1 - e^{-i(\lambda+t_c)(z_{1j}-z_{2j})}\right), \quad (35b)$$

$$-\lambda + T < t_c < \lambda \quad \Rightarrow \quad M_j/f = \frac{e^{-it_c z_{2j}}}{z_{1j} - z_{2j}} \left(1 - e^{-iT(z_{1j}-z_{2j})}\right), \quad (35c)$$

$$\lambda < t_c < \lambda + T \quad \Rightarrow \quad M_j/f = \frac{e^{-it_c z_{2j}}}{z_{1j} - z_{2j}} \left(e^{i(\lambda-t_c)(z_{1j}-z_{2j})} - e^{-iT(z_{1j}-z_{2j})}\right), \quad (35d)$$

$$\lambda + T < t_c < \infty \quad \Rightarrow \quad M_j/f = 0, \quad (35e)$$

$$\boxed{T > 2\lambda:}$$

$$-\infty < t_c < -\lambda \quad \Rightarrow \quad M_j/f = 0, \quad (36a)$$

$$-\lambda < t_c < \lambda \quad \Rightarrow \quad M_j/f = \frac{e^{-it_c z_{2j}}}{z_{1j} - z_{2j}} \left(1 - e^{-i(\lambda+t_c)(z_{1j}-z_{2j})}\right), \quad (36b)$$

$$\lambda < t_c < -\lambda + T \quad \Rightarrow \quad M_j/f = \frac{e^{-it_c z_{2j}}}{z_{1j} - z_{2j}} \left(e^{i(\lambda-t_c)(z_{1j}-z_{2j})} - e^{-i(\lambda+t_c)(z_{1j}-z_{2j})}\right), \quad (36c)$$

$$-\lambda + T < t_c < \lambda + T \quad \Rightarrow \quad M_j/f = \frac{e^{-it_c z_{2j}}}{z_{1j} - z_{2j}} \left(e^{i(\lambda-t_c)(z_{1j}-z_{2j})} - e^{-iT(z_{1j}-z_{2j})}\right), \quad (36d)$$

$$\lambda + T < t_c < \infty \quad \Rightarrow \quad M_j/f = 0. \quad (36e)$$

There is a common factor

$$f = -\frac{2\pi i}{(1 - \bar{\beta}_S)(1 - \bar{\beta}_D)} \quad (37)$$

in all cases. For the sake of brevity we have omitted in equations (35) and (36) the dependence on the variables t_c and $\Delta\bar{E}$ in M_j . It is easy to check that $M_j(t_c, \Delta\bar{E})$ is continuous in t_c for both $T < 2\lambda$ and $T > 2\lambda$. This is necessarily so because $I(c)$ in equation (B.4) is continuous.

4 Time correlation between neutrino production and detection

Equations (35) and (36) lead to a correlation between the times t_S and t_D [19] because a non-zero $M_j(t_c, \Delta\bar{E})$ is only possible for

$$-\lambda < t_c < \lambda + T. \quad (38)$$

Usually it is not known when the source particle and the neutrino are produced, but the time $t_D \pm \tau/2$ of neutrino detection is recorded. Therefore, equation (38) allows to infer the time t_S when the source particle is created:

$$t_D - \lambda - L - T < t_S < t_D + \lambda - L. \quad (39)$$

This is an automatic consequence of the formalism and lends credibility to our neutrino oscillation model [12].

Let us compare equation (39) with a purely classical consideration. We keep the measurement interval $t_D \pm \tau/2$ fixed and determine the earliest production time t_{S1} and the latest production time t_{S2} of the source particle such that a neutrino measurement is possible. Therefore, t_S lies in the interval

$$t_{S1} < t_S < t_{S2}. \quad (40)$$

Determination of t_{S1} : In this case the source particle decays at the latest possible time $t_{S1} + \Delta t$ at the point $x_{S1} = x_S + \bar{\beta}_S \Delta t$ and the neutrino arrives at the time $t_D - \tau/2$, which is the earliest possible measurement time, at the point $x_{D1} = x_D - \bar{\beta}_D \tau/2$.² Therefore, the condition for the determination of t_{S1} is given by

$$x_{D1} = x_{S1} + t_D - \frac{\tau}{2} - t_{S1} - \Delta t \quad \Rightarrow \quad t_{S1} = t_D - \lambda - L - T. \quad (41)$$

Determination of t_{S2} : Now the source particle decays at the time t_{S2} at the point $x_{S2} = x_S$ and the neutrino arrives at the time $t_D + \tau/2$ at the point $x_{D2} = x_D + \bar{\beta}_D \tau/2$. This leads to

$$x_{D2} = x_{S2} + t_D + \frac{\tau}{2} - t_{S2} \quad \Rightarrow \quad t_{S2} = t_D + \lambda - L. \quad (42)$$

In this way, taking into account positions and velocities of the source and detector particle and the time of flight of the neutrino, we obtain full agreement with equation (39) derived from a QFT formalism.

It is interesting that even an analogue to the “separation of neutrino wave packets” [19, 20] is contained in our model. To understand this point we have to take a look at equation (17). Up to now we have only taken into account the first three terms on the right-hand side of this equation. However, it is the fourth term that leads to this effect [12]. If we take it into account, the quantity t_c depends on the index j :

$$t_{cj} = t_D - t_S - L - L \frac{m_j^2}{2Q^2}. \quad (43)$$

²We assume that the neutrino travels with the speed of light.

How do we have to interpret of the last term? The velocity of the massive neutrino ν_j with energy \bar{Q} is

$$\beta_j = \frac{\sqrt{\bar{Q}^2 - m_j^2}}{\bar{Q}} \simeq 1 - \frac{m_j^2}{2\bar{Q}^2}. \quad (44)$$

Therefore, it covers the distance L in the time

$$\frac{L}{\beta_j} \simeq L \left(1 + \frac{m_j^2}{2\bar{Q}^2} \right) \equiv L + \delta t_j. \quad (45)$$

Consequently, δt_j is the time delay due to the finite neutrino mass. It has to be incorporated into equation (38):

$$-\lambda < t_{cj} < \lambda + T. \quad (46)$$

If we have two neutrino masses m_i and m_j ($m_i \neq m_j$), both t_{ci} and t_{cj} have to fulfill equation (46) in order to guarantee coherence. Assuming $m_i < m_j$ which in turn implies $t_{ci} > t_{cj}$, coherence between ν_i and ν_j is lost for

$$t_{ci} \in (-\lambda, \lambda + T), \quad t_{cj} < -\lambda \quad \text{or} \quad t_{cj} \in (-\lambda, \lambda + T), \quad t_{ci} > \lambda + T. \quad (47)$$

In practice, however, δt_j is so small that this decoherence effect is irrelevant. For instance, setting $m_j = 0.1 \text{ eV}$, $\bar{Q} = 0.5 \text{ MeV}$, $L = 12000 \text{ km}$, we find $t_j = 0.8 \times 10^{-15} \text{ s}$. This is roughly the maximal possible time delay δt_j for (future) neutrino oscillation experiments on earth.

5 The function $|M_j(t_c, \Delta \bar{E})|^2$

In typical neutrino-oscillation experiments the decay width Γ is very small. For instance, mean lives of fission products in a reactor are rather large. Assuming that the bulk of reactor neutrinos comes from mean lives larger than 1 second, we obtain extremely small decay widths:

$$\Gamma \lesssim \frac{\hbar}{1 \text{ s}} \simeq 6.6 \times 10^{-22} \text{ MeV}. \quad (48)$$

Accelerator neutrinos are produced by the decay of charged pions in which case

$$\Gamma \simeq 2.5 \times 10^{-14} \text{ MeV}. \quad (49)$$

Since the pions are relativistic, the effective width $\Gamma m_S / \bar{E}_S$ is even smaller by several orders of magnitude.

The smallness of Γ in the denominator in equation (10b) suggests a potential source of decoherence. Indeed, inspecting equation (28) we are tempted deduce a coherence condition

$$\left| (\bar{\beta}_S - \bar{\beta}_D) \frac{\Delta m^2}{2\bar{Q}} \right| \lesssim \frac{\Gamma m_S}{\bar{E}_S}. \quad (50)$$

This inequality could be violated even for thermal velocities $\bar{\beta}_S$, $\bar{\beta}_D$ and the small quantities

$$\frac{\Delta m_{\text{atm}}^2}{2\bar{Q}} \lesssim 2.5 \times 10^{-15} \text{ MeV} \quad \text{and} \quad \frac{\Delta m_{\text{sol}}^2}{2\bar{Q}} \lesssim 7 \times 10^{-17} \text{ MeV} \quad (51)$$

whose upper limits have been obtained from $\bar{Q} \gtrsim 0.5 \text{ MeV}$ and a three-neutrino fit [21, 22] to all available data yielding

$$\Delta m_{\text{atm}}^2 \sim 2.5 \times 10^{-3} \text{ eV}^2 \quad \text{and} \quad \Delta m_{\text{sol}}^2 \sim 7 \times 10^{-5} \text{ eV}^2. \quad (52)$$

However, the coherence condition above is fictitious.

To prove this claim we consider the dependence on $\Delta \bar{E}$ of the function

$$|M_j(t_c, \Delta \bar{E})|^2 = \frac{(2\pi)^2}{(1 - \bar{\beta}_S)^2(1 - \bar{\beta}_D)^2} \left| \frac{1 - e^{-i\mathcal{T}(t_c)(z_{1j} - z_{2j})}}{z_{1j} - z_{2j}} \right|^2 \quad (53)$$

that occurs in the event rate or total cross section of the compound process. With equations (35) and (36) it can easily be shown that

$$\mathcal{T}(t_c) = \text{length}(\mathcal{I}_c \cap \mathcal{I}_\lambda) \quad (54)$$

for both $T < 2\lambda$ and $T > 2\lambda$. The intervals \mathcal{I}_c and \mathcal{I}_λ are defined in equation (34). Therefore, \mathcal{T} rises linearly from zero to T if $T < 2\lambda$ or to 2λ if $T > 2\lambda$, then remains constant, and thereafter decreases linearly to zero.

Actually, if t_c is fixed, $|M_j(t_c, \Delta \bar{E})|^2$ is proportional to the probability distribution of $\Delta \bar{E}$. Its maximum will give the preferred value of $\Delta \bar{E}$. It will approximately be reached at $\text{Re}(z_{1j} - z_{2j}) = 0$ because Γ is extremely small. According to equation (28), the corresponding value of $\Delta \bar{E}$ depends on m_j . But whenever we choose a t_c such that $|M_j(t_c, \Delta \bar{E})|^2$ is safely away from zero, the order of magnitude of its width in the variable $\Delta \bar{E}$ is given by $1/T$ or $1/\tau$. Since it is reasonable to assume that both $1/T$ and $1/\tau$ are much larger than the quantities in equation (51), the peak of $|M_j(t_c, \Delta \bar{E})|^2$ as a function of $\Delta \bar{E}$ is so broad that its maximum is practically independent of m_j .

So from this consideration we conclude that $M_j(t_c, \Delta \bar{E})$ has nothing to do with coherence but it determines the goodness of total energy conservation, *i.e.* how well $\Delta \bar{E} = 0$ is fulfilled. Clearly, as discussed above, in terms of order of magnitude, energy conservation cannot be better than $1/\tau$ or $1/T$. In the further discussion we neglect all small quantities in $M_j(t_c, \Delta \bar{E})$. In addition, for practical purposes we introduce

Assumption 5: We assume that, apart from thermal motion, the detector particle is at rest.

Altogether we resort to the approximation

$$\bar{\beta}_D = 0, \quad \bar{E}_D = m_D, \quad z_{1j} - z_{2j} = \frac{\Delta \bar{E}}{1 - \bar{\beta}_S}, \quad \lambda = \frac{\tau}{2}. \quad (55)$$

Note that we do not fix the value of $\bar{\beta}_S$ because we want the following discussion to be valid for both reactor and accelerator neutrinos. With the above approximations, $M_j(t_c, \Delta \bar{E})$ is independent of m_j and we drop the index j . In this way, we obtain

$$|M(t_c, \Delta \bar{E})|^2 = (2\pi)^2 \times \left| \frac{1 - e^{-i\mathcal{T}(t_c)\Delta \bar{E}/(1 - \bar{\beta}_S)}}{\Delta \bar{E}} \right|^2. \quad (56)$$

When a neutrino event is recorded in the detector, it is unknown when the corresponding source particle has been produced. Suppose source particles are created at constant rate N_S . Then in a time interval dt_S the number of source particles created is $N_S dt_S$. Therefore, in order to compute the event rate in the detector, one has to integrate over t_S or, equivalently, over t_c . Thus we have to compute the integral

$$\int_{-\lambda}^{\lambda+T} dt_c |M(t_c, \Delta \bar{E})|^2 = 4 \times (2\pi)^2 \int_{-\lambda}^{\lambda+T} dt_c \left(\frac{\sin [\mathcal{T}(t_c) \Delta \bar{E} / (2(1 - \bar{\beta}_s))]}{\Delta \bar{E}} \right)^2. \quad (57)$$

In order to present the result of the integration we define the functions

$$\delta_1(E, t) = t f_1(Et) \quad \text{and} \quad \delta_2(E, t) = t f_2(Et) \quad (58)$$

with

$$f_1(y) = \frac{1}{\pi} \left(\frac{\sin y}{y} \right)^2 \quad \text{and} \quad f_2(y) = \frac{2}{\pi} \left(\frac{1}{y^2} - \frac{\sin y}{y^3} \right), \quad (59)$$

respectively. Note that

$$\int_{-\infty}^{\infty} dy f_1(y) = \int_{-\infty}^{\infty} dy f_2(y) = 1, \quad (60)$$

whence it follows that

$$\lim_{t \rightarrow \infty} \delta_1(E, t) = \lim_{t \rightarrow \infty} \delta_2(E, t) = \delta(E). \quad (61)$$

The integration yields

$$\int_{-\lambda}^{\lambda+T} dt_c |M(t_c, \Delta \bar{E})|^2 = (2\pi)^3 \Delta t [(\tau - T) \delta_1(\Delta \bar{E}, \Delta t/2) + T \delta_2(\Delta \bar{E}, \Delta t)] \quad \text{for } T < \tau \quad (62)$$

and

$$\begin{aligned} & \int_{-\lambda}^{\lambda+T} dt_c |M(t_c, \Delta \bar{E})|^2 \\ &= (2\pi)^3 \frac{\tau}{1 - \bar{\beta}_S} \left[(T - \tau) \delta_1 \left(\Delta \bar{E}, \frac{\tau}{2(1 - \bar{\beta}_S)} \right) + \tau \delta_2 \left(\Delta \bar{E}, \frac{\tau}{1 - \bar{\beta}_S} \right) \right] \\ & \text{for } T > \tau. \end{aligned} \quad (63)$$

We expect that for all practical purposes the approximation

$$\delta_1(\Delta \bar{E}, t/2) \simeq \delta_2(\Delta \bar{E}, t) \simeq \delta(\Delta \bar{E}) \quad \text{for } t = \Delta t, \quad \frac{\tau}{1 - \bar{\beta}_S} \quad (64)$$

holds. Consequently, both equations (62) and (63) approximately yield

$$\int_{-\lambda}^{\lambda+T} dt_c |M(t_c, \Delta \bar{E})|^2 \simeq (2\pi)^3 \tau \Delta t \delta(\Delta \bar{E}) \quad (65)$$

with a δ -function $\delta(\Delta \bar{E})$. Alternatively we may define

$$\int_{-\lambda}^{\lambda+T} dt_c |M(t_c, \Delta \bar{E})|^2 = (2\pi)^3 \tau \Delta t \bar{\delta}(\Delta \bar{E}), \quad (66)$$

where the definition of $\bar{\delta}(\Delta \bar{E})$ can be read off from equations (62) and (63). In $\bar{\delta}(\Delta \bar{E})$ we have suppressed the dependence on τ , Δt and $\bar{\beta}_S$ that disappears in the limit $\bar{\delta}(\Delta \bar{E}) \rightarrow \delta(\Delta \bar{E})$.

6 Reactor and accelerator neutrinos

In addition to equation (55), for reactor neutrinos a reasonable approximation is

$$\bar{\beta}_S = 0, \quad T = \Delta t, \quad \bar{E}_S = m_S, \quad (67)$$

while for accelerator neutrinos, where the typical source particle is a relativistic charged pion and $m_S = m_\pi$, we have

$$1 - \bar{\beta}_S \simeq \frac{m_S^2}{2\bar{E}_S^2}, \quad T \simeq \frac{m_S^2}{2\bar{E}_S^2} \Delta t. \quad (68)$$

In order to get feeling for the orders of magnitude, we choose some reasonable numbers for accelerator neutrinos: $m_S = m_\pi \simeq 140 \text{ MeV}$, $\bar{E}_S = 5 \text{ GeV}$, and for the length of the decay tunnel we assume 300 m. Since the pion is relativistic, with this length of the decay tunnel we obtain $\Delta t \simeq 10^{-6} \text{ s}$. Therefore, using equation (68), we find $T \simeq 4 \times 10^{-10} \text{ s}$. Moreover, with $\Gamma \simeq 2.5 \times 10^{-14} \text{ MeV}$ we obtain $\Gamma m_S / \bar{E}_S \simeq 0.7 \times 10^{-15} \text{ MeV}$.

Let us collect all terms in the event rate of the compound process we have discussed and include the factors that we have left out so far:

$$\frac{1}{(2\pi)^3 2\bar{E}_S} \times (2\pi)^6 |\psi_S|^2 |\mathcal{M}_S|^2 \prod_{i=1}^{n_S-1} \frac{d^3 p'_i}{(2\pi)^3 2E'_{pi}} \quad (69a)$$

$$\times \frac{1}{(2\pi)^8} \left(\frac{2\pi^2}{L} \right)^2 P_{\nu_\alpha \rightarrow \nu_\beta} \quad (69b)$$

$$\times \frac{1}{(2\pi)^3 2\bar{E}_D} \times (2\pi)^6 |\psi_D|^2 |\mathcal{M}_D|^2 \prod_{j=1}^{n_D} \frac{d^3 k'_j}{(2\pi)^3 2E'_{kj}} \quad (69c)$$

$$\times (2\pi)^3 \tau \Delta t \bar{\delta}(\bar{E}_S - \bar{Q} - E'_S). \quad (69d)$$

The symbol \mathcal{M}_S denotes the matrix element of the source process with n_S particles in the final state, including the neutrino. With analogous meanings, \mathcal{M}_D and n_D refer to the detection process. In particular, in equation (69) we have taken into account all factors π . Equation (69b) stems from the neutrino propagator in the asymptotic limit $L \rightarrow \infty$ [16] and is multiplied by the neutrino oscillation probability that originates in the phase $\exp(-im_j^2 L / (2\bar{Q}))$ of equation (29). Clearly, in the end we have to arrive at an expression of the form

$$\text{diff } \Gamma \times \frac{1}{L^2} P_{\nu_\alpha \rightarrow \nu_\beta} \times d\sigma_D \quad \text{with} \quad P_{\nu_\alpha \rightarrow \nu_\beta} = \left| \sum_j U_{\beta j} U_{\alpha j}^* e^{-iLm_j^2/(2\bar{Q})} \right|^2, \quad (70)$$

diff Γ being some differential decay rate and $d\sigma_D$ the infinitesimal cross section of the detection process. The following steps have to be performed to achieve this aim.

Neutrino momentum and propagator: In our approximation, neutrinos are massless in the source and detection process. Moreover, we have to insert the numerator of the neutrino propagator that we have left out so far, *cf.* appendix A, that is

$$\not{q} = \sum_s u(q, s) \bar{u}(q, s) \quad \text{with} \quad q = \begin{pmatrix} \bar{Q} \\ \vec{q} \end{pmatrix}, \quad \vec{q} = \bar{Q} \vec{\ell}. \quad (71)$$

In the sum over s only the negative helicity contributes and provides the spinors $\bar{u}(q, -)$ for the neutrino in the final state of the source process and $u(q, -)$ for the neutrino in the initial state of the detection process. These spinors are not visible in equation (69) because they are part of \mathcal{M}_S and \mathcal{M}_D , respectively.

Rates: In order to obtain a decay rate in the source process, we have to divide equation (69) by Δt . Similarly, we divide by τ to arrive at an event rate in the detection process.

Wave functions: The disturbing presence of the wave functions ψ_S and ψ_D can be remedied by assuming that they are strongly peaked around \vec{p}_{in} and $\vec{k}_{\text{in}} = \vec{0}$, respectively. Choosing the source process for definiteness, the usual trick [23]

$$\begin{aligned}
& (2\pi)^3 \int d^3p \int d^3p' \delta(\vec{p} - \vec{q} - \vec{p}'_S) \delta(\vec{p}' - \vec{q} - \vec{p}'_S) \psi_S(\vec{p}) \psi_S^*(\vec{p}') \cdots \\
&= (2\pi)^3 \int d^3p \int d^3p' \delta(\vec{p} - \vec{q} - \vec{p}'_S) \delta(\vec{p}' - \vec{p}) \psi_S(\vec{p}) \psi_S^*(\vec{p}') \cdots \\
&= (2\pi)^3 \int d^3p |\psi_S(\vec{p})|^2 \delta(\vec{p} - \vec{q} - \vec{p}'_S) \cdots \\
&\simeq (2\pi)^3 \delta(\vec{p}_{\text{in}} - \vec{q} - \vec{p}'_S) \cdots
\end{aligned} \tag{72}$$

can be applied. In this way we obtain δ -functions for the spatial momenta in the source and detection process. For the neutrino momentum \vec{q} see equation (71) and the energy of the source particle is given by

$$\bar{E}_S = \sqrt{m_S^2 + \vec{p}_{\text{in}}^2}. \tag{73}$$

Neutrino in the final state: In the source process we need the factor

$$\frac{d^3q}{(2\pi)^3 2\bar{Q}} = \frac{\bar{Q}^2 d\bar{Q} d\Omega}{(2\pi)^3 2\bar{Q}} \tag{74}$$

for obtaining a decay rate. Since the direction between source and decay process is fixed by $\vec{\ell}$, the result will not be the total decay rate but the differential decay rate with respect to the infinitesimal solid angle $d\Omega$. Note that $L^2 d\Omega$ corresponds to an infinitesimal area at the detection process. Similarly, as we will see below, the decay rate will also be differential with respect to $d\bar{Q}$. Thus, effectively a factor $\bar{Q}/(2(2\pi)^3)$ is inserted and, in the further discussion, we have to remember to compensate for it later.

Neutrino in the initial state: In the detection process we need a factor $1/(2\bar{Q})$. So the total compensating factor is now $4(2\pi)^3$.

Neutrino energy: The neutrino energy has to satisfy equation (20). This is achieved by inserting the integration $\int d\bar{Q} \delta(\bar{Q} + m_D - E'_D)$ into equation (69). For the differential decay rate with respect to $d\bar{Q}$ we simply drop $\int d\bar{Q}$ and arrive at $d^2\Gamma/(d\Omega d\bar{Q})$. The energy δ -function is attributed to the detection process.

Factors of 2π : Finally, we collect all numerical factors. These are the total compensating factor, the factor from equation (69b) and that from equation (69d), leading to

$$4(2\pi)^3 \times \frac{(2\pi^2)^2}{(2\pi)^8} \times (2\pi)^3 = (2\pi)^2. \quad (75)$$

These are just the two factors 2π that are needed in the context of energy conservation:

$$2\pi \bar{\delta}(\bar{E}_S - \bar{Q} - E'_S) \times 2\pi \delta(\bar{Q} + m_D - E'_D). \quad (76)$$

We find that our formalism automatically leads to the correct number of factors 2π .

In summary, we have obtained the differential decay rate

$$\begin{aligned} \text{diff } \Gamma \equiv \frac{d^2\Gamma}{d\Omega d\bar{Q}} &= \frac{1}{2\bar{E}_S} \int \prod_{i=1}^{n_S-1} \frac{d^3p'_i}{(2\pi)^3 2E'_{pi}} \\ &\times (2\pi)^4 \delta(\vec{p}_{\text{in}} - \vec{q} - \vec{p}'_S) \bar{\delta}(\bar{E}_S - \bar{Q} - E'_S) |\mathcal{M}_S|^2 \times \frac{\bar{Q}^2}{(2\pi)^3 2\bar{Q}} \end{aligned} \quad (77)$$

and the detection cross section

$$d\sigma_D = \frac{1}{2m_D 2\bar{Q}} \times (2\pi)^4 \delta(\vec{k}_{\text{in}} + \vec{q} - \vec{k}'_S) \delta(\bar{Q} + m_D - E'_D) |\mathcal{M}_D|^2 \prod_{j=1}^{n_D} \frac{d^3k'_j}{(2\pi)^3 2E'_{kj}}. \quad (78)$$

The correct energy dimensions of these expressions can be checked with

$$\dim |\mathcal{M}_S|^2 = 6 - 2n_S, \quad \dim |\mathcal{M}_D|^2 = 4 - 2n_D. \quad (79)$$

The integral symbol in equation (77) indicates that in the source process usually the final particles are not measured and, therefore, one has to sum over all final momenta except the neutrino momentum.

7 Conclusions

In this paper we have introduced a minimal QFT model of neutrino oscillations that attempts to take into account the environment in the neutrino production and detection process by assuming that production and detection are “interrupted” by collisions with surrounding atoms and molecules. In the case of reactor neutrino production and neutrino detection the collisions take place in a thermal environment while in the case of accelerator neutrinos the charged-pion decay is interrupted when the trajectory of the pion intersects the end of the decay tunnel. These “interruptions” are parameterized in time by Heaviside functions such that the source particle has a time interval of Δt for uninterrupted decay while the detector particle has a time interval τ for undisturbed measurement.³ Our main results are the following:

³In [15] Gaussian wave packets are used instead of Heaviside functions and the final states are described by Gaussians instead of plane waves. Otherwise the topics and results in the present paper have some overlap with [15].

- Oscillations occur in space, *not* in time.
- In realistic situations there is no decoherence in $\mathcal{A}_{\nu_\alpha \rightarrow \nu_\beta}$ —*cf.* section 4.⁴
- Notwithstanding that we work in a QFT model, the resulting correlation between the time of the production of the neutrino source particle and neutrino detection completely coincides with that of a classical consideration—*cf.* section 4.
- Our formalism correctly reproduces the factorization of the event rate of the compound production–detection process into the decay rate of the source particle, the neutrino oscillation probability and the detection cross section—see equations (70), (77) and (78).

The present notes are a substantial generalization of the discussion in [12] where roughly speaking the special case $T \ll \tau$, $\Delta t \Gamma \gg 1$ has been considered.

The following interesting observations concerning Δt , the time interval in which the source particle decays, and τ , the time interval when the measurement takes place, can be made:

- As discussed in section 5, these time intervals prevent decoherence in the amplitude.
- Since they are arbitrary or unknown, it is gratifying that, in good approximation, in the end they drop out—*cf.* section 5.

Acknowledgement: The author thanks Thomas Schwetz for clarifying discussions.

⁴Decoherence takes place only by “classical averaging” at the probability level in the rate of the compound process [15]. This averaging is caused by measurement inaccuracies or lack of knowledge of momenta and energies in the final state of the detection process and by inaccuracies in the determination of \vec{x}_S and \vec{x}_D .

A Integrations

For reasons discussed in the main body of the paper, we omit the matrix elements of the weak currents. We also assume that the decay of the neutrino source particle produced or present at time t_D is interrupted after a time Δt by collisions with particles in the environment. Therefore, the time window in which the source particle decays has a length of Δt . The decay width of the source particle is denoted by Γ . The uncertainty in the detection time is modelled by the function $\Theta_\tau(t_2 - t_D)$ defined in equation (4). With these assumptions the oscillation amplitude is proportional to

$$\mathcal{A}_{\nu_\alpha \rightarrow \nu_\beta} \propto \sum_j U_{\beta j} U_{\alpha j}^* \int_{t_S}^{t_S + \Delta t} dt_1 \int d^3 x_1 \int_{-\infty}^{\infty} dt_2 \int d^3 x_2 \quad (\text{A.1a})$$

$$\times \int d^3 p \psi_S(\vec{p}) \exp \left[-i p \cdot (x_1 - x_S) - \frac{\Gamma m_S}{2E_S} (t_1 - t_S) + i p'_S \cdot x_1 \right] \quad (\text{A.1b})$$

$$\times \int \frac{d^4 q}{(2\pi)^4} \frac{e^{-iq \cdot (x_2 - x_1)}}{q^2 - m_j^2 + i\epsilon} \quad (\text{A.1c})$$

$$\times \int d^3 k \psi_D(\vec{k}) \exp [-ik \cdot (x_2 - x_D) + i k'_D \cdot x_2] \Theta_\tau(t_2 - t_D). \quad (\text{A.1d})$$

In equation (A.1a), U is the CKM lepton mixing matrix and the propagator for a neutrino with mass m_j occurs in equation (A.1c).⁵ We simplify the amplitude by the variable transformation [24]

$$x'_1 = x_1 - x_S, \quad x'_2 = x_2 - x_D \quad (\text{A.2})$$

and obtain

$$\mathcal{A}_{\nu_\alpha \rightarrow \nu_\beta} \propto e^{i(p'_S \cdot x_S + k'_D \cdot x_D)} \sum_j U_{\beta j} U_{\alpha j}^* \int_0^{\Delta t} dt'_1 \int d^3 x'_1 \int_{-\infty}^{\infty} dt'_2 \int d^3 x'_2 \quad (\text{A.3a})$$

$$\times \int d^3 p \psi_S(\vec{p}) \exp \left[-i p \cdot x'_1 - \frac{\Gamma m_S}{2E_S} t'_1 + i p'_S \cdot x'_1 \right] \quad (\text{A.3b})$$

$$\times \int \frac{d^4 q}{(2\pi)^4} \frac{e^{-iq \cdot (x'_2 - x'_1)} \times e^{-iq \cdot (x_D - x_S)}}{q^2 - m_j^2 + i\epsilon} \quad (\text{A.3c})$$

$$\times \int d^3 k \psi_D(\vec{k}) \exp [-ik \cdot x'_2 + i k'_D \cdot x'_2] \Theta_\tau(t'_2). \quad (\text{A.3d})$$

Integration over \vec{x}'_1 and \vec{x}'_2 leads to the product of delta functions

$$\delta(\vec{p} - \vec{p}'_S - \vec{q}) \times \delta(\vec{k} - \vec{k}'_D + \vec{q}). \quad (\text{A.4})$$

The subsequent integrations over \vec{p} and \vec{k} result in the substitutions

$$\vec{p} = \vec{p}'_S + \vec{q} \quad \text{and} \quad \vec{k} = \vec{k}'_D - \vec{q}, \quad (\text{A.5})$$

⁵Since we do not consider the weak matrix elements for the time being, we also leave out the numerator $\not{q} + m_j$ of the neutrino propagator.

respectively. Integration over t'_1 leads to

$$\frac{\exp \left[\left(-i (E_S - E'_S - q^0) - \frac{\Gamma m_S}{2E_S} \right) \Delta t \right] - 1}{-i (E_S - E'_S - q^0) - \frac{\Gamma m_S}{2E_S}}. \quad (\text{A.6})$$

Defining

$$\tilde{\Theta}_\tau(E) = \int dt e^{-iEt} \Theta_\tau(t), \quad (\text{A.7})$$

we obtain

$$\tilde{\Theta}_\tau(E_D - E'_D + q^0) \quad (\text{A.8})$$

from the integration over t_2 . The energies E_S , E'_S , E_D , E'_D denote the time components of p , p' , k , k' , respectively. The remaining two integrations are discussed in the main body of the paper.

B A useful integral

We consider the integral

$$I = 2i \int dx \frac{e^{-icx}}{x - a - ib} \frac{\sin(\lambda x - d)}{\lambda x - d} \quad \text{with} \quad \lambda > 0, \quad b < 0. \quad (\text{B.1})$$

The parameters a , c and d are real as well but can be positive or negative. Defining

$$z_1 = a + ib \quad \text{and} \quad z_2 = \frac{d}{\lambda}, \quad (\text{B.2})$$

we note that the integrand has a pole at z_1 in the complex plane but is analytic at z_2 . Because of this, we can shift the integration path that goes along the x -axis into the complex plane in the vicinity of z_2 and use the residue theorem for the computation of I . Denoting the thus obtained path in the complex plane by C , we split I into

$$I = I_1 - I_2 \quad \text{with} \quad I_1 = \int_C dz \frac{e^{-icz}}{z - a - ib} \frac{e^{i(\lambda z - d)}}{\lambda z - d}, \quad I_2 = \int_C dz \frac{e^{-icz}}{z - a - ib} \frac{e^{-i(\lambda z - d)}}{\lambda z - d}. \quad (\text{B.3})$$

Now the application of the residue theorem is straightforward. For our purpose it is useful to conceive I as a function of c . The result is

$$I(c) = 0 \quad \text{for} \quad c < -\lambda, \quad (\text{B.4a})$$

$$I(c) = -\frac{2\pi i}{\lambda} \frac{e^{-icz_2}}{z_1 - z_2} (1 - e^{-i(\lambda+c)(z_1-z_2)}) \quad \text{for} \quad |c| < \lambda, \quad (\text{B.4b})$$

$$I(c) = -\frac{2\pi i}{\lambda} \frac{e^{-icz_2}}{z_1 - z_2} (e^{i(\lambda-c)(z_1-z_2)} - e^{-i(\lambda+c)(z_1-z_2)}) \quad \text{for} \quad c > \lambda. \quad (\text{B.4c})$$

We emphasize that $I(c)$ is continuous in c .

References

- [1] S. M. Bilenky and B. Pontecorvo, *Lepton mixing and neutrino oscillations*, Phys. Rept. **41** (1978) 225.
- [2] S. M. Bilenky and S. T. Petcov, *Massive neutrinos and neutrino oscillations*, Rev. Mod. Phys. **59** (1987) 671 [Erratum: Rev. Mod. Phys. **61** (1989) 169, Erratum: Rev. Mod. Phys. **60** (1988) 575].
- [3] M. Zrałek, *From kaons to neutrinos: Quantum mechanics of particle oscillations*, Acta Phys. Polon. B **29** (1998) 3925 [hep-ph/9810543].
- [4] S. M. Bilenky, C. Giunti and W. Grimus, *Phenomenology of neutrino oscillations*, Prog. Part. Nucl. Phys. **43** (1999) 1 [hep-ph/9812360].
- [5] M. Beuthe, *Oscillations of neutrinos and mesons in quantum field theory*, Phys. Rept. **375** (2003) 105 [hep-ph/0109119].
- [6] W. Grimus, *Neutrino Physics – Theory*, Lect. Notes Phys. **629** (2004) 169 [hep-ph/0307149].
- [7] E. Akhmedov, *Quantum mechanics aspects and subtleties of neutrino oscillations*, talk given at the International Conference on the History of the Neutrino, Paris, France, September 5-7, 2018, arXiv:1901.05232 [hep-ph].
- [8] M. Sajjad Athar, S. W. Barwick, T. Brunner, J. Cao, M. Danilov, K. Inoue, T. Kajita, M. Kowalski, M. Lindner and K. R. Long, *et al.*, *Status and perspectives of neutrino physics*, Prog. Part. Nucl. Phys. **124** (2022), 103947 [arXiv:2111.07586 [hep-ph]].
- [9] E. K. Akhmedov and A. Y. Smirnov, *Paradoxes of neutrino oscillations*, Phys. Atom. Nucl. **72** (2009) 1363 [arXiv:0905.1903 [hep-ph]].
- [10] D. Boyanovsky, *Short baseline neutrino oscillations: when entanglement suppresses coherence*, Phys. Rev. D **84** (2011) 065001 [arXiv:1106.6248 [hep-ph]].
- [11] L. Lello and D. Boyanovsky, *Searching sterile neutrinos from π and K decays*, Phys. Rev. D **87** (2013) 073017 [arXiv:1208.5559 [hep-ph]].
- [12] W. Grimus, *Revisiting the quantum field theory of neutrino oscillations in vacuum*, J. Phys. G **47** (2020) 085004 [arXiv:1910.13776 [hep-ph]].
- [13] E. Akhmedov and A. Y. Smirnov, *Damping of neutrino oscillations, decoherence and the lengths of neutrino wave packets*, JHEP **11** (2022) 082 [arXiv:2208.03736 [hep-ph]].
- [14] S. Kovalenko and F. Šimkovic, *Neutrino oscillations in quantum field theory*, arXiv:2212.13635 [hep-ph].
- [15] R. Krüger and T. Schwetz, *Decoherence effects in reactor and Gallium neutrino oscillation experiments – a QFT approach*, arXiv:2303.15524 [hep-ph].

- [16] W. Grimus and P. Stockinger, *Real oscillations of virtual neutrinos*, Phys. Rev. D **54** (1996) 3414 [hep-ph/9603430].
- [17] D. Karamitros and A. Pilaftsis, *Towards a localised S-matrix theory*, arXiv:2208.10425 [hep-th].
- [18] W. Grimus, P. Stockinger and S. Mohanty, *The field-theoretical approach to coherence in neutrino oscillations*, Phys. Rev. D **59** (1999), 013011 [arXiv:hep-ph/9807442].
- [19] C. Giunti, C. W. Kim, J. A. Lee and U. W. Lee, *On the treatment of neutrino oscillations without resort to weak eigenstates*, Phys. Rev. D **48** (1993) 4310 [hep-ph/9305276].
- [20] B. Kayser, *On the quantum mechanics of neutrino oscillation*, Phys. Rev. D **24** (1981) 110.
- [21] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, T. Schwetz and A. Zhou, *The fate of hints: updated global analysis of three-flavor neutrino oscillations*, JHEP **09** (2020), 178 [arXiv:2007.14792 [hep-ph]].
- [22] F. Capozzi, E. Di Valentino, E. Lisi, A. Marrone, A. Melchiorri and A. Palazzo, *Unfinished fabric of the three neutrino paradigm*, Phys. Rev. D **104** (2021) no.8, 083031 [arXiv:2107.00532 [hep-ph]].
- [23] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory* (Addison–Wesley Publishing Company, Reading, Massachusetts, 1995).
- [24] A. Ioannisian and A. Pilaftsis, *Neutrino oscillations in space within a solvable model*, Phys. Rev. D **59** (1999) 053003 [arXiv:hep-ph/9809503 [hep-ph]].