

# The Pseudoinverse of $A = CR$ is $A^+ = R^+C^+$ (?)

Michał P. Karpowicz  
michal.karpowicz@nask.pl  
NASK PIB

Gilbert Strang  
gilstrang@gmail.com  
MIT

## Abstract

This paper gives three formulas for the pseudoinverse of a matrix product  $A = CR$ . The first is sometimes correct, the second is always correct, and the third is almost never correct. But that third randomized pseudoinverse  $A_r^+$  may be very useful when  $A$  is a very large matrix.

1.  $A^+ = R^+C^+$  when  $A = CR$  and  $C$  has independent columns and  $R$  has independent rows.
2.  $A^+ = (C^+CR)^+(CRR^+)^+$  is always correct.
3.  $A_r^+ = (P^T CR)^+ P^T CR Q (CR Q)^+ = A^+$  only when  $\text{rank}(P^T A) = \text{rank}(AQ) = \text{rank}(A)$  with  $A = CR$ .

The statement in the title is not generally true, as the following example shows:

$$C = \begin{bmatrix} 1 & 0 \end{bmatrix} \quad R = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad (CR)^+ = \begin{bmatrix} 1 & 1 \end{bmatrix} \quad R^+C^+ = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \end{bmatrix}.$$

Here,  $C$  has a full row rank and  $R$  has a full column rank. We need to have it the other way around in Theorem 1. Then Theorem 2 will give a formula for the pseudoinverse of  $A = CR$  that applies in every case.

When the  $m$  by  $r$  matrix  $C$  has  $r$  independent columns (full column rank  $r$ ), and the  $r$  by  $n$  matrix  $R$  has  $r$  independent rows (full row rank  $r$ ), the pseudoinverse  $C^+$  is the left inverse of  $C$ , and the pseudoinverse  $R^+$  is the right inverse of  $R$ :

$$C^+ = (C^T C)^{-1} C^T \quad \text{has} \quad C^+ C = I_r \quad \text{and} \quad R^+ = R^T (R R^T)^{-1} \quad \text{has} \quad R R^+ = I_r.$$

In this case, the  $m$  by  $n$  matrix  $A = CR$  will also have rank  $r$  and the statement correctly claims that its  $n$  by  $m$  pseudoinverse is  $A^+ = R^+C^+$ .

**Theorem 1.** *The pseudoinverse  $A^+$  of a product  $A = CR$  is the product of the pseudoinverses  $R^+C^+$  (as for inverses) when  $C$  has full column rank and  $R$  has full row rank  $r$ .*

The simplest proof verifies the four Penrose identities [4] that determine the pseudoinverse  $A^+$  of any matrix  $A$ :

$$AA^+A = A \quad A^+AA^+ = A^+ \quad (A^+A)^T = A^+A \quad (AA^+)^T = AA^+ \quad (1)$$

Our goal is a different proof of  $A^+ = R^+C^+$ , starting from first principles. We begin with the “four fundamental subspaces” associated with any  $m$  by  $n$  matrix  $A$  of rank  $r$ . Those are the column space  $\mathbf{C}$  and nullspace  $\mathbf{N}$  of  $A$  and  $A^T$ .

Note that every matrix  $A$  gives an invertible map (Figure 1) from its row space  $\mathbf{C}(A^T)$  to its column space  $\mathbf{C}(A)$ . If  $\mathbf{x}$  and  $\mathbf{y}$  are in the row space and  $A\mathbf{x} = A\mathbf{y}$ , then  $A(\mathbf{x} - \mathbf{y}) = \mathbf{0}$ . Therefore  $\mathbf{x} - \mathbf{y}$  is in the nullspace  $\mathbf{N}(A)$  as well as the row space. So  $\mathbf{x} - \mathbf{y}$  is orthogonal to itself and  $\mathbf{x} = \mathbf{y}$ .

The pseudoinverse  $A^+$  in Figure 2 inverts the row space to column space map in Figure 1 when  $\mathbf{N}(A^T) = \mathbf{N}(A^+)$ . If  $\mathbf{b}_c$  and  $\mathbf{d}_c$  are in the column space  $\mathbf{C}(A)$  and  $\mathbf{x} = A^+\mathbf{b}_c = A^+\mathbf{d}_c$ , then  $A^+(\mathbf{b}_c - \mathbf{d}_c) = \mathbf{0}$  and  $\mathbf{b}_c - \mathbf{d}_c$  is in the nullspace  $\mathbf{N}(A^+)$ . Therefore, if  $\mathbf{N}(A^+) = \mathbf{N}(A^T)$ , then  $\mathbf{b}_c - \mathbf{d}_c$  is in the nullspace of  $A^T$  and the column space of  $A$ , it is orthogonal to itself and  $(\mathbf{b}_c - \mathbf{d}_c)^T(\mathbf{b}_c - \mathbf{d}_c) = 0$  with  $\mathbf{b}_c = \mathbf{d}_c$ . For all vectors  $\mathbf{b}_n$  in the orthogonal complement of  $\mathbf{C}(A)$ , we have  $A^T\mathbf{b}_n = \mathbf{0}$  and  $A^+\mathbf{b}_n = \mathbf{0}$ . Theorem 1 shows why  $\mathbf{N}(A^T) = \mathbf{N}(A^+)$  and  $A^+$  is the inverse map.

That upper map from the row space  $\mathbf{C}(A^T)$  to the column space  $\mathbf{C}(A)$  is inverted by the pseudoinverse  $A^+$  in Figure 2. And the nullspace of  $A^+$  is the same as the nullspace of  $A^T$ —the orthogonal complement of  $\mathbf{C}(A)$ . **Thus  $\mathbf{1}/\mathbf{0} = \mathbf{0}$  for  $A^+$ .** In the extreme case, the pseudoinverse of  $A = \text{zero matrix}$  ( $m$  by  $n$ ) is  $A^+ = \text{zero matrix}$  ( $n$  by  $m$ ).




Figure 1:  $A\mathbf{x}_r = \mathbf{b}$  is in the column space of  $A$  and  $A\mathbf{x}_n = \mathbf{0}$ . The complete solution to  $A\mathbf{x} = \mathbf{b}$  is  $\mathbf{x} = \text{one } \mathbf{x}_r + \text{any } \mathbf{x}_n$ .




Figure 2: The four subspaces for  $A^+$  are the four subspaces for  $A^T$ .

The proof of Theorem 1 begins with a simple Lemma.

**Lemma 1.** *We have  $\mathbf{C}(A) = \mathbf{C}(AB)$  if and only if  $\text{rank}(A) = \text{rank}(AB)$ , and  $\mathbf{N}(B) = \mathbf{N}(AB)$  if and only if  $\text{rank}(B) = \text{rank}(AB)$ .*

**Proof:** Always  $\mathbf{C}(A) \supset \mathbf{C}(AB)$  and  $\mathbf{N}(AB) \supset \mathbf{N}(B)$ . In each pair, equality of dimensions guarantees equality of spaces. The column space can only become smaller, and the nullspace larger.

**Proof of Theorem 1:** We will show that  $\mathbf{N}(A^T) \subset \mathbf{N}(A^+)$  and  $\mathbf{N}(A^T) \supset \mathbf{N}(A^+)$ , and that  $\mathbf{C}(A^T) \subset \mathbf{C}(A^+)$  and  $\mathbf{C}(A^T) \supset \mathbf{C}(A^+)$  when  $A^+ = R^+C^+$ .

**Step 1:**  $\mathbf{N}(A^T) \subset \mathbf{N}(A^+)$ . Every vector  $\mathbf{b}_n$  in the nullspace of  $A^T$  is also in the nullspace of  $A^+$ .

If  $A^T \mathbf{b}_n = 0$ , then  $R^T C^T \mathbf{b}_n = 0$ . Since  $C^T$  has full column rank, we conclude that  $C^T \mathbf{b}_n = (C^T C)^{-1} C^T \mathbf{b}_n = C^+ \mathbf{b}_n = 0$  and  $\mathbf{b}_n$  is in the nullspace of  $C^+$ . However, by Lemma 1, we also have  $\mathbf{N}(C^+) = \mathbf{N}(R^+ C^+) = \mathbf{N}(A^+)$ . Every  $\mathbf{b}_n$  in the nullspace of  $A^T$  is also in the nullspace of  $A^+$ .

**Step 2:**  $\mathbf{N}(A^T) \supset \mathbf{N}(A^+)$ . Every vector  $\mathbf{b}_n$  in the nullspace of  $A^+$  is also in the nullspace of  $A^T$ .

Suppose that  $A^+ \mathbf{b}_n = R^+ C^+ \mathbf{b}_n = 0$ . Since  $RR^+ = I_r$ , we have  $AA^+ \mathbf{b}_n = CRR^+ C^+ \mathbf{b}_n = C(C^T C)^{-1} C^T \mathbf{b}_n = 0$ . Since  $C(C^T C)^{-1}$  has full column rank, we see that  $C^T \mathbf{b}_n = 0$ , so  $\mathbf{b}_n$  is orthogonal to every column of  $C$ . Therefore  $\mathbf{b}_n \in \mathbf{N}(A^T)$ .

**Step 3:**  $\mathbf{C}(A^T) \subset \mathbf{C}(A^+)$ . Every vector  $\mathbf{x}$  in the row space of  $A$  is also in the column space of  $A^+$ .

We have  $\mathbf{x} = A^T \mathbf{y} = R^T C^T \mathbf{y} = R^T [(RR^T)^{-1} RR^T] C^T \mathbf{y} = R^+ R (R^T C^T \mathbf{y}) = R^+ R \mathbf{x}$ . So by Lemma 1, we conclude that  $\mathbf{x} \in \mathbf{C}(R^+) = \mathbf{C}(R^+ C^+) = \mathbf{C}(A^+)$ .

**Step 4:**  $\mathbf{C}(A^T) \supset \mathbf{C}(A^+)$ . Every vector  $\mathbf{x}$  in the column space of  $A^+$  is also in the row space of  $A$ .

Given  $\mathbf{b} = A\mathbf{x} = CR\mathbf{x}$ , we have

$$\mathbf{x} = A^+ \mathbf{b} = R^+ C^+ CR\mathbf{x} = R^T (RR^T)^{-1} (C^T C)^{-1} C^T CR\mathbf{x} = R^T (RR^T)^{-1} R\mathbf{x}$$

So again by Lemma 1, we conclude that  $\mathbf{x} \in \mathbf{C}(R^T) = \mathbf{C}(R^T C^T) = \mathbf{C}(A^T)$ .

An  $m$  by  $n$  matrix of rank  $r$  has many factorizations. The simplest form of  $A = CR$  fills  $C$  with the first  $r$  independent columns of  $A$ . Those columns are a basis for the column space of  $A$ . Then column  $j$  of  $R$  specifies the combination of columns of  $C$  that produces column  $j$  of  $A$ . This example has column 3 = column 1 + column 2:

$$A = \begin{bmatrix} 1 & 4 & 5 \\ 2 & 3 & 5 \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix} = CR.$$

That matrix  $R$  contains the nonzero rows of the **reduced row echelon form** of  $A$ . It also reveals nullspace  $\mathbf{N}(A) = \mathbf{N}(R)$ .

$A = CR$  expresses this classical “elimination by row operations” as a matrix factorization [5, 6]. It is slower and less stable numerically than the SVD, but rational  $A$  produces rational  $C$  and  $R$  and  $C^+$  and  $R^+$ . The following useful formula shows that explicitly:

$$A^+ = R^+ C^+ = R^T (RR^T)^{-1} (C^T C)^{-1} C^T = R^T (C^T A R^T)^{-1} C^T. \quad (2)$$

The inverses of square rational matrices  $RR^T$  and  $C^T C$  are formed by elementary operations and division by a rational determinant. Therefore, the pseudoinverse of  $A$  is rational when  $A$  is rational.

The ranks of  $A^+$  and  $A$  are equal when  $A^+$  is the inverse map. Then,

$$\mathbf{b} = A\mathbf{x} + (I_m - AA^+)\mathbf{w}$$

is a solution of  $A^+ \mathbf{b} = \mathbf{x}$  for arbitrary  $\mathbf{w}$ . We have  $A^+ \mathbf{b} = A^+ A\mathbf{x} = \mathbf{x}$ , since  $A^+ AA^+ = A^+$  when  $A^+$  and  $A$  have equal ranks.

That is not always the case and not always necessary. Consider the rank-deficient

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \end{bmatrix} = C_0 R_0.$$

When we complete  $C_0$  and  $R_0$  to get square invertible  $[C_0, C_1]$  and  $[R_0, R_1]^T$ , then the generalized  $A = CR$  becomes

$$A = \begin{bmatrix} C_0 & C_1 \end{bmatrix} \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} R_0 \\ R_1 \end{bmatrix} = \bar{C} \bar{R}$$

and its generalized inverse is

$$G = \begin{bmatrix} R_0 \\ R_1 \end{bmatrix}^{-1} \begin{bmatrix} I_r & Z_{11} \\ Z_{21} & Z_{22} \end{bmatrix} \begin{bmatrix} C_0 & C_1 \end{bmatrix}^{-1}$$

for arbitrary  $Z_{ij}$  of proper size [8]. The first Penrose identity holds,  $AGA = A$ , but the second one gives

$$GAG = \begin{bmatrix} R_0 \\ R_1 \end{bmatrix}^{-1} \begin{bmatrix} I_r & Z_{12} \\ Z_{21} & Z_{21}Z_{12} \end{bmatrix} \begin{bmatrix} C_0 & C_1 \end{bmatrix}^{-1}.$$

We have  $GAG = G$  only if  $Z_{22} = Z_{21}Z_{12}$ . Then  $A$  and  $G$  have equal ranks and  $G$  is the inverse map. Otherwise, the nullspaces  $\mathbf{N}(A^T)$  and  $\mathbf{N}(G)$  are different. For

$$A^T = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \end{bmatrix} = C_2 R_2$$

the nullspace  $\mathbf{N}(A^T) = \mathbf{N}(R_2)$  is spanned by  $(0, 1, 0)^T$  and  $(0, 0, 1)^T$ . For:

$$G = \begin{bmatrix} 1 & 3 & 2 \\ 3 & 3 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2/3 \end{bmatrix} = C_3 R_3,$$

we see that  $\mathbf{N}(G) = \mathbf{N}(R_3)$  is spanned by  $(0, -2/3, 0)^T$ . Here  $AGA = A$ , but  $GAG$  is not equal to  $G$  with  $1 = \text{rank}(A) < \text{rank}(G) = 2$ . Also, neither  $AG$  nor  $GA$  gives an identity matrix. Still,

$$\mathbf{x} = G\mathbf{b} + (I_n - GA)\mathbf{z}$$

solves  $A\mathbf{x} = AG\mathbf{b} + (A - AGA)\mathbf{z} = AG\mathbf{b} = \mathbf{b}$  for arbitrary  $\mathbf{z}$ .

The necessary and sufficient conditions for  $A^+ = (CR)^+ = R^+C^+$  are surprisingly complex. Greville formulated them in [7]. The reverse order law for pseudoinverse,  $(CR)^+ = R^+C^+$ , holds if and only if

$$\mathbf{C}(RR^TC^T) \subset \mathbf{C}(C^T) \quad \text{and} \quad \mathbf{C}(C^TCR) \subset \mathbf{C}(R). \quad (3)$$

Matrix  $R$  maps  $\mathbf{x}_r = R^TC^T\mathbf{y}$  from the row space  $\mathbf{C}(A^T)$  into the row space  $\mathbf{C}(C^T)$ . Then matrix  $C$  takes  $R\mathbf{x}_r$  to the column space  $\mathbf{C}(A)$ , so we have  $CR\mathbf{x}_r = A\mathbf{x}_r = \mathbf{b}_c$ . Matrix  $C^T$  maps  $\mathbf{b}_c$  into the column space  $\mathbf{C}(R)$ . Then  $R^TC^T\mathbf{b}_c$  is also in the row space  $\mathbf{C}(A^T)$ . The inverse map  $C^+$  identifies  $R\mathbf{x}_r$  in the column space  $\mathbf{C}(R)$  and then  $R^+$  gives  $\mathbf{x}_r = R^+C^+\mathbf{b}_c$  in Figure 3. The reverse order law demands that

$$C^+C(RA^T) = RA^T \quad \text{and} \quad RR^+(C^TA) = C^TA. \quad (4)$$

It follows that  $C$  and  $R$  solve the two-sided projection equation:

$$C^+C(RR^TC^T)RR^+ = RR^TC^TC. \quad (5)$$

For the full-rank factorization of Theorem 1 the equation holds with  $C^+C = I_r = RR^+$ .




Figure 3: Given a full rank decomposition  $A = CR$ , matrix  $R$  maps  $\mathbf{C}(A^T)$  into  $\mathbf{C}(C^T)$  and matrix  $C^T$  maps  $\mathbf{C}(A)$  into  $\mathbf{C}(R)$  when  $(CR)^+ = R^+C^+$ .




Figure 4: The pseudoinverse  $A^+ = (CR)^+ = (C^+CR)^+(CRR^+)^+$  decomposes into the product of the pseudoinverse of  $R$  projected on the row space of  $C^T$  and the pseudoinverse of  $C^T$  projected on the column space of  $R$ .

In general, the inverse map  $A^+ = (CR)^+$  takes  $b_c$  in the column space  $\mathbf{C}(C)$  and projects it into  $Rx_r$  in the part of the column space  $\mathbf{C}(R)$  intersecting with  $\mathbf{C}(C^T)$ . That is ensured by  $(C^+RR^+)^+$ , since by the first principles

$$\mathbf{C}((CRR^+)^+) = \mathbf{C}((CRR^+)^T) = RR^+\mathbf{C}(C^T) = RR^+\mathbf{C}(C^+).$$

Then  $(C^+CR)^+$  maps  $Rx_r$  into the row space  $\mathbf{C}(R^TC^T)$ . We have

$$\mathbf{C}((C^+CR)^+) = \mathbf{C}((C^+CR)^T) = R^T\mathbf{C}(C^+C) = R^T\mathbf{C}(C^T) = \mathbf{C}(R^TC^T).$$

Therefore,  $\mathbf{C}(R^TC^T) = \mathbf{C}(A^+)$  in Figure 4. Also, for any  $b_n$  in the nullspace of  $R^TC^T$  we have  $RR^+C^Tb_n = \mathbf{0}$ , which shows that  $b_n$  is in the nullspace

$$\mathbf{N}(RR^+C^T) = \mathbf{N}((CRR^+)^+) \subset \mathbf{N}((C^+CR)^+(CRR^+)^+) = \mathbf{N}(A^+).$$

Conversely, any  $\mathbf{b}_n$  in the nullspace  $\mathbf{N}((CRR^+)^+) = RR^+\mathbf{N}(C^T)$  is also in the nullspace of  $\mathbf{N}(C^T)$ . It is orthogonal to the columns of  $C$  spanning  $\mathbf{C}(A)$ . Therefore, it is also in the nullspace  $\mathbf{N}(A^T)$ . That proves the following formula for  $A^+$ :

**Theorem 2.** *The pseudoinverse  $A^+$  of a product  $A = CR$  is given by the product*

$$A^+ = (CR)^+ = (C^+CR)^+(CRR^+)^+. \quad (6)$$

The statement in the title of this paper is not generally true. But the statement of Theorem 2 corrects the mistake as the following example shows:

$$C = \begin{bmatrix} 1 & 0 \end{bmatrix} \quad R = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad C^+C = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \quad RR^+ = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$(C^+CR)^+(CRR^+)^+ = \begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = (CR)^+.$$

Theorem 3 gives an even more general statement (following from [3]).

**Theorem 3.** *The pseudoinverse  $A^+$  of a product  $A = CR$  is given by the product*

$$A^+ = ((P^T CR)^+ P^T C) (RQ(CRQ)^+), \quad (7)$$

for any  $P$  and  $Q$  satisfying

$$\text{rank}(P^T A) = \text{rank}(AQ) = \text{rank}(A). \quad (8)$$

**Proof:** When  $\text{rank}(P^T A) = \text{rank}(AQ) = \text{rank}(A)$  for  $P$  and  $Q$  of proper size, then

$$\mathbf{C}((P^T A)^T) = \mathbf{C}((A)^T) \quad \text{and} \quad \mathbf{C}(AQ) = \mathbf{C}(A).$$

In that case,  $(P^T A)^+(P^T A) = A^+A$  and  $(AQ)(AQ)^+ = AA^+$  and (7) is true:

$$A^+ = (P^T A)^+ P^T A Q (A Q)^+.$$

Matrices  $P$  and  $Q$  must be rank-preserving to reconstruct  $A^+$ . But apart from that requirement,  $P$  and  $Q$  can be random matrices. The pseudoinverse of

$$A = \begin{bmatrix} 1 & 4 & 5 \\ 2 & 3 & 5 \end{bmatrix} \text{ is equal to } A^+ = \frac{1}{15} \begin{bmatrix} -8 & 9 \\ 7 & -6 \\ -1 & 3 \end{bmatrix}.$$

Randomly selected matrices

$$P = \begin{bmatrix} 2 & 2 & 2 \\ 1 & 2 & 2 \end{bmatrix} \quad \text{and} \quad Q = \begin{bmatrix} 1 & 1 \\ 0 & 2 \\ 0 & 0 \end{bmatrix}$$

The Pseudoinverse of  $A = CR$  is  $A^+ = R^+C^+$  (?)

---




Figure 5: Computation time and relative approximation error of the randomized  $A_r^+ = \text{rpinv}(A)$ , randomized SVD-based  $A_s^+ = \text{rsvdi}(A)$  and direct  $A^+ = \text{pinv}(A)$  method of calculating the pseudoinverse of  $A$ . In the experiment, we generate a random test matrix  $A = \text{gallery}(\text{'randsvd'}, n, 1e100)$  with  $n$  ranging from 100 to 1000. Then, we calculate the approximation  $A_r^+$  taking random  $m$  by  $p$  matrix  $P$  and  $n$  by  $q$  matrix  $Q$  (in Theorem 3) with  $p = q = \lceil \alpha \cdot n \rceil$  for  $\alpha = 0.4$  (top) and  $\alpha = 0.1$  (bottom). Randomized SVD-based pseudoinverse is calculated based on  $\text{rank } s = \text{rank}(A)$  approximation  $A_s = QU_s\Sigma_sV_s^T$  for  $[Q, \sim] = \text{qr}(A * \text{randn}(n, s))$ .

preserve  $\text{rank}(A) = \text{rank}(P^T A) = \text{rank}(AQ) = 2$ . Therefore,

$$(P^T CR)^+ P^T C = \frac{1}{3} \begin{bmatrix} 2 & -1 \\ -1 & 2 \\ 1 & 1 \end{bmatrix} \quad \text{and} \quad RQ(CRQ)^+ = \frac{1}{5} \begin{bmatrix} -3 & 4 \\ 2 & -1 \end{bmatrix}$$

reconstruct

$$A^+ = \frac{1}{3} \begin{bmatrix} 2 & -1 \\ -1 & 2 \\ 1 & 1 \end{bmatrix} \cdot \frac{1}{5} \begin{bmatrix} -3 & 4 \\ 2 & -1 \end{bmatrix} = \frac{1}{15} \begin{bmatrix} -8 & 9 \\ 7 & -6 \\ -1 & 3 \end{bmatrix}.$$



Random sampling matrices  $P$  and  $Q$  that are not rank-preserving produce a low-rank approximation of  $A^+$  from the samples of bases for the column space and row space of  $A$ . Therefore, the formula in Theorem 3 describes a randomized algorithm approximating  $A^+$ :

```
function Arplus = rpinv(A,p,q)
% rpinv: randomized pseudoinverse of A
[m,n] = size(A);
P = randn(m,p);
Q = randn(n,q);
PTA = P'*A;
AQ = A*Q;
Arplus = pinv(PTA)*PTA*Q*pinv(AQ);
end
```

When  $P$  and  $Q$  are small ( $m$  by  $p$  and  $n$  by  $q$  matrices), it is more time-efficient to compute  $A_r^+$  using two pseudoinverses,  $(P^T A)^+$  and  $(AQ)^+$ , than to calculate  $A^+$  directly. Also, that approach may lead to smaller relative errors (under the circumstances to be investigated) than the randomized SVD method while maintaining similar time efficiency. Figure 5 demonstrates these points by comparing each method.

We have given three formulas for the pseudoinverse of a matrix product  $A = CR$  and derived them from the four fundamental subspaces of  $A$ . Every matrix  $A$  gives an invertible map from its row space to its column space. The pseudoinverse  $A^+$  inverts the row space to the column space map when the nullspace of  $A^T$  and  $A^+$  match. The first formula in Theorem 1 is correct if  $C$  has independent columns and  $R$  has independent rows or when Greville's conditions hold. The second in Theorem 2 is always correct. The third in Theorem 3 is correct when the sampling matrices  $P$  and  $Q$  preserve rank. Otherwise, it is not correct but potentially very useful, especially for large  $A$ .

## References

- [1] Adi Ben-Israel and Thomas N.E. Greville, **Generalized Inverses: Theory and Applications**. *Springer* (2003).
- [2] Keaton Hamm and Longxiu Huang, **Perspectives on CUR Decompositions**. *Applied and Computational Harmonic Analysis*, **48** (2020) 1088–1099.
- [3] Michał P. Karpowicz, **A Theory of Meta-factorization**. *arXiv*: 2111.14385 (2021).
- [4] Roger Penrose, **A Generalized Inverse for Matrices**. *Mathematical Proceedings of the Cambridge Philosophical Society*, **51** (1955) 406–413.
- [5] Gilbert Strang and Daniel Drucker, **Three Matrix Factorizations from the Steps of Elimination**. *Analysis and Applications*, **20** (2022) 1147–1157.

- [6] Gilbert Strang, **Introduction to Linear Algebra, 6th edition** (2023), *Wellesley-Cambridge Press*.
- [7] Thomas N.E. Greville, **Note on the Generalized Inverse of a Matrix Product**, *SIAM Review*, **8** (1966) 518–521.
- [8] George Marsaglia and George Styan, **Equalities and inequalities for ranks of matrices**, *Linear and Multilinear Algebra*, **3** (1974) 269–292.