$(\varepsilon, \delta) - \mbox{QUASI-NEGATIVE CURVATURE AND POSITIVITY OF THE CANONICAL BUNDLE}$

KYLE BRODER AND KAI TANG

ABSTRACT. A recent theorem of Diverio–Trapani and Wu–Yau asserts that a compact Kähler manifold with a Kähler metric of quasi-negative holomorphic sectional curvature is projective and canonically polarized. This confirms a long-standing conjecture of Yau. We consider the notion of (ε, δ) –quasi-negativity, generalizing quasi-negativity, and obtain gap-type theorems for $\int_X c_1(K_X)^n > 0$ in terms of the real bisectional curvature and weighted orthogonal Ricci curvature. These theorems are also a generalization of that results in [34] by Zhang-Zheng and in [11] by Chu-Lee-Tam.

1. INTRODUCTION

The Wu-Yau theorem [9, 26, 16, 27, 28, 25] asserts that a compact Kähler manifold with negative holomorphic sectional curvature is projective and canonically polarized. In particular, the canonical bundle K_X is ample and there is a unique invariant Kähler–Einstein metric [1, 31]. Diverio–Trapani [15] relaxed the strict negativity of the holomorphic sectional curvature to quasi-negativity (non-positive everywhere and negative at one point). The crux of the argument is the positivity of the top intersection number

$$\int_{X} c_1(K_X)^n > 0.$$
 (1.1)

Indeed, from [14, Theorem 0.5], (1.1) implies that the canonical bundle K_X is big if we assume that K_X is nef. Hence, X is Moishezon, and by Moishezon's theorem [19], X is projective and of general type. A projective manifold of general type with no rational curves has ample canonical bundle (see [15, Lemma 2.1] or [12, p. 219]).

Key words and phrases. Hermitian manifolds, Kähler manifolds, Wu–Yau theorem, hyperbolic complex manifolds, ample canonical bundle, holomorphic sectional curvature, real bisectional curvature, Moishezon manifolds, manifolds of general type.

The first named author was supported by the Australian Government through the Australian Research Council's Discovery Projects funding scheme (project DP220102530). The second named author was supported by National Natural Science Foundation of China (Grant No.12001490).

In their recent work, Zhang-Zheng [34] considered a natural notion of almost quasinegative holomorphic sectional curvature and extended these theorems to compact Kähler manifolds of almost quasi-negative holomorphic sectional curvature. They also obtained a gap-type theorem for the inequality $\int_X c_1(K_X)^n > 0$ in terms of the holomorphic sectional curvature on compact Kähler manifolds. We will extend these results of Diverio–Trapani [15] and Zhang-Zheng [34] to the Hermitian category. We remind the reader that for non-Kähler Hermitian metrics, the holomorphic sectional curvature is naturally replaced by the *real bisectional curvature*

$$\operatorname{RBC}_{\omega}(\xi) := \frac{1}{|\xi|^2} \sum_{i,j,k,\ell} R_{i\overline{j}k\overline{\ell}} \xi^{i\overline{j}} \xi^{k\overline{\ell}},$$

where ξ is a non-negative (1, 1)-tensor. The real bisectional curvature was introduced by Yang–Zheng [30] in a frame-dependent manner. The above definition is also due to Lee–Streets [17], where it is referred to as the 'complex curvature operator'.

When the metric is Kähler (or, more generally, Kähler-like [29]), the real bisectional curvature is equivalent to the altered holomorphic sectional curvature formally defined in [8], which is comparable to the familiar holomorphic sectional curvature in the sense that they always have the same sign. In general, however, the real bisectional curvature is stronger than the holomorphic sectional curvature. It is not strong enough, however, to dominate the Ricci curvatures (see [30]). For a discussion of these curvatures for connections more general than the Chern connection, namely, the Gauduchon connections, we invite the reader to see [6].

To state the main results, we introduce the following terminology (building from [34]): For a positive constant $\delta > 0$ and a non-empty open set $\mathcal{U} \subset X$ we say that a Hermitian metric η has (ε, δ) -quasi-negative real bisectional curvature (relative to \mathcal{U}) if there is a sufficiently small $\varepsilon > 0$ such that $\text{RBC}_{\eta} \leq \varepsilon$ on X and if $\text{RBC}_{\eta} \leq -\delta$ on \mathcal{U} . For positive constants $\delta_1, \delta_2 > 0$, we say that a Hermitian metric η on a compact Kähler manifold (X, ω) is (i) δ_1 -bounded if there is a smooth function $\psi : X \to \mathbf{R}$ such that $\eta \leq \delta_1 \omega + dd^c \psi$, and (ii) δ_2 -volume non-collapsed on an open set $\mathcal{U} \subset X$ if $\eta^n \geq \delta_2 \omega^n$. If a Hermitian metric is both δ_1 -bounded and δ_2 -volume non-collapsed, we say that η has (δ_1, δ_2) -bounded geometry.

The first main theorem of the present manuscript is the following:

Theorem 1.1. Let X be a compact Kähler manifold. Suppose there is a Hermitian metric η with (δ_1, δ_2) -bounded geometry and (ε, δ) -quasi-negative real bisectional curvature. Then

$$\int_X c_1(K_X)^n > 0$$

If one further assumes that K_X is nef, then K_X is big and X is projective.

The above theorem is a result in this direction that does not require the curvature of the Hermitian metric to have a sign. There are also many interesting results in [32, 33, 34, 23] without assuming a pointwise signed curvature condition. Further, let us emphasize that although the δ_1 -boundedness property is pointwise, it can be interpreted as a bound on the Bott-Chern class represented by the metric, not at the level of the metric. On the other hand, δ_2 -volume non-collapsing is a local weighted volume non-collapsing condition; for instance, it holds if $\eta^n \geq \delta_2 \omega_0^n$ on \mathcal{U} .

Given the perplexing relationship between the holomorphic sectional curvature and the Ricci curvature (c.f., [5]), one can attempt to interpolate between these curvatures by considering the following curvature constraint: In [7] (see also [11]), the authors introduced, for $\alpha, \beta \in \mathbf{R}$ the *weighted orthogonal Ricci curvature*

$$\operatorname{Ric}_{\alpha,\beta}^{\perp}(v) := \frac{\alpha}{|v|_{\omega_0}^2} \operatorname{Ric}_{\omega_0}(v,\overline{v}) + \beta \operatorname{HSC}_{\omega_0}(v),$$

for a (1,0)-tangent vector $v \in T^{1,0}X$.

From Siu's resolution [22] of the Grauert–Riemenschneider conjecture, a compact Kähler manifold with quasi-negative Ricci curvature has big canonical bundle. On the other hand, the Diverio–Trapani [15] extension of the Wu–Yau theorem shows that a compact Kähler manifold with quasi-negative holomorphic sectional curvature has big canonical bundle. The following theorem shows that these results can be interpolated via the weighted orthogonal Ricci curvature, and more generally, extended to the (ε, δ) –quasi-negative situation:

Theorem 1.2. Let (X, ω) be a compact Kähler manifold with (ε, δ) -quasi-negative weighted orthogonal Ricci curvature for $\alpha, \beta \in \mathbf{R}_{>0}$ with at least one positive. Then

$$\int_X c_1(K_X)^n > 0$$

If one further assumes that K_X is nef, then K_X is big and X is projective.

Theorem 1.2 also extends a result of Chu–Lee–Tam [11], where they showed that if a compact Kähler manifold X has a Kähler metric of quasi-negative weighted orthogonal

Ricci curvature (for $\alpha, \beta \in \mathbf{R}_{>0}$), then $\int_X c_1(K_X)^n > 0$.

An alternative means of interpolating between the holomorphic sectional curvature and the Ricci curvature was described by Ni [20]: The so-called k-Ricci curvature of a Kähler metric is the Ricci curvature of the k-dimensional holomorphic subspaces of $T^{1,0}X$. By making use of a result in [11], the method used to prove Theorem 1.1 and Theorem 1.2 also provides the following (see also [23]):

Theorem 1.3. Let (X, ω) be a compact Kähler manifold with (ε, δ) -quasi-negative k-Ricci curvature. Then

$$\int_X c_1(K_X)^n > 0$$

If one further assumes that K_X is nef, then K_X is big and X is projective.

Acknowledgements. The first named author would like to thank his former Ph.D. advisors Ben Andrews and Gang Tian for their support and encouragement. He would also like to thank Simone Diverio, Jeffrey Streets, Finnur Lárusson, Ramiro Lafuente, and James Stanfield for many valuable discussions and communications. The authors would like to thank Yashan Zhang for his support, and for many invaluable discussions and suggestions.

2. Proof of Theorem 1.1

To show that $\int_X c_1(K_X)^n > 0$, it suffices to obtain the estimate

$$\int_{X} (-\operatorname{Ric}_{\omega_0})^n = (2\pi)^n \int_{X} c_1(K_X)^n > 0.$$
(2.1)

Let $\rho : \mathbf{R} \to \mathbf{R}$ be the function defined by $\rho(t) = \frac{1}{n}$ for $t \leq 0$ and $\rho(t) = 1$ for t > 0. For the Hermitian metric η in the statement of Theorem 1.1, let $\kappa_{\eta} : X \to \mathbf{R}$ be the function

$$\kappa_{\eta}(x) := \rho\left(\max_{(\vartheta, v_x) \in \mathcal{F}_X \times \mathbf{R}^n} \operatorname{RBC}_{\eta}(\vartheta, v_x)\right) \cdot \max_{(\vartheta, v_x) \in \mathcal{F}_X \times \mathbf{R}^n} \operatorname{RBC}_{\eta}(\vartheta, v_x),$$

where ϑ is a unitary frame (i.e., a section of the unitary frame bundle \mathcal{F}_X and $v_x \in \mathbf{R}^n$).

Let us also introduce the notation

$$\mu_{\eta} := \max_{x \in X} \max_{(\vartheta, v_x) \in \mathcal{F}_X \times \mathbf{R}^n} \operatorname{RBC}_{\eta}(\vartheta, v_x).$$
(2.2)

($\varepsilon,\delta)\text{-}\textsc{Quasi-negative}$ curvature and positivity of the canonical bundle ~~5

To establish (2.1), we will show that there are constants ε , c_3 , $c_4 > 0$ such that

$$\int_{X} (-\operatorname{Ric}_{\omega_0})^n \geq \int_{X} (n\delta_1 \kappa_\eta \omega_0 - \operatorname{Ric}_{\omega_0})^n - c_4 \varepsilon$$
(2.3)

$$\geq c_3 - c_4 \varepsilon, \tag{2.4}$$

where $\varepsilon > 0$ can be chosen such that $c_3 - c_4 \varepsilon > 0$.

To this end, consider the twisted Wu–Yau continuity method, given by the complex Monge–Ampère equation

$$(t(\omega_0 + \delta_1^{-1} dd^c \psi) - \operatorname{Ric}_{\omega_0} + dd^c \varphi_t)^n = e^{\varphi_t} \omega_0^n.$$
(2.5)

From the assumption of δ_1 -boundedness, we see that

$$\omega_0 + \delta_1^{-1} dd^c \psi \geq \delta_1^{-1} \eta.$$
(2.6)

Set $\omega_t := t(\omega_0 + \delta_1^{-1} dd^c \psi) - \operatorname{Ric}_{\omega_0} + dd^c \varphi_t$, allowing us to write (2.5) as

$$\omega_t^n = e^{\varphi_t} \omega_0^n$$

From (2.6), the metrics ω_t afford the lower bound

$$\operatorname{Ric}_{\omega_t} \geq -\omega_t + t\delta_1^{-1}\eta$$

We will make use of the following Schwarz lemma for holomorphic maps between Hermitian manifolds due to Yang–Zheng [30] (see also [21, 3, 4]):

Lemma 2.1. (Yang–Zheng). Let $f : (X, \omega_g) \to (Y, \eta)$ be a holomorphic map from a Kähler manifold to a Hermitian manifold. If there are constants $C_1, C_2 \in \mathbf{R}$ such that $\operatorname{Ric}_{\omega_g} \geq -C_1\omega + C_2f^*\omega_h$ and $\operatorname{RBC}_{\eta} \leq \mu_{\eta}$, then

$$\Delta_{\omega_g} \log \operatorname{tr}_{\omega_g}(f^*\eta) \geq -C_1 + \left(-\mu_\eta + \frac{C_2}{n}\right) \operatorname{tr}_{\omega_g}(f^*\eta).$$

Taking f to be the identity map, $C_1 = 1$, and $C_2 = t\delta_1^{-1}$ in Lemma 2.1, we have

$$\Delta_{\omega_t} \log \operatorname{tr}_{\omega_t}(\eta) \geq (-\mu_\eta + t(n\delta_1)^{-1}) \operatorname{tr}_{\omega_t}(\eta) - 1.$$
(2.7)

By the maximum principle,

$$\sup_{X} \operatorname{tr}_{\omega_{t}}(\eta) \leq \frac{n\delta_{1}}{t - \mu_{\eta}n\delta_{1}}.$$
(2.8)

For $t > n\delta_1\mu_\eta$, the estimate (2.8) is independent of t. As a consequence, the continuity method admits a smooth solution for $t > n\delta_1\mu_\eta$ (c.f., [25, 32]). Introduce the potential $u_t := \varphi_t + t \delta_1^{-1} \psi$. The crux of the argument is to estimate $\sup_{\mathfrak{U}} u_t$ from below, and $\sup_X u_t$ from above. Indeed, the constant c_3 in (2.4) is given by

$$c_3 := \liminf_{t \to n\delta_1 \mu_\eta} \int_X e^{u_t} \omega_0^n,$$

where μ_{η} is defined in (2.2). The positivity of c_3 demands u_t to not be identically $-\infty$, while the finiteness of u_t requires an upper bound on $\sup_X u_t$. Before obtaining the upper bound on $\sup_X u_t$, we first note that in contrast to the situations considered by Wu–Yau [27], Tosatti–Yang [25], and Diverio–Trapani [15], we do not have a metric with signed curvature. To handle the positive curvature contributions, we make use of *Tian's* α -invariant [24] (see also [34]):

Proposition 2.2. (Tian's α -invariant). Let (X, ω) be a compact Kähler manifold. There is a positive number $\alpha = \alpha(X, [\omega])$ depending only on X and the Kähler class $[\omega]$ such that for any $\beta \in (0, \alpha)$ and any $u \in \mathbb{C}^{\infty}(X, \mathbf{R})$ with $\omega + dd^{c}u > 0$ and $\sup_{X} u = 0$, there is a constant $C_{\beta} = C_{\beta}(\omega)$ such that

$$\int_X e^{-\beta u} \omega^n \leq C_\beta.$$

We now obtain an upper bound on $\sup_X u_t$:

Lemma 2.3. For $t \in (n\delta_1\mu_\eta, 2n\delta_1\mu_\eta]$, we have

$$\sup_{x \in X} u_t(x) \leq \log(2n\delta_1\varepsilon + b_0)^n \leq \log(c_0 + b_0)^n, \tag{2.9}$$

where $c_0 > 0$ is such that Tian's α -invariant satisfies $\alpha(X, c_0\omega_0) \geq 2$, and $\varepsilon \leq c_0/(2n\delta_1)$.

Proof. Since $u_t = \varphi_t + t\delta_1^{-1}\psi$, (2.7) reads

$$e^{u_t - t\delta_1^{-1}\psi}\omega_0^n = e^{\varphi_t}\omega_0^n = (t\omega_0 - \operatorname{Ric}_{\omega_0} + dd^c u_t)^n.$$
 (2.10)

Since the metric ω_0 is smooth, and X is compact, there is some $b_0 \ge 0$ such that

$$\operatorname{Ric}_{\omega_0} \geq -b_0\omega_0. \tag{2.11}$$

From (2.10), (2.11), and the fact that $\psi \leq 0$, at the point $x \in X$ where u_t achieves its maximum, we have

$$\sup_{x \in X} u_t(x) \leq \sup_X \left(t \delta_1^{-1} \psi + \log \frac{(t\omega_0 - \operatorname{Ric}_{\omega_0} + dd^c u_t)^n}{\omega_0^n} \right) \leq \log(t_0 + b_0)^n$$

Since $t \leq 2n\delta_1\mu_\eta \leq 2n\delta_1\varepsilon$, this proves (2.9).

We now want to estimate $\sup_{\mathfrak{U}} u_t$ from below.

Lemma 2.4.

$$\sup_{\mathfrak{U}} u_t \geq n \log(n) + \log\left(\frac{-\int_{\mathfrak{U}} \kappa_{\eta} e^{t(n\delta_1)^{-1}\psi} \left(\frac{\eta^n}{\omega_0^n}\right)^{\frac{1}{n}} \omega_t^n}{\int_X \omega_t^n}\right).$$
(2.12)

Proof. Start by integrating (2.7) over X with respect to the volume form ω_t^n . By the divergence theorem,

$$\int_{X} \omega_{t}^{n} \geq \int_{X} (-\kappa_{\eta} + t(n\delta_{1})^{-1}) \operatorname{tr}_{\omega_{t}}(\eta) \omega_{t}^{n}$$

$$\geq \int_{\mathfrak{U}} (-\kappa_{\eta} + t(n\delta_{1})^{-1}) \operatorname{tr}_{\omega_{t}}(\eta) \omega_{t}^{n} \geq -\int_{\mathfrak{U}} \kappa_{\eta} \operatorname{tr}_{\omega_{t}}(\eta) \omega_{t}^{n},$$

using the fact that $(-\kappa_{\eta} + t(n\delta_1)^{-1}) \operatorname{tr}_{\omega_t}(\eta) > 0$. By the arithmetic-geometric mean inequality,

$$-\int_{\mathfrak{U}} \kappa_{\eta} \operatorname{tr}_{\omega_{t}}(\eta) \omega_{t}^{n} \geq -n \int_{\mathfrak{U}} \kappa_{\eta} \left(\frac{\eta^{n}}{\omega_{t}^{n}}\right)^{\frac{1}{n}} \omega_{t}^{n}$$
$$= -n \int_{\mathfrak{U}} \kappa_{\eta} \left(\frac{\omega_{0}^{n}}{\omega_{t}^{n}}\right)^{\frac{1}{n}} \left(\frac{\eta^{n}}{\omega_{0}^{n}}\right)^{\frac{1}{n}} \omega_{t}^{n}.$$
(2.13)

Using (2.10) in (2.13), we achieve the estimate

$$\int_{X} \omega_{t}^{n} \geq -n \int_{\mathfrak{U}} \kappa_{\eta} e^{-\frac{1}{n}(u_{t}-t\delta_{1}^{-1}\psi)} \left(\frac{\eta^{n}}{\omega_{0}^{n}}\right)^{\frac{1}{n}} \omega_{t}^{n} \\
\geq -n e^{-\frac{1}{n}\sup_{\mathfrak{U}} u_{t}} \int_{\mathfrak{U}} \kappa_{\eta} e^{t(n\delta_{1})^{-1}\psi} \left(\frac{\eta^{n}}{\omega_{0}^{n}}\right)^{\frac{1}{n}} \omega_{t}^{n}.$$
(2.14)

Then (2.12) follows from (2.14).

From the above lemma, it suffices to estimate

$$\frac{-\int_{\mathfrak{U}}\kappa_{\eta}e^{t(n\delta_{1})^{-1}\psi}\left(\frac{\eta^{n}}{\omega_{0}^{n}}\right)^{\frac{1}{n}}\omega_{t}^{n}}{\int_{X}\omega_{t}^{n}}$$
(2.15)

from below. The following lemma gives an estimate for the numerator:

Lemma 2.5. There are positive constants $c_1, \delta_2 > 0$ such that

$$-\int_{\mathfrak{U}}\kappa_{\eta}e^{t(n\delta_{1})^{-1}\psi}\left(\frac{\eta^{n}}{\omega_{0}^{n}}\right)^{\frac{1}{n}}\omega_{t}^{n} \geq \delta_{2}\delta\int_{\mathfrak{U}}e^{u_{t}^{*}}\omega_{0}^{n} \geq \frac{c_{1}\delta_{2}^{\frac{1}{n}}\delta}{n}.$$
 (2.16)

Proof. Let us write $u_t^* := u_t - \sup_X u_t$, so that $\sup_X u_t^* \leq 0$. In this notation, (2.15) reads

$$\frac{-\int_{\mathcal{U}} \kappa_{\eta} e^{t(n\delta_{1})^{-1}\psi} \left(\frac{\eta^{n}}{\omega_{0}^{n}}\right)^{\frac{1}{n}} \omega_{t}^{n}}{\int_{X} \omega_{t}^{n}} = \frac{-\int_{\mathcal{U}} \kappa_{\eta} e^{u_{t}} e^{-\left(1-\frac{1}{n}\right)t\delta_{1}^{-1}\psi} \left(\frac{\eta^{n}}{\omega_{0}^{n}}\right)^{\frac{1}{n}} \omega_{0}^{n}}{\int_{X} e^{u_{t}} e^{-t\delta_{1}^{-1}\psi} \omega_{0}^{n}}$$
$$= \frac{-\int_{\mathcal{U}} \kappa_{\eta} e^{u_{t}^{*}} e^{-\left(1-\frac{1}{n}\right)t\delta_{1}^{-1}\psi} \left(\frac{\eta^{n}}{\omega_{0}^{n}}\right)^{\frac{1}{n}} \omega_{0}^{n}}{\int_{X} e^{u_{t}^{*}} e^{-t\delta_{1}^{-1}\psi} \omega_{0}^{n}} \qquad (2.17)$$

Since $\psi \leq 0$, and $t \in (n\delta_1 \mu_\eta, 2n\delta_1 \mu_\eta]$, we have

$$-\int_{\mathfrak{U}} \kappa_{\eta} e^{u_{t}^{*}} e^{-\left(1-\frac{1}{n}\right)t\delta_{1}^{-1}\psi} \left(\frac{\eta^{n}}{\omega_{0}^{n}}\right)^{\frac{1}{n}} \omega_{0}^{n} \geq -\int_{\mathfrak{U}} \kappa_{\eta} e^{u_{t}^{*}} e^{(1-n)\kappa_{\eta}\psi} \left(\frac{\eta^{n}}{\omega_{0}^{n}}\right)^{\frac{1}{n}} \omega_{0}^{n}$$
$$\geq \delta \int_{\mathfrak{U}} e^{u_{t}^{*}} e^{(1-n)\kappa_{\eta}\psi} \left(\frac{\eta^{n}}{\omega_{0}^{n}}\right)^{\frac{1}{n}} \omega_{0}^{n}, \quad (2.18)$$

where the last inequality follows from the negative curvature estimate $\text{RBC}_{\eta} \leq -\delta_3$ on \mathcal{U} . Let

$$c_1 := \inf \left\{ \int_{\mathcal{U}} e^v \omega_0^n : v \in PSH_{(c_0+b_0)\omega_0}(X), \sup_X v = 0 \right\}.$$
 (2.19)

Then c_1 is a positive constant depending only on \mathcal{U} and ω_0 . Since η has (δ_1, δ_2) -bounded geometry, we have

$$-\int_{\mathfrak{U}}\kappa_{\eta}e^{u_{t}^{*}}e^{-(n-1)\mu_{\eta}\psi}\left(\frac{\eta^{n}}{\omega_{0}^{n}}\right)^{\frac{1}{n}}\omega_{0}^{n} \geq \frac{\delta_{2}^{\frac{1}{n}}\delta}{n}\int_{\mathfrak{U}}e^{u_{t}^{*}}\omega_{0}^{n} \geq \frac{c_{1}\delta_{2}^{\frac{1}{n}}\delta}{n}.$$

From (2.18) and (2.19), this gives the desired lower bound for the numerator in (2.17). $\hfill \Box$

We now complete the estimate for lower bound on $\sup_{\mathfrak{U}} u_t$:

Lemma 2.6. There are constants $c_1, c_2, \delta_1, \delta_2 > 0$ such that

$$\sup_{\mathfrak{U}} u_t \geq n \log(n) + \log\left(\frac{c_1 \delta_2 \delta}{c_2}\right).$$

Proof. For the denominator in (2.17), we will again use Tian's α -invariant [24]. Indeed, first observe that, since $0 < t \leq 2n\delta_1\mu_\eta \leq c_0$, (2.6) implies

$$c_0\omega_0 + t\delta_1^{-1}dd^c\psi \geq t\omega_0 + t\delta_1^{-1}dd^c\psi \geq t\delta_1^{-1}\eta > 0.$$

In other words, $t\delta_1^{-1}\psi$ is $c_0\omega_0$ -plurisubharmonic. Hence, since $c_0 > 0$ is chosen such that Tian's α -invariant satisfies $\alpha(X, c_0\omega_0) \ge 2$, we have

$$\int_{X} e^{u_{t}^{*}} e^{-t\delta_{1}^{-1}\psi} \omega_{0}^{n} \leq \int_{X} e^{-t\delta_{1}^{-1}\psi} \omega_{0}^{n} = c_{0}^{-n} \int_{X} e^{-t\delta_{1}^{-1}\psi} (c_{0}\omega_{0})^{n} \leq c_{2}, \quad (2.20)$$

for some constant $c_2 > 0$ depending only on c_0 and ω_0 . Combining (2.15), (2.17), (2.16), and (2.20), we see that

$$\frac{-\int_{\mathfrak{U}}\kappa_{\eta}e^{t(n\delta_{1})^{-1}\psi}\left(\frac{\eta^{n}}{\omega_{0}^{n}}\right)^{\frac{1}{n}}\omega_{t}^{n}}{\int_{X}\omega_{t}^{n}} \geq \frac{c_{1}\delta_{2}^{\frac{1}{n}}\delta}{nc_{2}}.$$
(2.21)

Inserting (2.21) into (2.12),

$$\sup_{\mathcal{U}} u_t \geq n \log(n) + \log\left(\frac{c_1 \delta_2^{\frac{1}{n}} \delta}{n c_2}\right).$$

We now complete the proof: Let $c_4 > 0$ be the constant such that

$$\int_X \sum_{k=1}^n \binom{n}{k} (2n\delta_1 \varepsilon \omega_0)^k \wedge (-\operatorname{Ric}_{\omega_0})^{n-k} \leq c_4 \varepsilon.$$

Then

$$\int_{X} (2\pi c_1(K_X))^n = \int_{X} (-\operatorname{Ric}_{\omega_0})^n \geq \int_{X} (2n\delta_1 \varepsilon \omega_0 - \operatorname{Ric}_{\omega_0})^n - c_4 \varepsilon$$
$$\geq \lim_{t \searrow n\delta_1 \mu_\eta} (t\omega_0 - \operatorname{Ric}_{\omega_0} + dd^c u_t)^n - c_4 \varepsilon$$
$$= \lim_{t \searrow n\delta_1 \mu_\eta} \int_{X} e^{u_t} e^{-t\delta_1^{-1}\psi} \omega_0^n - c_4 \varepsilon$$
$$\geq \liminf_{t \searrow n\delta_1 \mu_\eta} \int_{X} e^{u_t} \omega_0^n - c_4 \varepsilon$$
$$= c_3 - c_4 \varepsilon.$$

Taking $\varepsilon \leq \min\left\{\frac{c_3}{2c_4}, \frac{c_0}{2n\delta_1}\right\}$ completes the proof.

Recall that a compact complex manifold X is said to be *Moishezon* if X is bimeromorphic to a projective manifold. Any compact complex manifold with a big line bundle $\mathcal{L} \to X$ is Moishezon, with the linear system $|\mathcal{L}|$ furnishing the bimeromorphic map. In particular, a compact complex manifold X of general type (i.e., the canonical bundle K_X is big) is Moishezon. In particular, we have:

Corollary 2.7. Let (X, ω) be a compact Kähler manifold with (δ_1, δ_2) -bounded geometry and (ε, δ) -quasi-negative holomorphic sectional curvature. If one further assumes that K_X is nef, then X is of general type, and thus, in particular, Moishezon.

Moishezon's theorem [19] asserts that a Moishezon manifold that is Kähler, is projective. Further, from the recent developments in the minimal model program [2, Corollary 1.4.6], a Moishezon manifold without rational curves is projective. In particular, from [14, Theorem 0.5], we have the following:

Corollary 2.8. Let X be a compact Kähler manifold endowed with a Hermitian metric of (δ_1, δ_2) -bounded geometry and (ε, δ) -quasi-negative holomorphic sectional curvature. If one further assumes that K_X is nef and X is Kobayashi hyperbolic, then X is projective with ample canonical bundle.

Remark 2.9. It would be interesting to further explore the role played by Tian's α -invariant, given its place in algebraic geometry. Indeed, we remind the reader that Tian's α -invariant is an asymptotic version of the log canonical threshold (see [24, 13, 10]).

3. Proof of Theorem 1.2

To see Theorem 1.2, we first prove a useful lemma by employing a method of Li– Ni–Zhu [18]. Let

$$\lambda_{\omega_0} := \max_{x \in X} \max_{v \in T_x^{1,0} X} \operatorname{Ric}_{\alpha,\beta}^{\perp}(v)$$

Define

$$\tau_{\omega_0}(x) := \rho\left(\max_{v \in T_x^{1,0}X} \operatorname{Ric}_{\alpha,\beta}^{\perp}(v)\right) \cdot \max_{v \in T_x^{1,0}X} \operatorname{Ric}_{\alpha,\beta}^{\perp}(v),$$

where $\rho : \mathbf{R} \to \{n+1, 2n\}$ is a function with $\rho(s) = n+1$ for $s \leq 0$ and $\rho(s) = 2n$ for s > 0.

10

Lemma 3.1. Let (X^n, ω_0) be a compact Kähler manifold. We consider the following complex Monge-Ampère equation for φ_t :

$$(t\omega_0 - \operatorname{Ric}_{\omega_0} + dd^c \varphi_t)^n = e^{(1 + \frac{\alpha}{2\beta})\varphi_t} \omega_0^n$$
(3.1)

(a) If $\lambda_{\omega_0} \ge 0$, then the solution φ_t to equation (3.1) exists for $t \in \left(\frac{2n\lambda_{\omega_0}}{(n+1)\alpha}, +\infty\right)$;

(b) If $\lambda_{\omega_0} < 0$, then the solution φ_t to equation (3.1) exists for $t \in [0, +\infty)$.

Proof. We consider the following complex Monge-Ampère equation for φ_t :

$$(t\omega_0 - \operatorname{Ric}_{\omega_0} + dd^c \varphi_t)^n = e^{(1 + \frac{\alpha}{2\beta})\varphi_t} \omega_0^n.$$
(3.2)

if t is large enough, then $t\omega_0 - \operatorname{Ric}_{\omega_0} > 0$. By the Aubin–Yau theorem [1, 31], there is a unique smooth solution φ_t of (3.2) for $t \in (T, +\infty)$, where T is the minimum existence time of φ_t . We assume Lemma 3.1(a) fails, i.e. $T > \frac{2n\lambda_{\omega_0}}{(n+1)\alpha} \ge 0$. Without losing generality, we only consider the $t \in (T, A]$, where A is a fixed positive constant large enough. Set $\omega_t := t\omega_0 - \operatorname{Ric}_{\omega_0} + dd^c \varphi_t$. Then for $t \in (T, A]$, we have

$$\omega_t = t\omega_0 - \operatorname{Ric}_{\omega_0} + dd^c \varphi_t > 0, \qquad (3.3)$$

and (3.2) is equivalent to

$$\operatorname{Ric}_{\omega_t} = -\omega_t + t\omega_0 - \frac{\alpha}{2\beta} dd^c \varphi_t.$$
(3.4)

To simplify notation we write the components of ω_t as $g_{i\bar{j}}$ and the components of ω_0 as $h_{\alpha\bar{\beta}}$. Let $G = \operatorname{tr}_{\omega_t}\omega_0 = \operatorname{tr}_g h$, from Lemma 2.1 we have

$$\Delta_{\omega_t} \log G \geq \frac{1}{G} (\operatorname{Ric}^g_{i\overline{j}} g^{i\overline{q}} g^{p\overline{j}} h_{p\overline{q}} - g^{i\overline{j}} g^{k\overline{l}} R^h_{i\overline{j}k\overline{l}}).$$
(3.5)

By [18, Lemma 2.2], we get

$$\frac{1}{G}g^{i\overline{j}}g^{k\overline{l}}R^{h}_{i\overline{j}k\overline{l}} \leq \frac{\lambda}{2\beta}G + \frac{\lambda}{2\beta G}|h|_{g}^{2} - \frac{\alpha}{\beta}\mathrm{tr}_{g}\mathrm{Ric}^{h} + \frac{\alpha}{2\beta G}(G\cdot\mathrm{tr}_{g}\mathrm{Ric}^{h} - \langle\omega_{h},\mathrm{Ric}^{h}\rangle_{g}).$$
(3.6)

Choosing local coordinates such that $g_{i\overline{j}} = \delta_{i\overline{j}}$, $h_{i\overline{j}} = h_{i\overline{i}}\delta_{ij}$, then

$$G \cdot \operatorname{tr}_{g}\operatorname{Ric}^{h} - \langle \omega_{h}, \operatorname{Ric}^{h} \rangle_{g} = \sum_{i} \operatorname{Ric}_{i\overline{i}}^{h} (\sum_{j} h_{j\overline{j}} - h_{i\overline{i}}) = \sum_{i} (\operatorname{Ric}_{i\overline{i}}^{h} (\sum_{j \neq i} h_{j\overline{j}}))$$

$$\leq \sum_{i} (t \cdot h_{i\overline{i}} + \sqrt{-1}\varphi_{t,i\overline{i}}) (\sum_{j} h_{j\overline{j}} - h_{i\overline{i}})$$

$$= tG^{2} - t|h|_{g}^{2} + G\Delta_{g}\varphi_{t} - \langle dd^{c}\varphi_{t}, h\rangle_{g}.$$
(3.7)

Here we used (3.3). Combining (3.6) and (3.7), we get

$$\frac{1}{G}g^{i\overline{j}}g^{k\overline{l}}R^{h}_{i\overline{j}k\overline{l}} \leq \frac{\lambda + \alpha t}{2\beta}G + \frac{\lambda - \alpha t}{2\beta}\frac{|h|_{g}^{2}}{G} - \frac{\alpha}{\beta}\mathrm{tr}_{g}\mathrm{Ric}^{h} + \frac{\alpha}{2\beta}\Delta_{g}\varphi_{t} - \frac{\alpha}{2\beta}\frac{1}{G}\langle dd^{c}\varphi_{t}, h\rangle_{g}.$$
(3.8)

By using (3.4), we have

$$\frac{1}{G}\operatorname{Ric}_{i\overline{j}}^{g}g^{i\overline{q}}g^{p\overline{j}}h_{p\overline{q}} = \frac{1}{G}\langle\operatorname{Ric}^{g},h\rangle_{g} = \frac{1}{G}\langle-g+th-\frac{\alpha}{2\beta}dd^{c}\varphi_{t},h\rangle_{g}$$

$$= \frac{1}{G}\langle-g+th,h\rangle_{g} - \frac{1}{G}\frac{\alpha}{2\beta}\langle\varphi_{t},h\rangle_{g}.$$
(3.9)

Combining (3.5), (3.8) and (3.9), we get

$$\Delta_{\omega_t} \left(\log G - \frac{\alpha}{2\beta} \varphi_t \right) \geq -\frac{\lambda + \alpha t}{2\beta} G - \frac{\lambda - \alpha t}{2\beta} \frac{|h|_g^2}{G} + \frac{\alpha}{\beta} (\operatorname{tr}_g \operatorname{Ric}^h - \Delta_g \varphi_t) + \frac{1}{G} \langle -g + th, h \rangle_g.$$
(3.10)

If $\lambda \ge 0$, since t > T > 0, $n|h|_g^2 \ge G^2$, $|h|_g^2 \le G^2$,

$$\Delta_g \varphi_t = n - tG + \operatorname{tr}_g \operatorname{Ric}^h, \qquad (3.11)$$

and

$$\frac{1}{G}\langle -g + th, h \rangle_g = -1 + \frac{t}{G} |h|_g^2 \ge -1.$$
(3.12)

so we have

$$\Delta_{\omega_t} \left(\log G - \frac{\alpha}{2\beta} \varphi_t \right) \geq \left(\frac{(n+1)\alpha t - 2n\lambda}{2\beta n} \right) G - \left(\frac{\alpha n}{\beta} + 1 \right).$$
(3.13)

Now we apply the maximum principle to get a uniform lower estimate of ω_t . For any fixed $t \in (T, A]$, at the maximum point x_1 of the φ_t , we have $dd^c \varphi_t \leq 0$, so we can get

$$0 < g = \omega_t |_{x_1} \leq t\omega_0 - \operatorname{Ric}_{\omega_0}(x_1),$$
 (3.14)

and

$$e^{(1+\frac{\alpha}{2\beta})\sup_X\varphi_t(x)} = e^{(1+\frac{\alpha}{2\beta})\varphi_t(x_1)} \leq \frac{(t\omega_0 - \operatorname{Ric}_{\omega_0}))^n}{\omega_0^n} \leq C,$$
(3.15)

where C is independent of t. Hence φ_t has a uniform upper bound, and we also get

$$\omega_t^n \leq C' \omega_0^n. \tag{3.16}$$

12

(ε, δ)–QUASI-NEGATIVE CURVATURE AND POSITIVITY OF THE CANONICAL BUNDLE 13

Again applying maximum principle to $\log G - \frac{\alpha}{2\beta}\varphi_t$. At the point x_2 , where the maximum of $\log G - \frac{\alpha}{2\beta}\varphi_t$ is attained, since $t > T > \frac{2n\lambda\omega_0}{(n+1)\alpha} \ge 0$, by (3.13), we have that

$$G(x_2) \leq C'', \tag{3.17}$$

where C'' is also independent of t. We note that

$$G \cdot \left(\frac{\omega_0^n}{\omega_t^n}\right)^{\frac{\alpha}{2\beta + \alpha}} = G \cdot \left(e^{-(1 + \frac{\alpha}{2\beta})\varphi_t}\right)^{\frac{\alpha}{2\beta + \alpha}} = G \cdot e^{-\frac{\alpha}{2\beta}\varphi_t}.$$
 (3.18)

So $\sup_X G \cdot \left(\frac{\omega_0^n}{\omega_t^n}\right)^{\frac{\alpha}{2\beta+\alpha}}$ is also attained at x_2 . By the AM-GM inequality, at x_2 , we have $G \cdot \left(\frac{\omega_0^n}{\omega_t^n}\right)^{\frac{\alpha}{2\beta+\alpha}} \leq G \cdot \left(\frac{G}{n}\right)^{\frac{\alpha n}{2\beta+\alpha}}$. By (3.17), we have $\sup_X G \cdot \left(\frac{\omega_0^n}{\omega_t^n}\right)^{\frac{\alpha}{2\beta+\alpha}} \leq C'''$. Again by (3.16), we get

$$\omega_t \geq \widetilde{C}\omega_0, \tag{3.19}$$

where \widetilde{C} is a positive number and independent of t. Combining (3.16) and (3.19), for some positive C > 0 independent of t, we have

$$C^{-1}\omega_0 \leq \omega_t \leq C\omega_0. \tag{3.20}$$

Therefore we obtain the higher order estimates (see [25]). Moreover, ω_t converges to a smooth Kähler form as $t \to T$. This is a contradiction since T is the minimum existence time of φ_t . Therefore, there must holds $T \leq \frac{2n\lambda}{(n+1)\alpha}$. This completes the proof of Lemma 3.1 (a).

If $\lambda < 0$, by ((3.10)), it is easy to get that

$$\Delta_{\omega_t} \left(\log G - \frac{\alpha}{2\beta} \varphi_t \right) \geq \left(\frac{(n+1)\alpha t - (n+1)\lambda}{2\beta n} \right) G - \left(\frac{\alpha n}{\beta} + 1 \right).$$
(3.21)

Following the proof of Lemma 3.1 (a), we can easily prove Lemma 3.1 (b).

Proof of Theorem 1.2: The proof of Theorem 1.2 is similar to that of Theorem 1.1.
For completeness, we give a detailed proof. We consider the following complex Monge-
Ampère equation for
$$\varphi_t$$
:

$$(t\omega_0 - \operatorname{Ric}_{\omega_0} + dd^c \varphi_t)^n = e^{(1 + \frac{\alpha}{2\beta})\varphi_t} \omega_0^n.$$
(3.22)

let $\omega_t = t\omega_0 - \operatorname{Ric}_{\omega_0} + dd^c \varphi_t$. By Lemma 3.1, we have that

$$\Delta_{\omega_t} \left(\log(\operatorname{tr}_{\omega_t} \omega_0) - \frac{\alpha}{2\beta} \varphi_t \right) \geq \left(\frac{(n+1)\alpha t - \tau_{\omega_0}}{2\beta n} \right) \operatorname{tr}_{\omega_t}(\omega_0) - \left(\frac{\alpha n}{\beta} + 1 \right).$$
(3.23)

and there exists smooth solution φ_t for $t \in \left(\frac{2n\lambda_{\omega_0}}{(n+1)\alpha}, +\infty\right)$. In the following we just consider $t \in \left(\frac{2n\lambda_{\omega_0}}{(n+1)\alpha}, \frac{4n\lambda_{\omega_0}}{(n+1)\alpha}\right]$. Integrating (3.23) gives

$$\left(\frac{\alpha n}{\beta}+1\right)\int_{X}\omega_{t}^{n} \geq \int_{X}\left(\frac{(n+1)\alpha t}{2\beta n}-\frac{\tau_{\omega_{0}}}{2\beta n}\right)(\mathrm{tr}_{\omega_{t}}\omega_{0})\omega_{t}^{n}$$
$$\geq -\frac{1}{2\beta n}\int_{U}\tau_{\omega_{0}}(\mathrm{tr}_{\omega_{t}}\omega_{0})\omega_{t}^{n}$$
$$\geq -\frac{1}{2\beta}\int_{U}\tau_{\omega_{0}}\cdot\left(\frac{\omega_{0}^{n}}{\omega_{t}^{n}}\right)^{\frac{1}{n}}\omega_{t}^{n}$$
$$= -\frac{1}{2\beta}\int_{U}\tau_{\omega_{0}}\cdot e^{-\frac{1}{n}(1+\frac{\alpha}{2\beta})\varphi_{t}}\omega_{t}^{n}$$
$$\geq -\frac{1}{2\beta}e^{-\frac{1}{n}(1+\frac{\alpha}{2\beta})\sup_{X}\varphi_{t}}\int_{U}\tau_{\omega_{0}}\omega_{t}^{n} \qquad (3.24)$$

Now let us estimate the bounds of the following fractions:

$$\frac{-\int_{\mathfrak{U}}\tau_{\omega_0}\omega_t^n}{\int_X\omega_t^n} = \frac{-\int_{\mathfrak{U}}\tau_{\omega_0}e^{(1+\frac{\alpha}{2\beta})\varphi_t}\omega_0^n}{\int_Xe^{(1+\frac{\alpha}{2\beta})\varphi_t}\omega_0^n} = \frac{-\int_{\mathfrak{U}}\tau_{\omega_0}e^{(1+\frac{\alpha}{2\beta})\varphi_t^*}\omega_0^n}{\int_Xe^{(1+\frac{\alpha}{2\beta})\varphi_t^*}\omega_0^n},$$
(3.25)

where $\varphi_t^* := \varphi_t - \sup_X \varphi_t$. We fix a nonnegative number $b_0 \ge 0$ such that $\operatorname{Ric}_{\omega_0} \ge -b_0\omega_0$. Applying the maximum principle to φ_t , by (3.22), we get that

$$\sup_{X} \varphi_t \leq \frac{2\beta}{\alpha + 2\beta} \log\left(\frac{4n\varepsilon}{(n+1)\alpha} + b_0\right)^n \leq \frac{2\beta}{\alpha + 2\beta} \log(c_0 + b_0)^n$$
(3.26)

for $t \in \left(\frac{2n\lambda_{\omega_0}}{(n+1)\alpha}, \frac{4n\lambda_{\omega_0}}{(n+1)\alpha}\right]$, where c_0 is a positive number such that the α -invariant $\alpha(X, c_0\omega_0) \ge 2$ and we choose $\varepsilon \le \frac{(n+1)\alpha c_0}{4n}$. Set

$$c_1 := \inf \left\{ \int_{\mathfrak{U}} e^u \omega_0^n | u \in \mathrm{PSH}(X, (c_0 + b_0)\omega_0), \sup_X u = 0 \right\},$$

which is a positive number depending only on $\mathcal{U}, \omega_0, c_0, b_0$, and hence only on \mathcal{U}, ω_0 . Then, for $t \in \left(\frac{2n\lambda_{\omega_0}}{(n+1)\alpha}, \frac{4n\lambda_{\omega_0}}{(n+1)\alpha}\right]$, we have

$$-\int_{\mathfrak{U}}\tau_{\omega_0}e^{(1+\frac{\alpha}{2\beta})\varphi_t^*}\omega_0^n \geq (n+1)\delta\int_{\mathfrak{U}}e^{(1+\frac{\alpha}{2\beta})\varphi_t^*}\omega_0^n \geq (n+1)\delta c_1$$

Since $\sup_X \varphi_t^* = 0$, so we get

$$\int_X e^{(1+\frac{\alpha}{2\beta})\varphi_t^*} \omega_0^n \le \int_X \omega_0^n =: c_2,$$

where c_2 is a positive number depending only on ω_0 . In conclusion, we obtain that for $t \in \left(\frac{2n\lambda_{\omega_0}}{(n+1)\alpha}, \frac{4n\lambda_{\omega_0}}{(n+1)\alpha}\right]$,

$$\frac{-\int_{\mathcal{U}}\tau_{\omega_0}\omega_t^n}{\int_X\omega_t^n} \geq \frac{(n+1)\delta c_1}{c_2}.$$

with, combining with (3.24), concludes

$$\sup_{\mathcal{U}} \varphi_t \geq \frac{2\beta n}{\alpha + 2\beta} \log \frac{(n+1)\delta c_1}{(2\alpha n + 2\beta)c_2}.$$

Now we define

$$c_3 := \inf \left\{ \int_X e^v \omega_0^n \mid v \in \text{PSH}(X, c_0 b_0 \omega_0), \ n \log \frac{(n+1)\delta c_1}{(2\alpha n + 2\beta)c_2} \le \sup_X v \le \log(c_0 + b_0)^n \right\},$$

which is a positive number depending only on U, ω_0, δ . Therefore,

$$\int_{X} \left(\frac{2n\lambda_{\omega_{0}}}{(n+1)\alpha} \omega_{0} - \operatorname{Ric}_{\omega_{0}} \right)^{n} = \lim_{t \to \frac{2n\lambda_{\omega_{0}}}{(n+1)\alpha}} \int_{X} \left(t\omega_{0} - \operatorname{Ric}_{\omega_{0}} + dd^{c}\varphi_{t} \right)^{n}$$
$$= \lim_{t \to \frac{2n\lambda_{\omega_{0}}}{(n+1)\alpha}} \int_{X} e^{(1+\frac{\alpha}{2\beta})\varphi_{t}} \omega_{0}^{n}$$
$$\geq \liminf_{t \to \frac{2n\lambda_{\omega_{0}}}{(n+1)\alpha}} \int_{X} e^{(1+\frac{\alpha}{2\beta})\varphi_{t}} \omega_{0}^{n}$$
$$\geq c_{3} \qquad (3.27)$$

On the other hand,

$$\int_{X} \left(\frac{2n\lambda_{\omega_{0}}}{(n+1)\alpha} \omega_{0} - \operatorname{Ric}_{\omega_{0}} \right)^{n} = \lim_{t \to \frac{2n\lambda_{\omega_{0}}}{(n+1)\alpha}} \int_{X} (t\omega_{0} - \operatorname{Ric}_{\omega_{0}} + dd^{c}\varphi_{t})^{n}$$
$$\leq \int_{X} \left(\frac{4n\varepsilon}{(n+1)\alpha} \omega_{0} - \operatorname{Ric}_{\omega_{0}} \right)^{n}$$
$$\leq \int_{X} (-\operatorname{Ric}_{\omega_{0}})^{n} + c_{4}\varepsilon$$
$$= \int_{X} (2\pi c_{1}(K_{X}))^{n} + c_{4}\varepsilon \qquad (3.28)$$

where c_4 is positive number depending only on ω_0 . by (3.27) and (3.28), we have that

$$\int_X (2\pi c_1(K_X))^n \ge c_3 - c_4\varepsilon$$

Therefore, if $\varepsilon \leq \min\{\frac{c_3}{2c_4}, \frac{(n+1)\alpha c_0}{4n}\}$, we have $\int_X (2\pi c_1(K_X))^n \geq \frac{c_3}{2} > 0$.

Proof of Theorem 1.3. We note that the argument can be repeated more or less without change to prove Theorem 1.3. The only modification is that one requires [18, Lemma 2.1].

References

- Aubin, T., Équations du type Monge-Ampère sur les varietés kählériennes compactes, C. Rendus Acad. Sci. Paris 283 (1976), pp. 119–121. [Cited on pages 1 and 11.]
- [2] Birkar, C., Cascini, P., Hacon, C., McKernan, J., Existence of minimal models for varieties of log general type. J. Amer. Math. Soc., 23(2):405–468, 2010. [Cited on page 10.]
- Broder, K., The Schwarz lemma in Kähler and non-Kähler geometry, arXiv:2109.06331, to appear in the Asian J. Math. [Cited on page 5.]
- [4] Broder, K., The Schwarz lemma: An Odyssey, arXiv:2110.04989, Rocky Mountain J. Math. 52 (2022), no. 4, 1141–1155. [Cited on page 5.]
- [5] Broder, K., Remarks on the Wu-Yau theorem, (submitted). [Cited on page 3.]
- [6] Broder, K., Stanfield, J., On the Gauduchon Curvature of Hermitian Manifolds, Int. J. of Math., doi: https://doi.org/10.1142/S0129167X23500398 [Cited on page 2.]
- Broder, K., Tang, K., On the weighted orthogonal Ricci curvature, J. Geom. and Phys., 104783, doi: https://doi.org/10.1016/j.geomphys.2023.104783. [Cited on page 3.]
- [8] Broder, K., Tang, K., On the altered holomorphic curvatures of Hermitian manifolds, arXiv:2201.03666. [Cited on page 2.]
- Campana, F., Twistor spaces and nonhyperbolicity of certain symplectic Kähler manifolds, In Complex analysis (Wuppertal 1991), Aspects Math., E17, pp. 64–69. Friedr. Veiweg, Braunschweig, 1991. [Cited on page 1.]
- [10] Cheltsov, I., Shramov, C., Log canonical thresholds of smooth Fano threefolds (Russian), with an appendix by Jean-Pierre Demailly, Uspekhi Mat. Nauk 63 (2008), pp. 73–180; translation in Russian Math. Surveys 63 (2008), pp. 859–958. [Cited on page 10.]
- [11] Chu, J. C., Lee, M. C., Tam L.-F., Kähler manifolds and mixed curvature. Trans. Amer. Math. Soc. 375 (2022), no. 11, 7925-7944. [Cited on pages 1, 3, and 4.]
- [12] Debarre, O., Higher-dimensional algebraic geometry, Universitext, Springer-Verlag, New York, 2001. MR 1841091 [Cited on page 1.]
- [13] Demailly, J.-P., Kollár, J., Semi-continuity of complex singularity exponents and Kähler–Einstein metrics on Fano orbifolds, Ann. Sci. École Norm. Sup. (4) 34 (2001), 525–556. [Cited on page 10.]
- [14] Demailly, J.-P., Paun, M., Numerical characterization of the Kähler cone of a compact Kähler manifold, Ann. of Math. (2) 159 (2004), no. 3, 1247–1274. [Cited on pages 1 and 10.]

- [15] Diverio, S., Trapani, S., Quasi-negative holomorphic sectional curvature and positivity of the canonical bundle, J. Diff. Geom., **111** (2019), pp. 303–314. [Cited on pages 1, 2, 3, and 6.]
- [16] Heier, G., Lu, S., Wong, B., On the canonical line bundle and negative holomorphic sectional curvature, Math. Res. Lett. 17 (2010), no. 06, pp. 1101–1110 [Cited on page 1.]
- [17] Lee, M. C., Streets, J., Complex manifolds with negative curvature operator, Int. Math. Res. Not, arXiv: 1903.12645. [Cited on page 2.]
- [18] Li, C., Ni, L., Zhu, X., An application of a C²-estimate for a complex Monge-Ampère equation, Internat. J. Math. 32 (2021), no. 12, Paper No. 2140007, 13 pp. [Cited on pages 10, 11, and 16.]
- [19] Moishezon, B. G., On n-dimensional compact complex manifolds having n algebraically independent meromorphic functions. I. Izv. Akad. Nauk SSSR Ser. Mat., 30: 133–174, 1966. [Cited on pages 1 and 10.]
- [20] Ni, L., Liouville theorems and a Schwarz Lemma for holomorphic mappings between Kähler manifolds. Comm. Pure Appl. Math, 2021, 74: 1100–1126. [Cited on page 4.]
- [21] Royden, H. L., The Ahlfors-Schwarz lemma in several complex variables, Comment. Math. Helvetici 55 (1980), p. 547–558. [Cited on page 5.]
- [22] Siu, Y.-T., A vanishing theorem for semipositive line bundles over non-Kähler manifolds, J. Diff. Geom. 19 (1984), pp. 431–452. [Cited on page 3.]
- [23] Tang, K., On almost nonpositive k-Ricci curvature, J. Geom. Anal. 32 (2022), no. 12, Paper No. 306, 12 pp. [Cited on pages 3 and 4.]
- [24] Tian, G., On Kähler-Einstein metrics on certain Kähler manifolds with $C_1(M) > 0$, Invent. Math. 89 (1987), no. 2, 225–246 [Cited on pages 6, 9, and 10.]
- [25] Tosatti, V., Yang, X.-K., An extension of a theorem of Wu-Yau, J. Differential Geom. 107(3): 573-579 [Cited on pages 1, 5, 6, and 13.]
- [26] Wong, B., The uniformization of compact Kähler surfaces of negative curvature, J. Differential Geom. 16 (1981), no. 3, 407–420. [Cited on page 1.]
- [27] Wu, D., Yau, S.-T., Negative holomorphic sectional curvature and positive canonical bundle, Invent. Math. 204 (2016), no. 2, 595–604 [Cited on pages 1 and 6.]
- [28] Wu, D., Yau, S.-T., A remark on our paper "Negative holomorphic curvature and positive canonical bundle". Comm Anal Geom. 24 (2016) 901–912. [Cited on page 1.]
- [29] Yang, B., Zheng, F., On curvature tensors of Hermitian manifolds, Communications in Analysis and Geometry, vol. 26 (2018), no. 5, pp. 1195–1222 [Cited on page 2.]
- [30] Yang, X., Zheng, F., On the real bisectional curvature for Hermitian manifolds, Trans. Amer. Math. Soc. 371 (2019), no. 4, 2703–2718 [Cited on pages 2 and 5.]
- [31] Yau, S.-T., On the Ricci curvature of a compact Kähler manifold and the complex Monge–Ampère equation, I, Comm. Pure Appl. Math. 31 (1978), pp. 339–411. [Cited on pages 1 and 11.]
- [32] Zhang, Y., Integral inequalities for holomorphic maps and applications, Trans. Amer. Math. Soc. 374 (2021), no.4, 2341–2358 [Cited on pages 3 and 5.]
- [33] Zhang, Y., Holomorphic sectional curvature, nefness and Miyaoka-Yau type inequality, Math. Z. 298 (2021), no.3-4, 953–974 [Cited on page 3.]

[34] Zhang, Y., Zheng, T., On almost quasi-negative holomorphic sectional curvature, arXiv:2010.01314v4.

Cited on pages 1, 2, 3, and 6.

THE UNIVERSITY OF QUEENSLAND, ST. LUCIA, QLD 4067, AUSTRALIA *Email address*: k.broder@uq.edu.au

College of Mathematics and Computer Science, Zhejiang Normal University, Jinhua, Zhejiang, 321004, China

Email address: kaitang001@zjnu.edu.cn