

On torse-forming vector fields and biharmonic hypersurfaces in Riemannian manifolds

Ahmed Mohammed Cherif*

Abstract

In this paper, we give some properties of biharmonic hypersurface in Riemannian manifold has a torse-forming vector field.

Keywords: Torse-forming vector fields, Biharmonic hypersurfaces; Chen's conjecture.

Subclass: 53C20; 58E20.

1 Introduction

The energy functional of a smooth map $\varphi : (M, g) \longrightarrow (N, h)$ between two Riemannian manifolds is defined by

$$E(\varphi) = \frac{1}{2} \int_D |d\varphi|^2 v^g, \quad (1.1)$$

where D is compact domain of M , $|d\varphi|$ is the Hilbert-Schmidt norm of the differential $d\varphi$, and v^g is the volume element on (M, g) . A map φ is called harmonic if it is a critical point of the energy functional (1.1). The Euler Lagrange equation associated to (1.1) is given by (see [1, 5, 12])

$$\tau(\varphi) = \text{trace } \nabla d\varphi = \sum_{i=1}^m \nabla_{e_i}^\varphi d\varphi(e_i) - \sum_{i=1}^m d\varphi(\nabla_{e_i}^M e_i) = 0, \quad (1.2)$$

where $\{e_i\}_{i=1}^m$ is a local orthonormal frame field on (M, g) , ∇^M is the Levi-Civita connection of (M, g) , ∇^φ denote the pull-back connection on $\varphi^{-1}TN$, and m is the dimension of M . A natural generalization of harmonic maps

*University Mustapha Stambouli Mascara, Faculty of Exact Sciences, Mascara 29000, Algeria. Email: a.mohammedcherif@univ-mascara.dz

is given by integrating the square of the norm of the tension field. More precisely, the bienergy functional of a map $\varphi \in C^\infty(M, N)$ is defined by

$$E_2(\varphi) = \frac{1}{2} \int_D |\tau(\varphi)|^2 v^g. \quad (1.3)$$

A map $\varphi \in C^\infty(M, N)$ is called biharmonic if it is a critical point of the bienergy functional, that is, if it is a solution of the Euler Lagrange equation associated to (1.3)

$$\begin{aligned} \tau_2(\varphi) &= -\text{trace } R^N(\tau(\varphi), d\varphi)d\varphi - \text{trace } (\nabla^\varphi)^2 \tau(\varphi) \\ &= -\sum_{i=1}^m R^N(\tau(\varphi), d\varphi(e_i))d\varphi(e_i) - \sum_{i=1}^m \nabla_{e_i}^\varphi \nabla_{e_i}^\varphi \tau(\varphi) \\ &\quad + \sum_{i=1}^m \nabla_{\nabla_{e_i}^M e_i}^\varphi \tau(\varphi) = 0, \end{aligned} \quad (1.4)$$

where R^N is the curvature tensor of (N, h) defined by

$$R^N(X, Y)Z = \nabla_X^N \nabla_Y^N Z - \nabla_Y^N \nabla_X^N Z - \nabla_{[X, Y]}^N Z,$$

where ∇^N is the Levi-Civita connection of (N, h) and $X, Y, Z \in \Gamma(TN)$ (see [6, 12]). Clearly, it follows from (1.4) that any harmonic map is biharmonic and we call those non-harmonic biharmonic maps proper biharmonic maps. Let M be a submanifold in (N, \langle, \rangle) of dimension m , $\mathbf{i} : M \hookrightarrow (N, \langle, \rangle)$ the canonical inclusion, and let $\{e_i\}_{i=1}^m$ be a local orthonormal frame field with respect to induced Riemannian metric g on M by \langle, \rangle . We denote by $\overline{\nabla}$ (resp. ∇) the Levi-Civita connection of (N, \langle, \rangle) (resp. of (M, g)), by $\overline{\text{grad}}$ (resp. grad) the gradient operator on (N, \langle, \rangle) (resp. on (M, g)), by B the second fundamental form of the submanifold (M, g) , and by H the mean curvature vector field of (M, g) (see [1, 10]). The submanifold (M, g) is called a harmonic (resp. biharmonic) submanifold in (N, h) if $\tau(\mathbf{i}) = 0$ (resp. $\tau_2(\mathbf{i}) = 0$). The expressions assumed by the tension and bitension fields are given by

$$\tau(\mathbf{i}) = mH, \quad \tau_2(\mathbf{i}) = -m \sum_{i=1}^m \left\{ \overline{R}(H, e_i)e_i + \overline{\nabla}_{e_i} \overline{\nabla}_{e_i} H - \overline{\nabla}_{\nabla_{e_i} e_i} H \right\}, \quad (1.5)$$

where \overline{R} is the curvature tensor of (N, h) . In [11], Ye-Lin Ou proved that a hypersurface (M, g) in a Riemannian manifold (N, \langle, \rangle) with mean curvature

vector field $H = f\eta$, that is the dimension of N is $m + 1$, is biharmonic if and only if

$$\begin{cases} -\Delta(f) + f|A|^2 - f\overline{\text{Ric}}(\eta, \eta) &= 0; \\ 2A(\text{grad } f) + mf \text{grad } f - 2f(\overline{\text{Ricci}} \eta)^\top &= 0, \end{cases} \quad (1.6)$$

where $\overline{\text{Ric}}$ (resp. $\overline{\text{Ricci}}$) is the Ricci curvature (resp. Ricci tensor) of (N, \langle, \rangle) , f denote the mean curvature function of (M, g) , and A the shape operator with respect to the unit normal vector field η .

Let (N, \langle, \rangle) be a Riemannian manifold admits a torse-forming vector field P , that is P satisfies the following formula

$$\overline{\nabla}_X P = \mu X + \omega(X)P, \quad \forall X \in \Gamma(TN), \quad (1.7)$$

for some smooth function μ and 1-form on N . The 1-form ω is called the generating form and the function μ is called the conformal scalar. Let (M, g) be a hypersurface in (N, \langle, \rangle) . We consider the following decomposition of the torse-forming vector field

$$P = \phi \eta + V,$$

where V denote the tangential component of P and $\phi = \langle P, \eta \rangle$.

References

- [1] P. Baird and J. C. Wood, *Harmonic morphisms between Riemannian manifolds*, Clarendon Press Oxford (2003).
- [2] R. Caddeo and S. Montaldo, *C. Oniciuc. Biharmonic submanifolds in spheres*, Israel J. Math., **130** (2002), 109-123.
- [3] B-Y. Chen, *Some open problems and conjectures on submanifolds of finite type*, Soochow J. Math., **17** (1991), 169-188.
- [4] B.Y. Chen, *A report on submanifolds of finite type*, Soochow J. Math., **22** (1996), 117-337.
- [5] J. Eells and J. H. Sampson, *Harmonic mappings of Riemannian manifolds*, Amer. J. Math., **86** (1964), 109-160.
- [6] G. Y. Jiang, *2-Harmonic maps between Riemannian manifolds*, Annals of Math., China, **7A(4)** (1986), 389-402.

- [7] K. Kenmotsu, *A class of almost contact Riemannian manifolds*, Tohoku Math.J., **24** (1972), 93-103.
- [8] A. Mohammed Cherif and M. Djaa, *Harmonic maps and torse-forming vector fields*, International electronic journal of geometry, **13**(1) (2020), 87-93.
- [9] N. Nakauchi and H. Urakawa, *Biharmonic hypersurfaces in a Riemannian manifold with non-positive Ricci curvature*, Ann. Global Anal. Geom., **40** (2011), 125-131.
- [10] O'Neil, *Semi-Riemannian Geometry*, Academic Press, New York (1983).
- [11] Y-L. Ou, *Biharmonic hypersurfaces in Riemannian manifolds*, Pacific Journal of Mathematics, **248**(1) (2010), 217-232.
- [12] Y. Xin, *Geometry of harmonic maps*, Fudan University (1996).