

COURANT-NIJENHUIS ALGEBROIDS

HENRIQUE BURSZTYN, THIAGO DRUMMOND, AND CLARICE NETTO

ABSTRACT. We introduce Courant 1-derivations, which describe a compatibility between Courant algebroids and linear (1,1)-tensor fields and lead to the notion of Courant-Nijenhuis algebroids. We provide examples of Courant 1-derivations on exact Courant algebroids and show that holomorphic Courant algebroids can be viewed as special types of Courant-Nijenhuis algebroids. By considering Dirac structures, one recovers the Dirac-Nijenhuis structures of [5] (in the special case of the standard Courant algebroid) and obtains an equivalent description of Lie-Nijenhuis bialgebroids [9] via Manin triples.

CONTENTS

1. Introduction	1
2. 1-Derivations on vector bundles	3
2.1. Equivalence with linear (1,1)-tensors fields	3
2.2. The Nijenhuis condition	4
2.3. Duality	5
2.4. Compatibility with (pre-)Lie algebroids	6
3. 1-Derivations on Courant algebroids	8
3.1. Courant 1-derivations and Courant-Nijenhuis algebroids	8
3.2. Invariant Dirac structures	10
4. Courant 1-derivations on $TM \oplus T^*M$	11
4.1. Courant 1-derivations from pseudo-Riemmanian metrics	11
4.2. Nijenhuis 1-derivations and the Kähler condition	13
4.3. B-field transformations	15
5. Holomorphic Courant algebroids	16
6. Lagrangian splittings and doubles	19
6.1. Lagrangian splittings of Courant algebroids	19
6.2. Lagrangian splittings of Courant 1-derivations	20
References	22

1. INTRODUCTION

A 1-derivation on a vector bundle $E \rightarrow M$ is a connection-like object that codifies a linear (1,1)-tensor field on the total space of E in the same way that usual derivations of vector bundles correspond to linear vector fields. As it turns out, many examples of compatibility conditions involving structures of interest in Poisson geometry can be conveniently expressed in terms of such 1-derivations; in this context, a special role is played by “Nijenhuis 1-derivations”, i.e., 1-derivations whose corresponding linear (1,1)-tensor fields have vanishing Nijenhuis torsion.

A motivating example is that of Poisson-Nijenhuis structures [21], originally formulated in terms of an intricate notion of compatibility involving Poisson structures and Nijenhuis operators (that includes the vanishing of the so-called Magri-Morosi concomitant). From the recent

viewpoint of [4, 9], this is understood as follows: a Nijenhuis operator r on a manifold M canonically gives rise to a 1-derivation on TM and a dual one on T^*M , and the compatibility of r with a Poisson bivector field π is simply that $\pi^\sharp : TM \rightarrow T^*M$ intertwines these 1-derivations. This perspective leads the way to different generalizations, such as

- (a) Dirac-Nijenhuis structures [5],
- (b) Lie-Nijenhuis bialgebroids [9].

Just as Poisson structures are particular examples of both Dirac structures and Lie bialgebroids [8, 19], Poisson-Nijenhuis structures are special cases of the objects in (a) and (b). In this paper, we take a step further and introduce the closely related notion of *Courant-Nijenhuis algebroid*.

Dirac-Nijenhuis structures and Lie-Nijenhuis bialgebroids are motivated by the theory of Lie groupoids, in that they arise as infinitesimal counterparts of presymplectic-Nijenhuis and Poisson-Nijenhuis groupoids, respectively, generalizing the correspondence between Poisson-Nijenhuis structures and symplectic-Nijenhuis groupoids from [27], see [5, 9]. An important class of examples is given by holomorphic structures. Any holomorphic vector bundle can be seen as a smooth real vector bundle equipped with a special type of Nijenhuis 1-derivation, that we call a “Dolbeault 1-derivation”. In this particular context, (a) and (b) recover holomorphic Dirac structures and holomorphic Lie bialgebroids, respectively, and their integrations correspond to holomorphic presymplectic and holomorphic Poisson groupoids.

Courant algebroids are central ingredients in the theory of Dirac structures and Lie bialgebroids. The main object of study in this paper is a notion of compatibility between 1-derivations and Courant algebroids, described in Definition 3.3 by what we call a *Courant 1-derivation*. A Courant algebroid equipped with a Courant 1-derivation that is also Nijenhuis is called a *Courant-Nijenhuis algebroid*. Dirac structures therein generalize the Dirac-Nijenhuis structures of [5] and provide an alternative approach to the Lie-Nijenhuis bialgebroids of [9] via Manin triples.

The paper is structured as follows. In § 2 we recall 1-derivations on vector bundles, their main examples and properties, including the notion of duality, the Nijenhuis condition, and their compatibility with (pre-)Lie algebroid structures. Courant 1-derivations and Courant-Nijenhuis algebroids are introduced in § 3, along with their Dirac structures. As a basic example, we show that any (1,1)-tensor field on a manifold M canonically defines a Courant 1-derivation on the standard Courant algebroid $\mathbb{T}M = TM \oplus T^*M$ that underlies the Dirac-Nijenhuis structures studied in [5]. In § 4 we consider more general Courant 1-derivations on $\mathbb{T}M$, including a class of examples obtained through the additional choice of a (pseudo-)riemannian metric; in this case, the Nijenhuis condition is shown to be related to the Kähler compatibility condition (Prop. 4.5). In § 5 we show (Theorem 5.3) that Courant-Nijenhuis algebroids defined by Dolbeault 1-derivations coincide with holomorphic Courant algebroids. In § 6 we consider lagrangian splittings of Courant algebroids equipped with Courant 1-derivations (Theorem 6.1). We show in particular that the Drinfeld double of a Lie-Nijenhuis bialgebroid is a Courant-Nijenhuis algebroid, thereby establishing an equivalence between Lie-Nijenhuis bialgebroids and Courant-Nijenhuis algebroids equipped with splittings by Dirac-Nijenhuis structures.

Remark 1.1. Other works in the literature involve Nijenhuis structures and Courant algebroids but follow a different direction. Nijenhuis tensors on Courant algebroids have been considered by various authors [7, 10, 15], with applications to deformations, hierarchies and compatibilities of geometric structures, see e.g. [1, 2]. The fundamental objects of interest in these papers are vector bundle endomorphisms $N : E \rightarrow E$ with vanishing Nijenhuis torsion, where E is a Courant algebroid and Nijenhuis torsion is defined with respect to the Courant bracket. In contrast, the present paper studies a notion of compatibility between a (ordinary) linear Nijenhuis operator $K : TE \rightarrow TE$ on the total space of E and the Courant structure on E .

Acknowledgments. This project was partially supported by CNPq (National Council for Scientific and Technological Development), FAPERJ (Rio de Janeiro State Research Foundation) and by grant # 2022/06205-2 of FAPESP (São Paulo State Research Foundation).

2. 1-DERIVATIONS ON VECTOR BUNDLES

A central role in this paper is played by 1-derivations on vector bundles, so we start by briefly recalling their definition and basic properties.

Let $p : E \rightarrow M$ be a smooth, real vector bundle. A *1-derivation* on $E \rightarrow M$ is a triple $\mathcal{D} = (D, l, r)$, where $r : TM \rightarrow TM$ and $l : E \rightarrow E$ are vector-bundle maps covering the identity, and $D : \Gamma(E) \rightarrow \Gamma(T^*M \otimes E)$ is an \mathbb{R} -linear map satisfying the following Leibniz-type condition:

$$(2.1) \quad D_X(f\sigma) = fD_X(\sigma) + (\mathcal{L}_X f)l(\sigma) - (\mathcal{L}_{r(X)}f)\sigma,$$

where $X \in \mathfrak{X}(M)$, $\sigma \in \Gamma(E)$ and $f \in C^\infty(M)$, and we use the notation

$$D_X : \Gamma(E) \rightarrow \Gamma(E), \quad D_X(\sigma) = i_X(D(\sigma)).$$

Note that the linear combination of 1-derivations (defined componentwise) is a 1-derivation.

Let $F \subseteq E$ be a subbundle (over the same base, for simplicity).

Definition 2.1. *We say that F is \mathcal{D} -invariant if*

- $l(F) \subseteq F$,
- $D_X(\Gamma(F)) \subseteq \Gamma(F), \quad \forall X \in \mathfrak{X}(M)$.

When F is \mathcal{D} -invariant, \mathcal{D} restricts to a 1-derivation on F ,

$$(D|_{\Gamma(F)}, l|_F, r).$$

2.1. Equivalence with linear (1,1)-tensors fields. Just as usual derivations on a vector bundle $E \rightarrow M$ are equivalent to linear vector fields on E (see e.g. [20]), 1-derivations are in bijective correspondence with linear (1,1)-tensor fields, see [4, § 6.1] and [5, § 2.1].

Let $K \in \Omega^1(E, TE)$, i.e. K is a (1,1)-tensor field on the total space E , that we view as a vector-bundle morphism $K : TE \rightarrow TE$. We say that K is *linear* if $K : TE \rightarrow TE$ is also a vector-bundle morphism from the tangent prolongation bundle $TE \rightarrow TM$ to itself (not necessarily covering the identity on TM). Linear (1,1)-tensor fields form a linear subspace of $\Omega^1(E, TE)$. The reader can find a detailed treatment of linear tensors in [4], where the results stated in this section can be found.

Any linear (1,1)-tensor field $K : TE \rightarrow TE$ gives rise to a 1-derivation $\mathcal{D} = (D, l, r)$ on E as follows: $r : TM \rightarrow TM$ is the restriction of $K|_M$ to vectors tangent to the zero section, $l : E \rightarrow E$ is the restriction of $K|_M$ to vectors tangent to the p -fibers, and $D : \Gamma(E) \rightarrow \Gamma(T^*M \otimes E)$ is the \mathbb{R} -linear map defined by

$$D_X(\sigma) = (\mathcal{L}_{\sigma^\uparrow}K)(X), \quad \sigma \in \Gamma(E), X \in TM,$$

where $\sigma^\uparrow \in \mathfrak{X}(E)$ is the *vertical lift* of $\sigma \in \Gamma(E)$, given by $\sigma^\uparrow(e) = \frac{d}{dt}|_{t=0} e + t\sigma(p(e))$. This assignment establishes a linear bijection between linear (1,1)-tensor fields and 1-derivations on E with natural functorial properties, see [5, Thm. 2.1]. A direct consequence is that a subbundle $(F \rightarrow M) \subset (E \rightarrow M)$ is \mathcal{D} -invariant if and only if it is preserved by the corresponding linear (1,1) tensor field K ,

$$K(TF) \subseteq TF.$$

Let us give some examples.

Example 2.2 (Connections). A connection ∇ on E defines a 1-derivation with $r = 0$, $l = \text{id}_E$ and $D = \nabla$. The corresponding linear (1,1)-tensor field $K : TE \rightarrow TE$ is the projection operator on the vertical bundle $\ker(Tp) \subseteq TE$ along the horizontal bundle defined by ∇ . \diamond

Example 2.3 (Tangent and cotangent lifts). Given a (1,1)-tensor on M , $r : TM \rightarrow TM$, its tangent lift $r^{tg} : T(TM) \rightarrow T(TM)$ and cotangent lift $r^{ctg} : T(T^*M) \rightarrow T(T^*M)$ are linear (1,1)-tensor fields on TM and T^*M , respectively, defined as follows:

$$r^{tg} = \Theta \circ Tr \circ \Theta, \quad (\omega_{can})^b \circ r^{ctg} = (\varphi_r^* \omega_{can})^b,$$

where $\Theta : T(TM) \rightarrow T(TM)$ is the canonical involution of the double tangent bundle $T(TM)$, ω_{can} is the canonical symplectic form on T^*M , and $\varphi_r : T^*M \rightarrow T^*M$ is just r^* seen as a smooth map from T^*M to itself. The linearity of r^{tg} and r^{ctg} was proved in [9, Thm. 3.4], along with the identification of their corresponding 1-derivations as

$$\mathcal{D}^r = (D^r, r, r), \quad \text{and} \quad \mathcal{D}^{r,*} = (D^{r,*}, r^*, r),$$

where, for $X \in \mathfrak{X}(M)$, $D_X^r : \mathfrak{X}(M) \rightarrow \mathfrak{X}(M)$ and $D_X^{r,*} : \Omega^1(M) \rightarrow \Omega^1(M)$ are given by

$$(2.2) \quad D_X^r(Y) = (\mathcal{L}_Y r)(X) = [Y, r(X)] - r([Y, X]),$$

$$(2.3) \quad D_X^{r,*}(\alpha) = \mathcal{L}_X(r^*\alpha) - \mathcal{L}_{r(X)}\alpha.$$

◇

Example 2.4 (Holomorphic structures and Dolbeault 1-derivations). Let $r : TM \rightarrow TM$ be a complex structure on M . A holomorphic vector bundle $\mathcal{E} \rightarrow M$ can be regarded as a real vector bundle $E \rightarrow M$ equipped with a fibrewise complex structure $l \in \text{End}(E)$, $l^2 = -\text{Id}$, and a flat $T^{0,1}$ -connection $\bar{\partial}$ on the complex vector bundle (E, l) [22]. One can equivalently express the holomorphic structure on E determined by r , l and $\bar{\partial}$ as a 1-derivation $\mathcal{D}^{Dolb} = (D, l, r)$, where

$$(2.4) \quad D_X(\sigma) = l(\bar{\partial}_{X+ir(X)}\sigma).$$

Such 1-derivations arising from holomorphic structures will be referred to as *Dolbeault 1-derivations*, and they are characterized by the fact that the corresponding linear (1,1)-tensor fields $K : TE \rightarrow TE$ are complex structures on the total space of E (see Example 2.5 below). Subbundles of E which are \mathcal{D}^{Dolb} -invariant (in the sense of Def. 2.1) are holomorphic subbundles.

As special cases, the holomorphic structures on TM and T^*M induced by a complex structure $r : TM \rightarrow TM$ are given by the 1-derivations \mathcal{D}^r and $\mathcal{D}^{r,*}$ from the previous example, with corresponding linear complex structures r^{tg} and r^{ctg} , see [9, § 5.1]. ◇

2.2. The Nijenhuis condition. Let $T : TN \rightarrow TN$ be a (1,1)-tensor field on a manifold N . The Nijenhuis torsion of T is $\mathcal{N}_T \in \Omega^2(E, TE)$ given by

$$\mathcal{N}_T(X_1, X_2) = [T(X_1), T(X_2)] - T([X_1, X_2]_T), \quad X_1, X_2 \in \mathfrak{X}(N),$$

where $[X_1, X_2]_T = [T(X_1), X_2] + [X_1, T(X_2)] - T([X_1, X_2])$ is the deformed bracket. If $\mathcal{N}_T = 0$, T is called a *Nijenhuis operator*.

By the equivalence between 1-derivations and linear (1,1)-tensor fields, properties of the latter can be expressed in terms of the former. Let $\mathcal{D} = (D, l, r)$ be a 1-derivation with corresponding linear (1,1)-tensor field K on E . It is proven in [4, § 6] that

$$(2.5) \quad \mathcal{N}_K = 0 \iff \begin{cases} \mathcal{N}_r = 0, \\ D_X(l(\sigma)) - l(D_X(\sigma)) = 0, \\ l(D_{[X,Y]}(\sigma)) - [D_X, D_Y](\sigma) - D_{[X,Y]_r}(\sigma) = 0, \end{cases}$$

where $[D_X, D_Y]$ is the commutator of operators on $\Gamma(E)$. We refer to the equations on the right hand side as the *Nijenhuis equations* for the 1-derivation (D, l, r) , and 1-derivations satisfying them will be called *Nijenhuis 1-derivations*.

We will also consider the algebraic condition $K^2 = -\text{id}_{TE}$, saying that K is an almost complex structure on the total space of E . In this case, it is also proven in [4, Cor. 6.2] that

$$(2.6) \quad K^2 = -\text{id}_{TE} \iff \begin{cases} r^2 = -\text{id}_{TM}, \\ l^2 = -\text{id}_E, \\ D_{r(X)}(\sigma) + l(D_X(\sigma)) = 0. \end{cases}$$

Example 2.5 (Dolbeault 1-derivations revisited). Following Example 2.4, a Dolbeault 1-derivation on $E \rightarrow M$ is a 1-derivation (D, l, r) on E satisfying both the Nijenhuis equations (2.5) and the almost complex equations (2.6); note that these conditions exactly say that (M, r) is a complex manifold and

$$(2.7) \quad \bar{\partial}_{X+i_r(X)} := -l(D_X)$$

is a flat $T^{0,1}$ -connection on the complex vector bundle (E, l) . We will think of a holomorphic vector bundle \mathcal{E} as a pair

$$\mathcal{E} = (E, \mathcal{D}^{Dolb})$$

given by a real vector bundle $E \rightarrow M$ equipped with a Dolbeault 1-derivation \mathcal{D}^{Dolb} . \diamond

2.3. Duality. Just as usual connections, 1-derivations on a vector bundle $E \rightarrow M$ possess a duality operation that establishes a bijection between 1-derivations on E and E^* . Given a 1-derivation $\mathcal{D} = (D, l, r)$ on E , its dual 1-derivation is $\mathcal{D}^* = (D^*, l^*, r)$, where $D^* : \Gamma(E^*) \rightarrow \Gamma(T^*M \otimes E^*)$ is characterized by

$$(2.8) \quad \langle D_X^*(\mu), \sigma \rangle = \mathcal{L}_X \langle \mu, l(\sigma) \rangle - \mathcal{L}_{r(X)} \langle \mu, \sigma \rangle - \langle \mu, D_X(\sigma) \rangle.$$

The corresponding linear (1,1)-tensor on E^* is denoted by

$$K^\top : T(E^*) \rightarrow T(E^*)$$

and admits the following characterization. If $\langle \cdot, \cdot \rangle : TE \times_{TM} T(E^*) \rightarrow TM \times \mathbb{R}$ is the non-degenerate, symmetric, bilinear pairing obtained from differentiation of the natural pairing $\langle \cdot, \cdot \rangle : E \times_M E^* \rightarrow M \times \mathbb{R}$, then

$$\langle \langle K(U), K^\top(V) \rangle \rangle = \langle \langle Tl(U), V \rangle \rangle, \quad \forall (U, V) \in TE \times_{TM} T(E^*).$$

Applying duality in Examples 2.2, 2.3 and 2.4, one obtains the following (see [9, § 2.1.2] for details):

- For the 1-derivation associated to a connection ∇ , the dual 1-derivation corresponds to the dual connection ∇^* on E^* .
- For $r : TM \rightarrow TM$, the 1-derivations \mathcal{D}^r and $\mathcal{D}^{r,*}$ (corresponding to r^{tg} and r^{ctg}) are dual to each other.
- A 1-derivation \mathcal{D} is Nijenhuis if and only if so is \mathcal{D}^* , see [9]; in this case, \mathcal{D} satisfies the almost complex conditions if and only if so does \mathcal{D}^* . As a consequence, \mathcal{D} is a Dolbeault 1-derivation if and only if so is \mathcal{D}^* . For a holomorphic vector bundle \mathcal{E} , the dual to its Dolbeault 1-derivation is the Dolbeault 1-derivation corresponding to the holomorphic structure of the dual \mathcal{E}^* , see Example 2.7 below.

It will be useful to extend, for each $X \in \mathfrak{X}(M)$, the operator D_X^* to $\Gamma(\wedge^m E^*)$ as follows:

$$(2.9) \quad D_X^*(f) = -\mathcal{L}_{r(X)}f,$$

$$(2.10) \quad D_X^*(\mu)(\sigma_1, \dots, \sigma_m) = \mathcal{L}_X \mu(l(\sigma_1), \dots, \sigma_m) - \mathcal{L}_{r(X)} \mu(\sigma_1, \dots, \sigma_m) \\ - \sum_{k=1}^m (-1)^{k-1} \mu(D_X(\sigma_k), \sigma_1, \dots, \widehat{\sigma}_k, \dots, \sigma_m),$$

where $f \in C^\infty(M)$, and $m \geq 1$. Note that $D_X^*(\mu)$ does not define an element of $\Gamma(\wedge^m E^*)$ in general, since it is not $C^\infty(M)$ -multilinear. In general, $D_X(\mu)(\sigma_1, \cdot)$ is skew-symmetric as a map from $(m-1)$ copies of $\Gamma(E)$ to $C^\infty(M)$, and it satisfies

$$\begin{aligned} D_X^*(\mu)(f\sigma_1, \cdot) &= fD_X^*(\mu)(\sigma_1, \cdot), \\ D_X^*(\mu)(\sigma_1, f\sigma_2, \cdot) &= fD_X^*(\mu)(\sigma_1, \sigma_2, \cdot) + (\mathcal{L}_X f)(\mu(l(\sigma_1), \sigma_2, \cdot) - \mu(\sigma_1, l(\sigma_2), \cdot)). \end{aligned}$$

Let us consider

$$\Gamma_l(\wedge^m E^*) = \{\mu \in \Gamma(\wedge^m E^*) \mid \mu(l(\sigma_1), \sigma_2, \cdot) = \mu(\sigma_1, l(\sigma_2), \cdot), \forall \sigma_1, \sigma_2 \in \Gamma(E)\}.$$

The next result follows immediately.

Proposition 2.6. *For $\mu \in \Gamma(\wedge^m E^*)$, $D_X^*(\mu) \in \Gamma(\wedge^m E^*)$ if and only if $\mu \in \Gamma_l(\wedge^m E^*)$. Moreover, if $[D_X, l] = D_X \circ l - l \circ D_X = 0$, then $D_X^*(\Gamma_l(\wedge^\bullet E^*)) \subset \Gamma_l(\wedge^\bullet E^*)$.*

Recall from (2.5) that condition $[D_X, l] = 0$ holds if \mathcal{D} is Nijenhuis.

In the following, for $\mu \in \Gamma(\wedge^m E^*)$, we shall denote by μ_l the element of $\Gamma(E^* \otimes \wedge^{m-1} E^*)$ given by $\mu_l = 0$ if $m = 0$, and

$$\mu_l(\sigma; \sigma_1, \dots, \sigma_{m-1}) = \mu(l(\sigma), \sigma_1, \dots, \sigma_{m-1}),$$

for $m \geq 1$, in such a way that $\mu \in \Gamma_l(\wedge^m E^*)$ if and only if $\mu_l \in \Gamma(\wedge^m E^*)$.

Example 2.7. For a holomorphic vector bundle $\mathcal{E} = (E, \mathcal{D}^{Dolb})$, with $\mathcal{D}^{Dolb} = (D, l, r)$, the dual 1-derivation $\mathcal{D}^{Dolb,*} = (D^*, l^*, r)$ is a Dolbeault 1-derivation on E^* corresponding to the holomorphic structure on the complex vector bundle (E^*, l^*) given by the $T^{0,1}$ -connection $\bar{\partial}^*$ dual to $\bar{\partial}$, see (2.7). Note also that (E^*, l^*) is identified with the bundle of complex-linear functionals $(E, l) \rightarrow \mathbb{C}$ via

$$(2.11) \quad \mu \mapsto \mu - \mathbf{i}\mu_l.$$

Let us denote the complex exterior algebra bundle of (E^*, l^*) by $\wedge_{\mathbb{C}} E^*$, so that $\Gamma(\wedge_{\mathbb{C}}^m E^*)$ is the bundle of complex-multilinear alternating m -forms on (E, l) , that carries a natural extension of $\bar{\partial}^*$. The map (2.11) extends to an isomorphism

$$\Phi : \Gamma_l(\wedge^m E^*) \rightarrow \Gamma(\wedge_{\mathbb{C}}^m E^*), \quad \Phi(\mu) = \mu - \mathbf{i}\mu_l,$$

of $C^\infty(M, \mathbb{C})$ -modules (multiplication by \mathbf{i} on $\Gamma_l(\wedge^m E^*)$ corresponds to the operation $\mu \mapsto \mu_l$). This isomorphism satisfies

$$\Phi(D_X^*(\mu)) = \mathbf{i}\bar{\partial}_{X+\mathbf{i}r(X)}^* \Phi(\mu)$$

showing that the extension of D^* (as in (2.9) and (2.10)) matches that of $\bar{\partial}^*$. In particular, $\mu \in \Gamma_l(\wedge^m E^*)$ satisfies $D^*(\mu) = 0$ if and only if $\mu - \mathbf{i}\mu_l \in \Gamma(\wedge_{\mathbb{C}}^m E^*)$ is a holomorphic section. Hence, for each open subset $U \subseteq M$, elements in $\Gamma_l(\wedge^* E^*|_U)$ in the kernel of D^* characterize holomorphic sections of $\Gamma(\wedge_{\mathbb{C}}^* E^*|_U)$ in terms of their real parts. \diamond

2.4. Compatibility with (pre-)Lie algebroids. The discussion in this subsection will be used later in the context of Dirac structures and lagrangian splittings of Courant algebroids.

Let us consider a vector bundle $A \rightarrow M$ equipped with an anchor map $\rho : A \rightarrow TM$ (i.e., a vector bundle map over the identity map on M) and an \mathbb{R} -bilinear, skew-symmetric bracket $[\cdot, \cdot]$ on $\Gamma(A)$ such that

$$[a, fb] = f[a, b] + (\mathcal{L}_{\rho(a)} f)b,$$

for $a, b \in \Gamma(A)$ and $f \in C^\infty(M)$. The triple $(A, \rho, [\cdot, \cdot])$ is called a *pre-Lie algebroid* [11]. A Lie algebroid is a pre-Lie algebroid such that $[\cdot, \cdot]$ satisfies the Jacobi identity.

Just as commonly done for Lie algebroids (see e.g. [20]), on a pre-Lie algebroid the anchor ρ and bracket $[\cdot, \cdot]$ can be encoded in a degree-1 derivation d_A of $\Gamma(\wedge A^*)$, or, alternatively, in a

linear bivector field π_A on the total space of $A^* \rightarrow M$. In terms of d_A and π_A , Lie algebroids are characterized by the further conditions that $d_A^2 = 0$ or that π_A is a Poisson structure.

Definition 2.8. *A 1-derivation $\mathcal{D} = (D, l, r)$ on a vector bundle $A \rightarrow M$ is compatible with a pre-Lie algebroid structure $(\rho, [\cdot, \cdot])$ if the following equations hold:*

$$\begin{aligned} \text{(IM1)} \quad & \rho \circ l = r \circ \rho \\ \text{(IM2)} \quad & \rho(D_X(a)) = D_X^r(\rho(a)) \\ \text{(IM3)} \quad & l([a, b]) = [a, l(b)] - D_{\rho(b)}(a) \\ \text{(IM4)} \quad & D_X([a, b]) = [a, D_X(b)] + [D_X(a), b] + D_{[\rho(b), X]}(a) - D_{[\rho(a), X]}(b), \end{aligned}$$

for all $X \in \mathfrak{X}(M)$ and $a, b \in \Gamma(A)$.

These compatibility equations have the following geometric interpretations. Let $K : TA \rightarrow TA$ be the linear (1,1)-tensor field corresponding to \mathcal{D} .

- The conditions in Def. 2.8 hold if and only if the bivector field π_A on A^* is compatible with the (1,1)-tensor field $K^\top : TA^* \rightarrow TA^*$ in the sense of Magri-Morosi [21], see [9, § 4.3] and Example 3.10 below. In particular, when A is a Lie algebroid and \mathcal{D} is a Nijenhuis 1-derivation, they hold if and only if the pair (π_A, K^\top) is a Poisson-Nijenhuis structure.
- When A is a Lie algebroid, the compatibility in Def. 2.8 says that $K : TA \rightarrow TA$ is a Lie algebroid morphism with respect to the tangent prolongation Lie algebroid $TA \rightarrow TM$ [4, § 6]. Hence 1-derivations compatible with a Lie algebroid are the infinitesimal counterparts of multiplicative (1,1)-tensor fields on Lie groupoids; for this reason (IM1)–(IM4) above are called *IM equations* (where IM stands for “infinitesimally multiplicative”), and 1-derivations satisfying them are also referred to as *IM (1,1)-tensors* on Lie algebroids.

The compatibility of a 1-derivation \mathcal{D} with a pre-Lie algebroid structure on A can be encoded using the dual 1-derivation \mathcal{D}^* and the operator d_A on $\Gamma(\wedge A^*)$ as follows.

Proposition 2.9. *A 1-derivation $\mathcal{D} = (D, l, r)$ is compatible with a pre-Lie algebroid structure $(\rho, [\cdot, \cdot])$ on a vector bundle $A \rightarrow M$ if and only if, for all $\mu \in \Gamma_l(\wedge^m A^*)$, the following holds:*

$$(2.12) \quad D_{\rho(a)}^*(\mu) = i_a d_A(\mu_l) - i_{l(a)} d_A \mu,$$

$$(2.13) \quad i_a d_A D_X^*(\mu) = D_X^*(d_A \mu)(a; \cdot) + \mathcal{R}_X(\mu)(a; \cdot),$$

where $\mathcal{R}_X(\mu)(a; \cdot)$ is the \mathbb{R} -multilinear skew-symmetric map from m -copies of $\Gamma(A)$ to $C^\infty(M)$ defined by

$$\begin{aligned} \mathcal{R}_X(\mu)(a; \sigma_1, \dots, \sigma_m) &= \mathcal{L}_X(D_{\rho(a)}^*(\mu)(\sigma_1, \dots, \sigma_m)) + D_{[\rho(a), X]}^*(\mu)(\sigma_1, \dots, \sigma_m) \\ &\quad - \sum_{i=1}^m (-1)^{i+1} D_{[\rho(\sigma_i), X]}^*(\mu)(a, \sigma_1, \dots, \hat{\sigma}_i, \dots, \sigma_m). \end{aligned}$$

Proof. For $\mu = f \in C^\infty(M)$, making use of (2.3) and (2.9), one can check that (2.12) and (2.13) are equivalent to

$$\mathcal{L}_{l(\rho(a)) - \rho(r(a))} f = 0 \quad \text{and} \quad D_X^*(d_A f) = \rho^* D_X^{r*}(df),$$

respectively. Using (2.8), one can now check that (2.12) and (2.13) in degree 0 are equivalent to (IM1) and (IM2). Similarly, for $\mu \in \Gamma(A)$ and assuming that (IM1) holds, one can check that (2.12) is equivalent to (IM3). Finally, under the assumption that (IM1), (IM2) and (IM3) hold, one verifies that (2.13) is equivalent to (IM4). For higher degrees, (2.12) and (2.13) follow directly from the IM equations. \square

Define

$$\Gamma_{\mathcal{D}}(\wedge^m A^*) = \{\mu \in \Gamma_l(\wedge^m A^*) \mid D_X^*(\mu) = 0, \forall X \in \mathfrak{X}(M)\}.$$

A direct consequence of Proposition 2.9 is that, when \mathcal{D} is compatible with the a pre-Lie algebroid structure on A ,

$$d_A(\Gamma_{\mathcal{D}}(\wedge^m A^*)) \subseteq \Gamma_{\mathcal{D}}(\wedge^m A^*),$$

so $(\Gamma_{\mathcal{D}}(\wedge^\bullet A^*), d_A)$ is a subcomplex of $(\Gamma(\wedge^\bullet A^*), d_A)$.

Example 2.10 (Holomorphic Lie algebroids). Consider a Dolbeault 1-derivation \mathcal{D}^{Dolb} on a vector bundle $A \rightarrow M$, so that $\mathcal{A} = (A, \mathcal{D}^{Dolb})$ is a holomorphic vector bundle. It is shown in [4, § 6.4] that a Lie algebroid structure on $A \rightarrow M$ compatible with \mathcal{D}^{Dolb} (in the sense of Def. 2.8) is equivalent to a holomorphic Lie algebroid structure on \mathcal{A} (in the sense of [17, § 3.1]). Under this correspondence, following Example 2.7, we see that for each open subset $U \subseteq M$, the complex $(\Gamma_{\mathcal{D}}(\wedge^\bullet A^*|_U), d_A)$ is identified with the holomorphic Lie algebroid complex (see [17, § 4.4]) of \mathcal{A} over U . \diamond

3. 1-DERIVATIONS ON COURANT ALGEBROIDS

In this section we introduce a notion of compatibility between 1-derivations and Courant algebroids that is the main object of study in this paper.

3.1. Courant 1-derivations and Courant-Nijenhuis algebroids. We start by recalling Courant algebroids [19, 23].

Definition 3.1. A *Courant algebroid* over a manifold M is a vector bundle $E \rightarrow M$ together with a bundle map $\mathfrak{a} : E \rightarrow TM$ (called the *anchor*), a pseudo-euclidean metric $\langle \cdot, \cdot \rangle$ (i.e, a fibrewise nondegenerate symmetric bilinear form), and an \mathbb{R} -bilinear bracket $[[\cdot, \cdot]] : \Gamma(E) \times \Gamma(E) \rightarrow \Gamma(E)$ such that, for all $\sigma_1, \sigma_2, \sigma_3 \in \Gamma(E)$ and $f \in C^\infty(M)$, the following hold:

- (C1) $[[\sigma_1, [[\sigma_2, \sigma_3]]]] = [[[[\sigma_1, \sigma_2], \sigma_3]] + [[\sigma_2, [[\sigma_1, \sigma_3]]]]$
- (C2) $\mathfrak{a}([[\sigma_1, \sigma_2]]) = [\mathfrak{a}(\sigma_1), \mathfrak{a}(\sigma_2)]$
- (C3) $[[\sigma_1, f\sigma_2]] = f[[\sigma_1, \sigma_2]] + (\mathcal{L}_{\mathfrak{a}(\sigma_1)}f)\sigma_2$
- (C4) $[[\sigma_1, \sigma_2]] + [[\sigma_2, \sigma_1]] = \mathfrak{a}^*(d\langle \sigma_1, \sigma_2 \rangle)$
- (C5) $\mathcal{L}_{\mathfrak{a}(\sigma_1)}\langle \sigma_2, \sigma_3 \rangle = \langle [[\sigma_1, \sigma_2], \sigma_3 \rangle + \langle \sigma_2, [[\sigma_1, \sigma_3]] \rangle$

where $\mathfrak{a}^* : T^*M \rightarrow E^* \simeq E$ is the map dual to the anchor \mathfrak{a} , and the isomorphism $E^* \simeq E$ is given by $\langle \cdot, \cdot \rangle$. We refer to $[[\cdot, \cdot]]$ as the *Courant bracket*.

The following is an important class of examples [25, 26].

Example 3.2 (Exact Courant algebroids). Any closed 3-form $H \in \Omega^3(M)$ defines a Courant algebroid structure on $E = \mathbb{T}M := TM \oplus T^*M$, with anchor map $\mathfrak{a} = \text{pr}_{TM}$, symmetric pairing

$$\langle (X, \alpha), (Y, \beta) \rangle = \beta(X) + \alpha(Y),$$

and the *H-twisted Courant bracket*

$$[[(X, \alpha), (Y, \beta)]]] = ([X, Y], \mathcal{L}_X\beta - i_Y d\alpha + i_Y i_X H).$$

When $H = 0$, one refers to this structure as the *standard Courant algebroid* on $\mathbb{T}M$. \diamond

Let $(E, \mathfrak{a}, \langle \cdot, \cdot \rangle, [[\cdot, \cdot]])$ be a Courant algebroid.

Definition 3.3. A *Courant 1-derivation* is a 1-derivation $\mathcal{D} = (D, l, r)$ on the vector bundle $E \rightarrow M$ such that $\mathcal{D}^* = \mathcal{D}$ (under the identification $E \cong E^*$ given by the pairing) and the following compatibility equations are satisfied:

$$\begin{aligned}
 \text{(CN1)} \quad & \mathbf{a} \circ l = r \circ \mathbf{a}, \\
 \text{(CN2)} \quad & \mathbf{a}(D_X(\sigma)) = D_X^r(\mathbf{a}(\sigma)), \\
 \text{(CN3)} \quad & l([\sigma_1, \sigma_2]) = [\sigma_1, l(\sigma_2)] - D_{\mathbf{a}(\sigma_2)}(\sigma_1) - \mathbf{a}^*(C(\sigma_1, \sigma_2)), \\
 \text{(CN4)} \quad & D_X([\sigma_1, \sigma_2]) = [\sigma_1, D_X(\sigma_2)] - [\sigma_2, D_X(\sigma_1)] + D_{[\mathbf{a}(\sigma_2), X]}(\sigma_1) \\
 & \quad - D_{[\mathbf{a}(\sigma_1), X]}(\sigma_2) - \mathbf{a}^*(i_X dC(\sigma_1, \sigma_2)),
 \end{aligned}$$

for all $\sigma_1, \sigma_2 \in \Gamma(E)$ and $X \in \mathfrak{X}(M)$, where $C(\sigma_1, \sigma_2) := \langle D_{(\cdot)}(\sigma_1), \sigma_2 \rangle \in \Omega^1(M)$.

Equations (CN1)-(CN4) are called *Courant compatibility equations* for \mathcal{D} . Note that they impose a linear condition on 1-derivations, so linear combinations of Courant 1-derivations are still Courant 1-derivations.

Definition 3.4. A *Courant-Nijenhuis 1-derivation* is a Courant 1-derivation that is also a Nijenhuis 1-derivation, i.e., satisfies the Nijenhuis equations (2.5). A Courant algebroid equipped with a Courant-Nijenhuis 1-derivation is a *Courant-Nijenhuis algebroid*.

Example 3.5. Consider a 1-derivation \mathcal{D} defined by a connection ∇ on $E \rightarrow M$, as in Example 2.2. If it is a Courant 1-derivation, then the anchor \mathbf{a} must be trivial (by (CN1)), so that, as a Courant algebroid, E is a bundle of quadratic Lie algebras. In this case, ∇ is a Courant 1-derivation if and only if it is symmetric (i.e., $\nabla = \nabla^*$ with respect to the pseudo-euclidean metric) as well as compatible with the metric and fibrewise Lie bracket (by (CN4)):

$$\mathcal{L}_X \langle \sigma_1, \sigma_2 \rangle = \langle \nabla_X \sigma_1, \sigma_2 \rangle + \langle \sigma_1, \nabla_X \sigma_2 \rangle, \quad \nabla_X([\sigma_1, \sigma_2]) = [\nabla_X \sigma_1, \sigma_2] + [\sigma_1, \nabla_X \sigma_2],$$

for $X \in \mathfrak{X}(M)$.

The Nijenhuis condition on \mathcal{D} amounts to the flatness of ∇ . In this case, assuming that M is connected and viewing its universal cover \widetilde{M} as a $\pi_1(M)$ -principal bundle over M , E is of the form $(\widetilde{M} \times \mathfrak{d})/\pi_1(M)$, where \mathfrak{d} is a quadratic Lie algebra equipped with a representation of $\pi_1(M)$ that preserves bracket and pairing. \diamond

The next example from [5] is the original motivation for the Courant compatibility equations (CN1)-(CN4).

Example 3.6. For a (1,1)-tensor field $r : TM \rightarrow TM$, consider the 1-derivations \mathcal{D}^r and $\mathcal{D}^{r,*}$ from (2.2) and (2.3). Then, setting

$$\mathbb{D}^r := (D^r, D^{r,*}),$$

we obtain a 1-derivation

$$(3.1) \quad \mathcal{D}^r = (\mathbb{D}^r, (r, r^*), r)$$

on TM that is a Courant 1-derivation with respect to the standard Courant algebroid structure (this is verified in the proof of [5, Lem. 6.1]). More generally, one can check that \mathcal{D}^r is a Courant 1-derivation with respect to an H -twisted Courant bracket as long as the closed 3-form $H \in \Omega^3(M)$ is compatible with r in the sense of [5, Def. 5.1], i.e.,

- the tensor field $H_r \in \Gamma(T^*M \otimes \wedge^2 T^*M)$, $H_r(X_1, X_2, X_3) := H(r(X_1), X_2, X_3)$, is skew-symmetric, that is, $H_r \in \Omega^3(M)$;
- $dH_r = 0$.

Moreover, r is a Nijenhuis operator if and only if \mathcal{D}^r is a Nijenhuis 1-derivation [5, Lem. 6.1], in which case it defines a Courant-Nijenhuis structure on TM . One can also directly verify that $r^2 = -\text{Id}_{TM}$ if and only if \mathcal{D}^r satisfies the almost complex equations in (2.6). When r is a

complex structure on M , \mathcal{D}^r is the Dolbeault 1-derivation codifying the holomorphic structure on $\mathbb{T}M \rightarrow M$ (see Example 2.4), with corresponding linear complex structure given by

$$(3.2) \quad K_r := (r^{tg}, r^{ctg}) : T(\mathbb{T}M) \rightarrow T(\mathbb{T}M).$$

◇

We will describe in § 4 modifications of the last example yielding more general Courant 1-derivations on $\mathbb{T}M$.

As we will see in § 5, for Dolbeault 1-derivations (see Example 2.4), the compatibility with Courant structures yields *holomorphic Courant algebroids*.

Remark 3.7 (On the compatibility equations). A natural issue concerning Courant 1-derivations is whether the Courant compatibility equations for \mathcal{D} in Def. 3.3 admit a geometric interpretation in terms of the corresponding linear (1,1)-tensor field K (c.f. the discussion after Def. 2.8). Although a satisfactory answer does not seem evident (in contrast with Lie algebroids, one can check that, in general, K does not define a Courant morphism [6] of the tangent Courant algebroids [3]), some key properties of K will be presented in Prop. 3.11 below. From another perspective, it would be interesting to see if the super-geometric viewpoint on Courant algebroids [23] can shed light on the Courant compatibility equations. ◇

3.2. Invariant Dirac structures. Much of the importance of Courant algebroids lies in their Dirac structures. We now consider Dirac structures invariant by Courant 1-derivations.

Let $\mathcal{D} = (D, l, r)$ be a Courant 1-derivation on a Courant algebroid $E \rightarrow M$. We will be concerned with lagrangian subbundles $(L \rightarrow M) \subseteq (E \rightarrow M)$ (i.e., $L = L^\perp$) that are \mathcal{D} -invariant in the sense of Def. 2.1:

- $l(L) \subset L$, and
- $D_X(\Gamma(L)) \subset \Gamma(L)$, for all $X \in \mathfrak{X}(M)$.

For lagrangian subbundles, these conditions can be equivalently expressed as follows. Consider the symmetric 2-form $S \in \Gamma(S^2 E^*)$,

$$S(\sigma_1, \sigma_2) = \langle l(\sigma_1), \sigma_2 \rangle.$$

(Recall that $l = l^*$ since $\mathcal{D}^* = \mathcal{D}$.) Let $C : \Gamma(E) \times \Gamma(E) \rightarrow \Omega^1(M)$ be the map that appeared in the definition of Courant 1-derivations,

$$C(\sigma_1, \sigma_2) = \langle D_{(\cdot)}(\sigma_1), \sigma_2 \rangle.$$

For a lagrangian subbundle $L \subset E$, the restriction of S to L defines an element $S_L \in \Gamma(S^2 L^*)$ with the property that $l(L) \subseteq L$ if and only if $S_L = 0$. Assuming that this holds, the restriction of C to sections of L is tensorial, i.e., it defines an element

$$C_L \in \Gamma(\wedge^2 L^* \otimes T^*M)$$

called the *concomitant of L and \mathcal{D}* . The following result can be directly verified.

Lemma 3.8. *For a Courant 1-derivation \mathcal{D} , a lagrangian subbundle $L \subseteq E$ is \mathcal{D} -invariant if and only if*

$$S_L = 0, \quad \text{and} \quad C_L = 0.$$

Suppose that \mathcal{D} is a Courant-Nijenhuis 1-derivation on E , so that (E, \mathcal{D}) is a Courant-Nijenhuis algebroid.

Definition 3.9. A Dirac structure $L \subset E$ that is \mathcal{D} -invariant (equivalently, such that $S_L = 0$ and $C_L = 0$) is called a *Dirac-Nijenhuis structure*.

Example 3.10 (Poisson-Nijenhuis structures). Given a (1,1)-tensor field $r : TM \rightarrow TM$, consider the associated Courant 1-derivation on $\mathbb{T}M$ (with the standard Courant bracket) given by \mathcal{D}^r , as in Example 3.6. Its invariant lagrangian subbundles are precisely those considered in [5, § 3.3]. In particular, a lagrangian subbundle given by the graph of a bivector field $\pi \in \mathfrak{X}^2(M)$,

$$L_\pi = \{(\pi^\sharp(\xi), \xi) \in \mathbb{T}M \mid \xi \in T^*M\},$$

where $\pi^\sharp : T^*M \rightarrow TM$ is the map obtained via contraction, is \mathcal{D}^r -invariant if and only if π and r satisfy

$$(3.3) \quad r \circ \pi^\sharp = \pi^\sharp \circ r^*, \quad \text{and} \quad \pi^\sharp \circ D_X^{r,*}(\alpha) - D_X^r \circ \pi^\sharp(\alpha) = \pi^\sharp(\mathcal{L}_X r^* \alpha - \mathcal{L}_{r(X)} \alpha) - (\mathcal{L}_{\pi^\sharp(\alpha)} r)(X) = 0$$

for all $X \in \mathfrak{X}(M)$, $\alpha \in \Omega^1(M)$. The expression in the second condition is known as the *Magri-Morosi concomitant* of r and π [21], see [5, § 3.1]. From the viewpoint of Lemma 3.8, using the natural identification $L_\pi \cong T^*M$, we have that

$$S_{L_\pi}(\alpha, \beta) = \pi(r^* \alpha, \beta) - \pi(\alpha, r^* \beta), \quad \text{and} \quad C_{L_\pi}(\alpha, \beta) = \langle \beta, \pi^\sharp \circ D_{(\cdot)}^{r,*}(\alpha) - D_{(\cdot)}^r \circ \pi^\sharp(\alpha) \rangle.$$

(The alternative formulation of the vanishing of the Magri-Morosi concomitant in terms of C_{L_π} goes back to [16] and is now more frequent in the literature.)

Recall that π is Poisson if and only if L_π is a Dirac structure, and r is a Nijenhuis operator if and only if \mathcal{D}^r is a Courant-Nijenhuis 1-derivation, so L_π is a Dirac-Nijenhuis structure in $(\mathbb{T}M, \mathcal{D}^r)$ if and only if (π, r) is a Poisson-Nijenhuis structure [5, Ex. 3.11]. \diamond

We refer to [5] for more on Dirac-Nijenhuis structures in the specific Courant-Nijenhuis algebroid $\mathbb{T}M$ of the previous example.

It is well known that if L is a Dirac structure in a Courant algebroid E , then the restrictions of the anchor and Courant bracket make L into a Lie algebroid. In the presence of a Courant 1-derivation \mathcal{D} for which L is \mathcal{D} -invariant, it is a straightforward verification that the 1-derivation on L obtained by restriction of \mathcal{D} is compatible with its Lie algebroid structure, in the sense of Def. 2.8. Following the discussion in § 2.4, in terms of linear (1,1)-tensor fields we have

Proposition 3.11. *Consider a Courant 1-derivation \mathcal{D} on a Courant algebroid E , and let $K : TE \rightarrow TE$ be the corresponding linear (1,1)-tensor field. Then a lagrangian subbundle $L \subset E$ is \mathcal{D} -invariant if and only if $K(TL) \subset TL$. If L is, in addition, a Dirac structure then $K|_{TL} : TL \rightarrow TL$ is a Lie algebroid morphism, where $TL \rightarrow TM$ is the tangent prolongation Lie algebroid.*

Proof. As recalled in § 2.1, the fact that L is \mathcal{D} -invariant is equivalent to $K(TL) \subset TL$, see [5, Thm. 2.1]. When L is a Dirac structure, the restricted 1-derivation satisfies the IM equations from Def. 2.8, which are equivalent to $K|_{TL} : TL \rightarrow TL$ being a Lie-algebroid morphism [4, § 5.2]. \square

4. COURANT 1-DERIVATIONS ON $TM \oplus T^*M$

We now present other examples of Courant 1-derivations on $\mathbb{T}M = TM \oplus T^*M$ with respect to twisted Courant brackets, extending Example 3.6

4.1. Courant 1-derivations from pseudo-Riemmanian metrics. A direct calculation shows that a 1-derivation $\mathcal{D} = (D, l, r)$ on $\mathbb{T}M$ satisfying $\mathcal{D}^* = \mathcal{D}$ and equations (CN1) and (CN2) must have the form

$$(4.1) \quad l = (r, r^* + g^\flat), \quad D = (D^r, D^{r,*} + \Sigma),$$

where $g^b : TM \rightarrow T^*M$, $g^b(X) = i_X g$, is the map obtained by contraction of a symmetric bilinear form g , and $\Sigma : \Gamma(TM) \rightarrow \Gamma(T^*M \otimes T^*M)$ is an \mathbb{R} -linear map satisfying

$$(4.2) \quad \Sigma_X(fY) = f \Sigma_X(Y) + (\mathcal{L}_X f) g^b(Y),$$

$$(4.3) \quad \mathcal{L}_X g(Y, Z) = \langle \Sigma_X(Y), Z \rangle + \langle Y, \Sigma_X(Z) \rangle,$$

where $f \in C^\infty(M)$, $X, Y, Z \in \mathfrak{X}(M)$, and $\Sigma_X : \Gamma(TM) \rightarrow \Gamma(T^*M)$ is given by $\Sigma_X(Y) = i_X(\Sigma(Y))$. In particular, any Courant 1-derivation $\mathcal{D} = (D, l, r)$ on $\mathbb{T}M$ is determined by the data r , g and Σ as above, via (4.1). When g is non-degenerate, i.e., when g is a *pseudo-Riemannian metric*, note that $\nabla_X = (g^b)^{-1} \circ \Sigma_X$ defines a metric connection on M (i.e., $\nabla g = 0$).

In the following, we shall assume that g is a pseudo-Riemannian metric and ∇ is its Levi-Civita connection. In this case g and ∇ give rise to a 1-derivation $\mathcal{D}^g = (\mathbb{D}^g, (0, g^b), 0)$ of $\mathbb{T}M$, where

$$\mathbb{D}_X^g((Y, \beta)) = (0, g^b(\nabla_X Y)),$$

by setting $r = 0$ in (4.1). The corresponding linear (1,1)-tensor field on $q : \mathbb{T}M \rightarrow M$,

$$(4.4) \quad K_g : T(\mathbb{T}M) \rightarrow T(\mathbb{T}M),$$

is described as follows. The connection ∇ defines a horizontal distribution $\text{Hor} \subset T(\mathbb{T}M)$, complementary to the vertical distribution $\text{Ver} = \ker(Tq) = q^*\mathbb{T}M$. With respect to this splitting of $T(\mathbb{T}M)$, K_g vanishes on Hor and acts as $(0, g^b)$ on Ver .

Lemma 4.1. *The 1-derivation \mathcal{D}^g is a Courant 1-derivation of $\mathbb{T}M$ for any H -twisted Courant bracket.*

Proof. Since \mathcal{D}^g is symmetric, it remains to show that the Courant equations (CN1)–(CN4) are satisfied. The only non-trivial equations to check are (CN3) and (CN4). Proving (CN3) amounts to verifying that

$$(4.5) \quad g^b([X, Y]) = \mathcal{L}_X g^b(Y) - g^b(\nabla_Y X) - g(\nabla_{(\cdot)} X, Y), \quad \forall X, Y \in \mathfrak{X}(M).$$

This holds because ∇ is Levi-Civita: since ∇ is metric and has zero torsion, we have

$$\begin{aligned} i_Z \mathcal{L}_X g^b(Y) &= \mathcal{L}_X g(Y, Z) - g(Y, [X, Z]) = g(\nabla_Y X + [X, Y], Z) + g(Y, \nabla_Z X) \\ &= i_Z \left(g^b(\nabla_Y X) + g^b([X, Y]) + g(Y, \nabla_{(\cdot)} X) \right). \end{aligned}$$

Condition (CN4), in turn, follows from the first Bianchi identity. Indeed, one must show that

$$\begin{aligned} g(\nabla_Z[X, Y], W) &= \langle \mathcal{L}_X g^b(\nabla_Z Y), W \rangle - \langle \mathcal{L}_Y g^b(\nabla_Z X), W \rangle + g(\nabla_{[Y, Z]} X, W) - g(\nabla_{[X, Z]} Y, W) \\ &\quad - i_W i_Z dg(\nabla_{(\cdot)} X, Y) \end{aligned}$$

Let us denote the right-hand side of this last equation by Υ . Using that

$$i_W i_Z dg(\nabla_{(\cdot)} X, Y) = \mathcal{L}_Z g(\nabla_W X, Y) - \mathcal{L}_W g(\nabla_Z X, Y) - g(\nabla_{[Z, W]} X, Y)$$

and, once again, the fact that ∇ is metric and has zero torsion, one obtains that

$$\begin{aligned} \Upsilon &= g(R(X, Z)(Y), W) - g(R(Y, Z)(X), W) - g(R(Z, W)(X), Y) + g(\nabla_Z \nabla_X Y - \nabla_Z \nabla_Y X, W) \\ &= \underbrace{g(R(X, Z)(Y), W) + g(R(Z, Y)(X), W) + g(R(Y, X)(Z), W)}_{=0 \text{ (by the first Bianchi identity)}} + g(\nabla_Z[X, Y], W), \end{aligned}$$

where R is the curvature tensor, and we used its symmetries in the last equality. This concludes the proof. \square

Now let

$$\mathcal{D}^{r,g} = (\mathbb{D}^{r,g}, (r, r^* + g^\flat), r)$$

be the 1-derivation defined by (4.1) with $\Sigma = g^\flat(\nabla)$, i.e., the sum of \mathcal{D}^r (see Example 3.6) with the 1-derivation \mathcal{D}^g .

Proposition 4.2. *The 1-derivation $\mathcal{D}^{r,g}$ is a Courant 1-derivation on $\mathbb{T}M$ for any H -twisted Courant bracket such that H is compatible with r .*

(The compatibility of H and r is recalled in Example 3.6).

Proof. The result follows from the previous lemma, the property that the sum of Courant 1-derivations is a Courant 1-derivation, and the fact that \mathcal{D}^r is a Courant 1-derivation with respect to any H -twisted Courant bracket such that H is compatible with r . \square

Remark 4.3. More generally, one can replace the Levi-Civita connection by any other metric connection ∇ in the definition of $\mathbb{D}^{r,g}$. In this case, equations (CN3) and (CN4) are equivalent to the torsion T of ∇ being skew-symmetric and closed (i.e., $\varphi(X, Y, Z) := g(T(X, Y), Z)$ defines a closed 3-form). \diamond

Example 4.4. The lagrangian subbundles $L \subset \mathbb{T}M$ which are \mathcal{D}^g -invariant are characterized, following Lemma 3.8, by the vanishing of $S_L \in \Gamma(S^2 L^*)$ and $C_L \in \Gamma(\wedge^2 L^* \otimes T^* M)$ given by

$$S_L(\sigma_1, \sigma_2) = g(\text{pr}_{TM}(\sigma_1), \text{pr}_{TM}(\sigma_2)) \quad \text{and} \quad C_L(\sigma_1, \sigma_2, Z) = g(\nabla_Z \text{pr}_{TM}(\sigma_1), \sigma_2).$$

The vanishing of S_L is equivalent to the presymplectic distribution of L being isotropic with respect to g . In particular, the only 2-form whose graph is \mathcal{D}^g -invariant is the zero 2-form. Under the assumption that $S_L = 0$, a sufficient condition for the vanishing of C_L is the invariance of $\text{pr}_{TM}(L)$ with respect to the Levi-Civita connection (and these conditions are equivalent when $\text{pr}_{TM}(L)$ is maximally isotropic). \diamond

4.2. Nijenhuis 1-derivations and the Kähler condition. Let $r : TM \rightarrow TM$ be a (1,1)-tensor field, and let g be a pseudo-Riemannian metric with Levi-Civita connection ∇ . We now give conditions on the pair (r, g) ensuring that the 1-derivation $\mathcal{D}^{r,g}$ satisfies the Nijenhuis equations, thereby defining a Courant-Nijenhuis 1-derivation (by Prop. 4.2). We will denote the corresponding linear (1,1)-tensor field on $\mathbb{T}M$ by

$$K_{r,g} = K_r + K_g,$$

see (3.2) and (4.4).

As recalled in Example 3.6, the 1-derivation \mathcal{D}^r satisfies the Nijenhuis equations if and only if r is a Nijenhuis operator ($\mathcal{N}_r = 0$). On the other hand, it is a simple verification that \mathcal{D}^g is always a Nijenhuis 1-derivation. More generally, we have

Proposition 4.5. *For a pair (r, g) , suppose that*

- r is a Nijenhuis operator,
- $g^\flat \circ r = -r^* \circ g^\flat$,
- $\nabla r = 0$.

Then the 1-derivation $\mathcal{D}^{r,g}$ satisfies the Nijenhuis equations (equivalently, $K_{r,g}$ has vanishing Nijenhuis torsion).

Proof. Since $\mathcal{N}_r = 0$ by assumption, one only has to check the remaining two Nijenhuis equations in (2.5). Using that \mathcal{D}^r is already a Courant-Nijenhuis 1-derivation on $\mathbb{T}M$ (see Example 3.6),

it suffices to show that

$$(4.6) \quad g^b(D_Z^r(X)) - D_Z^{r,*}(g^b(X)) = g^b(\nabla_Z r(X)) - r^*(g^b(\nabla_Z X)),$$

$$(4.7) \quad D_Y^{r,*}(g^b(\nabla_X Z)) - D_X^{r,*}(g^b(\nabla_Y Z)) + r^*g^b(\nabla_{[X,Y]}Z) = g^b\left(\nabla_X D_Y^r(Z) - \nabla_Y D_X^r(Z) - D_{[X,Y]}^r(Z) - \nabla_{[X,Y],r}Z\right).$$

Using (2.2) and (2.3), the compatibility $g^b \circ r = -r^* \circ g^b$ and that ∇ is Levi-Civita, we can prove that (4.6) is equivalent to

$$g^b((\nabla_X r)(Z)) + g((\nabla_{(\cdot)} r)(Z), X) = 0,$$

which holds since $\nabla r = 0$.

Regarding (4.7), we can use that $\nabla r = 0$ and that ∇ is Levi-Civita to rewrite some of its terms as follows:

$$\begin{aligned} \nabla_X D_Y^r(Z) &= \nabla_X r(\nabla_Y Z) - \nabla_X \nabla_{r(Y)} Z, \\ D_{[X,Y]}^r(Z) &= r(\nabla_{[X,Y]}Z) - \nabla_{r([X,Y])} Z, \\ D_X^{r,*}(g^b(\nabla_Y Z)) &= -g^b(\nabla_X r(\nabla_Y Z)) - g^b(\nabla_{r(X)} \nabla_Y Z). \end{aligned}$$

Similar formulas hold for $\nabla_Y D_X^r(Z)$ and $D_Y^{r,*}(g^b(\nabla_X Z))$. After some cancellations and regroupings, one obtains that (4.7) is equivalent to

$$g(R(Z, \cdot)(r(X)) - r(R(Z, \cdot)(X)), Y) = 0,$$

where we have used the symmetries of the curvature tensor R . Therefore (4.7) holds, since $[R, r] = 0$. This concludes the proof. \square

A pair (r, g) , where r is a (1,1)-tensor field and g is a pseudo-Riemannian metric on M , defines a (*pseudo-*) *Kähler structure* if r and g satisfy the three conditions in Prop. 4.5 with the additional requirement that $r^2 = -\text{Id}_{TM}$ (so that r is a complex structure). Pseudo-Kähler structures admit the following characterization in terms of 1-derivations.

Proposition 4.6. *A pair (r, g) defines a pseudo-Kähler structure if and only if the 1-derivation $\mathcal{D}^{r,g}$ satisfies the almost-complex equations in (2.6) (equivalently, $K_{r,g}^2 = -\text{Id}$).*

Proof. Recall that the almost-complex equations are

$$r^2 = -\text{id}_{TM}, \quad (r, r^* + g^b)^2 = -\text{id}_{T(TM)}, \quad \mathbb{D}_{r(X)}^{r,g}(\sigma) + (r, r^* + g^b)(\mathbb{D}_X^{r,g}(\sigma)) = 0$$

By the first equation, r is an almost complex structure. The second equation is equivalent to $r^* \circ g^b = -g^b \circ r$ (i.e., (g, r) is almost Hermitian). Since $r^2 = -\text{id}_{TM}$, the 1-derivation $(\mathbb{D}^r, (r, r^*), r)$ satisfies the almost complex equations (see Example 3.6), and using this fact one verifies that

$$\mathbb{D}_{r(X)}^{r,g}(Y, \beta) + (r, r^* + g^b)(\mathbb{D}_X^{r,g}(Y, \beta)) = (0, \langle \beta, \mathcal{N}_r(X, \cdot) \rangle + g^b((\nabla_Y r)(X))).$$

By setting $Y = 0$ or $\beta = 0$ above, we see that the third almost-complex equation is equivalent to $\mathcal{N}_r = 0$ and $\nabla r = 0$. \square

It follows from Propositions 4.5 and 4.6 that, for a pair (r, g) , if $K_{r,g}$ is an almost complex structure then it is automatically integrable.

Corollary 4.7. *A pseudo-Kähler metric g on a complex manifold (M, r) defines a holomorphic vector bundle structure on $TM \rightarrow M$ with Dolbeault 1-derivation $\mathcal{D}^{r,g}$ and linear complex structure on the total space TM given by $K_{r,g}$. Moreover, for any holomorphic 3-form H , $\mathcal{D}^{r,g}$ is a Courant 1-derivation with respect to the H -twisted Courant bracket on TM .*

The last assertion follows from Prop. 4.2.

For a pseudo-Kähler structure (r, g) , we will see below that there is an explicit isomorphism relating the holomorphic structures on $\mathbb{T}M \rightarrow M$ defined by \mathcal{D}^r and $\mathcal{D}^{r,g}$.

4.3. B-field transformations. Given a closed 2-form $B \in \Omega^2(M)$, the map

$$\tau_B : \mathbb{T}M \rightarrow \mathbb{T}M, \quad \tau_B(X, \alpha) = (X, \alpha + i_X B),$$

is a Courant automorphism of $\mathbb{T}M$ for any H -twisted Courant bracket (if B is not closed, then τ_B intertwines Courant brackets twisted by H and $(H - dB)$). Such τ_B is called a *gauge transformation* [26], or a *B-field transform* [12].

We say that two Courant 1-derivations $\mathcal{D}_1 = (D_1, l_1, r_1)$ and $\mathcal{D}_2 = (D_2, l_2, r_2)$ on $\mathbb{T}M$ are *gauge equivalent* if $r_1 = r_2$ and there exists a closed 2-form B such that

$$D_2 = \tau_B \circ D_1 \circ \tau_B^{-1} \text{ and } l_2 = \tau_B \circ l_1 \circ \tau_B^{-1}.$$

The Nijenhuis and almost complex equations (2.5) and (2.6), respectively, are invariant by gauge equivalence.

Consider a Courant 1-derivation \mathcal{D} determined by r, g and Σ via (4.1); here we no longer assume that the symmetric bilinear form g is nondegenerate. Then \mathcal{D} is gauge equivalent to \mathcal{D}^r if and only if there exists a closed 2-form B such that

$$(4.8) \quad g^b = B^b \circ r - r^* \circ B^b, \quad \Sigma_X = B^b \circ D_X^r - D_X^{r,*} \circ B^b.$$

In particular, τ_B preserves the 1-derivation \mathcal{D}^r (i.e., $g = 0$ and $\Sigma = 0$) if and only if B and r are compatible in the sense of Magri-Morosi, see [5, § 5.2] (cf. Example 3.6).

Suppose that r is a complex structure. Since \mathcal{D}^r satisfies the almost complex equations (2.6), any Courant 1-derivation \mathcal{D} gauge equivalent to it must satisfy these equations as well. In this case, if \mathcal{D} is determined by r, g and Σ (as in (4.1)), then necessarily $g^b \circ r = -r^* \circ g^b$ (see the proof of Proposition 4.6). So the pair (r, g) gives rise to a 2-form $\omega \in \Omega^2(M)$ such that $\omega^b = g^b \circ r$.

We will now describe a natural class of Courant 1-derivations on $\mathbb{T}M$ that are gauge equivalent to \mathcal{D}^r .

Proposition 4.8. *Suppose that \mathcal{D} is a Courant-Nijenhuis 1-derivation on $\mathbb{T}M$ determined by r, g and Σ (as in (4.1)) satisfying the almost complex equations. If the 2-form ω defined by $\omega^b = g^b \circ r$ is closed, then \mathcal{D} is gauge equivalent to \mathcal{D}^r .*

Proof. Let $B = \frac{1}{2}\omega$, and let \mathcal{D}^B be the Courant-Nijenhuis 1-derivation obtained by conjugating \mathcal{D} with the B-field transform τ_B . Denote by r, g^B and Σ^B the data that determine \mathcal{D}^B . Then it is straightforward to check that

$$g_B = 0 \quad \text{and} \quad \Sigma^B = \Sigma + (B^b(D^r) - D^{r,*}(B^b)).$$

Using (4.2) and (4.3) for \mathcal{D}^B , we see that Σ^B defines an element $H \in \Gamma(\wedge^2 T^*M \otimes T^*M)$ by

$$H(X, Y; Z) = \langle \Sigma_X^B(Y), Z \rangle.$$

As a consequence of (CN3), H is totally skew-symmetric, i.e. $H \in \Omega^3(M)^1$. It now follows directly from the almost complex equations and Nijenhuis equations that H must satisfy

$$H(r(X), Y, Z) = -H(X, r(Y), Z) \quad \text{and} \quad H(r(X), Y, Z) = H(X, r(Y), Z).$$

Hence $H = 0$, which means that $\Sigma^B = 0$. Therefore $\mathcal{D}^B = \mathcal{D}^r$, as we wanted to prove. \square

Corollary 4.9. *For a pseudo-Kähler structure (r, g) , the Courant 1-derivations \mathcal{D}^r and $\mathcal{D}^{r,g}$ are gauge equivalent.*

¹Although not needed in the proof, one can check that (CN4) is equivalent to $dH = 0$. In fact, Courant 1-derivations \mathcal{D} determined by r, g and Σ on $\mathbb{T}M$ with $g = 0$ are in bijective correspondence with closed 3-forms.

It follows that the usual holomorphic structure on $\mathbb{T}M \rightarrow M$ (defined by \mathcal{D}^r , see Example 3.6) and the one modified by the metric g (defined by $\mathcal{D}^{r,g}$, see Cor. 4.7) are isomorphic through a gauge transformation τ_B . Since τ_B establishes a bijective correspondence between the sets of \mathcal{D}^r -invariant and $\mathcal{D}^{r,g}$ -invariant lagrangian subbundles, it is clear that a lagrangian subbundle $L \subset \mathbb{T}M$ is holomorphic with respect to the holomorphic structure modified by g if and only if $\tau_B(L)$ is holomorphic in the usual sense. (In particular, this shows that $\mathcal{D}^{r,g}$ -invariant Dirac structures are much less restrictive than those for \mathcal{D}^g , c.f. Example 4.4).

5. HOLOMORPHIC COURANT ALGEBROIDS

In this section we show that holomorphic Courant algebroids can be seen as special cases of Courant-Nijenhuis structures.

For a complex manifold (M, r) and holomorphic vector bundle $\mathcal{E} \rightarrow M$, we denote by \mathcal{O} the sheaf of holomorphic functions on M and by $\Gamma_{\mathcal{E}}$ the sheaf of holomorphic sections of E .

Definition 5.1. *A holomorphic Courant algebroid is a holomorphic vector bundle $\mathcal{E} \rightarrow M$ endowed with a holomorphic non-degenerate symmetric \mathcal{O} -bilinear pairing $\langle \cdot, \cdot \rangle : \Gamma_{\mathcal{E}} \times \Gamma_{\mathcal{E}} \rightarrow \mathcal{O}$, a holomorphic vector bundle map $\mathfrak{p} : E \rightarrow T^{1,0}M$ and a \mathbb{C} -bilinear bracket $[[\cdot, \cdot]] : \Gamma_{\mathcal{E}} \times \Gamma_{\mathcal{E}} \rightarrow \Gamma_{\mathcal{E}}$ satisfying axioms (C1), \dots , (C5) in Definition 3.1 with \mathfrak{p} in place of \mathfrak{a} , f a local holomorphic function, and σ_1, σ_2 and σ_3 local holomorphic sections.*

For the sake of completeness, we start by recalling that there exists a natural (real) smooth Courant algebroid underlying a holomorphic one (see e.g. [13]). For a holomorphic vector bundle \mathcal{E} , we denote its underlying real, smooth vector bundle by E .

Proposition 5.2. *Let $(\mathcal{E}, \mathfrak{p}, [[\cdot, \cdot]], \langle \cdot, \cdot \rangle)$ be a holomorphic Courant algebroid. There exists a unique (real) smooth Courant algebroid structure $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{C^\infty}, [[\cdot, \cdot]]_{C^\infty})$ on E such that*

- (1) $\mathfrak{p}(\sigma) = \frac{1}{2}(\mathfrak{a}(\sigma) - \mathbf{i}r(\mathfrak{a}(\sigma)))$
- (2) $[[\sigma_1, \sigma_2]]_{C^\infty} = [[\sigma_1, \sigma_2]]$
- (3) $\langle \sigma_1, \sigma_2 \rangle_{C^\infty} = \text{Re}(\langle \sigma_1, \sigma_2 \rangle)$

for all local holomorphic sections $\sigma, \sigma_1, \sigma_2 \in \Gamma_{\mathcal{E}}(U)$.

Proof. We outline the proof, split in two parts: uniqueness and existence.

Uniqueness. We will show that the restriction of the real smooth Courant algebroid structure (satisfying (1), (2) and (3)) to holomorphic sections is sufficient to completely determine it. Let $U \subset M$ be an open subset for which there exists a frame of holomorphic sections $\{\sigma_1, \dots, \sigma_n\} \subset \Gamma_{\mathcal{E}}(U)$. Any smooth section σ of E over U can be expressed uniquely as

$$(5.1) \quad \sigma = \sum_{k=1}^n f_k \sigma_k + g_k l(\sigma_k),$$

for $f_k, g_k \in C^\infty(U)$, where $l : E \rightarrow E$ the fibrewise complex structure on E . From the $C^\infty(U)$ -linearity of \mathfrak{a} , we see that \mathfrak{a} is completely characterized by $\mathfrak{p}(\sigma_k)$ (note that the \mathbb{C} -linearity of \mathfrak{p} implies that $\mathfrak{a} \circ l = r \circ \mathfrak{a}$). Also, the $C^\infty(U)$ -bilinearity of $\langle \cdot, \cdot \rangle_{C^\infty}$ implies that it is determined by $\langle \sigma_j, \sigma_k \rangle$, noticing that

$$(5.2) \quad \langle \sigma_j, l(\sigma_k) \rangle_{C^\infty} = -\text{Im}(\langle \sigma_j, \sigma_k \rangle).$$

Finally, the Leibniz equation (C3) implies that $[[\cdot, \cdot]]_{C^\infty}$ is completely determined by $[[\sigma_j, \sigma_k]]$ (note that $[[\sigma_j, l(\sigma_k)]]_{C^\infty} = l([[\sigma_j, \sigma_k]])$).

Existence. By uniqueness, it suffices to describe the real smooth Courant algebroid structure locally. We will use the isomorphism of $C^\infty(U)$ -modules

$$C^\infty(U, \mathbb{C}) \otimes_{\mathcal{O}(U)} \Gamma \mathcal{E}(U) \ni (f + \mathbf{i}g) \otimes \sigma \mapsto f\sigma + gl(\sigma) \in \Gamma(U, E)$$

to construct the Courant algebroid structure locally. The anchor and the bracket are given by

$$\mathbf{a}(\psi \otimes \sigma) = \operatorname{Re}(\psi)\mathbf{a}(\sigma) + \operatorname{Im}(\psi)r(\mathbf{a}(\sigma)), \quad \langle \psi_1 \otimes \sigma_1, \psi_2 \otimes \sigma_2 \rangle_{C^\infty} = \operatorname{Re}(\psi_1\psi_2 \langle \sigma_1, \sigma_2 \rangle).$$

Note that $\langle \cdot, \cdot \rangle_{C^\infty}$ is non-degenerate and, for $h \in \mathcal{O}(U)$,

$$(5.3) \quad \mathcal{L}_{\mathbf{a}(\psi \otimes \sigma)} h = \psi \mathcal{L}_{\mathbf{p}(\sigma)} h.$$

We can now define $\mathbf{a}^* : \Omega^1(U) \rightarrow C^\infty(U, \mathbb{C}) \otimes_{\mathcal{O}(U)} \mathcal{E}(U)$ naturally as

$$\langle \mathbf{a}^*(\alpha), \psi \otimes \sigma \rangle_{C^\infty} = i_{\mathbf{a}(\psi \otimes \sigma)} \alpha.$$

The Courant bracket on $C^\infty(U, \mathbb{C}) \otimes_{\mathcal{O}(U)} \mathcal{E}(U)$ is given by

$$(5.4) \quad \begin{aligned} \llbracket \psi_1 \otimes \sigma_1, \psi_2 \otimes \sigma_2 \rrbracket_{C^\infty} = & \psi_1\psi_2 \otimes \llbracket \sigma_1, \sigma_2 \rrbracket + \mathcal{L}_{\mathbf{a}(\psi_1 \otimes \sigma_1)}(\psi_2) \otimes \sigma_2 - \mathcal{L}_{\mathbf{a}(\psi_2 \otimes \sigma_2)}(\psi_1) \otimes \sigma_1 \\ & + \operatorname{Re}(\psi_2 \langle \sigma_1, \sigma_2 \rangle) \mathbf{a}^* d\operatorname{Re}(\psi_1) - \operatorname{Im}(\psi_2 \langle \sigma_1, \sigma_2 \rangle) \mathbf{a}^* d\operatorname{Im}(\psi_1). \end{aligned}$$

One can check that \mathbf{a} , $\langle \cdot, \cdot \rangle_{C^\infty}$ and $\llbracket \cdot, \cdot \rrbracket_{C^\infty}$ are well-defined in the sense that their definitions agree on $(h\psi) \otimes \sigma$ and $\psi \otimes (h\sigma)$, for $h \in \mathcal{O}(U)$. The axioms (C1)–(C5) follow by direct inspection. \square

Following Examples 2.4 and 2.5, we regard a holomorphic vector bundle \mathcal{E} as a pair $(E, \mathcal{D}^{\operatorname{Dolb}})$, where $E \rightarrow M$ is a real, smooth vector bundle endowed with a Dolbeault 1-derivation $\mathcal{D}^{\operatorname{Dolb}}$ defining its holomorphic structure. Our main result in this section is an equivalence between holomorphic Courant algebroid structures on \mathcal{E} and real Courant algebroid structures on E for which $\mathcal{D}^{\operatorname{Dolb}}$ is a Courant 1-derivation.

Theorem 5.3. *Consider a holomorphic vector bundle $\mathcal{E} = (E, \mathcal{D}^{\operatorname{Dolb}})$.*

- (a) *If $(\mathbf{p}, \langle \cdot, \cdot \rangle, \llbracket \cdot, \cdot \rrbracket)$ is a holomorphic Courant algebroid structure on \mathcal{E} , then $\mathcal{D}^{\operatorname{Dolb}}$ is a Courant 1-derivation of the underlying real Courant algebroid structure on E (so $\mathcal{D}^{\operatorname{Dolb}}$ makes E into a Courant-Nijenhuis algebroid).*
- (b) *If $(\mathbf{a}, \langle \cdot, \cdot \rangle, \llbracket \cdot, \cdot \rrbracket)$ is a Courant algebroid structure on E such that $\mathcal{D}^{\operatorname{Dolb}}$ is a Courant 1-derivation, then $\llbracket \cdot, \cdot \rrbracket$ restricts to a \mathbb{C} -bilinear bracket on $\Gamma \mathcal{E}$ in such a way that it defines a holomorphic Courant algebroid structure on \mathcal{E} together with*

$$\mathbf{p}(\sigma) = \frac{1}{2}(\mathbf{a}(\sigma) - \mathbf{i}r(\mathbf{a}(\sigma))), \quad \langle \sigma_1, \sigma_2 \rangle_{\operatorname{hol}} = \langle \sigma_1, \sigma_2 \rangle - \mathbf{i} \langle \sigma_1, l(\sigma_2) \rangle.$$

The constructions in (a) and (b) are inverses of one another.

Proof. To prove part (a), we must show that $\mathcal{D}^{\operatorname{Dolb}} = (D, l, r)$ is a Courant 1-derivation, i.e., that it satisfies (CN1)–(CN4) in Def. 3.3 with respect to the real Courant algebroid E . The \mathbb{C} -linearity of \mathbf{p} implies that $\mathbf{a} \circ l = r \circ \mathbf{a}$, so (CN1) holds. To verify conditions (CN2), (CN3) and (CN4), we must check, for each open subset $U \subset M$, the vanishing of the following expressions:

$$\begin{aligned} W_2(X, \sigma_1) &= \mathbf{a}(D_X(\sigma_1)) - D_X^r(\mathbf{a}(\sigma_1)), \\ W_3(\sigma_1, \sigma_2) &= l(\llbracket \sigma_1, \sigma_2 \rrbracket_{C^\infty}) - (\llbracket \sigma_1, l(\sigma_2) \rrbracket_{C^\infty} - D_{\mathbf{a}(\sigma_2)}(\sigma_1) - \mathbf{a}^*(C(\sigma_1, \sigma_2))), \\ W_4(X, \sigma_1, \sigma_2) &= D_X(\llbracket \sigma_1, \sigma_2 \rrbracket_{C^\infty}) - (\llbracket \sigma_1, D_X(\sigma_2) \rrbracket_{C^\infty} - \llbracket \sigma_2, D_X(\sigma_1) \rrbracket_{C^\infty} + D_{[\mathbf{a}(\sigma_2), X]}(\sigma_1) \\ &\quad - D_{[\mathbf{a}(\sigma_1), X]}(\sigma_2) - \mathbf{a}^*(i_X dC(\sigma_1, \sigma_2))), \end{aligned}$$

for $X \in \mathfrak{X}(U)$ and σ_1, σ_2 smooth sections of E over U .

The vanishing of W_2 follows from the fact that, since \mathbf{a} is holomorphic, it must intertwine $\mathcal{D}^{\operatorname{Dolb}}$ and \mathcal{D}^r (recalling that \mathcal{D}^r is the Dolbeault 1-derivation encoding the holomorphic structure on TM , see Example 2.4), so (CN2) holds.

Using that $\mathfrak{a} \circ l = r \circ \mathfrak{a}$ and $l = l^*$ (by (5.2)), one can show that W_3 is $C^\infty(U)$ -linear in each entry, whereas W_4 is $C^\infty(U)$ -linear in the first entry and satisfies

$$\begin{aligned} W_4(X, \sigma_1, f\sigma_2) - fW_4(X, \sigma_1, \sigma_2) &= (\mathcal{L}_X f) W_3(\sigma_1, \sigma_2) - (\mathcal{L}_{W_2(X, \sigma_1)} f) \sigma_2 \\ &= (\mathcal{L}_X f) W_3(\sigma_1, \sigma_2), \\ W_4(X, f\sigma_1, \sigma_2) - fW_4(X, \sigma_1, \sigma_2) &= -(\mathcal{L}_X f) W_3(\sigma_2, \sigma_1) - W_1(X, \sigma_1, \sigma_2) \mathfrak{a}^* df \end{aligned}$$

where

$$W_1(X, \sigma_1, \sigma_2) = \langle D_X(\sigma_1), \sigma_2 \rangle_{C^\infty} + \langle \sigma_1, D_X(\sigma_2) \rangle_{C^\infty} - \mathcal{L}_X \langle \sigma_1, l(\sigma_2) \rangle_{C^\infty} + \mathcal{L}_{r(X)} \langle \sigma_1, \sigma_2 \rangle_{C^\infty}.$$

Note that W_1 is $C^\infty(U)$ -linear in each entry due to (2.1).

We claim that W_1 , W_3 and W_4 vanish on holomorphic sections. Indeed, for W_3 and W_4 this follows from the facts that D_X is zero on holomorphic sections (by (2.4)) and that $[\cdot, \cdot]_{C^\infty}$ restricts to the \mathbb{C} -bilinear holomorphic Courant bracket on holomorphic sections. For W_1 , we use the additional fact that

$$\mathcal{L}_X \langle \sigma_1, l(\sigma_2) \rangle_{C^\infty} - \mathcal{L}_{r(X)} \langle \sigma_1, \sigma_2 \rangle_{C^\infty} = -\text{Im}(\mathcal{L}_{X+ir(X)} \langle \sigma_1, \sigma_2 \rangle) = 0,$$

for holomorphic $\sigma_1, \sigma_2 \in \Gamma_{\mathcal{E}}(U)$, since $\langle \sigma_1, \sigma_2 \rangle \in \mathcal{O}(U)$.

Since smooth sections can be locally expressed by means of a frame of holomorphic sections as in (5.1), the $C^\infty(U)$ -multilinearity of W_1 and W_3 implies that they vanish. This in turn implies that W_4 is also multilinear over $C^\infty(U)$, and hence also vanishes. This concludes the proof of (a).

Let us prove part (b). Since D vanishes on holomorphic sections, (CN4) implies that $[\Gamma_{\mathcal{E}}(U), \Gamma_{\mathcal{E}}(U)] \subset \Gamma_{\mathcal{E}}(U)$, and it follows from (CN3) that the restricted bracket on holomorphic sections is \mathbb{C} -bilinear. Regarding the anchor, (CN1) implies that \mathfrak{p} is \mathbb{C} -linear, and (CN2) says that it is holomorphic. It remains to check that $\langle \Gamma_{\mathcal{E}}(U), \Gamma_{\mathcal{E}}(U) \rangle \subset \mathcal{O}(U)$. This is a consequence of the fact that, for local holomorphic sections σ_1, σ_2 , the duality equation (2.8) implies that

$$\mathcal{L}_{X+ir(X)} \langle \sigma_1, \sigma_2 \rangle_{\text{hol}} = \langle \bar{\partial}_{X+ir(X)} \sigma_1, \sigma_2 \rangle_{\text{hol}} + \langle \sigma_1, \bar{\partial}_{X+ir(X)} \sigma_2 \rangle_{\text{hol}} = 0,$$

where $\bar{\partial}$ is the flat $T^{0,1}$ -connection (2.4). Therefore $\langle \sigma_1, \sigma_2 \rangle_{\text{hol}}$ is holomorphic, thus concluding the proof. \square

Example 5.4. Let (M, r) be a complex manifold and H a closed holomorphic 3-form. Then $\mathbb{T}M$, viewed as a holomorphic vector bundle, carries an H -twisted holomorphic Courant algebroid structure (analogous to Example 3.2). From the perspective of Theorem 5.3, this corresponds to the fact that \mathcal{D}^r is a Courant 1-derivation of the H -twisted Courant bracket on $\mathbb{T}M$, viewed as a real vector bundle (see Example 3.6). \diamond

Remark 5.5. Any holomorphic Courant algebroid whose underlying real Courant algebroid is $\mathbb{T}M$ with the H -twisted Courant bracket, for a closed 3-form H , is completely characterized by a Courant-Nijenhuis 1-derivation \mathcal{D} on $(\mathbb{T}M, H)$ determined by the data r, g and Σ (as in (4.1)), where r is the complex structure on M . Then g and Σ must satisfy the equations corresponding to the fact that \mathcal{D} is Courant-Nijenhuis and almost complex. If the 2-form $\omega^b = g^b \circ r$ is closed, Proposition 4.8 implies that the holomorphic Courant algebroid determined by \mathcal{D} is isomorphic to the one determined by \mathcal{D}^r . In general this is not the case, but one can always use ω to gauge away g leaving the construction of more general holomorphic Courant algebroids as a problem of choosing Σ suitably; this provides a different approach to the classification of holomorphic Courant algebroids in [13, Prop.1.3]. \diamond

6. LAGRANGIAN SPLITTINGS AND DOUBLES

6.1. Lagrangian splittings of Courant algebroids. Let $(E, \langle \cdot, \cdot \rangle, \mathfrak{a}, \llbracket \cdot, \cdot \rrbracket)$ be a Courant algebroid. By a *lagrangian splitting* of E we mean a decomposition $E = A \oplus B$, where A and B are lagrangian subbundles. In this case, there is an isomorphism $B \cong A^*$ via the pairing that yields an identification $E = A \oplus A^*$ as pseudo-euclidean vector bundles, where $A \oplus A^*$ is equipped with its canonical pairing

$$\langle (a, \alpha), (b, \beta) \rangle := \beta(a) + \alpha(b).$$

Let $p_A : E \rightarrow A$ and $p_{A^*} : E \rightarrow A^*$ the natural projections onto A and A^* , respectively. The anchor and bracket on E induce the following structures on A and A^* : an anchor $\rho := \mathfrak{a}|_A : A \rightarrow TM$, along with an \mathbb{R} -bilinear bracket $[\cdot, \cdot]$ on $\Gamma(A)$ and an element $\varphi \in \Gamma(\wedge^3 A^*)$ given by

$$[a, b] := p_A(\llbracket (a, 0), (b, 0) \rrbracket), \quad i_b i_a \varphi := p_{A^*}(\llbracket (a, 0), (b, 0) \rrbracket),$$

and, similarly, an anchor $\rho_* : A^* \rightarrow TM$, bracket $[\cdot, \cdot]_*$ on $\Gamma(A^*)$ and element $\chi \in \Gamma(\wedge^3 A)$. Note that A and A^* , endowed with their anchors and brackets, become pre-Lie algebroids (see § 2.4). We denote the corresponding operators by

$$d_A : \Gamma(\wedge^\bullet A^*) \rightarrow \Gamma(\wedge^{\bullet+1} A^*), \quad d_{A^*} : \Gamma(\wedge^\bullet A) \rightarrow \Gamma(\wedge^{\bullet+1} A).$$

With respect to these structures, the Courant bracket on $E = A \oplus A^*$ is given by

$$(6.1) \quad \llbracket (a, \alpha), (b, \beta) \rrbracket = ([a, b] + \mathcal{L}_\alpha b - i_\beta d_{A^*} a + i_\beta i_\alpha \chi, [\alpha, \beta]_* + \mathcal{L}_a \beta - i_b d_A \alpha + i_b i_a \varphi),$$

where $\mathcal{L}_\alpha = i_\alpha d_{A^*} + d_{A^*} i_\alpha$, similarly for \mathcal{L}_a .

Let $(A, \rho, [\cdot, \cdot])$ and $(A^*, \rho_*, [\cdot, \cdot]_*)$ be pre-Lie algebroids in duality, further equipped with sections $\varphi \in \Gamma(\wedge^3 A^*)$ and $\chi \in \Gamma(\wedge^3 A)$. The pair (A, A^*) is called a *proto bialgebroid* when the anchors, brackets and 3-sections satisfy compatibility conditions (spelled out in [24], see also [14]) saying that the bracket (6.1) on $\Gamma(A \oplus A^*)$ makes $A \oplus A^*$ into a Courant algebroid with anchor $\mathfrak{a} = \rho + \rho_*$ and canonical pairing, called the *double* of (A, A^*) .

We therefore obtain the following equivalence: any Courant algebroid equipped with a lagrangian splitting yields a proto bialgebroid, and any proto bialgebroid gives rise, by means of its double, to a Courant algebroid endowed with a lagrangian splitting.

The following are special cases of interest of this correspondence.

- When $\varphi = 0$ and $\chi = 0$, a proto bialgebroid (A, A^*) is a Lie bialgebroid, i.e., (A, A^*) is a pair of Lie algebroids $(A, \rho, [\cdot, \cdot])$ and $(A^*, \rho_*, [\cdot, \cdot]_*)$ in duality such that the Lie-algebroid differential d_{A^*} and the Schouten bracket $[\cdot, \cdot]$ on $\Gamma(\wedge A)$ satisfy

$$(6.2) \quad d_{A^*}[a, b] = [d_{A^*} a, b] + [a, d_{A^*} b], \quad \forall a, b \in \Gamma(A).$$

Lie bialgebroids are in correspondence with Courant algebroids equipped with a lagrangian splitting by Dirac structures [19], known as *Manin triples*.

- When $\chi = 0$, a proto bialgebroid (A, A^*) is a Lie quasi-bialgebroid [24], in which case A is a Lie algebroid, d_{A^*} satisfies (6.2), $d_{A^*}^2 = [\chi, \cdot]$, and $d_{A^*} \chi = 0$. Lie quasi-bialgebroids correspond to Courant algebroids equipped with a splitting given by a Dirac structure and a lagrangian complement, known as *Manin quasi-triples*.

For a Courant algebroid $E \rightarrow M$, a Lagrangian splitting $E = A \oplus A^*$ induces a bivector field π on M via

$$(6.3) \quad \pi^\sharp = \rho_* \circ \rho^* : T^*M \rightarrow TM$$

that satisfies

$$\frac{1}{2}[\pi, \pi] = \rho(\chi) + \rho_*(\varphi),$$

see [18, § 3.2 and 3.4]. In particular π is a Poisson structure when (A, A^*) is a Lie bialgebroid.

6.2. Lagrangian splittings of Courant 1-derivations. Consider a Courant algebroid E equipped with a lagrangian splitting, that we write as $E = A \oplus A^*$.

Assumption. We assume throughout this subsection that

$$\mathcal{D} = (\mathbb{D}, \ell, r)$$

is a 1-derivation on the vector bundle E that keeps the splitting invariant, i.e., the subbundles A and A^* are \mathcal{D} -invariant (as in Def. 2.1). We will further assume that \mathcal{D} is symmetric ($\mathcal{D} = \mathcal{D}^*$), in which case the restricted 1-derivations on A and A^* are dual to one another.

We denote the 1-derivation on A by $\mathcal{D} = (D, l, r)$, so that the 1-derivation on A^* is $\mathcal{D}^* = (D^*, l^*, r)$ and

$$\mathbb{D} = (D, D^*), \quad \ell = (l, l^*).$$

We keep the notation from § 6.1 for the pre-Lie algebroids $(A, \rho, [\cdot, \cdot])$ and $(A, \rho_*, [\cdot, \cdot]_*)$, with 3-sections $\chi \in \Gamma(\wedge^3 A)$ and $\varphi \in \Gamma(\wedge^3 A^*)$.

Theorem 6.1. *The 1-derivation $\mathcal{D} = (\mathbb{D}, \ell, r)$ is a Courant 1-derivation of $E = A \oplus A^*$ if and only if the following conditions hold:*

- \mathcal{D} is compatible with the pre-Lie algebroid $(A, \rho, [\cdot, \cdot])$, and

$$\varphi \in \Gamma_l(\wedge^3 A^*), \quad D^*(\varphi) = 0;$$

- \mathcal{D}^* is compatible with the pre-Lie algebroid $(A^*, \rho_*, [\cdot, \cdot]_*)$, and

$$\chi \in \Gamma_{l^*}(\wedge^3 A), \quad D(\chi) = 0.$$

Moreover, \mathcal{D} is Nijenhuis (resp. Dolbeault) if and only if so is \mathcal{D} .

Proof. We must show the equivalence between conditions (CN1)–(CN4) (in Definition 3.3) for \mathcal{D} and (IM1)–(IM4) (in Definition 2.8) for both \mathcal{D} and \mathcal{D}^* along with the conditions on φ and χ in the statement.

It directly follows from $\mathfrak{a} = \rho + \rho_*$ and $\ell = (l, l^*)$ that (CN1) for \mathcal{D} is equivalent to (IM1) for \mathcal{D} and \mathcal{D}^* . Similarly, the fact that $\mathbb{D} = (D, D^*)$ implies that (CN2) for \mathcal{D} is equivalent to (IM2) for \mathcal{D} and \mathcal{D}^* .

Claim 1. Assume that (IM1) holds for \mathcal{D} and \mathcal{D}^* . Then condition (CN3) for \mathcal{D} is equivalent to (IM3) for \mathcal{D} and \mathcal{D}^* , as well as $\varphi \in \Gamma_l(\wedge^3 A^*)$ and $\chi \in \Gamma_{l^*}(\wedge^3 A)$.

Let us verify the claim. For sections of type $\sigma_1 = (a, 0)$, $\sigma_2 = (b, 0)$, (CN3) becomes

$$\ell([a, b] + i_b i_a \varphi) = [a, l(b)] + i_{l(b)} i_a \varphi - D_{\rho(b)}(a),$$

which splits into

$$l([a, b]) = [a, l(b)] - D_{\rho(b)}(a) \quad \text{and} \quad l^*(i_b i_a \varphi) = i_{l(b)} i_a \varphi.$$

These conditions hold for all $a, b \in \Gamma(A)$ if and only if (IM3) holds for \mathcal{D} and $\varphi \in \Gamma_l(\wedge^3 A^*)$. Similarly, (CN3) for sections of type $\sigma_1 = (0, \alpha)$, $\sigma_2 = (0, \beta)$ is equivalent to (IM3) for \mathcal{D}^* and $\chi \in \Gamma_{l^*}(\wedge^3 A)$. For sections of type $\sigma_1 = (0, \alpha)$, $\sigma_2 = (b, 0)$, (CN3) amounts to two equations:

$$(6.4) \quad l^*(i_b d_A \alpha) = i_{l(b)} d_A \alpha + D_{\rho(b)}^*(\alpha) + \langle D_{\rho(\cdot)}^*(\alpha), b \rangle,$$

$$(6.5) \quad l(\mathcal{L}_\alpha b) = \mathcal{L}_\alpha l(b) - \langle D_{\rho_*(\cdot)}^*(\alpha), b \rangle.$$

We will see that (6.4) (resp. (6.5)) follows directly from (2.12) for \mathcal{D} (resp. \mathcal{D}^*) and $m = 1$, which is equivalent to (IM3) under the assumption that (IM1) holds. Indeed, note that (2.12) for $m = 1$ has the following alternative formulations

$$(6.6) \quad \langle D_{\rho(\cdot)}^*(\alpha), b \rangle = l^*(i_b d_A \alpha) - i_b d_A(l^*(\alpha)) \quad (\text{similarly, } \langle \alpha, D_{\rho_*(\cdot)}(b) \rangle = l(i_\alpha d_{A^*} b) - i_\alpha d_A(l(b)).)$$

So (6.4) is obtained from adding up (2.12) and (6.6). The second equation (6.5) follows from the Cartan formula $\mathcal{L}_\alpha = i_\alpha d_{A^*} + d_{A^*} i_\alpha$ together with (2.8) and (6.6). The equations corresponding to (CN3) for sections of type $\sigma_1 = (a, 0)$, $\sigma_2 = (0, \beta)$ are entirely analogous to (6.4) and (6.5), and hold for similar reasons. This proves the claim.

Claim 2. Assume that (IM1), (IM2) and (IM3) hold for \mathcal{D} and \mathcal{D}^* . Then condition (CN4) for \mathcal{D} is equivalent to (IM4) for \mathcal{D} and \mathcal{D}^* , as well as $D^*(\varphi) = 0$ and $D(\chi) = 0$.

To verify the claim, note that for sections of type $\sigma_1 = (a, 0)$, $\sigma_2 = (b, 0)$, (CN4) takes the form

$$\begin{aligned} D_X([a, b]) + D_X^*(i_b i_a \varphi) &= [a, D_X(b)] + i_{D_X(b)} i_a \varphi - [b, D_X(a)] - i_{D_X(a)} i_b \varphi \\ &\quad + D_{[\rho(b), X]}(a) - D_{[\rho(a), X]}(b). \end{aligned}$$

So in this case (CN4) amounts to (IM4) for \mathcal{D} together with the following condition on φ (using (2.8)):

$$\mathcal{L}_X \varphi(a, b, l(c)) - \mathcal{L}_{r(X)} \varphi(a, b, c) - \varphi(D_X a, b, c) - \varphi(a, D_X b, c) - \varphi(a, b, D_X c) = 0,$$

for all $a, b, c \in \Gamma(A)$. This last condition is the same as $D^*(\varphi) = 0$ when $\varphi \in \Gamma_l(\wedge^3 A^*)$ (see (2.10)). Similarly, (CN4) holds for sections of type $\sigma_1 = (0, \alpha)$, $\sigma_2 = (0, \beta)$ if and only if \mathcal{D}^* satisfies (IM4) and $D(\chi) = 0$.

For sections of type $\sigma_1 = (0, \alpha)$, $\sigma_2 = (b, 0)$, (CN4) is equivalent to

$$(6.7) \quad D_X(\mathcal{L}_\alpha b) = \mathcal{L}_\alpha(D_X b) + i_{D_X^*(\alpha)} d_{A^*} b - D_{[\rho^*(\alpha), X]} b - (\rho^*)^* i_X d \langle D_{(\cdot)}^* \alpha, b \rangle,$$

$$(6.8) \quad D_X^*(i_b d_{A^*} \alpha) = i_{D_X(b)} d_{A^*} \alpha + \mathcal{L}_b D_X^*(\alpha) - D_{[\rho(b), X]}^* \alpha + \rho^* i_X d \langle D_{(\cdot)}^* \alpha, b \rangle.$$

These equations follows directly from (2.13) for D and D^* and $m = 1$, which is equivalent to (IM4) under the assumption that (IM1), (IM2) and (IM3) hold. Indeed, first notice that using (2.8) and $\rho^* D_X^* (d \langle \alpha, b \rangle) = D_X(d_{A^*} \langle \alpha, b \rangle)$ together with Cartan formula $\mathcal{L}_\alpha = i_\alpha d_{A^*} + d_{A^*} i_\alpha$, one can see that both equations are exactly the same under the change $\alpha \leftrightarrow b$. Now, using that

$$\begin{aligned} \langle D_X^*(i_b d_{A^*} \alpha) - i_{D_X(b)} d_{A^*} \alpha, a \rangle &= -D_X^*(d_{A^*} \alpha)(a, b) \\ \langle \mathcal{L}_b D_X^*(\alpha) + \rho^* i_X d \langle D_{(\cdot)}^* \alpha, b \rangle, a \rangle &= -d_{A^*} D_X^*(\alpha)(a, b) + \mathcal{L}_X \langle D_{\rho(a)}^*(\alpha), b \rangle + \langle D_{[\rho(a), X]}^*(\alpha), b \rangle, \end{aligned}$$

one can check that (6.8) is exactly (2.13) in degree 1.

The situation for sections of type $\sigma_1 = (a, 0)$, $\sigma_2 = (0, \beta)$ is entirely analogous. This proves claim 2 and concludes the proof of the first part of the theorem.

The assertion about the Nijenhuis condition follows from the decompositions $\mathbb{D} = (D, D^*)$ and $\ell = (l, l^*)$, and the fact that D is Nijenhuis (resp. Dolbeault) if and only is so is D^* , see [9, Thm. 2.11]. \square

When $\mathcal{D} = (\mathbb{D}, \ell, r)$ is a Courant 1-derivation, there is also a compatibility with the bivector field $\pi \in \mathfrak{X}^2(M)$ in (6.3), defined by the lagrangian splitting $E = A \oplus A^*$, see [9, Prop. 4.6 (i)].

Corollary 6.2. *The pair (π, r) is compatible in the sense of (3.3). In particular, if the lagrangian splitting is by Dirac structures and \mathcal{D} is Nijenhuis, then (π, r) defines a Poisson-Nijenhuis structure.*

Proof. Using that $\pi^\sharp = \rho_* \circ \rho^*$, $\mathbb{D} = (D, D^*)$ and $\ell = (l, l^*)$, the first condition in (3.3) follows from condition (IM1) for \mathcal{D} and \mathcal{D}^* , while the second condition in (3.3) follows from (IM2) for \mathcal{D} and \mathcal{D}^* . \square

Recall from [9] that a *Lie-Nijenhuis bialgebroid* is a triple (A, A^*, \mathcal{D}) , where (A, A^*) is a Lie bialgebroid and \mathcal{D} is a Nijenhuis 1-derivation on A such that \mathcal{D} is compatible with the Lie algebroid structure on A , and \mathcal{D}^* is compatible with the Lie algebroid structure on A^* (in

the sense of Def. 2.8). These are the infinitesimal objects corresponding to Poisson-Nijenhuis groupoids, see [9, § 4.3].

By Theorem 6.1 we have the following enhancement of the known correspondence between Lie bialgebroids and Manin triples.

Corollary 6.3. *Lie-Nijenhuis bialgebroids are equivalent to Courant-Nijenhuis algebroids equipped with a splitting by Dirac-Nijenhuis structures.*

When \mathcal{D} is a Dolbeault 1-derivation, Theorem 6.1 gives the known correspondence between Lie (quasi-)bialgebroids and Manin (quasi-)triples in the holomorphic category.

REFERENCES

- [1] P. Antunes, C. Laurent-Gengoux, J. Nunes da Costa. Hierarchies and compatibility on Courant algebroid. *Pac. J. Math.* **261** (2013) 1-32.
- [2] P. Antunes, J. Nunes da Costa. Nijenhuis and Compatible Tensors on Lie and Courant algebroids. *J. Geom. Phys.* **65** (2013) 66-79.
- [3] M. Boumaiza, N. Zaalani. Relèvement d'une algébroïde de Courant. *C. R. Math. Acad. Sci. Paris*, **347** (2009), no. 3-4, 177–182.
- [4] H. Bursztyn and T. Drummond. Lie theory of multiplicative tensors. *Math. Ann.* **375** (2019), 1489–1554.
- [5] H. Bursztyn, T. Drummond, and C. Netto. Dirac structures and Nijenhuis operators. *Math. Z.* **302** (2022), 875–915.
- [6] H. Bursztyn, D. Iglesias-Ponte, P. Severa. Courant morphisms and moment maps. *Math. Res. Lett.* **16** (2009), no. 2, 215–232.
- [7] J. F. Carinena, J. Grabowski, G. Marmo. Courant algebroid and Lie bialgebroid contractions. *J. Phys.* **37** (2004), 5189–5202.
- [8] T. Courant. Dirac manifolds. *Trans. Amer. Math. Soc.* **319** (1990), 631–661.
- [9] T. Drummond. Lie-Nijenhuis bialgebroids. *Q. J. Math.* **73** (2022), 849–883.
- [10] J. Grabowski, Courant-Nijenhuis tensors and generalized geometries. in Groups, geometry and physics, Real Acad. Ci. Exact. Fis. Quim. Nat. Zaragoza. **29** (2006), 101–112.
- [11] J. Grabowski, P. Urbanski. Lie algebroids and Poisson-Nijenhuis structures. *Rep. Math. Phys.* **40** (1997), 195–208.
- [12] M. Gualtieri. Generalized complex geometry. *Ann. of Math.* **174** (2011), 75–123.
- [13] M. Gualtieri. Generalized Kähler Geometry. *Commun. Math. Phys.* **331** (2014), 297–331.
- [14] Y. Kosmann-Schwarzbach. Quasi, twisted, and all that... in Poisson geometry and Lie algebroid theory. The Breadth of Symplectic and Poisson Geometry. *Progr. Math.*, **232** (2005), 363–389.
- [15] Y. Kosmann-Schwarzbach. Nijenhuis structures on Courant algebroids, *Bull. Brazilian Math. Soc.* **42** (4) (2011), 625–649.
- [16] Y. Kosmann-Schwarzbach and F. Magri. Poisson-Nijenhuis structures. *Ann. Inst. H. Poincaré Phys. Théor.* **53** (1990), 35–81.
- [17] C. Laurent-Gengoux, M. Stiénon, P. Xu. Holomorphic Poisson Manifolds and Holomorphic Lie Algebroids, *Int. Math. Res. Not.* (2008), Art.ID rnn 088, 46 pp.
- [18] D. Li-Bland, E. Meinrenken. Courant algebroids and Poisson geometry. *Int. Math. Res. Not.* **11** (2009), 2106–2145.
- [19] Z. Liu, A. Weinstein, P. Xu. Manin triples for Lie bialgebroids. *J. Differential Geom.* **45** (1997), 547–574
- [20] K. Mackenzie, *General theory of Lie groupoids and Lie algebroids*. London Mathematical Society Lecture Note Series, **213**. Cambridge University Press, Cambridge, (2005).
- [21] F. Magri, C. Morosi. A geometrical characterization of integrable Hamiltonian systems through the theory of Poisson-Nijenhuis manifolds. *Quaderno, University of Milan*, **19** (1984).
- [22] J. Rawnsley. Flat partial connections and holomorphic structures in C^∞ vector bundles, *Proc. Amer. Math. Soc.* **73** (1979), 391–397.
- [23] D. Roytenberg. On the structure of graded symplectic supermanifolds and Courant algebroids. *Quantization, Poisson brackets and beyond (Manchester, 2001)*, 169–185, *Contemp. Math.*, **315**, Amer. Math. Soc., Providence, RI, 2002.
- [24] D. Roytenberg. Quasi-Lie Bialgebroids and Twisted Poisson Manifolds. *Lett. Math. Phys.* **61** (2022), 123–137.
- [25] P. Severa. Letters to Alan Weinstein about Courant algebroids. *arXiv:1707.00265*.
- [26] P. Severa, A. Weinstein. Poisson geometry with a 3-form background. Noncommutative geometry and string theory (Yokohama, 2001) *Progr. Theoret. Phys. Suppl.* **144** (2001), 145–154.

- [27] M. Stiénon, P. Xu. Poisson Quasi-Nijenhuis Manifolds. *Commun. Math. Phys.* **270** (2007), 709—725.

IMPA, ESTRADA DONA CASTORINA 110, RIO DE JANEIRO, 22460-320, BRAZIL.
Email address: henrique@impa.br

DEPARTAMENTO DE MATEMÁTICA, INSTITUTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO RIO DE JANEIRO,
CAIXA POSTAL 68530, RIO DE JANEIRO, RJ, 21941-909, BRASIL.
Email address: drummond@im.ufrj.br

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, RUA DO MATÃO 1010, CIDADE
UNIVERSITÁRIA, 05508-090 SÃO PAULO, BRASIL.
Email address: cnetto@ime.usp.br