# Discovery of a new lunar mineral rich in water and ammonium in lunar soils returned by Chang'e-5 mission

Authors: Shifeng Jin<sup>1,2</sup><sup>†\*</sup>, Munan Hao<sup>1,2</sup><sup>†</sup>, Zhongnan Guo<sup>3</sup>, Bohao Yin<sup>4</sup>, Yuxin Ma<sup>1</sup>, Lijun Deng<sup>5</sup>, Xu Chen<sup>1</sup>, Yanpeng Song<sup>1</sup>, Cheng Cao<sup>1,2</sup>, Congcong Chai<sup>1,2</sup>, Yunqi Ma<sup>6</sup>, Jiangang Guo<sup>1,7</sup>, Xiaolong Chen<sup>1,2,7\*</sup>

## Affiliations:

<sup>1</sup>Beijing National Laboratory for Condensed Matter Physics and Institute of Physics, Chinese Academy of Sciences; Beijing 100190, China.

<sup>2</sup>School of Physical Sciences, University of Chinese Academy of Sciences, Beijing 100190, China.

<sup>3</sup>School of Chemistry and Biological Engineering, University of Science & Technology Beijing, Beijing 100083, China.

<sup>4</sup>School of Earth System Science, Tianjin University, Tianjin 300072, China

<sup>5</sup>Center of Advanced Analysis & Gene Sequencing, Zhengzhou University, Zhengzhou 450001, China

<sup>6</sup>Qinghai Institute of Salt Lakes, Chinese Academy of Sciences, Xining 810008, China

<sup>7</sup>Songshan Lake Materials Laboratory, Dongguan 523808, China

\*Corresponding author. Email: shifengjin@iphy.ac.cn, chenx29@iphy.ac.cn

<sup>†</sup> These authors contributed equally to this work.

## ABSTRACT

The origin and distribution of lunar water are among the most important issues in understanding the earth-moon system. After more than half a century of laboratory research and remote detection, only hydroxyl contained minerals and lunar ice (H<sub>2</sub>O) are identified. Here we report the discovery of a hydrous mineral (NH<sub>4</sub>)MgCl<sub>3</sub>·6H<sub>2</sub>O in the lunar soil returned by Chang'e-5 mission, which contains ~417,000 parts per million H<sub>2</sub>O. The determined structure and composition are similar to novograblenovite—a terrestrial fumarole mineral formed by reaction of hot basalt in water-rich volcanic gases, whereas the measured isotope composition  $\delta^{37}$ Cl reached 20.4 ‰, a high value that only found in lunar minerals. We rule out the possibility that this hydrate originated from terrestrial contaminants or rocket exhaust through analysis of its chemical, isotopic compositions as well as the formation conditions. Our finding indicates that water can exist on some parts of the sunlit Moon in the form of hydrate compounds. Moreover, this hydrate is rich in ammonium, providing new information in understanding the origin of the Moon.

Keywords: Lunar water, lunar ammonium, lunar volcanism, lunar hydrate, Chang'e-5

## **INTRODUCTION**

The Earth and Moon are nonidentical twins that were presumably born in the same giant impact (1). On the Earth, water plays a crucial role in planetary evolution through the influence on melting (2), viscosity (3), magma crystallization (4) and volcanic eruption (5). However, the Moon was long thought to be anhydrous based on analyses of lunar samples returned by the Apollo and Luna missions (6-7), which places strict constraints on the Moon-forming giant impact (8), lunar magma ocean (9) and subsequent lunar volcanic eruptions (10). With advances in microanalysis techniques over the past decade, the notion of a "dry Moon" is challenged by the detection of small amount of 'water' (hydroxyl, OH<sup>-</sup>) in various lunar samples (11,12), including pyroclastic glass beads (13), melt inclusions (14-15), apatite (16,17), and anorthosite (18,19). The estimated water abundances for the mantle source regions span a wide range, from 0.3 to 1410 ppm (20,21), with the highest values comparable to the mid-oceanic ridge basalts (MORB) mantle on the Earth (14). Meanwhile, several remote sensing and direct impact missions have confirmed the existence of water molecules (H<sub>2</sub>O) in some permanently shadowed carters near lunar poles (22-28). The large number of water molecules (5.6 % wt.) detected is commonly believed to be water-ice (29-31) that results from exogenous sources (11), including solar wind interaction with the lunar regolith (22-24,32,33) and impact delivery of volatiles (28,34). Given the low oxygen fugacity f(O<sub>2</sub>) on the Moon, degassing of  $H_2O$  vapor in lunar magma is considered unfavorable (35), and it is unclear whether the gases-emitting volcanoes ever contributed to the lunar hydrosphere (36). Moreover, the recent confirmed presence of H<sub>2</sub>O on the high-latitudes sunlit region suggests lunar water may exist in a form other than water-ice to survive the thermal attack (37). Due to the lack of returned lunar samples from high-latitude and polar regions, neither the origin nor the chemical form of lunar water (H<sub>2</sub>O) have been determined.

#### RESULTS

The Chang'e-5 (CE5) mission successfully returned 1.731 kg of lunar soil samples from northern Oceanus Procellarum at 43.058 °N (*38,39*), the landing site is much higher in latitude than the Apollo and Luna missions. The CE5 soil samples have been confirmed as mare basalt erupted about 2.0 Ga ago (*40,41*). Both in-situ reflectance spectra and examination of the returned lunar regolith revealed the presence of trace water (OH<sup>-</sup>) (*42-44*), which is mostly attributed to solar wind implantation (*42*) or hydroxyl-containing apatite (*44*). We singled out more than 1000 mineral clasts (larger than 50 µm) from Sample CE5C0400 (1.5 g, Fig. 1A) by aid of an optic

microscope for Raman measurement, crystallographic study and chemical analyses. Pyroxene, anorthite, olivine, ilmenite and glassy bead as the major minerals along with a small amount of silica and spinel are ascertained. Noticeably, a prismatic, plate-like transparent crystal is identified to be an unknown lunar mineral (ULM-1) with a size of about 160×109×73 µm (Fig. 1B).

The chemical compositions of ULM-1 were analyzed on fragments of the crystal using both SEM energy dispersive spectroscopy (EDS) and Electron probe microanalysis (EPMA). As shown in Fig. 1C and S1, the  $K_{\alpha}$  lines of the EDS spectrum suggest the mineral contain Mg and abundant volatile elements N, Cl and O, see the measurement details and the choice of standards in Support Information. The analytical results are listed in table S1, revealing the ratios of major elements is close to N:Mg:Cl:O =1:1:3:6. The concentration of N, Cl, and O elements decreases rapidly in comparison to Mg when the crystal is measured repeatedly, indicating the crystal is decomposing under electron beam radiation. More detailed EDS measurement revealed the existence of a variety of minor species with concentrations about 1% or more, including K, Cs, Na, Si, Al, Ca, S, etc, many of which are also volatile elements. Upon examining different parts of the specimen, it was found that even at the micrometer scale, the spatial distribution of minor species within the crystal is quite inhomogeneous. The measurements suggest the crystallization of the ULM-1 crystal happed in a chemically complex and non-equilibrium environment. Taking the EPMA data into account gives a more accurate chemical compositions close to (N<sub>0.83</sub>Na<sub>0.009</sub>K<sub>0.022</sub>Cs<sub>0.012</sub>) (MgCa<sub>0.023</sub>Si<sub>0.034</sub>Al<sub>0.007</sub>)Cl<sub>2.98</sub>O<sub>6.17</sub> for ULM-1.

The presence of water and ammonium molecule in ULM-1 crystal are confirmed in both the Raman and Infrared (IR) spectra. The strong normal vibrational modes at 3430 cm<sup>-1</sup> (symmetric stretching  $v_1$ ) and at 1645 cm<sup>-1</sup> (bending O–H deformations  $v_2$ ) of H<sub>2</sub>O molecule (45) are clearly observed in the Raman spectrum of ULM-1 as shown in Fig. 1D. Normal modes of the NH<sup>4+</sup> ion at 3080 cm<sup>-1</sup> and 1415 cm<sup>-1</sup> are due to its  $v_1$  in phase stretching and  $v_4$  triply degenerate bending (46), respectively. The band at 3250 cm<sup>-1</sup> corresponds to NH<sup>4+</sup> stretching vibrations (47). The infrared spectrum of ULM-1 shows clearly the major modes for water molecule and NH<sup>4+</sup>, see Fig. 1E. The IR absorption at 1650 cm<sup>-1</sup> can be attributed to the bending mode of water molecule, which is recently observed in the sunlit moon by SOFIA remote sensing (37). Overall, the Raman and IR spectra as well as the determined chemical compositions on ULM-1 are very consistent to the observations in novograblenovite (NH<sub>4</sub>,K)MgCl<sub>3</sub>·6H<sub>2</sub>O (48,49), an earth mineral recently

found on basaltic lava from the 2012–2013 Tolbachik fissure eruption at the Plosky Tolbachik volcano (48).

The crystal structure of ULM-1 was determined by X-ray single crystal diffraction data (Table S2). As shown in Fig. 2A&B, the mineral crystallizes in monoclinic space group C2/c, with lattice parameters a=9.3222(10) Å, b=9.5731(10) Å, c= 13.3328(12) Å and  $\beta$ =90.101(5) °, Z=4. Notably, the six independent H atoms on the water molecules can be clearly revealed in difference-Fourier maps, see Fig. 2c. Meanwhile, the H atoms in ammonium groups were also revealed and found to be disordered. Subsequent structure refinements revealed nearly full occupancy of the major elements N, Mg, Cl and O, and the trace elements determined by EPMA were complemented to the structure during the last refinements. The final structure parameters and the selected bond lengths and angles are shown in Table S2-5, respectively. The chemical formula for the ULM-1 crystal can then be best represented as [(NH<sub>4</sub>)<sub>0.87</sub>Na<sub>0.009</sub>K<sub>0.021</sub>Cs<sub>0.012</sub>][Mg<sub>0.97</sub>Ca<sub>0.023</sub>Al<sub>0.007</sub>Si<sub>0.03</sub>] Cl<sub>3</sub>·6H<sub>2</sub>O according to the results of chemical analysis, structure refinement and the principle of charge neutrality.

In the determined crystal structure, each  $Mg^{2+}$  ion is coordinated by 6 O from water molecules to form a nearly regular octahedron  $[Mg(H_2O)_6]^{2+}$ ; each  $(NH_4)^+$  is connected by 6 Cl<sup>-</sup> anions through H-Cl hydrogen bonds to form corner shared octahedrons in a 3D framework (Fig. 2B). Therefore, the crystal structure of ULM-1 is topologically similar to that of perovskite *ABO*<sub>3</sub>, where  $[Mg(H_2O)_6]^{2+}$  serves as *A*,  $(NH_4)^+$  as *B* and Cl<sup>-</sup> as *O*. As shown in Fig. 2A, the main units of *A*, *B* and *O* in ULM-1 are inter-connected entirely by hydrogen bonds (short dotted line). Each chlorine atom accommodates six hydrogen bonds: four from O atoms at the vertices of Mg(H\_2O)\_6 octahedra and two from the ammonium ions. The O–H····Cl hydrogen bonds lie in the interval 2.325–2.412 Å, while N–H····Cl bonds are normally longer from 2.328 to 2.883 Å (table S5).

## DISCUSSION

Our chemical and structural analyses suggest that ULM-1 be an extremely water (41.7 % wt.) and ammonia (6.6 % wt.) rich mineral in CE5 lunar samples. The presence of lunar mineral hydrate is unexpected since previous studies show a main feature of lunar mineralogy is the total lack of minerals that contain water, such as clays, micas, and amphiboles (35). This hydrate, isolated from CE5 lunar soils, may come from three possible sources. First, it has an ingenious lunar origin as a rare fumarole mineral. Second, it is simply a contamination from terrestrial sources. Third, a

chemical reaction product from rocket exhaust and the lunar soils during the landing. We argue, based on chemical, structural and isotopic analysis, that this unusual CE5 hydrate crystal should have an ingenious lunar origin.

As well known, rocks from the Moon and Earth have very similar isotopic ratios for the most abundant elements, one of the most significant exceptions is lunar chlorine (50,51). Lunar  $\delta^{37}$ Cl values [defined as <sup>37</sup>Cl/<sup>35</sup>Cl relative to standard mean ocean chlorine] are determined to be in a range from 0% to +81% (52,53), far more spread and higher than observed in any other planetary bodies in solar system (50-52). The terrestrial  $\delta^{37}$ Cl values, for instance, only cluster around 0 ‰ with a typical spread of  $\pm 1.0 \%$  (54). So do for meteorites (55), comets (56) and Mars (51), see Fig. 3. To characterize the isotropic composition of ULM-1 crystal, a terrestrial standard is required. We synthesis the single crystal samples of (NH<sub>4</sub>)MgCl<sub>3</sub>·6H<sub>2</sub>O from aqueous solution based on methods descripted in Ref (70). The chemical reagents NH<sub>4</sub>Cl, MgCl<sub>2</sub>·6H<sub>2</sub>O were purchased from Sinopharm Co., Ltd. and used as received. The Cl isotopic ratios in the synthetic (NH<sub>4</sub>)MgCl<sub>3</sub>·6H<sub>2</sub>O standard is carefully measured based on high accuracy solid thermal ionization mass spectrometer, and the  $\delta^{37}$ Cl value is determined to be +0.08 ‰. Relative to such a small  $\delta^{37}$ Cl value in terrestrial sample, the measured  $\delta^{37}$ Cl value of ULM-1 is exceptionally high, up to +20.4 ‰ on average, see Fig. 3 and table S6 for measurement details. As shown in Figure 3, such a large  $\delta^{37}$ Cl value in ULM-1 indicates that this crystal couldn't be sourced from terrestrial environment (71), instead, this extremely high  $\delta^{37}$ Cl value well fall into the range of lunar minerals (72).

The determined structure of ULM-1 is isostructural to a terrestrial mineral novograblenovite,  $(K,NH_4)MgCl_3 \cdot 6H_2O$  (48,49), which shed light on the origin of this hydrate. On the Earth, novograblenovite is a rare fumarole mineral that was discovered in Tolbacik volcano only recently. In-depth analysis suggests this mineral should be formed by a reaction of basalt with hot volcanic gases enriched in H<sub>2</sub>O, HCl and NH<sub>3</sub> (48). Meanwhile, novograblenovite was also found as a sublimate around vents of hot gases in a few burning coal-dumps, where the crystallization temperature is slightly above 100 °C (49). Notably, the latter samples without a basaltic origin is free from K and has a simpler formula (NH<sub>4</sub>)MgCl<sub>3</sub>  $\cdot 6H_2O$  (49). Compared to the known terrestrial mineral, ULM-1 has much complicated chemical compositions, [(NH<sub>4</sub>)<sub>0.87</sub>Na<sub>0.009</sub>K<sub>0.021</sub>Cs<sub>0.012</sub>] [Mg<sub>0.97</sub>Ca<sub>0.023</sub>Al<sub>0.007</sub>Si<sub>0.03</sub>] Cl<sub>3</sub>  $\cdot 6H_2O$ . Occasionally, S is also detected on some parts of the crystal by EDS analysis. The presents of Ca, Al and Si along with various alkali metals in ULM-1 crystal,

is consist with the source of nature mineral formed by exposure of the host basalt to eruptive gas exhalations, ruling out the possibility of contamination in lab handling the lunar soils.

The oxyhydration of lunar soil in air atmosphere is well-known since the compound FeOOH, arising from the water vapor contamination of ingenious FeCl<sub>2</sub>, was found in the form of "rust" in lunar rocks from Apollo mission (*36*). However, the possibility of air contact that leads to the formation of water rich ULM-1 crystal is highly unlikely. The formation of ULM-1 can essentially be described by the following chemical equations,

$$6H_2O(g) + NH_3(g) + HCl(g) + MgCl_2(s) = (NH_4)MgCl_3 \cdot 6H_2O(s),$$
 (eq. 1)  
Or

 $6H_2O(l) + NH_4Cl(s) + MgCl_2(s) = (NH_4)MgCl_3 \cdot 6H_2O(s),$  (eq. 2)

In either case, the formation of ULM-1 needs abundant ammonia species (NH<sub>3</sub> or NH<sub>4</sub>Cl), which is absent in air. The high temperature above 100 °C in vapor phase reaction (eq. 1) or near room temperature aqueous condition (eq. 2) required for the crystallization of ULM-1 further help to rule out the air contact as the source (48, 49).

The formation condition of ULM-1 in hot vapor phases also lead to the possibility of rocket exhaust as the source. There are several reasons that the rocket exhaust scenario is highly unlikely. First, investigations on the lunar surface from Chang'E-5 in-situ spectra show hydroxyl contents on the weak end of lunar hydration features, no higher than the predictions from previous remote sensing (*44*). Additionally, the highest water content in the landing site is observed in a CE5-Rock that with potential basaltic origin, rather than the lunar regolith that are more subjective to rocket exhaust (*42*). Second, the formation of ULM-1 crystal on lunar surface requires the reaction of hot Mg-rich basalt with eruptive gas exhalations that enriched in HCl and NH<sub>3</sub>. However, both the eruptive HCl and NH<sub>3</sub> are exceedingly rare in rocket exhaust plume, regardless of the different fuel used in CE5 main rocket or the lander, e.g., liquid H<sub>2</sub>/O<sub>2</sub> and UDMH/N<sub>2</sub>O<sub>4</sub>. Third, the extraordinary high Cl isotopic composition of ULM-1 cannot be explained by a contamination of terrestrial sourced rocket exhaust as well.

Having shown that ULM-1 should have an ingenious lunar origin, it's important to analyze the stability of this hydrate crystal under the harsh lunar environment around the CE5 landing site. It's generally accepted that in the permanently shadowed carters, molecular water could be preserved in the form of ice at low enough temperatures. However, in the sunlit moon, where the lunar

condition has become too harsh for ice to survive, the actual chemical form of lunar water (H<sub>2</sub>O) is still unknown. CE5 spaceship landed at a relatively high latitude 43.058°, where the highest surface temperature in a lunar year is about 80 °C according to the Diviner Dataset on lunar surface temperatures (*57*), see fig. S5. Moreover, the thermal conductivity of lunar regolith is very low, so that the solar radiation only heats up the top layer of a few mm thickness (fig. S6). The lunar surface temperature decreases quickly, insulating deeper soil from heat attack (*36*). Compared to water ice, the decomposition temperature of novograblenovite is reported to be 90 ~ 150 °C higher (*49,58*). The increased thermally stability of this hydrate should be crucial to protect the lunar water (H<sub>2</sub>O) from the harsh sunlit lunar environment for billions of years. The existence of lunar hydrated water can naturally explain the water molecule signal recently detected in the sunlit moon (*37*), as well as the enhanced water (OH<sup>-</sup> or H<sub>2</sub>O) signals in some pyroclastic deposits (*62*).

Although the returned lunar basalt is generally highly depleted in water (35), the formation condition of this lunar hexahydrate suggests during eruption, some lunar fumaroles have exsolved considerable amount of water and hydrogen vapors, along with other H-bearing species including NH<sub>3</sub> and HCl. We performed the thermodynamic calculations of equilibrium constants in a temperature range from 50°C to 150°C (detailed in Methods and Supplementary materials), allowing us to constraint the fugacity of gaseous water  $f(H_2O)$  in the lunar volcanic fume. On the earth, novograblenovite was found to crystallize from vapors at temperatures slightly higher than 100°C (49). We note the maximum sum of the fugacity of NH<sub>3</sub> and HCl is further thermodynamically constrained by the chemical reaction:

$$NH_3 (g) + HCl (g) = NH_4Cl (s)$$
 (eq.3)

Thus the temperature dependent low-limit  $f(H_2O)$  to form ULM-1 can be obtained and delineated by the red curve in Fig. 4. Below this equilibrium line, the hexahydrate becomes unstable and gives way to a known dehydrate -  $(NH_4)MgCl_3 \cdot 2H_2O(59)$ . The upper-limit of  $f(H_2O)$  is set by the equilibrium between the liquid water and its saturated vapor (blue line). In this temperature range, the low-limit of  $f(H_2O)$  of lunar volcanic gas is constrained to be  $3.5\%\sim16.6\%$  of the saturated water vapor. With reference to the formation temperature of terrestrial novograblenovite (>100°C) (49), the  $f(H_2O)$  of the volcanic gas should be greater than 7.7% atm. As a comparison, this low-limit is around 3 times lower than the  $f(H_2O)$  of the driest Earth volcano – Lengai volcano (>24% atm) (60). It worth noting that in Tolbacik volcano, where the earth mineral is located, the measured  $f(H_2O)$  is higher than 0.92 atm (pentagon) (5), one lg unit higher

than the thermodynamic equilibrium value. Oversaturated vapor could be necessary to overcome the kinetic barriers during the crystal growth of this hydrous mineral. Moreover, by taking H<sub>2</sub> degassing into account, the estimated  $f(H_2O + H_2)$  values (purple doted line) become comparable to some 'hot spot' volcanoes on earth (detailed in Supplementary materials).

The existence of lunar volcanic gases rich in  $H_2O$  and  $H_2$  vapors also has implications for the origin of the volatiles detected in permanently shadowed carters. The remote sensing missions have shown that in addition to  $H_2O$ , there is significant amount of solid  $H_2$  and  $NH_3$  in the lunar polar regions (28,63), which is within the lunar volcanic gas inventory revealed here. It is conceivable that some of these volatiles, including water ( $H_2O$ ), could have originated from magmatic degassing during eruption of ancient lunar volcanos.

#### SUMMARY

After more than half a century of lunar sampling missions and laboratory research, molecular water (H<sub>2</sub>O) is firstly discovered in lunar soil in the form of a well-crystallized hydrate. The determined structure and composition of this hydrate is similar to a rare terrestrial fumarole mineral novograblenovite, suggesting the lunar mineral [(NH<sub>4</sub>)<sub>0.87</sub>Na<sub>0.009</sub>K<sub>0.021</sub>Cs<sub>0.012</sub>] [Mg<sub>0.97</sub>Ca<sub>0.023</sub>Al<sub>0.007</sub>Si<sub>0.03</sub>] Cl<sub>3</sub>·6H<sub>2</sub>O is formed by reaction of basaltic lava in H<sub>2</sub>O, HCl and NH<sub>3</sub> rich lunar volcanic gases. While lunar water has been commonly attributed to solar wind, cometary and meteoritic sources, the extremely high  $\delta^{37}$ Cl value of this hydrate suggest the lunar mineral itself and the molecular water in it are ingenious to the moon. The increased thermally stability of this hexahydrate compared with water-ice and the high latitude of its location may help to preserve the molecular water on the sunlit moon. This hydrate constraints the lowest water fugacity f(H<sub>2</sub>O) of lunar volcanic gases during its crystallization, the resulting f(H<sub>2</sub>O+H<sub>2</sub>) is on the low-end of some the driest volcanoes on Earth.

#### REFERENCES

- 1. Canup RM and Asphaug E. Origin of the Moon in a giant impact near the end of the Earth's formation. *Nature* 2001; **412**: 708–712.
- 2. Gaetani GA and Grove TL. The influence of water on melting of mantle peridotite. *Contrib Mineral Petrol* 1998; **131**: 323–346.
- 3. Hirth G, Kohlstedt D. Water in the oceanic upper mantle: implications for rheology, melt extraction and the evolution of the lithosphere. *Earth Planet Sci Lett* 1996; **144**: 93–108.
- 4. Asimow PD and Langmuir CH. The importance of water to oceanic mantle melting regimes. *Nature* 2003; **421**: 815–820.

- 5. Haraldur Sigurdsson, Ed. *Encyclopedia of volcanoes*, (Academic Press, New York, ed. 1, 1999).
- 6. Lunar Science Preliminary Examination Team. Preliminary examination of lunar samples from Apollo 12. *Science* 1970; **167**: 1325–1339.
- 7. Lucey P. "New Views of the Moon" in Reviews in Mineralogy and Geochemistry (Mineralogical Society of America, Chantilly, Virginia, 2006) pp. 83–219.
- 8. Albarède F. Volatile accretion history of the terrestrial planets and dynamic implications. *Nature* 2009; **461**: 1227–1233.
- Warren PH. The magma ocean concept and lunar evolution. *Annu Rev Earth Planet Sci* 1985; 13: 201–240.
- 10. Shearer CK, Hess PC and Wieczorek MA *et al*. Thermal and magmatic evolution of the moon. *Rev Mineral Geochem* 2006; **60**: 365–518.
- 11. Lucey PG. A Lunar Waterworld. Science 2010; **326**: 531-532.
- 12. Anand M. Lunar water: a brief review. Earth Moon Planets 2010; 107: 65-73.
- 13. Saal AE, Hauri EH and Cascio ML *et al.* Volatile content of lunar volcanic glasses and the presence of water in the Moon's interior. *Nature* 2008; **454**: 192–195.
- 14. Hauri EH, Weinreich T and Saal AE *et al*. High pre-eruptive water contents preserved in lunar melt inclusions. *Science* 2011; **333**: 213-215.
- 15. Ni P, Zhang YX and Chen S *et al*. A melt inclusion study on volatile abundances in the lunar mantle. *Geochim Cosmochim Acta* 2019; **249**: 17–41.
- 16. Boyce JW, Liu Y and Rossman GR *et al*. Lunar apatite with terrestrial volatile abundances. *Nature* 2010; **466**: 466–469.
- 17. Boyce JW, Tomlinson SM and McCubbin FM *et al*. The Lunar Apatite Paradox. *Science* 2014; **344**: 400-402.
- 18. Hui HJ, Peslier AH and Zhang YX *et al*. Water in lunar anorthosites and evidence for a wet early Moon. *Nat Geosci* 2013; **6**: 177–180.
- 19. Hui HJ, Guan YB and Chen Y *et al*. A heterogeneous lunar interior for hydrogen isotopes as revealed by the lunar highlands samples. *Earth Planet Sci Lett* 2017; **473**: 14–23.
- 20. Sharp ZD, Shearer CK and McKeegan KD *et al*. The chlorine isotope composition of the moon and implications for an anhydrous mantle. *Science* 2010; **329**: 1050-1053.
- McCubbin FM, Vander KE and Tartèse R *et al.* Magmatic volatiles (H, C, N, F, S, Cl) in the lunar mantle, crust, and regolith: abundances, distributions, processes, and reservoirs. *Am Mineral* 2015; **100**: 1668–1707.
- 22. Pieters CM, Goswami JN and Clark RN *et al*. Character and spatial distribution of OH/H<sub>2</sub>O on the surface of the moon seen by M3 on Chandrayaan-1. *Science* 2009; **326**: 568–572.
- 23. Sunshine JM, Farnham TL, Feaga LM *et al*. Temporal and spatial variability of lunar hydration as observed by the Deep Impact spacecraft. *Science* 2009; **326**: 565–568.
- 24. Clark RN. Detection of adsorbed water and hydroxyl on the moon. *Science* 2009; **326**: 562–564.
- 25. Harmon JK and Slade MA. Radar mapping of Mercury: Full disk images and polar anomalies. *Science* 1992; **258**: 640–643 (1992).
- 26. Feldman WC, Maurice S and Binder AB. Fluxes of fast and epithermal neutrons from lunar prospector: evidence for water ice at the lunar poles. *Science* 1998; **281**: 1496-1500.

- 27. Margot JL, Campbell DB and Jurgens RF *et al.* Topography of the lunar poles from radar interferometry: a survey of cold trap locations. *Science* 1999; **284**: 1658-1660.
- 28. Colaprete A, Schultz P and Heldmann J *et al*. Detection of Water in the LCROSS Ejecta Plume. *Science* 2010; **330**; 463-467.
- 29. Arnold JR. Ice at the lunar poles. J Geophys Res 1979; 84: 5659–5668.
- 30. Slade MA, Butler BJ and Muhleman DO. Mercury radar imaging: Evidence for polar ice. *Science* 1992; **258**: 635–640.
- 31. Campbell DB, Campbell BA and Carter LM *et al*. No evidence for thick deposits of ice at the lunar south pole. *Nature* 2006; **443**: 835–837.
- 32. Liu Y, Guan YB and Zhang YX *et al.* Direct measurement of hydroxyl in the lunar regolith and the origin of lunar surface water. *Nat Geosci* 2012; **5**: 779–782.
- 33. Chaussidon M. Lunar water from the solar wind. Nat Geosci 2012; 5: 766-767.
- 34. Greenwood JP, Itoh S and Sakamoto N *et al*. Hydrogen isotope ratios in lunar rocks indicate delivery of cometary water to the Moon. *Nat Geosci* 2011; **4**: 79–82.
- 35. Heiken GH, Vaniman DT and French BM. *The Lunar Source Book*. Cambridge University press, Cambridge, 1991.
- 36. Kerr RA. How Wet the Moon? Just Damp Enough to Be Interesting. *Science* 2010; **330**: 434-434.
- 37. Honniball CI, Lucey PG and Li S *et al*. Molecular water detected on the sunlit Moon by SOFIA. *Nat Astron* 2021; **5**: 121–127.
- Qian YQ, Xiao L and Zhao SY. Geology and scientific significance of the Rümker region in northern Oceanus Procellarum: China's Chang'E-5 landing region. *J Geophys Res* 2018; 123: 1407–1430.
- 39. Qian Y, Xiao L and Yin S *et al*. The regolith properties of the Chang'e-5 landing region and the ground drilling experiments using lunar regolith simulants. *Icarus* 2020; **337**: 113508.
- 40. Che X, Nemchin A and Liu D *et al*. Age and composition of young basalts on the Moon, measured from samples returned by Chang'e-5. *Science* 2021; **374**: 887–890.
- 41. Li QL, Zhou Q, Liu Y *et al.* Two-billion-year-old volcanism on the Moon from Chang'E-5 basalts. *Nature* 2021; **600**: 54–58.
- 42. Lin HL, Li S, Xu R *et al.* Wang, In situ detection of water on the Moon by the Chang'E-5 lander. *Sci Adv* 2022; **8**: eabl9174.
- 43. Hu S, He HC, Ji JL *et al*. A dry lunar mantle reservoir for young mare basalts of Chang'e-5. *Nature* 2021; **600**: 49–53.
- 44. Liu JJ, Liu B, Ren X *et al.* Evidence of water on the lunar surface from Chang'E-5 in-situ spectra and returned samples. *Nat Commun* 2022 **13**, 3119.
- 45. Haas C and Hornig D. Inter-and Intramolecular Potentials and the Spectrum of Ice. *The Journal of Chemical Physics* 1960; **32**: 1763–1769.
- 46. Nakamoto K, *Infrared and Raman Spectra of Inorganic and Coordination Compounds*. John Wiley & Sons, Inc., Hoboken, New Jersey, 2008.
- 47. Majzlan J, Schlicht H and Wierzbicka-Wieczorek M *et al*. A contribution to the crystal chemistry of the voltaite group: solid solutions, Mössbauer and infrared spectra, and anomalous anisotropy. *Mineralogy and Petrology* 2013; **107**: 221–233.
- 48. Okrugin VM, Kudaeva SS and Karimova OV *et al*. The new mineral novograblenovite, (NH<sub>4</sub>,K)MgCl<sub>3</sub>·6H<sub>2</sub>O from the Tolbachik volcano, Kamchatka, Russia: mineral description and crystal structure. *Mineralogical Magazine* 2019; **83**: 223–231.

- 49. Parafiniuk J, Stachowicz M and Woźniak K. Novograblenovite from Radlin, Upper Silesia, Poland and its relation to 'redikortsevite'. *Mineralogical Magazine* 2021; **85**: 132–141.
- 50. Sharp ZD, Shearer CK and McKeegan KD *et al*. The chlorine isotope composition of the Moon and implications for an anhydrous mantle. *Science* 2010; **329**: 1050–1053.
- 51. Barnes JD and Sharp ZD. Chlorine isotope geochemistry. *Reviews in Mineralogy and Geochemistry* 2017; 82: 345.
- 52. Boyce JW, Treiman AH and Guan YB *et al*. The chlorine isotope fingerprint of the lunar magma ocean. *Sci Adv* 2015; 1: e1500380.
- 53. Wang Y, Hsu W and Guan Y. An extremely heavy chlorine reservoir in the Moon: Insights from the apatite in lunar meteorites. *Sci Rep* 2019; **9**: 5727.
- 54. Sharp ZD, Barnes JD and Brearley AJ *et al*. Chlorine isotope homogeneity of the mantle, crust and carbonaceous chondrites. *Nature* 2007; **446**: 1062–1065.
- 55. Bonifacie M, Jendrzejewski N and Agrinier P et al. The chlorine isotope composition of Earth's mantle. Science 2008; **319**: 1518.
- 56. Dhooghe F, Keyser JD and Hänni N *et al*. Chlorine-bearing species and the <sup>37</sup>Cl/<sup>35</sup>Cl isotope ratio in the coma of comet 67P/Churyumov–Gerasimenko. *Monthly Notices of the Royal Astronomical Society* 2021; **508**: 1020-1032.
- 57. Williams JP, Paige DA and Greenhagen BT *et al*. The global surface temperatures of the Moon as measured by the Diviner Lunar Radiometer Experiment. *Icarus* 2017; **283**: 300-325.
- 58. Zolotarev AA, Zhitova ES and Krzhizhanovskaya MG *et al.* Crystal chemistry and high-temperature behaviour of ammonium phases NH<sub>4</sub>MgCl<sub>3</sub>·6H<sub>2</sub>O and (NH<sub>4</sub>)<sub>2</sub>Fe<sup>3+</sup>Cl<sub>5</sub>·H<sub>2</sub>O from the burned dumps of the chelyabinsk coal basin. *Minerals* 2019; **9**: 486.
- 59. Okrugin VM, Kudaeva SS and OV Karimova *et al.* Novograblenovite, IMA 2017-060. CNMNC Newsletter No. 39, October 2017, page 1284. *Mineralogical Magazine* 2017; **81**: 1279–1286.
- 60. Koepenick KW, Brantley SL and Thompson JM *et al.* Volatile emissions from the crater and flank of Oldoinyo Lengai volcano, Tanzania. *Journal of Geophysical Research* 1996; **101**: 13819-13830.
- 61. Sharp ZD, McCubbin FM and Shearer CK. A hydrogen-based oxidation mechanism relevant to planetary formation. *Earth and Planetary Science Letters* 2013; **380**: 88–97.
- 62. Milliken RE and Li S. Remote detection of widespread indigenous water in lunar pyroclastic deposits. *Nat Geosci* 2017 **10**, 561-656.
- 63. Gladstone GR, Hurley DM and Retherford KD *et al.* LRO-LAMP Observations of the LCROSS Impact Plume, *Science* 2010; **330**: 472.
- 64. Wetzel DT, Hauri EH and Saal AE *et al*. Carbon content and degassing history of the lunar volcanic glasses. *Nat Geosci* 2015; **8**: 755–758.
- 65. Dolomanov OV, Bourhis LJ, Gildea RJ et al. OLEX2: a complete structure solution, refinement and analysis program. J Appl Cryst 2009; 42: 339-341.
- 66. Sheldrick GM. SHELXT Integrated space-group and crystal structure determination. *Acta Cryst A* 2015; **71**: 3-8.
- 67. Sheldrick GM. Crystal Structure Refinement with SHELXL. Acta Cryst C 2015; 71: 3-8.
- 68. Solans X, Font-Altaba M, Aguilo M *et al.* Crystal Form and Structure of Ammonium Hexaaquamagnesium Trichloride, NH<sub>4</sub>(Mg(H<sub>2</sub>O)<sub>6</sub>)Cl<sub>3</sub>. *Acta Cryst C* 1983; **39**: 1488-1490.
- 69. Ball MC and Ladner NG. Dehydration of Ammonium Magnesium Chloride Hexahydrate (Ammonium Carnallite), *J Chem Soc, Dalton Trans* 1979; **2**: 330-333.

- 70. Solans X, Font-Altaba M, Aguilo M *et al*. Crystal Form and Structure of Ammonium Hexaaquamagnesium Trichloride, NH<sub>4</sub>(Mg(H<sub>2</sub>O)<sub>6</sub>)Cl<sub>3</sub>. *Acta Cryst C* 1983; **39**: 1488-1490.
- 71. Gargano, A, Sharp, Z. The chlorine isotope composition of iron meteorites: Evidence for the Cl isotope composition of the solar nebula and implications for extensive devolatilization during planet formation. *Meteoritics & Planetary Science* 2019; **54**: 1619-1631.
- 72. Gargano, A, Sharp, Z, Shearer, C, Simon, JI, Halliday, A, Buckley, W. The Cl isotope composition and halogen contents of Apollo-return samples. *Proc Natl Acad Sci. U S A* 2020;**117**: 23418-23425.

## ACKNOWLEDGMENTS

The Chang'e-5 lunar sample CE5C0400YJFM00507 (1.5 g) was provided by the China National Space Administration. We thank Ying Li, Qinghua Zhang, Xuefeng Wang, Junyan Zhou, Tianping Ying from IOP, CAS, Chuanqiang Sun from Tianjin University and Chunlai Li, Bin Liu from National Astronomical Observatories, CAS for their assistance in experiments and useful discussions.

## FUNDING

This work was supported by the Key Research Program of Chinese Academy of Sciences grant ZDBS-SSW-JSC007-2 and the Youth Innovation Promotion Association of CAS grant 2019005.

## **AUTHOR CONTRIBUTIONS**

XLC designed the research and supervised this project, SFJ, MNH, YXM, YPS, CCC, CC and GJG performed physical properties experiments, MNH determined the crystal structure, SFJ, ZNG performed thermodynamic calculations, BHY and YQM measured the isotopic composition, LGD performed the EPMA measurement, XLC and SFJ processed analytical data and wrote the manuscript, all authors contribute to the revision of the manuscript.

#### Conflict of interest statement. None declared.

**Data and materials availability**. The lunar sample, designation CE5C0400YJFM00507 was provided by the China National Space Administration (CNSA) under a materials transfer agreement a half-year loan (with an extension for another half-year), after which they will be returned to CNSA. Readers may request Chang'e-5 samples from CNSA through a standard procedure. Further details are provided in the supplementary materials. Our EDS and EPMA data are provided in data S1 to S4, the thermodynamic results in data S7-19, chemical composition in table S1-5, Cl isotropic data in table S6.



**Figure. 1. Photograph and composition of lunar hydrous mineral ULM-1.** (A) Chang'e-5 soil samples. (B) Photo of ULM-1 single crystal mounted on the top of a cactus thorn. (C) Energy-dispersive X-ray spectroscopy (EDX) spectrum, inset: Enlarged electron probe microanalysis (EPMA) spectra show the presence of trace elements. (D) Raman spectrum of ULM-1. (E) Infrared spectrum of ULM-1.



Figure. 2. Crystal structure and charge density of lunar hydrate ULM-1. (A) Crystal structure of lunar mineral (NH<sub>4</sub>)MgCl<sub>3</sub>·6H<sub>2</sub>O, the dotted lines present a network of hydrogen bond interactions around the H<sub>2</sub>O molecules and NH<sup>4+</sup> ions. (B) The mineral crystalized in a perovskite-like structure. (C) Differential charge density analysis around all the oxygen atoms shows clearly the H peaks. (D) The atoms in an asymmetric unit, where three independent H<sub>2</sub>O molecules in a [Mg(H<sub>2</sub>O)<sub>6</sub>]<sup>2+</sup> unit and the NH<sub>4</sub><sup>+</sup> motif are interconnected by Cl<sup>-</sup>.



**Figure. 3.** Chlorine isotope composition of solid materials in solar system. Inset: NanoSIMS scanning Cl isotope image of ULM-1 specimen. Samples from the Earth, Meteorites, Comets, the Mars and the Moon are shown as blue, green, olive, yellow and half red symbols, respectively. Chlorine isotope composition of ULM-1 is shown in green pentagram. The terrestrial and meteorite data are from Ref. (*51*) and therein. Error bars are smaller than the size of symbols.



Figure. 4. Constraints on water fugacity  $f(H_2O)$  by the crystallization of ULM-1. The hexahydrate is stable above its thermodynamic equilibrium line (red), which constitutes the lower limit of  $f(H_2O)$  for lunar volcanic gases. The vapor-liquid phase boundary of water forms the upper limits (blue). Between the red and gray lines, a dihydrate phase of NLM-1 become thermodynamically more stable. Reference to the formation temperature of the earth mineral (>100 °C), the lower limit of  $f(H_2O)$  reached 1/3 of the record value in the driest volcano on earth - Lengai volcano (diamond). Taking H<sub>2</sub> degassing into account, the  $f(H_2O + H_2)$  values (purple doted line) become comparable to some 'hot spot' volcanoes on earth.  $P^{\emptyset}$  is the standard pressure (1 atm).