PINNS ERROR ESTIMATES FOR NONLINEAR EQUATIONS IN \mathbb{R} -smooth Banach spaces

Jiexing Gao Moscow Research Center, 2012 Labs Huawei Technologies Co., Ltd gaojiexing@huawei.com Vurii Zakharian
 Moscow Research Center, 2012 Labs
 Huawei Technologies Co., Ltd
 zakharyan.yuriy1@huawei-partners.com

ABSTRACT

In the paper, we describe in operator form classes of PDEs that admit PINN's error estimation. Also, for L^p spaces, we obtain a Bramble-Hilbert type lemma that is a tool for PINN's residuals bounding.

1 Introduction

In 2017, M. Raissi et al. introduced the Physics-informed neural network (PINN) approximating solutions to partial differential equations (PDEs) [29, 30]. It reduces losses related to PDE and boundary/initial conditions. In recent years, the number of papers dedicated to deep learning methods for solving PDEs, including PINNs, is constantly increasing (see, for instance, [9, 20, 24, 21] for deep learning methods and [14, 16, 17, 18, 31, 32] for PINN). Consequently, a thorough exploration of the theoretical aspects associated with PINNs is of great significance.

For instance, the question arises as to why PINN's training algorithm leads us to an accurate approximation. In other words, is it possible to control total error for sufficiently small residuals/training error? In [25], S. Mishra and R. Molinaro presented an error estimation answering this question and offered an operator description of the sufficient conditions for applying such a method. Yet, these conditions, while initially outlined rather generally, in practice, require obtaining the estimate itself to verify them. Also, while these conditions were outlined for L^p spaces, examples in [25] were all related to the problems in L^2 .

In [13], the authors studied a similar issue and presented PINN's error estimation for a homogeneous initial value problem with a linear generator of a strongly continuous semigroup as an operator. Furthermore, they applied this method to several nonlinear equations. Yet, they used another approach for a narrower class of PDEs.

Another related question is whether there is any neural network with sufficiently small residuals. In [5], the authors answered this question and derived an L^2 -bound on residuals using the Bramble-Hilbert lemma. For this reason, they used the Bramble-Hilbert lemma [36]. The challenging moment for our research is to obtain the Bramble-Hilbert lemma for L^p to extend results in [5] to such spaces. Also, refer to [2, 6, 7].

Therefore, the paper aims to resolve the following issues:

- 1. Specify PDE classes suitable for PINN's error estimation due to S. Mishra et al.
- 2. At the same time, to consider a general framework, for a broad class of non-Hilbert spaces (e.g., L^p).
- 3. Extend the approach in [13] to a broader PDEs class. Compare the respective estimate to the one obtained by the estimation for the PDE classes of question 2.
- 4. Extend results in [5] on PINN's residuals bounding to the case of L^p spaces.

Answering the first question, we consider the following four natural equation types and define used operators.

• (Parabolic-type equation)

$$\frac{dw}{dt} = A(t)(w) \tag{1}$$

• (Generalized parabolic-type equation)

$$\frac{dw}{dt} + \frac{d}{dt} \sum_{k=1}^{K} U_k w = A(t)(w)$$
(2)

• (Hyperbolic-type equation)

$$\frac{d^2w}{dt^2} = -Uw + F(t)(w) + A(t)\left(\frac{dw}{dt}\right)$$
(3)

• (Elliptic-type equation)

$$A(y) = f_{eq} \tag{4}$$

Regarding the second question, in [2, 5, 25], the authors repeated one procedure: they estimated the time derivative of the total error squared L^2 -norm using the residuals. After that, they applied the Grönwall-Bellman lemma to obtain the desired estimate. To extend this idea to a Banach space X, we need an existence of the time derivative of the total error p-powered X-norm for some p > 1. For this reason, in subsection 2.1, we consider so-called \mathbb{R} -smooth Banach spaces and introduce a real p-form. In subsections 2.2–2.4, using this form, we define operators of (1)–(4) in such spaces. As a result, in subsections 3.1 and 3.3–3.5, we obtain a total error estimate in general (operator) form, answering the second question.

To answer the third question, in subsection 3.2, we expand the approach in [13] to semi-linear equations and compare the respective estimate with the estimate derived from differentiating the p-powered norm.

Finally, in section 4, since the Bramble-Hilbert lemma was a core tool for PINN's residuals bounding in [5], we derive a Bramble-Hilbert type lemma for L^p and outline obtaining L^p bounds on PINN's residuals.

2 Preliminary

In this section, to describe PDE classes (1)–(4), we first define admissible spaces. Then in such spaces, we introduce a *real p-form* that is a core tool in the paper and derive its properties. Using this form, we define four types of operators used in (1)–(4). Moreover, in these definitions, we use functions ψ , which allow us to factor in additional conditions (boundary conditions in most cases). Precisely, we introduce (p, ψ) -submonotone, (p, ψ) -powered, (p, ψ) -submonotone w.r.t. norm, and (p, ψ) -coercive operators and consider several examples. Also, we define (p, κ) -dissipative operators as one essential subtype.

In what follows, we will use the following denotations. Let \mathbb{K} be either \mathbb{C} or \mathbb{R} . Let \mathbb{R}^+ stand for $[0; +\infty)$. Let X be a Banach space over \mathbb{K} with norm $\|\cdot\|$, and $S = \{y \in X \mid \|y\| = 1\}$ be a unit sphere. Also, let I = (0; T) and $\overline{I} = [0; T]$ be time intervals for some T > 0.

2.1 R-smooth Banach space and real *p*-form

An estimation in [2, 5, 25] was based on the Grönwall-Bellman lemma. To apply the lemma, the authors estimated the time derivative of squared norm $\frac{d}{dt} ||w - \tilde{w}||_{L^2}^2$, where w is the solution of a PDE, and \tilde{w} is its approximation. To extend this idea to a Banach space X, we need an existence of the derivative $\frac{d}{dt} ||w - \tilde{w}||^p$ for some p > 1. For this reason, we consider \mathbb{R} -smooth Banach spaces as admissible spaces. In fact, this existence is necessary only for (1)–(3), yet \mathbb{R} -smooth Banach spaces appear to be useful also for (4).

Definition 1. ([12]) A real-valued function $\varphi : X \to \mathbb{R}$ is said to be *Gateaux* \mathbb{R} -differentiable at $y \in X$ if for every $\chi \in X$

$$\exists \lim_{\mathbb{R} \ni s \to 0} \frac{\varphi(y + s\chi) - \varphi(y)}{s} =: D(\varphi)(y;\chi)$$

called *Gateaux* \mathbb{R} -*derivative at* y *w.r.t. direction* χ .

Remark 1. We used the Gateaux derivative as in paper [12], since we don't need s to be complex. Let us note that the Gateaux \mathbb{R} -derivative still satisfies the chain rule: if $w : \mathbb{R}^+ \to X$ is differentiable, $D(\varphi)(y, \cdot)$ exists and continuous, for $\varphi : X \to \mathbb{R}$ and every $y \in X$, then

$$\frac{d\varphi(w)}{dt} = D(\varphi)\left(w;\frac{dw}{dt}\right) \tag{5}$$

Definition 2. ([12]) A Banach space X is said to be \mathbb{R} -smooth if its norm is Gateaux \mathbb{R} -differentiable at every $y \in S$. **Remark 2.** If X is a \mathbb{R} -smooth Banach space, then for every $y \in X \setminus \{0\}$, 1. $D(\|\cdot\|)(y;\cdot)$ is \mathbb{R} -linear, that is

$$D(\|\cdot\|)(y;\lambda_1\chi_1 + \lambda_2\chi_2) = \lambda_1 D(\|\cdot\|)(y;\chi_1) + \lambda_2 D(\|\cdot\|)(y;\chi_2)$$

for every $y_1, y_2 \in X$ and $\lambda_1, \lambda_2 \in \mathbb{R}$.

2. $D(\|\cdot\|)(y;\cdot)$ is bounded and

$$||D(||\cdot||)(y;\cdot)|| = 1$$

3. $D(\|\cdot\|)(y;y) = \|y\|.$

Furthermore, $D(\|\cdot\|)(y;\cdot)$ is a unique real-valued functional satisfying (1) - (3) [15]. On the other hand, we can define a semi-inner-product [22],

- 1. $[y_1 + y_2, y_3] = [y_1, y_3] + [y_2, y_3]$, and $[\lambda y_1, y_2] = \lambda [y_1, y_2]$, for $\lambda \in \mathbb{K}$.
- 2. [y, y] > 0, for $y \neq 0$,
- 3. $|[y_1, y_2]|^2 \le [y_1, y_1][y_2, y_2].$

which agrees with the norm, that is $[y, y] = ||y||^2$.

J. R. Giles showed that the semi-inner-product $[\cdot, \cdot] : X \times X \to \mathbb{K}$ satisfies

- 1. (homogeneity) $[y_1, \lambda y_2] = \overline{\lambda}[y_1, y_2]$, for every $x, y \in X$ and any $\lambda \in \mathbb{K}$.
- 2. (continuity) $\operatorname{Re}[y_1, y_2 + sy_1] \to \operatorname{Re}[y_1, y_2]$ for all real $s \to 0$.

if and only if the respective norm is Gateaux \mathbb{R} -differentiable at any point of unit sphere S [12]. Furthermore,

$$\operatorname{Re}([\chi, y]) = \|y\| D(\|\cdot\|)(y; \chi), \, \forall y \in X$$

Hence,

$$[\chi, y] = \|y\| \left(D(\|\cdot\|)(y;\chi) - iD(\|\cdot\|)(y;i\chi) \right).$$

In the paper, we don't need to use the Gateaux \mathbb{R} -derivative of the norm or even the semi-inner product. Instead, we need $||y_2||^{p-2} \operatorname{Re}[y_1, y_2]$ for some p > 1. For this reason, we introduce the following notion.

Definition 3. Let p > 1, X be a \mathbb{R} -smooth Banach space. We will call a form $\langle \cdot, \cdot \rangle_p : X \times X \to \mathbb{R}$, where $\langle y_1, y_2 \rangle_p := \|y_2\|^{p-2} \operatorname{Re}[y_1, y_2]$, a *real p-form*.

Remark 3. The real *p*-form is correctly defined. Really, even though at $y_2 = 0$, the Gateaux \mathbb{R} -derivative $D(\|\cdot\|)(y_2; y_1)$ does not exists, for p > 1, we have

$$\langle y_1, y_2 \rangle_p = ||y_2||^{p-1} D(||\cdot||) (y_2; y_1) \to 0, y_2 \to 0$$

Let us note that $||y_2||^{p-2}[y_1, y_2]$ (without real part) appears in literature as a *semi-inner product of type p* [26]. Lemma 1. A real p-form on a \mathbb{R} -smooth Banach space satisfies the following properties.

- $I. \ \langle y_1 + y_2, y_3 \rangle_p = \langle y_1, y_3 \rangle_p + \langle y_2, y_3 \rangle_p, \text{ and } \langle \lambda y_1, y_2 \rangle_p = (\operatorname{Re}\lambda) \langle y_1, y_2 \rangle_p, \text{ for } \lambda \in \mathbb{K}.$
- 2. $\langle y_1, \lambda y_2 \rangle_p = (\operatorname{Re}\lambda) |\lambda|^{p-2} \langle y_1, y_2 \rangle_p$, for any $\lambda \in \mathbb{K}$.
- 3. $\langle y, y \rangle_p = \|y\|^p$.
- 4. $|\langle y_1, y_2 \rangle_p| \le ||y_1|| ||y_2||^{p-1}$
- 5. $\|\cdot\|^p$ is Gateaux \mathbb{R} -differentiable at any point and

$$D(\|\cdot\|^p)(y_2;y_1) = p\langle y_1, y_2 \rangle_p,$$
(6)

for every $y_1, y_2 \in X$.

- 6. $\langle \chi_n, y \rangle_p \to \langle \chi, y \rangle_p$, if $\chi_n \to \chi$.
- 7. If $w: \overline{I} \to X$ is differentiable, then

$$\frac{d}{dt} \|w\|^p = p \left\langle \frac{dw}{dt}, w \right\rangle_p \tag{7}$$

8. Let $w : \overline{I} \to X$ be differentiable, $\mathcal{D}(U) \subset X$ be a subspace, $U : \mathcal{D}(U) \to X$ be a linear operator such that Uw is differentiable and $\frac{d}{dt}Uw = U\frac{dw}{dt}$. Then

$$\frac{d}{dt} \|Uw\|^p = p \left\langle U\frac{dw}{dt}, Uw \right\rangle_p \tag{8}$$

9. If $D(\|\cdot\|)(\cdot, \chi)$ is continuous on the ring $\{y \in X \mid r_0 < \|x\| < r_1\}$, for every $\chi \in X$ and $0 < r_0 < r_1$, then the real p-form is continuous w.r.t. the second variable, that is,

$$\langle \chi, y_n \rangle_p \to \langle \chi, y \rangle_p, \ \forall y \in X, \forall y_n \to y.$$

Proof. Properties 1–6 follow from the definition and semi-inner product properties. Property 7 is a consequence of 5, 6, and (5).

In property 8, we cannot apply the chain rule since $D(||U \cdot ||^p)(y; \chi) = p \langle U\chi, Uy \rangle_p$ is not continuous w.r.t. χ in general. However,

$$\begin{aligned} \frac{\|Uw(t+\Delta t)\|^p - \|Uw(t)\|^p}{\Delta t} &= \frac{\|Uw(t) + U(w(t+\Delta t) - w(t))\|^p - \|Uw(t)\|^p}{\Delta t} = \\ &= D(\|U\cdot\|^p) \left(w(t); U\frac{w(t+\Delta t) - w(t)}{\Delta t}\right) + \frac{o(\Delta t)}{\Delta t} = \\ &= p \left\langle U\frac{w(t+\Delta t) - w(t)}{\Delta t}, w(t) \right\rangle_p + \frac{o(\Delta t)}{\Delta t}. \end{aligned}$$

With $U \frac{w(t+\Delta t)-w(t)}{\Delta t} \rightarrow \frac{d}{dt}Uw = U \frac{dw}{dt}$ and property 6, we obtain (8).

To prove property 9, we need to consider two cases. If $y \neq 0$, then starting from some n, y_n all together with y contain in a ring, where the real p-form is continuous. If $y_n \to 0$, then $\langle \chi, y_n \rangle_p \to 0 = \langle \chi, 0 \rangle_p$ by property 4.

Let us describe two examples.

Example 1. Every Hilbert space is a \mathbb{R} -smooth space. The inner product coincides with the semi-inner product, and the real 2-form is

$$\langle y_1, y_2 \rangle_2 = \operatorname{Re} \langle y_1, y_2 \rangle$$

Example 2. If $X = L^p(\Omega; \mathbb{C}^n)$ is a space of vector-valued functions, in a case of σ -finite measure, p > 1, with the norm

$$\|\mathbf{y}\|_{L^p(\Omega;\mathbb{C}^n)} = \left(\sum_{i=1}^n \|y_i\|_{L^p(\Omega;\mathbb{C})}^p\right)^{\frac{1}{p}}$$

then one can show that

$$[\mathbf{y}_1, \mathbf{y}_2] = \frac{\sum_{i=1}^n \int_{\Omega} |y_{2,i}|^{p-1} \operatorname{sgn}(\overline{y_{2,i}}) y_{1,i} dx}{\|\mathbf{y}_2\|_{L^p(\Omega;\mathbb{C}^n)}^{p-2}}$$

and real *p*-form is

$$\langle \mathbf{y}_1, \mathbf{y}_2 \rangle_p = \operatorname{Re} \sum_{i=1}^n \int_{\Omega} |y_{2,i}|^{p-1} \operatorname{sgn}(\overline{y_{2,i}}) y_{1,i} dx$$

Let us note that it is continuous w.r.t. the second variable.

For other examples, see, for instance, [19].

2.2 Submonotone operators

First, we define operators A(t) used in (1) and (3). In fact, we require -A(t) instead (see remark below).

Definition 4. Let X be a \mathbb{R} -smooth Banach space, $\mathcal{D}(A) \subset X$ be a subspace. Let $A : \mathcal{D}(A) \to X$ be an operator on X and $\mathcal{M} \subset \mathcal{D}(A)$ be a subspace. We will say that A is (p, ψ) -submonotone on \mathcal{M} for some $\psi : \mathcal{D}(A) \times \mathcal{M} \to \mathbb{R}$ and p > 1 if it is true that

$$\langle A(\chi) - A(y), \chi - y \rangle_p \ge \psi(\chi, y) - \Lambda(\chi, y) \|\chi - y\|^p$$

for every $\chi \in \mathcal{D}(A)$, $y \in \mathcal{M}$, and some $\Lambda(\chi, y) \ge 0$.

Here, ψ and \mathcal{M} are related to some additional conditions (boundary in most cases). Precisely, $\psi(\chi, y)$ should vanish if χ satisfies these conditions, and \mathcal{M} contains functions that satisfy one of the conditions. Also, in the case of a solution w and some PINN approximation \tilde{w} , $\psi(\tilde{w}, w)$ should be estimated in terms of respective additional condition residual. For this reason, we give the following definition.

Definition 5. Let $\mathcal{M} \subset \mathcal{D}_{\psi} \subset \mathcal{D}(B) \subset X$ be subspaces. Let $B : \mathcal{D}(B) \to Y$ be an operator on X and $\psi : \mathcal{D}_{\psi} \times \mathcal{M} \to \mathbb{R}$ be a function. We say that ψ is *subordinate to B on* \mathcal{M} , if there exists $\rho \in C(Y; \mathbb{R}^+)$ such that $s \to 0 \Rightarrow \rho(s) \to 0$ and

$$|\psi(\chi, y)| \le \gamma(\chi, y)\rho(B(\chi) - B(y))$$

for every $\chi \in \mathcal{D}_{\psi}$, $y \in \mathcal{M}$, and some $\gamma(\chi, y) \ge 0$.

Remark 4. We use the name "submonotone" due to a similar term in the paper [33]. In what follows, we will use the fact that -A is submonotone, that is,

$$\langle A(\chi) - A(y), \chi - y \rangle_p \le \psi(\chi, y) + \Lambda(\chi, y) \|\chi - y\|^p.$$

Here we take $-\psi$ instead of ψ since we don't care about its sign.

Remark 5. In most cases, operator A is submonotone on $\mathcal{M} = \mathcal{D}(A)$, or function ψ is subordinate on $\mathcal{M} = \mathcal{D}_{\psi}$. For these cases, we will say (p, ψ) -submonotone and subordinate, respectively.

Let us consider an essential case of A, where -A is submonotone. Working with semi inner-product G. Lumer and R. S. Phillips [23] gave the following definition.

Definition 6. ([23]) Let $\mathcal{D}(A) \subset X$ be a subspace. $A : \mathcal{D}(A) \to X$ is said to be *dissipative* if

$$\operatorname{Re}[A(y), y] \le 0,$$

for every $y \in \mathcal{D}(A)$.

Let us note that in a \mathbb{R} -smooth Banach space one can replace $\operatorname{Re}[\cdot, \cdot]$ with the real *p*-form $\langle \cdot, \cdot \rangle_p$. We define an operator to be dissipative on the kernel of some function κ .

Definition 7. Let $\mathcal{D}(A) \subset X$ be a subspace of a \mathbb{R} -smooth Banach space X. An operator $A : \mathcal{D}(A) \to X$ is said to be (p, κ) -dissipative for some $\kappa : \mathcal{D}(A) \to \mathbb{R}$ and p > 1 if it is true that

$$\langle A(y), y \rangle_p \le \kappa(y)$$

for every $y \in \mathcal{D}(A)$.

Now, let $A = A_0 + F$, where $A_0 : \mathcal{D}(A_0) \to X$ is linear (p, κ) -dissipative, and $F : \mathcal{D}(F) \to X$ is conditionally Lipshitz. The last means that

$$||F(y_1) - F(y_2)|| \le \Lambda(y_1, y_2) ||y_1 - y_2||,$$

for every $y_1, y_2 \in \mathcal{D}(F)$ and some $\Lambda(y_1, y_2) \ge 0$. If we denote $\psi(\chi, y) := \kappa(\chi - y)$, then -A is (p, ψ) -submonotone.

Let us consider several examples. Not to complicate, we consider simple derivatives only. The following statement allows us to extend submonotonicity to a positive linear combination and a closure. The proof is standard. The second property follows from the real *p*-form continuous property (9 of lemma 1).

Statement 1. Let X be a \mathbb{R} -smooth Banach space. The following two properties hold.

1. If $A_{1,2}$ are $(p, \psi_{1,2})$ -submonotone on \mathcal{M} , then

$$A(t) := g_1 A_1 + g_2 A_2$$

is (p, ψ) -submonotone on \mathcal{M} , for $\psi = g_1\psi_1 + g_2\psi_2$ and every $g_{1,2} \ge 0$.

2. Moreover, let $D(\|\cdot\|)(\cdot, \chi)$ be continuous on the ring $\{y \in X \mid r_0 < \|y\| < r_1\}$, for every $0 < r_0 < r_1$ and $\chi \in X$. If A is (p, ψ) -submonotone, there exists a closure \overline{A} , ψ is continuous w.r.t. \overline{A} -norm, that is

$$y_n \to y, \ \chi_n \to \chi, \ Ay_n \to \overline{Ay}, \ A\chi_n \to \overline{A}\chi \ \Rightarrow \ \psi(\chi_n, y_n) \to \psi(\chi, y),$$

and Λ is continuous w.r.t. \overline{A} -norm, then \overline{A} is (p, ψ) -submonotone.

We start with (p, κ) -dissipative operators.

Example 3. Let $\Omega \subset \mathbb{R}^m$, $\Gamma = \partial \Omega$, $X = L^p(\Omega; \mathbb{C})$, $1 \le j \le m, p > 1$. Then we have a (p, κ) -dissipative operator ∂y

$$Ay = s \frac{\partial y}{\partial x_j}, \ s \in \mathbb{R}$$

Really, let $y = \eta + i\zeta$. Then

$$\begin{split} \langle Ay, y \rangle_p &= s \operatorname{Re} \int_{\Omega} \frac{\partial y}{\partial x_j} |y|^{p-1} \operatorname{sgn}(\overline{y}) dx = s \int_{\Omega} |y|^{p-2} \left(\frac{\partial \eta}{\partial x_j} \eta + \frac{\partial \zeta}{\partial x_j} \zeta \right) dx = \\ &= \frac{s}{2} \int_{\Omega} |y|^{p-2} \frac{\partial |y|^2}{\partial x_j} dx = \frac{s}{p} \int_{\Gamma} |y|^p (\mathbf{e}_j \cdot \mathbf{n}_{\Gamma}) d\Gamma \end{split}$$

Example 4. Let $\Omega \subset \mathbb{R}^m$, $\Gamma = \partial \Omega$, $X = L^p(\Omega; \mathbb{C})$, $1 \le j \le m, p \ge 4$ or p = 2. Then

$$Ay = \frac{\partial^2 y}{\partial x_j^2}$$

is (p, κ) -dissipative.

Really, let again $y = \eta + i\zeta$. Then

$$\begin{split} \langle Ay, y \rangle_p &= \operatorname{Re} \int_{\Omega} \frac{\partial^2 y}{\partial x_j^2} |y|^{p-1} \operatorname{sgn}(\overline{y}) dx = \int_{\Omega} |y|^{p-2} \left(\frac{\partial^2 \eta}{\partial x_j^2} \eta + \frac{\partial^2 \zeta}{\partial x_j^2} \zeta \right) dx = \\ &= \int_{\Gamma} |y|^{p-2} \left(\eta \frac{\partial \eta}{\partial x_j} + \zeta \frac{\partial \zeta}{\partial x_j} \right) (\mathbf{e}_j \cdot \mathbf{n}_{\Gamma}) d\Gamma - \int_{\Omega} \frac{\partial}{\partial x_j} \left(|y|^{p-2} \eta \right) \frac{\partial \eta}{\partial x_j} + \frac{\partial}{\partial x_j} \left(|y|^{p-2} \zeta \right) \frac{\partial \zeta}{\partial x_j} dx = \\ &= \int_{\Gamma} |y|^{p-2} \left(\eta \frac{\partial \eta}{\partial x_j} + \zeta \frac{\partial \zeta}{\partial x_j} \right) (\mathbf{e}_j \cdot \mathbf{n}_{\Gamma}) d\Gamma - \int_{\Omega} |y|^{p-2} \left(\left(\frac{\partial \eta}{\partial x_j} \right)^2 + \left(\frac{\partial \zeta}{\partial x_j} \right)^2 \right) dx - \\ &- \frac{p-2}{4} \int_{\Omega} |y|^{p-4} \left(\frac{\partial |y|^2}{\partial x_j} \right)^2 dx \leq \int_{\Gamma} |y|^{p-2} \left(\eta \frac{\partial \eta}{\partial x_j} + \zeta \frac{\partial \zeta}{\partial x_j} \right) (\mathbf{e}_j \cdot \mathbf{n}_{\Gamma}) d\Gamma = \\ &= \int_{\Gamma} |y|^{p-2} \operatorname{Re} \left(\overline{y} \frac{\partial y}{\partial x_j} \right) (\mathbf{e}_j \cdot \mathbf{n}_{\Gamma}) d\Gamma. \end{split}$$

Example 5. Let $\Omega \subset \mathbb{R}^m$, $\Gamma = \partial \Omega$, $X = L^2(\Omega; \mathbb{C})$, $1 \le j \le m$. Then

$$Ay = (s_1^2 + is_2)\frac{\partial^2 y}{\partial x_j^2},$$

where $s_1, s_2 \in \mathbb{R}$, is $(2, \kappa)$ -dissipative. Really, let again $y = \eta + i\zeta$. Then

$$\begin{split} \langle Ay, y \rangle_2 &= \operatorname{Re} \int_{\Omega} (s_1^2 + is_2) \frac{\partial^2 y}{\partial x_j^2} \overline{y} dx = s_1^2 \int_{\Omega} \left(\frac{\partial^2 \eta}{\partial x_j^2} \eta + \frac{\partial^2 \zeta}{\partial x_j^2} \zeta \right) dx + s_2 \int_{\Omega} \left(\frac{\partial^2 \eta}{\partial x_j^2} \zeta - \frac{\partial^2 \zeta}{\partial x_j^2} \eta \right) dx = \\ &= \int_{\Gamma} \left((s_1^2 \eta + s_2 \zeta) \frac{\partial \eta}{\partial x_j} + (s_1^2 \zeta - s_2 \eta) \frac{\partial \zeta}{\partial x_j} \right) (\mathbf{e}_j \cdot \mathbf{n}_{\Gamma}) d\Gamma - s_1^2 \int_{\Omega} \left(\left(\frac{\partial \eta}{\partial x_j} \right)^2 + \left(\frac{\partial \zeta}{\partial x_j} \right)^2 \right) dx - \\ &- s_2 \int_{\Omega} \left(\frac{\partial \eta}{\partial x_j} \frac{\partial \zeta}{\partial x_j} - \frac{\partial \eta}{\partial x_j} \frac{\partial \zeta}{\partial x_j} \right) dx \leq \int_{\Gamma} \left((s_1^2 \eta + s_2 \zeta) \frac{\partial \eta}{\partial x_j} + (s_1^2 \zeta - s_2 \eta) \frac{\partial \zeta}{\partial x_j} \right) (\mathbf{e}_j \cdot \mathbf{n}_{\Gamma}) d\Gamma = \\ &= \int_{\Gamma} \operatorname{Re} \left((s_1^2 + is_2) \overline{y} \frac{\partial y}{\partial x_j} \right) (\mathbf{e}_j \cdot \mathbf{n}_{\Gamma}) d\Gamma. \end{split}$$

Example 6. Let $\Omega \subset \mathbb{R}^m$, $\Gamma = \partial \Omega$, $X = L^2(\Omega; \mathbb{R}^n)$, $1 \le j \le m$. Then

$$A\mathbf{y} = \frac{\partial^{\sigma}}{\partial x_j^{\sigma}} Q\mathbf{y},$$

is $(2,\kappa)\text{-dissipative, where }\sigma\geq 0 \text{ and } Q\in \mathbb{R}^{n\times n}$ is a such matrix that

$$\begin{cases} (-1)^{\frac{\sigma}{2}}Q \leq 0, & \sigma = 2l \\ Q = Q^T, & \sigma = 2l+1 \end{cases}$$

Really, if $\sigma = 2l$,

$$\begin{split} \langle A\mathbf{y}, \mathbf{y} \rangle_2 &= \int_{\Omega} \left(\frac{\partial^{\sigma}}{\partial x_j^{\sigma}} Q \mathbf{y} \cdot \mathbf{y} \right) dx = \sum_{\nu=0}^{l-1} \int_{\Gamma} (-1)^{\nu} \left(\frac{\partial^{\sigma-1-\nu}}{\partial x_j^{\sigma-1-\nu}} Q \mathbf{y} \cdot \frac{\partial^{\nu}}{\partial x_j^{\nu}} \mathbf{y} \right) (\mathbf{e}_j \cdot \mathbf{n}_{\Gamma}) d\Gamma + \\ &+ \int_{\Omega} \left((-1)^l Q \frac{\partial^l}{\partial x_j^l} \mathbf{y} \cdot \frac{\partial^l}{\partial x_j^l} \mathbf{y} \right) dx \leq \sum_{\nu=0}^{l-1} \int_{\Gamma} (-1)^{\nu} \left(\frac{\partial^{\sigma-1-\nu}}{\partial x_j^{\sigma-1-\nu}} Q \mathbf{y} \cdot \frac{\partial^{\nu}}{\partial x_j^{\nu}} \mathbf{y} \right) (\mathbf{e}_j \cdot \mathbf{n}_{\Gamma}) d\Gamma \end{split}$$

If $\sigma = 2l + 1$,

$$\begin{split} \langle A\mathbf{y}, \mathbf{y} \rangle_2 &= \int_{\Omega} \left(\frac{\partial^{\sigma}}{\partial x_j^{\sigma}} Q \mathbf{y} \cdot \mathbf{y} \right) dx = \sum_{\nu=0}^{l-1} \int_{\Gamma} (-1)^{\nu} \left(\frac{\partial^{\sigma-1-\nu}}{\partial x_j^{\sigma-1-\nu}} Q \mathbf{y} \cdot \frac{\partial^{\nu}}{\partial x_j^{\nu}} \mathbf{y} \right) (\mathbf{e}_j \cdot \mathbf{n}_{\Gamma}) d\Gamma + \\ &+ (-1)^l \int_{\Omega} \left(\frac{\partial^{l+1}}{\partial x_j^l} Q \mathbf{y} \cdot \frac{\partial^l}{\partial x_j^l} \mathbf{y} \right) dx \end{split}$$

Since

$$\int_{\Omega} \left(\frac{\partial^{l+1}}{\partial x_j^{l+1}} Q \mathbf{y} \cdot \frac{\partial^l}{\partial x_j^l} \mathbf{y} \right) dx = \int_{\Gamma} \left(\frac{\partial^l}{\partial x_j^l} Q \mathbf{y} \cdot \frac{\partial^l}{\partial x_j^l} \mathbf{y} \right) \left(\mathbf{e}_j \cdot \mathbf{n}_{\Gamma} \right) d\Gamma - \int_{\Omega} \left(\frac{\partial^l}{\partial x_j^l} Q \mathbf{y} \cdot \frac{\partial^{l+1}}{\partial x_j^{l+1}} \mathbf{y} \right) dx$$

and $Q = Q^T$, then

$$\begin{split} \langle A\mathbf{y}, \mathbf{y} \rangle_2 &= \sum_{\nu=0}^{l-1} \int_{\Gamma} (-1)^{\nu} \left(\frac{\partial^{\sigma-1-\nu}}{\partial x_j^{\sigma-1-\nu}} Q\mathbf{y} \cdot \frac{\partial^{\nu}}{\partial x_j^{\nu}} \mathbf{y} \right) \left(\mathbf{e}_j \cdot \mathbf{n}_{\Gamma} \right) d\Gamma \\ &+ \frac{(-1)^l}{2} \int_{\Gamma} \left(\frac{\partial^l}{\partial x_j^l} Q\mathbf{y} \cdot \frac{\partial^l}{\partial x_j^l} \mathbf{y} \right) \left(\mathbf{e}_j \cdot \mathbf{n}_{\Gamma} \right) d\Gamma \end{split}$$

Now, let us consider other submonotone operators.

Example 7. Let $X = L^p((a; b); \mathbb{R}), p > 1$. We consider the following operator

$$A(y) = syy',$$

where $s \in \mathbb{R}$. Then -A is (p, ψ) -submonotone. Really, if $\hat{y} = y_1 - y_2$, then we have

$$A(y_1) - A(y_2) = s\hat{y}\hat{y}' + s\hat{y}y_2' + sy_2\hat{y}'$$

Hence, the desired estimate is

$$\begin{split} \langle A(y_1) - A(y_2), \hat{y} \rangle_p &= \int_a^b (s \hat{y} \hat{y}' + s \hat{y} y_2' + s y_2 \hat{y}') |\hat{y}|^{p-1} \operatorname{sgn}(\hat{y}) dx \leq \\ &\leq \frac{s}{p+1} |\hat{y}|^{p+1} \Big|_a^b + |s| \|y_2\|_{C^1([a;b];\mathbb{R})} \|\hat{y}\|_{L^p((a;b);\mathbb{R})}^p + \frac{s}{p} |\hat{y}|^p y_2 \Big|_a^b - \frac{s}{p} \int_a^b y_2' |y|^p dx \leq \\ &\leq \left(\frac{s}{p+1} |\hat{y}|^{p+1} + \frac{s}{p} |\hat{y}|^p y_2\right) \Big|_a^b + |s| \left(1 + \frac{1}{p}\right) \|y_2\|_{C^1([a;b];\mathbb{R})} \|\hat{y}\|_{L^p((a;b);\mathbb{R})}^p. \end{split}$$

The following example is a multidimensional case of example 7; yet it's negative is submonotone on a subspace. **Example 8.** Here we consider in $L^p(\Omega; \mathbb{R}^m)$, p > 1, the nonlinearity of the Navier-Stokes equation for an incompressible fluid. Let $\Omega \subset \mathbb{R}^m$, $\Gamma = \partial \Omega$, $X = L^p(\Omega; \mathbb{R}^m)$, and

$$A(\mathbf{y}) = s(\mathbf{y} \cdot \nabla)\mathbf{y}$$

where $s \in \mathbb{R}$. Then -A is (p, ψ) -submonotone on $\mathcal{M} = \{\mathbf{y} \mid \nabla \cdot \mathbf{y} = 0\}$ (continuity condition). Really, if we put $\hat{\mathbf{y}} = \chi - \mathbf{y}, \mathbf{y} \in \mathcal{M}$, then

$$A(\chi) - A(\mathbf{y}) = s(\hat{\mathbf{y}} \cdot \nabla)\hat{\mathbf{y}} + s(\hat{\mathbf{y}} \cdot \nabla)\mathbf{y} + s(\mathbf{y} \cdot \nabla)\hat{\mathbf{y}}$$

For the first term, we have

$$\begin{split} s\langle (\hat{\mathbf{y}} \cdot \nabla) \hat{\mathbf{y}}, \hat{\mathbf{y}} \rangle_{p} &= s \sum_{i=1}^{m} \int_{\Omega} |\hat{y}_{i}|^{p-1} \mathrm{sgn}(\hat{y}_{i}) \sum_{j=1}^{m} \hat{y}_{j} \frac{\partial \hat{y}_{i}}{\partial x_{j}} dx = \\ &= \frac{s}{p} \int_{\Gamma} |\hat{\mathbf{y}}|^{p} \left(\hat{\mathbf{y}} \cdot \mathbf{n}_{\Gamma} \right) d\Gamma - \frac{s}{p} \int_{\Omega} |\hat{\mathbf{y}}|^{p} \left(\nabla \cdot \hat{\mathbf{y}} \right) dx \leq \\ &\leq \frac{s}{p} \int_{\Gamma} |\hat{\mathbf{y}}|^{p} \left(\hat{\mathbf{y}} \cdot \mathbf{n}_{\Gamma} \right) d\Gamma + |s| \frac{\|\chi\|_{C(\overline{\Omega};\mathbb{R}^{m})} + \|\mathbf{y}\|_{C(\overline{\Omega};\mathbb{R}^{m})}}{p^{2}} \left((p-1) \|\hat{\mathbf{y}}\|_{L^{p}(\Omega;\mathbb{R}^{m})}^{p} + \|\nabla \cdot \hat{\mathbf{y}}\|_{L^{p}(\Omega;\mathbb{R})}^{p} \right) \end{split}$$

For the second term,

$$\begin{split} s\langle (\hat{\mathbf{y}} \cdot \nabla) \mathbf{y}, \hat{\mathbf{y}} \rangle_p &= s \sum_{i=1}^m \int_{\Omega} |\hat{y}_i|^{p-1} \mathrm{sgn}(\hat{y}_i) \sum_{j=1}^m \hat{y}_j \frac{\partial y_i}{\partial x_j} dx \leq \\ &\leq |s| \sum_{i=1}^m \sum_{j=1}^m \|\nabla \mathbf{y}\|_{L^{\infty}(\Omega; \mathbb{R}^{m \times m})} \int_{\Omega} |\hat{y}_i|^{p-1} |\hat{y}_j| dx \leq m |s| \|\nabla \mathbf{y}\|_{L^{\infty}(\Omega; \mathbb{R}^{m \times m})} \|\hat{\mathbf{y}}\|_{L^p(\Omega; \mathbb{R}^m)}^p \end{split}$$

Finally,

$$s\langle (\mathbf{y}\cdot\nabla)\hat{\mathbf{y}}, \hat{\mathbf{y}}\rangle_{p} = s\sum_{i=1}^{m} \int_{\Omega} |\hat{y}_{i}|^{p-1} \operatorname{sgn}(\hat{y}_{i}) \sum_{j=1}^{m} y_{j} \frac{\partial \hat{y}_{i}}{\partial x_{j}} dx =$$
$$= \frac{s}{p} \int_{\Gamma} |\hat{\mathbf{y}}|^{p} \left(\mathbf{y}\cdot\mathbf{n}_{\Gamma}\right) d\Gamma - \frac{s}{p} \int_{\Omega} |\hat{\mathbf{y}}|^{p} \left(\nabla\cdot\mathbf{y}\right) dx = \frac{s}{p} \int_{\Gamma} |\hat{\mathbf{y}}|^{p} \left(\mathbf{y}\cdot\mathbf{n}_{\Gamma}\right) d\Gamma$$

Let us note that ψ contains the term with $\nabla \cdot \hat{\mathbf{y}}$, which is subordinate to the continuity condition operator. **Example 9.** Let $\Omega \subset \mathbb{R}^m$, $\Gamma = \partial \Omega$, $X = L^p(\Omega; \mathbb{R})$, $p \ge 2$, and let A be a p-Laplace operator, that is

$$A(y) = \nabla \cdot \left(|\nabla y|^{p-2} \nabla y \right)$$

Then -A is (p, ψ) -submonotone. Really, let $\hat{y} = y_1 - y_2$. Then

$$\begin{aligned} \langle A(y_1) - A(y_2), \hat{y} \rangle_p &= \int_{\Omega} \nabla \cdot \left(|\nabla y_1|^{p-2} \nabla y_1 - |\nabla y_2|^{p-2} \nabla y_2 \right) |\hat{y}|^{p-1} \operatorname{sgn}(\hat{y}) dx = \\ &= \int_{\Gamma} |\hat{y}|^{p-1} \operatorname{sgn}(\hat{y}) \left(\left(|\nabla y_1|^{p-2} \nabla y_1 - |\nabla y_2|^{p-2} \nabla y_2 \right) \cdot \mathbf{n}_{\Gamma} \right) d\Gamma - \\ &- (p-1) \int_{\Omega} |\hat{y}|^{p-2} \left(|\nabla y_1|^{p-2} \nabla y_1 - |\nabla y_2|^{p-2} \nabla y_2 \right) \cdot (\nabla y_1 - \nabla y_2) dx \end{aligned}$$

With the Cauchy-Schwartz inequality in \mathbb{R}^m and the Young inequality,

$$- (|\nabla y_1|^{p-2} \nabla y_1 - |\nabla y_2|^{p-2} \nabla y_2) \cdot (\nabla y_1 - \nabla y_2) =$$

= $-|\nabla y_1|^p - |\nabla y_2|^p + (|\nabla y_1|^{p-2} + |\nabla y_2|^{p-2}) (\nabla y_1 \cdot \nabla y_2) \le$
= $-|\nabla y_1|^p - |\nabla y_2|^p + |\nabla y_1|^{p-1} |\nabla y_2| + |\nabla y_2|^{p-1} |\nabla y_1| \le 0$

Hence,

$$\langle A(y_1) - A(y_2), \hat{y} \rangle_p \le \int_{\Gamma} |\hat{y}|^{p-1} \operatorname{sgn}(\hat{y}) \left(\left(|\nabla y_1|^{p-2} \nabla y_1 - |\nabla y_2|^{p-2} \nabla y_2 \right) \cdot \mathbf{n}_{\Gamma} \right) d\Gamma$$

Example 10. Let $\Omega \subset \mathbb{R}^m$, $\Gamma = \partial \Omega$, $X = L^p(\Omega; \mathbb{R})$, p > 1, and let

$$A(y) = -y|y|^{p-2}$$

As in example 9, one can show that

$$\langle A(y_1) - A(y_2), y_1 - y_2 \rangle_p \le 0,$$

that is -A is (p, ψ) -submonotone.

2.3 Powered operators and submonotone operators w.r.t. norm

In [2], the authors considered the one-dimensional Camassa-Holm equation involving mixed derivative $\frac{\partial^3}{\partial t \partial x^2}$. For a sufficiently regular solution, one can take it as $-\frac{d}{dt}U$, where $U = -\frac{d^2}{dx^2}$. For such cases, we extend a concept of submonotonicity/subordination (for simplicity, we deal with the whole domain only). Precisely, we define operators U and A(t) used in (2). Remarkably, we can use such a definition of U in hyperbolic-type equation (3).

Definition 8. Let X be a \mathbb{R} -smooth Banach space, and $\mathcal{D}(U) \subset X$ be a subspace. Let $U : \mathcal{D}(U) \to X$ be a linear operator on X. We will say that U is (p, ψ) -powered for some $\psi : (\mathcal{D}(U))^4 \to \mathbb{R}$, if there exists an operator $U^{\frac{1}{p}} : \mathcal{D}\left(U^{\frac{1}{p}}\right) \to X, \mathcal{D}(U) \subset \mathcal{D}\left(U^{\frac{1}{p}}\right)$, satisfying

$$\langle U(\chi_1 - \chi_2), y_1 - y_2 \rangle_p = \psi(\chi_1, \chi_2, y_1, y_2) + \left\langle U^{\frac{1}{p}}(\chi_1 - \chi_2), U^{\frac{1}{p}}(y_1 - y_2) \right\rangle_p$$

for every $\chi_1, \chi_2, y_1, y_2 \in \mathcal{D}(U)$.

Now, we unveil the following extension of definition 4.

Definition 9. Let X be a \mathbb{R} -smooth Banach space, $\{U_k\}_{k=1}^K$ be a set of (p, ψ_k) -powered operators, and $\mathcal{D}(A) \subset \mathcal{D}(U_k) \subset \mathcal{D}\left(U_k^{\frac{1}{p}}\right)$ be subspaces. We say that A is (p, ψ) -submonotone, w.r.t. norm $\|\cdot\|_{\sum U_k^{\frac{1}{p}}}$ for some $\psi : (\mathcal{D}(A))^2 \to \mathbb{R}$, if

$$\langle A(\chi-y), \chi-y \rangle_p \le \psi(\chi,y) + \Lambda(\chi,y) \left(\|\chi-y\|^p + \sum_{k=1}^K \left\| U_k^{\frac{1}{p}} \chi - U_k^{\frac{1}{p}} y \right\|^p \right)$$

for all $\chi, y \in \mathcal{D}(A)$ and some $\Lambda(\chi, y) > 0$.

Also, we need to describe ψ and ψ_k as in definition 5.

Definition 10. Let $\mathcal{D}_{\psi} \subset \mathcal{D}(B) \subset X$ and $\mathcal{D}_{\psi} \subset \mathcal{D}(U_k) \subset \mathcal{D}\left(U_k^{\frac{1}{p}}\right) \subset X$ be subspaces, $1 \leq k \leq K$. Let $\{U_k\}_{k=1}^K$ be a set of (p, ψ_k) -powered operators, $\operatorname{Im}(U_k) \subset \mathcal{D}(B)$; $B : \mathcal{D}(B) \to Y$ be an operator on X, and $\psi : (\mathcal{D}_{\psi})^2 \to \mathbb{R}$

be a function. We say that ψ is subordinate to B w.r.t. U_1, \ldots, U_K if there exist $\rho_0, \ldots, \rho_K \in C(Y; \mathbb{R}^+)$, such that $s \to 0 \Rightarrow \rho_k(s) \to 0$ and

$$|\psi(\chi, y)| \le \gamma_0(\chi, y)\rho_0(B(\chi) - B(y)) + \sum_{k=1}^K \gamma_k(\chi, y)\rho_k(B(U_k\chi) - B(U_ky)),$$

for every $\chi, y \in \mathcal{D}_{\psi}$, and some $\gamma_k(\chi, y) \ge 0, 0 \le k \le K$.

Definition 11. Let $\mathcal{D}_{\psi} \subset \mathcal{D}(B) \subset X$ be subspaces. Let $B : \mathcal{D}(B) \to Y$ be an operator on X and $\psi : (\mathcal{D}_{\psi})^4 \to \mathbb{R}$ be a function. We say that ψ is *subordinate to* B if there exists $\rho \in C(Y; \mathbb{R}^+)$ such that $s \to 0 \Rightarrow \rho(s) \to 0$ and the following conditions holds,

$$|\psi(\chi_1, \chi_2, y_1, y_2)| \le \gamma(\chi_1, \chi_2, y_1, y_2)\rho(B(y_1) - B(y_2))$$

for every $\chi_1, \chi_2, y_1, y_2 \in \mathcal{D}_{\psi}$ and some $\gamma(\chi_1, \chi_2, y_1, y_2) \ge 0$.

Before considering the examples, we formulate an extension of (p, ψ) -powered property to a closure.

Statement 2. Let X be a \mathbb{R} -smooth Banach space. Moreover, let $D(\|\cdot\|)(\cdot,\chi)$ be continuous on the ring $\{y \in X \mid r_0 < \|y\| < r_1\}$ for every $0 < r_0 < r_1$ and $\chi \in X$. If U is (p, ψ) -powered, there exist closures $\overline{U^{\frac{1}{p}}}$ and \overline{U} with property

$$\chi_n \to \chi, \ U\chi_n \to \overline{U}\chi \ \Rightarrow \ U^{\frac{1}{p}}\chi_n \to \overline{U^{\frac{1}{p}}}\chi,$$

and ψ is continuous w.r.t. \overline{U} -norm, then \overline{U} is (p, ψ) -powered and $\overline{U}^{\frac{1}{p}} = U^{\frac{1}{p}}$.

Remark 6. One can also consider properties of submonotone w.r.t. norm operator, similar to statement 1.

We start with (p, ψ) -powered operators.

Example 11. Let $X = L^p(\Omega; \mathbb{R}^n)$, and $U = \text{diag}(\mu_1, \dots, \mu_n)$, $\mu_i > 0$, that is

$$U: \mathbf{y} = (y_1, \dots, y_n)^T \mapsto (\mu_1 y_1, \dots, \mu_n y_n)^T = U\mathbf{y}, \ \forall \mathbf{y} \in L^p(\Omega; \mathbb{R}^n).$$

Then U is (p, ψ) -powered with $U^{\frac{1}{p}} = \operatorname{diag}\left(\mu_1^{\frac{1}{p}}, \dots, \mu_n^{\frac{1}{p}}\right)$ and $\psi \equiv 0$. Such operators appear, for instance, in the Maxwell equation. Furthermore, if p = 2, U can be any symmetric, positively defined matrix.

Example 12. Let $X = L^2((a;b);\mathbb{R})$ (one can also consider \mathbb{C} -valued functions), $U = (-1)^{\sigma} \frac{\partial^{2\sigma}}{\partial x^{2\sigma}}$. Then U is $(2,\psi)$ -powered and $\sqrt{U} = \frac{\partial^{\sigma}}{\partial x^{\sigma}}$. Really,

$$\langle U\chi, y \rangle_2 = \left\langle (-1)^{\sigma} \chi^{(2\sigma)}, y \right\rangle_2 = \sum_{\nu=1}^{\sigma} (-1)^{\sigma+\nu-1} \chi^{(2\sigma-\nu)} y^{(\nu)} \Big|_a^b + \left\langle \chi^{(\sigma)}, y^{(\sigma)} \right\rangle_2$$

Moreover, respective \sqrt{U} -norm is the Sobolev $H^{\sigma}((a; b); \mathbb{R})$ -norm.

Now, let us turn to a submonotone w.r.t. a norm operator.

Example 13. Let $X = L^2((a; b); \mathbb{R})$. Then

$$A(y) = \frac{\frac{d}{dx} (|y|^2 y^{(\sigma)})}{y} = 2y' y^{(\sigma)} + y y^{(\sigma+1)},$$

where $\sigma \ge 0$, is $(2, \psi)$ -submonotone w.r.t. Sobolev $H^{\left\lceil \frac{\sigma}{2} \right\rceil}((a; b); \mathbb{R})$ -norm (see Example 12). In other words, let us show that

$$\langle A(y_1) - A(y_2), \hat{y} \rangle_2 \le \psi(\hat{y}) + \Lambda(y_1, y_2) \| \hat{y} \|_{H^{\lceil \frac{\sigma}{2} \rceil}((a;b);\mathbb{R})}^2, \ \hat{y} = y_1 - y_2,$$

for some ψ and some Λ .

We have

$$A(y_1) - A(y_2) = 2\left(\hat{y}'\hat{y}^{(\sigma)} + y'_2\hat{y}^{(\sigma)} + \hat{y}'y_2^{(\sigma)}\right) + \hat{y}\hat{y}^{(\sigma+1)} + y_2\hat{y}^{(\sigma+1)} + \hat{y}y_2^{(\sigma+1)} = = A(\hat{y}) + 2\left(y'_2\hat{y}^{(\sigma)} + \hat{y}'y_2^{(\sigma)}\right) + y_2\hat{y}^{(\sigma+1)} + \hat{y}y_2^{(\sigma+1)}$$

Hence, we need to estimate $\langle A(\hat{y}), \hat{y} \rangle_2$ and four terms of form $\langle \chi \hat{y}^{\tilde{\sigma}}, \hat{y} \rangle_2$, $0 \leq \tilde{\sigma} \leq \sigma + 1$, $\chi \in C^{\sigma+1-\left[\frac{\tilde{\sigma}}{2}\right]}([a;b];\mathbb{R})$. First,

$$\langle A(\hat{y}), \hat{y} \rangle_2 = \hat{y}^2 \hat{y}^{(\sigma)} \Big|_a^b,$$

where $y^2 y^{(\sigma)}|_a^b$ is subordinate, for instance, to the periodic boundary conditions operator $\left(y\Big|_a^b, \ldots, y^{(\sigma)}\Big|_a^b\right)^T$.

Let us show that

$$\left| \langle \chi \hat{y}^{(\tilde{\sigma})}, \hat{y} \rangle_2 \right| \le \psi(\hat{y}) + \Lambda(\|\chi\|_{C^{\sigma+1-\left[\frac{\tilde{\sigma}}{2}\right]}([a;b];\mathbb{R})}) \|\hat{y}\|_{H^{\left[\frac{\tilde{\sigma}}{2}\right]}((a;b);\mathbb{R})}^2$$

where $\psi(y)$ is subordinate to the periodic boundary conditions operator.

We consider two possible cases. First, let $\tilde{\sigma} = 2l$. Then

$$\left| \langle \chi \hat{y}^{(\tilde{\sigma})}, \hat{y} \rangle_2 \right| = \int_a^b \chi \hat{y}^{(2l)} \hat{y} dx = \sum_{\nu=1}^l (-1)^{\nu-1} \hat{y}^{(2l-\nu)} \left(\chi \hat{y} \right)^{(\nu)} \Big|_a^b + (-1)^l \int_a^b \hat{y}^{(l)} \left(\chi \hat{y} \right)^{(l)} dx$$

The first sum is subordinate to the periodic boundary conditions operator. It remains to consider

$$\left| \int_{a}^{b} \hat{y}^{(l)} \left(\chi \hat{y} \right)^{(l)} dx \right| = \left| \int_{a}^{b} \hat{y}^{(l)} \sum_{\nu=0}^{l} \binom{l}{\nu} \chi^{(l-\nu)} \hat{y}^{(\nu)} dx \right| \le 2^{l} \|\chi\|_{C^{l}([a;b];\mathbb{R})} \|\hat{y}\|_{H^{l}((a;b);\mathbb{R})}^{2}.$$

Now, let $\tilde{\sigma} = 2l + 1$. Then, similarly,

$$\begin{split} \left| \langle \chi \hat{y}^{(\tilde{\sigma})}, \hat{y} \rangle_2 \right| &= \int_a^b \chi \hat{y}^{(2l+1)} \hat{y} dx = \sum_{\nu=1}^{l+1} (-1)^{\nu-1} \hat{y}^{(2l+1-\nu)} \left(\chi \hat{y} \right))^{(\nu)} \Big|_a^b + \\ &+ (-1)^{l+1} \int_a^b \hat{y}^{(l)} \left(\chi \hat{y} \right)^{(l+1)} dx \end{split}$$

Again, we need to estimate

$$\left| \int_{a}^{b} \hat{y}^{(l)} \left(\chi \hat{y} \right)^{(l+1)} dx \right| = \left| \int_{a}^{b} \hat{y}^{(l)} \sum_{\nu=0}^{l+1} \binom{l+1}{\nu} \chi^{(l+1-\nu)} \hat{y}^{(\nu)} dx \right|.$$

If $\nu \leq l$, then we estimate the following expression as in the previous case,

$$\left| \int_{a}^{b} \hat{y}^{(l)} \chi^{(l+1-\nu)} \hat{y}^{(\nu)} dx \right| \leq \|\chi\|_{C^{l+1}([a;b];\mathbb{R})} \|\hat{y}\|_{H^{l}((a;b);\mathbb{R})}^{2}$$

It remains to estimate

$$\begin{split} \left| \int_{a}^{b} \hat{y}^{(l)} \chi \hat{y}^{(l+1)} dx \right| &\leq \left| \chi \frac{\left(\hat{y}^{(l)} \right)^{2}}{2} \right|_{a}^{b} \right| + \left| \int_{a}^{b} \frac{\left(\hat{y}^{(l)} \right)^{2}}{2} \chi' dx \right| \leq \\ &\leq \left| \chi \frac{\left(\hat{y}^{(l)} \right)^{2}}{2} \right|_{a}^{b} \right| + \frac{1}{2} \| \chi \|_{C^{1}([a;b];\mathbb{R})} \| \hat{y} \|_{H^{l}((a;b);\mathbb{R})}^{2} \end{split}$$

Hence, in both cases, we obtain the desired estimate.

Remark 7. One can consider an operator in $L^p((a; b); \mathbb{R}), p > 1$, similar to the one of example 13. Really, let

$$A(y) = \frac{\frac{d}{dx} \left(|y|^p y^{(\sigma)} \right)}{|y|^{p-1} \operatorname{sgn}(y)} = py' y^{(\sigma)} + yy^{(\sigma+1)},$$

where $\sigma \ge 0$, and if p is not an even integer, $\left\lceil \frac{\sigma}{2} \right\rceil \le p-1$.

With Faà di Bruno's formula, one can show that -A(y) is (p, ψ) submonotone w.r.t. $W^{\left\lceil \frac{\sigma}{2} \right\rceil, p}((a; b); \mathbb{R})$ norm, that is,

$$\langle A(y_1) - A(y_2), \hat{y} \rangle_p \le \psi(\hat{y}) + \Lambda(y_1, y_2) \| \hat{y} \|_{W^{\left\lceil \frac{\sigma}{2} \right\rceil, p}((a;b);\mathbb{R})}^p, \ \hat{y} = y_1 - y_2.$$

However, for the case $p \neq 2$, the (p, ψ) -power of the differentiation operator (as in example 12) cannot be built. In other words, we cannot connect $W^{\left\lceil \frac{\sigma}{2} \right\rceil, p}((a; b); \mathbb{R})$ norm with some (p, ψ) -powered operator U. Alternatively, we can try to consider operators U_1 and U_2 such that

$$\left\langle U_1\left(\frac{dU_2(y)}{dt}, y\right), y\right\rangle_p = \psi(\dots) + \left\langle \frac{dU^{\frac{1}{p}}(y)}{dt}, U^{\frac{1}{p}}(y)\right\rangle_p$$

however, such constructions lead us to the nonlinear and artificial operators. Really, one can check that even for an operator $U^{\frac{1}{p}} = \frac{\partial^{\sigma}}{\partial x^{\sigma}}$, in $L^{p}((a;b);\mathbb{R})$ space, we have

$$\left\langle \frac{\frac{d}{dt} \frac{\partial^{\sigma}}{\partial x^{\sigma}} (\operatorname{sgn}(y^{(\sigma)}) | y^{(\sigma)} | p^{-1})}{(-1)^{\sigma} (p-1) | y | p^{-2}}, y \right\rangle_{p} = \psi \left(y, \dots, y^{(\sigma)}, \frac{dy}{dt}, \dots, \frac{dy^{(\sigma)}}{dt} \right) + \left\langle \frac{dy^{(\sigma)}}{dt}, y^{(\sigma)} \right\rangle_{p}$$

2.4 Coercive operators

The last type of equations we consider is elliptic (4). In [25], the authors considered similar equations, and for this reason, they took an operator A to have a conditional Lipschitz inverse operator. Dealing with a \mathbb{R} -smooth Banach space, we consider instead an extended version. Precisely, we define the following type of operators. Again, we consider the case of the whole domain only.

Definition 12. Let X be a \mathbb{R} -smooth Banach space, $\mathcal{D}(A) \subset X$ be a subspace. We will say that an operator $A: \mathcal{D}(A) \to X$ is (p, ψ) -coercive, for some $\psi: \mathcal{D}(A) \times \mathcal{D}(A) \to \mathbb{R}$ and p > 1, if it is true that

$$\|\chi - y\|^p \le \psi(\chi, y) + \Lambda(\chi, y) \langle A(\chi) - A(y), \chi - y \rangle_p$$

for every $\chi, y \in \mathcal{D}(A)$, and some $\Lambda(\chi, y) \ge 0$.

Again, we can consider the following extention of (p, ψ) -coercivity to a positive linear combination and a closure. **Statement 3.** Let X be a \mathbb{R} -smooth Banach space. The following two properties hold. 1. If $A_{1,2}$ are $(p, \psi_{1,2})$ -coercive, then

$$A := s_1 A_1 + s_2 A_2$$

- is (p, ψ) -coercive for $\psi = s_1\psi_1 + s_2\psi_2$ and every $s_{1,2} \ge 0$.
- 2. Moreover, let $D(\|\cdot\|)(\cdot,\chi)$ be continuous on the ring $\{y \in X \mid r_0 < \|y\| < r_1\}$ for every $0 < r_0 < r_1$ and $\chi \in X$. If A is (p, ψ) -coercive, there exists a closure \overline{A} , and ψ , Λ are continuous w.r.t. \overline{A} -norm, then \overline{A} is (p, ψ) -coercive.

It is natural to consider an elliptic operator. For simplicity, we deal with its the first term only. **Example 14.** Let $\Omega \subset \mathbb{R}^m$ be a Lipschitz domain, $\Gamma = \partial \Omega$, $X = L^2(\Omega; \mathbb{R})$. Also, let for every $x \in \Omega$ and some $\varepsilon > 0$, matrix $Q(x) \in \mathbb{R}^{m \times m}$ satisfy

$$Q^T(x) = Q(x) \text{ and } Q(x)\xi \cdot \xi \ge \varepsilon \|\xi\|_{\mathbb{R}^m}^2, \ \xi \in \mathbb{R}^m.$$

If $Q_{ij} \in L^{\infty}(\Omega; \mathbb{R})$, for every $1 \leq i, j \leq m$, then an elliptic operator

$$Ay = -\nabla \cdot (Q(x)\nabla y)$$

is $(2, \psi)$ -coercive.

Really, let $\hat{y} = y_1 - y_2$. Then

$$\begin{split} \langle A\hat{y}, \hat{y} \rangle_2 &= \int_{\Omega} \left(-\nabla \cdot \left(Q(x) \nabla \hat{y} \right) \right) \hat{y} dx = -\int_{\Gamma} \left(\left(Q(x) \nabla \hat{y} \right) \cdot \mathbf{n}_{\Gamma} \right) \hat{y} dx + \int_{\Omega} \nabla \hat{y} \cdot Q(x) \nabla \hat{y} dx \geq \\ &= -\int_{\Gamma} \left(\left(Q(x) \nabla \hat{y} \right) \cdot \mathbf{n}_{\Gamma} \right) \hat{y} dx + \varepsilon \| \nabla \hat{y} \|_{L^2(\Omega; \mathbb{R}^m)}^2 \end{split}$$

With the Poincare inequality, for some q > 0,

$$\begin{split} \|\hat{y}\|_{L^{2}(\Omega;\mathbb{R})}^{2} &\leq 2\left\|\hat{y} - \frac{1}{|\Omega|}\int_{\Omega}\hat{y}dx\right\|_{L^{2}(\Omega;\mathbb{R})}^{2} + \frac{2}{|\Omega|}\left|\int_{\Omega}\hat{y}dx\right|^{2} \leq q\|\nabla\hat{y}\|_{L^{2}(\Omega;\mathbb{R}^{m})}^{2} + \frac{2}{|\Omega|}\left|\int_{\Omega}\hat{y}dx\right|^{2} \leq \frac{2}{|\Omega|}\left|\int_{\Omega}\hat{y}dx\right|^{2} + \frac{q}{\varepsilon} + \int_{\Gamma}\left(\left(Q(x)\nabla\hat{y}\right)\cdot\mathbf{n}_{\Gamma}\right)\hat{y}dx + \frac{q}{\varepsilon}\langle A\hat{y},\hat{y}\rangle_{2}. \end{split}$$

Here $\left|\int_{\Omega} \hat{y} dx\right|^2$ is subordinate to the constant mean value condition. Let us note, that in one-dimensional case, we can take Cauchy boundary conditions instead, as following inequality holds

$$|\hat{y}\|_{L^{2}((a;b);\mathbb{R})}^{2} \leq (b-a)|\hat{y}(a) + \hat{y}(b)|^{2} + (b-a)^{2}\|\hat{y}'\|_{L^{2}((a;b);\mathbb{R})}^{2}$$

Example 15. Let $\Omega \subset \mathbb{R}^m$, $\Gamma = \partial \Omega$, $X = L^p(\Omega; \mathbb{R})$, $p \ge 2$. Let

$$A_0(y) = -\nabla \cdot \left(|\nabla y|^{p-2} \nabla y \right)$$

be a minus p-Laplace operator, and

$$A_1(y) = |y|^{p-2}y.$$

Then

$$A(y) = A_0(y) + q_1 A_1(y) + q_2 y,$$

is (p, ψ) -coercive, for $q_1 \ge 0$, $q_2 > 0$. In examples 9 and 10, we showed that

$$\langle A_0(y_1) - A_0(y_2), y_1 - y_2 \rangle_p \ge -\int_{\Gamma} |\hat{y}|^{p-1} \operatorname{sgn}(\hat{y}) \left(\left(|\nabla y_1|^{p-2} \nabla y_1 - |\nabla y_2|^{p-2} \nabla y_2 \right) \cdot \mathbf{n}_{\Gamma} \right) d\Gamma,$$

and

$$\langle A_1(y_1) - A_1(y_2), y_1 - y_2 \rangle_p \ge 0,$$

where $\hat{y} = y_1 - y_2$. Hence,

$$\begin{split} \|y_1 - y_2\|^p &\leq \frac{1}{q_2} \int_{\Gamma} |\hat{y}|^{p-1} \operatorname{sgn}(\hat{y}) \left(\left(|\nabla y_1|^{p-2} \nabla y_1 - |\nabla y_2|^{p-2} \nabla y_2 \right) \cdot \mathbf{n}_{\Gamma} \right) d\Gamma + \\ &+ \frac{1}{q_2} \langle A(y_1) - A(y_2), y_1 - y_2 \rangle_p \end{split}$$

3 PINN's error estimation

In this section, we obtain PINN's error estimates for PDEs' types (1)–(4). Also, in the case of semilinear parabolic-type equations, we compare the estimation with an approach in [13].

Let us note that we describe in detail only the case of parabolic-type equations (1). In other cases, for simplicity, we omit some conditions. For instance, we give an estimate based on training error for the parabolic-type equations (1) only. Also, we consider the case of submonotonicity/subordination on a subspace for this type only. For other cases, we deal with the whole domain.

3.1 Parabolic-type equation

Consider the problem (9) in a \mathbb{R} -smooth Banach space X

$$\frac{dw}{dt} = A(t)(w)$$

$$w\Big|_{t=0} = w_0$$

$$B(t)(w) = h_b$$
(9)

where $A(t) : \mathcal{D}(A(t)) \to X$ and $B(t) : \mathcal{D}(B(t)) \to Y$ are nonlinear operators, $\mathcal{D}(A(t)) \subset \mathcal{D}(B(t)) \subset X$ are subspaces, Y is a Banach space. We also consider the following subspace of $\mathcal{D}(A(t))$

$$\mathcal{M}(A(t), B(t)) := (B(t))^{-1}(h_b(t)) \cap \mathcal{D}(A(t))$$

$$\tag{10}$$

For a solution of 9, we assume $w \in C(\overline{I}; X) \cap C^1(I; X)$ and $w(t) \in \mathcal{M}(A(t), B(t))$ for $t \in I$.

Let us consider a neural network w_{θ} with parameter θ , approximating solution w of (9), and the following PINN residuals

$$\mathcal{R}_{eq} = \frac{dw_{\theta}}{dt} - A(t)(w_{\theta}),$$

$$\mathcal{R}_{in} = w_{\theta}|_{t=0} - w_{0},$$

$$\mathcal{R}_{bn} = B(t)(w_{\theta}) - h_{b}.$$
(11)

We are interested in the following total error

$$\mathcal{E} := \|w_{\theta} - w\|_{L^q(I;X)}^q \tag{12}$$

Also, we need to consider approximating rules for norms of residuals (quadrature rules in particular).

1. Let $\mathcal{T}_{eq} \subset \overline{I} \times \mathbb{R}^m$, and let there exist $\mathcal{Q}_{N,eq} : L^p(I;X) \times \mathcal{T}_{eq}^N \to \mathbb{R}$ with

$$\left| \|w\|_{L^{p}(I;X)} - \mathcal{Q}_{N,eq}(w,(t_{1},x_{1}),\ldots,(t_{N},x_{N})) \right| \leq \beta_{eq}^{*}(w) N^{-\alpha_{eq}},$$

for some
$$\beta_{eq}^*(w) \ge 0$$
, $\alpha_{eq} > 0$ and every $N \in \mathbb{N}$, $w \in L^p(I; X)$, and $\{(t_i, x_i)\}_{i=1}^N \subset \mathcal{T}_{eq}$

2. Let $\mathcal{T}_{in} \subset \mathbb{R}^m$, and let there exist $\mathcal{Q}_{N,in} : X \times \mathcal{T}_{in}^N \to \mathbb{R}$ with

$$|||y||_X - \mathcal{Q}_{N,in}(y, x_1, \dots, x_N)| \le \beta_{in}^*(y) N^{-\alpha_{in}}$$

for some $\beta_{in}^*(y) \ge 0$, $\alpha_{in} > 0$ and every $N \in \mathbb{N}$, $y \in X$, and $\{x_i\}_{i=1}^N \subset \mathcal{T}_{in}$.

3. Let $\mathcal{T}_{bn} \subset \overline{I}$, and let there exist $\mathcal{Q}_{N,bn} : L^1(I; \mathbb{R}) \times \mathcal{T}_{bn}^N \to \mathbb{R}$ with

$$\left| \|g\|_{L^1(I;\mathbb{R})} - \mathcal{Q}_{N,bn}(g,t_1,\ldots,t_N) \right| \le \beta_{bn}^*(g) N^{-\alpha_{bn}}$$

for some $\beta_{bn}^*(g) \ge 0$, $\alpha_{bn} > 0$ and every $N \in \mathbb{N}$, $g \in L^1(I; \mathbb{R})$, and $\{t_i\}_{i=1}^N \subset \mathcal{T}_{bn}$.

Given training sets $\{(t_i, x_i)\}_{i=1}^{N_{eq}} \subset \mathcal{T}_{eq}, \{x_i\}_{i=1}^{N_{in}} \subset \mathcal{T}_{in}, \{t_i\}_{i=1}^{N_{bn}} \subset \mathcal{T}_{bn}$, we have the following training errors

$$\begin{aligned} \mathcal{E}_{T,eq} &= \mathcal{Q}_{N_{eq},eq}(\mathcal{R}_{eq},(t_1,x_1),\ldots,(t_{N_{eq}},x_{N_{eq}})),\\ \mathcal{E}_{T,in} &= \mathcal{Q}_{N_{in},in}(\mathcal{R}_{in},x_1,\ldots,x_{N_{in}}),\\ \mathcal{E}_{T,bn} &= \mathcal{Q}_{N_{bn},bn}(\rho(\mathcal{R}_{bn}),t_1,\ldots,t_{N_{bn}}) \end{aligned}$$

Theorem 1. Let us given a solution w of problem (9), and a neural network w_{θ} with residuals (11) and total error (12). Let, moreover, -A(t) be $(p, \psi(t))$ -submonotone on $\mathcal{M}(A(t), B(t))$, with $\psi(t)$ subordinate to B(t) on $\mathcal{M}(A(t), B(t))$, for every $t \in \overline{I}$. ($\mathcal{M}(A(t), B(t))$ is defined in (10)). Furthermore, let respective $\gamma(\cdot, w_{\theta}(\cdot), w(\cdot)) \in C(\overline{I}; \mathbb{R})$, $\rho(t) \equiv \rho$, and $\Lambda(\cdot, w_{\theta}(\cdot), w(\cdot)) \in L^{1}(I; \mathbb{R})$. Then

$$\mathcal{E} \leq \mathcal{C}^{\frac{q}{p}}\left(\frac{p(e^{\frac{q(p-1)T}{p}}-1)}{q(p-1)}\right) e^{q\|\Lambda(\cdot,w_{\theta}(\cdot),w(\cdot))\|_{L^{1}(I;\mathbb{R})}},$$

where

$$\mathcal{C} = \|\mathcal{R}_{in}\|^p + \|\mathcal{R}_{eq}\|_{L^p(I;X)}^p + p\|\gamma(\cdot, w_\theta(\cdot), w(\cdot))\|_{C(\overline{I};\mathbb{R})}\|\rho(\mathcal{R}_{bn})\|_{L^1(I;\mathbb{R})}$$

Proof. With a denotation $\hat{w} := w_{\theta} - w$ the total error and the residuals can be expressed as

$$\mathcal{E} = \|\hat{w}\|_{L^q(I;X)}^q \quad \text{and} \quad \begin{aligned} \mathcal{R}_{eq} &= \frac{dw}{dt} - A(t)(w_\theta) + A(t)(w) \\ \mathcal{R}_{in} &= \hat{w}|_{t=0}, \\ \mathcal{R}_{bn} &= B(t)(w_\theta) - B(t)(w). \end{aligned}$$

With (7), we have

$$\frac{d}{dt} \|\hat{w}\|^p = p \left\langle \frac{d\hat{w}}{dt}, \hat{w} \right\rangle_p = p \left\langle \mathcal{R}_{eq}, \hat{w} \right\rangle_p + p \left\langle A(t)(w_\theta) - A(t)(w), \hat{w} \right\rangle_p$$

With properties of *p*-form and the Young inequality,

$$p\left|\left\langle \mathcal{R}_{eq}, \hat{w} \right\rangle_{p}\right| \leq p \|\mathcal{R}_{eq}\| \|\hat{w}\|^{p-1} \leq \|\mathcal{R}_{eq}\|^{p} + (p-1)\|\hat{w}\|^{p}$$

Since -A(t) is $(p, \psi(t))$ -submonotone on $\mathcal{M}(A(t), B(t)), \psi(t)$ is subordinate to B(t) on $\mathcal{M}(A(t), B(t))$, then

$$\begin{aligned} \langle A(t)(w_{\theta}) - A(t)(w), \hat{w} \rangle_{p} &\leq \psi(t, w(t), w_{\theta}(t)) + \Lambda(t, w_{\theta}(t), w(t)) \|\hat{w}\|^{p} \leq \\ &\leq \gamma(t, w_{\theta}(t), w(t)) \rho(B(t)(w_{\theta}) - B(t)(w)) + \Lambda(t, w_{\theta}(t), w(t)) \|\hat{w}\|^{p} = \\ &= \gamma(t, w_{\theta}(t), w(t)) \rho(\mathcal{R}_{bn}) + \Lambda(t, w_{\theta}(t), w(t)) \|\hat{w}\|^{p} \end{aligned}$$

Hence,

$$\frac{a}{dt}\|\hat{w}\|^p \le \|\mathcal{R}_{eq}\|^p + p\gamma(t, w_{\theta}(t), w(t))\rho(\mathcal{R}_{bn}) + (p-1+p\Lambda(t, w_{\theta}(t), w(t)))\|\hat{w}\|^p$$

Integrating from 0 to $t \leq T$,

$$\begin{aligned} \|\hat{w}\|^{p} &\leq \|\mathcal{R}_{in}\|^{p} + \|\mathcal{R}_{eq}\|_{L^{p}(I;X)}^{p} + p\|\gamma(\cdot, w_{\theta}(\cdot), w(\cdot))\|_{C(\overline{I};\mathbb{R})} \|\rho(\mathcal{R}_{bn})\|_{L^{1}(I;\mathbb{R})} + \\ &+ \int_{0}^{t} (p - 1 + p\Lambda(t, w_{\theta}(t), w(t))) \|w\|^{p} d\tau \end{aligned}$$

With the Grönwall-Bellman lemma,

$$\|\hat{w}\|^p \leq \mathcal{C}e^{(p-1)t+p\int_0^t (\Lambda(t,w_\theta(t),w(t)))d\tau}$$

where

$$= \|\mathcal{R}_{in}\|^{p} + \|\mathcal{R}_{eq}\|_{L^{p}(I;X)}^{p} + p\|\gamma(\cdot, w_{\theta}(\cdot), w(\cdot))\|_{C(\overline{I};\mathbb{R})}\|\rho(\mathcal{R}_{bn})\|_{L^{1}(I;\mathbb{R})}$$

Taking q-powered L^q -norm on $t, 1 \le q < \infty$,

С

$$\mathcal{E} = \|\hat{w}\|_{L^{q}(I;X)}^{q} \leq \mathcal{C}^{\frac{q}{p}} \left(\frac{p(e^{\frac{q(p-1)T}{p}} - 1)}{q(p-1)} \right) e^{q\|\Lambda(\cdot, w_{\theta}(\cdot), w(\cdot))\|_{L^{1}(I;\mathbb{R})}}$$

Remark 8. In theorem 1, instead of $L^q(I; X)$ norm, one can take another one. Also, we assume that solution w and network w_θ are sufficiently regular to have finite L^q -norm.

Corollary 1. With conditions of theorem 1 and training errors, we have

$$\mathcal{E} \leq \tilde{\mathcal{C}}^{\frac{q}{p}} \left(\frac{p(e^{\frac{q(p-1)T}{p}} - 1)}{q(p-1)} \right) e^{q \|\Lambda(\cdot, w_{\theta}(\cdot), w(\cdot))\|_{L^{1}(I;\mathbb{R})}},$$

where

$$\tilde{\mathcal{C}} = \left(\mathcal{E}_{T,in} + \beta_{in}^*(w, w_{\theta})N^{-\alpha_{in}}\right)^p + \left(\mathcal{E}_{T,eq} + \beta_{eq}^*(w, w_{\theta})N^{-\alpha_{eq}}\right)_{L^p(I;X)}^p + p\|\gamma(\cdot, w_{\theta}(\cdot), w(\cdot))\|_{C(\overline{I};\mathbb{R})} \left(\mathcal{E}_{T,bn} + \beta_{bn}^*(w, w_{\theta})N^{-\alpha_{bn}}\right)$$

Remark 9. Approximation (quadrature) constants depend on some norm of residuals.

3.2 Parabolic-type equation, non-smooth Banach space

We briefly describe an extension of the approach presented in [13] to the semilinear problem 13 in arbitrary Banach space X, and compare obtained estimate with the one of the previous subsection.

$$\frac{dw}{dt} = Aw + F(t)(w)$$

$$w\Big|_{t=0} = w_0$$

$$B(w) = 0$$
(13)

where $A : \mathcal{D}(A) \to X$ and $B : \mathcal{D}(B) \to Y$ are linear operators, $\mathcal{D}(A) \subset \mathcal{D}(B) \subset X$ are subspaces, Y is a Banach space. Again, we consider the following subspace of $\mathcal{D}(A)$

$$\mathcal{M}(A,B) := \operatorname{Ker}(B) \cap \mathcal{D}(A).$$

Residuals are

$$\mathcal{R}_{eq} = \frac{dw_{\theta}}{dt} - Aw_{\theta} - F(t)(w_{\theta}),$$

$$\mathcal{R}_{in} = w_{\theta}|_{t=0} - w_0,$$

$$\mathcal{R}_{bn} = B(w_{\theta}).$$
(14)

Remark 10. For main results in the operator semigroup theory, see for instance, [28]. For generator A of a strongly continuous semigroup V(t), -A is "submonotone" in the following sense. In a \mathbb{R} -smooth Banach space, an operator of form $-(A_0 + \omega I)$, where A_0 is dissipative and $\omega \in \mathbb{R}$, is submonotone. Since A is a generator of strongly continuous semigroup, then for some $M \ge 1$ and $\omega \in \mathbb{R}$, it is true that

$$\|(sI - A)^{-1}\| \le \frac{M}{s - \omega}, \ \forall s > \omega$$

Then, for $A_0 := A - \omega I$, we have

$$||(sI - A_0)^{-1}|| \le \frac{M}{s}, \, \forall s > 0.$$

There exists an equivalent norm $\|\cdot\|_V$ for which

$$||(sI - A_0)^{-1}||_V \le \frac{1}{s}, \ \forall s > 0,$$

that is, A_0 is dissipative (in the meaning similar to the Banach space case).

Returning to problem (13), we have the following result.

Theorem 2. Let us given a solution w of problem (13), and a neural network w_{θ} with residuals (14) and total error (12). Let $A\Big|_{\mathcal{M}(A,B)}$ be a generator of a semigroup V(t), $||V(t)|| \leq Me^{\omega t}$. Let B be right invertible, and for a right inverse Θ , let $\left(A\Big|_{\mathcal{M}(A,B)}\right) \cdot \Theta$ be bounded. Also, let F(t) be conditionally Lipschitz for every $t \in \overline{I}$ and $\Lambda(\cdot, w_{\theta}(\cdot), w(\cdot)) \in L^{1}(I; \mathbb{R})$. Then we have an estimate

$$\mathcal{E} \le \mathcal{C} \int_0^T e^{\frac{M \|\Lambda(\cdot, w_\theta(\cdot), w(\cdot))\|_{L^1(I;\mathbb{R})} e^{\omega t}}{\omega}} dt$$
(15)

where

$$\mathcal{C} = \|\Theta\mathcal{R}_{bn}\| + Me^{\omega T} \|\mathcal{R}_{in}\| + Me^{\omega T} \|\Theta\mathcal{R}_{bn}(0)\| + Me^{\omega T} \|\mathcal{R}_{eq}\|_{L^{1}(I;X)} + Me^{\omega T} \|A\Theta\| \|\mathcal{R}_{bn}\|_{L^{1}(I;X)} + Me^{\omega T} \left\|\frac{d\Theta\mathcal{R}_{bn}}{dt}\right\|_{L^{1}(I;X)}$$

Proof. Given fixed w and w_{θ} , let again $\hat{w} = w_{\theta} - w$. Then $\tilde{w} := \hat{w} - \Theta \mathcal{R}_{bn} \in \mathcal{M}(A, B)$ is a solution to the following problem

$$\frac{d\tilde{w}}{dt} = A\tilde{w} + z(t),$$

$$\tilde{w}|_{t=0} = \mathcal{R}_{in} - \Theta \mathcal{R}_{bn}(0)$$

where

$$z = \mathcal{R}_{eq} + F(t)(w_{\theta}) - F(t)(w) + A\Theta \mathcal{R}_{bn} - \frac{d\Theta \mathcal{R}_{bn}}{dt}$$

Then \tilde{w} is also a "mild" solution, that is, it satisfies the following integral equation

$$\tilde{w}(t) = V(t)\tilde{w}|_{t=0} + \int_0^t V(t-\tau)z(\tau)d\tau$$

Hence,

$$\begin{split} \|\hat{w}\| &\leq \|\Theta\mathcal{R}_{bn}\| + Me^{\omega T} \|\mathcal{R}_{in}\| + Me^{\omega T} \|\Theta\mathcal{R}_{bn}(0)\| + Me^{\omega T} \|\mathcal{R}_{eq}\|_{L^{1}(I;X)} + \\ &+ Me^{\omega T} \|A\Theta\mathcal{R}_{bn}\|_{L^{1}(I;X)} + Me^{\omega T} \left\|\frac{d\Theta\mathcal{R}_{bn}}{dt}\right\|_{L^{1}(I;X)} + \\ &+ M \int_{0}^{t} \Lambda(t, w_{\theta}(t), w(t))e^{\omega(t-\tau)} \|\hat{w}\| d\tau \leq \|\Theta\mathcal{R}_{bn}\| + Me^{\omega T} \|\mathcal{R}_{in}\| + Me^{\omega T} \|\Theta\mathcal{R}_{bn}(0)\| + \\ &+ Me^{\omega T} \|\mathcal{R}_{eq}\|_{L^{1}(I;X)} + Me^{\omega T} \|A\Theta\| \|\mathcal{R}_{bn}\|_{L^{1}(I;X)} + Me^{\omega T} \left\|\frac{d\Theta\mathcal{R}_{bn}}{dt}\right\|_{L^{1}(I;X)} + \\ &+ M \int_{0}^{t} \Lambda(t, w_{\theta}(t), w(t))e^{\omega(t-\tau)} \|\hat{w}\| d\tau = \mathcal{C} + M \int_{0}^{t} \Lambda(t, w_{\theta}(t), w(t))e^{\omega(t-\tau)} \|\hat{w}\| d\tau, \end{split}$$

for a.e. $t \in \overline{I}$, where

$$\mathcal{C} = \|\Theta\mathcal{R}_{bn}\| + Me^{\omega T} \|\mathcal{R}_{in}\| + Me^{\omega T} \|\Theta\mathcal{R}_{bn}(0)\| + Me^{\omega T} \|\mathcal{R}_{eq}\|_{L^1(I;X)} + Me^{\omega T} \|A\Theta\| \|\mathcal{R}_{bn}\|_{L^1(I;X)} + Me^{\omega T} \left\|\frac{d\Theta\mathcal{R}_{bn}}{dt}\right\|_{L^1(I;X)}$$

With the Grönwall-Bellman lemma,

$$\|\hat{w}\| \leq \mathcal{C}e^{\int_0^t \Lambda(t, w_\theta(t), w(t))e^{\omega(t-\tau)}d\tau} \leq \mathcal{C}e^{\frac{M\|\Lambda(\cdot, w_\theta(\cdot), w(\cdot)\|_{L^1(I;\mathbb{R})}e^{\omega t}}{\omega}}$$

for a.e. $t \in \overline{I}$.

Finally,

$$\mathcal{E} \leq \mathcal{C} \int_0^T e^{\frac{M \|\Lambda(\cdot, w_\theta(\cdot), w(\cdot))\|_{L^1(I;\mathbb{R})} e^{\omega t}}{\omega}} dt$$

Remark 11. Let us compare this estimate with the one of theorem 1. (15) contains $W^{1,1}(I;Y)$ norm of the additional conditions residual and a double exponent on T if the semigroup is not uniformly bounded.

The assumption of right invertibility is natural for "good" additional conditions, as the following example demonstrates. **Example 16.** Let $y \in C^1([a;b];\mathbb{R})$ and $By = (y(b) - y(a), y'(b) - y'(a))^T \in \mathbb{R}^2$. Let us denote $f_1 := y(b) - y(a)$, $f_2 := y'(b) - y'(a)$. We need to construct such a linear operator $\Theta : \mathbb{R}^2 \to \mathcal{D}(B)$, that $A\Theta$ is bounded for some A. We just put Θ to be a polynomial of degree ≤ 2 , and

$$\Theta(f_1, f_2)(x) := \frac{f_2}{2(b-a)}x^2 + \frac{2f_1 - (b+a)f_2}{2(b-a)}x.$$

One can show that $B\Theta(f_1, f_2) = (f_1, f_2)^T$. Since $ran(\Theta)$ is finite dimensional, $A\Theta$ is bounded for every linear operator A.

3.3 Generalized parabolic-type equation

For simplicity, we deal with submonotonicity/subordination on the whole domain. Consider problem (16) in a \mathbb{R} -smooth Banach space X

$$\frac{dw}{dt} + \frac{d}{dt} \sum_{k=1}^{K} U_k w = A(t)(w)$$

$$w\Big|_{t=0} = w_0$$

$$B(t)(w) = h_{b,0}$$

$$B(t)(U_k w) = h_{k,b}, \ 1 \le k \le K,$$
(16)

where $A(t) : \mathcal{D}(A(t)) \to X$ and $B(t) : \mathcal{D}(B(t)) \to Y$ are nonlinear operators, $\mathcal{D}(A(t)) \subset \mathcal{D}(B(t)) \subset X$ are subspaces, Y is a Banach space. Also, U_k are (p, ψ_k) -powered, $\operatorname{Im}(U_k) \subset \mathcal{D}(B(t)), \mathcal{D}(A(t)) \subset \mathcal{D}(U_k) \subset \mathcal{D}\left(U_k^{\frac{1}{p}}\right)$, U_k and $U_k^{\frac{1}{p}}$ commute with $\frac{d}{dt}$, $U_k^{\frac{1}{p}}$ commute with $\Big|_{t=0}$, $1 \le k \le K$. For a solution of (16), we assume $w \in U_k^{\frac{1}{p}}$ $C(\overline{I}; X) \cap C^1(I; X).$

Consider a neural network w_{θ} with parameter θ , approximating solution w of (16), and the PINN residuals

$$\mathcal{R}_{eq} = \frac{dw_{\theta}}{dt} + \frac{d}{dt} \sum_{k=1}^{K} U_k w_{\theta} - A(t)(w_{\theta}),$$

$$\mathcal{R}_{in} = w_{\theta}|_{t=0} - w_0,$$

$$\mathcal{R}_{0,bn} = B(t)(w_{\theta}) - h_{b,0},$$

$$\mathcal{R}_{k,bn} = B(t)(U_k w_{\theta}) - h_{b,k}$$
(17)

Again, the total error is given by formula

$$\mathcal{E} := \|w_{\theta} - w\|_{L^q(I;X)}^q \tag{18}$$

Theorem 3. Let us given a solution w of problem (16), and a neural network w_{θ} with residuals (17) and total error (18). Let also -A(t) be a $(p, \psi(t))$ -submonotone w.r.t. norm $\|\cdot\|_{\sum U_{k}^{\frac{1}{p}}}$, with (p, ψ_{k}) -powered operators U_k , where ψ_k are subordinate to B(t), $\psi(t)$ is subordinate to B(t) w.r.t. U_1, \ldots, U_k , for every $t \in \overline{I}$. Also, let respective $\tilde{\gamma}_k \left(\frac{dw_\theta}{dt}(\cdot), \frac{dw}{dt}(\cdot), w_\theta(\cdot), w(\cdot)\right), \gamma_k(\cdot, w_\theta(\cdot), w(\cdot)) \in C(\overline{I}; \mathbb{R}), \ \rho_k(t) \equiv \rho_k, \ \tilde{\rho}_k(t) \equiv \tilde{\rho}_k$, and $\Lambda(\cdot, w_\theta(\cdot), w(\cdot)) \in L^1(I; \mathbb{R})$. Then we have an estimate

$$\mathcal{E} \leq \mathcal{C}^{\frac{q}{p}} \left(\frac{p(e^{\frac{q(p-1)T}{p}} - 1)}{q(p-1)} \right) e^{q \|\Lambda(\cdot, w_{\theta}(\cdot), w(\cdot))\|_{L^{1}(I;\mathbb{R})}},$$

where

$$\begin{aligned} \mathcal{C} &= \|\mathcal{R}_{in}\|^{p} + \sum_{k=1}^{K} \|U_{k}^{\frac{1}{p}} \mathcal{R}_{in}\|^{p} + \|\mathcal{R}_{eq}\|_{L^{p}(I;X)}^{p} + p\|\gamma_{0}(\cdot, w_{\theta}(\cdot), w(\cdot))\|_{C(\overline{I};\mathbb{R})} \|\rho_{0}\left(\mathcal{R}_{0,bn}\right)\|_{L^{1}(I;X)} + \\ &+ p \sum_{k=1}^{K} \|\gamma_{k}(\cdot, w_{\theta}(\cdot), w(\cdot))\|_{C(\overline{I};\mathbb{R})} \|\rho_{k}\left(\mathcal{R}_{k,bn}\right)\|_{L^{1}(I;X)} + \\ &+ p \left\|\tilde{\gamma}_{k}\left(\frac{dw_{\theta}}{dt}(\cdot), \frac{dw}{dt}(\cdot), w_{\theta}(\cdot), w(\cdot)\right)\right\|_{L^{1}(I;\mathbb{R})} \|\tilde{\rho}_{k}\left(\mathcal{R}_{0,bn}\right)\|_{L^{1}(I;X)} \end{aligned}$$

Proof. With a denotation $\hat{w} := w_{\theta} - w$ the total error and the residuals can be expressed as

$$\mathcal{R}_{eq} = \frac{d\hat{w}}{dt} + \frac{d}{dt} \sum_{k=1}^{K} U_k \hat{w} - A(t)(w_\theta) + A(t)(w),$$

$$\mathcal{E} = \|\hat{w}\|_{L^q(I;X)}^q \text{ and } \mathcal{R}_{in} = \hat{w}|_{t=0},$$

$$\mathcal{R}_{0,bn} = B(t)(w_\theta) - B(t)(w),$$

$$\mathcal{R}_{k,bn} = B(t)(U_k w_\theta) - B(t)(U_k w)$$

With (7) and (8), we have

$$\begin{aligned} \frac{d}{dt} \|\hat{w}\|^p + \sum_{k=1}^K \frac{d}{dt} \|U_k^{\frac{1}{p}} \hat{w}\|^p &= p \left\langle \frac{d\hat{w}}{dt}, \hat{w} \right\rangle_p + p \sum_{k=1}^K \left\langle U_k^{\frac{1}{p}} \frac{d\hat{w}}{dt}, U_k^{\frac{1}{p}} \hat{w} \right\rangle_p \\ &= p \left\langle \frac{d\hat{w}}{dt}, \hat{w} \right\rangle_p + p \sum_{k=1}^K \left\langle U_k \frac{d\hat{w}}{dt}, \hat{w} \right\rangle_p \\ &- p \sum_{k=1}^K \left[\psi_k \left(\frac{dw_\theta}{dt}(t), \frac{dw}{dt}(t), w_\theta(t), w(t) \right) \right] \\ &= p \left\langle \mathcal{R}_{eq}, \hat{w} \right\rangle_p + p \left\langle A(t)(w_\theta) - A(t)(w), \hat{w} \right\rangle_p \\ &- p \sum_{k=1}^K \left[\psi_k \left(\frac{dw_\theta}{dt}(t), \frac{dw}{dt}(t), w_\theta(t), w(t) \right) \right] \end{aligned}$$

Again,

$$p\left|\left\langle \mathcal{R}_{eq}, \hat{w} \right\rangle_{p}\right| \leq p \|\mathcal{R}_{eq}\| \|\hat{w}\|^{p-1} \leq \|\mathcal{R}_{eq}\|^{p} + (p-1)\|\hat{w}\|^{p}$$

Also,

$$\begin{split} \langle A(t)(w_{\theta}) - A(t)(w), \hat{w} \rangle_{p} &\leq \psi(t, w(t), w_{\theta}(t)) + \Lambda(t, w_{\theta}(t), w(t)) \left(\left\| \hat{w} \right\|^{p} + \sum_{k=1}^{K} \left\| U_{k}^{\frac{1}{p}} \hat{w} \right\|^{p} \right) \leq \\ &\leq \gamma_{0}(t, w_{\theta}(t), w(t)) \rho_{0}(B(t)(w_{\theta}) - B(t)(w)) + \sum_{k=1}^{K} \gamma_{k}(t, w_{\theta}(t), w(t)) \rho_{k}(B(t)(U_{k}w_{\theta}) - B(t)(U_{k}w)) + \\ &+ \Lambda(t, w_{\theta}(t), w(t)) \left(\left\| \hat{w} \right\|^{p} + \sum_{k=1}^{K} \left\| U_{k}^{\frac{1}{p}} \hat{w} \right\|^{p} \right) = \gamma_{0}(t, w_{\theta}(t), w(t)) \rho_{0}\left(\mathcal{R}_{0, bn}\right) + \sum_{k=1}^{K} \gamma_{k}(t, w_{\theta}(t), w(t)) \rho_{k}\left(\mathcal{R}_{k, bn}\right) + \\ &+ \Lambda(t, w_{\theta}(t), w(t)) \left(\left\| \hat{w} \right\|^{p} + \sum_{k=1}^{K} \left\| U_{k}^{\frac{1}{p}} \hat{w} \right\|^{p} \right) \end{split}$$

Moreover,

$$\left|\psi_k\left(\frac{dw_\theta}{dt}(t), \frac{dw}{dt}(t), w_\theta(t), w(t)\right)\right| \le \tilde{\gamma}_k\left(\frac{dw_\theta}{dt}(t), \frac{dw}{dt}(t), w_\theta(t), w(t)\right) \tilde{\rho}_k(B(t)(w_\theta) - B(t)(w)) = \tilde{\gamma}_k\left(\frac{dw_\theta}{dt}(t), \frac{dw}{dt}(t), w_\theta(t), w(t)\right) \tilde{\rho}_k\left(\mathcal{R}_{0,bn}\right)$$

Hence,

$$\begin{aligned} \frac{d}{dt} \|\hat{w}\|^{p} + \sum_{k=1}^{K} \frac{d}{dt} \|U_{k}^{\frac{1}{p}} \hat{w}\|^{p} &\leq \|\mathcal{R}_{eq}\|^{p} + p\gamma_{0}(t, w_{\theta}(t), w(t))\rho_{0}\left(\mathcal{R}_{0,bn}\right) + p\sum_{k=1}^{K} \gamma_{k}(t, w_{\theta}(t), w(t))\rho_{k}\left(\mathcal{R}_{k,bn}\right) + \\ &+ p\tilde{\gamma}_{k}\left(\frac{dw_{\theta}}{dt}(t), \frac{dw}{dt}(t), w_{\theta}(t), w(t)\right) \tilde{\rho}_{k}\left(\mathcal{R}_{0,bn}\right) + (p - 1 + p\Lambda(t, w_{\theta}(t), w(t)))\|\hat{w}\|^{p} + p\Lambda(t, w_{\theta}(t), w(t))\left(\sum_{k=1}^{K} \left\|U_{k}^{\frac{1}{p}} \hat{w}\right\|^{p}\right) \leq \\ &\leq \|\mathcal{R}_{eq}\|^{p} + p\gamma_{0}(t, w_{\theta}(t), w(t))\rho_{0}\left(\mathcal{R}_{0,bn}\right) + p\sum_{k=1}^{K} \gamma_{k}(t, w_{\theta}(t), w(t))\rho_{k}\left(\mathcal{R}_{k,bn}\right) + \\ &+ p\tilde{\gamma}_{k}\left(\frac{dw_{\theta}}{dt}(t), \frac{dw}{dt}(t), w_{\theta}(t), w(t)\right) \tilde{\rho}_{k}\left(\mathcal{R}_{0,bn}\right) + (p - 1 + p\Lambda(t, w_{\theta}(t), w(t)))\left(\left\|\hat{w}\|^{p} + \sum_{k=1}^{K} \left\|U_{k}^{\frac{1}{p}} \hat{w}\right\|^{p}\right) \end{aligned}$$

Integrating from 0 to $t \leq T$,

$$\|\hat{w}\|^{p} + \sum_{k=1}^{K} \|U_{k}^{\frac{1}{p}}\hat{w}\|^{p} \leq \mathcal{C} + \int_{0}^{t} (p-1+p\Lambda(t,w_{\theta}(t),w(t))) \left(\|\hat{w}\|^{p} + \sum_{k=1}^{K} \left\|U_{k}^{\frac{1}{p}}\hat{w}\right\|^{p} \right) d\tau,$$

where

$$\begin{aligned} \mathcal{C} &= \|\mathcal{R}_{in}\|^{p} + \sum_{k=1}^{K} \|U_{k}^{\frac{1}{p}} \mathcal{R}_{in}\|^{p} + \|\mathcal{R}_{eq}\|_{L^{p}(I;X)}^{p} + p\|\gamma_{0}(\cdot, w_{\theta}(\cdot), w(\cdot))\|_{C(\overline{I};\mathbb{R})} \|\rho_{0}\left(\mathcal{R}_{0,bn}\right)\|_{L^{1}(I;X)} + \\ &+ p \sum_{k=1}^{K} \|\gamma_{k}(\cdot, w_{\theta}(\cdot), w(\cdot))\|_{C(\overline{I};\mathbb{R})} \|\rho_{k}\left(\mathcal{R}_{k,bn}\right)\|_{L^{1}(I;X)} + \\ &+ p \left\|\tilde{\gamma}_{k}\left(\frac{dw_{\theta}}{dt}(\cdot), \frac{dw}{dt}(\cdot), w_{\theta}(\cdot), w(\cdot)\right)\right\|_{L^{1}(I;\mathbb{R})} \|\tilde{\rho}_{k}\left(\mathcal{R}_{0,bn}\right)\|_{L^{1}(I;X)} \end{aligned}$$

The rest of the proof is similar to the one of theorem 1.

3.4 Hyperbolic-type equation

We apply the classical technique to reduce hyperbolic-type equations to the parabolic-type (see, for instance, [4]). We consider the Hilbert space case only. One can deal with a smooth \mathbb{R} -smooth Banach space, but, as stated in remark 7, it seems unuseful in practice. For simplicity, we consider exactly one $(2, \psi)$ -powered operator and with submonotonicity/subordination on the whole domain. Let us given problem (19) in a Hilbert space X

$$\frac{d^2w}{dt^2} = -Uw + F(t)(w) + A(t)\left(\frac{dw}{dt}\right)$$

$$w\Big|_{t=0} = w_0, \ \frac{dw}{dt}\Big|_{t=0} = w_{t,0}$$

$$B_1(t)(w) = h_b, \ B_2(t)\left(\frac{dw}{dt}\right) = h_{t,b}$$
(19)

where $A(t): \mathcal{D}(A(t)) \to X$ and $B_{1,2}(t): \mathcal{D}(B_{1,2}(t)) \to Y_{1,2}$ are nonlinear operators, $\mathcal{D}(F(t)) \subset \mathcal{D}(B_1(t)) \subset X$ and $\mathcal{D}(A(t)) \subset \mathcal{D}(B_2(t)) \subset X$ are subspaces, $Y_{1,2}$ are Banach spaces. Also, U is $(2, \tilde{\psi})$ -powered, $\mathcal{D}(F(t)) \subset \mathcal{D}(U) \subset \mathcal{D}\left(U^{\frac{1}{2}}\right)$, $\mathcal{D}(A(t)) \subset \mathcal{D}\left(U^{\frac{1}{2}}\right)$, U and $U^{\frac{1}{2}}$ commute with $\frac{d}{dt}$, $U^{\frac{1}{2}}$ commute with $\Big|_{t=0}$. For a solution of (19), we assume $w \in C^1(\overline{I}; X) \cap C^2(I; X)$.

Consider a neural network w_{θ} with parameter θ , approximating solution w of (19), and the PINN residuals

$$\mathcal{R}_{eq} = \frac{d^2 w_{\theta}}{dt^2} + U w_{\theta} - F(t)(w_{\theta}) - A(t) \left(\frac{dw_{\theta}}{dt}\right)$$

$$\mathcal{R}_{in} = w_{\theta}\Big|_{t=0} - w_0$$

$$\mathcal{R}_{in,t} = \frac{dw_{\theta}}{dt}\Big|_{t=0} - w_{t,0}$$

$$\mathcal{R}_{bn} = B_1(t)(w_{\theta}) - h_b$$

$$\mathcal{R}_{bn,t} = B_2(t) \left(\frac{dw_{\theta}}{dt}\right) - h_{t,b}$$
(20)

Again, the total error is given by formula

$$\mathcal{E} := \|w_{\theta} - w\|_{L^q(I;X)}^q \tag{21}$$

Theorem 4. Let us given a solution w of problem (19), and a neural network w_{θ} with residuals (20) and total error (21). Let also -A(t) be $(2, \psi(t))$ -submonotone, F(t) be conditionally Lipschitz w.r.t. to $U^{\frac{1}{2}}$ -norm (see remark 12), with $(2, \tilde{\psi})$ -powered operator U, where $\tilde{\psi}$, $\psi(t)$ are subordinate to $B_2(t)$, for every $t \in \overline{I}$. Also, let respective $\tilde{\gamma}(w_{\theta}(\cdot), w(\cdot), \frac{dw_{\theta}}{dt}(\cdot), \frac{dw}{dt}(\cdot)), \gamma(\cdot, w_{\theta}(\cdot), w(\cdot)) \in C(\overline{I}; \mathbb{R}), \rho(t) \equiv \rho_k, \tilde{\rho}(t) \equiv \tilde{\rho}_k, \Lambda_A(\cdot, \frac{dw_{\theta}}{dt}(\cdot), \frac{dw}{dt}(\cdot)) \in L^1(I; \mathbb{R})$ and $\Lambda_F(\cdot, w_{\theta}(\cdot), w(\cdot)) \in L^2(I; \mathbb{R})$. Then we have an estimate

$$\mathcal{E} \leq \mathcal{C}^{\frac{q}{2}} \frac{(e^{qT}-1)}{q} e^{q\left(\left\|\Lambda_A\left(\cdot, \frac{dw_{\theta}}{dt}(\cdot), \frac{dw}{dt}(\cdot)\right)\right\|_{L^1(I;\mathbb{R})} + \left\|\Lambda_F\left(\cdot, w_{\theta}(\cdot), w(\cdot)\right)\right\|_{L^2(I;\mathbb{R})}^2\right)},$$

where

$$\begin{aligned} \mathcal{C} &= \|\mathcal{R}_{in}\|^2 + \|\mathcal{R}_{in,t}\|^2 + \|U^{\frac{1}{2}}\mathcal{R}_{in}\|^2 + \|\mathcal{R}_{eq}\|_{L^2(I;X)}^2 + \\ &+ 2\left\|\gamma\left(\cdot,\frac{dw_{\theta}}{dt}(\cdot),\frac{dw}{dt}(\cdot)\right)\right\|_{C(\overline{I};\mathbb{R})} \left\|\rho\left(\mathcal{R}_{bn,t}\right)\right\|_{L^1(I;X)} + \\ &+ 2\left\|\tilde{\gamma}\left(w_{\theta}(\cdot),w(\cdot),\frac{dw_{\theta}}{dt}(\cdot),\frac{dw}{dt}(\cdot)\right)\right\|_{C(\overline{I};\mathbb{R})} \|\tilde{\rho}\left(\mathcal{R}_{bn,t}\right)\|_{L^1(I;X)} \end{aligned}$$

Proof. With a denotation $\hat{w} := w_{\theta} - w$ the total error and the residuals can be expressed as

$$\mathcal{R}_{eq} = \frac{d^2 \hat{w}}{dt^2} + U \hat{w} - F(t)(w_\theta) + F(t)(w) - A(t) \left(\frac{dw_\theta}{dt}\right) + A(t) \left(\frac{dw}{dt}\right)$$
$$\mathcal{R}_{in} = \hat{w}\Big|_{t=0}$$
$$\mathcal{R}_{in,t} = \hat{w}\Big|_{t=0}$$
$$\mathcal{R}_{in,t} = \frac{d\hat{w}}{dt}\Big|_{t=0}$$
$$\mathcal{R}_{bn} = B_1(t)(w_\theta) - B_1(t)(w)$$
$$\mathcal{R}_{bn,t} = B_2(t) \left(\frac{dw_\theta}{dt}\right) - B_2(t) \left(\frac{dw}{dt}\right)$$

Let us consider $w_{\theta,1} = w_{\theta}$, $w_1 = w$, $\hat{w}_1 = \hat{w}$, $w_{\theta,2} = \frac{dw_{\theta}}{dt}$, $w_2 = \frac{dw}{dt}$, $\hat{w}_2 = \frac{d\hat{w}}{dt}$. Then we have a parabolic-type equation,

$$\begin{cases} \frac{d\hat{w}_1}{dt} = \hat{w}_2 \\ \frac{d\hat{w}_2}{dt} = \mathcal{R}_{eq} - U\hat{w}_1 + F(t)(w_{\theta,1}) - F(t)(w_1) + A(t)(w_{\theta,2}) - A(t)(w_2) \end{cases}$$

Hence, one can deal with the space $\tilde{X} = \mathcal{D}(U^{\frac{1}{2}}) \times X$ endowed the following norm

$$\left\| \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \right\|_{\tilde{X}}^2 := \|y_1\|^2 + \|y_2\|^2 + \|U^{\frac{1}{2}}y_1\|^2,$$

With (7) and (8), we have

$$\begin{split} \frac{d}{dt} \left\| \begin{pmatrix} \hat{w}_1 \\ \hat{w}_2 \end{pmatrix} \right\|_{\tilde{X}}^2 &= \frac{d}{dt} \| \hat{w}_1 \|^2 + \frac{d}{dt} \| \hat{w}_2 \|^2 + \frac{d}{dt} \| U^{\frac{1}{2}} \hat{w}_1 \|^2 = 2 \left\langle \frac{d\hat{w}_1}{dt}, \hat{w}_1 \right\rangle_2 + 2 \left\langle \frac{d\hat{w}_2}{dt}, \hat{w}_2 \right\rangle_2 + 2 \left\langle \frac{U^{\frac{1}{2}} d\hat{w}_1}{dt}, U^{\frac{1}{2}} \hat{w}_1 \right\rangle_2 = \\ &= 2 \langle \hat{w}_2, \hat{w}_1 \rangle_2 + 2 \langle \mathcal{R}_{eq}, \hat{w}_2 \rangle_2 - 2 \langle U \hat{w}_1, \hat{w}_2 \rangle_2 + 2 \langle F(t)(w_{\theta,1}) - F(t)(w_1), \hat{w}_2 \rangle_2 + 2 \langle A(t)(w_{\theta,2}) - A(t)(w_2), \hat{w}_2 \rangle_2 + \\ &+ 2 \langle U^{\frac{1}{2}} \hat{w}_1, U^{\frac{1}{2}} \hat{w}_2 \rangle_2 = 2 \langle \hat{w}_2, \hat{w}_1 \rangle_2 + 2 \langle \mathcal{R}_{eq}, \hat{w}_2 \rangle_2 + 2 \langle F(t)(w_{\theta,1}) - F(t)(w_1), \hat{w}_2 \rangle_2 + 2 \langle A(t)(w_{\theta,2}) - A(t)(w_2), \hat{w}_2 \rangle_2 - \\ &- 2 \tilde{\psi}(w_{1,\theta}(t), w_1(t), w_{2,\theta}(t), w_2(t)) \leq \| \hat{w}_1 \|^2 + 2 \| \hat{w}_2 \|^2 + \| \mathcal{R}_{eq} \|^2 + 2 \Lambda_F^2(t, w_{\theta,1}(t), w_1(t)) \left(\| \hat{w}_1 \|^2 + \| U^{\frac{1}{2}} \hat{w}_1 \|^2 \right) + \\ &+ 2 \psi(t, w_{2,\theta}(t), w_2(t)) + 2 \Lambda_A(t, w_{2,\theta}(t), w_2(t)) \| \hat{w}_2 \|^2 - 2 \tilde{\psi}(w_{1,\theta}(t), w_1(t), w_{2,\theta}(t), w_2(t)) \rangle \leq \\ &\leq \| \hat{w}_1 \|^2 + 2 \| \hat{w}_2 \|^2 + \| \mathcal{R}_{eq} \|^2 + 2 \Lambda_F^2(t, w_{\theta,1}(t), w_1(t)) \left(\| \hat{w}_1 \|^2 + \| U^{\frac{1}{2}} \hat{w}_1 \|^2 \right) \\ &+ 2 \gamma(t, w_{2,\theta}(t), w_2(t)) \rho(\mathcal{R}_{bn,t}) + 2 \Lambda_A(t, w_{2,\theta}(t), w_2(t)) \| \hat{w}_2 \|^2 + 2 \tilde{\gamma}(w_{1,\theta}(t), w_1(t), w_2(t), w_2(t)) \tilde{\rho}(\mathcal{R}_{bn,t}) \end{split}$$

Hence, after integrating from 0 to $t \leq T$, we obtain

$$\left\| \begin{pmatrix} \hat{w}_1 \\ \hat{w}_2 \end{pmatrix} \right\|_{\tilde{X}}^2 \le \mathcal{C} + 2 \int_0^t (1 + \Lambda_F^2(t, w_{\theta, 1}(t), w_1(t)) + \Lambda_A(t, w_{2, \theta}(t), w_2(t))) \left\| \begin{pmatrix} \hat{w}_1 \\ \hat{w}_2 \end{pmatrix} \right\|_{\tilde{X}}^2 d\tau$$

where

$$\begin{aligned} \mathcal{C} &= \|\mathcal{R}_{in}\|^2 + \|\mathcal{R}_{in,t}\|^2 + \|U^{\frac{1}{2}}\mathcal{R}_{in}\|^2 + \|\mathcal{R}_{eq}\|_{L^2(I;X)}^2 + 2\|\gamma(\cdot, w_{2,\theta}(\cdot), w_2(\cdot))\|_{C(\overline{I};\mathbb{R})}\|\rho(\mathcal{R}_{bn,t})\|_{L^1(I;X)} + \\ &+ 2\|\tilde{\gamma}(w_{1,\theta}(\cdot), w_1(\cdot), w_{2,\theta}(\cdot), w_2(\cdot))\|_{C(\overline{I};\mathbb{R})}\|\tilde{\rho}(\mathcal{R}_{bn,t})\|_{L^1(I;X)} \end{aligned}$$

The rest of the proof is similar to the one of theorem 1.

Remark 12. Conditionally Lipschitz continuity of F w.r.t. $U^{\frac{1}{2}}$ -norm stands for

$$\|F(\chi) - F(y)\| \le \Lambda_F(\chi, y) \left(\|\chi - y\|^2 + \|U^{\frac{1}{2}}\chi - U^{\frac{1}{2}}y\|^2 \right)^{\frac{1}{2}}$$

In fact, instead of this, one can consider a "weak $(2, \psi_F)$ -Lipschitz continuity w.r.t. $U^{\frac{1}{2}}$ -norm", that is

$$|\langle F(\chi_1) - F(y_1), \chi_2 - y_2 \rangle_2| \le \psi_F(\chi_1, y_1, \chi_2, y_2) + \Lambda_F^2(\chi_1, y_1, \chi_2, y_2) \left(\|\chi - y\|^2 + \left\| U^{\frac{1}{2}} \chi - U^{\frac{1}{2}} y \right\|^2 \right) + \|\chi_2 - y_2\|^2.$$

Remark 13. Let us note that obtained estimate does not depend on \mathcal{R}_{bn} . However, in practice, we can observe such dependence. For instance, let $U = \frac{\partial^2}{\partial x^2}$ in $L^2((a; b); \mathbb{R})$,

$$\left\langle -Uw, \frac{dw}{dt} \right\rangle_2 = \frac{\partial w}{\partial x} \frac{dw}{dt} \Big|_a^b + \left\langle U^{\frac{1}{2}}w, U^{\frac{1}{2}} \frac{dw}{dt} \right\rangle_2$$

Hence, if we have Neumann-type boundary conditions, then, substituting w with $\frac{dw}{dt}$, respective $\tilde{\psi}$ subordinates to such an operator.

3.5 Elliptic equation

Consider problem (22) in a \mathbb{R} -smooth Banach space X

$$\begin{aligned} A(y) &= f_{eq} \\ B(y) &= f_b, \end{aligned} \tag{22}$$

where $A : \mathcal{D}(A) \to X$ and $B : \mathcal{D}(B) \to Y$ are nonlinear operators, $\mathcal{D}(A) \subset \mathcal{D}(B) \subset X$ are subspaces, Y is a Banach space.

Let us consider a neural network y_{θ} with parameter θ , approximating solution y of (22) and the PINN residuals

$$\mathcal{R}_{eq} = A(y_{\theta}) - f_{eq},$$

$$\mathcal{R}_{bn} = B(y_{\theta}) - f_{b}.$$
(23)

Total error is

$$\mathcal{E} := \|y_{\theta} - y\|^{q}, \ q > 1.$$
(24)

Theorem 5. Let us given a solution y of problem (22), and a neural network y_{θ} with residuals (23) and total error (24). Let also A be (p, ψ) -coercive, with ψ subordinate to B. Then we have an estimate

$$\mathcal{E} \leq p^{\frac{q}{p}} \left(\gamma(y_{\theta}, y) \rho(\mathcal{R}_{bn}) + \frac{1}{p} \Lambda^{p}(y_{\theta}, y) \|\mathcal{R}_{eq}\|^{p} \right)^{\frac{q}{p}}$$

Proof. With a denotation $\hat{y} := y_{\theta} - y$ the total error and the residuals can be expressed as

$$\mathcal{E} = \|\hat{y}\|^q$$
 and $\begin{array}{c} \mathcal{R}_{eq} = A(y_{\theta}) - A(y), \\ \mathcal{R}_{bn} = B(y_{\theta}) - B(y) \end{array}$

With conditions of the theorem,

$$\begin{aligned} \|\hat{y}\|^{p} &\leq \psi(y_{\theta}, y) + \Lambda(y_{\theta}, y) \langle A(y_{\theta}) - A(y), \hat{y} \rangle_{p} = \psi(y_{\theta}, y) + \Lambda(y_{\theta}, y) \langle \mathcal{R}_{eq}, \hat{y} \rangle_{p} \leq \\ &\leq \gamma(y_{\theta}, y) \rho(\mathcal{R}_{bn}) + \Lambda(y_{\theta}, y) \|\mathcal{R}_{eq}\| \|\hat{y}\|^{p-1}. \end{aligned}$$

Then with the Young inequality,

$$\|\hat{y}\|^p \leq \gamma(y_{\theta}, y)\rho(\mathcal{R}_{bn}) + \frac{\Lambda^p(y_{\theta}, y)}{p} \|\mathcal{R}_{eq}\|^p + \frac{p-1}{p} \|\hat{y}\|^p.$$

Hence,

$$\mathcal{E} \le p^{\frac{q}{p}} \left(\gamma(y_{\theta}, y) \rho(\mathcal{R}_{bn}) + \frac{1}{p} \Lambda^{p}(y_{\theta}, y) \|\mathcal{R}_{eq}\|^{p} \right)^{\frac{q}{p}}.$$

4 PINN's residuals upper bound

In [5], the authors proved the existence of a two-layer tahn neural network approximation for a sufficiently regular solution of the Navier-Stokes equation. Also, they obtained L^2 upper bounds for the residuals depending on the neural network width. The core theorem underlying this result claims the existence of a tahn neural network, approximating a function from $H^{\sigma}(\Omega)$, where $\Omega \subset \mathbb{R}^m$ is an integer right parallelepiped and $\sigma \geq 3$. We obtain a similar theorem in $W^{\sigma,p}(\Omega)$.

Let $\Omega \subset \mathbb{R}^m$ be a convex bounded domain, and let $p \ge 1$. First, we need to obtain the Bramble-Hilbert lemma in $W^{\sigma,p}(\Omega)$, similar to the result of [36]. R. Verfürth in [36] noted that the core point for such extension is a Poincaré

inequality with a "good" constant in $W^{1,p}(\Omega)$ for zero mean value functions. However, for such functions (i.e. $\frac{1}{|\Omega|} \int_{\Omega} y dx = 0$), a constant is known only for one-dimensional cases [11]. We provide slightly another idea: the mean value in $L^2(\Omega)$ is a projection to the set of constant functions, and for a function with vanishing projection, there is a "good" Poincaré inequality. The projection operator is defined as follows.

Statement 4. Let $p \ge 1$, and $\Omega \subset \mathbb{R}^m$ be a convex bounded domain. Then there exists an operator $\mathcal{J}_p : L^p(\Omega; \mathbb{R}) \to \mathbb{R}$,

$$\mathcal{J}_p(y) := \begin{cases} \frac{1}{|\Omega|} \int_{\Omega} y dx, & 1 \le p < 2, \\ \arg\min_{s \in \mathbb{R}} \|y - s\|_{L^p(\Omega; \mathbb{R})}, & p \ge 2 \end{cases}$$

with the following properties:

$$egin{aligned} &\mathcal{J}_p(-y) = -\mathcal{J}_p(y), \ &\mathcal{J}_p(y-\mathcal{J}_p(y)) = 0. \end{aligned}$$

Proof. If $1 \le p < 2$, the result is clear. Let $p \ge 2$. Given fixed element $y \in L^p(\Omega; \mathbb{R})$, let us consider the following function

$$\varphi : \mathbb{R} \to \mathbb{R}, \ \varphi(s) := \|y - s\|_{L^p(\Omega;\mathbb{R})}$$

Then φ is a differentiable convex function. Since

$$\varphi(s) \ge |s| |\Omega|^{\frac{1}{p}} - ||y||$$

we have that $\varphi(+\infty) = \varphi(-\infty) = +\infty$, and, therefore, φ attends a global minimum. We have two possible cases.

If $y \equiv s_0$ for some $s_0 \in \mathbb{R}$, then we have an exactly one minimum $\varphi(s_0) = 0$. If $y \neq const$, then the equality condition in the Minkowski inequality implies that φ is strictly convex. Hence, φ attends its global minimum exactly at one point.

Thus, an operator

$$\mathcal{J}_p(y) := \arg\min_{s \in \mathbb{R}} \|y - s\|_{L^p(\Omega;\mathbb{R})}$$

is correctly defined.

Also, we have

$$\|-y-(-\mathcal{J}_p(y))\|_{L^p(\Omega;\mathbb{R})} = \|y-\mathcal{J}_p(y)\|_{L^p(\Omega;\mathbb{R})} \le \|-y-s\|_{L^p(\Omega;\mathbb{R})}$$

and

$$\|y - \mathcal{J}_p(y)\|_{L^p(\Omega;\mathbb{R})} \le \|y - \mathcal{J}_p(y) - s\|_{L^p(\Omega;\mathbb{R})}$$

for every $s \in \mathbb{R}$. Hence, \mathcal{J}_p has the desired properties.

We turn to the Poincaré inequality.

Lemma 2. (Poincaré inequality) For every $y \in W^{1,p}(\Omega; \mathbb{R})$ with $\mathcal{J}_p(y) = 0$, the following inequality holds

 $\|y\|_{L^p(\Omega;\mathbb{R})} \le \pi_p \operatorname{diam}(\Omega) \|\nabla y\|_{L^p(\Omega;\mathbb{R})},$

where

$$\pi_p = \begin{cases} \frac{1}{2}, & p = 1\\ \pi^{\frac{2}{p}-2}2^{1-\frac{2}{p}}, & 1 (25)$$

Proof. For p = 2, the inequality is a result from [27] which was correctly proved in [3]. For p = 1, it was proved in [1]. In [3], there is a remark that the proof in [1] also contains a similar mistake, but the authors of [1] corrected their text after the paper was posted.

If
$$p \ge 2$$
, then $\mathcal{J}_p(y) = 0$ implies $\int_{\Omega} y|y|^{p-2} dx = 0$. Really, since $s = \mathcal{J}_p(y) = 0$ is a stationary point of φ , we have
$$0 = \frac{d\varphi}{ds}\Big|_{s=0} = -p \int_{\Omega} y|y|^{p-2} dx$$

For such functions, the inequality was proved in [10].

Finally, in case $1 , the constant <math>\pi_p$ can be obtained by the Riecz-Thorin interpolation theorem between L^1 and L^2 .

As a result, we have

Statement 5. (Bramble-Hilbert lemma in $W^{\sigma+1,p}(\Omega;\mathbb{R})$) Let $\Omega \subset \mathbb{R}^m$ be a convex bounded domain, $p \geq 1$, $y \in W^{\sigma+1,p}(\Omega;\mathbb{R})$. Then there exists a polynomial P_y in m variables of degree at most σ such that

$$y - P_y|_{W^{\nu,p}(\Omega;\mathbb{R})} \le c_{\sigma+1,\nu,p} \left(\operatorname{diam}(\Omega)\right)^{\sigma+1-\nu} |y|_{W^{\sigma+1,p}(\Omega;\mathbb{R})}, \ \forall 0 \le \nu \le \sigma,$$

where

 $c_{\sigma,\nu,p} \le \pi_p^{\sigma-\nu} \binom{m+\nu-1}{\nu}^{\frac{1}{p}} \frac{\left((\sigma-\nu)!\right)^{\frac{1}{p}}}{\left(\left\lceil\frac{\sigma-\nu}{m}\right\rceil!\right)^{\frac{m}{p}}}$ (26)

and π_p is defined in (25).

Proof. As in [36], we define polynomials $P_{y,\sigma}, \ldots, P_{y,0} =: P_y$ recursively

$$P_{y,\sigma} := \sum_{\iota \in \mathbb{N}^m, |\iota| = \sigma} \frac{1}{\iota!} x^{\iota} \mathcal{J}_p(\partial^{\iota} y)$$

and

$$P_{y,\nu-1} := P_{y,\nu} + \sum_{\iota \in \mathbb{N}^m, |\iota| = \nu-1} \frac{1}{\iota!} x^{\iota} \mathcal{J}_p \left(\partial^{\iota} (y - \mathcal{J}_p(y)) \right), \ \sigma \ge \nu \ge 1.$$

As in [36], with properties of \mathcal{J}_p , one can show that

$$\partial^{\xi}(P_{y,\sigma-\tilde{\sigma}}) = P_{y,\sigma-\tilde{\sigma}-\nu}(\partial^{\xi}y)$$

and

$$\mathcal{J}_p\left(\partial^{\xi}(y - P_{y,\sigma-\tilde{\sigma}}(y))\right) = 0,$$

for all $y \in W^{\tilde{\sigma},p}(\Omega;\mathbb{R}), 0 \le \nu \le \tilde{\sigma}$, and $\xi \in \mathbb{N}^m$ with $|\xi| := \xi_1 + \cdots + \xi_m = \nu$.

The rest of the proof is similar to the one in [36].

As a corollary, arguing exactly as in [5], one can obtain a theorem on neural network approximation existence in $W^{\sigma,p}(\Omega;\mathbb{R})$.

Theorem 6. Let $p \ge 1$, $m \ge 2$, $\sigma \ge 3$, $\delta > 0$, $a_j, b_j \in \mathbb{Z}$, $a_j < b_j$, for $1 \le j \le m$, $\Omega = \prod_{j=1}^m [a_j, b_j]$ and $y \in W^{\sigma,p}(\Omega; \mathbb{R})$. Then for every $N \in \mathbb{N}$ with N > 5 there exists a tanh neural network $y_{\theta,N}$ with two hidden layers, one of width at most $3 \lceil \frac{\sigma}{2} \rceil \binom{\sigma+m-1}{m+1} + \sum_{j=1}^m (b_j - a_j)(N-1)$ and another of width at most $3 \lceil \frac{m+2}{2} \rceil \binom{2m+1}{m+1} N^{m+1} \prod_{i=1}^{m+1} (b_j - a_j)$, such that for $\nu \in \{0, 1, 2\}$ it holds that,

$$\|y - y_{\theta,N}\|_{W^{\nu,p}(\Omega;\mathbb{R})} \le 2^{\nu} 3^n \mathcal{A}_{\nu,\sigma,m,y}(1+\delta) \ln^{\nu} \left(\mathcal{B}_{\nu,\delta,d,y} N^{m+\sigma+2}\right) N^{-\sigma+\nu},\tag{27}$$

where

$$\mathcal{B}_{\nu,\delta,m,y} = \frac{5 \cdot 2^{\nu m} \max\left\{\prod_{j=1}^{m} (b_j - a_j), m\right\} \max\{\|y\|_{W^{\nu,\infty}(\Omega;\mathbb{R})}, 1\}}{3^m \delta \min\{1, \mathcal{A}_{\nu,\sigma,m,y}\}},$$
(28)

and

$$\mathcal{A}_{\nu,\sigma,m,y} = \max_{0 \le l \le \nu} c_{\sigma,l,p} \left(3\sqrt{m} \right)^{\sigma-l} |y|_{W^{\sigma,p}(\Omega;\mathbb{R})}, \tag{29}$$

where $c_{\sigma,l,p}$ is defined in (26).

Furthermore, the weights of $y_{\theta,N}$ scale as $O\left(N^{\max\left\{\frac{\sigma^2}{2},m\left(1+\frac{\sigma}{2}+\frac{m}{2}\right)\right\}}\right)$.

5 Conclusion

Hence, we gave an operator description of four PDE classes to which the error estimation, due to S. Mishra et al., can be applied. These classes cover a wide range of equations. Furthermore, we dealt with a more general real *p*-form instead of the inner product. For each class, we obtained an exact error estimate. Also, replacing the mean value with a projection, we extended the Bramble-Hilbert lemma to the non-Hilbert Sobolev spaces. Finally, we stated the theorem on neural network approximation existence in such a space.

However, obtained estimates remain a posteriori since we cannot guarantee that the training error will be small. It is still an open question under what conditions the gradient descent algorithm converges to a global minimum in the context of the PINN training.

References

- [1] G. Acosta, R. G. Durán. An optimal Poincaré inequality in L¹ for convex domains. *Proc. Amer. Math. Soc.*, 132(1):195–202, 2004.
- [2] G. Bai, U. Koley, S. Mishra, R. Molinaro. Physics informed neural networks (PINNs) for approximating nonlinear dispersive PDEs. J. Comp. Math., 39:816–847, 2022.
- [3] M. Bebendorf. A note on the Poincaré inequality for convex domains. J. Anal. Appl., 22:751–756, 2003.
- [4] H. Brezis. Functional analysis, Sobolev spaces and partial differential equations. Springer New York, NY, 2011.
- [5] T. De Ryck, A. D. Jagtap, S. Mishra. Error estimates for physics informed neural networks approximating the Navier-Stokes equations. *IMA J. Numer. Anal.*, drac085, 2023.
- [6] T. De Ryck, S. Langhaler, S. Mishra. On the approximation of functions by tanh neural networks. *Neural Netw.*, 143:732–750, 2021.
- [7] T. De Ryck, S. Mishra, R. Molinaro. wPINNs: Weak physics informed neural networks for approximating entropy solutions of hyperbolic conservation laws. arXiv preprint arXiv:2207.08483, 2022.
- [8] J. Diestel. Geometry of Banach spaces: selected topics. Springer Berlin, Heidelberg, 1975.
- [9] W. E, J. Han, A. Jentzen. Deep learning-based numerical methods for high-dimensional parabolic partial differential equations and backward stochastic differential equations. *Commun Math Stat.*, 5(4):349–380, 2017.
- [10] L. Esposito, C. Nitsch, C. Trombetti. Best constants in Poincaré inequalities for convex domains. J. Convex Anal., 20:253–264, 2013.
- [11] I. V. Gerasimov, A. I. Nazarov. Best constant in a three-parameter Poincaré inequality. J. Math. Sc., 179(1):80–99, 2011.
- [12] J. R. Giles. Classes of semi-inner-product spaces. Trans. Amer. Math. Soc., 129(3):436-446, 1967.
- [13] B. Hillebrecht, B. Unger. Certified machine learning: rigorous a posteriori error bounds for PDE defined PINNs. *arXiv preprint arXiv:2210.03426*, 2022.
- [14] Z. Hu, A. D. Jagtap, G. E. Karniadakis, K. Kawaguchi. When do extended physics-informed neural networks (XPINNs) improve generalization? SIAM J Sci Comput., 44(5):A3158–A3182, 2022.
- [15] H. Ishihara. On the constructive Hahn-Banach theorem. Bull. Lond. Math. Soc., 21:79–81, 1989.
- [16] A. D. Jagtap, G. E. Karniadakis. Extended physics-informed neural networks (XPINNs): A generalized space-time domain decomposition based deep learning framework for nonlinear partial differential equations. *Commun. Comput. Phys.*, 28(5):2002–2041, 2020.
- [17] A. D. Jagtap, E. Kharazmi, G. E. Karniadakis. Conservative physics-informed neural networks on discrete domains for conservation laws: Applications to forward and inverse problems. *Comput. Methods Appl. Mech. Eng.*, 365:113028, 2020.
- [18] A. D. Jagtap, Z. Mao, N. Adams, G. E. Karniadakis. Physics-informed neural networks for inverse problems in supersonic flows. J. Comput. Phys., 466:111402, 2022.
- [19] D. Johns, C. Gibson. A constructive approach to the duality theorem for certain Orlicz spaces. *Math. Proc. Camb. Philos. Soc.*, 89:49–69, 1981.
- [20] G. Kutyniok, P. Petersen, M. Raslan, R. Schneider. A Theoretical Analysis of Deep Neural Networks and Parametric PDEs. *Constr. Approx.*, 55:73–125, 2022.
- [21] L. Lu, P. Jin, G. E. Karniadakis. DeepONet: Learning nonlinear operators for identifying differential equations based on the universal approximation theorem of operators. *arXiv preprint arXiv:1910.03193*, 2019.
- [22] G. Lumer. Semi-inner-product spaces. Trans. Amer. Math. Soc., 100:29-43, 1961.
- [23] G. Lumer, R. S. Phillips. Dissipative operators in a Banach space. Pac. J. Math., 11(2):679–698, 1961.
- [24] K. O. Lye, S. Mishra, D. Ray. Deep learning observables in computational fluid dynamics. J. Comput. Phys., 410:109339, 2020.
- [25] S. Mishra, R. Molinaro. Estimates on the generalization error of physics-informed neural networks for approximating PDEs. *IMA J. Numer. Anal.*, 43(1):1–43, 2023.
- [26] E. Pap, R. Pavlović. On semi-inner product spaces of type p. Univ. u Novom Sadu, Zb. Rad. Prirod. Mat. Fak. Ser. Mat., 23(2):147–153, 1993.

- [27] L. E. Payne, H. F. Weinberger. An optimal Poincaré-inequality for convex domains. Arch. Rational Mech. Anal., 5:286–292, 1960.
- [28] A. Pazy. Semigroups of linear operators and applications to partial differential equations. Springer New York, NY, 1983.
- [29] M. Raissi, P. Perdikaris, G. Em Karniadakis. Physics informed deep learning (part I): data-driven solutions of nonlinear partial differential equations arXiv preprint arXiv:1711.10561, 2017.
- [30] M. Raissi, P. Perdikaris, G. Em Karniadakis. Physics informed deep learning (part II): data-driven discovery of nonlinear partial differential equations arXiv preprint arXiv:711.10566, 2017.
- [31] K. Shukla, A. D. Jagtap, J. L. Blackshire, D. Sparkman, G. E. Karniadakis. A physics- informed neural network for quantifying the microstructural properties of polycrystalline nickel using ultrasound data: A promising approach for solving inverse problems. IEEE Signal Process. Mag., 39(1):68–77, 2021.
- [32] K. Shukla, A. D. Jagtap, G. E. Karniadakis. Parallel physics-informed neural networks via domain decomposition. J. Comput. Phys., 447:110683, 2021.
- [33] J. E. Spingarn. Submonotone subdifferentials of lipschitz functions. Trans. Amer. Math. Soc., 264:77-89, 1981.
- [34] K. Sundaresan, S. Swaminathan. *Geometry and nonlinear analysis in Bahach spaces*, volume 1131 of *Lecture Notes in Mathematics* Springer Berlin, Heidelberg, 1985.
- [35] S. L. Troyanski. Gateaux differentiable norms in L^p. Math. Annal., 287(2):221–228, 1990.
- [36] R. Verfürth. A note on polynomial approximation in Sobolev spaces. *ESAIM: Math. Model. Numer. Anal.*, 33(4):715–719, 1999.