

Projective covers of the simple modules for the triplet W -algebra \mathcal{W}_{p_+,p_-}

Hiromu Nakano

Abstract

We study the structure of the abelian category of modules for the triplet W -algebra \mathcal{W}_{p_+,p_-} . Using the logarithmic deformation by Fjelstad et al.(2002), we construct logarithmic \mathcal{W}_{p_+,p_-} -modules that have L_0 nilpotent rank three or two. By using the structure of these logarithmic modules and the results on logarithmic Virasoro modules by Kytölä and Ridout(2009), we compute Ext^1 groups between certain indecomposable modules and simple modules. Based on these Ext^1 groups we determine the structure of the projective covers of all \mathcal{W}_{p_+,p_-} -simple modules.

Contents

1	Introduction	3
2	Bosonic Fock modules	4
2.1	Free field theory	5
2.2	The structure of Fock modules	6
2.3	Screening currents and Felder complex	8
3	The triplet W-algebra \mathcal{W}_{p_+,p_-}	12
3.1	The lattice vertex operator algebra and the vertex operator algebra \mathcal{W}_{p_+,p_-}	13
3.2	Simple \mathcal{W}_{p_+,p_-} -modules	14
3.3	The block decomposition of \mathcal{C}_{p_+,p_-}	19
4	Logarithmic \mathcal{W}_{p_+,p_-} modules	20
4.1	Logarithmic deformation	20
4.2	Logarithmic modules in the thick block	24
4.3	Logarithmic modules in the thin blocks	27

5	Logarithmic extension of Virasoro modules	29
5.1	Ext^1 -groups between simple Virasoro modules	29
5.2	Logarithmic extensions	33
6	The projective covers of simple \mathcal{W}_{p_+, p_-}-modules	43
6.1	The structure of the logarithmic modules $\mathcal{Q}(\mathcal{X}_{\bullet, \bullet}^{\pm})_{\bullet, \bullet}$	44
6.2	The Ext^1 -groups between all simple modules	48
6.3	The projective covers of the simple modules in the thin blocks	54
6.4	The projective covers of the simple modules $\mathcal{X}_{\bullet, \bullet}^{\pm}$ in the thick blocks	60
6.5	The projective covers of the minimal simple modules $L(h_{r,s})$.	77

1 Introduction

In 2006, Feigin, Gainutdinov, Semikhatov and Tipunin introduced the triplet W -algebra \mathcal{W}_{p_+, p_-} , where p_{\pm} are coprime integers such that $p_{\pm} \geq 2$ [12]. This W -algebra is an extension of the minimal Virasoro model

$$L(c_{p_+, p_-}, 0) \subset \mathcal{W}_{p_+, p_-}, \quad c_{p_+, p_-} := 1 - 6 \frac{(p_+ - p_-)^2}{p_+ p_-}$$

as a vertex operator algebra. The triplet W -algebra \mathcal{W}_{p_+, p_-} is irrational, C_2 -cofinite [3], [4], [33] and a natural generalization of the triplet W -algebra \mathcal{W}_p (cf. [1], [14], [15], [23], [25], [27]). Let \mathcal{C}_{p_+, p_-} be the abelian category of \mathcal{W}_{p_+, p_-} -modules. Similar to the case of \mathcal{W}_p , the abelian category \mathcal{C}_{p_+, p_-} has the block decomposition and each block is assigned to one of three groups: $\frac{(p_+ - 1)(p_- - 1)}{2}$ thick blocks, $p_+ + p_- - 2$ thin blocks and two semisimple blocks. The most complex groups are the thick blocks and each thick block contains five simple modules of which is the minimal Virasoro simple module. Each thick block contains certain logarithmic modules whose L_0 nilpotent rank three and the detailed structure of these rank three logarithmic modules has been studied from logarithmic conformal field theories and vertex operator algebras. In [31], Rasmussen examined the structure of indecomposable modules by using the method of solvable lattice models. In [18], [19], [34], Gaberdiel, Runkel and Wood examined the structure of rank three logarithmic modules from the direction of boundary conformal field theories and showed that the tensor category on \mathcal{C}_{p_+, p_-} is not rigid. In [6], Adamović and Milas constructed certain rank three logarithmic modules by using the method of the lattice construction. However, unlike the case of the triplet W -algebra \mathcal{W}_p [25], [27], there were few detailed studies of the structure of the abelian category \mathcal{C}_{p_+, p_-} , such as the properties of Ext^1 -groups.

The main goal of this paper is to determine the structure of these rank three logarithmic modules by using the logarithmic deformation in [17] and the center of the Zhu-algebra $A(\mathcal{W}_{p_+, p_-})$ [3], [4], [33], and to show that these logarithmic modules are projective.

This paper is organized as follows.

In Section 2, we review the structure of Fock modules and the Felder complex in accordance with [33]. The basic facts in this section are frequently used in later sections.

In Section 3, we introduce the vertex operator algebra \mathcal{W}_{p_+, p_-} and review some important results in [3], [4], [33] briefly. In Subsection 6.2, we introduce the block decomposition of \mathcal{C}_{p_+, p_-} . Each block of \mathcal{C}_{p_+, p_-} -mod is assigned to one of three groups: $\frac{(p_+ - 1)(p_- - 1)}{2}$ thick blocks $\mathcal{C}_{r, s}^{thick}$, $p_+ + p_- - 2$ thin

blocks $C_{r,p_-}^{thin}, C_{p_+,s}^{thin}$ and two semisimple blocks C_{p_+,p_-}^\pm . The most complex groups are the thick blocks and each thick block $C_{r,s}^{thick}$ contains five simple modules $\mathcal{X}_{r,s}^+, \mathcal{X}_{r^\vee,s^\vee}^+, \mathcal{X}_{r^\vee,s}^-, \mathcal{X}_{r,s^\vee}^-$ and $L(h_{r,s})$, where $L(h_{r,s})$ is the minimal simple module module of the Virasoro algebra. The thick blocks and the thin blocks contain logarithmic \mathcal{W}_{p_+,p_-} -modules on which the Virasoro zero-mode L_0 acts non-semisimply.

In Section 4, by gluing lattice simple modules $\mathcal{V}_{r,s}^\pm$ using the logarithmic deformation by [17], we construct logarithmic \mathcal{W}_{p_+,p_-} -modules $\mathcal{P}_{r,s}^\pm$ and $\mathcal{Q}(\mathcal{X}_{r,s}^\pm)_{\bullet,\bullet}$ whose L_0 nilpotent rank three and two, respectively.

In Section 5, we determine Ext^1 -groups between certain Virasoro modules, by using some important results by [24]. We also study the subquotient structure of some logarithmic Virasoro modules by using certain limit operations. These results are stated in Theorem 5.26 and Proposition 5.28.

In Section 6, we determine the structure of the projective covers of all \mathcal{W}_{p_+,p_-} -simple modules. In Subsections 6.1 and 6.2, we determine the structure of the socle series of indecomposable modules $\mathcal{Q}(\mathcal{X}_{\bullet,\bullet}^\pm)_{\bullet,\bullet}$ and the Ext^1 -groups between all simple modules. In Subsections 6.3 and 6.4, we study the structure of Ext^1 -groups in the thin blocks $C_{r,p_-}^{thin}, C_{p_+,s}^{thin}$ and the thick blocks $C_{r,s}^{thick}$. Based on the structure of the logarithmic Virasoro modules determined in Section 5, we compute Ext^1 groups between certain indecomposable \mathcal{W}_{p_+,p_-} -modules and the simple modules, and show that the logarithmic modules $\mathcal{Q}(\mathcal{X}_{\bullet,p_-}^\pm)_{\bullet,p_-}, \mathcal{Q}(\mathcal{X}_{p_+,\bullet}^\pm)_{p_+,\bullet}$ and $\mathcal{P}_{\bullet,\bullet}^\pm$ are the projective covers of the top composition factors. In Subsection 6.5, we determine the structure of the projective covers of the minimal simple modules $L(h_{r,s})$ by using the structure of the center of the Zhu-algebra $A(\mathcal{W}_{p_+,p_-})$ determined in [3],[4],[33].

This paper is partially based on our thesis [28].

2 Bosonic Fock modules

Recall that the Virasoro algebra \mathcal{L} is the Lie algebra over \mathbb{C} generated by $L_n (n \in \mathbb{Z})$ and C (the central charge) with the relation

$$[L_m, L_n] = (m - n)L_{m+n} + \frac{m^3 - m}{12}C\delta_{m+n,0}, \quad [L_n, C] = 0.$$

Fix two coprime integers p_+, p_- such that $p_- > p_+ \geq 2$, and let

$$c_{p_+,p_-} := 1 - 6 \frac{(p_+ - p_-)^2}{p_+ p_-}$$

be the central charge of the minimal model $M(p_+, p_-)$. In this section, we briefly review theories of Fock modules whose central charges are $C = c_{p_+,p_-}$.

id in accordance with [33]. As for the representation theory of the Virasoro algebra, see [11] and [22].

2.1 Free field theory

The Heisenberg Lie algebra

$$\mathcal{H} = \bigoplus_{n \in \mathbb{Z}} \mathbb{C}a_n \oplus \mathbb{C}K_{\mathcal{H}}$$

is the Lie algebra whose commutation is given by

$$[a_m, a_n] = m\delta_{m+n,0}K_{\mathcal{H}}, \quad [K_{\mathcal{H}}, \mathcal{H}] = 0.$$

Let

$$\mathcal{H}^{\pm} = \bigoplus_{n>0} \mathbb{C}a_{\pm n}, \quad \mathcal{H}^0 = \mathbb{C}a_0 \oplus \mathbb{C}K_{\mathcal{H}}, \quad \mathcal{H}^{\geq} = \mathcal{H}^+ \oplus \mathcal{H}^0.$$

For any $\alpha \in \mathbb{C}$, let $\mathbb{C}|\alpha\rangle$ be the one dimensional \mathcal{H}^{\geq} -module defined by

$$a_n|\alpha\rangle = \delta_{n,0}\alpha|\alpha\rangle \quad (n \geq 0), \quad K_{\mathcal{H}}|\alpha\rangle = |\alpha\rangle.$$

For any $\alpha \in \mathbb{C}$, the bosonic Fock module is defined by the induced module

$$F_{\alpha} = \text{Ind}_{\mathcal{H}^{\geq}}^{\mathcal{H}} \mathbb{C}|\alpha\rangle.$$

Let

$$a(z) = \sum_{n \in \mathbb{Z}} a_n z^{-n-1}$$

be the bosonic current. Then we have the following operator expansion

$$a(z)a(w) = \frac{1}{(z-w)^2} + \cdots,$$

where \cdots denotes the regular part in $z = w$. We define the energy-momentum tensor

$$T(z) := \frac{1}{2} : a(z)a(z) : + \frac{\alpha_0}{2} \partial a(z), \quad \alpha_0 := \sqrt{\frac{2p_-}{p_+}} - \sqrt{\frac{2p_+}{p_-}}.$$

where $: :$ is the normal ordered product. The energy-momentum tensor satisfies the following operator expansion

$$T(z)T(w) = \frac{c_{p_+, p_-}}{2(z-w)^4} + \frac{2T(w)}{(z-w)^2} + \frac{\partial T(w)}{z-w} + \cdots.$$

Then the Fourier modes of $T(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2}$ generate the Virasoro algebra whose central charge is c_{p_+, p_-} . Thus, by the energy-momentum tensor $T(z)$, each Fock module F_α has the structure of a Virasoro module whose central charge is c_{p_+, p_-} . Note that

$$L_0|\alpha\rangle = \frac{1}{2}\alpha(\alpha - \alpha_0)|\alpha\rangle.$$

Let us denote

$$h_\alpha := \frac{1}{2}\alpha(\alpha - \alpha_0). \quad (2.1)$$

For any $\alpha \in \mathbb{C}$, the Fock module F_α has the following L_0 weight decomposition

$$F_\alpha = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} F_\alpha[n], \quad F_\alpha[n] := \{v \in F_\alpha \setminus \{0\} \mid L_0 v = (h_\alpha + n)v\},$$

where each weight space $F_\alpha[n]$ has a basis

$$\{a_{-\lambda}|\alpha\rangle \mid \lambda \vdash n\}$$

with $a_{-\lambda} = a_{-\lambda_k} \cdots a_{-\lambda_1}$ for a partition $\lambda = (\lambda_1, \dots, \lambda_k)$.

We define the following conformal vector in F_0

$$T = \frac{1}{2}(a_{-1}^2 + \alpha_0 a_{-2})|0\rangle \in F_0[2].$$

Definition 2.1. *The Fock module F_0 carries the structure of a $\mathbb{Z}_{\geq 0}$ -graded vertex operator algebra, with*

$$Y(|0\rangle; z) = \text{id}, \quad Y(a_{-1}|0\rangle; z) = a(z), \quad Y(T; z) = T(z).$$

We denote this vertex operator algebra by \mathcal{F}_{α_0} .

2.2 The structure of Fock modules

We set

$$\alpha_+ = \sqrt{\frac{2p_-}{p_+}}, \quad \alpha_- = -\sqrt{\frac{2p_+}{p_-}}.$$

For $r, s, n \in \mathbb{Z}$ we introduce the following symbols

$$\alpha_{r,s;n} = \frac{1-r}{2}\alpha_+ + \frac{1-s}{2}\alpha_- + \frac{\sqrt{2p_+p_-}}{2}n, \quad \alpha_{r,s} = \alpha_{r,s;0}. \quad (2.2)$$

For $r, s, n \in \mathbb{Z}$, we use the following abbreviations

$$\begin{aligned} F_{r,s;n} &= F_{\alpha_{r,s;n}}, & F_{r,s} &= F_{\alpha_{r,s}}, \\ h_{r,s;n} &= h_{\alpha_{r,s;n}}, & h_{r,s} &= h_{\alpha_{r,s}}. \end{aligned}$$

Note that

$$h_{r,s;n} = h_{r-np_+,s} = h_{r,s+np_-}, \quad h_{r,s;n} = h_{-r,-s;-n}$$

for $r, s, n \in \mathbb{Z}$. For each $r, s, n \in \mathbb{Z}$, let $L(h_{r,s;n})$ be the irreducible Virasoro module whose highest weight is $h_{r,s;n}$ and the central charge $C = c_{p_+,p_-} \cdot \text{id}$.

Before describing the structure of Fock modules, let us introduce the notion of socle series.

Definition 2.2. *Let V be a vertex operator algebra or the Virasoro algebra, and let M be a finite length V -module. Let $\text{Soc}(M)$ be the socle of M , that is $\text{Soc}(M)$ is the maximal semisimple submodule of M . Since M is finite length, we have the sequence of the submodule*

$$\text{Soc}_1(M) \subsetneq \text{Soc}_2(M) \subsetneq \cdots \subsetneq \text{Soc}_n(M) = M$$

such that $\text{Soc}_1(M) = \text{Soc}(M)$ and $\text{Soc}_{i+1}(M)/\text{Soc}_i(M) = \text{Soc}(M/\text{Soc}_i(M))$. We call such a sequence of the submodules of M the socle series of M .

The following proposition is due to [11].

Proposition 2.3. *As the Virasoro modules, there are four cases of socle series for the Fock modules $F_{r,s;n} \in \mathcal{F}_{\alpha_0}\text{-mod}$:*

1. *For each $1 \leq r \leq p_+ - 1$, $1 \leq s \leq p_- - 1$, $n \in \mathbb{Z}$, we have*

$$\text{Soc}_1(F_{r,s;n}) \subsetneq \text{Soc}_2(F_{r,s;n}) \subsetneq \text{Soc}_3(F_{r,s;n}) = F_{r,s;n}$$

such that

$$\begin{aligned} \text{Soc}_1(F_{r,s;n}) &= \text{Soc}(F_{r,s;n}) = \bigoplus_{k \geq 0} L(h_{r,p_- - s;|n|+2k+1}), \\ \text{Soc}_2(F_{r,s;n})/\text{Soc}_1(F_{r,s;n}) &= \text{Soc}(F_{r,s;n}/\text{Soc}_1(F_{r,s;n})) \\ &= \bigoplus_{k \geq a} L(h_{r,s;|n|+2k}) \oplus \bigoplus_{k \geq 1-a} L(h_{p_+ - r, p_- - s;|n|+2k}), \\ \text{Soc}_3(F_{r,s;n})/\text{Soc}_2(F_{r,s;n}) &= \text{Soc}(F_{r,s;n}/\text{Soc}_2(F_{r,s;n})) = \bigoplus_{k \geq 0} L(h_{p_+ - r, s;|n|+2k+1}), \end{aligned}$$

where $a = 0$ if $n \geq 0$ and $a = 1$ if $n < 0$.

2. For each $1 \leq s \leq p_- - 1$, $n \in \mathbb{Z}$, we have

$$\text{Soc}_1(F_{p_+,s;n}) \subsetneq \text{Soc}_2(F_{p_+,s;n}) = F_{p_+,s;n}$$

such that

$$\text{Soc}_1(F_{p_+,s;n}) = \text{Soc}(F_{p_+,s;n}) = \bigoplus_{k \geq 0} L(h_{p_+,p_- - s;|n|+2k+1}),$$

$$\text{Soc}_2(F_{p_+,s;n})/\text{Soc}_1(F_{p_+,s;n}) = \bigoplus_{k \geq a} L(h_{p_+,s;|n|+2k})$$

where $a = 0$ if $n \geq 1$ and $a = 1$ if $n < 1$.

3. For each $1 \leq r \leq p_+ - 1$, $n \in \mathbb{Z}$, we have

$$\text{Soc}_1(F_{r,p_-;n}) \subsetneq \text{Soc}_2(F_{r,p_-;n}) = F_{r,p_-;n}$$

such that

$$\text{Soc}_1(F_{r,p_-;n}) = \text{Soc}(F_{r,p_-;n}) = \bigoplus_{k \geq 0} L(h_{r,p_-;|n|+2k}),$$

$$\text{Soc}_2(F_{r,p_-;n})/\text{Soc}_1(F_{r,p_-;n}) = \bigoplus_{k \geq a} L(h_{p_+ - r,p_-;|n|+2k-1})$$

where $a = 1$ if $n \geq 0$ and $a = 0$ if $n < 0$.

4. For each $n \in \mathbb{Z}$, the Fock module $F_{p_+,p_-;n}$ is semi-simple as a Virasoro module:

$$F_{p_+,p_-;n} = \text{Soc}(F_{p_+,p_-;n}) = \bigoplus_{k \geq 0} L(h_{p_+,p_-;|n|+2k}).$$

Let the Fock modules, whose socle length are three, be denoted by braided type, and let the Fock modules, whose length are two, be denoted by chain type.

2.3 Screening currents and Felder complex

We introduce a free scalar field $\phi(z)$, which is a formal primitive of $a(z)$

$$\phi(z) = \hat{a} + a_0 \log z - \sum_{n \neq 0} \frac{a_n}{n} z^{-n}$$

where \hat{a} is defined by

$$[a_m, \hat{a}] = \delta_{m,0} \text{id}. \quad (2.3)$$

The scalar field $\phi(z)$ satisfies the operator product expansion

$$\phi(z)\phi(w) = \log(z-w) + \dots$$

For any $\alpha \in \mathbb{C}$ we introduce the field $V_\alpha(z)$

$$\begin{aligned} V_\alpha(z) &=: e^{\alpha\phi(z)} := e^{\alpha\hat{a}} z^{\alpha a_0} \bar{V}_\alpha(z), \quad z^{\alpha a_0} = e^{\alpha a_0 \log z}, \\ \bar{V}_\alpha(z) &= e^{\alpha \sum_{n \geq 1} \frac{a-n}{n} z^n} e^{-\alpha \sum_{n \geq 1} \frac{an}{n} z^{-n}}. \end{aligned}$$

The fields $V_\alpha(z)$ satisfy the following operator product expansion

$$V_\alpha(z)V_\beta(w) = (z-w)^{\alpha\beta} : V_\alpha(z)V_\beta(w) :.$$

We introduce the following two screening currents $Q_+(z), Q_-(z)$

$$Q_\pm(z) = V_{\alpha_\pm}(z)$$

whose conformal weights are $h_{\alpha_\pm} = 1$:

$$\begin{aligned} T(z)Q_\pm(w) &= \frac{Q_\pm(w)}{(z-w)^2} + \frac{\partial_w Q_\pm(w)}{z-w} + \dots \\ &= \partial_w \left(\frac{Q_\pm(w)}{z-w} \right) + \dots \end{aligned}$$

Therefore the zero modes of the fields $Q_\pm(z)$

$$\begin{aligned} \text{Res}_{z=0} Q_+(z) dz &= Q_+ : F_{1,k} \rightarrow F_{-1,k}, \quad k \in \mathbb{Z} \\ \text{Res}_{z=0} Q_-(z) dz &= Q_- : F_{k,1} \rightarrow F_{k,-1}, \quad k \in \mathbb{Z} \end{aligned}$$

commute with every Virasoro mode.

For $r, s \geq 1$, we introduce more complicated screening currents

$$\begin{aligned} Q_+^{[r]}(z) &\in \text{Hom}_{\mathbb{C}}(F_{r,k}, F_{-r,k})[[z, z^{-1}]], \quad r \geq 1, k \in \mathbb{Z}, \\ Q_-^{[s]}(z) &\in \text{Hom}_{\mathbb{C}}(F_{k,s}, F_{k,-s})[[z, z^{-1}]], \quad s \geq 1, k \in \mathbb{Z}, \end{aligned}$$

constructed by [32] (cf. [22],[33]) as follows

$$\begin{aligned} Q_+^{[r]}(z) &= \int_{\bar{\Gamma}_r(\kappa_+)} Q_+(z) Q_+(zx_1) Q_+(zx_2) \cdots Q_+(zx_{r-1}) z^{r-1} dx_1 \cdots dx_{r-1}, \\ Q_-^{[s]}(z) &= \int_{\bar{\Gamma}_s(\kappa_-)} Q_-(z) Q_-(zx_1) Q_-(zx_2) \cdots Q_-(zx_{s-1}) z^{s-1} dx_1 \cdots dx_{s-1}, \end{aligned} \tag{2.4}$$

where $\bar{\Gamma}_n(\kappa_{\pm})$ is a certain regularized cycle constructed from the simplex

$$\Delta_{n-1} = \{ (x_1, \dots, x_{n-1}) \in \mathbb{R}^{n-1} \mid 1 > x_1 > \dots > x_{n-1} > 0 \}.$$

These fields satisfy the following operator product expansion

$$\begin{aligned} T(z)Q_+^{[r]}(w) &= \frac{Q_+^{[r]}(w)}{(z-w)^2} + \frac{\partial_w Q_+^{[r]}(w)}{z-w} + \dots, \\ T(z)Q_-^{[s]}(w) &= \frac{Q_-^{[s]}(w)}{(z-w)^2} + \frac{\partial_w Q_-^{[s]}(w)}{z-w} + \dots. \end{aligned}$$

In particular the following proposition holds

Proposition 2.4. *The zero modes*

$$\begin{aligned} \text{Res}_{z=0} Q_+^{[r]}(z)dz &= Q_+^{[r]} \in \text{Hom}_{\mathbb{C}}(F_{r,k}, F_{-r,k}), \quad r \geq 1, k \in \mathbb{Z}, \\ \text{Res}_{z=0} Q_-^{[s]}(z)dz &= Q_-^{[s]} \in \text{Hom}_{\mathbb{C}}(F_{k,s}, F_{k,-s}), \quad s \geq 1, k \in \mathbb{Z} \end{aligned}$$

commute with every Virasoro mode of \mathcal{F}_{α_0} -Mod. These zero modes are called screening operators.

For $1 \leq r \leq p_+ - 1$, $1 \leq s \leq p_- - 1$, we set

$$r^{\vee} := p_+ - r, \quad s^{\vee} := p_- - s.$$

For $1 \leq r \leq p_+$, $1 \leq s \leq p_-$ and $n \in \mathbb{Z}$, we define the following Virasoro modules :

1. For $1 \leq r < p_+$, $1 \leq s \leq p_-$, $n \in \mathbb{Z}$

$$\begin{aligned} K_{r,s;n;+} &= \ker Q_+^{[r]} : F_{r,s;n} \rightarrow F_{r^{\vee},s;n+1}, \\ X_{r^{\vee},s;n+1;+} &= \text{im} Q_+^{[r]} : F_{r,s;n} \rightarrow F_{r^{\vee},s;n+1}. \end{aligned}$$

2. For $1 \leq r \leq p_+$, $1 \leq s < p_-$, $n \in \mathbb{Z}$

$$\begin{aligned} K_{r,s;n;-} &= \ker Q_-^{[s]} : F_{r,s;n} \rightarrow F_{r,s^{\vee};n-1}, \\ X_{r,s^{\vee};n-1;-} &= \text{im} Q_-^{[s]} : F_{r,s;n} \rightarrow F_{r,s^{\vee};n-1}. \end{aligned}$$

The following propositions are due to [16].

Proposition 2.5. *The socle series of $K_{r,s;n;\pm}$ and $X_{r,s;n;\pm}$ are given by :*

1. For $1 \leq r \leq p_+ - 1$, $1 \leq s \leq p_- - 1$, we have

$$\begin{aligned} S_1(K_{r,s;n;\pm}) &= \text{Soc}(K_{r,s;n;\pm}) \subsetneq K_{r,s;n;\pm}, \\ S_1(X_{r,s;n;\pm}) &= \text{Soc}(X_{r,s;n;\pm}) \subsetneq X_{r,s;n;\pm} \end{aligned}$$

such that

$$\begin{aligned} n \geq 0 \qquad \qquad \qquad n \leq -1 \\ S_1(K_{r,s;n;+}) &= \bigoplus_{k \geq 1} L(h_{r,s^\vee;n+2k-1}), & S_1(K_{r,s;n;+}) &= \bigoplus_{k \geq 1} L(h_{r,s^\vee;-n+2k-1}), \\ K_{r,s;n;+}/S_1 &= \bigoplus_{k \geq 1} L(h_{r,s;n+2(k-1)}), & K_{r,s;n;+}/S_1 &= \bigoplus_{k \geq 1} L(h_{r,s;-n+2k}), \\ S_1(X_{r,s;n+1;+}) &= \bigoplus_{k \geq 1} L(h_{r,s^\vee;n+2k}), & S_1(X_{r,s;n+1;+}) &= \bigoplus_{k \geq 1} L(h_{r,s^\vee;-n+2(k-1)}), \\ X_{r,s;n+1;+}/S_1 &= \bigoplus_{k \geq 1} L(h_{r,s;n+2k-1}), & X_{r,s;n+1;+}/S_1 &= \bigoplus_{k \geq 1} L(h_{r,s;-n+2k-1}). \end{aligned}$$

$$\begin{aligned} n \geq 1 \qquad \qquad \qquad n \leq 0 \\ S_1(K_{r,s;n;-}) &= \bigoplus_{k \geq 1} L(h_{r,s^\vee;n+2k-1}), & S_1(K_{r,s;n;-}) &= \bigoplus_{k \geq 1} L(h_{r,s^\vee;-n+2k-1}), \\ K_{r,s;n;-}/S_1 &= \bigoplus_{k \geq 1} L(h_{r,s;n+2(k-1)}), & K_{r,s;n;-}/S_1 &= \bigoplus_{k \geq 1} L(h_{r,s;-n+2k}), \\ S_1(X_{r,s;n+1;-}) &= \bigoplus_{k \geq 1} L(h_{r,s^\vee;n+2(k-1)}), & S_1(X_{r,s;n+1;-}) &= \bigoplus_{k \geq 1} L(h_{r,s^\vee;-n+2k}), \\ X_{r,s;n+1;-}/S_1 &= \bigoplus_{k \geq 1} L(h_{r^\vee,s^\vee;n+2k-1}), & X_{r,s;n+1;-}/S_1 &= \bigoplus_{k \geq 1} L(h_{r^\vee,s^\vee;-n+2k-1}). \end{aligned}$$

2. For $1 \leq r \leq p_+ - 1$, $s = p_-$, $n \in \mathbb{Z}$, we have

$$X_{r,p_-;n} = \text{Soc}(F_{r,p_-;n}).$$

3. For $r = p_+$, $1 \leq s \leq p_- - 1$, $n \in \mathbb{Z}$, we have

$$X_{p_+,s;n} = \text{Soc}(F_{p_+,s;n}).$$

Proposition 2.6 ([16]).

1. For $1 \leq r < p_+$, $1 \leq s < p_-$ and $n \in \mathbb{Z}$ the screening operators $Q_+^{[r]}$ and $Q_+^{[r^\vee]}$ define the Felder complex

$$\cdots \xrightarrow{Q_+^{[r]}} F_{r^\vee, s; n-1} \xrightarrow{Q_+^{[r^\vee]}} F_{r, s; n} \xrightarrow{Q_+^{[r]}} F_{r^\vee, s; n+1} \xrightarrow{Q_+^{[r^\vee]}} \cdots$$

This complex is exact everywhere except in $F_{r, s} = F_{r, s; 0}$ where the cohomology is given by

$$\ker Q_+^{[r]} / \operatorname{im} Q_+^{[r^\vee]} \simeq L(h_{r, s; 0}).$$

2. For $1 \leq r < p_+$, $1 \leq s < p_-$ and $n \in \mathbb{Z}$ the screening operators $Q_-^{[s]}$ and $Q_-^{[s^\vee]}$ define the Felder complex

$$\cdots \xrightarrow{Q_-^{[s]}} F_{r, s^\vee; n+1} \xrightarrow{Q_-^{[s^\vee]}} F_{r, s; n} \xrightarrow{Q_-^{[s]}} F_{r, s^\vee; n-1} \xrightarrow{Q_-^{[s^\vee]}} \cdots$$

This complex is exact everywhere except in $F_{r, s} = F_{r, s; 0}$ where the cohomology is given by

$$\ker Q_-^{[s]} / \operatorname{im} Q_-^{[s^\vee]} \simeq L(h_{r, s; 0}).$$

3. For $1 \leq r < p_+$ and $n \in \mathbb{Z}$ the screening operators $Q_+^{[r]}$ and $Q_+^{[r^\vee]}$ define the Felder complex

$$\cdots \xrightarrow{Q_+^{[r]}} F_{r^\vee, p_-; n-1} \xrightarrow{Q_+^{[r^\vee]}} F_{r, p_-; n} \xrightarrow{Q_+^{[r]}} F_{r^\vee, p_-; n+1} \xrightarrow{Q_+^{[r^\vee]}} \cdots$$

and this complex is exact.

4. For $1 \leq s < p_-$ and $n \in \mathbb{Z}$ the screening operators $Q_-^{[s]}$ and $Q_-^{[s^\vee]}$ define the Felder complex

$$\cdots \xrightarrow{Q_-^{[s]}} F_{p_+, s^\vee; n+1} \xrightarrow{Q_-^{[s^\vee]}} F_{p_+, s; n} \xrightarrow{Q_-^{[s]}} F_{p_+, s^\vee; n-1} \xrightarrow{Q_-^{[s^\vee]}} \cdots$$

and this complex is exact.

3 The triplet W -algebra \mathcal{W}_{p_+, p_-}

In this section, we introduce a vertex operator algebra \mathcal{W}_{p_+, p_-} which is called the triplet W -algebra of type (p_+, p_-) and review some important results in [3], [4], [33] briefly. In Subsection 3.3, we introduce the abelian category of \mathcal{W}_{p_+, p_-} -modules and the block decomposition of this abelian category.

3.1 The lattice vertex operator algebra and the vertex operator algebra \mathcal{W}_{p_+, p_-}

Definition 3.1.

The lattice vertex operator algebra $\mathcal{V}_{[p_+, p_-]}$ is the tuple

$$(\mathcal{V}_{1,1}^+, |0\rangle, \frac{1}{2}(a_{-1}^2 - \alpha_0 a_{-2})|0\rangle, Y),$$

where underlying vector space of $\mathcal{V}_{[p_+, p_-]}$ is given by

$$\mathcal{V}_{1,1}^+ = \bigoplus_{n \in \mathbb{Z}} F_{1,1;2n} = \bigoplus_{n \in \mathbb{Z}} F_{n\sqrt{2p_+p_-}},$$

and $Y(|\alpha_{1,1;2n}\rangle; z) = V_{\alpha_{1,1;2n}}(z)$ for $n \in \mathbb{Z}$.

It is a known fact that simple $\mathcal{V}_{[p_+, p_-]}$ -modules are given by the following $2p_+p_-$ direct sum of Fock modules

$$\mathcal{V}_{r,s}^+ = \bigoplus_{n \in \mathbb{Z}} F_{r,s;2n}, \quad \mathcal{V}_{r,s}^- = \bigoplus_{n \in \mathbb{Z}} F_{r,s;2n+1},$$

where $1 \leq r \leq p_+, 1 \leq s \leq p_-$.

Note that the two screening operators Q_+ and Q_- act on $\mathcal{V}_{1,1}^+$. We define the following vector subspace of $\mathcal{V}_{1,1}^+$:

$$\mathcal{K}_{1,1} = \ker Q_+ \cap \ker Q_- \subset \mathcal{V}_{1,1}^+.$$

Definition 3.2. The triplet W -algebra

$$\mathcal{W}_{p_+, p_-} = (\mathcal{K}_{1,1}, |0\rangle, T, Y)$$

is a sub vertex operator algebra of $\mathcal{V}_{[p_+, p_-]}$, where the vacuum vector, conformal vector and vertex operator map are those of $\mathcal{V}_{[p_+, p_-]}$.

Definition 3.3. Let W^\pm, W^0 be the following singular vectors

$$W^+ = Q_-^{[p_- - 1]} |\alpha_{1, p_- - 1; 3}\rangle, \quad W^- = Q_+^{[p_+ - 1]} |\alpha_{p_+ - 1, 1; -3}\rangle, \quad W^0 = Q_+^{[2p_+ - 1]} |\alpha_{p_+ - 1, 1; -3}\rangle.$$

Proposition 3.4. \mathcal{W}_{p_+, p_-} is strongly generated by the fields $T(z), Y(W^\pm; z), Y(W^0; z)$.

Theorem 3.5 ([3, 4, 33]). \mathcal{W}_{p_+, p_-} is C_2 -cofinite.

3.2 Simple \mathcal{W}_{p_+, p_-} -modules

For each $1 \leq r \leq p_+$, $1 \leq s \leq p_-$, let $\mathcal{X}_{r,s}^\pm$ be the following vector subspace of $\mathcal{V}_{r,s}^\pm$:

1. For $1 \leq r \leq p_+ - 1$, $1 \leq s \leq p_- - 1$,

$$\mathcal{X}_{r,s}^+ = Q_+^{[r^\vee]}(\mathcal{V}_{r^\vee, s}^-) \cap Q_-^{[s^\vee]}(\mathcal{V}_{r, s^\vee}^-), \quad \mathcal{X}_{r,s}^- = Q_+^{[r^\vee]}(\mathcal{V}_{r^\vee, s}^+) \cap Q_-^{[s^\vee]}(\mathcal{V}_{r, s^\vee}^+).$$

2. For $1 \leq r \leq p_+ - 1$, $s = p_-$,

$$\mathcal{X}_{r, p_-}^+ = Q_+^{[r^\vee]}(\mathcal{V}_{r^\vee, p_-}^-), \quad \mathcal{X}_{r, p_-}^- = Q_+^{[r^\vee]}(\mathcal{V}_{r^\vee, p_-}^+).$$

3. For $r = p_+$, $1 \leq s \leq p_- - 1$,

$$\mathcal{X}_{p_+, s}^+ = Q_-^{[s^\vee]}(\mathcal{V}_{p_+, s^\vee}^-), \quad \mathcal{X}_{p_+, s}^- = Q_-^{[s^\vee]}(\mathcal{V}_{p_+, s^\vee}^+).$$

4. $r = p_+$, $s = p_-$,

$$\mathcal{X}_{p_+, p_-}^+ = \mathcal{V}_{p_+, p_-}^+, \quad \mathcal{X}_{p_+, p_-}^- = \mathcal{V}_{p_+, p_-}^-.$$

Definition 3.6.

1. We define the interior Kac table K_{p_+, p_-} as the following quotient set

$$K_{p_+, p_-} = \{(r, s) \mid 1 \leq r < p_+, 1 \leq s < p_-\} / \sim$$

where $(r, s) \sim (r', s')$ if and only if $r' = p_+ - r$, $s' = p_- - s$. Note that $\#K_{p_+, p_-} = \frac{(p_+ - 1)(p_- - 1)}{2}$.

2. For each $1 \leq r \leq p_+$, $1 \leq s \leq p_-$, $n \geq 0$, we define the following symbols

$$\Delta_{r,s;n}^+ = \begin{cases} h_{r^\vee, s; -2n-1} & r \neq p_+, s \neq p_- \\ h_{p_+, s; -2n} & r = p_+, s \neq p_- \\ h_{r, p_-; 2n} & r \neq p_+, s = p_- \\ h_{p_+, p_-; -2n} & r = p_+, s = p_- \end{cases}, \quad \Delta_{r,s;n}^- = \begin{cases} h_{r^\vee, s; -2n-2} & r \neq p_+, s \neq p_- \\ h_{p_+, s; -2n-1} & r = p_+, s \neq p_- \\ h_{r, p_-; 2n+1} & r \neq p_+, s = p_- \\ h_{p_+, p_-; -2n-1} & r = p_+, s = p_- \end{cases}.$$

Proposition 3.7 ([3, 4, 33]). For each $\mathcal{X}_{r,s}^\pm$ ($1 \leq r \leq p_+$, $1 \leq s \leq p_-$), we have the following decompositions as the Virasoro modules

$$\mathcal{X}_{r,s}^+ = \bigoplus_{n \geq 0} (2n+1)L(\Delta_{r,s;n}^+), \quad \mathcal{X}_{r,s}^- = \bigoplus_{n \geq 0} (2n+2)L(\Delta_{r,s;n}^-).$$

Theorem 3.8 ([3, 4, 33]). *The $\frac{(p_+-1)(p_--1)}{2} + 2p_+p_-$ vector spaces*

$$L(h_{r,s}), (r, s) \in K_{p_+, p_-}, \quad \mathcal{X}_{r,s}^\pm, \quad 1 \leq r \leq p_+, \quad 1 \leq s \leq p_-$$

become simple \mathcal{W}_{p_+, p_-} -modules and give all simple \mathcal{W}_{p_+, p_-} -modules.

Proposition 3.9 ([3, 4, 33]). *Each $2p_+p_-$ simple $\mathcal{V}_{[p_+, p_-]}$ -module becomes \mathcal{W}_{p_+, p_-} -module and has the socle series*

1. *For each $1 \leq r < p_+, 1 \leq s < p_-, \mathcal{V}_{r,s}^+$ has the following socle series*

$$\text{Soc}_1(\mathcal{V}_{r,s}^+) \subsetneq \text{Soc}_2(\mathcal{V}_{r,s}^+) \subsetneq \text{Soc}_3(\mathcal{V}_{r,s}^+) = \mathcal{V}_{r,s}^+$$

such that

$$\begin{aligned} \text{Soc}_1(\mathcal{V}_{r,s}^+) &= \text{Soc}(\mathcal{V}_{r,s}^+) = \mathcal{X}_{r,s}^+, \\ \text{Soc}(\mathcal{V}_{r,s}^+ / \text{Soc}_1(\mathcal{V}_{r,s}^+)) &= \mathcal{X}_{r^\vee, s}^- \oplus \mathcal{X}_{r, s^\vee}^- \oplus L(h_{r,s}), \\ \text{Soc}(\mathcal{V}_{r,s}^+ / \text{Soc}_2(\mathcal{V}_{r,s}^+)) &= \mathcal{X}_{r^\vee, s^\vee}^+. \end{aligned}$$

2. *For each $1 \leq r < p_+, 1 \leq s < p_-, \mathcal{V}_{r,s}^-$ has the socle series*

$$\text{Soc}_1(\mathcal{V}_{r,s}^-) \subsetneq \text{Soc}_2(\mathcal{V}_{r,s}^-) \subsetneq \text{Soc}_3(\mathcal{V}_{r,s}^-) = \mathcal{V}_{r,s}^-$$

such that

$$\begin{aligned} \text{Soc}_1(\mathcal{V}_{r,s}^-) &= \text{Soc}(\mathcal{V}_{r,s}^-) = \mathcal{X}_{r,s}^-, \\ \text{Soc}(\mathcal{V}_{r,s}^- / \text{Soc}_1(\mathcal{V}_{r,s}^-)) &= \mathcal{X}_{r^\vee, s}^+ \oplus \mathcal{X}_{r, s^\vee}^+, \\ \text{Soc}(\mathcal{V}_{r,s}^- / \text{Soc}_2(\mathcal{V}_{r,s}^-)) &= \mathcal{X}_{r^\vee, s^\vee}^-. \end{aligned}$$

3. *For each $1 \leq r < p_+, \mathcal{V}_{r, p_-}^+$ and $\mathcal{V}_{r^\vee, p_-}^-$ have the following socle series*

$$\mathcal{V}_{r, p_-}^+ / \mathcal{X}_{r, p_-}^+ \simeq \mathcal{X}_{r^\vee, p_-}^-, \quad \mathcal{V}_{r^\vee, p_-}^- / \mathcal{X}_{r^\vee, p_-}^- \simeq \mathcal{X}_{r, p_-}^+.$$

4. *For each $1 \leq s < p_-, \mathcal{V}_{p_+, s}^+$ and $\mathcal{V}_{p_+, s^\vee}^-$ have the following socle series*

$$\mathcal{V}_{p_+, s}^+ / \mathcal{X}_{p_+, s}^+ \simeq \mathcal{X}_{p_+, s^\vee}^-, \quad \mathcal{V}_{p_+, s^\vee}^- / \mathcal{X}_{p_+, s^\vee}^- \simeq \mathcal{X}_{p_+, s}^+.$$

5. *For $r = p_+, s = p_-$, we have*

$$\mathcal{V}_{p_+, p_-}^+ = \mathcal{X}_{p_+, p_-}^+, \quad \mathcal{V}_{p_+, p_-}^- = \mathcal{X}_{p_+, p_-}^-.$$

Let $A(\mathcal{W}_{p_+, p_-})$ be the Zhu-algebra [36] of \mathcal{W}_{p_+, p_-} .

Proposition 3.10 ([3, 4, 33]). *In $A(\mathcal{W}_{p_+, p_-})$, the following relations hold*

$$\begin{aligned} [W^0] * [W^-] - [W^-] * [W^0] &= -2f([T])[W^-], \\ [W^0] * [W^+] - [W^+] * [W^0] &= 2f([T])[W^+], \\ [W^+] * [W^-] - [W^-] * [W^+] &= 2f([T])[W^0], \\ [W^0] * [W^0] &= g([T]), \\ [W^+] * [W^+] &= 0, \\ [W^-] * [W^-] &= 0, \end{aligned}$$

where $f([T])$ and $g([T])$ are non-trivial polynomials of $[T]$.

Proposition 3.11 ([3, 4, 33]).

1. $\mathcal{X}_{1,1}^+$ acts trivially on $L(h_{r,s})$, ($1 \leq r \leq p_+ - 1$, $1 \leq s \leq p_- - 1$).
2. For each $1 \leq r \leq p_+$, $1 \leq s \leq p_-$, the highest weight space of $\mathcal{X}_{r,s}^+$ is a one dimensional $A(\mathcal{W}_{p_+, p_-})$ -module.
3. For each $1 \leq r \leq p_+$, $1 \leq s \leq p_-$, the highest weight space of $\mathcal{X}_{r,s}^-$ is a two dimensional irreducible $A(\mathcal{W}_{p_+, p_-})$ -module.

Proposition 3.12 ([3, 4, 33]). *For any $1 \leq r \leq p_+$, $1 \leq s \leq p_-$,*

$$f(\Delta_{r,s;0}^-) \neq 0.$$

In particular, the highest weight space of $\mathcal{X}_{r,s}^-$ has the structure of a two dimensional irreducible sl_2 -module with respect to the following elements

$$E = \frac{1}{\sqrt{2}f(\Delta_{r,s;0}^-)}[W^+], \quad F = -\frac{1}{\sqrt{2}f(\Delta_{r,s;0}^-)}[W^-], \quad H = \frac{1}{f(\Delta_{r,s;0}^-)}[W^0].$$

For $1 \leq r \leq p_+$, $1 \leq s \leq p_-$, we define

$$\begin{aligned} G(\Delta_{r,s;n}^+) &:= \begin{cases} \bigoplus_{n-1 \geq i \geq 0} (2i+1)L(\Delta_{r,s;i}^+) & n \geq 1 \\ 0 & n = 0, \end{cases} \\ G(\Delta_{r,s;n}^-) &:= \begin{cases} \bigoplus_{n-1 \geq i \geq 0} (2i+2)L(\Delta_{r,s;i}^-) & n \geq 1 \\ 0 & n = 0. \end{cases} \end{aligned}$$

As an extension of Proposition 3.11, the following propositions holds (see the proof of Proposition 5.6 in [33]).

Proposition 3.13 ([3, 4, 33]).

1. With respect to the actions of the zero-modes of the fields $Y(W^+; z)$, $Y(W^-; z)$ and $Y(W^0; z)$, the Virasoro highest weight space of the vector subspace $(2n+1)L(\Delta_{r,s;n}^+) \subset \mathcal{X}_{r,s}^+$ becomes a $(2n+1)$ -dimensional irreducible sl_2 -module modulo $G(\Delta_{r,s;n}^+)$.
2. With respect to the actions of the zero-modes of the fields $Y(W^+; z)$, $Y(W^-; z)$ and $Y(W^0; z)$, the Virasoro highest weight space of the vector subspace $(2n+2)L(\Delta_{r,s;n}^-) \subset \mathcal{X}_{r,s}^-$ becomes a $(2n+2)$ -dimensional irreducible sl_2 -module modulo $G(\Delta_{r,s;n}^-)$.

For $W = W^\pm, W^0$, let $W[n]$ be the n -th mode of the field $Y(W; z)$ defined by

$$W[n] = \oint_{z=0} Y(W; z) z^{h_{4p+}-1, 1+n-1} dz.$$

Proposition 3.14 ([3, 4, 33]).

1. For $n \geq 0$, let $\{w_i^{(n)}\}_{i=-n}^n$ be the basis of the Virasoro highest weight space of the vector subspace $(2n+1)L(\Delta_{r,s;n}^+) \subset \mathcal{X}_{r,s}^+$ such that

$$W^\pm[0]w_i^{(n)} \in \mathbb{C}^\times w_{i\pm 1}^{(n)} + G(\Delta_{r,s;n}^+), \quad \text{for } -n \leq i \leq n,$$

where $w_{-n-1}^{(n)} = w_{n+1}^{(n)} = 0$ and $W^\pm[0]$ is the zero mode of the field $Y(W^\pm; z)$. Then we have

$$\begin{aligned} W^\pm[\Delta_{r,s;n}^+ - \Delta_{r,s;n-1}^+]w_i^{(n)} &\in \mathbb{C}^\times w_{i\pm 1}^{(n-1)} + G(\Delta_{r,s;n-1}^+), \\ W^\pm[\Delta_{r,s;n}^+ - \Delta_{r,s;n+1}^+]w_i^{(n)} &\in \mathbb{C}^\times w_{i\pm 1}^{(n+1)} + G(\Delta_{r,s;n+1}^+), \\ W^0[\Delta_{r,s;n}^+ - \Delta_{r,s;n-1}^+]w_i^{(n)} &\in \mathbb{C}^\times w_i^{(n-1)} + G(\Delta_{r,s;n-1}^+), \\ W^0[\Delta_{r,s;n}^+ - \Delta_{r,s;n+1}^+]w_i^{(n)} &\in \mathbb{C}^\times w_i^{(n+1)} + G(\Delta_{r,s;n+1}^+), \end{aligned}$$

where $w_i^{(-1)} = 0$.

2. For $n \geq 0$, let $\{v_{\frac{i}{2}}^{(n)}, v_{\frac{-i}{2}}^{(n)}\}_{i=1}^{n+1}$ be the basis of the Virasoro highest weight space of the vector subspace $(2n+2)L(\Delta_{r,s;n}^-) \subset \mathcal{X}_{r,s}^-$ such that

$$W^\pm[0]v_{\frac{i}{2}}^{(n)} \in \mathbb{C}^\times v_{\frac{i}{2}\pm 1}^{(n)} + G(\Delta_{r,s;n}^-), \quad \text{for } -n-1 \leq i \leq n+1 \wedge i \neq 0,$$

where $v_{\frac{-n-2}{2}}^{(n)} = v_{\frac{n+2}{2}}^{(n)} = 0$. Then we have

$$\begin{aligned} W^\pm[\Delta_{r,s;n}^- - \Delta_{r,s;n-1}^-]v_{\frac{i}{2}}^{(n)} &\in \mathbb{C}^\times v_{\frac{i}{2}\pm 1}^{(n-1)} + G(\Delta_{r,s;n-1}^-), \\ W^\pm[\Delta_{r,s;n}^- - \Delta_{r,s;n+1}^-]v_{\frac{i}{2}}^{(n)} &\in \mathbb{C}^\times v_{\frac{i}{2}\pm 1}^{(n+1)} + G(\Delta_{r,s;n+1}^-), \\ W^0[\Delta_{r,s;n}^- - \Delta_{r,s;n-1}^-]v_{\frac{i}{2}}^{(n)} &\in \mathbb{C}^\times v_{\frac{i}{2}}^{(n-1)} + G(\Delta_{r,s;n-1}^-), \\ W^0[\Delta_{r,s;n}^- - \Delta_{r,s;n+1}^-]v_{\frac{i}{2}}^{(n)} &\in \mathbb{C}^\times v_{\frac{i}{2}}^{(n+1)} + G(\Delta_{r,s;n+1}^-), \end{aligned}$$

where $v_i^{(-1)} = 0$.

The following results for the Zhu-algebra $A(\mathcal{W}_{p_+,p_-})$ will be used to determine the structure of the projective covers of the minimal simple modules in Subsection 6.5.

Theorem 3.15 ([3, 4, 33]). *The center of the Zhu-algebra $A(\mathcal{W}_{p_+,p_-})$ is generated by $[T]$ and isomorphic to*

$$\mathbb{C}[x]/f_{p_+,p_-}(x),$$

where

$$\begin{aligned} f_{p_+,p_-}(x) &= \prod_{(i,j) \in \mathcal{T}} (x - h_{r,s})^3 \\ &\times \prod_{i=1}^{p_+-1} \prod_{j=1}^{p_--1} (x - \Delta_{i,j;0}^+)^2 \prod_{i=1}^{p_+-1} \prod_{j=1}^{p_--1} (x - \Delta_{i,j;0}^-) \\ &\times \prod_{i=1}^{p_+-1} (x - \Delta_{i,p_-;0}^+)^2 \prod_{i=1}^{p_+-1} (x - \Delta_{i,p_-;0}^-) \\ &\times \prod_{j=1}^{p_--1} (x - \Delta_{p_+,j;0}^+)^2 \prod_{j=1}^{p_--1} (x - \Delta_{p_+,j;0}^-) \\ &\times (x - \Delta_{p_+,p_-;0}^+)(x - \Delta_{p_+,p_-;0}^-). \end{aligned}$$

Corollary 3.16. *The Zhu algebra $A(\mathcal{W}_{p_+,p_-})$ has three dimensional indecomposable modules on which $[T]$ acts as*

$$\begin{pmatrix} h_{r,s} & 1 & 0 \\ 0 & h_{r,s} & 1 \\ 0 & 0 & h_{r,s} \end{pmatrix},$$

where $(r, s) \in K_{p_+,p_-}$.

3.3 The block decomposition of \mathcal{C}_{p_+, p_-}

Definition 3.17. Let M be a \mathcal{W}_{p_+, p_-} -module that satisfies the following properties:

1. For the central charge, $C = c_{p_+, p_-} \cdot \text{id}$ on M .
2. M has the following decomposition $M = \sum_{h \in H(M)} M[h]$:
 - For some finite subset $H_0(M)$ of \mathbb{C} , $H(M) = H_0(M) + \mathbb{Z}_{\geq 0}$.
 - For $h \in H(M)$, $M[h] = \{m \in M : \exists n \geq 0 \text{ s.t. } (L_0 - h)^n m = 0\}$.
 - $0 < \dim_{\mathbb{C}} M[h] < \infty$.

Let \mathcal{C}_{p_+, p_-} be the abelian category of \mathcal{W}_{p_+, p_-} -modules that satisfies the above two properties. We call any $M \in \mathcal{C}_{p_+, p_-}$ on which L_0 acts non-semisimply logarithmic module.

Since \mathcal{W}_{p_+, p_-} is C_2 -cofinite, any M in \mathcal{C}_{p_+, p_-} has finite length. For any M in \mathcal{C}_{p_+, p_-} , let M^* be the contragredient of M . Note that \mathcal{C}_{p_+, p_-} is closed under contragredient.

Definition 3.18. In the following, we define $\frac{(p_+-1)(p_--1)}{2}$ thick blocks, $p_+ + p_- - 2$ thin blocks and two semi-simple blocks.

1. For each $(r, s) \in K_{p_+, p_-}$, we denote by $C_{r, s}^{\text{thick}} = C_{p_+ - r, p_- - s}^{\text{thick}}$ the full abelian subcategory of \mathcal{C}_{p_+, p_-} such that

$$\begin{aligned} M \in C_{r, s}^{\text{thick}} \\ \Leftrightarrow \text{all composition factors of } M \text{ are given by } \mathcal{X}_{r, s}^+, \mathcal{X}_{r^\vee, s^\vee}^+, \\ \mathcal{X}_{r^\vee, s}^-, \mathcal{X}_{r, s^\vee}^- \text{ and } L(h_{r, s}). \end{aligned}$$

2. For each $1 \leq s \leq p_- - 1$, we denote by $C_{p_+, s}^{\text{thin}}$ the full abelian subcategory of \mathcal{C}_{p_+, p_-} such that

$$\begin{aligned} M \in C_{p_+, s}^{\text{thin}} \\ \Leftrightarrow \text{all composition factors of } M \text{ are given by } \mathcal{X}_{p_+, s}^+ \text{ and } \mathcal{X}_{p_+, s^\vee}^-. \end{aligned}$$

3. For each $1 \leq r \leq p_+ - 1$, we denote by C_{r, p_-}^{thin} the full abelian subcategory of \mathcal{C}_{p_+, p_-} such that

$$\begin{aligned} M \in C_{r, p_-}^{\text{thin}} \\ \Leftrightarrow \text{all composition factors of } M \text{ are given by } \mathcal{X}_{r, p_-}^+ \text{ and } \mathcal{X}_{r^\vee, p_-}^-. \end{aligned}$$

4. We denote by C_{p_+, p_-}^\pm the full abelian subcategory of \mathcal{C}_{p_+, p_-} such that

$$\begin{aligned} M &\in C_{p_+, p_-}^\pm \\ \Leftrightarrow \text{all composition factors of } M &\text{ are given by } \mathcal{X}_{p_+, p_-}^\pm. \end{aligned}$$

By using Theorem 5.6 in Section 5, we can prove the block decomposition of \mathcal{C}_{p_+, p_-} in the same way as Theorem 4.4 in [1]. The following proposition can be proved in the same way as Theorem 4.4 in [1].

Proposition 3.19. *The abelian category \mathcal{C}_{p_+, p_-} has the following block decomposition*

$$\mathcal{C}_{p_+, p_-} = \bigoplus_{(r,s) \in K_{p_+, p_-}} C_{r,s}^{thick} \oplus \bigoplus_{r=1}^{p_+-1} C_{r,p_-}^{thin} \oplus \bigoplus_{s=1}^{p_--1} C_{p_+,s}^{thin} \oplus C_{p_+, p_-}^+ \oplus C_{p_+, p_-}^-.$$

In the next section we will construct logarithmic modules in the thick blocks $C_{r,s}^{thick}$ and the thin blocks C_{r,p_-}^{thin} , $C_{p_+,s}^{thin}$.

4 Logarithmic \mathcal{W}_{p_+, p_-} modules

In this section, by using the logarithmic deformation by [17], we construct certain logarithmic \mathcal{W}_{p_+, p_-} -modules which correspond to the projective covers of all simple \mathcal{W}_{p_+, p_-} -modules $\mathcal{X}_{\bullet, \bullet}^\pm$ in the thick blocks and the thin blocks, and we introduce indecomposable modules $\mathcal{Q}(\mathcal{X}_{\bullet, \bullet}^\pm)_{\bullet, \bullet}$ which become important after this section. These logarithmic modules are closely related to certain indecomposable modules of the quantum group \mathfrak{g}_{p_+, p_-} at roots of unity [7],[13]. In the one thick block $C_{1,1}^{thick}$, these logarithmic modules are constructed by [6].

4.1 Logarithmic deformation

Proposition 4.1. *For $r, s \geq 1$, we have the following relation*

$$\alpha_-[Q_+^{[r]}, Q_-^{[s]}(z)] = \alpha_+[Q_-^{[s]}, Q_+^{[r]}(z)].$$

Proof. In the following, we use the notation

$$dx = dx_1 \cdots dx_{r-1}, \quad dy = dy_1 \cdots dy_{s-1}.$$

Recall the definition of the screening currents $Q_{\pm}^{[\bullet]}$ of (2.4).

$$\begin{aligned}
& \text{Res}_{z=w} Q_+^{[r]}(z) Q_-^{[s]}(w) \\
&= \text{Res}_{z=w} \int_{\bar{\Gamma}_r(\kappa_+)} d\mathbf{x} \int_{\bar{\Gamma}_s(\kappa_-)} d\mathbf{y} \frac{1}{(z-w)^2} : e^{\alpha_+ \phi(z) + \alpha_- \phi(w)} : \\
&\quad \times Q_-(zx_1) \cdots Q_+(zx_{r-1}) Q_-(wy_1) \cdots Q_-(wy_{s-1}) z^{r-1} w^{s-1} \\
&= \int_{\bar{\Gamma}_r(\kappa_+)} d\mathbf{x} \int_{\bar{\Gamma}_s(\kappa_-)} d\mathbf{y} \frac{\alpha_+}{\alpha_+ + \alpha_-} \left(\frac{\partial}{\partial w} V_{\alpha_+ + \alpha_-}(w) \right) \\
&\quad \times Q_-(wx_1) \cdots Q_+(wx_{r-1}) Q_-(wy_1) \cdots Q_-(wy_{s-1}) w^{r+s-2} \\
&+ \int_{\bar{\Gamma}_r(\kappa_+)} d\mathbf{x} \int_{\bar{\Gamma}_s(\kappa_-)} d\mathbf{y} z^{r-1} V_{\alpha_+ + \alpha_-}(z) \\
&\quad \times \frac{\partial}{\partial z} (Q_+(zx_1) \cdots Q_+(zx_{r-1})) Q_-(wy_1) \cdots Q_-(wy_{s-1}) w^{s-1} \Big|_{z=w} \\
&+ (r-1) \int_{\bar{\Gamma}_r(\kappa_+)} d\mathbf{x} \int_{\bar{\Gamma}_s(\kappa_-)} d\mathbf{y} \\
&\quad V_{\alpha_+ + \alpha_-}(w) Q_+(wx_1) \cdots Q_+(wx_{r-1}) Q_-(wy_1) \cdots Q_-(wy_{s-1}) w^{r+s-3}.
\end{aligned} \tag{4.1}$$

Since

$$\begin{aligned}
& \frac{\partial}{\partial z} (Q_+(zx_1) \cdots Q_+(zx_{r-1})) \\
&= \sum_{i=1}^{r-1} Q_+(zx_1) \cdots \frac{\partial}{\partial z} Q_+(zx_i) \cdots Q_+(zx_{r-1}) \\
&= \sum_{i=1}^{r-1} Q_+(zx_1) \cdots \left(\frac{1}{z} x_i \frac{\partial}{\partial x_i} Q_+(zx_i) \right) \cdots Q_+(zx_{r-1}) \\
&= \frac{1}{z} \sum_{i=1}^{r-1} Q_+(zx_1) \cdots \left(\frac{\partial}{\partial x_i} x_i Q_+(zx_i) \right) \cdots Q_+(zx_{r-1}) \\
&\quad - \frac{r-1}{z} Q_+(zx_1) \cdots Q_+(zx_{r-1}),
\end{aligned}$$

the second term of (4.1) becomes

$$\begin{aligned}
& - (r-1) \int_{\bar{\Gamma}_r(\kappa_+)} d\mathbf{x} \int_{\bar{\Gamma}_s(\kappa_-)} d\mathbf{y} \\
& \quad V_{\alpha_+ + \alpha_-}(w) Q_+(wx_1) \cdots Q_+(wx_{r-1}) Q_-(wy_1) \cdots Q_-(wy_{s-1}) w^{r+s-3} \\
& + \int_{\bar{\Gamma}_s(\kappa_-)} d\mathbf{y} \int_{\bar{\Gamma}_r(\kappa_+)} d\mathbf{x} V_{\alpha_+ + \alpha_-}(w) \\
& \quad \times d_x \left(\sum_{i=1}^{r-1} x_i Q_+(wx_1) \cdots Q_+(wx_{r-1}) dx_1 \cdots \widehat{dx_i} \cdots dx_{r-1} \right) \\
& \quad \times Q_-(wy_1) \cdots Q_-(wy_{s-1}).
\end{aligned} \tag{4.2}$$

The first term of (4.2) cancels with the third term of (4.1) and the second term of this equation becomes zero because $\bar{\Gamma}_r(\kappa_+)$ is the twisted cycle. Thus $[Q_+^{[r]}, Q_-^{[s]}(w)]$ becomes

$$\begin{aligned}
& \frac{\alpha_+}{\alpha_+ + \alpha_-} \int_{\bar{\Gamma}_r(\kappa_+)} d\mathbf{x} \int_{\bar{\Gamma}_s(\kappa_-)} d\mathbf{y} \\
& \quad \left(\frac{\partial}{\partial w} V_{\alpha_+ + \alpha_-}(w) \right) Q(w x_1) \cdots Q_+(wx_{r-1}) Q_-(wy_1) \cdots Q_-(wy_{s-1}) w^{r+s-2}.
\end{aligned}$$

In the same way, we have

$$\begin{aligned}
& [Q_-^{[s]}, Q_+^{[r]}(w)] \\
& = \frac{\alpha_-}{\alpha_+ + \alpha_-} \int_{\bar{\Gamma}_r(\kappa_+)} d\mathbf{x} \int_{\bar{\Gamma}_s(\kappa_-)} d\mathbf{y} \\
& \quad \left(\frac{\partial}{\partial w} V_{\alpha_+ + \alpha_-}(w) \right) Q(w x_1) \cdots Q_+(wx_{r-1}) Q_-(wy_1) \cdots Q_-(wy_{s-1}) w^{r+s-2}.
\end{aligned}$$

Therefore we obtain

$$\alpha_- [Q_+^{[r]}, Q_-^{[s]}(z)] = \alpha_+ [Q_-^{[s]}, Q_+^{[r]}(z)].$$

□

Proposition 4.2. *For $r, s \geq 1$ the screening operators $Q_+^{[r]}$ and $Q_-^{[s]}$ are \mathcal{W}_{p_+, p_-} -homomorphism, that is, for $A \in \mathcal{W}_{p_+, p_-}$ we have*

$$[Q_+^{[r]}, Y(A; z)] = 0, \quad [Q_-^{[s]}, Y(A; z)] = 0.$$

Proof. For each generator of \mathcal{W}_{p_+, p_-} , we have the following two expressions

$$\begin{aligned} W^+ &= Q_-^{[p_- - 1]} |\alpha_{1, p_- - 1; 3}\rangle = Q_+^{[3p_+ - 1]} |\alpha_{p_+ - 1, 1; -3}\rangle, \\ W^- &= Q_+^{[p_+ - 1]} |\alpha_{p_+ - 1, 1; -3}\rangle = Q_-^{[3p_- - 1]} |\alpha_{1, p_- - 1; 3}\rangle, \\ W^0 &= Q_+^{[2p_+ - 1]} |\alpha_{p_+ - 1, 1; -3}\rangle = Q_-^{[2p_- - 1]} |\alpha_{1, p_- - 1; 3}\rangle, \end{aligned}$$

up to non-zero constants. Thus, by the proof of Proposition 4.1, we obtain

$$[Q^{[r]}, Y(A; z)] = [Q_-^{[s]}, Y(A; z)] = 0.$$

□

We introduce the following logarithmic deformation introduced by [17].

Definition 4.3 ([17]). *1. Let $E(z)$ and $A(z)$ be any mutually local fields. We define the logarithmic deformation of $A(z)$ by $E(z)$ as follows*

$$\Delta_E(A(z)) = \log z (E[0]A)(z) + \sum_{n \geq 1} \frac{(-1)^{n+1}}{n} \frac{(E[n]A)(z)}{z^n},$$

where

$$(E[n]A)(w) = \oint_{z=w} (z-w)^n E(z)A(w)dz.$$

2. Let $E(z)$, $A(z)$ and $B(z)$ be any mutually local fields. We define

$$\Delta_E(A(z)B(w)) = \sum_{n \in \mathbb{Z}} \frac{\Delta_E((A[n]B)(w))}{(z-w)^{n+1}}.$$

Theorem 4.4 ([17]). *Let $E(z)$, $A(z)$ and $B(z)$ be any mutually local fields. Then the operator Δ_E satisfies the following derivation property*

$$\Delta_E(A(z)B(w)) = \Delta_E(A(z))B(w) + A(z)\Delta_E(B(w)).$$

In the next subsection, we consider the logarithmic deformations by the screening currents $Q_{\pm}^{[\bullet]}(z)$. We set

$$\Delta_+^{[r]} := \Delta_{Q_+^{[r]}}, \quad \Delta_-^{[s]} := \Delta_{Q_-^{[s]}}.$$

Note that, for the energy-momentum tensor, we have

$$\Delta_+^{[r]}(T(z)) = T(z) + \frac{Q_+^{[r]}(z)}{z}, \quad \Delta_-^{[s]}(T(z)) = T(z) + \frac{Q_-^{[s]}(z)}{z}. \quad (4.3)$$

By Proposition 4.2, each $\Delta_{\pm}^{[\bullet]}(Y(A; z))$, $A \in \mathcal{W}_{p_+, p_-}$ do not contain log terms in z .

4.2 Logarithmic modules in the thick block

For each $1 \leq r \leq p_+ - 1$, $1 \leq s \leq p_- - 1$ we set

$$\mathcal{P}_{r,s} = \mathcal{V}_{r,s}^+ \oplus \mathcal{V}_{r^\vee, s^\vee}^+ \oplus \mathcal{V}_{r, s^\vee}^- \oplus \mathcal{V}_{r^\vee, s}^-$$

where $r^\vee = p_+ - r$ and $s^\vee = p_- - s$. Note that $\mathcal{V}_{r,s}^+$, $\mathcal{V}_{r^\vee, s^\vee}^+$, $\mathcal{V}_{r, s^\vee}^-$, $\mathcal{V}_{r^\vee, s}^- \in C_{r,s}^{thick}$. Let $(\mathcal{P}_{r,s}, Y_{\mathcal{P}_{r,s}})$ be the ordinary \mathcal{W}_{p_+, p_-} -module. Fix any element $\tau = (a, b, \epsilon)$ in

$$\{(a, b, \epsilon)\} = \{(r, s, +), (r^\vee, s^\vee, +), (r^\vee, s, -), (r, s^\vee, -)\}.$$

For $A \in \mathcal{W}_{p_+, p_-}$, we define the following operators on $\mathcal{P}_{r,s}$:

$$\begin{aligned} & \tilde{\Delta}_\tau(Y_{\mathcal{P}_{r,s}}(A; z)) \\ = & \begin{cases} (\alpha_- - \alpha_+)(\Delta_+^{[a]} + \Delta_-^{[b]})(Y_{\mathcal{P}_{r,s}}(A; z)) \\ + (-\alpha_+ \Delta_-^{[b]} \circ \Delta_+^{[a]} + \alpha_- \Delta_+^{[a]} \circ \Delta_-^{[b]})(Y_{\mathcal{P}_{r,s}}(A; z)) & \text{on } \mathcal{V}_{a,b}^\epsilon \\ 0 & \text{on } \mathcal{P}_{r,s} \setminus \mathcal{V}_{a,b}^\epsilon \end{cases} \end{aligned}$$

$$\Delta_\tau^-(Y_{\mathcal{P}_{r,s}}(A; z)) = \begin{cases} \Delta_-^{[b]}(Y_{\mathcal{P}_{r,s}}(A; z)) & \text{on } \mathcal{V}_{a^\vee, b}^{-\epsilon} \\ 0 & \text{on } \mathcal{P}_{r,s} \setminus \mathcal{V}_{a^\vee, b}^{-\epsilon} \end{cases}$$

$$\Delta_\tau^+(Y_{\mathcal{P}_{r,s}}(A; z)) = \begin{cases} \Delta_+^{[a]}(Y_{\mathcal{P}_{r,s}}(A; z)) & \text{on } \mathcal{V}_{a, b^\vee}^{-\epsilon} \\ 0 & \text{on } \mathcal{P}_{r,s} \setminus \mathcal{V}_{a, b^\vee}^{-\epsilon} \end{cases}$$

By the following lemma, we can see that the operators $\tilde{\Delta}_\tau$ does not contain a $\log z$ terms.

Lemma 4.5. *For each $1 \leq r \leq p_+ - 1$, $1 \leq s \leq p_- - 1$ and $A \in \mathcal{W}_{p_+, p_-}$,*

$$-\alpha_+ \Delta_-^{[s]}(\Delta_+^{[r]}(Y(A; z))) + \alpha_- \Delta_+^{[r]}(\Delta_-^{[s]}(Y(A; z)))$$

does not contain log terms in z .

Proof. The $\log z$ terms of $\Delta_-^{[s]}(\Delta_+^{[r]}(Y(A; z)))$ and $\Delta_+^{[r]}(\Delta_-^{[s]}(Y(A; z)))$ are given by

$$[Q_-^{[s]}, \Delta_+^{[r]}(Y(A; z))]\log z, \quad [Q_+^{[r]}, \Delta_-^{[s]}(Y(A; z))]\log z.$$

By using Proposition 4.1 we have

$$\begin{aligned}
& [Q_-^{[s]}, \Delta_+^{[r]}(Y(A; z))] \\
&= \left[Q_-^{[s]}, \sum_{n \geq 1} \frac{(-1)^{n+1}}{n} \oint_{w=z} \frac{(w-z)^n}{z^n} Q_+^{[r]}(w) Y(A; z) dw \right] \\
&= \frac{\alpha_-}{\alpha_+} \left[Q_+^{[r]}, \sum_{n \geq 1} \frac{(-1)^{n+1}}{n} \oint_{w=z} \frac{(w-z)^n}{z^n} Q_-^{[s]}(w) Y(A; z) dw \right] \\
&= \frac{\alpha_-}{\alpha_+} [Q_+^{[r]}, \Delta_-^{[s]}(Y(A; z))].
\end{aligned}$$

Therefore

$$-\alpha_+ \Delta_-^{[s]}(\Delta_+^{[r]}(Y(A; z))) + \alpha_- \Delta_+^{[r]}(\Delta_-^{[s]}(Y(A; z)))$$

does not contain log terms in z . \square

Using Theorem 4.4, we can define logarithmic deformations of the ordinary \mathcal{W}_{p_+, p_-} -module $(\mathcal{P}_{r,s}, Y_{\mathcal{P}_{r,s}})$ as follows.

Theorem 4.6. *Fix any $\tau = (a, b, \epsilon)$ in*

$$\{(a, b, \epsilon)\} = \{(r, s, +), (r^\vee, s^\vee, +), (r^\vee, s, -), (r, s^\vee, -)\}.$$

We can define a logarithmic \mathcal{W}_{p_+, p_-} -module $\mathcal{P}_{a^\vee, b^\vee}^\epsilon$ as follows. As the vector space $\mathcal{P}_{a^\vee, b^\vee}^\epsilon = \mathcal{P}_{r,s}$ and the module actions is defined by

$$J_{a^\vee, b^\vee}^\epsilon(A; z) = Y_{\mathcal{P}_{r,s}}(A; z) + (\tilde{\Delta}_\tau + \Delta_\tau^+ + \Delta_\tau^-)(Y_{\mathcal{P}_{r,s}}(A; z)),$$

for any $A \in \mathcal{W}_{p_+, p_-}$.

Proof. By Lemma 4.5, we have $J_{a^\vee, b^\vee}^\epsilon : \mathcal{W}_{p_+, p_-} \rightarrow \text{End} \mathcal{P}_{a^\vee, b^\vee}^\epsilon[[z, z^{-1}]]$. The condition that $J_{a^\vee, b^\vee}^\epsilon(|0\rangle; z) = \text{id}_{\mathcal{P}_{a^\vee, b^\vee}^\epsilon}$ is trivial from the definition of logarithmic deformation. In the following we prove the compatibility condition

$$J_{a^\vee, b^\vee}^\epsilon(A; z) J_{a^\vee, b^\vee}^\epsilon(B; w) = J_{a^\vee, b^\vee}^\epsilon(Y(A; z - w)B; w)$$

for $A, B \in \mathcal{W}_{p_+, p_-}$. Fix any non-zero vector $v \in \mathcal{P}_{r,s}$ and write v be as follows

$$v = k_1 v_{a,b}^\epsilon + k_2 v_{a^\vee, b}^{-\epsilon} + k_3 v_{a, b^\vee}^{-\epsilon} + k_4 v_{a^\vee, b^\vee}^\epsilon,$$

where $v_{a,b}^\epsilon \in \mathcal{V}_{a,b}^\epsilon$, $v_{a^\vee,b}^{-\epsilon} \in \mathcal{V}_{a^\vee,b}^{-\epsilon}$, $v_{a,b^\vee}^{-\epsilon} \in \mathcal{V}_{a,b^\vee}^{-\epsilon}$, $v_{a^\vee,b^\vee}^\epsilon \in \mathcal{V}_{a^\vee,b^\vee}^\epsilon$, and $k_i (i = 1, \dots, 4)$ are constants. By using Theorem 4.4 we have

$$\begin{aligned}
& J_{a^\vee,b^\vee}^\epsilon(A; z) J_{a^\vee,b^\vee}^\epsilon(B; z) v_{a,b}^\epsilon \\
&= Y(A; z) Y(B; w) v_{a,b}^\epsilon \\
&\quad + (\alpha_- - \alpha_+) [\Delta_+^{[a]}(Y(A; z)) + \Delta_-^{[b]}(Y(A; z))] Y(B; w) v_{a,b}^\epsilon \\
&\quad + [-\alpha_+ \Delta_-^{[b]}(\Delta_+^{[a]}(Y(A; z))) + \alpha_- \Delta_+^{[a]}(\Delta_-^{[b]}(Y(A; z)))] Y(B; w) v_{a,b}^\epsilon \\
&\quad + (\alpha_- - \alpha_+) Y(A; z) [\Delta_+^{[a]}(Y(B; w)) + \Delta_-^{[b]}(Y(B; w))] v_{a,b}^\epsilon \\
&\quad + (\alpha_- - \alpha_+) [\Delta_-^{[b]}(Y(A; z)) \Delta_+^{[a]}(Y(B; w)) + \Delta_+^{[a]}(Y(A; z)) \Delta_-^{[b]}(Y(B; w))] v_{a,b}^\epsilon \\
&\quad + Y(A; z) [-\alpha_+ \Delta_-^{[b]}(\Delta_+^{[a]}(Y(B; w))) + \alpha_- \Delta_+^{[a]}(\Delta_-^{[b]}(Y(B; w)))] v_{a,b}^\epsilon \\
&= Y(A; z) Y(B; w) v_{a,b}^\epsilon \\
&\quad + (\alpha_- - \alpha_+) [\Delta_+^{[a]}(Y(A; z) Y(B; w)) + \Delta_-^{[b]}(Y(A; z) Y(B; w))] v_{a,b}^\epsilon \\
&\quad - \alpha_+ \Delta_-^{[b]} \circ \Delta_+^{[a]}(Y(A; z) Y(B; w)) v_{a,b}^\epsilon + \alpha_- \Delta_+^{[a]} \circ \Delta_-^{[b]}(Y(A; z) Y(B; w)) v_{a,b}^\epsilon \\
&= Y(Y(A; z - w) B; w) v_{a,b}^\epsilon + [(\tilde{\Delta}_\tau + \Delta_\tau^+ + \Delta_\tau^-)(Y(Y(A; z - w) B; w))] v_{a,b}^\epsilon \\
&= J_{a^\vee,b^\vee}^\epsilon(Y(A; z - w) B; w) v_{a,b}^\epsilon.
\end{aligned}$$

In the same way, we can prove

$$\begin{aligned}
J_{a^\vee,b^\vee}^\epsilon(A; z) J_{a^\vee,b^\vee}^\epsilon(B; z) v_{a^\vee,b}^{-\epsilon} &= J_{a^\vee,b^\vee}^\epsilon(Y(A; z - w) B; w) v_{a^\vee,b}^{-\epsilon}, \\
J_{a^\vee,b^\vee}^\epsilon(A; z) J_{a^\vee,b^\vee}^\epsilon(B; z) v_{a,b^\vee}^{-\epsilon} &= J_{a^\vee,b^\vee}^\epsilon(Y(A; z - w) B; w) v_{a,b^\vee}^{-\epsilon}, \\
J_{a^\vee,b^\vee}^\epsilon(A; z) J_{a^\vee,b^\vee}^\epsilon(B; z) v_{a^\vee,b^\vee}^\epsilon &= J_{a^\vee,b^\vee}^\epsilon(Y(A; z - w) B; w) v_{a^\vee,b^\vee}^\epsilon.
\end{aligned}$$

Therefore we obtain

$$J_{a^\vee,b^\vee}^\epsilon(A; z) J_{a^\vee,b^\vee}^\epsilon(B; z) v = J_{a^\vee,b^\vee}^\epsilon(Y(A; z - w) B; w) v.$$

□

By Proposition 2.5 and (4.3), we can see that the four logarithmic modules $\mathcal{P}_{\bullet,\bullet}^\pm \in C_{r,s}^{thick}$ have L_0 nilpotent rank three.

Remark 4.7. We will give the socle series of the logarithmic modules $\mathcal{P}_{\bullet,\bullet}^\pm$ in Subsection 6.4 (see Proposition 6.36). These logarithmic modules $\mathcal{P}_{r,s}^+$, $\mathcal{P}_{r^\vee,s^\vee}^+$, $\mathcal{P}_{r^\vee,s}^-$ and \mathcal{P}_{r,s^\vee}^- correspond to the projective covers of $\mathcal{X}_{r,s}^+$, $\mathcal{X}_{r^\vee,s^\vee}^+$, $\mathcal{X}_{r^\vee,s}^-$ and \mathcal{X}_{r,s^\vee}^- , respectively.

Proposition 4.8. By taking quotients of $\mathcal{P}_{r,s}^+$, $\mathcal{P}_{r^\vee,s^\vee}^+$, $\mathcal{P}_{r^\vee,s}^-$ and \mathcal{P}_{r,s^\vee}^- , we can define eight logarithmic modules $\mathcal{Q}(\mathcal{X}_{a,b}^\epsilon)_{b,c}$ where

$$\begin{aligned}
\{(\epsilon, a, b, c, d)\} = \{ & (+, r, s, r^\vee, s), (+, r, s, r, s^\vee), (+, r^\vee, s^\vee, r^\vee, s), (+, r^\vee, s^\vee, r, s^\vee), \\
& (-, r^\vee, s, r, s), (-, r^\vee, s, r^\vee, s^\vee), (-, r, s^\vee, r, s), (-, r, s^\vee, r^\vee, s^\vee) \},
\end{aligned}$$

and each composition series is given by:

1. For $\mathcal{Q}(\mathcal{X}_{a,b}^+)_{c,d}$, we have

$$G_1 \subsetneq G_2 \subsetneq G_3 \subsetneq G_4 = \mathcal{Q}(\mathcal{X}_{a,b}^+)_{c,d}$$

such that

$$\begin{aligned} G_1 &= \mathcal{X}_{a,b}^+, \\ G_2/G_1 \oplus G_3/G_2 &= \mathcal{X}_{c,d}^- \oplus L(h_{a,b}) \oplus \mathcal{X}_{c,d}^-, \\ \mathcal{Q}(\mathcal{X}_{a,b}^+)_{c,d}/G_3 &= \mathcal{X}_{a,b}^+. \end{aligned}$$

2. For $\mathcal{Q}(\mathcal{X}_{a,b}^-)_{c,d}$, we have

$$G_1 \subsetneq G_2 \subsetneq G_3 \subsetneq G_4 = \mathcal{Q}(\mathcal{X}_{a,b}^-)_{c,d}$$

such that

$$\begin{aligned} G_1 &= \mathcal{X}_{a,b}^-, \\ G_2/G_1 \oplus G_3/G_2 &= \mathcal{X}_{c,d}^+ \oplus \mathcal{X}_{c,d}^+, \\ \mathcal{Q}(\mathcal{X}_{a,b}^-)_{c,d}/G_3 &= \mathcal{X}_{a,b}^-. \end{aligned}$$

Remark 4.9. We will give the socle series of these logarithmic modules in Subsection 6.1.

4.3 Logarithmic modules in the thin blocks

For each $1 \leq r \leq p_+ - 1$, $1 \leq s \leq p_- - 1$, we set

$$\mathcal{P}_{r,p_-} = \mathcal{V}_{r,p_-}^+ \oplus \mathcal{V}_{r^\vee,p_-}^- \in C_{r,p_-}^{thin}, \quad \mathcal{P}_{p_+,s} = \mathcal{V}_{p_+,s}^+ \oplus \mathcal{V}_{p_+,s^\vee}^- \in C_{p_+,s}^{thin}.$$

Let $(\mathcal{P}_{r,p_-}, Y_{\mathcal{P}_{r,p_-}})$ and $(\mathcal{P}_{p_+,s}, Y_{\mathcal{P}_{p_+,s}})$ be the ordinary \mathcal{W}_{p_+,p_-} -module. Similar to Theorem 4.6, we can construct the following logarithmic modules.

Theorem 4.10.

1. For $1 \leq r \leq p_+ - 1$, we can define logarithmic modules $(\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}, J_{r,p_-}^+)$ and $(\mathcal{Q}(\mathcal{X}_{r^\vee,p_-}^-)_{r,p_-}, J_{r^\vee,p_-}^-)$ as follows. As the vector spaces

$$\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-} = \mathcal{Q}(\mathcal{X}_{r^\vee,p_-}^-)_{r,p_-} = \mathcal{P}_{r,p_-}$$

and the module actions are defined by

$$J_{r,p_-}^+(A; z) = \begin{cases} Y_{\mathcal{P}_{r,p_-}}(A; z) + \Delta_+^{[r^\vee]}(Y_{\mathcal{P}_{r,p_-}}(A; z)) & \text{on } \mathcal{V}_{r^\vee,p_-}^- \\ Y_{\mathcal{P}_{r,p_-}}(A; z) & \text{on } \mathcal{V}_{r,p_-}^+, \end{cases}$$

$$J_{r^\vee,p_-}^-(A; z) = \begin{cases} Y_{\mathcal{P}_{r,p_-}}(A; z) + \Delta_+^{[r]}(Y_{\mathcal{P}_{r,p_-}}(A; z)) & \text{on } \mathcal{V}_{r,p_-}^+ \\ Y_{\mathcal{P}_{r,p_-}}(A; z) & \text{on } \mathcal{V}_{r^\vee,p_-}^-, \end{cases}$$

for $A \in \mathcal{W}_{p_+,p_-}$.

2. For $1 \leq s \leq p_- - 1$, we can define logarithmic modules $(\mathcal{Q}(\mathcal{X}_{p_+,s}^+)_{p_+,s^\vee}, J_{p_+,s}^+)$ and $(\mathcal{Q}(\mathcal{X}_{p_+,s^\vee}^-)_{p_+,s}, J_{p_+,s}^-)$ as follows. As the vector spaces

$$\mathcal{Q}(\mathcal{X}_{p_+,s}^+)_{p_+,s^\vee} = \mathcal{Q}(\mathcal{X}_{p_+,s^\vee}^-)_{p_+,s} = \mathcal{P}_{p_+,s}$$

and the module actions are defined by

$$J_{p_+,s}^+(A; z) = \begin{cases} Y_{\mathcal{P}_{p_+,s}}(A; z) + \Delta_-^{[s^\vee]}(Y_{\mathcal{P}_{p_+,s}}(A; z)) & \text{on } \mathcal{V}_{p_+,s^\vee}^- \\ Y_{\mathcal{P}_{p_+,s}}(A; z) & \text{on } \mathcal{V}_{p_+,s}^+, \end{cases}$$

$$J_{p_+,s^\vee}^-(A; z) = \begin{cases} Y_{\mathcal{P}_{p_+,s}}(A; z) + \Delta_-^{[s]}(Y_{\mathcal{P}_{p_+,s}}(A; z)) & \text{on } \mathcal{V}_{p_+,s}^+ \\ Y_{\mathcal{P}_{p_+,s}}(A; z) & \text{on } \mathcal{V}_{p_+,s^\vee}^-. \end{cases}$$

for $A \in \mathcal{W}_{p_+,p_-}$.

Proposition 4.11. *The length of the composition series of the logarithmic modules $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}$, $\mathcal{Q}(\mathcal{X}_{r^\vee,p_-}^-)_{r,p_-}$, $\mathcal{Q}(\mathcal{X}_{p_+,s}^+)_{p_+,s^\vee}$ and $\mathcal{Q}(\mathcal{X}_{p_+,s^\vee}^-)_{p_+,s}$ are four, and each composition series is given by:*

1. The composition series of $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}$ is given by

$$G_1 = \mathcal{X}_{r,p_-}^+,$$

$$G_2/G_1 \oplus G_3/G_2 = \mathcal{X}_{r^\vee,p_-}^- \oplus \mathcal{X}_{r^\vee,p_-}^-,$$

$$\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}/G_3 = \mathcal{X}_{r,p_-}^+.$$

2. The composition series of $\mathcal{Q}(\mathcal{X}_{r^\vee,p_-}^-)_{r,p_-}$ is given by

$$G_1 = \mathcal{X}_{r^\vee,p_-}^-,$$

$$G_2/G_1 \oplus G_3/G_2 = \mathcal{X}_{r,p_-}^+ \oplus \mathcal{X}_{r,p_-}^+,$$

$$\mathcal{Q}(\mathcal{X}_{r^\vee,p_-}^-)_{r,p_-}/G_3 = \mathcal{X}_{r^\vee,p_-}^-.$$

3. The composition series of $\mathcal{Q}(\mathcal{X}_{p_+,s}^+)_{p_+,s^\vee}$ is given by

$$\begin{aligned} G_1 &= \mathcal{X}_{p_+,s}^+, \\ G_2/G_1 \oplus G_3/G_2 &= \mathcal{X}_{p_+,s^\vee}^- \oplus \mathcal{X}_{p_+,s^\vee}^-, \\ \mathcal{Q}(\mathcal{X}_{p_+,s}^+)_{p_+,s^\vee}/G_3 &= \mathcal{X}_{p_+,s}^+. \end{aligned}$$

4. The composition series of $\mathcal{Q}(\mathcal{X}_{p_+,s^\vee}^-)_{p_+,s}$ is given by

$$\begin{aligned} G_1 &= \mathcal{X}_{p_+,s^\vee}^-, \\ G_2/G_1 \oplus G_3/G_2 &= \mathcal{X}_{p_+,s}^+ \oplus \mathcal{X}_{p_+,s}^+, \\ \mathcal{Q}(\mathcal{X}_{p_+,s^\vee}^-)_{p_+,s}/G_3 &= \mathcal{X}_{p_+,s^\vee}^-. \end{aligned}$$

Remark 4.12. We will give the socle series of these logarithmic modules in Subsection 6.1. These logarithmic modules are projective modules in the thin blocks (see Subsection 6.3). The structure of these projective modules are similar to the projective modules of the triplet W -algebra \mathcal{W}_p determined by [25] and [27].

5 Logarithmic extension of Virasoro modules

In this section, we determine Ext^1 -groups between simple Virasoro modules and certain indecomposable modules in the abelian category of generalized Virasoro modules, by using the results in [24] and the structure of Fock modules. The results of this section will be crucial in analyzing the complex structure of logarithmic \mathcal{W}_{p_+,p_-} -modules. From this section, we identify any Virasoro modules that are isomorphic among each other.

5.1 Ext^1 -groups between simple Virasoro modules

We set

$$\begin{aligned} A_{p_+,p_-} &:= \{ \alpha_{r,s;n} \mid r, s, n \in \mathbb{Z} \}, \\ H_{p_+,p_-} &:= \left\{ h_\alpha \mid \alpha \in A_{p_+,p_-} \right\} \end{aligned}$$

(for the definition of symbols $\alpha_{r,s;n}$ and h_α , see (2.2) and (2.1), respectively). Let $U(\mathcal{L})$ be the universal enveloping algebra of the Virasoro algebra \mathcal{L} .

Definition 5.1. Let $\mathcal{L}_{c_{p_+,p_-}}\text{-Mod}$ be the abelian category of left generalized $U(\mathcal{L})$ -modules whose morphisms are Virasoro-homomorphisms and whose objects are left $U(\mathcal{L})$ -modules that satisfy the following conditions:

1. For the central charge, $C = c_{p_+, p_-} \cdot \text{id}$ on M .
2. Every object M has the following decomposition $M = \sum_{h \in H(M)} M[h]$:
 - For some finite subset $H_0(M)$ of \mathbb{C} , $H(M) = H_0(M) + \mathbb{Z}_{\geq 0}$.
 - For $h \in H(M)$, $M[h] = \{m \in M : \exists n \geq 0 \text{ s.t. } (L_0 - h)^n m = 0\}$.
 - $0 < \dim_{\mathbb{C}} M[h] < \infty$.
3. For every object $M \in \mathcal{L}_{c_{p_+, p_-}}\text{-Mod}$, there exists the contragredient object $M^* \in \mathcal{L}_{c_{p_+, p_-}}\text{-Mod}$ on which the anti-involution $\sigma(L_n) = L_{-n}$ induces the structure of a left $U(\mathcal{L})$ -module by

$$\langle L_n \phi, u \rangle = \langle \phi, \sigma(L_n) u \rangle, \quad \phi \in M^*, \quad u \in M.$$

4. We call any $M \in \mathcal{L}_{c_{p_+, p_-}}\text{-Mod}$ on which L_0 acts non-semisimply logarithmic.

Definition 5.2. We define $\mathcal{L}_{c_{p_+, p_-}}\text{-mod}$ to be the full subcategory of $\mathcal{L}_{c_{p_+, p_-}}\text{-Mod}$ such that all objects in $\mathcal{L}_{c_{p_+, p_-}}\text{-mod}$ satisfy the following conditions:

1. The socle series of M has finite length.
2. The highest weights h of the simple modules $L(h)$, appearing in the composition factors of M , are elements of H_{p_+, p_-} .

For $M_1, M_2 \in \mathcal{L}_{c_{p_+, p_-}}\text{-mod}$, we denote by $\text{Ext}_{\mathcal{L}}^1(M_1, M_2)$ the Ext^1 -group in $\mathcal{L}_{c_{p_+, p_-}}\text{-mod}$ consisting of $E \in \mathcal{L}_{c_{p_+, p_-}}\text{-mod}$ satisfying the exact sequence

$$0 \rightarrow M_2 \rightarrow E \rightarrow M_1 \rightarrow 0.$$

For $M_1, M_2 \in \mathcal{L}_{c_{p_+, p_-}}\text{-Mod}$, we denote by $\text{Hom}_{U(\mathcal{L})}(M_1, M_2)$ the set of the $U(\mathcal{L})$ -homomorphisms from M_1 to M_2 . Note that each of $\text{Ext}_{\mathcal{L}}^1(\bullet, \bullet)$ and $\text{Hom}_{U(\mathcal{L})}(\bullet, \bullet)$ has the structure of a \mathbb{C} -vector space.

Let $M(h, c_{p_+, p_-})$ be the Verma module of the Virasoro algebra whose highest weight is $h \in \mathbb{C}$ and the central charge $C = c_{p_+, p_-} \cdot \text{id}$. Let $|h\rangle$ be the highest weight vector of $M(h, c_{p_+, p_-})$. Note that, for $r, s \geq 1$, $M(h_{r,s}, c_{p_+, p_-})$ has the singular vector whose L_0 -weight is $h_{r,s} + rs$. Let $S_{r,s} \in U(\mathcal{L})$ be the Shapovalov element corresponding to this singular vector, normalized as

$$S_{r,s}|h_{r,s}\rangle = (L_{-1}^{rs} + \cdots)|h_{r,s}\rangle.$$

For $r, s \geq 1$ and $h \in \mathbb{C}$, let us consider the following value

$$\langle h | \sigma(S_{r,s}) S_{r,s} | h \rangle,$$

where we choose a norm of the highest weight vector $|h\rangle \in M(h, c_{p_+, p_-})$ as $\langle h|h\rangle = 1$. We can see that this value is a polynomial of h and is divisible by $(h - h_{r,s})$. A more detailed value is given by the following proposition.

Proposition 5.3 ([35]). *For $r, s \geq 1$ and $h \in \mathbb{C}$,*

$$\langle h|\sigma(S_{r,s})S_{r,s}|h\rangle = R_{r,s}(h - h_{r,s}) + O((h - h_{r,s})^2),$$

where $R_{r,s}$ is given by

$$R_{r,s} = 2 \prod_{\substack{(k,l) \in \mathbb{Z}^2, \\ 1-r \leq k \leq r, 1-s \leq l \leq s, \\ (k,l) \neq (0,0), (r,s)}} \left(k \left(\frac{p_+}{p_-} \right)^{-\frac{1}{2}} + l \left(\frac{p_+}{p_-} \right)^{\frac{1}{2}} \right).$$

Remark 5.4. *In this paper, it is important that $R_{r,s}$ be non-zero, specific value is not necessary. In fact, the non-triviality of $R_{r,s}$ can be shown using the Jantzen-filtration of the Fock module $F_{r,s}$.*

By using Proposition 5.3, we obtain the following proposition (cf.[20]).

Proposition 5.5. *For $h \in H_{p_+, p_-}$, we have*

$$\text{Ext}_{\mathcal{L}}^1(L(h), L(h)) = 0.$$

Proof. We prove only in the case of $h = h_{r,s}$ ($1 \leq r < p_+, 1 \leq s < p_-$). The other cases can be proved in the same way.

Assume $\text{Ext}_{\mathcal{L}}^1(L(h_{r,s}), L(h_{r,s})) \neq 0$. Fix a non-trivial extension

$$0 \rightarrow L(h_{r,s}) \xrightarrow{\iota} E \xrightarrow{\pi} L(h_{r,s}) \rightarrow 0.$$

Let $\{u_0, u_1\}$ be a basis of the highest weight space of E such that

$$\begin{aligned} \pi(u_0) &= |h_{r,s}\rangle, & \iota(|h_{r,s}\rangle) &= u_1, \\ (L_0 - h_{r,s})u_0 &= cu_1, \end{aligned}$$

where c is a non-zero constant and $|h_{r,s}\rangle$ is the highest weight vector of $L(h_{r,s})$. Then, by Proposition 5.3, we have

$$\begin{aligned} \sigma(S_{r,s})S_{r,s}u_0 &= f(c)u_1, \\ \frac{f(c)}{c} \Big|_{c=0} &\neq 0, \end{aligned}$$

where $f(c)$ is a polynomial of c . Thus, we see that $S_{r,s}u_0$ is non-zero and

$$S_{r,s}u_0 \in \iota(L(h_{r,s})).$$

On the other hand, by the irreducibility of $L(h_{r,s})$, we have

$$\sigma(S_{r,s})S_{r,s}u_0 = 0.$$

But this is a contradiction. □

The following proposition is due to [8],[11],[22].

Proposition 5.6 ([8, 11, 22]). *For any $r, s, \in \mathbb{Z}$ and $h \in \mathbb{C}$, the Ext^1 -group $\text{Ext}_{\mathcal{L}}^1(L(h_{r,s;n}), L(h))$ is given by:*

1. For $1 \leq r < p_+, 1 \leq s < p_-$ and $n = 0$, we have

$$\text{Ext}_{\mathcal{L}}^1(L(h_{r,s}), L(h)) = \begin{cases} \mathbb{C} & \text{for } h = h_{r^\vee, s; -1} \text{ or } h_{r^\vee, s; 1} \\ 0 & \text{otherwise} \end{cases}.$$

2. For $1 \leq r < p_+, 1 \leq s < p_-$ and $n \geq 1$, we have

$$\text{Ext}_{\mathcal{L}}^1(L(h_{r,s;n}), L(h)) = \begin{cases} \mathbb{C} & \text{for } h = h_{r^\vee, s; n \pm 1} \text{ or } h_{r, s^\vee; n \pm 1} \\ 0 & \text{otherwise} \end{cases}.$$

3. For $1 \leq r < p_+, s = p_-$ and $n = 0$, we have

$$\text{Ext}_{\mathcal{L}}^1(L(h_{r, p_-}), L(h)) = \begin{cases} \mathbb{C} & \text{for } h = h_{r^\vee, p_-; 1} \\ 0 & \text{otherwise} \end{cases}.$$

4. For $1 \leq r < p_+, s = p_-$ and $n \geq 1$, we have

$$\text{Ext}_{\mathcal{L}}^1(L(h_{r, p_-; n}), L(h)) = \begin{cases} \mathbb{C} & \text{for } h = h_{r^\vee, p_-; n+1} \text{ or } h_{r^\vee, p_-; n-1} \\ 0 & \text{otherwise} \end{cases}.$$

5. For $r = p_+, 1 \leq s < p_-$ and $n = 0$, we have

$$\text{Ext}_{\mathcal{L}}^1(L(h_{p_+, s}), L(h)) = \begin{cases} \mathbb{C} & \text{for } h = h_{p_+, s^\vee; -1} \\ 0 & \text{otherwise} \end{cases}.$$

6. For $r = p_+, 1 \leq s < p_-$ and $n \leq -1$, we have

$$\text{Ext}_{\mathcal{L}}^1(L(h_{p_+, s; n}), L(h)) = \begin{cases} \mathbb{C} & \text{for } h = h_{p_+, s^\vee; n-1} \text{ or } h_{p_+, s^\vee; n+1} \\ 0 & \text{otherwise} \end{cases}.$$

7. For $r = p_+, s = p_-, n \in \mathbb{Z}$, we have

$$\text{Ext}_{\mathcal{L}}^1(L(h_{p_+, p_-; n}), L(h)) = 0.$$

5.2 Logarithmic extensions

In the following, we identify any Virasoro modules that are isomorphic among each other. Let us define the following indecomposable modules in \mathcal{L}_{p_+, p_-} -mod as quotient modules of certain Virasoro Verma modules.

Definition 5.7. For $h, h' \in H_{p_+, p_-}$ such that $\text{Ext}_{\mathcal{L}}^1(L(h), L(h')) \simeq \mathbb{C}$ and $h < h'$, we identify the indecomposable modules in $\text{Ext}_{\mathcal{L}}^1(L(h), L(h'))$ and denote them as $L(h, h')$.

Definition 5.8. Given $1 \leq r < p_+$, $1 \leq s < p_-$, $n \geq 0$, by Proposition 5.6, we have

$$\text{Ext}_{\mathcal{L}}^1(L(h_{r,s;n}, h_{r^\vee, s; n+1}), L(h_{r, s^\vee; n+1})) \simeq \mathbb{C}.$$

We identify the indecomposable modules in this $\text{Ext}_{\mathcal{L}}^1$ -group and denote them as $\tilde{L}(h_{r,s;n})$.

The following theorems are due to [24] (see also [9]).

Theorem 5.9 ([24]). Fix any $h_1, h_2, h_3 \in H_{p_+, p_-}$ satisfying $h_1 < h_2 < h_3$, $\text{Ext}_{\mathcal{L}}^1(L(h_1), L(h_2)) \neq 0$ and $\text{Ext}_{\mathcal{L}}^1(L(h_2), L(h_3)) \neq 0$. Let us assume that there exists a logarithmic extension in

$$\text{Ext}_{\mathcal{L}}^1(L(h_2, h_3), L(h_1, h_2)).$$

Fix any logarithmic module E in this $\text{Ext}_{\mathcal{L}}^1$ -group. Then the quotient module $E/L(h_2)$ is indecomposable.

Theorem 5.10 ([24]). Given $1 \leq r < p_+$, $1 \leq s < p_-$, $n \geq 2$, let us assume that there exists a logarithmic extension in

$$\text{Ext}_{\mathcal{L}}^1(\tilde{L}(h_{r,s;n}), L(h_{a,b;n-1}, h_{r,s;n})),$$

where $(a, b) = (r^\vee, s)$ or (r, s^\vee) . Fix any logarithmic module E in this $\text{Ext}_{\mathcal{L}}^1$ -group. Then the quotient module $E/L(h_{r,s;n})$ is indecomposable.

Remark 5.11. The non-triviality of the logarithmic couplings of certain rank two Virasoro modules is proved in [24] (see Theorem 6.15 in [24]). The two theorems above are their consequences.

In the following, we introduce indecomposable modules $K(\tau)$ and $\tilde{K}(h_{r,s;n})$, and determine the $\text{Ext}_{\mathcal{L}}^1$ -groups of types

$$\text{Ext}_{\mathcal{L}}^1(K(\tau), L(h_{\alpha_2})), \quad \text{Ext}_{\mathcal{L}}^1(\tilde{K}(h_{r,s;n}), L(h_{r,s;n})).$$

Definition 5.12. We define \mathcal{T}_{p_+, p_-} to be the subset of A_{p_+, p_-}^3 such that every element $(\alpha_1, \alpha_2, \alpha_3) \in A_{p_+, p_-}^3$ satisfies the following conditions:

1. $h_{\alpha_1} \leq h_{\alpha_2} < h_{\alpha_3}$.
2. The three Fock modules F_{α_1} , F_{α_2} and F_{α_3} are contained in the same Felder complex in Proposition 2.6 and are adjacent to each other as

$$\cdots \rightarrow F_{\alpha_1} \xrightarrow{Q_\epsilon^{[\bullet]}} F_{\alpha_2} \xrightarrow{Q_\epsilon^{[\bullet]}} F_{\alpha_3} \rightarrow \cdots.$$

Definition 5.13. We define the following subsets of \mathcal{T}_{p_+, p_-} :

1. We define

$$\mathcal{T}_{p_+, p_-}^{\text{ch}} := \{(\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-} \mid F_{\alpha_1} \text{ is chain type}\}.$$

2. We define

$$\mathcal{T}_{p_+, p_-}^{\text{br}} := \{(\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-} \mid F_{\alpha_1} \text{ is braided type}\}.$$

3. We define

$$\mathcal{T}_{p_+, p_-}^0 := \{(\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}^{\text{ch}} \mid h_{\alpha_1} = h_{\alpha_2}\}.$$

4. We define

$$\mathcal{T}_{p_+, p_-}^{\text{Min}} = \{(\alpha_r, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}^{\text{br}} \mid \alpha_1 = \alpha_{r,s}, 1 \leq r < p_+, 1 \leq s < p_-\}.$$

Remark 5.14.

- Note that $\#\mathcal{T}_{p_+, p_-}^0 = (p_+ - 1)(p_- - 1)$ and every element of \mathcal{T}_{p_+, p_-}^0 is given by

$$(\alpha_{r, p_-; -1}, \alpha_{r, p_-; 0}, \alpha_{r, p_-; 1}), \quad (\alpha_{p_+, s^\vee; 1}, \alpha_{p_+, s; 0}, \alpha_{p_+, s^\vee; -1})$$

for $1 \leq r < p_+$, $1 \leq s < p_-$.

- Note that $\#\mathcal{T}_{p_+, p_-}^{\text{Min}} = (p_+ - 1)(p_- - 1)$ and every element of $\mathcal{T}_{p_+, p_-}^{\text{Min}}$ is given by

$$(\alpha_{r,s}, \alpha_{r^\vee, s; 1}, \alpha_{r,s; 2}), \quad (\alpha_{r,s}, \alpha_{r, s^\vee; -1}, \alpha_{r,s; -2})$$

for $1 \leq r < p_+$, $1 \leq s < p_-$.

Definition 5.15.

1. For any $\tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}^0$, we define $K(\tau) = L(h_{\alpha_2}, h_{\alpha_3})$.
2. Given $\tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}$ such that $h_{\alpha_1} \neq h_{\alpha_2}$, by Proposition 5.6, we have

$$\text{Ext}_{\mathcal{L}}^1(L(h_{\alpha_2}, h_{\alpha_3}), L(h_{\alpha_1})) \simeq \mathbb{C}.$$

We identify the indecomposable modules in this Ext^1 -group and denote them as $K(\tau)$.

The following proposition is a consequence of Theorem 5.9.

Proposition 5.16. For any $\tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}$, we have

$$\text{Ext}_{\mathcal{L}}^1(K(\tau), L(h_{\alpha_2})) \simeq \mathbb{C}. \quad (5.1)$$

Proof. Fix any $\tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}$. From the Virasoro subquotient structure of the logarithmic \mathcal{W}_{p_+, p_-} -modules $\mathcal{Q}(\mathcal{X}_{r,s}^{\pm})_{\bullet, \bullet}$ defined in Section 4 (see also logarithmic modules given in Definition 5.24), we have a logarithmic Virasoro module in

$$\text{Ext}_{\mathcal{L}}^1(K(\tau), L(h_{\alpha_2})) \setminus \{0\}.$$

Fix any logarithmic Virasoro module E_1 in this Ext^1 -group. If $\tau \in \mathcal{T}_{p_+, p_-}^0$, then we obtain the claim of theorem by Propositions 5.5 and 5.6, and thus let $\tau \notin \mathcal{T}_{p_+, p_-}^0$. It is sufficient to show that $\text{Ext}_{\mathcal{L}}^1(E_1, L(h_{\alpha_2})) = 0$. Let us assume that

$$\text{Ext}_{\mathcal{L}}^1(E_1, L(h_{\alpha_2})) \neq 0. \quad (5.2)$$

Note that, by Theorem 5.9, E_1 has $L(h_{\alpha_1}, h_{\alpha_2})$ as a submodule. Then, by the exact sequence

$$0 \rightarrow L(h_{\alpha_1}, h_{\alpha_2}) \rightarrow E_1 \rightarrow L(h_{\alpha_2}, h_{\alpha_3}) \rightarrow 0$$

and by the assumption (5.2), we have

$$\text{Ext}_{\mathcal{L}}^1(L(h_{\alpha_2}, h_{\alpha_3}), L(h_{\alpha_2})) \neq 0.$$

Let E_2 be any non-trivial extension in $\text{Ext}_{\mathcal{L}}^1(L(h_{\alpha_2}, h_{\alpha_3}), L(h_{\alpha_2}))$. Then, by Proposition 5.5, E_2 must have $L(h_{\alpha_2}, h_{\alpha_3})^*$ as a submodule. By the exact sequence

$$0 \rightarrow L(h_{\alpha_2}, h_{\alpha_3})^* \rightarrow E_2 \rightarrow L(h_{\alpha_2}) \rightarrow 0,$$

we have the following exact sequence

$$0 \rightarrow \mathbb{C} \rightarrow \text{Ext}_{\mathcal{L}}^1(E_2, L(h_{\alpha_1})).$$

Thus we have $\text{Ext}_{\mathcal{L}}^1(E_2, L(h_{\alpha_1})) \neq 0$. Let E_3 be any non-trivial extension of $\text{Ext}_{\mathcal{L}}^1(L(h_{\alpha_1}), E_2^*)$. By Proposition 5.3, we see that E_3 is logarithmic, that is, E_3 has L_0 nilpotent rank two. Note that

$$E_3/L(h_{\alpha_2}) = L(h_{\alpha_1}) \oplus L(h_{\alpha_2}, h_{\alpha_3}).$$

But this contradicts Theorem 5.9. \square

Definition 5.17. Given $\tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}$, we identify the indecomposable modules in the Ext^1 -group (5.1) and denote them as $P(\tau)$.

Remark 5.18. Given $\tau \in \mathcal{T}_{p_+, p_-}^0$, by Propositions 5.5 and 5.16, we see that the logarithmic Virasoro module $P(\tau)$ is self-contragredient.

The following indecomposable modules $\tilde{K}(h_{r,s;n})$ can be realized as the quotients of certain Fock modules.

Definition 5.19. Fix $(r, s) \in \mathbb{Z}^2$ such that $1 \leq r \leq p_+ - 1$, $1 \leq s \leq p_- - 1$.

1. By Proposition 5.6, indecomposable modules in the extension

$$\text{Ext}_{\mathcal{L}}^1(L(h_{r,s;1}), L(h_{r,s;0}) \oplus L(h_{r^\vee,s;2}) \oplus L(h_{r,s^\vee;2})) \setminus \{0\}$$

are unique up to isomorphism. We identify these indecomposable modules and denote them as $\tilde{K}(h_{r,s;1})$.

2. For $n \geq 2$, by Proposition 5.6, indecomposable modules in the extension

$$\text{Ext}_{\mathcal{L}}^1(L(h_{r,s;n}), \bigoplus_{i=-1,1} L(h_{r,s^\vee;n+i}) \oplus \bigoplus_{j=-1,1} L(h_{r^\vee,s;n+j})) \setminus \{0\}$$

are unique up to isomorphism. We identify these indecomposable modules and denote them as $\tilde{K}(h_{r,s;n})$.

The following proposition is a consequence of Theorem 5.10.

Proposition 5.20. For $1 \leq r \leq p_+ - 1$, $1 \leq s \leq p_- - 1$, $n \geq 2$, we have

$$\text{Ext}_{\mathcal{L}}^1(\tilde{K}(h_{r,s;n}), L(h_{r,s;n})) \simeq \mathbb{C}^2 \quad (5.3)$$

and there exists an indecomposable module in

$$\text{Ext}_{\mathcal{L}}^1(\tilde{K}(h_{r,s;n}), L(h_{r,s;n}) \oplus L(h_{r,s;n}))$$

such that has quotients isomorphic to $P(\tau_1)$ and $P(\tau_2)$, where

$$\begin{aligned} \tau_1 &= (\alpha_{r^\vee,s;n-1}, \alpha_{r,s;n}, \alpha_{r^\vee,s;n+1}), \\ \tau_2 &= (\alpha_{r^\vee,s;-n+1}, \alpha_{r^\vee,s^\vee;-n}, \alpha_{r^\vee,s;-n-1}). \end{aligned}$$

Proof. Noting Proposition 5.16, from the Virasoro subquotient structure of the \mathcal{W}_{p_+, p_-} -modules $\mathcal{P}_{\bullet, \bullet}^{\pm}$, we obtain an indecomposable module in

$$\text{Ext}_{\mathcal{L}}^1(\tilde{K}(h_{r,s;n}), L(h_{r,s;n}) \oplus L(h_{r,s;n}))$$

We denote by $\tilde{P}(h_{r,s;n})$ this indecomposable module. Note that, by Proposition 5.16, $\tilde{P}(h_{r,s;n})$ has $P(\tau_1)$ and $P(\tau_2)$ as quotients, where

$$\begin{aligned}\tau_1 &= (\alpha_{r^\vee, s; n-1}, \alpha_{r, s; n}, \alpha_{r^\vee, s; n+1}), \\ \tau_2 &= (\alpha_{r^\vee, s; -n+1}, \alpha_{r^\vee, s^\vee; -n}, \alpha_{r^\vee, s; -n-1}).\end{aligned}$$

Similar to the proof of Proposition 5.16, we can show

$$\text{Ext}_{\mathcal{L}}^1(\tilde{P}(h_{r,s;n}), L(h_{r,s;n})) = 0$$

by using Theorem 5.9 and Theorem 5.10. Thus, by Proposition 5.5, we obtain (5.3). \square

So far we have studied properties for Ext^1 -groups. In the following, we will investigate the subquotient structure of logarithmic Virasoro modules in the Ext^1 -groups

$$\text{Ext}_{\mathcal{L}}^1(K(\tau), L(h_{\alpha_2})), \quad \tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}^{\text{Min}}$$

and

$$\text{Ext}_{\mathcal{L}}^1(K(\tau), L(h_{\alpha_2}, h_{\alpha_3})^*), \quad \tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}^0 \sqcup \mathcal{T}_{p_+, p_-}^{\text{Min}}.$$

For that purpose, we will construct a family of infinite length logarithmic Virasoro modules by using certain limit operations. A similar construction is given by [10], [29].

Recall that \hat{a} be the dual of the zero mode a_0 defined by

$$[a_m, \hat{a}] = \delta_{m,0} \text{id}. \quad (5.4)$$

For any $\alpha, \beta \in \mathbb{C}$, let us identify

$$e^{\beta \hat{a}} |\alpha\rangle = |\alpha + \beta\rangle.$$

Let F_α ($\alpha \in A_{p_+, p_-}$) be any braided or chain type Fock module and let $Q_\epsilon^{[k]}$ be a screening operator acting on F_α , where $\epsilon = +$ or $-$. Let v be any L_0 -homogeneous vector of F_α and let $A \in U(\mathcal{L})$ be any L_0 -homogeneous

element. Let n_1 and n_2 be the L_0 -weight of v and A , respectively. For any $x \in \mathbb{C}^\times$, let us consider the following operator

$$[Q_\epsilon^{[k]}, e^{-x\hat{a}} A e^{x\hat{a}}] = Q_\epsilon^{[k]} e^{-x\hat{a}} A e^{x\hat{a}} - e^{-x\hat{a}} A e^{x\hat{a}} Q_\epsilon^{[k]}$$

on v , where $F_{\alpha+x}, F_{\alpha+k\alpha_\epsilon+x} \in \mathcal{F}_{\alpha_0}$ -mod. Let us write $[Q_\epsilon^{[k]}, e^{-x\hat{a}} A e^{x\hat{a}}]v$ as

$$[Q_\epsilon^{[k]}, e^{-x\hat{a}} A e^{x\hat{a}}]v = \sum_{\lambda \vdash N} f_\lambda(x) a_{-\lambda} |\alpha + k\alpha_\epsilon\rangle,$$

where $N = n_1 + n_2 + h_\alpha - h_{\alpha+k\alpha_\epsilon}$, and $f_\lambda(x)$ are some polynomials of x . Since $[Q_\epsilon^{[k]}, A] = 0$, we can see that every $f_\lambda(x)$ is divisible by x . Then we define

$$\lim_{\substack{x \rightarrow 0 \\ x \neq 0}} \frac{1}{x} [Q_\epsilon^{[k]}, e^{-x\hat{a}} A e^{x\hat{a}}]v := \sum_{\lambda \vdash N} (x^{-1} f_\lambda(x)|_{x=0}) a_{-\lambda} |\alpha + k\alpha_\epsilon\rangle.$$

We introduce the following \mathbb{C} -linear operators.

Definition 5.21. Let $F_\alpha (\alpha \in A_{p_+, p_-})$ be any braided or chain type Fock module and let $Q_\epsilon^{[k]}$ be a screening operator acting on F_α , where $\epsilon = +$ or $-$. We define the \mathbb{C} -linear operator $\Lambda_{Q_\epsilon^{[k]}} : U(\mathcal{L}) \rightarrow \text{Hom}_{\mathbb{C}}(F_\alpha, F_{\alpha+k\alpha_\epsilon})$ as follows

$$\Lambda_{Q_\epsilon^{[k]}}(A) = \lim_{\substack{x \rightarrow 0 \\ x \neq 0}} \frac{1}{x} [Q_\epsilon^{[k]}, e^{-x\hat{a}} A e^{x\hat{a}}], \quad \text{for } A \in U(\mathcal{L}),$$

where $F_{\alpha+x}, F_{\alpha+k\alpha_\epsilon+x} \in \mathcal{F}_{\alpha_0}$ -mod.

Proposition 5.22. The two operators $\Lambda_{Q_\epsilon^{[k]}}$ satisfy the following derivation property

$$\Lambda_{Q_\epsilon^{[k]}}(AB) = \Lambda_{Q_\epsilon^{[k]}}(A)B + A\Lambda_{Q_\epsilon^{[k]}}(B), \quad A, B \in U(\mathcal{L}).$$

Proof. For any $A, B \in U(\mathcal{L})$, we have

$$\begin{aligned} & [Q_\epsilon^{[k]}, e^{-x\hat{a}} A B e^{x\hat{a}}] \\ &= [Q_\epsilon^{[k]}, e^{-x\hat{a}} A e^{x\hat{a}} \cdot e^{-x\hat{a}} B e^{x\hat{a}}] \\ &= [Q_\epsilon^{[k]}, e^{-x\hat{a}} A e^{x\hat{a}}] e^{-x\hat{a}} B e^{x\hat{a}} + e^{-x\hat{a}} A e^{x\hat{a}} [Q_\epsilon^{[k]}, e^{-x\hat{a}} B e^{x\hat{a}}] \\ &= [Q_\epsilon^{[k]}, e^{-x\hat{a}} A e^{x\hat{a}}] B + A [Q_\epsilon^{[k]}, e^{-x\hat{a}} B e^{x\hat{a}}] \\ &\quad + [Q_\epsilon^{[k]}, e^{-x\hat{a}} A e^{x\hat{a}}] (e^{-x\hat{a}} B e^{x\hat{a}} - B) + (e^{-x\hat{a}} A e^{x\hat{a}} - A) [Q_\epsilon^{[k]}, e^{-x\hat{a}} B e^{x\hat{a}}]. \end{aligned}$$

Dividing both sides by x and taking the limit, we have the derivation property. \square

We define the following symbols for some screening operators.

Definition 5.23. 1. For any $\tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}$, we denote by Q_τ the screening operator from F_{α_1} to F_{α_2} , and we define $\Lambda_\tau = \Lambda_{Q_\tau}$.
 2. Fix any $\tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}^{\text{br}}$. Let $(r_\tau, s_\tau) \in \mathbb{Z}^2$ be a unique integer pair satisfying

$$(1 \leq r_\tau < p_+ \wedge 1 \leq s_\tau < p_-) \wedge (Q_+^{[r_\tau]} \text{ and } Q_-^{[s_\tau]} \text{ are acting on } F_{\alpha_1})$$

(note that $F_{\alpha_2} = F_{\alpha_1 + r_\tau \alpha_+}$ or $F_{\alpha_1 + s_\tau \alpha_-}$). We define $\tilde{Q}_\tau = Q_+^{[r_\tau]} + Q_-^{[s_\tau]}$ and $\tilde{\Lambda}_\tau = \Lambda_{Q_+^{[r_\tau]}} + \Lambda_{Q_-^{[s_\tau]}}$.

We define the following indecomposable Virasoro modules by gluing certain Fock modules.

Definition 5.24. 1. Fix any $\tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}$. We set

$$F(\tau) = F_{\alpha_1} \oplus F_{\alpha_2}.$$

For $A \in U(\mathcal{L})$, we define the following operator $J_\tau(A)$ on $F(\tau)$

$$J_\tau(A) = \begin{cases} A + \Lambda_\tau(A) & \text{on } F_{\alpha_1}, \\ A & \text{on } F_{\alpha_2}. \end{cases}$$

Then, by Proposition 5.22, we have

$$J_\tau(AB) = J_\tau(A)J_\tau(B), \quad \text{for any } A, B \in U(\mathcal{L}).$$

Thus we see that J_τ defines a structure of Virasoro module on $F(\tau)$. We denote this Virasoro module by $(F(\tau), J_\tau)$.

2. Fix any $\tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}^{\text{br}}$. We set

$$\tilde{F}(\tau) = F_{\alpha_1} \oplus F_{\alpha_1 + r_\tau \alpha_+} \oplus F_{\alpha_1 + s_\tau \alpha_-}.$$

For $A \in U(\mathcal{L})$, we define the following operator $\tilde{J}_\tau(A)$ on $\tilde{F}(\tau)$

$$\tilde{J}_\tau(A) = \begin{cases} A + \tilde{\Lambda}_\tau(A) & \text{on } F_{\alpha_1}, \\ A & \text{on } F_{\alpha_1 + r_\tau \alpha_+} \oplus F_{\alpha_1 + s_\tau \alpha_-}. \end{cases}$$

Then, by Proposition 5.22, we have

$$\tilde{J}_\tau(AB) = \tilde{J}_\tau(A)\tilde{J}_\tau(B), \quad \text{for any } A, B \in U(\mathcal{L}).$$

Thus we see that \tilde{J}_τ defines a structure of Virasoro module on $\tilde{F}(\tau)$. We denote this Virasoro module by $(\tilde{F}(\tau), \tilde{J}_\tau)$.

The Virasoro modules $(F(\tau), J_\tau)$ and $(\tilde{F}(\tau), \tilde{J}_\tau)$ have L_0 -nilpotent rank two. In fact we have the following lemma.

Lemma 5.25. 1. Fix any $\tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}$. The $J_\tau(L_n)$ action on the subspace $F_{\alpha_1} \subset F(\tau)$ is given by

$$J_\tau(L_n) = L_n + [Q_\tau, a_n]$$

2. Fix any $\tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}^{\text{br}}$. The $\tilde{J}_\tau(L_n)$ action on the subspace $F_{\alpha_1} \subset \tilde{F}(\tau)$ is given by

$$\tilde{J}_\tau(L_n) = L_n + [\tilde{Q}_\tau, a_n]$$

Proof. We only prove the first case. The second case can be proved in the same way.

Note that the ordinary action of L_n on the Fock modules in \mathcal{F}_{α_0} -mod is given by

$$L_n = \frac{1}{2} \sum_{m \in \mathbb{Z}} : a_m a_{n-m} : - \frac{1}{2} \alpha_0 (n+1) a_n. \quad (5.5)$$

Then, by (5.4), (5.5) and $[Q_\tau, L_n] = 0$, we have

$$\begin{aligned} J_\tau(L_n) - L_n &= \Lambda_\tau(L_n) \\ &= \lim_{x \rightarrow 0} \frac{1}{x} Q_\tau e^{-x\hat{a}} L_n e^{x\hat{a}} - \lim_{x \rightarrow 0} \frac{1}{x} e^{-x\hat{a}} L_n e^{x\hat{a}} Q_\tau \\ &= \lim_{x \rightarrow 0} \frac{1}{x} Q_\tau x a_n - \lim_{x \rightarrow 0} \frac{1}{x} x a_n Q_\tau \\ &= [Q_\tau, a_n]. \end{aligned}$$

□

Theorem 5.26. For any $\tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}^{\text{Min}}$, we have

$$\text{Soc}(P(\tau)) = L(h_{\alpha_2}).$$

In particular $P(\tau)$ is self-contragredient.

Proof. Note that $\mathcal{T}_{p_+, p_-}^{\text{Min}}$ is given by the following set

$$\{(\alpha_{r^\vee, s^\vee}, \alpha_{r, s^\vee; 1}, \alpha_{r^\vee, s^\vee; 2}), (\alpha_{r^\vee, s^\vee}, \alpha_{r^\vee, s; -1}, \alpha_{r^\vee, s^\vee; -2}) \mid 1 \leq r < p_+, 1 \leq s < p_-\}.$$

Fix any $1 \leq r < p_+, 1 \leq s < p_-$ and set

$$\tau = (\alpha_{r^\vee, s^\vee}, \alpha_{r, s^\vee; 1}, \alpha_{r^\vee, s^\vee; 2}), \quad \tau' = (\alpha_{r^\vee, s^\vee}, \alpha_{r^\vee, s; -1}, \alpha_{r^\vee, s^\vee; -2}).$$

Let us show

$$\text{Soc}(P(\tau)) = \text{Soc}(P(\tau')) = L(\Delta_{r,s;0}^+) \quad (5.6)$$

(for the definition of $\Delta_{r,s;0}^+$, see Definition 3.6). Let us consider the vector $e^{-x\hat{a}} S_{r^\vee, s^\vee} e^{x\hat{a}} |\alpha_{r^\vee, s^\vee}\rangle \in F_{r^\vee, s^\vee}$ for $x \in \mathbb{C}^\times$, where $F_{\alpha_{r^\vee, s^\vee} + x} \in \mathcal{F}_{\alpha_0}$ -mod. Let us write

$$e^{-x\hat{a}} S_{r^\vee, s^\vee} e^{x\hat{a}} |\alpha_{r^\vee, s^\vee}\rangle = \sum_{\lambda \vdash r^\vee s^\vee} f_\lambda(x) a_{-\lambda} |\alpha_{r^\vee, s^\vee}\rangle,$$

where $f_\lambda(x)$ are some polynomials of x . Since $S_{r^\vee, s^\vee} |\alpha_{r^\vee, s^\vee}\rangle = 0$, we can see that every $f_\lambda(x)$ is divisible by x . Then we define

$$u_\tau := \sum_{\lambda \vdash r^\vee s^\vee} (x^{-1} f_\lambda(x) |_{x=0}) a_{-\lambda} |\alpha_{r^\vee, s^\vee}\rangle \in F_{r^\vee, s^\vee}[r^\vee s^\vee].$$

By the Jantzen filtration of the Fock module F_{r^\vee, s^\vee} (cf. [11, 22]), we can see that this vector is non-zero, and

$$Q_+^{[r^\vee]}(u_\tau) \in \mathbb{C}^\times |\alpha_{r, s^\vee; 1}\rangle, \quad Q_-^{[s^\vee]}(u_\tau) \in \mathbb{C}^\times |\alpha_{r^\vee, s; -1}\rangle$$

(cf. [16]).

Let us consider the logarithmic module $(\tilde{F}(\tau), \tilde{J}_\tau)$. By Lemma 5.25, we have

$$(\tilde{J}_\tau(L_0) - h_{r, s^\vee; 1})u_\tau = -r^\vee Q_+^{[r^\vee]}(u_\tau) - s^\vee Q_-^{[s^\vee]}(u_\tau). \quad (5.7)$$

in $(\tilde{F}(\tau), \tilde{J}_\tau)$. By using Proposition 5.3 we have

$$\tilde{J}_\tau(S_{r^\vee, s^\vee} \sigma(S_{r^\vee, s^\vee}))u_\tau = \frac{1}{2}(2\alpha_{r^\vee, s^\vee} - \alpha_0)R_{r^\vee, s^\vee}(Q_+^{[r^\vee]}(u_\tau) + Q_-^{[s^\vee]}(u_\tau)) \quad (5.8)$$

in $(\tilde{F}(\tau), \tilde{J}_\tau)$.

Let us consider the following finite length submodule of $(\tilde{F}(\tau), \tilde{J}_\tau)$

$$E_\tau := \tilde{J}_\tau(U(\mathcal{L})).u_\tau.$$

By (5.7) and (5.8), E_τ has a indecomposable quotient in

$$\text{Ext}_{\mathcal{L}}^1(\tilde{K}(\Delta_{r,s;0}^+), L(\Delta_{r,s;0}^+) \oplus L(\Delta_{r,s;0}^+))$$

(for the definition of $\tilde{K}(\Delta_{r,s;0}^+)$, see Definition 5.19). We denote this quotient by $\tilde{P}(\Delta_{r,s;0}^+)$. Note that by Proposition 5.16, $\tilde{P}(\Delta_{r,s;0}^+)$ has $P(\tau)$ and $P(\tau')$, as quotients.

Let us consider the quotient module $E_\tau/\tilde{J}_\tau(U(\mathcal{L})).|\alpha_{\tau^\vee,s^\vee}\rangle$. By (5.7) and (5.8), this quotient module has L_0 nilpotent rank two and has a logarithmic quotient isomorphic to $\tilde{P}(\Delta_{r,s;0}^+)/L(h_{r,s}, \Delta_{r,s;0}^+)$. We set $R(\Delta_{r,s;0}^+) = \tilde{P}(\Delta_{r,s;0}^+)/L(h_{r,s}, \Delta_{r,s;0}^+)$. Then, by Propositions 5.5 and 5.16, we have

$$\text{Soc}(R(\Delta_{r,s;0}^+)) = L(\Delta_{r,s;0}^+).$$

Therefore we obtain the claim (5.6). \square

Remark 5.27. By Proposition 5.16, we have

$$\text{Ext}_{\mathcal{L}}^1(R(\Delta_{r,s;0}^+), L(\Delta_{r,s;0}^+)) = 0.$$

From this, we have $\text{Ext}_{\mathcal{L}}^1(\tilde{P}(\Delta_{r,s;0}^+), L(\Delta_{r,s;0}^+)) = 0$.

Fix any element $\tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}^0 \sqcup \mathcal{T}_{p_+, p_-}^{\text{Min}}$. By Proposition 5.16, we see that

$$\text{Ext}_{\mathcal{L}}^1(K(\tau), L(h_{\alpha_2}, h_{\alpha_3})^*) \simeq \mathbb{C}.$$

We identify the indecomposable modules in this Ext^1 -group and denote them as $\hat{P}(\tau)$. From the definition of $\hat{P}(\tau)$, we can easily see that

$$\begin{aligned} \dim_{\mathbb{C}} \text{Hom}_{U(\mathcal{L})}(L(h_{\alpha_2}, h_{\alpha_3})^*, \hat{P}(\tau)) &\leq 2, \\ \dim_{\mathbb{C}} \text{Hom}_{U(\mathcal{L})}(\hat{P}(\tau), K(\tau)) &\leq 2. \end{aligned} \tag{5.9}$$

Note that this logarithmic module $\hat{P}(\tau)$ can be realized as the quotient of $(F(\tau), J_\tau)$ or $(\tilde{F}(\tau), \tilde{J}_\tau)$. The following proposition is trivial from the self-dualities $P(\tau)^* \simeq P(\tau)$ and $\hat{P}(\tau)^* \simeq \hat{P}(\tau)$, but we give a direct proof using the structure of $(F(\tau), J_\tau)$ and $(\tilde{F}(\tau), \tilde{J}_\tau)$.

Proposition 5.28. Fix any $\tau = (\alpha_1, \alpha_2, \alpha_3) \in \mathcal{T}_{p_+, p_-}^0 \sqcup \mathcal{T}_{p_+, p_-}^{\text{Min}}$. Then $\hat{P}(\tau)$ satisfies the following properties:

1. We have

$$\text{Hom}_{U(\mathcal{L})}(L(h_{\alpha_2}, h_{\alpha_3})^*, \hat{P}(\tau)) \simeq \mathbb{C}^2, \tag{5.10}$$

and we can choose two injections as a basis of (5.10).

2. We have

$$\text{Hom}_{U(\mathcal{L})}(\hat{P}(\tau), K(\tau)) \simeq \mathbb{C}^2, \tag{5.11}$$

and we can choose two surjections as a basis of (5.11).

Proof. We only prove in the case of $\tau = (\alpha_{r^\vee, p_-; -1}, \alpha_{r, p_-}, \alpha_{r^\vee, p_-; 1}) \in \mathcal{T}_{p_+, p_-}^0$. The case where $\tau \in \mathcal{T}_{p_+, p_-}^{\text{Min}}$ can be proved similarly by considering the structure of $(\tilde{F}(\tau), \tilde{J}_\tau)$.

Let us consider the logarithmic module $(F(\tau), J_\tau)$. Set $u_0 = |\alpha_{r^\vee, p_-; -1}\rangle$ and $u_1 = |\alpha_{r, p_-}\rangle$. Fix any homogeneous vector $v_1 \in F_{r^\vee, p_-; -1}[rp_-]$ such that $Q_+^{[r]}(v_1) \in \mathbb{C}^\times |\alpha_{r^\vee, p_-; 1}\rangle$, and let $v_2 = Q_+^{[r]}(|\alpha_{r, p_-; -2}\rangle)$. Note that

$$\sigma(S_{r, p_-})v_1 \in \mathbb{C}^\times |\alpha_{r, p_-}\rangle. \quad (5.12)$$

Then by the Jantzen filtration of the Fock modules $F_{r^\vee, p_-; -1}$ and F_{r, p_-} , we obtain

$$\begin{aligned} J_\tau(\sigma(S_{r, p_-}))v_2 &\in \mathbb{C}^\times u_1, \\ J_\tau(S_{r, p_-})u_0 &\in \mathbb{C}^\times v_2 + \mathbb{C}^\times v_1 + U(\mathcal{L}).u_1. \end{aligned} \quad (5.13)$$

Note that the submodule

$$I = J_\tau(U(\mathcal{L})).u_0 + J_\tau(U(\mathcal{L})).v_1$$

has a quotient isomorphic to $\hat{P}(\tau)$. Let π be the surjection from I to $\hat{P}(\tau)$, and define the following injections and surjections

$$\begin{aligned} \iota_i &: L(h_{r, p_-}, h_{r^\vee, p_-; 1})^* \xrightarrow{\cong} \pi(J_\tau(U(\mathcal{L})).v_i) \subset \hat{P}(\tau), \quad i = 1, 2, \\ p_j &: \hat{P}(\tau) \rightarrow \hat{P}(\tau)/\pi(J_\tau(U(\mathcal{L})).v_j), \quad j = 1, 2. \end{aligned}$$

Then, from (5.9), (5.12) and (5.13), we can see that $\{\iota_1, \iota_2\}$ and $\{p_1, p_2\}$ give a basis of (5.10) and (5.11), respectively. \square

Remark 5.29. Figure 5.1 represents the embedding structure of the logarithmic modules $\hat{P}(\tau)$ ($\tau = (\alpha_{r^\vee, p_-; -1}, \alpha_{r, p_-}, \alpha_{r^\vee, p_-; 1}) \in \mathcal{T}_{p_+, p_-}^0$). The symbols correspond to those introduced in the proof of Proposition 5.28.

6 The projective covers of simple \mathcal{W}_{p_+, p_-} -modules

Since \mathcal{W}_{p_+, p_-} is C_2 -cofinite, so by [21], every simple \mathcal{W}_{p_+, p_-} -module has the projective cover. In this section, we determine some Ext^1 -groups between certain indecomposable modules and simple modules. Based on these Ext^1 groups, we determine the structure of projective covers of the simple modules in each thick block and thin block. From this section, we denote the Ext^1 -groups in \mathcal{C}_{p_+, p_-} as $\text{Ext}^1(\bullet, \bullet)$ simply and identify any \mathcal{W}_{p_+, p_-} -modules that are isomorphic among each other, unless otherwise stated.




Figure 5.1: The schematic diagram of the logarithmic modules $\widehat{P}(\tau)$ given in the proof of Proposition 5.28.

6.1 The structure of the logarithmic modules $\mathcal{Q}(\mathcal{X}_{\bullet,\bullet}^{\pm})_{\bullet,\bullet}$

Fix any $(r, s) \in \mathbb{Z}_{\geq 1}^2$ such that $1 \leq r \leq p_+ - 1$ and $1 \leq s \leq p_- - 1$. In this subsection we consider the structure of the indecomposable modules $\mathcal{Q}(\mathcal{X}_{\bullet,\bullet}^{\pm})_{\bullet,\bullet}$ in the blocks C_{r,p_-}^{thin} , $C_{p_+,s}^{thin}$ and $C_{r,s}^{thick}$. First let us consider the structure of the logarithmic module $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-} \in C_{r,p_-}^{thin}$. Recall that as the vector space $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-} = \mathcal{V}_{r^\vee,p_-}^- \oplus \mathcal{V}_{r,p_-}^+$. Fix any $v_1^+ \in F_{r,p_-}[rp_-]$ satisfying $Q_+^{[r]}(v_1^+) \in \mathbb{C}^\times |\alpha_{r^\vee,p_-;1}\rangle$, and set

$$\begin{aligned} u_0 &= |\alpha_{r^\vee,p_-;-1}\rangle, & u_1 &= |\alpha_{r,p_-}\rangle, \\ v_2^+ &= |\alpha_{r^\vee,p_-;1}\rangle, & v_1^- &= |\alpha_{r,p_-;-2}\rangle, & v_2^- &= Q_+^{[r]}(|\alpha_{r,p_-;-2}\rangle). \end{aligned} \quad (6.1)$$

Let us consider the following Virasoro submodule of $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}$

$$I = U(\mathcal{L}).u_0 + U(\mathcal{L}).v_1^+ \in \mathcal{L}_{p_+,p_-}\text{-mod},$$

where we omit the symbol J_{r,p_-}^+ associated to the module action defined in Theorem 4.10. Note that

$$\begin{aligned} (L_0 - \Delta_{r,p_-;0}^+)u_0 &\in \mathbb{C}^\times u_1, \\ \sigma(S_{r,p_-})v_1^+ &\in \mathbb{C}^\times u_1 \end{aligned} \quad (6.2)$$

in $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}$. Then, by (6.2), we see that I has a quotient isomorphic to $\widehat{P}(\tau)$, where $\tau = (\alpha_{r^\vee,p_-;-1}, \alpha_{r,p_-}, \alpha_{r^\vee,p_-;1})$. Thus, by Proposition 5.28, we

have

$$\begin{aligned} S_{r,p_-} u_0 &\in \mathbb{C}^\times v_1^+ + \mathbb{C}^\times v_2^- + U(\mathcal{L}).u_1, \\ \sigma(S_{r,p_-}) v_2^- &\in \mathbb{C}^\times u_1, \end{aligned} \quad (6.3)$$

in $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}$. By (6.3) we see that $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}$ has two submodules $\mathcal{W}_{p_+,p_-}.v_1^+$ and $\mathcal{W}_{p_+,p_-}.v_2^-$ satisfying

$$\begin{aligned} [\mathcal{W}_{p_+,p_-}.v_1^+] &\in \text{Ext}^1(\mathcal{X}_{r^\vee,p_-}^-, \mathcal{X}_{r,p_-}^+) \setminus \{0\}, \\ [\mathcal{W}_{p_+,p_-}.v_2^-] &\in \text{Ext}^1(\mathcal{X}_{r^\vee,p_-}^-, \mathcal{X}_{r,p_-}^+) \setminus \{0\}. \end{aligned} \quad (6.4)$$

By (6.4), we see that $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}$ is generated from the top composition factor \mathcal{X}_{r,p_-}^+ and $\text{Soc}(\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}) = \mathcal{X}_{r,p_-}^+$.

We have similar results for the other indecomposable module $\mathcal{Q}(\mathcal{X}_{p_+,s}^+)_{p_+,s^\vee}$ in $C_{p_+,s}^{\text{thin}}$. Thus we obtain the following proposition.

Proposition 6.1. *The logarithmic modules $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}$ and $\mathcal{Q}(\mathcal{X}_{p_+,s}^+)_{p_+,s^\vee}$ are generated from the top composition factors and have the following length three socle series:*

1. *The socle series of $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}$ is given by*

$$\begin{aligned} \text{Soc}_1 &= \mathcal{X}_{r,p_-}^+, \\ \text{Soc}_2/\text{Soc}_1 &= \mathcal{X}_{r^\vee,p_-}^- \oplus \mathcal{X}_{r^\vee,p_-}^-, \\ \mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}/\text{Soc}_2 &= \mathcal{X}_{r,p_-}^+. \end{aligned}$$

2. *The socle series of $\mathcal{Q}(\mathcal{X}_{p_+,s}^+)_{p_+,s^\vee}$ is given by*

$$\begin{aligned} \text{Soc}_1 &= \mathcal{X}_{p_+,s}^+, \\ \text{Soc}_2/\text{Soc}_1 &= \mathcal{X}_{p_+,s^\vee}^- \oplus \mathcal{X}_{p_+,s^\vee}^-, \\ \mathcal{Q}(\mathcal{X}_{p_+,s}^+)_{p_+,s^\vee}/\text{Soc}_2 &= \mathcal{X}_{p_+,s}^+. \end{aligned}$$

Remark 6.2. *Figure 6.1 represents relations between generators of the composition factors of $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}$. The symbols correspond to those introduced in (6.1).*

Next let us consider the logarithmic module $\mathcal{Q}(\mathcal{X}_{r^\vee,p_-}^-)_{r,p_-}$. Recall that as the vector space $\mathcal{Q}(\mathcal{X}_{r^\vee,p_-}^-)_{r,p_-} = \mathcal{V}_{r^\vee,p_-}^- \oplus \mathcal{V}_{r,p_-}^+$. Fix any $\tilde{v}_+ \in F_{r,p_-}[rp_-]$ satisfying $Q_+^{[r]}(\tilde{v}_+) \in \mathbb{C}^\times |\alpha_{r^\vee,p_-;-1}\rangle$, and set

$$\begin{aligned} u_- &= |\alpha_{r^\vee,p_-;-1}\rangle, \quad u_+ = |\alpha_{r,p_-}\rangle, \\ v_+ &= |\alpha_{r^\vee,p_-;1}\rangle, \quad v_- = Q_+^{[r]}(|\alpha_{r,p_-;-2}\rangle), \quad \tilde{v}_- = |\alpha_{r,p_-;-2}\rangle. \end{aligned} \quad (6.5)$$




Figure 6.1: The schematic diagram of the indecomposable module $\mathcal{Q}(\mathcal{X}_{r,p-}^+)_{r^\vee,p-}$, where $W_0^\pm = W^\pm[0]$.

Let us consider the indecomposable Virasoro modules

$$I^\pm = U(\mathcal{L}).\tilde{v}_\pm.$$

Note that

$$(L_0 - \Delta_{r^\vee,p-;0}^-)\tilde{v}_\pm \in \mathbb{C}^\times v_\pm, \quad (6.6)$$

in $\mathcal{Q}(\mathcal{X}_{r^\vee,p-}^-)_{r,p-}$ (we omit the symbol $J_{r^\vee,p-}^-$ associated the module action defined in Theorem 4.10). Then, noting Theorem 5.9, Proposition (5.16) and (6.6), we see that I^ϵ ($\epsilon = \pm$) has a quotient isomorphic to $P(\tau)$, where $\tau = (\alpha_{r,p-}, \alpha_{r^\vee,p-;1}, \alpha_{r,p-;2})$ (for the definition of $P(\tau)$, see Definition 5.17). Then, from the structure of $P(\tau)$, we obtain

$$S_{r,p-}\sigma(S_{r,p-})\tilde{v}_\pm \in \mathbb{C}^\times v_\pm. \quad (6.7)$$

By (6.7) we see that $\mathcal{Q}(\mathcal{X}_{r^\vee,p-}^-)_{r,p-}$ has the submodules $\mathcal{W}_{p+,p-}.\sigma(S_{r,p-})\tilde{v}_\pm$ satisfying

$$[\mathcal{W}_{p+,p-}.\sigma(S_{r,p-})\tilde{v}_\pm] \in \text{Ext}^1(\mathcal{X}_{r,p-}^+, \mathcal{X}_{r^\vee,p-}^-) \setminus \{0\}.$$

In particular we see that $\mathcal{Q}(\mathcal{X}_{r^\vee,p-}^-)_{r,p-}$ is generated from the top composition factor $\mathcal{X}_{r^\vee,p-}^-$. We have similar results for the other indecomposable module $\mathcal{Q}(\mathcal{X}_{p+,s^\vee}^-)_{p+,s}$ in $C_{p+,s}^{thin}$. Thus we obtain the following theorem.

Proposition 6.3. *The logarithmic modules $\mathcal{Q}(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}$ and $\mathcal{Q}(\mathcal{X}_{p_+, s^\vee}^-)_{p_+, s}$ are generated from the top composition factors and have the following length three socle series:*

1. *The socle series of $\mathcal{Q}(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}$ is given by*

$$\begin{aligned}\text{Soc}_1 &= \mathcal{X}_{r^\vee, p_-}^-, \\ \text{Soc}_2/\text{Soc}_1 &= \mathcal{X}_{r, p_-}^+ \oplus \mathcal{X}_{r, p_-}^+, \\ \mathcal{Q}(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}/\text{Soc}_2 &= \mathcal{X}_{r^\vee, p_-}^-.\end{aligned}$$

2. *The socle series of $\mathcal{Q}(\mathcal{X}_{p_+, s^\vee}^-)_{p_+, s}$ is given by*

$$\begin{aligned}\text{Soc}_1 &= \mathcal{X}_{p_+, s^\vee}^-, \\ \text{Soc}_2/\text{Soc}_1 &= \mathcal{X}_{p_+, s}^+ \oplus \mathcal{X}_{p_+, s}^+, \\ \mathcal{Q}(\mathcal{X}_{p_+, s^\vee}^-)_{p_+, s}/\text{Soc}_2 &= \mathcal{X}_{p_+, s^\vee}^-.\end{aligned}$$




Figure 6.2: The schematic diagram of the structure of the indecomposable module $\mathcal{Q}(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}$.

Remark 6.4. *Figure 6.2 represents relations between generators of the composition factors of $\mathcal{Q}(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}$. The symbols correspond to those introduced in (6.5).*

Similar to Theorems 6.1 and 6.3, by using Proposition 5.16 and Proposition 5.28, we obtain the following theorems for the indecomposable modules $\mathcal{Q}(\mathcal{X}_{\bullet, \bullet}^\pm)_{\bullet, \bullet}$ in $C_{r, s}^{thick}$. We omit the proofs.

Proposition 6.5. *Let (a, b, c, d) be any element in*

$$\{(a, b, c, d)\} = \{(r, s, r^\vee, s), (r, s, r, s^\vee), (r^\vee, s^\vee, r^\vee, s), (r^\vee, s^\vee, r, s^\vee)\}.$$

Then the socle series of $\mathcal{Q}(\mathcal{X}_{a,b}^+)_{c,d}$ is given by

$$\begin{aligned} \text{Soc}_1 &= \text{Socle} = \mathcal{X}_{a,b}^+, \\ \text{Soc}_2/\text{Soc}_1 &= \mathcal{X}_{c,d}^- \oplus L(h_{a,b}) \oplus \mathcal{X}_{c,d}^-, \\ \mathcal{Q}(\mathcal{X}_{a,b}^+)_{c,d}/\text{Soc}_2 &= \mathcal{X}_{a,b}^+. \end{aligned}$$

Moreover, $\mathcal{Q}(\mathcal{X}_{a,b}^+)_{c,d}$ is generated from the top composition factor $\mathcal{X}_{a,b}^+$.

Proposition 6.6. *Let (a, b, c, d) be any element in*

$$\{(a, b, c, d)\} = \{(r^\vee, s, r, s), (r^\vee, s, r^\vee, s^\vee), (r, s^\vee, r, s), (r, s^\vee, r^\vee, s^\vee)\}.$$

Then the socle series of $\mathcal{Q}(\mathcal{X}_{a,b}^-)_{c,d}$ is given by

$$\begin{aligned} \text{Soc}_1 &= \text{Socle} = \mathcal{X}_{a,b}^-, \\ \text{Soc}_2/\text{Soc}_1 &= \mathcal{X}_{c,d}^+ \oplus \mathcal{X}_{c,d}^+, \\ \mathcal{Q}(\mathcal{X}_{a,b}^-)_{c,d}/\text{Soc}_2 &= \mathcal{X}_{a,b}^-. \end{aligned}$$

Moreover, $\mathcal{Q}(\mathcal{X}_{a,b}^-)_{c,d}$ is generated from the top composition factor $\mathcal{X}_{a,b}^-$.

6.2 The Ext^1 -groups between all simple modules

Fix any $(r, s) \in \mathbb{Z}_{\geq 1}^2$ such that $1 \leq r \leq p_+ - 1$ and $1 \leq s \leq p_- - 1$. In this subsection, we determine the Ext^1 -groups between all simple modules.

Definition 6.7. *Fix any (a, b, c, d) in*

$$\begin{aligned} \{(a, b, c, d)\} = \{ & (r^\vee, s, r, s), (r^\vee, s, r^\vee, s^\vee), (r, s^\vee, r, s), (r, s^\vee, r^\vee, s^\vee), \\ & (r^\vee, p_-, r, p_-), (p_+, s^\vee, p_+, s)\}. \end{aligned}$$

1. *For $\mathcal{Q}(\mathcal{X}_{a,b}^-)_{c,d}$, let $\{v_+, v_-\}$ be a basis of the highest weight space of the submodule $\mathcal{X}_{a,b}^- \subset \mathcal{Q}(\mathcal{X}_{a,b}^-)_{c,d}$ such that*

$$W^\pm[0]v_\pm = 0, \quad W^\pm[0]v_\mp \in \mathbb{C}^\times v_\pm,$$

and let u_\pm be the vectors in $\mathcal{Q}(\mathcal{X}_{a,b}^-)_{c,d}[\Delta_{c,d;0}^+]$ satisfying $v_\pm \in U(\mathcal{L}).u_\pm$. Then we define

$$\mathcal{E}^+(\mathcal{X}_{c,d}^+)_{a,b} := \mathcal{W}_{p_+, p_-}.u_+, \quad \mathcal{E}^-(\mathcal{X}_{c,d}^+)_{a,b} := \mathcal{W}_{p_+, p_-}.u_-,$$

which give different extensions in $\text{Ext}^1(\mathcal{X}_{c,d}^+, \mathcal{X}_{a,b}^-) \setminus \{0\}$.

2. We define

$$\mathcal{E}^\pm(\mathcal{X}_{a,b}^-)_{c,d} := \mathcal{Q}(\mathcal{X}_{a,b}^-)_{c,d} / \mathcal{E}^\mp(\mathcal{X}_{c,d}^+)_{a,b}.$$

3. As the quotient of $\mathcal{Q}(\mathcal{X}_{c,d}^+)_{a,b}$, we define the indecomposable module $\tilde{\mathcal{E}}(\mathcal{X}_{c,d}^+)_{a,b}$ satisfying the exact sequence

$$0 \rightarrow \mathcal{X}_{a,b}^- \oplus \mathcal{X}_{a,b}^- \rightarrow \tilde{\mathcal{E}}(\mathcal{X}_{c,d}^+)_{a,b} \rightarrow \mathcal{X}_{c,d}^+ \rightarrow 0.$$

4. We define $\tilde{\mathcal{E}}(\mathcal{X}_{a,b}^-)_{c,d} := \mathcal{Q}(\mathcal{X}_{a,b}^-)_{c,d} / \mathcal{X}_{a,b}^-$, which satisfies the exact sequence

$$0 \rightarrow \mathcal{X}_{c,d}^+ \oplus \mathcal{X}_{c,d}^+ \rightarrow \tilde{\mathcal{E}}(\mathcal{X}_{a,b}^-)_{c,d} \rightarrow \mathcal{X}_{a,b}^- \rightarrow 0.$$

Definition 6.8. Given a non-logarithmic Virasoro module M , any nonzero vector $v \in M$ is called a primary vector when the following satisfied

$$L_n v = 0, \quad n \geq 1.$$

Similar to the arguments in Section 9.3 of [22], we have the following proposition (see also [26],[30]). We omit the proofs.

Proposition 6.9 ([11],[22]). Let M_1 , M_2 and M_3^* be non-logarithmic Virasoro modules which have primary vectors $v_1 \in M_1$, $v_2 \in M_2$ and $v_3^* \in M_3^*$ whose L_0 weights are h_{r_1,s_1} , h_{r_2,s_2} and h_{r_3,s_3} , respectively, where $r_i \geq 1$ and $s_i \geq 1$ ($i = 1, 2, 3$). Assume that there exists a non-logarithmic intertwining operator \mathcal{Y} of type $\begin{pmatrix} M_3 \\ M_1 \ M_2 \end{pmatrix}$. Then we have

$$\begin{aligned} \langle v_3^*, \mathcal{Y}(v_1, z) S_{r_2,s_2} v_2 \rangle &= \prod_{i=1}^{r_2} \prod_{j=1}^{s_2} (h_{r_1,s_1} - h_{r_2+r_3-2i+1,s_2+s_3-2j+1}) \langle v_3^*, \mathcal{Y}(v_1, z) v_2 \rangle, \\ \langle S_{r_3,s_3} v_3^*, \mathcal{Y}(v_1, z) v_2 \rangle &= \prod_{i=1}^{r_3} \prod_{j=1}^{s_3} (h_{r_1,s_1} - h_{r_2+r_3-2i+1,s_2+s_3-2j+1}) \langle v_3^*, \mathcal{Y}(v_1, z) v_2 \rangle. \end{aligned}$$

Proposition 6.10 ([11],[22]). For $h \in \mathbb{C}$, $1 \leq r_1, r_2 < p_+$, $1 \leq s_1, s_2 < p_-$ and $n_1, n_2 \in \mathbb{Z}_{\geq 0}$, we have

$$\mathcal{N}_{L(h_{r_1,s_1;n_1}), L(h_{r_2,s_2;n_2})}^{L(h)} \leq 1,$$

where $\mathcal{N}_{L(h_2), L(h_1)}^{L(h_3)}$ is the dimension of the space of Virasoro intertwining operators of type $\begin{pmatrix} L(h_3) \\ L(h_2) \ L(h_1) \end{pmatrix}$. If $\mathcal{N}_{L(h_{r_1,s_1;n_1}), L(h_{r_2,s_2;n_2})}^{L(h)} \neq 0$, then h is the common

solution of the following equations

$$\begin{aligned}
& \prod_{i=1}^{r_1} \prod_{j=1}^{s_1+n_1p_-} (h - h_{r_1+r_2-2i+1, s_1+s_2-2j+1; n_1+n_2}) = 0, \\
& \prod_{i=1}^{(n_1+1)p_+-r_1} \prod_{j=1}^{p_--s_1} (h - h_{2p_+-r_1-r_2-2i+1, 2p_--s_1-s_2-2j+1; -n_1-n_2}) = 0, \\
& \prod_{i=1}^{r_2} \prod_{j=1}^{s_2+n_2p_-} (h - h_{r_1+r_2-2i+1, s_1+s_2-2j+1; n_1+n_2}) = 0, \\
& \prod_{i=1}^{(n_2+1)p_+-r_2} \prod_{j=1}^{p_--s_2} (h - h_{2p_+-r_1-r_2-2i+1, 2p_--s_1-s_2-2j+1; -n_1-n_2}) = 0.
\end{aligned}$$

From this subsection, we use the following notation.

Definition 6.11. For any simple \mathcal{W}_{p_+, p_-} -module \mathcal{X} and any $n \geq 1$, we define

$$n\mathcal{X} := \overbrace{\mathcal{X} \oplus \cdots \oplus \mathcal{X}}^n.$$

Lemma 6.12. Let $n \geq 1$. Fix any (a, b, c, d) in

$$\begin{aligned}
\{(a, b, c, d)\} = \{ & (r, s, r^\vee, s), (r^\vee, s^\vee, r^\vee, s), (r, s, r, s^\vee), (r^\vee, s^\vee, r, s^\vee), \\
& (r, p_-, r^\vee, p_-), (p_+, s, p_+, s^\vee)\}.
\end{aligned}$$

Any extension $[E] \in \text{Ext}^1(\mathcal{X}_{a,b}^+, n\mathcal{X}_{c,d}^-)$ is trivial if E satisfies

$$\text{Hom}_{U(\mathcal{L})}(L(\Delta_{a,b;0}^+), E) \neq 0. \quad (6.8)$$

Proof. We only prove in the case of $(a, b, c, d) = (r, s, r^\vee, s)$ and $n = 1$. The other cases can be proved in the same way.

Fix any non-trivial extension $[E] \in \text{Ext}(\mathcal{X}_{r,p_-}^+, \mathcal{X}_{r^\vee,p_-}^-) \setminus \{0\}$. Let u be the highest weight vector in $E[\Delta_{r,s;0}^+]$. Assume (6.8). Then we have

$$S_{r,s^\vee+p_-}u = 0. \quad (6.9)$$

Let $\{v_+, v_-\}$ be a basis of the highest weight space of the submodule $\mathcal{X}_{r^\vee,s}^- \subset E$ such that

$$W^\pm[0]v_\pm = 0, \quad W^\pm[0]v_\mp \in \mathbb{C}^\times v_\pm.$$

Let v_+^* and v_-^* be L_0 -homogeneous vectors of E^* such that $\langle v_\pm^*, v_\pm \rangle \neq 0$, and $L_k v_\pm^* = 0$ for $k \geq 1$. Assume that for any $W = W^\pm, W^0$

$$W[k]v_\pm^* = 0, \quad k \geq 1.$$

Then the vector space $\mathbb{C}v_+^* + \mathbb{C}v_-^*$ becomes a $A(\mathcal{W}_{p_+, p_-})$ -module and this vector space is isomorphic to the highest weight space of $\mathcal{X}_{r^\vee, s}^-$ as a $A(\mathcal{W}_{p_+, p_-})$ -module. Thus E^* has the submodule $\mathcal{W}_{p_+, p_-}(\mathbb{C}v_+^* + \mathbb{C}v_-^*) \simeq \mathcal{X}_{r^\vee, s}^-$ and thus $E^* = \mathcal{X}_{r, s}^+ \oplus \mathcal{X}_{r^\vee, s}^-$. But this contradicts the assumption that E is non-trivial. Therefore we have

$$\langle v_\pm^*, Y_E(W^\bullet; z)u \rangle \neq 0,$$

where W^\bullet is one of W^+ , W^0 or W^- . On the other hand, using Proposition 6.9, we have

$$\begin{aligned} & \langle v_\pm^*, Y_E(W^\bullet; z)S_{r, s^\vee + p_-}u \rangle \\ &= \prod_{i=1}^r \prod_{j=1}^{s^\vee + p_-} (h_{4p_+ - 1, 1} - h_{r+r+2p_+ - 2i+1, s^\vee + p_- + s - 2j+1}) \langle v_\pm^*, Y_E(W^\bullet; z)u \rangle \\ &\neq 0. \end{aligned}$$

But this contradicts (6.9). □

Similar to the triplet W -algebras \mathcal{W}_p [2],[15],[25],[27], we have the following proposition.

Proposition 6.13.

1. In the thin block C_{r, p_-}^{thin} , we have

$$\text{Ext}^1(\mathcal{X}_{r, p_-}^+, \mathcal{X}_{r^\vee, p_-}^-) = \text{Ext}^1(\mathcal{X}_{r^\vee, p_-}^-, \mathcal{X}_{r, p_-}^+) = \mathbb{C}^2.$$

The other extensions between the simple modules in this block are trivial.

2. In the thin block $C_{p_+, s}^{thin}$, we have

$$\text{Ext}^1(\mathcal{X}_{p_+, s}^+, \mathcal{X}_{p_+, s^\vee}^-) = \text{Ext}^1(\mathcal{X}_{p_+, s^\vee}^-, \mathcal{X}_{p_+, s}^+) = \mathbb{C}^2.$$

The other extensions between the simple modules in this block are trivial.

Proof. We will only prove

$$\begin{aligned}\mathrm{Ext}^1(\mathcal{X}_{r^\vee, p_-}^-, \mathcal{X}_{r^\vee, p_-}^-) &= 0, & \mathrm{Ext}^1(\mathcal{X}_{r, p_-}^+, \mathcal{X}_{r^\vee, p_-}^-) &= \mathbb{C}^2, \\ \mathrm{Ext}^1(\mathcal{X}_{r, p_-}^+, \mathcal{X}_{r, p_-}^+) &= 0.\end{aligned}$$

The other Ext^1 -groups can be proved in a similar way, so we omit the proofs.

First let us prove $\mathrm{Ext}^1(\mathcal{X}_{r^\vee, p_-}^-, \mathcal{X}_{r^\vee, p_-}^-) = 0$. Fix any extension $[E_1] \in \mathrm{Ext}^1(\mathcal{X}_{r^\vee, p_-}^-, \mathcal{X}_{r^\vee, p_-}^-)$. By Proposition 5.5, we see that L_0 acts semisimply on E_1 . Let \overline{E}_1 be the highest weight space of E_1 . Note that E_1 is generated from \overline{E}_1 . Let \widetilde{E}_1 be the \mathcal{W}_{p_+, p_-} -module induced from \overline{E}_1 . Then we have $\widetilde{E}_1 = E_1$. By Proposition 3.12, we see that as the $A(\mathcal{W}_{p_+, p_-})$ -module

$$\overline{E}_1 \simeq \overline{\mathcal{X}_{r^\vee, p_-}^-} \oplus \overline{\mathcal{X}_{r^\vee, p_-}^-},$$

where $\overline{\mathcal{X}_{r^\vee, p_-}^-}$ is the highest weight space of $\mathcal{X}_{r^\vee, p_-}^-$. Note that the \mathcal{W}_{p_+, p_-} -module induced from $\overline{\mathcal{X}_{r^\vee, p_-}^-}$ is isomorphic to $\mathcal{X}_{r^\vee, p_-}^-$. Thus we have $\widetilde{E}_1 \simeq \mathcal{X}_{r^\vee, p_-}^- \oplus \mathcal{X}_{r^\vee, p_-}^-$.

Next, we prove $\mathrm{Ext}^1(\mathcal{X}_{r, p_-}^+, \mathcal{X}_{r^\vee, p_-}^-) = \mathbb{C}^2$. Since $\mathrm{Ext}^1(\mathcal{X}_{r^\vee, p_-}^-, \mathcal{X}_{r^\vee, p_-}^-) = 0$, it is sufficient to show that

$$\mathrm{Ext}^1(\widetilde{\mathcal{E}}(\mathcal{X}_{r, p_-}^+)_{r^\vee, p_-}, \mathcal{X}_{r^\vee, p_-}^-) = 0,$$

where $\widetilde{\mathcal{E}}(\mathcal{X}_{r, p_-}^+)_{r^\vee, p_-}$ is given in Definition 6.7. Assume that

$$\mathrm{Ext}^1(\widetilde{\mathcal{E}}(\mathcal{X}_{r, p_-}^+)_{r^\vee, p_-}, \mathcal{X}_{r^\vee, p_-}^-) \neq 0$$

and fix any non-trivial extension E_2 in this Ext^1 -group. By $\mathrm{Ext}^1(\mathcal{X}_{r^\vee, s}^-, \mathcal{X}_{r^\vee, s}^-) = 0$, we see that

$$\mathrm{Soc}(E_2) = \mathcal{X}_{r^\vee, p_-}^- \oplus \mathcal{X}_{r^\vee, p_-}^- \oplus \mathcal{X}_{r^\vee, p_-}^-. \quad (6.10)$$

Let u be the highest weight vector of E_2 and let us consider the submodule $\mathcal{W}_{p_+, p_-} \cdot S_{r, p_-} u$ of E_2 . By Theorems 3.10 and 3.11, we can see that

$$\mathcal{W}_{p_+, p_-} \cdot S_{r, p_-} u = \mathcal{X}_{r^\vee, p_-}^- \text{ or } \mathcal{X}_{r^\vee, p_-}^- \oplus \mathcal{X}_{r^\vee, p_-}^-. \quad (6.11)$$

Thus by (6.10) and (6.11), we have

$$[E_2 / \mathcal{W}_{p_+, p_-} \cdot S_{r, p_-} u] \in \mathrm{Ext}^1(\mathcal{X}_{r, p_-}^+, n\mathcal{X}_{r^\vee, p_-}^-) \setminus \{0\},$$

where n is 1 or 2. But this contradicts Lemma 6.12. Therefore we obtain

$$\mathrm{Ext}^1(\mathcal{X}_{r, p_-}^+, \mathcal{X}_{r^\vee, p_-}^-) = \mathbb{C}^2.$$

Finally let us prove $\text{Ext}^1(\mathcal{X}_{r,p_-}^+, \mathcal{X}_{r,p_-}^+) = 0$. Let $\overline{\mathcal{X}_{r,p_-}^+}$ be the highest weight space of \mathcal{X}_{r,p_-}^+ . Since

$$\text{Ext}^1(\mathcal{X}_{r^\vee,p_-}^-, \mathcal{X}_{r^\vee,p_-}^-) = 0, \quad \text{Ext}^1(\mathcal{X}_{r,p_-}^+, \mathcal{X}_{r^\vee,p_-}^-) = \mathbb{C}^2,$$

we see that the induced \mathcal{W}_{p_+,p_-} -module from $\overline{\mathcal{X}_{r,p_-}^+}$ is isomorphic to $\tilde{\mathcal{E}}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}$. Let E_3 be any extension in $\text{Ext}^1(\mathcal{X}_{r,p_-}^+, \mathcal{X}_{r,p_-}^+)$. By Proposition 5.5, we see that L_0 acts semisimply on E_3 . Let \overline{E}_3 be the highest weight space of E_3 . Note that E_3 is generated from \overline{E}_3 . Let us assume $\overline{E}_3 \not\cong \overline{\mathcal{X}_{r,p_-}^+} \oplus \overline{\mathcal{X}_{r,p_-}^+}$ as a $A(\mathcal{W}_{p_+,p_-})$ -module. Then, from the \mathcal{W}_{p_+,p_-} -module action on E_3 , we have a non-trivial non-logarithmic Virasoro intertwining operator of type

$$\begin{pmatrix} L(\Delta_{r,p_-}^+; 0) \\ L(h_{4p_+-1,1}) & L(\Delta_{r,p_-}^+; 0) \end{pmatrix}.$$

But we can see the contradiction by using Proposition 6.10. Thus, as the $A(\mathcal{W}_{p_+,p_-})$ -module, $\overline{E}_3 \simeq \overline{\mathcal{X}_{r,p_-}^+} \oplus \overline{\mathcal{X}_{r,p_-}^+}$. Let \tilde{E}_3 be the induced module from \overline{E}_3 . Then we have

$$\tilde{E}_3 \simeq \tilde{\mathcal{E}}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-} \oplus \tilde{\mathcal{E}}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}.$$

Therefore we obtain

$$E_3 \simeq \mathcal{X}_{r,p_-}^+ \oplus \mathcal{X}_{r,p_-}^+.$$

□

Proposition 6.14. *In the thick block $C_{r,s}^{thick}$, we have*

$$\text{Ext}^1(\mathcal{X}^\pm, \mathcal{X}^\mp) = \mathbb{C}^2, \quad \text{Ext}^1(L(h_{r,s}), \mathcal{X}^+) = \text{Ext}^1(\mathcal{X}^+, L(h_{r,s})) = \mathbb{C},$$

where $\mathcal{X}^+ = \mathcal{X}_{r,s}^+$ or $\mathcal{X}_{r^\vee,s^\vee}^+$ and $\mathcal{X}^- = \mathcal{X}_{r^\vee,s}^-$ or \mathcal{X}_{r,s^\vee}^- . The other extensions between the simple modules in $C_{r,s}^{thick}$ are trivial.

Proof. We will only prove $\text{Ext}^1(L(h_{r,s}), \mathcal{X}_{r,s}^+) = \mathbb{C}$. The other Ext^1 -groups can be proved in a similar way as Proposition 6.13.

Note that

$$[\mathcal{W}_{p_+,p_-} \cdot |\alpha_{r,s}\rangle] \in \text{Ext}^1(L(h_{r,s}), \mathcal{X}_{r,s}^+) \setminus \{0\},$$

and, as the Virasoro module

$$\mathcal{W}_{p_+,p_-} \cdot |\alpha_{r,s}\rangle = L(h_{r,s}, \Delta_{r,s}^+; 0) \oplus \bigoplus_{n \geq 1} (2n+1)L(\Delta_{r,s}^+; n).$$

Fix any extension $[E] \in \text{Ext}^1(L(h_{r,s}), \mathcal{X}_{r,s}^+)$. Let t be the highest weight vector of E and assume $S_{r^\vee, s^\vee} t \neq 0$. Then, as the Virasoro module

$$E = L(h_{r,s}, \Delta_{r,s;0}^+) \oplus \bigoplus_{n \geq 1} (2n+1)L(\Delta_{r,s;n}^+).$$

Since $\text{Ext}_{\mathcal{L}}^1(L(h_{r,s}), L(\Delta_{r,s;0}^+)) = \mathbb{C}$, as the Baer sum of extensions obtained from the indecomposable modules E and $\mathcal{W}_{p_+, p_-} \cdot |\alpha_{r,s}\rangle$, we have a extension $[E'] \in \text{Ext}^1(L(h_{r,s}), \mathcal{X}_{r,s}^+)$ such that $S_{r^\vee, s^\vee} t' = 0$, where t' is the highest weight vector of E' . Thus, by Theorem 5.6, we have the following decomposition as the Virasoro module

$$E' = L(h_{r,s}) \oplus \bigoplus_{n \geq 0} (2n+1)L(\Delta_{r,s;n}^+).$$

Assume $[E'] \neq 0$. Then, from the \mathcal{W}_{p_+, p_-} -module action on E' , we must have a non-trivial Virasoro intertwining operator of type

$$\begin{pmatrix} L(\Delta_{r,s;n}^+) \\ L(h_{4p_+-1,1}) & L(h_{r,s}) \end{pmatrix}$$

for some $n \geq 0$. But, by using Proposition 6.10, we can see the contradiction. In case $S_{r^\vee, s^\vee} t = 0$, we see that $[E] = 0$ as shown above. \square

6.3 The projective covers of the simple modules in the thin blocks

Fix any two thin blocks $C_{r,p_-}^{\text{thin}}, C_{p_+,s}^{\text{thin}}$ ($1 \leq r < p_+, 1 \leq s < p_-$). In this subsection, we will show the logarithmic modules $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee, p_-}, \mathcal{Q}(\mathcal{X}_{r,p_-}^-)_{r, p_-} \in C_{r,p_-}^{\text{thin}}$ and $\mathcal{Q}(\mathcal{X}_{p_+,s}^+)_{p_+, s^\vee}, \mathcal{Q}(\mathcal{X}_{p_+,s}^-)_{p_+, s} \in C_{p_+,s}^{\text{thin}}$ are projective.

The following lemma can be proved in the same way as Proposition 5.28, by using the structure of the logarithmic modules $(F(\tau), J_\tau)$.

Lemma 6.15. *For $\tau = (\alpha_{r^\vee, p_-;1}, \alpha_{r, p_-;2}, \alpha_{r^\vee, p_-;3}) \in \mathcal{T}_{p_+, p_-}$, we have*

$$\text{Ext}_{\mathcal{L}}^1(K(\tau), L(\Delta_{r^\vee, p_-;0}^-, \Delta_{r, p_-;1}^+)) \simeq \mathbb{C}. \quad (6.12)$$

Any indecomposable Virasoro module E in the Ext^1 -group (6.12) has a submodule isomorphic to $P(\tau)$, and satisfies

$$\text{Hom}_{U(\mathcal{L})}(E, K(\tau)) \simeq \mathbb{C}^2. \quad (6.13)$$

Furthermore, we can choose two surjections as a basis of (6.13).

From the structure of $\mathcal{E}^\pm(\mathcal{X}_{r^\vee, p_-}^\pm)_{r, p_-}$, we obtain the following lemma.

Lemma 6.16. *Fix any indecomposable module E in $\text{Ext}^1(\mathcal{X}_{r^\vee, p_-}^-, \mathcal{X}_{r, p_-}^+)$. For the surjection $\pi : E \rightarrow \mathcal{X}_{r^\vee, p_-}^-$, let $(v^-, v^+) \in E[\Delta_{r^\vee, p_-; 0}^-]^2$ be any generators of E satisfying*

$$W^\pm[0]\pi(v^\pm) = 0, \quad W^\pm[0]\pi(v^\mp) \in \mathbb{C}^\times \pi(v^\pm).$$

Then, for $\epsilon = \pm$, the Virasoro module $U(\mathcal{L}).v^\epsilon$ has a quotient isomorphic to $L(\Delta_{r^\vee, p_-; 0}^-, \Delta_{r, p_-; 1}^+)$.

Proposition 6.17.

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r, p_-}^+)_{r^\vee, p_-}, \mathcal{X}_{r, p_-}^+) = \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{p_+, s}^+)_{p_+, s^\vee}, \mathcal{X}_{p_+, s}^+) = 0.$$

Proof. We only prove the first equality. The second equality can be proved in the same way.

Assume that $\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r, p_-}^+)_{r^\vee, p_-}, \mathcal{X}_{r, p_-}^+) \neq 0$ and fix any non-trivial extension E in this Ext^1 -group.

First let us introduce some symbols. Let u be the highest weight vector of \mathcal{X}_{r, p_-}^+ . From Proposition 3.13, we can choose a basis $\{w_-, w_0, w_+\}$ of the Virasoro highest weight spaces of $3L(\Delta_{r, p_-; 1}^+) \subset \mathcal{X}_{r, p_-}^+$ satisfying

$$\begin{aligned} W^\pm[0]w_\pm &\equiv 0 \pmod{U(\mathcal{L}).u}, \\ W^\pm[0]w_\mp &\in \mathbb{C}^\times w_0 + U(\mathcal{L}).u, \\ W^\pm[0]w_0 &\in \mathbb{C}^\times w_\pm + U(\mathcal{L}).u. \end{aligned}$$

Let ι_1 and ι_2 be injections from \mathcal{X}_{r, p_-}^+ to E such that

$$\iota_1(\mathcal{X}_{r, p_-}^+) \oplus \iota_2(\mathcal{X}_{r, p_-}^+) = \text{Soc}(E).$$

For the surjection $\pi : E \rightarrow \mathcal{X}_{r, p_-}^+$, we fix L_0 -homogeneous vectors $\tilde{u}_0, \tilde{w}_0 \in E$ such that

$$\pi(\tilde{u}_0) = u, \quad \pi(\tilde{w}_0) = w_0.$$

Set $v^\pm = W^\pm[0]W^\mp[0]S_{r, p_-}\tilde{u}_0$. Note that $\{v^-, W^+[0]v^-\}$ and $\{W^-[0]v^+, v^+\}$ correspond to bases of the highest weight spaces of $\mathcal{X}_{r^\vee, p_-}^- \oplus \mathcal{X}_{r^\vee, p_-}^-$.

Let us consider the Virasoro module

$$M = U(\mathcal{L}).\tilde{w}_0 + U(\mathcal{L}).v^+ + U(\mathcal{L}).v^-.$$

From the Virasoro structure of $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}$, we see that the Virasoro module M has an indecomposable quotient in the Ext^1 -group (6.12), and the quotient module

$$M/(U(\mathcal{L}).v^+ + U(\mathcal{L}).v^-)$$

does not contain $L(\Delta_{r^\vee,p_-;0}^-)$ as the composition factors. Thus by Lemma 6.15, we have

$$\sigma(S_{r^\vee,2p_-})\tilde{w}_0 \in \mathbb{C}^\times v^+ + \mathbb{C}^\times v^- + U(\mathcal{L}).\iota_1(u) + U(\mathcal{L}).\iota_2(u). \quad (6.14)$$

By the structure of $P(\tau)$, we have

$$S_{r^\vee,2p_-}\sigma(S_{r^\vee,2p_-})\tilde{w}_0 \not\equiv 0 \bmod U(\mathcal{L}).\iota_1(u) + U(\mathcal{L}).\iota_2(u). \quad (6.15)$$

and

$$\begin{aligned} S_{r^\vee,2p_-}\sigma(S_{r^\vee,2p_-})\tilde{w}_0 \in & \mathbb{C}\iota_1(w_0) + \mathbb{C}\iota_2(w_0) + \sum_{i=1,2} \sum_{\epsilon=\pm} \mathbb{C}\iota_i(w_\epsilon) \\ & + U(\mathcal{L}).\iota_1(u) + U(\mathcal{L}).\iota_2(u). \end{aligned} \quad (6.16)$$

Assume that the coefficient of $\iota_i(w_\pm)$ in $S_{r^\vee,2p_-}\sigma(S_{r^\vee,2p_-})\tilde{w}_0$ is nonzero. Then, from the structure of $P(\tau)$, we see that the coefficient of $\iota_i(w_\pm)$ in $(L_0 - \Delta_{r,p_-;1}^+)\tilde{w}_0$ is also nonzero. Thus, noting Propositions 3.13 and 3.14, multiplying $(L_0 - \Delta_{r,p_-;1}^+)\tilde{w}_0$ by $(W^\mp[0])^2$, we have $\iota_i(w_\mp) \in U(\mathcal{L}).\iota_1(u) + U(\mathcal{L}).\iota_2(u)$. But this is a contradiction. Therefore, from (6.16), we have

$$S_{r^\vee,2p_-}\sigma(S_{r^\vee,2p_-})\tilde{w}_0 \equiv k_1\iota_1(w_0) + k_2\iota_2(w_0) \bmod \sum_{i=1}^2 U(\mathcal{L}).\iota_i(u), \quad (6.17)$$

where $(k_1, k_2) (\neq (0, 0))$ are some constants. Since $\text{Ext}^1(\mathcal{X}_{r,p_-}^+, \mathcal{X}_{r,p_-}^+) = 0$, we have

$$\mathcal{W}_{p_+,p_-} \cdot (k_1\iota_1(w_0) + k_2\iota_2(w_0)) \simeq \mathcal{X}_{r,p_-}^+. \quad (6.18)$$

We set

$$E' = \frac{E}{\mathcal{W}_{p_+,p_-} \cdot (k_1\iota_1(w_0) + k_2\iota_2(w_0))}.$$

Let ϕ be the surjection from E to E' . Then, from (6.15), (6.17) and (6.18), E' satisfies

$$[E'] \in \text{Ext}^1(\tilde{\mathcal{E}}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}, \mathcal{X}_{r,p_-}^+) \setminus \{0\}$$

and

$$S_{r^\vee, 2p_-} \sigma(S_{r^\vee, 2p_-}) \phi(\tilde{w}_0) \in \sum_{i=1,2} U(\mathcal{L}).\phi \circ \iota_i(u). \quad (6.19)$$

By (6.19), we see that $\phi(M)$ does not have $P(\tau)$ as any subquotient. Then, from Lemma 6.15 and (6.14), we have

$$S_{r^\vee, 2p_-} \phi(v^\pm) \in \sum_{i=1,2} U(\mathcal{L}).\phi \circ \iota_i(u). \quad (6.20)$$

By Lemma 6.16 and (6.20), E' has $\mathcal{X}_{r^\vee, p_-}^- \oplus \mathcal{X}_{r^\vee, p_-}^-$ as submodules. Thus E' has a indecomposable quotient in $\text{Ext}^1(\mathcal{X}_{r, p_-}^+, \mathcal{X}_{r, p_-}^+)$. But this contradicts Proposition 6.13. □

Proposition 6.18.

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}, \mathcal{X}_{r, p_-}^+) = \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{p_+, s^\vee}^-)_{p_+, s}, \mathcal{X}_{p_+, s}^+) = 0.$$

Proof. We only prove the first equality. The second equality can be proved in the same way.

By Proposition 6.17, we have

$$\text{Ext}^1(\mathcal{E}^+(\mathcal{X}_{r, p_-}^+)_{r^\vee, p_-}, \mathcal{X}_{r, p_-}^+) = 0. \quad (6.21)$$

From the structure of $\mathcal{Q}(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}$, we have the following exact sequence

$$0 \rightarrow \mathcal{E}^+(\mathcal{X}_{r, p_-}^+)_{r^\vee, p_-} \rightarrow \mathcal{Q}(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-} \rightarrow \mathcal{E}^-(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-} \rightarrow 0.$$

By this exact sequence and (6.21), we have the following exact sequence

$$0 \rightarrow \mathbb{C} \rightarrow \text{Ext}^1(\mathcal{E}^-(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}, \mathcal{X}_{r, p_-}^+) \rightarrow \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}, \mathcal{X}_{r, p_-}^+) \rightarrow 0.$$

By Proposition 6.13, we have $\text{Ext}^1(\mathcal{E}^-(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}, \mathcal{X}_{r, p_-}^+) \simeq \mathbb{C}$. Therefore we obtain

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}, \mathcal{X}_{r, p_-}^+) = 0. \quad \square$$

Lemma 6.19.

$$\text{Ext}^1(\mathcal{E}^\pm(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}, \mathcal{X}_{r^\vee, p_-}^-) = \text{Ext}^1(\mathcal{E}^\pm(\mathcal{X}_{p_+, s^\vee}^-)_{p_+, s}, \mathcal{X}_{p_+, s^\vee}^-) = 0.$$

Proof. We only prove $\text{Ext}^1(\mathcal{E}^+(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}, \mathcal{X}_{r^\vee, p_-}^-) = 0$. The other equalities can be proved in the same way.

Assume that

$$\text{Ext}^1(\mathcal{E}^+(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}, \mathcal{X}_{r^\vee, p_-}^-) \neq 0$$

and fix any non-trivial extension

$$0 \rightarrow \mathcal{X}_{r^\vee, p_-}^- \xrightarrow{\iota} E \rightarrow \mathcal{E}^+(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-} \rightarrow 0.$$

Let $\{v_+, v_-\}$ be a basis of the highest weight space of $\mathcal{X}_{r^\vee, p_-}^-$ such that

$$W^\pm[0]v_\pm = 0, \quad W^\pm[0]v_\mp \in \mathbb{C}^\times v_\pm.$$

For the surjection $\pi : E \rightarrow \mathcal{X}_{r^\vee, p_-}^-$, let \tilde{v}_\pm be any L_0 homogeneous vectors of E such that $\pi(\tilde{v}_\pm) = v_\pm$. Since $\text{Ext}^1(\mathcal{X}_{r^\vee, p_-}^-, \mathcal{X}_{r^\vee, p_-}^-) = 0$, we see that E has an indecomposable submodule in $\text{Ext}^1(\mathcal{X}_{r^\vee, p_-}^+, \mathcal{X}_{r^\vee, p_-}^-)$. Then, noting Lemma 6.12, we see that the Virasoro module $U(\mathcal{L}) \cdot \tilde{v}_+$ is a quotient isomorphic to $P(\tau)$, where $\tau = (\alpha_{r, p_-}, \alpha_{r^\vee, p_-; 1}, \alpha_{r, p_-; 2})$ (for the definition of the logarithmic modules $P(\tau)$, see Definition 5.17). Note that, from the structure of $\mathcal{E}^+(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}$, $L_n \tilde{v}_- = 0$ for $n \geq 1$. Thus, by Proposition 5.16, we have

$$(L_0 - \Delta_{r^\vee, p_-; 0}^-) \tilde{v}_+ = k_+ \iota(v_+) + k_- \iota(v_-), \quad (6.22)$$

$$(L_0 - \Delta_{r^\vee, p_-; 0}^-) \tilde{v}_- = 0, \quad (6.23)$$

where $(k_+, k_-) \neq (0, 0)$. Assume $k_+ \neq 0$. Then, multiplying both sides of (6.22) by $W^-[0]$, we have

$$(L_0 - \Delta_{r^\vee, p_-; 0}^-) W^-[0] \tilde{v}_+ \in \mathbb{C}^\times \iota(v_-).$$

But this contradicts (6.23). Next assume $k_- \neq 0$. Then, multiplying both sides of (6.22) by $W^+[0]$, we have

$$(L_0 - \Delta_{r^\vee, p_-; 0}^-) W^+[0] \tilde{v}_+ \in \mathbb{C}^\times \iota(v_+).$$

On the other hand, by the definition of \tilde{v}_+ , we have $(L_0 - \Delta_{r^\vee, p_-; 0}^-) W^+[0] \tilde{v}_+ = 0$. Thus we have a contradiction. \square

Proposition 6.20.

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}, \mathcal{X}_{r^\vee, p_-}^-) = \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{p_+, s^\vee}^-)_{p_+, s}, \mathcal{X}_{p_+, s^\vee}^-) = 0$$

Proof. We only prove the first equality. The second equality can be proved in the same way.

Assume that

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}, \mathcal{X}_{r^\vee, p_-}^-) \neq 0$$

and fix any non-trivial extension

$$0 \rightarrow \mathcal{X}_{r^\vee, p_-}^- \xrightarrow{\iota} E \xrightarrow{\phi} \mathcal{Q}(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-} \rightarrow 0.$$

Since $\text{Ext}^1(\mathcal{X}_{r^\vee, p_-}^-, \mathcal{X}_{r^\vee, p_-}^-) = 0$, E has a submodule in

$$\text{Ext}^1(\tilde{\mathcal{E}}(\mathcal{X}_{r^\vee, p_-}^-)_{r, p_-}^*, \mathcal{X}_{r^\vee, p_-}^-) \setminus \{0\}.$$

Then, from Propositions 6.13 and Lemma 6.19, we see that the following sequence of submodules holds

$$\iota(\mathcal{X}_{r^\vee, p_-}^-) \subset \tilde{\mathcal{E}}(\mathcal{X}_{r^\vee, p_-}^+)_{r^\vee, p_-} \subset E. \quad (6.24)$$

Let $\{v_+, v_-\}$ be a basis of the highest weight space of $\mathcal{X}_{r^\vee, p_-}^-$ such that

$$W^\pm[0]v_\pm = 0, \quad W^\pm[0]v_\mp \in \mathbb{C}^\times v_\pm.$$

Let $\{v_+^0, v_-^0\}$ be a basis of the highest weight space of the submodule $\mathcal{X}_{r^\vee, p_-}^- \subset E$ such that $\phi(v_\pm^0) \neq 0$ and

$$W^\pm[0]v_\pm^0 = 0, \quad W^\pm[0]v_\mp^0 \in \mathbb{C}^\times v_\pm^0.$$

For the surjection $\pi : E \rightarrow \mathcal{X}_{r^\vee, p_-}^-$, we fix any L_0 -homogeneous vectors $\tilde{v}_-, \tilde{v}_+ \in E$ such that $\pi(\tilde{v}_\pm) = v_\pm$. Note that the Virasoro module $U(\mathcal{L}).\tilde{v}_\pm$ has a quotient isomorphic to $P(\tau)$, where $\tau = (\alpha_{r, p_-}, \alpha_{r^\vee, p_-}; 1, \alpha_{r, p_-}; 2)$. Thus, by Proposition 6.13 and (6.24), we have

$$\begin{aligned} S_{r, p_-} \sigma(S_{r, p_-}) \tilde{v}_- &\in \mathbb{C}^\times \iota(v_+) + \mathbb{C} \iota(v_-) + \mathbb{C}^\times v_-^0, \\ S_{r, p_-} \sigma(S_{r, p_-}) \tilde{v}_+ &\in \mathbb{C} \iota(v_+) + \mathbb{C}^\times \iota(v_-) + \mathbb{C}^\times v_-^0. \end{aligned} \quad (6.25)$$

Then, by Proposition 5.16 and (6.25), we see that one of the followings holds

$$\begin{aligned} (L_0 - \Delta_{r^\vee, p_-; 0}^-) \tilde{v}_- &= k_+ \iota(v_+) + k_- \iota(v_-) + \mathbb{C}^\times v_-^0, \quad k_+ \neq 0, \\ (L_0 - \Delta_{r^\vee, p_-; 0}^-) \tilde{v}_+ &= l_- \iota(v_-) + l_+ \iota(v_+) + \mathbb{C}^\times v_+^0, \quad l_- \neq 0. \end{aligned}$$

Assume that the first equality is true. Multiplying the first equation by $W^-[0]$, we have

$$(L_0 - \Delta_{r^\vee, p_-; 0}^-) W^-[0] \tilde{v}_- = k_+ \iota(v_-).$$

By the definition of \tilde{v}_- , the left hand side becomes zero. But this is a contradiction. Similarly, assuming the second equality, we can show the contradiction. \square

Proposition 6.21.

$$\mathrm{Ext}^1(\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}, \mathcal{X}_{r^\vee,p_-}^-) = \mathrm{Ext}^1(\mathcal{Q}(\mathcal{X}_{p_+,s}^+)_{p_+,s^\vee}, \mathcal{X}_{p_+,s^\vee}^-) = 0.$$

Proof. We only prove the first equality. The second equality can be proved in the same way.

Note that $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}$ satisfies the following exact sequence

$$0 \rightarrow \mathcal{E}^-(\mathcal{X}_{r^\vee,p_-}^-)_{r,p_-} \rightarrow \mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-} \rightarrow \mathcal{E}^+(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-} \rightarrow 0.$$

By this exact sequence and Proposition 6.20, we have the following exact sequence

$$0 \rightarrow \mathbb{C} \rightarrow \mathrm{Ext}^1(\mathcal{E}^+(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}, \mathcal{X}_{r^\vee,p_-}^-) \rightarrow \mathrm{Ext}^1(\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}, \mathcal{X}_{r^\vee,p_-}^-) \rightarrow 0.$$

By Proposition 6.13 we have $\mathrm{Ext}^1(\mathcal{E}^+(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}, \mathcal{X}_{r^\vee,p_-}^-) \simeq \mathbb{C}$. Therefore we obtain $\mathrm{Ext}^1(\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}, \mathcal{X}_{r^\vee,p_-}^-) = 0$. □

By Propositions 6.17, 6.18, 6.20 and 6.21, we obtain the following theorem.

Theorem 6.22.

1. $\mathcal{Q}(\mathcal{X}_{r,p_-}^+)_{r^\vee,p_-}$ and $\mathcal{Q}(\mathcal{X}_{r^\vee,p_-}^-)_{r,p_-}$ are the projective covers of \mathcal{X}_{r,p_-}^+ and $\mathcal{X}_{r^\vee,p_-}^-$, respectively.
2. $\mathcal{Q}(\mathcal{X}_{p_+,s}^+)_{p_+,s^\vee}$ and $\mathcal{Q}(\mathcal{X}_{p_+,s^\vee}^-)_{p_+,s}$ are the projective covers of $\mathcal{X}_{p_+,s}^+$ and $\mathcal{X}_{p_+,s^\vee}^-$, respectively.

6.4 The projective covers of the simple modules $\mathcal{X}_{\bullet,\bullet}^\pm$ in the thick blocks

In this subsection, we fix any thick block $C_{r,s}^{thick}$ and compute Ext^1 groups between certain indecomposable \mathcal{W}_{p_+,p_-} -modules and the simple modules in this block. Based on these Ext^1 -groups, we prove that the logarithmic modules $\mathcal{P}_{\bullet,\bullet}^\pm$ are projective \mathcal{W}_{p_+,p_-} -modules.

Definition 6.23. Let (a,b) be (r,s) or (r^\vee, s^\vee) . We identify the indecomposable modules in the Ext^1 -group

$$\mathrm{Ext}^1(L(h_{r,s}), \mathcal{X}_{a,b}^+) \simeq \mathbb{C}$$

(see Proposition 6.14) and denote them as $\mathcal{K}_{a,b}$.

The following proposition can be proved in the same way as Proposition 6.17, but we give a different proof.

Proposition 6.24. *Let (a, b) be (r, s) or (r^\vee, s^\vee) . Then we have*

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^+)_{a^\vee, b}, \mathcal{X}_{a,b}^+) = \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^+)_{a, b^\vee}, \mathcal{X}_{a,b}^+) = 0.$$

Proof. We will only prove $\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^+)_{a^\vee, b}, \mathcal{X}_{a,b}^+) = 0$ in the case $(a, b) = (r, s)$. The other cases can be proved in the same way.

Assume $\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee, s}, \mathcal{X}_{r,s}^+) \neq 0$. Then, by Theorem 6.14, we have

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee, s}/\mathcal{K}_{r,s}, \mathcal{X}_{r,s}^+) \neq 0.$$

Fix any non trivial extension E in this Ext^1 -group. By Theorem 5.16, we see that L_0 acts semisimply on the highest weight space of E . Thus, by Propositions 3.13 and 3.14, we see that L_0 acts semisimply on E . Let $u_0 \in E[\Delta_{r,s;0}^+]$ be a generator of E , and let u_1 be the highest weight vector of the submodule $\mathcal{X}_{r,s}^+ \subset E$. Fix any homogeneous vector $u_1^* \in E^*$ such that $\langle u_1^*, u_1 \rangle \neq 0$. Since $[E] \neq 0$, E has one of $\mathcal{E}^+(\mathcal{X}_{r^\vee, s}^-)_{r,s}$ or $\mathcal{E}^-(\mathcal{X}_{r^\vee, s}^-)_{r,s}$, as a submodule. Thus, by the structure of $\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee, s}/\mathcal{K}_{r,s}$ and $\mathcal{E}^\pm(\mathcal{X}_{r^\vee, s}^-)_{r,s}$, we see that

$$\langle u_1^*, \sigma(S_{r, s^\vee + p_-})Y_E(W^\epsilon; z)S_{r, s^\vee + p_-}u_0 \rangle \neq 0,$$

where W^ϵ is one of W^+ , W^0 or W^- . In particular, we have

$$\langle u_1^*, Y_E(W^\epsilon; z)u_0 \rangle \neq 0. \quad (6.26)$$

Note that $S_{r^\vee + p_+, s}u_0 = 0$. Thus, by Proposition 6.9, we have

$$\begin{aligned} 0 &= \langle u_1^*, Y_E(W^\epsilon; z)S_{r^\vee + p_+, s}u_0 \rangle \\ &= \prod_{i=1}^{r^\vee + p_+} \prod_{j=1}^s (h_{4p_+ - 1, 1} - h_{2r^\vee + 2p_+ - 2i + 1, 2s - 2j + 1}) \langle u_1^*, Y_E(W^\epsilon; z)u_0 \rangle. \end{aligned}$$

The coefficient in the above equation is nonzero, so we have $\langle u_1^*, Y_E(W^\epsilon; z)u_0 \rangle = 0$. But this contradicts (6.26). \square

The following three propositions can be proved in the same way as Propositions 6.18, 6.20 and 6.21, so we omit the proofs.

Proposition 6.25. *Let (a, b) be (r^\vee, s) or (r, s^\vee) . Then we have*

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^-)_{a^\vee, b}, \mathcal{X}_{a,b}^+) = \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^-)_{a, b^\vee}, \mathcal{X}_{a,b}^+) = 0.$$

Proposition 6.26. *Let (a, b) be (r^\vee, s) or (r, s^\vee) . Then we have*

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^-)_{a^\vee, b}, \mathcal{X}_{a,b}^-) = \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^-)_{a, b^\vee}, \mathcal{X}_{a,b}^-) = 0.$$

Proposition 6.27. *Let (a, b) be (r, s) or (r^\vee, s^\vee) . Then we have*

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^+)_{a^\vee, b}, \mathcal{X}_{a^\vee, b}^-) = \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^+)_{a, b^\vee}, \mathcal{X}_{a, b^\vee}^-) = 0.$$

Proposition 6.28. *Let (a, b, ϵ) be any element in*

$$\{(r, s, +), (r^\vee, s^\vee, +), (r^\vee, s, -), (r, s^\vee, -)\}.$$

Then we have

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^\epsilon)_{a^\vee, b}, \mathcal{X}_{a^\vee, b^\vee}^\epsilon) = \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^\epsilon)_{a, b^\vee}, \mathcal{X}_{a^\vee, b^\vee}^\epsilon) = 0.$$

Proof. We will prove only

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee, s}, \mathcal{X}_{r^\vee, s^\vee}^+) = 0, \quad \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r^\vee, s}^-)_{r, s}, \mathcal{X}_{r, s^\vee}^-) = 0.$$

The other equality can be proved in the same way.

First we prove $\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee, s}, \mathcal{X}_{r^\vee, s^\vee}^+) = 0$. Recall that $\tilde{K}(\Delta_{r,s;0}^+)$ is the indecomposable Virasoro module satisfying the following exact sequence

$$0 \rightarrow L(\Delta_{r,s^\vee;0}^-) \rightarrow \tilde{K}(\Delta_{r,s;0}^+) \rightarrow K(\tau) \rightarrow 0,$$

where $\tau = (\alpha_{r^\vee, s^\vee}, \alpha_{r, s^\vee;1}, \alpha_{r^\vee, s^\vee;2})$. By the structure of Virasoro Verma modules ([8],[11],[22]), we see that

$$\text{Ext}_{\mathcal{L}}^1(\tilde{K}(\Delta_{r,s;0}^+)/L(h_{r,s}), L(\Delta_{r^\vee, s^\vee;0}^+)) = 0.$$

Thus, by the structure of the Fock module F_{r^\vee, s^\vee} , we have

$$\text{Ext}_{\mathcal{L}}^1(\tilde{K}(\Delta_{r,s;0}^+), L(\Delta_{r^\vee, s^\vee;0}^+)) \simeq \mathbb{C}. \quad (6.27)$$

We identify the indecomposable modules in this Ext^1 -group and denote them as $V(\Delta_{r,s;0}^+)$. By (6.27), we have

$$\text{Ext}_{\mathcal{L}}^1(V(\Delta_{r,s;0}^+), L(\Delta_{r^\vee, s^\vee;0}^+)) = 0. \quad (6.28)$$

Note that $V(\Delta_{r,s;0}^+)$ satisfies the exact sequence

$$0 \rightarrow L(\Delta_{r^\vee, s^\vee;0}^+, \Delta_{r, s^\vee;0}^-)^* \rightarrow V(\Delta_{r,s;0}^+) \rightarrow K(\tau) \rightarrow 0.$$

By this exact sequence and (6.28), we obtain

$$\mathrm{Ext}_{\mathcal{L}}^1(K(\tau), L(\Delta_{r^\vee, s^\vee; 0}^+)) = 0. \quad (6.29)$$

Assume $\mathrm{Ext}^1(\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee, s}, \mathcal{X}_{r^\vee, s^\vee}^+) \neq 0$. Then, by Proposition 6.14, we have

$$\mathrm{Ext}^1(\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee, s} / \mathcal{X}_{r,s}^+, \mathcal{X}_{r^\vee, s^\vee}^+) \neq 0.$$

Fix a non-trivial extension $[E] \in \mathrm{Ext}^1(\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee, s} / \mathcal{X}_{r,s}^+, \mathcal{X}_{r^\vee, s^\vee}^+)$. Note that L_0 acts semisimply on E . By (6.29), we can choose a L_0 -homogeneous generator u_0 of E satisfying

$$U(\mathcal{L}).u_0 \simeq K(\tau). \quad (6.30)$$

Let u_1 be the highest weight vector of the submodule $\mathcal{X}_{r^\vee, s^\vee}^+ \subset E$. By (6.30), we see that there exists a homogeneous vector $u_1^* \in E^*$ such that $\langle u_1^*, u_1 \rangle \neq 0$ and

$$L_n u_1^* = 0, \quad \text{for } n \geq 1. \quad (6.31)$$

Since E is non-trivial, by the structure of $\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee, s} / \mathcal{X}_{r,s}^+$, $\mathcal{E}^\pm(\mathcal{X}_{r,s}^+)_{r^\vee, s}$ and $\mathcal{E}^\pm(\mathcal{X}_{r^\vee, s^\vee}^+)_{r^\vee, s}$, we have

$$\langle u_1^*, Y_E(W^\bullet; z)u_0 \rangle \neq 0, \quad (6.32)$$

where W^\bullet is one of W^\pm or W^0 . Note that by (6.30),

$$S_{r^\vee + p_+, s} u_0 \in L(h_{r,s}).$$

Then by Proposition 6.9 and (6.31), we have

$$\begin{aligned} 0 &= \langle u_1^*, Y_E(W^\bullet; z)S_{r^\vee + p_+, s} u_0 \rangle \\ &= \prod_{i=1}^{r^\vee + p_+} \prod_{j=1}^s (h_{4p_+ - 1, 1} - h_{r^\vee + r + 2p_+ - 2i + 1, s + s^\vee - 2j + 1}) \langle u_1^*, Y_E(W^\bullet; z)u_0 \rangle. \end{aligned}$$

The coefficient in the above equation is nonzero, so we have $\langle u_1^*, Y_E(W^\bullet; z)u_0 \rangle = 0$. But this contradicts (6.32).

Next we prove $\mathrm{Ext}^1(\mathcal{Q}(\mathcal{X}_{r^\vee, s}^-)_{r, s}, \mathcal{X}_{r, s^\vee}^-) = 0$. Note that, by the structure of Virasoro Verma modules and by the structure of the Fock module $F_{r, s^\vee; 1}$,

$$\mathrm{Ext}_{\mathcal{L}}^1(\tilde{K}(\Delta_{r^\vee, s; 0}^-) / L(\Delta_{r^\vee, s^\vee; 0}^+), L(\Delta_{r, s^\vee; 0}^-)) \simeq \mathbb{C} \quad (6.33)$$

(see Definitions 5.19 for the definitions of Virasoro module $\tilde{K}(\Delta_{r^\vee, s; 0}^-)$). We identify the indecomposable modules in this Ext^1 -group and denote them as $V(\Delta_{r^\vee, s; 0}^-)$. By (6.33), we have

$$\text{Ext}_{\mathcal{L}}^1(V(\Delta_{r^\vee, s; 0}^-), L(\Delta_{r, s^\vee; 0}^-)) = 0. \quad (6.34)$$

Note that $V(\Delta_{r^\vee, s; 0}^-)$ satisfies the exact sequence

$$0 \rightarrow L(\Delta_{r, s^\vee; 0}^-, \Delta_{r^\vee, s^\vee; 1}^+)^* \rightarrow V(\Delta_{r^\vee, s; 0}^-) \rightarrow K(\tau') \rightarrow 0,$$

where $\tau' = (\alpha_{r, s^\vee; 1}, \alpha_{r^\vee, s^\vee; 2}, \alpha_{r, s^\vee; 3})$. By this exact sequence and (6.34), we obtain

$$\text{Ext}_{\mathcal{L}}^1(K(\tau'), L(\Delta_{r, s^\vee; 0}^-)) = 0. \quad (6.35)$$

Let us assume that

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r^\vee, s}^-)_{r, s}, \mathcal{X}_{r, s^\vee}^-) \neq 0.$$

Then, since $\text{Ext}^1(\mathcal{X}_{r^\vee, s}^-, \mathcal{X}_{r, s^\vee}^-) = 0$, we have

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r^\vee, s}^-)_{r, s} / \mathcal{X}_{r^\vee, s}^-, \mathcal{X}_{r, s^\vee}^-) \neq 0.$$

Note that L_0 acts semisimply on any extensions of this Ext^1 -group. Fix any non-trivial extension $[E] \in \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r^\vee, s}^-)_{r, s} / \mathcal{X}_{r^\vee, s}^-, \mathcal{X}_{r, s^\vee}^-)$. Then, noting Lemma 6.12 and Proposition 6.14, by the Virasoro module structure of E we have

$$\text{Ext}_{\mathcal{L}}^1(K(\tau'), L(\Delta_{r, s^\vee; 0}^-)) \neq 0.$$

But this contradicts (6.35). □

Proposition 6.29. *Let (a, b) be (r, s) or (r^\vee, s^\vee) . Then we have*

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a, b}^+)_{a^\vee, b}, L(h_{a, b})) = \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a, b}^+)_{a, b^\vee}, L(h_{a, b})) = 0.$$

Proof. By the exact sequence

$$0 \rightarrow \mathcal{K}_{r, s} \rightarrow \mathcal{Q}(\mathcal{X}_{r, s}^+)_{r^\vee, s} \rightarrow \mathcal{Q}(\mathcal{X}_{r, s}^+)_{r^\vee, s} / \mathcal{K}_{r, s} \rightarrow 0,$$

we have the exact sequence

$$0 \rightarrow \mathbb{C} \rightarrow \mathbb{C} \rightarrow \text{Ext}(\mathcal{Q}(\mathcal{X}_{r, s}^+)_{r^\vee, s}, L(h_{r, s})) \rightarrow \text{Ext}^1(\mathcal{K}_{r, s}, L(h_{r, s})) \rightarrow 0.$$

Thus we have $\text{Ext}(\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee,s}, L(h_{r,s})) \simeq \text{Ext}^1(\mathcal{K}_{r,s}, L(h_{r,s}))$. Assume

$$\text{Ext}^1(\mathcal{K}_{r,s}, L(h_{r,s})) \neq 0,$$

and fix a non-trivial extension $[E] \in \text{Ext}^1(\mathcal{K}_{r,s}, L(h_{r,s}))$. Since

$$\text{Ext}^1(L(h_{r,s}), L(h_{r,s})) = 0,$$

E has a submodule isomorphic to $\mathcal{K}_{r,s}^*$. Thus, by Proposition 5.3, we see that E has L_0 -nilpotent rank two. Let $\{u_0, u_1\}$ be a basis of the highest weight space of E such that

$$(L_0 - h_{r,s})u_0 \in \mathbb{C}^\times u_1. \quad (6.36)$$

Then, by Proposition 5.3 and (6.36), we have

$$\sigma(S_{r,s})S_{r,s}u_0 \in \mathbb{C}^\times u_1.$$

In particular we have $S_{r,s}u_0 \neq 0$. Thus E has $\mathcal{X}_{r^\vee,s^\vee}^+$ as a composition factor. But this is a contradiction. Thus we obtain $\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee,s}, L(h_{r,s})) = 0$. The other equality can be proved in the same way, so we omit the proofs. \square

Proposition 6.30. *Let (a, b) be (r^\vee, s) or (r, s^\vee) . Then we have*

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^-)_{a^\vee,b}, L(h_{a^\vee,b})) = \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^-)_{a,b^\vee}, L(h_{a,b^\vee})) = 0.$$

Proof. By Proposition 6.14, we have

$$\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r^\vee,s}^-)_{r,s}, L(h_{r,s})) \simeq \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r^\vee,s}^-)_{r,s}/\mathcal{X}_{r^\vee,s}^-, L(h_{r,s})).$$

Assume $\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r^\vee,s}^-)_{r,s}/\mathcal{X}_{r^\vee,s}^-, L(h_{r,s})) \neq 0$. Then any non-trivial extension of $\text{Ext}^1(\mathcal{Q}(\mathcal{X}_{r^\vee,s}^-)_{r,s}/\mathcal{X}_{r^\vee,s}^-, L(h_{r,s}))$ has an indecomposable submodule in $\text{Ext}^1(\mathcal{X}_{r^\vee,s}^-, \mathcal{K}_{r,s}^*)$. Thus we have

$$\text{Ext}^1(\mathcal{K}_{r,s}, \mathcal{X}_{r^\vee,s}^-) \neq 0.$$

Since $\text{Ext}^1(L(h_{r,s}), \mathcal{X}_{r^\vee,s}^-) = 0$, any non-trivial extension of $\text{Ext}^1(\mathcal{K}_{r,s}, \mathcal{X}_{r^\vee,s}^-)$ has a submodule in $\text{Ext}^1(\mathcal{X}_{r,s}^+, \mathcal{X}_{r^\vee,s}^-) \setminus \{0\}$. In particular, we have

$$\text{Ext}_{\mathcal{L}}^1(L(h_{r,s}, \Delta_{r,s;0}^+), L(\Delta_{r^\vee,s;0}^-)) \neq 0.$$

On the other hand, by the structure of Virasoro Verma modules ([8],[11],[22]), we see that

$$\text{Ext}_{\mathcal{L}}^1(L(h_{r,s}, \Delta_{r,s;0}^+), L(\Delta_{r^\vee,s;0}^-)) = 0.$$

Thus we have a contradiction. The other equality can be proved in the same way. \square

The following is a summary of Proposition 6.24, 6.25, 6.26, 6.27, 6.28, 6.29 and 6.30.

Proposition 6.31. *Let (ϵ, a, b, c, d) be any element in*

$$\{(\epsilon, a, b, c, d)\} = \{(+, r, s, r^\vee, s), (+, r, s, r, s^\vee), (+, r^\vee, s^\vee, r^\vee, s), (+, r^\vee, s^\vee, r, s^\vee), \\ (-, r^\vee, s, r, s), (-, r^\vee, s, r^\vee, s^\vee), (-, r, s^\vee, r, s), (-, r, s^\vee, r^\vee, s^\vee)\}.$$

Then we have

$$\begin{aligned} \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^\epsilon)_{c,d}, L(h_{r,s})) &= 0, & \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^\epsilon)_{c,d}, \mathcal{X}_{a,b}^\epsilon) &= 0, \\ \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^\epsilon)_{c,d}, \mathcal{X}_{a^\vee, b^\vee}^\epsilon) &= 0, & \text{Ext}^1(\mathcal{Q}(\mathcal{X}_{a,b}^\epsilon)_{c,d}, \mathcal{X}_{c,d}^{-\epsilon}) &= 0. \end{aligned}$$

In the following, let us introduce the socle structure of the logarithmic modules $\mathcal{P}_{\bullet, \bullet}^\pm$. By Propositions 6.14, 6.24 and 6.26, we have the following lemma.

Lemma 6.32. *Fix any (ϵ, a, b, c, d) in*

$$\{(\epsilon, a, b, c, d)\} = \{(+, r, s, r^\vee, s), (+, r, s, r, s^\vee), (+, r^\vee, s^\vee, r^\vee, s), (+, r^\vee, s^\vee, r, s^\vee), \\ (-, r^\vee, s, r, s), (-, r^\vee, s, r^\vee, s^\vee), (-, r, s^\vee, r, s), (-, r, s^\vee, r^\vee, s^\vee)\}.$$

Then, any indecomposable module whose composition factors are the same as those of $\mathcal{Q}(\mathcal{X}_{a,b}^\epsilon)_{c,d}$ is isomorphic to $\mathcal{Q}(\mathcal{X}_{a,b}^\epsilon)_{c,d}$.

Proposition 6.33. *Fix any (a, b, ϵ) in*

$$\{(r, s, +), (r^\vee, s^\vee, +), (r^\vee, s, -), (r, s^\vee, -)\}.$$

Then the logarithmic module $\mathcal{P}_{a,b}^\epsilon$ has the following sequences of submodules

$$\begin{aligned} U_1(\mathcal{P}_{a,b}^\epsilon) &\subset U_2(\mathcal{P}_{a,b}^\epsilon) \subset U_3(\mathcal{P}_{a,b}^\epsilon) \subset U_4(\mathcal{P}_{a,b}^\epsilon) = \mathcal{P}_{a,b}^\epsilon \\ V_1(\mathcal{P}_{a,b}^\epsilon) &\subset V_2(\mathcal{P}_{a,b}^\epsilon) \subset V_3(\mathcal{P}_{a,b}^\epsilon) \subset V_4(\mathcal{P}_{a,b}^\epsilon) = \mathcal{P}_{a,b}^\epsilon, \end{aligned}$$

such that

$$\begin{aligned} U_1 &= \mathcal{Q}(\mathcal{X}_{a,b}^\epsilon)_{a^\vee, b}, \quad U_2/U_1 = U_3/U_2 = \mathcal{Q}(\mathcal{X}_{a, b^\vee}^{-\epsilon})_{a^\vee, b^\vee}, \quad U_4/U_3 = \mathcal{Q}(\mathcal{X}_{a,b}^\epsilon)_{a^\vee, b} \\ V_1 &= \mathcal{Q}(\mathcal{X}_{a,b}^\epsilon)_{a, b^\vee}, \quad V_2/V_1 = V_3/V_2 = \mathcal{Q}(\mathcal{X}_{a^\vee, b}^{-\epsilon})_{a^\vee, b^\vee}, \quad V_4/V_3 = \mathcal{Q}(\mathcal{X}_{a,b}^\epsilon)_{a, b^\vee}. \end{aligned}$$

Proof. Let us show that $\mathcal{P}_{r,s}^+$ has a sequence

$$U_1(\mathcal{P}_{r,s}^+) \subset U_2(\mathcal{P}_{r,s}^+) \subset U_3(\mathcal{P}_{r,s}^+) \subset U_4(\mathcal{P}_{r,s}^+) = \mathcal{P}_{r,s}^+$$

such that

$$U_1 = \mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee,s}, \quad U_2/U_1 = U_3/U_2 = \mathcal{Q}(\mathcal{X}_{r,s^\vee}^-)_{r^\vee,s^\vee}, \quad U_4/U_3 = \mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee,s}.$$

The other cases can be proved in the same way.

By Theorem 4.6, we see that $\mathcal{P}_{r,s}^+$ has a logarithmic submodule M satisfying

$$\mathcal{V}_{r,s}^+ \subset M, \quad M/\mathcal{V}_{r,s}^+ \simeq \mathcal{V}_{r^\vee,s}^-.$$

From the structure of $\mathcal{V}_{r,s}^+$ and $\mathcal{V}_{r^\vee,s}^-$, we see that M has two indecomposable subquotients whose composition factors are the same as those of $\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee,s}$ and $\mathcal{Q}(\mathcal{X}_{r,s^\vee}^-)_{r^\vee,s^\vee}$, respectively. Then, from Lemma 6.32, we see that M satisfies

$$\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee,s} \subset M, \quad M/\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee,s} \simeq \mathcal{Q}(\mathcal{X}_{r,s^\vee}^-)_{r^\vee,s^\vee}. \quad (6.37)$$

Set $N = \mathcal{P}_{r,s}^+/M$. From Theorem 4.6, we see that N is logarithmic and satisfies

$$\mathcal{V}_{r,s^\vee}^- \subset N, \quad N/\mathcal{V}_{r,s^\vee}^- \simeq \mathcal{V}_{r^\vee,s}^+.$$

From the structure of \mathcal{V}_{r,s^\vee}^- and $\mathcal{V}_{r^\vee,s}^+$, we see that N has two indecomposable subquotients whose composition factors are the same as those of $\mathcal{Q}(\mathcal{X}_{r,s^\vee}^-)_{r^\vee,s^\vee}$ and $\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee,s}$, respectively. Then, from Lemma 6.32, we see that N satisfies

$$\mathcal{Q}(\mathcal{X}_{r,s^\vee}^-)_{r^\vee,s^\vee} \subset N, \quad N/\mathcal{Q}(\mathcal{X}_{r,s^\vee}^-)_{r^\vee,s^\vee} \simeq \mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee,s}. \quad (6.38)$$

By (6.37) and (6.38), we obtain the claim. \square

By Propositions 6.31 and 6.33, we obtain the following theorems.

Theorem 6.34.

$$\begin{aligned} \text{Ext}^1(\mathcal{P}_{r,s}^+, L(h_{r,s})) &= \text{Ext}^1(\mathcal{P}_{r^\vee,s^\vee}^+, L(h_{r,s})) = 0, \\ \text{Ext}^1(\mathcal{P}_{r^\vee,s}^-, L(h_{r,s})) &= \text{Ext}^1(\mathcal{P}_{r,s^\vee}^-, L(h_{r,s})) = 0. \end{aligned}$$

Theorem 6.35. *Let (a, b, ϵ) be any element in*

$$\{(r, s, +), (r^\vee, s^\vee, +), (r^\vee, s, -), (r, s^\vee, -)\}.$$

Then we have

$$\begin{aligned} \text{Ext}^1(\mathcal{P}_{a,b}^\epsilon, \mathcal{X}_{a^\vee,b^\vee}^\epsilon) &= \text{Ext}^1(\mathcal{P}_{a^\vee,b^\vee}^\epsilon, \mathcal{X}_{a,b}^\epsilon) = 0, \\ \text{Ext}^1(\mathcal{P}_{a,b}^\epsilon, \mathcal{X}_{a^\vee,b}^{-\epsilon}) &= \text{Ext}^1(\mathcal{P}_{a,b}^\epsilon, \mathcal{X}_{a,b^\vee}^{-\epsilon}) = 0, \\ \text{Ext}^1(\mathcal{P}_{a^\vee,b^\vee}^\epsilon, \mathcal{X}_{a^\vee,b}^{-\epsilon}) &= \text{Ext}^1(\mathcal{P}_{a^\vee,b^\vee}^\epsilon, \mathcal{X}_{a,b^\vee}^{-\epsilon}) = 0. \end{aligned}$$

By Proposition 6.33, we obtain the following proposition.

Proposition 6.36. *Let $(a, b) = (r, s)$ or (r^\vee, s^\vee) . Each of the logarithmic modules $\mathcal{P}_{a,b}^+, \mathcal{P}_{a,b^\vee}^- \in C_{r,s}^{thick}$ is generated from the top composition factor and has the following length five socle series:*

1. *For the socle series of $\mathcal{P}_{a,b}^+$, we have*

$$\begin{aligned} S_1 &= \mathcal{X}_{a,b}^+, \\ S_2/S_1 &= \mathcal{X}_{a,b^\vee}^- \oplus \mathcal{X}_{a,b^\vee}^- \oplus L(h_{r,s}) \oplus \mathcal{X}_{a^\vee,b}^- \oplus \mathcal{X}_{a^\vee,b}^-, \\ S_3/S_2 &= \mathcal{X}_{a,b}^+ \oplus \mathcal{X}_{a^\vee,b^\vee}^+ \oplus \mathcal{X}_{a^\vee,b^\vee}^+ \oplus \mathcal{X}_{a^\vee,b^\vee}^+ \oplus \mathcal{X}_{a^\vee,b^\vee}^+ \oplus \mathcal{X}_{a,b}^+, \\ S_4/S_3 &= \mathcal{X}_{a^\vee,b}^- \oplus \mathcal{X}_{a^\vee,b}^- \oplus L(h_{r,s}) \oplus \mathcal{X}_{a,b^\vee}^- \oplus \mathcal{X}_{a,b^\vee}^-, \\ \mathcal{P}_{a,b}^+/S_4 &= \mathcal{X}_{a,b}^+, \end{aligned}$$

where $S_i = \text{Soc}_i$.

2. *For the socle series of \mathcal{P}_{a,b^\vee}^- , we have*

$$\begin{aligned} S_1 &= \mathcal{X}_{a,b^\vee}^-, \\ S_2/S_1 &= \mathcal{X}_{a,b}^+ \oplus \mathcal{X}_{a,b}^+ \oplus \mathcal{X}_{a^\vee,b^\vee}^+ \oplus \mathcal{X}_{a^\vee,b^\vee}^+, \\ S_3/S_2 &= \mathcal{X}_{a,b^\vee}^- \oplus \mathcal{X}_{a^\vee,b}^- \oplus \mathcal{X}_{a^\vee,b}^- \oplus L(h_{r,s}) \oplus L(h_{r,s}) \oplus \mathcal{X}_{a^\vee,b}^- \oplus \mathcal{X}_{a^\vee,b}^- \oplus \mathcal{X}_{a,b^\vee}^-, \\ S_4/S_3 &= \mathcal{X}_{a^\vee,b^\vee}^+ \oplus \mathcal{X}_{a^\vee,b^\vee}^+ \oplus \mathcal{X}_{a,b}^+ \oplus \mathcal{X}_{a,b}^+, \\ \mathcal{P}_{a,b^\vee}^-/S_4 &= \mathcal{X}_{a,b^\vee}^-. \end{aligned}$$

Remark 6.37. *Figure 6.3 is the embedding structure of the logarithmic \mathcal{W}_{p_+,p_-} -modules defined in Theorem 4.6. The structure of these logarithmic modules are studied by [18],[19] in the case $(p_+, p_-) = (2, 3)$. In one block $C_{1,1}^{thick}$, explicit realizations are given by [6].*

We define the following notation.

Definition 6.38. *Given a \mathcal{W}_{p_+,p_-} -module M with*

$$\text{Soc}_1(M) \subsetneq \cdots \subsetneq \text{Soc}_n(M) = M,$$

fix i satisfying $0 \leq i < n$ and fix a simple module $X \subset \text{Soc}_{n-i}(M)/\text{Soc}_{n-i-1}(M)$. Let k be the multiplicity of X in $\text{Soc}_{n-i}(M)/\text{Soc}_{n-i-1}(M)$. Then we denote by $(M : X : i)$ the submodule of $\text{Soc}_{n-i}(M)$ generated from the composition factors $kX \subset \text{Soc}_{n-i}(M)/\text{Soc}_{n-i-1}(M)$, where we use the notation given in Definition 6.11.




Figure 6.3: The embedding structure of logarithmic \mathcal{W}_{p_+, p_-} -modules $\mathcal{P}_{\bullet, \bullet}^\pm$. The triangle \triangle corresponds to the simple module $L(h_{r,s})$, \heartsuit to $\mathcal{X}_{r,s}^+$, \diamond to $\mathcal{X}_{r^\vee, s^\vee}^+$, \spadesuit to $\mathcal{X}_{r, s^\vee}^-$ and \clubsuit to $\mathcal{X}_{r^\vee, s}^-$.

Definition 6.39. Let (a, b) be (r, s) or (r^\vee, s^\vee) . We define the following indecomposable modules:

1. We define

$$\mathcal{P}_{a,b}^{+d} := (\mathcal{P}_{a,b}^+ : \mathcal{X}_{a,b}^+ : 2).$$

2. We define

$$\mathcal{P}_{a,b}^{+u} := \mathcal{P}_{a,b}^+ / (\mathcal{P}_{a,b}^+ : \mathcal{X}_{a^\vee, b^\vee}^+ : 2).$$




Figure 6.4: The embedding structure of the logarithmic \mathcal{W}_{p_+, p_-} -modules $\mathcal{P}_{\bullet, \bullet}^{+u}$ and $\mathcal{P}_{\bullet, \bullet}^{+d}$. The triangle \triangle corresponds to the simple module $L(h_{r,s})$, \heartsuit to $\mathcal{X}_{r,s}^+$, \diamondsuit to $\mathcal{X}_{r^\vee, s^\vee}^+$, \spadesuit to $\mathcal{X}_{r,s}^-$ and \clubsuit to $\mathcal{X}_{r^\vee, s^\vee}^-$.

Remark 6.40. Figure 6.4 represents the embedding structure of the logarithmic \mathcal{W}_{p_+, p_-} -modules given in Definition 6.39.

Proposition 6.41.

$$\text{Ext}^1(\mathcal{P}_{r,s}^{+d}, \mathcal{X}_{r,s}^+) = \text{Ext}^1(\mathcal{P}_{r^\vee, s^\vee}^{+d}, \mathcal{X}_{r^\vee, s^\vee}^+) = 0.$$

Proof. By Proposition 6.33, we see that $\mathcal{P}_{r,s}^{+d}$ has $\mathcal{Q}(\mathcal{X}_{r,s}^+)_{r,s^\vee}$ as a submodule. Then by the exact sequence

$$0 \rightarrow \mathcal{Q}(\mathcal{X}_{r,s}^+)_{r,s^\vee} \rightarrow \mathcal{P}_{r,s}^{+d} \rightarrow \mathcal{Q}(\mathcal{X}_{r,s}^+)_{r^\vee,s}/\mathcal{K}_{r,s} \rightarrow 0$$

and by Proposition 6.24, we obtain

$$\text{Ext}^1(\mathcal{P}_{r,s}^{+d}, \mathcal{X}_{r,s}^+) = 0.$$

The second equation can be proved in the same way, so we omit the proofs. \square

The following lemma can be proved in the same way as Proposition 5.28, by using the structure of the logarithmic Virasoro modules $(\tilde{F}(\tau), \tilde{J}_\tau)$.

Lemma 6.42. *Any indecomposable Virasoro module E in*

$$\text{Ext}_{\mathcal{L}}^1(\tilde{K}(\Delta_{r,s;1}^+), L(\Delta_{r^\vee,s;0}^-, \Delta_{r,s;1}^+) \oplus L(\Delta_{r,s^\vee;0}^-, \Delta_{r,s;1}^+)) \quad (6.39)$$

has a indecomposable submodule isomorphic to $\tilde{P}(\Delta_{r,s;1}^+)$ (for the definition of $\tilde{P}(\Delta_{r,s;1}^+)$, see the proof of 5.20), and satisfies

$$\text{Hom}_{U(\mathcal{L})}(E, \tilde{K}(\Delta_{r,s;1}^+)) \simeq \mathbb{C}^3. \quad (6.40)$$

Furthermore, we can choose three surjections as a basis of (6.40).

From the structure of $\mathcal{E}^\pm(\mathcal{X}_{\bullet,\bullet}^-)_{\bullet,\bullet}$, we obtain the following lemma.

Lemma 6.43. *Let (a, b, c, d) be any element in*

$$\{(r^\vee, s, r, s), (r^\vee, s, r^\vee, s^\vee), (r, s^\vee, r, s), (r, s^\vee, r^\vee, s^\vee)\},$$

and fix any indecomposable module E in $\text{Ext}^1(\mathcal{X}_{a,b}^-, \mathcal{X}_{c,d}^+)$. For the surjection $\pi : E \rightarrow \mathcal{X}_{a,b}^-$, let $(v^-, v^+) \in E[\Delta_{a,b;0}^-]^2$ be any generators of E satisfying

$$W^\pm[0]\pi(v^\pm) = 0, \quad W^\pm[0]\pi(v^\mp) \in \mathbb{C}^\times \pi(v^\pm).$$

Then, for $\epsilon = \pm$, the Virasoro module $U(\mathcal{L}).v^\epsilon$ has a quotient isomorphic to $L(\Delta_{a,b;0}^-, \Delta_{c,d;1}^+)$.

Proposition 6.44.

$$\text{Ext}^1(\mathcal{P}_{r,s}^{+u}/(2\mathcal{X}_{r,s}^+), \mathcal{X}_{r,s}^+) = \text{Ext}^1(\mathcal{P}_{r^\vee,s^\vee}^{+u}/(2\mathcal{X}_{r^\vee,s^\vee}^+), \mathcal{X}_{r^\vee,s^\vee}^+) = \mathbb{C}^2.$$

Proof. Let us prove the first equality. The second equality can be proved in the same way. Since $\text{Ext}^1(\mathcal{X}_{r,s}^+, \mathcal{X}_{r,s}^+) = 0$, it is sufficient to show that

$$\text{Ext}^1(\mathcal{P}_{r,s}^{+u}, \mathcal{X}_{r,s}^+) = 0.$$

Assume $\text{Ext}^1(\mathcal{P}_{r,s}^{+u}, \mathcal{X}_{r,s}^+) \neq 0$ and fix any non-trivial extension E in this Ext^1 -group.

First let us introduce some symbols. Let t be a nonzero vector of the one dimensional space $E[h_{r,s}]$, and let u be the highest weight vector of $\mathcal{X}_{r,s}^+$. From Proposition 3.13, we can choose a basis $\{w_-, w_0, w_+\}$ of the Virasoro highest weight spaces of $3L(\Delta_{r,s;1}^+) \subset \mathcal{X}_{r,s}^+$ satisfying

$$\begin{aligned} W^\pm[0]w_\pm &\equiv 0 \pmod{U(\mathcal{L}).u}, \\ W^\pm[0]w_\mp &\in \mathbb{C}^\times w_0 + U(\mathcal{L}).u, \\ W^\pm[0]w_0 &\in \mathbb{C}^\times w_\pm + U(\mathcal{L}).u. \end{aligned}$$

Let ι_1, ι_2 and ι_3 be injections from $\mathcal{X}_{r,s}^+$ to E such that

$$\iota_1(\mathcal{X}_{r,s}^+) \oplus \iota_2(\mathcal{X}_{r,s}^+) \oplus \iota_3(\mathcal{X}_{r,s}^+) = \text{Soc}(E).$$

For the surjection $\pi : E \rightarrow \mathcal{X}_{r,s}^+$, we fix L_0 -homogeneous vectors $\tilde{u}_0, \tilde{w}_0 \in E$ such that

$$\pi(\tilde{u}_0) = u, \quad \pi(\tilde{w}_0) = w_0.$$

Set $v_{r^\vee,s}^\pm = W^\pm[0]W^\mp[0]S_{r,s^\vee+p_-}\tilde{u}_0$ and $v_{r^\vee,s}^\pm = W^\pm[0]W^\mp[0]S_{r^\vee+p_+,s}\tilde{u}_0$. Note that the sets

$$\{v_{r^\vee,s}^-, W^+[0]v_{r^\vee,s}^-\}, \quad \{W^-[0]v_{r^\vee,s}^+, v_{r^\vee,s}^+\}$$

correspond to bases of the highest weight spaces of the composition factors $\mathcal{X}_{r^\vee,s}^- \oplus \mathcal{X}_{r^\vee,s}^-$, and

$$\{v_{r,s^\vee}^-, W^+[0]v_{r,s^\vee}^-\}, \quad \{W^-[0]v_{r,s^\vee}^+, v_{r,s^\vee}^+\}$$

to bases of the highest weight spaces of the composition factors $\mathcal{X}_{r,s^\vee}^- \oplus \mathcal{X}_{r,s^\vee}^-$.

Let us consider the Virasoro module

$$M = U(\mathcal{L}).\tilde{w}_0 + \sum_{\epsilon=\pm} U(\mathcal{L}).v_{r^\vee,s}^\epsilon + \sum_{\epsilon'=\pm} U(\mathcal{L}).v_{r,s^\vee}^{\epsilon'}.$$

From the Virasoro structure of $\mathcal{P}_{r,s}^{+u}$, we see that the Virasoro module M has an indecomposable quotient in the Ext^1 -group (6.39), and the quotient module

$$M / \left(\sum_{\epsilon=\pm} U(\mathcal{L}).v_{r^\vee,s}^\epsilon + \sum_{\epsilon'=\pm} U(\mathcal{L}).v_{r,s^\vee}^{\epsilon'} \right)$$

does not contain $L(\Delta_{r^\vee, s; 0}^-)$ and $L(\Delta_{r, s^\vee; 0}^-)$ as the composition factors. Thus by Lemma 6.42, we have

$$\begin{aligned}\sigma(S_{r^\vee, s^\vee+2p_-})\tilde{w}_0 &\in \mathbb{C}^\times v_{r^\vee, s}^+ + \mathbb{C}^\times v_{r^\vee, s}^- + \sum_{i=1}^3 U(\mathcal{L}).\iota_i(u) + U(\mathcal{L}).t, \\ \sigma(S_{r^\vee+2p_+, s^\vee})\tilde{w}_0 &\in \mathbb{C}^\times v_{r, s^\vee}^+ + \mathbb{C}^\times v_{r, s^\vee}^- + \sum_{i=1}^3 U(\mathcal{L}).\iota_i(u) + U(\mathcal{L}).t.\end{aligned}\tag{6.41}$$

Similar to the argument in the proof of Proposition 6.17, from the structure of $\tilde{P}(\Delta_{r, s; 1}^+)$, we have

$$\begin{aligned}S_{r^\vee, s^\vee+2p_-}\sigma(S_{r^\vee, s^\vee+2p_-})\tilde{w}_0 &\not\equiv 0 \pmod{\sum_{i=1}^3 U(\mathcal{L}).\iota_i(u) + U(\mathcal{L}).t}, \\ S_{r^\vee+2p_+, s^\vee}\sigma(S_{r^\vee+2p_+, s^\vee})\tilde{w}_0 &\not\equiv 0 \pmod{\sum_{i=1}^3 U(\mathcal{L}).\iota_i(u) + U(\mathcal{L}).t}, \\ S_{r^\vee, s^\vee+2p_-}\sigma(S_{r^\vee, s^\vee+2p_-})\tilde{w}_0 &\equiv \sum_{i=1}^3 k_i \iota_i(w_0) \pmod{\sum_{j=1}^3 U(\mathcal{L}).\iota_j(u) + U(\mathcal{L}).t}, \\ S_{r^\vee+2p_+, s^\vee}\sigma(S_{r^\vee+2p_+, s^\vee})\tilde{w}_0 &\equiv \sum_{i=1}^3 l_i \iota_i(w_0) \pmod{\sum_{j=1}^3 U(\mathcal{L}).\iota_j(u) + U(\mathcal{L}).t},\end{aligned}\tag{6.42}$$

where (k_i, l_i) ($i = 1, 2, 3$) are some constants. Since $\text{Ext}^1(\mathcal{X}_{r, s}^+, \mathcal{X}_{r, s}^+) = 0$, we have

$$\begin{aligned}\mathcal{W}_{p_+, p_-} \cdot \left(\sum_{i=1}^3 k_i \iota_i(w_0) \right) &\simeq \mathcal{X}_{r, s}^+, \\ \mathcal{W}_{p_+, p_-} \cdot \left(\sum_{i=1}^3 l_i \iota_i(w_0) \right) &\simeq \mathcal{X}_{r, s}^+.\end{aligned}\tag{6.43}$$

We set

$$E' = \frac{E}{\mathcal{W}_{p_+, p_-} \cdot \left(\sum_{i=1}^3 k_i \iota_i(w_0) \right) \oplus \mathcal{W}_{p_+, p_-} \cdot \left(\sum_{i=1}^3 l_i \iota_i(w_0) \right)}.$$

Let ϕ be the surjection from E to E' . Then, from (6.42) and (6.43), E' satisfies

$$[E'] \in \text{Ext}^1(\mathcal{P}_{r, s}^{+u} / (\mathcal{X}_{r, s}^+ \oplus \mathcal{X}_{r, s}^+), \mathcal{X}_{r, s}^+ \setminus \{0\})$$

and

$$\begin{aligned}
S_{r^\vee, s^\vee + 2p_-} \sigma(S_{r^\vee, s^\vee + 2p_-}) \phi(\tilde{w}_0) &\in \sum_{i=1}^3 U(\mathcal{L}).\phi \circ \iota_i(u) + U(\mathcal{L}).\phi(t), \\
S_{r^\vee + 2p_+, s^\vee} \sigma(S_{r^\vee + 2p_+, s^\vee}) \phi(\tilde{w}_0) &\in \sum_{i=1}^3 U(\mathcal{L}).\phi \circ \iota_i(u) + U(\mathcal{L}).\phi(t).
\end{aligned} \tag{6.44}$$

From Lemma 6.42, (6.41) and (6.44), we have

$$\begin{aligned}
S_{r^\vee, s^\vee + 2p_-} \phi(v_{r^\vee, s}^\pm) &\in \sum_{i=1}^3 U(\mathcal{L}).\phi \circ \iota_i(u) + U(\mathcal{L}).\phi(t), \\
S_{r^\vee + 2p_+, s^\vee} \phi(v_{r^\vee, s^\vee}^\pm) &\in \sum_{i=1}^3 U(\mathcal{L}).\phi \circ \iota_i(u) + U(\mathcal{L}).\phi(t).
\end{aligned} \tag{6.45}$$

Then, by Lemma 6.43 and (6.45), E' has $2\mathcal{X}_{r^\vee, s}^- \oplus 2\mathcal{X}_{r^\vee, s^\vee}^-$ as submodules. Thus E' has an indecomposable quotient in $\text{Ext}^1(\mathcal{K}_{r, s}^*, \mathcal{X}_{r, s}^+)$. But this contradicts Proposition 6.24. \square

Theorem 6.45. *For $\mathcal{P}_{r, s}^+$, $\mathcal{P}_{r^\vee, s^\vee}^+ \in C_{r, s}^{thick}$, we have*

$$\text{Ext}^1(\mathcal{P}_{r, s}^+, \mathcal{X}_{r, s}^+) = \text{Ext}^1(\mathcal{P}_{r^\vee, s^\vee}^+, \mathcal{X}_{r^\vee, s^\vee}^+) = 0.$$

Proof. From the exact sequence

$$0 \rightarrow \mathcal{P}_{r, s}^{+d} \rightarrow \mathcal{P}_{r, s}^+ \rightarrow \mathcal{P}_{r, s}^+ / \mathcal{P}_{r, s}^{+d} \rightarrow 0$$

and Proposition 6.41, we have the following exact sequence

$$0 \rightarrow \mathbb{C} \rightarrow \mathbb{C} \rightarrow \mathbb{C}^2 \rightarrow \text{Ext}^1(\mathcal{P}_{r, s}^+ / \mathcal{P}_{r, s}^{+d}, \mathcal{X}_{r, s}^+) \rightarrow \text{Ext}^1(\mathcal{P}_{r, s}^+, \mathcal{X}_{r, s}^+) \rightarrow 0.$$

By Proposition 6.44, we have

$$\text{Ext}^1(\mathcal{P}_{r, s}^+ / \mathcal{P}_{r, s}^{+d}, \mathcal{X}_{r, s}^+) = \mathbb{C}^2.$$

Thus we have

$$\text{Ext}^1(\mathcal{P}_{r, s}^+, \mathcal{X}_{r, s}^+) = 0.$$

The second equality can be proved in the same way. \square

Definition 6.46. *Let (a, b) be (r^\vee, s) or (r, s^\vee) . We define*

$$\mathcal{P}_{a, b}^{-u} := \frac{\mathcal{P}_{a, b}^-}{(\mathcal{P}_{a, b}^- : \mathcal{X}_{a^\vee, b^\vee}^- : 2) + (\mathcal{P}_{a, b}^- : L(h_{r, s}) : 2)}.$$




Figure 6.5: The embedding structure of the \mathcal{W}_{p_+,p_-} -modules in Definition 6.46. The \heartsuit corresponds to the simple module $\mathcal{X}_{r,s}^+$, \diamond to $\mathcal{X}_{r^\vee,s^\vee}^+$, \spadesuit to \mathcal{X}_{r,s^\vee}^- and \clubsuit to $\mathcal{X}_{r^\vee,s}^-$.

Remark 6.47. Figure 6.5 represents the embedding structure of the quotient modules $\mathcal{P}_{r,s^\vee}^{-u}$ and $\mathcal{P}_{r^\vee,s}^{-u}$ given in Definition 6.46.

Proposition 6.48.

$$\text{Ext}^1(\mathcal{P}_{r^\vee,s}^{-u}/(2\mathcal{X}_{r^\vee,s}^-), \mathcal{X}_{r^\vee,s}^-) = \text{Ext}^1(\mathcal{P}_{r,s^\vee}^{-u}/(2\mathcal{X}_{r,s^\vee}^-), \mathcal{X}_{r,s^\vee}^-) = \mathbb{C}^2.$$

Proof. Let us prove the first equality. The second equality can be proved in the same way. Since $\text{Ext}^1(\mathcal{X}_{r^\vee,s}^-, \mathcal{X}_{r^\vee,s}^-) = 0$, it is sufficient to show that

$$\text{Ext}^1(\mathcal{P}_{r^\vee,s}^{-u}, \mathcal{X}_{r^\vee,s}^-) = 0.$$

Assume $\text{Ext}^1(\mathcal{P}_{r^\vee,s}^{-u}, \mathcal{X}_{r^\vee,s}^-) \neq 0$. Fix a non-trivial extension

$$0 \rightarrow \mathcal{X}_{r^\vee,s}^- \xrightarrow{\iota} E \xrightarrow{\phi} \mathcal{P}_{r^\vee,s}^{-u} \rightarrow 0.$$

By Propositions 6.14 and 6.26, we see that

$$\text{Soc}(E) = \iota(\mathcal{X}_{r^\vee,s}^-) \oplus 2\mathcal{X}_{r^\vee,s}^-.$$

Let ι_1 and ι_2 be injections from $\mathcal{X}_{r^\vee,s}^-$ to E such that

$$\phi \circ \iota_1(\mathcal{X}_{r^\vee,s}^-) \oplus \phi \circ \iota_2(\mathcal{X}_{r^\vee,s}^-) = \mathcal{X}_{r^\vee,s}^- \oplus \mathcal{X}_{r^\vee,s}^- = \text{Soc}(\mathcal{P}_{r^\vee,s}^{-u}),$$

and let $\{v_+, v_-\}$ be a basis of the highest weight space of $\mathcal{X}_{r^\vee,s}^-$ such that

$$W^\pm[0]v_\pm = 0, \quad W^\pm[0]v_\mp \in \mathbb{C}^\times v_\pm.$$

For the canonical surjection $\pi : E \rightarrow \mathcal{X}_{r^\vee, s}^-$, we fix any L_0 -homogeneous vectors $\tilde{v}_-, \tilde{v}_+ \in E$ such that $\pi(\tilde{v}_\pm) = v_\pm$. Note that, by Proposition 5.20, the Virasoro module $U(\mathcal{L}).\tilde{v}_\pm$ has $\tilde{P}(\Delta_{r^\vee, s; 0}^-)$ as the quotient (for the definition of $\tilde{P}(\Delta_{r^\vee, s; 0}^-)$, see the proof of 5.20). Let

$$E_1 = (E : \mathcal{X}_{r, s}^+ : 1), \quad E_2 = (E : \mathcal{X}_{r^\vee, s^\vee}^+ : 1).$$

Then, by Proposition 6.26, we see that E_1 and E_2 must contain $\iota(\mathcal{X}_{r^\vee, s}^-)$ as the submodule:

$$(\iota(\mathcal{X}_{r^\vee, s}^-) \subset E_1) \wedge (\iota(\mathcal{X}_{r^\vee, s}^-) \subset E_2). \quad (6.46)$$

Noting the structure of $\tilde{P}(\Delta_{r^\vee, s; 0}^-)$, $\tilde{\mathcal{E}}(\mathcal{X}_{r, s}^+)_{r^\vee, s}$ and $\tilde{\mathcal{E}}(\mathcal{X}_{r^\vee, s^\vee}^+)_{r^\vee, s}$, from (6.46), we have

$$\begin{aligned} (L_0 - \Delta_{r^\vee, s; 0}^-)\tilde{v}_+ &\in \mathbb{C}^\times \iota(v_-) + \mathbb{C} \iota(v_+) + \mathbb{C}^\times \iota_1(v_+) + \mathbb{C}^\times \iota_2(v_+), \\ (L_0 - \Delta_{r^\vee, s; 0}^-)\tilde{v}_- &\in \mathbb{C}^\times \iota(v_+) + \mathbb{C} \iota(v_-) + \mathbb{C}^\times \iota_1(v_-) + \mathbb{C}^\times \iota_2(v_-). \end{aligned} \quad (6.47)$$

Multiplying $(L_0 - \Delta_{r^\vee, s; 0}^-)\tilde{v}_\pm$ by $W^\pm[0]$, from (6.47), we have $\iota(v_\pm) = 0$. But this is a contradiction. \square

Theorem 6.49.

$$\text{Ext}^1(\mathcal{P}_{r^\vee, s}^-, \mathcal{X}_{r^\vee, s}^-) = \text{Ext}^1(\mathcal{P}_{r, s^\vee}^-, \mathcal{X}_{r, s^\vee}^-) = 0.$$

Proof. By the exact sequence

$$0 \rightarrow \mathcal{Q}(\mathcal{X}_{r^\vee, s}^-)_{r^\vee, s^\vee} \rightarrow \mathcal{P}_{r^\vee, s}^- \rightarrow \mathcal{P}_{r^\vee, s}^- / \mathcal{Q}(\mathcal{X}_{r^\vee, s}^-)_{r^\vee, s^\vee} \rightarrow 0$$

and Proposition 6.26, we have the following exact sequence

$$0 \rightarrow \mathbb{C} \rightarrow \mathbb{C} \rightarrow \mathbb{C} \rightarrow \text{Ext}^1(\mathcal{P}_{r^\vee, s}^- / \mathcal{Q}(\mathcal{X}_{r^\vee, s}^-)_{r^\vee, s^\vee}, \mathcal{X}_{r^\vee, s}^-) \rightarrow \text{Ext}^1(\mathcal{P}_{r^\vee, s}^-, \mathcal{X}_{r^\vee, s}^-) \rightarrow 0. \quad (6.48)$$

Note that $\mathcal{P}_{r^\vee, s}^- / \mathcal{Q}(\mathcal{X}_{r^\vee, s}^-)_{r^\vee, s^\vee}$ has a submodule isomorphic to $\mathcal{K}_{r, s} \oplus \mathcal{K}_{r, s}$. Let \mathcal{M} be the quotient module defined by the following exact sequence

$$0 \rightarrow \mathcal{K}_{r, s} \oplus \mathcal{K}_{r, s} \rightarrow \mathcal{P}_{r^\vee, s}^- / \mathcal{Q}(\mathcal{X}_{r^\vee, s}^-)_{r^\vee, s^\vee} \rightarrow \mathcal{M} \rightarrow 0.$$

By this exact sequence and by $\text{Ext}^1(\mathcal{K}_{r, s}, \mathcal{X}_{r^\vee, s}^-) = 0$ (see the proof of Proposition 6.30), we have

$$\text{Ext}^1(\mathcal{M}, \mathcal{X}_{r^\vee, s}^-) \simeq \text{Ext}^1(\mathcal{P}_{r^\vee, s}^- / \mathcal{Q}(\mathcal{X}_{r^\vee, s}^-)_{r^\vee, s^\vee}, \mathcal{X}_{r^\vee, s}^-). \quad (6.49)$$

Note that \mathcal{M} satisfies the exact sequence

$$0 \rightarrow \mathcal{X}_{r^\vee, s}^- \oplus 4\mathcal{X}_{r, s^\vee}^- \rightarrow \mathcal{M} \rightarrow \mathcal{P}_{r^\vee, s}^{-u} / (2\mathcal{X}_{r^\vee, s}^-) \rightarrow 0.$$

By this exact sequence and Proposition 6.48, we have the following exact sequence

$$0 \rightarrow \mathbb{C} \rightarrow \mathbb{C} \rightarrow \mathbb{C} \rightarrow \mathbb{C}^2 \rightarrow \text{Ext}^1(\mathcal{M}, \mathcal{X}_{r^\vee, s}^-) \rightarrow 0.$$

Thus, by this exact sequence, we have $\text{Ext}^1(\mathcal{M}, \mathcal{X}_{r^\vee, s}^-) \simeq \mathbb{C}$. Therefore, by (6.48) and (6.49), we obtain

$$\text{Ext}^1(\mathcal{P}_{r^\vee, s}^-, \mathcal{X}_{r^\vee, s}^-) = 0.$$

The second equality can be proved in the same way. \square

Since all logarithmic modules $\mathcal{P}_{\bullet, \bullet}^\pm$ in $C_{r, s}^{thick}$ are generated from the top composition factors, by Theorems 6.34, 6.35, 6.45 and 6.49, we obtain the following theorem.

Theorem 6.50. *Let (a, b, ϵ) be any element in*

$$\{(r, s, +), (r^\vee, s^\vee, +), (r^\vee, s, -), (r, s^\vee, -)\}.$$

Then the indecomposable module $\mathcal{P}_{a, b}^\epsilon$ is the projective cover of $\mathcal{X}_{a, b}^\epsilon$.

6.5 The projective covers of the minimal simple modules $L(h_{r, s})$

Fix any thick block $C_{r, s}^{thick}$. Let $\mathcal{P}(h_{r, s})$ be the projective cover of the minimal simple module $L(h_{r, s})$. By Corollary 3.16, we can see that $\mathcal{P}(h_{r, s})$ has L_0 nilpotent rank three. In this subsection, we determine the structure of $\mathcal{P}(h_{r, s})$.

Definition 6.51. *Let (a, b) be (r, s) or (r^\vee, s^\vee) . We define*

$$\mathcal{N}_{a, b} := (\mathcal{P}_{a, b}^+ : L(h_{r, s}) : 1).$$

For $(a, b) = (r, s)$ or (r^\vee, s^\vee) , $\mathcal{N}_{a, b}$ has the following length four socle series

$$\begin{aligned} \text{Soc}_1(\mathcal{N}_{a, b}) &= \mathcal{X}_{a, b}^+, \\ \text{Soc}_2(\mathcal{N}_{a, b}) / \text{Soc}_1(\mathcal{N}_{a, b}) &= 2\mathcal{X}_{a^\vee, b}^- \oplus L(h_{r, s}) \oplus 2\mathcal{X}_{a, b^\vee}^-, \\ \text{Soc}_3(\mathcal{N}_{a, b}) / \text{Soc}_2(\mathcal{N}_{a, b}) &= \mathcal{X}_{a, b}^+ \oplus \mathcal{X}_{a^\vee, b^\vee}^+, \\ \mathcal{N}_{a, b} / \text{Soc}_3(\mathcal{N}_{a, b}) &= L(h_{r, s}). \end{aligned}$$




Figure 6.6: The embedding structure of the logarithmic \mathcal{W}_{p_+, p_-} -modules $\mathcal{N}_{r,s}$ and $\mathcal{N}_{r^\vee, s^\vee}$. The triangle \triangle corresponds to the simple module $L(h_{r,s})$, \heartsuit to $\mathcal{X}_{r,s}^+$, \diamond to $\mathcal{X}_{r^\vee, s^\vee}^+$, \spadesuit to $\mathcal{X}_{r,s}^-$ and \clubsuit to $\mathcal{X}_{r^\vee, s^\vee}^-$.

Remark 6.52. Figure 6.6 represents the embedding structure of the logarithmic \mathcal{W}_{p_+, p_-} -modules $\mathcal{N}_{r,s}$ and $\mathcal{N}_{r^\vee, s^\vee}$.

We define

$$\mathcal{Q}(h_{r,s}) := \frac{\mathcal{N}_{r,s}}{(\mathcal{N}_{r,s} : \mathcal{X}_{r^\vee, s^\vee}^- : 2) + (\mathcal{N}_{r,s} : \mathcal{X}_{r,s}^- : 2)}.$$

By Propositions 6.14 and by the proof of Proposition 6.29, we obtain the following lemmas.

Lemma 6.53. The socle series of the indecomposable module $\mathcal{Q}(h_{r,s})$ is given by

$$\text{Soc}_1 = L(h_{r,s}), \quad \text{Soc}_2/\text{Soc}_1 = \mathcal{X}_{r,s}^+ \oplus \mathcal{X}_{r^\vee, s^\vee}^+, \quad \text{Soc}_3/\text{Soc}_2 = L(h_{r,s}).$$

Lemma 6.54.

$$\text{Ext}^1(\mathcal{Q}(h_{r,s}), L(h_{r,s})) = 0.$$

Recall the indecomposable modules $\mathcal{P}_{r,s}^{+u}$ and $\mathcal{P}_{r^\vee, s^\vee}^{+u}$ given in Definition 6.39. Note that $\mathcal{P}_{r,s}^{+u}$ and $\mathcal{P}_{r^\vee, s^\vee}^{+u}$ have $\mathcal{K}_{r,s}$ and $\mathcal{K}_{r^\vee, s^\vee}$ as submodules respectively. We define the indecomposable modules

$$\mathcal{R}_{r,s} := \mathcal{P}_{r,s}^{+u} / \mathcal{K}_{r,s}, \quad \mathcal{R}_{r^\vee, s^\vee} := \mathcal{P}_{r^\vee, s^\vee}^{+u} / \mathcal{K}_{r^\vee, s^\vee}.$$

By Proposition 5.3 and by Proposition 6.14, we obtain the following lemma.

Lemma 6.55. *Let (a, b) be (r, s) or (r^\vee, s^\vee) . Then the indecomposable module $\mathcal{R}_{a,b}$ has L_0 nilpotent rank two and has the following length three socle series*

$$\begin{aligned}\text{Soc}_1 &= \mathcal{X}_{a,b}^+, \\ \text{Soc}_2/\text{Soc}_1 &= \mathcal{X}_{r^\vee, s}^- \oplus \mathcal{X}_{r^\vee, s}^- \oplus \mathcal{X}_{r, s^\vee}^- \oplus \mathcal{X}_{r, s^\vee}^-, \\ \mathcal{R}_{a,b}/\text{Soc}_2 &= \mathcal{X}_{a,b}^+.\end{aligned}$$

By Propositions 6.14, 6.44, we obtain the following lemma.

Lemma 6.56.

$$\text{Ext}^1(\mathcal{R}_{r,s}, \mathcal{X}_{r,s}^+) = \text{Ext}^1(\mathcal{R}_{r^\vee, s^\vee}, \mathcal{X}_{r^\vee, s^\vee}^+) = 0.$$

Lemma 6.57.

$$\text{Ext}^1(\mathcal{N}_{r,s}/\mathcal{K}_{r,s}, \mathcal{X}_{r,s}^+) = \text{Ext}^1(\mathcal{N}_{r^\vee, s^\vee}/\mathcal{K}_{r^\vee, s^\vee}, \mathcal{X}_{r^\vee, s^\vee}^+) = 0.$$

Proof. We only prove $\text{Ext}^1(\mathcal{N}_{r,s}/\mathcal{K}_{r,s}, \mathcal{X}_{r,s}^+) = 0$. The second equality can be proved in the same way.

Assume that $\text{Ext}^1(\mathcal{N}_{r,s}/\mathcal{K}_{r,s}, \mathcal{X}_{r,s}^+) \neq 0$ and fix any non-trivial extension E in this Ext^1 -group. Let $t \in E$ be the highest weight vector in the one dimensional space $E[h_{r,s}]$. By Lemma 6.56, we see that E has a submodule isomorphic to $\mathcal{R}_{r,s}$. Note that by Lemma 6.55, $\mathcal{R}_{r,s}$ has L_0 nilpotent rank two. Then we have a basis $\{u_0, u_1\}$ of the highest weight space of the submodule $\mathcal{R}_{r,s} \subset E$ satisfying

$$(L_0 - \Delta_{r,s;0}^+)u_0 \in \mathbb{C}^\times u_1. \quad (6.50)$$

On the other hand, by

$$(L_0 - h_{r,s})t = 0, \quad S_{r^\vee, s^\vee}t \in \mathbb{C}^\times u_0 + \mathbb{C}u_1,$$

we have $(L_0 - \Delta_{r,s;0}^+)u_0 = 0$. But this contradicts (6.50). \square

Lemma 6.58.

$$\text{Ext}^1(\mathcal{N}_{r,s}, \mathcal{X}_{r^\vee, s^\vee}^+) = \mathbb{C}. \quad (6.51)$$

Proof. By the exact sequence

$$0 \rightarrow \mathcal{K}_{r,s} \rightarrow \mathcal{N}_{r,s} \rightarrow \mathcal{N}_{r,s}/\mathcal{K}_{r,s} \rightarrow 0,$$

we have the exact sequence

$$0 \rightarrow \text{Ext}^1(\mathcal{N}_{r,s}/\mathcal{K}_{r,s}, \mathcal{X}_{r^\vee, s^\vee}^+) \rightarrow \text{Ext}^1(\mathcal{N}_{r,s}, \mathcal{X}_{r^\vee, s^\vee}^+) \rightarrow \mathbb{C}. \quad (6.52)$$

By Proposition 6.14, we have $\mathcal{N}_{r,s}/\mathcal{K}_{r,s} \simeq \mathcal{N}_{r^\vee, s^\vee}/\mathcal{K}_{r^\vee, s^\vee}$. Then, by Lemma 6.57 and (6.52), we obtain

$$\dim_{\mathbb{C}} \text{Ext}^1(\mathcal{N}_{r,s}, \mathcal{X}_{r^\vee, s^\vee}^+) \leq 1. \quad (6.53)$$

Note that by Proposition 6.14 and Lemma 6.54,

$$\mathcal{N}_{r,s}/\mathcal{X}_{r,s}^+ \simeq \mathcal{N}_{r^\vee, s^\vee}/\mathcal{X}_{r^\vee, s^\vee}^+.$$

Then, we have $\text{Ext}^1(\mathcal{N}_{r,s}, \mathcal{X}_{r^\vee, s^\vee}^+) \neq 0$. Thus, by (6.53), we obtain (6.51). \square

We identify the indecomposable modules in the Ext^1 -group (6.51) and denote them as $\tilde{\mathcal{N}}(h_{r,s})$. By Lemma 6.57, we obtain the following lemma.

Lemma 6.59.

$$\text{Ext}^1(\tilde{\mathcal{N}}(h_{r,s}), \mathcal{X}_{r,s}^+) = \text{Ext}^1(\tilde{\mathcal{N}}(h_{r,s}), \mathcal{X}_{r^\vee, s^\vee}^+) = 0.$$

Lemma 6.60.

$$\text{Ext}^1(\tilde{\mathcal{N}}(h_{r,s}), \mathcal{X}_{r^\vee, s}^-) = \text{Ext}^1(\tilde{\mathcal{N}}(h_{r,s}), \mathcal{X}_{r, s^\vee}^-) = 0.$$

Proof. We only prove the first equality. The second equality can be proved in the same way. Assume

$$\text{Ext}^1(\tilde{\mathcal{N}}(h_{r,s}), \mathcal{X}_{r^\vee, s}^-) \neq 0$$

and fix any non-trivial extension E in this Ext^1 -group. Then we see that E has an indecomposable submodule in

$$\text{Ext}^1(\tilde{\mathcal{E}}(\mathcal{X}_{r,s}^+)_{r^\vee, s}^*, \mathcal{E}^\pm(\mathcal{X}_{r^\vee, s^\vee}^+)_{r^\vee, s}) \quad \text{or} \quad \text{Ext}^1(\tilde{\mathcal{E}}(\mathcal{X}_{r^\vee, s^\vee}^+)_{r^\vee, s}^*, \mathcal{E}^\pm(\mathcal{X}_{r,s}^+)_{r^\vee, s})$$

(for the definition of $\tilde{\mathcal{E}}(\mathcal{X}_{r,s}^+)_{r^\vee, s}$ and $\tilde{\mathcal{E}}(\mathcal{X}_{r^\vee, s^\vee}^+)_{r^\vee, s}$, see Definition 6.7). From Propositions 6.24 and 6.26, we see that these Ext^1 -groups are trivial, and thus we have a contradiction. \square

Let us assume that $\text{Ext}^1(\tilde{\mathcal{N}}(h_{r,s}), L(h_{r,s})) = 0$. Then, by Lemmas 6.59 and 6.60, we see that $\tilde{\mathcal{N}}(h_{r,s})$ is the projective cover of $L(h_{r,s})$. But this contradicts Corollary 3.16. Thus we obtain

$$\text{Ext}^1(\tilde{\mathcal{N}}(h_{r,s}), L(h_{r,s})) \neq 0.$$

Fix any indecomposable module in this Ext^1 -group and denote it by $\mathcal{P}'(h_{r,s})$. $\mathcal{P}'(h_{r,s})$ has the following length five socle series

$$\begin{aligned}\text{Soc}_1 &= L(h_{r,s}), \\ \text{Soc}_2/\text{Soc}_1 &= \mathcal{X}_{r,s}^+ \oplus \mathcal{X}_{r^\vee, s^\vee}^+, \\ \text{Soc}_3/\text{Soc}_2 &= 2\mathcal{X}_{r^\vee, s}^- \oplus L(h_{r,s}) \oplus 2\mathcal{X}_{r, s^\vee}^-, \\ \text{Soc}_4/\text{Soc}_3 &= \mathcal{X}_{r,s}^+ \oplus \mathcal{X}_{r^\vee, s^\vee}^+, \\ \text{Soc}_5/\text{Soc}_4 &= L(h_{r,s}).\end{aligned}$$

Let us consider the submodule $(\mathcal{P}'(h_{r,s}) : L(h_{r,s}) : 2)$. By Lemm 6.54, we see that this submodule is isomorphic to $\mathcal{Q}(h_{r,s})$.

Lemma 6.61.

$$\text{Ext}^1(\mathcal{P}'(h_{r,s}), L(h_{r,s})) = 0.$$

Proof. By Lemm 6.54 and by the exact sequence

$$0 \rightarrow \mathcal{Q}(h_{r,s}) \rightarrow \mathcal{P}'(h_{r,s}) \rightarrow \mathcal{P}'(h_{r,s})/\mathcal{Q}(h_{r,s}) \rightarrow 0,$$

we obtain the exact sequence

$$0 \rightarrow \mathbb{C} \rightarrow \mathbb{C} \rightarrow \mathbb{C} \rightarrow \mathbb{C} \rightarrow \text{Ext}^1(\mathcal{P}'(h_{r,s}), L(h_{r,s})) \rightarrow 0.$$

From this, we obtain

$$\text{Ext}^1(\mathcal{P}'(h_{r,s}), L(h_{r,s})) = 0.$$

□

Lemma 6.62.

$$\text{Ext}^1(\mathcal{P}'(h_{r,s}), \mathcal{X}_{r,s}^+) = \text{Ext}^1(\mathcal{P}'(h_{r,s}), \mathcal{X}_{r^\vee, s^\vee}^+) = 0.$$

Proof. We only prove the first equality. The second equality can be proved in the same way. By Proposition 6.24, we see that

$$\text{Ext}^1(\mathcal{Q}(h_{r,s}), \mathcal{X}_{r,s}^+) = 0. \tag{6.54}$$

Note that $\mathcal{P}'(h_{r,s})/\mathcal{Q}(h_{r,s}) \simeq \mathcal{N}_{r,s}/\mathcal{K}_{r,s}$. Thus, by Lemma 6.57 and by (6.54), we obtain $\text{Ext}^1(\mathcal{P}'(h_{r,s}), \mathcal{X}_{r,s}^+) = 0$. □

By Proposition 6.14 and Lemma 6.60, we obtain the following lemma.

Lemma 6.63.

$$\text{Ext}^1(\mathcal{P}'(h_{r,s}), \mathcal{X}_{r^\vee, s}^-) = \text{Ext}^1(\mathcal{P}'(h_{r,s}), \mathcal{X}_{r, s^\vee}^-) = 0.$$

By Lemmas 6.61, 6.62 and 6.63, we have $\mathcal{P}(h_{r,s}) \simeq \mathcal{P}'(h_{r,s})$. Therefore we obtain the following theorem.

Theorem 6.64. *The projective module $\mathcal{P}(h_{r,s})$ has the following length five socle series:*

$$\begin{aligned} \text{Soc}_1(\mathcal{P}(h_{r,s})) &= L(h_{r,s}), \\ \text{Soc}_2(\mathcal{P}(h_{r,s}))/\text{Soc}_1(\mathcal{P}(h_{r,s})) &= \mathcal{X}_{r,s}^+ \oplus \mathcal{X}_{r^\vee, s^\vee}^+, \\ \text{Soc}_3(\mathcal{P}(h_{r,s}))/\text{Soc}_2(\mathcal{P}(h_{r,s})) &= 2\mathcal{X}_{r^\vee, s}^- \oplus L(h_{r,s}) \oplus 2\mathcal{X}_{r, s^\vee}^-, \\ \text{Soc}_4(\mathcal{P}(h_{r,s}))/\text{Soc}_3(\mathcal{P}(h_{r,s})) &= \mathcal{X}_{r,s}^+ \oplus \mathcal{X}_{r^\vee, s^\vee}^+, \\ \mathcal{P}(h_{r,s})/\text{Soc}_4(\mathcal{P}(h_{r,s})) &= L(h_{r,s}). \end{aligned}$$




Figure 6.7: The embedding structure of the logarithmic \mathcal{W}_{p_+, p_-} -module $\mathcal{P}(h_{r,s})$. The triangle \triangle corresponds to the simple module $L(h_{r,s})$, \heartsuit to $\mathcal{X}_{r,s}^+$, \diamondsuit to $\mathcal{X}_{r^\vee, s^\vee}^+$, \spadesuit to $\mathcal{X}_{r^\vee, s}^-$ and \clubsuit to $\mathcal{X}_{r, s^\vee}^-$.

Remark 6.65. *Figure 6.7 represents the embedding structure of the projective module $\mathcal{P}(h_{r,s})$. This embedding structure is given by [19].*

Acknowledgement

We would like to thank Akihiro Tsuchiya, Koji Hasegawa, Gen Kuroki and Masaru Sugawara for useful discussions.

References

- [1] D. Adamović and A. Milas, “On the triplet vertex algebra $W(p)$ ”, *Advances in Mathematics* **217** (2008), 2664-2699.
- [2] D. Adamović and A. Milas, “Lattice construction of logarithmic modules for certain vertex algebras”, *Selecta Math. (N.S.)* **15** (2009), 535-561; arXiv:0902.3417.
- [3] D. Adamović and A. Milas, “On \mathcal{W} -algebras associated to $(2, p)$ minimal models for certain vertex algebras”, *International Mathematics Research Notices* 2010 (2010) 20 : 3896-3934, arXiv:0908.4053.
- [4] D. Adamović and A. Milas, “On W -algebra extensions of $(2, p)$ minimal models: $p > 3$ ”, *Journal of Algebra* **344** (2011) 313-332. arXiv:1101.0803.
- [5] D. Adamović and A. Milas, “The structure of Zhu’s algebras for certain W -algebras”, *Advances in Math* 227 (2011) 2425-2456; arXiv:1006.5134.
- [6] D. Adamović and A. Milas, “An explicit realization of logarithmic modules for the vertex operator algebra $W_{p+, p-}$ ”, *J. Math. Phys.* **53**, (2012), 16pp.
- [7] Y. Arike, “A matrix realization of the quantum group $\mathfrak{g}_{p,q}$ ”, *International Journal of Mathematics*, 22.03 (2011): 345-398.
- [8] B. Boe, D. Nakano and E. Wiesner, “Category \mathcal{O} for the Virasoro algebra: cohomology and Koszulity”, *Pacific J. Math.* **234** (2008), no. 1, 1-21.
- [9] T. Creutzig and D. Ridout. “Logarithmic conformal field theory: Beyond an introduction”, *J. Phys. A* 46: 494006, 2013. arXiv:1303.0847 [hep-th].
- [10] M. Cromer, “Free field realisations of staggered modules in 2D logarithmic CFTs”, arXiv preprint arXiv:1612.02909 (2016).

- [11] B. Feigin and D.B. Fuchs. “Representations of the Virasoro algebra”, in *Representations of infinite-dimensional Lie groups and Lie algebras*, Gordon and Breach, New York (1989).
- [12] B. L. Feigin, A.M. Gainutdinov, A.M. Semikhatov, and I. Yu Tipunin, “Logarithmic extensions of minimal models: characters and modular transformation”, *Nuclear Phys. B* 757(2006),303-343.
- [13] B.L. Feigin, A.M. Gainutdinov, A.M. Semikhatov, and I. Yu Tipunin, “Kazhdan-Lusztig-dual quantum group for logarithmic extensions of Virasoro minimal models”, *J. Math. Phys.* 48:032303, 2007.
- [14] B. L. Feigin, A.M. Gainutdinov, A.M. Semikhatov, and I. Yu Tipunin, “Modular group representations and fusion in logarithmic conformal field theories and in the quantum group center”, *Comm. Math. Phys.* 265 (2006), 47–93.
- [15] B. L. Feigin, A.M. Gainutdinov, A.M. Semikhatov, and I. Yu Tipunin, “Kazhdan-Lusztig correspondence for the representation category of the triplet W-algebra in logarithmic CFT”, *Theor. Math. Phys.* 148 (2006) 1210-1235; *Teor. Mat. Fiz.* **148** (2006) 398-427.
- [16] G. Felder, “BRST approach to minimal models”, *Nuc. Phys. B* **317** (1989) 215-236.
- [17] J. Fjelstad, J. Fuchs, S. Hwang, A. M. Semikhatov, and I. Yu. Tipunin, “Logarithmic conformal field theories via logarithmic deformation”, *Nuclear Phys. B* 633, 379-413 (2002).
- [18] M. Gaberdiel, I. Runkel, and S. Wood, “Fusion rules and boundary conditions in the $c = 0$ triplet model”, *J.Phys. A* **42** (2009) 325403, arXiv:0905.0916 [hep-th].
- [19] M. Gaberdiel, I. Runkel, and S. Wood, “A modular invariant bulk theory for the $c = 0$ triplet model”, *J.Phys. A:math. Theor.* 44 (2011) 015204, arXiv:1008.0082v1.
- [20] M. Gorelik and V. Kac, “On complete reducibility for infinite-dimensional Lie algebras”, *Adv. Math.* **226** (2011), no. 2, 1911-1972.
- [21] Y. Z. Huang, “Cofiniteness conditions, projective covers and the logarithmic tensor product theory”, *J. Pure Appl. Algebra*, 213(4):458-475, 2009.

- [22] K. Iohara and Y. Koga, *Representation Theory of the Virasoro Algebra*, Springer Monographs in Mathematics, Berlin, Springer 2011.
- [23] H. G. Kausch, “Extended conformal algebras generated by multiplet of primary fields”, *Phys. Lett. B*, **259** (1991), 448-455.
- [24] K. Kytölä and D. Ridout, “On staggered indecomposable Virasoro modules”, *J. Math. Phys.* 50 (2009) 123503, arXiv:0905.0108[math-ph].
- [25] R. McRae and J. Yang, “Structure of Virasoro tensor categories at central charge $13 - 6p - 6p^{-1}$ for integers $p > 1$ ”, arXiv:2011.02170 (2020).
- [26] A. Milas, “Fusion rings for degenerate minimal models”, *J. Algebra* **254** (2002), no. 2, 300-335.
- [27] K. Nagatomo and A. Tsuchiya, “The Triplet Vertex Operator Algebra $W(p)$ and Restricted Quantum Group at Root of Unity”, *Adv. Stud. in Pure Math., Exploring new Structures and Natural Constructions in Mathematical Physics*, Amer. Math. Soc. **61** (2011) 1–49, arXiv:0902.4607.
- [28] H. Nakano, “The category of modules of the triplet W-algebras associated to the Virasoro minimal models”, *the Doctor Thesis, Mathematical Institute, Tohoku University* (2023).
- [29] H. Nakano, “Explicit formulas of the logarithmic couplings of certain staggered Virasoro modules”, *Letters in Mathematical Physics*, 113(2), 44 (2023).
- [30] X. Lin, “Fusion rules of Virasoro vertex operator algebras”, *Proceedings of the American Mathematical Society* 143.9 (2015):3765-3776.
- [31] J. Rasmussen, “W-extended logarithmic minimal models”, *Nucl. Phys. B* **807** (2009) 495 [0805.2991 [hep-th]].
- [32] A. Tsuchiya and Y. Kanie, “Fock space representations of the Virasoro algebra - Intertwining operators”, *Publ. RIMS, Kyoto Univ.* 22(1986) 259-327.
- [33] A. Tsuchiya and S. Wood, “On the extended W-algebra of type sl_2 at positive rational level” *International Mathematics Research Notices*, Volume 2015, Issue 14, 1 January 2015, Pages 5357-5435.

- [34] S. Wood, “Fusion Rules of the $\mathcal{W}_{p,q}$ Triplet Models”, *J. Phys. A* **43** (2010) 045212.
- [35] S. Yanagida. “Norm of logarithmic primary of Virasoro algebra”, *Letters in Mathematical Physics* 98.2 (2011): 133-156.
- [36] Y. Zhu, “Modular invariance of characters of of vertex operator algebras”, *J. Amer. Math. Soc.* **9** (1996), 237-302.

H. Nakano, OSAKA CITY UNIVERSITY ADVANCED MATHEMATICAL
INSTITUTE
E-mail address : `hiromutakati@gmail.com`