# NATURAL COORDINATES AND HORIZONTAL APPROXIMATIONS IN TWO-FIELD COSMOLOGICAL MODELS

CALIN IULIU LAZAROIU

ABSTRACT. We construct natural local coordinate systems on the phase space of two-field cosmological models with orientable target space, which allow for a description of cosmological flows through quantities of direct physical interest. Such coordinates are induced by the *fundamental observables* of the model, which we formulate geometrically using the tautological bundle of the tangent bundle of the scalar manifold. We also describe a large class of geometric dynamical approximations induced by the choice of an Ehresmann connection in the tangent bundle of the scalar manifold. Such approximations take a conceptually simple form in natural coordinates and we illustrate one of them as an application.

#### INTRODUCTION

Cosmological models with more than one scalar field and arbitrary target space geometry provide natural generalizations of classical one-field inflationary cosmology and are expected to play a crucial role in cosmological applications of fundamental theories of gravity and matter such as superstring theory and its low energy limit (which to leading order is described by supergravity). Such *multifield models* arise naturally in supersymmetric string compactifications (which typically produce a large number of moduli fields) and might be preferred or even required in fundamental theories which allow for a consistent quantization of gravity [1, 2, 3, 4]. This led to renewed interest in such models, which traditionally were much less studied than their one-field counterparts. Multifield models allow for rich behavior which is relevant to various questions in phenomenological cosmology [5, 6, 7, 8, 9, 10, 11, 12].

Despite their crucial conceptual and theoretical importance, multifield cosmological models are poorly understood — mainly because their dynamics (even in the absence of cosmological perturbations) can be extremely nontrivial already in the two-field case, as illustrated for example in [13, 14, 15]. Even when resorting to numerical methods, the identification of solutions with various desirable features is unlikely to be feasible in generic multifield models without sophisticated mathematical tools<sup>1</sup> pertaining to the geometric theory of dynamical systems [19, 20]. It is precisely for this reason that such models are of interest in mathematical physics and require the development of a systematic theory. A few steps towards this were were made in [21, 22].

In the absence of cosmological perturbations, a multifield cosmological model is defined by a nonlinear geometric second order ODE associated to a *scalar triple*  $(\mathcal{M}, \mathcal{G}, \Phi)$ , where  $(\mathcal{M}, \mathcal{G})$  is a connected (but generally non-compact) smooth and complete Riemannian manifold without boundary (called the *scalar manifold* of the model) and  $\Phi$  is a smooth real-valued function defined on  $\mathcal{M}$ , which we assume to be non-constant and strictly positive. To such data one associates the *cosmological* 

 $<sup>^{1}</sup>$ A notable exception is provided by models admitting "hidden symmetries" [16, 17, 18], for which exact solutions can sometimes be found .

equation of the scalar triple:

$$\nabla_t \dot{\varphi}(t) + \frac{1}{M_0} \left[ || \dot{\varphi}(t) ||^2 + 2\Phi(\varphi(t)) \right]^{1/2} \dot{\varphi}(t) + (\text{grad}\Phi)(\varphi(t)) = 0 ,$$

where  $\varphi: I \to \mathcal{M}$  is a smooth curve in  $\mathcal{M}$  defined on a nondegenerate interval Iand the dot stands for derivation with respect to  $t \in I$ . Here  $|| || : T\mathcal{M} \to \mathbb{R}_{\geq 0}$ and grad :  $\mathcal{C}^{\infty}(\mathcal{M}) \to \mathcal{X}(\mathcal{M})$  are the norm function and gradient operator defined by  $\mathcal{G}$ ,  $\nabla$  is the Levi-Civita connection of  $(\mathcal{M}, \mathcal{G})$  and  $\nabla_t \stackrel{\text{def.}}{=} \nabla_{\dot{\varphi}(t)}$ . The rescaled Planck mass  $M_0 > 0$  is a parameter which can be absorbed by redefining  $\mathcal{G}$  -though it is more convenient for various arguments to treat it as a free parameter. The problem of mathematical and physical interest is to study the solutions of this geometric ODE for general scalar triples  $(\mathcal{M}, \mathcal{G}, \Phi)$  and to develop geometric methods for extracting the qualitative features of solutions and for approximating them *efficiently*. Among other questions, one is interested in the asymptotic behavior of solutions which (depending on the asymptotics of  $\Phi$ ) might "escape towards infinity" at the Freudenthal ends of  $\mathcal{M}$ . We refer the reader to [13, 21, 22] for a conceptual discussion of certain aspects of multifield cosmological models with topologically non-trivial target spaces  $\mathcal{M}$ .

The first nontrivial case of interest arises for two-field cosmological models, which comprise the case dim  $\mathcal{M} = 2$  – i.e. when  $(\mathcal{M}, \mathcal{G})$  is a (generally non-compact) Riemann surface. Such models can already display intricate behavior which depends strongly on the topology of  $\mathcal{M}$  and on the asymptotic behavior of  $\Phi$  and  $\mathcal{G}$  towards its Freudenthal ends (see [13, 14, 15]). Despite being the focus of much recent interest, little is known about the behavior of solutions of the cosmological equation for such models, except for special examples which are generally studied using naive approximations or by numerical means. In particular, no systematic theory exists which provides geometric approximation schemes that allow for efficient control of the error terms. Moreover, no global geometric description of quantities of physical interest was developed in the physics literature.

In the present paper, we address some of these issues for two-field models with orientable target space  $\mathcal{M}$  by developing the geometric theory of natural cosmological observables and associated notions while introducing a wide class of approximation schemes (which we call "horizontal approximations") for the cosmological equation, which are defined by the choice of an Ehresmann connection on an open subset of  $T\mathcal{M}$ . As applications, we write the first order form of the cosmological equation in certain "universal" local coordinate systems on  $T\mathcal{M}$  which have direct physical interpretation and describe the horizontal approximation and some of its attending objects defined by one of of these. This turns out to correspond to a proposal made previously in the physics literature, some aspects of which were partially clarified in [23]. Given space limitations, we postpone the analysis of two-field cosmological dynamics in the natural coordinates identified in this paper (as well as an analysis of the error of horizontal approximations) for further work.

The paper is organized as follows. In Section 1, we describe the concept of horizontal approximation of a second order geometric ODE which is induced by a choice of an Ehresmann connection in the tangent bundle of the underlying manifold. We also discuss fiberwise and "special" coordinate systems on the tangent bundle, which naturally define such approximations and will arise in latter sections. In Section 2, we summarize the mathematical theory of two-field cosmological models in the absence of perturbations and give a geometric treatment of local cosmological observables and their dynamical reduction. Section 3 studies a certain local frame of  $T\mathcal{M}$  which is naturally defined by a scalar triple. Section 4 gives a geometric formulation of the fundamental observables of a scalar triple, which are dynamical reductions of certain natural observables used in the physics literature. Section

5 discusses natural fiberwise coordinates on  $T\mathcal{M}$  which are induced by the fundamental observables. In Section 6, we discuss local coordinate systems of  $T\mathcal{M}$ induced by natural fiberwise coordinates and compute the semispray of the cosmological equation in those coordinate systems. Section 7 discusses constant roll (and in particular slow roll and ultra slow roll) manifolds while Section 8 discusses the quasi-conservative and strongly dissipative regimes, giving some insight into the meaning of the conservative function introduced in Section 4. In Section 9, we illustrate the general ideas of Section 1 by discussing the horizontal approximation in adapted coordinates. Appendix A recalls some geometric notions associated to curves in oriented Riemann surfaces and introduces the universal Frenet frame which plays a crucial role in Section 4. Appendix B gives the relation between various quantities used in this paper and certain quantities considered in the physics literature.

0.1. Notations and conventions. All manifolds considered in this paper are smooth and paracompact. For any manifold  $\mathcal{M}$ , we let  $\dot{T}\mathcal{M}$  denote the slit tangent bundle of  $\mathcal{M}$ , which is defined as the complement of the image of the zero section in  $T\mathcal{M}$ . We denote by  $\mathbb{R}_{\mathcal{M}}$  the trivial real line bundle on  $\mathcal{M}$ . For any smooth vector bundle E over  $\mathcal{M}$  and any  $k \in \mathbb{Z}_{\geq 0} \sqcup \{\infty\}$ , we denote by  $\mathcal{C}^k(E)$  the space of those sections of E over  $\mathcal{M}$  which are of class  $\mathcal{C}^k$ . For ease of notation, we let  $\Gamma(E) \stackrel{\text{def.}}{=} \mathcal{C}^{\infty}(E)$  denote the space of smooth sections of E. We denote by  $\mathfrak{X}(\mathcal{M}) \stackrel{\text{def.}}{=} \Gamma(T\mathcal{M})$  the space of smooth vector fields defined on  $\mathcal{M}$ . We denote by  $\pi: T\mathcal{M} \to \mathcal{M}$  the bundle projection of  $T\mathcal{M}$  and by  $F \stackrel{\text{def.}}{=} \pi^*(T\mathcal{M})$  the tautological vector bundle over  $T\mathcal{M}$  (sometimes called the "Finsler bundle" of  $\mathcal{M}$ , see [24]). Let:

$$C(\mathcal{M}) \stackrel{\text{def.}}{=} \sqcup_{I \in \text{Int}} \mathcal{C}^{\infty}(I, \mathcal{M})$$

be the set of all smooth curves in  $\mathcal{M}$ , where Int is the set of all nondegenerate intervals on the real axis and let  $C(T\mathcal{M})$  be the set of all smooth curves in  $T\mathcal{M}$ . The *canonical lift* of curves from  $\mathcal{M}$  to  $T\mathcal{M}$  is the map  $c : C(\mathcal{M}) \to C(T\mathcal{M})$  defined through:

$$c(\varphi) \stackrel{\text{def.}}{=} \dot{\varphi} \ \forall \varphi \in C(\mathcal{M}) \ ,$$

while the *canonical projection* of curves from  $T\mathcal{M}$  to  $\mathcal{M}$  is the map  $p: C(T\mathcal{M}) \to C(\mathcal{M})$  defined through:

$$p(\gamma) \stackrel{\text{def.}}{=} \pi \circ \gamma \ \forall \gamma \in C(T\mathcal{M}) \ .$$

We have  $p \circ c = \operatorname{id}_{C(\mathcal{M})}$ , which means that c is a section of p. We denote by  $C_0(T\mathcal{M})$ the image of the map c. The set  $C_0(T\mathcal{M})$  consists of those curves  $\gamma : I \to T\mathcal{M}$ which have the property that  $\dot{\gamma}(t) \in TT\mathcal{M}$  is a second order tangent vector for all  $t \in I$  (i.e. an element of  $TT\mathcal{M}$  which is invariant under the flip involution of  $TT\mathcal{M}$ ). Then the maps p and c restrict to mutually-inverse bijections between  $C(\mathcal{M})$  and  $C_0(T\mathcal{M})$ .

For any Riemannian manifold  $(\mathcal{M}, \mathcal{G})$ , we denote  $\mathcal{N} : T\mathcal{M} \to \mathbb{R}_{\geq 0}$  the norm function induced by  $\mathcal{G}$  on  $T\mathcal{M}$ :

$$\mathcal{N}(u) \stackrel{\text{def.}}{=} ||u|| \ \forall u \in T\mathcal{M} \ . \tag{1}$$

**Definition 0.1.** A curve  $\varphi : I \to N$  (where N is any manifold) is called pointed if  $0 \in \mathring{I}$ , where  $\mathring{I}$  is the interior of the interval I.

1. HORIZONTAL APPROXIMATIONS FOR SECOND ORDER GEOMETRIC ODES

1.1. Approximations of a geometric dynamical system induced by complementary distributions. Let N be a manifold and  $S \in \mathcal{X}(N)$  a vector field defined on N. We view (N, S) as an autonomous dynamical system whose flow curves  $\gamma: I \to N$  are the solutions of the integral curve equation of S:

$$\dot{\gamma}(t) = S(\gamma(t)) \ \forall t \in I$$
 .

Two Frobenius distributions  $V, W \in TN$  are called *complementary* if  $TN = V \oplus W$ . We think of V as being "vertical" and of W as being "horizontal" and use notation justified by this for reasons which will become clear in the next subsection. Given a pair of complementary distributions (V, W) on N, any vector field  $X \in \mathcal{X}(N)$ decomposes uniquely as  $X = X_V \oplus X_H$  with  $X_V \in \Gamma(V)$  and  $X_H \in \Gamma(W)$ . This defines projectors  $P_V \in \text{Hom}(TN, V)$  and  $P_H \in \text{Hom}(TN, W)$  such that:

$$P_V X = X_V$$
,  $P_H X = X_H$ .

We will use terminology which doesn't refer to V explicitly, for reasons which will become clear in our discussion of second order geometric ODEs below.

**Definition 1.1.** The *W*-flow curves of *S* are the integral curves of the vector field  $S_H$ , i.e. the solutions  $\lambda : I \to N$  of the equation:

$$\lambda(t) = S_H(\lambda(t)) \ \forall t \in I \ .$$

In the situation above, the *W*-approximation of the dynamical system (N, S) consists of replacing it with the dynamical system  $(N, S_H)$ , which we call the *W*-approximant of (N, S). Accordingly, one approximates the integral curves  $\gamma$  of S by the integral curves  $\lambda$  of  $S_H$ . The approximation can be accurate only for those integral curves  $\gamma$  of S which have the property that  $S_V(\gamma(t))$  is "small" for all  $t \in I$  with respect to some appropriate notion of smallness for sections of W; this requires the choice of an appropriate topology on  $\Gamma(W)$  (for example, the Whitney topology or the distance topology induced by a metric on TN). Intuitively, the approximation is good if  $\gamma$  "stays close" to the zero locus of  $S_V$  in N. One can of course also consider the "complementary" approximation in which one replaces S by  $S_V$ .

**Definition 1.2.** The *W*-mean field locus  $\mathcal{Z}_W$  is the zero set of  $S_V$ :

$$\mathcal{Z}_W \stackrel{\text{def.}}{=} \{ n \in N \mid S_V(n) = 0 \} .$$

Notice that  $\mathcal{Z}_W$  is a manifold if  $S_V$  is transverse to the zero section of V. Let  $n_0$  be a point of  $Z_W$  and  $\gamma : I \to N$  be a pointed<sup>2</sup> integral curve of S such that  $\gamma(0) = n_0$ . Further restricting I if necessary, let  $\lambda : I \to N$  be an integral curve of  $S_H$  such that  $\lambda(0) = n_0$ ; of course,  $\lambda$  and  $\gamma$  are uniquely determined by  $n_0$  and the interval I (which need not be the maximal interval of definition of any of these two integral curves). Then the approximation  $\gamma(t) \approx \lambda(t)$  is exact at t = 0 and remains accurate for  $t \in (-\epsilon, \epsilon) \subset I$  for some sufficiently small  $\epsilon > 0$ .

An important special case arises if the distribution W is Frobenius integrable. In this case, we have  $W = T\mathcal{F}$ , where  $\mathcal{F}$  is the foliation of N which integrates W. Since  $S_H \in \Gamma(W)$  is tangent to the leaves of  $\mathcal{F}$ , each integral curve  $\lambda$  of  $S_H$  lies on a leaf  $\mathcal{L}$  of this foliation. Let dim N = d and dim  $\mathcal{L} = \operatorname{rk} W = r$ . If  $(U, (x^i, y^j)_{i=1,\dots,r,j=1,\dots,d-r})$  is a foliated chart of  $\mathcal{F}$  with  $x^i$  and  $y^j$  corresponding respectively to the leaf and transverse directions, then the plaques of  $\mathcal{F}$  in this chart are locally described the condition  $y^i = \operatorname{constant}$  and the integral curve equation of  $S_H$  amounts to the system of ODEs:

$$\dot{x}^{i}(t) = S^{i}_{H}(x(t), y(t)) , \ \dot{y}^{j}(t) = 0 \ (i = 1, \dots, r \ , \ j = 1, \dots d - r) \ ,$$

where  $S_H = S_H^i \frac{\partial}{\partial x^i}$ . In particular, the "vertical speeds"  $\dot{y}^j$  are neglected in the W-approximation. This parallels the separation into "slow" and "fast variables"

<sup>&</sup>lt;sup>2</sup>See Definition 0.1.

(where the time variation of the "slow variables" is neglected) which is common in various physics approximations such as the Born-Oppenheimer approximation for the multiparticle Schrödinger equation.

1.2. Formulation for second order geometric ODEs. Let  $\mathcal{M}$  be a connected manifold and  $S \in \mathcal{X}(T\mathcal{M})$  be a second order vector field [24] (a.k.a. "semispray") defined on the total space of the tangent bundle  $\pi : T\mathcal{M} \to \mathcal{M}$ ; this means that S is invariant under the "flip" involution of the double tangent bundle  $TT\mathcal{M}$ . It is well-known (see loc. cit.) that such a vector field describes the first order formulation of an autonomous geometric second order ODE defined on  $\mathcal{M}$  whose second order term separates. In any local coordinate system  $(x^i)$  on  $\mathcal{M}$ , this corresponds to a system of n second order ODEs of the form:

$$\ddot{x}^i = f^i(x, \dot{x})$$

The adjective "geometric" means that this second order system is invariant under coordinate transformations on  $\mathcal{M}$ ; equivalently, it means that it can be formulated as the condition  $j^2(\varphi) = \mathfrak{s}(j^1(\varphi))$  for curves  $\varphi : I \to \mathcal{M}$ , where  $j^1(\varphi) = \dot{\varphi} : I \to J^1(\mathcal{M}) = T\mathcal{M}$  and  $j^2(\varphi) : I \to J^2(\mathcal{M})$  are the first and second jet prolongations of  $\varphi$  and  $\mathfrak{s}$  is a section of the jet bundle fibration  $\pi_{21} : J^2(\mathcal{M}) \to J^1(\mathcal{M})$ . Here  $J^k(\mathcal{M}) \to \mathcal{M}$  denotes the k-th jet bundle of curves in  $\mathcal{M}$  (see Subsection 2.3).

By definition, a flow curve of S is an integral curve  $\gamma : I \to T\mathcal{M}$  of S while a solution curve of S is a solution  $\varphi : I \to \mathcal{M}$  of the corresponding second order geometric ODE defined on  $\mathcal{M}$ . Let  $C_s(\mathcal{M}) \subset C(\mathcal{M})$  be the set of solution curves and  $C_f(T\mathcal{M}) \subset C(T\mathcal{M})$  be the set of flow curves, where C(N) denotes the set of smooth curves in a manifold N. Since S is a semispray, the maps  $c : C(\mathcal{M}) \to C(T\mathcal{M})$  and  $p : C(T\mathcal{M}) \to C(\mathcal{M})$  defined through  $c(\varphi) := \dot{\varphi}$  and  $p(\gamma) := \pi \circ \gamma$ (see Subsection 0.1) restrict to mutually-inverse bijections between  $C_s(\mathcal{M})$  and  $C_f(T\mathcal{M})$ .

Let  $\pi : T\mathcal{M} \to \mathcal{M}$  be the projection of  $T\mathcal{M}, V := \ker(\mathrm{d}\pi) \subset TT\mathcal{M}$  be the vertical distribution of  $T\mathcal{M}$  and W be a complete Ehresmann connection on  $T\mathcal{M}$ , i.e. a distribution  $W \subset TT\mathcal{M}$  on  $T\mathcal{M}$  which is complementary to V in  $TT\mathcal{M}$ :

$$TT\mathcal{M} = V \oplus W$$

and which defines a horizontal lift of any curve in  $\mathcal{M}$  through any point of  $T\mathcal{M}$  lying above that curve. We denote by  $X_V, X_H$  the vertical and horizontal projections of any vector field field  $X \in \Gamma(TT\mathcal{M})$  defined on  $T\mathcal{M}$ . For any pointed curve  $\varphi: I \to \mathcal{M}$  and any  $u \in T\mathcal{M}$  with  $\pi(u) = \varphi(0)$ , let  $\varphi_u: I \to T\mathcal{M}$  be the horizontal lift of  $\varphi$  through u at t = 0, which is uniquely determined by the conditions:

$$\dot{\varphi}_u(t) \in W_{\varphi_u(t)} \ \forall t \in I \ , \ \varphi_u(0) = u \text{ and } \pi \circ \varphi_u = \varphi \ .$$

1.3. The mean field locus defined by W. The mean field locus  $\mathcal{Z}_W$  of S relative to the Ehresmann connection W is the W-mean field locus of S defined by the complementary distributions V and W on  $T\mathcal{M}$ , i.e. the zero set of  $S_V$  in  $T\mathcal{M}$ :

$$\mathcal{Z}_W \stackrel{\text{def.}}{=} \{ u \in T\mathcal{M} \mid S_V(u) = 0 \} \subset T\mathcal{M} .$$

**Definition 1.3.** We say that S is regular with respect to W if  $S_V$  is transverse to the zero section of the vertical bundle  $V \to T\mathcal{M}$ .

When S is regular with respect to W, the mean field locus  $\mathcal{Z}_W$  is a submanifold of  $T\mathcal{M}$  of dimension equal to that of  $\mathcal{M}$ . In this case,  $\mathcal{Z}_W$  is called the *mean field manifold* of S relative to W. In this case, the implicit function theorem implies that  $\mathcal{Z}_W$  is locally a multisection of the fiber bundle  $\pi : T\mathcal{M} \to \mathcal{M}$  (because the restriction of  $S_V$  to each fiber F of  $T\mathcal{M}$  is a vector field tangent to F which is transverse to the zero section of TF and hence its vanishing locus is a discrete set of points). In particular,  $T\mathcal{Z}_W$  is complementary to the distribution  $V|_{\mathcal{Z}_W}$  inside  $TT\mathcal{M}|_{\mathcal{Z}_W}$ . We have  $S(u) = S_H(u) \in W$  for all  $u \in \mathcal{Z}_W$ . From now on, we assume that S is regular relative to W.

Remark 1.4. Notice that  $S|_{\mathcal{Z}_W} = S_H|_{\mathcal{Z}_W}$  need not be tangent to  $\mathcal{Z}_W$  and hence the integral curves of  $S_H$  starting from a point of the mean field manifold need not be contained in that manifold.

### 1.4. The horizontal approximation defined by W.

**Definition 1.5.** A curve  $\lambda : I \to T\mathcal{M}$  is called a *phase space W*-curve of *S* if it is an integral curve of the vector field  $S_H$ , i.e. if it satisfies:

$$\lambda(t) = S_H(\lambda(t)) \ \forall t \in I \ . \tag{2}$$

Notice that any phase space W-curve  $\lambda : I \to T\mathcal{M}$  is horizontal. Let  $C_H(T\mathcal{M})$  be the set of curves in  $T\mathcal{M}$  which are horizontal with respect to W and  $C_W(T\mathcal{M}) \subset C_H(T\mathcal{M})$  be the subset of phase space W-curves. Recall that  $C_f(T\mathcal{M})$  denotes the set of integral curves of S.

**Definition 1.6.** A curve  $\psi : I \to \mathcal{M}$  is called a *W*-curve of *S* if  $\psi = \pi \circ \lambda$  for some phase space *W*-curve  $\lambda : I \to T\mathcal{M}$  of *S*, i.e. if there exists a horizontal lift of  $\psi$  which is a phase space *W*-curve.

Let  $C_W(\mathcal{M})$  be the set of *W*-curves of *S* and recall that  $C_s(\mathcal{M})$  denotes the set of solution curves of the second order geometric ODE defined by *S*. Since a horizontal lift  $\lambda$  of  $\psi$  is a horizontal curve which satisfies  $\pi \circ \lambda = \psi$ , we have  $C_W(\mathcal{M}) = p(C_W(T\mathcal{M}))$ , where  $p \stackrel{\text{def.}}{=} \pi \circ (-) : C(T\mathcal{M}) \to C(\mathcal{M})$ . Since  $\pi \circ \lambda = \psi$ , we have  $(d\pi) \circ \dot{\lambda} = \dot{\psi}$ . When  $\lambda$  is a phase space *W*-curve, this gives  $(d\pi) \circ S \circ \lambda = \dot{\psi}$ , where we used the relation  $\pi_*(S_H) = \pi_*(S)$ .

Since the horizontal lift a pointed curve  $\psi$  at t = 0 is uniquely determined by its value  $u \in T_{\psi(0)}$  at zero time, this gives a first order ODE for  $\psi$  parameterized by u which characterizes W-curves. More precisely, a pointed curve  $\psi : I \to \mathcal{M}$  with  $\psi(0) = m \in \mathcal{M}$  is a W-curve iff there exists  $u \in T_m \mathcal{M}$  such that the horizontal lift  $\psi_u : I \to T\mathcal{M}$  of  $\psi$  through u at t = 0 is a phase space W-curve. Equation (2) for  $\psi_u$  takes the form:

$$\boldsymbol{\psi}_u(t) = S_H(\boldsymbol{\psi}_u(t)) \; \forall t \in I \tag{3}$$

and can be viewed as a first order geometric ODE for  $\psi$  with initial condition  $\psi(0) = m$  and parameter  $u \in T_m \mathcal{M}$ . This ODE has a unique maximal solution for each parameter  $u \in T_{\psi(0)} \mathcal{M}$  and  $\psi$  can be recovered by projecting that solution to  $\mathcal{M}$ . Let  $I_u$  be the interval of definition of the maximal solution of (3) at a fixed value of u. Then the maximal solution of (3) for parameter u projects to a W-curve  $\psi^{(u)} : I_u \to \mathcal{M}$  which satisfies  $\psi^{(u)}(0) = m$ . This gives a family of W-curves passing through each point m of  $\mathcal{M}$  and parameterized by the vectors tangent to  $\mathcal{M}$  at that point. Notice that this parameterization need not be single-valued. A specific W-curve  $\psi^{(u)}$  which passes through m at t = 0 can be specified by selecting a tangent vector  $u \in T_m \mathcal{M}$ . This determines the speed of  $\psi^{(u)}$  at t = 0 through the relation:

$$\dot{\psi}^{(u)}(0) = (\mathrm{d}\pi)(S(u))$$

Remark 1.7. Two phase space W-curves  $\lambda_1, \lambda_2 : I \to T\mathcal{M}$  determine the same W-curve  $\psi : I \to \mathcal{M}$  iff both of them project to  $\psi$ , which requires  $(d\pi) \circ S \circ \lambda_1 = (d\pi) \circ S \circ \lambda_2 = \dot{\psi}$ .

The horizontal approximation defined by the Ehresmann connection W consists of approximating integral curves of S by phase space W-curves of S, i.e. approximating the flow of S on  $T\mathcal{M}$  by the flow of  $S_H$ . As explained below, this amounts to approximating solution curves  $\varphi$  of the second order geometric ODE defined by S on  $\mathcal{M}$  by W-curves of S whose speed approximates the projection of the acceleration of  $\varphi$ . The approximation is accurate for those integral curves  $\gamma : I \to T\mathcal{M}$  of S which are almost horizontal in the sense that  $S_V \circ \gamma$  is close to zero (where "closeness to zero" is measured, for example, by the fiberwise norm induced on  $TT\mathcal{M}$  by a metric on  $T\mathcal{M}$ ).

Let  $\varphi : I \to \mathcal{M}$  be a pointed solution curve such that the vector  $u := \dot{\varphi}(0)$  lies on the mean field manifold  $\mathcal{Z}_W$ . Then  $S(u) \in W$  and the acceleration of  $\varphi$  at t = 0is horizontal:

$$\ddot{\varphi}(0) = S(\dot{\varphi}(0)) = S(u) \in W \text{ in } TT\mathcal{M} .$$

After passing to a subinterval of I if necessary, let  $\psi = \pi \circ \lambda : I \to \mathcal{M}$  be the W-curve which is the projection of the unique phase space W-curve  $\lambda : I \to T\mathcal{M}$  which satisfies  $\lambda(0) = \dot{\varphi}(0) = u$ . The curve  $\lambda$  and the canonical lift  $\gamma = \dot{\varphi}$  of  $\varphi$  satisfy:

$$\lambda(t) = S_H(\lambda(t)) , \ \dot{\gamma}(t) = S(\gamma(t)) \ \forall t \in I$$

with the initial conditions:

$$\gamma(0) = \lambda(0) = u$$
 .

Thus:

$$\psi(0) = \pi(u) = \varphi(0)$$
 and  $\dot{\psi}(0) = (d\pi)(S(u)) = (d\pi)(\ddot{\varphi}(0))$ 

and we have:

$$\dot{\psi}(t) = (\mathrm{d}\pi)(\dot{\lambda}(t)) = (\mathrm{d}\pi)(S(\lambda(t))) \ \forall t \in I \ .$$

The approximation  $\gamma(t) \approx \lambda(t)$  implies  $S(\lambda(t)) \approx S(\gamma(t)) = \dot{\gamma}(t) = \ddot{\varphi}(t)$  and is equivalent with the joint approximation  $\varphi(t) \approx \psi(t) \& \dot{\varphi}(t) \approx \dot{\lambda}(t)$ , which implies  $\varphi(t) \approx \psi(t) \& (d\pi)(\ddot{\varphi}(t)) \approx \dot{\psi}(t)$ . Notice that the "mean field vector"  $u = \dot{\varphi}(0) \in \mathcal{Z}_W$  determines the W-curve  $\psi$  which passes through the point  $\varphi(0)$  at t = 0 and is used to approximate the solution curve  $\varphi$ . Intuitively, the approximation neglects the vertical component of the acceleration  $\ddot{\varphi}$  of  $\varphi$ . This approximation is exact at t = 0 and remains accurate on a small enough neighborhood of the origin inside the interval I. In particular, the phase space W-approximation of flow curves is exact on the mean field manifold for all flow curves which intersect  $\mathcal{Z}_W$ .

**Definition 1.8.** A *mean field* for S relative to W is a locally defined vector field  $X \in \Gamma(TU_0)$  such that:

$$X(m) \in \mathcal{Z}_W \ \forall m \in U_0 \ , \tag{4}$$

where  $U_0$  is an open subset of  $\mathcal{M}$ .

If U is connected, then the image X(U) of a mean field  $X \in \Gamma(U, T\mathcal{M})$  is a subset of a "branch" of the mean field manifold  $\mathcal{Z}_W$  (when the latter is viewed as a multisection of  $\pi$ ). This allows one to give local parameterizations of that branch of the mean field manifold by choosing local coordinates defined on open subsets  $U \subset \mathcal{M}$ . The name "mean field" was chosen by analogy with similar objects which appear in condensed matter physics.

1.5. The mean field approximation defined by W. Let  $S_W \in \Gamma(TW)$  be the projection of  $S|_{\mathcal{Z}_W}$  to the integrable distribution  $T\mathcal{Z}_W$ , where the projection is taken parallel to V.

**Definition 1.9.** A curve  $\rho : I \to \mathcal{Z}_W$  is called a *phase mean field curve* of S relative to W if it is an integral curve of the vector field  $S_W \in \Gamma(TW)$ , i.e. if it satisfies:

$$\dot{\rho}(t) = S_W(\rho(t)) \ \forall t \in I \ . \tag{5}$$

Notice that any phase mean field curve lies on  $\mathcal{Z}_W$ 

The mean field approximation consists of approximating solution curves of the geometric ODE defined by S by phase mean field curves of S relative to W. This is a singular approximation in the sense that it restricts the phase space of the dynamical system from  $T\mathcal{M}$  to  $T\mathcal{Z}_W$ . The approximation is accurate for those solution curves  $\varphi$  whose flow curve  $\gamma = \dot{\varphi}$  lies very close to the mean field manifold  $\mathcal{Z}_W$ , provided that  $S|_{\mathcal{Z}_W}$  is well-approximated by  $S_W$ .

Let  $\varphi: I \to \mathcal{M}$  be a pointed solution curve of S such that the vector  $u := \dot{\varphi}(0)$ lies on the mean field manifold  $\mathcal{Z}_W$ . Let  $\gamma := \dot{\varphi}: I \to T\mathcal{M}$  be the canonical lift of  $\varphi$ . Then:

$$\dot{\gamma}(0) = S(\gamma(0)) = S(u) \in W_u$$

After passing to a subinterval of I if necessary, let  $\rho: I \to \mathcal{Z}_W$  be the unique phase mean field curve which satisfies:

$$\rho(0) = u \; .$$

Then:

$$\dot{\rho}(t) = S_W(\rho(t)) \ \forall t \in I$$

and we have:

$$\dot{\rho}(0) = S_W(u) \; .$$

Let  $\mu \stackrel{\text{def.}}{=} \pi \circ \rho : I \to \mathcal{M}$ . The approximation  $\gamma(t) \approx \rho(t)$  implies  $\varphi(t) \approx \mu(t)$ and  $(d\pi)(S_W(\dot{\varphi}(t))) \approx \dot{\mu}(t)$  and is accurate for t close to zero only if  $\dot{\gamma}(0) \approx \dot{\rho}(0)$ , i.e. only if the horizontal vector S(u) is almost tangent to  $\mathcal{Z}_W$  and hence is well approximated by  $S_W(u)$ . This approximation is much coarser than the horizontal approximation and is not guaranteed to be accurate for any flow curve or nondegenerate time interval.

Remark 1.10. The discussion above can be adapted easily to the restriction of S and W to a non-empty open subset of  $T\mathcal{M}$ . We leave the details of this generalization to the reader.

1.6. Fiberwise coordinates and special coordinate systems on  $T\mathcal{M}$ . Suppose for simplicity that dim  $\mathcal{M} = 2$ , which is the case relevant for the remainder of the paper. Let U be a non-empty open subset of  $T\mathcal{M}$ . Then  $U_0 \stackrel{\text{def.}}{=} \pi(U)$  is an open subset of  $\mathcal{M}$  since  $\pi$  is a submersion and hence an open map.

**Definition 1.11.** A system of fiberwise coordinates defined on U is a (generally non-surjective) submersion  $y = (y^1, y^2) : U \to \mathbb{R}^2$  whose restriction to every non-empty intersection of U with a fiber of  $T\mathcal{M}$  is a system of coordinates on that fiber. That is, the map:

$$y|_{\pi^{-1}(m)\cap U}:\pi^{-1}(m)\cap U\to\mathbb{R}^2$$

is a diffeomorphism to its image  $y(\pi^{-1}(m)\cap U)$  for all  $m\in U_0$  .

A particular case arises when  $y(\pi^{-1}(m) \cap U) = y(U)$  for all  $m \in U$ .

**Proposition 1.12.** Let  $y : U \to \mathbb{R}^2$  be a smooth map. Then the following statements are equivalent:

- (a) y is a fiberwise coordinate system such that  $y(\pi^{-1}(m) \cap U) = y(U)$  for all  $m \in U_0 \stackrel{\text{def.}}{=} \pi(U)$ .
- (b) The restriction  $\pi_U \stackrel{\text{def.}}{=} \pi|_U : U \to U_0$  is a topologically trivial fiber bundle with typical fiber y(U) and the map  $(\pi_U, y) : U \to U_0 \times y(U)$  is a trivialization of this fiber bundle, where  $U_0 \times y(U)$  is viewed as trivial fiber bundle over  $U_0$  using projection on the first factor.
- In this case,  $\pi_U$  is fiber sub-bundle of the vector bundle  $T\mathcal{U}_0$ .

8

*Proof.* To see that (a) implies (b), notice that all fibers of the surjective submersion  $\pi_U$  are diffeomorphic with y(U) and  $(\pi_U, y)$  is an isomorphism of fiber bundles to the trivial y(U)-bundle over  $U_0$ . The converse implication follows from the definition of trivializations of fiber bundles.

An interesting class of examples arises from local frames of  $T\mathcal{M}$ .

**Definition 1.13.** Let  $U_0$  be a non-empty open subset of  $\mathcal{M}$ . A surjective system of fiberwise coordinates  $y = (y^1, y^2) : \pi^{-1}(U_0) = TU_0 \to \mathbb{R}^2$  is called *linear* if the restriction of y to every fiber of  $TU_0$  is a linear isomorphism.

**Proposition 1.14.** The following statements are equivalent for a map  $y: TU_0 \rightarrow \mathbb{R}^2$ .

- (a) y is a linear system of fiberwise coordinates.
- (b) There exists a frame  $(e_1, e_2)$  of  $TU_0$  such that:

 $u = y^{1}(u)e_{1}(\pi(u)) + y^{2}(u)e_{2}(\pi(u)) \ \forall u \in TU_{0}$ 

Notice that the frame  $(e_1, e_2)$  need not be holonomic (i.e. it need not be induced by a local coordinate system on  $\mathcal{M}$ ).

*Proof.* Suppose that (a) holds. For any  $m \in U_0$ , the restrictions  $y_m^i : T_m \mathcal{M} \to \mathbb{R}$  give linear coordinates on  $T_m \mathcal{M}$ . If  $e_1(m), e_2(m) \in T_m \mathcal{M}$  are the unique vectors which satisfy  $y(e_1(m)) = (1,0)$  and  $y(e_2(m)) = (0,1)$  then any  $u \in T_m \mathcal{M}$  can be written as:

$$u = y_m^1(u)e_1(m) + y_m^2(u)e_2(m)$$

and hence  $(e_1(m), e_2(m))$  forms a basis of the fiber  $T_m \mathcal{M}$ . This gives a frame  $(e_1, e_2)$  of  $TU_0$  and hence (b) holds. Now suppose that (b) holds. Then we have  $y_m^i(u) = u^i$  for  $m \in U_0$  and  $u \in T_m U_0$ , where  $u = u^1 e_1(m) + u^2 e_2(m)$  is the frame expansion of u. It is clear that  $y = (y^1, y^2)$  is a linear system of fiberwise coordinates.  $\Box$ 

**Definition 1.15.** A coordinate system  $x = (y^1, y^2, \xi^1, \xi^2) : U \to \mathbb{R}^4$  defined on an open subset  $U \subset T\mathcal{M}$  is called *special* if it satisfies the following two conditions: 1.  $(y^1, y^2) : U \to \mathbb{R}^2$  is a system of fiberwise coordinates defined on U.

2. There exists a system of coordinates  $x = (x^1, x^2) : U_0 \stackrel{\text{def.}}{=} \pi(U) \to \mathbb{R}^2$  such that  $\xi^1 = (x^1)^{\text{v}} \stackrel{\text{def.}}{=} x^1 \circ \pi_U$  and  $\xi^2 = (x^2)^{\text{v}} \stackrel{\text{def.}}{=} x^2 \circ \pi_U$ , where  $\pi_U \stackrel{\text{def.}}{=} \pi|_U$ .

Given any system of fiberwise coordinates  $(y^1, y^2)$  on  $U \subset T\mathcal{M}$  and any system of coordinates  $(x^1, x^2)$  on  $U_0 = \pi(U)$ , it is clear that  $(y^1, y^2, \xi^1 \stackrel{\text{def.}}{=} (x^1)^{\text{v}}, \xi^2 \stackrel{\text{def.}}{=} (x^2)^{\text{v}})$  is a special coordinate system on U.

1.7. The horizontal approximation defined by a special system of coordinates. Let  $(y^1, y^2, \xi^1 = (x^1)^{\vee}, \xi^{\vee} = (x^2)^{\vee}) : U \to \mathbb{R}^4$  be a special system of coordinates defined on the open subset  $U \subset T\mathcal{M}$  and let  $U_0 \stackrel{\text{def.}}{=} \pi(U)$ . The rank two integrable distribution W spanned by the vector fields  $\frac{\partial}{\partial \xi^1}$  and  $\frac{\partial}{\partial \xi^2}$  on U is complementary to the vertical distribution V of the submersion  $\pi_U : U \to U_0$  and defines a complete *flat* Ehresmann connection on U. As in the previous subsection, this gives a decomposition  $S = S_V + S_H$  of any semispray S defined on U. Write:

$$S = S^{y^i} \frac{\partial}{\partial y^i} + S^{\xi^i} \frac{\partial}{\partial \xi^i}$$

where we use implicit summation over repeated indices and  $i \in \{1, 2\}$ . Since the vector fields  $\frac{\partial}{\partial u^i}, \frac{\partial}{\partial \xi^i} \in \mathcal{X}(U)$  are respectively vertical and horizontal, we have:

$$S_V = S^{y^i} \frac{\partial}{\partial y^i}$$
 and  $S_H = S^{\xi^i} \frac{\partial}{\partial \xi^i}$ .

To identify the horizontal part of S, let  $\zeta = (\zeta^1, \zeta^2) : TU_0 \to \mathbb{R}^2$  be the linear system of fiberwise coordinates defined by the frame  $\left(\frac{\partial}{\partial x^1}, \frac{\partial}{\partial x^2}\right)$  of  $TU_0$ . This is known as the *tangent chart* of the chart  $(x^1, x^2) : U_0 \to \mathbb{R}^2$  of  $\mathcal{M}$ . The change of coordinates from  $(\zeta, \xi)$  to  $(y, \xi)$  and its inverse have the forms:

$$y^{i} = y^{i}(\zeta, \xi) , \ \xi^{i} = \xi^{i} \text{ and } \zeta^{i} = \zeta^{i}(y, \xi) , \ \xi^{i} = \xi^{i} ,$$
 (6)

where  $\mathbf{y}^i$  and  $\zeta^i$  are functions. In particular, we have  $\frac{\partial \xi^i}{\partial y^j} = 0$  and hence:

$$\frac{\partial}{\partial y^i} = \frac{\partial \zeta^j}{\partial y^i} \frac{\partial}{\partial \zeta^j}$$

Since S is a semispray, its horizontal part is given by the following formula in the tangent chart  $(\zeta, \xi)$  (see [24]):

$$S_H = \zeta^i \frac{\partial}{\partial \xi^i}$$

Thus:

$$S^{\xi^i} = \zeta^i$$

For any curve  $\gamma: I \to U$ , we have:

$$\dot{\gamma}(t) = \dot{y}^{i}(t) \frac{\partial}{\partial y^{i}} \Big|_{\gamma(t)} + \dot{\xi}^{i}(t) \frac{\partial}{\partial \xi^{i}} \Big|_{\gamma(t)} ,$$

where  $y^i(t) \stackrel{\text{def.}}{=} y^i(\gamma(t))$  and  $\xi^i(t) \stackrel{\text{def.}}{=} \xi^i(\gamma(t))$ . Hence:

$$\dot{\gamma}(t)_V = \dot{y}^j(t) \frac{\partial}{\partial y^i} \Big|_{\gamma(t)} , \ \dot{\gamma}(t)_H = \dot{\xi}^i(t) \frac{\partial}{\partial \xi^i} \Big|_{\gamma(t)}$$

and the integral curve equation of S amounts to the following system of ODEs:

$$\dot{y}^{i}(t) = S^{y^{i}}(y(t),\xi(t)) , \ \dot{\xi}^{i}(t) = S^{\xi^{i}}(y(t),\xi(t)) = \zeta^{i}(y(t),\xi(t)) .$$
(7)

The mean field locus  $\mathcal{Z}_W$  of S relative to W has equations:

$$S^{y_1}(y,\xi) = S^{y_2}(y,\xi) = 0$$

As before, we assume that S is regular relative to W, i.e.  $S_V$  is transverse to the zero section of V. This amounts to the condition that the map  $(S^{y^1}, S^{y^2}) : U \to \mathbb{R}^2$  restricts to a submersion on a tubular neighborhood of  $\mathcal{Z}_W$  in U. Writing a vector field  $X \in \mathcal{X}(U_0)$  as  $X = X^i \frac{\partial}{\partial x^i}$  with  $X^i \in \mathcal{C}^{\infty}(U_0)$ , we have  $\xi^i \circ X = x^i$  and  $\zeta^i \circ X = X^i$ . Hence the mean field equation (4) reads:

$$S^{y^{i}}(\mathbf{y}^{1}(X^{1}(x), X^{2}(x), x), \mathbf{y}^{2}(X^{1}(x), X^{2}(x), x), x) = 0 \ (i = 1, 2) ,$$

where  $y^i$  are the functions appearing in the coordinate transformations (6). This system of functional equations locally determines  $X^i$  as functions of x, though the solution need not be globally unique.

Let  $\lambda : I \to U$  be a curve in U. Setting  $y^i(t) \stackrel{\text{def.}}{=} y^i(\lambda(t))$  and  $\xi^i(t) \stackrel{\text{def.}}{=} \xi^i(\lambda(t))$ , the phase space W-curve equation (2) amounts to the system:

$$\dot{y}^i(t)=0 \text{ and } \dot{\xi}^i(t)=S^{\xi^i}(y(t),\xi(t))=\zeta^i(y(t),\xi(t)) \; \forall t\in I$$
 .

Intuitively, the horizontal approximation defined by the special coordinates  $(y, \xi)$  amounts to treating the variables  $y^1$  and  $y^2$  as 'slow' in the sense that  $\dot{y}^1$  and  $\dot{y}^2$  are neglected. This corresponds to neglecting the following combinations:

$$\frac{\partial \mathbf{y}^{i}}{\partial \zeta^{j}}(\dot{x}(t), x(t)) \ddot{x}^{j}(t) + \frac{\partial \mathbf{y}^{i}}{\partial \xi^{j}}(\dot{x}(t), x(t)) \dot{x}^{j}(t) \ (i = 1, 2)$$

of the components of the speed and acceleration of a solution curve  $\varphi$  of the geometric second order ODE  $\ddot{\varphi}(t) = S(\dot{\varphi})(t)$  defined by S on  $U_0$ . Here  $x^i(t) \stackrel{\text{def.}}{=} x^i(\varphi(t))$  and we noticed that  $\xi^i(\dot{\varphi}(t)) = x^i(t)$  and  $\zeta^i(\dot{\varphi}(t)) = \dot{x}^i$ .

#### 2. Two-field cosmological models

By a *two-field cosmological model* we mean a classical cosmological model with two real scalar fields derived from the following Lagrangian on a spacetime with topology  $\mathbb{R}^4$ :

$$\mathcal{L}[g,\varphi] = \left[\frac{M^2}{2}\mathrm{R}(g) - \frac{1}{2}\mathrm{Tr}_g\varphi^*(\mathcal{G}) - \Phi\circ\varphi\right]\mathrm{vol}_g .$$
(8)

Here M is the reduced Planck mass, g is the spacetime metric on  $\mathbb{R}^4$  (taken to be of "mostly plus" signature) and  $\operatorname{vol}_g$  and  $\operatorname{R}(g)$  are the volume form and scalar curvature of g. The scalar fields are described by a smooth map  $\varphi : \mathbb{R}^4 \to \mathcal{M}$ , where  $\mathcal{M}$  is a (generally non-compact) connected, smooth and paracompact surface without boundary which we assume to be oriented and endowed with a smooth Riemannian metric  $\mathcal{G}$ , while  $\Phi : \mathcal{M} \to \mathbb{R}$  is a smooth function which plays the role of potential for the scalar fields. One usually requires that  $\mathcal{G}$  is complete to ensure conservation of energy. For simplicity, we also assume that  $\Phi$  is strictly positive and non-constant on  $\mathcal{M}$ . The triplet  $(\mathcal{M}, \mathcal{G}, \Phi)$  is called the *scalar triple* of the model, while  $(\mathcal{M}, \mathcal{G})$  is called the *scalar manifold*. The model is parameterized by the quadruplet  $\mathfrak{M} \stackrel{\text{def.}}{=} (M_0, \mathcal{M}, \mathcal{G}, \Phi)$ , where:

$$M_0 \stackrel{\text{def.}}{=} M \sqrt{\frac{2}{3}}$$

is the *rescaled Planck mass*. We denote by:

$$\operatorname{Crit} \Phi \stackrel{\text{def.}}{=} \{ m \in \mathcal{M} \mid (\mathrm{d}\Phi)(m) = 0 \} \text{ and } \mathcal{M}_0 \stackrel{\text{def.}}{=} \mathcal{M} \setminus \operatorname{Crit} \Phi$$

the critical and noncritical subsets of  $\mathcal{M}$ . Notice that  $\mathcal{M}_0$  is open in  $\mathcal{M}$ . It is also non-empty since we assume that  $\Phi$  is not constant.

The two-field model parameterized by the quadruplet  $\mathfrak{M}$  is obtained by assuming that g is an FLRW metric with flat spatial section:

$$ds_g^2 = -dt^2 + a(t)^2 \sum_{i=1}^3 dx_i^2$$
(9)

(where a is a smooth and positive function of t) and that  $\varphi$  depends only on the cosmological time  $t \stackrel{\text{def.}}{=} x^0$ . Here  $(x^0, x^1, x^2, x^3)$  are the Cartesian coordinates on  $\mathbb{R}^4$ . Define the Hubble parameter through:

$$H(t) \stackrel{\text{def.}}{=} \frac{\dot{a}(t)}{a(t)} ,$$

where the dot indicates derivation with respect to t.

2.1. The cosmological equation. Let  $\pi : T\mathcal{M} \to \mathcal{M}$  be bundle projection. Let  $\mathcal{H} : T\mathcal{M} \to \mathbb{R}_{>0}$  be the rescaled Hubble function of the scalar triple  $(\mathcal{M}, \mathcal{G}, \Phi)$ , which is defined through:

$$\mathcal{H}(u) \stackrel{\text{def.}}{=} \frac{1}{M_0} \left[ ||u||^2 + 2\Phi(\pi(u)) \right]^{1/2} \quad \forall u \in T\mathcal{M}$$
(10)

and let  $\nabla$  be the Levi-Civita connection of  $(\mathcal{M}, \mathcal{G})$ . When H > 0 (which we assume for the remainder of the paper), the Euler-Lagrange equations of (8) reduce (see [13, 16, 17]) to the *cosmological equation*:

$$\nabla_t \dot{\varphi}(t) + \mathcal{H}(\dot{\varphi}(t))\dot{\varphi}(t) + (\operatorname{grad}\Phi)(\varphi(t)) = 0 , \qquad (11)$$

(where  $\nabla_t \stackrel{\text{def.}}{=} \nabla_{\dot{\varphi}(t)}$ ) together with the condition:

$$H(t) = \frac{1}{3} \mathcal{H}(\dot{\varphi}(t)) .$$
(12)

The smooth solutions  $\varphi : I \to \mathcal{M}$  of (11) (where *I* is a non-degenerate interval) are called *cosmological curves*, while their images in  $\mathcal{M}$  are called *cosmological curves*, while their images in  $\mathcal{M}$  are called *cosmological curve*  $\varphi$ , relation (12) determines the function *a* up to a multiplicative constant.

A cosmological curve  $\varphi : I \to \mathcal{M}$  need not be an immersion. Accordingly, we define the singular and regular parameter sets of  $\varphi$  through:

$$I_{\text{sing}} \stackrel{\text{def.}}{=} \{t \in I \mid \dot{\varphi}(t) = 0\} , \ I_{\text{reg}} \stackrel{\text{def.}}{=} I \setminus I_{\text{sing}} = \{t \in I \mid \dot{\varphi}(t) \neq 0\} .$$

When  $\varphi$  is not constant, it can be shown that  $I_{\text{sing}}$  is a discrete subset of I. The sets of *critical and noncritical times* of  $\varphi$  are defined through:

$$I_{\text{crit}} \stackrel{\text{def.}}{=} \{t \in I \mid \varphi(t) \in \text{Crit}\Phi\} , \ I_{\text{noncrit}} \stackrel{\text{def.}}{=} I \setminus I_{\text{crit}} \stackrel{\text{def.}}{=} \{t \in I \mid \varphi(t) \notin \text{Crit}\Phi\} .$$

The cosmological curve  $\varphi$  is called *nondegenerate* if  $I_{\text{sing}} = \emptyset$  and *noncritical* if  $I_{\text{crit}} = \emptyset$ . Noncriticality means that the orbit of  $\varphi$  does not meet the critical set of  $\Phi$ , i.e.  $\varphi(I)$  is contained in  $\mathcal{M}_0$ . It is easy to see that a cosmological curve  $\varphi : I \to \mathcal{M}$  is constant iff its image coincides with a critical point of  $\Phi$ , which in turn happens iff there exists some  $t \in I_{\text{sing}} \cap I_{\text{crit}}$ . Hence for any non-constant cosmological curve we have:

$$I_{\text{sing}} \cap I_{\text{crit}} = \emptyset$$

2.2. The cosmological dynamical system. The cosmological equation (11) is geometric in the sense that it can be written as the condition  $j^2(\varphi) = \mathfrak{s}(j^1(\varphi))$ , where  $\mathfrak{s}$  is a section of the fiber bundle  $J^2(\mathcal{M}) \xrightarrow{\pi_{21}} J^1(\mathcal{M})$  (see Subsection 2.3). This statement follows from the fact that (11) is expressed in terms of diffeomorphism-covariant operators and from the fact that the double derivative term in (11) separates from the lower order terms in any local system of coordinates.

The geometric character of (11) implies that the cosmological equation is invariant under changes of local coordinates on  $\mathcal{M}$  and hence its first order form defines a geometric dynamical system on the total space of  $T\mathcal{M}$  in the sense of [19, 20], which we call the *cosmological dynamical system* defined by the scalar triple  $(\mathcal{M}, \mathcal{G}, \Phi)$ . As explained for example in [24], the defining vector field  $S \in \mathcal{X}(T\mathcal{M})$  of this dynamical system is a second-order vector field which we call the *cosmological semispray*. We have (see [21]):

$$S = \mathbb{S} - Q$$

where S is the geodesic spray of the Riemannian manifold  $(\mathcal{M}, \mathcal{G})$  and  $Q \in \mathcal{X}(T\mathcal{M})$  is the cosmological correction, which is the vertical vector field:

$$Q = \mathcal{H}C + (\operatorname{grad}\Phi)^v$$
.

Here  $C \in \mathcal{X}(T\mathcal{M})$  is the Euler vector field of  $T\mathcal{M}$  and  $(\operatorname{grad}\Phi)^v \in T\mathcal{M}$  is the vertical lift of the vector field  $\operatorname{grad}\Phi \in \mathcal{X}(\mathcal{M})$  to a vector field defined on  $T\mathcal{M}$ . The cosmological equation (11) is equivalent with the integral curve equation of S in the following sense (see [21]):

**Proposition 2.1.** Let  $\varphi : I \to \mathcal{M}$  be a curve in  $\mathcal{M}$ . Then the following statements are equivalent:

- (a)  $\varphi$  satisfies the cosmological equation (11)
- (b) The canonical lift  $\gamma \stackrel{\text{def.}}{=} \dot{\varphi} : I \to T\mathcal{M}$  of  $\varphi$  is an integral curve of the vector field S, i.e. it satisfies the equation:

$$\dot{\gamma}(t) = S(\gamma(t)) \ \forall t \in I \ . \tag{13}$$

The flow of S on the total space of  $T\mathcal{M}$  is called the *cosmological flow* of the model. Following dynamical system terminology, we will refer to  $T\mathcal{M}$  as the *phase space* of the cosmological dynamical system. The integral curves of S are called *cosmological*  flow curves while their images in  $T\mathcal{M}$  are called cosmological flow orbits. It can be shown that any cosmological flow orbit is aperiodic and without self-intersections. This follows from the fact that the cosmological dynamical system is dissipative (see Proposition 2.12 below).

2.3. Local cosmological observables. For any  $k \in \mathbb{Z}_{\geq 0}$ , let  $E^k \stackrel{\text{def.}}{=} J^k(\mathcal{M})$  be the k-th jet bundle of curves in  $\mathcal{M}$ , whose projection to  $\mathcal{M}$  we denote by  $\pi_k$ . By definition, the fiber of  $E^k$  at  $m \in \mathcal{M}$  is the k-th jet space  $E_m^k = J_m^k(\mathcal{M})$  of curves in  $\mathcal{M}$  which pass through the point m at time t = 0. Let  $F^k \stackrel{\text{def.}}{=} \pi_k^*(T\mathcal{M})$  be the k-th Finsler bundle of  $\mathcal{M}$ , whose projection to  $E^k$  we denote by  $p_k$ . We have  $E^0 = \mathcal{M}, E^1 = T\mathcal{M}$  and  $F^0 = T\mathcal{M}, F^1 = F$ , where  $F = \pi^*(T\mathcal{M})$  is the ordinary Finsler bundle of  $\mathcal{M}$ , i.e. the tautological bundle of  $T\mathcal{M}$ . For any  $k > l \ge 0$ , let  $\pi_{kl} : E^k \to E^l$  be the natural projection, which is a fiber bundle. We have  $\pi_{k,0} = \pi_k$ and  $\pi_{l,m} \circ \pi_{k,l} = \pi_{l,m}$  for k > l > m. Notice that  $\pi_{k,k-1} : E^k \to E^{k-1}$  is an affine bundle of rank 2 which is modeled on the vector bundle  $F^{k-1}$  (see [26]). For any smooth curve  $\varphi : I \to \mathcal{M}$ , let  $j^k(\varphi) : I \to J^k(\mathcal{M})$  denote the k-th jet prolongation of  $\varphi$ . We have  $j^0(\varphi) = \varphi$  and  $j^1(\varphi) = \dot{\varphi}$ .

*Remark* 2.2. The product  $\mathbb{R} \times E^k \to \mathbb{R}$  identifies with the jet bundle of sections of the trivial bundle  $\mathbb{R} \times \mathcal{M} \to \mathbb{R}$ , where the projection to  $\mathbb{R}$  is the first projection of the Cartesian product.

Let  $\mathcal{E}$  be a vector bundle over  $\mathcal{M}$  and set  $\mathcal{E}^k \stackrel{\text{def.}}{=} \pi_k^*(\mathcal{E})$  for all  $k \in \mathbb{Z}_{>0}$ .

**Definition 2.3.** A (continuous and off-shell)  $\mathcal{E}$ -valued local cosmological observable of order  $k \in \mathbb{Z}_{\geq 0}$  is a continuous section  $s \in \mathcal{C}^0(U, \mathcal{E}^k)$ , where U is a non-empty open subset of  $E^k = J^k(\mathcal{M})$ . A local cosmological observable of order one is called *basic*; in this case, s is a section of  $\pi^*(\mathcal{E})$  defined on an open subset of  $J^1(\mathcal{M}) = T\mathcal{M}$ .

Local cosmological observables valued in the trivial real line bundle  $\mathbb{R}_{\mathcal{M}}$  of  $\mathcal{M}$  are called *scalar local observables*, while  $T\mathcal{M}$ -valued local observables are called *vector local observables*. In the first case,  $\mathcal{E}^k$  is isomorphic with the trivial line bundle  $\mathbb{R}_{E^k}$  over  $E^k$ , so a scalar local observable is a function  $f: U \to \mathbb{R}$ , where U is an open subset of  $E^k$ . In the second case, we have  $\mathcal{E}^k = F^k$ , so a local vector observable is a section  $s \in \Gamma(U, F^k)$ , where U is an open subset of  $E^k$ . Notice that  $\mathcal{E}$ -valued local observables of order zero are local sections of the vector bundle  $\mathcal{E}$ .

Example 2.4. A simple example of scalar basic cosmological observable is the norm function  $|| || : T\mathcal{M} \to \mathbb{R}_{\geq 0}$  of  $(\mathcal{M}, \mathcal{G})$ , which is continuous everywhere but smooth only on the slit tangent bundle  $\dot{T}\mathcal{M}$ . Another simple example of such is the rescaled Hubble function  $\mathcal{H} = \frac{1}{M_0} \sqrt{\mathcal{N}^2 + 2\Phi \circ \pi}$ , which is smooth on  $T\mathcal{M}$ . Here  $\mathcal{N} : T\mathcal{M} \to \mathbb{R}_{\geq 0}$  is the norm function induced by  $\mathcal{G}$  (see Subsection 0.1).

**Definition 2.5.** Let  $U \subset T\mathcal{M}$  be a non-empty open set and  $U_0 \stackrel{\text{def.}}{=} \pi(U)$ . A basic scalar local observable  $f : U \to \mathbb{R}$  is called *norm reducible* if there exists a map  $g: U_0 \times \mathbb{R}_{\geq 0} \to \mathbb{R}$  such that:  $f = g \circ (\pi \times \mathcal{N})$ , i.e.:

$$f(u) = g(\pi(u), ||u||) \ \forall u \in U .$$

In this case, we have:

$$f(u) = g_m(||u||) \ \forall m \in U_0 \ \forall u \in U \cap T_m \mathcal{M} ,$$

where  $g_m : \mathbb{R}_{\geq 0} \to \mathbb{R}$  is the *m*-section of *g*:

$$g_m(u) \stackrel{\text{def.}}{=} g(m, ||u||) \ \forall u \in U \cap T_m \mathcal{M} \ \forall m \in U_0 \ .$$

*Example 2.6.* The rescaled Hubble function  $\mathcal{H}$  is norm reducible.

**Definition 2.7.** Given an  $\mathcal{E}$ -valued local cosmological observable  $s \in \mathcal{C}^0(U, \mathcal{E}^k)$ with  $U \subset E^k$  and a curve  $\varphi : I \to \mathcal{M}$  such that  $j^k(\varphi)(I) \subset U$ , the *evaluation* of salong  $\varphi$  is defined though:

$$s_{\varphi} \stackrel{\text{def.}}{=} j^k(\varphi)^*(s) \in \mathcal{C}^0(I, \varphi^*(\mathcal{E}))$$

In the definition above, we used the isomorphism  $(j^k(\varphi))^*(\mathcal{E}^k) = (j^k(\varphi))^*(\pi_k^*(\mathcal{E})) \simeq \varphi^*(\mathcal{E})$ , which follows from the relation  $\pi_k \circ j^k(\varphi) = \varphi$ . Notice that  $s_{\varphi}$  can be viewed as a map  $s_{\varphi} : I \to \mathcal{E}$  which satisfies  $s_{\varphi}(t) \in \mathcal{E}_{\varphi(t)}$  for all  $t \in I$ .

2.3.1. The cosmological section, cosmological shell and dynamical reduction of local observables. The cosmological equation (11) defines the cosmological section  $\mathfrak{s}$ :  $E^1 \to E^2$  of the fiber bundle  $\pi_{21}: E^2 \to E^1$ , which has the following expression in the canonical local coordinates  $(x_0^i = x^i, x_1^i = \dot{x}^i, x_2^i = \ddot{x}^i)_{i=1,...,d}$  on  $E^2$  induced by local coordinates  $x = (x^i)_{i=1,2}$  on  $\mathcal{M}$  (here the dots denote formal derivatives).

$$\mathfrak{s}^{x_{2}^{i}}(x,x_{1}) = -\Gamma_{jk}^{i}(x)x_{1}^{j}x_{1}^{k} - \frac{1}{M_{0}} \left[\mathcal{G}_{kl}(x)x_{1}^{k}x_{1}^{l} + 2\Phi(x)\right]^{1/2}x_{1}^{i} - \mathcal{G}^{ij}(x)(\partial_{j}\Phi)(x)$$

$$\mathfrak{s}^{x_{1}^{i}}(x,x_{1}) = x_{1}^{i}$$

$$\mathfrak{s}^{x_{0}^{i}}(x,x_{1}) = x_{1}^{i}$$
(14)

where  $\mathfrak{s}^{x_k^i} \stackrel{\text{def.}}{=} x_k^i \circ \mathfrak{s}$  for k = 0, 1, 2 and i = 1, 2. Here  $\Gamma_{jk}^i$  are the Christoffel symbols of  $\mathcal{G}$  in the local coordinates  $x^i$  on  $\mathcal{M}$ . The image of this section is the closed submanifold  $\mathfrak{S}$  (called the *cosmological shell*) of the total space of  $E^2 = J^2(\mathcal{M})$ defined by the cosmological equation. The cosmological shell has equations:

$$x_2^i = \hat{\mathfrak{s}}_1^i(x, x_1) , \qquad (15)$$

where we set  $\hat{\mathfrak{s}}_1^i \stackrel{\text{def.}}{=} \mathfrak{s}_2^{x_2^i}$ . Formal differentiation of (15) with respect to time gives:

 $x_3^i = (\partial_{x_0^j} \hat{\mathfrak{s}}_1^i)(x_0, x_1) x_1^j + (\partial_{x_1^j} \hat{\mathfrak{s}}_1^i)(x_0, x_1) \hat{\mathfrak{s}}_1^j(x_0, x_1) \stackrel{\text{def.}}{=} \hat{\mathfrak{s}}_2^i(x_0, x_1)$ 

and generally:

$$x_{k+1}^{i} = \hat{\mathfrak{s}}_{k}^{i}(x_{0}, x_{1}) \ \forall k \ge 1 ,$$
(16)

for certain maps  $\hat{\mathfrak{s}}_k^i: E^1 \to \mathbb{R}$ . The map  $\mathfrak{s}_k: E_1 \to E^{k+1}$  with components:

$$\begin{split} & \mathfrak{s}_k^{x_0^i}(x_0, x_1) = x_0^i \ , \ \mathfrak{s}_k^{x_1^i}(x_0, x_1) = x_1^i \\ & \mathfrak{s}_k^{x_{l+1}^i}(x_0, x_1) = \hat{\mathfrak{s}}_l^i(x_0, x_1) \ \forall l \in \{1, \dots, k\} \end{split}$$

defines a section of the fiber bundle  $E^{k+1} \to E^1$  and we have  $\mathfrak{s}_1 = \mathfrak{s}$ . Notice that  $\mathfrak{s}_k$  depends on the k-th jets of  $\mathcal{G}$  and  $\Phi$ .

For any  $k \ge 1$ , the image of  $\mathfrak{s}_k$  is a closed submanifold  $\mathfrak{S}_k$  of  $E^{k+1}$  which gives the (k-1)-th prolongation of the cosmological shell and we have  $\mathfrak{S}_1 = \mathfrak{S}$ . The equations:

$$x_l^i = \mathfrak{s}_k^{x_l}(x_0, x_1) \ (l = 0, \dots, k+1, i = 1, 2)$$

give the (k-1)-th prolongation of the cosmological equation. A curve  $\varphi : I \to \mathcal{M}$ satisfies the cosmological equation iff  $j^2(\varphi)(I) \subset \mathfrak{S}$ , which amounts to the condition  $j^2(\varphi)(t) = \mathfrak{s}(j^1(\varphi)(t))$  for all  $i \in I$ . In this case, we also have  $j^{k+1}(\varphi)(I) \subset \mathfrak{S}_k$  for all  $k \geq 1$ , which amounts to

$$j^{k+1}(\varphi)(t) = \mathfrak{s}_k(j^1(\varphi)(t)) \ \forall t \in I \ .$$
(17)

**Definition 2.8.** The dynamical reduction (or on-shell reduction) of an  $\mathcal{E}$ -valued cosmological observable  $s \in \mathcal{C}^0(U, \mathcal{E}^k)$  of order  $k \geq 2$  (where U is an open subset of  $E^k$ ) is the basic cosmological observable:

$$s^{\operatorname{red det.}} \stackrel{\text{det.}}{=} \mathfrak{s}_{k-1}^*(s) \in \mathcal{C}^0(\mathfrak{s}_{k-1}^{-1}(U), \mathcal{E}^1)$$
.

In the definition above, we used the relation  $\mathfrak{s}_k^*(\mathcal{E}^k) = \mathfrak{s}_k^*(\pi_k^*(\mathcal{E})) \simeq (\pi_k \circ \mathfrak{s}_k)^*(\mathcal{E}) = \pi^*(\mathcal{E}) = \mathcal{E}^1$ , which follows from the relation  $\pi_k = \pi \circ \pi_{k,1}$  and from the fact that  $\mathfrak{s}_k$  is a section of the fiber bundle  $\pi_{k,1} : E^k \to E^1$ .

**Proposition 2.9.** Consider a  $\mathcal{E}$ -valued local cosmological observable  $\mathfrak{s} \in \mathcal{C}^0(U, \mathcal{E}^k)$ of order  $k \geq 2$ , where U is a non-empty open subset of  $E^k$  which satisfies  $(\mathfrak{s}_{k-1} \circ \pi_{k,1})(U) \subset U$ . Then for any cosmological curve  $\varphi : I \to \mathcal{M}$  such that  $j^k(\varphi)(I) \subset U$ , we have  $\dot{\varphi}(I) \subset \mathfrak{s}_{k-1}^{-1}(U)$  and:

$$s_{\varphi} = s_{\varphi}^{\text{red}} = \dot{\varphi}^*(s^{\text{red}}) \in \mathcal{C}^0(I, \varphi^*(\mathcal{E})) .$$
(18)

*Proof.* Since  $\dot{\varphi} = \pi_{k,1} \circ j^k(\varphi)$ , we have  $\mathfrak{s}_{k-1}(\dot{\varphi}(t)) = (\mathfrak{s}_{k-1} \circ \pi_{k,1})(j^k(\varphi)(t)) \in (\mathfrak{s}_{k-1} \circ \pi_{k,1})(U) \subset U$ , which implies  $\dot{\varphi}(I) \subset \mathfrak{s}_{k-1}^{-1}(U)$ . Thus  $\dot{\varphi}^*(s^{\text{red}})$  is defined on the entire interval I. By the definition of  $s_{\varphi}$ , we have:

$$s_{\varphi} = j^{k}(\varphi)^{*}(s) = j_{1}^{*}(\varphi)(\mathfrak{s}_{k-1}^{*}(s)) = j_{1}^{*}(\varphi)(s^{\text{red}}) = \dot{\varphi}^{*}(s^{\text{red}}) ,$$

where we used (17) and the fact that  $j_1(\varphi) = \dot{\varphi}$ .

The condition that s be continuous (though natural for most observables of physical interest) is inessential in the discussion above. It is easy to see that the results of this subsection also apply to discontinuous sections of  $\mathcal{E}^k$ , which define "discontinuous local observables of order k". We will encounter some examples of such in latter sections.

Remark 2.10. Since  $E_k = J^k(\mathcal{M})$  consists of k-jets of pointed curves at t = 0, any  $\mathcal{E}$ -valued observable  $s \in \mathcal{C}^0(U, \mathcal{E}^k)$  of order k is completely determined by specifying its evaluation  $s_{\varphi}$  on arbitrary curves  $\varphi : I \to U$ . We will use this fact in latter sections to describe second order observables through their evaluation on arbitrary curves instead of giving their jet bundle description.

2.3.2. The cosmological energy function.

**Definition 2.11.** The cosmological energy function is the scalar basic local cosmological observable  $E: T\mathcal{M} \to \mathbb{R}_{>0}$  defined through:

$$E(u) \stackrel{\text{def.}}{=} \frac{1}{2} ||u||^2 + \Phi(\pi(u)) \ \forall u \in T\mathcal{M} \ .$$

The cosmological kinetic and potential energy functions and the cosmological potential energy function are the basic scalar local cosmological observables  $E_{\rm kin}, E_{\rm pot}$ :  $T\mathcal{M} \to \mathbb{R}_{\geq 0}$  defined through:

$$E_{\text{kin}}(u) \stackrel{\text{def.}}{=} \frac{1}{2} ||u||^2 , \ E_{\text{pot}}(u) \stackrel{\text{def.}}{=} \Phi(\pi(u)) \ \forall u \in T\mathcal{M} .$$

With these definitions, we have:

$$E = E_{\rm kin} + E_{\rm pot}$$
.

Notice that  $E_{\text{pot}}$  coincides with the natural lift of  $\Phi$  to  $T\mathcal{M}$ :

$$E_{\rm pot} = \Phi^{\rm v} \stackrel{\rm def.}{=} \Phi \circ \pi$$
 .

Also notice the relation:

$$\mathcal{H} = \frac{1}{M_0} \sqrt{2E} \; .$$

**Proposition 2.12.** The evaluation of the cosmological energy along any cosmological curve  $\varphi : I \to \mathcal{M}$  satisfies the cosmological dissipation equation:

$$\frac{\mathrm{d}E_{\varphi}(t)}{\mathrm{d}t} = -\frac{\sqrt{2E_{\varphi}(t)}}{M_0} ||\dot{\varphi}(t)||^2 \iff \frac{\mathrm{d}\mathcal{H}_{\varphi}(t)}{\mathrm{d}t} = -\frac{||\dot{\varphi}(t)||^2}{M_0^2} \ \forall t \in I \ . \tag{19}$$

In particular,  $E_{\varphi}$  is a strictly decreasing function of t if  $\varphi$  is not constant.

*Proof.* Follows immediately by using the cosmological equation (11).

The cosmological dissipation equation is equivalent with:

$$\frac{\mathrm{d}\mathcal{H}_{\varphi}(t)}{\mathrm{d}t} = -||\dot{\varphi}(t)||^2 \ . \tag{20}$$

Integrating (20) from  $t_1$  to  $t_2$  along the cosmological curve  $\varphi$  gives:

$$\mathcal{H}_{\varphi}(t_1) - \mathcal{H}_{\varphi}(t_2) = \int_{t_1}^{t_2} \mathrm{d}t ||\dot{\varphi}(t)||^2$$

Since the integral in the right hand side is minimized by the smallest length geodesic which connects the points  $\varphi(t_1)$  to  $\varphi(t_2)$  in  $\mathcal{M}$ , we have:

$$\int_{t_0}^{t_1} \mathrm{d}t ||\dot{\varphi}(t)||^2 \ge \mathrm{d}(\varphi(t_1), \varphi(t_2))^2$$

where d is the distance function [25] of  $(\mathcal{M}, \mathcal{G})$ . This gives the Hubble inequality:

$$\mathcal{H}_{\varphi}(t_1) - \mathcal{H}_{\varphi}(t_2) \ge \mathrm{d}(\varphi(t_1), \varphi(t_2))^2 .$$
(21)

## 3. The adapted frame of the non-critical submanifold

In this section, we discuss a certain frame of  $T\mathcal{M}_0$  which is induced by  $\mathcal{G}$  and  $\Phi$ . Let:

$$n \stackrel{\text{def.}}{=} \frac{\text{grad}\Phi}{||\text{grad}\Phi||} \in \mathfrak{X}(\mathcal{M}_0)$$
(22)

be the normalized gradient vector field of  $\Phi$ , which is well-defined on the noncritical submanifold  $\mathcal{M}_0$ . Let  $\tau \in \mathfrak{X}(\mathcal{M}_0)$  be the normalized vector field which is orthogonal to n and chosen such that the frame  $(n, \tau)$  of  $T\mathcal{M}_0$  is positively oriented at every point of  $\mathcal{M}_0$ . In the notations of Appendix A, we have  $\tau = N_n$ . Thus (see equation (145)):

$$\tau = (\iota_n \operatorname{vol}|_{\mathcal{M}_0})^{\sharp} , \qquad (23)$$

where vol is the volume form of  $(\mathcal{M}, \mathcal{G})$ .

**Definition 3.1.** The oriented orthonormal frame  $(n, \tau)$  of  $T\mathcal{M}_0$  is called the *adapted frame* of the oriented scalar triple  $(\mathcal{M}, \mathcal{G}, \Phi)$ .

Recall that the gradient flow and level set curves of  $\Phi$  are mutually orthogonal on the noncritical locus  $\mathcal{M}_0$ . Hence these curves determine an orthogonal mesh on the Riemannian manifold  $(\mathcal{M}_0, \mathcal{G}|_{\mathcal{M}_0})$ .

**Definition 3.2.** The *potential mesh* of the oriented scalar triple  $(\mathcal{M}, \mathcal{G}, \Phi)$  is the oriented mesh determined by the gradient flow and level set curves of  $\Phi$  on the noncritical set  $\mathcal{M}_0$ , where we orient gradient flow lines towards increasing values of  $\Phi$  and level set curves such that their normalized tangent vectors generate the vector field  $\tau$ .

3.1. The characteristic functions of a scalar triple. With Definition 3.2, the frame  $(n, \tau)$  is the oriented orthonormal frame defined by the potential mesh. For any point  $m \in \mathcal{M}_0$ , let  $\mu(m)$  and  $\lambda(m)$  respectively be the *signed* curvatures (see Definition A.10 in Appendix A) at m of the unique gradient and level set curves of  $\Phi$  which pass through m. This gives smooth functions  $\mu, \lambda \in \mathcal{C}^{\infty}(\mathcal{M}_0)$ .

**Definition 3.3.** The functions  $\mu, \lambda \in C^{\infty}(\mathcal{M}_0)$  are called the *first and second* characteristic functions of the oriented scalar triple  $(\mathcal{M}, \mathcal{G}, \Phi)$ .

Since  $N_n = \tau$  and  $N_\tau = -n$ , the Frenet-Serret relations (148), (149) for the gradient flow and level set curves give the *characteristic equations* of the oriented scalar triple  $(\mathcal{M}, \mathcal{G}, \Phi)$ :

$$\nabla_n n = \mu \tau , \ \nabla_n \tau = -\mu n$$
  
$$\nabla_\tau \tau = -\lambda n , \ \nabla_\tau n = \lambda \tau .$$
(24)

For any vector field  $X \in \mathfrak{X}(\mathcal{M}_0)$ , we denote by:

$$\Phi_X \stackrel{\text{def.}}{=} (\mathrm{d}\Phi)(X) = X(\Phi) \in \mathcal{C}^{\infty}(\mathcal{M})$$

the directional derivative of  $\Phi$  with respect to X. Then the gradient of  $\Phi$  expands as follows in any orthonormal frame (X, Y) of  $T\mathcal{M}$ :

$$\operatorname{grad}\Phi = \mathcal{G}(X, \operatorname{grad}\Phi)X + \mathcal{G}(Y, \operatorname{grad}\Phi)Y = (\mathrm{d}\Phi)(X)X + (\mathrm{d}\Phi)(Y)Y = \Phi_X X + \Phi_Y Y.$$
(25)

For the adapted frame, we have:

$$\Phi_n = n(\Phi) = (\mathrm{d}\Phi)(n) = \mathcal{G}(\mathrm{grad}\Phi, n) = ||\mathrm{grad}\Phi|| = ||\mathrm{d}\Phi||$$
  
$$\Phi_\tau = \tau(\Phi) = (\mathrm{d}\Phi)(\tau) = \mathcal{G}(\mathrm{grad}\Phi, \tau) = 0.$$
(26)

For any vector fields X, Y defined on  $\mathcal{M}$ , we set:

$$\Phi_{XY} \stackrel{\text{def.}}{=} \text{Hess}(\Phi)(X,Y) = (\nabla_X d\Phi)(Y) = X(Y(\Phi)) - (d\Phi)(\nabla_X Y) = X(Y(\Phi)) - (\nabla_X Y)(\Phi)$$
(27)

Notice that  $\Phi_{XY} = \Phi_{YX}$  since the Levi-Civita connection  $\nabla$  of  $(\mathcal{M}, \mathcal{G})$  is torsion-free.

**Definition 3.4.** The function  $\Phi_{nn} \in \mathcal{C}^{\infty}(\mathcal{M}_0)$  is called the *third characteristic* function of the scalar triple  $(\mathcal{M}, \mathcal{G}, \Phi)$ .

**Proposition 3.5.** The following relations hold on the non-critical submanifold  $\mathcal{M}_0$ :

$$\Phi_{nn} = n(||d\Phi||) \quad , \quad \Phi_{\tau\tau} = \lambda ||d\Phi|| \Phi_{n\tau} = \mu ||d\Phi|| = \tau(||d\Phi||) = \Phi_{\tau n}$$
(28)

In particular, the following relations hold on  $\mathcal{M}_0$ :

$$\lambda = \frac{\Phi_{\tau\tau}}{||\mathrm{d}\Phi||} \ , \ \mu = \frac{\Phi_{n\tau}}{||\mathrm{d}\Phi||} \ . \tag{29}$$

Remark 3.6. The last two relations in (28) imply:

$$\mu = \tau(\log ||\mathbf{d}\Phi||) \ . \tag{30}$$

*Proof.* Equations (26) give:

$$n(\Phi) = ||d\Phi|| , \ \tau(\Phi) = 0 .$$
 (31)

Relations (28) follow immediately from (24) and (27) upon using (31). The second and third relation in (28) give (29).  $\hfill \Box$ 

## 3.2. Adapted coordinates on the nondegenerate set.

**Definition 3.7.** The *nondegenerate set* of  $(\mathcal{M}, \mathcal{G}, \Phi)$  is the following open subset of  $\mathcal{M}_0$ :

$$\mathcal{U} \stackrel{\text{def.}}{=} \{ m \in \mathcal{M}_0 \mid \mu(m) \neq 0 \} = \{ m \in \mathcal{M}_0 \mid \Phi_{n\tau}(m) \neq 0 \} .$$
(32)

Notice that  $\mathcal{U}$  is empty iff all gradient lines of  $\Phi$  are geodesics. In this case, we say the scalar triple  $(\mathcal{M}, \mathcal{G}, \Phi)$  is *degenerate*. For the remainder of this paper, we assume that  $(\mathcal{M}, \mathcal{G}, \Phi)$  is a nondegenerate scalar triple. Consider the non-negative function:

$$\Psi \stackrel{\text{def.}}{=} ||d\Phi|| \in \mathcal{C}^{\infty}(\mathcal{M}, \mathbb{R}_{\geq 0}) ,$$

whose vanishing locus coincides with  $\operatorname{Crit}\Phi$  and let  $\beta \stackrel{\text{def.}}{=} (\Phi, \Psi) : \mathcal{M} \to \mathbb{R}_{>0} \times \mathbb{R}_{>0}$ .

**Proposition 3.8.** The following relations hold on the non-critical submanifold  $\mathcal{M}_0$ :

$$grad\Phi = ||d\Phi||n$$
$$grad||d\Phi|| = \Phi_{nn}n + \mu ||d\Phi||\tau$$

*Proof.* The first relation is obvious. The second follows from:

 $\mathcal{G}(\operatorname{grad}||\mathrm{d}\Phi||, n) = n(||\mathrm{d}\Phi||) = \Phi_{nn} , \ \mathcal{G}(\operatorname{grad}||\mathrm{d}\Phi||, \tau) = \tau(||\mathrm{d}\Phi||) = \mu ||\mathrm{d}\Phi|| ,$ 

where we used (28). on  $\mathcal{U}$ .

**Corollary 3.9.** If the map  $\beta_{\mathcal{U}}$  is proper, then it is a covering map.

*Proof.* Follows immediately from [28, Lemma 2] using the fact that  $\mathbb{R}^2$  and  $\mathcal{M}$  are locally compact space when endowed with the manifold topology.

**Corollary 3.10.** The map  $\beta$  has rank 2 on  $\mathcal{U}$ , rank 1 on  $\mathcal{M}_0 \setminus \mathcal{U}$  and is locally constant on  $\operatorname{Crit}\Phi = \mathcal{M} \setminus \mathcal{M}_0$ . In particular the restriction  $\beta_{\mathcal{U}} \stackrel{\text{def.}}{=} (\Phi, \Psi) : \mathcal{U} \to \mathcal{B}$  is a surjective local diffeomorphism, where  $\Psi \stackrel{\text{def.}}{=} ||\mathrm{d}\Phi||$  and we defined  $\mathcal{B} \stackrel{\text{def.}}{=} \beta(\mathcal{U})$ .

Since  $\beta_{\mathcal{U}}$  is a local diffeomorphism, it provides local coordinates on a small enough neighborhood (contained in  $\mathcal{U}$ ) of each point of  $\mathcal{U}$ .

**Definition 3.11.** The coordinates  $(\Phi, \Psi = ||d\Phi||)$  defined on a small enough neighborhood of a point  $m \in \mathcal{U}$  of the non-degenerate set are called *adapted coordinates*.

## 3.3. Basic observables defined by the adapted frame.

3.3.1. The signed characteristic angle and characteristic signature. The adapted frame defines two natural discontinuous local observables which are related to each other. We refer the reader to Appendix A for the definition of the signed angle and relative chirality of pairs of nonzero vectors on an oriented Riemannian manifold, which are used in the definition below.

**Definition 3.12.** Let  $u \in T\mathcal{M}_0$  be a non-zero vector tangent to  $\mathcal{M}$  at a noncritical point  $m \in \mathcal{M}_0$ .

• The signed angle:

Jaf

$$\theta(u) \stackrel{\text{def.}}{=} \theta_{n(m)}(u) \in (-\pi, \pi]$$

of u relative to the unit gradient vector  $n(m) \in T_m \mathcal{M}$  is called the *signed* characteristic angle of u

• The chirality of u with respect to n(m):

$$\sigma(u) \stackrel{\text{def.}}{=} \sigma_{n(m)}(u) = \operatorname{sign} \left[ \sin \theta(u) \right] = \operatorname{sign} \mathcal{G}(\tau(m), u) \in \{-1, 0, 1\}$$

is called the *characteristic signature* of u.

The signed characteristic angle and characteristic signature define functions  $\theta$ :  $\dot{T}\mathcal{M}_0 \rightarrow (-\pi,\pi]$  and  $\sigma$ :  $\dot{T}\mathcal{M}_0 \rightarrow \{-1,0,1\}$  which we call the signed characteristic angle function and characteristic signature function of  $(\mathcal{M}, \mathcal{G}, \Phi)$ . These are discontinuous basic local observables of the model.

3.3.2. The lift of the adapted frame. Recall the abstract oriented Frenet frame (T, N) of  $F|_{\dot{T}\mathcal{M}}$  defined in Appendix A, which gives a universal geometric description of the oriented Frenet frame of curves trough Finsler vector fields defined on the slit tangent bundle of  $\mathcal{M}$ .

**Definition 3.13.** The *tautological lift* of a vector field  $X \in \mathfrak{X}(\mathcal{M})$  is the Finsler vector field  $X^{\mathsf{v}} \in \Gamma(T\mathcal{M}, F)$  given by:

$$X^{\mathrm{v}}(u) \stackrel{\mathrm{def.}}{=} X(\pi(u)) \ \forall u \in T\mathcal{M} \ .$$

The adapted frame  $(n, \tau)$  of  $T\mathcal{M}_0$  lifts to a frame  $(n^{\mathrm{v}}, \tau^{\mathrm{v}})$  of the restricted tautological bundle  $F_0 \stackrel{\text{def.}}{=} F|_{\mathcal{M}_0} = \pi^*(T\mathcal{M}_0)$ . Let  $T_0 \stackrel{\text{def.}}{=} T|_{\dot{T}\mathcal{M}_0}$ . The signed angle and chirality maps  $\theta : \dot{T}\mathcal{M} \times_{\mathcal{M}} \dot{T}\mathcal{M} \to (-\pi, \pi]$  and  $\sigma : \dot{T}\mathcal{M} \times_{\mathcal{M}} \dot{T}\mathcal{M} \to \{-1, 0, 1\}$  of the oriented Riemann surface  $(\mathcal{M}, \mathcal{G})$  (see Definition A.6) restrict to maps  $\theta_0$  and  $\sigma_0$  defined on  $\mathcal{M}_0$ , which are the signed angle and chirality maps of the oriented Riemannian manifold  $(\mathcal{M}_0, \mathcal{G}|_{\mathcal{M}_0})$ . The following immediate statement expresses  $\theta$ and  $\sigma$  through  $n^{\mathrm{v}}$ ,  $T_0$  and  $\theta_0, \sigma_0$ .

**Proposition 3.14.** The signed characteristic angle and characteristic signature functions of  $(\mathcal{M}, \mathcal{G}, \Phi)$  satisfy:

$$\theta = \theta_0 \circ (n^{\mathrm{v}}, T_0) , \ \sigma \stackrel{\text{def.}}{=} \sigma_0 \circ (n^{\mathrm{v}}, T_0) , \tag{33}$$

where  $n^{v}$  and  $T_{0}$  are viewed as maps from  $\dot{T}\mathcal{M}_{0}$  to  $\dot{T}\mathcal{M}_{0}$  and the map  $(n^{v}, T_{0})$ :  $\dot{T}\mathcal{M}_{0} \rightarrow \dot{T}\mathcal{M}_{0} \times_{\mathcal{M}_{0}} \dot{T}\mathcal{M}_{0}$  is defined through:

$$(n^{\mathbf{v}}, T_0)(u) = (n^{\mathbf{v}}(u), T(u)) = \left(n(\pi(u)), \frac{u}{||u||}\right) \in \dot{T}_{\pi(u)}\mathcal{M} \times \dot{T}_{\pi(u)}\mathcal{M} \ \forall u \in \dot{T}\mathcal{M}_0 \ .$$

*Remark* 3.15. For simplicity of notation, we will omit the composition sign in relations such as (33) from now on.

The following result follows immediately from Proposition A.4 of Appendix A.

**Proposition 3.16.** The following relations hold on  $T\mathcal{M}_0$ :

$$T = n^{v} \cos \theta + \tau^{v} \sin \theta$$
  

$$N = -n^{v} \sin \theta + \tau^{v} \cos \theta .$$
(34)

3.3.3. The adapted speed components. Let  $I \in \Gamma(T\mathcal{M}, F)$  be the tautological Finsler field, which is defined though:

$$I(u) \stackrel{\text{def.}}{=} u \ \forall u \in T\mathcal{M}$$

and let  $I_0 \stackrel{\text{def.}}{=} I|_{T\mathcal{M}_0}$ . Since  $u = \mathcal{N}(u)T(u)$  for all  $u \in \dot{T}\mathcal{M}$ , we have:

$$I|_{\dot{T}\mathcal{M}} = \mathcal{N}|_{\dot{T}\mathcal{M}}T . \tag{35}$$

**Definition 3.17.** The adapted speed component functions are the basic local observables  $v_n, v_\tau : T\mathcal{M}_0 \to \mathbb{R}$  defined through:

$$I_0 = v_n n^{\mathsf{v}} + v_\tau \tau^{\mathsf{v}} . \tag{36}$$

Relation (36) amounts to:

$$u = v_n(u)n(\pi(u)) + v_\tau(u)\tau(\pi(u)) \ \forall u \in T\mathcal{M}_0$$
(37)

and we have:

$$v_n(u) = \mathcal{G}(n^{\mathsf{v}}(u), u) , \ v_\tau(u) = \mathcal{G}(\tau^{\mathsf{v}}(u), u) \ \forall u \in T\mathcal{M}_0 .$$
(38)

Notice that  $(v_n, v_\tau) : T\mathcal{M}_0 \to \mathbb{R}^2$  is the linear system of fiberwise coordinates defined by the adapted frame  $(n, \tau)$  of  $T\mathcal{M}_0$  in the sense of Subsection 1.6.

**Proposition 3.18.** The following relations hold on  $T\mathcal{M}_0$ :

$$\mathcal{N} = \sqrt{v_n^2 + v_\tau^2} , \ v_n = \mathcal{N}\cos\theta , \ v_\tau = \mathcal{N}\sin\theta ,$$
 (39)

where  $\mathcal{N}: T\mathcal{M} \to \mathbb{R}_{\geq 0}$  is the norm function. The first of these relations holds on all of  $T\mathcal{M}_0$ .

*Proof.* Follows immediately from (37), (38) and the first equation in (34) using (35).  $\Box$ 

3.3.4. The tangential and positive normal derivative of the scalar potential. For any function  $f \in \mathcal{C}^{\infty}(\mathcal{M})$ , we define:

$$f^{\mathrm{v}} \stackrel{\mathrm{def.}}{=} \pi^*(f) = f \circ \pi \in \mathcal{C}^{\infty}(T\mathcal{M})$$
.

For any Finsler vector field  $W \in \Gamma(T\mathcal{M}, F)$ , define  $\Phi_W \in \mathcal{C}^{\infty}(T\mathcal{M})$  through:

 $\Phi_W(u) \stackrel{\text{def.}}{=} (\mathrm{d}\Phi)(\pi(u))(W(u)) = \mathcal{G}((\mathrm{grad}\Phi)^{\mathrm{v}}(u), W(u)) \ \forall u \in T\mathcal{M} \ .$ 

With these definitions, we have:

$$\Phi_{X^{\mathbf{v}}} = (\Phi_X)^{\mathbf{v}} \; \forall X \in T\mathcal{M} \; .$$

**Definition 3.19.** The basic local observables  $\Phi_T, \Phi_N \in \mathcal{C}^{\infty}(\dot{T}\mathcal{M})$  are called the *tangential derivative* and *positive normal derivative* of  $\Phi$ .

**Proposition 3.20.** The following relations hold on  $T\mathcal{M}_0$ :

$$\Phi_T = \mathcal{G}((\operatorname{grad}\Phi)^{\mathrm{v}}, T) = + ||\mathrm{d}\Phi||^{\mathrm{v}} \cos\theta$$
  
$$\Phi_N = \mathcal{G}((\operatorname{grad}\Phi)^{\mathrm{v}}, N) = - ||\mathrm{d}\Phi||^{\mathrm{v}} \sin\theta .$$
(40)

*Proof.* Follows immediately using (34) upon noticing that  $(\operatorname{grad} \Phi)^{\mathsf{v}}(u) = ||\mathrm{d} \Phi||^{\mathsf{v}} n(u)$  for all  $u \in T\mathcal{M}_0$ .  $\Box$ 

3.3.5. Evaluation along a curve. The formulation given above encodes in a universal manner certain objects and relations which are generally presented using a given curve in  $\mathcal{M}_0$ . This universal formulation is made possible by the fact that jets of curves generate the bundles  $J^k(\mathcal{M})$  (see Remark 2.10). The usual formulation is recovered by considering the evaluation of the basic local observables defined above a long a curve.

Given a noncritical curve  $\varphi: I \to \mathcal{M}_0$  and a regular time  $t \in I_{\text{reg}}$ , we define:

$$n_{\varphi}(t) \stackrel{\text{def.}}{=} n(\varphi(t)) \in T_{\varphi(t)}\mathcal{M}_0 \ , \ \tau_{\varphi}(t) \stackrel{\text{def.}}{=} \tau(\varphi(t)) \in T_{\varphi(t)}\mathcal{M}_0 \ .$$

Let  $T_{\varphi}(t)$  be the unit tangent vector to  $\varphi$  at time t and  $N_{\varphi}(t)$  be the positive normal vector to  $\varphi$  at time t, which is chosen such that the orthonormal basis  $(T_{\varphi}(t), N_{\varphi}(t))$  of  $T_{\varphi(t)}\mathcal{M}_0$  is positively oriented. We have (see Appendix A):

$$N_{\varphi}(t) = N(\dot{\varphi}(t)) , \ T_{\varphi}(t) = T(\dot{\varphi}(t)) .$$

**Definition 3.21.** Consider a noncritical curve  $\varphi: I \to \mathcal{M}_0$ . Then:

• The signed characteristic angle:

θ

$$\varphi(t) \stackrel{\text{def.}}{=} \theta(T_{\varphi}(t)) = \theta_{n_{\varphi}(t)}(T_{\varphi}(t)) \in (-\pi, \pi] \ (t \in I_{\text{reg}})$$

of  $T_{\varphi}(t)$  is called the *signed characteristic angle* of  $\varphi$  at the regular time t. • The characteristic signature:

 $\sigma_{\varphi}(t) \stackrel{\text{def.}}{=} \sigma(T_{\varphi}(t)) = \operatorname{sign}\left[\sin \theta_{\varphi}(t)\right] = \operatorname{sign} \mathcal{G}(\tau_{\varphi}(t), T_{\varphi}(t)) \ (t \in I_{\operatorname{reg}})$ 

of  $T_{\varphi}(t)$  is called the *characteristic signature* of  $\varphi$  at the regular time t.

• The functions  $v_n, v_\tau : I \to \mathbb{R}$  defined through:

$$v_n^{\varphi} = v_n \circ \dot{\varphi} , \ v_{\tau}^{\varphi} = v_{\tau} \circ \dot{\varphi} \ (t \in I).$$

are called the *adapted speed components of*  $\varphi$ .

For any noncritical curve  $\varphi: I \to \mathcal{M}_0$ , relation (36) gives:

$$\dot{\varphi} = v_n^{\varphi} n_{\varphi} + v_{\tau}^{\varphi} \tau_{\varphi} \ \forall t \in I \tag{41}$$

and we have:

$$v_n^{\varphi} \stackrel{\text{def.}}{=} \mathcal{G}(n_{\varphi}, \dot{\varphi}) = ||\dot{\varphi}|| \cos \theta_{\varphi} , \ v_{\tau}^{\varphi} \stackrel{\text{def.}}{=} \mathcal{G}(\tau_{\varphi}, \dot{\varphi}) = ||\dot{\varphi}|| \sin \theta_{\varphi}$$
(42)

Propositions 3.16, 3.18 and 3.20 imply:

20

**Proposition 3.22.** Consider a noncritical curve  $\varphi : I \to \mathcal{M}_0$ . Then the following relations hold for  $t \in I_{reg}$ :

$$T_{\varphi}(t) = n_{\varphi}(t) \cos \theta_{\varphi}(t) + \tau_{\varphi}(t) \sin \theta_{\varphi}(t) N_{\varphi}(t) = -n_{\varphi}(t) \sin \theta_{\varphi}(t) + \tau_{\varphi}(t) \cos \theta_{\varphi}(t) .$$
(43)

In particular, we have:

$$\mathcal{G}(n_{\varphi}(t), T_{\varphi}(t)) = \cos \theta_{\varphi}(t) , \ \mathcal{G}(n_{\varphi}(t), N_{\varphi}(t)) = -\sin \theta_{\varphi}(t) \ \forall t \in I_{\text{reg}} .$$
(44)

Moreover, we have:

$$||\dot{\varphi}|| = \sqrt{(v_n^{\varphi})^2 + (v_\tau^{\varphi})^2} \quad \forall t \in I$$
(45)

and:

$$(\mathrm{d}\Phi)(T_{\varphi}(t)) = \Phi_T(\dot{\varphi}(t)) , \ (\mathrm{d}\Phi)(N_{\varphi}(t)) = \Phi_N(\dot{\varphi}(t)) \ \forall t \in I_{\mathrm{reg}} .$$

3.4. Adapted and modified phase space coordinates. Consider the following functions defined on  $T\mathcal{M}$ :

$$\Phi \stackrel{\text{def.}}{=} \Phi^{\mathbf{v}}$$
,  $\Psi \stackrel{\text{def.}}{=} \Psi^{v} = ||\mathbf{d}\Phi||^{\mathbf{v}}$ .

Notice that  $\Phi$  is smooth on  $T\mathcal{M}$  while  $\Psi$  is smooth on  $T\mathcal{M}_0$ . Recall that  $\mathcal{H}$  denotes the rescaled Hubble function of  $\mathcal{M}$ .

**Lemma 3.23.** The functions  $(v_n, v_\tau, \Phi, \Psi)$  and  $(v_n, v_\tau, \mathcal{H}, \Psi)$  form systems of local coordinates on a small enough neighborhood of any point of the open set  $T\mathcal{U} \subset T\mathcal{M}_0$ , where  $\mathcal{U}$  is the nondegenerate set of  $(\mathcal{M}, \mathcal{G}, \Phi)$  (see Definition 3.7).

*Proof.* Follows immediately from Proposition 3.8 upon using the fact that  $v_n, v_\tau$  give (linear) fiber coordinates on  $T\mathcal{M}_0$ .

**Definition 3.24.** The local coordinates  $(v_n, v_\tau, \Phi, \Psi)$  defined on a small enough neighborhood of any point of  $T\mathcal{U}$  are called *adapted phase space coordinates*.

If  $u \in T\mathcal{U}$  and  $U \subset \mathcal{U}$  is a neighborhood of  $\pi(u)$  on which  $\Phi$  and  $\Psi$  are coordinates, then  $(v_n, v_\tau, \Phi, \Psi)$  are coordinates on the neighborhood TU of u. Adapted phase space coordinates give a special coordinate system defined on TU whose associated fiberwise coordinate system  $(v_n, v_\tau)$  is linear; the latter is defined by the frame  $(n, \tau)$  of  $T\mathcal{M}_0$  (which generally is *not* holonomic).

**Proposition 3.25.** The functions  $(v_n, v_\tau, \mathcal{H}, \Phi)$  give local coordinates on a small enough vicinity of any point of  $T\mathcal{U}$ .

*Proof.* Since  $\Phi$  is strictly positive, the relation:

$$\mathcal{H} = \frac{1}{M_0} (v_n^2 + v_\tau^2 + 2\Phi)^{1/2} \Longleftrightarrow \Phi = \frac{1}{2} (M_0^2 \mathcal{H}^2 - v_n^2 - v_\tau^2)$$
(46)

gives a smooth change of coordinates  $(v_n, v_\tau, \Phi, \Psi) \leftrightarrow (v_n, v_\tau, \mathcal{H}, \Psi)$  on a small enough vicinity of every point of  $T\mathcal{U}$ . Hence  $(v_n, v_\tau, \mathcal{H}, \Psi)$  is a local coordinate system on a small enough vicinity of any point of  $T\mathcal{U}$ .

**Definition 3.26.** The local coordinates  $(v_n, v_\tau, \mathcal{H}, \Psi)$  defined on a small enough neighborhood of any point of  $T\mathcal{U}$  are called *modified phase space coordinates*.

Remark 3.27. Relation (46) gives:

$$\kappa = \frac{v_n^2 + v_\tau^2}{M_0^2 \mathcal{H}^2 - (v_n^2 + v_\tau^2)} = \frac{1}{\frac{M_0^2 \mathcal{H}^2}{v_n^2 + v_\tau^2} - 1} .$$
(47)

3.5. The cosmological semispray in adapted phase space coordinates. In this subsection, we determine the form of the cosmological equation (and hence of the cosmological semispray) in adapted coordinates. We will first write the cosmological equation in modified phase space coordinates and then change to adapted phase space coordinates. Let  $\varphi : I \to \mathcal{M}_0$  be a noncritical cosmological curve. Recall that:

$$\Phi_{\varphi} \stackrel{\text{def.}}{=} \Phi \circ \varphi \ , \ \Psi_{\varphi} \stackrel{\text{def.}}{=} \Psi \circ \varphi \ , \ \mathcal{H}_{\varphi} \stackrel{\text{def.}}{=} \mathcal{H} \circ \varphi = \frac{1}{M_0} [(v_n^{\varphi})^2 + (v_\tau^{\varphi})^2 + 2\Phi_{\varphi}]^{1/2}$$

and:

$$v_n^{\varphi} \stackrel{\text{def.}}{=} v_n \circ \dot{\varphi} \ , \ v_{\tau}^{\varphi} \stackrel{\text{def.}}{=} v_{\tau} \circ \dot{\varphi} \ , \ n_{\varphi} \stackrel{\text{def.}}{=} n \circ \varphi \ , \ \tau_{\varphi} \stackrel{\text{def.}}{=} \tau \circ \varphi \ .$$
(48)

and define:

$$\lambda_{\varphi} \stackrel{\text{def.}}{=} \lambda \circ \varphi \ , \ \mu_{\varphi} \stackrel{\text{def.}}{=} \mu \circ \varphi$$

We remind the reader that  $\Psi \stackrel{\text{def.}}{=} ||d\Phi||$ . When  $\varphi$  is both noncritical and nondegenerate, we have (see (43)):

$$T_{\varphi} = n_{\varphi} \cos \theta_{\varphi} + \tau_{\varphi} \sin \theta_{\varphi}$$
$$N_{\varphi} = -n_{\varphi} \sin \theta_{\varphi} + \tau_{\varphi} \cos \theta_{\varphi} ,$$

where  $\theta_{\varphi}: I \to (-\pi, \pi]$  is the signed characteristic angle of  $\varphi$ .

**Lemma 3.28.** The cosmological equation for a curve  $\varphi : I \to \mathcal{U}$  whose image is contained in  $\mathcal{U}$  is equivalent with:

$$\dot{v}_{n}^{\varphi} - \lambda_{\varphi} (v_{\tau}^{\varphi})^{2} - \mu_{\varphi} v_{n}^{\varphi} v_{\tau}^{\varphi} + \mathcal{H}_{\varphi} v_{n}^{\varphi} + \Psi_{\varphi} = 0$$
  
$$\dot{v}_{\tau}^{\varphi} + \mu_{\varphi} (v_{n}^{\varphi})^{2} + \lambda_{\varphi} v_{n}^{\varphi} v_{\tau}^{\varphi} + \mathcal{H}_{\varphi} v_{\tau}^{\varphi} = 0 , \qquad (49)$$

together with the following two equations:

$$\frac{\mathrm{d}\mathcal{H}_{\varphi}}{\mathrm{d}t} = -\frac{1}{M_0^2} [(v_n^{\varphi})^2 + (v_{\tau}^{\varphi})^2] 
\frac{\mathrm{d}\Psi_{\varphi}}{\mathrm{d}t} = \frac{1}{2} (\Phi_{nn\varphi} v_n^{\varphi} + \mu_{\varphi} \Psi_{\varphi} v_{\tau}^{\varphi}) ,$$
(50)

where  $\Phi_{nn\varphi} \stackrel{\text{def.}}{=} \Phi_{nn} \circ \varphi$ .

*Proof.* The speed  $\dot{\varphi}: I \to T\mathcal{U}$  expands in the adapted frame as:

$$\dot{\varphi} = v_n^{\varphi} n_{\varphi} + v_{\tau}^{\varphi} \tau_{\varphi} \ . \tag{51}$$

We have:

$$\nabla_t n_{\varphi} = (\mu_{\varphi} v_n^{\varphi} + \lambda_{\varphi} v_{\tau}^{\varphi}) \tau_{\varphi} \ , \ \nabla_t \tau_{\varphi} = -(\mu_{\varphi} v_n^{\varphi} + \lambda_{\varphi} v_{\tau}^{\varphi}) n_{\varphi} \ ,$$

where we used (24). Thus:

$$\nabla_t \dot{\varphi} = [\dot{v}_n^{\varphi} - \lambda_{\varphi} (v_{\tau}^{\varphi})^2 - \mu_{\varphi} v_n^{\varphi} v_{\tau}^{\varphi}] n_{\varphi} + [\dot{v}_{\tau}^{\varphi} + \mu_{\varphi} (v_n^{\varphi})^2 + \lambda_{\varphi} v_n^{\varphi} v_{\tau}^{\varphi}] \tau_{\varphi}$$

Hence the cosmological equation (11) implies (49). The first equation in (50) is the dissipation equation (19). To derive the second equation in (50), we compute:

$$\begin{aligned} \frac{\mathrm{d}||(\mathrm{d}\Phi)(\varphi(t))||}{\mathrm{d}t} &= \frac{\mathcal{G}(\nabla_t[(\mathrm{grad}\Phi)(\varphi(t))], (\mathrm{grad}\Phi)(\varphi(t)))}{2||(\mathrm{d}\Phi)(\varphi(t))||} \\ &= \frac{\mathrm{Hess}(\Phi)(\varphi(t))(\dot{\varphi}(t), (\mathrm{grad}\Phi)(\varphi(t)))}{2||(\mathrm{d}\Phi)(\varphi(t))||} = \frac{1}{2}\mathrm{Hess}(\Phi)(\varphi(t))(\dot{\varphi}(t), n_{\varphi}(t)) \\ &= \frac{1}{2}\left[\Phi_{nn}(\varphi(t))v_n^{\varphi}(t) + \Phi_{n\tau}(\varphi(t))v_{\tau}^{\varphi}(t)\right] \;, \end{aligned}$$

where we used (41). The second equation in (50) now follows from the relation above upon using (28). Hence the cosmological equation for  $\varphi$  implies (49) and (50). The system formed by these four first order ODEs is equivalent with the cosmological

22

equation on  $T\mathcal{U}$  since  $(v_n, v_\tau, \mathcal{H}, \Psi)$  is a local coordinate system around every point of  $T\mathcal{U}$ .

The previous Lemma can be reformulated as follows.

**Proposition 3.29.** The cosmological equation for curves whose image is contained in  $\mathcal{U}$  is equivalent with the following system written in local coordinates  $(v_n, v_\tau, \mathcal{H}, \Psi)$ in a small enough vicinity of any point  $u \in T\mathcal{U}$ :

$$\begin{split} \dot{v}_n - \lambda^{\mathsf{v}} v_\tau^2 &- \mu^{\mathsf{v}} v_n v_\tau + \mathcal{H} v_n + \Psi = 0 \\ \dot{v}_\tau + \mu^{\mathsf{v}} v_n^2 + \lambda^{\mathsf{v}} v_n v_\tau + \mathcal{H} v_\tau = 0 \\ \dot{\mathcal{H}} &= -\frac{1}{M_0^2} (v_n^2 + v_\tau^2) \\ \dot{\Psi} &= \frac{1}{2} (\Phi_{nn}^{\mathsf{v}} v_n + \mu^{\mathsf{v}} \Psi v_\tau) , \end{split}$$
(52)

where  $\Phi_{nn}$ ,  $\lambda$  and  $\mu$  are expressed locally as functions of  $v_n$ ,  $v_\tau$ ,  $\mathcal{H}$  and  $\Psi$  using the fact that  $(\Phi, \Psi)$  are local coordinates on a vicinity of  $\pi(u) \in \mathcal{U}$  and relations (46).

More precisely, the quantities  $\Phi_{nn}$ ,  $\lambda$  and  $\mu$  can be expressed locally around  $\pi(u)$  as functions of  $\Phi$  and  $\Psi$  and hence locally around u as functions of  $\Phi$  and  $\Psi$ . Using relations (46), their lifts to  $\mathcal{U}$  locally become functions of  $\Phi$ ,  $\mathcal{H}$  and  $v_n$ ,  $v_{\tau}$ .

**Proposition 3.30.** The cosmological equation for curves whose image is contained in  $\mathcal{U}$  is equivalent with the following system written in adapted phase space coordinates  $(v_n, v_\tau, \Phi, \Psi)$  on a small enough vicinity of any point  $u \in T\mathcal{U}$ :

$$\dot{v}_{n} = \lambda^{v} v_{\tau}^{2} + \mu^{v} v_{n} v_{\tau} - \frac{1}{M_{0}} (v_{n}^{2} + v_{\tau}^{2} + 2\Phi)^{1/2} v_{n} - \Psi$$
  

$$\dot{v}_{\tau} = -\mu^{v} v_{n}^{2} - \lambda^{v} v_{n} v_{\tau} - \frac{1}{M_{0}} (v_{n}^{2} + v_{\tau}^{2} + 2\Phi)^{1/2} v_{\tau}$$
  

$$\dot{\Phi} = \Psi v_{n}$$
  

$$\dot{\Psi} = \frac{1}{2} (\Phi_{nn}^{v} v_{n} + \mu^{v} \Psi v_{\tau}) ,$$
(53)

where  $\Phi_{nn}$ ,  $\lambda$  and  $\mu$  are expressed locally around  $\pi(u)$  in terms of  $\Phi$  and  $\Psi$  (and hence  $\Phi_{nn}^{v}$ ,  $\lambda^{v}$  and  $\mu^{v}$  are expressed locally around u in terms of  $\Phi$  and  $\Psi$ ).

*Proof.* The first two equations in (52) imply:

$$v_n \dot{v}_n + v_\tau \dot{v}_\tau = -\mathcal{H}(v_n^2 + v_\tau^2) - \Psi v_n .$$
(54)

Relation (46) implies:

$$\dot{\Phi} = M_0^2 \mathcal{H} \dot{\mathcal{H}} - (v_n \dot{v}_n + v_\tau \dot{v}_\tau) = -\mathcal{H}(v_n^2 + v_\tau^2) - (v_n \dot{v}_n + v_\tau \dot{v}_\tau) = \Psi v_n ,$$

where we used Proposition 2.12 and (54). This shows that the third equations of (52) and (53) are equivalent modulo the first two equations.

**Corollary 3.31.** The cosmological semispray has the following form in adapted phase space coordinates around every point  $u \in TU$ :

$$S = S^{v_n} \frac{\partial}{\partial v^n} + S^{v_\tau} \frac{\partial}{\partial v^\tau} + S^{\Phi} \frac{\partial}{\partial \Phi} + S^{\Psi} \frac{\partial}{\partial \Psi} \ ,$$

where:

$$S^{v_n} = \lambda^{v} v_{\tau}^2 + \mu^{v} v_n v_{\tau} - \frac{1}{M_0} (v_n^2 + v_{\tau}^2 + 2\Phi)^{1/2} v_n - \Psi$$

$$S^{v_{\tau}} = -\mu^{v} v_n^2 - \lambda^{v} v_n v_{\tau} - \frac{1}{M_0} (v_n^2 + v_{\tau}^2 + 2\Phi)^{1/2} v_{\tau}$$

$$S^{\Phi} = \Psi v_n$$

$$S^{\Psi} = \frac{1}{2} (\Phi_{nn}^{v} v_n + \mu^{v} \Psi v_{\tau}) ,$$
(55)

where  $\Phi_{nn}$ ,  $\lambda$  and  $\mu$  are expressed locally around  $\pi(u)$  in terms of  $\Phi$  and  $\Psi$  (and hence  $\Phi_{nn}^{v}$ ,  $\lambda^{v}$  and  $\mu^{v}$  are expressed locally around u in terms of  $\Phi$  and  $\Psi$ ).

These results show that the functions  $\lambda(\Phi, \Psi)$ ,  $\mu(\Phi, \Psi)$  and  $\Phi_{nn}(\Phi, \Psi)$  (which can be defined on a small enough vicinity of  $\pi(u)$ ) characterize the cosmological semispray on a small enough vicinity of any point of  $u \in T\mathcal{U}$ . Hence cosmological dynamics on  $T\mathcal{U}$  is locally determined by these three functions. This opens the possibility of studying the local cosmological dynamics of two-field models using the "universal form" (53) of their first order equations. In the next section, we rewrite (53) in certain special local coordinate systems which are induced by the so-called "fundamental basic observables" of the model, which have more direct physical interpretation.

#### 4. Fundamental basic observables

4.1. The fundamental parameters of a smooth curve in  $\mathcal{M}$ . Let  $\varphi: I \to \mathcal{M}$  be a smooth curve and  $T_{\varphi}(t)$ ,  $N_{\varphi}(t)$  be the unit tangent and positive unit normal vectors to  $\varphi$  at a regular time  $t \in I_{\text{reg}}$  (see Appendix A).

4.1.1. Fundamental first order parameters. We start by considering first order parameters, whose value at  $t \in I$  depends only only the tangent vector  $\dot{\varphi}(t)$ .

**Definition 4.1.** The fundamental first order parameters of a curve  $\varphi : I \to \mathcal{M}$  are defined as follows:

• The first IR parameter of  $\varphi$  at time  $t \in I$  is the non-negative real number:

$$\kappa_{\varphi}(t) \stackrel{\text{def.}}{=} \frac{||\dot{\varphi}(t)||^2}{2\Phi(\varphi(t))} \in \mathbb{R}_{\geq 0} .$$
(56)

• The rescaled conservative parameter of  $\varphi$  at non-critical time  $t \in I_{\text{noncrit}}$  is the non-negative real number:

$$\hat{c}_{\varphi}(t) \stackrel{\text{def.}}{=} \frac{\mathcal{H}_{\varphi}(t)||\dot{\varphi}(t)||}{||(\mathrm{d}\Phi)(\varphi(t))||} \in \mathbb{R}_{\geq 0} .$$
(57)

• The signed characteristic angle of  $\varphi$  at a regular and noncritical time  $t \in I_{\text{reg}} \cap I_{\text{noncrit}}$  is the characteristic angle of  $T_{\varphi(t)}$ , i.e. the signed angle of the tangent vector  $T_{\varphi}(t)$  relative to the unit gradient vector  $n_{\varphi}(t)$  (see Definition 3.12):

$$\theta_{\varphi}(t) \stackrel{\text{def.}}{=} \theta(T_{\varphi}(t)) = \theta_{n_{\varphi}(t)}(T_{\varphi}(t)) \in (-\pi, \pi] .$$

The first IR parameter equals the ratio between the kinetic and potential energy of  $\varphi$  and plays a crucial role in reference [21]. As we recall in Appendix B, this parameter carries the same information as the ordinary first slow roll parameter used in the physics literature. The conservative parameter is the ratio of the norms of the friction and gradient terms of the cosmological equation and controls what we call the quasi-conservative and strongly dissipative approximations (see Section 8).

24

4.1.2. Fundamental second order parameters. We next consider parameters which also depend on the covariant acceleration of  $\varphi$ . For this, we start by introducing a vector which compares the acceleration term of the cosmological equation to the norm of the friction term at a regular time  $t \in I_{\text{reg}}$ . The fundamental second order parameters will be defined as the components of this vector along  $T_{\varphi}(t)$  and  $N_{\varphi}(t)$ .

**Definition 4.2.** The rescaled opposite relative acceleration of a curve  $\varphi : I \to \mathcal{M}$  at the regular time  $t \in I_{\text{reg}}$  is the vector:

$$\hat{\boldsymbol{\eta}}_{\varphi}(t) \stackrel{\text{def.}}{=} -\frac{\nabla_t \dot{\varphi}(t)}{\mathcal{H}_{\varphi}(t) || \dot{\varphi}(t) ||} \in T_{\varphi(t)} \mathcal{M} .$$
(58)

Let

$$\Omega_{\varphi}(t) \stackrel{\text{def.}}{=} -\frac{\mathcal{G}(N_{\varphi}(t), \nabla_t \dot{\varphi}(t))}{||\dot{\varphi}(t)||} = -\mathcal{G}(N_{\varphi}(t), \nabla_t T_{\varphi}(t)) \; \forall t \in I_{\text{reg}}$$

be the signed turning rate of  $\varphi$  at the regular time t (see Appendix A.2).

Remark 4.3. Consider the following open submanifold of  $E^2 = J^2(\mathcal{M})$ :

$$J_{\text{reg}}^2(\mathcal{M}) \stackrel{\text{def.}}{=} \{ \phi \in J^2(\mathcal{M}) \mid \pi_{21}(\phi) \neq 0 \} ,$$

where  $\pi_{21} : J^2(\mathcal{M}) \to J^1(\mathcal{M})$  is the natural projection. Since the curve  $\varphi$  is arbitrary, we have:

$$\hat{\boldsymbol{\eta}}_{\varphi}(t) = \underline{\hat{\boldsymbol{\eta}}}(j^2(\varphi)(t)) \ \forall t \in I_{\text{reg}}$$

where  $\underline{\hat{\eta}} \in \Gamma(J^2_{\text{reg}}(\mathcal{M}), F^2)$  is a  $T\mathcal{M}$ -valued second order local observable in the sense of Subsection 2.3. When  $\varphi$  is a regular curve, we have  $j^2(\varphi)(I) \subset J^2_{\text{reg}}(\mathcal{M})$  and  $\hat{\eta}_{\varphi}$  is the evaluation of  $\underline{\hat{\eta}}$  along  $\varphi$ . In this case,  $\Omega_{\varphi}$  is the evaluation along  $\varphi$  of a second order scalar observable  $\underline{\Omega} \in \mathcal{C}^{\infty}(J^2_{\text{reg}}(\mathcal{M}))$ .

**Proposition 4.4.** For any curve  $\varphi : I \to \mathcal{M}$  and any  $t \in I_{reg}$ , we have:

$$\hat{\boldsymbol{\eta}}_{\varphi}(t) = -\frac{1}{\mathcal{H}_{\varphi}(t)} \frac{\mathrm{d}\log||\dot{\varphi}(t)||}{\mathrm{d}t} T_{\varphi}(t) + \frac{\Omega_{\varphi}(t)}{\mathcal{H}_{\varphi}(t)} N_{\varphi}(t) \in T_{\varphi(t)} \mathcal{M} .$$
(59)

*Proof.* Follows by projecting the left hand side on the vectors  $T_{\varphi}(t)$  and  $N_{\varphi}(t)$ .  $\Box$ 

**Definition 4.5.** The fundamental second order parameters of a curve  $\varphi : I \to \mathcal{M}$  are defined as follows:

• The rescaled second slow roll parameter of  $\varphi$  at a regular time  $t \in I_{\text{reg}}$  is the real number:

$$\hat{\eta}_{\varphi}^{\parallel}(t) \stackrel{\text{def.}}{=} \mathcal{G}(T_{\varphi}(t), \hat{\eta}_{\varphi}(t)) = -\frac{\mathrm{d}\log||\dot{\varphi}(t)||}{\mathcal{H}_{\varphi}(t)\mathrm{d}t} \in \mathbb{R} .$$
(60)

• The rescaled signed relative turning rate of  $\varphi$  at a regular time  $t \in I_{\text{reg}}$  is the real number:

$$\hat{\omega}_{\varphi}(t) \stackrel{\text{def.}}{=} \mathcal{G}(N_{\varphi}(t), \hat{\eta}_{\varphi}(t)) = \frac{\Omega_{\varphi}(t)}{\mathcal{H}_{\varphi}(t)} \in \mathbb{R} .$$
(61)

The fundamental second order local observables are the corresponding functions  $\hat{\eta}^{\parallel}, \hat{\underline{\omega}} \in \mathcal{C}^{\infty}(J^2_{\text{reg}}(\mathcal{M}))$  (see Remark 4.3).

We have:

$$\hat{\boldsymbol{\eta}}_{\varphi}(t) = \hat{\eta}_{\varphi}^{\parallel}(t)T_{\varphi}(t) + \hat{\omega}_{\varphi}(t)N_{\varphi}(t)$$

Relation (150) of Appendix A gives:

$$\hat{\omega}_{\varphi}(t) = -\frac{||\dot{\varphi}(t)||}{\mathcal{H}_{\varphi}(t)} \xi_{\varphi}(t) \; \forall t \in I_{\text{reg}} \; ,$$

where  $\xi_{\varphi}$  is the *signed* curvature of  $\varphi$  defined in Appendix A.2.

Let us extract the on-shell reduction of  $\underline{\hat{\eta}}^{\parallel}$  and  $\underline{\hat{\omega}}$  using the cosmological equation for  $\varphi$ . Projecting the cosmological equation (11) for a curve  $\varphi : I \to \mathcal{M}$  on the vectors  $T_{\varphi}(t)$  and  $N_{\varphi}(t)$  at a regular time  $t \in I_{\text{reg}}$  gives respectively the *adiabatic* equation:

$$\mathcal{G}(T_{\varphi}(t), \nabla_t \dot{\varphi}(t)) + \mathcal{H}(\dot{\varphi}(t)) || \dot{\varphi}(t) || + \Phi_T(\dot{\varphi}(t)) = 0 \ \forall t \in I_{\text{reg}}$$

and the *entropic equation*:

$$\mathcal{G}(N_{\varphi}(t), \nabla_t \dot{\varphi}(t)) + \Phi_N(\dot{\varphi}(t)) = 0 \ \forall t \in I_{\text{reg}} .$$

These equations can be written as:

$$\hat{\eta}^{\parallel}(t) = 1 + \frac{\Phi_T(\dot{\varphi}(t))}{\mathcal{H}(\dot{\varphi}(t))||\dot{\varphi}(t)||} , \ \hat{\omega}_{\varphi}(t) = \frac{\Phi_N(\dot{\varphi}(t))}{\mathcal{H}(\dot{\varphi}(t))||\dot{\varphi}(t)||} \ \forall t \in I_{\text{reg}} .$$
(62)

These relations show that the dynamical reductions of  $\hat{\eta}^{\parallel}$  and  $\underline{\hat{\omega}}$  are given by:

$$\underline{\hat{\eta}}^{\parallel^{\text{red}}} = 1 + \frac{\Phi_T}{\mathcal{H}\mathcal{N}} , \ \underline{\hat{\omega}}^{\text{red}} = \frac{\Phi_N}{\mathcal{H}\mathcal{N}} .$$
(63)

Together with Definition 4.1, this leads to the following.

**Definition 4.6.** The *fundamental basic observables* of the oriented scalar triple  $(\mathcal{M}, \mathcal{G}, \Phi)$  are the basic local scalar observables defined as follows:

• The first IR function of  $(\mathcal{M}, \mathcal{G}, \Phi)$  is the map  $\kappa : T\mathcal{M} \to \mathbb{R}_{\geq 0}$  defined through:

$$\kappa \stackrel{\text{def.}}{=} \frac{\mathcal{N}^2}{2\Phi} \ . \tag{64}$$

• The rescaled conservative function of  $(\mathcal{M}, \mathcal{G}, \Phi)$  is the map  $\hat{c} : T\mathcal{M}_0 \to \mathbb{R}_{\geq 0}$  defined through:

$$\hat{c} \stackrel{\text{def.}}{=} \frac{\mathcal{N}\mathcal{H}}{\Psi} = \frac{\mathcal{N}\sqrt{\mathcal{N}^2 + 2\Phi}}{M_0\Psi} \ . \tag{65}$$

• The signed characteristic angle function of  $(\mathcal{M}, \mathcal{G}, \Phi)$  is the map  $\theta : \dot{T}\mathcal{M}_0 \to (-\pi, \pi]$  defined through (see Definition 3.12 and Proposition 3.14):

$$\theta \stackrel{\text{def.}}{=} \theta_0(n^{\mathrm{v}}, T_0) \ . \tag{66}$$

• The rescaled second slow roll function of  $(\mathcal{M}, \mathcal{G}, \Phi)$  is the map  $\hat{\eta}^{\parallel} : \dot{T}\mathcal{M} \to \mathbb{R}$  defined through:

$$\hat{\eta}^{\parallel} \stackrel{\text{def.}}{=} 1 + \frac{\Phi_T}{\mathcal{HN}} \ . \tag{67}$$

• The rescaled turn rate function is the map  $\hat{\omega} : \dot{T}\mathcal{M} \to \mathbb{R}$  defined through:

$$\hat{\omega} = \frac{\Phi_N}{\mathcal{H}\mathcal{N}} \ . \tag{68}$$

Remark 4.7. Since  $\Phi$  is strictly positive, the function  $\hat{c}$  vanishes only on the zero section of  $T\mathcal{M}_0$  and hence is strictly positive on  $\dot{T}\mathcal{M}_0$ . This function tends to  $+\infty$  on the topological frontier of  $T\mathcal{M}_0$  in  $T\mathcal{M}$ , which coincides with the  $\pi$ -preimage of the critical locus:

$$\operatorname{Fr}(T\mathcal{M}_0) = \overline{T\mathcal{M}_0} \setminus T\mathcal{M}_0 = \pi^{-1}(\operatorname{Crit}\Phi) = \{u \in T\mathcal{M} \mid \pi(u) \in \operatorname{Crit}\Phi\}.$$

Also notice that the limits of  $\hat{\eta}^{\parallel}$  and  $\hat{\omega}$  at a point which lies on the zero section of  $T\mathcal{M}_0$  depend on the direction from which one approaches that point inside  $T\mathcal{M}_0$  and the directional limits equal plus or minus infinity when the directional derivatives of  $\Phi$  at that point with in that direction (respectively in its positive normal direction) are nonzero. **Proposition 4.8.** Suppose that  $\varphi: I \to \mathcal{M}$  is a cosmological curve. Then we have:

$$\kappa_{\varphi}(t) = \kappa(\dot{\varphi}(t)) \ \forall t \in I \tag{69}$$

and:

$$\hat{c}_{\varphi}(t) = \hat{c}(\dot{\varphi}(t)) \ \forall t \in I_{\text{noncrit}}$$

$$\tag{70}$$

as well as:

$$\hat{\eta}_{\varphi}^{\parallel}(t) = \hat{\eta}(\dot{\varphi}(t)) , \ \hat{\omega}_{\varphi}(t) = \hat{\omega}(\dot{\varphi}(t)) \ \forall t \in I_{\text{reg}}.$$

$$(71)$$

*Proof.* Relations (69) and (70) are obvious, while (71) follows from (63) and the results of Subsection 2.3.  $\Box$ 

The basic fundamental observables introduced above are not independent. Below, we discuss some obvious relations between them. More interesting relations will arise in the next section.

4.2. Relations between  $\mathcal{N}$ ,  $\mathcal{H}$ ,  $\hat{c}$  and  $\kappa$ . The basic observables  $\kappa$  and  $\hat{c}$  are norm reducible in the sense of Definition 2.5. We can express  $\mathcal{N}$  in terms of  $\kappa$  as:

$$\mathcal{N} = \sqrt{2\Phi\kappa} \,\,, \tag{72}$$

which gives:

$$\mathcal{H} = \frac{\sqrt{2\bar{\Phi}}}{M_0} (1+\kappa)^{1/2} .$$
 (73)

To express  $\hat{c}$  in terms of  $\kappa$ , we define:

**Definition 4.9.** The *characteristic one-form* of the model  $(M_0, \mathcal{M}, \mathcal{G}, \Phi)$  is the following one-form defined on  $\mathcal{M}$ :

$$\Xi \stackrel{\text{def.}}{=} \frac{M_0 \mathrm{d}\Phi}{2\Phi} \in \Omega^1(\mathcal{M}) \ . \tag{74}$$

**Proposition 4.10.** The following relations hold on  $T\mathcal{M}_0$ :

$$\mathcal{NH} = \Psi \hat{c} \tag{75}$$

$$\hat{c} = \frac{[\kappa(1+\kappa)]^{1/2}}{||\Xi||^{v}}$$
(76)

$$\kappa = \frac{1}{2} \left[ -1 + \sqrt{1 + 4(||\Xi||^{\mathbf{v}})^2 \hat{c}^2} \right] .$$
(77)

*Proof.* The first relation follows from the definition (65) of  $\hat{c}$ , while the second relation follows from the first using (72), (73) and the definition (74) of  $\Xi$ . The last relation follows by solving (76) for  $\kappa$ .

#### 5. NATURAL FIBERWISE COORDINATES

The natural basic observables introduced in the previous section allow us to define fiberwise coordinate systems on certain open subsets of  $T\mathcal{M}_0$ .

5.0.1. Fiberwise norm-angle coordinates. Let  $L_n^{\pm} \to \mathcal{M}_0$  be the closed half line bundles generated by the vector fields  $\pm n$ :

$$L^{+} \stackrel{\text{def.}}{=} \{ u \in T\mathcal{M}_{0} \mid \theta(u) = 0 \} = \sqcup_{m \in \mathcal{M}_{0}} \mathbb{R}_{\geq 0} n(m)$$
(78)

$$L^{-} \stackrel{\text{def.}}{=} \{ u \in T\mathcal{M}_0 \mid \theta(u) = \pi \} = \sqcup_{m \in \mathcal{M}_0} \mathbb{R}_{>0}(-n(m))$$
(79)

and notice that  $(\mathcal{N}, \theta)$  is a fiberwise coordinate system on  $T\mathcal{M}_0 \setminus L^-$  in the sense of Definition 1.11.

**Definition 5.1.** The pair  $(\mathcal{N}, \theta)$  is called the system of *fiberwise norm-angle coor*dinates on  $T\mathcal{M}_0 \setminus L^- \subset T\mathcal{M}_0$ .

These fiberwise coordinates define a polar coordinate system in each fiber of  $T\mathcal{M}_0$ .

5.0.2. Fiberwise conservative coordinates.

**Lemma 5.2.** The following relations hold on  $T\mathcal{M}_0$ :

$$\mathcal{N} = \sqrt{-\Phi + \sqrt{\Phi^2 + M_0^2 \Psi^2 \hat{c}^2}} .$$
 (80)

$$\mathcal{H} = \frac{1}{M_0} \left[ \Phi + \sqrt{\Phi^2 + M_0^2 \Psi^2 \hat{c}^2} \right] \,. \tag{81}$$

*Proof.* Follows by combining (77) with with (72) and (73).

**Proposition 5.3.** The pair of functions  $(\hat{c}, \theta)$  is a fiberwise coordinate system on  $T\mathcal{M}_0 \setminus L^-$ , which we call fiberwise conservative coordinates.

*Proof.* The values of  $\theta(u)$  and  $\hat{c}(u)$ , determine the direction and norm of the nonzero tangent vector  $u \in \dot{T}\mathcal{M}_0$  in each fiber of  $\dot{T}\mathcal{M}_0$ , where the second statement follows from equation (80).

Notice that  $\hat{c}$  can be expressed in terms of  $\mathcal{N}$  and  $\Phi, \Psi$  as:

$$\hat{c} = \frac{1}{M_0 \Psi} \sqrt{(\mathcal{N}^2 + \Phi)^2 - \Phi^2} = \frac{1}{M_0 \Psi} \sqrt{\mathcal{N}^2 (\mathcal{N}^2 + 2\Phi)} \; .$$

Corollary 5.4. The functions:

$$q_1 \stackrel{\text{def.}}{=} \hat{c} \cos \theta \ , \ q_2 \stackrel{\text{def.}}{=} \hat{c} \sin \theta \ .$$
 (82)

are fiberwise coordinates on  $\dot{T}\mathcal{U}$ , which we call fiberwise Cartesian conservative coordinates.

5.0.3. Fiberwise roll-turn coordinates.

**Lemma 5.5.** The following relations hold on  $\dot{T}\mathcal{M}_0$ :

$$\hat{\eta}^{\parallel} = 1 + \frac{\cos\theta}{\hat{c}} , \ \hat{\omega} = -\frac{\sin\theta}{\hat{c}}$$
(83)

$$(1 - \hat{\eta}^{\parallel})^2 + \hat{\omega}^2 = \frac{1}{\hat{c}^2}$$
(84)

$$\hat{c} = \frac{1}{\sqrt{(1-\hat{\eta}^{\parallel})^2 + \hat{\omega}^2}} , \ \sin\theta = -\frac{\hat{\omega}}{\sqrt{(1-\hat{\eta}^{\parallel})^2 + \hat{\omega}^2}} , \ \cos\theta = -\frac{1-\hat{\eta}^{\parallel}}{\sqrt{(1-\hat{\eta}^{\parallel})^2 + \hat{\omega}^2}} .$$
(85)

Moreover, we we have  $\operatorname{sign}(\hat{\omega}) = -\sigma$  and:

$$\mathcal{NH} = \frac{\Psi}{\sqrt{(1-\hat{\eta}^{\parallel})^2 + \hat{\omega}^2}} .$$
(86)

as well as:

$$\hat{\omega}\cos\theta = (1 - \hat{\eta}^{\parallel})\sin\theta . \tag{87}$$

Remark 5.6. Notice that  $\hat{\eta}^{\parallel}(u) = 1$  iff  $|\theta(u)| = \frac{\pi}{2}$  i.e. iff the non-zero vector  $u \in \dot{T}\mathcal{M}_0$  is orthogonal to the unit gradient vector  $n(\pi(u))$ . In this case, we have  $\hat{\omega}(u) = \pm \frac{1}{\hat{c}(u)} \neq 0$ .

*Proof.* Relations (83) follow from Proposition 3.20 and the definitions of  $\hat{\eta}^{\parallel}$  and  $\hat{\omega}$ . These relations give:

$$\cos\theta = -(1 - \hat{\eta}^{\parallel})\hat{c} , \ \sin\theta = -\hat{\omega}\hat{c} .$$
(88)

Using these equations, the identity  $1 = \cos^2 \theta + \sin^2 \theta$  implies (84) and (85). The second relation in (83) gives  $\operatorname{sign}(\hat{\omega}) = -\sigma$ , where we remind the reader that  $\sigma = \operatorname{sign}(\sin \theta)$ . Relation (75) and the first relation in (85) imply (86). The last two relations in (85) imply (87).

**Proposition 5.7.** The pair of functions  $(\hat{\eta}^{\parallel}, \hat{\omega})$  is a fiberwise coordinate system on  $\dot{T}\mathcal{M}_0$  in the sense of Definition 1.11, which we call fiberwise roll-turn coordinates.

*Proof.* The values of  $\hat{\eta}^{\parallel}(u)$  and  $\hat{\omega}(u)$  determine and are determined by those of  $\theta(u)$  and  $\hat{c}(u)$  by Lemma 5.5, so the conclusion follows from Proposition 5.3.

**Corollary 5.8.** The functions  $\zeta \stackrel{\text{def.}}{=} 1 - \hat{\eta}^{\parallel}$  and  $\hat{\omega}$  give fiberwise coordinates defined on  $\dot{T}\mathcal{M}_0$ , which we call modified roll-turn coordinates.

The relation between fiberwise roll-turn and conservative coordinates can be described geometrically as follows. Recall the Cartesian fiberwise conservative coordinates of Corollary 5.4.

**Proposition 5.9.** In any fiber of  $T\mathcal{M}_0$ , the point with Cartesian conservative coordinates  $(q_1, q_2)$  is the opposite of the inversion of the point with modified roll-turn coordinates  $(\zeta, \hat{\omega})$  with respect to the origin.

*Proof.* The quantities  $\hat{c} = \sqrt{q_1^2 + q_2^2}$  and  $\theta$  are the radial distance from the origin and polar angle in the  $(q_1, q_2)$  plane while  $\sqrt{\zeta^2 + \hat{\omega}^2}$  is the radial distance from the origin in the  $(\zeta, \hat{\omega})$  plane. Relations (85) are equivalent with:

$$q_1 = -\frac{\zeta}{\zeta^2 + \hat{\omega}^2} , \ q_2 = -\frac{\hat{\omega}}{\zeta^2 + \hat{\omega}^2} .$$
 (89)

Remark 5.10. In every fiber of  $T\mathcal{M}_0$ , the zero tangent vector corresponds to the origin of the  $(q_1, q_2)$  plane and to the point at infinity in the  $(\zeta, \hat{\omega})$  and  $(\hat{\eta}^{\parallel}, \hat{\omega})$  planes.

**Proposition 5.11.** The following relations hold on  $T\mathcal{M}_0$ :

$$\mathcal{N} = \sqrt{-\Phi + \sqrt{\Phi^2 + \frac{M_0^2 \Psi^2}{(1 - \hat{\eta}^{\parallel})^2 + \hat{\omega}^2}}}$$
(90)

$$\mathcal{H} = \frac{1}{M_0} \sqrt{\Phi + \sqrt{\Phi^2 + \frac{M_0^2 \Psi^2}{(1 - \hat{\eta}^{\parallel})^2 + \hat{\omega}^2}}} .$$
(91)

$$\kappa = \frac{1}{2} \left[ -1 + \sqrt{1 + \frac{4(||\Xi||^{\mathbf{v}})^2}{(1 - \hat{\eta}^{\parallel})^2 + \hat{\omega}^2}} \right] .$$
(92)

Moreover,  $\kappa, \hat{\omega}$  and  $\hat{\eta}^{\parallel}$  satisfy:

$$(1 - \hat{\eta}^{\parallel})^2 + \hat{\omega}^2 = \frac{(||\Xi||^{\mathbf{v}})^2}{\kappa(1+\kappa)} \text{ on } \dot{T}\mathcal{M}_0 .$$
 (93)

*Proof.* The first three relations follow from Lemma 5.2 and equation (77) using the first identity in (85) of Lemma 5.5. Relation (93) follows from (76) and (84).  $\Box$ 

Remark 5.12. Relation (93) shows that fixing  $\kappa, \hat{\eta}^{\parallel}$  and  $\hat{\omega}$  determines  $||\Xi|| = \frac{M_0}{2} ||d \log \Phi||$ . This gives an Eikonal equation for  $\log \Phi$  (with index of refraction  $\frac{2}{M_0} ||\Xi||$ ) on the connected Riemannian manifold  $(\mathcal{M}, \mathcal{G})$ , which can be used to determine the homothety class of  $\Phi$  when the critical locus is known.

5.0.4. Fiberwise slow roll coordinates. Consider the following open subsets of  $\dot{T}\mathcal{M}_0$ :

$$T^{\pm}\mathcal{M}_{0} \stackrel{\text{def.}}{=} \{ u \in \dot{T}\mathcal{M}_{0} \mid \sigma(u) = \pm 1 \} = \{ u \in \dot{T}\mathcal{M}_{0} \mid \operatorname{sign}\theta(u) = \pm 1 \}$$

where  $\sigma(u)$  is the characteristic signature of u (see Definition 3.12). We have:

$$T^{\pm}\mathcal{M}_0 = \{ u \in T\mathcal{M}_0 \mid \operatorname{sign}(q_1(u)) = \pm 1 \} ,$$

where  $(q_1, q_2)$  are the Cartesian conservative coordinates of Corollary 5.4. Moreover:

$$T^+\mathcal{M}_0 \cup T^-\mathcal{M}_0 = T\mathcal{M}_0 \setminus L$$
,

where  $L = L^+ \cup L^- \subset T\mathcal{M}_0$  is the line bundle generated by the vector field n. The sets  $T^+\mathcal{M}_0$  and  $T^-\mathcal{M}_0$  are the total spaces of fiber sub-bundle of  $T\mathcal{M}_0$  whose typical fiber is an open half-plane.

**Lemma 5.13.** The following relations hold on  $T^{\lambda}\mathcal{M}_0$  for each  $\lambda \in \{-1, 1\}$ :

$$\hat{\eta}^{\parallel} = 1 + \frac{\cos\theta}{\hat{c}} \ , \ \kappa = \frac{1}{2} \left[ -1 + \sqrt{1 + \frac{M_0^2 \Psi^2}{\Phi^2} \hat{c}^2} \right]$$
(94)

$$\hat{c} = \frac{2\Phi}{M_0\Psi}\sqrt{\kappa(1+\kappa)} , \ \theta = \lambda \arccos\left[\frac{2\Phi}{M_0\Psi}(\hat{\eta}^{\parallel} - 1)\sqrt{\kappa(1+\kappa)}\right] .$$
(95)

*Proof.* The first relation in (94) is the first equation in (83), while the second follows immediately from (77) and the definition (74) of  $\Xi$ . The first relation in (95) follows from (76). The second relation in (95) follows from the first relations in (94) and (95) and the fact that sign $\theta = \lambda$  on  $T^{\lambda}\mathcal{M}_0$ .

**Proposition 5.14.** For any  $\lambda \in \{-1, 1\}$ , the pair of functions  $(\kappa, \hat{\eta}^{\parallel})$  is a fiberwise coordinate system on  $T^{\lambda}\mathcal{M}_0$  in the sense of Definition 1.11, which we call fiberwise slow roll coordinates.

*Proof.* Follows immediately from Lemma 5.13 and Proposition 5.3.

*Remark* 5.15. Recall the Cartesian conservative coordinates of Corollary 5.4. Relations (94) can be written as:

$$\hat{\eta}^{\parallel} = 1 + \frac{q_1}{q_1^2 + q_2^2} \ , \ \kappa = \frac{1}{2} \left[ -1 + \sqrt{1 + \frac{M_0^2 \Psi^2}{\varPhi^2} (q_1^2 + q_2^2)} \right]$$

and define a branched double covering map  $\mathbb{R}^2 \to \mathbb{R}^2$  which is invariant under the transformation  $(q_1, q_2) \to (q_1, -q_2)$ . The ramification set is the line with equation  $q_2 = 0$ , while the branching locus is the line with equation  $\eta = 1$ . In each fiber of  $T\mathcal{M}_0$ , the ramification set corresponds to the intersection of L with that fiber, whose complement in the fiber is the union of the corresponding fibers of  $T^{\pm}\mathcal{M}_0$ .

Corollary 5.16. Let

$$h \stackrel{\text{def.}}{=} 2\sqrt{\kappa(1+\kappa)} : T\mathcal{M} \to \mathbb{R}_{\geq 0} .$$
(96)

For all  $\lambda \in \{-1,1\}$ , the pair  $(h, \hat{\eta}^{\parallel})$  is a system of fiberwise coordinates on  $T^{\lambda}\mathcal{M}_0$ , which we call modified slow roll coordinates.

*Proof.* Since  $\kappa$  is positive on  $T\mathcal{M}_0$ , relation (96) determines  $\kappa$  in terms of h as:

$$\kappa = \frac{1}{2} \left[ -1 + \sqrt{1 + h^2} \right] \; .$$

The conclusion follows using Proposition 5.14.

#### 6. NATURAL PHASE SPACE COORDINATES

Recall the notion of special local coordinate system on  $T\mathcal{M}$  introduced in Definition 1.15. In Sections 3 and 5, we showed that each of the pairs  $(v_n, v_\tau)$ ,  $(\mathcal{N}, \theta)$ ,  $(\hat{c}, \theta), (\hat{\eta}^{\parallel}, \hat{\omega}), (\kappa, \hat{\eta}^{\parallel})$  and  $(h, \hat{\eta}^{\parallel})$  is a system fiberwise coordinates on an open submanifold of  $T\mathcal{M}_0$ .

**Definition 6.1.** A special phase space coordinate system  $(x^1, x^2, x^3, x^4)$  defined on an open subset of  $T\mathcal{U}$  is called *natural* if  $(x^3, x^4) = (\Phi, \Psi)$ .

A natural coordinate system is completely specified by its two fiberwise coordinates. In the following, we consider the following natural phase space local coordinate systems:

- The phase space adapted coordinates  $(v_n, v_\tau, \Phi, \Psi)$  on neighborhoods of  $T\mathcal{U}$ , which already appeared in Section 3. These coordinates are fiberwise linear.
- The norm-angle coordinates  $(\mathcal{N}, \theta, \Phi, \Psi)$  on neighborhoods of  $\dot{T}\mathcal{U}$
- The conservative coordinates  $(c, \theta, \Phi, \Psi)$  on neighborhoods of  $T\mathcal{U}$
- The roll-turn coordinates  $(\hat{\eta}^{\parallel}, \hat{\omega}, \Phi, \Psi)$  on neighborhoods of  $\dot{T}\mathcal{U}$
- The slow roll coordinates  $(\kappa, \hat{\eta}^{\parallel}, \Phi, \Psi)$  on neighborhoods of  $T^{\pm}\mathcal{U}$
- The modified slow roll coordinates  $(h, \hat{\eta}^{\parallel}, \Phi, \Psi)$  on neighborhoods of  $T^{\pm}\mathcal{U}$ .

The cosmological semispray in adapted phase space coordinates was given in Corollary 3.31. Below, we give the components of S in norm-angle, conservative, roll-turn and modified slow roll coordinates. These follow by applying the corresponding changes of variables to components of S in adapted coordinates.

## 6.1. The cosmological semispray in norm-angle coordinates.

**Proposition 6.2.** The cosmological semispray has the following form in normangle coordinates:

$$S = S^{\mathcal{N}} \frac{\partial}{\partial \mathcal{N}} + S^{\theta} \frac{\partial}{\partial \theta} + S^{\Phi} \frac{\partial}{\partial \Phi} + S^{\Psi} \frac{\partial}{\partial \Psi} ,$$

where:

$$S^{\mathcal{N}} = -\Psi \cos \theta - \frac{\mathcal{N}\sqrt{\mathcal{N}^2 + 2\Phi}}{M_0}$$

$$S^{\theta} = \frac{\Psi \sin \theta}{\mathcal{N}} - \mu \mathcal{N} \cos \theta - \lambda \mathcal{N} \sin \theta$$

$$S^{\Phi} = \mathcal{N}\Psi \cos \theta \qquad (97)$$

$$S^{\Psi} = \frac{1}{2} \mathcal{N}(\mu \Psi \sin \theta + \Phi_{nn}^{\mathsf{v}} \cos \theta) .$$

*Proof.* Follows from Proposition 6.5 using the relations above.

## 6.2. The cosmological semispray in conservative coordinates.

**Proposition 6.3.** The restriction of the cosmological semispray to TU has the following form in conservative coordinates:

$$S = S^c \frac{\partial}{\partial c} + S^\theta \frac{\partial}{\partial \theta} + S^\Phi \frac{\partial}{\partial \Phi} + S^\Psi \frac{\partial}{\partial \Psi} \ ,$$

where:

$$S^{c} = -\frac{2}{M_{0}^{2}\Psi}\sqrt{\left(\Phi^{2} + M_{0}^{2}\Psi^{2}\hat{c}^{2}\right)\left(-\Phi + \sqrt{\Phi^{2} + M_{0}^{2}\Psi^{2}\hat{c}^{2}}\right)} - \frac{1}{2M_{0}\Psi^{2}}\sqrt{\Phi + \sqrt{\Phi^{2} + M_{0}^{2}\Psi^{2}\hat{c}^{2}}} \left[2\Psi^{2}\cos\theta + \left(-\Phi + \sqrt{\Phi^{2} + M_{0}^{2}\Psi^{2}\hat{c}^{2}}\right)(\mu\Psi\sin\theta + \Phi_{nn}^{v}\cos\theta)\right]}$$

$$S^{\theta} = \frac{\Psi\sin\theta}{\sqrt{-\Phi + \sqrt{\Phi^{2} + M_{0}^{2}\Psi^{2}\hat{c}^{2}}}} - (\lambda\sin\theta + \mu\cos\theta)\sqrt{-\Phi + \sqrt{\Phi^{2} + M_{0}^{2}\Psi^{2}\hat{c}^{2}}}$$

$$S^{\Phi} = \Psi\cos\theta\sqrt{-\Phi + \sqrt{\Phi^{2} + M_{0}^{2}\Psi^{2}\hat{c}^{2}}}$$

$$(98)$$

$$S^{\Psi} = \frac{1}{2}(\mu\Psi\sin\theta + \Phi_{nn}^{v}\cos\theta)\sqrt{-\Phi + \sqrt{\Phi^{2} + M_{0}^{2}\Psi^{2}\hat{c}^{2}}} .$$

# 6.3. The cosmological semispray in roll-turn coordinates.

**Proposition 6.4.** The restriction of the cosmological semispray to  $\dot{T}U$  has the following form in roll-turn coordinates:

$$S = S^{\hat{\eta}^{\parallel}} \frac{\partial}{\partial \hat{\eta}^{\parallel}} + S^{\hat{\omega}} \frac{\partial}{\partial \hat{\omega}} + S^{\Phi} \frac{\partial}{\partial \Phi} + S^{\Psi} \frac{\partial}{\partial \Psi} ,$$

where:

$$\begin{split} S^{\hat{\eta}^{\|}} &= -\frac{\Psi\hat{\omega}^{2}}{\sqrt{(1-\hat{\eta}^{\|})^{2}+\hat{\omega}^{2}}\sqrt{-\Phi + \sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\|})^{2}+\hat{\omega}^{2}}}}} + \hat{\omega}[\lambda\hat{\omega} + (1-\hat{\eta}^{\|})\mu]\sqrt{\frac{-\Phi + \sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\|})^{2}+\hat{\omega}^{2}}}{(1-\hat{\eta}^{\|})^{2}+\hat{\omega}^{2}}} \\ &+ 4|1-\hat{\eta}^{\|}|\Psi\sqrt{(1-\hat{\eta}^{\|})^{2} + \hat{\omega}^{2}}\sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\|})^{2}+\hat{\omega}^{2}}}}\sqrt{-\Phi + \sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\|})^{2}+\hat{\omega}^{2}}}} \\ &+ M_{0}\sqrt{\Phi + \sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\|})^{2}+\hat{\omega}^{2}}}}\left[2(1-\hat{\eta}^{\|})^{2}\Psi^{2} - (1-\hat{\eta}^{\|})(\mu\hat{\omega}\Psi + (1-\hat{\eta}^{\|})\Phi_{nn}^{*})\left(\Phi - \sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\|})^{2}+\hat{\omega}^{2}}}} \\ &+ M_{0}\sqrt{\Phi + \sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\|})^{2}+\hat{\omega}^{2}}}}\left[2(1-\hat{\eta}^{\|})^{2}\Psi^{2} - (1-\hat{\eta}^{\|})(\mu\hat{\omega}\Psi + (1-\hat{\eta}^{\|})\Phi_{nn}^{*})\left(\Phi - \sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\|})^{2}+\hat{\omega}^{2}}}} \\ &- \frac{(1-\hat{\eta}^{\|})\hat{\omega}\Psi}{M_{0}^{2}\Psi}\sqrt{(1-\hat{\eta}^{\|})^{2} + \hat{\omega}^{2}}}\sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\|})^{2} + \hat{\omega}^{2}}}}\sqrt{-\Phi + \sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\|})^{2} + \hat{\omega}^{2}}}}} \\ &- \frac{(1-\hat{\eta}^{\|})\hat{\omega}\Psi}{\sqrt{(1-\hat{\eta}^{\|})^{2} + \hat{\omega}^{2}}}\sqrt{-\Phi + \sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\|})^{2} + \hat{\omega}^{2}}}}}} + (1-\hat{\eta}^{\|})(\lambda\hat{\omega} + (1-\hat{\eta}^{\|})\mu)\sqrt{\frac{-\Phi + \sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\|})^{2} + \hat{\omega}^{2}}}}} \\ &+ \frac{\hat{\omega}(\mu\hat{\omega}\Psi + (1-\hat{\eta}^{\|})\Phi_{nn}^{*})}{2M_{0}\Psi^{2}}\sqrt{\Phi + \sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\|})^{2} + \hat{\omega}^{2}}}}} \left(\sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\|})^{2} + \hat{\omega}^{2}}} - \Phi - 2(1-\hat{\eta}^{\|})\Psi^{2}\right)} \\ S^{\Phi} = |1 - \hat{\eta}^{\|}|\Psi\sqrt{\frac{-\Phi + \sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\|})^{2} + \hat{\omega}^{2}}}{(1-\hat{\eta}^{\|})^{2} + \hat{\omega}^{2}}}}} . \tag{99}$$

# 6.4. The cosmological semispray in modified slow roll coordinates.

**Proposition 6.5.** The restriction of the cosmological semispray to  $T^+\mathcal{U}$  has the following form in modified slow roll coordinates:

$$S = S^{\hat{\eta}^{\parallel}} \frac{\partial}{\partial \hat{\eta}^{\parallel}} + S^{h} \frac{\partial}{\partial h} + S^{\varPhi} \frac{\partial}{\partial \varPhi} + S^{\varPsi} \frac{\partial}{\partial \varPsi} ,$$

32

$$\begin{split} S^{\dagger} &= -\frac{1}{2M_0 h \phi^4 \Psi^2 \sqrt{-1 + \sqrt{1 + h^2 \Psi^4}}} \Big[ 2M_0^2 \phi^{5/2} \Psi^2 + 2\phi^{3/2} \Psi \left( -1 + \sqrt{1 + h^2 \Psi^4} \right) \left( h^2 (1 - \bar{\eta}^{\parallel})^2 \Psi^2 - M_0^2 \phi^2 \right) \lambda - \\ &- (1 - \bar{\eta}^{\parallel}) \left( -4\phi^{5/2} - 4h^2 \sqrt{\phi} (-1 + \bar{\eta}^{\parallel} + \phi^2) \Psi^4 + 4\sqrt{\phi^5 (1 + h^2 \Psi^4)} \right) + \\ &+ 3(1 - \bar{\eta}^{\parallel}) h \Psi^2 \left( -\phi^{3/2} \sqrt{-h^2 (-1 + \bar{\eta}^{\parallel})^2 \Psi^2 + M_0^2 \Phi^2} + \sqrt{\phi^3 (1 + h^2 \Psi^4)} \right) - \\ &- h^2 \Psi^2 (-1 + \bar{\eta}^{\parallel})^2 \left( -\phi^{3/2} + \sqrt{\phi^3 (1 + h^2 \Psi^4)} \right) \Phi_{nn}^{\vee} \Big] \\ S^h &= \frac{\Psi^2}{2M_0 \sqrt{\frac{x^3 (1 + \sqrt{1 + h^2} (1 + h^2 \Psi^4)}{1 + h^2}}} \left[ -4\phi^2 \sqrt{\left( 1 + \sqrt{1 + h^2} \right) (1 + h^2 \Psi^4) \left( -1 + \sqrt{1 + h^2 \Psi^4} \right)} - \\ &- 2(-1 + \bar{\eta}^{\parallel}) h^2 \Psi^4 \sqrt{\left( 1 + \sqrt{1 + h^2} \right) \left( -1 + \sqrt{1 + h^2 \Psi^4} \right)} + \sqrt{\left( 1 + h^2 \right) \left( -1 + \sqrt{1 + h^2} \right) \left( 1 + \sqrt{1 + h^2 \Psi^4} \right)} + \\ &+ 2(-1 + \bar{\eta}^{\parallel}) \Psi^2 \left( \sqrt{\left( -1 + \sqrt{1 + h^2} \right) \left( 1 + \sqrt{1 + h^2 \Psi^4} \right)} + \sqrt{\left( 1 + h^2 \right) \left( -1 + \sqrt{1 + h^2 \Psi^4} \right)} \right) + \\ &+ \psi \left( \sqrt{-\left( 1 + \sqrt{1 + h^2} \right) \left( 1 + \sqrt{1 + h^2 \Psi^4} \right) \left( h^2 (-1 + \bar{\eta}^{\parallel})^2 \Psi^2 - M_0^2 \Phi^2 \right)} - \\ &- \sqrt{-\left( (1 + \sqrt{1 + h^2} \right) \left( 1 + \sqrt{1 + h^2 \Psi^4} \right) \left( h^2 (-1 + \bar{\eta}^{\parallel})^2 \Psi^2 - M_0^2 \Phi^2 \right)} + \\ &+ \Psi^2 \sqrt{-\left( -1 + \sqrt{1 + h^2} \right) \left( -1 + \sqrt{1 + h^2 \Psi^4} \right) \left( h^2 (-1 + \bar{\eta}^{\parallel})^2 \Psi^2 - \Phi^2 M_0^2 \right)} + \\ &+ \psi^2 \sqrt{-\left( -1 + \sqrt{1 + h^2} \right) \left( -1 + \sqrt{1 + h^2 \Psi^4} \right) \left( h^2 (-1 + \bar{\eta}^{\parallel})^2 \Psi^2 - \Phi^2 M_0^2 \right)} + \\ &+ \sqrt{\left( 1 + h^2 \right) \left( -1 + \sqrt{1 + h^2} \right) \left( -1 + \sqrt{1 + h^2 \Psi^4} \right) \left( h^2 (-1 + \bar{\eta}^{\parallel})^2 \Psi^2 - \Phi^2 M_0^2 \right)} \right) \mu + \\ &- (1 - \bar{\eta}^{\parallel}) \Phi \left( h^2 \Psi^2 \sqrt{\left( 1 + \sqrt{1 + h^2} \right) \left( -1 + \sqrt{1 + h^2 \Psi^4} \right)} - \sqrt{\left( -1 + \sqrt{1 + h^2} \right) \left( 1 + \sqrt{1 + h^2 \Psi^4} \right)} \right) - \\ &- \sqrt{\left( 1 + h^2 \right) \left( -1 + \sqrt{1 + h^2} \right) \left( 1 + \sqrt{1 + h^2 \Psi^4} \right) \left( 1 + \sqrt{1 + h^2 \Psi^4} \right)} \right) \Phi_{nn}^{\vee} \Big]} \right] S^{\Phi} = - \frac{h(1 - \bar{\eta}^{\parallel}) \Psi}{M_0} \sqrt{\frac{-1 + \sqrt{1 + h^2 \Psi^4}}{\Phi}}} \left( \sqrt{M_0^2 \Phi^2 - h^2 (1 - \bar{\eta}^{\parallel})^2 \Psi^2} - h(1 - \bar{\eta}^{\parallel}) \Phi_{nn}^{\vee} \right)}. \quad (100)$$

The formidable expressions given above in roll-turn and modified slow roll coordinates can be expanded to first order in various regions to extract much simpler expressions which can be used to perform, for example, a general study of the second order slow roll regime or of the slow roll-rapid turn regime.

## 7. Constant roll conditions and constant roll manifolds

The basic observable  $\kappa$  and  $\hat{\eta}^{\parallel}$  allow one to describe various loci in  $T\mathcal{M}$  which are relevant to the slow roll and constant roll approximations used in cosmology. Below, we give a geometric description of some of these regions.

**Definition 7.1.** The first slow roll condition is the condition  $\kappa(u) \ll 1$  on the tangent vector  $u \in T\mathcal{M}$ .

Since  $\kappa(u) = 0$  iff u = 0 (see eq. (72)), the first slow roll condition is satisfied iff u lies close to the image Z of the zero section of  $T\mathcal{M}$ . Hence Z plays the role of first slow roll manifold. Notice that a cosmological curve  $\varphi : I \to \mathcal{M}$  satisfies  $\dot{\varphi}(t) \in Z$  iff  $t \in I_{\text{sing}}$ .

**Definition 7.2.** The *(adiabatic) constant roll condition* with parameter  $C \in \mathbb{R}$  is the condition  $|\hat{\eta}^{\parallel}(u) - C| \ll 1$  on the nonzero tangent vector  $u \in \dot{T}\mathcal{M}$ .

The constant roll condition with parameter zero is called the *second slow roll condition*, while the constant roll condition with parameter one is called the *ultra slow roll condition*. The combination of the first and second slow roll conditions is called the *second order slow roll condition*. By (88), the second slow roll condition  $|\hat{\eta}^{\parallel}(u)| \ll 1$  holds for  $u \in \dot{T}\mathcal{M}_0$  iff  $\cos\theta(u) \approx -\hat{c}(u)$ , which in particular requires that u points towards decreasing values of  $\Phi$ .

**Definition 7.3.** We say that a cosmological curve  $\varphi : I \to \mathcal{M}$  satisfies the first slow (resp. second slow, ultra, constant with parameter C) roll condition at time  $t \in I$  if the tangent vector  $\dot{\varphi}(t) \in T_{\varphi(t)}\mathcal{M}$  satisfies the corresponding condition.

**Proposition 7.4.** The first slow roll condition  $\kappa \ll 1$  is equivalent on  $\dot{T}\mathcal{M}_0$  with:

$$\sqrt{(1-\hat{\eta}^{\parallel})^2 + \hat{\omega}^2 \gg 2||\Xi||^{\mathrm{v}}}$$
 (101)

Moreover, the second order slow roll conditions are equivalent on  $T\mathcal{M}_0$  with:

$$|\hat{\eta}^{\parallel}| \ll 1 \text{ and } \sqrt{1 + \hat{\omega}^2} \gg 2||\Xi||^{\mathrm{v}}$$
 (102)

*Proof.* Follows immediately from relation (92) of Proposition 5.11.  $\Box$ 

## 7.1. Constant roll manifolds.

**Definition 7.5.** The (adiabatic) constant roll manifold of  $(\mathcal{M}, \mathcal{G}, \Phi)$  at parameter C is the following set:

$$\mathcal{Z}(C) \stackrel{\text{def.}}{=} \{ u \in \dot{T}\mathcal{M}_0 \mid \hat{\eta}^{\parallel}(u) = C \} ,$$

where the overline denotes closure in  $T\mathcal{M}_0$ .

The constant roll manifolds at parameters  $\mathcal{Z}(0)$  and  $\mathcal{Z}(1)$  are called the *second* slow roll and ultra slow roll manifolds, respectively. The intersection of  $\mathcal{Z}(C)$  with a fiber of  $T\mathcal{M}_0$  is called the *constant roll curve* at parameter C of that fiber. For C = 0, 1 this is called respectively the slow roll curve or ultra slow roll curve.

**Lemma 7.6.** The following relations hold on  $\dot{T}\mathcal{M}_0$ , where in the second equality we assume  $\hat{\eta}^{\parallel} \neq 1$ :

$$\mathcal{N} = \sqrt{\Phi \left[ -1 + \sqrt{1 + 4\hat{c}^2(||\Xi||^{\nu})^2} \right]} = \sqrt{\Phi \left[ -1 + \sqrt{1 + \frac{4(||\Xi||^{\nu})^2}{(1 - \hat{\eta}^{\parallel})^2} \cos^2 \theta} \right]} .$$
(103)

*Proof.* The first equality follows from (80) and (74). The second follows the first and (83).  $\Box$ 

Recall the first relation in (83), which can be written as:

$$\cos\theta = \hat{c}(\hat{\eta}^{\parallel} - 1) . \tag{104}$$

**Proposition 7.7.** For any  $C \neq 1$ , the constant roll curve of parameter C in a fiber of  $T\mathcal{M}_0$  is closed simple curve which contains the origin of that fiber. Hence  $\mathcal{Z}(C)$  is a 3-manifold which contains the image of the zero section of  $T\mathcal{M}_0$ .

*Proof.* Setting  $\hat{\eta}^{\parallel} = C$  in Lemma 7.6 shows that the constant roll curve  $\Gamma_C(m)$  of parameter C in the fiber  $T_m \mathcal{M}$  (where  $m \in \mathcal{M}_0$ ) satisfies the implicit equation:

$$||u||^{2} = \Phi(m) \left[ -1 + \sqrt{1 + \frac{4||\Xi(m)||^{2}}{(C-1)^{2}} \cos^{2}\theta(u)} \right] \ (u \in T_{m}\mathcal{M})$$
(105)

and hence is a portion of the curve  $\Gamma'_C(m) \subset T_m \mathcal{M}$  which consists of all solutions of this equation. Since  $||u||^2 = v_n(u)^2 + v_\tau(u)^2$  and  $\theta$  is the signed angle between u and n(m), relation (105) is the equation of  $\Gamma'_C(m)$  in the polar coordinates associated to

the Cartesian coordinate system defined on  $T_m \mathcal{M}$  by  $v_n$  and  $v_{\tau}$ , whose coordinate axes are the lines generated by the vectors n(m) and  $\tau(m)$ . The curve  $\Gamma'_C(m)$ passes through the origin of  $T_m \mathcal{M}$  at  $\theta = \pm \frac{\pi}{2}$ . The equation is invariant under the transformations  $\theta \to -\theta$  and  $\theta \to \pi - \theta$  (with *u* unchanged). The second of these symmetries shows that  $\Gamma'_C(m) \setminus \{0\}$  has two connected components which lie on the two sides of the  $\tau$ -axis. The closure of each connected component (which is obtained by adding the origin) is a closed simple curve. Relation (104) gives:

$$\cos\theta(u) = \hat{c}(u)(C-1) ,$$

which implies:

$$\operatorname{sign}(\frac{\pi}{2} - |\theta(u)|) = \operatorname{sign}[\cos \theta(u)] = \operatorname{sign}(C - 1)$$

since c(u) > 0. This shows that  $\Gamma_C(m) \setminus \{0\}$  coincides with one of the connected components of  $\Gamma'_C(m) \setminus \{0\}$ , which implies that  $\Gamma_C(m)$  is a closed simple curve.  $\Box$ 

Remark 7.8. The constant roll curve  $\Gamma_C(m)$  degenerates to the line  $\mathbb{R}\tau(m)$  orthogonal to n(m) inside  $T_m\mathcal{M}$  in the ultra slow roll limit  $C \to 1$ . In this case, equation (90) gives:

$$||u|| = \sqrt{\Phi(m) \left[ -1 + \sqrt{1 + \frac{4||\Xi(m)||^2}{\hat{\omega}(u)^2}} \right]},$$

where  $\hat{\omega}(u) \in \mathbb{R}^{\times}$ . When  $\hat{\eta}^{\parallel}(u) \to \pm \infty$ , the curve degenerates to the origin of  $T_m \mathcal{M}$ .

Consider now the special case of slow roll and ultra-slow roll curves. Recall the fiberwise Cartesian conservative coordinates  $q_1 \stackrel{\text{def.}}{=} \hat{c} \cos \theta, q_2 \stackrel{\text{def.}}{=} \hat{c} \sin \theta : \dot{T} \mathcal{M}_0 \to \mathbb{R}$  of Corollary 5.4.


**Lemma 7.9.** We have  $\hat{\eta}^{\parallel} = 1$  in  $\dot{T}\mathcal{M}_0$  iff  $q_1 = 0$ . Moreover, we have  $\hat{\eta}^{\parallel} = 0$  in  $\dot{T}\mathcal{M}_0$  iff the value of  $(q_1, q_2)$  lies on the circle of radius 1/2 centered at the point (-1/2, 0) in the  $(q_1, q_2)$  plane. In this case, the position of  $(q_1, q_2)$  on this circle determines the value of  $\hat{\omega}$ .

*Proof.* The first statement follows immediately from eqs. (89) (where we remind the reader that  $\zeta \stackrel{\text{def.}}{=} 1 - \hat{\eta}^{\parallel}$ ). The same relations show that  $\hat{\eta}^{\parallel} = 0$  is equivalent with:

$$q_1 = -\frac{1}{1+\hat{\omega}^2}, \ q_2 = -\frac{\hat{\omega}}{1+\hat{\omega}^2}$$

with  $\omega \in \mathbb{R}$ . This implies  $\hat{\omega} = \frac{q_2}{q_1}$  and:

$$(q_1 + \frac{1}{2})^2 + q_2^2 = \frac{1}{4}$$
.



(A) Five points in the  $(\hat{\eta}^{\parallel}, \hat{\omega})$  plane. The green and red points lie on the slow roll and ultra slow roll curves (shown respectively in red and green).

(B) The image of the same five points in the  $(q_1, q_2)$  plane. In these coordinates, the slow roll curve (shown in red) becomes a circle while the ultra slow roll curve (shown in green) coincides with the vertical axis.

FIGURE 1. The slow roll and ultra slow roll curves in the  $(\hat{\eta}^{\parallel}, \hat{\omega})$  and  $(q_1, q_2)$  planes.

**Corollary 7.10.** The second slow roll manifold  $\mathcal{Z}(0)$  is a closed three-dimensional submanifold of  $T\mathcal{M}_0$  which intersects each fiber of  $T\mathcal{M}_0$  in a simple closed curve. This submanifold contains the image  $Z_0 \stackrel{\text{def.}}{=} Z \cap T\mathcal{M}_0$  of the zero section of  $T\mathcal{M}_0$ . The ultra slow roll manifold  $\mathcal{Z}(1)$  is a closed three-dimensional manifold of  $T\mathcal{M}$  which intersects each fiber along a line and is tangent to  $\mathcal{Z}(0)$  along  $Z_0$ .

## 7.2. Simplifications under slow roll.

7.2.1. Simplifications in the second slow roll regime.

**Proposition 7.11.** Suppose that  $|\hat{\eta}^{\parallel}| \ll 1$ . Then the following relations hold on  $\dot{T}\mathcal{M}_0$ :

$$\hat{\omega} \approx \tan \theta \approx -\sigma \sqrt{\frac{1}{\hat{c}^2} - 1}$$
 (106)

and:

$$\cos\theta \approx -\frac{1}{\sqrt{1+\hat{\omega}^2}} \approx -\hat{c} \ , \ \sin\theta \approx -\frac{\hat{\omega}}{\sqrt{1+\hat{\omega}^2}} \approx -\hat{\omega}\hat{c} \approx \sigma\sqrt{1-\hat{c}^2} \ . \tag{107}$$

In particular, we have  $\cos \theta < 0$  and hence a vector  $u \in \dot{T}\mathcal{M}_0$  which satisfies  $|\hat{\eta}^{\parallel}(u)| \ll 1$  points towards decreasing values of  $\Phi$ . Moreover, we have  $|\theta(u)| \to \frac{\pi}{2}$  in the rapid turn limit  $|\hat{\omega}| \to \infty$ .

*Proof.* Follows immediately from Lemma 5.5.  $\Box$ 

Hence in the second slow roll regime the reduced turning rate  $\hat{\omega}_{\varphi}(t)$  of a noncritical cosmological curve  $\varphi: I \to \mathcal{M}_0$  at a regular time  $t \in I_{\text{reg}}$  determines and is determined by the signed characteristic angle  $\theta_{\varphi}(t)$  and in this regime we have  $|\theta_{\varphi}(t)| > \frac{\pi}{2}$ , i.e. the tangent vector to the curve at time t points towards decreasing values of  $\Phi$ . **Proposition 7.12.** Suppose that  $|\hat{\eta}^{\parallel}| \ll 1$ . Then the following relations hold on  $\dot{T}\mathcal{M}_0$ :

$$\mathcal{N} \approx \sqrt{-\Phi + \sqrt{\Phi^2 + \frac{M_0^2 \Psi^2}{1 + \hat{\omega}^2}}} = \sqrt{-\Phi + \sqrt{\Phi^2 + M_0^2 \Psi^2 \cos^2 \theta}}$$
(108)  
$$\mathcal{H} \approx \frac{1}{M_0} \sqrt{\Phi + \sqrt{\Phi^2 + \frac{M_0^2 \Psi^2}{1 + \hat{\omega}^2}}} = \frac{1}{M_0} \sqrt{\Phi + \sqrt{\Phi^2 + M_0^2 \Psi^2 \cos^2 \theta}}$$

and:

$$\mathcal{NH} \approx \frac{\Psi}{\sqrt{1+\hat{\omega}^2}} \approx -\Psi\cos\theta$$
 (109)

Moreover, we have:

$$\begin{split} v_n &\approx -\frac{1}{\sqrt{1+\hat{\omega}^2}}\sqrt{-\varPhi + \sqrt{\varPhi^2 + \frac{M_0^2 \Psi^2}{1+\hat{\omega}^2}}} = \sqrt{-\varPhi + \sqrt{\varPhi^2 + M_0^2 \Psi^2 \cos^2 \theta}} \cos \theta \\ v_\tau &\approx -\frac{\hat{\omega}}{\sqrt{1+\hat{\omega}^2}}\sqrt{-\varPhi + \sqrt{\varPhi^2 + \frac{M_0^2 \Psi^2}{1+\hat{\omega}^2}}} = \sqrt{-\varPhi + \sqrt{\varPhi^2 + M_0^2 \Psi^2 \cos^2 \theta}} \sin \theta \ . \end{split}$$

Remark 7.13. Notice that in the second slow roll regime we have  $\cos \theta \approx -\hat{c}$  and  $\sin \theta = \sigma \sqrt{1 - \hat{c}^2}$ . This allows us to express the quantities in Proposition 7.12 in terms of  $\hat{c}$ .

*Proof.* Propositions 5.11 and 7.11 imply relations (108), while (86) implies (109). The last two relations in the statement follow from the first equation in (108) by using (39) and (107).  $\Box$ 

7.2.2. Simplifications in the second order slow roll regime. The second equation in (108) of Proposition 7.12 shows that, in the second slow roll regime, the first slow roll condition  $\kappa \ll 1$  (which is equivalent with  $\mathcal{H} \approx \frac{\sqrt{2\Phi}}{M_0}$ ) amounts to the requirement:

$$-M_0 \frac{\Psi}{\Phi} \cos \theta \ll 1 \iff -2||\Xi||^{\mathsf{v}} \cos \theta \ll 1 .$$

Let:

$$V = M_0 \sqrt{2\Phi}$$

be the classical effective potential of the model  $(M_0, \mathcal{M}, \mathcal{G}, \Phi)$  (see reference [21]).

**Proposition 7.14.** In the second order slow roll regime  $\kappa \ll 1$  and  $|\hat{\eta}^{\parallel}| \ll 1$ , we have:

$$\mathcal{N} \approx -||\mathrm{d}V||^{\mathrm{v}} \cos\theta \approx ||\mathrm{d}V||^{\mathrm{v}} \hat{c} , \qquad (110)$$

and:

$$v_n \approx -||\mathrm{d}V||^{\mathrm{v}} \cos^2 \theta , \ v_\tau \approx -||\mathrm{d}V||^{\mathrm{v}} \cos \theta \sin \theta .$$
 (111)

Proof. Relation (109) gives:

$$\mathcal{N} \approx -\frac{\Psi}{\mathcal{H}}\cos\theta$$
 (112)

Since we also impose the first slow roll condition, we have  $\mathcal{H} \approx \frac{\sqrt{2\Phi}}{M_0}$  and (112) becomes (110). Combining this with (39) gives (111).

#### CALIN IULIU LAZAROIU

#### 8. The quasi-conservative and strongly dissipative regimes

**Definition 8.1.** The quasi-conservative condition on a vector  $u \in T\mathcal{M}_0$  is the condition  $c(u) \ll 1$ . The strong dissipation condition on u is the condition  $c(u) \gg 1$ .

**Definition 8.2.** We say that a noncritical cosmological curve  $\varphi : I \to \mathcal{M}_0$  satisfies the quasi-conservative or strong dissipation condition at time  $t \in I$  when  $c_{\varphi}(t) = c(\dot{\varphi}(t))$  is much smaller (respectively much larger) than one.

The cosmological equation (11) can be written as:

$$\nabla_t \dot{\varphi}(t) + ||\mathrm{d}\Phi|| \left[ c_{\varphi}(t) T_{\varphi}(t) + n_{\varphi}(t) \right] = 0 \; .$$

When  $\varphi$  satisfies the quasi-conservative condition at time  $t_0$ , this can be approximated by the *conservative equation*:

$$\nabla_t \dot{\varphi}(t) + (\operatorname{grad}\Phi)(\varphi(t)) = 0$$

for t close to  $t_0$ . The conservative equation describes motion of a particle in the Riemannian manifold  $(\mathcal{M}, \mathcal{G})$  under the influence of the potential  $\Phi$  (which corresponds to a conservative dynamical system). When  $\varphi$  satisfies the strong dissipation condition at time  $t_0$ , the cosmological equation can be approximated by the *modified* geodesic equation:

$$\nabla_t \dot{\varphi}(t) + \mathcal{H}_{\varphi}(t) \dot{\varphi}(t) = 0$$

for t close to  $t_0$ . The solutions of this equation are reparameterized geodesics of  $(\mathcal{M}, \mathcal{G})$ . Some more detail about the quasi-conservative and strongly dissipative approximations can be found in [23].

Remark 8.3. Recall equation (84):

$$(\hat{\eta}^{\parallel} - 1)^2 + \hat{\omega}^2 = \frac{1}{\hat{c}^2} .$$
(113)

This shows that the rapid turn condition  $|\hat{\omega}| \gg 1$  is equivalent with  $\frac{1}{\hat{c}} \gg \sqrt{1 + (\hat{\eta}^{\parallel} - 1)^2}$ , which reduces to the quasi-conservative condition  $\hat{c} \ll 1$  in the second slow roll regime  $|\hat{\eta}^{\parallel}| \ll 1$ . Hence the rapid turn and quasi-conservative limits are equivalent within the second slow roll regime. Recall that the condition  $\hat{\eta}^{\parallel} \approx 1$  defines the adiabatic ultra slow roll regime. In this case, we have  $\hat{\omega} \approx -\frac{\sigma}{\hat{c}} \neq 0$  and  $\theta \approx \sigma \frac{\pi}{2}$ .

The first relation in (83) requires:

$$\hat{\eta}^{\parallel} \in \left[1 - \frac{1}{\hat{c}}, 1 + \frac{1}{\hat{c}}\right]$$

and can be written as (104). In particular, the interval within which  $\hat{\eta}^{\parallel}$  can take values is centered at 1 and constrained by the value of  $\hat{c}$ . This interval is very large in the quasi-conservative regime  $\hat{c} \ll 1$  and becomes narrow in the strongly dissipative regime  $\hat{c} \gg 1$ , when  $\hat{\eta}^{\parallel}$  is forced to be close to one. In particular, the adiabatic ultra slow roll approximation  $\hat{\eta}^{\parallel} \approx 1$  is accurate in the strongly dissipative regime.

## 9. The horizontal approximation in adapted coordinates

In this section, we study the horizontal approximation defined by the adapted local coordinates on a neighborhood of a point of  $T\mathcal{U}$ , where  $\mathcal{U}$  is the non-degenerate set of  $(\mathcal{M}, \mathcal{G}, \Phi)$ .

# 9.1. The adapted speed component functions.

**Lemma 9.1.** The following relations hold on  $T\mathcal{M}_0$ :

$$v_n = -\frac{\Psi}{\mathcal{H}} \frac{1 - \hat{\eta}^{\|}}{(1 - \hat{\eta}^{\|})^2 + \hat{\omega}^2} , \ v_\tau = -\frac{\Psi}{\mathcal{H}} \frac{\hat{\omega}}{(1 - \hat{\eta}^{\|})^2 + \hat{\omega}^2} .$$
(114)

and:

$$\kappa = \frac{1}{2\Phi} \left(\frac{\Psi}{\mathcal{H}}\right)^2 \frac{1}{(1-\hat{\eta}^{\parallel})^2 + \hat{\omega}^2} .$$
(115)

*Proof.* Combining (39) with (85) and (86) gives (114), which implies (115) upon using  $\kappa = \frac{N^2}{2\Phi} = \frac{v_n^2 + v_\tau^2}{2\Phi}$ .

**Proposition 9.2.** The following relations hold on  $T\mathcal{M}_0$ :

$$v_{n} = -\frac{M_{0}\Psi}{\sqrt{\Phi + \sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\parallel})^{2} + \hat{\omega}^{2}}}}} \frac{1-\hat{\eta}^{\parallel}}{(1-\hat{\eta}^{\parallel})^{2} + \hat{\omega}^{2}}$$
$$v_{\tau} = -\frac{M_{0}\Psi}{\sqrt{\Phi + \sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1-\hat{\eta}^{\parallel})^{2} + \hat{\omega}^{2}}}}} \frac{\hat{\omega}}{(1-\hat{\eta}^{\parallel})^{2} + \hat{\omega}^{2}} .$$
(116)

*Proof.* Equations (114) give (116) upon using Proposition 5.11.

9.2. The adapted parameters and adapted functions. Let:

 $Q \stackrel{\text{def.}}{=} \{ u \in T\mathcal{M}_0 \mid v_n(u)v_\tau(u) = 0 \} = \{ u \in T\mathcal{M}_0 \mid \theta(u) \notin \{-\frac{\pi}{2}, 0, \frac{\pi}{2}, \pi\} \} = T\mathcal{M}_0 \setminus (L_n \cup L_\tau) \}$ 

and notice that  $Z \subset Q$ . Here  $L_n$  and  $L_{\tau}$  are the line bundles generated by the vector fields n and  $\tau$  on  $\mathcal{M}_0$ .

**Definition 9.3.** A noncritical curve  $\varphi : I \to \mathcal{M}_0$  is called *admissible* if its canonical lift  $\dot{\varphi} : I \to T\mathcal{M}_0$  does not intersect Q.

Notice that any admissible curve is nondegenerate.

**Definition 9.4.** The *adapted parameters* of an admissible curve  $\varphi : I \to \mathcal{M}_0$  are defined through:

$$f_{\varphi}^{n} \stackrel{\text{def.}}{=} -\frac{\dot{v}_{n}^{\varphi}}{\mathcal{H}_{\varphi} v_{n}^{\varphi}} \text{ and } f_{\varphi}^{r} \stackrel{\text{def.}}{=} -\frac{\dot{v}_{\tau}^{\varphi}}{\mathcal{H}_{\varphi} v_{\tau}^{\varphi}} .$$
(117)

**Definition 9.5.** The *adapted functions*  $f^n, f^r : T\mathcal{M}_0 \setminus Q \to \mathbb{R}$  of  $(\mathcal{M}, \mathcal{G}, \Phi)$  are the basic local scalar observables defined through:

$$f^{n} \stackrel{\text{def.}}{=} -\frac{S^{v_{n}}}{\mathcal{H}v_{n}} = 1 + \frac{\Psi - \lambda^{v}v_{\tau}^{2} - \mu^{v}v_{n}v_{\tau}}{\mathcal{H}v_{n}} = 1 + \frac{\Psi}{\mathcal{H}\mathcal{N}\cos\theta} - \frac{\mathcal{N}(\lambda^{v}\sin\theta + \mu^{v}\cos\theta)}{\mathcal{H}}\tan\theta$$

$$f^{\tau} \stackrel{\text{def.}}{=} -\frac{S^{v_{\tau}}}{\mathcal{H}v_{\tau}} = 1 + \frac{\mu^{v}v_{n}^{2} + \lambda^{v}v_{n}v_{\tau}}{\mathcal{H}v_{\tau}} = 1 + \frac{\mathcal{N}(\lambda^{v}\sin\theta + \mu^{v}\cos\theta)}{\mathcal{H}\tan\theta}, \qquad (118)$$

where  $S^{v_n}$  and  $S^{v_{\tau}}$  are the vertical components of the cosmological semispray in adapted phase space coordinates (see Corollary 3.31).

**Proposition 9.6.** Suppose that  $\varphi$  is an admissible cosmological curve. Then we have:

$$\mathfrak{f}_{\varphi}^n=\mathfrak{f}^n\circ\dot{\varphi}$$
 and  $\mathfrak{f}_{\varphi}^{\tau}=\mathfrak{f}^{\tau}\circ\dot{\varphi}$  .

Proof. Follows immediately from Lemma 3.28.

9.3. Relation to roll-turn fiberwise coordinates.

**Proposition 9.7.** The following relations hold on  $T\mathcal{M}_0 \setminus Q$ :

$$f^n \cos^2 \theta + f^\tau \sin^2 \theta = 1 + \frac{\cos \theta}{\hat{c}} = \hat{\eta}^{\parallel} .$$
(119)

and:

$$\hat{\eta}^{\parallel} = \frac{\left[1 + 2(\mathfrak{f}^n - \mathfrak{f}^{\tau})\hat{c}^2\right] \pm \sqrt{\left[1 + 2(\mathfrak{f}^n - \mathfrak{f}^{\tau})\hat{c}^2\right]^2 - 4(\mathfrak{f}^n - \mathfrak{f}^{\tau})\hat{c}^2[\mathfrak{f}^{\tau} + (\mathfrak{f}^n - \mathfrak{f}^{\tau})\hat{c}^2]}{2(\mathfrak{f}^n - \mathfrak{f}^{\tau})\hat{c}^2} .$$
(120)

*Proof.* Relation (119) follows immediately from (118) upon using (75) and the first relation in (83). Equation (104) implies  $\sin^2 \theta = 1 - \cos^2 \theta = 1 - \hat{c}^2 (\hat{\eta}^{\parallel} - 1)^2$ . Thus (119) is equivalent with:

$$\hat{\eta}^{\|} = \mathfrak{f}^{\tau} + (\mathfrak{f}^{n} - \mathfrak{f}^{\tau})(\hat{\eta}^{\|} - 1)^{2} \hat{c}^{2} , \qquad (121)$$

which amounts to a quadratic equation for  $\hat{\eta}^{\parallel} \colon$ 

$$(\mathbf{f}^{n} - \mathbf{f}^{\tau})\hat{c}^{2}(\hat{\eta}^{\parallel})^{2} - [1 + 2(\mathbf{f}^{n} - \mathbf{f}^{\tau})\hat{c}^{2}]\hat{\eta}^{\parallel} + \mathbf{f}^{\tau} + (\mathbf{f}^{n} - \mathbf{f}^{\tau})\hat{c}^{2} = 0.$$

The solutions of this equation are (120).

*Remark* 9.8. When  $|\mathfrak{f}^n - \mathfrak{f}^{\tau}| \cos^2 \theta \ll |\mathfrak{f}^{\tau}|$ , relation (120) reduces to:

$$\hat{\eta}^{\parallel} \approx \mathbf{f}^{\tau}$$

as follows from (121).

**Proposition 9.9.** The following relations hold on  $T\mathcal{M}_0 \setminus Q$ :

$$f^{n} = \hat{\eta}^{\parallel} - \frac{\hat{\omega}^{2}}{1 - \hat{\eta}^{\parallel}} + \frac{1}{\mathcal{H}^{2}} \frac{\hat{\omega}}{1 - \hat{\eta}^{\parallel}} \frac{\Phi^{v}_{\tau\tau}\omega + \Phi^{v}_{n\tau}(1 - \hat{\eta}^{\parallel})}{(1 - \hat{\eta}^{\parallel})^{2} + \hat{\omega}^{2}}$$

$$f^{\tau} = 1 - \frac{1}{\mathcal{H}^{2}} \frac{1 - \hat{\eta}^{\parallel}}{\hat{\omega}} \frac{\Phi^{v}_{\tau\tau}\omega + \Phi^{v}_{n\tau}(1 - \hat{\eta}^{\parallel})}{(1 - \hat{\eta}^{\parallel})^{2} + \hat{\omega}^{2}} , \qquad (122)$$

where:

$$\mathcal{H} = \frac{1}{M_0} \sqrt{\Phi + \sqrt{\Phi^2 + \frac{M_0^2 \Psi^2}{(1 - \hat{\eta}^{\parallel})^2 + \hat{\omega}^2}}} .$$
(123)

Moreover, we have:

$$\hat{\eta}^{\parallel} = \frac{(1 - \hat{\eta}^{\parallel})^2 \mathfrak{f}^n + \hat{\omega}^2 \mathfrak{f}^\tau}{(1 - \hat{\eta}^{\parallel})^2 + \hat{\omega}^2} .$$
(124)

*Proof.* Using relation (75) and Lemma 5.5, we compute:

$$1 + \frac{\Psi}{\mathcal{HN}\cos\theta} = \frac{1}{\hat{c}\cos\theta} = 1 + \frac{(1-\hat{\eta}^{\|})^2 + \hat{\omega}^2}{\hat{\eta}^{\|} - 1} = \hat{\eta}^{\|} + \frac{\hat{\omega}^2}{\hat{\eta}^{\|} - 1}$$
(125)

and:

$$-\frac{\mathcal{N}(\lambda^{\mathrm{v}}\sin\theta + \mu^{\mathrm{v}}\cos\theta)}{\mathcal{H}} = \frac{\mathcal{N}}{\mathcal{H}}\frac{\lambda^{\mathrm{v}}\hat{\omega} + \mu^{\mathrm{v}}(1-\hat{\eta}^{\parallel})}{\sqrt{(1-\hat{\eta}^{\parallel})^{2} + \hat{\omega}^{2}}} = \frac{1}{\mathcal{H}^{2}}\frac{\Phi_{\tau\tau}^{\mathrm{v}}\hat{\omega} + \Phi_{n\tau}^{\mathrm{v}}(1-\hat{\eta}^{\parallel})}{(1-\hat{\eta}^{\parallel})^{2} + \hat{\omega}^{2}}$$
(126)

where we used (29) and the relation:

$$\frac{\mathcal{N}}{\mathcal{H}} = \frac{\Psi}{\mathcal{H}^2} \hat{c} = \frac{\Psi}{\mathcal{H}^2} \frac{1}{\sqrt{(1-\hat{\eta}^{\parallel})^2 + \hat{\omega}^2}} ,$$

which follows from (75) and Lemma 5.5. The latter also gives:

$$\tan \theta = \frac{\hat{\omega}}{1 - \hat{\eta}^{\parallel}} . \tag{127}$$

Using this, (125) and (126) in the definition of  $\mathfrak{f}^n$  gives the first equation in (122). The second equation in (122) follows from (126), (127) and the definition of  $\mathfrak{f}^\tau$ . Relation (123) was proved before (see (91)). Relation (124) follows from (119) by using Lemma 5.5.

Remark 9.10. Suppose that  $\mathfrak{f}^n$  and  $\mathfrak{f}^\tau$  are fixed and view (122) as a system of equations for  $\hat{\eta}^{\parallel}$  and  $\hat{\omega}$ , where  $\mathcal{H}$  is expressed in terms of the latter using (123). Then this system is equivalent with the equation:

$$f^{n} = \hat{\eta}^{\parallel} - \frac{\hat{\omega}^{2}}{1 - \hat{\eta}^{\parallel}} + M_{0}^{2} \frac{\hat{\omega}}{1 - \hat{\eta}^{\parallel}} \frac{\Phi_{\tau\tau}^{v} \omega + \Phi_{n\tau}^{v} (1 - \hat{\eta}^{\parallel})}{\left[(1 - \hat{\eta}^{\parallel})^{2} + \hat{\omega}^{2}\right] \left[\Phi + \sqrt{\Phi^{2} + \frac{M_{0}^{2}\Psi^{2}}{(1 - \hat{\eta}^{\parallel})^{2} + \hat{\omega}^{2}}}\right]} \quad (128)$$

(which is obtained using (123) from the first equation in (122)) together with equation (124), which can be written as:

$$\hat{\omega}^{2} = -(1 - \hat{\eta}^{\parallel})^{2} \frac{\mathfrak{f}^{n} - \hat{\eta}^{\parallel}}{\mathfrak{f}^{\tau} - \hat{\eta}^{\parallel}}$$
(129)

and can be used to eliminate  $\hat{\omega}$  in terms of  $\hat{\eta}^{\parallel}$ . Substituting this in (128) gives an algebraic equation for  $\hat{\eta}^{\parallel}$  whose coefficients depend on  $\mathfrak{f}^n$  and  $\mathfrak{f}^{\tau}$ . Picking a solution of this algebraic equation, relation (129) determines  $\hat{\omega}$  as a function of the adapted parameters.

**Corollary 9.11.** The intersection of the second slow roll manifold  $\mathcal{Z}(0)$  with  $T\mathcal{M}_0 \setminus Q$  has implicit equations:

$$\mathbf{f}^{n} = -\hat{\omega}^{2} \mathbf{f}^{\tau} , \ \mathbf{f}^{\tau} = 1 - \frac{M_{0}^{2}}{\hat{\omega} \left[ \Phi + \sqrt{\Phi^{2} + \frac{M_{0}^{2} \Psi^{2}}{1 + \hat{\omega}^{2}}} \right]} \frac{\Phi_{\tau\tau}^{\mathbf{v}} \hat{\omega} + \Phi_{n\tau}^{\mathbf{v}}}{1 + \hat{\omega}^{2}} .$$
(130)

where  $\hat{\omega} \in \mathbb{R}^{\times}$ .

**Corollary 9.12.** The following relations hold on  $T\mathcal{M}_0 \setminus Q$  in the slow roll approximation  $|\hat{\eta}^{\parallel}| \ll 1$ :

$$\mathfrak{f}^n \approx -\hat{\omega}^2 + \frac{\hat{\omega}}{\mathcal{H}^2} \frac{\Phi^{\mathsf{v}}_{\tau\tau} \hat{\omega} + \Phi^{\mathsf{v}}_{n\tau}}{1 + \hat{\omega}^2} , \ \mathfrak{f}^\tau \approx -\frac{\mathfrak{f}^n}{\hat{\omega}^2} = 1 - \frac{1}{\mathcal{H}^2 \hat{\omega}} \frac{\Phi^{\mathsf{v}}_{\tau\tau} \hat{\omega} + \Phi^{\mathsf{v}}_{n\tau}}{1 + \hat{\omega}^2} , \qquad (131)$$

where:

$$\mathcal{H} \approx \frac{1}{M_0} \sqrt{\Phi + \sqrt{\Phi^2 + \frac{M_0^2 \Psi^2}{1 + \hat{\omega}^2}}}$$
 (132)

Moreover, we have:

$$\hat{\eta}^{\parallel} \approx \frac{\mathfrak{f}^n + \hat{\omega}^2 \mathfrak{f}^\tau}{1 + \hat{\omega}^2} . \tag{133}$$

**Corollary 9.13.** The following relations hold on  $T\mathcal{M}_0 \setminus Q$  in the second order slow roll approximation  $\kappa \ll 1$  and  $|\hat{\eta}^{\parallel}| \ll 1$ :

$$f^{n} \approx -\hat{\omega}^{2} + \frac{M_{0}^{2}}{2\Phi}\hat{\omega}\frac{\Phi_{\tau\tau}^{v}\omega + \Phi_{n\tau}^{v}}{1 + \hat{\omega}^{2}} , \ f^{\tau} \approx -\frac{f^{n}}{\hat{\omega}^{2}} = 1 - \frac{M_{0}^{2}}{2\Phi\hat{\omega}}\frac{\Phi_{\tau\tau}^{v}\omega + \Phi_{n\tau}^{v}}{1 + \hat{\omega}^{2}} .$$
(134)

*Proof.* Follows from Corollary 9.12 upon recalling that the first slow roll condition is equivalent with  $\mathcal{H} \approx \frac{\sqrt{2\Phi}}{M_0}$ .

9.4. The adapted manifold.

**Definition 9.14.** The *adapted manifold* is the submanifold  $\mathcal{Z}_a$  of  $T\mathcal{M}_0 \setminus Q$  defined by the *strict adapted conditions*:

$$\mathbf{f}^n = \mathbf{f}^\tau = 0 \ . \tag{135}$$

Let  $W_a$  be the flat Ehresmann connection defined by the adapted phase space coordinates in the vicinity of a point of  $TU \setminus Q$ , which we shall call the adapted connection. Definition 9.5 shows that the adapted manifold coincides with the mean field manifold of S relative to  $W_a$ , which has equations  $S^{v_n} = S^{v_\tau} = 0$ . Relation (124) shows that we have  $\mathcal{Z}_a \subset \mathcal{Z}(0)$ .

**Proposition 9.15.** The adapted manifold  $\mathcal{Z}_a$  has the following implicit equations in fiberwise phase space roll-turn coordinates  $(\hat{\eta}^{\parallel}, \hat{\omega}, \Phi, \Psi)$ :

$$\begin{aligned} \hat{\eta}^{\parallel} &= 0 \\ (2\lambda^{\mathsf{v}}\Phi + \Psi)\omega^4 + 2\mu^{\mathsf{v}}\Phi\omega^3 + (2\lambda^{\mathsf{v}}\Phi + \Psi - M_0^2(\lambda^{\mathsf{v}})^2\Psi)\omega^2 \\ &+ 2\mu^{\mathsf{v}}\left(\Phi - M_0^2\lambda^{\mathsf{v}}\Psi\right)\omega - M_0^2(\mu^{\mathsf{v}})^2\Psi = 0 . \end{aligned}$$

In particular,  $\mathcal{Z}_a$  is a multisection of  $T\mathcal{M}_0$  which is contained in the second slow roll manifold  $\mathcal{Z}(0)$ .

*Proof.* When  $f^n = f^{\tau} = 0$ , Proposition 9.9 gives  $\hat{\eta}^{\parallel} = 0$  and:

$$\frac{1}{\mathcal{H}^2} \frac{\Phi^{\mathbf{v}}_{\tau\tau} \hat{\omega} + \Phi^{\mathbf{v}}_{n\tau}}{1 + \hat{\omega}^2} = \hat{\omega} , \qquad (137)$$

where:

$$\mathcal{H} = \frac{1}{M_0} \sqrt{\Phi + \sqrt{\Phi^2 + \frac{M_0^2 \Psi^2}{1 + \hat{\omega}^2}}} .$$
(138)

Combining these equations gives:

$$M_0^2 \frac{\Phi_{\tau\tau}^{\rm v} \hat{\omega} + \Phi_{n\tau}^{\rm v}}{1 + \hat{\omega}^2} = \hat{\omega} \left[ \varPhi + \sqrt{\varPhi^2 + \frac{M_0^2 \varPsi^2}{1 + \hat{\omega}^2}} \right] \;,$$

which implies:

$$\left(M_0^2 \frac{\Phi_{\tau\tau}^{\rm v} \hat{\omega} + \Phi_{n\tau}^{\rm v}}{\hat{\omega}(1+\hat{\omega}^2)} - \Phi\right)^2 = \Phi^2 + \frac{M_0^2 \Psi^2}{1+\hat{\omega}^2}$$

The latter amounts to the quartic equation:

$$\begin{split} M_0^4 (\varPhi_{\tau\tau}^{\rm v} \hat{\omega} + \varPhi_{n\tau}^{\rm v})^2 &- 2M_0^2 \varPhi(\varPhi_{\tau\tau}^{\rm v} \hat{\omega} + \varPhi_{n\tau}^{\rm v}) \hat{\omega} (1 + \hat{\omega}^2) - M_0^2 \Psi^2 \hat{\omega}^2 (1 + \hat{\omega}^2) = 0 \ , \ (139) \end{split}$$
which is equivalent to the second equation in (136) upon using (28).

**Corollary 9.16.** When  $\hat{\eta}^{\parallel} = 0$ , the first slow roll regime occurs when

$$\sqrt{1+\hat{\omega}^2} \gg M_0 \frac{\Psi}{\Phi} \ . \tag{140}$$

In this regime, the implicit equations of the adapted manifold  $Z_a$  reduce to:

$$\hat{\eta}^{\parallel} = 0 \ , \ \hat{\omega}^3 + \left(1 - \frac{M_0^2}{2\Phi} \varPhi_{\tau\tau}^{\rm v}\right) \hat{\omega} - \frac{M_0^2}{2\Phi} \varPhi_{n\tau}^{\rm v} = 0 \ . \tag{141}$$

In particular, the first slow roll regime can be realized along  $Z_a$  iff the cubic equation in (141) admits a real solution  $\hat{\omega}$  which satisfies (140).

$$\sqrt{1+\hat{\omega}^2} \gg 2||\Xi||^{\mathrm{v}} = M_0 \frac{\Psi}{\Phi}$$
.

In this case, we have  $\mathcal{H} \approx \frac{\sqrt{2\Phi}}{M_0}$  and (137) becomes:

$$\frac{M_0^2}{2\Phi} (\Phi_{\tau\tau}^{\rm v} \hat{\omega} + \Phi_{n\tau}^{\rm v}) \approx \hat{\omega} (1 + \hat{\omega}^2) ,$$

which is equivalent with the second equation in (141).

The cubic equation in (141) is in depressed form:

$$\hat{\omega}^3 + p\hat{\omega} + q = 0 ,$$

where:

$$p \stackrel{\mathrm{def.}}{=} 1 - \frac{M_0^2}{2\Phi} \varPhi_{\tau\tau}^{\mathrm{v}} \ , \ q \stackrel{\mathrm{def.}}{=} - \frac{M_0^2}{2\Phi} \varPhi_{n\tau}^{\mathrm{v}} < 0 \ .$$

The equation has discriminant  $\Delta$  given by:

$$-\Delta = 4p^3 + 27q^2 = 4\left(1 - \frac{M_0^2}{2\Phi}\Phi_{\tau\tau}^{\rm v}\right)^3 + \frac{27}{4}\frac{M_0^4}{\Phi^2}(\Phi_{n\tau}^{\rm v})^2 \ .$$

Distinguish the cases:

1.  $\Delta < 0$ . Then the equation has one real root and two complex roots (which are conjugate to each other). In this case, the real root is given by Cardano's formula:

$$\hat{\omega} = \sqrt[3]{-\frac{q}{2} + \sqrt{\frac{q^2}{4} + \frac{p^3}{27}} + \sqrt[3]{-\frac{q}{2} - \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}}$$

and condition (140) requires this root to satisfy:

$$\hat{\omega}^2 \gg M_0^2 \frac{\Psi^2}{\Phi^2} - 1$$

which gives a complicated condition on  $\Phi$ .

2.  $\Delta \geq 0$ . Then the equation has three real roots (some of which may coincide) and the condition that one of these roots satisfies (140) amounts to the requirement that the largest root satisfies that condition. Since  $\Delta \geq 0$  and q < 0, we have p < 0 (i.e.  $\frac{M_0^2}{2\Phi} \Phi_{\tau\tau}^{\rm v} > 1$ ) and  $\frac{27q^2}{4|p|^3} \leq 1$ . Let:

$$A \stackrel{\text{def.}}{=} 2\sqrt{\frac{|p|}{3}} = \sqrt{\frac{4|p|}{3}} \ge 0$$
.

and:

$$B \stackrel{\text{def.}}{=} \frac{3|q|}{A|p|} = \sqrt{\frac{27q^2}{4|p|^3}} \in [0,1] \; .$$

Setting  $\phi \stackrel{\text{def.}}{=} \frac{1}{3} \arccos B \in [0, \frac{\pi}{9}]$ , the three real roots are given by Viete's formulas:

$$\hat{\omega}_k = A\cos(\phi + \frac{2\pi}{3}k) \ (k = 0, 1, 2)$$

and correspond to the projection on the x axis of three equidistant points of the unit circle. It is clear that the largest of the roots belongs to the interval  $[A\cos(\frac{\pi}{3}), A] = [\frac{A}{2}, A]$ . Hence the requirement that a root satisfies (140) is equivalent with the condition:

$$1 + \frac{A^2}{4} \gg M_0^2 \frac{\Psi^2}{\Phi^2} \Longleftrightarrow 2 + M_0^2 \frac{\Phi_{\tau\tau}}{2\Phi} \gg 3M_0^2 \frac{\Psi^2}{\Phi^2} \ ,$$

which amounts to:

$$M_0^2 \left(\frac{\Psi^2}{\Phi^2} - \frac{\Phi_{\tau\tau}}{6\Phi}\right) \ll \frac{2}{3}$$

9.5. The horizontal approximation defined by the adapted connection. Let  $\mathcal{U} \subset \mathcal{M}_0$  be the non-critical set of  $(\mathcal{M}, \mathcal{G}, \Phi)$  (see Definition 3.7) and recall that the adapted phase space coordinates  $(v_n, v_\tau, \Phi = \Phi, \Psi = \Psi)$  give a special coordinate system on  $T\mathcal{U}$ . On this open subset of  $T\mathcal{M}_0$ , the horizontal projection of the cosmological semispray has the form:

$$S_H = S^{\Phi} \frac{\partial}{\partial \Phi} + S^{\Psi} \frac{\partial}{\partial \Psi} ,$$

where (see Corollary 3.31):

$$S^{\Phi} = \Psi v_n \tag{142}$$
$$S^{\Psi} = \frac{1}{2} (\Phi_{nn}^{\mathrm{v}} v_n + \mu^{\mathrm{v}} \Psi v_{\tau}) \; .$$

Hence the horizontal approximation defined by the adapted connection replaces locally the cosmological equation for curves whose canonical lift lies in  $TU \setminus Q$  with the system:

$$\dot{v}_n = \dot{v}_\tau = 0 , \ \dot{\Phi} = \Psi v_n , \ \dot{\Psi} = \frac{1}{2} (\Phi_{nn}^{\mathsf{v}} v_n + \mu^{\mathsf{v}} \Psi v_\tau) .$$
 (143)

The first two equations show that  $v_n$  and  $v_{\tau}$  are constant while the remaining equations give a system for the time evolution of  $\Phi$  and  $\Psi$ . In these equations, it is understood that the quantities  $\Phi_{nn}$  and  $\mu$  are expressed as functions of  $\Phi$  and  $\Psi$ and hence  $\Phi_{nn}^{v}$  and  $\mu^{v}$  are expressed as functions of  $\Phi$  and  $\Psi$ . The solutions of (143) are the  $W_a$ -curves which serve as horizontal approximants of cosmological curves contained in  $\mathcal{U}$  in the sense of Section 1. As explained there, the approximation is accurate for cosmological curves which are close to the adapted manifold  $\mathcal{Z}_a$ .

# 9.6. The adapted approximation.

**Definition 9.17.** The adapted approximation is defined by the adapted conditions:

$$|\mathbf{f}^n| \ll 1 \text{ and } |\mathbf{f}^\tau| \ll 1 . \tag{144}$$

A vector  $u \in T\mathcal{M}_0 \setminus Q$  satisfies the adapted conditions iff it is very close to the adapted manifold  $\mathcal{Z}_a$ . Notice from relation (119) that the adapted conditions imply  $|\hat{\eta}^{\parallel}| \ll 1$ , so the adapted approximation implies the second slow roll approximation; in particular, it implies the relation (see (133)):

$$\mathfrak{f}^\tau\approx -\frac{\mathfrak{f}^n}{\hat{\omega}^2}~.$$

However, the second slow roll condition  $|\hat{\eta}^{\parallel}| \ll 1$  does *not* imply the adapted conditions (as made manifest by the fact that  $\mathcal{Z}_a$  is a codimension one submanifold of the second slow roll manifold  $\mathcal{Z}(0)$ ).

## Acknowledgments

This work was supported by national grant PN 19060101/2019-2022 and by a Maria Zambrano Excellency Fellowship as well by the COST Action COSMIC WISPers CA21106, supported by COST (European Cooperation in Science and Technology).

# Appendix A. Signed angles, curvatures and turning rates on an oriented Riemann surface

Let  $(\mathcal{M}, \mathcal{G})$  be an oriented, connected and borderless Riemann surface, whose volume form we denote by vol. In a positively-oriented local coordinate system  $(x^1, x^2)$  on  $\mathcal{M}$ , we have:

$$\operatorname{vol} = \sqrt{\det \hat{\mathcal{G}}} \mathrm{d}x^1 \wedge \mathrm{d}x^2 = \frac{1}{2} \sqrt{\det \hat{\mathcal{G}}} \epsilon_{ij} \mathrm{d}x^i \wedge \mathrm{d}x^j$$

where  $\epsilon_{ij}$  is the Levi-Civita symbol and we use Einstein summation over repeated indices. We have  $vol(e_j, e_j) = \epsilon_{ij}$  in any orthonormal and positive local frame of  $T\mathcal{M}$ .

A.1. The signed angle between vectors. Let  $v, v_0 \in T_m \mathcal{M}$  be two non-zero tangent vectors to  $\mathcal{M}$  at a point  $m \in \mathcal{M}$ . Let  $\Theta(v, v_0) \in [0, \pi]$  be the ordinary angle between v and  $v_0$  in the Euclidean space  $(T_m \mathcal{M}, \mathcal{G}_m)$ , which is defined through the relation:

$$\mathcal{G}(v, v_0) = ||v||||v_0||\cos\Theta(v, v_0)$$

Notice that  $\Theta(v_0, v) = \Theta(v, v_0)$ .

**Definition A.1.** Suppose that v and  $v_0$  are linearly independent, i.e. v is not a multiple of  $v_0$ . Then the *chirality* of v relative to  $v_0$  is the signature of the basis  $(v_0, v)$  of  $T_m \mathcal{M}$ , which is defined through:

$$\sigma_{v_0}(v) \stackrel{\text{def.}}{=} \operatorname{sign}[\operatorname{vol}(v_0, v)] \in \{-1, 1\}$$
.

When  $v_0$  and v are linearly dependent, we define  $\sigma_{v_0}(v) = 0$ .

When  $v_0$  and v are linearly independent, we have  $\sigma_{v_0}(v) = +1$  iff the basis  $(v_0, v)$  of  $T_m \mathcal{M}$  is positively oriented and  $\sigma_{v_0}(v) = -1$  iff this basis is negatively oriented. Notice that  $\sigma_{v_0}(v) = -\sigma_v(v_0)$ .

**Definition A.2.** The signed angle of v relative to  $v_0$  is the real number  $\theta_{v_0}(v) \in (-\pi, \pi]$  defined as follows:

- (1) If v is not proportional to  $v_0$ , then we set  $\theta_{v_0}(v) \stackrel{\text{def.}}{=} \sigma_{v_0}(v)\Theta(v_0,v) \in (-\pi,0)\cup(0,\pi).$
- (2) If  $v = \alpha v_0$  with  $\alpha > 0$ , then we set  $\theta_{v_0}(v) \stackrel{\text{def.}}{=} 0$ .
- (3) If  $v = \alpha v_0$  with  $\alpha < 0$ , then we set  $\theta_{v_0}(v) \stackrel{\text{def.}}{=} \pi$ .

Notice that  $\theta_v(v_0) = -\theta_{v_0}(v)$  when v and  $v_0$  are linearly independent and that  $\theta_v(v_0) = \theta_{v_0}(v) \in \{0, \pi\}$  iff v and  $v_0$  are linearly dependent. In all cases, we have  $\theta_v(v_0) = -\theta_{v_0}(v) \mod 2\pi$  and hence:

$$\cos \theta_v(v_0) = \cos \theta_{v_0}(v) , \ \sin \theta_v(v_0) = -\sin \theta_{v_0}(v) .$$

We have:

$$\operatorname{sign}(\sin \theta_{v_0}(v)) = \sigma_{v_0}(v) \;,$$

with the convention sign(0) = 0.

**Definition A.3.** Given any non-zero vector  $v \in T_m \mathcal{M} \setminus \{0\}$ , the *positive unit* normal to v is the unique unit norm vector  $N_v \in T_m \mathcal{M}$  which is orthogonal to vand satisfies  $\sigma_v(N_v) = +1$ , i.e. the orthonormal basis  $(v, N_v)$  of  $T_m \mathcal{M}$  is positively oriented. The negative unit normal to v is the vector  $-N_v$ .

We have:

$$N_v = \frac{1}{||v||} (\iota_v \operatorname{vol})^{\sharp} \quad \forall v \in \dot{T}\mathcal{M} .$$
(145)

Notice that:

$$N_{\alpha v} = \operatorname{sign}(\alpha) N_v \ \forall \alpha \in \mathbb{R}^{\times}$$

We have  $\theta_v(N_v) = +\frac{\pi}{2}$  and  $\theta_v(-N_v) = -\frac{\pi}{2}$ . The following statement is immediate.

**Proposition A.4.** Let  $v_0, v \in T_m \mathcal{M}$  be two non-vanishing vectors. Then  $(v, N_v)$  is obtained from  $(v_0, N_{v_0})$  by a rotation of angle  $\theta_{v_0}(v)$ , i.e. we have:

$$v = [\cos \theta_{v_0}(v)]v_0 + [\sin \theta_{v_0}(v)]N_{v_0}$$
  
$$N_v = -[\sin \theta_{v_0}(v)]v_0 + [\cos \theta_{v_0}(v)]N_{v_0}$$

Remark A.5. We have:

$$\sigma_{v_0}(v) = \operatorname{sign}(\cos \theta_v(N_{v_0})) = \operatorname{sign}(\mathcal{G}(N_{v_0}, v))$$

with the convention sign(0) = 0.

To give a universal geometric description of the notions introduced above, let  $\pi: T\mathcal{M} \to \mathcal{M}$  be the projection of the tangent bundle of  $\mathcal{M}$  and  $F \stackrel{\text{def.}}{=} \pi^*(T\mathcal{M})$  be the tautological vector bundle over  $T\mathcal{M}$  (which is also known [24] as the Finsler bundle of  $\mathcal{M}$ ). A local section of F defined on an open set  $U \subset T\mathcal{M}$  can be viewed as a map  $f: U \to T\mathcal{M}$  which satisfies  $f(u) \in T_{\pi(u)}\mathcal{M}$  for all  $u \in U$ . Thus f is a map which associates to a tangent vector  $u \in T\mathcal{M}$  another vector, tangent to  $\mathcal{M}$  at the same point as u. Notice that F carries the natural Euclidean pairing  $\mathcal{G}_F: F \times_{T\mathcal{M}} F \to \mathbb{R}_{T\mathcal{M}}$  given by the bundle pull-back of  $\mathcal{G}$ . It also carries an orientation described by the bundle pullback of the volume form of  $(\mathcal{M}, \mathcal{G})$ , which is a nowhere-vanishing section of the determinant line bundle of F. We will denote  $\mathcal{G}_F$  by  $\mathcal{G}$  for ease of notation.

**Definition A.6.** The normalization map and positive normal map of  $(\mathcal{M}, \mathcal{G})$  are the sections  $T, N \in \Gamma(\dot{T}\mathcal{M}, F)$  defined through:

$$T(u) \stackrel{\text{def.}}{=} \frac{u}{||u||} , \ N(u) \stackrel{\text{def.}}{=} N_u \ \forall u \in \dot{T}\mathcal{M} .$$

The chirality map and signed angle map of  $(\mathcal{M}, \mathcal{G})$  are the maps  $\sigma : \dot{T}\mathcal{M} \times_{\mathcal{M}} \dot{T}\mathcal{M} \rightarrow \{-1, 0, 1\}$  and  $\theta : \dot{T}\mathcal{M} \times_{\mathcal{M}} \dot{T}\mathcal{M} \rightarrow (-\pi, \pi]$  defined through:

$$\sigma(v_0, v) \stackrel{\text{def.}}{=} \sigma_{v_0}(v) , \ \theta(v_0, v) \stackrel{\text{def.}}{=} \theta_{v_0}(v) .$$

for all  $(v_0, v) \in \dot{T}\mathcal{M} \times_{\mathcal{M}} \dot{T}\mathcal{M}$ .

Notice that the maps  $\sigma$  and  $\theta$  are discontinuous. The following notion encodes oriented Frenet frames of nondegenerate curves in a universal manner.

**Definition A.7.** The abstract oriented Frenet frame of  $(\mathcal{M}, \mathcal{G})$  is the orthononormal frame (T, N) of the vector bundle  $F|_{\dot{T}\mathcal{M}}$ .

As we show in the next subsection, the Finsler vector fields T and N give a universal geometric description of the oriented Frenet frame of nondegenerate curves. With the definitions above, we have (see Remark A.5):

$$\sigma = \operatorname{sign}(\mathcal{G} \circ (N \times_{\mathcal{M}} T)) , \qquad (146)$$

where we view  $\mathcal{G}$  as a map  $\mathcal{G} : T\mathcal{M} \times_{\mathcal{M}} T\mathcal{M} \to \mathbb{R}$  and N, T as maps from  $\dot{T}\mathcal{M}$  to  $\dot{T}\mathcal{M}$ . Here  $N \times_{\mathcal{M}} T : \dot{T}\mathcal{M} \times_{\mathcal{M}} \dot{T}\mathcal{M} \to \dot{T}\mathcal{M} \times_{\mathcal{M}} \dot{T}\mathcal{M}$  denotes the map defined through:

$$(N \times_{\mathcal{M}} T)(u, v) \stackrel{\text{def.}}{=} (N(u), T(v)) \in \dot{T}_m \mathcal{M} \times \dot{T}_m \mathcal{M} \ \forall (u, v) \in \dot{T}_m \mathcal{M} \times \dot{T}_m \mathcal{M} .$$

A.2. The signed curvature and signed turning rate of a curve in  $\mathcal{M}$ . Let  $(\mathcal{M}, \mathcal{G})$  be an oriented Riemann surface and  $\varphi : I \to \mathcal{M}$  be a smooth curve in  $\mathcal{M}$ . For any  $t \in I_{reg}$ , let:

$$T_{\varphi}(t) \stackrel{\text{def.}}{=} \frac{\varphi(t)}{||\dot{\varphi}(t)||} \in T_{\varphi(t)}\mathcal{M}$$

be the unit tangent vector to  $\varphi$  at time t.

**Definition A.8.** The positive normal vector to the curve  $\varphi$  at time  $t \in I_{\text{reg}}$  is the positive unit normal  $N_{\varphi}(t) \stackrel{\text{def.}}{=} N_{T_{\varphi}(t)} \in T_{\varphi(t)} \mathcal{M}$  to  $T_{\varphi}(t)$ . The positive orthonormal basis  $(T_{\varphi}(t), N_{\varphi}(t))$  of  $T_{\varphi(t)} \mathcal{M}$  is called the positive Frenet basis at time  $t \in I_{\text{reg}}$ . The oriented Frenet frame of  $\varphi$  is the frame  $(T_{\varphi}, N_{\varphi})$  of the vector bundle  $(\varphi|_{I_{\text{reg}}})^*(T\mathcal{M})$  defined by  $T_{\varphi}(t)$  and  $N_{\varphi}(t)$  as t varies in  $I_{\text{reg}}$ .

For any  $t \in I_{reg}$ , we have:

$$T_{\varphi}(t) = T(\dot{\varphi}(t)) , \ N_{\varphi}(t) = N(\dot{\varphi}(t)) ,$$

where T and N are the normalization and positive normal maps of  $(\mathcal{M}, \mathcal{G})$  defined in the previous subsection. Accordingly, we have:

$$T_{\varphi} = (\dot{\varphi}|_{I_{\text{reg}}})^*(T) , \ N_{\varphi} = (\dot{\varphi}|_{I_{\text{reg}}})^*(N) ,$$

where  $\dot{\varphi} = c(\varphi) : I \to T\mathcal{M}$  is the canonical lift of  $\varphi$  to  $T\mathcal{M}$ . Since  $\pi \circ \dot{\varphi} = \varphi$ , we have  $\dot{\varphi}^*(F) = \varphi^*(T\mathcal{M})$ .

Let s be an increasing proper length parameter along  $\varphi$  with image interval J (thus s gives an orientation-preserving diffeomorphism  $\hat{s}$  from I to J) and let  $J_{\text{reg}} \stackrel{\text{def.}}{=} \hat{s}(I_{\text{reg}})$ . We have  $\dot{s}(t) = ||\dot{\varphi}(t)||$  for all  $t \in I$ , which implies:

$$T_{\varphi}(s) = \varphi'(s) \ \forall s \in I_{\text{reg}}$$
,

where the prime denotes derivative with respect to s. The vector  $K_{\varphi}(s) \stackrel{\text{def.}}{=} \nabla_s T_{\varphi}(s)$  is the classical *curvature vector* of  $\varphi$ . The relation  $\mathcal{G}(T_{\varphi}(s), T_{\varphi}(s)) = 1$  implies:

$$K_{\varphi}(s) \perp T_{\varphi}(s) \ \forall s \in J_{\text{reg}}$$

Recall that the ordinary curvature of  $\varphi$  at  $s \in J_{reg}$  is the *strictly positive* quantity  $\chi_{\varphi}(s)$  defined through:

$$\chi_{\varphi}(s) \stackrel{\text{def.}}{=} ||K_{\varphi}(s)|| = ||\nabla_s T_{\varphi}(s)|| > 0 \ \forall s \in I_{\text{reg}} \ .$$

**Definition A.9.** The *turning sign* (or *chirality*) of the curve  $\varphi$  at  $s \in J_{\text{reg}}$  is the chirality of the vector  $K_{\varphi}(s)$  with respect to  $T_{\varphi}(s)$ :

$$\sigma_{\varphi}(s) \stackrel{\text{def.}}{=} \sigma_{T_{\varphi}(s)}(K_{\varphi}(s)) = \operatorname{sign} \mathcal{G}(N_{\varphi}(s), K_{\varphi}(s)) \in \{-1, 0, 1\} ,$$

where we set  $\sigma_{\varphi}(s) = 0$  when  $K_{\varphi}(s) = 0$ .

1.4

Notice that  $\sigma_{\varphi}(s) = 0$  iff  $K_{\varphi}(s) = 0$ . When  $K_{\varphi}(s) \neq 0$ , the oriented angle  $\theta_{T_{\varphi}(s)}(K_{\varphi}(s))$  equals  $\sigma_{\varphi}(s)\frac{\pi}{2}$ . In all cases, we have  $K_{\varphi}(s) = \sigma_{\varphi}(s)\chi_{\varphi}(s)N_{\varphi}(s)$ .

**Definition A.10.** The signed curvature of  $\varphi$  at  $s \in J_{reg}$  is defined through:

$$\xi_{\varphi}(s) \stackrel{\text{def.}}{=} \sigma_{\varphi}(s) \chi_{\varphi}(s) \in \mathbb{R} .$$
(147)

Thus  $K_{\varphi}(s) = \xi_{\varphi}(s)N_{\varphi}(s)$ . Notice that  $\xi_{\varphi}$  coincides with the ordinary signed curvature of plane curves when  $(\mathcal{M}, \mathcal{G})$  is the Euclidean plane. The Frenet-Serret relations of  $\varphi$  take the form:

$$K_{\varphi}(s) \stackrel{\text{def.}}{=} \nabla_s T_{\varphi}(s) = +\xi_{\varphi}(s) N_{\varphi}(s) \tag{148}$$

$$\nabla_s N_{\varphi}(s) = -\xi_{\varphi}(s) T_{\varphi}(s) \ \forall s \in J_{\text{reg}} .$$
(149)

**Definition A.11.** The rescaled signed turning rate of  $\varphi$  at  $t \in I_{reg}$  is the quantity:

$$\Omega_{\varphi}(t) \stackrel{\text{def.}}{=} -\frac{\mathcal{G}(N_{\varphi}(t), \nabla_t \dot{\varphi}(t))}{||\dot{\varphi}(t)||} = -||\dot{\varphi}(t)||\xi_{\varphi}(t) \in \mathbb{R} .$$
(150)

**Proposition A.12.** The Fenet-Serret relations of  $\varphi$  are equivalent with:

$$\nabla_t T_{\varphi}(t) = -\Omega_{\varphi}(t) N_{\varphi}(t) , \ \nabla_t N_{\varphi}(t) = +\Omega_{\varphi}(t) T_{\varphi}(t) \ \forall t \in I_{\text{reg}} .$$
(151)

*Proof.* Follows by direct computation from (148) and (149) upon using (150).  $\Box$ 

The first Frenet-Serret relation can also be written as:

$$\nabla_t \dot{\varphi}(t) = \frac{\mathrm{d}||\dot{\varphi}(t)||}{\mathrm{d}t} T_{\varphi}(t) - ||\dot{\varphi}(t)||\Omega_{\varphi}(t)N_{\varphi}(t) \ \forall t \in I_{\mathrm{reg}} \ . \tag{152}$$

Note that  $K_{\varphi}$ ,  $\sigma_{\varphi}$ ,  $\chi_{\varphi}$ ,  $\xi_{\varphi}$  and  $\Omega_{\varphi}$  are second order quantities which cannot be expressed through objects defined on  $T\mathcal{M}$  unless  $\varphi$  satisfies a geometric second order ODE on  $\mathcal{M}$ .

APPENDIX B. RELATION BETWEEN PARAMETERS USED IN THIS PAPER AND PARAMETERS USED IN THE PHYSICS LITERATURE

B.0.1. *The first slow roll parameter*. It is customary to define another second order parameter of a curve as follows:

**Definition B.1.** The *first slow roll parameter* of a curve  $\varphi : I \to \mathcal{M}$  is the function  $\varepsilon_{\varphi} : I \to \mathbb{R}$  defined through:

$$\boldsymbol{\varepsilon}_{\varphi}(t) \stackrel{\text{def.}}{=} -\frac{H_{\varphi}(t)}{H_{\varphi}(t)^2} = -\frac{1}{H_{\varphi}(t)} \frac{\mathrm{d}}{\mathrm{d}t} \log H_{\varphi}(t) = 3\hat{\boldsymbol{\varepsilon}}_{\varphi}(t) \ ,$$

where:

$$\hat{\boldsymbol{\varepsilon}}_{\varphi}(t) \stackrel{\text{def.}}{=} -\frac{1}{\mathcal{H}_{\varphi}(t)} \frac{\mathrm{d}}{\mathrm{d}t} \log \mathcal{H}_{\varphi}(t)$$

is the *rescaled* first slow roll parameter of  $\varphi$ .

For cosmological curves, the first slow roll parameter reduces to a function of the first IR parameter and hence does not give an independent basic observable.

**Proposition B.2.** Suppose that  $\varphi : I \to \mathcal{M}$  is a cosmological curve. Then we have:

$$\hat{\boldsymbol{\varepsilon}}_{\varphi}(t) = rac{\kappa_{\varphi}(t)}{1 + \kappa_{\varphi}(t)} \ \forall t \in I \ .$$

*Proof.* Follows immediately from the cosmological equation (11).

Remark B.3. When  $\varphi$  is a cosmological curve, we have  $\varepsilon_{\varphi}(t) = 0$  iff  $\kappa_{\varphi}(t) = 0$  and  $\varepsilon_{\varphi}(t) = 1$  (i.e.  $\hat{\varepsilon}_{\varphi}(t) = 1/3$ ) iff  $\kappa(t) = 1/2$ .

The definition of other parameters (beyond the first slow roll parameter) used in the physics literature (in the conventions of reference [29]) is as follows, where for ease of notation we don't indicate the curve  $\varphi$  on which s is an increasing length parameter:

$$\begin{split} \delta \stackrel{\text{def.}}{=} & \frac{\ddot{s}}{H\dot{s}} , \ \eta \stackrel{\text{def.}}{=} & \frac{\dot{\varepsilon}}{H\varepsilon} = 2\delta + 2\varepsilon \ , \ \xi \stackrel{\text{def.}}{=} & \frac{\ddot{s}}{H^2\ddot{s}} \\ \nu \stackrel{\text{def.}}{=} & \frac{1}{\Omega} \frac{\mathrm{d}}{\mathrm{d}t} \left(\frac{\Omega}{H}\right) \ , \ \varepsilon_V \stackrel{\text{def.}}{=} & \frac{M^2}{2} \frac{||\mathrm{d}\Phi||^2}{\Phi^2} \ . \end{split}$$

Here  $\eta$  is the ordinary second slow roll parameter. We have  $\eta = -\delta$ . The parameters used in this paper are:

$$M_0 \stackrel{\text{def.}}{=} M \sqrt{\frac{2}{3}} \ , \ \mathcal{H} \stackrel{\text{def.}}{=} 3H \ , \ \kappa \stackrel{\text{def.}}{=} \frac{\dot{s}^2}{2\Phi} \ , \ \hat{\eta}^{\parallel} \stackrel{\text{def.}}{=} \frac{\eta}{3} = -\frac{1}{3}\delta \ , \ \Xi \stackrel{\text{def.}}{=} \frac{M_0}{2} \frac{\mathrm{d}\Phi}{\Phi}$$

and  $\hat{\boldsymbol{\varepsilon}} \stackrel{\text{def.}}{=} -\frac{\boldsymbol{\varepsilon}}{3}$ , which was defined above. We have:

$$\mathcal{H} = \sqrt{2\Phi} (1+\kappa)^{1/2} , \ \kappa = \frac{\hat{\varepsilon}}{1-\hat{\varepsilon}} , \ \kappa(\kappa+1) = \frac{\hat{\varepsilon}}{(1-\hat{\varepsilon})^2} , \ ||\Xi||^2 = \hat{\varepsilon}_V .$$

B.0.2. A certain formula found in the physics literature. Relations (65) and (68) give:

$$\Omega = -\mathcal{H} \frac{\sin \theta}{\hat{c}} = \frac{||\mathrm{d}\Phi||^{\mathrm{v}}}{\mathcal{N}} \sin \theta \; .$$

As mentioned above, the parameter  $\delta$  of [29] is related to our parameters by  $\delta = -3\hat{\eta}^{\parallel}$ , while the ordinary second slow roll parameter used in loc. cit. is given by  $\eta = \frac{\dot{\epsilon}}{H\epsilon} = 2\delta + 2\epsilon$ . Using this and the relation  $\varepsilon = 3\hat{\varepsilon}$ , we compute:

$$1 + \frac{\eta}{2(3-\varepsilon)} = \frac{3+\delta_{\varphi}}{3-\epsilon} = \frac{1-\hat{\eta}^{\parallel}}{1-\hat{\varepsilon}}$$

and:

$$\frac{\boldsymbol{\varepsilon}}{3(1-\boldsymbol{\varepsilon}/3)^2} = \frac{\hat{\boldsymbol{\varepsilon}}}{(1-\hat{\boldsymbol{\varepsilon}})^2} = \kappa(1+\kappa) \ .$$

Moreover, we have  $||d\Xi||^2 = \frac{1}{3}\varepsilon_V$ . This allows one to write our relation (113) (which is a consequence (83)) in the form:

$$\frac{\varepsilon_V}{\varepsilon} = \left(1 + \frac{\eta}{2(3-\varepsilon)}\right)^2 + \frac{\Omega^2}{9H^2} \frac{1}{(1-\frac{\varepsilon}{3})^2} , \qquad (153)$$

which was originally given in [3, 30]. We stress that this formidable looking equation is a rewriting of relations (83) (the first of which which can be taken as the geometric definition of  $\hat{\eta}^{\parallel}$ ), being a complicated re-formulation of the latter in terms of the parameters  $\boldsymbol{\varepsilon}, \boldsymbol{\varepsilon}_{V}, \Omega$  and  $\mathcal{H}$ . The comparative simplicity of (83) and (113) underscores the conceptual advantage of using parameters which are geometrically meaningful.

## References

- S. Garg, C. Krishnan, Bounds on Slow Roll and the de Sitter Swampland, JHEP 11 (2019) 075, arXiv:1807.05193 [hep-th].
- [2] H. Ooguri, E. Palti, G. Shiu, C. Vafa, Distance and de Sitter Conjectures on the Swampland, Phys. Lett. B 788 (2019) 180, arXiv:1810.05506 [hep-th].
- [3] A. Achucarro, G. Palma, The string swampland constraints require multi-field inflation, JCAP 02 (2019) 041, arXiv:1807.04390 [hep-th].
- [4] R. Bravo, G. A. Palma, S. Riquelme, A Tip for Landscape Riders: Multi-Field Inflation Can Fulfill the Swampland Distance Conjecture, JCAP 02 (2020) 004, arXiv:1906.05772 [hep-th].
- G. A. Palma, S. Sypsas, C. Zenteno, Seeding primordial black holes in multi-field inflation, Phys. Rev. Lett. 125 (2020) 121301, arXiv:2004.06106 [astro-ph.CO].
- [6] J. Fumagalli, S. Renaux-Petel, J. W. Ronayne, L. T. Witkowski, *Turning in the landscape: a new mechanism for generating Primordial Black Holes*, Phy. Lett. B 841 (2023) 137921, arXiv:2004.08369 [hep-th].
- [7] L. Anguelova, On Primordial Black Holes from Rapid Turns in Two-field Models, JCAP 06 (2021) 004, arXiv:2012.03705 [hep-th].
- [8] Y. Akrami, M. Sasaki, A. Solomon and V. Vardanyan, Multi-field dark energy: cosmic acceleration on a steep potential, Phys. Lett. B 819 (2021) 136427, arXiv:2008.13660 [astroph.CO].
- [9] L. Anguelova, J. Dumancic, R. Gass, L.C.R. Wijewardhana, *Dark Energy from Inspiraling in Field Space*, JCAP 03 (2022) 018, arXiv:2111.12136 [hep-th].
- [10] R. Schimmrigk, Modular Inflation Observables and j-Inflation Phenomenology, JHEP 09 (2017) 043, arXiv:1612.09559 [hep-th].
- [11] R. Schimmrigk, Multifield Reheating after Modular j-Inflation, Phys. Lett. B 782 (2018) 193-197, arXiv:1712.09961 [hep-ph].
- [12] R. Schimmrigk, Large and small field inflation from hyperbolic sigma models, Phys. Rev. D 105, 063541, arXiv:2108.05400 [hep-th].

### CALIN IULIU LAZAROIU

- [13] C. I. Lazaroiu, C. S. Shahbazi, Generalized two-field α-attractor models from geometrically finite hyperbolic surfaces, Nucl. Phys. B 936 (2018) 542-596.
- [14] E. M. Babalic and C. I. Lazaroiu, Generalized α-attractor models from elementary hyperbolic surfaces, Adv. Math. Phys. 2018 (2018) 7323090, arXiv:1703.01650.
- [15] E. M. Babalic and C. I. Lazaroiu, Generalized α-attractors from the hyperbolic triplypunctured sphere, Nucl. Phys. B 937 (2018) 434-477, arXiv:1703.06033 [hep-th].
- [16] L. Anguelova, E. M. Babalic, C. I. Lazaroiu, Two-field Cosmological α-attractors with Noether Symmetry, JHEP 04 (2019) 148, arXiv:1809.10563 [hep-th].
- [17] L. Anguelova, E. M. Babalic, C. I. Lazaroiu, Hidden symmetries of two-field cosmological models, JHEP 09 (2019) 007, arXiv:1905.01611 [hep-th].
- [18] C. I. Lazaroiu, Hesse manifolds and Hessian symmetries of multifield cosmological models, Rev. Roumaine Math. Pures Appl. 66 (2021) 2, 329-345, arXiv:2009.05117 [hep-th].
- [19] J. Palis Jr. and W. De Melo, Geometric theory of dynamical systems: an introduction, Springer, New York, U.S.A. (2012).
- [20] A. Katok, B. Hasselblatt, Introduction to the modern theory of dynamical systems, Cambridge U.P., 1995.
- [21] C. I. Lazaroiu, Dynamical renormalization and universality in classical multifield cosmological models, Nucl. Phys. B 983 (2022) 115940, arXiv:2202.13466 [hep-th].
- [22] E. M. Balabic, C. I. Lazaroiu, The infrared behavior of tame two-field cosmological models , Nucl. Phys. B 983 (2022) 115929, arXiv:2203.02297 [gr-qc].
- [23] L. Anguelova, C. I. Lazaroiu, Dynamical consistency conditions for rapid turn inflation, JCAP 05 (2023) 020, arXiv:2210.00031 [hep-th].
- [24] J. Szilasi, R. L. Lovas, D. C. Kertesz, Connections, sprays and Finsler structures, World Scientific, 2014.
- [25] P. Petersen, Riemannian geometry, Graduate Texts in Mathematics, 3rd ed., 2016.
- [26] D. J. Saunders, The Geometry of Jet Bundles, Cambridge University Press, 1989.
- [27] C. I. Lazaroiu, Natural observables and dynamical approximations in multifield cosmological models, arXiv:2305.04367 [gr-qc].
- [28] C.-Wu Ho, A note on proper maps, Proc. AMS., 51 (1975) 1.
- [29] V. Aragam, R. Chiovoloni, S. Paban, R. Rosati, I. Zavala, Rapid-turn inflation in supergravity is rare and tachyonic, JCAP 03 (2022) 002, arXiv:2110.05516 [hep-th].
- [30] A. Hetz, G. A. Palma, Sound Speed of Primordial Fluctuations in Supergravity Inflation, Phys. Rev. Lett. 117 (2016) 101301, arXiv:1601.05457 [hep-th].

DEPARTAMENTO DE MATEMATICAS, UNIVERSIDAD UNED - MADRID, CALLE DE JUAN DEL Rosal 10, 28040, Madrid, Spain, clazaroiu@mat.uned.es, and, Horia Hulubei National Institute of Physics and Nuclear Engineering, Reactorului 30, Bucharest-Magurele, 077125, Romania, lcalin@theory.nipne.ro