The Spectral Geometry of the Mesh Matrices of Graphs

Sylvain E. Cappell and Edward Y. Miller Dedicated to the memory of Jacob Eli Goodman

Abstract

The mesh matrix $Mesh(G, T_0)$ of a connected finite graph G = (V(G), E(G)) =(Rectices, edges) of G of with respect to a choice of a spanning tree $T_0 \subset G$ is defined and studied. It was introduced by Trent [30],[31]. Its characteristic polynomial $\det(X \cdot Id - Mesh(G, T_0))$ is shown to equal $\sum_{j=0}^{N} (-1)^j ST_j(G, T_0) (X-1)^{N-j} (\star)$ where $ST_i(G, T_0)$ is the number of spanning trees of G meeting $E(G - T_0)$ in j edges and $N = |E(G - T_0)|$. As a consequence, there are Tutte-type deletion-contraction Firmulae for computing this polynomial. Additionally, $Mesh(G, T_0) - Id$ is of the special form $Y^t \cdot Y$; so the eigenvalues of the mesh matrix $Mesh(G, T_0)$ are all real and are furthermore be shown to be $\geq +1$. It is shown that $Y \cdot Y^t$, called the mesh Laplacian, is a generalization of the standard graph Kirchhoff Laplacian $\Delta(H) = Deg - Adj$ of a graph H. For example, (\star) generalizes the all minors matrix tree theorem for graphs H and gives a deletion-contraction formula for the characteristic polynomial of $\Delta(H)$. This generalization is explored in some detail. The smallest positive eigenvalue of the mesh Laplacian, a measure of flux, is estimated, thus extending the classical inequality for the Rirchoff Laplacian of graphs.

The Spectral Geometry of the Mesh Matrices of Graphs

Sylvain E. Cappell and Edward Y. Miller Dedicated to the memory of Jacob Eli Goodman

Abstract

The mesh matrix $Mesh(G, T_0)$ of a connected finite graph G = (V(G), E(G)) = (vertices, edges) of G of with respect to a choice of a spanning tree $T_0 \subset G$ is defined and studied. It was introduced by Trent [30, 31]. Its characteristic polynomial $det(X \cdot Id - Mesh(G, T_0))$ is shown to equal $\sum_{j=0}^{N} (-1)^j ST_j(G, T_0) (X-1)^{N-j}$ (*) where $ST_j(G, T_0)$ is the number of spanning trees of G meeting $E(G-T_0)$ in f edges and f edges and f edges are a consequence, there are Tutte-type deletion-contraction formulae for computing this polynomial. Additionally, f edges are a consequence, there are Tutte-type deletion-contraction formulae for computing this polynomial. Additionally, f edges are a consequence, there are Tutte-type deletion-contraction formulae for computing this polynomial. Additionally, f edges are a consequence, there are Tutte-type deletion-contraction formulae for computing this polynomial. Additionally, f edges are a consequence, there are Tutte-type deletion-contraction formulae for computing this polynomial. Additionally, f edges are a consequence, there are Tutte-type deletion-contraction formulae for computing this polynomial. Additionally, f edges are f edges and f edges are f edges and f edges are f edges and f edges are f edges are f edges are f edges and f edges are f edges

1 Introduction

For a graph H = (V(H), E(H)) with vertices V(H) and edges E(H), two distinct vertices, say P, Q, are called adjacent if there is an edge e of H with P, Q as end points. This is written as $P \sim Q$. The graph H is called connected if for any two distinct vertices, P, Q there is a simple path from P to Q in H, i.e., a sequence of distinct vertices R_1, R_2, \dots, R_K with $P = R_1, Q = R_K$ and $R_k \sim R_{k+1}, k = 1, \dots, K-1$. The graph H is called a tree if H is connected and any two distinct vertices are connected by a unique simple path. A subgraph $T \subset G$ of a graph G is called a spanning tree of G if the subgraph T is a tree and V(T) = V(G), i.e., T and G contain the same vertices.

The context of the mesh matrix studied here is that of a pair (G, T_0) with G a connected finite graph and the chosen subgraph $T_0 \subset G$ a spanning tree of G. It is called $Mesh(G, T_0)$ and is a real, symmetric, $|E(G - T_0)| \times |E(G - T_0)|$ matrix.

Before giving the precise definition of the mesh matrix $Mesh(G, T_0)$ of a connected graph G with a choice $T_0 \subset G$ of spanning tree of G (see section 2), it will be helpful to recall two relevant items and state two theorems:

Item 1: An interesting and classically studied invariant of a graph G is the number of spanning trees in G [32]:

$$ST(G) := \# \ spanning \ trees \ of \ G.$$

Naturally, ST(G) = 0 if G is not connected. This invariant satisfies Tutte-type deletion-contraction formulas. More generally, if $H \subset G$ is a subgraph and N = |E(G - H)|, one may introduce a mild generalization, the polynomial

$$ST(G, H)(X) = \sum_{j=0}^{N} (-1)^{j} ST_{j}(G, H) X^{N-j}$$

with $ST_j(G, H)$ the number of spanning trees of G which contain precisely j edges of E(G - H). Again, see section 4, there are Tutte-type deletion-contraction formulas for

computing these polynomials.

Theorem 1.1 Let G be a connected graph and $T_0 \subset G$ a spanning tree in G, then the characteristic polynomial of the mesh matrix $Mesh(G, T_0)$ is specified by

$$det(X \cdot Id - Mesh(G, T_0)) = ST(G, T_0)(X - 1).$$

See theorem 2.2 for the complete theorem.

In view of the above remarks, this polynomial can be computed directly by deletioncontraction methods.

Item 2: It is claimed that $Mesh(G, T_0)$ is of the special form $Id + Y^t \cdot Y$ and

$$Y \cdot Y^t$$
, where Y is defined via (G, T_0) ,

is a natural generalization of the graph Laplacian $\Delta(H) = Deg - Adj$ of a graph H. This will be seen as follows:

Let H be a graph and W be a disjoint point. Form C(V(H)), the cone on the vertices V(H) of H from cone point W, by attaching an edge PW for each vertex P of H; in particular, the vertices of C(V(H)) is the union $V(H) \sqcup \{W\}$. Let the "H with cone on vertices added", be the union

$$C'(H) = H \sqcup C(V(H)),$$

i.e., the graph C'(H) has vertices $V(C'(H)) = V(H) \sqcup \{W\}$ and edges $E(C'(H)) = E(H) \sqcup \{PW \text{ for } P \text{ a vertex of } H\}$. Note that C'(H) is connected and C(V(H)) is a spanning tree of C'(H).

Theorem 1.2 Let H be a finite graph. The characteristic polynomials of the Kirchhoff graph Laplacian $\Delta(H) = Deg - Adj$ of a graph H and that of the mesh matrix of the pair (C'(H), C(V(H))) are related by

$$Mesh(C'(H),C(V(H))) = Id + Y^t \cdot Y \quad with \quad \Delta(H) = Y \cdot Y^t = \Delta(C'(H),C(V(H))).$$

In particular, $\Delta(H)$ has the same non-vanishing eigenvalues counted with multiplicities as $Y^t \cdot Y = Mesh(C'(H), C(V(H))) - Id$ and as $Y \cdot Y^t = \Delta(C'(H), C(V(H)))$.

See section 2.

By these means, one sees that the mesh Laplacian defined by $\Delta(G, T_0) = Y \cdot Y^t$ generalizes to the mesh setting the well known and studied Kirchhoff graph Laplacian [23] of a graph H, $\Delta(H)$.

In particular, in view of item # 1, the characteristic polynomial of the graph Laplacian of a graph H, namely $det(T \cdot Id - \Delta(H))$, acquires a Tutte-type deletion-contraction method of computation.

Recently, an elegant paper of F. Aliniaeifard, V. Wang, and S. Van Willigenburg, "Deletion-Contraction for a Unified Laplacian and Applications", [2], has appeared with

a deletion-contraction method for computing the characteristic polynomial of the graph Laplacian of a graph H among other valuable results. It proceeds by finding such a formula in terms of an extension $det(T \cdot \epsilon - \Delta(H))$ where ϵ is a diagonal matrix of vertex weights. The computation stays in this restricted context as opposed to the above which stays in the context defined for pairs (G, H) with G a graph and H a subgraph. The methods for these cases are analogous, but different.

The definition of the mesh matrix $Mesh(G, T_0)$ appears in section 2 along with the proof of theorem 1.2 in a more explicit form in theorem 2.3. Section 3 contains the proofs of theorems 1.1, 2.2, 2.4. In section 4, the Tutte-type deletion-contraction formulas for ST(G) and the polynomial ST(G, H)(X) are described. In section 5, by specializing theorem 2.2 to the case (C'(H), C(V(H))) and employing theorem 1.2, one recovers the all minors theorem for graphs; see theorem 5.2. Section 6 proves that $det(Mesh(G, T_0)) = ST(G, T_0)$ is the order of the torsion group $C_1(G; \mathbb{Z})/[Z_1(G; \mathbb{Z}) \oplus \Pi(B^1(G; \mathbb{Z}))]$ defined there. Section 7 introduces eigenvalue estimates for the smallest positive eigenvalue of $\Delta(G, T_0)$, a measure of the flux from T_0 to T_0 through $G - T_0$. The present authors' earlier work on mesh matrices on general C-W complexes appeared in [8]; section 8 below concerns a higher dimensional generalization of theorem 2.2 with a correction of a slightly misstated theorem of that paper.

2 Definition of the mesh matrix $Mesh(G, T_0)$:

The context of the mesh matrix is that of a pair (G, T_0) with G a connected graph and $T_0 \subset G$ a spanning tree of G.

Definition of the mesh matrix : $Mesh(G, T_0)$.

Now let T_0 be a choice of spanning tree of G. Let $e_1, \dots, e_{|E(G-T_0)|}$ be a listing of edges of $E(G-T_0)$ and $f_1, f_2, \dots, f_{|E(T_0)|}$ comprise the edges of $E(T_0)$.

Choose for each edge of G a direction, giving a representation $\vec{e_j} = \vec{e_j}[P_j, Q_j]$ with $\vec{e_j}$ going from vertex P_j to vertex Q_j and $\vec{f_k} = \vec{f_k}[R_k, S_k]$ going from vertex R_k to vertex S_k .

Consider the directed edge $\vec{e_j} \in E(G-T_0)$ with end points P_j, Q_j . Then P_j, Q_j are vertices of the spanning tree T_0 ; so if $P_j \neq Q_j$ there is a unique directed simple path from Q_j to P_j in the spanning tree T_0 , given by distinct vertices say $Q_j = R_{j,1}, R_{j,2}, \cdots, R_{j,n[j]} = P_j$ with $R_{j,t} \sim R_{j,t+1}$ and the edge $[R_{j,t} \to R_{j,t+1}]$ chosen in T_0 . Let $N = |E(G-T_0)|$.

Now define the mapping

$$D: C_{1}(E(G - T_{0}); \mathbb{R}) \to C_{1}(T_{0}; \mathbb{R})$$

$$by \ D(1 \cdot \vec{e_{j}}) = 0 \ if \ 1 \leq j \leq N \ and \ e_{j} \ is \ a \ loop$$

$$by \ D(1 \cdot \vec{e_{j}}[P_{j}, Q_{j}]) = \sum_{t=1}^{n[j]-1} \ 1 \cdot [R_{j,t} \to R_{j,t+1}] \ if \ 1 \leq j \leq N \ and \ P_{j} \neq Q_{j}.$$

As seen the 1-chain $D(1 \cdot \vec{e_j}[P_j, Q_k])$ has boundary $\partial(D(1 \cdot \vec{e_j})) = 1 \cdot P_j - 1 \cdot Q_j$ as the directed path starts at Q_j and ends at P_j .

In this manner one has a uniquely defined 1-chain for G, namely:

$$Z[j] = 1 \cdot \vec{e_j} + D(1 \cdot \vec{e_j}) \text{ for } j = 1, \dots, N = |E(G - T_0)|.$$

This 1-chain is a 1-cycle, it has boundary 0, and $D(1 \cdot \vec{e_j})$ supported in T_0 . If e_j is not a loop, this 1-cycle is a directed simple closed curve, it goes from P_j to Q_j in $E(G - T_0)$, then back through the spanning tree T_0 from Q_j to P_j .

Since Z[j] restricted to $E(G - T_0)$ becomes $1 \cdot \vec{e_j}$, the linear mapping defined by $1 \cdot \vec{e_j} \mapsto Z[j]$ is one to one. It maps $C_1(E(G - T_0); \mathbb{Z})$ into the integral 1-cycles, $Z_1(G; \mathbb{Z})$. Indeed, this gives an isomorphism:

$$C_1(E(G-T_0);\mathbb{Z}) \stackrel{\cong}{\to} Z_1(G;\mathbb{Z}).$$

In particular, these 1-cycles $\{Z[j]\}$ form an integral basis for the integral 1-cycles of G.

[Proof: For any 1-cycle $z = \sum_{j=1}^{N} b_j \cdot \vec{e}_j + \sum_{k=1}^{|E(T_0)|} c_k \cdot \vec{f}_k$ of G, the 1-chain $z - \sum_{j=1}^{N} b_j \cdot Z[j]$ is supported in T_0 and is a 1-cycle, so equals the zero chain.] The mesh matrix of Trent [30, 31] may be defined using this integral basis, $\{Z[j]\}$, of 1-cycles of G.

Having chosen a spanning tree T_0 of the graph G, and having made these directed edge choices, one defines the mesh matrix for the pair (G, T_0) via the intersection pairing on $C_1(G; \mathbb{R})$:

$$Mesh(G, T_0) = \{ \langle Z[j_1], Z[j_2] \rangle \mid 1 \leq j_1, j_2 \leq N = |E(G - T_0)| \}.$$

The basic theorem about the mesh matrix $Mesh(G, T_0)$ is the theorem of Trent evaluating its determinant. This is studied in more detail in [8] along with higher dimensional generalizations.

Theorem 2.1 (Trent [30, 31]) For a connected graph G, the determinant $det(Mesh(G, T_0))$ equals the number of spanning trees of G, or as in item # 1 equals ST(G).

More explicitly, expand the 1-cycle Z[k] as a column $|E(G)| \times 1$ vector, say z[k], in terms of the basis $\{1 \cdot \vec{e_j}\}$ and then $\{1 \cdot \vec{f_k}\}$ of $C_1(G; \mathbb{Z})$ and form the $|E(G)| \times N$ matrix of column vectors $Y[+] := (z[1], \cdots, z[N])$, then $Mesh(G, T_0) = Y[+]^t \cdot Y[+]$, As seen Y[+] has the special form $Y[+] = \begin{pmatrix} Id \\ Y \end{pmatrix}$ where Id denotes the $N \times N$ identity matrix and Y is the $|E(T_0)| \times |E(G - T_0)|$ matrix representing the linear mapping D.

As a consequence,

$$Mesh(G, T_0) = Y[+]^t \cdot Y[+] = Id + (Y^t \cdot Y) \text{ and}$$

 $charactistic polynomial of $Mesh(G, T_0) = det(X Id - Y[+]^t \cdot Y[+])$
 $= det((X - 1) Id - Y^t \cdot Y)$
 $= characteristic polynomial of (Y^t \cdot Y) \text{ evaluated at } (X - 1).$$

In particular, the eigenvalues of the reduced mesh matrix $Mesh^{\#}(G, T_0) \stackrel{def.}{:=} Y^t \cdot Y$ are real and non-negative, while those of the mesh matrix $Mesh(G, T_0)$ are obtained by adding +1 to those of the reduced mesh matrix.

The following theorem specifies the coefficients of the characteristic polynomial of the mesh matrix and the reduced mesh matrix of (G, T_0) . It is a restatement of theorem 1.1.

Theorem 2.2 Let G be a connected graph and T_0 a spanning tree of G. For $j = 1, \dots, N = |E(G - T_0)|$, denote by $ST_j(G, F)$ the number of spanning trees T of G such that $|E(T) \cap (E(G - T_0))| = j$. Then

$$det((U+1)Id - Mesh(G, T_0)) = det(U Id - Mesh^{\#}(G, T_0))$$

= $U^N + \sum_{j=1}^{N} (-1)^j ST_j(G, T_0) U^{N-j} = ST(G, T_0)(U).$

Setting U = -1 gives $det(Mesh(G, T_0)) = \sum_{j=0}^{N} ST_j(G, T_0) = ST(G)$ in agreement with Trent's theorem 2.1. Here $Y^t \cdot Y$ represents the reduced mesh matrix $Mesh^{\#}(G, T_0)$ and $Mesh(G, T_0) = Id + Y^t \cdot Y$.

In terms of the previously defined polynomials for H a subgraph of G listed in item # 1, section 1 ST(G,H)(U), this theorem evaluates the characteristic polynomials of the reduced mesh matrix $Y^t \cdot Y$ as equal to $ST(G,T_0)(U)$. In this larger context of pairs of graphs, (G,H) one has, see section 4, deletion-contraction formulas which may be used to compute this invariant.

It is natural to define the $|E(T_0)| \times |E(T_0)|$ matrix

$$Y \cdot Y^t$$
 as the Mesh Laplacian of $(G, T_0) = \Delta(G, T_0) = D \cdot D^*$.

This terminology is consistent with the standard notations in view of the following theorem. [See [15, 16] for a discussion of $\Delta(G) = Deg - Adj$ and its normalized version.]

As in item # 2, section 1, for a graph H, let $C'(H) = H \cup C(V(H))$ be H with the cone on vertices of H added,

Theorem 2.3 For a graph H, consider C'(H) and its subcone C(V(H)) with cone point W. Then C'(H) is connected and C(V(H)) is a spanning tree for C'(H).

Following the above construction, for each edge of H once directed, say

$$\vec{e_j} = \vec{e_j}[P_j \to Q_j], j = 1, \cdots, |E(H)|$$
 associate the 1-cycle, $Z[j]$

 $=1\cdot\vec{e_j}+1\cdot(Q_j\to W)-1\cdot(P_j\to W)$ of C'(H), then the associated matrix Y defined by

$$1 \cdot \vec{f_j} \mapsto D(1 \cdot \vec{f_j}) = 1 \cdot (Q_j \to W) - 1 \cdot (P_j \to W) \in C_1(C(V(H)); Z)$$

under the identification $C_1(C(V(H)); Z) \cong C_0(V(H); Z)$ given by $1 \cdot (P \to W) \mapsto 1 \cdot P$

becomes [under this identification] precisely the standard boundary mapping $\partial: C_1(H; Z) \to C_0(H; Z), \ \partial(1 \cdot \vec{f_j}) = 1 \cdot Q_j - 1 \cdot P_j \ of the graph H.$

Consequently, the mesh Laplacian of $(G, T_0) = \Delta(G, T_0)$ represented by $Y \cdot Y^t$ for this special case (C'(H), C(V(H))) becomes exactly the graph Laplacian matrix $\Delta(H) = \partial \cdot \partial^t = Deg - Adj$ of H.

Also
$$Mesh(C'(H), C(V(H)) = Id + Y^t \cdot Y = Id + Mesh^{\#}(C'(H), C(V(H))).$$

The significance of this theorem is that the mesh Laplacian matrix $Y \cdot Y^t$ representing $D \cdot D^*$ may be viewed as a natural generalization of the Kirchhoff graph Laplacian $\Delta(H)$ to the context of mesh matrices for pairs (G, T_0) .

Since eigenvalues of $Y \cdot Y^t$ and $Y^t \cdot Y$ are non-negative, and their non-vanishing eigenvalues are equal [counted with multiplicities] the characteristic polynomials of $Y \cdot Y^t$ and $= Y^t \cdot Y$ are related by

$$U^{|E(T_0)|} \cdot det(U \ Id - Y^t \cdot Y) = U^{|E(G-T_0)|} \cdot det(U \ Id - Y \cdot Y^t).$$

Applied to the case of C'(H) theorem 1.2 of item # 2 follows.

Additionally, the mesh Laplacian $\Delta(G, T_0) = Y \cdot Y^t$ has a nice direct description:

Theorem 2.4 Let the edges of $E(T_0)$ be directed and labeled as $\vec{f_k}$, $k = 1, \dots, |E(T_0)|$. Then the mesh Laplacian $\Delta(G, T_0) = Y \cdot Y^t$ is specified by

$$(Y \cdot Y^t)_{\vec{f_k}, \vec{f_k}} = \#\{e \in E(G - T_0) \text{ with } f_k \text{ in the support of } D(1 \cdot \vec{e})\},$$

for $k \neq l$

$$(Y \cdot Y^t)_{\vec{f_k}, \vec{f_l}} = Sign(k, l) \cdot \#\{e \in E(G - T_0) \text{ with } f_k \text{ and } f_l \text{ in the support of } D(1 \cdot \vec{e})\}$$

Here $Sign(k,l) = \pm 1$ for $k \neq l$ is defined by taking the unique directed path, γ , in T_0 from the edge f_k to the edge f_l . Let $Sign(k,l) = <1 \cdot \vec{f_k}, \vec{\gamma} > \cdot <1 \cdot \vec{f_l}, \vec{\gamma} > \cdot$

3 Proof of theorems 1.1, 2.2, 2.4:

Now the characteristic polynomial of $Mesh(G, T_0)$ represented by $Y[+]^t \cdot Y[+]$ is easily determined by the following method: Let

$$det(X \ Id - Y[+]^t \cdot Y[+]) = X^N + \sum_{j=1}^N (-1)^j \ b_j \ X^{N-j}.$$

Firstly, by expanding to get the coefficient of X^{N-j} , b_j equals the sum over $j \times j$ diagonal minors of $Y[+]^t \cdot Y[+]$ of the determinant det(minor). This sum may be organized as summing over a choice of j indices say $1 \le k_1 < k_2 < \cdots < k_j \le N$ [specifying the rows and corresponding columns of the diagonal block chosen].

More explicitly, let $Y[+][k_1, \dots, k_j]$ denote the $|E(G)| \times j$ matrix from Y[+] obtained by deleting all but these j specified columns. Then the desired block determinant equals

$$det(Y[+][k_1, \cdots, k_j]^t \cdot Y[+][k_1, \cdots, k_j]).$$

Now compare this last to the determinant of the mesh matrix for the subgraph $G[k_1, k_2, \dots, k_j]$ obtained from the spanning tree T_0 by adding in the j edges $e_{k_a}^{\vec{j}}$, a =

 $1, \dots, j$ of $E(G - T_0)$ to T_0 . Each of the 1-cycles $Z[k_a]$ lie in $G[k_1, k_2, \dots, k_j]$. As seen, $Y[+][k_1, \dots, k_j]^t \cdot Y[+][k_1, \dots, k_j] = Mesh(G[k_1, k_2, \dots, k_j]; T_0)$. Consequently, by the results of Trent:

#spanning trees of
$$G[k_1, \dots, k_j] = det(Y[+][k_1, \dots, k_j]^t \cdot Y[+][k_1, \dots, k_j])$$
.

So in total, the coefficient b_i is expressed in terms of spanning trees. More explicitly:

Lemma 3.1 Let G be a connected graph and T_0 be a spanning tree of G and $N = |E(G - T_0)|$. Then the characteristic polynomial of the mesh matrix $Mesh(G, T_0)$ is given by

$$det(X \ Id - (Y[+]^t \cdot Y[+]) = X^N + \sum_{j=0}^N (-1)^j \ b_j \ X^{N-j}$$

with b_j equal to the sum over j distinct indices $1 \le k_1 < k_2 < \cdots k_j \le N$ of the number of spanning trees in the subgraph $T_0 \cup (e_{k_1} \cup e_{k_2} \cup \cdots \cup e_{k_j})$.

Now consider this sum b_i in detail with reference to the integers

$$ST_j(M, T_0) = \#\{ \text{ spanning trees } T \text{ of } G \text{ with } |E(T) \cap E(G - T_0)| = j \}.$$

Each spanning tree T counted in $ST_j(G, T_0)$ appears exactly once in the counting of b_j as it fills in all the j added edges of $E(G - T_0)$.

Each spanning tree T of $ST_{j-1}(G, T_0)$ fills in j-1 edges of $E(G-T_0)$ so to get j one needs to chose 1 from the remaining N-(j-1) edges. Thus each spanning tree T of $ST_{j-1}(G, T_0)$ appears exactly $\binom{N-(j-1)}{1} = \binom{N-(j-1)}{N-j}$ times in b_j . Each spanning tree T of $ST_{j-2}(G, T_0)$ fills in j-2 edges of $E(G-T_0)$ so to get j one

Each spanning tree T of $ST_{j-2}(G, T_0)$ fills in j-2 edges of $E(G-T_0)$ so to get j one needs to chose 2 from the remaining N-(j-2) edges. Thus each spanning tree T of $ST_{j-2}(G, T_0)$ appears exactly $\binom{N-(j-2)}{2} = \binom{N-(j-2)}{N-j}$ times in b_j .

Combining these observations, proves the formula for $N \geq j \geq 1$:

$$b_j = \sum_{t=0}^{j} {N - (j-t) \choose N-j} ST_{j-t}(G, T_0).$$

It also holds for j = 0 since both sides equal +1. Inserting [and letting k = j - t] this gives:

$$\begin{split} \det(X\ Id - Mesh(G, T_0)) &= \Sigma_{j=0}^N\ (-1)^j\ b_j\ X^{N-j} \\ &= \Sigma_{j=0}^N\ (-1)^j \big[\Sigma_{t=0}^j\ \binom{N-(j-t)}{N-j}\ ST_{j-t}(G, T_0)\big]\ X^{N-j} \\ &= \Sigma_{k=0}^N\ (\Sigma_{j=k}^N\ (-1)^j\ \binom{N-k}{N-j}X^{N-j}\)\ ST_k(G, T_0) \\ &= \Sigma_{k=0}^N\ (-1)^k\ (\Sigma_{j=k}^N(-1)^{j+k}\ \binom{N-k}{N-j}X^{N-j})\ ST_k(G, T_0) \\ &= \Sigma_{k=0}^N\ (-1)^k\ (X-1)^{N-k}\ ST_k(G, T_0) = \det((X-1)\ Id - Mesh^\#(G, T_0)). \end{split}$$

Thus

$$det((U+1) Id - Mesh(G, T_0)) = det(U Id - Mesh^{\#}(G, T_0))$$

= $U^N + \sum_{i=1}^{N} (-1)^{i} ST_i(G, T_0) U^{N-j}$

as claimed.

Proof of theorem 2.4:

For directed edge e_j with end points directed by $P_j \to Q_j$ let $C[Q_j, P_j]$ be the unique simple path from Q_j to P_j in T_0 . By definition $D(1 \cdot \vec{e}_j[P_j, Q_j])$ is then the sum of these directed edges of $C[P_j, Q_j]$ regarded as a 1-chain.

Tautologically, the mapping D and adjoint D^* have the properties:

$$\begin{array}{ll} If \ P_j \neq Q_j, \ D(1 \cdot \vec{e_j}[P_j,Q_j]) = \sum_{f_k \ in \ support \ of \ C[Q_j,P_j]} &< 1 \cdot \vec{f_k}, D(1 \cdot \vec{e_j}[P_j,Q_j]) > \cdot \vec{f_k}; \\ D(1 \cdot \vec{e_j}[P_j,Q_j]) = 0 \ if \ e_j \ is \ a \ loop \ i.e., P_j = Q_j. \\ Similarly, \end{array}$$

if
$$f_k$$
 not a loop, $D^*(1 \cdot \vec{f_k}) = \sum_{f_k \text{ in support of } C[Q_j, P_j]} < 1 \cdot \vec{f_k}, D(1 \cdot \vec{e_j}[P_j, Q_j]) > \cdot \vec{e_j};$
 $D^*(1 \cdot \vec{f_k}) = 0$ if f_k is a loop.

Hence, by $\langle f, \Delta(G, T_0)(f) \rangle = \langle D^*(f), D^*(f) \rangle$, the mesh Laplacian matrix is defined by

$$\Delta(G, T_0)_{f_k, f_l} = \sum_{\substack{e_j \text{ with } f_k \text{ and } f_k \text{ in the support of } C[Q_j, P_j]} < 1 \cdot \vec{f_k}, D(1 \cdot \vec{e_j}[P_j, Q_j]) > \cdot < 1 \cdot \vec{f_l}, D(1 \cdot \vec{e_j}[P_j, Q_j]) > \cdot$$

For k = l, this give the desired result. For $k \neq l$, one may change the ordering of Q_j, P_j if necessary so that the unique path $C[Q_j, P_j]$ running from Q_j to P_j in T_0 containing the edges f_k , f_l gives the unique directed path, say $\vec{\gamma}$, in T_0 from f_k to f_l . In particular, for any e_j with f_k , f_l in $C[Q_j, P_j]$ one has the equality independent of j

$$<1 \cdot \vec{f_k}, D(1 \cdot \vec{e_j}[P_j, Q_j]) > \cdot <1 \cdot \vec{f_l}, D(1 \cdot \vec{e_j}[P_j, Q_j]) > = <1 \cdot \vec{f_k}, \vec{\gamma} > \cdot <1 \cdot \vec{f_l}, \vec{\gamma} >$$
, as desired.

4 Properties of ST(G, H)(X); Computation by deletion contraction methods:

It is well known that the number of spanning trees, SF(G), obeys nice formulas for deletion and contraction [32].

For a graph G and a edge e, let $G \setminus e$ be the graph obtained by deleting the open edge e preserving the end points, i.e, $V(G \setminus e) = V(G)$ and $E(G \setminus e) = E(G) - \{e\}$. Let G/e denote the graph obtained by contracting the end points of e along with all of the edge e to a single point.

As ST(G) counts spanning trees, ST(G) = 0 if G is disconnected.

As no tree contains a loop, $ST(G) = ST(G \setminus e)$ if e is a loop.

If e is an edge with distinct end points, then the spanning trees of G counted in SF(G) either do not contain e and are faithfully counted in $ST(G \setminus e)$ or contain the edge e. In this second case, the spanning trees of G/e are exactly the result of collapsing the edge e in a spanning tree of G containing e. In this manner, one gets the well known relations:

$$ST(G) = ST(G \setminus e) + ST(G/e)$$
 if e an edge of G is not a loop.

If e is an isthmus, that is all trees of G contain the edge e, then $G \setminus e$ is disconnected, so

$$ST(G) = ST(G \setminus e) + ST(G/e) = ST(G/e)$$
 if e is an isthmus.

By using these relations in the context of a graph G and a subgraph H with

$$ST(G,H)(X) = \sum_{j=0}^{N} (-1)^{j} ST_{j}(G,H) X^{N-j} \text{ and } N = |E(G-H)|$$

where $ST_j(G, H)$ is the number of spanning trees T of G with $|E(T) \cap (E(G) - E(H))| = j$, one obtains by counting:

$$ST(G,H)(X) = ST(G,H \setminus e)(X)$$
 if $e \in H$ is a loop;
 $ST(G,H)(X) = ST(G,H \setminus e)(X) + ST(G,H/e)(X)$ if $e \in H$ is not a loop;
 $ST(G,H)(X) = ST(G \setminus e,H)(X)$ if $e \in G$ is a loop;
 $ST(G,H)(X) = ST(G \setminus e,H)(X) - X \cdot ST(G/e,H)(X)$
if $e \in G$ is not a loop.

These deletion-contraction formulas give an inductive method for computing the polynomials SF(G, H).

By theorem 2.2, $det(X \cdot Id - Mesh^{\#}(G, T_0)) = ST(G, T_0)(X)$ so these formulas give a deletion-contraction method for computing the characteristic polynomial of the reduced mesh matrix $Mesh^{\#}(G, T_0)$ and the mesh matrix $Mesh(G, T_0) = Mesh^{\#}(G, T_0) + Id$.

As a special case, by theorem 2.3, the characteristic polynomial of the graph Laplacian $\Delta(H)$ of a graph H equals that of $\Delta(C'(H), C(V(H)))$ so these deletion-contraction formulas give a method that also applies to the graph Laplacian $\Delta(H)$.

Recall from item # 2, section 1, the paper [2] gives an alternative method to get deletion-contraction formulas for computation in these graph Laplacian cases.

5 Recapturing the All Minors Matrix Tree Theorem

The all minors matrix tree theorem, see [12, 13, 27], asserts:

Theorem 5.1 Let H be a finite graph. Let a spanning forest F in H consist of the union of K disjoint trees, say $F = \bigsqcup_{j=1}^K T[j], \subset H$ which spans, i.e., V(F) = V(H). Let a spanning forest F of K components have multiplicity mult $(F) = \prod_{j=1}^K |V(T[j])|$. Then the characteristic polynomial of the graph Laplacian $\Delta(H) = \partial \cdot \partial^* = Deg - Adj$ is given by

$$det(T \cdot Id - \Delta(H)) = T^{|V(H)|} + \sum_{j=1}^{|V(H)|-1} (-1)^{j} b_{j}(H) T^{|V(H)|-j}$$
with $b_{j}(H) = \sum_{F \ a \ spanning \ forest \ of \ H \ with \ |V(H)|-j \ components} \ mult(F)$.

Note: Let a spanning rooted forest consist of a spanning forest F and a choice of a vertex, say $P_j \in V(T[j])$, called roots for each j. Then

$$b_j(H) = \#$$
 spanning rooted forests in H with $|V(H)| - j$ components

Now by contrast consider the spanning trees T of $C'(H) = H \cup C(V(H))$, obtained from the graph H with the cone on the vertices added. Necessarily, the spanning tree T must contains the cone point, say W, and all vertices of H. Also, the intersection $F = T \cap H$ is a spanning forest in H, say $F = \bigsqcup_{j=1}^K T[j]$ with K components and each T[j] a tree in H. To complete the tree T in C'(H) each component of $F = \bigsqcup_{j=1}^K T[j]$ must be joined by exactly one edge of the form $P_j W$ with $P_j \in V(T[j])$. So one has the equality

```
#\{spanning trees T in C'(H) with | E(T) \cap C(V(H))| = K\}
#\{spanning rooted forests F in H with K components\}
= \sum_{spanning forests F in H with K components} mult(F).
```

By these means the all minors matrix tree theorem can be recast in the form

Theorem 5.2 (All minors matrix tree theorem):

```
\begin{split} \det(T \cdot Id - \Delta(H)) &= T^{|V(H)|} + \Sigma_{j=1}^{|V(H)|-1} \ (-1)^j \ b_j(H) \ T^{|V(H)|-j} \\ with \ b_j(H) &= \# \{spanning \ trees \ T \ in \ C'(H) \ with \ |E(T) \cap C(V(H))| = |V(H)| - j \} \\ &= \# \{spanning \ trees \ T \ in \ C'(H) \ with \ |E(T) \cap E(H)| = j \} \\ &= ST_j(C'(H), C(V(H))) \end{split}
```

The last equalities use the fact that a spanning tree T of C'(H) will have |V(C'(H))|-1 = |V(H)| edges.

Hence, one sees that the all minors matrix tree theorem is the special case with $(G, T_0) = (C'(H), C(V(H)))$ of theorem 2.2 in view of theorem 2.3.

6 # spanning trees as the order of a torsion group associated to G:

Let G be a connected graph with choices of directed edges as above. Let $Z_1(G; \mathbb{Z})$ be the integral 1-cycles, i.e., the kernel of the boundary mapping $\partial: C_1(G; \mathbb{Z}) \to C_0(G; \mathbb{Z})$; let $B^1(G; \mathbb{Z})$ be the 1-coboundaries i.e., the image of the dual coboundary mapping $\partial^*: C^0(G; \mathbb{Z}) = Hom(C_0(G; \mathbb{Z}), \mathbb{Z}) \stackrel{Hom(.,\partial)}{\to} C^1(G; \mathbb{Z}) = Hom(C_1(G; \mathbb{Z}), \mathbb{Z})$; let

 $\Pi: C^1(G; \mathbb{Z}) \to C_1(G; \mathbb{Z})$ be the isomorphism defined by $[F: E(G) \to \mathbb{Z}] \mapsto [\Sigma_{\sigma \in E(G)} F(\sigma) \cdot \vec{\sigma}]$ with chosen direction $\vec{\sigma}$ on an edge σ of G.

Theorem 6.1 The inclusion of integral chains $[Z_1(G; \mathbb{Z}) \oplus \Pi(B^1(G; \mathbb{Z}))] \subset C_1(G; \mathbb{Z})$ is a rational equivalence; so the quotient $U = C_1(G; \mathbb{Z}) / [Z_1(G; \mathbb{Z}) \oplus \Pi(B^1(G; \mathbb{Z}))]$ is a finite abelian group. The order of this group is

$$\mid U \mid = \mid C_1(G; \mathbb{Z}) \mid [Z_1(G; \mathbb{Z}) \oplus \Pi(B^1(G; \mathbb{Z}))] \mid = \# \text{ spanning trees of } G = ST(G)$$

First, observe that the orthogonal complement of $Z_1(G; \mathbb{R})$ is $\Pi(B^1(G; \mathbb{R}))$, so one gets an isomorphism of real vector spaces $Z_1(G; \mathbb{R}) \oplus \Pi(B^1(X; \mathbb{R})) = C_1(G; \mathbb{R})$.

Proof: If $\langle ., . \rangle$ is the standard inner product on $C_1(X; \mathbb{R})$, then one has for $a \in Z_1(G, \mathbb{R}), b \in C_0(G; \mathbb{R}), \langle a, \Pi(\delta(b)) \rangle = \delta(b)(a) = b(\partial a)$. Hence, $\partial a = 0$ if and only if a is orthogonal to $\Pi(B^1(G; \mathbb{R}))$.

By this result, the associated embedding of lattices has finite quotient,

$$U = C_1(G; \mathbb{Z}) / [Z_1(G; \mathbb{Z}) \oplus \Pi(B^1(G; \mathbb{Z}))].$$

Next, let T_0 be a choice of spanning tree for G. Following the notation of section 2, let the directed edges of $G-T_0$ be denoted by $\vec{e_j}, j=1,\cdots,|E(G-T_0)|$. Let the directed edges of T_0 be denoted by $\vec{f_k}, k=1,\cdots,|E(T_0)|$.

As above, consider the integer 1-cycles $Z[j], j = 1, \dots, |E(G - T_0)|$. Here the 1-cycle

$$Z[j] = 1 \cdot \vec{e}_j + D(\vec{e}_j), \ j = 1, \dots, |E(G - T_0)|$$

has $D(\vec{e}_j)$ supported in the tree T_0 . As above, define the $|E(T_0)| \times |E(G - T_0)|$ matrix $Y = \{A[k, j]\}$ by

$$Z[j] = 1 \cdot \vec{e_j} + \sum_{k=1}^{|E(T_0)|} Y[k,j] \vec{f_k}, \quad j = 1, \dots, |E(G - T_0)|.$$

Here the coefficients Y[k, j] are either 0 or ± 1 . As proved above, these Z[j]'s form an integral basis for $Z_1(G; \mathbb{Z})$. That is, part a) below is known.

Lemma 6.2 a) The 1-chains

$$Z[\vec{e}_j] = 1 \cdot \vec{e}_j + D(\vec{e}_j) = \vec{e}_j + \sum_{k=1}^{|E(T_0)|} A[k,j] \vec{f}_k, \quad j = 1, \dots, |E(G - T_0)|$$

form an integral basis for $Z_1(G; \mathbb{Z})$.

b) The 1-chains

$$b[\vec{f_k}] = 1 \cdot \vec{f_k} - \sum_{j=1}^{|E(G-T_0)|} Y[j, k] \ \vec{e_j}, \ k = 1, \dots, |E(T_0)|$$

form an integral basis for $\Pi(B^1(G;\mathbb{Z}))$.

Using the direct sum decomposition $C_1(G; \mathbb{Z}) = C_1(G - T_0; \mathbb{Z}) \oplus C_1(T_0; \mathbb{Z})$, write elements in column form: $C_1(G; \mathbb{Z}) = \begin{pmatrix} C_1(G - T_0; \mathbb{Z}) \\ C_1(T_0; \mathbb{Z}) \end{pmatrix}$. As noted, the $\vec{e}_j \mapsto Z[\vec{e}_j]$ giving an integral basis of $Z_1(G; Z)$ is represented by the

As noted, the $\vec{e}_j \mapsto Z[\vec{e}_j]$ giving an integral basis of $Z_1(G;Z)$ is represented by the matrix $\begin{pmatrix} Id \\ Y \end{pmatrix}$ with Y an integer valued matrix above. Granting lemma 6.2 part b), the mapping $\vec{f}_k \mapsto b[\vec{f}_k]$ giving an integral basis of $\Pi(B^1(G;\mathbb{Z}))$ is represented by the matrix $\begin{pmatrix} -Y^t \\ Id \end{pmatrix}$.

Thus,
$$|U| = det \begin{pmatrix} Id & -Y^t \\ Y & Id \end{pmatrix} = det \begin{pmatrix} Id + Y^t & Y & -Y^t + Y^t \\ Y & Id \end{pmatrix} = det(Id + Y^t & Y) = det(Mesh(G, T_0) = ST(G))$$
 by Trent's theorem 2.1.

Proof of lemma 6.2 part b):

The first part about the 1-cycles Z[j] has already been proved in section 2.

Pick a directed edge in T_0 , say $\vec{f_k} = \vec{f_k}[R_k \to S_k]$. Recall that the spanning tree T_0 is connected.

Decompose the set of vertices of T_0 into two disjoint non-empty sets $V(T_0) = A[k] \sqcup B[k]$ where A[k] consists of the vertices of T_0 which can be connected to the initial vertex R_k by a simple path not including f_k and B[k] consists of the vertices of T_0 which can be connected to the final vertex S_k by a simple path not including f_k . For convenience, chose the directions of the edges \vec{e}_j [which by assumption range over $E(G - T_0)$] so that if \vec{e}_j has one end point in A[k] and the other in B[k], then \vec{e}_j is directed from the point in A[k] to the point in B[k].

Now recall that for a vertex P of G, the coboundary $\delta(1 \cdot P)$ has the property that $\Pi(\delta(X))$ is the 1-chain consisting of all edges with distinct end points containing P as a end point and directed [inwards] towards P.

One easily sees that $\Pi(\delta(\Sigma_{b \in B[k]} \ 1 \cdot b))$ consists of a sum of two parts. Firstly $1 \cdot \vec{f_k}$; and secondly a sum over all edges $\vec{e_j}$ in $G - T_0$ with distinct end points and having one end in A[k] and the other in B[k]. This is usually called the cut defined by $\vec{f_k}$. The edges $\vec{e_j}$ with this property have been directed as going from A[k] to B[k].

Now consider the 1-cycle Z[j] and its evaluation $\langle Z[j], 1 \cdot \vec{f_k} \rangle$; it equals zero unless $\vec{e_j}$ has one end in A[k] and the other in B[k]. It has sign -1 as $\vec{e_j}$ goes from A[k] to B[j]. Hence,

$$\Sigma_j < z[j], 1 \cdot \vec{f_k} > \vec{e_j} = -\Sigma_{summed \ as \ below} \ Y[j, k] \ \vec{e_j}$$

with the above sum is only over the directed edges \vec{e}_j of $G - T_0$ going from A[k] to B[k] with direction from A[k] to B[k]. Note Y[j,k] = 0 otherwise.

In total then

$$\Pi(\delta(\Sigma_{b \in B[k]} \ 1 \cdot b) = 1 \cdot \vec{f_k} - \Sigma_j \ Y[j, k] \ \vec{e_j}.$$

Thus there is a well defined mapping

$$F: C_d(T_0; Z) \to \Pi(B^d(X; \mathbb{Z})), \quad \vec{f_k} \mapsto 1 \cdot \vec{f_k} - \Sigma_j Y[j, k] \vec{e_j}.$$

Now let a be any element of $\Pi(B^1(X; Z))$ and note that

$$< a, Z[j] >= 0 \text{ as } Z[j] \in Z_1(X; \mathbb{Z}) \text{ and } \Pi(B^1(X; \mathbb{R}))^{\perp} = Z_1(G; \mathbb{R}).$$

Say $a = (\Sigma_j x_j \vec{e}_j) + (\Sigma_k y_k \vec{f}_k)$. Then the difference $Diff = a - (\Sigma_k y_k F(\vec{f}_k))$ lies in $\Pi(B^d(X;\mathbb{Z}))$ and is supported in $G - T_0$. But then $0 = \langle Z[e_j], Diff \rangle = x_j$ for each $j = 1, \dots, |E(G - T_0)|$ proving that Diff = 0. Hence, the integral chains $F(\vec{f}_k)$ generate $\Pi(B^d(X;\mathbb{Z}))$ They are independent since projecting to $C_1(T_0;\mathbb{Z})$ each goes to $1 \cdot \vec{f}_k$, part of the basis of $C_1(T_0;\mathbb{Z})$.

Hence, the lemma is proved.

7 Flux and Eigenvalue Inequalities

For the Kirchoff Laplacian $\Delta(G)$ of a graph G there is the foundational Rayleigh-Ritz inequality providing an upper estimate of the first positive eigenvalue. Following the principle that the mesh Laplacian is a natural generalization of the Kirchoff Laplacian, it is natural to discuss an extension of such inequalities to the mesh context. The flux of the of the graph G is a measured by the eigenvalues of the Kirchoff Laplacian.

The mesh Laplacian $\Delta(G, T_0) = D \cdot D^*$ is represented by $Y \cdot Y^t$ which is real symmetric with non-negative eigenvalues. Let Λ be the smallest positive eigenvalue of $Y \cdot Y^t$, equivalently of $\Delta(G, T_0)$. This Λ is also the smallest positive eigenvalue of $Y^t \cdot Y$ which represents the reduced mesh matrix $Mesh^{\#}(G, T_0)$.

By the standard Rayleigh-Ritz theorem, there is the associated equality giving upper estimates for Λ . Namely,

$$\Lambda = \min_{f \neq 0, f \in [kernel \ C_1(T_0; \mathbb{R})]^{D^{\star}} C_1(G - T_0; \mathbb{R})]^{\perp}} \frac{\langle D^{\star}(f), D^{\star}(f) \rangle}{\langle f, f \rangle}$$

where $[kernel\ C_1(T_0;\mathbb{R}) \xrightarrow{D^*} C_1(G-T_0;\mathbb{R})]^{\perp}$ is the orthogonal complement of this kernel in $C_1(T_0;\mathbb{R})$ endowed with the standard cellular inner product. Here $< D^*(f), D^*(f) > = < f, D \cdot D^*(f) > = < f, \Delta(G,T_0)(f) >$. Hence, the kernel consists precisely of the eigenvectors of $\Delta(G,t_0)$ with eigenvalue zero.

One may regard Λ as a measure of the **flux** of $\Delta(G, T_0)$ through $G - T_0$ from T_0 to T_0 .

Let the edges in $E(G-T_0)$ be enumerated and directed as before. Then there are three different types of directed edges. Firstly $\vec{e}_j[P_j,Q_j]$ may be a loop, i.e., $P_j=Q_j$. Let these be called type 1 edges. Secondly, the edge \vec{e}_j may have distinct end points, but these end points P_j, Q_j [which lie in $V(T_0) = V(G)$] are adjacent in the subgraph T_0 . Call these type 2 edges. Note that in the first case $D(1 \cdot \vec{e}_j[P_j,Q_j]) = 0$ and in the second case $D(1 \cdot \vec{e}_j[P_j,Q_j])$ is the unique directed edge in T_0 from Q_j to P_j . Finally and thirdly, the directed edge $\vec{e}_j[P_j,Q_j]$ is called of type 3, if it has a directed simple path in T_0 from Q_j to P_j of length ≥ 2 .

In the third type 3 case, $\vec{e_j}[P_j, Q_j]$, P_j, Q_j are vertices of the spanning tree T_0 so by $P_j \neq Q_j$ there is a unique directed simple path from Q_j to P_j in the spanning tree T_0 , given by distinct vertices say $Q_j = R_{j,1}, R_{j,2}, \cdots, R_{j,n[j]} = P_j$ with $R_{j,t} \sim R_{j,t+1}$ [adjacent] and the edge $[R_{j,t} \to R_{j,t+1}]$ chosen in T_0 . Here $n[j] \geq 2$, so one has two distinct well defined directed edges, initial and final directed inward into T_0 given by

$$[Q_j \to R_{j,2}] = [R_{j,1} \to R_{j,2}] \in E(T_0) \text{ and } [R_{j,n[j]} \to R_{j,n[j]-1}] = [P_j \to R_{j,n[j]-1}] \in E(T_0)$$

For a directed edge $\vec{e}_j[P_j, Q_j] \in E(G - T_0)$ of type 3, let these directed two directed edges be denoted by $F_1(\vec{e}_j[P_j, Q_j]) = [Q_j \to R_{j,2}] \in E(T_0)$, the initial edge of $D(\vec{e}_j[P_j, Q_j])$, and

¹An inequality giving a lower estimate on the eigenvalues is due to Cheeger for Riemannian manifolds [14]. Bounds either way in the setting of graph theory are often referred to as Cheeger type inequalities inspired by his methods, i.e., see [15, 16].

 $F_2(\vec{e_j}[P_j,Q_j]) = [P_j \to R_{j,n[j]-1}] \in E(T_0)$, the final edge of $D(\vec{e_j}[P_j,Q_j])$ both directed inward to T_0 .

Let $\vec{e}_j[P_j,Q_j]$ range over all type 3 directed edges of $E(G-T_0)$, set X to be the set of all these inward directed edges so obtained:

$$X = \{F_1(\vec{e_j}[P_j, Q_j])\} \cup \{F_2(\vec{e_j}[P_j, Q_j])\}.$$

Naturally a given such directed edge may arise in several ways; multiplicities are neglected in this representation. The construction provides a set of distinct type 3 inward directed edges of G.

Now form a graph W with vertices the elements of the set X of the type 3 edges. Declare that two distinct vertices, say with representatives $[F_r(\vec{e}_s[P_s,Q_s])], [F_t(\vec{e}_u[P_u,Q_u])],$ having an edge in the graph W between them for each type 3 directed edge $\vec{e}_j[P_j,Q_j]$ with ingoing edges equal to these two, i.e., $\{F_1(\vec{e}_j[P_j,Q_j]), F_2(\vec{e}_j[P_j,Q_j])\}$ = $\{F_r(\vec{e}_s[P_s,Q_s]), F_t(\vec{e}_u[P_u,Q_u])\}.$

Theorem 7.1 Let λ be the smallest positive eigenvalue of the graph Laplacian $\Delta(W)$ of the graph W derived from G. Let Λ be the smallest positive eigenvalue of the mesh Laplacian $\Delta(G, T - 0)$. Then the following inequality holds:

$$\Lambda \leq \lambda$$

By the standard methods, [15, 16] λ has upper estimates of the Cheeger type. These then imply upper estimates for Λ .

In detail, let the graph W have M distinct connected components, say

$$W = \sqcup_{k=1}^{M} W[k]$$
 and correspondingly vertex decomposition $X = V(W) = \sqcup_{k=1}^{M} X[k]$.

For each k make a choice of proper subset $C[k] \subset X[k]$ and denote the complement in X[k] by D[k] = X[k] - C[k], then

$$\Lambda/2 \leq \lambda/2$$

 $\leq min\{\frac{\#\vec{e_j}[P_j,Q_j] \text{ of type 3 with } F_1(\vec{e_j}[P_j,Q_j]) \in C[k] \text{ and } F_2(\vec{e_j}[P_j,Q_j]) \in D[k] \text{ or visa versa}}{min(|A[k]|,|B[k])}, k = 1, \cdots, M\}$

Proof: Let $j: R[X] \subset C_1(T_0; \mathbb{R})$ denote the inclusion of the real sums of multiples of the [inward directed] edges in X. Let $\overline{D}^* = D^* \cdot j: R[X] \to C_1(G - T_0; \mathbb{R})$ with adjoint $(\overline{D}^*)^*$. Let μ be the smallest positive eigenvalue of the composite $\overline{\Delta} = (\overline{D}^*)^* \cdot \overline{D}^*$. Tautologously, one has the inequality [restricting the minimum over fewer cases]:

$$\Lambda \leq \min_{g \neq 0, g \in R[X] \cap [kernel C_1(T_0; \mathbb{R}) \xrightarrow{D^{\star}} C_1(G - T_0; \mathbb{R})]^{\perp}} \frac{\langle D^{\star}(g), D^{\star}(g) \rangle}{\langle g, g \rangle} = \mu$$

Now here $R[X] \cap [kernel\ C_1(T_0; \mathbb{R}) \xrightarrow{D^*} C_1(G - T_0; \mathbb{R})]$ consists of the sums $Sum = \sum_{x \in X} A[x] x$ for which $D^*(Sum) = 0$. That is, for each $\vec{e_j}[P_j, Q_j]$ the equation $< D(\vec{e_j}[P_j, Q_j]), Sum >= 0$ holds. But $< D(\vec{e_j}[P_j, Q_j]), Sum >= < F_2(\vec{e_j}[P_j, Q_j]) -$

 $F_1(\vec{e}_j[P_j,Q_j]), Sum >= A[F_2(\vec{e}_j[P_j,Q_j])] - A[F_1(\vec{e}_j[P_j,Q_j])].$ This means that if the vertices in W, x, y are adjacent, then A[x] = A[y]. Consequently, $R[X] \cap [kernel\ C_1(T_0;\mathbb{R}) \xrightarrow{D^*} C_1(G-T_0;\mathbb{R})]$ consists of the sum of multiples of the entries one for each component

$$S[k] = (\Sigma_{x \in W[k]} \ 1 \cdot x) \in C_1(T_0; \mathbb{R}), \ k = 1, \dots, M$$

and the mapping $(\hat{D}^*)^*$: $C_1(G - T_0; \mathbb{R}) \to \{\Sigma_{x \in X} \ A[x] \ x\}$ sends $1 \cdot \vec{e_j}(P_j, Q_j) = F_1(\vec{e_j}[P_j, Q_j]) - F_2(\vec{e_j}[P_j, Q_j])$.

That is, under this identification, $(\overline{D}^*)^*$ is identified with $\partial: C_1(W;\mathbb{R}) \to C_0(W;\mathbb{R})$ and \overline{D}^* with $\partial^*: C_0(W;\mathbb{R}) \to C_1(W;\mathbb{R})$ and the kernel of ∂^* with the sums of multiples of the S[k]'s. The orthogonal complement of this kernel is identified with the span of the eigenspaces with positive eigenvalues. In particular, $\mu = \lambda$ is given as the least positive eigenvalue of the graph Laplacian of the graph W in the above inequality.

The standard Cheeger inequality [15, 16] now applies to the graph W to give the above estimate.

8 Appendix; A Correction to the statement of theorem 7.1 of [8] and extension of theorem 2.2 to the higher dimensional context:

The paper [8] considers generalizations of the mesh matrix to the context of d-dimensional finite CW-complexes X. The theorem 7.1 of [8], once stated correctly, is a generalization of lemma 3.1. Once lemma 3.1 is generalized, the methods of this paper combined with [8] also give a generalization of theorem 2.2 above. This is carried out in this section. In the context of the higher dimensional Kirchoff Laplacian for CW-complexes some relevant papers are [1, 3, 4, 2, 9, 10, 11, 17, 18, 19, 24, 25, 29] and others.

Let X be a finite d-dimensional CW-complex. Let S_d be the d-cells of X with chosen orientations, i.e., X is obtained from the d-1 skeleton of X, i.e., X^{d-1} , by attaching the oriented d-cells of S_d . For $V \subset S_d$ define the subcomplex

$$X_V = X^{d-1} \cup V \subset X^{d-1} \cup S_d = X.$$

For a finite cell complex, say X, let $t_d(X)$ denote the order of the torsion subgroup of the $(d-1)^{th}$ integral homology of X, $H_{d-1}(X;\mathbb{Z})$.

In this higher dimensional context a subset $V \subset S_d$ is called a **spanning forest** of X if the composition

$$C_d(X_V; \mathbb{R}) \stackrel{\partial_d}{\to} B_{d-1}(X_V; \mathbb{R}) \subset B_{d-1}(X; \mathbb{R})$$

is an isomorphism. In the case that G is a graph and the graph G is connected, the spanning forests in the above sense are exactly the spanning trees of the graph G. In the case that a graph G has K connected components, say $G = \bigsqcup_{j=1}^K G[j]$, a spanning forest is a choice for each connected component G[j] of a spanning tree T[j] of G[j]. It is shown in [8] that all d-dimensional CW-complexes have spanning forests.

Now the integral d-cycles $Z_d(X; \mathbb{Z})$ is a free abelian group, so one may pick a integral basis, say $\{z_i\}$. Via the natural pairing

$$Z_d(X; \mathbb{Z}) \times Z_d(X; \mathbb{Z}) \subset C_d(X; \mathbb{Z}) \times C_d(X; \mathbb{Z}) \stackrel{\langle \dots, \rangle}{\to} \mathbb{Z},$$

one may form the mesh matrix for this integral choice of basis

$$Mesh(X; \{z_r\}) = \{\langle z_a, z_b \rangle\}$$

On the other hand, if one chooses a spanning forest, say V_0 , of X, for each e in $S_d - V_0$, the boundary $\partial_d(e)$ is **uniquely expressed** as $\partial_d(D(e))$ for a unique d-chain $D(e) \in C_d(X_{V_0}; \mathbb{R})$. In particular, for each $e \in S_d - V_0$ there the unique d-cycle

$$z(e) = 1 \cdot e - D(e) \in Z_d(X; \mathbb{R})$$

with D(e) supported exactly in X_{V_0} . Since $Z_d(X_{V_0}; \mathbb{R}) = 0$ by V_0 a spanning tree, one sees that these $\{z(e) \mid e \in S_d - V_0\}$ form a "geometric" basis for $Z_d(X; \mathbb{R})$ with associated mesh matrix

$$Mesh(X; \{z(e) \mid e \in S_d - V_0\}, geometric)$$

Note that z(e) is supported in $X_e = X^{d-1} \cup e$ for $e \in S_d - V_0$. Let $N = |S_d - V_0|$.

Let the *d*-cells in $S_d - V_0$, be enumerated via e_1, \dots, e_N . Consider a choice of $j \times j$ diagonal minor, say $Mesh(\{z(e)\}, geometric)_{k_1,k_2,\dots,k_j}$ with $1 \leq k_1 < k_2 < \dots < k_j \leq N$ by picking these j rows and columns. The entries of this minor all are of the form $< z(e_{k_a}), z(e_{k_b}) >$ with each $z(e_{k_a}) \in Z_d(X_{V_0 \cup \{e_{k_1}, e_{k_2}, \dots, e_{k_j}\}}); \mathbb{R})$ for $1 \leq 1 \leq j$. Consequently, as in §3,

$$det(Mesh(\{X(e)\}), geometric)_{k_1, k_i, \cdots, k_j}) = det(Mesh(X_{V_0 \cup \{e_{k_1}, \cdots, e_{k_j}\}}), geometric).$$

Consequently, modeled on the proof of lemma 3.1, one has for the geometric basis the equality

$$det(X \cdot Id - Mesh(X; \{z(e) \mid in \ S_d - V_0\}, geometric) = X^N + \sum_{j=1}^N (-1)^j \ b_j \ X^{N-j})$$

with $b_j = \sum_{1 \leq k_1 < k_2, \dots, k_j \leq N} det(Mesh(X_{V_0 \cup \{e_{k_1}, \dots, e_{k_j}\}}))$, geometric)). Here the choice of k_1, k_2, \dots, k_j in $S_d - V_0$ may be recorded as a choice of subset $U = \{k_1, \dots, k_j\} \subset S_d - V_0$ with |U| = j.

In these terms the slightly corrected version of theorem 7.1 of [8] is: [It is the high dimensional version of lemma 3.1.]

Theorem 8.1 Let $V_0 \subset S_d$ be a d-dimensional spanning forest of a d-dimensional CW-complex X with oriented d-cells S_d , then $\sigma_j(X; V_0) =$ the coefficient of $(-1)^j T^{N-j}$ in the characteristic polynomial

$$det(Mesh(X, \{z(e) \mid e \in (S_d - V_0)\}), geometric\ basis)$$

is given by

$$\sigma_j(X; V_0) = \Sigma_{U \subset (S_d - V_0) \text{ with } |U| = j} \quad \left(\Sigma_{V \text{ a spanning forest of } X_{V_0 \cup U}} \left(\frac{t_{d-1}(X_V)}{t_{d-1}(V_0)} \right)^2 \right)$$

For d=1, there is no torsion in the homologies, so sum is over entries all equal to +1. So, this is the high dimensional version of lemma 3.1. The proof is a consequence of the above argument and the following equality: For Y a d-dimensional finite CW-complex, and a choice of spanning forest $V_0 \subset S_d$ for Y.

$$det(Mesh(Y, \{z(e)\}, geometric) = \sum_{V \ a \ spanning \ forest \ of \ Y} \left(\frac{t_{d-1}(Y_V)}{t_{d-1}(Y_{V_0})}\right)^2. \quad (\star)$$

applied to the various $Y = X_{V_0 \cup U}$.

This last is the analog of Trent's theorem 2.1 above. In the paper [8] theorem 2.1 asserts the evaluation of the determinant of the mesh matrix for the **integral basis** $\{z_r\}$ as:

$$det(Mesh(X; \{z_r\})) = \sum_{V \text{ a spanning forest of } X} \left(\frac{t_{d-1}(X_V)}{t_{d-1}(X)}\right)^2 \quad (\star \star)$$

This equality $(\star \star)$ is to be modified to get (\star) which utilizes the geometric basis rather than the geometric basis.

Here under the mapping $Z_d(X;\mathbb{R}) \subset C_d(X;\mathbb{R}) \to C_d(X,X_{V_0};\mathbb{R})$ [which by V_0 a spanning forest is an isomorphism] the lattice of the geometric basis $Z\{z(e) \mid e \in S_d - V_0\}$ has image equal to the image of the integral classes $C_d(X;\mathbb{Z})$, [as $z(e) = 1 \cdot e + D(e)$ with $D[e] \in C_d(X_{V_0};\mathbb{R})$]; and the lattice of the integral basis for $Z_d(X;\mathbb{Z})$ has image equal of course to the image of the integral basis of $Z_d(X;\mathbb{Z})$. As V_0 is a spanning forest, these images provide two real bases for the $C_d(X,X_{V_0};\mathbb{R})$ with one lattice inside the other. Hence, the change of basis from integral to geometric has determinant equal to the order of the finite abelian group:

$$\frac{C_d(X, X_{V_0}; \mathbb{Z})}{image \ Z_d(X; \mathbb{Z})}.$$

The order of this finite abelian group is identified on page 14 of [8] as

$$\left| \frac{C_d(X, X_{V_0}; \mathbb{Z})}{Z_d(X; \mathbb{Z})} \right| = \frac{t_{d-1}(X_{V_0})}{t_{d-1}(X)}$$

The respective determinants of mesh matrices are related by the square of this quotient. This relation demonstrates that $(\star \star)$ implies (\star) so the above forms an independent proof of theorem 8.1.

Now as above in the present paper, the geometric basis is of the form: $z(e_j) = 1 \cdot e_j + D(e_j) \in Z_d(X; \mathbb{R})$ for $e_j \in S_d - V_0$ with $D(e_j)$ supported in X_{V_0} . This implies

$$\langle z(e_i), z(e_k) \rangle = Id_{i,k} + \langle D(e_i), D(e_k) \rangle.$$

That is, the geometric mesh matrix is of the form,

$$Mesh(X; \{z(e_j)\}, geometric) = Id + Mesh^{\#}(X; \{z(e_j)\}, geometric)$$

with the reduced mesh matrix $Mesh^{\#}(X; \{z(e_j)\}, geometric)$ defined by $\{\langle D(e_j), D(e_k) \rangle\}$. Hence, the counting argument given for proving theorem 2.2 of this paper applies without essential change to yield a higher dimensional generalization of this paper's theorem 2.2. Namely:

Theorem 8.2

$$\begin{split} &\det((U+1)\ Id - Mesh(X; \{z(e_a)\}, geometric)) \\ &= \det(U\ Id - Mesh^{\#}(X; \{z(e_a)\}, geometric) = U^N + \Sigma_{j=1}^N\ (-1)^j\ c_j\ U^{N-j} \\ &with \\ &c_j = \Sigma_{U\ a\ spanning\ forest\ of\ X\ with\ |U\cap(S_d-V_0)| = j\ \left(\frac{t_{d-1}(X_U)}{t_{d-1}(X_{V_0})}\right)^2 \end{split}$$

References

- [1] R. M. Adin, Counting colorful multi-dimensional trees, Combinatorica 12 (1992), no. 3, pp. 247-260.
- [2] F. Aliniaeifard, V. Wang, S. Van Willigenburg, Deletion-Contraction for a Unified Laplacian and Applications, https://arxiv.org/abs/2110.13949, to appear.
- [3] O. Bernardi, C. J. Klivans, Directed rooted forests in higher dimension, Electron. J. Combin. 23 (2016), no. 4, 20 pp.
- [4] E. D. Bolker, Simplicial goemetry and transportation polytopes, Trrans. Amere. Math. Soc, 217 (1976), pp. 121-142
- [5] P. R. Bryant, Graph theory applied to electrical networks in F. Harary (ed.), Graph theory and theoretical physics 1967, Academic Press, N.Y., pp. 111-137.
- [6] Y. Burman, B. Shapiro, Around matrix-tree theorem. Math. Res. Lett. 13 (2006), no. 5-6, pp. 761-774.
- [7] Y. Burman, A. Ploskonosov, A. Trofimova, Matrix-tree theorems and discrete path integration. Linear Algebra Appl. 466 (2015), pp. 64-82.
- [8] S. Cappell, E. Miller, Enumerative Combinatorics of Simplicial and Cell Complexes: Kirchoff and Trent Type Theorems, Discrete and Combinatorial Geometry (2019), vol 61, 1-41.
- [9] M. Catanzaro, V. Chernyak, and J. Klein, On Kirchhoff's theorems with coefficients in a line bundle, Homology, Homotopy and Applications, vol. 15 no. 2 (2013), 267 280.

- [10] M. Catanzaro, V. Chernyak, and J. Klein, Kirchhoff's Theorems in Higher Dimensions and Reidemeister Torsion, Homology Homotopy Appl. (2015), vol 17, # 1, 165-189.
- [11] M. J. Catanzaro, V. Y. Chernyak, and J. R. Klein, A higher Boltzmann Distribution, J. Appl. Comp. Topol. 1, (2017), 215-240.
- [12] S. Chaiken, A Combinatorial Proof of the All Minors Matrix Tree Theorem, SIAM J. Alg. Disc. Maht, (1982), vol 3, No. 3, pp. 319-339.
- [13] S. Chaiken and D. Kleitman, Matrix tree theorems, J. of Combinatorial Theory, Series A, 24 (1978), pp. 377-381.
- [14] Cheeger, Jeff, A lower bound for the smallest eigenvalue of the Laplacian, Problems in analysis (Sympos. in honor of Salomon Bochner, Princeton Univ., Princeton, N.J., 1969), pp. 195–199. Princeton Univ. Press, Princeton, N.J., 1970.
- [15] F. Chung, Spectral Graph Theory, American Mathematical Society, Regional Conference Series in Mathematics, (1997) # 92.
- [16] F. Chung, Four Proofs of the Cheeger Inequality and graph partition algorithms, Proceeding of ICCM (2007).
- [17] A. Duval, C. Klivans, and J. Martin, Simplicial Matrix Tree Theorems, Trans. Amer. Math. Soc., 361 (2009), pp. 6073-6114.
- [18] A. Duval, C. Klivans, J. Martin, Simplicial and cellular trees. Recent trends in combinatorics, IMA Vol. Math. Appl., 159 (2016), pp. 713-752, Springer Press.
- [19] G. Kalai, Enumeration of Q-acyclic Simplicial Complexes, Israel J. Math., vol 45 (1983), pp. 337-351.
- [20] R. Kenyon, The asymptotic determinant of the discrete Laplacian, Acta Math., 185 (2000), pp. 239-286.
- [21] Kenyon, R. The Laplacian and Dirac operators on critical planar graphs, Invent. Math. 150 (2002), no. 2, pp. 409-439.
- [22] R. Kenyon, Spanning forests and the vector bundle laplacian, The Annals of Probability, 39 no. 5 (2011), pp. 1983-2017.
- [23] G. Kirchhoff, Uber die Aufläung der Gleichungen, auf welche man bei der Untersuchung der linearen Verteilung galvanischer Strome gerfuhrt wird, Poggendorf's Ann. Phys. Chem., vol. 72 (1847), pp. 497-508; English translation, IRE Trans. Circuit Theory, vol. 5 (1958), pp. 4-8.
- [24] R. Lyons, Random Complexes and L^2 -Betti numbers,, J. Topol. Anal. vol. 1 (2009), pp. 153-175.
- [25], R. Lyons, and Y.Peres, Probability on trees and networks. Cambridge Series in Statistical and Probabilistic Mathematics, 42. Cambridge University Press, New York, 2016.

- [26] G. Masbaum, Gregor, A. Vaintrob, A new matrix-tree theorem. Int. Math. Res. Not. 27 (2002), pp. 1397-1426.
- [27] S. Maurer, Matrix Generalizations of Some Theorems on Trees, Cycles, and Cocycles in Graphs, SIAM J. Math., vol 30 (1976), # 1, 143-148.
- [28] J. W. Moon, Counting Labeled Trees, Canadian Mathematical Congress, From lectures delivered to the Twelfth Biennial Seminar of the Canadian Mathematical Congress (Vancouver, 1969). Canadian Mathematical Monographs, No. 1 Canadian Mathematical Congress, Montreal, Que. 1970.
- [29] A. Peterson, Enumerating Spanning Trees in Simplicial Complexes, Masters Thesis Uppsala University, 2009.
- [30] H. M. Trent, A Note on the Enumeration and Listing of All Possible Trees in a Connected Linear Graph, Proc. National Acad. Sci,(1954) vol 40, # 10, 1004-1007.
- [31] H. Trent, A note on the enumeration and listing of all maximal trees of a connected linear graph, Proc. Nat. Acad. Sci. U.S.A., 40 (1954), pp. 1004-1007.
- [32] W. Tutte, On the spanning trees of self-dual maps, in Second International Conference on Combinatorial Mathematics (A. Gewirtz and L/V. Quintas, eds.), The New York Academy of Science, 1979, pp. 540-548.
- Sylvain E. Cappell, cappell@courant.nyu.edu, Courant Institute of Mathematical Sciences, NYU.
- Edward Y. Miller, emiller@cims.nyu.edu, , Courant Institute of Mathematical Sciences, NYU.