

Modified Friedmann Equations from Maxwell-Weyl Gauge Theory

Salih Kibaroglu*

Maltepe University, Faculty of Engineering and Natural Sciences, 34857, Istanbul, Turkey and

Institute of Space Sciences (IEEC-CSIC) C. Can Magrans s/n, 08193 Barcelona, Spain

(Dated: May 24, 2023)

This study investigates the possibility of a homogeneous and isotropic cosmological solution within the context of the Maxwell-Weyl gauge theory of gravity. To achieve this, we utilize the Einstein-Yang-Mills theory as an analogy and represent the Maxwell gauge field in terms of two time-dependent scalar fields. Then, we present the modified Friedmann equations, which incorporate the contributions of the Maxwell gauge field, as well as the effective cosmological constant, which is a function of the Dirac scalar field.

PACS numbers: 02.20.Sv, 11.15.-q, 98.80.-k, 95.36.+x

Keywords: Cosmology, Maxwell symmetry, Scale invariance

I. INTRODUCTION

The general theory of relativity, proposed by Albert Einstein, is a fundamental theory of gravity that forms the basis for our current understanding of cosmology. One of the most successful theories in cosmology is the Λ CDM model which describes the universe as being composed of dark matter, dark energy, and baryonic matter. It assumes that the universe is homogeneous and isotropic on large scales and that it began with a hot and dense Big Bang. However, there are several reasons we need to generalize current cosmological models. Addressing some of the most fundamental questions in physics, such as the nature of dark matter and dark energy, the origin and evolution of the universe, and the potential existence of other universes or dimensions beyond our own. By exploring these questions and developing more sophisticated models, we can gain deeper insights into the nature of reality and our place in the cosmos.

In this context, the gauge theory of gravity provides a powerful framework to reach a more comprehensive cosmological model by extending the gravitation theory to include additional fields and symmetries. In the 1950s and 1960s, physicists such as Yang-Mills, Utiyama, and Kibble began exploring the application of gauge theory to the study of fundamental forces, including gravity [1–3]. They developed the concept of gauge invariance, which played a key role in the development of the Standard Model of particle physics. In this framework, gravity is described as a curvature of space-time, and the force of gravity is seen as a manifestation of the curvature. This has important implications for cosmology, as it implies that the large-scale structure of the universe is intimately connected to the curvature of space-time. For example, the gauge theory of gravity, which is based on the Weyl group, constitutes a potent framework for understanding the properties and behavior of the universe on its largest scales, and scale invariance plays a key role in this framework [4–8].

In the pursuit of constructing cosmological models rooted in the gauge theory of gravity, one of the main challenges is the suitable implementation of gauge fields. The literature has previously examined gauge field configurations in the Friedmann-Lemaître-Robertson-Walker (FLRW) cosmology [9–16]. One area of focus is the non-Abelian Yang-Mills (YM) theories, where the coupling of a non-Abelian YM field to the scalar curvature is analyzed to determine the cosmological implications of nonminimal gravitational coupling [15, 17–19] (for a comprehensive review, see [20]). In

* salihkibaroglu@maltepe.edu.tr

addition to its use in inflationary cosmology, YM theory is applied in models for dark energy [13, 14, 21–26] and dark matter [27–29].

An intriguing extension of the theory of gravity can be derived from the gauge theory of the Maxwell algebra. The Maxwell algebra can be interpreted as a modified version of the Poincaré algebra, incorporating six additional tensorial abelian generators that render the four-momenta non-commutative [30–32],

$$[P_a, P_b] = iZ_{ab}. \quad (1)$$

The Maxwell algebra provides a useful background for constructing a gauge theory of gravity, resulting in a generalized theory that encompasses the cosmological constant and an additional energy-momentum tensor term [33–46]. This gravitational model is referred to as the Maxwell gauge theory of gravity or Maxwell gravity, and while the energy-momentum term remains less explored, on the other hand, it is known that such an additional term may be related to dark energy [47, 48]. A minimal cosmological model with Maxwell symmetry is introduced in the literature [49], and it is also suggested that the gauge fields of this symmetry group may provide a geometric background for describing vector inflatons in cosmological models [50]. Furthermore, there is another study that proposes a cosmological model for the Maxwell gravity which is constructed by gauging the semi-simple extended Poincaré algebra [51].

In this work, we extend our previous study [51] to include the scale invariance which arises naturally in the Maxwell-Weyl gauge theory of gravity [37] to find more general cosmological solutions. Because the concept of scale invariance is crucial to our understanding of cosmology, it provides a foundation for many of the theoretical frameworks and models that are used to describe the universe on its largest scales [8]. Our objective is to achieve this goal by utilizing a method that is similar in nature to the one used for the realization of gauge fields in terms of time-dependent scalar fields. This method relies on the application of the Einstein-Yang-Mills theory and its cosmological implications.

The present manuscript is structured as follows. In Section II, we present a concise overview of the Maxwell-Weyl gauge theory of gravity and the geometric interpretation of the Weyl transformation. In Section III, we investigate the potential solutions for the gravitational field equations arising from the framework of Maxwell-Weyl gravity and give the modified version of the Friedmann equations. Finally, the last section provides conclusive remarks and discussions.

II. MAXWELL-WEYL GAUGE THEORY OF GRAVITY

This section provides a concise overview of the findings from our prior investigation [37]. The mathematical framework of the Maxwell-Weyl algebra, denoted as \mathcal{MW} , may be regarded as a Weyl extension of the Maxwell algebra that encompasses a dilatation generator [52].

The non-zero commutation relations of \mathcal{MW} algebra are given by,

$$\begin{aligned} [M_{ab}, M_{cd}] &= i(\eta_{ad}M_{bc} + \eta_{bc}M_{ad} - \eta_{ac}M_{bd} - \eta_{bd}M_{ac}), \\ [P_a, P_b] &= i\lambda Z_{ab}, \\ [M_{ab}, P_c] &= i(\eta_{bc}P_a - \eta_{ac}P_b), \\ [M_{ab}, Z_{cd}] &= i(\eta_{ad}Z_{bc} + \eta_{bc}Z_{ad} - \eta_{ac}Z_{bd} - \eta_{bd}Z_{ac}), \\ [P_a, D] &= iP_a, \\ [Z_{ab}, D] &= 2iZ_{ab}, \end{aligned} \quad (2)$$

where η_{ab} is the Minkowski metric which has $\text{diag}(\eta_{ab}) = (+, -, -, -)$, the constant λ has the unit of L^2 which will be related to the cosmological constant where L is considered as the unit of length and the indices $a, b, \dots = 0, \dots, 3$. Here, M_{ab} is the Lorentz generator, P_a is the translation generator, D is the dilatation generator, and Z_{ab} is the anti-symmetric Maxwell symmetry generator. In addition to the Weyl algebra, \mathcal{MW} algebra contains six new additional

tensorial generators Z_{ab} .

Let us start to construct the gauge theory of gravity based on \mathcal{MW} algebra. We first introduce a one-form gauge field that possesses the following form,

$$\mathcal{A} = e^a P_a + B^{ab} Z_{ab} + \chi D - \frac{1}{2} \omega^{ab} M_{ab}, \quad (3)$$

where the coefficient fields $e^a(x) = e_\mu^a(x) dx^\mu$, $B^{ab}(x) = B_\mu^{ab}(x) dx^\mu$, $\chi(x) = \chi_\mu(x) dx^\mu$ and $\omega^{ab}(x) = \omega_\mu^{ab}(x) dx^\mu$ are the one-form gauge fields for the corresponding generators, respectively. The transformation of these fields under infinitesimal gauge transformations is given by $\delta\mathcal{A} = -d\zeta - i[\mathcal{A}, \zeta]$ with the gauge generator,

$$\zeta(x) = y^a(x) P_a + \varphi^{ab}(x) Z_{ab} + \sigma(x) D - \frac{1}{2} \tau^{ab}(x) M_{ab}, \quad (4)$$

where $y^a(x)$ are space-time translations, $\varphi^{ab}(x)$ are translations in tensorial space, $\sigma(x)$ dilatation parameter and $\tau^{ab}(x)$ are the Lorentz transformation parameters. Thus we get,

$$\begin{aligned} \delta e^a &= -dy^a - \omega_b^a y^b - \chi y^a + \sigma e^a + \tau_b^a e^b, \\ \delta B^{ab} &= -d\varphi^{ab} - \omega_c^{[a} \varphi^{cb]} - 2\chi \varphi^{ab} + \frac{1}{2} e^{[a} y^{b]} + 2\sigma B^{ab} + \tau_c^{[a} B^{cb]}, \\ \delta \chi &= -d\sigma, \\ \delta \omega^{ab} &= -d\tau^{ab} - \omega_c^{[a} \tau^{cb]}, \end{aligned} \quad (5)$$

where the antisymmetrization and symmetrization of the objects are defined by $A_{[a} B_{b]} = A_a B_b - A_b B_a$ and $A_{(a} B_{b)} = A_a B_b + A_b B_a$, respectively.

Using the structure equation $\mathcal{F} = d\mathcal{A} + \frac{i}{2} [\mathcal{A}, \mathcal{A}]$ and defining the curvatures as $\mathcal{F} = \mathcal{F}^a P_a + \mathcal{F}^{ab} Z_{ab} + f D - \frac{1}{2} \mathcal{R}^{ab} M_{ab}$, we find the associated two-form curvatures as,

$$\begin{aligned} \mathcal{F}^a &= de^a + \omega_b^a \wedge e^b + \chi \wedge e^a, \\ \mathcal{F}^{ab} &= dB^{ab} + \omega_c^{[a} \wedge B^{c]b]} + 2\chi \wedge B^{ab} - \frac{\lambda}{2} e^a \wedge e^b, \\ f &= d\chi, \\ \mathcal{R}^{ab} &= d\omega^{ab} + \omega_c^a \wedge \omega^{cb}. \end{aligned} \quad (6)$$

The infinitesimal gauge transformation of the curvatures can be found by the help of the equation $\delta\mathcal{F} = i[\zeta, \mathcal{F}]$ as follows

$$\begin{aligned} \delta\mathcal{F}^a &= \omega_b^a \mathcal{F}^b + \sigma \mathcal{F}^a - \mathcal{R}_b^a y^b - f y^a, \\ \delta\mathcal{F}^{ab} &= \omega_c^{[a} \mathcal{F}^{c]b]} + 2\sigma \mathcal{F}^{ab} + \frac{\lambda}{2} \mathcal{F}^{[a} y^{b]} - \mathcal{R}_c^{[a} \varphi^{c]b]} - \frac{\lambda}{2} f \varphi^{ab}, \\ \delta f &= 0, \\ \delta\mathcal{R}^{ab} &= \omega_c^{[a} \mathcal{R}^{c]b]}. \end{aligned} \quad (7)$$

By performing the covariant derivative of the given curvatures, one can derive the corresponding Bianchi identities in the following manner

$$\mathcal{D}\mathcal{F}^a = \mathcal{R}_b^a \wedge e^b + f \wedge e^a,$$

$$\begin{aligned}
\mathcal{D}\mathcal{F}^{ab} &= \mathcal{R}^{[a}_c \wedge B^{c|b]} + 2f \wedge B^{ab} - \frac{\lambda}{2}\mathcal{F}^{[a} \wedge e^{b]}, \\
\mathcal{D}f &= 0, \\
\mathcal{D}\mathcal{R}^{ab} &= 0,
\end{aligned} \tag{8}$$

where $\mathcal{D}\Phi = [d + \omega + w(\Phi)\chi]\Phi$ is the Weyl gauge covariant derivative and w being the Weyl weight of the corresponding field.

A. Geometrical interpretation of Weyl's gauge theory

In our scenario, constructing locally invariant Lagrangians it is necessary to consider the scale invariance and its connection to the dilatation subgroup of \mathcal{MW} . Achieving localization of the dilatation symmetry leads us back to the Weyl gauge theory (for more details, see [8, 53–55]). For this reason, we want to give a brief discussion of the geometrical interpretation of the Weyl gauge theory.

The Weyl space and the Riemann space are two different mathematical structures used to describe the geometry of space-time in the context of general relativity and related theories. One of the key differences is the conformal invariance. The Weyl space has conformal invariance, which means that its geometry is invariant under changes in scale. In contrast, the Riemann space does not have conformal invariance and its geometry is not invariant under scale transformations.

The main point of the geometric interpretation is the identification of $e^\mu_a(x)$ as the components of a vierbein system in the Weyl space-time. This identification implies that the components of the space-time metric tensor and the vierbein field are related by the requirement that

$$g_{\mu\nu} = \eta_{ab} e^\mu_a e^\nu_b, \tag{9}$$

where $\mu, \nu = 0, \dots, 3$ are space-time indices and η_{ab} is the Minkowski metric which satisfies the following relation,

$$\eta_{ab} = g_{\mu\nu} e^\mu_a e^\nu_b. \tag{10}$$

If one utilizes the gauge transformation for $e^a(x)$ as given in Eq.(5), it can be demonstrated that the transformation of the space-time metric tensor satisfies the subsequent expression,

$$\delta g_{\mu\nu}(x) = 2\sigma(x) g_{\mu\nu}(x), \tag{11}$$

and the Minkowski metric satisfies $\delta\eta_{ab} = 0$ under infinitesimal gauge transformations. Moreover, it is known that the infinitesimal scale transformations which act on a local generic field $\Phi(x)$ can be expressed as,

$$\delta\Phi(x) = w\sigma(x)\Phi(x), \tag{12}$$

where $\sigma(x)$ is local parameter for the scale transformation and w is the “Weyl weight” (or simply “weight”) of scalar field $\Phi(x)$ (for more detail about the Weyl transformation see [4, 5, 7, 53, 56, 57]). Thus, the space-time metric and vierbein field have the Weyl weights $w(g_{\mu\nu}) = 2$, $w(g^{\mu\nu}) = -2$, $w(\sqrt{-g}) = 4$, and $w(e^\mu_a) = 1$, respectively, and so (9) and (10) imply that $w(\eta_{ab}) = 0$, as expected.

In this geometry, the affine connection in the Weyl-Cartan space-time is defined by

$$\Gamma_{\mu\nu}^{*\lambda} = \tilde{\Gamma}_{\mu\nu}^{\lambda} + \tilde{K}_{\mu\nu}^{\lambda}, \quad (13)$$

where $\tilde{\Gamma}_{\mu\nu}^{\lambda} = \tilde{\Gamma}_{\nu\mu}^{\lambda}$ is the Weyl connection which reads

$$\begin{aligned} \tilde{\Gamma}_{\mu\nu}^{\lambda} &= \frac{1}{2}g^{\lambda\rho} \left(\tilde{\partial}_{\mu}g_{\nu\rho} + \tilde{\partial}_{\nu}g_{\mu\rho} - \tilde{\partial}_{\rho}g_{\mu\nu} \right) \\ &= \Gamma_{\mu\nu}^{\lambda} + \delta_{\nu}^{\lambda}\chi_{\mu} + \delta_{\mu}^{\lambda}\chi_{\nu} - g_{\mu\nu}\chi^{\lambda}, \end{aligned} \quad (14)$$

where $\tilde{\partial}_{\mu} = \partial_{\mu} + w\chi_{\mu}$ and $\Gamma_{\mu\nu}^{\lambda} = \frac{1}{2}g^{\lambda\rho} (\partial_{\mu}g_{\nu\rho} + \partial_{\nu}g_{\mu\rho} - \partial_{\rho}g_{\mu\nu})$ is the well-known Lorentz connection (or the Riemann-Christoffel connection). The term $\tilde{K}_{\mu\nu}^{\lambda} = -\tilde{K}_{\nu\mu}^{\lambda}$ in (13) is the Weyl-Cartan contortion tensor, which is given in terms of the Weyl-Cartan torsion $\tilde{T}_{\mu\nu}^{\lambda} = \Gamma_{[\nu\mu]}^{*\lambda}$ by

$$\tilde{K}_{\mu\nu}^{\lambda} = -\frac{1}{2} \left(\tilde{T}_{\mu\nu}^{\lambda} - \tilde{T}_{\mu}^{\lambda}{}_{\nu} + \tilde{T}_{\mu\nu}^{\lambda} \right). \quad (15)$$

It is clear from (13)–(15) that setting the torsion to zero ($\tilde{T}_{\mu\nu}^{\lambda} = 0$) then the Weyl-Cartan space-time reduces to the Weyl space-time. Together with this condition, if dilatation gauge field $\chi_{\mu} \rightarrow 0$, then $\tilde{\Gamma}_{\mu\nu}^{\lambda} \rightarrow \Gamma_{\mu\nu}^{\lambda}$ and the Weyl geometry becomes Riemannian and the whole geometry is completely defined by the metric tensor $g_{\mu\nu}(x)$.

Within the context of Weyl space-time, wherein torsion is absent, the covariant differentiation of a vector A^{ν} formed with Weyl connection may be expressed in the following manner,

$$\begin{aligned} \tilde{\nabla}_{\mu}A^{\alpha} &= \partial_{\mu}A^{\alpha} + \tilde{\Gamma}_{\mu\beta}^{\alpha}A^{\beta} \\ &= \nabla_{\mu}A^{\alpha} + \delta_{\mu}^{\alpha}\chi_{\beta}A^{\beta} + A^{\alpha}\chi_{\mu} - A_{\mu}\chi^{\alpha}, \end{aligned} \quad (16)$$

where $\nabla_{\mu}A^{\alpha} = \partial_{\mu}A^{\alpha} + \Gamma_{\mu\beta}^{\alpha}A^{\beta}$ is the covariant derivative for diffeomorphisms in the Riemann geometry. With the help of Eq.(16), the covariant derivative of the metric tensor is conjectured to be nonzero

$$\tilde{\nabla}_{\rho}g_{\mu\nu} = -2\chi_{\rho}g_{\mu\nu} = Q_{\rho\mu\nu}, \quad (17)$$

where the new geometric quantity $Q_{\rho\mu\nu}$ is called the nonmetricity and Eq.(17) is also called “semi-metricity” condition. Moreover, if we make use of the total covariant derivative applied to the vierbein field $e_{\mu}^a(x)$, we get

$$\mathcal{D}_{\mu} \left(w + \tilde{\Gamma} \right) e_{\nu}^a = \tilde{\partial}_{\mu}e_{\nu}^a - \tilde{\Gamma}_{\nu\mu}^{\sigma}e_{\sigma}^a + \omega_{b\mu}^a e_{\nu}^b = 0, \quad (18)$$

in which this relation is sometimes known as the “tetrad postulate” and demonstrates a connection between $\omega_{b\mu}^a$, $\tilde{\Gamma}_{\mu\nu}^{\lambda}$ and χ_{μ} . Now, it is straightforward to show that $\omega_{b\mu}^a$ or $\tilde{\Gamma}_{\mu\nu}^{\lambda}$ may be written explicitly in terms of the other by using Eq.(18) as follows,

$$\tilde{\Gamma}_{\mu\nu}^{\lambda} = e_a^{\lambda} \left(\tilde{\partial}_{\mu}e_{\nu}^a + \omega_{b\mu}^a e_{\nu}^b \right), \quad (19)$$

and

$$\omega_{b\mu}^a = e_{\lambda}^a \left(\tilde{\partial}_{\mu}e_b^{\lambda} + \tilde{\Gamma}_{\nu\mu}^{\lambda}e_b^{\nu} \right), \quad (20)$$

where we used $e_\lambda^a \tilde{\partial}_\mu e_c^\lambda = -e_c^\lambda \tilde{\partial}_\mu e_\lambda^a$ because $\tilde{\partial}_\mu (e_\lambda^a e_c^\lambda) = \tilde{\partial}_\mu (\delta_c^a) = 0$. If we substitute (14) in (20),

$$\omega_{b\mu}^a = \tilde{\omega}_{b\mu}^a + (e_\mu^a e_b^\nu - e^{a\nu} e_{b\mu}) \chi_\nu, \quad (21)$$

we get the Weyl spin connection in terms of e_μ^a , χ_ν and $\tilde{\omega}_{b\mu}^a$. Here $\tilde{\omega}_{b\mu}^a$ represents the standard spin connection for a torsionless theory which is defined as

$$\tilde{\omega}_{b\mu}^a = e_\lambda^a (\partial_\mu e_b^\lambda + \Gamma_{\nu\mu}^\lambda e_b^\nu), \quad (22)$$

or by using the definition of the Lorentz connection $\Gamma_{\nu\mu}^\lambda$, it can be also written as

$$\tilde{\omega}_{\mu}^{ab} = \frac{1}{2} [e^{\nu a} (\partial_\mu e_\nu^b - \partial_\nu e_\mu^b) - e^{\nu b} (\partial_\mu e_\nu^a - \partial_\nu e_\mu^a) - e_{\mu c} e^{\nu a} e^{\lambda b} (\partial_\nu e_\lambda^c - \partial_\lambda e_\nu^c)]. \quad (23)$$

Subsequently, by utilizing Eq.(21) and (22), one can express the gauge covariant derivative for a vector A^a of weight w as follows.

$$\begin{aligned} \mathcal{D}_\mu A^a &= \partial_\mu A^a + w \chi_\mu A^a + \omega_{b\mu}^a A^b \\ &= \partial_\mu A^a + w \chi_\mu A^a + e_\lambda^a (\partial_\mu e_b^\lambda + \Gamma_{\nu\mu}^\lambda e_b^\nu) A^b + (e_\mu^a e_b^\nu - e^{a\nu} e_{b\mu}) \chi_\nu A^b, \end{aligned} \quad (24)$$

thus, we have formulated the covariant derivative in terms of e_μ^a , $\Gamma_{\mu\nu}^\lambda$ and χ_μ .

From this theoretical background, just as in the Riemannian geometry, we can use the Weyl connection to compute a Weyl invariant curvature tensor as follows

$$\begin{aligned} \mathcal{R}_{\mu\nu\rho}{}^\sigma &= \partial_\nu \tilde{\Gamma}_{\mu\rho}^\sigma - \partial_\mu \tilde{\Gamma}_{\nu\rho}^\sigma + \tilde{\Gamma}_{\mu\rho}^\alpha \tilde{\Gamma}_{\alpha\nu}^\sigma - \tilde{\Gamma}_{\nu\rho}^\alpha \tilde{\Gamma}_{\alpha\mu}^\sigma \\ &= R_{\mu\nu\rho}{}^\sigma + \left(\delta_{[\mu}^\sigma \nabla_{\nu]} \chi_\rho - g_{[\mu\rho} \nabla_{\nu]} \chi^\sigma - \delta_\rho^\sigma f_{\mu\nu} \right) \\ &\quad + \left(g_{\rho[\mu} \chi_{\nu]} \chi^\sigma - \delta_{[\mu}^\sigma \chi_{\nu]} \chi_\rho + \delta_{[\mu}^\sigma g_{\nu]\rho} \chi^\alpha \chi_\alpha \right), \end{aligned} \quad (25)$$

where $R_{\mu\nu\rho}{}^\sigma$ is the usual Riemann tensor and $f_{\mu\nu} = \partial_\mu \chi_\nu - \partial_\nu \chi_\mu$ is the Weyl invariant field strength. From this equation, it is possible to establish a Weyl invariant Ricci tensor as,

$$\mathcal{R}_{\mu\rho} = \mathcal{R}_{\mu\sigma\rho}{}^\sigma = R_{\mu\rho} - (g_{\mu\rho} \nabla_\sigma \chi^\sigma + 2 \nabla_\mu \chi_\rho + f_{\mu\rho}) + 2 (\chi_\mu \chi_\rho - g_{\mu\rho} \chi^\alpha \chi_\alpha), \quad (26)$$

where $R_{\mu\rho}$ is the Ricci tensor for the Riemannian geometry. Finally, one finds that the scalar curvatures \mathcal{R} of Weyl geometry and R of the Riemannian geometry are related by

$$\mathcal{R} = g^{\mu\rho} \mathcal{R}_{\mu\rho} = R - 6 \nabla_\mu \chi^\mu - 6 \chi_\mu \chi^\mu. \quad (27)$$

B. Gravitational action

The formulation of a Lagrangian based on \mathcal{MW} algebra differs somewhat from the ordinary Maxwell algebra because \mathcal{MW} algebra contains the scale transformations. In the gauge theory of gravity based on \mathcal{MW} algebra, the well-known action $S = \int d^4x \sqrt{-g} R$ is not invariant under the local scale transformations. To overcome this difficulty, Weyl introduced quadratic terms into the action to establish a consistent theory [58]. Furthermore, Brans and Dicke constructed a Lagrangian with the combination of the scalar curvature \mathcal{R} and a new compensating scalar field $\phi(x)$

[59]. Subsequently, Dirac employed a scalar field $\phi(x)$ with a Weyl weight of -1 and constructed the action using $\phi^2 \mathcal{R}$ rather than \mathcal{R} [6] (for more information, see [60, 61]).

These theoretical foundations were used to introduce the gravitational action in the Maxwell-Weyl gauge theory in [37], following Dirac's convention. In this study, a scalar field $\phi(x)$ (Dirac scalar field) is introduced which transforms as $\delta\phi = -\sigma\phi$ to make the Einstein-Hilbert action to be local scale invariant (for more detail, see [6, 37]).

In the present context, it is possible to introduce a shifted curvature as,

$$\mathcal{J}^{ab} = \mathcal{R}^{ab} + 2\mu\phi^2 \mathcal{F}^{ab}, \quad (28)$$

where μ is an arbitrary constant. Using this definition, one can express a gravitational action with zero Weyl weight in the following manner [37],

$$S = \int \frac{1}{2\kappa\lambda\mu} \mathcal{J} \wedge^* \mathcal{J} + \frac{1}{4} f \wedge^* f - \frac{1}{2} \mathcal{D}\phi \wedge^* \mathcal{D}\phi + \frac{\varsigma}{4} \phi^4 * 1, \quad (29)$$

where $\kappa = 8\pi G$ and ς are constants and $*$ represents the Hodge duality operator. The equations of motion can be found by taking variations of Eq.(29) with respect to gauge fields $\omega^{ab}(x)$, $e^a(x)$, $B^{ab}(x)$, $\chi(x)$, and $\phi(x)$, respectively,

$$\varepsilon_{abcd} \mathcal{D}\mathcal{J}^{ab} - 2\mu\phi^2 \varepsilon_{abcd} \mathcal{J}_e^{[a} \wedge B^{eb]} = 0, \quad (30)$$

$$\begin{aligned} & -\frac{\phi^2}{\kappa} \varepsilon_{abcd} \mathcal{J}^{ab} \wedge e^d + \frac{1}{2} [\mathcal{D}_c \phi^* \mathcal{D}\phi + \mathcal{D}\phi \wedge^* (e_a \wedge e_c) \mathcal{D}^a \phi] \\ & - \frac{1}{2} (f_{cb} e^b \wedge^* f - \frac{1}{2} \varepsilon_{abcd} f^{ab} e^d \wedge f) + \frac{\varsigma\phi^4}{2} * e_c = 0, \end{aligned} \quad (31)$$

$$\varepsilon_{abcd} \mathcal{D}(\phi^2 \mathcal{J}^{ab}) = 0, \quad (32)$$

$$\frac{\phi^2}{\kappa\lambda} \varepsilon_{abcd} \mathcal{J}^{ab} \wedge B^{cd} + \frac{1}{2} \mathcal{D}^* f + \phi^* \mathcal{D}\phi = 0, \quad (33)$$

$$\frac{\phi}{\kappa\lambda} \varepsilon_{abcd} \mathcal{J}^{ab} \wedge \mathcal{F}^{cd} + \mathcal{D}^* \mathcal{D}\phi + \varsigma\phi^3 * 1 = 0. \quad (34)$$

These equations are invariant with respect to the local Maxwell-Weyl transformation considered above. By a straightforward calculation, passing from tangent space to world indices for Eq.(31), one gets the following field equation with cosmological term depending on the dilaton field,

$$\mathcal{R}_\alpha^\mu - \frac{1}{2} \delta_\alpha^\mu \mathcal{R} + 3\mu\lambda\phi^2 \delta_\alpha^\mu = -2\mu\phi^2 T_{B\alpha}^\mu - \kappa\phi^{-2} \left(T_{\phi\alpha}^\mu + \frac{1}{2} T_{f\alpha}^\mu \right), \quad (35)$$

where

$$T_{B\alpha}^\mu = e_a^\mu e_b^\beta \mathcal{D}_{[\alpha} B_{\beta]}^{ab} - \frac{1}{2} \delta_\alpha^\mu \left(e_a^\rho e_b^\sigma \mathcal{D}_{[\rho} B_{\sigma]}^{ab} \right),$$

$$\begin{aligned}
T_{\phi}^{\mu}{}_{\alpha} &= \mathcal{D}^{\mu}\phi\mathcal{D}_{\alpha}\phi - \frac{1}{2}\delta_{\alpha}^{\mu}\left(\mathcal{D}_{\gamma}\phi\mathcal{D}^{\gamma}\phi - \frac{\varsigma}{2}\phi^4\right), \\
T_f^{\mu}{}_{\alpha} &= f^{\mu\beta}f_{\beta\alpha} + \frac{1}{4}\delta_{\alpha}^{\mu}f_{\gamma\delta}f^{\gamma\delta},
\end{aligned} \tag{36}$$

are the energy-momentum tensors for the B -gauge field, Dirac scalar field and dilatation curvature, respectively. Note that one can unify the energy-momentum tensor $T_B^{\mu}{}_{\alpha}$ and the term $3\mu\lambda\phi^2\delta_{\alpha}^{\mu}$ because both of these terms originated from the Maxwell curvature. But we prefer to use these quantities separately to show the cosmological constant explicitly.

Moreover, in Eq.(35), $\mathcal{R}_{\alpha}^{\mu}$ and \mathcal{R} are the Ricci tensor and scalar curvatures in Weyl space-time which are given in Eq.(26) and Eq.(27), so if substitute these equations into Eq.(35), the field equation becomes,

$$G_{\alpha}^{\mu} + 3\mu\lambda\phi^2\delta_{\alpha}^{\mu} = T_{\chi}^{\mu}{}_{\alpha} - 2\mu\phi^2T_B^{\mu}{}_{\alpha} - \kappa\phi^{-2}\left[T_{\phi}^{\mu}{}_{\alpha} + \frac{1}{2}T_f^{\mu}{}_{\alpha}\right], \tag{37}$$

where $G_{\alpha}^{\mu} = R_{\alpha}^{\mu} - \frac{1}{2}\delta_{\alpha}^{\mu}R$ is the well-known Einstein tensor and

$$T_{\chi}^{\mu}{}_{\alpha} = 2\left[\nabla^{\mu}\chi_{\alpha} - \chi^{\mu}\chi_{\alpha} + \frac{1}{2}g^{\mu\beta}(\partial_{\beta}\chi_{\alpha} - \partial_{\alpha}\chi_{\beta})\right] - \delta_{\alpha}^{\mu}(2\nabla_{\sigma}\chi^{\sigma} + \chi_{\sigma}\chi^{\sigma}). \tag{38}$$

is the energy-momentum tensor with respect to the dilatation gauge field.

To utilize the field equations for cosmological models, let's examine the following constraint which was discussed in [8];

$$\mathcal{D}_{\mu}\phi = \partial_{\mu}\phi - \chi_{\mu}\phi = 0. \tag{39}$$

By using this constraint, we can establish the dilatation gauge field by expressing it as the gradient of the Dirac scalar field in the following manner

$$\chi_{\mu} = \partial_{\mu}\ln\phi. \tag{40}$$

This constraint leads that the dilatation curvature $f_{\mu\nu} = \partial_{\mu}\chi_{\nu} - \partial_{\nu}\chi_{\mu}$ in Eq.(6) to be zero. This is another way of expressing the inverse Higgs effect [62, 63]. Clearly, this also shows the equivalence of the inverse Higgs constraint to equations of motion. As a result of this constraint, the field equation (37) is reduced to the following form,

$$G_{\alpha}^{\mu} + \Lambda\phi^2\delta_{\alpha}^{\mu} = T_{\chi}^{\mu}{}_{\alpha} - 2\mu\phi^2T_B^{\mu}{}_{\alpha}, \tag{41}$$

where it is worth noting that the term $\Lambda\phi^2$ can be seen as the effective cosmological constant and therefore this term can also be interpreted as the effective dark energy density [64]. Here, the constant Λ is given by

$$\Lambda = \left(3\mu\lambda + \frac{\kappa\varsigma}{4}\right). \tag{42}$$

On the other hand, in the presence of matter the action which is invariant under the Weyl transformation, the gravitational field is given by $S = S_g + S_M$ similar to [8]. Thus, we can write the field equation as follows,

$$G_{\alpha}^{\mu} + \Lambda\phi^2\delta_{\alpha}^{\mu} = \frac{\kappa}{\phi^2}T_M^{\mu}{}_{\alpha} + T_{\chi}^{\mu}{}_{\alpha} - 2\mu\phi^2T_B^{\mu}{}_{\alpha}, \tag{43}$$

where $T_M^{\mu}{}_{\alpha}$ is the ordinary matter energy-momentum tensor which is covariantly conserved, $\nabla_{\mu}T_M^{\mu}{}_{\alpha} = 0$.

At this point, we assume that the matter-energy constituents of the Universe are governed by a perfect fluid, which may be characterized by two thermodynamic variables: the energy density $\rho(t)$ and the pressure $P(t)$. Accordingly, the complete energy-momentum tensor generated by the Maxwell-Weyl gauge theory of gravity takes the following form:

$$T^\mu_\alpha = \text{diag}(\rho, -P, -P, -P), \quad (44)$$

which can be defined as follows by using Eq.(43),

$$T^\mu_\alpha = \frac{\kappa}{\phi^2} T_M^\mu{}_\alpha + T_\chi^\mu{}_\alpha - 2\mu\phi^2 T_B^\mu{}_\alpha + T_\Lambda^\mu{}_\alpha. \quad (45)$$

Thus, Eq.(43) can be written in a compact form as $G^\mu_\alpha = T^\mu_\alpha$. In this case, the total energy density and total pressure are defined by the combination of the matter, dilatation, Maxwell gauge field and the effective cosmological constant contributions as follows

$$\rho = \rho_M + \rho_\chi + \rho_B + \rho_\Lambda, \quad (46)$$

and

$$P = P_M + P_\chi + P_B + P_\Lambda, \quad (47)$$

where we assume that all energy densities are positive definite. The conservation of the energy-stress tensor in the FLRW background is represented by the equation $\nabla_\mu T^{\mu\nu} = 0$, which consequently gives rise to the continuity equation

$$\dot{\rho} + 3H(\rho + P) = 0, \quad (48)$$

where H the Hubble parameter.

III. MODIFIED FRIEDMANN EQUATIONS

The Einstein-Yang-Mills cosmology is a theoretical model that combines the principles of general relativity and Yang-Mills theory to study the evolution of the universe as a whole. The model proposes that the gravitational force can be described in terms of a connection between space-time and a non-Abelian gauge field. This model employed a non-abelian gauge field to study the behavior of the universe during the early stages of its evolution, including the inflationary period and the formation of cosmic structures such as galaxies and galaxy clusters. Similar to this theory, instead of the non-abelian Yang-Mills fields, we consider finding potential cosmological consequences of the Maxwell gauge fields B_μ^{ab} which are present in the gravitational field equation in Eq.(35).

In general, for most cosmological models, the metric is assumed to be a flat FLRW metric which based on Einstein's equations of general relativity, provides a framework for describing the evolution of the universe by assuming homogeneity and isotropy on large scales. So we start from the following line element,

$$ds^2 = dt^2 - a(t)^2 (dx^2 + dy^2 + dz^2), \quad (49)$$

where $a(t)$ is the scale factor. In accordance with our assumptions, let us take the scalar field as a function of the cosmic time $\phi(x) \rightarrow \phi(t)$. Thus, by using Eq.(40), the dilatation gauge field can be identified with only one nonzero

component;

$$\chi_\mu = \left(\frac{\dot{\phi}}{\phi}, 0, 0, 0 \right). \quad (50)$$

Moreover, according to [50], we can define the Maxwell gauge fields $B_\mu^{ab}(x)$ in terms of one-dimensional time dependent fields $\psi(t)$ and $\zeta(t)$,

$$B_\mu^{0s}(x) \rightarrow B_\mu^{0s}(t) = (0, \delta_i^s \psi(t)), \quad (51)$$

$$B_\mu^{rs}(x) \rightarrow B_\mu^{rs}(t) = (0, \epsilon_i^{rs} \zeta(t)), \quad (52)$$

where $r, s = 1, 2, 3$ are the tangent indices and $i, j = 1, 2, 3$ are the space-time indices. Here, δ_i^s and ϵ_i^{rs} are three dimensional $so(3)$ tensors.

Having established the necessary background, we can now proceed to derive the Friedmann equations in the torsion-free condition by taking into account of the gauge in Eq.(39) and $\mathcal{DF}^{ab} = 0$. Therefore, by employing Eqs.(51), (52) and (22) and subsequently applying them to the field equation given in Eq.(35), the following equation can be obtained by focusing on the $(0, 0)$ component of Eq.(35),

$$\left(\frac{\dot{a}}{a} \right)^2 = \frac{\kappa \rho_M}{3\phi^2} - \frac{2\dot{a}\dot{\phi}}{a\phi} - \left(\frac{\dot{\phi}}{\phi} \right)^2 - \frac{4\mu\phi^2}{a^2} \left(\dot{a} + \frac{a\dot{\phi}}{\phi} \right) \psi - \frac{\Lambda\phi^2}{3}, \quad (53)$$

where the dot denotes the derivative with respect to the cosmological time t . The (i, i) components of Eq.(35) lead to,

$$\frac{2\ddot{a}}{a} + \left(\frac{\dot{a}}{a} \right)^2 = -\frac{\kappa P_M}{\phi^2} - \frac{2\ddot{\phi}}{\phi} - \frac{4\dot{a}\dot{\phi}}{a\phi} + \left(\frac{\dot{\phi}}{\phi} \right)^2 - \frac{4\mu\phi^2}{a^2} \left(\dot{a}\psi + a\dot{\psi} + 5\frac{a\dot{\phi}\psi}{\phi} \right) - \Lambda\phi^2. \quad (54)$$

Then by making use of Eqs.(53) and (54), the acceleration equation can be obtained as,

$$\frac{\ddot{a}}{a} = -\frac{\kappa}{6\phi^2} (3P_M + \rho_M) - \frac{\dot{a}\dot{\phi}}{a\phi} + \left(\frac{\dot{\phi}}{\phi} \right)^2 - \frac{\ddot{\phi}}{\phi} - \frac{2\mu\phi^2}{a} \left(\dot{\psi} + \frac{4\dot{\phi}\psi}{\phi} \right) - \frac{\Lambda\phi^2}{3}, \quad (55)$$

which can also be obtained by considering the trace of Eq.(35).

Furthermore, by using Eqs.(53) and (54), one can find the energy density and the pressure expressions for the corresponding fields, as presented below;

$$\rho_\chi = -\frac{3\phi^2}{\kappa} \left[\frac{2\dot{a}\dot{\phi}}{a\phi} + \left(\frac{\dot{\phi}}{\phi} \right)^2 \right], \quad P_\chi = \frac{\phi^2}{\kappa} \left[\left(\frac{4\dot{a}\dot{\phi}}{a\phi} - \left(\frac{\dot{\phi}}{\phi} \right)^2 \right) + \frac{2\ddot{\phi}}{\phi} \right], \quad (56)$$

$$\rho_\Lambda = -\frac{\Lambda\phi^4}{\kappa}, \quad P_\Lambda = \frac{\Lambda\phi^4}{\kappa}, \quad (57)$$

$$\rho_B = -\frac{12\mu\phi^4\psi}{\kappa a^2} \left(\dot{a} + \frac{a\dot{\phi}}{\phi} \right), \quad P_B = \frac{4\mu\phi^4}{\kappa a^2} \left(\dot{a}\psi + a\dot{\psi} + 5\frac{a\dot{\phi}\psi}{\phi} \right). \quad (58)$$

For completeness, if we combine Eqs.(53) and (55), we also derive the time derivative of the Hubble parameter ($H(t) = \dot{a}(t)/a(t)$) as

$$\begin{aligned} \dot{H} &= -\frac{\kappa}{2\phi^2} (\rho_M + P_M) + \frac{\dot{a}\dot{\phi}}{a\phi} + 2 \left(\frac{\dot{\phi}}{\phi} \right)^2 - \frac{\ddot{\phi}}{\phi} + \frac{2\mu\phi^2}{a^2} \left(2\dot{a}\psi - a\dot{\psi} - \frac{2a\dot{\phi}\psi}{\phi} \right) \\ &= -\frac{\kappa}{2\phi^2} (\rho + P). \end{aligned} \quad (59)$$

By setting $\psi(t) = 0$, the equations (53), (54) and (55) can be reduced to the forms presented in Rosen's paper [8]. Additionally, imposing the compensating field $\phi(t) = 1$ yields the equations derived in [51] for the Maxwell algebra. Finally, the conventional cosmological equations of general relativity, along with the cosmological term, can be obtained by setting both $\psi(t) = 0$ and $\phi(t) = 1$ as follows,

$$\begin{aligned} \left(\frac{\dot{a}}{a} \right)^2 &= \frac{\kappa\rho_M}{3} - \frac{\Lambda}{3}, \\ \frac{2\ddot{a}}{a} + \left(\frac{\dot{a}}{a} \right)^2 &= -\kappa P_M - \Lambda, \\ \frac{\ddot{a}}{a} &= -\frac{\kappa}{6} (\rho_M + 3P_M) - \frac{\Lambda}{3}, \end{aligned} \quad (60)$$

with a small difference in the definition of the cosmological constant which is given by Eq.(42).

IV. CONCLUSION

In this paper, we have explored the gauge theory of gravity based on the Maxwell-Weyl group and its potential for a more comprehensive cosmological model. We carried out our investigation by applying an analogous methodology to that employed in the Einstein-Yang-Mills cosmology, we have established a cosmological model that incorporates generalized Friedmann equations including the effective cosmological constant and the contribution of additional scalar fields originating from the Maxwell symmetry in Eqs.(53)-(55).

The obtained result extends the conventional cosmological solution derived from the Weyl gauge theory of gravity, as described in Rosen [8], by introducing the Maxwell symmetry under a particular gauge constraint expressed in Eq.(39). Moreover, the present findings suggest that the Maxwell gauge field may play a significant role in the acceleration of the universe, as indicated in Eq.(55), and these new terms may be interpreted as a kind of dark energy. Furthermore, it is worth noting that in a specific limit, the modified Friedmann equations presented in this study reduce to the well-known cosmological equations of general relativity with the cosmological constant, (see Eq.(60)).

Consequently, our results highlight the potential of the Maxwell-Weyl gauge theory for advancing our understanding of the universe's acceleration and cosmological evolution and have the potential to shape the future direction of theoretical physics research in this area.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

DATA AVAILABILITY STATEMENT

No new data were created or analysed in this study.

ACKNOWLEDGMENTS

This study is supported by the Scientific and Technological Research Council of Turkey (TUBİTAK) under grant number 2219.

-
- [1] C.-N. Yang and R. L. Mills, Conservation of Isotopic Spin and Isotopic Gauge Invariance, *Phys. Rev.* **96**, 191 (1954).
 - [2] R. Utiyama, Invariant theoretical interpretation of interaction, *Phys. Rev.* **101**, 1597 (1956).
 - [3] T. W. B. Kibble, Lorentz invariance and the gravitational field, *J. Math. Phys.* **2**, 212 (1961).
 - [4] A. Bregman, Weyl transformations and Poincare gauge invariance, *Prog. Theor. Phys.* **49**, 667 (1973).
 - [5] J. M. Charap and W. Tait, A gauge theory of the Weyl group, *Proc. Roy. Soc. Lond. A* **340**, 249 (1974).
 - [6] P. A. M. Dirac, Long range forces and broken symmetries, *Proc. Roy. Soc. Lond. A* **333**, 403 (1973).
 - [7] M. Kasuya, On the Gauge Theory in the Einstein-Cartan-Weyl Space-Time, *Nuovo Cim. B* **28**, 127 (1975).
 - [8] N. Rosen, Weyl's geometry and physics, *Found. Phys.* **12**, 213 (1982).
 - [9] J. Cervero and L. Jacobs, Classical Yang-Mills Fields in a Robertson-walker Universe, *Phys. Lett. B* **78**, 427 (1978).
 - [10] M. Henneaux, Remarks on space-time symmetries and nonabelian gauge fields, *J. Math. Phys.* **23**, 830 (1982).
 - [11] D. V. Galtsov and M. S. Volkov, Yang-Mills cosmology: Cold matter for a hot universe, *Phys. Lett. B* **256**, 17 (1991).
 - [12] P. V. Moniz, J. M. Mourao, and P. M. Sa, The dynamics of a flat friedmann-robertson-walker inflationary model in the presence of gauge fields, *Class. Quant. Grav.* **10**, 517 (1993).
 - [13] K. Bamba, S. Nojiri, and S. D. Odintsov, Inflationary cosmology and the late-time accelerated expansion of the universe in non-minimal Yang-Mills-F(R) gravity and non-minimal vector-F(R) gravity, *Phys. Rev. D* **77**, 123532 (2008).
 - [14] D. V. Gal'tsov, Non-Abelian condensates as alternative for dark energy, in *43rd Rencontres de Moriond on QCD and High Energy Interactions* (2008) [arXiv:0901.0115 \[gr-qc\]](https://arxiv.org/abs/0901.0115).
 - [15] A. Maleknejad and M. M. Sheikh-Jabbari, Gauge-flation: Inflation From Non-Abelian Gauge Fields, *Phys. Lett. B* **723**, 224 (2013).
 - [16] A. Guarnizo, J. B. Orjuela-Quintana, and C. A. Valenzuela-Toledo, Dynamical analysis of cosmological models with non-Abelian gauge vector fields, *Phys. Rev. D* **102**, 083507 (2020).
 - [17] A. Maleknejad and M. M. Sheikh-Jabbari, Non-abelian gauge field inflation, *Phys. Rev. D* **84**, 043515 (2011).
 - [18] M. M. Sheikh-Jabbari, Gauge-flation Vs Chromo-Natural Inflation, *Phys. Lett. B* **717**, 6 (2012).
 - [19] J. Bielefeld and R. R. Caldwell, Cosmological consequences of classical flavor-space locked gauge field radiation, *Phys. Rev. D* **91**, 124004 (2015).
 - [20] A. Maleknejad, M. M. Sheikh-Jabbari, and J. Soda, Gauge Fields and Inflation, *Phys. Rept.* **528**, 161 (2013).
 - [21] W. Zhao and Y. Zhang, The state equation of the Yang-Mills field dark energy models, *Class. Quant. Grav.* **23**, 3405 (2006).
 - [22] D. V. Gal'tsov and E. A. Davydov, Yang-Mills condensates in cosmology, in *Int. J. Mod. Phys. Conf. Ser.*, Vol. 14 (2012) pp. 316–325.
 - [23] E. Elizalde, A. J. Lopez-Revelles, S. D. Odintsov, and S. Y. Vernov, Cosmological models with Yang-Mills fields, *Phys. Atom. Nucl.* **76**, 996 (2013).
 - [24] M. R. Setare and V. Kamali, Warm Vector Inflation, *Phys. Lett. B* **726**, 56 (2013).
 - [25] M. Rinaldi, Dark energy as a fixed point of the Einstein Yang-Mills Higgs Equations, *JCAP* **10**, 023.
 - [26] A. Mehrabi, A. Maleknejad, and V. Kamali, Gaugessence: a dark energy model with early time radiation-like equation of state, *Astrophys. Space Sci.* **362**, 53 (2017).
 - [27] H. Zhang, C. S. Li, Q.-H. Cao, and Z. Li, A Dark Matter Model with Non-Abelian Gauge Symmetry, *Phys. Rev. D* **82**, 075003 (2010).

- [28] M. A. Buen-Abad, G. Marques-Tavares, and M. Schmaltz, Non-Abelian dark matter and dark radiation, *Phys. Rev. D* **92**, 023531 (2015).
- [29] C. Gross, O. Lebedev, and Y. Mambrini, Non-Abelian gauge fields as dark matter, *JHEP* **08**, 158.
- [30] H. Bacry, P. Combe, and J. L. Richard, Group-theoretical analysis of elementary particles in an external electromagnetic field. 1. the relativistic particle in a constant and uniform field, *Nuovo Cim. A* **67**, 267 (1970).
- [31] R. Schrader, The maxwell group and the quantum theory of particles in classical homogeneous electromagnetic fields, *Fortschr. Phys.* **20**, 701 (1972).
- [32] D. V. Soroka and V. A. Soroka, Tensor extension of the Poincare' algebra, *Phys. Lett. B* **607**, 302 (2005).
- [33] J. A. de Azcarraga, K. Kamimura, and J. Lukierski, Generalized cosmological term from Maxwell symmetries, *Phys. Rev. D* **83**, 124036 (2011).
- [34] R. Durka, J. Kowalski-Glikman, and M. Szczachor, Gauged AdS-Maxwell algebra and gravity, *Mod. Phys. Lett. A* **26**, 2689 (2011).
- [35] D. V. Soroka and V. A. Soroka, Gauge semi-simple extension of the Poincaré group, *Phys. Lett. B* **707**, 160 (2012).
- [36] J. A. de Azcarraga and J. M. Izquierdo, Minimal $D = 4$ supergravity from the superMaxwell algebra, *Nucl. Phys. B* **885**, 34 (2014).
- [37] O. Cebecioglu and S. Kibaroğlu, Gauge theory of the Maxwell-Weyl group, *Phys. Rev. D* **90**, 084053 (2014).
- [38] O. Cebecioglu and S. Kibaroğlu, Maxwell-affine gauge theory of gravity, *Phys. Lett. B* **751**, 131 (2015).
- [39] P. K. Concha, E. K. Rodríguez, and P. Salgado, Generalized supersymmetric cosmological term in $N = 1$ Supergravity, *JHEP* **08**, 009.
- [40] S. Kibaroğlu, M. Şenay, and O. Cebecioglu, $D = 4$ topological gravity from gauging the Maxwell-special-affine group, *Mod. Phys. Lett. A* **34**, 1950016 (2019).
- [41] S. Kibaroğlu and O. Cebecioglu, $D = 4$ supergravity from the Maxwell-Weyl superalgebra, *Eur. Phys. J. C* **79**, 898 (2019).
- [42] S. Kibaroğlu and O. Cebecioglu, Generalized cosmological constant from gauging Maxwell-conformal algebra, *Phys. Lett. B* **803**, 135295 (2020).
- [43] S. Kibaroğlu and O. Cebecioglu, Gauge theory of the Maxwell and semi-simple extended (anti) de Sitter algebra, *Int. J. Mod. Phys. D* **30**, 2150075 (2021).
- [44] O. Cebecioglu and S. Kibaroğlu, Maxwell-modified metric affine gravity, *Eur. Phys. J. C* **81**, 900 (2021).
- [45] O. Cebecioglu, A. Saban, and S. Kibaroğlu, Maxwell extension of $f(R)$ gravity, *Eur. Phys. J. C* **83**, 95 (2023).
- [46] S. Kibaroğlu, O. Cebecioglu, and A. Saban, Gauging the Maxwell extended $GL(n, R)$ and $SL(n+1, R)$ algebras, *Symmetry* **15**, 464 (2023).
- [47] J. Frieman, M. Turner, and D. Huterer, Dark Energy and the Accelerating Universe, *Ann. Rev. Astron. Astrophys.* **46**, 385 (2008).
- [48] T. Padmanabhan, Dark Energy and its Implications for Gravity, *Adv. Sci. Lett.* **2**, 174 (2009).
- [49] R. Durka and J. Kowalski-Glikman, Local Maxwell symmetry and gravity, arXiv preprint [arXiv:1110.6812](https://arxiv.org/abs/1110.6812) [hep-th].
- [50] J. A. de Azcarraga, K. Kamimura, and J. Lukierski, Maxwell symmetries and some applications, in *Int. J. Mod. Phys. Conf. Ser.*, Vol. 23 (2013) pp. 350–356.
- [51] S. Kibaroğlu, Cosmological application of the Maxwell gravity, arXiv preprint [arXiv:2208.06641](https://arxiv.org/abs/2208.06641) [hep-th].
- [52] S. Bonanos and J. Gomis, Infinite Sequence of Poincare Group Extensions: Structure and Dynamics, *J. Phys. A* **43**, 015201 (2010).
- [53] M. Blagojevic, *Gravitation and gauge symmetries* (CRC Press, 2002).
- [54] G. K. Karananas and A. Monin, Weyl and Ricci gauging from the coset construction, *Phys. Rev. D* **93**, 064013 (2016).
- [55] M. P. Hobson and A. N. Lasenby, Weyl gauge theories of gravity do not predict a second clock effect, *Phys. Rev. D* **102**, 084040 (2020).
- [56] M. Omote, Scale transformations of the second kind and the weyl space-time, *Lett. Nuovo Cim.* **2S2**, 58 (1971).
- [57] O. V. Babourova, B. N. Frolov, and V. C. Zhukovsky, Gauge field theory for Poincare-Weyl group, *Phys. Rev. D* **74**, 064012 (2006).
- [58] H. Weyl, *Space-time-matter* (Dutton, 1922).
- [59] C. Brans and R. H. Dicke, Mach's principle and a relativistic theory of gravitation, *Phys. Rev.* **124**, 925 (1961).
- [60] A. G. Agnese and P. Calvini, Gauge Fields Arising from Space-Time Symmetries and Gravitational Theories. 2., *Phys. Rev. D* **12**, 3804 (1975).

- [61] T. Dereli and R. W. Tucker, Weyl scalings and spinor matter interactions in scalar-tensor theories of gravitation, *Phys. Lett. B* **110**, 206 (1982).
- [62] E. A. Ivanov and V. I. Ogievetsky, The Inverse Higgs Phenomenon in Nonlinear Realizations, *Teor. Mat. Fiz.* **25**, 164 (1975).
- [63] I. Low and A. V. Manohar, Spontaneously broken space-time symmetries and Goldstone's theorem, *Phys. Rev. Lett.* **88**, 101602 (2002).
- [64] O. V. Babourova, B. N. Frolov, and K. N. Lipkin, Gravitation theory with a Dirac scalar field in the exterior form formalism and the cosmological constant problem, *Grav. Cosmol.* **18**, 225 (2012).