

ON WEAK AMENABILITY OF FOURIER ALGEBRAS

VIKTOR LOSERT

1. INTRODUCTION AND MAIN RESULTS

Let A be a Banach algebra, X a Banach A -bimodule. A linear map $D: A \rightarrow X$ is called a derivation if $D(ab) = aD(b) + D(a)b$ for $a, b \in A$. By A' we denote the dual space of A with the dual action (also called the dual module of A). A is called *weakly amenable* if every bounded derivation $D: A \rightarrow A'$ is inner (i.e. there exists $f \in A'$ such that $D(a) = af - fa$ for $a \in A$).

From now, we assume that A is commutative. X is called a symmetric A -bimodule if $ax = xa$ for $a \in A, x \in X$. Then ([Da] Th. 2.8.63) A is weakly amenable iff there is no non-zero bounded derivation from A to any symmetric Banach A -bimodule X .

Our main subject will be $A = A(G)$ the Fourier algebra of a locally compact group G .

Theorem. *Let G be a connected locally compact group. If $A(G)$ is weakly amenable then G is abelian.*

Corollary. *Let G be a locally compact group with identity component G^0 . $A(G)$ is weakly amenable if and only if G^0 is abelian.*

Some History

The notion of weak amenability was introduced by Bade, Curtis and Dales for commutative Banach algebras and then extended by Johnson, see [Da] p. 306 for more details and references. For the Fourier algebra it was shown in [J] that $A(G)$ is not weakly amenable for $G = SO(3)$ and then in [Pl] for G any non-abelian compact connected Lie group. For non-compact groups, the case of the $ax + b$ group was settled in [CG1] and then the Heisenberg group in [CG2]. Finally in [LLSS] it was shown that $A(G)$ is not weakly amenable for any non-abelian connected Lie group G .

Proof of the Corollary. If G^0 is abelian it was shown by Forrest and Runde ([FR] Th. 3.3) that $A(G)$ is weakly amenable. There is a heredity for subgroups ([FR]

L. 3.1): if G is any locally compact group and $A(G)$ is weakly amenable then $A(H)$ is weakly amenable for every closed subgroup H of G . Hence the converse in the Corollary follows from the Theorem (and this was already conjectured in [FR]). \square

Outline of Proof of the Theorem. The proof of the theorem will take most of the remaining sections. The basic strategy follows the approach of the earlier authors. By the heredity for subgroups (see above) it will be enough to prove the theorem for “small” non-abelian connected groups. In Section 2 we present lists $(\mathcal{M}), (\mathcal{M}+)$ of the minimal non-abelian connected groups (extending that in [LLSS]). Using the results from Appendix A on connected dense subgroups we show (Proposition 1) that every non-abelian connected locally compact group has a closed subgroup isomorphic to one of the groups in $(\mathcal{M}) \cup (\mathcal{M}+)$.

For the treatment of $A(G)$ for specific groups, [LLSS] used techniques from spectral synthesis. Put $\check{\Delta}_G = \{(g, g^{-1}) : g \in G\}$ (the anti-diagonal). For a connected Lie group G they show ([LLSS] Th. 3.2) that $A(G)$ is weakly amenable iff $\check{\Delta}_G$ is of local synthesis for $G \times G$. This synthesis property is inherited by locally isomorphic groups, hence it was enough to check the simply connected groups from (\mathcal{M}) . But, so far, this approach works only for Lie groups.

We will use the method of Johnson, Choi, Ghandehari who consider explicit derivations arising from differentiation operators. Differentiation operators are prototypes of derivations on function spaces and their continuity properties express some kind of smoothness for the functions. It is a basic construction for a Lie group G with Lie algebra \mathfrak{g} that the elements of \mathfrak{g} correspond to point derivations (at e) on the algebra $C^\infty(G)$ of C^∞ -functions. These point derivations extend to left invariant (or right invariant) derivations on $C^\infty(G)$ ([Va] Sec. 2.3 and 1.1). Norm estimates for such derivations are done using Fourier transforms. In Appendix B we review the relevant properties of the non-commutative Fourier transform and also the applications to Fourier algebras. It turns out that in some cases the derivations that we consider do not map to the dual $VN(G)$ of $A(G)$. Then (taking up constructions from [CG2]) we consider appropriate $A(G)$ -modules which needs some knowledge about the dual convolution for these groups. In the case of the group $M(2)$ (and the related groups $\widetilde{M(2)}$ and $M(2)_K$) this becomes somewhat more involved. \square

Notations

On semidirect products $G = H \ltimes K$: in those examples where the notation should (hopefully) make it easy to distinguish elements of H and K we will use the internal

presentation (see [HR] I.2.6). Thus H and K are considered as (closed) subgroups of G , with H normal, $H \cap K = \{e\}$ (e denoting the unit element of G), $G=HK$ ($(h, k) \mapsto hk$ defining a homeomorphism $H \times K \rightarrow G$). There is given a continuous (left) action of K on H , denoted by \circ , (equivalently, a continuous homomorphism $K \rightarrow \text{Aut}(H)$) and the group multiplication of G is governed by the rule $kh = (k \circ h)k$ for $k \in K, h \in H$. Then a mapping $\varphi: G \rightarrow G_1$ (G_1 some other topological group) is a continuous homomorphism iff its restrictions to H and K are continuous homomorphisms and it preserves the action, i.e. $\varphi(k \circ h) = \varphi(k)\varphi(h)\varphi(k)^{-1}$. In Section 4 we use a slightly modified notation.

2. MINIMAL NON-ABELIAN CONNECTED GROUPS

In [LLSS] Prop. 3.1 the minimal non-abelian real Lie algebras are listed. The corresponding simply connected (and connected) Lie groups are

$$(\mathcal{M}) \quad SU(2), \quad \mathbb{R}^n \rtimes \mathbb{R} \quad (n \in \{1, 2\})$$

with the actions $(t, x, y, z \in \mathbb{R}, v \in \mathbb{R}^2)$

$$n = 1 \quad t \circ x = e^t x$$

$ax + b$ group

$$n = 2 \quad t \circ (y, z) = (y, z + ty)$$

\mathbb{H} Heisenberg group

$$t \circ v = A_t v \quad \text{with } A_t = \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix} \quad \widetilde{M(2)} \text{ universal covering of the} \\ \text{Euclidian motion group of the plane}$$

$$t \circ v = A_t v \quad \text{with } A_t = e^{at} \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix} \quad \mathbb{G}_a \text{ Grélaud groups, } a > 0.$$

Their Lie algebras do not have a proper non-abelian subalgebra and any non-abelian finite dimensional real Lie algebra contains a copy of one of them.

Lemma 1. *Let G be a non-abelian connected locally compact group. Then there exists a group H from (\mathcal{M}) and a continuous homomorphism $\varphi: H \rightarrow G$ with discrete kernel.*

Proof. We represent G using an Iwasawa pair, $G \cong (L \times K)/D$ where L is a connected Lie group, K a compact group (not necessarily connected), D a discrete central subgroup ([G1] Prop. 5, see also [PW] sec. 2 for more properties and [L] p. 120 for further references). This gives homomorphisms $j_L: L \rightarrow G, j_K: K \rightarrow G$ with discrete kernel (in fact, it is easy to modify the representation so that j_L, j_K are injective). G being connected, the image of $L \times K^0$ must be dense. Hence if G is non-abelian, then either L or K^0 must be non-abelian. If L is non-abelian there exists (using e.g. [Va] Th. 2.7.5 and Cor. 2.7.4) a group H from (\mathcal{M}) and a

homomorphism $\varphi_0: H \rightarrow L$ with discrete kernel. Then we may take $\varphi = j_L \circ \varphi_0$. Similarly, if K^0 is non-abelian then (using e.g. [HM] Cor. 9.6, Th. 9.19(i),(ii) and Prop. 6.46(IV)) it follows that there exists a homomorphism $\varphi_0: SU(2) \rightarrow K$ with discrete kernel. \square

We see that for arbitrary connected groups one has to consider also closures of non-abelian homomorphic images of the groups from (\mathcal{M}) . This gives in addition the groups

$$(\mathcal{M}+) \quad SO(3), \quad \mathbb{H}_{\mathcal{K}}, \quad M(2)_{\mathcal{K}}$$

where \mathcal{K} is a (non-zero) solenoidal group i.e. \mathcal{K} compact and there exists a continuous homomorphism $\varphi_0: \mathbb{R} \rightarrow \mathcal{K}$ with dense image, \mathcal{K} is written additively.

$$\mathbb{H}_{\mathcal{K}} = (\mathbb{R} \times \mathcal{K}) \rtimes \mathbb{R}, \quad \text{action } t \circ (y, k) = (y, k + \varphi_0(ty)) \quad (t, y \in \mathbb{R}, k \in \mathcal{K}).$$

$M(2)_{\mathcal{K}} = \mathbb{R}^2 \rtimes \mathcal{K}$ with the additional assumption that there exists a proper closed subgroup \mathcal{K}_0 of \mathcal{K} such that $\varphi_0(2\pi\mathbb{Z}) = \mathcal{K}_0 \cap \varphi_0(\mathbb{R})$. Then $\mathcal{K}/\mathcal{K}_0 \cong \mathbb{R}/2\pi\mathbb{Z}$ and the action of \mathbb{R} on \mathbb{R}^2 (defining $\widetilde{M(2)}$) induces an action of \mathcal{K} on \mathbb{R}^2 .

Special cases: For $\mathcal{K} = \mathbb{R}/\mathbb{Z}$ (φ_0 the quotient mapping) $\mathbb{H}_{\mathcal{K}}$ is the *reduced Heisenberg group*. For $\mathcal{K} = \mathbb{R}/2\pi\mathbb{Z}$ ($\mathcal{K}_0 = \{0\}$) $M(2)_{\mathcal{K}}$ equals $M(2)$ the *Euclidian motion group of the plane*.

Proposition 1. *Every non-abelian connected locally compact group G has a closed subgroup isomorphic to one of the groups in $(\mathcal{M}) \cup (\mathcal{M}+)$.*

Proof. By Lemma 1 there exists a group H from (\mathcal{M}) and a continuous homomorphism $\varphi: H \rightarrow G$ with discrete (hence central) kernel. Replacing G by $\overline{\varphi(H)}$ we want to show that G is isomorphic to one of the groups in $(\mathcal{M}) \cup (\mathcal{M}+)$.

For $H = SU(2)$, $\varphi(H)$ is closed, hence $G \cong H / \ker \varphi$ ([HR] Th. 5.29 and Th. 5.27). Since $Z(SU(2)) = \{\pm I\}$ (centre), it follows that $G \cong SU(2)$ or $G \cong SU(2)/\{\pm I\}$. It is classical (see e.g. [HM] Ex. E1.2(ii)) that $SU(2)/\{\pm I\} \cong SO(3)$.

For the other groups in (\mathcal{M}) we use the results and constructions from Appendix A. For the $ax+b$ -group and the Grélaud groups one can check that the centre is trivial, hence φ must be injective. A necessary condition for $x \in H$ to generate a relatively compact subgroup in $\text{Aut}(H)$ is relative compactness of the orbit $\{x^n y x^{-n} : n \in \mathbb{Z}\}$ for every $y \in H$. Again one can check that this does not happen for $x \neq e$. In the notation of Theorem A it follows that the vector group V is trivial, hence $\varphi(H)$ is always closed and as above it follows that $\varphi(H) \cong H$ (i.e. these are examples of “absolutely closed” groups, [G2] p. 728).

For $H = \mathbb{H}$ the Heisenberg group an element generates a relatively compact subgroup in $\text{Aut}(H)$ iff it belongs to the centre $Z(H) = \{(0, z) : z \in \mathbb{R}\}$ (notation as in (\mathcal{M})). Hence the only non-trivial vector group as in Theorem A is $V = Z(H)$. Then $\ker \varphi$ is trivial and the action α is trivial on $\varphi(V)$. Thus the action α_0 in Proposition 6 (one has $H_1 = H$) is trivial on $\mathcal{K} = \overline{\varphi(V)}$. By Proposition 6(b) it follows that $G \cong (H \times \mathcal{K})/D$. It is easy to see that the homomorphism $((y, z)t, k) \mapsto (y, \varphi_0(z) + k)t$ (mapping $H \times \mathcal{K}$ onto $\mathbb{H}_{\mathcal{K}}$) induces an isomorphism of $(H \times \mathcal{K})/D$ and $\mathbb{H}_{\mathcal{K}}$. These gives the cases where $\ker \varphi_0 = \ker \varphi$ is trivial. But the construction of Proposition 6 works more generally for \mathcal{K} as in $(\mathcal{M}+)$ with $\ker \varphi_0$ discrete (see also Remark (c) in Appendix A). In Theorem A this adds the cases where V is trivial. Then φ is surjective, hence as above $G \cong H/\ker \varphi$. When $\ker \varphi_0$ is a non-trivial subgroup of $Z(H) \cong \mathbb{R}$, φ_0 is also surjective, thus $\mathcal{K} \cong \mathbb{R}/\ker \varphi_0$. It gives the reduced Heisenberg group and some variants that are isomorphic to it.

For $H = \widetilde{M(2)} = \mathbb{R}^2 \rtimes \mathbb{R}$ the universal covering of the Euclidian motion group of the plane, $h \in H$ generates a relatively compact subgroup in $\text{Aut}(H)$ iff $h \notin \mathbb{R}^2$ and then h is conjugate to an element of \mathbb{R} . Hence, in Theorem A, if V is non-trivial we may assume that $V = \mathbb{R}$ (the second factor in the semidirect product). The centre $Z(H)$ is $2\pi\mathbb{Z}$ (subgroup of V), thus $\ker \varphi$ is trivial. Put $\mathcal{K} = \overline{\varphi(V)}$, $\mathcal{K}_0 = \overline{\varphi(2\pi\mathbb{Z})}$, $\varphi_0 = \varphi|_V$. By Proposition 6(b) (splitting case, $H_2 = \mathbb{R}^2$), $G \cong \mathbb{R}^2 \rtimes \mathcal{K}$. The action α is trivial on \mathcal{K}_0 and non-trivial on $\varphi(\mathbb{R} \setminus 2\pi\mathbb{Z})$ (using that $\alpha \circ \varphi = \iota_H$). It follows that $\varphi_0(2\pi\mathbb{Z}) = \mathcal{K}_0 \cap \varphi_0(\mathbb{R})$ and one can check that the action used in Proposition 6(b) (restricting α_0 to \mathbb{R}^2) coincides with that described in $(\mathcal{M}+)$. Again there are the additional cases where V is trivial. Then φ is surjective, $G \cong H/\ker \varphi \cong \mathbb{R}^2 \rtimes (\mathbb{R}/\ker \varphi)$. $\ker \varphi$ has to be a subgroup of $2\pi\mathbb{Z}$. Thus $\mathcal{K} \cong \mathbb{R}/2k\pi\mathbb{Z}$ $k = 0, 1, \dots$, giving $M(2)$ ($k = 1$) and its finite covering groups ($k > 0$), and of course $\widetilde{M(2)}$ ($k = 0$). Conversely, if \mathcal{K} is a compact group as in $(\mathcal{M}+)$ it is easy to see that φ_0 extends to a homomorphism $\varphi: \widetilde{M(2)} \rightarrow M(2)_{\mathcal{K}}$ with dense image and discrete kernel. \square

Remarks. (a) For an alternative argument, one can compute the adjoint action of the groups. It turns out that all the groups H in (\mathcal{M}) are CA -groups, i.e. $\text{Int}(H)$ is closed in $\text{Aut}(H)$. By [G2] 4.1, if the connected Lie group H is a CA group, $\varphi: H \rightarrow G$ an injective (continuous) homomorphism with dense image, G a Hausdorff topological group, then $G = \varphi(H) \overline{\varphi(Z)}$ ($Z = Z(H)$ denoting the centre of H). This holds more generally when H is a connected locally compact group

and a CA group, $\ker \varphi$ totally disconnected (compare [PW] Cor.4.2). It follows that G is a central extension of H/Z .

(b) As explained in Appendix A, Rem. (c), for a (necessarily abelian) compact group \mathcal{K} as in $(\mathcal{M}+)$, the dual group $\widehat{\mathcal{K}}$ is isomorphic to a subgroup of \mathbb{R} . For the Heisenberg group \mathbb{H} any non-zero subgroup Γ of \mathbb{R} defines in this way a group $\mathbb{H}_{\mathcal{K}}$ (the trivial subgroup would give $\mathbb{R}^2 \cong \mathbb{H}/Z(\mathbb{H})$, this is a case where $\ker \varphi$ is non-discrete, as mentioned in Appendix A, Rem. (c); it follows that there is no minimal extension $\mathbb{H}_{\mathcal{K}}$). One can show that two subgroups Γ, Γ' define isomorphic groups $\mathbb{H}_{\mathcal{K}}, \mathbb{H}_{\mathcal{K}'}$ iff $\Gamma' = t\Gamma$ for some $t > 0$.

For $H = \widetilde{M(2)}$ a restriction comes through the adjoint action. Put $\chi_t(x) = e^{itx}$ and use $\chi_t \leftrightarrow t$ for the identification of $\widehat{\mathbb{R}}$ and \mathbb{R} . One computes that (in the notation of Appendix A, Rem. (c)) $\widehat{\iota}_0(\Gamma_0) = \mathbb{Z}$, hence $\widehat{\varphi}_0(\widehat{\mathcal{K}})$ must be a subgroup of \mathbb{R} containing \mathbb{Z} and this is also equivalent to the condition $\varphi_0(2\pi\mathbb{Z}) = \mathcal{K}_0 \cap \varphi_0(\mathbb{R})$ of (\mathcal{M}) (and then $\widehat{\mathcal{K}}_0 \cong \widehat{\varphi}_0(\widehat{\mathcal{K}})/\mathbb{Z}$). Different groups $\Gamma, \Gamma' \subseteq \mathbb{R}$ (containing \mathbb{Z}) give non-isomorphic groups $M(2)_{\mathcal{K}}, M(2)_{\mathcal{K}'}$. The minimal case is $\Gamma = \mathbb{Z}$ with $\mathcal{K} = \mathbb{R}/2\pi\mathbb{Z}$, $M(2)_{\mathcal{K}} = M(2)$. The maximal case is $\Gamma = \mathbb{R}$ (discrete topology) with $\mathcal{K} = b(\mathbb{R})$ (Bohr compactification), this is also mentioned in [LLSS] Ex.3.4 (denoted as \mathbb{R}^{ap}). $M(2)_{\mathcal{K}}$ is a Lie group iff Γ is finitely generated and it is second countable iff Γ is countable.

3. THE GROUPS $SO(3)$ AND $SU(2)$

For $G = SU(2)$ Johnson's method has been elaborated in [CG1] sec.3. We add here the aspect of (generalized) convolution operators that will be useful in the following sections.

The Lie algebra \mathfrak{g} of G consists of the complex skew-hermitian 2×2 matrices with trace 0. For $X \in \mathfrak{g}$ we consider the distribution $d = \lim_{t \rightarrow 0} \frac{1}{t}(\delta_{\exp(-tX)} - \delta_I)$ (directional derivative for the tangent vector $-X$; δ_x point measure at x , I identity matrix). We put $D(f) = f \star d$. Classically, d and D are defined for C^1 -functions, d gives a point derivation at the unit I of G , D defines a left invariant derivation on $C^\infty(G)$ (i.e. $D(\delta_x \star f) = \delta_x \star D(f)$ for all $x \in G$). The example of [CG1] (related to that of Johnson [J] for $SO(3)$) is obtained for $X_0 = \begin{pmatrix} \frac{i}{2} & 0 \\ 0 & -\frac{i}{2} \end{pmatrix}$.

We take normalized Haar measure on G , then the Plancherel measure on the discrete space \widehat{G} gives the weights $\nu(\{\pi\}) = \dim(\pi)$. Thus for $f \in A(G)$ ($\subseteq L^1(G)$ in the compact case) $\|f\|_A = \sum_{\pi \in \widehat{G}} \dim(\pi) \|\pi(f)\|_1$ and $\|f\|_{VN} = \sup_{\pi \in \widehat{G}} \|\pi(f)\|$ for $f \in L^1(G)$. If f is a C^1 -function such that $f, f \star d \in L^1(G)$ it is easy to see that $\pi(f \star d) = \pi(f) \circ \pi(d)$ (using the extension of π to compactly supported

distributions). For X_0 as above, $\pi(d) = -F_\pi$ (for the standard realization of π) as given in [CG1] p. 6508. As explained there, one has $\|F_\pi\| = \frac{\dim(\pi)-1}{2}$. It follows that $\|f \star d\|_{VN} \leq \frac{1}{2}\|f\|_A$. By [E] Prop. 3.26 $A(G)$ contains the space $\mathcal{D}(G)$ of compactly supported C^∞ -functions as a dense subspace (for any Lie group G). It follows that D extends to a bounded operator $A(G) \rightarrow VN(G)$ which is still a derivation.

This construction of derivations on $C^\infty(G)$ can be applied on any Lie group G . For a compact semisimple Lie group, similar arguments as above show that this gives derivations $A(G) \rightarrow VN(G)$ for any $X \in \mathfrak{g}$. But the same estimate applies also to the corresponding right invariant derivations $D^{(r)}f = d \star f$. Combined the method produces a subspace of derivations $A(G) \rightarrow VN(G)$ with the dimension $2 \dim(G)$ (for any compact semisimple Lie group, with the extended theory of [HM] Th. 9.49 this extends to some extent even to general compact connected semisimple groups). In any case the guess made in [LLSS] Rem. 2.4 needs some revision.

4. THE $ax + b$ GROUP

$G = \mathbb{R} \rtimes \mathbb{R}$, the (proper) affine transformations of the real line. For clarity, we use the notation $G = \beta(\mathbb{R}) \alpha(\mathbb{R})$, i.e. $(x, t) \mapsto \beta(x) \alpha(t)$ defines a homeomorphism $\mathbb{R}^2 \rightarrow G$, $\alpha, \beta : \mathbb{R} \rightarrow G$ are homomorphisms and group multiplication is governed by the relation $\alpha(t) \beta(x) = \beta(e^t x) \alpha(t)$ ($t, x \in \mathbb{R}$; hence $\beta(\mathbb{R})$ is the normal factor). For (left) Haar measure we take $e^{-t} dx dt$. G is not unimodular, $\Delta(\beta(x) \alpha(t)) = e^{-t}$. Irreducible representations are described in numerous texts (and with numerous variations), starting with Gelfand, Naimark. We keep close to [CG1] sec. 4. Take $\mathcal{H} = L^2(\mathbb{R}^+, \frac{1}{\psi} d\psi)$ (with $\mathbb{R}^+ = [0, \infty[$) and for $f \in \mathcal{H}$, $(\pi^+(\beta(x))f)(\psi) = e^{-ib\psi} f(\psi)$, $(\pi^-(\beta(x))f)(\psi) = e^{ib\psi} f(\psi)$, $(\pi^\pm(\alpha(t))f)(\psi) = f(e^t \psi)$, $(Kf)(\psi) = \psi f(\psi)$ ($t, x \in \mathbb{R}$, $\psi \in \mathbb{R}^+$). π^+, π^- define non-equivalent representations and up to equivalence these are the only infinite dimensional irreducible representations of G . K ($= K_+, K_-$) is a Duflo–Moore operator for π^\pm , the Plancherel measure is concentrated on $\{\pi^+, \pi^-\}$ with $\nu(\{\pi^+\}) = \nu(\{\pi^-\}) = \frac{1}{2\pi}$. Thus \mathcal{F} maps $VN(G)$ isomorphically to the l^∞ -sum $\mathcal{B}(\mathcal{H}) \oplus \mathcal{B}(\mathcal{H})$. $\mathcal{F}_A(f) = ([\pi^\pm(f)K])$, $\|f\|_A = \frac{1}{2\pi}(\|[\pi^+(f)K]\|_1 + \|[\pi^-(f)K]\|_1)$ for $f \in A(G) \cap L^1(G)$. \mathcal{F}_A extends to an isometry of $A(G)$ onto the l^1 -sum $\mathcal{S}^1(\mathcal{H}) \oplus \mathcal{S}^1(\mathcal{H})$ weighted by ν .

We consider the distribution $d = \lim_{x \rightarrow 0} \frac{1}{x}(\delta_{\beta(-x)} - \delta_{\beta(0)})$ and $D(f) = f \star d$ (taking a direction from the normal factor; up to a constant factor this is the derivation D_b considered in [CG1] p. 6510). It follows that $(\pi^+(d)g)(\psi) = i\psi g(\psi)$, $(\pi^-(d)g)(\psi) = -i\psi g(\psi)$ hence $\pi^\pm(d) = \pm iK$. If f is a C^1 -function on G such that $f, f \star d \in L^1(G)$

one has $\pi(f \star d) = [\pi(f) \circ \pi(d)]$. Hence, if in addition $f \in A(G)$ holds, it follows that $\|f \star d\|_{VN} \leq 2\pi\|f\|_A$, D extends to a bounded derivation $A(G) \rightarrow VN(G)$.

If d involves a direction from the non-normal factor one can show that this does not give a bounded mapping $A(G) \rightarrow VN(G)$, the same when considering right invariant derivations $D^{(r)}f = d \star f$. Hence for the $ax+b$ group the method produces just a one dimensional space of derivations $A(G) \rightarrow VN(G)$.

5. THE HEISENBERG GROUP \mathbb{H}

We use the parametrization given in Section 2, $\mathbb{H} = \mathbb{R}^2 \rtimes \mathbb{R}$, considering \mathbb{R}, \mathbb{R}^2 as subgroups (i.e. omitting embeddings α, β as in Section 4) elements are written as v with $x \in \mathbb{R}$, $v \in \mathbb{R}^2$, with multiplication rule $xv = (x \circ v)x$ where $x \circ v = (y, z + xy)$ for $v = (y, z)$. For Haar measure we take standard Lebesgue measure of \mathbb{R}^3 . \mathbb{H} is unimodular. The representation theory is again classical. Take $\mathcal{H} = L^2(\mathbb{R})$. For real $t \neq 0$ an irreducible representation π_t is defined for $f \in \mathcal{H}$ by $(\pi_t(v)f)(\psi) = e^{it(z-y\psi)}f(\psi)$, $(\pi_t(x)f)(\psi) = f(\psi - x)$ ($x, \psi \in \mathbb{R}$, $v = (y, z) \in \mathbb{R}^2$). The representations π_t are pairwise non-equivalent and exhaust (up to equivalence) all the infinite dimensional irreducible representations of \mathbb{H} . The image of $\{\pi_t : t \neq 0\}$ (with the topology inherited from $\mathbb{R} \setminus \{0\}$) defines a locally compact (but non-closed) subset of $\widehat{\mathbb{H}}$. The Plancherel measure is concentrated on this subset and with the given parametrization of the representations we have $d\nu = \frac{1}{4\pi^2}|t|dt$. As explained in Appendix B, it follows that $\mathcal{B}_1^\oplus(\widehat{\mathbb{H}}, \nu) = L^1(\widehat{\mathbb{H}}, \nu, \mathcal{S}_1(\mathcal{H})) \cong L^1(\mathbb{R}, \frac{1}{4\pi^2}|t|dt, \mathcal{S}_1(\mathcal{H}))$ and $\|f\|_A = \frac{1}{4\pi^2} \int_{\mathbb{R}} \|\pi_t(f)\|_1 |t| dt$ for $f \in L^1(\mathbb{H}) \cap A(\mathbb{H})$. General elements of $\mathcal{B}_1^\oplus(\widehat{\mathbb{H}}, \nu)$ will be denoted as $T = (T_t)_{t \in \mathbb{R}}$.

We review now the approach of [CG2] for \mathbb{H} . $Z = \{(0, z) : z \in \mathbb{R}\}$ is the centre of \mathbb{H} and we consider the distribution $d = \lim_{z \rightarrow 0} \frac{1}{z}(\delta_{(0, -z)} - \delta_{(0, 0)})$ and $D(f) = f \star d$ (up to a constant factor this is the derivation ∂_Z considered in [CG2] (4.1)). It follows that $(\pi_t(d)g)(\psi) = -itg(\psi)$ (for $g \in \mathcal{H}$) and it is easy to see that now D does *not* produce a bounded mapping $A(\mathbb{H}) \rightarrow VN(\mathbb{H})$. Using distributional derivatives we have a linear mapping $D : A(\mathbb{H}) \rightarrow \mathcal{D}(\mathbb{H})'$. The idea is to construct a symmetric Banach $A(G)$ -bimodule containing the range of D and making D bounded. The module operation should extend pointwise multiplication. The obstacle is that the product with functions is not defined for general distributions and we should show that the distributions in the image of D are sufficiently tame to allow multiplication with functions from $A(\mathbb{H})$. Technically, this becomes practicable when working on the dual side.

Put $\mathcal{W} = L^1(\mathbb{R}, dt, \mathcal{K}(\mathcal{H}))$ and for $T = (T_t) \in \mathcal{B}_1^\oplus(\widehat{\mathbb{H}}, \nu)$ we put $\widehat{D}(T) = (-itT_t)$. Obviously, $\widehat{D}: \mathcal{B}_1^\oplus(\widehat{\mathbb{H}}, \nu) \rightarrow \mathcal{W}$ and $\|\widehat{D}(T)\|_{\mathcal{W}} = \int_{\mathbb{R}} |t| \|T_t(f)\|_1 dt = 2\pi \|T\|_{\mathcal{B}_1^\oplus}$. For a C^1 -function $f \in A(\mathbb{H})$ with $D(f) \in L^1(\mathbb{H})$ we have (using the extension of \mathcal{F} to $A(\mathbb{H})$) $\mathcal{F}(D(f)) = \widehat{D}(\mathcal{F}(f))$. $\mathcal{W} \cap \mathcal{B}_1^\oplus$ is dense in \mathcal{W} . The main observation of [CG2] is now that dual convolution \sharp can be extended to make \mathcal{W} a (Banach) \mathcal{B}_1^\oplus -module (hence also an $A(\mathbb{H})$ -module). D defines certainly a derivation on the space of C^1 -functions, thus by density and continuity $\widehat{D} \circ \mathcal{F}: A(\mathbb{H}) \rightarrow \mathcal{W}$ becomes a derivation.

Products of representation coefficients are coefficients of the tensor product of the representations $(\pi_1(x)f_1|g_1)(\pi_2(x)f_2|g_2) = ((\pi_1 \otimes \pi_2)(x)(f_1 \otimes f_2)|g_1 \otimes g_2)$. Formulas for \sharp are obtained by considering decompositions of $\pi_1 \otimes \pi_2$ into irreducible representations. For the Heisenberg group \mathbb{H} this was done in [CG2] sec. 5 (their notation is slightly different, defining \sharp as a product on $L^1(\widehat{\mathbb{H}}, dt, \mathcal{S}_1(\mathcal{H}))$ which is isomorphic to $\mathcal{B}_1^\oplus(\widehat{\mathbb{H}}, \nu)$ by the mapping $(\mathcal{T}F)_t = |t|^{-1}F_t$ — this simplifies some formulas). More abstractly, π_t can be obtained as an induced representation from the subgroup \mathbb{R}^2 and then one can apply e.g. [KL] Th. 1.4.11 (a special case of Mackey's tensor product theorem). This gives $\pi_{t_1} \otimes \pi_{t_2} \simeq \infty \pi_{t_1+t_2}$. But we need an explicit description of the equivalence.

$\mathcal{H} \otimes_2 \mathcal{H}$ denotes the (completed) Hilbert space tensor product and we use the identification with $L^2(\mathbb{R}^2)$ (be aware that [Fu] p. 9f. uses a different notion of tensor product which is quite non-standard, compare [Fo] 7.3). For $t_1 + t_2 \neq 0$, we consider a unimodular coordinate transformations giving rise to a unitary operator $(W_{t_1, t_2} f)(\psi_1, \psi_2) = f(\psi_1 - \frac{t_2}{t_1+t_2}\psi_2, \psi_1 + \frac{t_1}{t_1+t_2}\psi_2)$ for $f \in L^2(\mathbb{R}^2)$. Then $W_{t_1, t_2} \circ ((\pi_{t_1} \otimes \pi_{t_2})(h)) = (\pi_{t_1+t_2}(h) \otimes 1) \circ W_{t_1, t_2}$ for $h \in \mathbb{H}$ (1 denoting the identity operator) and the mapping $(t_1, t_2) \mapsto W_{t_1, t_2}$ is continuous for the strong operator topology. Let $1 \otimes \text{tr}: \mathcal{S}^1(\mathcal{H} \otimes_2 \mathcal{H}) \rightarrow \mathcal{S}^1(\mathcal{H})$ be the dual operator for the embedding $A \mapsto A \otimes 1$ ($\mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H} \otimes_2 \mathcal{H})$), i.e. $\text{tr}(A \circ (1 \otimes \text{tr})(B)) = \text{tr}((A \otimes 1) \circ B)$ for $A \in \mathcal{B}(\mathcal{H})$, $B \in \mathcal{B}(\mathcal{H} \otimes_2 \mathcal{H})$. For $A_1, A_2 \in \mathcal{S}^1(\mathcal{H})$ put $\theta(A_1, A_2, t_1, t_2) = (1 \otimes \text{tr})(W_{t_1, t_2}(A_1 \otimes A_2)W_{t_1, t_2}^*) \in \mathcal{S}^1(\mathcal{H})$. This describes the products of coefficients of π_{t_1}, π_{t_2} , i.e. $\text{tr}(A_1 \pi_{t_1}(h)^*) \text{tr}(A_2 \pi_{t_2}(h)^*) = \text{tr}(\theta(A_1, A_2, t_1, t_2) \pi_{t_1+t_2}(h)^*)$ for $h \in \mathbb{H}$. θ is continuous in all variables (for $\|\cdot\|_1$ and $t_1 + t_2 \neq 0$). It follows that (in our notation) $|t| (S \sharp T)_t = \frac{1}{2\pi} \int_{\mathbb{R}} \theta(|s|S_s, |t-s|T_{t-s}, s, t-s) ds$ for $S, T \in \mathcal{B}_1^\oplus(\widehat{\mathbb{H}}, \nu)$.

Recall that $\mathcal{F}(uv) = \mathcal{F}(u) \sharp \mathcal{F}(v)$ for $u, v \in A(\mathbb{H})$ and that \mathcal{F} extends to an isometric isomorphism $VN(\mathbb{H}) \rightarrow L^\infty(\mathbb{R}, \nu, \mathcal{B}(\mathcal{H}))$. This provides estimates for the operator norm $\|S \sharp T\| \leq \|S\|_1 \|T\|$ for $S \in \mathcal{B}_1^\oplus$, $T \in \mathcal{B}_1^\oplus \cap L^\infty (= \mathcal{F}(A(\mathbb{H}) \cap VN(\mathbb{H})))$.

Considering this for $S_t = f(t)A_1$, $T_t = g(t)A_2$ with continuous f, g it follows by standard arguments that $\|\theta(A_1, A_2, t_1, t_2)\| \leq \frac{|t_2|}{|t_1+t_2|} \|A_1\|_1 \|A_2\|$ for $A_1, A_2 \in \mathcal{S}^1(\mathcal{H})$, $t_1 + t_2 \neq 0$ (it does not look easy to prove this more directly, alternatively one can also use arguments as in the proof of [CG2] Th. 4.3 on p. 2458). This easily implies $\|S \sharp T\|_{\mathcal{W}} \leq \|S\|_1 \|T\|_{\mathcal{W}}$ for $S \in \mathcal{B}_1^\oplus$, $T \in \mathcal{W} \cap \mathcal{B}_1^\oplus$ and this allows to extend \sharp , making \mathcal{W} a \mathcal{B}_1^\oplus -module.

6. THE GROUPS $\mathbb{H}_{\mathcal{K}}$

The reduced Heisenberg group has been treated in [CG1] sec. 6 and for general groups $\mathbb{H}_{\mathcal{K}}$ the method works similarly. We take up the notations of Section 2, $\mathbb{H}_{\mathcal{K}} = (\mathbb{R} \times \mathcal{K}) \rtimes \mathbb{R}$, \mathcal{K} is a non-trivial compact group and we have given a continuous homomorphism $\varphi_0: \mathbb{R} \rightarrow \mathcal{K}$ with dense image, \mathcal{K} is written additively. The action of \mathbb{R} on $\mathbb{R} \times \mathcal{K}$ is defined by $x \circ (y, k) = (y, k + \varphi_0(xy))$ ($x, y \in \mathbb{R}$, $k \in \mathcal{K}$). Then $\varphi(y, z) = (y, \varphi_0(z))$ extends to a continuous homomorphism $\varphi: \mathbb{H} \rightarrow \mathbb{H}_{\mathcal{K}}$ with dense image. $j_{\mathcal{K}}(k) = (0, k)$ ($j_{\mathcal{K}}: \mathcal{K} \rightarrow \mathbb{H}_{\mathcal{K}}$) maps \mathcal{K} isomorphically onto the centre of $\mathbb{H}_{\mathcal{K}}$, similarly with $j_Z(z) = (0, z)$ ($j_Z: \mathbb{R} \rightarrow \mathbb{H}$), and we have $\varphi \circ j_Z = j_{\mathcal{K}} \circ \varphi_0$. For Haar measure we take the product of standard Lebesgue measure on \mathbb{R}^2 and normalized Haar measure on \mathcal{K} . $\mathbb{H}_{\mathcal{K}}$ is unimodular. The dual group $\Gamma = \widehat{\mathcal{K}}$ is considered as a subgroup of \mathbb{R} in such a way that for $t \in \Gamma$ the corresponding character χ_t of \mathcal{K} satisfies $\chi_t(\varphi_0(z)) = e^{itz}$ for $z \in \mathbb{R}$ (see Rem. (b) in Section 2 and Rem. (c) in Appendix A).

If π is an irreducible representation of $\mathbb{H}_{\mathcal{K}}$ then $\pi \circ \varphi$ must be irreducible. Hence if π is infinite dimensional it follows that $\pi \circ \varphi \simeq \pi_t$ for some $t \neq 0$. Conversely, looking at the definition of π_t in Sec. 5, it is easy to see that π_t can be written as $\pi \circ \varphi$ iff the mapping $\varphi_0(z) \mapsto e^{itz}$ is well defined on $\varphi_0(\mathbb{R})$ and extends continuously to \mathcal{K} (then defining a character of \mathcal{K}). By our identification this is equivalent to $t \in \Gamma \setminus \{0\}$. The corresponding representation induced on $\mathbb{H}_{\mathcal{K}}$ will be denoted again by π_t . The finite dimensional irreducible representations of \mathbb{H} are trivial on Z and one dimensional. Hence they are given by characters on $\mathbb{H}/Z \cong \mathbb{R}^2$ (projecting $(y, z)x \in \mathbb{H}$ to $(x, y) \in \mathbb{R}^2$) and every character of \mathbb{R}^2 defines also a one dimensional representation of $\mathbb{H}_{\mathcal{K}}$. As above we identify $(\mathbb{R}^2)^\wedge$ with \mathbb{R}^2 associating to $w \in \mathbb{R}^2$ the character $\chi_w(v) = e^{i(v|w)}$ ($v \in \mathbb{R}^2$). The corresponding representation of $\mathbb{H}_{\mathcal{K}}$ (and its extension to L^1 and distributions) will also be denoted by χ_w . With these notations and parametrizations we get $(\mathbb{H}_{\mathcal{K}})^\wedge = (\Gamma \setminus \{0\}) \cup \mathbb{R}^2$. It is not hard to see that both parts are open, on \mathbb{R}^2 the dual induces the euclidian topology and on

$\Gamma \setminus \{0\}$ the discrete topology (in particular $(\mathbb{H}_{\mathcal{K}})^{\wedge}$ is Hausdorff; $\mathbb{H}_{\mathcal{K}}$ are FC^{-} -groups, general results for the dual of FC^{-} -groups have been shown in [Li]).

Lemma 2. *Using Haar measure and parametrization as above, the Plancherel measure ν of $\mathbb{H}_{\mathcal{K}}$ satisfies $d\nu = \frac{1}{4\pi^2}dw$ on \mathbb{R}^2 and $\nu(\{t\}) = \frac{|t|}{2\pi}$ for $t \in \Gamma \setminus \{0\}$.*

Proof. This follows by similar computations as for the Heisenberg group ([Fo] 7.6.1). \square

If \mathcal{K} is not metrizable (equivalently, if Γ is uncountable) this gives an example of the Plancherel theorem for a group that is non-compact, σ -compact, but not second countable – of course the Plancherel measure is not σ -finite in that case (see also Rem. (d) in Appendix B). It follows that $\mathcal{B}_1^{\oplus}((\mathbb{H}_{\mathcal{K}})^{\wedge}, \nu)$ is the sum of (complex valued) $L^1(\mathbb{R}^2, \frac{1}{4\pi^2}dw)$ and a $(\mathcal{S}^1(\mathcal{H})$ -valued) weighted l^1 -space. $\|T\|_1 = \frac{1}{4\pi^2} \int_{\mathbb{R}^2} |T_w| dw + \frac{1}{2\pi} \sum_{t \in \Gamma \setminus \{0\}} |t| \|T_t\|_1$.

We project now the distribution from Section 5 to $\mathbb{H}_{\mathcal{K}}$, putting $d = \lim_{z \rightarrow 0} \frac{1}{z}(\delta_{\varphi(0,-z)} - \delta_{\varphi(0,0)})$ and $D(f) = f \star d$. Recall that $\varphi(0, z) = \varphi_0(z)$. If \mathcal{K} is a Lie group (equivalently, if Γ is finitely generated), φ_0 is analytic ([Va] L. 2.11.1 or more directly, using a duality argument) and it follows that d and D are defined for $f \in C^1(\mathbb{H}_{\mathcal{K}})$ ($\mathbb{H}_{\mathcal{K}}$ being again a Lie group). In the general case, \mathcal{K} is the projective limit of Lie groups \mathcal{K}/K_1 (e.g. taking all finitely generated subgroups of Γ) and $\mathbb{H}_{\mathcal{K}/K_1} \cong \mathbb{H}_{\mathcal{K}}/j_{\mathcal{K}}(K_1)$. With the usual embeddings (see also Rem. (d) in Appendix B) d and D are defined for $f \in \bigcup \{C^1(\mathbb{H}_{\mathcal{K}/K_1}) : K_1 \text{ a closed subgroup of } \mathcal{K} \text{ with } \mathcal{K}/K_1 \text{ Lie}\}$. It is easy to see that $\chi_w(d) = 0$ for $w \in \mathbb{R}^2$ and (as in Section 5) $(\pi_t(d)g)(\psi) = -itg(\psi)$ for $g \in \mathcal{H}$ and $t \in \Gamma \setminus \{0\}$. Then (similar to Section 3) our formula for the Plancherel measure (Lemma 2) implies that $\|f \star d\|_{VN} \leq 2\pi \|f\|_A$ for $f \in A(\mathbb{H}_{\mathcal{K}}) \cap L^1(\mathbb{H}_{\mathcal{K}}) \cap \bigcup C^1(\mathbb{H}_{\mathcal{K}/K_1})$ with $f \star d \in L^1$ (e.g. f of compact support). It follows that D extends to a derivation $A(G) \rightarrow VN(G)$.

Remarks. (a) We recall from [CG1] 2.1, for a Banach algebra A , a derivation $D : A \rightarrow A'$ is called cyclic if $(Da, b) = -(Db, a)$ holds for $a, b \in A$ ((\cdot, \cdot) denoting the evaluation of functionals). Grønbæk called A cyclically amenable if there are no non-zero bounded cyclic derivations. For $SO(3)$, $SU(2)$, $ax + b$, $\mathbb{H}_{\mathcal{K}}$ the derivations that we considered are cyclic. By [CG1] Prop. 2.5, for any locally compact group G that has one of these groups as a closed subgroup the Fourier algebra $A(G)$ is not cyclically amenable. But the situation is not clear for the groups \mathbb{H} , $\widetilde{M(2)}$, $M(2)_{\mathcal{K}}$, \mathbb{G}_a .

(b) The distribution d that we used for $\mathbb{H}_{\mathcal{K}}$ is central, thus $D^{(r)} = D$ (D is left and right invariant). For derivatives involving a non-central direction one does not get

a bounded mapping $A(\mathbb{H}_{\mathcal{K}}) \rightarrow VN(\mathbb{H}_{\mathcal{K}})$. Hence the present method produces just a one dimensional space of derivations $A(\mathbb{H}_{\mathcal{K}}) \rightarrow VN(\mathbb{H}_{\mathcal{K}})$.

7. THE EUCLIDIAN MOTION GROUP $M(2)$

$M(2)$ is not covered by [CG1],[CG2]. In [LLSS] sec.2.2 it was shown that $A(M(2))$ is not weakly amenable, using spectral synthesis. We develop here our method in detail, since the constructions are basic for the treatment of $M(2)_{\mathcal{K}}$ in the next section. Defining derivations as before, it turns out that they do not give bounded mappings $A(M(2)) \rightarrow VN(M(2))$. Working on the dual side, we are constructing a \mathcal{B}_1^{\oplus} -module \mathcal{W} such that $\widehat{D} : \mathcal{B}_1^{\oplus} \rightarrow \mathcal{W}$ is bounded. First we will derive a formula for the dual convolution \sharp . For $M(2)$ it turns out that \sharp behaves quite differently compared to the case of \mathbb{H} . We did not succeed to find estimates making it possible to define \mathcal{W} similarly as in Section 5 as a weighted L^1 -space of $\mathcal{K}(\mathcal{H})$ -valued (or $\mathcal{S}^1(\mathcal{H})$ -valued) functions. Recall that if $\mathcal{H} = L^2(\Omega)$, trace class operators are integral operators and for a finite measure space this gives embeddings $\mathcal{S}^1(\mathcal{H}) \subseteq \mathcal{S}^2(\mathcal{H}) = L^2(\Omega^2) \subseteq L^1(\Omega^2)$. In our case, for the infinite dimensional representations we may choose $\Omega = \mathbb{T} = \mathbb{R}/(2\pi\mathbb{Z})$ with normalized Haar measure $\frac{1}{2\pi}d\psi$. We will define \mathcal{W} as a subspace of the weighted L^1 -space $\overline{\mathcal{W}} = L^1([0, \infty[, w(\rho) d\rho, L^1(\mathbb{T}^2))$ with the weight function $w(\rho) = \frac{\rho}{1 + \rho^2}$, leading (partially) outside $\mathcal{B}(\mathcal{H})$.

Similar to Section 5 we use the parametrization $M(2) = \mathbb{R}^2 \rtimes \mathbb{T}$. With the rotation matrices $A_{\alpha} = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix} = \exp(\alpha \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix})$ ($\alpha \in \mathbb{R}$, but depending only on the coset mod 2π), the action is defined by $\alpha \circ v = A_{\alpha}v$ ($v \in \mathbb{R}^2$). For Haar measure we take $\frac{1}{(2\pi)^2} dv d\alpha$ ([Su] Ch. IV, (3.1)). $M(2)$ is unimodular.

The representation theory is again classical. Take $\mathcal{H} = L^2(\mathbb{R}, \frac{1}{2\pi}d\psi)$. For real $\rho > 0$ an irreducible representation π_{ρ} is defined for $f \in \mathcal{H}$ by $(\pi_{\rho}(v)f)(\psi) = e^{i\rho r \cos(\phi-\psi)} f(\psi)$, $(\pi_{\rho}(\alpha)f)(\psi) = f(\psi - \alpha)$ ($\alpha, \psi \in \mathbb{T}$, $v = r \begin{pmatrix} \cos \phi \\ \sin \phi \end{pmatrix} \in \mathbb{R}^2$, polar coordinates). $M(2)$ is of type I. The representations π_{ρ} are pairwise non-equivalent and exhaust (up to equivalence) all the infinite dimensional irreducible representations of $M(2)$. For usage in the next section, we discuss also the finite dimensional representations. We have $M(2)/\mathbb{R}^2 \cong \mathbb{T}$. For $n \in \mathbb{Z}$, $\chi_n(\psi) = e^{in\psi}$ ($\psi \in \mathbb{T}$) lifts to a one dimensional representation of $M(2)$ (again denoted by χ_n). We have $\widehat{M(2)} = \{\chi_n : n \in \mathbb{Z}\} \cup \{\pi_{\rho} : \rho > 0\}$ (more precisely: this gives a complete set of representatives). $\{\pi_{\rho} : \rho > 0\}$ is an open subset and $\widehat{M(2)}$ induces

the standard topology of $]0, \infty[$. On $\{\chi_n : n \in \mathbb{Z}\}$ we get the discrete topology, $\overline{\{\pi_\rho\}} = \{\pi_\rho\} \cup \{\chi_n : n \in \mathbb{Z}\}$ for each $\rho > 0$. The Plancherel measure is concentrated on $\{\pi_\rho : \rho > 0\}$ and with the given parametrization of the representations we have $d\nu = \rho d\rho$ ([Su] IV, Th. 4.2, [Vi] Ch. IV §5 uses a different normalization of Haar measure, resulting in a compensating factor in the Plancherel measure). Hence $\mathcal{B}_1^\oplus(\widehat{M(2)}, \nu) \cong L^1(]0, \infty[, \rho d\rho, \mathcal{S}_1(\mathcal{H}))$, general elements of $\mathcal{B}_1^\oplus(\widehat{M(2)}, \nu)$ will be denoted as $T = (T_\rho)_{\rho>0}$.

For the direction $e_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \in \mathbb{R}^2$ we will consider, similarly as before, the distribution $d = \lim_{y \rightarrow 0} \frac{1}{y}(\delta_{-ye_1} - \delta_{0e_1})$ and $D(f) = f \star d$. It follows that $(\pi_\rho(d)g)(\psi) = -i\rho \cos(\psi) g(\psi)$ (for $g \in \mathcal{H}$) and it is easy to see that D does *not* produce a bounded mapping $A(M(2)) \rightarrow VN(M(2))$. As sketched above, we will next investigate the dual convolution.

For the decomposition of tensor products, we follow [Vi] Ch. IV, §6. $\mathcal{H} \otimes_2 \mathcal{H}$ (completed Hilbert space tensor product) is identified with $L^2(\mathbb{T}^2, \frac{1}{(2\pi)^2} d(\psi_1, \psi_2))$. For $f \in \mathcal{H} \otimes_2 \mathcal{H}$ define $(Wf)(\mu)(\psi) = f(\psi, \psi + \mu)$. It follows from standard measure theory that $(Wf)(\mu) \in \mathcal{H}$ for a.e. μ and $W : \mathcal{H} \otimes \mathcal{H} \rightarrow L^2(\mathbb{T}, \frac{1}{2\pi} d\mu, \mathcal{H})$ gives an isometric isomorphism.

For $\varphi \in \mathbb{T}$, $g \in \mathcal{H}$, we define $(U_\varphi g)(\psi) = g(\psi - \varphi)$, giving a unitary operator on \mathcal{H} , depending strongly continuously on φ . Fix now $\rho_1, \rho_2 > 0$ and for $\mu \in \mathbb{R}$ put $\rho_1 + \rho_2 e^{i\mu} = \rho e^{i\varphi}$, defining functions $\rho = \rho(\rho_1, \rho_2, \mu) = |\rho_1 + \rho_2 e^{i\mu}|$ and $\varphi = \varphi(\rho_1, \rho_2, \mu)$ (using only the cosets of $\mu, \varphi \bmod 2\pi$). Finally, we define $(W_{\rho_1, \rho_2} f)(\mu) = U_\varphi((Wf)(\mu))$ ($f \in \mathcal{H} \otimes_2 \mathcal{H}$), giving again an isometric isomorphism $W_{\rho_1, \rho_2} : \mathcal{H} \otimes_2 \mathcal{H} \rightarrow L^2(\mathbb{T}, \frac{1}{2\pi} d\mu, \mathcal{H})$, $(\rho_1, \rho_2) \mapsto W_{\rho_1, \rho_2}$ is strongly continuous. For $n \in \mathbb{Z}$ we define $M_n g = \chi_n \cdot g$ giving also a unitary operator on \mathcal{H} .

Proposition 2. *For $\rho_1, \rho_2 > 0$, $n \in \mathbb{Z}$, $x \in M(2)$, we have*

$$W_{\rho_1, \rho_2} \circ (\pi_{\rho_1} \otimes \pi_{\rho_2})(x) \circ W_{\rho_1, \rho_2}^* = \int_{\mathbb{T}}^{\oplus} \pi_\rho(x) d\mu$$

$$(\chi_n \otimes \pi_\rho)(x) = M_n \pi_\rho(x) M_n^*.$$

The first formula gives a decomposable operator on $L^2(\mathbb{T}, \frac{1}{2\pi} d\mu, \mathcal{H})$ ([T1] Ch. IV, Def. 7.9). Both generalize to the extensions of the representations to $L^1(M(2))$.

Proof. Take $f \in \mathcal{H} \otimes_2 \mathcal{H}$ and put $g = W_{\rho_1, \rho_2} f$. Then $f = W_{\rho_1, \rho_2}^* g$ and $g(\mu)(\psi) = (Wf)(\mu)(\psi - \varphi) = f(\psi - \varphi, \psi - \varphi + \mu)$. It is enough to consider the case $f = f_1 \otimes f_2$, then $f(\psi - \varphi, \psi - \varphi + \mu) = f_1(\psi - \varphi) f_2(\psi - \varphi + \mu)$ and $(\pi_{\rho_1} \otimes \pi_{\rho_2})(x) f = (\pi_{\rho_1}(x) f_1) \otimes (\pi_{\rho_2}(x) f_2)$. For $\alpha \in \mathbb{T}$ an easy computation gives $[(W_{\rho_1, \rho_2} \circ (\pi_{\rho_1} \otimes \pi_{\rho_2})(\alpha)) f](\mu)(\psi) = f_1(\psi - \varphi - \alpha) f_2(\psi + \mu - \varphi - \alpha) = [\pi_\rho(\alpha)(g(\mu))](\psi)$, and vor $v = r \begin{pmatrix} \cos \phi \\ \sin \phi \end{pmatrix} \in \mathbb{R}^2$, using

$\rho_1 \left(\frac{\cos \psi}{\sin \psi} \right) + \rho_2 \left(\frac{\cos(\psi+\mu)}{\sin(\psi+\mu)} \right) = \rho \left(\frac{\cos(\psi+\varphi)}{\sin(\psi+\varphi)} \right)$ one gets $[(W_{\rho_1, \rho_2} \circ (\pi_{\rho_1} \otimes \pi_{\rho_2})(v))f](\mu)(\psi) = e^{i\rho r \cos(\phi-\psi)} f_1(\psi - \varphi) f_2(\psi + \mu - \varphi) = [\pi_\rho(v)(g(\mu))](\psi)$ showing the first formula.

We identify $\mathbb{C} \otimes \mathcal{H}$ with \mathcal{H} (via $\gamma \otimes h \mapsto \gamma h$) and consider $\chi_n \otimes \pi_\rho$ as a representation on \mathcal{H} . The second formula is checked easily (it implements the equivalence $\chi_n \otimes \pi_\rho \simeq \pi_\rho$). \square

The embedding of the subalgebra $L^\infty(\mathbb{T}) \overline{\otimes} \mathcal{B}(\mathcal{H})$ of decomposable operators into the algebra $\mathcal{B}(L^2(\mathbb{T}, \mathcal{H}))$ of all bounded operators defines a projection $\text{pr}_\mathcal{S} : \mathcal{S}^1(L^2(\mathbb{T}, \mathcal{H})) \rightarrow L^1(\mathbb{T}, \mathcal{S}^1(\mathcal{H}))$ of the preduals (using [T1] Ch. IV, Th. 7.17). For $A_1, A_2 \in \mathcal{S}^1(\mathcal{H})$ define

$$\theta_0(A_1, A_2) = \text{pr}_\mathcal{S}(W_{\rho_1, \rho_2} \circ (A_1 \otimes A_2) \circ W_{\rho_1, \rho_2}^*) \quad (\in L^1(\mathbb{T}, \frac{1}{2\pi} d\mu, \mathcal{S}^1(\mathcal{H})))$$

(a more explicit description in terms of integration kernels will be given in Lemma 3 below).

Corollary 1. *For $A_1, A_2 \in \mathcal{S}^1(\mathcal{H})$ consider coefficients $u_j(x) = \text{tr}(A_j \pi_{\rho_j}(x)^*)$ ($j = 1, 2$). Then*

$$u_1(x) u_2(x) = \frac{1}{2\pi} \int_0^{2\pi} \text{tr}(\pi_\rho(x)^* \theta_0(A_1, A_2)(\mu)) d\mu.$$

Proof. It is enough to show this for $A_j h = (h|h_j)k_j$, where $h_j, k_j \in \mathcal{H}$ ($j = 1, 2$). Then $u_j(x) = (k_j | \pi_{\rho_j}(x) h_j)$, $u_1(x) u_2(x) = (k_1 \otimes k_2 | [(\pi_{\rho_1} \otimes \pi_{\rho_2})(x)](h_1 \otimes h_2))$ and the formula follows by a short computation from Proposition 2. \square

By a change of coordinates, this can be expressed in terms of the Plancherel measure. We have $\cos \mu = \frac{\rho^2 - \rho_1^2 - \rho_2^2}{2\rho_1 \rho_2}$, hence $\mu \mapsto \rho(\rho_1, \rho_2, \mu)$ (briefly $\rho(\mu)$) is injective on $[0, \pi]$ and on $[\pi, 2\pi]$ and it maps both intervals onto $[|\rho_1 - \rho_2|, \rho_1 + \rho_2]$. In addition $\rho(\mu) = \rho(2\pi - \mu)$. For $A_1, A_2 \in \mathcal{S}^1(\mathcal{H})$, $\rho = \rho(\mu)$, we define

$$\theta(A_1, A_2)(\rho) = \theta_0(A_1, A_2)(\mu) + \theta_0(A_1, A_2)(2\pi - \mu)$$

(θ_0, θ depend on ρ_1, ρ_2 as well, sometimes we will write $\theta(A_1, A_2, \rho_1, \rho_2)$). For $|\rho_1 - \rho_2| < \rho < \rho_1 + \rho_2$ put $K(\rho_1, \rho_2, \rho) = \frac{2}{\pi \sqrt{4\rho_1^2 \rho_2^2 - (\rho^2 - \rho_1^2 - \rho_2^2)^2}}$ (this is the kernel defining the Bessel-Kingman hypergroup – [BH] p. 235 with $\alpha = 0$).

Corollary 2. *For A_1, A_2, u_1, u_2 as in Corollary 1, we have $\theta(A_1, A_2, \rho_1, \rho_2) \in L^1([|\rho_1 - \rho_2|, \rho_1 + \rho_2], \frac{1}{2} K(\rho_1, \rho_2, \rho) \rho d\rho, \mathcal{S}^1(\mathcal{H}))$ and*

$$u_1(x) u_2(x) = \int_{|\rho_1 - \rho_2|}^{\rho_1 + \rho_2} \frac{1}{2} \text{tr}(\pi_\rho(x)^* \theta(A_1, A_2)(\rho)) K(\rho_1, \rho_2, \rho) d\nu(\rho).$$

Thus $u_1 \cdot u_2 = \mathcal{F}^{-1}\left(\left(\frac{1}{2}\theta(A_1, A_2)(\rho) K(\rho_1, \rho_2, \rho)\right)_{|\rho_1 - \rho_2| < \rho < \rho_1 + \rho_2}\right) \in A(M(2))$ (the operator field is meant to be extended by 0 for ρ outside the interval). In particular, one gets that $\pi_{\rho_1} \otimes \pi_{\rho_2} \simeq 2 \int_{|\rho_1 - \rho_2|}^{\rho_1 + \rho_2} \pi_\rho d\rho$, but the formula above gives more detailed information for the product of coefficients.

Proof. Again, an elementary computation shows that under the substitution, $\frac{1}{2\pi}d\mu = \frac{1}{2}K(\rho_1, \rho_2, \rho) \rho d\rho$ holds for $\mu \in [0, \pi]$ and similarly for $\mu \in [\pi, 2\pi]$. \square

Corollary 3. For $S = (S_\rho), T = (T_\rho) \in \mathcal{B}_1^\oplus(\widehat{M(2)}, \nu)$, we get

$$(S \sharp T)_\rho = \iint_{D_\rho} \frac{1}{2} \theta(S_{\rho_1}, T_{\rho_2})(\rho) K(\rho_1, \rho_2, \rho) \rho_1 \rho_2 d(\rho_1, \rho_2)$$

where D_ρ denotes the half-strip $\{(\rho_1, \rho_2) : \rho_1, \rho_2 > 0, |\rho_1 - \rho_2| < \rho < \rho_1 + \rho_2\}$.

It follows that if $S_\rho = 0$ for $\rho \notin [a_1, b_1]$, $T_\rho = 0$ for $\rho \notin [a_2, b_2]$ then $(S \sharp T)_\rho = 0$ for $\rho \notin]|a_1 - a_2|, b_1 + b_2[$ (see also Lemma 6). One can give similar expressions for products of functions from the Fourier-Stieltjes algebra, using arbitrary finite measures on $]0, \infty[$ and adding contributions from the one dimensional representations.

Proof. For fixed $\rho_1, \rho_2 > 0$, $A_1, A_2 \in \mathcal{S}^1(\mathcal{H})$, we have noted after Corollary 2 that $\rho \mapsto \frac{1}{2}\theta(A_1, A_2)(\rho) K(\rho_1, \rho_2, \rho)$ (extended by 0 outside $]|\rho_1 - \rho_2|, \rho_1 + \rho_2[$) defines an element $\text{dc}(A_1, A_2, \rho_1, \rho_2)$ of $\mathcal{B}_1^\oplus(\widehat{M(2)})$ of norm $\leq \|A_1\|_1 \|A_2\|_1$. Once we can show that for $S, T \in \mathcal{B}_1^\oplus(\widehat{M(2)})$ $(\rho_1, \rho_2) \mapsto \text{dc}(S_{\rho_1}, T_{\rho_2}, \rho_1, \rho_2)$ defines a measurable mapping $]0, \infty[^2 \rightarrow \mathcal{B}_1^\oplus(\widehat{M(2)})$, its integrability (with respect to $\nu \otimes \nu$) follows immediately, hence the integral in the Corollary defines an element $\text{dc}(S, T)$ of $\mathcal{B}_1^\oplus(\widehat{M(2)})$. Then, by Fubini's theorem and Corollary 2 it is not hard to see that $\mathcal{F}^{-1}(\text{dc}(S, T)) = \mathcal{F}^{-1}(S) \mathcal{F}^{-1}(T)$, hence $\text{dc}(S, T) = S \sharp T$ proving the Corollary.

Since $(\rho_1, \rho_2) \mapsto W_{\rho_1, \rho_2}$ is strongly continuous it follows that $(\rho_1, \rho_2) \mapsto W_{\rho_1, \rho_2} \circ (A_1 \otimes A_2) \circ W_{\rho_1, \rho_2}^*$ is norm continuous for fixed $A_1, A_2 \in \mathcal{S}^1(\mathcal{H})$. Consequently, $(\rho_1, \rho_2) \mapsto \theta_0(A_1, A_2, \rho_1, \rho_2) \in L^1(\mathbb{T}, \mathcal{S}^1(\mathcal{H}))$ is norm continuous. From basic properties of the Bochner integral (see e.g. [DS] Th. III.11.17) it follows that $(\rho_1, \rho_2, \mu) \mapsto \theta_0(A_1, A_2, \rho_1, \rho_2)(\mu) \in \mathcal{S}^1(\mathcal{H})$ is (a.e.) measurable on $]0, \infty[^2 \times \mathbb{T}$. The substitution $\rho = \rho(\mu)$ being continuous, we get that $(\rho_1, \rho_2, \rho) \mapsto \theta(A_1, A_2, \rho_1, \rho_2)(\rho)$ is measurable on $\{(\rho_1, \rho_2, \rho) : |\rho_1 - \rho_2| < \rho < \rho_1 + \rho_2\}$. Then by standard measure theoretical approximation techniques it follows that $(\rho_1, \rho_2, \rho) \mapsto \theta(S_{\rho_1}, T_{\rho_2}, \rho_1, \rho_2)(\rho)$ is measurable for $S = (S_\rho), T = (T_\rho) \in \mathcal{B}_1^\oplus(\widehat{M(2)})$. $(\rho_1, \rho_2, \rho) \mapsto K(\rho_1, \rho_2, \rho)$ being continuous, we get measurability of $(\rho_1, \rho_2) \mapsto \text{dc}(S_{\rho_1}, T_{\rho_2}, \rho_1, \rho_2)$. \square

We represent now as integral operators $(Ah)(\psi) = \frac{1}{2\pi} \int_0^{2\pi} k_A(\psi, \psi_1) h(\psi_1) d\psi_1$ ($\psi \in \mathbb{T}$, $h \in L^2(\mathbb{T})$; $k_A \in L^2(\mathbb{T}^2)$ being called the integration kernel for A).

By classical results, $A \mapsto k_A$ defines an isometric isomorphism $\mathcal{S}^2(\mathcal{H}) \rightarrow L^2(\mathbb{T}^2, \frac{1}{4\pi^2}d(\psi, \psi_1))$. Recall that $\mathcal{S}^1(\mathcal{H}) \subseteq \mathcal{S}^2(\mathcal{H})$, hence this applies for $A \in \mathcal{S}^1(\mathcal{H})$. For one dimensional operators $Ah = (h|h_0)k_0$, $h_0, k_0 \in \mathcal{H}$ (as in the proof of Corollary 1 above), one has $k_A(\psi, \psi_1) = k_0(\psi) \overline{h_0(\psi_1)}$. In general, k_A can also be obtained by considering $A \in \mathcal{S}^1(\mathcal{H})$ as an absolutely convergent sum of one dimensional operators. The operators M_n ($n \in \mathbb{Z}$) defined before Proposition 2 induce operators $\mathbf{M}_n: \mathcal{S}_1(\mathcal{H}) \rightarrow \mathcal{S}_1(\mathcal{H})$ by $\mathbf{M}_n(A) = M_n \circ A \circ M_n^*$.

Lemma 3. *For $\rho_1, \rho_2 > 0$, $A_1, A_2 \in \mathcal{S}^1(\mathcal{H})$ we have for almost all $\mu \in \mathbb{T}$*
 $k_{\theta_0(A_1, A_2)(\mu)}(\psi, \psi_1) = k_{A_1}(\psi - \varphi, \psi_1 - \varphi) k_{A_2}(\psi - \varphi + \mu, \psi_1 - \varphi + \mu)$.
 $k_{\mathbf{M}_n(A)}(\psi, \psi_1) = \chi_n(\psi - \psi_1) k_A(\psi, \psi_1)$ for $A \in \mathcal{S}^1(\mathcal{H})$, $n \in \mathbb{Z}$.

$\varphi = \varphi(\rho_1, \rho_2, \mu)$ is defined before Proposition 2. Adding the kernels obtained above for μ and $2\pi - \mu$ (observe that $\varphi(2\pi - \mu) = -\varphi(\mu)$) gives a corresponding representation for $\theta(A_1, A_2)(\rho)$. For A_1, A_2 one dimensional, $\theta(A_1, A_2)(\rho)$ will (in general) be a two dimensional operator. It is not hard to construct examples of (e.g. one dimensional) $A_1, A_2 \in \mathcal{S}^1(\mathcal{H})$ such that (using the formula above) $k_{\theta_0(A_1, A_2)(\mu)} \notin L^2$ for some μ . Then, in general, $\theta(A_1, A_2)(\rho) \in \mathcal{S}^1(\mathcal{H})$ holds just ρ -a.e. and $\rho \mapsto \|\theta(A_1, A_2)(\rho)\|_1$ is unbounded.

Proof. This follows again from the definitions by direct computation, starting with one dimensional A_1, A_2 . \square

We extend the operators \mathbf{M}_n to operator fields, defining for $T = (T_\rho)$: $\mathbf{M}_n(T) = (\mathbf{M}_n(T_\rho))$ ($n \in \mathbb{Z}$).

Corollary. $\theta_0(A_1, \mathbf{M}_n(A_2)) = \mathbf{M}_n(\theta_0(A_1, A_2))$ for $A_1, A_2 \in \mathcal{S}^1(\mathcal{H})$.
 $S \sharp \mathbf{M}_n(T) = \mathbf{M}_n(S \sharp T)$ for $S, T \in \widehat{\mathcal{B}_1^\oplus(\widehat{M}(2), \nu)}$.

We use the embedding $A \mapsto k_A$ to consider \mathcal{S}^1 as a subset of $L^1(\mathbb{T}^2, \frac{1}{(2\pi)^2}d(\psi, \psi_1))$. For better distinction, we denote the L^1 -norm as $\|\cdot\|_{L^1}$, $\|\cdot\|_1$ shall refer (as before) to the \mathcal{S}^1 -norm. Using the formula of Lemma 3 we define $\theta(a_1, a_2)(\rho)$ for arbitrary measurable functions a_1, a_2 on \mathbb{T}^2 (thus $\theta(k_{A_1}, k_{A_2})(\rho) = k_{\theta(A_1, A_2)(\rho)}$). Also the operators \mathbf{M}_n ($n \in \mathbb{Z}$) extend to measurable functions on \mathbb{T}^2 , giving isometrical isomorphisms of $L^1(\mathbb{T}^2)$. As above, they extend further to $L^1(\mathbb{T}^2)$ -valued vector fields.

- Lemma 4.** (a) $\|k_A\|_{L^1} \leq \|A\|_1$
 (b) $\|\theta(k_{A_1}, k_{A_2})(\rho)\|_{L^1} \leq 2\|A_1\|_1\|A_2\|_1$
 (c) $\|\theta(a_1, a_2)(\rho)\|_{L^1} \leq 2\|a_1\|_{L^\infty}\|a_2\|_{L^1}$
 (d) $\|\theta(a_1, a_2)(\rho)\|_{L^\infty} \leq 2\|a_1\|_{L^\infty}\|a_2\|_{L^\infty}$.

Note also that $L^\infty(\mathbb{T}^2) \cap \mathcal{S}^1$ is dense in $\mathcal{S}^1(\mathcal{H})$ for the \mathcal{S}^1 -norm, \mathcal{S}^1 is dense in $L^1(\mathbb{T}^2)$ for the L^1 -norm.

Proof. For (a): $\|k_A\|_{L^1} \leq \|k_A\|_{L^2} = \|A\|_2 \leq \|A\|_1$. For (b) use Lemma 3 and the Cauchy-Schwarz inequality. (c),(d) are easy too. \square

For a weight function w on $]0, \infty[$ (i.e. $w:]0, \infty[\rightarrow [0, \infty[$ is a measurable function; it will be specified later, see Proposition 3 and the Remark (a) below), let

$$\overline{\mathcal{W}} = L^1\left(]0, \infty[, w(\rho) d\rho, L^1(\mathbb{T}^2, \frac{1}{(2\pi)^2} d(\psi, \psi_1))\right)$$

be the $L^1(\mathbb{T}^2)$ -valued *weighted* L^1 -space on $]0, \infty[$. It consists of (equivalence classes of) Bochner-measurable functions $t = (t_\rho)$ with $t_\rho \in L^1(\mathbb{T}^2)$ and $\|t\|_{\overline{\mathcal{W}}} = \int_0^\infty \|t_\rho\|_{L^1} w(\rho) d\rho < \infty$.

\mathcal{W}_0 shall consist of the (classes of) bounded, measurable, L^∞ -valued and boundedly supported functions (i.e., $t_\rho \in L^\infty(\mathbb{T}^2)$, $\sup_\rho \|t_\rho\|_{L^\infty} < \infty$ and there exists $b > 0$ such that $t_\rho = 0$ for $\rho > b$; in analogy to [T1] Ch.IV, Def. 7.7 we do not require that the range of t is a.e. separable for the norm topology of L^∞).

$\overline{\overline{\mathcal{W}}}$ shall consist of the (classes of) Bochner-measurable functions $t = (t_\rho)$ with $t_\rho \in L^1(\mathbb{T}^2)$ and $\int_0^b \|t_\rho\|_{L^1} \rho d\rho < \infty$ for each $b > 0$.

Lemma 5. *We have $\mathcal{W}_0, \mathcal{B}_1^\oplus(\widehat{M(2)}, \nu) \subseteq \overline{\overline{\mathcal{W}}}$.*

$\mathcal{W}_0 \cap \mathcal{B}_1^\oplus$ is dense in \mathcal{B}_1^\oplus (for the \mathcal{B}_1^\oplus -norm).

If $w(\rho) \leq \rho$ for all $\rho > 0$, then $\mathcal{W}_0, \mathcal{B}_1^\oplus \subseteq \overline{\mathcal{W}}$, both are dense (for the $\overline{\mathcal{W}}$ -norm).

If $\frac{\rho}{w(\rho)}$ is bounded on $]0, b]$ for each $b > 0$, then $\overline{\mathcal{W}} \subseteq \overline{\overline{\mathcal{W}}}$.

Proof. Considering \mathcal{S}^1 as a subset of $L^1(\mathbb{T}^2, \frac{1}{(2\pi)^2} d(\psi, \psi_1))$ defines (using Lemma 4(a)) a contractive embedding of $\mathcal{B}_1^\oplus(\widehat{M(2)}, \nu)$ into $L^1(]0, \infty[, \nu, L^1(\mathbb{T}^2))$ and we will identify now \mathcal{B}_1^\oplus with this subspace of L^1 -valued functions. The other properties use standard estimates and approximation techniques (see also the comment after Lemma 4). \square

Lemma 6. *Let $f, g:]0, \infty[\rightarrow [0, \infty[$ be measurable functions such that $f(\rho) = 0$ for $\rho > b$. Then for all $\rho > 0$*

$$\iint_{D_\rho} f(\rho_1) g(\rho_2) K(\rho_1, \rho_2, \rho) \rho_1 \rho_2 d(\rho_1, \rho_2) \leq \sup_{\rho_1} f(\rho_1) \int_{\max(\rho-b, 0)}^{\rho+b} g(\rho_2) \rho_2 d\rho_2.$$

Proof. See Proposition 2 for the definition of D_ρ . This is a special case (called "straightforward" in [BH] p. 27) of convolvability in the Bessel-Kingman hypergroup. More explicitly, one uses that (for fixed ρ, ρ_2) $\rho_1 \mapsto K(\rho_1, \rho_2, \rho) \rho_1$ is a

probability distribution concentrated on $]|\rho - \rho_2|, \rho + \rho_2[$ and if f vanishes outside $]0, b]$, then the domain of integration in the left integral can be restricted to $\max(\rho - b, 0) < \rho_2 < \rho + b$. \square

Lemma 6 and Lemma 4(c) imply that if $s \in \mathcal{W}_0$, $t \in \overline{\mathcal{W}}$, then $(s \sharp t)_\rho$ can be defined for each $\rho > 0$ by an integral as in Cor. 3 to Prop. 2, giving a measurable function $s \sharp t :]0, \infty[\rightarrow L^1(\mathbb{T}^2)$. This extends the dual convolution of \mathcal{B}_1^\oplus .

Lemma 7. (a) For $s \in \mathcal{W}_0$, $t \in \overline{\mathcal{W}}$, we have $s \sharp t \in \overline{\mathcal{W}}$. If $s_\rho = 0$ for $\rho > b$ we have $\|(s \sharp t)_\rho\|_{L^1} \leq 2 \sup_{\rho_1} \|s_{\rho_1}\|_{L^\infty} \int_0^{\rho+b} \|t_{\rho_2}\|_{L^1} \rho_2 d\rho_2$ for each $\rho > 0$.

(b) For $s, t \in \mathcal{W}_0$, we have $s \sharp t \in \mathcal{W}_0$.

(c) For $s^{(1)}, s^{(2)} \in \mathcal{W}_0 \cap \mathcal{B}_1^\oplus$, $t \in \overline{\mathcal{W}}$, we have $(s^{(1)} \sharp s^{(2)}) \sharp t = s^{(1)} \sharp (s^{(2)} \sharp t)$.

(d) For $n \in \mathbb{Z}$, $t \in \overline{\mathcal{W}}$ we have $\mathbf{M}_n(t) \in \overline{\mathcal{W}}$, $\mathcal{W}_0, \overline{\mathcal{W}}$ are \mathbf{M}_n -invariant subspaces, $\|\mathbf{M}_n(t)\|_{\overline{\mathcal{W}}} = \|t\|_{\overline{\mathcal{W}}}$, for $s \in \mathcal{W}_0$ $s \sharp \mathbf{M}_n(t) = \mathbf{M}_n(s \sharp t)$.

Proof. (a) follows from Lemma 6 and Lemma 4(c). Similarly, Lemma 4(d) is used for (b). We know that (c) holds for $t \in \mathcal{B}_1^\oplus$. By (b) there exists $b > 0$ such that $s_\rho^{(1)}, s_\rho^{(2)}, (s^{(1)} \sharp s^{(2)})_\rho = 0$ for $\rho > b$. Consider on $\overline{\mathcal{W}}$ the seminorms $\|t\|_{b'} = \int_0^{b'} \|t_{\rho_2}\|_{L^1} \rho_2 d\rho_2$. If $\rho > 0$ is fixed, $b' \geq \rho + b$, it follows from (a) that $t \mapsto (s^{(2)} \sharp t)_\rho$ is continuous for $\|\cdot\|_{b'}$. Then for each $b' > 0$, we get that $t \mapsto s^{(2)} \sharp t$ is a continuous mapping $(\overline{\mathcal{W}}, \|\cdot\|_{b'+b}) \rightarrow (\overline{\mathcal{W}}, \|\cdot\|_{b'})$. As in Lemma 5, \mathcal{B}_1^\oplus is dense in $(\overline{\mathcal{W}}, \|\cdot\|_{b'})$ for each $b' > 0$. It follows by approximation that (c) holds pointwise (for each $\rho > 0$) for general $t \in \overline{\mathcal{W}}$.

The last statement of (d) follows from the Corollary to Lemma 3, the others are easy. \square

From now we assume that $w(\rho) \leq \rho$ for all $\rho > 0$ and that $\frac{\rho}{w(\rho)}$ is bounded on $]0, b]$ for each $b > 0$ (see Lemma 5). We define

$$\mathcal{W} = \{t \in \overline{\mathcal{W}} : s \sharp t \in \overline{\mathcal{W}} \text{ for all } s \in \mathcal{W}_0 \cap \mathcal{B}_1^\oplus \text{ and there exists } c \geq 0$$

$$\text{such that } \|s \sharp t\|_{\overline{\mathcal{W}}} \leq c \|s\|_{\mathcal{B}_1^\oplus} \text{ for all } s \in \mathcal{W}_0 \cap \mathcal{B}_1^\oplus\} \quad \text{and}$$

$$\|t\|_{\mathcal{W}} = \sup \{\|\alpha t + s \sharp t\|_{\overline{\mathcal{W}}} : \alpha \geq 0, s \in \mathcal{W}_0 \cap \mathcal{B}_1^\oplus, \alpha + \|s\|_{\mathcal{B}_1^\oplus} \leq 1\} \quad (\text{see also Remark (c) below}).$$

Lemma 8. Dual convolution \sharp extends to an action of $\mathcal{B}_1^\oplus(\widehat{M(2)})$ on \mathcal{W} . \mathcal{W} is a Banach \mathcal{B}_1^\oplus -module and \mathbf{M}_n -invariant, $\|\mathbf{M}_n(t)\|_{\mathcal{W}} = \|t\|_{\mathcal{W}}$ ($n \in \mathbb{Z}$, $t \in \mathcal{W}$).

Proof. It follows immediately from the definition that \mathcal{W} is a subspace of $\overline{\mathcal{W}}$, $\|\cdot\|_{\mathcal{W}}$ a norm, $\|t\|_{\overline{\mathcal{W}}} \leq \|t\|_{\mathcal{W}}$, $\|s \sharp t\|_{\overline{\mathcal{W}}} \leq \|s\|_{\mathcal{B}_1^\oplus} \|t\|_{\mathcal{W}}$ for $s \in \mathcal{W}_0 \cap \mathcal{B}_1^\oplus$, $t \in \mathcal{W}$. Let $(t^{(n)})$ be a sequence in \mathcal{W} with $\|t^{(n)}\|_{\mathcal{W}} \leq 1$ for all n and converging in $\overline{\mathcal{W}}$, $t^{(0)} =$

$\overline{\mathcal{W}}$ - $\lim_{n \rightarrow \infty} t^{(n)}$. Let $s \in \mathcal{W}_0 \cap \mathcal{B}_1^\oplus$, $b' > 0$ be fixed. It follows from Lemma 7(a) (use that $\frac{\rho}{w(\rho)}$ is bounded on bounded intervals) that there exists $c > 0$ (depending only on s, b', w) such that $\int_0^{b'} \|(s \sharp t)_\rho\|_{L^1} w(\rho) d\rho \leq c \|t\|_{\overline{\mathcal{W}}}$ for all $t \in \overline{\mathcal{W}}$. This gives $\int_0^{b'} \|(s \sharp (t^{(n)} - t^{(0)}))_\rho\|_{L^1} w(\rho) d\rho \rightarrow 0$ for $n \rightarrow \infty$, hence $\int_0^{b'} \|(s \sharp t^{(0)})_\rho\|_{L^1} w(\rho) d\rho \leq \|s\|_{\mathcal{B}_1^\oplus}$. It follows that $\|s \sharp t^{(0)}\|_{\overline{\mathcal{W}}} \leq \|s\|_{\mathcal{B}_1^\oplus}$ for all $s \in \mathcal{W}_0 \cap \mathcal{B}_1^\oplus$, thus $t^{(0)} \in \mathcal{W}$. If, in addition, $(t^{(n)})$ is a Cauchy sequence in \mathcal{W} , we get similarly that $t^{(n)} \rightarrow t^{(0)}$ in \mathcal{W} , hence \mathcal{W} is complete.

It follows easily from Lemma 7(c) that $s \sharp t \in \mathcal{W}$ for $s \in \mathcal{W}_0 \cap \mathcal{B}_1^\oplus$, $t \in \mathcal{W}$ and $\|s \sharp t\|_{\mathcal{W}} \leq \|s\|_{\mathcal{B}_1^\oplus} \|t\|_{\mathcal{W}}$. By density (Lemma 5) \sharp extends to $s \in \mathcal{B}_1^\oplus$ and \mathcal{W} becomes a Banach \mathcal{B}_1^\oplus -module. Concerning \mathbf{M}_n the properties follow from Lemma 7(d). \square

\mathcal{W} is the maximal \mathcal{B}_1^\oplus -submodule of $\overline{\mathcal{W}}$ (for \sharp). By Lemma 5, $\mathcal{B}_1^\oplus \subseteq \mathcal{W}$ but the abstract definition does not give more information about the size of \mathcal{W} (which depends also on w).

Proposition 3. For $w(\rho) = \frac{\rho}{1 + \rho^2}$, we have $L^1(]0, \infty[, d\rho, \mathcal{S}_1(\mathcal{H})) \subseteq \mathcal{W}$.

Since $\widehat{D} = (\pi_\rho(D))$ maps $\mathcal{B}_1^\oplus(\widehat{M(2)}, \nu)$ to $L^1(]0, \infty[, d\rho, \mathcal{S}_1(\mathcal{H}))$, we have found the desired \mathcal{B}_1^\oplus -module containing the image of the derivation. This will finish our proof that $A(M(2))$ is not weakly amenable.

Proof. By direct evaluation, $\int_{|\rho_1 - \rho_2|}^{\rho_1 + \rho_2} K(\rho_1, \rho_2, \rho) w(\rho) d\rho = \frac{1}{\pi} \int_0^\pi \frac{1}{1 + \rho^2} d\mu = \frac{1}{\sqrt{(1 + (\rho_1 - \rho_2)^2)(1 + (\rho_1 + \rho_2)^2)}} \leq \frac{1}{\rho_2}$. Since $\|(s \sharp t)_\rho\|_{L^1} \leq \iint_{D_\rho} \frac{1}{2} \|\theta(s_{\rho_1}, t_{\rho_2})(\rho)\|_{L^1} K(\rho_1, \rho_2, \rho) \rho_1 \rho_2 d(\rho_1, \rho_2) \leq \iint_{D_\rho} \|s_{\rho_1}\|_1 \|t_{\rho_2}\|_1 K(\rho_1, \rho_2, \rho) \rho_1 \rho_2 d(\rho_1, \rho_2)$, this gives $\int_0^\infty \|(s \sharp t)_\rho\|_{L^1} w(\rho) d\rho \leq \int_0^\infty \int_0^\infty \|s_{\rho_1}\|_1 \|t_{\rho_2}\|_1 \rho_1 d(\rho_1, \rho_2) \leq \|s\|_{\mathcal{B}_1^\oplus} \int_0^\infty \|t_{\rho_2}\|_1 d\rho_2$. \square

Remarks. (a) By additional estimates, one can show that more generally the conclusion of Proposition 3 holds for a (positive, measurable) weight function w on $]0, \infty[$, if w is bounded and $\int_0^\infty \frac{w(\rho)}{\rho} d\rho < \infty$. These conditions are necessary as well. For the extension of \sharp we needed that $\frac{\rho}{w(\rho)}$ is bounded on every bounded interval $]0, b]$, $b > 0$. If one wants that $\mathcal{B}_1^\oplus \subseteq \mathcal{W}$, then $\frac{w(\rho)}{\rho}$ should be bounded (on $]0, \infty[$) as well.

(b) Further derivations $\mathcal{B}_1^\oplus \rightarrow \mathcal{W}$ are obtained if the distribution d is defined using some other direction from the normal subgroup \mathbb{R}^2 . But, again this does not work if d involves a direction from the non-normal factor.

(c) We used the unitization of \mathcal{B}_1^\oplus to define the norm on \mathcal{W} . One can show (using an appropriate approximate unit for \mathcal{B}_1^\oplus) that it is also given by $\|t\|_{\mathcal{W}} = \sup \{\|s\sharp t\|_{\overline{\mathcal{W}}} : s \in \mathcal{W}_0 \cap \mathcal{B}_1^\oplus, \|s\|_{\mathcal{B}_1^\oplus} \leq 1\}$.

Example. We consider the trigonometric basis $\chi_n(\psi) = e^{in\psi}$ ($n \in \mathbb{Z}$) of \mathcal{H} . For $m, n \in \mathbb{Z}$ there are one dimensional operators $A_{(m,n)}h = (h|\chi_n)\chi_m$ ($h \in \mathcal{H}$). Then $A_{(m,n)} \in \mathcal{S}_1(\mathcal{H})$ with $k_{A_{(m,n)}}(\psi, \psi_1) = \chi_m(\psi) \overline{\chi_n(\psi_1)}$. For the corresponding coefficients $u(x) = \text{tr}(A_{(m,n)} \pi_\rho(x)^*)$ ($x \in M(2)$) one gets $u(\alpha x \alpha') = u(x) \chi_m(\alpha) \chi_n(\alpha')$ for $\alpha, \alpha' \in \mathbb{T}$ and $u(v) = e^{i(m-n)\phi} J_{n-m}(\rho r)$ for $v = r \begin{pmatrix} \cos \phi \\ \sin \phi \end{pmatrix} \in \mathbb{R}^2$ where J_{n-m} denotes the classical (J-) Bessel functions (see also [Va] p. 206 (9)). Then Lemma 3 implies $\theta_0(A_{(m_1, n_1)}, A_{(m_2, n_2)}, \rho_1, \rho_2)(\mu) = \overline{k_{A_{(m_1, n_1)}}(\varphi, \varphi)} k_{A_{(m_2, n_2)}}(\mu - \varphi, \mu - \varphi) A_{(m_1+m_2, n_1+n_2)}$ and $\theta(\dots)(\mu) = 2\Re(\overline{k_{A_{(m_1, n_1)}}(\varphi, \varphi)} k_{A_{(m_2, n_2)}}(\mu - \varphi, \mu - \varphi)) A_{(m_1+m_2, n_1+n_2)}$ (this is an example where $\theta(A_1, A_2)(\mu)$ is one dimensional). For $m_1 = m_2 = 0$, $n = n_1 + n_2$ one gets from Cor. 2 to Prop. 2 the multiplication formula

$$J_{n-n_2}(\rho_1 r) J_{n_2}(\rho_2 r) = \frac{2}{\pi} \int_{|\rho_1 - \rho_2|}^{\rho_1 + \rho_2} \frac{\cos(n\varphi - n_2\mu) \rho J_n(\rho r)}{\sqrt{4\rho_1^2 \rho_2^2 - (\rho^2 - \rho_1^2 - \rho_2^2)^2}} d\rho$$

for $r, \rho_1, \rho_2 > 0$, $n, n_2 \in \mathbb{Z}$ with $\mu = \arccos(\frac{\rho^2 - \rho_1^2 - \rho_2^2}{2\rho_1 \rho_2})$, $\varphi = \arccos(\frac{\rho_1 + \rho_2 \cos \mu}{\rho})$ (taking the principal branches of \arccos , i.e. $\mu, \varphi \in [0, \pi]$; in fact the formula extends to $r \in \mathbb{R}$). Observe that there is an error in [Va], the corresponding formula (3) in [Va] p. 210 is not correct for all m, n .

For $m_j = n_j = 0$ (this corresponds to the “radial” part of $A(M(2))$, $B(M(2))$, i.e. the functions that are both left and right \mathbb{T} -periodic) one gets the multiplication of the Bessel-Kingman hypergroup, [BH] p. 235 with $\alpha = 0$.

8. THE GROUPS $M(2)_{\mathcal{K}}$ AND $\widetilde{M(2)}$

$\widetilde{M(2)}$ is covered by [LLSS] Th. 1.6 (using spectral synthesis). $M(2)_{\mathcal{K}}$ was the main open case for our Theorem (see [LLSS] Ex. 3.4).

We take up the notations of Section 2, $M(2)_{\mathcal{K}} = \mathbb{R}^2 \rtimes \mathcal{K}$, \mathcal{K} is a non-trivial compact group and we have given a continuous homomorphism $\varphi_0: \mathbb{R} \rightarrow \mathcal{K}$ with dense image, \mathcal{K} is written additively and there exists a proper closed subgroup \mathcal{K}_0 of \mathcal{K} such that $\varphi_0(2\pi\mathbb{Z}) = \mathcal{K}_0 \cap \varphi_0(\mathbb{R})$. Then $\mathcal{K}/\mathcal{K}_0 \cong \mathbb{T} (= \mathbb{R}/2\pi\mathbb{Z})$ giving a continuous surjective homomorphism $\alpha_{\mathcal{K}}: \mathcal{K} \rightarrow \mathbb{T}$ with $\ker \alpha_{\mathcal{K}} = \mathcal{K}_0$, $\alpha_{\mathcal{K}} \circ \varphi_0$ is the canonical projection $\mathbb{R} \rightarrow \mathbb{T}$. The action of \mathcal{K} on \mathbb{R}^2 is given by $k \circ v = A_{\alpha_{\mathcal{K}}(k)} v$ ($k \in \mathcal{K}$, $v \in \mathbb{R}^2$), using the rotations A_α from Section 7. Then $\alpha_{\mathcal{K}}$ extends to a continuous surjective homomorphism $M(2)_{\mathcal{K}} \rightarrow M(2)$, again denoted by $\alpha_{\mathcal{K}}$, with $\ker \alpha_{\mathcal{K}} = \mathcal{K}_0$. For

Haar measure we take $\frac{1}{2\pi} dv dk$ (with normalized Haar measure on \mathcal{K}). $\mathbb{H}_{\mathcal{K}}$ is unimodular. As in Section 6, the dual group $\Gamma = \widehat{\mathcal{K}}$ is considered as a subgroup of \mathbb{R} in such a way that for $t \in \Gamma$ the corresponding character χ_t of \mathcal{K} satisfies $\chi_t(\varphi_0(z)) = e^{itz}$ for $z \in \mathbb{R}$ (see Rem. (b) in Section 2 and Rem. (c) in Appendix A). Under this identification $\mathcal{K}_0^\perp = \mathbb{Z}$ (hence $\mathbb{Z} \subseteq \Gamma$). Conversely, for any subgroup Γ of \mathbb{R} with $\Gamma \supset \mathbb{Z}$, $\mathcal{K} = \widehat{\Gamma}$ gives a compact group satisfying the conditions above (with φ_0 the dual of the inclusion $\Gamma \rightarrow \mathbb{R}$, i.e. restricting the characters of \mathbb{R} to Γ ; $\mathcal{K}_0 = \mathbb{Z}^\perp$; $\alpha_{\mathcal{K}}$: restricting the characters of Γ to \mathbb{Z}).

$\widetilde{M(2)} = \mathbb{R}^2 \rtimes \mathbb{R}$ (universal covering group of $M(2)$). For Haar measure we take $\frac{1}{(2\pi)^2} dv d\alpha$ (now $\alpha \in \mathbb{R}$). $\widetilde{M(2)}$ is unimodular. $\varphi_0: \mathbb{R} \rightarrow \mathcal{K}$ extends to a continuous homomorphism $\varphi: \widetilde{M(2)} \rightarrow M(2)_{\mathcal{K}}$ with dense image. In particular for $\mathcal{K} = \mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$ (φ_0 the quotient mapping), $\varphi_1: \widetilde{M(2)} \rightarrow M(2)$ gives the covering homomorphism, for general \mathcal{K} $\varphi_1 = \alpha_{\mathcal{K}} \circ \varphi$. The representations π_ρ of $M(2)$ (Section 7) lift to $\widetilde{M(2)}$ and $M(2)_{\mathcal{K}}$, we denote these again by π_ρ ($\rho > 0$). \mathbb{R}^2 is the commutator subgroup of $\widetilde{M(2)}$, hence the one dimensional representations of $\widetilde{M(2)}$ are given by the characters $\{\chi_t: t \in \mathbb{R}\}$ of $\mathbb{R} \cong \widetilde{M(2)}/\mathbb{R}^2$. $2\pi\mathbb{Z} (\subseteq \mathbb{R})$ is the centre of $\widetilde{M(2)}$. If π is an irreducible representation of $\widetilde{M(2)}$ its restriction to $2\pi\mathbb{Z}$ is given by a character which extends to a character χ_t of \mathbb{R} (for some $t \in \mathbb{R}$). Then $\overline{\chi_t} \otimes \pi$ is trivial on $2\pi\mathbb{Z}$, hence it induces an irreducible representation of $M(2) \cong \widetilde{M(2)}/(2\pi\mathbb{Z})$. If π is infinite dimensional, it follows that $\overline{\chi_t} \otimes \pi \simeq \pi_\rho$, hence $\pi \simeq \chi_t \otimes \pi_\rho$ for some $\rho > 0$. Put $\mathcal{R} = [0, 1[$. χ_t and χ_{t+n} ($n \in \mathbb{Z}$) coincide on $2\pi\mathbb{Z}$, hence we may choose $t \in \mathcal{R}$. We put $\boldsymbol{\rho} = (t, \rho)$ and use the notation $\pi_{\boldsymbol{\rho}} = \chi_t \otimes \pi_\rho$ (considered again as a representation on \mathcal{H} , see the proof of Proposition 2). It is not hard to see that for $\boldsymbol{\rho} \neq \boldsymbol{\rho}' \in \mathcal{R} \times]0, \infty[$ the representations $\pi_{\boldsymbol{\rho}}, \pi_{\boldsymbol{\rho}'}$ are not equivalent. Thus we get

$$\left(\widetilde{M(2)}\right)^\wedge = \{\chi_t: t \in \mathbb{R}\} \cup \{\chi_t \otimes \pi_\rho: 0 \leq t < 1, \rho > 0\}$$

(as a complete set of representatives; one might get this also from Mackey's theory). One can show that $\{\pi_{\boldsymbol{\rho}}\}$ is open and (taking on \mathcal{R} the quotient topology from \mathbb{R}/\mathbb{Z}) it carries the product topology of $\mathcal{R} \times]0, \infty[$. On $\{\chi_t\}$ one gets the standard topology of \mathbb{R} , $\overline{\{\chi_t \otimes \pi_\rho\}} = \{\chi_t \otimes \pi_\rho\} \cup \{\chi_{t+n}: n \in \mathbb{Z}\}$ for each $\rho > 0, t \in \mathcal{R}$.

For $M(2)_{\mathcal{K}}$ the centre is \mathcal{K}_0 . Let \mathcal{R}_Γ be a set of representatives for the cosets in Γ/\mathbb{Z} (e.g. $\mathcal{R}_\Gamma = \Gamma \cap [0, 1[$). Then a similar argument as above gives

$$(M(2)_{\mathcal{K}})^\wedge = \{\chi_t: t \in \Gamma\} \cup \{\chi_t \otimes \pi_\rho: t \in \mathcal{R}_\Gamma, \rho > 0\}$$

On the open subset $\{\pi_\rho\}$ one has the product topology of $\mathcal{R}_\Gamma \times]0, \infty[$ (taking the discrete topology on \mathcal{R}_Γ) and on $\{\chi_t\}$ the discrete topology. The closure $\overline{\{\chi_t \otimes \pi_\rho\}}$ is as above.

Lemma 9. (a) *Using Haar measure and parametrization as above, the Plancherel measure ν of $\widetilde{M(2)}$ is concentrated on $\{\chi_t \otimes \pi_\rho : 0 \leq t < 1, \rho > 0\}$. It is the product of standard Lebesgue measure on $[0, 1[$ and $d\nu = \rho d\rho$ on $]0, \infty[$.*

(b) *Plancherel measure ν of $M(2)_\mathcal{K}$ is concentrated on $\{\chi_t \otimes \pi_\rho : t \in \mathcal{R}_\Gamma, \rho > 0\}$. It is the product of counting measure on \mathcal{R}_Γ and $d\nu = \rho d\rho$ on $]0, \infty[$.*

If \mathcal{K} is not metrizable (equivalently, if Γ is uncountable) (b) gives again an example of the Plancherel theorem for a group that is non-compact, σ -compact, but not second countable (see also Rem. (d) in Appendix B).

Proof. Again, this follows by direct computations (compare Lemma 2), using the Plancherel theorem for \mathbb{R}^2, \mathbb{R} and \mathcal{K} . \square

The distribution d and the corresponding derivation D can be defined as in Section 7. $\widetilde{M(2)}$ is a Lie group, d and D are defined on $C^1(\widetilde{M(2)})$. For $M(2)_\mathcal{K}$ we can use approximation by Lie groups as in Section 6, d and D are defined on $\bigcup\{C^1(M(2)_{\mathcal{K}/K_1})\}$ ranging over the closed subgroups K_1 of \mathcal{K}_0 such that \mathcal{K}_0/K_1 is Lie (equivalently, K_1^\perp is finitely generated and $K_1^\perp \supseteq \mathbb{Z}$). We have $\chi_t(d) = 0$ and for $\rho = (t, \rho)$ this gives $\pi_\rho(d) = \pi_\rho(d)$ where again $(\pi_\rho(d)g)(\psi) = -i\rho \cos(\psi)g(\psi)$ (for $g \in \mathcal{H} = L^2(\mathbb{R}, \frac{1}{2\pi}d\psi)$). In both cases D does *not* produce a bounded mapping $A \rightarrow VN$. We will next investigate the dual convolution and then use the construction for $M(2)$ to find a suitable \mathcal{B}_1^\oplus -module \mathcal{W} .

First, we consider now $M(2)_\mathcal{K}$. We assume that $\mathcal{R}_\Gamma = \Gamma \cap [0, 1[$. We start with the decomposition of tensor products making a similar procedure as in Section 7. For $t \in \mathbb{R}$, $[t]$ denotes the integer part, $\{t\}$ the fractional part. For $\rho_j = (t_j, \rho_j) \in \mathcal{R}_\Gamma \times]0, \infty[$ ($j = 1, 2$) we define $(W_{\rho_1, \rho_2} f)(\mu) = M_{-[t_1+t_2]} \circ ((W_{\rho_1, \rho_2} f)(\mu))$ ($W_{\rho_1, \rho_2} : \mathcal{H} \otimes_2 \mathcal{H} \rightarrow L^2(\mathbb{T}, \frac{1}{2\pi}d\mu, \mathcal{H})$). With the notations made before Proposition 2 we get

Proposition 4. *For $\rho_j = (t_j, \rho_j) \in \mathcal{R}_\Gamma \times]0, \infty[$ ($j = 1, 2$), $x \in M(2)_\mathcal{K}$, we have*

$$W_{\rho_1, \rho_2} \circ (\pi_{\rho_1} \otimes \pi_{\rho_2})(x) \circ W_{\rho_1, \rho_2}^* = \int_{\mathbb{T}}^{\oplus} \pi_{(\{t_1+t_2\}, \rho)}(x) d\mu.$$

Proof. Remembering our identifications, we have $(\pi_{\rho_1} \otimes \pi_{\rho_2})(x) = \chi_{t_1+t_2}(x)$ ($\pi_{\rho_1} \otimes \pi_{\rho_2})(x)$. Using $t = [t] + \{t\}$, the result can be obtained from Proposition 2. \square

We put $\theta_0(A_1, A_2, \rho_1, \rho_2) = \text{pr}_{\mathcal{S}}(W_{\rho_1, \rho_2} \circ (A_1 \otimes A_2) \circ W_{\rho_1, \rho_2}^*)$ and $\theta(A_1, A_2)(\rho) = \theta_0(A_1, A_2)(\mu) + \theta_0(A_1, A_2)(2\pi - \mu)$ ($A_1, A_2 \in \mathcal{S}^1(\mathcal{H})$).

Corollary 1. *For $A_1, A_2 \in \mathcal{S}^1(\mathcal{H})$, $\rho_j = (t_j, \rho_j) \in \mathcal{R}_{\Gamma} \times]0, \infty[$ consider coefficients $u_j(x) = \text{tr}(A_j \pi_{\rho_j}(x)^*)$ ($j = 1, 2, x \in M(2)_{\mathcal{K}}$). Then $\theta(A_1, A_2, \rho_1, \rho_2) = \mathbf{M}_{-[t_1+t_2]}(\theta(A_1, A_2, \rho_1, \rho_2)) \in L^1([|\rho_1 - \rho_2|, \rho_1 + \rho_2], \frac{1}{2}K(\rho_1, \rho_2, \rho) \rho d\rho, \mathcal{S}^1(\mathcal{H}))$ and*

$$u_1(x) u_2(x) = \int_{|\rho_1 - \rho_2|}^{\rho_1 + \rho_2} \frac{1}{2} \text{tr}(\pi_{(\{t_1+t_2\}, \rho)}(x)^* \theta(A_1, A_2)(\rho)) K(\rho_1, \rho_2, \rho) d\nu(\rho).$$

In particular, $\pi_{\rho_1} \otimes \pi_{\rho_2} \simeq 2 \int_{|\rho_1 - \rho_2|}^{\rho_1 + \rho_2} \pi_{(\{t_1+t_2\}, \rho)} d\rho$. Similarly, there is an analogue of Corollary 1 to Proposition 2 in terms of θ_0 .

Proof. An expression for θ_0 and then for θ can be obtained as in Lemma 3. For the product formula compare the Corollaries 1,2 to Proposition 2, using now Proposition 4. \square

For an operator field $T = (T_{\rho}) \in \mathcal{B}_1^{\oplus}((M(2)_{\mathcal{K}})^{\wedge}, \nu)$ and $t \in \mathcal{R}_{\Gamma}$ we put $T^{(t)} = (T_{(t, \rho)})_{\rho > 0}$. Then $T^{(t)} \in \mathcal{B}_1^{\oplus}(\widehat{M(2)}, \nu)$ and $\|T\|_{\mathcal{B}_1^{\oplus}} = \sum_{t \in \mathcal{R}_{\Gamma}} \|T^{(t)}\|_{\mathcal{B}_1^{\oplus}}$. Thus we get an isomorphism $\mathcal{B}_1^{\oplus}((M(2)_{\mathcal{K}})^{\wedge}, \nu) \cong l^1(\mathcal{R}_{\Gamma}, \mathcal{B}_1^{\oplus}(\widehat{M(2)}, \nu))$.

Corollary 2. *For $S, T \in \mathcal{B}_1^{\oplus}((M(2)_{\mathcal{K}})^{\wedge}, \nu)$, $t \in \mathcal{R}_{\Gamma}$, we have*

$$(S \sharp T)^{(t)} = \sum_{t_1 \in \mathcal{R}_{\Gamma}} \mathbf{M}_{[t-t_1]}(S^{(t_1)} \sharp T^{(\{t-t_1\})})$$

Thus one gets a sort of twisted convolution.

Proof. This follows from Corollary 3 to Proposition 2 and Corollary 1 above. \square

Let $\mathcal{W}(M(2))$ be the Banach module defined before Lemma 8 with w as in Proposition 3 (see also Remark (a) in Section 7). We consider now

$$\mathcal{W} = l^1(\mathcal{R}_{\Gamma}, \mathcal{W}(M(2))) = \{(w^{(t)}) : w^{(t)} \in \mathcal{W}(M(2)), \sum_{t \in \mathcal{R}_{\Gamma}} \|w^{(t)}\|_{\mathcal{W}} < \infty\}.$$

Proposition 5. *We have $\mathcal{B}_1^{\oplus}((M(2)_{\mathcal{K}})^{\wedge}, \nu) \subseteq \mathcal{W}$ and $l^1(\mathcal{R}_{\Gamma}, L^1(]0, \infty[, d\rho, \mathcal{S}_1(\mathcal{H}))) \subseteq \mathcal{W}$. Dual convolution extends to an action of \mathcal{B}_1^{\oplus} on \mathcal{W} . \mathcal{W} is a Banach \mathcal{B}_1^{\oplus} -module.*

This finishes our proof that $A(M(2)_{\mathcal{K}})$ is not weakly amenable.

Proof. This follows from the properties of $\mathcal{W}(M(2))$ obtained in Lemma 8, Proposition 3 and Lemma 5, and using Corollary 2 above and Lemma 7 (d) to get the extension of \sharp . \square

The construction works quite similarly for $\widetilde{M(2)}$. With $\mathcal{R} = [0, 1[$, Proposition 4 holds in the same way for $\rho_j \in \mathcal{R} \times]0, \infty[$, $x \in \widetilde{M(2)}$ and also Corollary 1. As above, $\mathcal{B}_1^\oplus(\widetilde{M(2)}^\wedge, \nu) \cong L^1([0, 1[, dt, \mathcal{B}_1^\oplus(\widetilde{M(2)}, \nu))$ by transforming operator fields. Similarly, in Corollary 2 one gets $(S \sharp T)^{(t)} = \int_{\mathcal{R}} \mathbf{M}_{[t-t_1]}(S^{(t_1)} \sharp T^{\{t-t_1\}}) dt_1$. Finally, one can take $\mathcal{W} = L^1(\mathcal{R}, dt, \mathcal{W}(M(2)))$.

Remarks. The formulas given above depend on the choice of \mathcal{R}_Γ . If $\mathcal{R}_\Gamma (\subseteq \Gamma)$ is an arbitrary set of representatives for Γ/\mathbb{Z} there is a section $\sigma : \Gamma \rightarrow \mathcal{R}_\Gamma$ and a corresponding cocycle $n(t_1, t_2) = \sigma(t_1) + \sigma(t_2) - \sigma(t_1 + t_2) (\in \mathbb{Z} \text{ for } t_1, t_2 \in \Gamma)$. Then in the definition of W_{ρ_1, ρ_2} one has to replace $[t_1 + t_2]$ by $n(t_1, t_2)$ and in Proposition 4 replace $\{t_1 + t_2\}$ by $\sigma(t_1 + t_2)$, the same in Corollary 1. In Corollary 2 one replaces $\{t - t_1\}$ by $\sigma(t - t_1)$ and $[t - t_1]$ by $-n(t_1, t - t_1)$.

9. THE GRÉLAUD GROUPS \mathbb{G}_a

They were treated in [LLSS] sec. 2.3, using spectral synthesis. We present also (slightly more briefly) our method for this case.

We use the complex presentation $\mathbb{G}_a = \mathbb{C} \rtimes \mathbb{R}$ with the action $t \circ z = e^{wt}z$ ($t \in \mathbb{R}$, $z \in \mathbb{C}$), putting $w = a + i$ for $a \neq 0$. It is not hard to see that $\mathbb{G}_{-a} \cong \mathbb{G}_a$ hence we will always assume that $a > 0$ (similarly for the version used in [FL] 4.4, $\mathbb{G}_{-a} \cong G_\theta$ with $\theta = -\frac{1}{a}$). Using the homeomorphism with $\mathbb{C} \times \mathbb{R}$ given by $(z, t) \mapsto zt$ we take for (left) Haar measure $e^{-2at} dz dt$. \mathbb{G}_a is not unimodular, $\Delta(zt) = e^{-2at}$.

The irreducible representations can again be obtained by Mackey's theory. We take here $\mathcal{H} = L^2(\mathbb{R})$ (as in Section 5). For real $b \neq 0$ an irreducible representation π_b is defined for $f \in \mathcal{H}$ by $(\pi_b(z)f)(\psi) = \exp(ib \Re(e^{-w\psi}z))f(\psi)$, $(\pi_b(t)f)(\psi) = f(\psi - t)$ (\Re denoting the real part; in the notation of [FL] this corresponds to π_μ with $\mu = -b$). For $t = 2\pi \in \mathbb{R}$, $\pi_b(t)\pi_b(\mathbf{x})\pi_b(-t) = \pi_{be^{2\pi a}}(\mathbf{x})$ ($\mathbf{x} \in \mathbb{G}_a$) gives $\pi_b \simeq \pi_{be^{2\pi a}}$ and for $t = \pi$, $\pi_b(t)\pi_b(\mathbf{x})\pi_b(-t) = \pi_{-be^{\pi a}}(\mathbf{x})$ implies $\pi_b \simeq \pi_{-be^{\pi a}}$. It follows that one can restrict to $b \in [1, e^{2\pi a}[$ and in this domain one can show that the representations are pairwise non-equivalent, they exhaust (up to equivalence) all the infinite dimensional irreducible representations of \mathbb{G}_a . $\{\pi_b : b \in [1, e^{2\pi a}[$ defines an open, compact (but non-closed) subset of $\widehat{\mathbb{G}_a}$ (again, the complement consists of the one dimensional representations, they are given by $\widehat{\mathbb{R}}$). It carries the quotient topology of $[1, e^{2\pi a}]$, obtained by identifying the end points. Formulas for closures are as in Section 7.

\mathbb{G}_a is type I, Plancherel measure is concentrated on $\{\pi_b : b \in [1, e^{2\pi a}[$. Put $(Kf)(\psi) = e^{-2a\psi}f(\psi)$ (densely defined on $\{f \in \mathcal{H} : Kf \in \mathcal{H}\}$). K is positive

definite, self-adjoint and satisfies the semi-invariance relations for all π_b . Hence it gives Duflo–Moore operators (independent of b , see Appendix B). The corresponding Plancherel measure is given by $d\nu = \frac{1}{(2\pi)^2} b db$ (that can again be verified by direct calculation). Thus $\mathcal{F}_A(f) = ([\pi_b(f)K])$ ($[\cdot]$ denoting now the closure of the operator), $\mathcal{B}_1^\oplus(\widehat{\mathbb{G}}_a, \nu) \cong L^1([1, e^{2\pi a}[, \frac{1}{(2\pi)^2} b db, \mathcal{S}_1(\mathcal{H}))$, general elements of $\mathcal{B}_1^\oplus(\widehat{\mathbb{G}}_a, \nu)$ will be denoted as $T = (T_b)_{1 \leq b < e^{2\pi a}}$.

As before, we consider distributions defined using directional derivatives from the normal subgroup \mathbb{C} . For example, writing $z = x + iy$, put $d = \lim_{x \rightarrow 0} \frac{1}{x}(\delta_{-x} - \delta_0)$ and $D(f) = f \star d$, defined on $C^1(\mathbb{G}_a)$. Put $\widehat{D}_b = \pi_b(d)$ and for $T = (T_b)$, $\widehat{D}(T) = ([T_b \widehat{D}_b])$. Then $(\widehat{D}_b g)(\psi) = -ibe^{-a\psi} \cos(\psi) g(\psi)$ (for appropriate $g \in \mathcal{H}$) and $\mathcal{F}_A(D(f)) = \widehat{D}(\mathcal{F}_A(f))$ for $f \in A(\mathbb{G}_a) \cap C^1(\mathbb{G}_a)$ with $D(f) \in A(\mathbb{G}_a)$. One can check that $\{([T_b \widehat{D}_b K^{-1}]) : T = (T_b) \in \mathcal{B}_1^\oplus\}$ is not contained in $L^\infty(\widehat{\mathbb{G}}_a, \nu, \mathcal{B}(\mathcal{H}))$, hence D does not produce a bounded mapping $A(\mathbb{G}_a) \rightarrow VN(\mathbb{G}_a)$.

For the decomposition of tensor products, we proceed similarly as in Section 7. $\mathcal{H} \otimes_2 \mathcal{H}$ is identified with $L^2(\mathbb{R}^2)$. For $f \in \mathcal{H} \otimes_2 \mathcal{H}$ define $(Wf)(\mu)(\psi) = f(\psi, \psi + \mu)$ giving an isometric isomorphism $W : \mathcal{H} \otimes \mathcal{H} \rightarrow L^2(\mathbb{R}, \mathcal{H})$. For $\varphi \in \mathbb{R}$, $g \in \mathcal{H}$, we define $(U_\varphi g)(\psi) = g(\psi - \varphi)$, giving a unitary operator on \mathcal{H} , depending strongly continuously on φ . Fix now $b_1, b_2 \neq 0$ and for $\mu \in \mathbb{R}$ put $b_1 + b_2 e^{-w\mu} = r e^{-i\varphi}$ (polar coordinates) and $b = r e^{a\varphi}$, defining functions $r = r(b_1, b_2, \mu)$, $\varphi = \varphi(b_1, b_2, \mu)$ (see below) and $b = b(b_1, b_2, \mu)$. Finally, we define $(W_{b_1, b_2} f)(\mu) = U_\varphi((Wf)(\mu))$ ($f \in \mathcal{H} \otimes_2 \mathcal{H}$), giving again an isometric isomorphism $W_{b_1, b_2} : \mathcal{H} \otimes_2 \mathcal{H} \rightarrow L^2(\mathbb{R}, \mathcal{H})$, $(b_1, b_2) \mapsto W_{b_1, b_2}$ is strongly continuous. As an analogue of Proposition 2 one gets $W_{b_1, b_2} \circ (\pi_{b_1} \otimes \pi_{b_2})(\mathbf{x}) \circ W_{b_1, b_2}^* = \int_{\mathbb{R}}^\oplus \pi_b(\mathbf{x}) d\mu$ for $\mathbf{x} \in \mathbb{G}_a$. $\text{pr}_{\mathcal{S}} : \mathcal{S}^1(L^2(\mathbb{R}, \mathcal{H})) \rightarrow L^1(\mathbb{R}, \mathcal{S}^1(\mathcal{H}))$ shall be dual to the embedding of the subalgebra $L^\infty(\mathbb{R}) \overline{\otimes} \mathcal{B}(\mathcal{H})$ into $\mathcal{B}(L^2(\mathbb{R}, \mathcal{H}))$ and for $A_1, A_2 \in \mathcal{S}^1(\mathcal{H})$ we define again $\theta_0(A_1, A_2) = \text{pr}_{\mathcal{S}}(W_{b_1, b_2} \circ (A_1 \otimes A_2) \circ W_{b_1, b_2}^*) \in L^1(\mathbb{R}, \mathcal{S}^1(\mathcal{H}))$. Then for $u_j(\mathbf{x}) = \text{tr}(A_j \pi_{b_j}(\mathbf{x})^*)$ ($j = 1, 2$, $\mathbf{x} \in \mathbb{G}_a$) it results as in Corollary 1 to Proposition 2 that $u_1(\mathbf{x}) u_2(\mathbf{x}) = \int_{\mathbb{R}} \text{tr}(\pi_b(\mathbf{x})^* \theta_0(A_1, A_2)(\mu)) d\mu$.

The definition of the function $b(\mu)$ and the operators W_{b_1, b_2}, θ_0 depends on the choice of φ . If φ is replaced by $\varphi + 2\pi$, then b transforms to $b e^{2\pi a}$ and similarly for W_{b_1, b_2}, θ_0 . The expression for $u_1(\mathbf{x}) u_2(\mathbf{x})$ above is valid for any measurable choice of $\varphi(\mu)$. We will now assume that φ is chosen so that $b \in [1, e^{2\pi a}[$ holds. This determines φ uniquely for all μ where $r \neq 0$. For fixed b_1, b_2 , $r(b_1, b_2, \mu) = 0$ can happen just for one μ (see Remark (b)). We define now $\theta(A_1, A_2)(b) = \frac{(2\pi)^2}{b} \sum_{b(\mu)=b} \theta_0(A_1, A_2)(\mu) |\frac{db}{d\mu}(\mu)|^{-1}$. This gives $u_1(\mathbf{x}) u_2(\mathbf{x}) =$

$\int_1^{e^{2\pi a}} \text{tr}(\pi_b(\mathbf{x})^* \theta(A_1, A_2)(b)) d\nu(b)$. Finally, it follows as in Corollary 3 of Proposition 2 that for $S = (S_b), T = (T_b) \in \mathcal{B}_1^\oplus(\widehat{\mathbb{G}_a}, \nu)$, we have

$$(S \sharp T)_b = \iint \theta(S_{b_1}, T_{b_2})(b) d\nu(b_1) d\nu(b_2) \quad \text{a.e.}$$

As in Lemma 3 there is an expression in terms of integration kernels, for $\mu \in \mathbb{R}$ one has $k_{\theta_0(A_1, A_2)(\mu)}(\psi, \psi_1) = k_{A_1}(\psi - \varphi, \psi_1 - \varphi) k_{A_2}(\psi - \varphi + \mu, \psi_1 - \varphi + \mu)$. It allows extensions of θ_0, θ and \sharp . For a Borel measurable function $a_0: \mathbb{R}^2 \rightarrow \mathbb{C}$ and $\xi_1, \xi_2 \in \mathbb{R}$ we put $\|a_0\|_{L^1, \xi_1, \xi_2} = \int_{\xi_1}^{\xi_1+1} \int_0^\infty |a_0(\xi + t, \xi_2 + t)| dt d\xi$. We consider now measurable fields $t = (t_b)_{1 \leq b \leq e^{2\pi a}}$ of functions, i.e. $(b, \psi, \psi_1) \rightarrow t_b(\psi, \psi_1)$ shall be Borel measurable and use $\overline{\mathcal{W}} = \{t : \int \|t_b\|_{L^1, \xi_1, \xi_2} db < \infty \text{ for all } \xi_1, \xi_2 \in \mathbb{R}\}$ (identifying fields which coincide a.e.), $\mathcal{W}_0 = \{t : \sup_b \|t_b\|_\infty < \infty \text{ and there exists a compact set } C \subseteq \mathbb{R}^2 \text{ such that for all } b \text{ } t_b(\psi, \psi_1) = 0 \text{ holds when } (\psi, \psi_1) \notin C\}$. Then, using the formula above, $s \sharp t$ can be defined for $s \in \mathcal{W}_0, t \in \overline{\mathcal{W}}$. The case $b_2 = e^{\pm \pi a} b_1$ causes some technical complications, it is not always true that $s_1 \sharp s_2 \in \mathcal{W}_0$ for $s_1, s_2 \in \mathcal{W}_0$.

Finally, we consider $\overline{\mathcal{W}} = \{t : \sup_{\xi_1, \xi_2} \frac{1}{1+e^{-\xi_2}} \int \|t_b\|_{L^1, \xi_1, \xi_2} db < \infty\}$. Clearly $\overline{\mathcal{W}} \subseteq \overline{\mathcal{W}}$ and one can define now \mathcal{W} as in Section 7. Similarly as in Lemma 8, dual convolution \sharp extends to an action of $\mathcal{B}_1^\oplus(\widehat{\mathbb{G}_a})$ on \mathcal{W} , making \mathcal{W} a Banach \mathcal{B}_1^\oplus -module. One can show that $\{T : \int \| [T_b K^{\frac{1}{2}}] \|_1 d\nu(b) < \infty\} \subseteq \mathcal{W}$ hence $\widehat{\mathcal{D}}(\mathcal{B}_1^\oplus) \subseteq \mathcal{W}$ finishing our proof for $A(\mathbb{G}_a)$.

Remarks. (a) It follows from the arguments above that $\pi_{b_1} \otimes \pi_{b_2} \simeq \int_1^{e^{2\pi a}} n(b) \pi_b db$ for some multiplicities $n(b, b_1, b_2) \in \mathbb{N}_0 \cup \{\infty\}$. The behaviour of $n(b)$ depends on the relative sizes of a, b_1, b_2 . For $b_2 \neq e^{\pm \pi a} b_1, 1 \leq b_1, b_2 \leq e^{2\pi a}$ one has that $n(b)$ is finite for all $b \neq b_1, b_2$, but $n(b) \rightarrow \infty$ when $b \rightarrow b_1$ or $b \rightarrow b_2$ (and $n(b)$ is bounded on every closed interval not containing b_1, b_2). In the exceptional cases $b_2 = e^{\pm \pi a} b_1$ one gets $n(b) = \infty$ for all b .

(b) Geometrically, $\beta(\mu) = b_1 + b_2 e^{-w\mu}$ describes a logarithmic spiral. For $1 \leq b_1, b_2 \leq e^{2\pi a}, \beta(\mu) = 0$ (equivalently $r(\mu) = 0$) happens only for $b_2 = e^{\pm \pi a} b_1$, and for $b_2 = e^{\pi a} b_1$ it happens only for $\mu = \pi$. With some computations one can show that $\frac{db}{d\mu} = -e^{a(\varphi-\mu)} b_1 b_2 (a^2 + 1) \frac{\sin(\mu)}{r}$ at every continuity point of φ . It follows that $b(\mu)$ is monotonic on every subinterval of $[k\pi, (k+1)\pi]$ where φ is chosen as a continuous function of μ ($k \in \mathbb{Z}$). This justifies also our definition of θ above (in particular that $\theta(A_1, A_2)(b) \in \mathcal{B}_1^\oplus(\mathbb{G}_a)$), based on the coordinate transformation $b = b(\mu)$.

(c) Our representations π_b are basically those of [FL] (more precisely, $\pi_b \simeq \pi_\mu$ with π_μ as in [FL] p.108, $\mu = -b$). Another version is given by $(\pi'_\alpha(z)f)(\psi) =$

$\exp(i \Re(e^{-w\psi} z \alpha)) f(\psi)$, $(\pi'_\alpha(t)f)(\psi) = f(\psi - t)$ where $\alpha \in \mathbb{C}$, $|\alpha| = 1$ and as before $f \in \mathcal{H}$. Then $\pi'_\alpha \simeq \pi_b$ with $\alpha = b^{-\frac{i}{a}}$, $b > 0$. The family $\{\pi'_\alpha : |\alpha| = 1\}$ better reflects the periodicity in the infinite dimensional representations of \mathbb{G}_a .

APPENDIX A. CONNECTED DENSE SUBGROUPS

We elaborate (and slightly extend) results of Malcev, Goto, Poguntke.

Theorem A. *Let G be a locally compact group, H a connected Lie group, $\varphi: H \rightarrow G$ a continuous homomorphism such that $\ker \varphi$ is discrete. Then there exists a closed vector subgroup V of H (i.e. $V \cong \mathbb{R}^n$ for some $n \geq 0$) such that $\overline{\varphi(H)} = \varphi(H) \overline{\varphi(V)}$, $\varphi(V)$ is relatively compact in G , $\overline{\varphi(V)} \cap \varphi(H) = \varphi(V)$ and the image $\iota_H(V)$ of V in the group of inner automorphisms $\text{Int}(H)$ is relatively compact in $\text{Aut}(H)$. Furthermore, $V \cap \ker \varphi$ is trivial, $V \ker \varphi$ closed. $\varphi(H)$ is a normal subgroup of $\overline{\varphi(H)}$ and there exists a continuous homomorphism $\alpha: \overline{\varphi(H)} \rightarrow \text{Aut}(H/\ker \varphi)$ (necessarily unique) such that $\alpha \circ \varphi = \iota_{H/\ker \varphi} \circ \pi_1$ and this lifts also to an action of $\overline{\varphi(H)}$ on H .*

Writing $\iota_x(y) = xyx^{-1}$ ($x, y \in H$) for inner automorphisms defines the homomorphism $\iota_H: H \rightarrow \text{Int}(H)$. $\pi_1: H \rightarrow H/\ker \varphi$ denotes the quotient map. We do not exclude the case that $\varphi(H)$ is closed. Then V must be trivial.

Proof. Clearly, we can assume that $\overline{\varphi(H)} = G$. First, we consider the case where G is a Lie group. We follow the method of [Po]. Put $H_1 = H/\ker \varphi$ and let $\varphi_1: H_1 \rightarrow G$ be the induced homomorphism, $\pi_1: H \rightarrow H_1$ denotes the quotient mapping. Thus $\varphi = \varphi_1 \circ \pi_1$ and φ_1 is injective. By [Po] Prop. 1.10 there exists a vector subgroup V_1 of H_1 such that $G = \varphi_1(H_1) \overline{\varphi_1(V_1)}$, $\varphi_1(V_1)$ is relatively compact in G and $\varphi_1^{-1}(\overline{\varphi_1(V_1)}) = V_1$ (hence V_1 is closed). Put $V = \pi_1^{-1}(V)^0$ (identity component). $\pi_1^{-1}(V)$ is a Lie group, locally isomorphic to V (using that $\ker \varphi$ is discrete, see [Va] 2.6). It follows that $\pi_1(V) = V_1$, hence $\varphi(V) = \varphi_1(V_1)$ and (V_1 being simply connected) $\pi_1|_V$ is an isomorphism. Consequently, V is a vector subgroup of H , $V \cap \ker \varphi$ is trivial, $V \ker \varphi = \pi_1^{-1}(V)$ closed. $\varphi(V)$ is relatively compact. $\varphi_1^{-1}(\overline{\varphi_1(V_1)}) = V_1$ is equivalent to $\overline{\varphi_1(V_1)} \cap \varphi_1(H_1) = \varphi_1(V_1)$ and this is also equivalent to the corresponding property for V . $\varphi_1(H_1)$ is normal in G by [Po] 1.2. The construction of a homomorphism $\alpha: G \rightarrow \overline{\text{Int}(H_1)}$ is explained in [Po] p. 636 (using the notation A_φ and Ad_H for our ι_H). H and H_1 are locally isomorphic, hence as explained in [Po] p. 634, $\overline{\text{Int}(H)}$ and $\overline{\text{Int}(H_1)}$ are both isomorphic to the same subgroup of automorphisms of the Lie algebra of H and it follows

that α lifts to a homomorphism $\alpha^+ : G \rightarrow \overline{\text{Int}(H)}$ with $\alpha^+ \circ \varphi = \iota_H$, in particular $\overline{\iota_H(V)} = \alpha^+(\overline{\varphi(V)})$ is compact.

In the general case, G is a connected locally compact group, hence a projective limit of Lie groups. Let K be a compact normal subgroup of G such that G/K is a Lie group and the dimension of the Lie group $D = \varphi^{-1}(K)$ becomes minimal. If D would be non-discrete, take $x \in D^0 \setminus \ker \varphi$. Then $\varphi(x) \neq e$ and there would exist a compact normal subgroup $K_1 \subseteq K$ of G such that G/K_1 is a Lie group and $\varphi(x) \notin K_1$. Then $x \notin \varphi^{-1}(K_1)$ contradicting the minimality condition for K . Thus D must be discrete. Let $\pi_K : G \rightarrow G/K$ be the quotient mapping and put $\varphi_K = \pi_K \circ \varphi$. Then $\ker \varphi_K = D$ and we can apply the special case to $G/K, \varphi_K$. Let V be a vector subgroup of H satisfying the properties of Theorem A for φ_K . As mentioned above, this implies that $\varphi_K^{-1}(\overline{\varphi_K(V)}) = V$ and then the same property for φ follows easily. Compactness of K implies that $\varphi(V)$ is relatively compact in G . For $T = \overline{\varphi(V)}K$ we get that $G = \varphi(H)T$. Then by [Po] L. 1.1(ii), $\overline{\varphi(\varphi^{-1}(T))} = T$. We have $\pi_K(T) \subseteq \overline{\varphi_K(V)}$ and this implies $\varphi^{-1}(T) \subseteq \varphi_K^{-1}(\overline{\varphi_K(V)}) = V$ hence $T = \overline{\varphi(V)}$. Let $\alpha_K : G/K \rightarrow \overline{\text{Int}(H/\ker \varphi_K)}$ be the homomorphism with $\alpha_K \circ \varphi_K = \iota_{H/\ker \varphi_K}$. Then $\alpha^- = \alpha_K \circ \pi_K : G \rightarrow \overline{\text{Int}(H/\ker \varphi_K)}$ and as above this lifts to $\alpha : G \rightarrow \overline{\text{Int}(H/\ker \varphi)}$ and $\alpha^+ : G \rightarrow \overline{\text{Int}(H)}$ with the required properties. In particular $\overline{\iota_H(V)} = \alpha^+(\overline{\varphi(V)})$ is compact. For $y \in G, x \in H$ it follows by approximation that $y\varphi(x)y^{-1} = \varphi(\alpha^+(y)(x))$ hence $\varphi(H)$ is normal in G . \square

Proposition 6. (a) Let H_1, \mathcal{K} be locally compact groups, V_1 a closed subgroup of H_1 , $\varphi_0 : V_1 \rightarrow \mathcal{K}$, $\alpha_0 : \mathcal{K} \rightarrow \text{Aut}(H_1)$ continuous homomorphisms such that $\alpha_0 \circ \varphi_0 = \iota_{H_1}|_{V_1}$, $\mathcal{K} = \overline{\varphi_0(V_1)}$. Put $D = \{x^{-1}\varphi_0(x) : x \in V_1\}$ (in $H_1 \rtimes_{\alpha_0} \mathcal{K}$), $G_1 = (H_1 \rtimes_{\alpha_0} \mathcal{K})/D$. Let $\pi : H_1 \rtimes_{\alpha_0} \mathcal{K} \rightarrow G_1$ be the quotient map, $\varphi_1 = \pi|_{H_1}$.

Then $\overline{\varphi_1(V_1)} = \pi(\mathcal{K}) \cong \mathcal{K}$ (topologically), $\varphi_1(H_1) \cap \overline{\varphi_1(V_1)} = \varphi_1(V_1) = \pi(\text{im } \varphi_0)$, $\varphi_1 : H_1 \rightarrow G_1$, $\ker \varphi_1 = \ker \varphi_0$, $\overline{\varphi_1(H_1)} = G_1 = \varphi_1(H_1) \overline{\varphi_1(V_1)}$ (“attaching \mathcal{K} to H_1 along V_1 ”).

(b) Conversely, assume H, φ, V, α are as in Theorem A with $G = \overline{\varphi(H)}$, $\mathcal{K} = \overline{\varphi(V)}$. Put $H_1 = H/\ker \varphi$ and let $\varphi^- : H_1 \rightarrow G$ be the induced homomorphism, $\pi_1 : H \rightarrow H_1$ the quotient mapping, $V_1 = \pi_1(V)$, $\varphi_0 = \varphi^-|_{V_1}$, $\alpha_0 = \alpha|_{\mathcal{K}}$. Then $xk \mapsto \varphi(x)k$ ($x \in H_1, k \in \mathcal{K}$) induces a (topological) isomorphism of $G_1 = (H_1 \rtimes_{\alpha_0} \mathcal{K})/D$ and G . With $\varphi_1^+ = \varphi_1 \circ \pi_1 : H \rightarrow G_1$ it follows that (G, φ) and (G_1, φ_1^+) are equivalent in the sense corresponding to [Po] Def. 2.1.

If there exists a closed normal subgroup H_2 of H_1 such that $H_1 = H_2 \rtimes V_1$ splits

(equivalently, there exists a closed normal subgroup H_2^+ of H such that $H_2^+ \supseteq \ker \varphi$ and $H = H_2^+ \rtimes V$ splits), then $G_1 \cong H_2 \rtimes \mathcal{K}$ (restricting the action α_0 to H_2).

As usual, we identify H, \mathcal{K} with the corresponding subgroups of the semidirect product.

Proof. Using density of $\varphi_0(V)$ in \mathcal{K} one can show that D is a normal subgroup (if V_1 is abelian then D is even central). The properties of G_1 follow by routine computations (see also [Po] p.637f. for a related construction). In the situation of Theorem A, H is connected, \mathcal{K} compact and it follows that G_1 is σ -compact. By [HR] Th. 5.29 the isomorphism between G_1 and G is also a homeomorphism. A condition for the splitting case is [Po] Th. 1.12 and [Po] 1.12 an example of non-splitting. \square

Remarks. (a) A related result was shown in [G2] Th. 3 for the case where G is an arbitrary Hausdorff topological group. But then $\varphi(V)$ need not be relatively compact in G . [Po] gives a more direct argument for the case where G is a Lie group and adds various refinements, using the notion (from [G2]) of a “generalized maximal torus” in H . This can also be extended to the case of locally compact G , similarly as in the proof of Theorem A (but we do not need it for our applications in Section 2).

(b) In [PW] extensions to [G2] are given for the case where H is locally compact connected group. One can also extend most of Theorem A to the case where H, G are locally compact groups, H connected, $\varphi: H \rightarrow G$ a continuous homomorphism such that $\ker \varphi$ is totally disconnected, except for the property $\overline{\varphi(V)} \cap \varphi(H) = \varphi(V)$.

(c) For a connected Lie group H and a discrete central subgroup D_1 put $H_1 = H/D_1$. Given a closed vector subgroup V_1 of H_1 such that $\iota_{H_1}(V_1)$ is relatively compact in $\text{Aut}(H_1)$ the possible choices of compact groups \mathcal{K} and continuous homomorphisms $\varphi_0: V_1 \rightarrow \mathcal{K}$, $\alpha_0: \mathcal{K} \rightarrow \text{Aut}(H_1)$ with $\mathcal{K} = \overline{\varphi_0(V_1)}$ (hence \mathcal{K} must be abelian), $\alpha_0 \circ \varphi_0 = \iota_{H_1}|_{V_1}$ can be obtained by duality. Put $\Gamma = \widehat{\mathcal{K}}$, $\Gamma_0 = (\overline{\iota_H(V_1)})^\wedge$ (dual groups), $\iota_0 = \iota_{H_1}|_{V_1}$. The dual homomorphisms $\widehat{\varphi}_0: \Gamma \rightarrow \widehat{V_1}$, $\widehat{\iota}_0: \Gamma_0 \rightarrow \widehat{V_1}$ (called the adjoint homomorphism in [HR] 24.37) must be injective and satisfy $\widehat{\varphi}_0 \circ \widehat{\alpha}_0 = \widehat{\iota}_0$. Hence, up to isomorphism, the discrete group $\widehat{\mathcal{K}}$ must be a subgroup (algebraically) of $\widehat{V_1} (\cong \mathbb{R}^n)$ containing $\widehat{\iota}_0(\Gamma_0)$. Conversely, for any such subgroup $\widehat{\mathcal{K}}$ of $\widehat{V_1}$ (equipped with discrete topology), taking $\widehat{\varphi}_0$ the inclusion, one can apply Proposition A, producing a continuous homomorphism $\varphi_1: H_1 \rightarrow G_1$ with $\overline{\varphi_1(H_1)} = G_1 = \varphi_1(H_1) \overline{\varphi_1(V_1)}$, $\overline{\varphi_1(V_1)} \cong \mathcal{K}$ compact. Composition with

the quotient mapping $\pi_1: H \rightarrow H_1$ gives $\varphi = \varphi_1 \circ \pi_1: H \rightarrow G_1$. The maximal case is $\Gamma = V_1$ (discrete topology) giving $\mathcal{K} = b(V_1)$ (Bohr compactification). By [HR] 24.41, φ_0 is injective iff $\widehat{\mathcal{K}}$ is dense in \widehat{V}_1 and then $\ker \varphi = D_1$ giving (up to isomorphism all) examples for Theorem A. If φ_0 is not injective then $\ker \varphi_0$ contributes to $\ker \varphi$ and this does not fit with the setting of Theorem A (if $\ker \varphi_0$ is non-discrete one even gets cases where $\ker \varphi$ is non-discrete, see also Remark (b) in Section 2 on $H = \mathbb{H}$).

It is easy to see that if V satisfies the properties of Theorem A, the same is true for xVx^{-1} , $x \in H$.

The saturation condition $\overline{\varphi(V)} \cap \varphi(H) = \varphi(V)$ (going back to [Po]) is essential for the representation in Proposition 6(b) as a quotient of a semidirect product. Often there exist proper subgroups V_0 of V with $\overline{\varphi(V_0)} = \overline{\varphi(V)}$. For example, if G is second countable there always exists a one dimensional subgroup with this property ([G1] Prop. 12). But one cannot directly apply Proposition 6 with V_0 . The representation in [Po] p.637 (based on [Po] Th. 2.5) is closely related to our Proposition 6(b), but in the notation of [Po] V may also contain a compact part. As indicated in (b), for general locally compact connected groups H (e.g. when H is not arcwise connected) there are examples where it is not possible to choose V with the saturation condition. Then the classification of the extensions G gets more involved.

APPENDIX B. FOURIER TRANSFORM, INVERSION AND EXTENSIONS

[KL] does not touch the topic of the non-commutative Fourier transform. We will recall here some basic definitions and review the general background. See also [Fu] for a slightly more detailed survey.

Let G be a locally compact group. We take a fixed (left) Haar measure, in integrals we write simply dx etc. $L^p(G)$ denotes the standard (complex) L^p -spaces, $M(G)$ the space of bounded complex Radon measures, convolution is denoted by \star (see [HR] for basic properties). Representations π of G will always be strongly continuous and act unitarily on Hilbert spaces with inner product denoted by $(|)$. The extension to the algebra $L^1(G)$, $\pi(f) = \int_G f(x)\pi(x)dx$ ($f \in L^1(G)$) is denoted by the same letter, similarly for $M(G)$. The Fourier-Stieltjes algebra $B(G)$ consists of all coefficient functions $x \mapsto (\pi(x)f|g)$ of all representations of G . The Fourier algebra $A(G)$ consists of the coefficients of the left (or right) regular representation of G on $L^2(G)$ (see [E]). Both are Banach algebras under pointwise multiplication. $VN(G)$ denotes the von Neumann algebra on $L^2(G)$ generated by the left regular

representation. It is isometrically isomorphic to the dual space $A(G)'$.

For \mathcal{H} a Hilbert space, $\mathcal{B}(\mathcal{H})$ denotes the space of bounded linear operators, $\mathcal{K}(\mathcal{H})$ the subspace of compact operators, $\mathcal{S}_2(\mathcal{H})$ the Hilbert–Schmidt operators (with norm $\|\cdot\|_2$) and $\mathcal{S}_1(\mathcal{H})$ the trace class (norm $\|\cdot\|_1$).

\widehat{G} (the dual object of G) consists of the equivalence classes of irreducible representations of G , it has a topology and a Borel structure ([Di] Ch. 18). If G is type I and second countable there exists a Plancherel measure on \widehat{G} which we denote by ν . There exists a measurable selection of representatives from the equivalence classes. We fix such a selection (defined up to a set of ν -measure 0) and will use for simplicity the same letter π for a representation on the space \mathcal{H}_π selected in this way from the class π . This gives a (ν -) measurable field of Hilbert spaces (\mathcal{H}_π) providing a decomposition of the left regular representation as a direct integral of factorial representations, $VN(G) \cong \int_{\widehat{G}}^{\oplus} \mathcal{B}(\mathcal{H}_\pi) d\nu(\pi)$. Following [Fu] we will use the abbreviation $\mathcal{B}_1^{\oplus}(\widehat{G}, \nu)$ for the space of measurable operator fields (T_π) satisfying $T_\pi \in \mathcal{S}_1(\mathcal{H}_\pi)$ ν -a.e. and $\|(T_\pi)\|_1 = \int_0^\infty \|T_\pi\|_1 d\nu(\pi) < \infty$ (with the usual equivalence classes with respect to ν ; in the notation of [T1] p. 285 this is $\int_{\widehat{G}}^{\oplus} \mathcal{S}_1(\mathcal{H}_\pi) d\nu(\pi)$ describing the predual of $VN(G)$). Similarly $\mathcal{B}_2^{\oplus}(\widehat{G}, \nu)$ denotes the direct integral of Hilbert spaces $\int_{\widehat{G}}^{\oplus} \mathcal{S}_2(\mathcal{H}_\pi) d\nu(\pi)$ (i.e. $\|(T_\pi)\|_2^2 = \int_{\widehat{G}} \|T_\pi\|_2^2 d\nu(\pi)$).

When G is abelian, \widehat{G} coincides with the dual group of G , ν is a (suitably normalized – see below) left Haar measure, $\mathcal{B}_1^{\oplus}(\widehat{G}, \nu) = L^1(\widehat{G}, \nu)$, $\mathcal{B}_2^{\oplus}(\widehat{G}, \nu) = L^2(\widehat{G}, \nu)$. If G is a compact group, \widehat{G} is discrete, $\mathcal{B}_1^{\oplus}(\widehat{G}, \nu)$ is a weighted l^1 -sum, $\mathcal{B}_2^{\oplus}(\widehat{G}, \nu)$ a weighted l^2 -sum. In the examples of the Sections 4, 5, 7, we have that $\mathcal{H}_\pi = \mathcal{H}$ is a fixed Hilbert space for ν -almost all π . Then $\mathcal{B}_1^{\oplus}(\widehat{G}, \nu) = L^1(\widehat{G}, \nu, \mathcal{S}_1(\mathcal{H}))$ (Bochner integral for $\mathcal{S}_1(\mathcal{H})$ -valued functions, [T1] Ch. IV, Sec. 7 uses the notation $L_{\mathcal{S}_1(\mathcal{H})}^1(\widehat{G}, \nu)$), $\mathcal{B}_2^{\oplus}(\widehat{G}, \nu) = L^2(\widehat{G}, \nu, \mathcal{S}_2(\mathcal{H}))$. Similarly, $VN(G) \cong L^\infty(\widehat{G}, \nu, \mathcal{B}(\mathcal{H})) \cong L^\infty(\widehat{G}, \nu) \overline{\otimes} \mathcal{B}(\mathcal{H})$ when using a slightly weaker notion of measurability for $\mathcal{B}(\mathcal{H})$ -valued functions ([T1] Ch. IV, Def. 7.7). A slightly different example is in Section 6.

For $f \in L^1(G)$ we write $\mathcal{F}(f) = (\pi(f))$ for the (non-commutative) Fourier transform of f (similarly for $\mu \in M(G)$). Let $\lambda(f)(g) = f \star g$ ($g \in L^2(G)$) be the left convolution operator on $L^2(G)$. Then λ defines an embedding of $L^1(G)$ to a w^* -dense subalgebra of $VN(G)$ and \mathcal{F} extends to a w^* -continuous isomorphism of von Neumann algebras $VN(G) \rightarrow L^\infty(\widehat{G}, \nu)$ (again denoted by \mathcal{F}).

Assume now that G is unimodular. Then the crucial property of the Plancherel measure is that for $f \in L^1(G) \cap L^2(G)$ we have $\mathcal{F}(f) \in \mathcal{B}_2^{\oplus}(\widehat{G}, \nu)$ and $\|f\|_2 = \|\mathcal{F}(f)\|_2$ (and this determines ν uniquely when fixing a Haar measure on G). \mathcal{F} extends to an isometric isomorphism of Hilbert spaces $L^2(G) \rightarrow \mathcal{B}_2^{\oplus}(\widehat{G}, \nu)$ (again

denoted by \mathcal{F} ; [Fu] Th.3.31 uses the notation \mathcal{P}), it follows also that $(f|g) = \int_{\widehat{G}} \text{tr}(\pi(g)^* \pi(f)) d\nu(\pi)$ for $f, g \in L^1(G) \cap L^2(G)$ ($\text{tr}(\cdot)$ denoting the trace of the operator). For $f \in L^2(G) \cap VN(G)$ (i.e. the functions defining a convolution operator on $L^2(G)$; called the Hilbert algebra of G , [Di] 13.10) the two extensions of \mathcal{F} coincide. For $f \in L^1(G) \cap A(G)$ we have $\mathcal{F}(f) \in \mathcal{B}_1^\oplus(\widehat{G}, \nu)$ and $\|f\|_A = \|\mathcal{F}(f)\|_1$ (note that $\|\cdot\|_A$ does not depend Haar measure, but $\mathcal{F}(f)$ does and this is compensated by the Plancherel measure). \mathcal{F} extends to an isometric isomorphism of Banach spaces $A(G) \rightarrow \mathcal{B}_1^\oplus(\widehat{G}, \nu)$ (again denoted by \mathcal{F}). For $f \in L^2(G) \cap A(G)$ the two extensions of \mathcal{F} coincide (recall also that for unimodular G , $VN(G) \cap A(G) \subseteq L^2(G)$). The inverse mapping, for $T = (T_\pi) \in \mathcal{B}_1^\oplus(\widehat{G}, \nu)$, $u = \mathcal{F}^{-1}(T) \in A(G)$ is given by $u(x) = \int_{\widehat{G}} \text{tr}(\pi(x)^* T_\pi) d\nu(\pi)$ ($x \in G$).

In the general case (if G is not necessarily unimodular), let Δ be the modular function for G . In [DM] Th.5 it is shown that there exists a measurable family (K_π) of unbounded (densely defined) operators (called Duflo–Moore operators) such that for ν -almost all π , K_π is positive definite, self-adjoint (on \mathcal{H}_π) and satisfies the semi-invariance relations $\pi(x)^* K_\pi \pi(x) = \Delta(x) K_\pi$ for all $x \in G$ ([Fu] Th.3.48 uses $C_\pi = K_\pi^{-\frac{1}{2}}$). This gives rise to a modified Fourier transform $\mathcal{F}_2(f) = ([\pi(f) K_\pi^{\frac{1}{2}}])$ ($[\cdot]$ denoting the closure of the operator – if it exists). For $f \in L^1(G) \cap L^2(G)$ one has $\mathcal{F}_2(f) \in \mathcal{B}_2^\oplus(\widehat{G}, \nu)$ and $\|f\|_2 = \|\mathcal{F}_2(f)\|_2$, \mathcal{F}_2 extends to an isometric isomorphism of Hilbert spaces $L^2(G) \rightarrow \mathcal{B}_2^\oplus(\widehat{G}, \nu)$ (again denoted by \mathcal{F}_2). The modification $\mathcal{F}_A(f) = ([\pi(f) K_\pi])$ satisfies for $f \in L^1(G) \cap A(G)$ that $\mathcal{F}_A(f) \in \mathcal{B}_1^\oplus(\widehat{G}, \nu)$ and $\|f\|_A = \|\mathcal{F}_A(f)\|_1$, \mathcal{F}_A extends to an isometric isomorphism of Banach spaces $A(G) \rightarrow \mathcal{B}_1^\oplus(\widehat{G}, \nu)$ (again denoted by \mathcal{F}_A). Similar to the unimodular case, the extensions are compatible. For example, when $f \in L^2(G) \cap VN(G)$ (the left bounded elements of L^2) one has $\mathcal{F}_2(f)_\pi = [\mathcal{F}(f)_\pi K_\pi^{\frac{1}{2}}]$ ν -a.e. The inverse mapping, for $T = (T_\pi) \in \mathcal{B}_1^\oplus(\widehat{G}, \nu)$, $u = \mathcal{F}_A^{-1}(T) \in A(G)$ is again given by $u(x) = \int_{\widehat{G}} \text{tr}(\pi(x)^* T_\pi) d\nu(\pi)$ ($x \in G$). The Plancherel measure ν depends in the general case also on the choice of the family (K_π) , different Plancherel measures are equivalent (as noted in [Fu] Th.4.4, the formula for \mathcal{F}_A^{-1} does not involve the Duflo–Moore operators). The formulas given before for the unimodular case amount to take for K_π identity operators.

The isomorphism \mathcal{F}_A (resp. \mathcal{F}) can be used to transfer pointwise multiplication on $A(G)$ to define a multiplication \sharp on $\mathcal{B}_1^\oplus(\widehat{G}, \nu)$ called *dual convolution*. Thus for $S, T \in \mathcal{B}_1^\oplus(\widehat{G}, \nu)$, $S \sharp T = \mathcal{F}_A(\mathcal{F}_A^{-1}(S) \mathcal{F}_A^{-1}(T))$.

For a Lie group G one can also extend a representation π to the algebra of distributions of compact support. Then $\pi(d)$ will in general be an unbounded operator,

defined e.g. on the Gårding subspace of \mathcal{H}_π . This applies in particular to the elements of the Lie algebra of G which can be seen as distributions supported by $\{e\}$ (see [W] Sec. 4.4.1).

We want to add here some comments about a statement of [Fu]. In proving results on pointwise inversion of the Plancherel transform Führr asserts about a "gap" in a density argument used by earlier authors and introduces a new method based on positivity and bounded vectors (see also [Fu] Rem. 4.16(a)). The author seems to be unaware of Stinespring's paper [St] (in particular Th. 9.17 from there and its proof). Terminology and notation of [St] is quite different. For the unimodular case, we give here an argument for the (critical) "only if"-part of [Fu] Th. 4.15, in the spirit of [St] Rem. 9.4, adapted to our notation above (which is close to [Fu]). The general (non-unimodular) case is sketched in Remark (b) below.

For clarity, we denote here the extension of \mathcal{F} to $L^2(G)$ by \mathcal{P} (Plancherel transform). For $A \in B_1^\oplus$, $a = \mathcal{F}_A^{-1}(A)$, G unimodular (and type I), we want to show that $a \in L^2(G)$ implies $A \in B_2^\oplus$ and $A = \mathcal{P}(a)$.

First, we treat the special case where, in addition, $\mathcal{P}(a) \in B_1^\oplus$ holds. Put $A^{(0)} = A - \mathcal{P}(a)$, $a_0 = \mathcal{F}_A^{-1}(A^{(0)})$. Then $A^{(0)} \in B_1^\oplus$ and for $g \in (L^1 \cap L^2)(G)$ an easy computation, using the Parseval equality ([Fu] Th. 4.4) and unitarity of \mathcal{P} ([Fu] Th. 3.31), shows that $\int_{\widehat{G}} \text{tr}(\pi(g) A_\pi^{(0)*}) d\nu(\pi) = 0$.

Recall that \mathcal{F} extends to a w^* -continuous isomorphism of the von Neumann algebras $VN(G)$ and $\int_{\widehat{G}}^\oplus \mathcal{B}(\mathcal{H}_\pi) d\nu(\pi)$. B_1^\oplus is the predual of $\int_{\widehat{G}}^\oplus \mathcal{B}(\mathcal{H}_\pi) d\nu(\pi)$ ([T1] Ch. IV Prop. 8.34) and the Parseval equality of [Fu] Th. 4.4 says that \mathcal{F}_A^{-1} is the dual mapping for this isomorphism (for the sesquilinear duality). It follows that $\{\mathcal{F}(g) : g \in (L^1 \cap L^2)(G)\}$ is w^* -dense in $\int_{\widehat{G}}^\oplus \mathcal{B}(\mathcal{H}_\pi) d\nu(\pi)$, hence $A^{(0)} = 0$, i.e., $A_\pi = \mathcal{P}(a)_\pi$ holds ν -a.e.

For the general case, take $h \in L^1(G)$. Then by Fubini's theorem (see [Fu] p. 81) and continuity of \mathcal{P} , we have $\mathcal{P}(h \star a) = \mathcal{F}(h) \circ \mathcal{P}(a)$ and similarly, $\mathcal{F}_A^{-1}(\mathcal{F}(h) \circ A) = h \star a$. If $h \in (L^1 \cap L^2)(G)$ then $\mathcal{F}(h) = \mathcal{P}(h) \in B_2^\oplus$, hence $\mathcal{F}(h) \circ \mathcal{P}(a) \in B_1^\oplus$. Thus $\mathcal{F}(h) \circ A$ satisfies the additional requirement of the special case and it follows that $\pi(h) \circ A_\pi = \pi(h) \circ \mathcal{P}(a)_\pi$ holds ν -a.e. (with an exceptional set depending possibly on h).

For $c > 0$, put $M_c = \{\pi \in \widehat{G} : \|A_\pi\|_1 \leq c\}$ and $A_\pi^{(c)} = A_\pi - \mathcal{P}(a)_\pi$ for $\pi \in M_c$, $A_\pi^{(c)} = 0$ else, $A^{(c)} = (A_\pi^{(c)})$. Then $\|A_\pi\|_2 \leq \|A_\pi\|_1$ implies $A^{(c)} \in B_2^\oplus$ and we get $\int_{\widehat{G}} \text{tr}(\pi(h) A_\pi^{(c)}) d\nu(\pi) = 0$ for all $h \in (L^1 \cap L^2)(G)$. Since $\{\mathcal{P}(h) : h \in (L^1 \cap L^2)(G)\}$ is dense in B_2^\oplus , it follows that $A^{(c)} = 0$ in B_2^\oplus , thus $A_\pi = \mathcal{P}(a)_\pi$ a.e. on M_c . $c > 0$

was arbitrary, hence we arrive at $A = \mathcal{P}(a) \in B_2^\oplus$.

A similar argument shows that just assuming $a \in VN(G)$ already implies $a \in L^2(G)$. Also one gets that $\mathcal{P}(Bg) = \mathcal{F}(B) \circ \mathcal{P}(g)$ for $B \in VN(G)$, $g \in L^2(G)$ and then that \mathcal{P} coincides with \mathcal{F} on $VN(G) \cap L^2(G)$.

Remarks. (a) Some care is needed when using the duality between $A(G)$ and $VN(G)$. Traditionally it is obtained from the extension of the pairing $\langle f, g \rangle = \int_G f(x) g(x) dx$ ($f \in A(G)$, $g \in L^1(G)$) (and then the dual $A(G)$ -module structure on $VN(G)$ becomes the extension of pointwise multiplication). Under Fourier transform there is an extended Parseval equality $\int_G f(x) \overline{g(x)} dx = \int_{\widehat{G}} \text{tr}(\pi(g)^* \mathcal{F}_A(f)_\pi) d\nu(\pi)$. To get expressions for $\langle f, g \rangle$ (and more generally for $\langle f, T \rangle$ where $T \in VN(G)$) one has to analyze the effect of the conjugation $g \mapsto \bar{g}$ (which extends also to a semi-linear automorphism of $VN(G)$) on the disintegration (see also [Fu] p.110).

(b) The density argument above works similarly (with some more technicalities) for non-unimodular groups G (of type I and second countable). Let $K = (K_\pi)$ be the field of Duflo-Moore operators, and similar to [Fu], $[T] = ([T_\pi])$ shall denote the field built from the closures of a field of densely defined operators $T = (T_\pi)$. Then for A, a as above, the aim is to show that $a \in L^2(G)$ implies $[A \circ K^{-\frac{1}{2}}] = \mathcal{F}_2(a) \in B_2^\oplus$. If in addition $[\mathcal{F}_2(a) \circ K^{\frac{1}{2}}] \in B_1^\oplus$ holds, this can be done as in the unimodular case above. For $u \in L^1(G)$ one has $\mathcal{F}_A^{-1}(A \circ \mathcal{F}(u)) = a \star (\Delta u)$. Then observe that for those π where the semi-invariance relations hold (thus for ν -almost all π) one gets for $(1 + \Delta^{-\frac{1}{2}}) f \in L^1(G)$ that $\pi(f) \circ K_\pi^{\frac{1}{2}} \subseteq K_\pi^{\frac{1}{2}} \pi(\Delta^{-\frac{1}{2}} f)$ (in the sense of unbounded operators, i.e., the right operator is an [in general proper] extension of the left one). It follows also that $[K^{-\frac{1}{2}} \circ \mathcal{F}_2(f) \circ K^{\frac{1}{2}}] = \mathcal{F}_2(\Delta^{-\frac{1}{2}} f)$ for $(1 + \Delta^{-\frac{1}{2}}) f \in L^2(G)$ (compare [Fu] L.4.14). This implies $\mathcal{F}_2(f) \circ \mathcal{F}_2(v) = [\mathcal{F}_2(f \star (\Delta^{\frac{1}{2}} v)) \circ K^{\frac{1}{2}}]$ for $f \in L^2(G)$, $v \in (L^1 \cap L^2)(G)$ and using this, one can treat the general case similar as above.

(c) The Plancherel theorem was shown by Duflo-Moore [DM] for an arbitrary second countable group G for which the left regular representation λ is of type I (in particular, for all second countable groups of type I). The restrictions in the earlier work of Tatsuuma, Kleppner, Lipsman are unnecessary: if λ is of type I, then the same is true for the regular representation of $N = \ker \Delta$ ([DM] p.240); "N regularly embedded" (used also in [Fu]) can be replaced by a weaker property which as they show is always satisfied when λ is of type I.

(d) The Plancherel theorem (and the consequences above) extends to σ -compact G (instead of second countable): for $\mathfrak{K} = \{K : \text{compact normal subgroup of } G \text{ with}$

G/K second countable} the Kakutani-Kodaira theorem (see [LNPS] L. 24) says that G is the projective limit of $(G/K)_{K \in \mathfrak{K}}$. This implies that $\widehat{G} = \bigcup_{K \in \mathfrak{K}} \widehat{G/K}$ (more precisely, $\widehat{G/K}$ is homeomorphic to an open-closed subset of \widehat{G}). Similarly to [LNPS] L. 1 there are embeddings of $L^2(G/K)$ into $L^2(G)$ and $L^2(G) = \bigcup_{K \in \mathfrak{K}} L^2(G/K)$. $VN(G/K)$ is isomorphic to a von Neumann subalgebra of $VN(G)$ and $\bigcup_{K \in \mathfrak{K}} VN(G/K)$ is w^* -dense in $VN(G)$. $A(G/K)$ is isomorphic to a subalgebra of $A(G)$ and $A(G) = \bigcup_{K \in \mathfrak{K}} A(G/K)$ (see [KL] Prop. 2.4.2). When G is not second countable (but σ -compact) the Plancherel measure is no longer σ -finite.

(e) In [St] a duality theory (including Plancherel theorem and Fourier inversion) is developed for arbitrary unimodular locally compact groups (with no topological countability assumptions and no restrictions on the type). This extends to the non-unimodular case using the Plancherel weight ([T2] p. 67). It gives rather abstract statements in terms of unbounded operators on $L^2(G)$. If the group is second countable (or just σ -compact), there is a disintegration of the weight built up on the central disintegration of the left regular representation. Explicit examples can be found in a paper of Kajiwara (ref. [69] in [Fu]). In the non-type I case, non-trivial subfactors of $\mathcal{B}(\mathcal{H}_\sigma)$ (and non-trivial weights) arise. In the examples these are often related to ergodic theory.

REFERENCES

- [BH] W.R. Bloom, H. Heyer, *Harmonic Analysis of Probability Measures on Hypergroups*, de Gruyter, Berlin, New York, 1995.
- [CG1] Y. Choi, M. Ghandehari, *Weak and cyclic amenability for Fourier algebras of connected Lie groups*, J. Funct. Anal. **266** (2014), 6501–6530.
- [CG2] Y. Choi, M. Ghandehari, *Weak amenability for Fourier algebras of 1-connected nilpotent Lie groups*, J. Funct. Anal. **268** (2015), 2440–2463.
- [Da] H.G. Dales, *Banach algebras and automatic continuity*, London Math. Soc. Monographs, New Series 24, Oxford University Press, New York, 2000.
- [Di] J. Dixmier, *C^* -Algebras*, North Holland, Amsterdam–New York–Oxford, 1977.
- [DM] M. Duflo, C.C. Moore, *On the regular representation of a nonunimodular locally compact group*, J. Functional Analysis **21** (1976), 209–243.
- [DS] N. Dunford, J. T. Schwartz, *Linear Operators I*, Interscience, New York, London, 1958.
- [E] P. Eymard, *L 'algèbre de Fourier d'une groupe localement compact*, Bull. Soc. Math. France **92** (1964), 181–236.
- [Fo] G.B. Folland, *A course in abstract harmonic analysis*, CRC Press, Boca Raton, FL, 1995
- [Fu] H. Führ, *Abstract harmonic analysis of continuous wavelet transforms*, Lecture Notes Math. 1863, Springer, Berlin-Heidelberg-New York, 2005.
- [FL] H. Fujiwara, J. Ludwig, *Harmonic analysis on exponential solvable Lie groups*, Springer Monographs in Mathematics, Springer, Tokyo, 2015.

- [FR] B. E. Forrest, V. Runde, *Amenability and weak amenability of the Fourier algebra*, Math. Z. **250** (2005), 731–744.
- [G1] M. Goto, *Dense imbeddings of locally compact connected groups*, Ann. of Math. (2) **61** (1955), 154–169.
- [G2] M. Goto, *Immersion of Lie groups*, J. Math. Soc. Japan **32** (1980), 727–749.
- [HM] K. H. Hofmann, S. A. Morris, *The Structure of Compact Groups*, Walter de Gruyter, Berlin–New York, 1998.
- [HR] E. Hewitt, K. A. Ross, *Abstract harmonic analysis I*, Springer, Berlin, 1979.
- [J] B. E. Johnson, *Non-amenability of the Fourier algebra of a compact group*, J. London Math. Soc. (2) **50** (1994), 361–374.
- [KL] E. Kaniuth, A.T.-M. Lau, *Fourier and Fourier-Stieltjes algebras on locally compact groups*, Mathematical Surveys and Monographs 231. American Mathematical Society, Providence, RI, 2018.
- [Li] J.R. Liukkonen, *Dual spaces of groups with precompact conjugacy classes*, Trans. Amer. Math. Soc. **180** (1973), 85–108.
- [L] V. Losert, *On the center of group C^* -algebras*, J. Reine Angew. Math. **554** (2003), 105–138.
- [LLSS] H. H. Lee, J. Ludwig, E. Samei, N. Spronk, *Weak amenability of Fourier algebras and local synthesis of the anti-diagonal*, Adv. Math. **292** (2016), 11–41.
- [LNPS] V. Losert, M. Neufang, J. Pachl, J. Steprāns, *Proof of the Ghahramani–Lau conjecture*, Adv. Math. **290** (2016), 709–738.
- [Pl] R. J. Plymen, *Fourier algebra of a compact Lie group*, unpublished, see arXiv:math/0104018, 2001.
- [Po] D. Poguntke, *Dense Lie group homomorphisms*, J. Algebra **169** (1994), 625–647.
- [PW] W. H. Previts, T. S. Wu, *Immersion of locally compact connected groups*, Bull. Inst. Math. Acad. Sinica **31** (2003), 225–242.
- [St] W.F. Stinespring, *Integration theorems for gages and duality for unimodular groups*, Trans. Amer. Math. Soc. **90** (1959), 15–56.
- [Su] M. Sugiura, *Unitary representations and harmonic analysis. An introduction*, Second edition, North-Holland Publishing Co., Amsterdam 1990.
- [T1] M. Takesaki, *Theory of operator algebras I*, Springer, New York–Heidelberg 1979.
- [T2] M. Takesaki, *Theory of operator algebras II*, Springer, Berlin 2003.
- [Va] V. S. Varadarajan, *Lie Groups, Lie Algebras and Their Representations*, Springer, Berlin–Heidelberg–New York, 1984.
- [Vi] N.Ja. Vilenkin, *Special functions and the theory of group representations*, Transl. of Math. Monographs, American Mathematical Society, Providence, R.I. 1968.
- [W] G. Warner, *Harmonic analysis on semi-simple Lie groups. I*, Springer, New York–Heidelberg, 1972.

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT WIEN, STRUDLHOFG. 4, A 1090 WIEN, AUSTRIA

Email address: Viktor.Losert@UNIVIE.AC.AT