Guaranteed estimation of Hadamard shape derivative for clustered eigenvalues

Ryoki Endo^{*}, Xuefeng Liu[†]

May 24, 2023

Abstract

This paper proposes a guaranteed computation method to evaluate the Hadamard shape derivative for repeated eigenvalues. The proposed method enables the investigation of the behavior of eigenvalue variations around repeated eigenvalues, and provides rigorous estimation for the range of the Hadamard shape derivative in the case of clustered eigenvalues.

MSC:35P15, 65N25

Keywords: Laplacian eigenvalue, Hadamard shape derivative, guaranteed computation.

1 Introduction

While numerous studies have been conducted over a century on the Hadamard shape derivative for eigenvalues, the task of rigorously evaluating its concrete values still remains difficult. For well-separated Laplacian eigenvalues, the authors have proposed a method that guarantees computation for the evaluation of the first-order shape derivative [1]. In this context, the recently developed eigenvector error estimation from [2, 3] plays a significant role.

This paper will discuss the Hadamard shape derivative for repeated and clustered eigenvalues. Generally, the numerical computation of eigenfunctions for clustered eigenvalues poses an ill-posed problem. Instead of rigorously computing eigenfunctions, we utilize the guaranteed computation of eigenspaces and propose a stable method to evaluate the Hadamard shape derivative for repeated eigenvalues. The proposed method enables the investigation of the behaviour of eigenvalue variations around a repeated eigenvalue and provides a rigorous estimation of the Hadamard shape derivative's range in the case of clustered eigenvalues.

First, let us quote a fundamental result about the shape derivative from Henrot [4] in Theorem 1.1, which is originally stated in Rousselet [5]. For an open set Ω of \mathbb{R}^2 , we consider a family of function $\Phi(t)$ satisfying

$$\Phi: t \in [0,T) \to W^{1,\infty}(\mathbb{R}^2,\mathbb{R}^2)$$
 differentiable at 0 with $\Phi(0) = I, \Phi'(0) = V$

where $W^{1,\infty}(\mathbb{R}^2, \mathbb{R}^2)$ is the set of bounded Lipschitz maps from \mathbb{R}^2 to itself, I is the identity and V a vector field. Let $\Omega_t = \Phi(t)(\Omega)$ for $t \in [0, T)$.

Theorem 1.1. [4, Theorem 2.5.1, 2.5.8] Let Ω be a bounded open set with C^2 boundary. Assume that $\lambda_k(\Omega)$ is a multiple eigenvalue with multiplicity $m \in \mathbb{N}$. Let us denote by u_1, \dots, u_m an L^2 -orthonormal system of eigenfunctions corresponding to λ_k . If m = 1, then $t \mapsto \lambda_k(\Omega_t)$ is differentiable at t = 0, and we have

$$\left. \frac{d}{dt} \lambda_k(\Omega_t) \right|_{t=0} = -\int_{\partial\Omega} \left(\frac{\partial u}{\partial n} \right) \left(V \cdot n \right) \, ds$$

If m > 1, then $t \mapsto \lambda_k(\Omega_t)$ has a directional derivative at t = 0 which is one of the eigenvalues of the $m \times m$ matrix M defined by

$$M_{ij} = -\int_{\partial\Omega} \left(\frac{\partial u_i}{\partial n}\right) \left(\frac{\partial u_j}{\partial n}\right) (V \cdot n) \ ds. \ i, j = 1, \cdots, m.$$

^{*}Graduate School of Science and Technology, Niigata University, Niigata, Japan (endo@m.sc.niigata-u.ac.jp).

[†]Faculty of Science, Niigata University, Niigata, Japan (xfliu@math.sc.niigata-u.ac.jp).

The above shape derivative results are depicted through the integration of the eigenfunction across the domain boundary. Such a formulation requires higher regularities (e.g., H^2) of the involved eigenfunctions and poses difficuties in practical evaluation. Additionally, these results persist as a theoretical analysis under the presupposition of eigenvalue simplicity or multiplicity. However, in addressing practical problems involving clustered eigenvalues, the simplicity of eigenvalues is usually unknown. Consequently, the results of Theorem 1.1 cannot be directly utilized. In contrast, this paper introduces a new formulation that only necessitates the first derivative of the eigenfunction and can be applied to clustered eigenvalues.

The remainder of this paper is organized as follows: In Section 2, we review theorems on eigenvalue and eigenvector error estimations for the computation of the directional derivatives. In Section 3, we revisit results on eigenvalue perturbations and analyze the difference quotient of eigenvalues. In Chapter 4, details of the computation scheme for the first variations are provided. Finally, in Section 5, we state our conclusions.

2 Preliminary

We shape our discussion within the framework of Sobolev spaces. For a triangular domain $T \subset \mathbb{R}^2$, the function space $L^2(T)$ is constructed by all the real square integrable functions over T, and $H^1(T)$ is the first order Sobolev function space that the function has the first derivative to be in $L^2(T)$. We further introduce subspace $H_0^1(T)$ associated with the homogeneous Dirichlet boundary conditions.

Denote by $||v||_T$ the L^2 -norm of $v \in L^2(T)$. Integration $(\cdot, \cdot)_T$ is the inner product in $L^2(T)$ or $(L^2(T))^2$. Note that due to the boundary condition of the space $H_0^1(T)$, $(\nabla \cdot, \nabla \cdot)_T$ is an inner product for $H_0^1(T)$. Using such notation, the variational form of the Laplacian eigenvalue problem $-\Delta u = \lambda u$ reads as

Find
$$u \in H_0^1(T) \setminus \{0\}$$
 and $\lambda > 0$ s.t. $(\nabla u, \nabla v) = \lambda(u, v) \quad \forall v \in H_0^1(T).$ (1)

Here, ∇ is the gradient operator. As the inverse of the Laplacian is a compact self-adjoint operator, the spectral theorem shows that problem (1) has a spectrum of infinitely many eigenvalues, $0 < \lambda_1(T) < \lambda_2(T) \leq \lambda_3(T) \leq \cdots$.

The finite element method (FEM) will be utilized to evaluate the eigenvalue over triangles. Let us introduce the FEM approximation to the eigenvalue problem (1). Let $T \subset \mathbb{R}^2$ be a triangular domain. Denote by \mathcal{T}^h a regular triangulation of T; that is, any two edges e_i and e_j of elements of \mathcal{T}^h satisfy $e_i \cap e_j = e_i = e_j$ or $\mu(e_i \cap e_j) = 0$, where $\mu(\cdot)$ is the 1-dimensional Hausdorff measure. Let h be the maximal edge length of \mathcal{T}^h .

Let us introduce the following finite element spaces V_h^{CG} and V_h^{CR} over \mathcal{T}^h :

• The Lagrange FEM space V_h^{CG} :

 $V_h^{\text{CG}} := \{v_h : v_h \text{ is a continuous piecewise linear polynomial on } \mathcal{T}^h.\}$

• The Crouzeix–Raviart FEM space V_h^{CR} :

 $V_h^{CR} := \{ v_h : v_h \text{ is a piecewise linear polynomial on } \mathcal{T}^h; \}$

 v_h is continuous on the midpoint of each inter-element edge e; }.

Let $V_{h,0}^{CG}$ and $V_{h,0}^{CR}$ be the subspaces of the above finite element spaces defined by

$$V_{h,0}^{\scriptscriptstyle \mathrm{CG}} := H_0^1(T) \cap V_h^{\scriptscriptstyle \mathrm{CG}},$$

$$V_{h,0}^{\scriptscriptstyle \mathrm{CR}} := \{ v_h \in V_h^{\scriptscriptstyle \mathrm{CR}} : \int_e v_h = 0 \text{ for each boundary edge } e \text{ of } \mathcal{T}^h \} .$$

To estimate upper and lower bounds of the exact eigenvalues $\lambda_k(T)$, the following two eigenvalue problems are considered:

(a) Find $u_h \in V_{h,0}^{\scriptscriptstyle CG}(T) \setminus \{0\}$ and $\lambda_h > 0$ such that

$$(\nabla u_h, \nabla v_h)_T = \lambda_h(u_h, v_h)_T \quad \forall v_h \in V_{h,0}^{\scriptscriptstyle \mathrm{CG}}(T)$$

(b) Find $u_h \in V_{h,0}^{CR}(T) \setminus \{0\}$ and $\lambda_h > 0$ such that

$$(\nabla u_h, \nabla v_h)_T = \lambda_h(u_h, v_h)_T \quad \forall v_h \in V_{h,0}^{CR}(T).$$

Let $N_1 = \dim(V_{h,0}^{\scriptscriptstyle CG}(T)), N_2 = \dim(V_{h,0}^{\scriptscriptstyle CR}(T))$. The eigenvalues of (a) are denoted by

$$(0 <) \lambda_{1,h}^{\rm CG}(T) \leq \lambda_{2,h}^{\rm CG}(T) \leq \cdots \leq \lambda_{N_1,h}^{\rm CG}(T) ,$$

and the eigenvalues of (b) are denoted by

$$(0 <) \lambda_{1,h}^{CR}(T) \leq \lambda_{2,h}^{CR}(T) \leq \cdots \leq \lambda_{N_2,h}^{CR}(T) .$$

The energy projector $P_h^{\rm\scriptscriptstyle CG}: H^1_0(T) \to V_{h,0}^{\rm\scriptscriptstyle CG}$ is defined by

$$(\nabla(u - P_h u), \nabla v_h) = 0$$
 for all $v_h \in V_{h,0}^{\text{CG}}$.

Also, the projector $P_h^{\rm CR}: H_0^1(T) + V_{h,0}^{\rm CR} \to V_{h,0}^{\rm CR}$ is defined by

$$(\nabla(u - P_h u), \nabla v_h) = 0 \text{ for all } v_h \in V_{h,0}^{CR}.$$

Let $C_h^{\scriptscriptstyle \mathrm{CG}}$ and $C_h^{\scriptscriptstyle \mathrm{CR}}$ be the projection error constants defined by

$$C_h^{\rm \scriptscriptstyle CG} := \sup_{u \in H_0^1(T)} \frac{\|u - P_h^{\rm \scriptscriptstyle CG} u\|_T}{\|\nabla u - \nabla (P_h^{\rm \scriptscriptstyle CG} u)\|_T}, \ \ C_h^{\rm \scriptscriptstyle CR} := \sup_{u \in H_0^1(T) + V_{h,0}^{\rm \scriptscriptstyle CR}} \frac{\|u - P_h^{\rm \scriptscriptstyle CR} u\|_T}{\|\nabla u - \nabla (P_h^{\rm \scriptscriptstyle CR} u)\|_T}.$$

Remark 2.1. Let *h* be the mesh size of \mathcal{T}^h . The computable projection error estimation for $P_h^{\scriptscriptstyle CG}$ and $P_h^{\scriptscriptstyle CR}$ has been investigated extensively. For example, Liu and Kikuchi [6] obtained the estimation $C_h^{\scriptscriptstyle CG} \leq 0.493h$ for a uniform mesh with right triangle elements. The estimate for $C_h^{\scriptscriptstyle CR}$ given in Liu [7] is $C_h^{\scriptscriptstyle CR} \leq 0.1893h$.

By using FEM approximation, we can evaluate eigenvalues with rigorous upper and lower bounds:

Lemma 2.2. Let $N_1 = \dim V_{h,0}^{\scriptscriptstyle CG}(T)$ and $N_2 = \dim V_{h,0}^{\scriptscriptstyle CR}(T)$. We have

$$\underline{\lambda}_k := \frac{\lambda_{k,h}^{\scriptscriptstyle \mathrm{CR}}(T)}{1 + (C_h^{\scriptscriptstyle \mathrm{CR}})^2 \lambda_{k,h}^{\scriptscriptstyle \mathrm{CR}}(T)} \le \lambda_k(T) \le \lambda_{k,h}^{\scriptscriptstyle \mathrm{CG}}(T) =: \overline{\lambda}_k \quad \text{ for } k = 1, 2, ..., \min\{N_1, N_2\} .$$

Proof. The lower eigenvalue bound is provided in Liu [7], while the upper eigenvalue bound is from the min-max principle since $V_{h,0}^{cG}(T) \subset H_0^1(T)$.

The eigenfunction of the eigenvalue problem can also be well approximated by FEM solutions. Let us introduce distances to measure the error of approximated eigenfunctions. Given two subspaces E and \hat{E} of $H_0^1(T)$, the directed distances $\delta_a, \delta_b, \bar{\delta}_a$ and $\bar{\delta}_b$ are defined by

$$\delta_{a}(E, \widehat{E}) := \max_{\substack{v \in E \\ \|\nabla v\|_{T} = 1}} \min_{\hat{v} \in \widehat{E}} \|\nabla v - \nabla \hat{v}\|_{T}, \quad \delta_{b}(E, \widehat{E}) := \max_{\substack{v \in E \\ \|v\|_{T} = 1}} \min_{\hat{v} \in \widehat{E}} \|v - \hat{v}\|_{T},$$
$$\bar{\delta}_{a}(E, \widehat{E}) := \max_{\substack{v \in E \\ \|v\|_{T} = 1}} \min_{\substack{v \in E \\ \|v\|_{T} = 1}} \|\nabla v - \nabla \hat{v}\|_{T}, \quad \bar{\delta}_{b}(E, \widehat{E}) := \max_{\substack{v \in E \\ \|v\|_{T} = 1}} \min_{\substack{v \in E \\ \|v\|_{T} = 1}} \|v - \hat{v}\|_{T}$$

The directed distances δ_a and δ_b are not symmetric in general, but if the two subspaces are of the same finite dimension, then δ_a and δ_b are symmetric.

For L^2 -orthonormal systems $\mathcal{B} := (v_1, \cdots, v_n)$ and $\mathcal{B}' := (v'_1, \cdots, v'_n)$, let us introduce the distances δ^*_a and δ^*_b to measure the error between \mathcal{B} and \mathcal{B}' .

$$\delta_a^*(\mathcal{B}, \mathcal{B}') := \max_{1 \le i \le n} \|\nabla v_i - \nabla v_i'\|_T, \quad \delta_b^*(\mathcal{B}, \mathcal{B}') := \max_{1 \le i \le n} \|v_i - v_i'\|_T.$$

In order to formulate the bound on eigenfunctions, a notation for clusters of eigenvalues is introduced. Let n_k and N_k stand for indices of the first and the last eigenvalue in the k-th cluster; see Figure 1. Note that eigenvalues in a cluster need not be equal to each other. We consider the k-th cluster to be of interest, and set $n = n_k$ and $N = N_k$ to simplify the notation.

Figure 1: Clusters of eigenvalues

Let E_k be the space of exact eigenfunctions associated to kth cluster of eigenvalues:

$$E_k = \operatorname{span}\{u_n, u_{n+1}, \cdots, u_N\}.$$

Similarly, approximations $u_{h,i} \in H_0^1(T)$ of exact eigenfunctions u_i , for $i = n, n + 1, \dots, N$, form the corresponding approximate space:

$$\widehat{E}_k = \operatorname{span}\{\widehat{u}_n, \widehat{u}_{n+1}, \cdots, \widehat{u}_N\}.$$

Let $\hat{\lambda}_i := \|\nabla \hat{u}_i\|^2 / \|\hat{u}_i\|^2$ for $i = n, n + 1, \dots, N$. Let us introduce measures of non-orthogonality between finitedimensional subspaces E and E' of $H_0^1(T)$

$$\hat{\varepsilon}_a(E, E') = \max_{\substack{v \in E \\ \|\nabla v\| = 1 \ \|\nabla v'\| = 1}} \max_{\substack{v' \in E' \\ \|\nabla v'\| = 1}} (\nabla v, \nabla v'), \quad \hat{\varepsilon}_b(E, E') = \max_{\substack{v \in E \\ \|v\| = 1 \ \|v'\| = 1}} \max_{\substack{v' \in E' \\ \|v\| = 1 \ \|v'\| = 1}} (v, v').$$

Lemma 2.3. [Theorem 1 of Liu-Vejchodskỳ [2]] Take ρ such that $\lambda_n < \rho \leq \lambda_{N+1}$. Then, we have

$$\begin{split} \delta_a^2(E_k, \widehat{E}_k) &\leq \frac{\rho(\widehat{\lambda}_N - \lambda_n) + \lambda_n \widehat{\lambda}_N^{(k)} \theta_a^{(k)}}{\widehat{\lambda}_N^{(k)}(\rho - \lambda_n)} \\ \delta_b^2(E_k, \widehat{E}_k) &\leq \frac{\widehat{\lambda}_N^{(k)} - \lambda_n + \theta_b^{(k)}}{\rho - \lambda_n}, \end{split}$$

where

$$\theta_a^{(k)} = \sum_{l=1}^{k-1} \frac{\rho - \lambda_{n_l}}{\lambda_{n_l}} \left[\hat{\varepsilon}_a(\widehat{E}_l, \widehat{E}_k) + \delta_a(E_l, \widehat{E}_l) \right]^2,$$

$$\theta_b^{(k)} = \sum_{l=1}^{k-1} (\rho - \lambda_{n_l}) \left[\hat{\varepsilon}_b(\widehat{E}_l, \widehat{E}_k) + \delta_b(E_l, \widehat{E}_l) \right]^2.$$

The following summarizes the results of Lemma 1 and Theorem 1 of Liu-Vejchodsky [3]:

Lemma 2.4 (Estimation for eigenspace approximation). Suppose that $\widehat{E}_k \subset V_h^{\scriptscriptstyle CG}(T)$. Let ξ be a quantity that satisfies

$$\tau_h \xi < 1 - |\mathcal{J}|^{-1/2},$$

where

$$\tau := \max_{j \in \mathcal{J}} \max_{i \in \mathcal{I} \setminus \mathcal{J}} \frac{\lambda_j}{|\hat{\lambda}_i - \lambda_j|}, \ \tau_h := \max_{j \in \mathcal{J}} \max_{i \in \mathcal{I} \setminus \mathcal{J}} \frac{\hat{\lambda}_i}{|\hat{\lambda}_i - \lambda_j|},$$
$$\mathcal{I} := \{1, \cdots, \dim(V_h^{\text{CG}})\}, \ \mathcal{J} := \{n, n+1, \cdots, N\}, \ |\mathcal{J}| := N - n + 1$$

Let

$$\beta := \frac{\tau}{1 - \tau_h \xi}.$$

Then, for an arbitrary cluster of eigenvalues, the following estimate holds:

$$\delta_b(E_k, \widehat{E}_k) \le (1+\beta)\lambda_N (C_h^{\rm CG})^2.$$

For the distances of subspaces, we have the following properties:

Lemma 2.5. [estimations (19) and (49) of Liu-Vejchodsky [2]] For the distances δ_a and $\bar{\delta}_a$, it holds that

$$\delta_a^2(E_k, \widehat{E}_k) \le 2 - 2\lambda_n \left(\frac{1 - \delta_b^2(E_k, \widehat{E}_k)}{\lambda_N \widehat{\lambda}_N}\right)^{1/2},$$

$$\overline{\delta}_a^2(E_k, \widehat{E}_k) \le \lambda_N + \widehat{\lambda}_N - 2\lambda_n \sqrt{1 - \delta_b^2(E_k, \widehat{E}_k)}.$$

Lemma 2.6. For the distance $\overline{\delta}_b$, it holds that

$$\overline{\delta}_b^2(E_k, \widehat{E}_k) \le 2 - 2\sqrt{1 - \delta_b^2(E_k, \widehat{E}_k)}.$$

Proof. By the definition of $\overline{\delta}_b^2(E_k, \widehat{E}_k)$, we have

$$\overline{\delta}_{b}^{2}(E_{k},\widehat{E}_{k}) = \max_{\substack{u \in E \\ \|u\|_{T}=1}} \min_{\substack{\hat{u} \in \hat{E} \\ \|\hat{u}\|_{T}=1}} \left(\|u\|_{T}^{2} + \|\hat{u}\|_{T}^{2} - 2(u,\hat{u})_{T} \right)$$
$$\leq 2 - 2 \max_{\substack{u \in E \\ \|u\|_{T}=1}} \min_{\hat{u} \in \hat{E}} (u,\hat{u})_{T}$$
$$= 2 - 2\sqrt{1 - \delta_{b}^{2}(E_{k},\widehat{E}_{k})}.$$

-	_	

We will utilize the following lemma to validate the orthogonality of eigenfunctions:

Lemma 2.7. [Lemma 2 of [8]] Let A be a $n \times n$ matrix. If ||A - I|| < 1 for some fixed matrix norm, then we have det(A) > 0, where I denotes the $n \times n$ identity matrix.

3 Main theories

For each $p = (r, \theta)$ with $r > 0, \theta \in (0, \pi]$, vertices of a triangular domain T^p are represented as follows: O = (0, 0), For each p = (r, o) with $r > 0, v \in (0, \pi]$, vertices of a change at a change in the represented as followed 0 = (0, 0), $A = (1, 0), B = (r \cos \theta, r \sin \theta)$. In case $B = (r \cos \theta, r \sin \theta)$, let $\lambda_k^p := \lambda_k(T^p)$ (k = 1, 2, ...). For t > 0 and a normalized vector $e \in \mathbb{R}^2$, let $p_t := p + te$ be a perturbation of p. Let $\nabla_e^t \lambda_k^p := (\lambda_k^{p_t} - \lambda_k^p)/t$ be the difference quotient of the k-the eigenvalue at p. If λ_k^p has the directional derivative in the direction of the normalized vector $e \in \mathbb{R}^2$ at $p \in C$, denote the directional derivative by $\nabla_e \lambda_k^p$. For

the simplicity of notations, for $e^r = (-1, 0)$ and $e^{\theta} = (0, -1)$, write

$$\nabla_r \lambda_k^p := \nabla_{e^r} \lambda_k^p, \quad \nabla_\theta \lambda_k^p := \nabla_{e^\theta} \lambda_k^p \quad \nabla_r^t \lambda_k^p := \nabla_{e^r}^t \lambda_k^p, \quad \nabla_\theta^t \lambda_k^p := \nabla_{e^\theta}^t \lambda_k^p,$$

respectively.

In case of the regular triangle T^{p_0} $(p_0 = (1, \pi/3))$, note that $\lambda_2^{p_0}, \lambda_3^{p_0}$ are multiple, i.e., $\lambda_2^{p_0} = \lambda_3^{p_0}$; for the part of such a fact, see [9].

Perturbation of functions with respect to variation of triangles 3.1

Let T be the triangle with vertices O(0,0), A(1,0) and B(a,b). Let us introduce the perturbation of T by linear transform $\Phi: T \to T$:

$$\begin{pmatrix} \tilde{x} \\ \tilde{y} \end{pmatrix} = Q \begin{pmatrix} x \\ y \end{pmatrix}, \quad Q = \begin{pmatrix} 1 & \alpha \\ 0 & \beta \end{pmatrix} \quad (\beta > 0).$$

For u over T, define \tilde{u} over \tilde{T} by $\tilde{u} = u \circ \Phi$. The transpose of Q is denoted by Q^{\intercal} . Let $\widetilde{\nabla}\tilde{u} := (\frac{\partial}{\partial \tilde{x}}\tilde{u}, \frac{\partial}{\partial \tilde{u}}\tilde{u})^{\intercal}$ be the gradient of \tilde{u} . It holds that $\widetilde{\nabla}\tilde{u}(\tilde{x},\tilde{y}) = Q^{-\intercal}\nabla u(x,y)$. Let $\lambda_{\min}(\cdot)$ and $\lambda_{\max}(\cdot)$ denote the minimum and the maximum eigenvalues of a given square symmetric matrix, respectively.

Below, let us confirm the properties for the perturbation of u; see [10] for a detailed proof.

Lemma 3.1. Given function u over T, define $\tilde{u} = u \circ Q^{-1}$ over \tilde{T} .

(a) For L^2 -norm, we have

$$\|\tilde{u}\|_{\widetilde{T}}^2 = \beta \|u\|_T^2$$

(b) For H^1 -norm, we have

$$\lambda_{\min}(QQ^{\mathsf{T}}) \|\nabla \tilde{u}\|_{\tilde{T}}^2 \le \beta \|\nabla u\|_T^2 \le \lambda_{\max}(QQ^{\mathsf{T}}) \|\nabla \tilde{u}\|_{\tilde{T}}^2 \quad .$$

$$\tag{2}$$

Let $\gamma = \alpha^2 + \beta^2 + 1$. The eigenvalues of QQ^{\intercal} are given by

$$\lambda_{\min}(QQ^{\mathsf{T}}) = \frac{\gamma - \sqrt{\gamma^2 - 4\beta^2}}{2}, \quad \lambda_{\max}(QQ^{\mathsf{T}}) = \frac{\gamma + \sqrt{\gamma^2 - 4\beta^2}}{2}.$$

(c) For quantities involving the first derivative, we have

$$\begin{pmatrix} (\tilde{u}_{\tilde{x}}, \tilde{u}_{\tilde{x}})_{\widetilde{T}} \\ (\tilde{u}_{\tilde{x}}, \tilde{u}_{\tilde{y}})_{\widetilde{T}} \\ (\tilde{u}_{\tilde{y}}, \tilde{u}_{\tilde{y}})_{\widetilde{T}} \end{pmatrix} = \begin{pmatrix} \beta & 0 & 0 \\ -\alpha & 1 & 0 \\ \alpha^2 \beta^{-1} & -2\alpha \beta^{-1} & \beta^{-1} \end{pmatrix} \begin{pmatrix} (u_x, u_x)_T \\ (u_x, u_y)_T \\ (u_y, u_y)_T \end{pmatrix} .$$
(3)

Proof. The equality of (a) is evident. Since $\nabla u = Q^{\intercal} \nabla \tilde{u}$, we have

$$\lambda_{\min}(QQ^{\mathsf{T}}) \cdot (\tilde{u}_{\tilde{x}}^2 + \tilde{u}_{\tilde{y}}^2) \le u_x^2 + u_y^2 \le \lambda_{\max}(QQ^{\mathsf{T}}) \cdot (\tilde{u}_{\tilde{x}}^2 + \tilde{u}_{\tilde{y}}^2) \ .$$

Noting that $d\tilde{x}d\tilde{y} = \beta dxdy$ holds for the integrates over T and \tilde{T} , we obtain (2). The relation of (3) can be shown with an analogous argument.

Next, let us consider a concrete transformation $\Phi_{p,\tilde{p}}$ that maps T^p to $T^{\tilde{p}}$: for $p = (r, \theta), \tilde{p} = (\tilde{r}, \tilde{\theta}) \in \mathcal{C}$, the transformation matrix is given by

$$S_{p,\tilde{p}} := \begin{pmatrix} 1 & (\tilde{r}\cos\hat{\theta} - r\cos\theta)/(r\sin\theta) \\ 0 & (\tilde{r}\sin\hat{\theta})/(r\sin\theta) \end{pmatrix}$$

In case $\tilde{p} = p + te$ with ||e|| = 1, we shall write $S_t := S_{p,\tilde{p}}$.

Lemma 3.2 (Eigenvalue perturbation; Extension of Theorem 4.2 of [7]). For $p = (r, \theta), \tilde{p} = (\tilde{r}, \tilde{\theta}) \in C$, let $\tilde{B}(\tilde{r}\cos\tilde{\theta}, \tilde{r}\sin\tilde{\theta})$ be a perturbation of $B(r\cos\theta, r\sin\theta)$. Then, for the *i*-th $(i = 1, 2, \cdots)$ eigenvalue, we have

$$\lambda_{\min}\left(S_{p,\tilde{p}}S_{p,\tilde{p}}^{\mathsf{T}}\right)\cdot\lambda_{i}^{\tilde{p}}\leq\lambda_{i}^{p}\leq\lambda_{\max}\left(S_{p,\tilde{p}}S_{p,\tilde{p}}^{\mathsf{T}}\right)\cdot\lambda_{i}^{\tilde{p}}.$$

Proof. This result is an extension of the estimation of Theorem 4.2 in [7], where only the first eigenvalue is considered. The detailed proof for the general *i*-th eigenvalue is provided in the appendix. \Box

For t > 0, and a normal vector $e \in \mathbb{R}^2$, let P_t^e be the 2×2 matrix defined by

$$P_t^e := \left(S_t^{-1} S_t^{-\mathsf{T}} - I \right) / t.$$

Let $e^r = (-1, 0)$ and $e^{\theta} = (0, -1)$. In case $e = e^r$, we have

$$P_t^e := \frac{1}{(r-t)^2} \begin{pmatrix} t \cot^2 \theta & r \cot \theta \\ r \cot \theta & 2r - t. \end{pmatrix}$$

In case $e = e^{\theta}$, we have

$$P_t^e = \frac{1}{t\sin^2(\theta - t)} \begin{pmatrix} (\cos\theta - \cos(\theta - t))^2 & (\cos\theta - \cos(\theta - t))\sin\theta\\ (\cos\theta - \cos(\theta - t))\sin\theta & \sin^2\theta - \sin^2(\theta - t) \end{pmatrix}.$$

Note that, with a constant $\alpha \in [\theta - t, \theta]$ such that $\cos \theta = \cos(\theta - t) + t \sin \alpha$, the matrix P_t^e can be written as

$$P_t^e = \sin \alpha \cdot \begin{pmatrix} \frac{\cos \theta - \cos(\theta - t)}{\sin^2(\theta - t)} & \frac{\sin \theta}{\sin^2(\theta - t)} \\ \frac{\sin \theta}{\sin^2(\theta - t)} & \frac{\cos \theta + \cos(\theta - t)}{\sin^2(\theta - t)} \end{pmatrix}.$$

Let $P^e := \lim_{t \to 0+} P_t^e$. For a 2 × 2 symmetric matrix P, introduce the symmetric bilinear form $F_P : H_0^1(T^p) \times$ $H^1_0(T^p) \to \mathbb{R}$ defined by

$$F_P(u,v) := (P\nabla u, \nabla v)_{T^P}.$$
(4)

In case $P = P^e$, we shall write F_{P^e} as F_e . In particular, for e^r and e^{θ} , we have

$$F_{e^{r}}(u,v) = \frac{1}{r\tan\theta} \{ (u_{x}, v_{y})_{T} + (v_{x}, u_{y})_{T^{p}} \} + \frac{2}{r} (u_{y}, v_{y})_{T^{p}},$$

$$F_{e^{\theta}}(u,v) = \frac{2}{\tan\theta} (u_{y}, v_{y})_{T^{p}} - (u_{x}, v_{y})_{T^{p}} - (v_{x}, u_{y})_{T^{p}}.$$

3.2Analysis for the differential quotients of eigenvalues

By estimating the error between the eigenfunctions in the perturbed domain and the eigenfunctions in the original domain using the error estimations in Lemma 2.3, we derive the Hadamard shape derivative formula for simple eigenvalues and the directional derivative formula for repeated eigenvalues.

For the simplicity of the notations, we shall denote by λ_i the *i*-th eigenvalue λ_i^p at $p \in \mathcal{C}$. Let u_n, \dots, u_N be L²-orthonormal eigenfunctions corresponding to the multiple eigenvalues $\lambda_n = \cdots = \lambda_N (=: \lambda)$ at p. Let $\mathcal{B} := (u_n, u_{n+1}, \cdots, u_N)$ and $E := \operatorname{span}(\mathcal{B}).$

Let e be a normalized vector in \mathbb{R}^2 . For a perturbation $p_t := p + te \in \mathcal{C}$ (t > 0), we shall denote by λ_i^t the *i*-th eigenvalue λ_i^t . Let u_n^t, \dots, u_N^t be L^2 -orthonormal eigenfunctions corresponding to eigenvalues $\lambda_n^t \leq \dots \leq \lambda_N^t$, respectively. Introduce $\tilde{u}_i = u_i^t \circ \Phi_{p,p_t} (\in H^1_0(T^p))$ $(i = n, \cdots, N)$.

From the definitions of $\overline{\delta}_a$ and $\overline{\delta}_b$, we can take a system $\tilde{u}_n^*, \cdots, \tilde{u}_N^* \in \widetilde{E}_t$ such that

$$\|\nabla u_i - \nabla \tilde{u}_i^*\|_{T^p} \le \bar{\delta}_a(E, E_t), \quad \|u_i - \tilde{u}_i^*\|_{T^p} \le \bar{\delta}_b(E, E_t), \quad \|\tilde{u}_i^*\|_{T^p} = 1 \quad (i = n, \cdots, N).$$

Let $\tilde{\lambda}_i^* := \|\nabla \tilde{u}_i^*\|_{T^p}^2$ $(i = n, \cdots, N)$. Let M_t^* and N_t^* be $(N - m + 1) \times (N - m + 1)$ matrices defined by

$$M_t^* := \left(F_{P_t^e} \left(\tilde{u}_i^*, u_j \right) \right), \quad N_t^* := \left(\left(\tilde{u}_i^*, u_j \right)_{T^p} \right), \quad i, j = n, \cdots, N.$$

Assume that the system $\widetilde{\mathcal{B}}_t^* := (\tilde{u}_n^*, \tilde{u}_{n+1}^*, \cdots, \tilde{u}_N^*)$ is linearly independent. Such an assumption is reasonable when $\overline{\delta}(E, \widetilde{E}_t)$ is small enough and can be verified by utilizing Lemma 3.6. Then, each \widetilde{u}_i $(i = n, \dots, N)$ is uniquely represented as

$$\tilde{u}_i = s_{ni}\tilde{u}_n^* + \dots + s_{Ni}\tilde{u}_N^*, \quad \text{where } s_{ni}, \dots, s_{Ni} \in \mathbb{R}.$$
(5)

Lemma 3.3. Let $\sigma_i = (s_{ni}, \dots, s_{Ni})^{\intercal}$ $(i = n, \dots, N)$ be the coefficient vector. Then, the pair $((\lambda_i^t - \lambda)/t, \sigma_i)$ becomes the (i - n + 1)-th eigenpair of the following generalized matrix eigenvalue problem: Find $\mu \in \mathbb{R}$ and $\sigma \in \mathbb{R}^{N-n+1} \setminus \{0\}$ such that

$$M_t^* \sigma = \mu N_t^* \sigma. \tag{6}$$

Proof. Let us calculate difference quotient $(\lambda_j^t - \lambda)/t$ $(j = n, \dots, N)$. For each $j = n, \dots, N$, the following variational equation holds:

$$(\nabla u_j^t, \nabla \tilde{v})_{T^{p_t}} = \lambda_j^t (u_j^t, \tilde{v})_{T^t} \quad \forall \tilde{v} \in H_0^1(T^t),$$

Note that $\widetilde{\nabla} u_j^t = (S_t^{-\intercal}) \nabla \widetilde{u}_j$, we have

$$\left((S_t^{-\mathsf{T}}) \nabla \tilde{u}_j, (S_t^{-\mathsf{T}}) \nabla v \right)_{T^p} = \lambda_j^t (\tilde{u}_j, v)_{T^p} \quad \forall v \in H^1_0(T^p)$$

Substituting $v = u_i$ $(i = n, \dots, N)$, it follows that

$$\left(S_t^{-1}S_t^{-\mathsf{T}}\nabla\tilde{u}_j, \nabla u_i\right)_{T^p} = \lambda_j^t(\tilde{u}_j, u_i)_{T^p}.$$
(7)

Also, for the eigenpair (λ^p, u_i) , it holds that

 $(\nabla u_i, \nabla v)_{T^p} = \lambda(u_i, v)_{T^p} \quad \forall v \in H^1_0(T^p).$

Take $v = \tilde{u}_j$ in the above variational equation, then we have

$$(\nabla u_i, \nabla \tilde{u}_j)_{T^p} = \lambda(u_i, \tilde{u}_j)_{T^p}.$$
(8)

Recall that the symmetric bilinear form $F_{P_t^e}$ is defined by (4). From (7) and (8), we have

$$\left(\frac{1}{t}(S_t^{-1}S_{p,p_t}^{-\intercal}-I)\nabla\tilde{u}_j,\nabla u_i\right)_{T^p}=\frac{\lambda_j^t-\lambda}{t}(\tilde{u}_j,u_i)_{T^p},$$

i.e.,

$$F_{P_t^e}(\tilde{u}_j, u_i) = \frac{\lambda_j^t - \lambda}{t} (\tilde{u}_j, u_i)_{T^p}.$$

Substituting (5), we have

$$\sum_{k=n}^{N} s_{kj} F_{P_t^e}(\tilde{u}_k^*, u_i) = \frac{\lambda_j^t - \lambda}{t} \sum_{k=n}^{N} s_{kj}(\tilde{u}_k^*, u_i)_T \text{ for } i, j = n, \cdots, N.$$
(9)

Introduce the matrices M_t^*, N_t^* and the orthogonal vectors σ_j $(j = n, \dots, N)$ by

$$M_t^* := \left(F_{P_t^e} \left(\tilde{u}_k^*, u_i \right) \right), \quad N_t^* := \left(\left(\tilde{u}_k^*, u_i \right)_{T^p} \right), \quad \sigma_j := \left(s_{kj} \right), \quad i, k = n, \cdots, N.$$

Then the relation (9) is transformed into

$$M_t^* \sigma_j = \left(\frac{\lambda_j^t - \lambda}{t}\right) \left(N_t^* \sigma_j\right).$$
⁽¹⁰⁾

Since we have $(\lambda_n^t - \lambda)/t \leq \cdots, \leq (\lambda_N^t - \lambda)/t$, the difference quotient $((\lambda_i^t - \lambda)/t, \sigma_i)$ becomes the (i - n + 1)-th eigenvpair of the matrix pair (M_t^*, N_t^*) .

From (6) of Lemma 3.3, one can evaluate the quantity $(\lambda_i^t - \lambda)/t$ for all $t \in (0, t_0]$ by solving the generalized matrix eigenvalue problem (6) with the matrices represented by intervals. Since $\lim_{t\to 0+} (\lambda_i^t - \lambda)/t$ provides shape derivative, we can obtain an explicit estimate of this derivative.

Corollary 3.4. For each $t \in (0, t_0]$, suppose that the concerned cluster of eigenvalues is separated from the other clusters, i.e., $\lambda_{n-1}^t < \lambda_n^t, \lambda_N^t < \lambda_{N+1}^t$. Also, suppose that the concerned eigenvalues are multiple at p, i.e., $\lambda_n = \lambda_{n+1} = \cdots = \lambda_N(=:\lambda)$. Then, the eigenvalues $\mathcal{C} \ni \tilde{p} \mapsto \lambda_i^{\tilde{p}}$ $(i = n, \cdots, N)$ are directionally differentiable with respect to the normalized vector $e \in \mathbb{R}^2$ at p.

Moreover, the directional derivative $\nabla_e \lambda_i$ $(k = n, \dots, N)$ coincides with the (i - n + 1)-th eigenvalue of the symmetric matrix M defined by

$$M = (F_{P^e}(u_i, u_j)), \quad \text{where} \quad i, j = n, \cdots, N.$$

$$\tag{11}$$

Proof. By the continuity of eigenvalues with respect to domain deformations and the estimations in Lemma 2.5 and Lemma 2.6, the values of $\overline{\delta}_a(E, \widetilde{E}_t)$ and $\overline{\delta}_b(E, \widetilde{E}_t)$ converge to 0 as $t \to 0+$. Thus, letting $t \to 0+$, the matrices M_t^* and N_t^* in (10) converge to the matrix M in (11) and the $(N - n + 1) \times (N - n + 1)$ identity matrix. Therefore, the value of $(\lambda_i^t - \lambda)/t$ converges to the (i - n + 1) eigenvalue of M.

Corollary 3.5. If the *i*-th $(i = n, \dots, N)$ eigenvalue λ_i is simple, then λ_i is partially differentiable w.r.t. r and θ at p. Moreover, we have

$$\frac{\partial \lambda_i}{\partial r} = F_r(u, u), \quad \frac{d\lambda_i}{d\theta} = F_\theta(u, u).$$

Proof. Let $e = e^r$ (resp. $e = e^{\theta}$). In case the concerned eigenvalue λ_k is simple, the matrix M defined in (11) becomes 1×1 -matrix. Hence, it follows that $\nabla_{e^r} \lambda_k = -\nabla_{e^{-r}} \lambda_k$ (resp. $\nabla_{e^{\theta}} \lambda_k = -\nabla_{e^{-\theta}} \lambda_k$). Therefore, λ_k is partially differentiable with respect to r (resp. θ).

By the following Lemma, we can validate the linear independency of the system $\widetilde{\mathcal{B}}_t^* := (\widetilde{u}_n^*, \widetilde{u}_{n+1}^*, \cdots, \widetilde{u}_N^*)$:

Lemma 3.6. Let N, \widetilde{N}^* be the $(N - m + 1) \times (N - m + 1)$ matrices defined by

$$N := \left((u_i, u_j)_T \right), \quad \widetilde{N}^* := \left(\left(\widetilde{u}_i^*, \widetilde{u}_j^* \right)_T \right), \quad i, j = n, \cdots, N.$$

If $2(N - n + 1) \cdot \overline{\delta}_b(E, \widetilde{E}_t) < 1$, then the system $\widetilde{\mathcal{B}}_t^* = (\tilde{u}_n^*, \tilde{u}_{n+1}^*, \cdots, \tilde{u}_N^*)$ is linearly independent.

Proof. Let us compare each element of N and \widetilde{N}^* . For each $i, j = n, \dots, N$, we have

$$\begin{aligned} \left| (u_i, u_j) - \left(\tilde{u}_i^*, \tilde{u}_j^* \right) \right| &= \left| (u_i, u_j)_T - \left(\tilde{u}_i^*, u_j^* \right)_T \right| \\ &\leq \left| (u_i, u_j) - (u_i, \tilde{u}_j^*) \right| + \left| (u_i, \tilde{u}_j^*) - (\tilde{u}_i^*, \tilde{u}_j^*) \right| \\ &\leq \left\| u_i \right\|_T \left\| u_j - \tilde{u}_j^* \right\|_T + \left\| \tilde{u}_j^* \right\|_T \left\| u_i - \tilde{u}_i^* \right\|_T \\ &\leq 2 \overline{\delta}_b(E, \widetilde{E}_t). \end{aligned}$$

Therefore, it follows that

$$\left\|\widetilde{N}^* - I\right\|_F = \left\|\widetilde{N}^* - N\right\|_F \le 2(N - n + 1) \cdot \overline{\delta}_b(E, \widetilde{E}_t).$$

Suppose that $2(N-n+1)\cdot \overline{\delta}_b(E, \widetilde{E}_t) < 1$. Then, we have $\left\| \widetilde{N}^* - I \right\|_F < 1$. By Lemma 2.7, it holds that $\det(\widetilde{N}^*) > 0$, that is, the system $\widetilde{\mathcal{B}}_t^*$ is linearly independent.

3.3 Estimation for difference quotients of eigenvalues

In this subsection, we estimate the error between the matrices M_t^*, N_t^* in the eigenvalue problem (6) and their approximate matrices calculated by numerical schemes.

Let us continue to use the notations introduced in the previous subsection. Let $\hat{u}_n, \dots, \hat{u}_N (\in H_0^1(T^p))$ be L^2 -orthonormal approximations to u_n, \dots, u_N obtained by numerical schemes, i.e., finite element method. Let $\hat{\lambda}_i := \|\nabla \hat{u}_i\|^2$ for $i = n, \dots, N$. Set $\hat{\mathcal{B}} := (\hat{u}_n, \hat{u}_{n+1}, \dots, \hat{u}_N)$ and $\hat{E} := \operatorname{span}(\hat{\mathcal{B}})$.

Define matrices $M_t, \widehat{M}_t, M_t^*$ and N_t^* by

$$M_t := \left(F_{P_t^e}(u_i, u_j) \right), \quad \widehat{M}_t = \left(F_{P_t^e}(\hat{u}_i, \hat{u}_j) \right), \quad M_t^* := \left(F_{P_t^e}(\tilde{u}_i^*, u_j) \right), \quad N_t^* := \left((\tilde{u}_i^*, u_j)_{T^p} \right), \quad i, j = n, \cdots, N.$$
(12)

Note that \widehat{M}_t will be constructed explicitly in the following numerical computation, with other matrices just for the purpose of theoretical discussion.

By Lemma 4.1, we can take an L^2 -orthonormal system $\widehat{\mathcal{B}}^* = (\hat{u}_n^*, \cdots, \hat{u}_N^*)$ such that

$$\delta_a^*(\mathcal{B},\widehat{\mathcal{B}}^*) \leq \operatorname{Err}_a^*(\mathcal{B},\widehat{\mathcal{B}}^*).$$

Define matrices \widehat{M}_t^* and \widehat{N}_t^* by

$$\widehat{M}_{t}^{*} := \left(F_{P_{t}^{e}}(\hat{u}_{i}^{*}, \hat{u}_{j}^{*}) \right), \quad \widehat{N}_{t}^{*} := \left((\hat{u}_{i}^{*}, \hat{u}_{j}^{*})_{T^{p}} \right), \quad i, j = n, \cdots, N$$

Note that the matrices \widehat{M}_t^* and \widehat{N}_t^* are similar to \widehat{M}_t and the identity matrix I, respectively.

Lemma 3.7. For the matrices M_t^*, N_t^* , we have

$$\left\| M_t^* - \widehat{M}_t^* \right\|_F \le \operatorname{Err}(M_t^*, \widehat{M}_t^*), \quad \|N_t^* - I\|_F \le \operatorname{Err}(N_t^*, I),$$
(13)

where

$$\operatorname{Err}(M_t^*, \widehat{M}_t^*) := (N - n + 1) \sqrt{\widehat{\lambda}_N} \|P_t^e\|_2 \cdot \left\{ \overline{\delta}_a(E, \widetilde{E}_t) + 2\operatorname{Err}_a^*(\mathcal{B}, \widehat{\mathcal{B}}^*) \right\},$$
$$\operatorname{Err}(N_t^*, I) = (N - n + 1) \cdot \overline{\delta}_b(E, \widetilde{E}_t).$$

Proof. Let us compare each element of N_t^* and the identity matrix I. For each $i, j = n, \dots, N$, since we have

$$|(u_i, u_j) - (\tilde{u}_i^*, u_j)| \le ||u_i - \tilde{u}_i^*||_{T^p} ||u_j||_{T^p} \le \overline{\delta}_b(E, E_t)$$

it holds that

$$\|N_t^* - I\|_F \le \operatorname{Err}(N_t^*, I).$$

Let us compare each element of M_t^* and \widehat{M}_t^* . For each $i, j = n, \dots, N$, we have

$$\begin{aligned} \left| F_{P_{t}^{e}} \left(\tilde{u}_{i}^{*}, u_{j} \right) - F_{P_{t}^{e}} \left(\hat{u}_{i}^{*}, \hat{u}_{j}^{*} \right) \right| &= \left| (P_{t}^{e} \nabla \tilde{u}_{i}^{*}, \nabla u_{j})_{T^{p}} - (P_{t}^{e} \nabla u_{i}, \nabla u_{j})_{T^{p}} \right| \\ &\leq \left| (P_{t}^{e} \nabla \tilde{u}_{i}^{*}, \nabla u_{j})_{T^{p}} - (P_{t}^{e} \nabla u_{i}, \nabla u_{j})_{T^{p}} \right| \\ &+ \left| (P_{t}^{e} \nabla u_{i}, \nabla u_{j})_{T^{p}} - (P_{t}^{e} \nabla \tilde{u}_{i}, \nabla \tilde{u}_{j})_{T^{p}} \right| \\ &+ \left| (P_{t}^{e} \nabla u_{i}, \nabla \tilde{u}_{j}^{*})_{T^{p}} - (P_{t}^{e} \nabla \tilde{u}_{i}, \nabla \tilde{u}_{j})_{T^{p}} \right| \\ &\leq \left\| P_{t}^{e} \right\|_{2} \left\| \nabla \tilde{u}_{i}^{*} - \nabla u_{i} \right\|_{T^{p}} \left\| \nabla u_{j} \right\|_{T^{p}} \\ &+ \left\| P_{t}^{e} \right\|_{2} \left\| \nabla u_{i} \right\|_{T^{p}} \left\| \nabla u_{j} - \nabla \tilde{u}_{j}^{*} \right\|_{T^{p}} \\ &+ \left\| P_{t}^{e} \right\|_{2} \left\| \nabla u_{i} - \nabla \tilde{u}_{i}^{*} \right\|_{T^{p}} \\ &\leq \sqrt{\hat{\lambda}_{N}} \left\| P_{t}^{e} \right\|_{2} \cdot \left\{ \overline{\delta}_{a}(E, \widetilde{E}_{t}) + 2\delta_{a}^{*}(\mathcal{B}, \widehat{\mathcal{B}}^{*}) \right\}. \end{aligned}$$
(14)

By the estimations (14), we have

$$\|M_t - M_t^*\|_F \le \operatorname{Err}(M_t^*, \widehat{M}_t^*).$$

Remark 3.8. Let $(\hat{u}, \hat{\sigma})$ be an eigenpair of the following generalized matrix eigenvalue problem: Find $\hat{\sigma} \in \mathbb{R}$ and $\hat{u} \in \mathbb{R}^{N-n+1} \setminus \{0\}$ such that

$$\widehat{M}_t^* \widehat{\sigma} = \widehat{\mu} \widehat{N}_t^* \widehat{\sigma}. \tag{15}$$

Note that the matrices $\widehat{M}_t^*, \widehat{N}_t^*$ are similar to \widehat{M}_t, I , respectively, where I denotes the $(N - n + 1) \times (N - n + 1)$ identity matrix. Thus, it is easy to see that the eigenvalues of the eigenvalue problem (15) coincide with the eigenvalues of the following generalized matrix eigenvalue problem: Find $\hat{\sigma} \in \mathbb{R}$ and $\hat{u} \in \mathbb{R}^{N-n+1} \setminus \{0\}$ such that

$$\widehat{M}_t \widehat{\sigma} = \widehat{\mu} \widehat{N}_t \widehat{\sigma}. \tag{16}$$

Denote by μ_k $(k = 1, \dots, N - n + 1)$ the k-the eigenvalue of the eigenvalue problem (6). In order to calculate the value of $\hat{\mu}_k$ $(k = 1, \dots, N - n + 1)$, it suffices to solve the eigenvalue problem (16).

Denote by $\hat{\mu}_k$ $(k = 1, \dots, N - n + 1)$ the k-the eigenvalue of the eigenvalue problem (15). Then, the following estimation tells us an error bound between μ_k and $\hat{\mu}_k$:

$$|\mu_k - \hat{\mu}_k| \le \left\| M_t^* - \widehat{M}_t^* (\widehat{N}_t^*)^{-1} \right\|_2 \le \operatorname{Err}(M_t^*, \widehat{M}_t^*) + \left\| \widehat{M}_t \right\|_2 \left\| I - (\widehat{N}_t^*)^{-1} \right\|_2 := \eta(M_t^*, \widehat{M}_t^*, \widehat{N}_t^*)$$
(17)

Note that each term of the error bound in (17) can be explicitly estimated utilizing Lemma 3.7.

3.4 Estimation for the range of directional derivative of eigenvalues

In this subsection, we estimate the range of directional derivatives of eigenvalues when the value of the eigenvalue of interest is very close to, or coincides with, other eigenvalues.

Let u_n, \dots, u_N be and L^2 -orthonormal eigenfunctions corresponding to the simple or multiple eigenvalues $\lambda_n \leq \dots \leq \lambda_N$ at $p \in \mathcal{C}$. Let \mathcal{B} be an L^2 -orthonormal system defined by $\mathcal{B} := (u_n, \dots, u_N)$, and let $E := \operatorname{span}(\mathcal{B})$. For a normalized vector $e \in \mathbb{R}^2$, define a $(N - n + 1) \times (N - n + 1)$ matrix M by

$$M = (F_{P^e}(u_i, u_j)) \quad \text{where } i, j = n, \cdots, N.$$

$$(18)$$

In case $e = e^r$, e^{θ} and all the eigenvalues are simple, i.e., $\lambda_n < \cdots < \lambda_N$, from Corollary 3.5, it follows that the (i - n + 1)-th $(i = n, \dots, N)$ diagonal element of the matrix M coincides with the partial derivative of λ_i with respect to r, θ , respectively.

Let $\hat{u}_n, \dots, \hat{u}_N (\in H_0^1(T^p))$ be L^2 -orthonormal approximations to $u_n, \dots, u_N (\in H_0^1(T^p))$ obtained by numerical schemes. Let $\hat{\lambda}_i := \|\nabla \hat{u}_i\|_{T^p}^2$ for $i = n, \dots, N$. Set $\hat{\mathcal{B}} := (\hat{u}_n, \dots, \hat{u}_N)$ and $\hat{E} := \operatorname{span}(\hat{\mathcal{B}})$. Define a $(N - n + 1) \times (N - n + 1)$ matrix \widehat{M} by

$$M = (F_{P^e}(\hat{u}_i, \hat{u}_j)) \quad \text{where } i, j = n, \cdots, N.$$
(19)

By Lemma 4.1, we can take an L^2 -orthonormal system $\widehat{\mathcal{B}}^* := (\hat{u}_n^*, \cdots, \hat{u}_N^*)$ such that

$$\delta_a^*(\mathcal{B},\widehat{\mathcal{B}}^*) \le \operatorname{Err}_a^*(\mathcal{B},\widehat{\mathcal{B}}^*), \quad \delta_b^*(\mathcal{B},\widehat{\mathcal{B}}^*) \le \operatorname{Err}_b^*(\mathcal{B},\widehat{\mathcal{B}}^*).$$
(20)

Since \widehat{B} and \widehat{B}^* both form L^2 -orthonormal system of \widehat{E} , there exists a $(N - n + 1) \times (N - n + 1)$ orthogonal matrix R such that

$$(\hat{u}_n^*,\cdots,\hat{u}_N^*)^\mathsf{T} = R(\hat{u}_n,\cdots,\hat{u}_N)^\mathsf{T}$$

By appropriately taking the sign of approximate eigenfunctions $\hat{u}_n, \dots, \hat{u}_N$, we can take $\hat{u}_n, \dots, \hat{u}_N$ so that the matrix R becomes a rotation matrix, i.e., $\det(R) = 1$.

By comparing each element of the matrices M and \widehat{M}^* utilizing (20), we obtain the following error estimation.

$$\left\|M - \widehat{M}^*\right\|_F \le 2(N - n + 1)\sqrt{\widehat{\lambda}_N} \|P^e\|_2 \cdot \delta_a^*(\mathcal{B}, \widehat{\mathcal{B}}^*).$$

Let us summarize the above results in the following Lemma:

Lemma 3.9. For the matrices M, \widehat{M}^* and \widehat{M} defined in (18) and (19), there exists a $(N - n + 1) \times (N - n + 1)$ rotation matrix R such that

$$\widehat{M}^* = R^{\mathsf{T}} \widehat{M} R \text{ and } \left\| M - \widehat{M}^* \right\|_F \le \operatorname{Err}(M, \widehat{M}^*),$$
(21)

where

$$\operatorname{Err}(M,\widehat{M}^*) := 2(N-n+1)\sqrt{\widehat{\lambda}_N} \|P^e\|_2 \cdot \delta_a^*(\mathcal{B},\widehat{\mathcal{B}}^*).$$

Remark 3.10. When the length of the concerned cluster is 1, and the concerned eigenvalue is well separated from the other clusters, i.e., $\lambda_{n-1} < \lambda_n < \lambda_{n+1}$ and n = N, the rotation matrix R in Lemma 3.9 becomes R = (1). Therefore, in this case, it is possible to calculate the value of the derivative without the influence of computational instability caused by the closeness of eigenvalues.

4 Computation scheme for the first variation of the eigenvalues λ_2 , λ_3 over triangles

Recall that the second and third Dirichlet eigenvalues coincide over the equilateral triangle, i.e. $\lambda_2^{p_0} = \lambda_3^{p_0}$ for $p_0 = (1, \pi/3)$. Let T^p be a perturbed triangle of the equilateral triangle T^{p_0} . In this section, we will explicitly estimate the range of the difference quotient $\nabla_r^t \lambda_k^p, \nabla_\theta^t \lambda_k^p$ (k = 2, 3), as well as the range of all possible values of the directional derivatives $\nabla_r \lambda_k^p, \nabla_\theta \lambda_k^p$ (k = 2, 3).

4.1 Estimations for the errors δ_a^* and δ_b^*

We formulate the error estimations to bound the error between systems of eigenfunctions and approximate eigenfunctions.

For $p \in C$, Let $\mathcal{B} := (u_2, u_3)$ be a system of L^2 -orthonormal eigenfunctions corresponding to $\lambda_2^p \leq \lambda_3^p$, respectively. Let $E := \operatorname{span}(\mathcal{B})$. Let E^h be a 2-dimensional subspace of $H_0^1(T^p)$. Let

$$\lambda_i^h := \min_{E^{(i)} \subset H_0^1(T^p)} \max_{v \in E^{(i)}} \frac{\|\nabla v\|_{T^p}^2}{\|v\|_{T^p}^2} \quad \text{for } i = 2, 3,$$

where $E^{(i)}$ denotes an arbitrary (i-1)-dimensional subspace of E^h . Let $\delta_b := \delta_b(E, E^h)$.

Lemma 4.1. Suppose that $\delta_b < 1/2$, which is reasonable since δ_b is regarded to be small. Then, there exists an L^2 -orthonormal basis $\mathcal{B}^h := (v_2, v_3)$ of E^h such that

$$\delta_a^*(\mathcal{B}, \mathcal{B}^h) \leq \operatorname{Err}_a^*(\mathcal{B}, \mathcal{B}^h), \quad \delta_b^*(\mathcal{B}, \mathcal{B}^h) \leq \operatorname{Err}_b^*(\mathcal{B}, \mathcal{B}^h),$$

where

$$\operatorname{Err}_{b}^{*}(\mathcal{B}, \mathcal{B}^{h}) := \frac{2\delta_{b}(2-\delta_{b})}{1-2\delta_{b}}, \quad \operatorname{Err}_{a}^{*}(\mathcal{B}, \mathcal{B}^{h}) := \left\{\lambda_{3}^{p} \cdot \operatorname{Err}_{b}^{*}(\mathcal{B}, \mathcal{B}^{h}) + \lambda_{3}^{h} - \lambda_{2}^{p}\right\}^{\frac{1}{2}}$$

Proof. Let $P_h: E \to E^h$ be the orthogonal projector defined by

$$(u - P_h u, v)_{T^p} = 0$$
 for all $u \in E, v \in E^h$.

Let us inductively define orthonormal system $v_2, v_3 \in E^h$ by the following relations:

$$v_2 := P_h u_2 / s_2, \quad s_2 := \|P_h u_2\|_{T^p},$$
$$v_3 := \frac{1}{s_3} (P_h u_3 - (P_h u_3, v_2)_{T^p} v_2), \quad s_3 := \|P_h u_3 - (P_h u_3, v_2)_{T^p} v_2\|_{T^p}.$$

Note that

$$1 - s_2 \le ||u_2 - s_2 v_2|| \le \delta_b, \quad \text{i.e., } 1 - \delta_b \le s_2 \le 1.$$
(22)

By the Schwarz inequality, we have

$$|(P_{h}u_{2}, P_{h}u_{3})_{T^{p}}| = |(P_{h}u_{2} - u_{2}, P_{h}u_{3})_{T^{p}} + (u_{2}, P_{h}u_{3})_{T^{p}}|$$

$$= |(u_{2}, P_{h}u_{3})_{T^{p}}|$$

$$= |(u_{2}, P_{h}u_{3})_{T^{p}} - (P_{h}u_{2}, P_{h}u_{3} - u_{3})_{T^{p}} - (u_{2}, u_{3})_{T^{p}}|$$

$$= |(u_{2} - P_{h}u_{2}, P_{h}u_{3} - u_{3})_{T^{p}}|$$

$$\leq ||u_{2} - P_{h}u_{2}||_{T^{p}} ||P_{h}u_{3} - u_{3}||_{T^{p}} \leq \delta_{b}^{2}.$$
(23)

From the estimations (22) and (23), it follows that

$$|(P_h u_3, v_2)| = |(P_h u_3, P_h u_2 / s_2)| \le \delta_b^2 / (1 - \delta_b).$$
⁽²⁴⁾

Thus, we have

$$1 - s_3 \le ||u_3 - s_3 v_3|| \le ||u_3 - P_h u_3|| + ||P_h u_3 - s_3 v_3|| \le \delta_b + \delta_b^2 / (1 - \delta_b).$$
(25)

By the estimations (22) and (25), we deduce

$$1 - \delta_b - \delta_b^2 / (1 - \delta_b) \le s_i \le 1 \text{ for } i = 2, 3.$$
(26)

Therefore, from the estimations (24) and (26)

$$\begin{aligned} \|u_{i} - v_{i}\|_{T^{p}} &\leq \|u_{i} - s_{i}^{-1}u_{i}\| + \|s_{i}^{-1}u_{i} - s_{i}^{-1}P_{h}u_{i}\| + \|s_{i}^{-1}P_{h}u_{i} - v_{i}\| \\ &\leq s_{i}^{-1}|1 - s_{i}| + s_{i}^{-1}\delta_{b} + s_{i}^{-1}\delta_{b}^{2}/(1 - \delta_{b}) \\ &\leq \frac{2\delta_{b}(2 - \delta_{b})}{1 - 2\delta_{b}} (= \operatorname{Err}_{a}^{*}(\mathcal{B}, \mathcal{B}^{h})). \end{aligned}$$

$$(27)$$

for i = 2, 3. That is,

$$\delta_b^*(\mathcal{B}, \mathcal{B}^h) \leq \operatorname{Err}_b^*(\mathcal{B}, \mathcal{B}^h).$$

By the following fundamental formula (see e.g. [11, page 55]), for i = 2, 3, we have

$$\begin{aligned} \|\nabla u_i - \nabla v_i\|_{T^p}^2 &= \|\nabla u_i\|_{T^p}^2 \|u_i - v_i\|_{T^p}^2 - (\|\nabla u_i\|_{T^p}^2 - \|\nabla v_i\|_{T^p}^2) \|v_i\|_{T^p}^2 \\ &\leq \lambda_3^p \|u_i - v_i\|_{T^p}^2 + \lambda_3^h - \lambda_2^p. \end{aligned}$$
(28)

From the estimations (27) and (28), it follows that

$$\delta_a^*(\mathcal{B}, \mathcal{B}^h) \leq \operatorname{Err}_a^*(\mathcal{B}, \mathcal{B}^h).$$

In the numerical results reported in the rest of this section, the FEM spaces are set up over a uniform triangulation of the triangle domain. Denote by N the subdivision number of the triangulation along the base edge. Below is the detailed setting for the FEM spaces:

$$V_h^{\rm CG}: N = 512, \ {\rm DOF} = 261121; \quad V_h^{\rm CR}: N = 512, \ {\rm DOF} = 785408$$

 \square

Estimation for the difference quotient $\nabla_r^t \lambda_i^p$, $\nabla_\theta^t \lambda_i^p$ (i=2,3)4.2

Let e be either of the normalized vectors e^r or e^{θ} . Utilizing Algorithm 1, we aim to estimate the value of the difference quotient $\nabla_r^t \lambda_i^{p_0}$, $\nabla_{\theta}^t \lambda_i^{p_0}$, (i = 2, 3) for $t \in (0, \varepsilon]$, $\varepsilon = 10^{-7}$. The values of several quantities are also provided in Table 1 and Table 2.

Algorithm 1 Estimation for the value of the difference quotients over $(0, \varepsilon)$

Data: Interval $(0, \varepsilon]$ **Result:** $[\underline{F}_i, \overline{F}_i]$ as the estimation of range of $\nabla_e^t \lambda_i^{p_0}$ over $(0, \varepsilon]$ (i = 2, 3)**Procedure:**

- 1. Evaluate $\lambda_i^{p_{\varepsilon}}$ (i = 1, 2, 3, 4) for triangular domain $T^{p_{\varepsilon}}$ $(p_{\varepsilon} = p_0 + \varepsilon e)$ by Lemma 2.2.
- 2. Evaluate the range of λ_i^p (i = 2, 3, 4) over $(0, \varepsilon]$ by Lemma 3.2.
- 3. Calculate $\hat{u}_2, \hat{u}_3 \in H_0^1(T^p)$ as L^2 -orthonormal approximations to u_2, u_3 and construct an approximation matrix $\widehat{M}_t = (F_{P_t^e}(\hat{u}_i, \hat{u}_j))$ to $M_t^* = (F_{P_t^e}(\tilde{u}_i^*, u_j))$ in (12).
- 4. Estimate upper bounds of the errors $\operatorname{Err}(M_t^*, \widehat{M}_t^*)$ and $\operatorname{Err}(N_t^*, I)$ over $(0, \varepsilon]$ in the estimation (13).
- 5. Estimate an upper bound of the error $\eta(M_t^*, \widehat{M}_t^*, \widehat{N}_t^*)$ over $(0, \varepsilon]$ in the estimation (17).
- 6. Evaluate the range of the eigenvalue μ_k over $(0, \varepsilon]$ as the output $[\underline{F}_i, \overline{F}_i]$ utilizing the estimation (17).

$[\underline{F}_2, \overline{F}_2]$	[59.425, 110.46]	$[\underline{F}_2, \overline{F}_2]$	[12.525, 53.538]
$[\underline{F}_3, \overline{F}_3]$	[135.18, 186.23]	$[\underline{F}_3, \overline{F}_3]$	[88.287, 129.30]
$\hat{\mu}_2$	≈ 84.943	$\hat{\mu}_2$	≈ 33.032
$\hat{\mu}_3$	≈ 160.71	$\hat{\mu}_3$	≈ 108.79
$\eta(M^*_t,\widehat{M}^*_t,\widehat{N}^*_t)$	≤ 25.517	$\eta(M_t^*,\widehat{M}_t^*,\widehat{N}_t^*)$	≤ 20.506
$\operatorname{Err}(M_t^*, \widehat{M}_t^*)$	≤ 25.466	$\operatorname{Err}(M_t^*, \widehat{M}_t^*)$	≤ 20.472
$\operatorname{Err}(N_t^*, I)$	$\leq 1.5658 \cdot 10^{-4}$	$\operatorname{Err}(N_t^*, I)$	$\leq 1.5658 \cdot 10^{-4}$

related quantities

Table 1: The obtained range of $\nabla_r^t \lambda_2^{p_0}$, $\nabla_r^t \lambda_3^{p_0}$ and Table 2: The obtained range of $\nabla_{\theta}^t \lambda_3^{p_0}$, $\nabla_{\theta}^t \lambda_3^{p_0}$ and related quantities

Let us consider the case $e = e^r$. Since we have $\nabla_r^t \lambda_2^{p_0} < \nabla_r^t \lambda_3^{p_0}$, it follows that $\lambda_2^{p_t} < \lambda_3^{p_t}$ for $t \in (0, \varepsilon]$. In case $e = e^{\theta}$, we similarly have $\lambda_2^{p_t} < \lambda_3^{p_t}$ for $t \in (0, \varepsilon]$. The range of λ_4 is estimated utilizing Theorem 2.2 and 3.2. Indeed, we have $\lambda_3^{p_t} \leq 124.078 < 210.04 \leq \lambda_4^{p_t}$ ($t \in (0, \varepsilon]$) for both cases $e = e^r$ and $e = e^{\theta}$. Thus, the eigenvalues $\lambda_4^{p_t} < t \in (0, \varepsilon]$. $\lambda_2^{p_t}, \lambda_3^{p_t}$ are simple for $t \in (0, \varepsilon]$. Note that, from Corollary 3.5, the values of directional derivatives $\nabla_e \lambda_i^{p_t}$ (i = 2, 3)coincide the value of the diagonal elements of the matrix $M = (F_{P^e}(u_i, u_j))$ (i, j = 2, 3).

Estimation for the directional derivatives $\nabla_r \lambda_i^{p_t}, \nabla_\theta \lambda_i^{p_t}$ (i = 2, 3)4.3

Let e be either of the normalized vectors e^r or e^{θ} . For the perturbation $p_{\varepsilon} = p_0 + \varepsilon e \ (\varepsilon = 10^{-7})$, we aim to estimate the range of all possible values of the directional derivative $\nabla_e \lambda_i^{p_t}$ (i=2,3). Recall that, we can evaluate diagonal elements of the matrix $M = (F_{P^e}(u_i, u_j))$ (i, j = 2, 3) utilizing the estimation (21):

$$\left\| M - R^{\mathsf{T}} \widehat{M} R \right\|_F \le \operatorname{Err}(M, \widehat{M}^*),$$

where \widehat{M} is an approximation matrix to M, and R is an 2×2 rotation matrix.

By taking FEM solutions $\hat{u}_2, \hat{u}_3 \in H^1_0(T^{p_{\varepsilon}})$ as L^2 -orthonormal approximations to u_2, u_3 , we can construct an

approximation matrix $\widehat{M} = (F_{P^e}(\hat{u}_i, \hat{u}_j))$. Then, in case $e = e^r$, we obtain

$$\widehat{M} = \begin{pmatrix} 89.1793 & 17.4075\\ 17.4075 & 156.4683 \end{pmatrix}, \operatorname{Err}(M, \widehat{M}^*) \le 20.1898.$$

In case $e = e^{\theta}$, we have

$$\widehat{M} = \begin{pmatrix} 53.5043 & -33.6434 \\ -33.6434 & 88.3205 \end{pmatrix}, \ \operatorname{Err}(M, \widehat{M}^*) \le 16.2295.$$

5 Summary

In this paper, we proposed a method for estimating the range of the first-order variations of eigenvalues, based on eigenvector perturbations with respect to domain deformations and the corresponding error estimations. This approach enables us to establish the simplicity of the second and third Dirichlet eigenvalues over nearly regular triangles. Moreover, we evaluated the range of all possible values for the directional derivatives of clustered eigenvalues.

6 Appendix

Denote by T a triangular domain in \mathbb{R}^2 , by S an invertible linear transform on \mathbb{R}^2 . Let $(\tilde{x}, \tilde{y}) = S(x, y)$ for $(x, y) \in T$, and \tilde{T} the triangle obtained by applying S to T. For v over T, define \tilde{v} over \tilde{T} by $\tilde{v}(\tilde{x}, \tilde{y}) := v(x, y)$. For $V = H_0^1(T)$, let $\tilde{V} := S(V)$ be the space obtained by applying S to the functions of V.

Denote R(T; v) and $R(\widetilde{T}; \widetilde{v})$ by

$$R(T;v) := \frac{\|\nabla v\|_T^2}{\|v\|_T^2}, \quad R(\widetilde{T}; \widetilde{v}) := \frac{\|\widetilde{\nabla} \widetilde{v}\|_{\widetilde{T}}^2}{\|\widetilde{v}\|_{\widetilde{T}}^2}$$

Lemma 6.1. Denote by $\lambda_{\min}(SS^{\intercal})$ and $\lambda_{\max}(SS^{\intercal})$ be the minimum and maximum eigenvalues of SS^{\intercal} , respectively. Then, we have

$$\lambda_{\min}(SS^{\mathsf{T}}) \cdot \lambda_k(T) \le \lambda_k(T) \le \lambda_{\max}(SS^{\mathsf{T}}) \cdot \lambda_k(T).$$

Proof. Since we have $\nabla v = S^{\intercal} \widetilde{\nabla} \widetilde{v}$, it holds that

$$\lambda_{\min}(SS^{\mathsf{T}}) \cdot |\widetilde{\nabla}\widetilde{v}|^2 \le |\nabla v|^2 \le \lambda_{\max}(SS^{\mathsf{T}}) \cdot |\widetilde{\nabla}\widetilde{v}|^2.$$

Therefore, we have

$$\lambda_{\min}(SS^{\mathsf{T}}) \|\nabla \tilde{v}\|_{\tilde{T}}^2 \cdot |\det S^{-1}| \le \|\nabla v\|_T^2 \le \lambda_{\max}(SS^{\mathsf{T}}) \|\nabla \tilde{v}\|_{\tilde{T}}^2 \cdot |\det S^{-1}|.$$

Note that $||v||_T^2 = ||\widetilde{v}||_{\widetilde{T}}^2 \cdot |\det S^{-1}|$. Hence, for any $v \in V(T)$,

)

$$\lambda_{\min}(SS^{\mathsf{T}}) \cdot R(\widetilde{T}; \widetilde{v}) \le R(T; v) \le \lambda_{\max}(SS^{\mathsf{T}}) \cdot R(\widetilde{T}; \widetilde{v})$$

The mapping $S: V(T) \ni v \mapsto \tilde{v} \in V(\tilde{T})$ is injective; see, Theorem 3.41 of [12]. By applying Lemma ?? and the above inequality, we have

$$\operatorname{A_{\min}}(SS^{\mathsf{T}}) \cdot \min_{\widetilde{V}^{k} \subset \widetilde{V}(\widetilde{T})} \max_{\widetilde{v} \in \widetilde{V}^{k}} R(\widetilde{T}; \widetilde{v}) \leq \min_{V^{k} \subset V(T)} \max_{v \in V^{k}} R(T; v),$$

and

$$\min_{V^k \subset V(T)} \max_{v \in V^k} R(T; v) \le \lambda_{\max}(SS^{\mathsf{T}}) \cdot \min_{\widetilde{V}^k \subset \widetilde{V}(\widetilde{T})} \max_{\widetilde{v} \in \widetilde{V}^k} R(\widetilde{T}; \widetilde{v}).$$

where \widetilde{V}^k and V^k are k-dimensional linear subspaces of $\widetilde{V}(\widetilde{T})$ and V(T), respectively. Thus, by the min-max principle, we have

$$\lambda_{\min}(SS^{\intercal}) \cdot \lambda_k(T) \le \lambda_k(T) \le \lambda_{\max}(SS^{\intercal}) \cdot \lambda_k(T).$$

Acknowledgement

The last author is supported by Japan Society for the Promotion of Science: Fund for the Promotion of Joint International Research (Fostering Joint International Research (A)) 20KK0306, Grant-in-Aid for Scientific Research (A) 22H00512, Grant-in-Aid for Scientific Research (B) 20H01820, 21H00998, and Grant-in-Aid for Scientific Research (C) 21K03367.

References

- R. Endo, X. Liu, Shape optimization for the laplacian eigenvalue over triangles and its application to interpolation error analysis, Available at SSRN 4296056.
- [2] X. Liu, T. Vejchodský, Fully computable a posteriori error bounds for eigenfunctions, Numer. Math. (2022) 1–39.
- [3] X. Liu, T. Vejchodský, Projection-based guaranteed l2 error bounds for finite element approximations of laplace eigenfunctions, Journal of Computational and Applied Mathematics 429 (2023) 115164.
- [4] A. Henrot, Extremum problems for eigenvalues of elliptic operators, Springer Science & Business Media, 2006.
- [5] B. Rousselet, Shape design sensitivity of a membrane, J. Optim. Theory Appl. 40 (4) (1983) 595–623.
- [6] X. Liu, F. Kikuchi, Analysis and estimation of error constants for P₀ and P₁ interpolations over triangular finite elements, J. Math. Sci. Univ. Tokyo 17 (2010) 27–78.
- [7] X. Liu, A framework of verified eigenvalue bounds for self-adjoint differential operators, Appl. Math. Comput. 267 (2015) 341–355.
- [8] H. Brönnimann, C. Burnikel, S. Pion, Interval arithmetic yields efficient dynamic filters for computational geometry, in: Proceedings of the fourteenth annual symposium on computational geometry, 1998, pp. 165–174.
- [9] B. J. McCartin, Eigenstructure of the equilateral triangle, part i: The dirichlet problem, Siam Review 45 (2) (2003) 267–287.
- [10] S. Liao, Y. Shu, X. Liu, Optimal estimation for the fujino-morley interpolation error constants, Jpn. J. Ind. Appl. Math. 36 (2019) 521–542.
- [11] D. Boffi, Finite element approximation of eigenvalue problems, Acta Numerica 19 (2010) 1–120.
- [12] R. A. Adams, J. J. Fournier, Sobolev spaces, Elsevier, 2003.