arXiv:2305.15366v1 [hep-ex] 24 May 2023

Weak decays of heavy-quark baryons

Janina Nicolini
On behalf of the LHCb collaboration

TU Dortmund University, Dortmund, Germany
Université Paris-Saclay, Orsay, France

Weak decays of heavy-quark baryons offer an attractive laboratory to search for effects beyond the Standard Model (SM), complementary to searches in meson decays. However, the properties of these baryons are not well understood. In particular, the Ω_b^- baryon is the least studied weakly decaying b baryon. Therefore, the most precise determination of the Ω_b^- baryon mass is reported, using $\Omega_b^- \to J/\psi\Omega^-$ decays at the LHCb experiment. In addition, the relative production rate of the Ω_b^- baryon in pp collisions at the LHC is determined for the first time. These quantities are crucial to predict and measure branching fractions of Ω_b^- baryon decays.

1 Precision measurements of weakly decaying b baryons

Weak decays of heavy-quark baryons provide a complementary laboratory to search for effects beyond the SM and offer rich angular structures compared to meson decays. These are particularly interesting as they allow to test spin dependencies of possible new physics scenarios. The quark model predicts four weakly decaying b baryons with one heavy quark: the lightest one, the $\Lambda_b^0 = |bud\rangle$, the isospin-doublet of $\Xi_b^0 = |bsu\rangle$ and $\Xi_b^- = |bsd\rangle$ and the heaviest one, the $\Omega_b^- = |bss\rangle$. The experimental precision of the measured b baryon properties scales largely with the production fraction f_i of each b hadron and therefore the available statistics. While the probability f_i of a $b\bar{b}$ pair hadronising into an Λ_b^0 baryon is about $f_{\Lambda_b^0} \sim 18\%$ [1], for each Ξ_b baryon the production fraction is only approximately $f_{\Xi_b} \sim 1.5\%$ [1]. For the Ω_b^- baryon, the production fraction is assumed to be $f_{\Omega_b^-} \sim 0.5\%$ [1], but this assumption is based on a measurement by the CDF experiment [2] with only 16 Ω_b^- events. The low expected production probability makes the Ω_b^- difficult to study with respect to other weakly decaying b baryons.
In order to search for effects beyond the SM, the properties of the b baryons have to be known with high precision. The production fractions of these baryons need to be known to measure the absolute branching fractions of their decays. As a first step, the relative production fraction of the Ω_b^- baryon has been determined relatively to the Ξ_b^- baryon for the first time at the Large Hadron Collider (LHC) by the LHCb experiment:

$$R = \frac{f_{\Omega_b^-}}{f_{\Xi_b^-}} \times \frac{\mathcal{B}(\Omega_b^- \to J/\psi\Omega^-)}{\mathcal{B}(\Xi_b^- \to J/\psi\Xi^-)}. \tag{1}$$

These decay modes were chosen as they both provide a decay chain of weakly decaying particles that cannot be mimicked by any mesonic decays, leading to very low background contamination. The two final states of the Ω_b^- and Ξ_b^- decay only differ by one particle since the Ω^- and the

Ξ^- are reconstructed in their decays to, ΛK^-) and $\Lambda\pi^-$, respectively. Given this particular topology, many systematic uncertainties cancel in the relative production fraction ratio R (Eq. 1). In addition, both decays have rather low energy releases making them excellent candidates to measure the Ω_b^- baryon mass through the mass difference with the Ξ_b^- baryon. Since the Ω^- baryon has shorter lifetime compared to that of the Ξ^-, more Ω_b^- decays happen within the LHCb acceptance. Therefore, it has a slightly better detection efficiency compared to the Ξ_b^- baryon.
The measurement [3] to determine the relative production fraction ratio R and the Ω_b^- mass has been performed with a fully cut-based selection in a fiducial phase-space.

2 The Ω_b^- mass measurement

The full Run 1 and Run 2 LHCb dataset corresponding to an integrated luminosity of 9 fb^{-1} is used to determine the Ω_b^- baryon mass. Instead of directly extracting the mass, the mass difference $m(\Omega_b^-) - m(\Xi_b^-)$ is measured. This choice allows to cancel the dominant systematic uncertainty due to the knowledge of the absolute momentum scale. The mass difference is kept as a free parameter and extracted from a simultaneous likelihood fit shown in Fig. 1.




Figure 1 – Invariant-mass distributions in the (top) $\Xi_b^- \to J/\psi\Xi^-$ and (bottom) $\Omega_b^- \to J/\psi\Omega^-$ datasets, in (left) Run 1 and (right) Run 2 data-taking periods used for the mass-difference measurement. (Taken from Ref.[3])

The resulting value is

$$m(\Omega_b^-) - m(\Xi_b^-) = 248.54 \pm 0.51(\text{stat}) \pm 0.38(\text{syst})\ \text{MeV}/c^2. \quad (2)$$

The dominant source for the systematic uncertainty is related to the poor knowledge of the hyperon masses. Together with the most precise determination of the Ξ_b^- baryon mass [4] using the

decay $\Xi_b^- \to \Xi_c^0 \pi^-$ from the LHCb experiment, with a value of $m(\Xi_b^-) = 5797.33 \pm 0.24$ (stat.) $\pm$ 0.29 (syst.) MeV/c^2, the Ω_b^- mass is determined to be

$$m(\Omega_b^-) = 6045.9 \pm 0.5 \text{ (stat.)} \pm 0.6 \text{ (syst.) MeV}/c^2. \quad (3)$$

This measurement provides an improvement of more than a factor of two in precision with respect to the previous best measurement [5].

3 The relative production fraction R

Due to the energy dependence of the production fractions and the limited statistical power of the Run 1 dataset, the relative production fraction R (see Eq. 1) is only determined through a simultaneous likelihood fit to the Run 2 dataset. This sample corresponds to an integrated luminosity of 6 fb^{-1} recorded at a centre-of-mass energy of 13 TeV. The selection is optimised to improve the sensitivity for the production fraction, and the corresponding mass fits are shown in Fig. 2.




Figure 2 – Invariant-mass distributions in the (left) $\Xi_b^- \to J/\psi \Xi^-$ and (right) $\Omega_b^- \to J/\psi \Omega^-$ Run 2 datasets used for the relative production estimation. (Taken from Ref.[3])

The relative production fraction is kept as a free fit parameter and measured to be

$$R = 0.120 \pm 0.008(\text{stat}) \pm 0.008(\text{syst}). \quad (4)$$

The dominant systematic uncertainties are due to simulation calibration linked to the small size of the Ω_b^- dataset and the poor knowledge of the b baryon lifetimes. The measurement is in agreement within 1.2σ with the CDF measurement [6] but the LHCb measurement suggests a larger relative production. In addition to the statistical limitation of the CDF measurement with 16 Ω_b^- events, the larger nominal value can be related to the different production environments at the Tevatron and the LHC, leading to differing production cross-sections.

4 Conclusion

These proceedings present the most precise measurement of the Ω_b^- baryon mass using the LHCb dataset corresponding to an integrated luminosity of 9 fb^{-1}. The measurement agrees with the world average and previous measurements, which are displayed in Fig. 3.

In addition, the first determination of the relative production fraction of the Ω_b^- baryon at the LHC at a centre-of-mass energy of $\sqrt{s} = 13$ TeV was presented

$$\frac{f_{\Omega_b^-}}{f_{\Xi_b^-}} \times \frac{\mathcal{B}(\Omega_b^- \to \Omega_b^- J/\psi)}{\mathcal{B}(\Xi_b^- \to \Xi_b^- J/\psi)} = 0.120 \pm 0.008(\text{stat.}) \pm 0.008(\text{syst.}), \quad (5)$$




Figure 3 – Overview of the Ω_b^- mass measurements to date (red points) [2,5,6,7,8], with corresponding statistical and systematic uncertainties marked in red and blue, respectively, as well as the current world average by the Particle Data Group [9] (light-blue band). The superseded measurements are shown with lighter colours. The new LHCb average is presented in orange, showing the total uncertainty. (Taken from Ref. [3])

using a dataset of 6 fb^{-1}. Further theoretical input on the branching fractions is needed to disentangle the ratio of production fractions from the ratio of branching fractions and enable to perform a first absolute branching fraction measurement of Ω_b^- baryon decays.

Acknowledgments

J.N. acknowledges support from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under grant agreement No. 714536: PRECISION, the German Federal Ministry of Education and Research (BMBF, grant no. 05H21PECL1) within ErUM-FSP T04, and the German Academic Scholarship Foundation.

References

1. Aaij, R. et al. (LHCb collaboration), Measurement of the b-Quark Production Cross Section in 7 and 13 TeV pp Collisions, *Phys. Rev. Lett.* **118**, 052002 (2017).
2. Aaltonen, T. et al. (CDF1 collaboration), Observation of the Ω_b^- baryon and measurement of the properties of the Ξ_b^- and Ω_b^- baryons. *Phys. Rev.* D **80**, 7 (072003)2009.
3. Aaij, R. et al. (LHCb collaboration), Measurement of the mass difference and relative production rate of the Ω_b^- and Ξ_b^- baryons. LHCb-PAPER-2022-053 (in preparation).
4. Aaij, R. et al. (LHCb collaboration), Observation of a new Ξ_b^0 state. *Phys. Rev.* D **103**, 012004 (2021).
5. Aaij, R. et al. (LHCb collaboration), Observation of excited Ω_c^0 baryons in $\Omega_b^- \to \Xi_c^+K^-\pi^-$ decays. *Phys. Rev.* D **104**, L091102 (2021).
6. Aaltonen, T. A. et al. (CDF2 collaboration), Mass and lifetime measurements of bottom and charm baryons in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV. *Phys. Rev.* D **89**, 7 (2014).
7. Aaij, R. et al. (LHCb collaboration), Measurements of the Λ_b^0, Ξ_b^-, and Ω_b^- baryon masses. *Phys. Rev. Lett.* **110**, 182001 (2013).
8. Aaij, R. et al. (LHCb collaboration), Measurements of the mass and lifetime of the Ω_b^- baryon. *Phys. Rev.* D **93**, 092007 (2016).
9. Workman, R. L. et al., PDG - Review of particle physics. *Prog. Theor. Exp. Phys.* **2022**, 8 (2022).