

Tests of general relativity in the nonlinear regime: a parametrized plunge-merger-ringdown waveform model

E. Maggio

Max Planck Institute for Gravitational Physics (Albert Einstein Institute), D-14476 Potsdam, Germany

Gravitational waves (GWs) provide a unique opportunity to test gravity in the dynamical and nonlinear regime. We present a parametrized test of general relativity (GR) that introduces generic deviations to the plunge, merger and ringdown stages of binary–black-hole coalescences. The novel feature of the model is that it can capture signatures of beyond-GR physics in the plunge-merger phase. We use the model to provide constraints on the plunge-merger parameters from the analysis of GW150914. Alarming, we find that GW200129 shows a strong violation of GR. We interpret this result as a false violation of GR either due to waveform systematics (mismodeling of spin precession) or data-quality issues.

1 Parametrized plunge-merger-ringdown model

We use as our baseline model the waveform constructed in the effective-one-body formalism for quasi-circular and spin-aligned binaries, which is calibrated to numerical relativity simulations and contains high-order modes, i.e., `SEOBNRv4HM_PA1,2,3,4,5,6`. The waveform is constructed by attaching the merger-ringdown waveform to the inspiral-plunge waveform at a matching time,

$$h_{\ell m}(t) = h_{\ell m}^{\text{insp-plunge}}(t) \Theta(t_{\text{match}}^{\ell m} - t) + h_{\ell m}^{\text{merger-RD}}(t) \Theta(t - t_{\text{match}}^{\ell m}), \quad (1)$$

where $\Theta(t)$ is the Heaviside function, and the matching time is defined as

$$t_{\text{match}}^{\ell m} = \begin{cases} t_{\text{peak}}^{22}, & (\ell, m) = (2, 2), (3, 3), (2, 1), (4, 4), \\ t_{\text{peak}}^{22} - 10M, & (\ell, m) = (5, 5), \end{cases} \quad (2)$$

where t_{peak}^{22} is the time in which the amplitude of the $(2, 2)$ mode peaks. The amplitude and the orbital frequency peak at different times, therefore we introduce a time-lag parameter $\Delta t_{\ell m}^{\text{GR}} = t_{\text{peak}}^{\Omega} - t_{\text{peak}}^{22}$. We develop a parametrized waveform model that allows for deviations from GR in the plunge-merger-ringdown stage⁷. We introduce fractional deviations to the NR-informed amplitude and frequency at the matching time and the time-lag parameter as,

$$|h_{\ell m}^{\text{NR}}| \rightarrow |h_{\ell m}^{\text{NR}}| (1 + \delta A_{\ell m}), \quad (3)$$

$$\omega_{\ell m}^{\text{NR}} \rightarrow \omega_{\ell m}^{\text{NR}} (1 + \delta \omega_{\ell m}), \quad (4)$$

$$\Delta t_{\ell m}^{\text{GR}} \rightarrow \Delta t_{\ell m}^{\text{GR}} (1 + \delta \Delta t_{\ell m}). \quad (5)$$

We also introduce deformations to the quasi-normal modes following the same strategy applied in Refs.^{8,9} for the oscillation frequency and damping time of the fundamental modes,

$$f_{\ell m 0} \rightarrow f_{\ell m 0} (1 + \delta f_{\ell m 0}),$$

$$\tau_{\ell m 0} \rightarrow \tau_{\ell m 0} (1 + \delta \tau_{\ell m 0}).$$

For simplicity, we assume that the plunge-merger parameters have the same values across different modes (i.e., $\delta A_{\ell m} = \delta A$, $\delta \omega_{\ell m} = \delta \omega$, $\delta \Delta t_{\ell m} = \delta \Delta t$), and the ringdown parameters are nonzero only for the fundamental $(2, 2)$ mode. Figure 1 shows the waveform morphology as a function of time where the plunge-merger parameters vary in the range $[-0.5, 0.5]$ for δA (top panel), $\delta \omega$ (middle panel) and $\delta \Delta t$ (bottom panel). The GR waveform is shown by the black solid line.

2 Bayesian parameter estimation

We analyse the events GW150914¹⁰ and GW200129¹¹, which are among the loudest binary–black-hole signals to date, by varying the binary parameters and a subset of the deviation parameters, i.e., $\{\delta A, \delta \omega\}$,




Figure 1: Time evolution of the GW strain for non-zero values of the plunge-merger parameters.

to perform a “plunge-merger test of GR”. From the analysis of GW150914, we provide constraints on the deviations from GR, i.e., $\delta A = -0.01^{+0.27}_{-0.19}$ and $\delta\omega = 0.00^{+0.17}_{-0.12}$. Deviations in the intermediate region of the IMRPhenom waveform were constrained in the TIGER test¹². From the analysis of GW200129, the inferred value of $\delta\omega$ ($\delta\omega = -0.002^{+0.097}_{-0.082}$) is consistent with GR, however the inferred value of δA ($\delta A = 0.44^{+0.38}_{-0.28}$) exhibits a gross violation of GR. The apparent deviation from GR could be due to systematic errors in the GW modeling or data-quality issues as highlighted by the analyses of Refs.^{13,14}. We explore the former possibility by performing synthetic-data studies where the injected GW signal is generated with spin-precessing waveform models. We conclude that the presence of spin-precession could bias us to find a false evidence for beyond-GR effects when using a nonprecessing non-GR model.

Acknowledgments

We thank Alessandra Buonanno and Hector O. Silva for discussions, and Gregorio Carullo for comments. We acknowledge funding from the Deutsche Forschungsgemeinschaft (DFG) - project number: 386119226. The material presented in this paper is based upon work supported by National Science Foundation’s (NSF) LIGO Laboratory, which is a major facility fully funded by the NSF.

References

1. A. Buonanno and T. Damour, *Phys. Rev. D* **59**, 084006 (1999).
2. A. Buonanno and T. Damour, *Phys. Rev. D* **62**, 064015 (2000).
3. A. Bohé *et al*, *Phys. Rev. D* **95**, 044028 (2017).
4. R. Cotesta *et al*, *Phys. Rev. D* **98**, 084028 (2018).
5. A. Nagar and P. Retegno, *Phys. Rev. D* **99**, 021501 (2019).
6. D. Mihaylov, S. Ossokine, A. Buonanno and A. Ghosh, *Phys. Rev. D* **104**, 124087 (2021).
7. E. Maggio, H. O. Silva, A. Buonanno and A. Ghosh, arXiv:2212.09655 (2022).
8. R. Brito, A. Buonanno and V. Raymond, *Phys. Rev. D* **98**, 084038 (2018).
9. A. Ghosh, R. Brito and A. Buonanno, *Phys. Rev. D* **103**, 124041 (2021).
10. B. P. Abbott *et al*, *Phys. Rev. Lett.* **116**, 061102 (2016).
11. B. P. Abbott *et al*, arXiv:2111.03606 (2021).
12. B. P. Abbott *et al*, *Phys. Rev. Lett.* **116**, 221101 (2016).
13. M. Hannam *et al*, *Nature* **610**, 7933 (2022).
14. E. Payne *et al*, *Phys. Rev. D* **106**, 104017 (2022).