

STRONG DUALITY DATA OF TYPE A AND EXTENDED T -SYSTEMS

KATSUYUKI NAOI

ABSTRACT. The extended T -systems are a number of relations in the Grothendieck ring of the category of finite-dimensional modules over the quantum affine algebras of types $A_n^{(1)}$ and $B_n^{(1)}$, introduced by Mukhin and Young as a generalization of the T -systems. In this paper we establish the extended T -systems for more general modules, which are constructed from an arbitrary strong duality datum of type A . Our approach does not use the theory of q -characters, and so also provides a new proof to the original Mukhin–Young’s extended T -systems.

1. INTRODUCTION

The T -systems are remarkable functional relations appearing in solvable lattice models (see [KNS09] and references therein). Let $\mathcal{C}_{\mathfrak{g}}$ denote the category of integrable finite-dimensional modules over a quantum affine algebra $U'_q(\mathfrak{g})$. It has been proved by Nakajima [Nak03] and Hernandez [Her06, Her10] that the q -characters (or equivalently, the classes in the Grothendieck ring) of Kirillov–Reshetikhin (KR) modules in $\mathcal{C}_{\mathfrak{g}}$ satisfy the T -systems. These T -system relations of KR modules play an essential role in the recent developments of the theory of monoidal categorifications of cluster algebras ([HL10, HL16, KKOP20, KKOP22]).

Snake modules are a relatively large family of simple modules in $\mathcal{C}_{\mathfrak{g}}$ of types $A_n^{(1)}$ and $B_n^{(1)}$ introduced in [MY12b], which contain all minimal affinizations ([Cha95]) of these types. Via the monomial parametrization of simple modules in $\mathcal{C}_{\mathfrak{g}}$ (see [FR99]), each snake module is expressed as $L(\prod_r Y_{i_r, aq^{k_r}})$ with a sequence $((i_1, k_1), \dots, (i_p, k_p)) \in (I_{\mathfrak{g}_0} \times \mathbb{Z})^p$ satisfying some combinatorial conditions, where $I_{\mathfrak{g}_0}$ denotes the index set of the simple Lie subalgebra \mathfrak{g}_0 of \mathfrak{g} .

Mukhin and Young introduced in [MY12a] a number of relations satisfied by the classes of snake modules as generalizations of the T -systems. They call them the *extended T -systems*. Let $L(\prod_{r=1}^p Y_{i_r, aq^{k_r}})$ be a prime snake module (recall that a simple module having no nontrivial tensor factorization is called prime). The corresponding extended T -system is the relation in the Grothendieck ring of the form

$$\left[L\left(\prod_{r=1}^{p-1} Y_{i_r, aq^{k_r}}\right) \right] \left[L\left(\prod_{r=2}^p Y_{i_r, aq^{k_r}}\right) \right] = \left[L\left(\prod_{r=1}^p Y_{i_r, aq^{k_r}}\right) \right] \left[L\left(\prod_{r=2}^{p-1} Y_{i_r, aq^{k_r}}\right) \right] + [M][N],$$

where M and N are other snake modules. In [MY12a], these relations were proved using the formula for the q -characters of snake modules established in [MY12b].

2010 *Mathematics Subject Classification.* 17B37, 81R50, 17B10.

Key words and phrases. quantum affine algebra, T -system, duality datum.

It is a natural problem to find extended T -systems (containing, at least, all the minimal affinizations) in other types (see [MY12a, Appendix A]). Such a family of relations was found for type G_2 in [LM13] and for type C_3 in [Li15], but in general types this is still open as far as the author knows. One essential obstacle is that the q -characters of minimal affinizations in other types are more complicated: they are not thin or special in the terminology of q -characters, while the q -characters of all the snake modules (in types $A_n^{(1)}$ and $B_n^{(1)}$) have these properties. To overcome this difficulty, one possible way is to establish an approach to extended T -systems not relying on the theory of q -characters.

For the T -systems, such an approach was developed in [KKOP22] using a *strong duality datum* and *affine cuspidal modules* introduced in [KKOP23]. Let \mathfrak{g} be a simple Lie algebra of type ADE . A strong duality datum associated with \mathfrak{g} is a family of simple $U'_q(\mathfrak{g})$ -modules $\mathcal{D} = \{\mathbb{L}_i\}_{i \in I_{\mathfrak{g}}} \subseteq \mathcal{C}_{\mathfrak{g}}$, characterized using the invariant \mathfrak{d} introduced in [KKOP20]. Given a strong duality datum \mathcal{D} and a reduced word $\mathbf{i} = (i_1, \dots, i_N)$ of the longest element of the Weyl group of \mathfrak{g} , one can construct the associated affine cuspidal modules $\{S_k^{\mathcal{D}, \mathbf{i}}\}_{k \in \mathbb{Z}} \subseteq \mathcal{C}_{\mathfrak{g}}$. When the pair $(\mathcal{D}, \mathbf{i})$ is associated with a Q -datum ([FO21]), the affine cuspidal modules $\{S_k^{\mathcal{D}, \mathbf{i}}\}_{k \in \mathbb{Z}}$ consist of fundamental modules. In [KKOP22], the authors showed for any pair $(\mathcal{D}, \mathbf{i})$ that *affine determinantal modules*, each of which is constructed as the head of the tensor product of some $S_k^{\mathcal{D}, \mathbf{i}}$'s, satisfy exact sequences corresponding to the T -systems. When $(\mathcal{D}, \mathbf{i})$ is associated with a Q -datum, affine determinantal modules coincide with KR modules, and the classical T -systems are recovered. This result was obtained by applying several properties of strong duality data and affine cuspidal modules, instead of q -characters.

In this paper, by applying a similar approach, we generalize the extended T -systems of type $A_n^{(1)}$ and $B_n^{(1)}$ to more general modules constructed from an arbitrary strong duality datum of type A , which we hope to be a first step toward extended T -systems of general types.

In the first part of this paper, we give a sufficient condition for the head of the tensor product of affine cuspidal modules of general type to satisfy a short exact sequence similar to the T -system. More explicitly, for an arbitrary pair $(\mathcal{D}, \mathbf{i})$ of a strong duality datum and a reduced word and the associated affine cuspidal modules $S_k := S_k^{\mathcal{D}, \mathbf{i}}$ ($k \in \mathbb{Z}$), the following theorem is proved.

Theorem A (Theorem 4.2.1). *Let $\mathbf{k} = (k_1 < \dots < k_p)$ be an increasing sequence of integers with $p \geq 2$. For $1 \leq a < b \leq p$, write $\mathbb{S}_{\mathbf{k}}[a, b] = \text{hd}(S_{k_a} \otimes S_{k_{a+1}} \otimes \dots \otimes S_{k_b})$, and set $\mathbb{S}_{\mathbf{k}} = \mathbb{S}_{\mathbf{k}}[1, p]$. Assume that the following two conditions are satisfied:*

- (a) *for any $1 \leq a < b \leq p$, we have $\mathfrak{d}(S_{k_a}, \mathbb{S}_{\mathbf{k}}[a+1, b]) = 1$, and*
- (b) *for any $1 \leq a < b \leq p$, we have $\mathfrak{d}(\mathbb{S}_{\mathbf{k}}[a, b-1], S_{k_b}) = 1$.*

Then there exists a short exact sequence

$$0 \rightarrow \text{hd} \left(\bigotimes_{a=1}^{p-1} (S_{k_a} \Delta S_{k_{a+1}}) \right) \rightarrow \mathbb{S}_{\mathbf{k}}[1, p-1] \otimes \mathbb{S}_{\mathbf{k}}[2, p] \rightarrow \mathbb{S}_{\mathbf{k}} \otimes \mathbb{S}_{\mathbf{k}}[2, p-1] \rightarrow 0,$$

where $M \Delta N$ denotes the socle of $M \otimes N$. Moreover, the first and third terms are both simple.

We also give sufficient conditions for $\mathbb{S}_{\mathbf{k}}$ to be prime or real (Propositions 4.1.1 and 4.1.3).

In the latter half of this paper, we focus on cuspidal modules $S_k^{\mathcal{D}, \mathbf{i}}$ such that \mathcal{D} is associated with \mathfrak{sl}_{n+1} and \mathbf{i} belongs to either of two special families: *reduced words adapted to a height function* and *adapted to a twisted height function* (see Subsection 5.1 for the definitions). Let \mathbf{i}^{hf} (resp. \mathbf{i}^{tw}) be a reduced word adapted to a height (resp. twisted height) function (these notations are for this introduction only). For an arbitrary strong duality datum \mathcal{D} associated with \mathfrak{sl}_{n+1} and $\mathbf{i} \in \{\mathbf{i}^{\text{hf}}, \mathbf{i}^{\text{tw}}\}$, we define an associated *snake module* by the head of the tensor product of $S_k^{\mathcal{D}, \mathbf{i}}$'s satisfying some conditions. When $(\mathcal{D}, \mathbf{i}^{\text{hf}})$ (resp. $(\mathcal{D}, \mathbf{i}^{\text{tw}})$) is associated with a Q -datum corresponding to $U'_q(\mathfrak{g})$ of type $A_n^{(1)}$ (resp. $B_{n_0}^{(1)}$ with $n = 2n_0 - 1$), these modules coincide with the Mukhin–Young's snake modules (recall that in both cases corresponding Q -data are of type A_n , see [FO21]). We show that these snake modules satisfy short exact sequences of the form in Theorem A, and moreover give more concrete description to the first terms (Theorems 6.2.4 and 7.2.5). When associated with a Q -datum, these recover the Mukhin–Young's extended T -systems. We also show that snake modules are real, and give a necessary and sufficient condition for them to be prime.

We explain our strategy for the proof of the short exact sequences. Given a strong duality datum \mathcal{D} , through the *quantum affine Schur–Weyl duality functor* $\mathcal{F}_{\mathcal{D}}$ ([KKK18]), we can define a crystal structure on a subset of the isomorphism classes of simple modules in $\mathcal{C}_{\mathfrak{g}}$, which is isomorphic to the crystal base of $U_q^-(\mathfrak{g})$. Letting $\varepsilon_i, \varepsilon_i^*$ ($i \in I_{\mathfrak{g}}$) be the functions on this crystal, the conditions (a) and (b) in Theorem A can be rephrased in terms of the values of these functions $\varepsilon_i, \varepsilon_i^*$ (see Proposition 3.3.3 (c)) at certain elements expressed in the Lusztig's parametrization ([Lus90]) with respect to \mathbf{i} . Reineke introduced in [Rei97] a useful algorithm to calculate the values of ε_i and ε_i^* at an element expressed in the Lusztig's parametrization with respect to a reduced word of some type. We can apply this algorithm to the word \mathbf{i}^{hf} , and show that snake modules associated with \mathbf{i}^{hf} satisfy the conditions (a) and (b) of Theorem A. Hence the existence of the short exact sequences in the case $\mathbf{i} = \mathbf{i}^{\text{hf}}$ is shown.

The Reineke's algorithm cannot be applied in the case $\mathbf{i} = \mathbf{i}^{\text{tw}}$. Instead, we give a detailed description of the connection between the Lusztig's parametrizations with respect to \mathbf{i}^{hf} and \mathbf{i}^{tw} . A similar work was previously done in [HO19, Section 12], but our result is more involved: we give a transition formula for all elements corresponding to snake modules (Proposition 7.1.3). Using this formula and the results of the previous case $\mathbf{i} = \mathbf{i}^{\text{hf}}$, we can show the existence of the short exact sequences in the case $\mathbf{i} = \mathbf{i}^{\text{tw}}$ as well.

Note that in [MY12a], the extended T -systems are given in terms of the relations in the Grothendieck ring, and therefore there are two possibilities of short exact sequences. As another advantage of our approach, we can determine which one is correct.

Theorem A holds for a strong duality datum of a general type, not only of type A , and we hope that it will help us to study extended T -systems of other types. One difficulty is that, not in type A , the Reineke's algorithm cannot be applied in full generality for any reduced word. We hope to return this problem in the future.

This paper is organized as follows. In Section 2, we recall basic notions concerning with a simple Lie algebra of type ADE , such as the upper global and dual PBW bases,

and crystals. In Section 3, we recall the basic notions and several properties on quantum affine algebras, the invariants Λ and \mathfrak{d} , and affine cuspidal modules. In Section 4, we give sufficient conditions for the head of the tensor product of affine cuspidal modules to be prime or real, and prove Theorem A. In Section 5 we give the definition of snake modules associated with \mathcal{D} of type A and \mathbf{i}^{hf} or \mathbf{i}^{tw} , and several related notions. In Section 6, we show that snake modules associated with \mathbf{i}^{hf} satisfy the extended T -systems, and at the same time we discuss their reality and primeness. In Section 7, we show analogous assertions for the snake modules associated with \mathbf{i}^{tw} .

2. PRELIMINARIES ON SIMPLE LIE ALGEBRAS OF TYPE ADE

Conventions.

- (i) For a base field \mathbf{k} , we write \otimes for $\otimes_{\mathbf{k}}$ when no confusion is likely.
- (ii) For $a, b \in \mathbb{Z}$ such that $a \leq b$, we denote by $[a, b]$ the set $\{k \in \mathbb{Z} \mid a \leq k \leq b\}$. We set $[a, b] = \emptyset$ if $a > b$.

2.1. Basic notation. Let \mathfrak{g} be a complex simple Lie algebra of type ADE , with an index set I and a Cartan matrix $A = (a_{ij})_{i,j \in I}$. Let α_i ($i \in I$) be the simple roots, R the root system, R^+ the set of positive roots, P the weight lattice, W the Weyl group with simple reflections $\{s_i \mid i \in I\}$, and $w_0 \in W$ the longest element. Denote by $\ell(w)$ for $w \in W$ the length of w , and set $N = \ell(w_0)$. For $i \in I$, define $i^* \in I$ by $w_0(\alpha_i) = -\alpha_{i^*}$. Let

$$R(w_0) = \{\mathbf{i} = (i_1, \dots, i_N) \in I^N \mid s_{i_1} \cdots s_{i_N} = w_0\}$$

denote the set of reduced words of w_0 . For two words \mathbf{i}, \mathbf{i}' , we say \mathbf{i} and \mathbf{i}' are *commutation equivalent* if \mathbf{i}' is obtained from \mathbf{i} by applying a sequence of operations which transform some adjacent components (i, j) such that $a_{ij} = 0$ into (j, i) . An equivalence class for this relation is called a *commutation class*.

2.2. Dual PBW bases and crystals. Let \mathbf{k} be a base field containing $\mathbb{Q}(q)$, and $U_q(\mathfrak{g})$ the quantized enveloping algebra associated with \mathfrak{g} over \mathbf{k} with generators $\{e_i, f_i, q^{h_i} \mid i \in I\}$. Let $U_q^-(\mathfrak{g})$ be the \mathbf{k} -subalgebra of $U_q(\mathfrak{g})$ generated by f_i ($i \in I$), and denote by \mathbf{B}^{up} the *upper global basis* (or *dual canonical basis*) of $U_q^-(\mathfrak{g})$ (see [Kas91]). Let $\mathcal{A}_0 \subseteq \mathbb{Q}(q)$ be the subring of rational functions that are regular at $q = 0$, and set $\mathcal{L} \subseteq U_q^-(\mathfrak{g})$ to be the \mathcal{A}_0 -span of \mathbf{B}^{up} .

We briefly recall dual PBW bases of $U_q^-(\mathfrak{g})$. Let $T_i = T_{i,+}''$ ($i \in I$) denote the algebra automorphism of $U_q(\mathfrak{g})$ given in [Lus93, Chapter 37], and take a reduced word $\mathbf{i} = (i_1, \dots, i_N) \in R(w_0)$. For each $1 \leq k \leq N$, set $\beta_k = s_{i_1} \cdots s_{i_{k-1}}(\alpha_{i_k}) \in R^+$,

$$F_{\text{low}}^{\mathbf{i}}(\beta_k) = T_{i_1} \cdots T_{i_{k-1}}(f_{i_k}), \quad \text{and} \quad F^{\mathbf{i}}(\beta_k) = \frac{F_{\text{low}}^{\mathbf{i}}(\beta_k)}{(F_{\text{low}}^{\mathbf{i}}(\beta_k), F_{\text{low}}^{\mathbf{i}}(\beta_k))}, \quad (2.2.1)$$

where $(\ , \)$ is the bilinear form on $U_q^-(\mathfrak{g})$ given in [Kas91, Section 3.4]. $F^{\mathbf{i}}(\beta_k)$ is called a *dual root vector*. We have $F^{\mathbf{i}}(\beta_k) \in \mathbf{B}^{\text{up}}$ for any \mathbf{i} and k . For $\mathbf{c} = (c_1, \dots, c_N) \in \mathbb{Z}_{\geq 0}^N$, set

$$F^{\mathbf{i}}(\mathbf{c}) = q^{\frac{1}{2} \sum_k c_k(c_k-1)} F^{\mathbf{i}}(\beta_1)^{c_1} \cdots F^{\mathbf{i}}(\beta_N)^{c_N}.$$

Then $\{F^{\mathbf{i}}(\mathbf{c}) \mid \mathbf{c} \in \mathbb{Z}_{\geq 0}^N\}$ forms a basis of $U_q^-(\mathfrak{g})$, and we call this the *dual PBW basis* associated with \mathbf{i} . For each $\mathbf{c} \in \mathbb{Z}_{\geq 0}^N$, there is a unique element $B^{\mathbf{i}}(\mathbf{c}) \in \mathbf{B}^{\text{up}}$ such that

$$F^{\mathbf{i}}(\mathbf{c}) \equiv B^{\mathbf{i}}(\mathbf{c}) \pmod{q\mathcal{L}},$$

see [Kim12, Theorem 4.29].

Let $*$ be the \mathbf{k} -algebra anti-involution on $U_q^-(\mathfrak{g})$ defined by $*f_i = f_i$ ($i \in I$). This $*$ preserves \mathbf{B}^{up} ([Kas93], [Kim12, Lemma 3.5]).

Lemma 2.2.1 ([Lus90, Subsection 2.11]). *For $\mathbf{i} = (i_1, \dots, i_N) \in R(w_0)$ and $\mathbf{c} = (c_1, \dots, c_N) \in \mathbb{Z}_{\geq 0}^N$, set*

$$\mathbf{i}^\vee = (i_N^*, i_{N-1}^*, \dots, i_1^*) \in R(w_0) \quad \text{and} \quad \mathbf{c}^\vee = (c_N, c_{N-1}, \dots, c_1).$$

*Then we have $*F^{\mathbf{i}}(\mathbf{c}) = F^{\mathbf{i}^\vee}(\mathbf{c}^\vee)$ and $*B^{\mathbf{i}}(\mathbf{c}) = B^{\mathbf{i}^\vee}(\mathbf{c}^\vee)$.*

Proposition 2.2.2. *For any $\mathbf{i} \in R(w_0)$ and $\mathbf{c} \in \mathbb{Z}_{\geq 0}^N$, we have*

$$F^{\mathbf{i}}(\mathbf{c}) \in B^{\mathbf{i}}(\mathbf{c}) + \sum_{\mathbf{c}' \prec \mathbf{c}} q\mathbb{Z}[q]B^{\mathbf{i}}(\mathbf{c}'),$$

where \prec is the bi-lexicographic order on $\mathbb{Z}_{\geq 0}^N$, namely, $(a_1, \dots, a_N) \prec (b_1, \dots, b_N)$ if and only if there are $1 \leq k \leq l \leq N$ satisfying $a_k < b_k$, $a_l < b_l$, and $a_j = b_j$ for all j such that $j < k$ or $l < j$.

Proof. Except for the triangularity with respect to the bi-lexicographic order, the assertion follows from [Kim12, Theorem 4.29]. The triangularity is proved by applying $*$ and using Lemma 2.2.1. \square

Proposition 2.2.3 ([Lus90, Subsection 2.3]). *Let $\mathbf{i}, \mathbf{i}' \in R(w_0)$, and $\mathbf{c}, \mathbf{c}' \in \mathbb{Z}_{\geq 0}^N$.*

- (i) *Assume for some $1 \leq k < N$ that $a_{i_k i_{k+1}} = 0$, $i'_k = i_{k+1}$, $i'_{k+1} = i_k$, and $i_l = i'_l$ for all $l \neq k, k+1$. Then $B^{\mathbf{i}}(\mathbf{c}) = B^{\mathbf{i}'}(\mathbf{c}')$ holds if and only if*

$$c'_k = c_{k+1}, \quad c'_{k+1} = c_k, \quad \text{and} \quad c'_l = c_l \quad \text{for all } l \neq k, k+1.$$

- (ii) *Let $i, j \in I$ be such that $a_{ij} = -1$, and assume for some $1 < k < N$ that $(i_{k-1}, i_k, i_{k+1}) = (i, j, i)$, $(i'_{k-1}, i'_k, i'_{k+1}) = (j, i, j)$, and $i_l = i'_l$ for all $l \notin \{k, k \pm 1\}$. Then $B^{\mathbf{i}}(\mathbf{c}) = B^{\mathbf{i}'}(\mathbf{c}')$ holds if and only if $c'_l = c_l$ for all $l \notin \{k, k \pm 1\}$, and*

$$c'_{k-1} = c_k + c_{k+1} - c_0, \quad c'_k = c_0, \quad c'_{k+1} = c_{k-1} + c_k - c_0, \quad (2.2.2)$$

where we set $c_0 = \min(c_{k-1}, c_{k+1})$.

By identifying \mathbf{B}^{up} with its image under the projection $\mathcal{L} \rightarrow \mathcal{L}/q\mathcal{L}$, we define the canonical (abstract) crystal structure on \mathbf{B}^{up} (see [Kas93]). Let

$$\text{wt}: \mathbf{B}^{\text{up}} \rightarrow P, \quad \varepsilon_i, \varphi_i: \mathbf{B}^{\text{up}} \rightarrow \mathbb{Z} \quad \text{and} \quad \tilde{e}_i, \tilde{f}_i: \mathbf{B}^{\text{up}} \rightarrow \mathbf{B}^{\text{up}} \sqcup \{0\} \quad \text{for } i \in I$$

be the maps giving this crystal structure. We also define maps $\varepsilon_i^*, \varphi_i^*, \tilde{e}_i^*, \tilde{f}_i^*$ ($i \in I$) on \mathbf{B}^{up} by

$$\varepsilon_i^* = \varepsilon_i \circ *, \quad \varphi_i^* = \varphi_i \circ *, \quad \tilde{e}_i^* = * \circ \tilde{e}_i \circ *, \quad \tilde{f}_i^* = * \circ \tilde{f}_i \circ *.$$

For $\mathbf{i} = (i_1, \dots, i_N) \in R(w_0)$ and $\mathbf{c} = (c_1, \dots, c_N) \in \mathbb{Z}_{\geq 0}^N$, it follows from [Lus93] and Lemma 2.2.1 that

$$\varepsilon_{i_1}(B^{\mathbf{i}}(\mathbf{c})) = c_1 \quad \text{and} \quad \varepsilon_{i_N}^*(B^{\mathbf{i}}(\mathbf{c})) = c_N. \quad (2.2.3)$$

3. PRELIMINARIES ON QUANTUM AFFINE ALGEBRAS

3.1. Basic notation. Let \mathfrak{g} be an affine Kac–Moody Lie algebra with index set $I_{\mathfrak{g}}$ and simple roots $\{\alpha_i^{\mathfrak{g}} \mid i \in I_{\mathfrak{g}}\}$. Denote by $0 \in I_{\mathfrak{g}}$ the special element prescribed in [Kac90, Section 4], except $A_{2n}^{(2)}$ -type in which we set $\alpha_0^{\mathfrak{g}}$ to be the longest simple root. Let $I_{\mathfrak{g}}^0 = I_{\mathfrak{g}} \setminus \{0\}$, $P^{\mathfrak{g}}$ be the weight lattice of \mathfrak{g} , and $P_{\text{cl}}^{\mathfrak{g}} = P^{\mathfrak{g}} / (P^{\mathfrak{g}} \cap \mathbb{Q}\delta)$, where δ is the indivisible imaginary positive root.

Now we fix the base field \mathbf{k} to be the algebraic closure of $\mathbb{C}(q)$ in $\bigcup_{m>0} \mathbb{C}((q^{1/m}))$, and denote by $U'_q(\mathfrak{g})$ the quantized enveloping algebra over \mathbf{k} associated with \mathfrak{g} with generators $\{e_i, f_i, q^h \mid i \in I_{\mathfrak{g}}, h \in P_{\text{cl}}^{\mathfrak{g}, \vee} := \text{Hom}_{\mathbb{Z}}(P_{\text{cl}}^{\mathfrak{g}}, \mathbb{Z})\}$ (here, by abuse of notation, we use the same symbols with the generators of $U_q(\mathfrak{g})$). We call $U'_q(\mathfrak{g})$ a *quantum affine algebra* in the sequel. Let $\Delta: U'_q(\mathfrak{g}) \rightarrow U'_q(\mathfrak{g}) \otimes U'_q(\mathfrak{g})$ denote the coproduct (we follow the convention in [Kas02, Section 7]).

A $U'_q(\mathfrak{g})$ -module M is said to be *integrable* if $M = \bigoplus_{\lambda \in P_{\text{cl}}^{\mathfrak{g}}} M_{\lambda}$ with $M_{\lambda} = \{v \in M \mid q^h v = q^{\langle h, \lambda \rangle} v \text{ } (h \in P_{\text{cl}}^{\mathfrak{g}, \vee})\}$, and e_i, f_i ($i \in I_{\mathfrak{g}}$) act nilpotently on M . We denote by $\mathcal{C}_{\mathfrak{g}}$ the category of integrable finite-dimensional $U'_q(\mathfrak{g})$ -modules. Let $\mathbf{1} \in \mathcal{C}_{\mathfrak{g}}$ denote the trivial module. For $M, N \in \mathcal{C}_{\mathfrak{g}}$, the tensor product $M \otimes N$ is also an object of $\mathcal{C}_{\mathfrak{g}}$ via the coproduct Δ , and this gives a monoidal category structure on $\mathcal{C}_{\mathfrak{g}}$ with unit object $\mathbf{1}$. Moreover, the monoidal category $\mathcal{C}_{\mathfrak{g}}$ is rigid, namely, every object $M \in \mathcal{C}_{\mathfrak{g}}$ has its right dual $\mathscr{D}M$ and left dual $\mathscr{D}^{-1}M$. There are isomorphisms

$$\begin{aligned} \text{Hom}_{\mathcal{C}_{\mathfrak{g}}}(M \otimes X, Y) &\cong \text{Hom}_{\mathcal{C}_{\mathfrak{g}}}(X, \mathscr{D}M \otimes Y), \quad \text{and} \\ \text{Hom}_{\mathcal{C}_{\mathfrak{g}}}(X \otimes M, Y) &\cong \text{Hom}_{\mathcal{C}_{\mathfrak{g}}}(X, Y \otimes \mathscr{D}^{-1}M), \end{aligned}$$

which are functorial in $X, Y \in \mathcal{C}_{\mathfrak{g}}$.

For simple modules M and N in $\mathcal{C}_{\mathfrak{g}}$, we say that M and N *commute* if $M \otimes N \cong N \otimes M$, and *strongly commute* if $M \otimes N$ is simple. Note that, if M and N strongly commute then they commute, since the Grothendieck ring of $\mathcal{C}_{\mathfrak{g}}$ is commutative [FR99]. We say M is *real* if M strongly commutes with itself.

Proposition 3.1.1 ([KKKO15]).

- (i) Let M_j ($j = 1, 2, 3$) be a module in $\mathcal{C}_{\mathfrak{g}}$, and assume that M_2 is simple. If $f: L \rightarrow M_2 \otimes M_3$ and $g: M_1 \otimes M_2 \rightarrow L'$ are nonzero homomorphisms, then the composition

$$M_1 \otimes L \xrightarrow{M_1 \otimes f} M_1 \otimes M_2 \otimes M_3 \xrightarrow{g \otimes M_3} L' \otimes M_3$$

does not vanish.

- (ii) Let M and N be simple modules in $\mathcal{C}_{\mathfrak{g}}$, and assume that one of them is real. Then both $M \otimes N$ and $N \otimes M$ have simple socles and simple heads.

For $M \in \mathcal{C}_{\mathfrak{g}}$, we denote by $\text{hd}(M)$ (resp. $\text{soc}(M)$) the head (resp. socle) of M . For $M, N \in \mathcal{C}_{\mathfrak{g}}$, we also use the notation $M \nabla N$ (resp. $M \Delta N$) to denote $\text{hd}(M \otimes N)$ (resp. $\text{soc}(M \otimes N)$).

Proposition 3.1.2 ([KKKO15, Corollary 3.14]). Let M, N be simple modules in $\mathcal{C}_{\mathfrak{g}}$, and assume that M is real. Then we have

$$\mathscr{D}^{-1}M \nabla (N \nabla M) \cong N \quad \text{and} \quad (M \nabla N) \nabla \mathscr{D}M \cong N.$$

3.2. R-matrices and invariants. In this subsection we briefly recall the definitions and properties of some invariants on pairs of modules in $\mathcal{C}_{\mathfrak{g}}$, which were introduced in [KKOP20]. For more details, see [KKOP20, KKOP23, KKOP22].

For any simple module M in $\mathcal{C}_{\mathfrak{g}}$, there is unique $\lambda \in P_{\text{cl}}^{\mathfrak{g}}$ such that $M_{\lambda} \neq 0$ and $M_{\mu} = 0$ unless $\mu \in \lambda - \sum_{i \in I_{\mathfrak{g}} \setminus \{0\}} \mathbb{Z}_{\geq 0} \text{cl}(\alpha_i^{\mathfrak{g}})$, where $\text{cl}: P^{\mathfrak{g}} \rightarrow P_{\text{cl}}^{\mathfrak{g}}$ is the canonical projection. We call a nonzero vector $u \in M_{\lambda}$ an ℓ -highest weight vector, which is unique up to a scalar multiplication.

For $M \in \mathcal{C}_{\mathfrak{g}}$ and an indeterminate z , denote by M_z the $U'_q(\mathfrak{g})$ -module $\mathbf{k}[z^{\pm 1}] \otimes M$ defined by

$$e_i(g(z) \otimes u) = z^{\delta_{i0}} g(z) \otimes e_i u, \quad f_i(g(z) \otimes u) = z^{-\delta_{i0}} g(z) \otimes f_i u, \quad q^h(g(z) \otimes u) = g(z) \otimes q^h u$$

for $i \in I_{\mathfrak{g}}$, $h \in P_{\text{cl}}^{\mathfrak{g}, \vee}$, $g(z) \in \mathbf{k}[z^{\pm 1}]$ and $u \in M$. We write $u_z = 1 \otimes u \in M_z$.

For simple modules M, N in $\mathcal{C}_{\mathfrak{g}}$ with ℓ -highest weight vectors $u \in M$ and $v \in N$, there exists a unique $\mathbf{k}(z) \otimes U'_q(\mathfrak{g})$ -module isomorphism

$$R_{M, N_z}^{\text{norm}}: \mathbf{k}(z) \otimes_{\mathbf{k}[z^{\pm 1}]} (M \otimes N_z) \xrightarrow{\sim} \mathbf{k}(z) \otimes_{\mathbf{k}[z^{\pm 1}]} (N_z \otimes M)$$

satisfying $R_{M, N_z}^{\text{norm}}(u \otimes v_z) = v_z \otimes u$ ([Kas02]). R_{M, N_z}^{norm} is called the *normalized R-matrix* of M and N . Let $d_{M, N}(z) \in \mathbf{k}[z]$ be the monic polynomial of the smallest degree satisfying

$$d_{M, N}(z) R_{M, N_z}^{\text{norm}}(M \otimes N_z) \subseteq N_z \otimes M.$$

The polynomial $d_{M, N}(z)$ is called the *denominator* of R_{M, N_z}^{norm} . We denote by $\mathbf{r}_{M, N}$ the specialization at $z = 1$:

$$\mathbf{r}_{M, N} = \left(d_{M, N}(z) R_{M, N_z}^{\text{norm}} \right) \Big|_{z=1} : M \otimes N \rightarrow N \otimes M.$$

We call this nonzero homomorphism $\mathbf{r}_{M, N}$ the *R-matrix* of M and N . If either M or N is real, then the image of $\mathbf{r}_{M, N}$ is isomorphic to $M \nabla N$ and $N \Delta M$ ([KKKO15]), and in particular we have

$$M \nabla N \cong N \Delta M. \quad (3.2.1)$$

Definition 3.2.1 ([KKOP20]). Let M, N be simple modules in $\mathcal{C}_{\mathfrak{g}}$.

- (i) We define $\mathfrak{d}(M, N) \in \mathbb{Z}_{\geq 0}$ by the order of the zero of the polynomial $d_{M, N}(z) d_{N, M}(z)$ at $z = 1$.
- (ii) We define $\Lambda(M, N) \in \mathbb{Z}$ by

$$\Lambda(M, N) = \sum_{k \in \mathbb{Z}_{\geq 0}} (-1)^k \mathfrak{d}(M, \mathcal{D}^k N) - \sum_{k \in \mathbb{Z}_{< 0}} (-1)^k \mathfrak{d}(M, \mathcal{D}^k N).$$

Remark 3.2.2. These definitions are different from [KKOP20, Definitions 3.6 and 3.14], but equivalent to those by [KKOP20, Propositions 3.16 and 3.22]. In [KKOP20], the invariant $\Lambda(M, N)$ are defined for not necessarily simple M and N as well, but we do not need this since in the present paper we will only treat the cases where M and N are simple.

Let us list several properties of the invariants.

Proposition 3.2.3 ([KKOP20, Lemma 3.7 and Corollaries 3.19 and 3.17]). *Let M and N be simple modules in $\mathcal{C}_{\mathfrak{g}}$.*

(i) We have

$$\mathfrak{d}(M, N) = \frac{1}{2}(\Lambda(M, N) + \Lambda(N, M)) = \mathfrak{d}(\mathcal{D}M, \mathcal{D}N).$$

(ii) Assume further that either M or N is real. Then M and N strongly commute if and only if $\mathfrak{d}(M, N) = 0$.

Proposition 3.2.4 ([KKOP20, Proposition 4.2 and Lemma 3.10]). *Let X, Y and Z be simple modules in $\mathcal{C}_{\mathfrak{g}}$.*

(i) For any simple subquotient S of $X \otimes Y$, we have

$$\mathfrak{d}(S, Z) \leq \mathfrak{d}(X, Z) + \mathfrak{d}(Y, Z).$$

(ii) Assume further that X and Y strongly commute. Then we have

$$\mathfrak{d}(X \otimes Y, Z) = \mathfrak{d}(X, Z) + \mathfrak{d}(Y, Z).$$

Lemma 3.2.5. *Let $X, Y, Z \in \mathcal{C}_{\mathfrak{g}}$ be simple modules, and assume that Z is real.*

- (i) If $\mathfrak{d}(X, Z) = \mathfrak{d}(X, \mathcal{D}^{-1}Z) = 0$, then we have $\mathfrak{d}(X, Y \nabla Z) = \mathfrak{d}(X, Y)$.
- (ii) If $\mathfrak{d}(X, Z) = \mathfrak{d}(X, \mathcal{D}Z) = 0$, then we have $\mathfrak{d}(X, Z \nabla Y) = \mathfrak{d}(X, Y)$.

Proof. (i) We have

$$\mathfrak{d}(X, Y \nabla Z) \leq \mathfrak{d}(X, Y) + \mathfrak{d}(X, Z) = \mathfrak{d}(X, Y)$$

by Proposition 3.2.4. On the other hand, we have

$$\mathfrak{d}(X, Y) = \mathfrak{d}(X, \mathcal{D}^{-1}Z \nabla (Y \nabla Z)) \leq \mathfrak{d}(X, \mathcal{D}^{-1}Z) + \mathfrak{d}(X, Y \nabla Z) = \mathfrak{d}(X, Y \nabla Z)$$

by Proposition 3.1.2. Hence the assertion is proved. The proof of (ii) is similar. \square

Proposition 3.2.6 ([KKOP20, Proposition 4.7]). *Let M and N be simple modules in $\mathcal{C}_{\mathfrak{g}}$, and assume that one of them is real and $\mathfrak{d}(M, N) = 1$. Then the composition length of $M \otimes N$ is 2, and we have an exact sequence*

$$0 \rightarrow N \nabla M \rightarrow M \otimes N \rightarrow M \nabla N \rightarrow 0.$$

Proposition 3.2.7 ([KKOP22, Lemma 2.22]). *Let M, N be real simple modules in $\mathcal{C}_{\mathfrak{g}}$ such that $\mathfrak{d}(M, N) \leq 1$. Then $M \nabla N$ is real.*

Proposition 3.2.8 ([KKOP23, Proposition 2.17]). *Let M, N be simple modules in $\mathcal{C}_{\mathfrak{g}}$, and assume that N is real. If $\mathfrak{d}(M, N) > 0$, then we have*

$$\mathfrak{d}(S, N) < \mathfrak{d}(M, N)$$

for any simple subquotient S of $M \otimes N$ and also for any simple subquotient S of $N \otimes M$.

Proposition 3.2.9 ([KKOP22, Proposition 2.25]). *Let X, Y and Z be simple modules in $\mathcal{C}_{\mathfrak{g}}$ such that Y is real. Assume that*

- (i) $\mathfrak{d}(\mathcal{D}X, Y) = \mathfrak{d}(\mathcal{D}Y, Z) = 0$, and
- (ii) $X \otimes Y \otimes Z$ has a simple head.

Then we have

$$\mathfrak{d}(Y, \text{hd}(X \otimes Y \otimes Z)) = \mathfrak{d}(Y, X \nabla Y) + \mathfrak{d}(Y, Y \nabla Z).$$

Following [KKOP20, Definition 4.14], we say a sequence (M_1, \dots, M_r) of real simple modules in $\mathcal{C}_{\mathfrak{g}}$ is a *normal sequence* if the composition of the R-matrices

$$(\mathbf{r}_{M_{r-1}, M_r}) \circ \dots \circ (\mathbf{r}_{M_2, M_r} \circ \dots \circ \mathbf{r}_{M_2, M_3}) \circ (\mathbf{r}_{M_1, M_r} \circ \dots \circ \mathbf{r}_{M_1, M_2}) \\ : M_1 \otimes \dots \otimes M_r \rightarrow M_r \otimes \dots \otimes M_1 \quad (3.2.2)$$

does not vanish.

Proposition 3.2.10 ([KKOP20, Lemma 4.15]). *If (M_1, \dots, M_r) is a normal sequence of real simple modules in $\mathcal{C}_{\mathfrak{g}}$, then $\text{hd}(M_1 \otimes \dots \otimes M_r)$ and $\text{soc}(M_r \otimes \dots \otimes M_1)$ are simple and isomorphic to the image of the composition (3.2.2). Moreover both (M_2, \dots, M_r) and (M_1, \dots, M_{r-1}) are normal sequences, and we have*

$$\Lambda(M_1, \text{hd}(M_2 \otimes \dots \otimes M_r)) = \sum_{k=2}^r \Lambda(M_1, M_k) \quad \text{and} \\ \Lambda(\text{hd}(M_1 \otimes \dots \otimes M_{r-1}), M_r) = \sum_{k=1}^{r-1} \Lambda(M_k, M_r).$$

Proposition 3.2.11 ([KKOP20, Lemma 4.17]). *For real simple modules X, Y and Z in $\mathcal{C}_{\mathfrak{g}}$, the triple (X, Y, Z) is a normal sequence if $\mathcal{D}X$ and Z strongly commute.*

Lemma 3.2.12 ([KKOP23, Lemma 2.24]). *For real simple modules X, Y, Z in $\mathcal{C}_{\mathfrak{g}}$, $\mathfrak{d}(X, Y \nabla Z) = \mathfrak{d}(X, Y) + \mathfrak{d}(X, Z)$ holds if and only if both (X, Y, Z) and (Y, Z, X) are normal sequences.*

Following [KKOP22, Definition 2.16], we say a sequence (M_1, \dots, M_r) of real simple modules in $\mathcal{C}_{\mathfrak{g}}$ is *unmixed* (resp. *strongly unmixed*) if for all $1 \leq j < k \leq r$ we have

$$\mathfrak{d}(\mathcal{D}M_j, M_k) = 0 \quad (\text{resp. } \mathfrak{d}(\mathcal{D}^l M_j, M_k) = 0 \text{ for all } l \in \mathbb{Z}_{>0}).$$

Proposition 3.2.13 ([KKOP23, Lemma 5.3]). *Any unmixed sequence of real simple modules is a normal sequence.*

Lemma 3.2.14 ([KKOP22, Lemma 4.26]). *Let X, Y, Z be real simple modules in $\mathcal{C}_{\mathfrak{g}}$, and assume the following two conditions:*

- (i) (X, Y, Z) is a normal sequence, and
- (ii) $\Lambda(Y, X) + \Lambda(Y, Z) - \Lambda(Y, X \nabla Z) = 2\mathfrak{d}(X, Y)$.

Then the composition

$$X \otimes Y \otimes Z \xrightarrow{\mathbf{r}_{X,Y} \otimes \mathbf{r}_{Y,Z}} Y \otimes X \otimes Z \twoheadrightarrow Y \nabla (X \nabla Z)$$

is surjective and induces an isomorphism $\text{hd}(X \otimes Y \otimes Z) \xrightarrow{\sim} Y \nabla (X \nabla Z)$.

3.3. Strong duality data. Let \mathfrak{g} be a simple Lie algebra of type *ADE* with a Cartan matrix $A = (a_{ij})_{i,j \in I}$ as in Section 2. We freely use the notation in the section. Recall that we mainly use plain symbols such as I, α_i , not for \mathfrak{g} but \mathfrak{g} .

A module $L \in \mathcal{C}_{\mathfrak{g}}$ is called a *root module* if L is a real simple module such that

$$\mathfrak{d}(L, \mathcal{D}^k L) = \delta_{k,-1} + \delta_{k,1} \quad \text{for any } k \in \mathbb{Z},$$

see [KKOP23, Section 3].

Definition 3.3.1 ([KKOP23, Definition 4.7]). Let $\mathcal{D} = \{\mathbf{L}_i\}_{i \in I}$ be a family of simple modules in $\mathcal{C}_{\mathfrak{g}}$.

- (i) We say \mathcal{D} is a *duality datum* associated with \mathfrak{g} if the following conditions are satisfied:
 - (a) \mathbf{L}_i is real for all $i \in I$, and
 - (b) $\mathfrak{d}(\mathbf{L}_i, \mathbf{L}_j) = -a_{ij}$ for all $i, j \in I$ such that $i \neq j$.
- (ii) We say \mathcal{D} is a *strong duality datum* associated with \mathfrak{g} if the following conditions are satisfied:
 - (c) \mathbf{L}_i is a root module for all $i \in I$, and
 - (d) $\mathfrak{d}(\mathbf{L}_i, \mathcal{D}^k \mathbf{L}_j) = -\delta_{k,0} a_{ij}$ for all $k \in \mathbb{Z}$ and $i, j \in I$ such that $i \neq j$.

Let $Q_+ = \sum_{i \in I} \mathbb{Z}_{\geq 0} \alpha_i$, and for $\beta \in Q_+$ let $R(\beta) = R^{\mathfrak{g}}(\beta)$ be a *symmetric quiver Hecke algebra* at β associated with \mathfrak{g} (see [KKK18]). If $\mathcal{D} = \{\mathbf{L}_i\}_{i \in I}$ is a duality datum associated with \mathfrak{g} , then there exists an exact monoidal functor (*quantum affine Schur–Weyl duality functor*)

$$\mathcal{F}_{\mathcal{D}}: \bigoplus_{\beta \in Q_+} R(\beta)\text{-gmod} \rightarrow \mathcal{C}_{\mathfrak{g}},$$

satisfying $\mathcal{F}_{\mathcal{D}}(qM) \cong \mathcal{F}_{\mathcal{D}}(M)$ for any $M \in \bigoplus_{\beta} R(\beta)\text{-gmod}$ and $\mathcal{F}_{\mathcal{D}}(L(i)) = \mathbf{L}_i$ for all $i \in I$ ([KKK18]). Here $\bigoplus_{\beta} R(\beta)\text{-gmod}$ is the direct sum of the categories of finite-dimensional graded $R(\beta)$ -modules equipped with a monoidal structure via the convolution product, q the grading shift of degree 1, and $L(i)$ the 1-dimensional simple module over $R(\alpha_i)$. In the sequel, we write $R\text{-gmod} = \bigoplus_{\beta} R(\beta)\text{-gmod}$. By [KL09, Rou08], there is a unique $\mathbb{Z}[q, q^{-1}]$ -algebra isomorphism $U_{\mathbb{Z}}^-(\mathfrak{g})^{\text{up}} \xrightarrow{\sim} K(R\text{-gmod})$ mapping f_i to $[L(i)]$, where $U_{\mathbb{Z}}^-(\mathfrak{g})^{\text{up}}$ denotes the $\mathbb{Z}[q, q^{-1}]$ -subalgebra of $U_q^-(\mathfrak{g})$ spanned by \mathbf{B}^{up} , and $K(R\text{-gmod})$ the Grothendieck ring of $R\text{-gmod}$. This isomorphism induces a bijection between the upper global basis and the set of the not necessarily degree preserving isomorphism classes of simple modules in $R\text{-gmod}$ ([VV11, Rou12]).

Given a duality datum $\mathcal{D} = \{\mathbf{L}_i\}_{i \in I}$, define a ring homomorphism $\mathcal{L}_{\mathcal{D}}$ from $U_{\mathbb{Z}}^-(\mathfrak{g})^{\text{up}}$ to the Grothendieck ring $K(\mathcal{C}_{\mathfrak{g}})$ by the composition

$$\mathcal{L}_{\mathcal{D}}: U_{\mathbb{Z}}^-(\mathfrak{g})^{\text{up}} \xrightarrow{\sim} K(R\text{-gmod}) \rightarrow K(\mathcal{C}_{\mathfrak{g}}), \quad (3.3.1)$$

where the second one is induced from $\mathcal{F}_{\mathcal{D}}$. By the properties of the isomorphism $U_{\mathbb{Z}}^-(\mathfrak{g})^{\text{up}} \xrightarrow{\sim} K(R\text{-gmod})$ stated above and [KKOP23, Corollary 4.14], we obtain the following lemma (hereafter, we occasionally identify the isomorphism class of a simple module in $\mathcal{C}_{\mathfrak{g}}$ with its class in $K(\mathcal{C}_{\mathfrak{g}})$).

Lemma 3.3.2. *If $\mathcal{D} = \{\mathbf{L}_i\}_{i \in I}$ is a strong duality datum associated with \mathfrak{g} , $\mathcal{L}_{\mathcal{D}}$ induces an injection, which we also denote by $\mathcal{L}_{\mathcal{D}}$, from \mathbf{B}^{up} to the set $\text{Irr}(\mathcal{C}_{\mathfrak{g}})$ of isomorphism classes of simple modules in $\mathcal{C}_{\mathfrak{g}}$.*

The bicrystal structure on \mathbf{B}^{up} is described in terms of $U'_q(\mathfrak{g})$ -modules as follows.

Proposition 3.3.3 ([KP22, Lemma 3.2]). *For a strong duality datum $\mathcal{D} = \{\mathbf{L}_i\}_{i \in I}$, $b \in \mathbf{B}^{\text{up}}$ and $i \in I$, we have*

- (a) $\mathcal{L}_{\mathcal{D}}(\tilde{e}_i b) \cong \mathcal{L}_{\mathcal{D}}(b) \nabla \mathcal{D} \mathbf{L}_i$ if $\varepsilon_i(b) \neq 0$, $\mathcal{L}_{\mathcal{D}}(\tilde{e}_i^* b) \cong \mathcal{D}^{-1} \mathbf{L}_i \nabla \mathcal{L}_{\mathcal{D}}(b)$ if $\varepsilon_i^*(b) \neq 0$,
- (b) $\mathcal{L}_{\mathcal{D}}(\tilde{f}_i b) \cong \mathbf{L}_i \nabla \mathcal{L}_{\mathcal{D}}(b)$, $\mathcal{L}_{\mathcal{D}}(\tilde{f}_i^* b) \cong \mathcal{L}_{\mathcal{D}}(b) \nabla \mathbf{L}_i$,

(c) $\varepsilon_i(b) = \mathfrak{d}(\mathcal{D}\mathbf{L}_i, \mathcal{L}_{\mathcal{D}}(b))$ and $\varepsilon_i^*(b) = \mathfrak{d}(\mathcal{D}^{-1}\mathbf{L}_i, \mathcal{L}_{\mathcal{D}}(b))$.

Assume that $\mathcal{D} = \{\mathbf{L}_i\}_{i \in I}$ is a strong duality datum associated with \mathfrak{g} , and fix a reduced word $\mathbf{i} = (i_1, \dots, i_N) \in R(w_0)$. For each $1 \leq k \leq N$, setting $\beta_k = s_{i_1} \cdots s_{i_{k-1}}(\alpha_{i_k})$, we denote by $S_k = S_k^{\mathcal{D}, \mathbf{i}}$ the simple module $\mathcal{L}_{\mathcal{D}}(F^{\mathbf{i}}(\beta_k)) \in \mathcal{C}_{\mathfrak{g}}$ (see (2.2.1)). Note that we have

$$S_k \cong \mathbf{L}_i \text{ if and only if } \beta_k = \alpha_i \text{ for } 1 \leq k \leq N \text{ and } i \in I. \quad (3.3.2)$$

It follows from the construction that

$$\mathcal{L}_{\mathcal{D}}(F^{\mathbf{i}}(\mathbf{c})) = [S_1^{\otimes c_1} \otimes \cdots \otimes S_N^{\otimes c_N}] \in K(\mathcal{C}_{\mathfrak{g}}) \quad (3.3.3)$$

for any $\mathbf{c} = (c_1, \dots, c_N) \in \mathbb{Z}_{\geq 0}^N$. Moreover, we also have the following.

Lemma 3.3.4. *For any $\mathbf{c} = (c_1, \dots, c_N) \in \mathbb{Z}_{\geq 0}^N$, we have*

$$\mathcal{L}_{\mathcal{D}}(B^{\mathbf{i}}(\mathbf{c})) \cong \text{hd}(S_1^{\otimes c_1} \otimes \cdots \otimes S_N^{\otimes c_N}).$$

Proof. This follows from (3.3.3) and [Kat14, Corollary 4.8]. \square

Following [KKOP23], we extend the above definition of $S_k = S_k^{\mathcal{D}, \mathbf{i}}$ to all $k \in \mathbb{Z}$ by

$$S_{k-N} = \mathcal{D}S_k \text{ for all } k \in \mathbb{Z}. \quad (3.3.4)$$

These modules S_k ($k \in \mathbb{Z}$) are called the *affine cuspidal modules* corresponding to \mathcal{D} and \mathbf{i} .

Remark 3.3.5. Our convention of affine cuspidal modules is different from that of [KKOP23]. Setting $\mathbf{i}^{\vee} = (i_N^*, \dots, i_1^*) \in R(w_0)$, our $S_k^{\mathcal{D}, \mathbf{i}}$ coincides with “ $S_{N+1-k}^{\mathcal{D}, \mathbf{i}^{\vee}}$ ” in [KKOP23, Definition 5.6].

Proposition 3.3.6 ([KKOP23, Propositions 5.7]). *The modules $\{S_k\}_{k \in \mathbb{Z}}$ satisfy the following.*

- (i) S_k is a root module for all $k \in \mathbb{Z}$.
- (ii) For any $a, b \in \mathbb{Z}$ with $a < b$, the pair (S_a, S_b) is strongly unmixed.
- (iii) For any increasing sequence $k_1 < k_2 < \cdots < k_p$ of integers and $(a_1, \dots, a_p) \in \mathbb{Z}_{>0}^p$, $(S_{k_1}^{\otimes a_1}, \dots, S_{k_p}^{\otimes a_p})$ is a normal sequence and $\text{hd}(S_{k_1}^{\otimes a_1} \otimes \cdots \otimes S_{k_p}^{\otimes a_p})$ is simple.

We have $(i_1, \dots, i_N) \in R(w_0)$ if and only if $(i_2, \dots, i_N, i_1^*) \in R(w_0)$. The following lemma is proved from [KKOP23, Propositions 5.9 and 5.10].

Lemma 3.3.7. *Let $\{S_k\}_{k \in \mathbb{Z}}$ be the sequence of affine cuspidal modules corresponding to a strong duality datum \mathcal{D} associated with \mathfrak{g} and $(i_1, \dots, i_N) \in R(w_0)$. We extend i_k to all $k \in \mathbb{Z}$ by*

$$i_{k-N} = i_k^* \text{ for all } k \in \mathbb{Z}.$$

Fix $a \in \mathbb{Z}$, and set $i'_k = i_{a+k}$ and $S'_k = S_{a+k}$ for all $k \in \mathbb{Z}$. Set $\beta'_k = s_{i'_1} \cdots s_{i'_{k-1}}(\alpha_{i'_k})$ for $1 \leq k \leq N$, and let $k(i) \in [1, N]$ ($i \in I$) be the unique integer satisfying $\beta'_{k(i)} = \alpha_i$. Then the family $\mathcal{D}' = \{S'_{k(i)}\}_{i \in I}$ forms a strong duality datum associated with \mathfrak{g} , and $\{S'_k\}_{k \in \mathbb{Z}}$ is the sequence of affine cuspidal modules corresponding to \mathcal{D}' and $(i'_1, \dots, i'_N) \in R(w_0)$.

The following lemma is easily proved from the construction and [Lus93, Subsection 39.2.5].

Lemma 3.3.8. *Let $\{S_k\}_{k \in \mathbb{Z}}$ be the sequence of affine cuspidal modules corresponding to a strong duality datum \mathcal{D} and $\mathbf{i} = (i_1, \dots, i_N) \in R(w_0)$. Assume that $a_{i_l i_{l+1}} = 0$ for some $1 \leq l < N$, and let $\mathbf{i}' \in R(w_0)$ be such that $i'_l = i_{l+1}$, $i'_{l+1} = i_l$ and $i'_k = i_k$ for $k \neq l, l+1$. For $k \in \mathbb{Z}$, set*

$$S'_k = \begin{cases} S_{k+1} & \text{if } k \equiv l \pmod{N}, \\ S_{k-1} & \text{if } k \equiv l+1 \pmod{N}, \\ S_k & \text{otherwise.} \end{cases}$$

Then $\{S'_k\}_{k \in \mathbb{Z}}$ is the affine cuspidal modules corresponding to \mathcal{D} and \mathbf{i}' .

4. PRIMENESS, REALITY, AND SHORT EXACT SEQUENCES

Throughout this section, fix a strong duality datum $\mathcal{D} = \{\mathbf{L}_i\}_{i \in I} \subseteq \mathcal{C}_{\mathfrak{g}}$ associated with \mathfrak{g} , a reduced word $\mathbf{i} = (i_1, \dots, i_N) \in R(w_0)$, and an increasing sequence $\mathbf{k} = (k_1 < k_2 < \dots < k_p)$ of integers. Let $S_k = S_k^{\mathcal{D}, \mathbf{i}}$ ($k \in \mathbb{Z}$) be the affine cuspidal modules corresponding to \mathcal{D} and \mathbf{i} . For $a, b \in [1, p]$ with $a \leq b$, we write

$$\mathbb{S}_{\mathbf{k}}[a, b] = \text{hd}(S_{k_a} \otimes S_{k_{a+1}} \otimes \dots \otimes S_{k_b}),$$

which is simple by Proposition 3.3.6. Set $\mathbb{S}_{\mathbf{k}} = \mathbb{S}_{\mathbf{k}}[1, p]$, and $\mathbb{S}_{\mathbf{k}}[a, b] = \mathbf{1}$ if $a > b$.

4.1. Primeness and reality. A simple module $M \in \mathcal{C}_{\mathfrak{g}}$ is said to be *prime* if any tensor decomposition $M \cong N_1 \otimes N_2$ satisfies $N_1 \cong \mathbf{1}$ or $N_2 \cong \mathbf{1}$.

Proposition 4.1.1. *Assume that one of the following two conditions is satisfied:*

- (i) *For all $1 \leq a < p$, we have $\mathfrak{d}(S_{k_a}, \mathbb{S}_{\mathbf{k}}[a+1, p]) > 0$.*
- (ii) *For all $1 < b \leq p$, we have $\mathfrak{d}(\mathbb{S}_{\mathbf{k}}[1, b-1], S_{k_b}) > 0$.*

Then $\mathbb{S}_{\mathbf{k}}$ is prime.

Proof. We will show the assertion by the induction on p , assuming (ii) (the case (i) is similarly proved). In this proof, we write k for k_p . First assume that $p = 1$ (namely, $\mathbb{S}_{\mathbf{k}} = S_k$), and $M, N \in \mathcal{C}_{\mathfrak{g}}$ satisfies $S_k \cong M \otimes N$. Recall that S_k is a root module by Proposition 3.3.6. Since S_k is real, so are M and N . By Proposition 3.2.4 (ii), we have

$$1 = \mathfrak{d}(\mathcal{D}S_k, S_k) = \mathfrak{d}(\mathcal{D}N \otimes \mathcal{D}M, M \otimes N) \geq \mathfrak{d}(\mathcal{D}M, M) + \mathfrak{d}(\mathcal{D}N, N),$$

which implies that either $\mathfrak{d}(\mathcal{D}M, M) = 0$ or $\mathfrak{d}(\mathcal{D}N, N) = 0$ holds. Hence we have $M \cong \mathbf{1}$ or $N \cong \mathbf{1}$, as required.

Let $p > 1$, and assume that $\mathbb{S}_{\mathbf{k}} \cong M \otimes N$. By Lemma 3.2.5 (ii) and Proposition 3.3.6 (ii), we have

$$\mathfrak{d}(\mathcal{D}^{-1}S_k, \mathbb{S}_{\mathbf{k}}) = \mathfrak{d}(\mathcal{D}^{-1}S_k, S_k) = 1.$$

From this, we see that either $\mathfrak{d}(\mathcal{D}^{-1}S_k, M) = 0$ or $\mathfrak{d}(\mathcal{D}^{-1}S_k, N) = 0$ holds, and we may assume the former. It follows from Proposition 3.1.2 that $\mathcal{D}^{-1}S_k \nabla \mathbb{S}_{\mathbf{k}} \cong \mathbb{S}_{\mathbf{k}}[1, p-1]$. Since $\mathfrak{d}(\mathcal{D}^{-1}S_k \nabla N, M) = 0$, we have

$$\mathbb{S}_{\mathbf{k}}[1, p-1] \cong \mathcal{D}^{-1}S_k \nabla (M \otimes N) \cong M \otimes (\mathcal{D}^{-1}S_k \nabla N).$$

Since $\mathbb{S}_{\mathbf{k}}[1, p-1]$ is prime by the induction hypothesis, we have $M \cong \mathbf{1}$ or $\mathcal{D}^{-1}S_k \nabla N \cong \mathbf{1}$. If the latter occurs, we have $N \cong S_k$ and $M \cong \mathbb{S}_{\mathbf{k}}[1, p-1]$ by Proposition 3.1.2, which contradicts the assumption $\mathfrak{d}(\mathbb{S}_{\mathbf{k}}[1, p-1], S_k) > 0$. Hence $M \cong \mathbf{1}$ holds, and the proof is complete. \square

Lemma 4.1.2. *Let $a, b, c \in [1, p]$ with $a \leq b \leq c$, and assume that*

$$\mathfrak{d}(S_{k_b}, \mathbb{S}_{\mathbf{k}}[a, b-1]) \leq 1 \quad \text{and} \quad \mathfrak{d}(S_{k_b}, \mathbb{S}_{\mathbf{k}}[b+1, c]) \leq 1. \quad (4.1.1)$$

Then S_{k_b} and $\mathbb{S}_{\mathbf{k}}[a, c]$ strongly commute.

Proof. Set

$$X = \mathbb{S}_{\mathbf{k}}[a, b-1], \quad Y = S_{k_b}, \quad Z = \mathbb{S}_{\mathbf{k}}[b+1, c].$$

By Proposition 3.3.6 (ii), we have $\mathfrak{d}(\mathcal{D}X, Y) = 0$ and $\mathfrak{d}(\mathcal{D}Y, Z) = 0$. This and Proposition 3.3.6 (iii) imply that (X, Y, Z) satisfies the assumptions of Proposition 3.2.9, and hence we have

$$\mathfrak{d}(S_{k_b}, \mathbb{S}_{\mathbf{k}}[a, c]) = \mathfrak{d}(Y, \text{hd}(X \otimes Y \otimes Z)) = \mathfrak{d}(Y, X \nabla Y) + \mathfrak{d}(Y, Y \nabla Z).$$

By Proposition 3.2.8, our assumption (4.1.1) implies that $\mathfrak{d}(Y, X \nabla Y) = \mathfrak{d}(Y, Y \nabla Z) = 0$. Hence we have $\mathfrak{d}(S_{k_b}, \mathbb{S}_{\mathbf{k}}[a, c]) = 0$. Now the assertion follows from Proposition 3.2.3 (ii). \square

Proposition 4.1.3. *Let $a, b \in [1, p]$ with $a \leq b$, and assume that*

$$\mathfrak{d}(S_{k_c}, \mathbb{S}_{\mathbf{k}}[a, c-1]) \leq 1 \quad \text{and} \quad \mathfrak{d}(S_{k_c}, \mathbb{S}_{\mathbf{k}}[c+1, b]) \leq 1 \quad (4.1.2)$$

for all $c \in [a, b]$.

- (i) $\mathbb{S}_{\mathbf{k}}[a, b]$ is a real simple module.
- (ii) For any $a', b' \in [a, b]$ with $a' \leq b'$, $\mathbb{S}_{\mathbf{k}}[a, b]$ and $\mathbb{S}_{\mathbf{k}}[a', b']$ strongly commute.

Proof. The assertion (i) is proved inductively using Proposition 3.2.7, and then (ii) follows from Lemma 4.1.2 and Propositions 3.2.3 (ii) and 3.2.4 (i). \square

4.2. A sufficient condition to have a short exact sequence.

Theorem 4.2.1. *Assume that $p \geq 2$, and both of the following two conditions are satisfied:*

- (a) *for any $1 \leq a < b \leq p$, we have $\mathfrak{d}(S_{k_a}, \mathbb{S}_{\mathbf{k}}[a+1, b]) = 1$, and*
- (b) *for any $1 \leq a < b \leq p$, we have $\mathfrak{d}(\mathbb{S}_{\mathbf{k}}[a, b-1], S_{k_b}) = 1$.*

Then there exists a short exact sequence

$$0 \rightarrow \text{hd} \left(\bigotimes_{a=1}^{p-1} (S_{k_a} \Delta S_{k_{a+1}}) \right) \rightarrow \mathbb{S}_{\mathbf{k}}[1, p-1] \otimes \mathbb{S}_{\mathbf{k}}[2, p] \rightarrow \mathbb{S}_{\mathbf{k}} \otimes \mathbb{S}_{\mathbf{k}}[2, p-1] \rightarrow 0, \quad (4.2.1)$$

and the first and third terms are simple.

Our proof goes along a similar line as the one of [KKOP22, Theorem 4.25]. First we note the following lemma.

Lemma 4.2.2. *Let $l, m \in \mathbb{Z}$ be such that $l < m$, and assume that $\mathfrak{d}(S_l, S_m) > 0$. Then we have*

$$S_l \Delta S_m \cong \text{hd} (S_{l+1}^{\otimes c_{l+1}} \otimes S_{l+2}^{\otimes c_{l+2}} \otimes \cdots \otimes S_{m-1}^{\otimes c_{m-1}}) \quad \text{for some } c_{l+1}, \dots, c_{m-1} \in \mathbb{Z}_{\geq 0}.$$

Proof. Since $\mathfrak{d}(S_l, S_m) > 0$, we have $m \leq l + N$ by (3.3.4) and Proposition 3.3.6 (ii). If the equality holds, we have $S_l \Delta S_m \cong \mathbf{1}$, which implies the assertion with $c_j = 0$ for all j . Assume that $m < l + N$. Extend $i_k \in I$ (fixed at the beginning of this section) to all $k \in \mathbb{Z}$ by $i_{k-N} = i_k^*$, and set $\mathbf{i}' = (i_l, \dots, i_{l+N-1}) \in R(w_0)$. By (3.3.3) and Lemmas 3.3.4 and 3.3.7, there is a strong duality datum \mathcal{D}' such that the algebra homomorphism $\mathcal{L}_{\mathcal{D}'} : U_{\mathbb{Z}}^-(\mathfrak{g})^{\text{up}} \rightarrow K(\mathcal{C}_{\mathfrak{g}})$ satisfies

$$\begin{aligned}\mathcal{L}_{\mathcal{D}'}(F^{\mathbf{i}'}(\mathbf{d})) &= [S_l^{\otimes d_l} \otimes \cdots \otimes S_{l+N-1}^{\otimes d_{l+N-1}}], \text{ and} \\ \mathcal{L}_{\mathcal{D}'}(B^{\mathbf{i}'}(\mathbf{d})) &= [\text{hd}(S_l^{\otimes d_l} \otimes \cdots \otimes S_{l+N-1}^{\otimes d_{l+N-1}})]\end{aligned}$$

for all $\mathbf{d} = (d_l, \dots, d_{l+N-1}) \in \mathbb{Z}_{\geq 0}^N$. The assertion is now proved by applying this $\mathcal{L}_{\mathcal{D}'}$ to $F^{\mathbf{i}'}(\mathbf{d})$, where

$$\mathbf{d} = (d_l, \dots, d_{l+N-1}) \quad \text{with} \quad d_r = \delta_{r,l} + \delta_{r,m} \quad (r \in [l, l+N-1]),$$

and using Proposition 2.2.2. \square

We devote the rest of this section to the proof of Theorem 4.2.1. Until the end of the proof, we assume that the sequence $(S_{k_1}, \dots, S_{k_p})$ satisfies the assumptions (a) and (b) of the theorem. Note that then the first and third terms of (4.2.1) are both simple by Lemma 4.2.2 and Proposition 4.1.3.

Lemma 4.2.3.

- (i) We have $\mathfrak{d}(\mathbb{S}_{\mathbf{k}}[1, p-1], \mathbb{S}_{\mathbf{k}}[2, p]) \leq 1$.
- (ii) We have $\mathbb{S}_{\mathbf{k}}[1, p-1] \nabla \mathbb{S}_{\mathbf{k}}[2, p] \cong \mathbb{S}_{\mathbf{k}} \otimes \mathbb{S}_{\mathbf{k}}[2, p-1]$.

Proof. (i) The assertion holds since

$$\mathfrak{d}(\mathbb{S}_{\mathbf{k}}[1, p-1], \mathbb{S}_{\mathbf{k}}[2, p]) \leq \mathfrak{d}(S_{k_1}, \mathbb{S}_{\mathbf{k}}[2, p]) + \mathfrak{d}(\mathbb{S}_{\mathbf{k}}[2, p-1], \mathbb{S}_{\mathbf{k}}[2, p]) = 1$$

by Proposition 4.1.3 and the assumption (a). (ii) There are homomorphisms

$$\mathbb{S}_{\mathbf{k}}[1, p-1] \otimes \mathbb{S}_{\mathbf{k}}[2, p] \hookrightarrow \mathbb{S}_{\mathbf{k}}[1, p-1] \otimes S_{k_p} \otimes \mathbb{S}_{\mathbf{k}}[2, p-1] \twoheadrightarrow \mathbb{S}_{\mathbf{k}} \otimes \mathbb{S}_{\mathbf{k}}[2, p-1]$$

by (3.2.1), and the last term is simple. By Proposition 3.1.1 (i), we obtain a surjection $\mathbb{S}_{\mathbf{k}}[1, p-1] \otimes \mathbb{S}_{\mathbf{k}}[2, p] \twoheadrightarrow \mathbb{S}_{\mathbf{k}} \otimes \mathbb{S}_{\mathbf{k}}[2, p-1]$, and the proof is complete. \square

Lemma 4.2.4. We have $\mathbb{S}_{\mathbf{k}}[2, p] \nabla \mathbb{S}_{\mathbf{k}}[1, p-1] \cong \text{hd}\left(\bigotimes_{a=1}^{p-1} (S_{k_a} \Delta S_{k_{a+1}})\right)$.

Proof. We shall show the assertion by the induction on p . The case $p = 2$ is obvious. Assume that $p > 2$. In this proof, we write k for k_p and k^- for k_{p-1} .

Claim 1. We have $\text{hd}(\mathbb{S}_{\mathbf{k}}[1, p-2] \otimes S_k \otimes S_{k^-}) \cong S_k \nabla \mathbb{S}_{\mathbf{k}}[1, p-1]$.

We apply Lemma 3.2.14 to $X = \mathbb{S}_{\mathbf{k}}[1, p-2]$, $Y = S_k$, and $Z = S_{k^-}$. Since $\mathcal{D}X$ and Z strongly commute, (X, Y, Z) is a normal sequence by Proposition 3.2.11 and the assumption (i) of the lemma follows. The assumption (ii) is equivalent to

$$\Lambda(X, Y) - \Lambda(Y, Z) + \Lambda(Y, X \nabla Z) = 0 \tag{4.2.2}$$

by Proposition 3.2.3 (i). It follows from our assumption (b) that

$$\begin{aligned} 2 &= 2\mathfrak{d}(\mathbb{S}_{\mathbf{k}}[1, p-1], S_k) = \Lambda(X \nabla Z, Y) + \Lambda(Y, X \nabla Z), \quad \text{and} \\ 2 &= 2\mathfrak{d}(S_{k-}, S_k) = \Lambda(Y, Z) + \Lambda(Z, Y), \end{aligned}$$

from which we have $(\Lambda(X \nabla Z, Y) - \Lambda(Z, Y)) - \Lambda(Y, Z) + \Lambda(Y, X \nabla Z) = 0$. Since (X, Z, Y) is a normal sequence, we have $\Lambda(X \nabla Z, Y) = \Lambda(X, Y) + \Lambda(Z, Y)$ by Proposition 3.2.10. Hence (4.2.2) holds, and Claim 1 follows from Lemma 3.2.14.

Therefore, we have the following homomorphisms:

$$\begin{aligned} \mathbb{S}_{\mathbf{k}}[2, p-1] \otimes S_k \otimes \mathbb{S}_{\mathbf{k}}[1, p-1] &\rightarrow \mathbb{S}_{\mathbf{k}}[2, p-1] \otimes (S_k \nabla \mathbb{S}_{\mathbf{k}}[1, p-1]) \\ &\xrightarrow{\sim} \mathbb{S}_{\mathbf{k}}[2, p-1] \otimes \text{hd}(\mathbb{S}_{\mathbf{k}}[1, p-2] \otimes S_k \otimes S_{k-}) \\ &\xrightarrow{\sim} \mathbb{S}_{\mathbf{k}}[2, p-1] \otimes (\mathbb{S}_{\mathbf{k}}[1, p-2] \nabla (S_{k-} \Delta S_k)). \end{aligned} \quad (4.2.3)$$

We see from Lemma 4.2.2 that $\mathcal{D}\mathbb{S}_{\mathbf{k}}[2, p-1]$ and $S_{k-} \Delta S_k$ strongly commute, and thus $(\mathbb{S}_{\mathbf{k}}[2, p-1], \mathbb{S}_{\mathbf{k}}[1, p-2], S_{k-} \Delta S_k)$ is a normal sequence by Proposition 3.2.11. Hence we obtain a surjection

$$\mathbb{S}_{\mathbf{k}}[2, p-1] \otimes S_k \otimes \mathbb{S}_{\mathbf{k}}[1, p-1] \rightarrow (\mathbb{S}_{\mathbf{k}}[2, p-1] \nabla \mathbb{S}_{\mathbf{k}}[1, p-2]) \nabla (S_{k-} \Delta S_k).$$

By the induction hypothesis, we have $\mathbb{S}_{\mathbf{k}}[2, p-1] \nabla \mathbb{S}_{\mathbf{k}}[1, p-2] \cong \text{hd}\left(\bigotimes_{a=1}^{p-2} (S_{k_a} \Delta S_{k_{a+1}})\right)$, and hence we obtain a surjection

$$\mathbb{S}_{\mathbf{k}}[2, p-1] \otimes S_k \otimes \mathbb{S}_{\mathbf{k}}[1, p-1] \rightarrow \text{hd}\left(\bigotimes_{a=1}^{p-1} (S_{k_a} \Delta S_{k_{a+1}})\right). \quad (4.2.4)$$

Since $\mathfrak{d}(\mathbb{S}_{\mathbf{k}}[2, p-1], S_k) = 1$, we have a short exact sequence

$$0 \rightarrow S_k \nabla \mathbb{S}_{\mathbf{k}}[2, p-1] \rightarrow \mathbb{S}_{\mathbf{k}}[2, p-1] \otimes S_k \rightarrow \mathbb{S}_{\mathbf{k}}[2, p] \rightarrow 0 \quad (4.2.5)$$

by Proposition 3.2.6.

Claim 2. $\mathbb{S}_{\mathbf{k}}[1, p-1]$ and $S_k \nabla \mathbb{S}_{\mathbf{k}}[2, p-1]$ strongly commute.

In the proof of this claim, set $X = \mathbb{S}_{\mathbf{k}}[1, p-1]$, $Y = S_k$ and $Z = \mathbb{S}_{\mathbf{k}}[2, p-1]$. We have

$$\mathfrak{d}(X, Y \nabla Z) \leq \mathfrak{d}(X, Y) + \mathfrak{d}(X, Z) = 1 + 0 = 1,$$

and hence it suffices to show that $\mathfrak{d}(X, Y \nabla Z) \neq \mathfrak{d}(X, Y) + \mathfrak{d}(X, Z)$. If the equality holds, then Lemma 3.2.12 implies that (X, Y, Z) is a normal sequence. Hence by Proposition 3.2.10, we have

$$\Lambda(X \nabla Y, Z) = \Lambda(X, Z) + \Lambda(Y, Z).$$

Moreover, (Z, X, Y) is also a normal sequence since $\mathcal{D}Z$ and Y strongly commute, and hence we have

$$\Lambda(Z, X \nabla Y) = \Lambda(Z, X) + \Lambda(Z, Y).$$

On the other hand, since $\mathfrak{d}(X \nabla Y, Z) = 0$ and $\mathfrak{d}(X, Z) = 0$, it follows from Proposition 3.2.3 (i) that

$$\Lambda(X \nabla Y, Z) = -\Lambda(Z, X \nabla Y) \quad \text{and} \quad \Lambda(X, Z) = -\Lambda(Z, X).$$

Now by combining them, we have

$$\begin{aligned} 0 &= \Lambda(X \nabla Y, Z) - \Lambda(X, Z) - \Lambda(Y, Z) = -\Lambda(Z, X \nabla Y) + \Lambda(Z, X) - \Lambda(Y, Z) \\ &= -\Lambda(Z, X) - \Lambda(Z, Y) + \Lambda(Z, X) - \Lambda(Y, Z) \\ &= -\Lambda(Z, Y) - \Lambda(Y, Z) = -2\mathfrak{d}(S_k, \mathbb{S}_k[2, p-1]), \end{aligned}$$

which contradicts the assumption (b). The proof is complete.

Write $\mathbb{H} = \text{hd} \left(\bigotimes_{a=1}^{p-1} (S_{k_a} \Delta S_{k_{a+1}}) \right)$. The simple modules $(S_k \nabla \mathbb{S}_k[2, p-1]) \otimes \mathbb{S}_k[1, p-1]$ and \mathbb{H} are not isomorphic. Indeed, this follows since we have

$$\mathfrak{d}(\mathcal{D}S_{k_1}, (S_k \nabla \mathbb{S}_k[2, p-1]) \otimes \mathbb{S}_k[1, p-1]) \geq \mathfrak{d}(\mathcal{D}S_{k_1}, \mathbb{S}_k[1, p-1]) = 1$$

by Lemma 3.2.5, and, on the other hand, $\mathfrak{d}(\mathcal{D}S_{k_1}, \mathbb{H}) = 0$ by Lemma 4.2.2. Hence the composition

$$(S_k \nabla \mathbb{S}_k[2, p-1]) \otimes \mathbb{S}_k[1, p-1] \hookrightarrow \mathbb{S}_k[2, p-1] \otimes S_k \otimes \mathbb{S}_k[1, p-1] \twoheadrightarrow \mathbb{H}$$

vanishes, where the second homomorphism is (4.2.4). Hence we obtain a surjection $\mathbb{S}_k[2, p] \otimes \mathbb{S}_k[1, p-1] \twoheadrightarrow \mathbb{H}$ by (4.2.5), which completes the proof of the lemma. \square

Proof of Theorem 4.2.1. By Proposition 3.2.6 and Lemmas 4.2.3 and 4.2.4, it suffices to show that $\mathbb{S}_k \otimes \mathbb{S}_k[2, p-1]$ and \mathbb{H} are not isomorphic, which follows from

$$\mathfrak{d}(\mathcal{D}S_{k_1}, \mathbb{S}_k \otimes \mathbb{S}_k[2, p-1]) = 1 \quad \text{and} \quad \mathfrak{d}(\mathcal{D}S_{k_1}, \mathbb{H}) = 0.$$

\square

5. SNAKE MODULES ASSOCIATED WITH A STRONG DUALITY DATUM OF TYPE A

5.1. Quivers and reduced words. *In the remainder of this paper, we assume that $\mathfrak{g} = \mathfrak{sl}_{n+1}$, namely of type A_n , whose index set is $I = [1, n]$ and the Dynkin diagram Δ is given by*

$$\begin{array}{ccccccc} \circ & \text{---} & \circ & \text{---} & \circ & \cdots & \circ & \text{---} & \circ \\ 1 & & 2 & & 3 & & n-1 & & n \end{array}$$

We have

$$N = \frac{n(n+1)}{2},$$

and $i^* = n+1-i$ for $i \in I$. For $i, j \in I$ with $i \leq j$, we write $\alpha_{i,j} = \alpha_i + \alpha_{i+1} + \cdots + \alpha_j \in R^+$. If $i > j$, we set $\alpha_{i,j} = 0$. We still assume that \mathfrak{g} is an arbitrary affine Lie algebra.

Definition 5.1.1.

- (i) A *height function* (or *untwisted height function*) on I is a function $\xi: I \rightarrow \mathbb{Z}$ satisfying

$$|\xi_i - \xi_{i+1}| = 1 \quad \text{for } 1 \leq i < n,$$

where we set $\xi_i = \xi(i)$ for simplicity. We denote by HF the set of height functions.

- (ii) Assume that $n = 2n_0 - 1$ for some $n_0 \in \mathbb{Z}_{\geq 2}$. A *twisted height function* on I is a function $\xi: I \rightarrow \mathbb{Z}/2$ satisfying

$$\begin{aligned} \xi_i \in \mathbb{Z} \text{ for } i \in I \setminus \{n_0\}, \quad |\xi_i - \xi_{i+1}| = 1 \text{ for } i \in I \setminus \{n_0 - 1, n_0, n\}, \\ |\xi_{n_0-1} - \xi_{n_0+1}| = 1 \text{ and } |\xi_{n_0} - \min(\xi_{n_0-1}, \xi_{n_0+1})| = 1/2. \end{aligned}$$

We denote by HF^{tw} the set of twisted height functions.

Remark 5.1.2. In [FO21], a notion of a height function on a pair of a Dynkin diagram and a diagram automorphism was defined. In this terminology, (i) is a height function on (Δ, id) , and (ii) is that on $(\Delta, (\cdot)^*)$, up to conventions.

Here are examples of untwisted and twisted height functions when $n = 5$, where the numbers are the values of each function:

$$\begin{array}{ccccccccc} 2 & 1 & 2 & 3 & 4 & & -1 & 0 & -1/2 & 1 & 2 \\ \circ & \circ & \circ & \circ & \circ & & \circ & \circ & \circ & \circ & \circ \end{array} \quad (5.1.1)$$

Following [FO21], we will define several notions associated with an untwisted or a twisted height function. Let $\xi \in \text{HF}^{\flat}$ with $\flat \in \{\emptyset, \text{tw}\}$, where $\text{HF}^{\emptyset} := \text{HF}$. We say $i \in I$ is a *sink* (resp. *source*) of ξ if $\xi_i < \xi_j$ (resp. $\xi_i - d_i > \xi_j - d_j$) for all $j \in I$ such that $|i - j| = 1$, where we set

$$d_i = 2 \text{ for all } i \in I \text{ if } \xi \in \text{HF}, \text{ and } d_i = \begin{cases} 2 & \text{for } i \in I \setminus \{n_0\} \\ 1 & \text{for } i = n_0 \end{cases} \text{ if } \xi \in \text{HF}^{\text{tw}}.$$


If $i \in I$ is a sink (resp. source) of ξ , we define a new function $s_i \xi \in \text{HF}^{\flat}$ by


$$(s_i \xi)_j = \xi_j + d_i \delta_{i,j} \quad (\text{resp. } (s_i \xi)_j = \xi_j - d_i \delta_{i,j}) \quad \text{for all } j \in I.$$

We say a sequence (i_1, \dots, i_r) of elements of I is *adapted* (or *sink-adapted*) to ξ if i_k is a sink of $s_{i_{k-1}} \cdots s_{i_1} \xi$ for all $k \in [1, r]$. The *repetition quiver* \widehat{Q}^{ξ} associated with ξ is a quiver whose vertex set \widehat{Q}_0^{ξ} and arrow set \widehat{Q}_1^{ξ} are given respectively by

$$\begin{aligned} \widehat{Q}_0^{\xi} &= \{(i, k) \in I \times \mathbb{Z}/2 \mid k - \xi_i \in d_i \mathbb{Z}\}, \\ \widehat{Q}_1^{\xi} &= \{(i, k) \rightarrow (j, l) \mid (i, k), (j, l) \in \widehat{Q}_0^{\xi}, |i - j| = 1, l - k = \min(d_i, d_j)/2\}. \end{aligned}$$

For example, \widehat{Q}^{ξ} for ξ in (5.1.1) are, respectively, as follows:





For $\xi, \xi' \in \text{HF}^b$ with $b \in \{\emptyset, \text{tw}\}$, we have $\widehat{Q}^\xi = \widehat{Q}^{\xi'}$ if $\xi_i - \xi'_i \in 2\mathbb{Z}$ for some (or any) $i \in I$ such that $d_i = 2$. Let $\mathcal{D}_\xi: \widehat{Q}^\xi \rightarrow \widehat{Q}^\xi$ denote the unique quiver automorphism satisfying

$$\mathcal{D}_\xi(i, k) = (i^*, k - \tilde{n}) \quad \text{for all } (i, k) \in \widehat{Q}_0^\xi, \quad \text{where we set } \tilde{n} = \begin{cases} n+1 & \text{if } \xi \in \text{HF}, \\ n & \text{if } \xi \in \text{HF}^{\text{tw}}. \end{cases}$$

We also write \mathcal{D} for \mathcal{D}_ξ when ξ is obvious. Define a partial order \preceq on \widehat{Q}_0^ξ by $(i, k) \preceq (j, l)$ if and only if there is an oriented path from (i, k) to (j, l) in \widehat{Q}^ξ .

For $\xi \in \text{HF}^b$ with $b \in \{\emptyset, \text{tw}\}$, let Γ^ξ denote the full subquiver of \widehat{Q}^ξ whose vertex set Γ_0^ξ is given by

$$\Gamma_0^\xi = \{(i, k) \in \widehat{Q}_0^\xi \mid \xi_i \leq k \leq n-1 + \xi_{i^*}\}.$$

We easily see that the number of the vertices of Γ^ξ is N . We say a total ordering $\Gamma_0^\xi = \{(i_1, k_1), \dots, (i_N, k_N)\}$ of vertices of Γ^ξ is a *compatible reading* of Γ^ξ if $r < s$ holds whenever there is an arrow $(i_r, k_r) \rightarrow (i_s, k_s)$ in Γ^ξ .


Proposition 5.1.3 ([FO21, Section 3]). *Assume that $\xi \in \text{HF}^b$ with $b \in \{\emptyset, \text{tw}\}$.*


- (i) *The set of reduced words of w_0 adapted to ξ forms a single commutation class in $R(w_0)$.*
- (ii) *If $\{(i_1, k_1), \dots, (i_N, k_N)\}$ is a compatible reading of Γ^ξ , then (i_1, \dots, i_N) is a reduced word of w_0 adapted to ξ . Conversely, any reduced word of w_0 adapted to ξ is obtained from a compatible reading of Γ^ξ in this way.*

Let $\xi \in \text{HF}^b$ with $b \in \{\emptyset, \text{tw}\}$, and take a compatible reading $\{(i_1, k_1), \dots, (i_N, k_N)\}$ of Γ^ξ . Define a bijection $\phi_\xi: \Gamma_0^\xi \rightarrow R^+$ by

$$\phi_\xi(i_l, k_l) = \beta_l \quad \text{with } \beta_l = s_{i_1} \cdots s_{i_{l-1}}(\alpha_{i_l}) \quad \text{for } l \in [1, N].$$

Here and below we write $\phi_\xi(i, k)$ instead of $\phi_\xi((i, k))$ for simplicity. The map ϕ_ξ does not depend on the choice of the compatible reading by Proposition 5.1.3. If we take ξ as in (5.1.1), Γ^ξ and ϕ_ξ are given, respectively, as follows:





If ξ is an untwisted height function, Γ^ξ is isomorphic to the Auslander-Reiten quiver of the category of finite-dimensional modules over the path algebra of type A_n associated with ξ (see [Gab80]).

5.2. Snake modules. Let $\mathcal{D} = \{\mathbf{L}_i\}_{i \in I} \subseteq \mathcal{C}_{\mathfrak{g}}$ be a strong duality datum associated with \mathfrak{sl}_{n+1} , and assume that $\xi \in \text{HF}^{\mathfrak{b}}$ with $\mathfrak{b} \in \{\emptyset, \text{tw}\}$. Choose an arbitrary compatible reading $\Gamma_0^\xi = \{(i_1, k_1), \dots, (i_N, k_N)\}$, and set $\mathbf{i} = (i_1, \dots, i_N) \in R(w_0)$. Let $S_k^{\mathcal{D}, \mathbf{i}}$ ($k \in \mathbb{Z}$) be the corresponding affine cuspidal modules. These modules are labeled by \mathbb{Z} , and we shall relabel them by \widehat{Q}_0^ξ as follows; set

$$S_{i_r, k_r}^{\mathcal{D}, \xi} := S_r^{\mathcal{D}, \mathbf{i}} \quad \text{for } r \in [1, N],$$

and extend this to all $(i, k) \in \widehat{Q}_0^\xi$ by $S_{\mathcal{D}_\xi(i, k)}^{\mathcal{D}, \xi} = \mathcal{D}S_{i, k}^{\mathcal{D}, \xi}$. It follows that

$$S_{\mathcal{D}_\xi^t(i_r, k_r)}^{\mathcal{D}, \xi} = S_{r-tN}^{\mathcal{D}, \mathbf{i}} \quad \text{for } 1 \leq r \leq N, t \in \mathbb{Z}. \quad (5.2.1)$$

It is easily seen from Lemma 3.3.8 that these $S_{i, k}^{\mathcal{D}, \xi}$ do not depend on the choice of the compatible reading (though $S_k^{\mathcal{D}, \mathbf{i}}$ do). We will write $S_{i, k}$ or $S_{i, k}^\xi$ for $S_{i, k}^{\mathcal{D}, \xi}$ when no confusion is likely.

Remark 5.2.1. Assume that $\xi \in \text{HF}$ (resp. $\xi \in \text{HF}^{\text{tw}}$) and \mathfrak{g} is of type $A_n^{(1)}$ (resp. $B_{n_0}^{(1)}$). For each $i \in I$, set

$$\mathbf{L}_i = L(Y_{j, q^{-l}}) \quad (\text{resp. } \mathbf{L}_i = L(Y_{\hat{j}, q^{-2l}}) \text{ with } \hat{j} = \min(j, j^*)),$$

where we put $(j, l) = \phi_\xi^{-1}(\alpha_i) \in \Gamma_0^\xi$. Here $L(Y_{j, a})$ denotes the fundamental module with highest weight monomial $Y_{j, a}$, and we refer the readers to [FR99, FM01] for the definition. Then $\mathcal{D} = \{\mathbf{L}_i\}_{i \in I}$ forms a strong duality datum associated with \mathfrak{sl}_{n+1} , and we have

$$S_{i, k}^{\mathcal{D}, \xi} \cong L(Y_{i, q^{-k}}) \quad (\text{resp. } S_{i, k}^{\mathcal{D}, \xi} \cong L(Y_{\hat{i}, q^{-2k}})) \quad \text{for all } (i, k) \in \widehat{Q}_0^\xi,$$

see [HL15, KKK15, FO21, KKOP23] (note that source-adapted reduced words are used by convention in these papers).

Lemma 5.2.2. Let $\mathcal{D} = \{\mathbf{L}_i\}_{i \in I}$ be a strong duality datum associated with \mathfrak{sl}_{n+1} , and assume that $\xi, \xi' \in \text{HF}^{\mathfrak{b}}$ with $\mathfrak{b} \in \{\emptyset, \text{tw}\}$.

- (i) If $\xi' - \xi$ is a constant function whose value is $p \in \mathbb{Z}$, then we have $S_{i, k}^{\mathcal{D}, \xi'} \cong S_{i, k-p}^{\mathcal{D}, \xi}$ for all $(i, k) \in \widehat{Q}_0^{\xi'}$.

- (ii) Assume that $\widehat{Q}^\xi = \widehat{Q}^{\xi'}$, and set $\mathbf{L}'_i = S_{\phi_{\xi'}^{-1}(\alpha_i)}^{\mathcal{D}, \xi}$ for $i \in I$. Then $\mathcal{D}' = \{\mathbf{L}'_i\}_{i \in I}$ forms a strong duality datum associated with \mathfrak{sl}_{n+1} , and we have

$$S_{i,k}^{\mathcal{D}, \xi} \cong S_{i,k}^{\mathcal{D}', \xi'} \quad \text{for all } (i, k) \in \widehat{Q}_0^\xi. \quad (5.2.2)$$

Proof. (i) This is obvious from the construction. (ii) It is easily seen that there is a sequence j_1, j_2, \dots, j_r of elements of I such that j_t is a sink or a source of $s_{j_{t-1}} \cdots s_{j_1} \xi$ for all $1 \leq t \leq r$, and $s_{j_r} \cdots s_{j_1} \xi = \xi'$. Hence it suffices to show the assertion for $\xi' = s_j \xi$, with j being a sink or a source of ξ . We show this for a sink j (the other case is proved similarly). Take a compatible reading $\{(i_1, k_1), \dots, (i_N, k_N)\}$ of Γ^ξ such that $(i_1, k_1) = (j, \xi_j)$, and set

$$(i'_r, k'_r) = (i_{r+1}, k_{r+1}) \quad \text{for } 1 \leq r < N \quad \text{and} \quad (i'_N, k'_N) = \mathcal{D}_\xi^{-1}(i_1, k_1).$$

We easily check that $\{(i'_1, k'_1), \dots, (i'_N, k'_N)\}$ is a compatible reading of $\Gamma^{\xi'}$. Set $\mathbf{i} = (i_1, \dots, i_N)$, $\mathbf{i}' = (i'_1, \dots, i'_N)$, and $S'_k = S_{k+1}^{\mathcal{D}, \mathbf{i}}$ for all $k \in \mathbb{Z}$. For $i \in I$ and $r \in [1, N]$,

$$s_{i'_1} \cdots s_{i'_{r-1}}(\alpha_{i'_r}) = \alpha_i \quad \text{holds if and only if} \quad S_{\phi_{\xi'}^{-1}(\alpha_i)}^{\mathcal{D}, \xi} = S_{i'_r, k'_r}^{\mathcal{D}, \xi} = S'_r.$$

Therefore by Lemma 3.3.7, $\mathcal{D}' = \{S_{\phi_{\xi'}^{-1}(\alpha_i)}^{\mathcal{D}, \xi}\}_{i \in I}$ forms a strong duality datum, and $\{S'_k\}_{k \in \mathbb{Z}}$ are the affine cuspidal modules corresponding to \mathcal{D}' and \mathbf{i}' . Now (5.2.2) for $(i, k) \in \Gamma_0^\xi$ is proved by

$$\begin{aligned} S_{i_1, k_1}^{\mathcal{D}, \xi} &\cong S_1^{\mathcal{D}, \mathbf{i}} \cong S'_0 \cong \mathcal{D}S'_N \cong \mathcal{D}S_{i'_N, k'_N}^{\mathcal{D}', \xi'} \cong S_{i_1, k_1}^{\mathcal{D}', \xi'}, \quad \text{and} \\ S_{i_r, k_r}^{\mathcal{D}, \xi} &\cong S_r^{\mathcal{D}, \mathbf{i}} \cong S'_{r-1} \cong S_{i'_{r-1}, k'_{r-1}}^{\mathcal{D}', \xi'} \cong S_{i_r, k_r}^{\mathcal{D}', \xi'} \quad \text{for } r \in [2, N]. \end{aligned}$$

Then (5.2.2) for general $(i, k) \in \widehat{Q}_0^\xi$ also follows from the construction. \square

Lemma 5.2.3. *Let $\mathcal{D} = \{\mathbf{L}_i\}$ be a strong duality datum associated with \mathfrak{sl}_{n+1} , and ξ a height function or a twisted height function.*

- (a) *Each $S_{i,k}$ is a root module.*
- (b) *Let $(i, k), (i', k') \in \widehat{Q}_0^\xi$ be such that $(i, k) \not\preceq (i', k')$. Then the pair $(S_{i,k}, S_{i',k'})$ is strongly unmixed.*
- (c) *Let $(i_1, k_1), \dots, (i_p, k_p)$ be a sequence of elements of \widehat{Q}_0^ξ , and assume that $(i_r, k_r) \not\preceq (i_s, k_s)$ for all $r, s \in [1, p]$ such that $r < s$. Then for any sequence a_1, \dots, a_p of positive integers, the head of $S_{i_1, k_1}^{\otimes a_1} \otimes \cdots \otimes S_{i_p, k_p}^{\otimes a_p}$ is simple.*
- (d) *If $(i, k), (i', k') \in \widehat{Q}_0^\xi$ are incomparable, then $S_{i,k}$ and $S_{i',k'}$ strongly commute.*

Proof. (a) This follows from Proposition 3.3.6 (i). (b) Let r, r' be the integers satisfying

$$S_{i,k} = S_r^{\mathcal{D}, \mathbf{i}} \quad \text{and} \quad S_{i',k'} = S_{r'}^{\mathcal{D}, \mathbf{i}},$$

where $\mathbf{i} \in R(w_0)$ is an element coming from a compatible reading of Γ^ξ . By replacing (\mathcal{D}, ξ) and \mathbf{i} using Lemma 5.2.2 (ii), if necessary, we may assume $r < r'$. Then the assertion follows from Proposition 3.3.6 (ii). (c) This follows from (b) and Proposition 3.2.13. (d) As above, we may assume that $S_{i,k} = S_r^{\mathcal{D}, \mathbf{i}}$ and $S_{i',k'} = S_{r+1}^{\mathcal{D}, \mathbf{i}}$ for some \mathbf{i} and $r \in \mathbb{Z}$. If $S_{i,k}$ and $S_{i',k'}$ do not strongly commute, it follows from Lemma 4.2.2 that

$S_{i,k} \Delta S_{i',k'} \cong \mathbf{1}$, which implies $S_{i,k} \cong \mathcal{D}S_{i',k'}$, or equivalently $(i', k') = \mathcal{D}^{-1}(i, k)$. This obviously contradicts the assumption, and the assertion is proved. \square

Let \mathcal{D} be a strong duality datum associated with \mathfrak{sl}_{n+1} and ξ an untwisted or a twisted height function. For a sequence $\mathbf{P} = ((i_1, k_1), \dots, (i_p, k_p))$ of elements of \widehat{Q}_0^ξ satisfying $(i_r, k_r) \not\preceq (i_s, k_s)$ for all $1 \leq r < s \leq p$, set


$$\mathbb{S}^{\mathcal{D}, \xi}(\mathbf{P}) = \text{hd}(S_{i_1, k_1}^{\mathcal{D}, \xi} \otimes \dots \otimes S_{i_p, k_p}^{\mathcal{D}, \xi}),$$

which is simple by Lemma 5.2.3. We often write $\mathbb{S}(\mathbf{P})$ for $\mathbb{S}^{\mathcal{D}, \xi}(\mathbf{P})$.

When ξ is a twisted height function, we define four subsets $\widehat{Q}_0^{\xi, \geq}$, $\widehat{Q}_0^{\xi, D}$ and $\widehat{Q}_0^{\xi, U}$ of \widehat{Q}_0^ξ by

$$\begin{aligned} \widehat{Q}_0^{\xi, \geq} &= \{(i, k) \in \widehat{Q}_0^\xi \mid i \geq n_0\}, \quad \widehat{Q}_0^{\xi, D} = \left\{ (n_0, k) \in \widehat{Q}_0^\xi \mid (n_0, k) \rightarrow \left(n_0 + 1, k + \frac{1}{2}\right) \in \widehat{Q}_1^\xi \right\}, \\ \widehat{Q}_0^{\xi, U} &= \left\{ (n_0, k) \in \widehat{Q}_0^\xi \mid (n_0, k) \rightarrow \left(n_0 - 1, k + \frac{1}{2}\right) \in \widehat{Q}_1^\xi \right\}. \end{aligned} \quad (5.2.3)$$

Here “D” (resp. “U”) stands for “downward” (resp. “upward”), the direction of all the arrows incident to the vertices belonging to the subset. When $n = 5$, these are illustrated as follows, where \circ (resp. \heartsuit , \spadesuit , \bullet) belong to $\widehat{Q}_0^{\xi, <}$ (resp. $\widehat{Q}_0^{\xi, U}$, $\widehat{Q}_0^{\xi, D}$, $\widehat{Q}_0^{\xi, >}$):



Note that $\mathcal{D}: \widehat{Q}^\xi \rightarrow \widehat{Q}^\xi$ maps $\widehat{Q}_0^{\xi, <}$ (resp. $\widehat{Q}_0^{\xi, D}$) bijectively onto $\widehat{Q}_0^{\xi, >}$ (resp. $\widehat{Q}_0^{\xi, U}$), and vice versa.

Definition 5.2.4 ([MY12b]).

- (1) Assume that ξ is a height function, and $(i, k), (i', k') \in \widehat{Q}_0^\xi$.
 - (i) We say (i', k') is *in snake position* with respect to (i, k) if $(i, k + 2) \preceq (i', k')$.
 - (ii) We say (i', k') is *in prime snake position* with respect to (i, k) if


$$(i, k + 2) \preceq (i', k') \preceq \mathcal{D}^{-1}(i, k).$$
- (2) Assume that ξ is a twisted height function, and $(i, k), (i', k') \in \widehat{Q}_0^\xi$.
 - (i) We say (i', k') is *in snake position* with respect to (i, k) if $(i, k + 2 - \delta_{i, n_0}) \preceq (i', k')$,

$$(i', k') \in \widehat{Q}_0^{\xi, <} \sqcup \widehat{Q}_0^{\xi, D} \quad \text{when } (i, k) \in \widehat{Q}_0^{\xi, <} \sqcup \widehat{Q}_0^{\xi, U}, \quad \text{and}$$

$$(i', k') \in \widehat{Q}_0^{\xi, >} \sqcup \widehat{Q}_0^{\xi, U} \quad \text{when } (i, k) \in \widehat{Q}_0^{\xi, >} \sqcup \widehat{Q}_0^{\xi, D}.$$
 - (ii) We say (i', k') is *in prime snake position* with respect to (i, k) if (i', k') is in snake position with respect to (i, k) , and $(i', k') \preceq \mathcal{D}^{-1}(i, k)$.

When we would like to emphasize that ξ is untwisted (resp. twisted), we say (i', k') is in snake position *of untwisted type* (resp. *of twisted type*) with respect to (i, k) .

When $n = 5$ these are illustrated as follows, where \bullet and \circ are in snake position with respect to \star , and \bullet are in prime snake position with respect to \star .



The following definition (with (\mathcal{D}, ξ) taken as in Remark 5.2.1) was introduced in [MY12b].

Definition 5.2.5. Let ξ be a height function or a twisted height function, and $\mathbf{P} = ((i_1, k_1), \dots, (i_p, k_p))$ a sequence of elements of \widehat{Q}_0^ξ .

- (i) We say \mathbf{P} is a *snake* (resp. *prime snake*) if (i_{s+1}, k_{s+1}) is in snake position (resp. in prime snake position) with respect to (i_s, k_s) for all $1 \leq s < p$. We also say \mathbf{P} is a snake of *untwisted* or *twisted type*, when we would like to emphasize the type of ξ . For a subset Ω of \widehat{Q}_0^ξ , we say \mathbf{P} is a snake in Ω if \mathbf{P} is a snake and all the elements of \mathbf{P} belong to Ω .
- (ii) Let \mathcal{D} be a strong duality datum associated with \mathfrak{sl}_{n+1} . If \mathbf{P} is a snake, we call $\mathbb{S}(\mathbf{P}) = \mathbb{S}^{\mathcal{D}, \xi}(\mathbf{P})$ a *snake module* (of *untwisted* or *twisted type*) associated with \mathcal{D} and ξ .

Lemma 5.2.6. If $\mathbf{P} = ((i_1, k_1), \dots, (i_p, k_p))$ is a snake, then so are

$$\mathcal{D}^{\pm 1} \mathbf{P} = (\mathcal{D}^{\pm 1}(i_1, k_1), \dots, \mathcal{D}^{\pm 1}(i_p, k_p)),$$

and we have $\mathcal{D}^{\pm 1} \mathbb{S}(\mathbf{P}) \cong \mathbb{S}(\mathcal{D}^{\pm 1} \mathbf{P})$. Moreover, if \mathbf{P} is prime then so are $\mathcal{D}^{\pm 1} \mathbf{P}$.

Proof. The first and last assertions are easily checked from the definition, and the second follows from Proposition 3.2.10, since

$$\begin{aligned} \mathcal{D}^{\pm 1} \mathbb{S}(\mathbf{P}) &\cong \text{soc}(S_{\mathcal{D}^{\pm 1}(i_p, k_p)} \otimes \cdots \otimes S_{\mathcal{D}^{\pm 1}(i_1, k_1)}) \\ &\cong \text{hd}(S_{\mathcal{D}^{\pm 1}(i_1, k_1)} \otimes \cdots \otimes S_{\mathcal{D}^{\pm 1}(i_p, k_p)}) = \mathbb{S}(\mathcal{D}^{\pm 1} \mathbf{P}). \end{aligned}$$

□

For a sequence $\mathbf{P} = ((i_1, k_1), \dots, (i_p, k_p))$ of elements of \widehat{Q}_0^ξ and $a, b \in [1, p]$ with $a \leq b$, set $\mathbf{P}_{[a, b]} = ((i_a, k_a), (i_{a+1}, k_{a+1}), \dots, (i_b, k_b))$.

Lemma 5.2.7.

- (i) If $(i, k), (i', k') \in \widehat{Q}_0^\xi$ satisfy $\mathcal{D}^{-1}(i, k) \not\subseteq (i', k')$, then $\mathcal{D}^\ell S_{i,k}$ and $S_{i',k'}$ strongly commute for all $\ell \in \mathbb{Z}_{\geq 0}$.
- (ii) Let $\mathbf{P} = ((i_1, k_1), \dots, (i_p, k_p)) \in (\widehat{Q}_0^\xi)^p$ be a snake. If (i_{r+1}, k_{r+1}) is not in prime snake position with respect to (i_r, k_r) for some r , then we have

$$\mathbb{S}(\mathbf{P}) \cong \mathbb{S}(\mathbf{P}_{[1,r]}) \otimes \mathbb{S}(\mathbf{P}_{[r+1,p]}).$$

Proof. Since $\mathcal{D}^{\ell+1} S_{\mathcal{D}^{-1}(i,k)} \cong \mathcal{D}^\ell S_{i,k}$, the assertion (i) follows from Lemma 5.2.3 (b). The assertion (ii) easily follows from (i). \square

Later, we will prove that $\mathbb{S}(\mathbf{P})$ is prime if \mathbf{P} is a prime snake (Theorems 6.2.4 (ii) and 7.2.5 (ii)).


6. THE CASE OF UNTWISTED HEIGHT FUNCTIONS

6.1. Reineke's algorithm. In this section, we focus on (untwisted) height functions. We will prove that, if ξ is a height function and \mathbf{P} is a prime snake in \widehat{Q}_0^ξ , the module $\mathbb{S}(\mathbf{P})$ satisfies the assumptions of Theorem 4.2.1, using Proposition 3.3.3 (c). For that, we need to calculate $\varepsilon_i(b)$ and $\varepsilon_i^*(b)$ for some $b \in \mathbf{B}^{\text{up}}$. An algorithm for this was introduced by Reineke in [Rei97], which we recall in this subsection.

For $i \in I$, let $\bar{i} \in \{0, 1\}$ be such that $\bar{i} \equiv i \pmod{2}$. For $\delta \in \{0, 1\}$, let $\xi^{(\delta)} \in \text{HF}$ denote the unique height function satisfying $\xi_i^{(\delta)} \in \{0, 1\}$ for all $i \in I$ and $\xi_1^{(\delta)} = \delta$. We have $\xi_i^{(\bar{i})} = 1$ for all $i \in I$. We write $\Gamma^{(\delta)}$ for $\Gamma^{\xi^{(\delta)}}$, and $\phi_{(\delta)}$ for $\phi_{\xi^{(\delta)}}: \Gamma_0^{(\delta)} \rightarrow R^+$. For each $i \in I$, define a subset $\Omega_i \subseteq \Gamma_0^{(\bar{i})}$ by

$$\Omega_i = \{(j, k) \in \Gamma_0^{(\bar{i})} \mid (i, 1) \preceq (j, k) \preceq (i^*, n)\} = \{\phi_{(\bar{i})}^{-1}(\alpha_{j,l}) \mid 1 \leq j \leq i \leq l \leq n\}.$$

When $n = 5$, $\Gamma^{(\delta)}$ and $\phi_{(\delta)}$ are as follows:



Here the vertices belonging to Ω_2 in $\Gamma_0^{(0)}$ and Ω_3 in $\Gamma_0^{(1)}$ are boxed. The proof of the following lemma is straightforward.

Lemma 6.1.1. *Let $\delta \in \{0, 1\}$. For $(i, k) \in \Gamma_0^{(\delta)}$, we have $\phi_{(\delta)}(i, k) = \alpha_{x,y}$, where*

$$x = \begin{cases} i - k & \text{if } i - k > 0, \\ k - i + 1 & \text{if } i - k \leq 0, \end{cases} \quad \text{and} \quad y = \begin{cases} i + k & \text{if } i + k \leq n, \\ 2n + 1 - i - k & \text{if } i + k > n. \end{cases}$$

Let $\delta \in \{0, 1\}$. Take a compatible reading $\{(i_1, k_1), \dots, (i_N, k_N)\}$ of $\Gamma^{(\delta)}$, and set $\mathbf{i} = (i_1, \dots, i_N) \in R(w_0)$. For a $\Gamma_0^{(\delta)}$ -tuple $\mathbf{c} = (c_{i,k})_{(i,k) \in \Gamma_0^{(\delta)}}$ of nonnegative integers, we set

$$B^{(\delta)}(\mathbf{c}) := B^{\mathbf{i}}(\mathbf{c}) \in \mathbf{B}^{\text{up}}, \quad (6.1.1)$$

where in the right-hand side \mathbf{c} is regarded as an element of $\mathbb{Z}_{\geq 0}^N$ via the bijection $[1, N] \rightarrow \Gamma_0^{(\delta)} : r \mapsto (i_r, k_r)$. We easily see from Propositions 2.2.3 (i) and 5.1.3 that $B^{(\delta)}(\mathbf{c})$ does not depend on the choice of the compatible reading. If $\bar{i} \neq \delta$, it follows from (2.2.3) that $\varepsilon_i(B^{(\delta)}(\mathbf{c})) = c_{i,0}$.

The other case is described as follows. For $i \in I$, let \mathcal{U}_i be the set of lower closed subsets of Ω_i . That is, a subset $\Sigma \subseteq \Omega_i$ belongs to \mathcal{U}_i if and only if for any $P, Q \in \Omega_i$, $P \in \Sigma$ and $Q \preceq P$ imply $Q \in \Sigma$.

Theorem 6.1.2 ([Rei97, Theorem 7.1]). *Let $j \in I$, and set $\delta = \bar{j} \in \{0, 1\}$. For any $\mathbf{c} = (c_{i,k}) \in \mathbb{Z}_{\geq 0}^{\Gamma_0^{(\delta)}}$, we have*

$$\varepsilon_j(B^{(\delta)}(\mathbf{c})) = \max_{\Sigma \in \mathcal{U}_j} \left(\sum_{(i,k) \in \Sigma} (c_{i,k} - c_{i,k-2}) \right),$$

where we set $c_{i,k} = 0$ if $k < 0$.

For $\delta \in \{0, 1\}$, let $\delta^\vee = \overline{\delta + n} \in \{0, 1\}$. For $\mathbf{c} \in \mathbb{Z}_{\geq 0}^{\Gamma_0^{(\delta)}}$, define $\mathbf{c}^\vee \in \mathbb{Z}_{\geq 0}^{\Gamma_0^{(\delta^\vee)}}$ by $\mathbf{c}^\vee = (c_{i^*, n-k})_{(i,k) \in \Gamma_0^{(\delta^\vee)}}$. By Lemma 2.2.1 we have $*B^{(\delta)}(\mathbf{c}) = B^{(\delta^\vee)}(\mathbf{c}^\vee)$, and hence

$$\varepsilon_j^*(B^{(\delta)}(\mathbf{c})) = \varepsilon_j(B^{(\delta^\vee)}(\mathbf{c}^\vee)) \quad (6.1.2)$$

holds. Using this, we can also calculate the values of ε_i^* 's.

6.2. Snake modules associated with height functions. Fix a strong duality datum $\mathcal{D} = \{\mathbf{L}_i\}_{i \in I}$ associated with \mathfrak{sl}_{n+1} , and a height function ξ . We write \widehat{Q} for \widehat{Q}^ξ , and $S_{i,k}$ for $S_{i,k}^{\mathcal{D}, \xi}$ ($(i, k) \in \widehat{Q}_0$).

Lemma 6.2.1. *Assume that a sequence $\mathbf{P} = ((i_1, k_1), \dots, (i_p, k_p))$ of elements of \widehat{Q}_0 is a snake, and $(j, l) \in \widehat{Q}_0$.*

- (1) *Suppose that $(j, l) \prec (i_1, k_1)$.*
 - (a) *If (i_1, k_1) is in prime snake position with respect to (j, l) , we have $\mathfrak{d}(S_{j,l}, \mathbb{S}(\mathbf{P})) = 1$.*
 - (b) *If (i_1, k_1) is not in snake position with respect to (j, l) (that is, $i_1 \in \{j \pm r\}$ and $k_1 = l + r$ hold for some $r \in \mathbb{Z}_{>0}$), then we have $\mathfrak{d}(S_{j,l}, \mathbb{S}(\mathbf{P})) = 0$.*
- (2) *Suppose that $(i_p, k_p) \prec (j, l)$.*
 - (c) *If (j, l) is in prime snake position with respect to (i_p, k_p) , we have $\mathfrak{d}(\mathbb{S}(\mathbf{P}), S_{j,l}) = 1$.*
 - (d) *If (j, l) is not in snake position with respect to (i_p, k_p) , we have $\mathfrak{d}(\mathbb{S}(\mathbf{P}), S_{j,l}) = 0$.*

Proof. We will use the notation in the previous subsection freely. (1) By replacing the pair (\mathcal{D}, ξ) using Lemma 5.2.2, we may assume that

$$l = -1,$$

and (\mathcal{D}, ξ) is the following specific one:

$$\xi = \xi^{(\delta)} \quad \text{and} \quad \mathcal{D} = \{\mathbf{L}_i\}_{i \in I} \quad \text{with} \quad \mathbf{L}_i = \begin{cases} S_{i,0} & \text{if } \bar{i} \neq \delta \\ S_{i^*,n} & \text{if } \bar{i} = \delta \end{cases} \quad \text{for } i \in I,$$

where we set $\delta = \bar{j} \in \{0, 1\}$. For any $(i, k) \in \widehat{Q}_0$ satisfying $\mathcal{D}^{-1}(j, l) \not\preceq (i, k)$, we have $\mathfrak{d}(S_{j,l}, \mathcal{D}^{-r} S_{i,k}) = 0$ for all $r \geq 0$ by Lemma 5.2.7 (i). Hence by Lemma 3.2.5 (i), we may further assume that $(i_p, k_p) \preceq \mathcal{D}^{-1}(j, l)$, which implies $(i_s, k_s) \in \Gamma_0^{(\delta)}$ for all $s \in [1, p]$. We have $S_{j,l} = \mathcal{D} S_{j^*,n} = \mathcal{D} \mathbf{L}_j$. Let $\{\mathbf{e}_{i,k} \mid (i, k) \in \Gamma_0^{(\delta)}\}$ be the standard basis of $\mathbb{Z}^{\Gamma_0^{(\delta)}}$, and set $\mathbf{c} = \sum_{s=1}^p \mathbf{e}_{i_s, k_s}$. It follows from Lemma 3.3.4 that

$$\mathcal{L}_{\mathcal{D}}(B^{(\delta)}(\mathbf{c})) \cong \mathbb{S}(\mathbf{P}), \quad (6.2.1)$$

and hence we have

$$\mathfrak{d}(S_{j,l}, \mathbb{S}(\mathbf{P})) = \mathfrak{d}(\mathcal{D} \mathbf{L}_j, \mathbb{S}(\mathbf{P})) = \varepsilon_j(B^{(\delta)}(\mathbf{c})) \quad (6.2.2)$$

by Proposition 3.3.3 (c). We see from Theorem 6.1.2 that

$$\varepsilon_j(B^{(\delta)}(\mathbf{c})) = \max_{\Sigma \in \mathcal{U}_j} (\#\{s \in [1, p] \mid (i_s, k_s) \in \Sigma\} - \#\{s \in [1, p] \mid (i_s, k_s + 2) \in \Sigma\}). \quad (6.2.3)$$

Define $P_1, P_2, \dots, P_{2p} \in \widehat{Q}_0$ by

$$P_{2s-1} = (i_s, k_s) \quad \text{and} \quad P_{2s} = (i_s, k_s + 2) \quad \text{for } s \in [1, p].$$

Note that $P_1 \prec P_2 \preceq P_3 \prec \dots \preceq P_{2p-1} \prec P_{2p}$ holds by the definition of the snake position. Now we show the assertion (a). In this case, we have $P_1 \in \Omega_j$. For any $\Sigma \in \mathcal{U}_j$, we easily see from the lower closedness that there is some $t \in [0, p]$ such that $P_s \in \Sigma$ if and only if $s \in [1, t]$, and then we have

$$\#\{s \in [1, p] \mid P_{2s-1} \in \Sigma\} - \#\{s \in [1, p] \mid P_{2s} \in \Sigma\} = \begin{cases} 0 & \text{if } t \in 2\mathbb{Z}, \\ 1 & \text{if } t \in 2\mathbb{Z} - 1. \end{cases}$$

Hence (a) follows from (6.2.2) and (6.2.3). In the case of (b), we have $P_1 \notin \Omega_j$ and $P_2 \in \Omega_j$, and the assertion is proved similarly.

(2) Let $\delta = \overline{n-j} \in \{0, 1\}$. Similarly as above, we may assume that

$$l = n + 1, \quad \xi = \xi^{(\delta)}, \quad \text{and} \quad \mathcal{D} = \{\mathbf{L}_i\}_{i \in I} \quad \text{with} \quad \mathbf{L}_i = \begin{cases} S_{i,0} & \text{if } \bar{i} \neq \delta \\ S_{i^*,n} & \text{if } \bar{i} = \delta \end{cases} \quad \text{for } i \in I.$$

By a similar argument as above, we may further assume that $(i_s, k_s) \in \Gamma_0^{(\delta)}$ for all $s \in [1, p]$. It follows from Proposition 3.3.3 that

$$\mathfrak{d}(\mathbb{S}(\mathbf{P}), S_{j,l}) = \mathfrak{d}(\mathbb{S}(\mathbf{P}), \mathcal{D}^{-1} \mathbf{L}_{j^*}) = \varepsilon_{j^*}^*(B^{(\delta)}(\mathbf{c})),$$

where we set $\mathbf{c} = \sum_{s=1}^p \mathbf{e}_{i_s, k_s} \in \mathbb{Z}_{\geq 0}^{\Gamma_0^{(\delta)}}$. We easily see that if \mathbf{P} and (j, l) satisfy the assumption of (c) (resp. (d)), then

$$\mathbf{P}^\vee = ((i_p^*, n - k_p), (i_{p-1}^*, n - k_{p-1}), \dots, (i_1^*, n - k_1))$$


and $(j^*, n - l)$ do that of (a) (resp. (b)). Hence the assertions (c) and (d) are proved from (the proof of) (a) and (b) by using (6.1.2). \square

Let $(i, k), (i', k') \in \widehat{Q}_0$, and suppose that (i', k') is in prime snake position with respect to (i, k) . We define $Q_{i,k}^{i',k'}$ and $R_{i,k}^{i',k'}$, each of which is an element of \widehat{Q}_0 or the empty set, by

$$Q_{i,k}^{i',k'} = \begin{cases} (\frac{1}{2}(i+i'+k-k'), \frac{1}{2}(i-i'+k+k')) & \text{if } k'-k < i+i', \\ \emptyset & \text{if } k'-k = i+i', \end{cases}$$

$$R_{i,k}^{i',k'} = \begin{cases} (\frac{1}{2}(i+i'-k+k'), \frac{1}{2}(-i+i'+k+k')) & \text{if } k'-k < 2n+2-i-i', \\ \emptyset & \text{if } k'-k = 2n+2-i-i'. \end{cases}$$

When $n = 5$, these are illustrated as follows, where (i, k) (resp. (i', k') , $Q_{i,k}^{i',k'}$, $R_{i,k}^{i',k'}$) are shown as \circ (resp. \bullet , $*$, \star):



Lemma 6.2.2. *Let $(j, l), (j', l') \in \widehat{Q}_0$, and assume that (j', l') is in prime snake position with respect to (j, l) . Then we have $S_{j,l} \Delta S_{j',l'} \cong S_{Q_{j,l}^{j',l'}} \otimes S_{R_{j,l}^{j',l'}}$, where we set $S_\emptyset = \mathbf{1}$.*

Proof. Essentially, this is a formula for the product of two dual root vectors, which has previously been known (see [Oh19]). For the reader's convenience, we give a proof.

As above, we may assume that

$$l = -1, \quad \xi = \xi^{(\delta)} \quad \text{with } \delta = \bar{j}, \quad \text{and } \mathcal{D} = \{\mathbf{L}_i\}_{i \in I} \quad \text{with } \mathbf{L}_i = \begin{cases} S_{i,0} & \text{if } \bar{i} \neq \delta, \\ S_{i^*,n} & \text{if } \bar{i} = \delta. \end{cases}$$

Set

$$\Pi := \{(i, k) \in \widehat{Q}_0 \mid (j, l) \prec (i, k) \prec (j', k')\} \subseteq \Gamma_0^{(\delta)}.$$

It follows from Lemma 6.2.1 that $\mathfrak{d}(S_{j,l}, S_{j',l'}) = 1$, and thus by Lemma 4.2.2 we have

$$S_{j,l} \Delta S_{j',l'} \cong \text{hd} \left(\bigotimes_{(i,k) \in \Pi} S_{i,k}^{\otimes a_{i,k}} \right) \quad (6.2.4)$$

for some $a_{i,k} \in \mathbb{Z}_{\geq 0}$, where the factors are ordered compatibly with \preceq . Set $\mathbf{a} = (a_{i,k}) \in \mathbb{Z}_{\geq 0}^{\Gamma_0^{(\delta)}}$, where $a_{i,k} = 0$ if $(i, k) \notin \Pi$. As in (6.2.1), the image of $B^{(\delta)}(\mathbf{a})$ under $\mathcal{L}_{\mathcal{D}}: \mathbf{B}^{\text{up}} \rightarrow \text{Irr}(\mathcal{C}_{\mathcal{D}})$ is isomorphic to the right-hand side of (6.2.4), and we have

$$\text{wt}(B^{(\delta)}(\mathbf{a})) = - \sum_{(i,k) \in \Pi} a_{i,k} \phi_{(\delta)}(i, k).$$

On the other hand, we have

$$\mathbf{L}_j \nabla (S_{j,l} \Delta S_{j',l'}) \cong \mathbf{L}_j \nabla (S_{j',l'} \nabla \mathcal{D} \mathbf{L}_j) \cong S_{j',l'},$$

and since \mathbf{L}_j and $S_{j',l'}$ are the images under $\mathcal{L}_{\mathcal{D}}$ of the dual root vectors of weight $-\alpha_j$ and $-\phi_{(\delta)}(j', l')$ respectively, we see that the weight of $\mathcal{L}_{\mathcal{D}}^{-1}(S_{j,l} \Delta S_{j',l'}) \in \mathbf{B}^{\text{up}}$ is $-\phi_{(\delta)}(j', l') + \alpha_j$. Hence it follows from (6.2.4) that

$$\sum_{(i,k) \in \Pi} a_{i,k} \phi_{(\delta)}(i, k) = \alpha_{x,y} - \alpha_j = \alpha_{x,j-1} + \alpha_{j+1,y}, \quad (6.2.5)$$

where we set $\phi_{(\delta)}(j', l') = \alpha_{x,y}$. We easily check from Lemma 6.1.1 that

$$\phi_{(\delta)}(\Pi) \subseteq \{\alpha_{r,s} \mid r \leq j \leq s\} \sqcup \{\alpha_{r,j-1} \mid r \leq j-1\} \sqcup \{\alpha_{j+1,s} \mid j+1 \leq s\},$$

and from this we see at once that (6.2.5) holds only when \mathbf{a} is given as follows:

$$a_{i,k} = 1 \text{ if } \phi_{(\delta)}(i, k) \in \{\alpha_{x,j-1}, \alpha_{j+1,y}\} \text{ and } a_{i,k} = 0 \text{ otherwise.}$$

It is easily seen from Lemma 6.1.1 that $x = j$ if $Q_{j,l}^{j',l'} = \emptyset$ and $\phi_{(\delta)}(Q_{j,l}^{j',l'}) = \alpha_{x,j-1}$ otherwise. Similarly, we also see that $y = j$ if $R_{j,l}^{j',l'} = \emptyset$ and $\phi_{(\delta)}(R_{j,l}^{j',l'}) = \alpha_{j+1,y}$ otherwise. Now, since $Q_{j,l}^{j',l'}$ and $R_{j,l}^{j',l'}$ are incomparable when they are nonzero, (6.2.4), together with Lemma 5.2.3 (d), completes the proof. \square

The following lemma is proved by inspection.

Lemma 6.2.3 ([MY12a, Proposition 3.2]). *Let $\mathbf{P} = ((i_1, k_1), \dots, (i_p, k_p)) \in (\widehat{Q}_0)^p$ be a prime snake with $p \geq 2$, and set*

$$\mathbf{Q} = (Q_{i_1, k_1}^{i_2, k_2}, \dots, Q_{i_{p-1}, k_{p-1}}^{i_p, k_p}) \quad \text{and} \quad \mathbf{R} = (R_{i_1, k_1}^{i_2, k_2}, \dots, R_{i_{p-1}, k_{p-1}}^{i_p, k_p}),$$

where \emptyset are ignored. Then \mathbf{Q} and \mathbf{R} are snakes with no elements in common.

Now we give the main theorem of this section, which is a generalization of [DLL19, Theorem 3.4] and [MY12a, Proposition 3.1, Theorem 4.1] in type A.

Theorem 6.2.4. *Let $\mathbf{P} = ((i_1, k_1), \dots, (i_p, k_p)) \in (\widehat{Q}_0)^p$ be a snake.*

- (i) *The simple module $\mathbb{S}(\mathbf{P})$ is real.*
- (ii) *If \mathbf{P} is prime, then $\mathbb{S}(\mathbf{P})$ is prime.*
- (iii) *Assume that \mathbf{P} is prime with $p \geq 2$, and set*

$$\mathbf{Q} = (Q_{i_1, k_1}^{i_2, k_2}, \dots, Q_{i_{p-1}, k_{p-1}}^{i_p, k_p}) \quad \text{and} \quad \mathbf{R} = (R_{i_1, k_1}^{i_2, k_2}, \dots, R_{i_{p-1}, k_{p-1}}^{i_p, k_p}).$$

Then $\mathbb{S}(\mathbf{Q})$ and $\mathbb{S}(\mathbf{R})$ strongly commute, and there is a short exact sequence

$$0 \rightarrow \mathbb{S}(\mathbf{Q}) \otimes \mathbb{S}(\mathbf{R}) \rightarrow \mathbb{S}(\mathbf{P}_{[1,p-1]}) \otimes \mathbb{S}(\mathbf{P}_{[2,p]}) \rightarrow \mathbb{S}(\mathbf{P}) \otimes \mathbb{S}(\mathbf{P}_{[2,p-1]}) \rightarrow 0.$$

Proof. Using Lemma 6.2.1, the assertion (i) follows from Proposition 4.1.3 and Lemma 5.2.7 (i), and (ii) from Proposition 4.1.1. For (iii), it suffices to show by Theorem 4.2.1 that

$$\text{hd} \left(\bigotimes_{a=1}^{p-1} (S_{i_a, k_a} \Delta S_{i_{a+1}, k_{a+1}}) \right) \cong \mathbb{S}(\mathbf{Q}) \otimes \mathbb{S}(\mathbf{R}), \quad (6.2.6)$$

which we prove by the induction on p . The case $p = 2$ is just Lemma 6.2.2. Assume that $p > 2$, and write Q for $Q_{i_{p-1}, k_{p-1}}^{i_p, k_p}$ and R for $R_{i_{p-1}, k_{p-1}}^{i_p, k_p}$. By the induction hypothesis, we have

$$\text{hd} \left(\bigotimes_{a=1}^{p-1} (S_{i_a, k_a} \Delta S_{i_{a+1}, k_{a+1}}) \right) \cong (\mathbb{S}(\mathbf{Q}') \otimes \mathbb{S}(\mathbf{R}')) \nabla (S_Q \otimes S_R),$$

where we set

$$\mathbf{Q}' = (Q_{i_1, k_1}^{i_2, k_2}, \dots, Q_{i_{p-2}, k_{p-2}}^{i_{p-1}, k_{p-1}}) \quad \text{and} \quad \mathbf{R}' = (R_{i_1, k_1}^{i_2, k_2}, \dots, R_{i_{p-2}, k_{p-2}}^{i_{p-1}, k_{p-1}}).$$

To prove (6.2.6), it is enough to show that $\mathfrak{d}(\mathbb{S}(\mathbf{Q}'), S_R) = \mathfrak{d}(\mathbb{S}(\mathbf{R}'), S_Q) = 0$. Indeed if this holds, then we have

$$\mathfrak{d}(\mathbb{S}(\mathbf{Q}), \mathbb{S}(\mathbf{R})) = \mathfrak{d}(\mathbb{S}(\mathbf{Q}') \nabla S_Q, \mathbb{S}(\mathbf{R}') \nabla S_R) = 0,$$

and there is a surjection

$$\begin{aligned} (\mathbb{S}(\mathbf{Q}') \otimes \mathbb{S}(\mathbf{R}')) \otimes (S_Q \otimes S_R) &\cong (\mathbb{S}(\mathbf{Q}') \otimes S_Q) \otimes (\mathbb{S}(\mathbf{R}') \otimes S_R) \\ &\twoheadrightarrow \mathbb{S}(\mathbf{Q}) \otimes \mathbb{S}(\mathbf{R}), \end{aligned}$$

which implies (6.2.6). Let us prove $\mathfrak{d}(\mathbb{S}(\mathbf{Q}'), S_R) = 0$ (the other is proved similarly). We may assume that $R \neq \emptyset$. Set $Q' = Q_{i_{p-2}, k_{p-2}}^{i_{p-1}, k_{p-1}}$. If $Q' \neq \emptyset$, then $\mathfrak{d}(\mathbb{S}(\mathbf{Q}'), S_R) = 0$ follows from Lemma 6.2.1 (d). Assume that $Q' = \emptyset$, which implies that $k_{p-1} - k_{p-2} = i_{p-1} + i_{p-2}$. We easily see that every element Q'' appearing in \mathbf{Q}' satisfies

$$Q'' \preceq (i_{p-2} - 1, k_{p-2} - 1) \text{ if } i_{p-2} \neq 1 \quad \text{and} \quad Q'' \preceq (i_{p-2}, k_{p-2} - 2) \text{ if } i_{p-2} = 1. \quad (6.2.7)$$

Hence by Lemma 5.2.7, it suffices to show that $\mathcal{D}^{-1}Q'' \not\preceq R$ holds if Q'' satisfies (6.2.7). We show this in the case $i_{p-2} \neq 1$ (the case $i_{p-2} = 1$ is proved similarly). We have

$$\mathcal{D}^{-1}Q'' \preceq (-i_{p-2} + n + 2, k_{p-2} + n) = (i_{p-1} + s, k_{p-1} - 2 + s)$$

for suitable $s \in \mathbb{Z}$. On the other hand, $R = (i_{p-1} + r, k_{p-1} + r)$ holds for some $r \in \mathbb{Z}_{>0}$, and therefore we have $\mathcal{D}^{-1}Q'' \not\preceq R$, as required. The proof is complete. \square

7. THE CASE OF TWISTED HEIGHT FUNCTIONS

7.1. Change of a reduced word. Throughout this section, we assume that $n = 2n_0 - 1$ with $n_0 \in \mathbb{Z}_{\geq 2}$. Unlike the untwisted case, the results in [Rei97] cannot be applied to reduced expressions adapted to twisted height functions. In this subsection, we will describe the connection between untwisted and twisted cases, which enable us to apply the results in the previous section to twisted cases.


Let θ be the untwisted height function defined by

$$\theta_i = \begin{cases} i & \text{for } i \in [1, n_0], \\ i - 2 & \text{for } i \in [n_0 + 1, n], \end{cases}$$

and Θ the twisted height function defined by

$$\Theta_i = \begin{cases} i & \text{for } i \in [1, n_0 - 1], \\ n_0 - 1/2 & \text{for } i = n_0, \\ n_0 - 1 & \text{for } i \in [n_0 + 1, n]. \end{cases}$$

In this section, we will treat these specific functions only. When $n = 7$, Γ^θ and Γ^Θ are given as follows:





Similarly as in (6.1.1), given a Γ_0^θ -tuple $\mathbf{c} = (c_{i,k})_{(i,k) \in \Gamma_0^\theta}$ of nonnegative integers, define $B^\theta(\mathbf{c}) \in \mathbf{B}^{\text{up}}$ by taking a compatible reading $\Gamma_0^\theta = \{(i_1, k_1), \dots, (i_N, k_N)\}$, putting $\mathbf{i} = (i_1, \dots, i_N) \in R(w_0)$, and setting $B^\theta(\mathbf{c}) = B^{\mathbf{i}}(\mathbf{c})$. We also define $B^\Theta(\mathbf{c}') \in \mathbf{B}^{\text{up}}$ similarly for a Γ_0^Θ -tuple $\mathbf{c}' = (c'_{i,k})_{(i,k) \in \Gamma_0^\Theta}$ of nonnegative integers. Given \mathbf{c}' , the goal of this subsection is to give a formula for \mathbf{c} satisfying $B^\theta(\mathbf{c}) = B^\Theta(\mathbf{c}')$ using Proposition 2.2.3 (ii), under certain conditions.

We prepare several notation. For $j \in [n_0 + 1, n]$, let $V\langle j \rangle$ be the subset of $I \times \mathbb{Z}/2$ defined by

$$\begin{aligned} V\langle j \rangle = & \{(i, i + 2k) \mid i < n_0, k \in [0, n - i - 1]\} \\ & \sqcup \{(n_0, n_0 + 2k) \mid k \in [0, n_0 - 1]\} \sqcup \{(i, i - 2 + 2k) \mid n_0 < i < j, k \in [0, n - i + 1]\} \\ & \sqcup \{(j, j - 3/2 + k) \mid k \in [0, 2n - 2j + 1]\} \sqcup \{(i, i - 1 + 2k) \mid i > j, k \in [0, n - j]\}. \end{aligned}$$

We also set $V\langle n_0 \rangle = \Gamma_0^\Theta$ and $V\langle n + 1 \rangle = \Gamma_0^\theta$. When $n = 7$, these are given as follows, where \star in $V\langle j \rangle$ denote the points $(j, j + 2r - 1/2)$, $(j + 1, j + 2r)$, $(j, j + 2r + 1/2)$ with $r \in [0, n - j - 1]$ appearing in Lemma 7.1.1 below (the dotted lines connecting points are for illustrative purposes only):





(7.1.1)

We have $\sharp V\langle j \rangle = N$ for all $j \in [n_0, n+1]$. For each $j \in [n_0, n+1]$, take and fix a total ordering $V\langle j \rangle = \{(i_1, k_1), \dots, (i_N, k_N)\}$ such that $r < s$ holds whenever $k_r < k_s$, and set $\mathbf{i}\langle j \rangle = (i_1, \dots, i_N) \in I^N$. For $(i, k) \in V\langle j \rangle$, denote by $i^{(k)}$ the letter i_r in $\mathbf{i}\langle j \rangle$, where $r \in [1, N]$ is such that $(i_r, k_r) = (i, k)$. Obviously, $\mathbf{i}\langle n_0 \rangle$ (resp. $\mathbf{i}\langle n+1 \rangle$) is adapted to the twisted height function Θ (resp. the height function θ), and we easily see that $\mathbf{i}\langle n \rangle$ and $\mathbf{i}\langle n+1 \rangle$ are commutation equivalent. Let $[\mathbf{i}\langle j \rangle]$ denote the commutation class containing $\mathbf{i}\langle j \rangle$.

Lemma 7.1.1.


(i) For each $j \in [n_0, n-1]$, there is a word $\mathbf{i} \in [\mathbf{i}\langle j \rangle]$ containing consecutive subwords

$$(j^{(j+2r-1/2)}, (j+1)^{(j+2r)}, j^{(j+2r+1/2)}) \quad \text{for all } r \in [0, n-j-1].$$

Let \mathbf{i}' be the word obtained from \mathbf{i} by transforming all these subwords into $(j+1, j, j+1)$. Then we have $\mathbf{i}' \in [\mathbf{i}\langle j+1 \rangle]$.

(ii) We have $\mathbf{i}\langle j \rangle \in R(w_0)$ for all $j \in [n_0, n+1]$.

Proof. The first assertion of (i) is easily checked, and the second is also checked directly by noting that the transformation $(j, j+1, j) \rightarrow (j+1, j, j+1)$ is expressed pictorially as follows:



See (7.1.1). Now (ii) is obvious since $\mathbf{i}\langle n_0 \rangle \in R(w_0)$. □

For each $j \in [n_0, n]$, define the map

$$\rho_{\langle j \rangle}: \mathbb{Z}_{\geq 0}^{V\langle j \rangle} \rightarrow \mathbb{Z}_{\geq 0}^{V\langle j+1 \rangle}, \quad \mathbf{c} = (c_{i,k})_{(i,k) \in V\langle j \rangle} \mapsto \mathbf{c}' = (c'_{i,k})_{(i,k) \in V\langle j+1 \rangle}$$

as follows: for $r \in [0, n-j-1]$, $(c'_{j+1, j+2r-1/2}, c'_{j, j+2r}, c'_{j+1, j+2r+1/2})$ is obtained by applying the transformation (2.2.2) to $(c_{j, j+2r-1/2}, c_{j+1, j+2r}, c_{j, j+2r+1/2})$, and the other

$c'_{i,k}$'s are determined by $c'_{i,k} = c_{i,k+t}$, where

$$t = \begin{cases} 1/2 & \text{if } (i, k) = (j, j-2), \\ -1/2 & \text{if } (i, k) = (j, 2n-j), \\ 0 & \text{if } i \neq j, j+1. \end{cases}$$

Set

$$\rho = \rho_{\langle n \rangle} \circ \rho_{\langle n-1 \rangle} \circ \cdots \circ \rho_{\langle n_0 \rangle} : \mathbb{Z}_{\geq 0}^{\Gamma_0^\Theta} \rightarrow \mathbb{Z}_{\geq 0}^{\Gamma_0^\theta}.$$

For any $\mathbf{c} \in \mathbb{Z}_{\geq 0}^{\Gamma_0^\Theta}$, it follows from Lemma 7.1.1 and Proposition 2.2.3 (ii) that

$$B^\Theta(\mathbf{c}) = B^\theta(\rho(\mathbf{c})). \quad (7.1.2)$$

Let $\{\mathbf{e}_{i,k} \mid (i,k) \in \Gamma_0^\Theta\}$ denote the standard basis of $\mathbb{Z}^{\Gamma_0^\Theta}$. We would like to give an explicit formula for $\rho(\mathbf{c})$ when \mathbf{c} is expressed as $\sum_{r=1}^p \mathbf{e}_{i_r, k_r}$ with $((i_1, k_1), \dots, (i_p, k_p))$ being a snake in $\Gamma_0^\Theta \subseteq \widehat{Q}_0^\Theta$. For this, the following remark is useful.

Remark 7.1.2. Under the transformation $(c_{k-1}, c_k, c_{k+1}) \mapsto (c'_{k-1}, c'_k, c'_{k+1})$ in (2.2.2), we have

$$\begin{aligned} \text{(i)} \quad & (1, 0, 0) \mapsto (0, 0, 1), & \text{(ii)} \quad & (0, 0, 1) \mapsto (1, 0, 0), \\ \text{(iii)} \quad & (0, 1, 0) \mapsto (1, 0, 1), & \text{(iv)} \quad & (1, 0, 1) \mapsto (0, 1, 0). \end{aligned}$$

Pictorially, these are expressed as follows:

$$\begin{aligned} \text{(i)} \quad & \begin{array}{c} \star \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \end{array} \Rightarrow \begin{array}{c} \diagdown \quad \diagup \\ \bullet \end{array} \quad \text{(ii)} \quad \begin{array}{c} \bullet \quad \star \\ \diagdown \quad \diagup \\ \bullet \end{array} \Rightarrow \begin{array}{c} \diagdown \quad \diagup \\ \star \end{array} \\ \text{(iii)} \quad & \begin{array}{c} \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \star \end{array} \Rightarrow \begin{array}{c} \diagdown \quad \diagup \\ \star \end{array} \quad \text{(iv)} \quad \begin{array}{c} \star \quad \star \\ \diagdown \quad \diagup \\ \bullet \end{array} \Rightarrow \begin{array}{c} \diagdown \quad \diagup \\ \bullet \end{array} \end{aligned}$$

Therefore, (i) (resp. (ii)) is seen as moving to lower right (resp. lower left), (iii) as splitting in two points, and (iv) as combining two points into one.

For $\flat \in \{<, >, \text{U}, \text{D}\}$, we write $\Gamma_0^{\Theta, \flat} := \Gamma_0^\Theta \cap \widehat{Q}_0^{\Theta, \flat}$. We also write $\Gamma_0^{\Theta, \geq} := \Gamma_0^\Theta \setminus \Gamma_0^{\Theta, <}$. For $(i, k) \in \Gamma_0^{\Theta, \geq}$, define $X_{i,k}^-$ and $X_{i,k}^+$, each of which is an element of Γ_0^θ (not Γ_0^Θ) or the empty set, as follows:

$$X_{i,k}^- = \begin{cases} \emptyset & \text{if } (i, k) \in \Gamma_0^{\Theta, \text{D}}, \\ (\frac{1}{2}(\Theta_i + k + 2), \frac{1}{2}(\Theta_i + k - 2)) & \text{if } (i, k) \in \Gamma_0^{\Theta, >} \sqcup \Gamma_0^{\Theta, \text{U}}, \end{cases}$$


and

$$X_{i,k}^+ = \begin{cases} \emptyset & \text{if } (i, k) \in \Gamma_0^{\Theta, \text{U}}, \\ (\frac{1}{2}(\Theta_i - k + 2n), \frac{1}{2}(-\Theta_i + k + 2n)) & \text{if } (i, k) \in \Gamma_0^{\Theta, >} \sqcup \Gamma_0^{\Theta, \text{D}}. \end{cases}$$

In addition, for $(i, k), (i', k') \in \Gamma_0^{\Theta, \geq}$ such that (i', k') is in snake position (of twisted type) with respect to (i, k) , define $X_{i,k}^{i',k'}$, which is an element of Γ_0^θ or the empty set, by

$$X_{i,k}^{i',k'} = \begin{cases} \emptyset & \text{if } (i, k) \in \Gamma_0^{\Theta, \text{U}}, \\ \left(\frac{1}{2}(\Theta_i + \Theta_{i'} - k + k'), \frac{1}{2}(\Theta_i - \Theta_{i'} + k + k')\right) & \text{if } (i, k) \in \Gamma_0^{\Theta, >} \sqcup \Gamma_0^{\Theta, \text{D}}. \end{cases}$$

When $n = 7$, these are illustrated as follows. Here $\Gamma_0^{\Theta, <}$ is omitted, (i, k) and (i', k') are shown as \circ whose first coordinates are read from the left-hand scale, and $X_{i,k}^\pm$ and $X_{i,k}^{i',k'}$ are shown as \star whose first coordinates are read from the right-hand scale:



For a sequence $\mathbf{P} = ((i_1, k_1), \dots, (i_p, k_p))$ of elements of Γ_0^ξ with $\xi \in \{\theta, \Theta\}$, set $e(\mathbf{P}) = \sum_r e_{i_r, k_r} \in \mathbb{Z}_{\geq 0}^{\Gamma_0^\xi}$.

Proposition 7.1.3. *Assume that $\mathbf{P} = ((i_1, k_1), \dots, (i_p, k_p))$ is a snake of twisted type in $\Gamma_0^\Theta \subseteq \widehat{Q}_0^\Theta$. Let $1 = r_0 \leq r_1 < \dots < r_{t-1} < r_t < r_{t+1} = p+1$ be the unique increasing sequence satisfying the following for all $a \in [0, t]$:*

- (i) *if $a \in 2\mathbb{Z}$, then $(i_s, k_s) \in \Gamma_0^{\Theta, <}$ holds for all $s \in [r_a, r_{a+1} - 1]$, and*
- (ii) *if $a \in 2\mathbb{Z} + 1$, then $(i_s, k_s) \in \Gamma_0^{\Theta, \geq}$ holds for all $s \in [r_a, r_{a+1} - 1]$.*


For each $a \in [1, t]$ with $a \in 2\mathbb{Z} + 1$, set

$$\mathbf{P}^{(a)} = ((i_r, k_r), \dots, (i_{r'-1}, k_{r'-1})) \in (\Gamma_0^{\Theta, \geq})^{r'-r}, \quad \text{and}$$

$$\mathbf{Q}^{(a)} = (X_{i_r, k_r}^-, X_{i_{r+1}, k_{r+1}}^{i_{r+1}, k_{r+1}}, \dots, X_{i_{r'-2}, k_{r'-2}}^{i_{r'-1}, k_{r'-1}}, X_{i_{r'-1}, k_{r'-1}}^+) \in (\Gamma_0^\theta)^{r''} \text{ for some } r'' \in \mathbb{Z}_{>0},$$

where $r = r_a$ and $r' = r_{a+1}$, and \emptyset are ignored. Let \mathbf{P}^\dagger denote the sequence of elements of Γ_0^θ obtained from \mathbf{P} by replacing the subsequences $\mathbf{P}^{(a)}$ with $\mathbf{Q}^{(a)}$ for all $a \in [1, t]$ such that $a \in 2\mathbb{Z} + 1$. Then \mathbf{P}^\dagger is a snake of untwisted type in Γ_0^θ and we have $\rho(e(\mathbf{P})) = e(\mathbf{P}^\dagger)$.

Proof. By tracing the transformations appearing in ρ using Remark 7.1.2, $\rho(e(\mathbf{P})) = e(\mathbf{P}^\dagger)$ is proved directly. See the following example, where \circ forms \mathbf{P} and \star does \mathbf{P}^\dagger .



It is also checked directly by inspection that \mathbf{P}^\dagger is a snake of untwisted type. \square

7.2. Snake modules associated with twisted height functions. Fix a strong duality datum $\mathcal{D} = \{\mathbf{L}_i\}_{i \in I} \subseteq \mathcal{C}_g$ associated with \mathfrak{sl}_{n+1} . In the sequel, we usually omit \mathcal{D} and write $S_{i,k}^\Theta$ for $S_{i,k}^{\mathcal{D},\Theta}$, etc. All the assertions in this subsection will be stated for $S_{i,k}^\Theta$, but no generality is lost by this specific choice of a twisted height function by Lemma 5.2.2.

By Proposition 7.1.3, (7.1.2) and Lemma 3.3.4, the following lemma is immediate.

Lemma 7.2.1. *Assume that $\mathbf{P} = ((i_1, k_1), \dots, (i_p, k_p))$ is a snake in Γ_0^Θ , and let \mathbf{P}^\dagger be the snake in Γ_0^Θ given in Proposition 7.1.3. Then the snake module $\mathbb{S}^\Theta(\mathbf{P})$ of twisted type is isomorphic to the snake module $\mathbb{S}^\Theta(\mathbf{P}^\dagger)$ of untwisted type.*

Lemma 7.2.2. *Let $\mathbf{P} = ((i_1, k_1), \dots, (i_p, k_p))$ be a snake in \widehat{Q}_0^Θ , and $(j, l) \in \widehat{Q}_0^\Theta$.*

- (1) *Suppose that $(j, l) \prec (i_1, k_1)$.*
 - (a) *If (i_1, k_1) is in prime snake position with respect to (j, l) , we have $\mathfrak{d}(S_{j,l}^\Theta, \mathbb{S}^\Theta(\mathbf{P})) = 1$.*
 - (b) *Assume that $\Theta_{i_1} \in \{\Theta_j \pm r\}$ and $k_1 = l + r$ hold for some $r \in \mathbb{Z}_{>0}/2$. Then we have $\mathfrak{d}(S_{j,l}^\Theta, \mathbb{S}^\Theta(\mathbf{P})) = 0$.*
 - (c) *If $(j, l) \in \widehat{Q}_0^{\Theta, \cup}$ and $(i_1, k_1) \in \widehat{Q}_0^{\Theta, >} \sqcup \widehat{Q}_0^{\Theta, \cup}$, or $(j, l) \in \widehat{Q}_0^{\Theta, \mathcal{D}}$ and $(i_1, k_1) \in \widehat{Q}_0^{\Theta, <} \sqcup \widehat{Q}_0^{\Theta, \mathcal{D}}$, then we have $\mathfrak{d}(S_{j,l}^\Theta, \mathbb{S}^\Theta(\mathbf{P})) = 0$.*
- (2) *Suppose that $(i_p, k_p) \prec (j, l)$.*
 - (d) *If (j, l) is in prime snake position with respect to (i_p, k_p) , we have $\mathfrak{d}(\mathbb{S}^\Theta(\mathbf{P}), S_{j,l}^\Theta) = 1$.*
 - (e) *Assume that $\Theta_j \in \{\Theta_{i_p} \pm r\}$ and $l = k_p + r$ hold for some $r \in \mathbb{Z}_{>0}/2$. Then we have $\mathfrak{d}(\mathbb{S}^\Theta(\mathbf{P}), S_{j,l}^\Theta) = 0$.*
 - (f) *If $(i_p, k_p) \in \widehat{Q}_0^{\Theta, <} \sqcup \widehat{Q}_0^{\Theta, \cup}$ and $(j, l) \in \widehat{Q}_0^{\Theta, \cup}$, or $(i_p, k_p) \in \widehat{Q}_0^{\Theta, >} \sqcup \widehat{Q}_0^{\Theta, \mathcal{D}}$ and $(j, l) \in \widehat{Q}_0^{\Theta, \mathcal{D}}$, then we have $\mathfrak{d}(\mathbb{S}^\Theta(\mathbf{P}), S_{j,l}^\Theta) = 0$.*

Proof. (1) By applying Lemma 5.2.6, if necessary, we may (and do) assume in the proof of each assertion that

- (a) $(j, l) \in \widehat{Q}_0^{\Theta, <} \sqcup \widehat{Q}_0^{\Theta, \cup}$,
- (b) $\Theta_{i_1} = \Theta_j - r$ and $k_1 = l + r$ for some r , and in particular, $j \notin \widehat{Q}_0^{\Theta, \mathcal{D}}$,
- (c) $(j, l) \in \widehat{Q}_0^{\Theta, \cup}$ and $(i_1, k_1) \in \widehat{Q}_0^{\Theta, >} \sqcup \widehat{Q}_0^{\Theta, \cup}$,

respectively. In all these three cases, we may further assume by a similar argument as in Lemma 6.2.1 that

$$l = \Theta_j - d_j \quad \text{and} \quad (i_s, k_s) \in \Gamma_0^\Theta \quad \text{for all } s \in [1, p], \quad (7.2.1)$$

where we set $d_i = 2 - \delta_{i, n_0}$ for $i \in I$. It follows from Lemma 7.2.1 that

$$\begin{aligned} \mathfrak{d}(S_{j,l}^\Theta, \mathbb{S}^\Theta(\mathbf{P})) &= \mathfrak{d}(\mathcal{D}S_{j^*, \Theta_j - d_j + n}^\Theta, \mathbb{S}^\Theta(\mathbf{P})) \\ &= \mathfrak{d}(\mathcal{D}\mathbb{S}^\theta((j^*, \Theta_j - d_j + n)^\dagger), \mathbb{S}^\theta(\mathbf{P}^\dagger)), \end{aligned} \quad (7.2.2)$$

and it is directly checked from Proposition 7.1.3 that

$$(j^*, \Theta_j - d_j + n)^\dagger = \begin{cases} ((n, -2 + n), (j^*, j - 1 + n)) & \text{if } j < n_0, \\ ((n_0, n_0 - 1 + n)) & \text{if } (j, l) = (n_0, n_0 - 3/2), \\ ((j^*, j - 3 + n)) & \text{if } j > n_0, \end{cases}$$

which implies

$$\mathcal{D}\mathbb{S}^\theta((j^*, \Theta_j - d_j + n)^\dagger) \cong \begin{cases} S_{1,-3}^\theta \nabla S_{j,j-2}^\theta & \text{if } j < n_0, \\ S_{n_0, n_0-2}^\theta & \text{if } j = n_0, \\ S_{j,j-4}^\theta & \text{if } j > n_0. \end{cases}$$

On the other hand, we can also show using the same proposition that

$$\text{FT}(\mathbf{P}^\dagger) = \begin{cases} (i_1, k_1) & \text{if } (i_1, k_1) \in \Gamma_0^{\Theta, <}, \\ (i_1 + s, k_1 + s + 1/2) \text{ for some } s & \text{if } (i_1, k_1) \in \Gamma_0^{\Theta, \text{D}}, \\ (\frac{1}{2}(\Theta_{i_1} + k_1 + 2), \frac{1}{2}(\Theta_{i_1} + k_1 - 2)) & \text{if } (i_1, k_1) \in \Gamma_0^{\Theta, >} \sqcup \Gamma_0^{\Theta, \text{U}}, \end{cases} \quad (7.2.3)$$

where we denote by $\text{FT}(\mathbf{P}^\dagger)$ the first term of the sequence \mathbf{P}^\dagger .

Now let us prove the assertion (a), where we are assuming that $(j, l) \in \widehat{Q}_0^{\Theta, <} \sqcup \widehat{Q}_0^{\Theta, \text{U}}$ and $l = \Theta_j - d_j$. By the above calculations, we have

$$\mathfrak{d}(S_{j,l}^\Theta, \mathbb{S}^\Theta(\mathbf{P})) = \begin{cases} \mathfrak{d}(S_{1,-3}^\theta \nabla S_{j,j-2}^\theta, \mathbb{S}^\theta(\mathbf{P}^\dagger)) & \text{if } j < n_0, \\ \mathfrak{d}(S_{n_0, n_0-2}^\theta, \mathbb{S}^\theta(\mathbf{P}^\dagger)) & \text{if } j = n_0. \end{cases} \quad (7.2.4)$$

By the definition of the snake position, we have $(i_1, k_1) \in \Gamma_0^{\Theta, <} \sqcup \Gamma_0^{\Theta, \text{D}}$, and then we easily see from (7.2.3) that

$$\mathcal{D}_\theta^{-1}(1, -3) = (n, n - 2) \not\leq \text{FT}(\mathbf{P}^\dagger),$$

which implies that $\mathcal{D}^s S_{1,-3}^\theta$ and $\mathbb{S}^\theta(\mathbf{P}^\dagger)$ strongly commute for all $s \in \mathbb{Z}_{\geq 0}$ by Lemma 5.2.7 (i). Therefore (7.2.4), together with Lemma 3.2.5, gives

$$\mathfrak{d}(S_{j,l}^\Theta, \mathbb{S}^\Theta(\mathbf{P})) = \mathfrak{d}(S_{j,j-2}^\theta, \mathbb{S}^\theta(\mathbf{P}^\dagger)) \quad \text{for all } j \leq n_0.$$

It is easy to check using (7.2.3) that, if (i_1, k_1) is in prime snake position with respect to (j, l) in \widehat{Q}_0^Θ , then $\text{FT}(\mathbf{P}^\dagger)$ is in prime snake position with respect to $(j, j - 2)$ in \widehat{Q}_0^θ . Since \mathbf{P}^\dagger is a snake in \widehat{Q}_0^θ by Proposition 7.1.3, the assertion (a) now follows from Lemma 6.2.1 (a).

Next let us prove (b), where we are assuming that

$$\Theta_{i_1} = \Theta_j - r, \quad k_1 = l + r \quad \text{for some } r, \quad \text{and} \quad l = \Theta_j - d_j. \quad (7.2.5)$$

If $j \leq n_0$, then $i_1 < n_0$ holds and by the same calculation as above we have

$$\mathfrak{d}(S_{j,l}^\Theta, \mathbb{S}^\Theta(\mathbf{P})) = \mathfrak{d}(S_{j,j-2}^\Theta, \mathbb{S}^\Theta(\mathbf{P}^\dagger)) \quad \text{and} \quad \text{FT}(\mathbf{P}^\dagger) = (i_1, k_1).$$

The assertion (b) then follows from Lemma 6.2.1 (b). If $j > n_0$, on the other hand, we have

$$\mathfrak{d}(S_{j,l}^\Theta, \mathbb{S}^\Theta(\mathbf{P})) = \mathfrak{d}(S_{j,j-4}^\Theta, \mathbb{S}^\Theta(\mathbf{P}^\dagger)).$$

Noting that $(i_1, k_1) \notin \Gamma_0^{\Theta, \text{D}}$, it is easy to show from (7.2.3) and (7.2.5) that $\text{FT}(\mathbf{P}^\dagger) = (j - r', j - 4 + r')$ holds for some r' , and hence the assertion follows from Lemma 6.2.1 (b) in this case as well.

Finally let us prove (c), where we are assuming that

$$(j, l) = (n_0, n_0 - 3/2) \in \widehat{Q}_0^{\Theta, \text{U}} \quad \text{and} \quad (i_1, k_1) \in \Gamma_0^{\Theta, >} \sqcup \Gamma_0^{\Theta, \text{U}}.$$

As calculated above, we have $S_{j,l}^\Theta \cong S_{n_0, n_0-2}^\Theta$ and $\text{FT}(\mathbf{P}^\dagger) = (n_0 + s, n_0 - 2 + s)$ for suitable s . Hence the assertion follows from Lemma 6.2.1 (b).


All the assertions of (2) are proved similarly. \square

Let $(i, k), (i', k') \in \widehat{Q}_0^\Theta$, and suppose that (i', k') is in prime snake position with respect to (i, k) . We define $\mathbf{Q}_{i,k}^{i',k'}$ and $\mathbf{R}_{i,k}^{i',k'}$, each of which is a sequence of zero, one or two elements of \widehat{Q}_0^Θ , as follows: if $(i, k) \in \widehat{Q}_0^{\Theta, <} \sqcup \widehat{Q}_0^{\Theta, \text{U}}$, we set

$$\mathbf{Q}_{i,k}^{i',k'} = \begin{cases} \emptyset & (k' - k = \Theta_i + \Theta_{i'}), \\ ((\frac{1}{2}(\Theta_i + \Theta_{i'} + k - k'), \frac{1}{2}(\Theta_i - \Theta_{i'} + k + k'))) & (k' - k < \Theta_i + \Theta_{i'}), \end{cases}$$

$$\mathbf{R}_{i,k}^{i',k'} = \begin{cases} ((\frac{1}{2}(i + i' - k + k'), \frac{1}{2}(-i + i' + k + k'))) & (i, i' < n_0, k' - k < 2n_0 - i - i'), \\ ((n_0, -i + k + n_0 - \frac{1}{2}), (n_0, i' + k' - n_0 + \frac{1}{2})) & (i, i' < n_0, k' - k \geq 2n_0 - i - i'), \\ ((n_0, -i + k + n_0 - \frac{1}{2})) & (i < n_0, i = n_0), \\ ((n_0, i' + k' - n_0 + \frac{1}{2})) & (i = n_0, i < n_0), \\ \emptyset & (i = i' = n_0). \end{cases}$$

If $(i, k) \in \widehat{Q}_0^{\Theta, >} \sqcup \widehat{Q}_0^{\Theta, \text{D}}$, putting $(j, l) = \mathcal{D}(i, k)$ and $(j', l') = \mathcal{D}(i', k')$, we set $\mathbf{Q}_{i,k}^{i',k'} = \mathcal{D}^{-1} \mathbf{R}_{j,l}^{j',l'}$ and $\mathbf{R}_{i,k}^{i',k'} = \mathcal{D}^{-1} \mathbf{Q}_{j,l}^{j',l'}$. These are illustrated as follows, where (i, k) (resp. (i', k') , $\mathbf{Q}_{i,k}^{i',k'}$, $\mathbf{R}_{i,k}^{i',k'}$) are shown as \circ (resp. \bullet , $*$, \star):



Lemma 7.2.3. *Let $(i, k), (i', k') \in \widehat{Q}_0^\Theta$, and assume that (i', k') is in prime snake position with respect to (i, k) . Then we have*

$$S_{i,k}^\Theta \Delta S_{i',k'}^\Theta \cong \mathbb{S}^\Theta(\mathbf{Q}_{i,k}^{i',k'}) \otimes \mathbb{S}^\Theta(\mathbf{R}_{i,k}^{i',k'}).$$

Proof. Since

$$\mathcal{D}(S_{i,k}^\Theta \Delta S_{i',k'}^\Theta) \cong S_{\mathcal{D}(i',k')}^\Theta \nabla S_{\mathcal{D}(i,k)}^\Theta \cong S_{\mathcal{D}(i,k)}^\Theta \Delta S_{\mathcal{D}(i',k')}^\Theta,$$

we may assume by using Lemma 5.2.6 that $(i, k) \in \widehat{Q}_0^{\Theta, <} \sqcup \widehat{Q}_0^{\Theta, \cup}$. First let us consider the case $(i, k) \in \widehat{Q}_0^{\Theta, \cup}$. We may further assume by Lemma 5.2.2 that $(i, k) = (n_0, n_0 - 3/2)$, which forces $(i', k') \in \Gamma_0^{\Theta, <} \sqcup \Gamma_0^{\Theta, \text{D}}$. If $(i', k') \in \Gamma_0^{\Theta, <}$, it is proved directly from Proposition 7.1.3 and Lemma 6.2.2 that

$$\begin{aligned} S_{i,k}^\Theta \Delta S_{i',k'}^\Theta &\cong S_{n_0, n_0-2}^\Theta \Delta S_{i',k'}^\Theta \cong S_{Q_{n_0, n_0-2}^{i',k'}}^\Theta \otimes S_{R_{n_0, n_0-2}^{i',k'}}^\Theta \\ &\cong \mathbb{S}^\Theta(\mathbf{Q}_{i,k}^{i',k'}) \otimes \mathbb{S}^\Theta(\mathbf{R}_{i,k}^{i',k'}), \end{aligned}$$

and the assertion is proved. The case $(i', k') \in \Gamma_0^{\Theta, \text{D}}$ is proved by a similar calculation.

Next let us consider the case $(i, k) \in \widehat{Q}_0^{\Theta, <}$. We may assume that $k = i$, and then by the definition of the prime snake position we have $(i', k') \in \Gamma_0^{\Theta, <} \sqcup \Gamma_0^{\Theta, \text{D}}$. Now the assertion is proved by a similar argument as above. \square

Similarly to Lemma 6.2.3, the following is proved by inspection.

Lemma 7.2.4 ([MY12a, Proposition 3.2]). *Let $p \in \mathbb{Z}_{\geq 2}$, and $\mathbf{P} = ((i_1, k_1), \dots, (i_p, k_p))$ be a prime snake in \widehat{Q}_0^Θ . Set*

$$\mathbf{Q} = \mathbf{Q}_{i_1, k_1}^{i_2, k_2} * \dots * \mathbf{Q}_{i_{p-1}, k_{p-1}}^{i_p, k_p} \quad \text{and} \quad \mathbf{R} = \mathbf{R}_{i_1, k_1}^{i_2, k_2} * \dots * \mathbf{R}_{i_{p-1}, k_{p-1}}^{i_p, k_p},$$

where $*$ denotes the concatenation. Then \mathbf{Q} and \mathbf{R} are snakes with no elements in common.

Now we give the main theorem of this section, which is a generalization of [DLL19, Theorem 3.4] and [MY12a, Proposition 3.1, Theorem 4.1] in type B .

Theorem 7.2.5. *Let $\mathbf{P} = ((i_1, k_1), \dots, (i_p, k_p))$ be a snake in \widehat{Q}_0^Θ .*

- (i) *The simple module $\mathbb{S}^\Theta(\mathbf{P})$ is real.*
- (ii) *If \mathbf{P} is prime, then $\mathbb{S}^\Theta(\mathbf{P})$ is prime.*
- (iii) *Assume that \mathbf{P} is prime with $p \geq 2$, and set*

$$\mathbf{Q} = \mathbf{Q}_{i_1, k_1}^{i_2, k_2} * \dots * \mathbf{Q}_{i_{p-1}, k_{p-1}}^{i_p, k_p} \quad \text{and} \quad \mathbf{R} = \mathbf{R}_{i_1, k_1}^{i_2, k_2} * \dots * \mathbf{R}_{i_{p-1}, k_{p-1}}^{i_p, k_p}.$$

Then $\mathbb{S}^\Theta(\mathbf{Q})$ and $\mathbb{S}^\Theta(\mathbf{R})$ strongly commute, and there is a short exact sequence

$$0 \rightarrow \mathbb{S}^\Theta(\mathbf{Q}) \otimes \mathbb{S}^\Theta(\mathbf{R}) \rightarrow \mathbb{S}^\Theta(\mathbf{P}_{[1, p-1]}) \otimes \mathbb{S}^\Theta(\mathbf{P}_{[2, p]}) \rightarrow \mathbb{S}^\Theta(\mathbf{P}) \otimes \mathbb{S}^\Theta(\mathbf{P}_{[2, p-1]}) \rightarrow 0.$$

Proof. The proof is similar to that of Theorem 6.2.4.

Using Lemma 7.2.2, the assertions (i) and (ii) are proved from Propositions 4.1.3 and 4.1.1, respectively. For (iii), by the same argument as in the proof of Theorem 6.2.4, it suffices to show for $p \geq 3$ that

$$\mathfrak{d}(\mathbb{S}^\Theta(\mathbf{Q}'), \mathbb{S}^\Theta(\mathbf{R}_{i_{p-1}, k_{p-1}}^{i_p, k_p})) = 0 = \mathfrak{d}(\mathbb{S}^\Theta(\mathbf{R}'), \mathbb{S}^\Theta(\mathbf{Q}_{i_{p-1}, k_{p-1}}^{i_p, k_p})),$$

where we set

$$\mathbf{Q}' = \mathbf{Q}_{i_1, k_1}^{i_2, k_2} * \dots * \mathbf{Q}_{i_{p-2}, k_{p-2}}^{i_{p-1}, k_{p-1}} \quad \text{and} \quad \mathbf{R}' = \mathbf{R}_{i_1, k_1}^{i_2, k_2} * \dots * \mathbf{R}_{i_{p-2}, k_{p-2}}^{i_{p-1}, k_{p-1}}.$$

Let us show the former equality (the latter is proved similarly). Set $\tilde{\mathbf{R}} = \mathbf{R}_{i_{p-1}, k_{p-1}}^{i_p, k_p}$, which we may assume not to be the empty set. If $\mathbf{Q}_{i_{p-2}, k_{p-2}}^{i_{p-1}, k_{p-1}} \neq \emptyset$, then the equality is proved from Lemma 7.2.2 (e) and (f). Assume that $\mathbf{Q}_{i_{p-2}, k_{p-2}}^{i_{p-1}, k_{p-1}} = \emptyset$. If $(i_{p-2}, k_{p-2}) \in \widehat{Q}_0^{\Theta, <} \sqcup \widehat{Q}_0^{\Theta, \cup}$, then the same argument as in the proof of Theorem 6.2.4 shows that

$$R \not\subseteq \mathcal{D}^{-1}Q \quad \text{for all } Q \in \mathbf{Q}' \text{ and } R \in \tilde{\mathbf{R}}, \quad (7.2.6)$$

and hence the equality holds. If $(i_{p-2}, k_{p-2}) \in \widehat{Q}_0^{\Theta, >} \sqcup \widehat{Q}_0^{\Theta, \mathcal{D}}$, on the other hand, $\mathbf{Q}_{i_{p-2}, k_{p-2}}^{i_{p-1}, k_{p-1}} = \emptyset$ implies

$$(i_{p-2}, k_{p-2}) \in \widehat{Q}_0^{\Theta, \mathcal{D}}, \quad (i_{p-1}, k_{p-1}) \in \widehat{Q}_0^{\Theta, \cup} \quad \text{and} \quad \tilde{\mathbf{R}} \subseteq \widehat{Q}_0^{\Theta, \cup}.$$

If $p = 3$ or $\mathbf{Q}_{i_{p-3}, k_{p-3}}^{i_{p-2}, k_{p-2}} = \emptyset$, then (7.2.6) holds as well, and the equality follows. If $\mathbf{Q}_{i_{p-3}, k_{p-3}}^{i_{p-2}, k_{p-2}} \neq \emptyset$, on the other hand, we have $\mathbf{Q}_{i_{p-3}, k_{p-3}}^{i_{p-2}, k_{p-2}} \subseteq \widehat{Q}_0^{\Theta, <}$ and therefore the equality follows from Lemma 7.2.2 (f). The proof is complete. \square

Acknowledgments. The author would like to thank Yoshiyuki Kimura for helpful comments. He was supported by JSPS Grant-in-Aid for Scientific Research (C) No. 20K03554.

REFERENCES

- [Cha95] V. Chari. Minimal affinizations of representations of quantum groups: the rank 2 case. *Publ. Res. Inst. Math. Sci.*, 31(5):873–911, 1995.
- [DLL19] B. Duan, J.-R. Li, and Y.-F. Luo. Cluster algebras and snake modules. *J. Algebra*, 519:325–377, 2019.
- [FM01] E. Frenkel and E. Mukhin. Combinatorics of q -characters of finite-dimensional representations of quantum affine algebras. *Comm. Math. Phys.*, 216(1):23–57, 2001.
- [FO21] R. Fujita and S.-j. Oh. Q -data and representation theory of untwisted quantum affine algebras. *Commun. Math. Phys.*, 384(2):1351–1407, 2021.
- [FR99] E. Frenkel and N. Reshetikhin. The q -characters of representations of quantum affine algebras and deformations of W -algebras. In *Recent developments in quantum affine algebras and related topics (Raleigh, NC, 1998)*, volume 248 of *Contemp. Math.*, pages 163–205. Amer. Math. Soc., Providence, RI, 1999.
- [Gab80] P. Gabriel. Auslander-Reiten sequences and representation-finite algebras. Representation theory I, Proc. Workshop, Ottawa 1979, Lect. Notes Math. 831, 1-71 (1980)., 1980.
- [Her06] D. Hernandez. The Kirillov-Reshetikhin conjecture and solutions of T -systems. *J. Reine Angew. Math.*, 596:63–87, 2006.
- [Her10] D. Hernandez. Kirillov-Reshetikhin conjecture: the general case. *Int. Math. Res. Not. IMRN*, (1):149–193, 2010.
- [HL10] D. Hernandez and B. Leclerc. Cluster algebras and quantum affine algebras. *Duke Math. J.*, 154(2):265–341, 2010.
- [HL15] D. Hernandez and B. Leclerc. Quantum Grothendieck rings and derived Hall algebras. *J. Reine Angew. Math.*, 2015(701):77–126, 2015.
- [HL16] D. Hernandez and B. Leclerc. A cluster algebra approach to q -characters of Kirillov-Reshetikhin modules. *J. Eur. Math. Soc. (JEMS)*, 18(5):1113–1159, 2016.

- [HO19] D. Hernandez and H. Oya. Quantum Grothendieck ring isomorphisms, cluster algebras and Kazhdan-Lusztig algorithm. *Adv. Math.*, 347:192–272, 2019.
- [Kac90] V.G. Kac. *Infinite-dimensional Lie algebras*. Cambridge University Press, Cambridge, third edition, 1990.
- [Kas91] M. Kashiwara. On crystal bases of the Q -analogue of universal enveloping algebras. *Duke Math. J.*, 63(2):465–516, 1991.
- [Kas93] M. Kashiwara. The crystal base and Littelmann’s refined Demazure character formula. *Duke Math. J.*, 71(3):839–858, 1993.
- [Kas02] M. Kashiwara. On level-zero representations of quantized affine algebras. *Duke Math. J.*, 112(1):117–175, 2002.
- [Kat14] S. Kato. Poincaré–Birkhoff–Witt bases and Khovanov–Lauda–Rouquier algebras. *Duke Math. J.*, 163(3):619–663, 2014.
- [Kim12] Y. Kimura. Quantum unipotent subgroup and dual canonical basis. *Kyoto J. Math.*, 52(2):277–331, 2012.
- [KKK15] S.-J. Kang, M. Kashiwara, and M. Kim. Symmetric quiver Hecke algebras and R -matrices of quantum affine algebras, II. *Duke Math. J.*, 164(8):1549–1602, 2015.
- [KKK18] S.-J. Kang, M. Kashiwara, and M. Kim. Symmetric quiver Hecke algebras and R -matrices of quantum affine algebras. *Invent. Math.*, 211(2):591–685, 2018.
- [KKKO15] S.-J. Kang, M. Kashiwara, M. Kim, and S.-J. Oh. Simplicity of heads and socles of tensor products. *Compos. Math.*, 151(2):377–396, 2015.
- [KKOP20] M. Kashiwara, M. Kim, S.-j. Oh, and E. Park. Monoidal categorification and quantum affine algebras. *Compos. Math.*, 156(5):1039–1077, 2020.
- [KKOP22] M. Kashiwara, M. Kim, S.-j. Oh, and E. Park. Monoidal categorification and quantum affine algebras II. *arXiv preprint arXiv:2103.10067*, 2022.
- [KKOP23] M. Kashiwara, M. Kim, S.-j. Oh, and E. Park. PBW theory for quantum affine algebras. *J. Eur. Math. Soc.*, 2023.
- [KL09] M. Khovanov and A. Lauda. A diagrammatic approach to categorification of quantum groups I. *Represent. Theory*, 13(14):309–347, 2009.
- [KNS09] A. Kuniba, T. Nakanishi, and J. Suzuki. T -systems and Y -systems for quantum affinizations of quantum Kac-Moody algebras. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 5:Paper 108, 23, 2009.
- [KP22] M. Kashiwara and E. Park. Categorical crystals for quantum affine algebras. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 2022(792):223–267, 2022.
- [Li15] J.-R. Li. On the extended T -system of type C_3 . *J. Algebr. Comb.*, 41(2):577–617, 2015.
- [LM13] J.-R. Li and E. Mukhin. Extended T -system of type G_2 . *SIGMA, Symmetry Integrability Geom. Methods Appl.*, 9:paper 054, 28, 2013.
- [Lus90] G. Luszt. Canonical bases arising from quantized enveloping algebras. *J. Amer. Math. Soc.*, 3(2):447–498, 1990.
- [Lus93] G. Luszt. *Introduction to quantum groups*, volume 110 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1993.
- [McN15] P. J McNamara. Finite dimensional representations of Khovanov–Lauda–Rouquier algebras I: finite type. *J. Reine Angew. Math.*, 2015(707):103–124, 2015.
- [MY12a] E. Mukhin and C. A. S. Young. Extended T -systems. *Selecta Math. (N.S.)*, 18(3):591–631, 2012.
- [MY12b] E. Mukhin and C. A. S. Young. Path description of type B q -characters. *Adv. Math.*, 231(2):1119–1150, 2012.
- [Nak03] H. Nakajima. t -analogs of q -characters of Kirillov-Reshetikhin modules of quantum affine algebras. *Represent. Theory*, 7:259–274 (electronic), 2003.
- [Oh19] S.-j. Oh. Auslander-Reiten quiver and representation theories related to KLR-type Schur-Weyl duality. *Math. Z.*, 291(1-2):499–554, 2019.
- [Rei97] M. Reineke. On the coloured graph structure of Luszt’s canonical basis. *Math. Ann.*, 307(4):705–723, 1997.
- [Rou08] R. Rouquier. 2-Kac-Moody algebras. *arXiv preprint arXiv:0812.5023*, 2008.

- [Rou12] R. Rouquier. Quiver Hecke algebras and 2-Lie algebras. In *Algebra colloquium*, volume 19, pages 359–410. World Scientific, 2012.
- [VV11] M. Varagnolo and E. Vasserot. Canonical bases and KLR-algebras. *J. Reine Angew. Math.*, 2011(659):67–100, 2011.

(K. Naoi) INSTITUTE OF ENGINEERING, TOKYO UNIVERSITY OF AGRICULTURE AND TECHNOLOGY,
2-24-16 NAKA-CHO, KOGANEI-SHI, TOKYO 184-8588, JAPAN

Email address: naoik@cc.tuat.ac.jp