

ON THE LEFSCHETZ PRINCIPLE FOR $GL(n, \mathbb{C})$ AND $GL(m, \mathbb{Q}_p)$

KEI YUEN CHAN AND KAYUE DANIEL WONG

ABSTRACT. We construct an exact functor from the category of Harish-Chandra modules of $GL_n(\mathbb{C})$ to the category of finite-dimensional modules of a graded Hecke algebra of type A. We show that the functor preserves parabolically induced modules, standard modules, irreducible modules, unitary modules and Dirac series. We also use the functor to connect a Bernstein-Zelevinsky type functor for graded Hecke algebra side to the tensor product for $GL_n(\mathbb{C})$ side. Some applications are also discussed.

1. INTRODUCTION

The Harish-Chandra Lefschetz principle predicts that the representation theory for real, p -adic and automorphic ones should be put in equal footing. This principle has been pursued in various situations. For example, the unitary dual problem has been exemplified by Barbasch in [Ba10], in which the correspondence between relevant K -types (real side) and relevant W -types (p -adic side) are established to match up actions of intertwining operators.

Another instance is the work of Ciubotaru-Trapa [CT11, CT12], which builds functorial connections between the categories for some real and p -adic groups, based on some Schur-Weyl type constructions of Arakawa-Suzuki [AS98] as well as the work of Oda [Od07] and Etingof-Freund-Ma [EFM09]. This allows one to transfer information between these two categories, including the unitarity of representations.

The main goal of this paper is to establish a new Schur-Weyl type duality between the category of Harish-Chandra modules of $GL_n(\mathbb{C})$ and the category of modules of graded Hecke algebra of type A. While the work is inspired by [AS98, Od07, EFM09, CT11, CT12], there are two important aspects in our work:

- Firstly, the construction for $GL_n(\mathbb{C})$ carries other subtleties such as finding a suitable choice of the Casimir element in constructing Hecke algebra actions and a suitable choice of a standard representation in defining our functor.
- Secondly, other than the unitary dual problem – which is much known now from the work of Barbasch [Ba89] for complex case (also see [St67] and [Vo86]) and Tadić [Ta86] for p -adic case, we also explore other problems of recent interest such as Dirac series [DW20], parabolic inductions [LM16] and Bernstein-Zelevinsky derivatives [Ch22+, Ch22+d] (related to branching laws).

Date: April 3, 2024.

We need more notations to explain our constructions and main results. Let \mathcal{HC}_n be the category of Harish-Chandra modules of $\mathrm{GL}_n(\mathbb{C})$. For the p -adic side, the classical result of Borel-Casselman [Bo76] allows to reduce the study of the Iwahori component of p -adic groups to the module category of affine Hecke algebras. Lusztig [Lu89] further shows that one can further reduce to study the module category of their infinitesimal version – graded Hecke algebras. In our context, we need the graded Hecke algebra of type A , see Definition 2.3. Let \mathcal{H}_m be the category of \mathbb{H}_m -modules.

Let V be the *conjugate* standard representation of $\mathrm{GL}_n(\mathbb{C})$ (see Lemma 3.1). Let K be the maximal compact subgroup in $\mathrm{GL}_n(\mathbb{C})$. In Section 3.3, we construct a functor $\Gamma_{n,m} : \mathcal{HC}_n \rightarrow \mathcal{H}_m$ whose underlying space takes the form:

$$\Gamma_{n,m}(X) = \mathrm{Hom}_K(\mathrm{triv}, X \otimes V^{\otimes m}).$$

Such functor has several nice behaviours:

Theorem 1.1. *The functor $\Gamma_{n,m}$ satisfies the following properties:*

- (Theorem 5.8) $\Gamma_{n,m}$ preserves parabolic inductions.
- (c.f. Theorem 6.4) $\Gamma_{n,m}$ sends a standard representation to a standard \mathbb{H}_m -module or zero;
- (c.f. Theorem 9.4) $\Gamma_{n,m}$ sends an irreducible module to an irreducible \mathbb{H}_m -module or zero;
- (Theorem 7.5) $\Gamma_{n,m}$ preserves unitarity.

We refer the reader to the corresponding statements for the precise meaning of the preservations in Theorem 1.1. We also remark that up to a thickening trick (see Section 9), for any irreducible Harish-Chandra module X of $\mathrm{GL}_n(\mathbb{C})$, we can find a unique m such that $\Gamma_{n,m}(X)$ is non-zero (and so irreducible). This nature is specific to $\mathrm{GL}_n(\mathbb{C})$, compared to the $\mathrm{GL}_n(\mathbb{R})$ case in [CT12].

In view of the recent development of parabolic induction on p -adic side (see e.g. [LM16]), one can transfer those properties using our functor in an effortless way (see Section 12). This avoids at least some reworking for the parabolic induction in the Harish-Chandra category, not mentioning techniques in these two categories are different. Indeed, our functor also reveals what the analogue should be, which is sometimes not completely trivial.

A refinement on the unitary dual is the Dirac series, which carries deeper structure between K -types and W -types. Combining the classification of the Dirac series in [BP11], [DW20] and [BC14], we show that up to thickening, $\Gamma_{n,m}$ also exhibits the Lefschetz principle for the Dirac series in Section 10. This can be regarded as an answer to a question posted in [BCT12, Page 200] for our setting. Extensions to some non-unitary representations with non-zero Dirac cohomology are also discussed.

One classical application of the Schur-Weyl duality is to link up the branching problem for symmetric groups S_i to the tensoring problem for $\mathfrak{gl}_n(\mathbb{C})$. To pose such analogous linkage in our context, we need to introduce two more ingredients. Firstly, using a natural subalgebra $\mathbb{C}[S_i]$ in \mathbb{H}_m , we can define a functor: for an irreducible representation τ of S_i ,

$$\mathbf{BZ}_\tau : \mathcal{H}_n \rightarrow \mathcal{H}_{n-i}; \quad \mathbf{BZ}_\tau(\pi) = \mathrm{Hom}_{S_i}(\tau, \pi),$$

see Section 8.1 for precise descriptions. Secondly, we define the Schur functor $\mathbb{S}_\tau(V) = \text{Hom}_{S_i}(\tau, V^{\otimes i})$ as a $GL_n(\mathbb{C})$ -representation, which is irreducible and can be computed explicitly from the classical Schur-Weyl duality. (Here we use the convention that S_i acts on $V^{\otimes i}$ by a sign permutation.) This gives the tensoring functor $T_\tau : \mathcal{HC}_n \rightarrow \mathcal{HC}_n$; $X \mapsto X \otimes \mathbb{S}_\tau(V)$. Our main result is to relate these two functors:

Theorem 1.2. (*=Theorem 8.2*) *Let $\tau \in \text{Irr}(S_i)$. For $X \in \mathcal{HC}_n$, there is a natural isomorphism:*

$$\mathbf{BZ}_\tau \circ \Gamma_{n,m+i}(X) \cong \Gamma_{n,m} \circ T_\tau(X).$$

We call \mathbf{BZ}_τ to be a generalized Bernstein-Zelevinsky functor, reflecting our original viewpoint from p -adic groups in [CS19, Ch22+]. One can then translate recent results in [Ch22+, Ch22+c] to \mathcal{HC}_n , see Section 11. It is an interesting problem to extend Theorem 1.2 to constructions of other classical groups such as [CT11] and [Ca22].

Another problem of our interest is the branching laws, whose Lefschetz principle is also investigated in [Ch23]. However, the branching law is dealt with in the category of Casselman-Wallach representations and irreducible representations could become non-admissible after restriction. This makes harder for using the functor $\Gamma_{n,m}$ to study branching law directly. On the other hand, as seen in this article, results on parabolic inductions and Jacquet functors in [Ch22+b, Ch22+c, Ch22+d] can be transferred in some form and this opens up possible applications to some branching problems, see e.g. Remarks 11.5 and 12.4.

1.1. Acknowledgements. This project is supported by Seed Fund Basic Research for New Staff (Project Code: 2201102282), the HKU startup grant, and the National Natural Science Foundation of China (grant 12371033).

2. PRELIMINARIES

In this section, we introduce some basic notations. We also review the representation theory for $GL_n(\mathbb{C})$ and \mathbb{H}_m , which also demonstrates some similarities between their representation theories.

2.1. Complex groups. Let $G = GL_n(\mathbb{C})$ be a complex Lie group treated as a real group, with a maximal compact subgroup $K = U(n)$ consisting of unitary matrices i.e.

$$U(n) := \left\{ E \in G : \overline{E}^t E = I_n \right\},$$

where \overline{E} is the complex conjugation of E and E^t is the transpose of E . Let B be the Borel subgroup of upper triangular matrices in $GL_n(\mathbb{C})$. Let H be the subgroup of diagonal matrices in $GL_n(\mathbb{C})$. Let $H = TA$ be the Cartan decomposition of H so that $T \cong (S^1)^{\times n}$ and $A \cong (\mathbb{R}^\times)^{\times n}$. Write $\mathfrak{g}_0 = \mathfrak{gl}_n(\mathbb{C})$, \mathfrak{k}_0 , \mathfrak{h}_0 , \mathfrak{t}_0 and \mathfrak{a}_0 as their Lie algebras, and remove the subscripts for their complexifications. (Those notions for Lie algebras depend on n , but it will be clear from the context.) The group S_n acts on \mathfrak{h}_0 by permuting the coordinates. We use j to denote the action of $\sqrt{-1}$ coming from the complexification (for instance, one has $\mathfrak{g} = \{P + jQ \mid P, Q \in \mathfrak{g}_0\}$).

Following [Du75] and [Vo81, Section 7.1], we make the following identifications for \mathfrak{g} and its subalgebras: Define $\phi^L, \phi^R : \mathfrak{g}_0 \rightarrow \mathfrak{g}$ by:

$$(1) \quad \phi^L(E) := \frac{1}{2}(E - j i E), \quad \phi^R(E) := \frac{1}{2}(\overline{E} + j i \overline{E}),$$

where i is $\sqrt{-1}I_n$ in \mathfrak{g}_0 . Then one can easily check there is an isomorphism:

$$(2) \quad \mathfrak{g}_0 \oplus \mathfrak{g}_0 \cong \mathfrak{g}.$$

given by $(E, E') \mapsto \phi^L(E) + \phi^R(E')$. Using the isomorphism, all elements in \mathfrak{g} will be written as $(E, E') \in \mathfrak{g}_0 \oplus \mathfrak{g}_0$ from now on. In particular, for $A, B \in \mathfrak{k}_0$, $A + jB = \phi^L(A + iB) + \phi^R(\overline{A - iB}) \in \mathfrak{k}$ corresponds to

$$(A + iB, \overline{A - iB}) = (A + iB, \overline{A} + i\overline{B}) = (A + iB, -A^t - iB^t) = (A + iB, -(A + iB)^t),$$

since $A^t = -\overline{A}$ and $B^t = -\overline{B}$. So we have the identifications

$$(3) \quad \mathfrak{k} \cong \{(E, -E^t) : E \in \mathfrak{g}_0\}, \quad \mathfrak{t} \cong \{(H, -H) : H \in \mathfrak{h}_0\}, \quad \mathfrak{a} \cong \{(H, H) : H \in \mathfrak{h}_0\}.$$

2.2. Representation theory of $\mathrm{GL}_n(\mathbb{C})$. Recall that \mathcal{HC}_n is the category of Harish-Chandra modules (a.k.a. (\mathfrak{g}, K) -modules) of $\mathrm{GL}_n(\mathbb{C})$ in the sense of [Vo81, Definiton 0.3.8]. Note that for any Harish-Chandra module X in \mathcal{HC}_n , the actions of $\phi^L(E)$ and $\phi^L(iE)$ are differed by the scalar $\sqrt{-1}$. The same holds if we replace ϕ^L by ϕ^R .

Let $U(\mathfrak{g}_0 \oplus \mathfrak{g}_0)$ be the universal enveloping algebra of $\mathfrak{g}_0 \oplus \mathfrak{g}_0$ over \mathbb{C} . Using the above discussion, we can and shall regard X as a $U(\mathfrak{g}_0 \oplus \mathfrak{g}_0)$ -module determined by the action:

$$(E_1, E_2) \cdot v = (\phi^L(E_1) + \phi^R(E_2)) \cdot v$$

for $v \in X$ and $(E_1, E_2) \in \mathfrak{g}_0 \oplus \mathfrak{g}_0$.

For $n_1 + n_2 = n$, let $G = \mathrm{GL}_{n_1+n_2}(\mathbb{C})$ and let $P = P_{n_1, n_2}$ be the parabolic subgroup containing matrices of the form

$$\begin{pmatrix} g_1 & * \\ & g_2 \end{pmatrix}$$

for $g_i \in \mathrm{GL}_{n_i}(\mathbb{C})$ ($i = 1, 2$). For a Casselman-Wallach representation τ of P , let $\mathrm{Ind}_P^G(\tau)$ be the space of smooth functions from G to τ satisfying: for $p \in P$ and $k \in K$,

$$f(pk) = \delta^{1/2}(p)(p \cdot f(k)),$$

where δ is the modular character of P . For a Harish Chandra module τ of P , by abuse of notations, we also write $\mathrm{Ind}_P^G(\tau)$ to be $\mathrm{Ind}_P^G(\tilde{\tau})$ for some Fréchet globalization $\tilde{\tau}$ of τ .

For Y_1 in \mathcal{HC}_{n_1} and Y_2 in \mathcal{HC}_{n_2} , define the real parabolic induction as:

$$Y_1 \times Y_2 = \mathrm{Ind}_P^G(Y_1 \boxtimes Y_2)_{K\text{-finite}},$$

which is regarded as a representation in $\mathcal{HC}_{n_1+n_2}$. It follows from [Knp86, Page 179] that the product is an associative operation.

Definition 2.1. For any $a, b \in \mathbb{C}$ satisfying $a - b \in \mathbb{Z}$, $\chi_{a,b}$ is a $GL_1(\mathbb{C})$ -character given by

$$(4) \quad \chi_{a,b}(z) = z^a \bar{z}^b.$$

Let $\lambda_L = (\lambda_{L,1}, \dots, \lambda_{L,n})$, $\lambda_R = (\lambda_{R,1}, \dots, \lambda_{R,n}) \in \mathfrak{h}_0^* \cong \mathbb{C}^n$ be such that $\lambda_{L,i} - \lambda_{R,i} \in \mathbb{Z}$ for all i .

The **principal series (representation)** $X(\lambda_L, \lambda_R)$ is defined as:

$$(5) \quad X(\lambda_L, \lambda_R) = \chi_{\lambda_{L,1}, \lambda_{R,1}} \times \chi_{\lambda_{L,2}, \lambda_{R,2}} \times \cdots \times \chi_{\lambda_{L,n}, \lambda_{R,n}}.$$

Write $\mu := \lambda_L - \lambda_R$ and $\nu := \lambda_L + \lambda_R$. Using the identification (3), it can also be written as:

$$X(\lambda_L, \lambda_R) = \text{Ind}_B^G(\mathbb{C}_\mu \boxtimes \mathbb{C}_\nu \boxtimes 1)_{K\text{-finite}},$$

where we write $B = TAN$ with N being the subgroup of unipotent upper triangular matrices.

The K -type with extremal weight μ occurs exactly once in $X(\lambda_L, \lambda_R)$. Let $J(\lambda_L, \lambda_R)$ be the unique irreducible subquotient of $X(\lambda_L, \lambda_R)$ containing this K -type (see [PRV67]).

The principal series plays a more prominent role in the representation theory of complex Lie groups since it constructs all irreducible Harish-Chandra modules in a nice manner:

Theorem 2.2 ([Zh74]). *Retain the above notations. Then the following statements hold:*

- (a) *Every irreducible Harish-Chandra module is of the form $J(\lambda_L, \lambda_R)$.*
- (b) *Two such modules $J(\lambda_L, \lambda_R)$ and $J(\lambda'_L, \lambda'_R)$ are equivalent if and only if there exists $w \in S_n$ such that $w\lambda_L = \lambda'_L$ and $w\lambda_R = \lambda'_R$.*

We say $X(\lambda_L, \lambda_R)$ is a **standard module** if $\text{Re}(\nu) = \text{Re}(\lambda_L + \lambda_R)$ is dominant i.e.

$$\text{Re}(\lambda_{L,1} + \lambda_{R,1}) \geq \dots \geq \text{Re}(\lambda_{L,n} + \lambda_{R,n}).$$

It is well-known (see e.g. [Du75, Théorème I.4.2], [Knp86, Theorem 8.54]) that if $X(\lambda_L, \lambda_R)$ is standard, then it has a unique maximal proper submodule, so that $J(\lambda_L, \lambda_R)$ appears as a unique quotient of $X(\lambda_L, \lambda_R)$. The notion of standard representations is more convenient (than principal series) when later we have to compare with the modules of graded Hecke algebras (defined in next section).

2.3. Graded Hecke algebras.

Definition 2.3. The graded Hecke algebra \mathbb{H}_m of type A is the associative unital algebra over \mathbb{C} with generators y_1, \dots, y_m and s_1, \dots, s_{m-1} such that

- $y_i y_j = y_j y_i$ for any i, j ;
- $s_i^2 = 1$ for all i ;
- $s_i s_j = s_j s_i$ for $|i - j| > 1$;
- $s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$;
- $s_i y_i - y_{i+1} s_i = 1$;
- $s_i y_j - y_j s_i = 0$ if $j \neq i, i + 1$.

Let S_m be the symmetric group permuting m elements. Note that s_1, \dots, s_{m-1} generate the group algebra $\mathbb{C}[S_m]$. Thus, for $w \in S_m$, we also regard as an element in \mathbb{H}_m via the natural embedding from $\mathbb{C}[S_m]$ to \mathbb{H}_m .

There is a natural embedding of $\mathbb{H}_{m_1} \otimes \mathbb{H}_{m_2}$ to $\mathbb{H}_{m_1+m_2}$ via the maps: for $i = 1, \dots, m_1$ and $j = 1, \dots, m_2$,

$$y_i \otimes 1 \mapsto y_i, \quad 1 \otimes y_j \mapsto y_{m_1+j},$$

for $i = 1, \dots, m_1 - 1$ and $j = 1, \dots, m_2 - 1$,

$$s_i \otimes 1 \mapsto s_i, \quad 1 \otimes s_j \mapsto s_{m_1+j}.$$

For \mathbb{H}_{m_1} -module π_1 and \mathbb{H}_{m_2} -module π_2 , we write

$$\pi_1 \times \pi_2 := \mathbb{H}_{m_1+m_2} \otimes_{\mathbb{H}_{m_1} \otimes \mathbb{H}_{m_2}} (\pi_1 \boxtimes \pi_2).$$

The associativity of this product follows from the associativity of tensor products and that standard fact that the tensoring $\mathbb{A} \otimes_{\mathbb{A}}$ is the identity functor for any algebra with an unit.

2.4. Representation theory of \mathbb{H}_m . The classification for irreducible \mathbb{H}_m -modules is known for long time, see e.g. [Ze80, Ro86]. Let \mathcal{H}_m be the category of finite-dimensional \mathbb{H}_m -modules.

A segment is of the form $[a, b]$ for some $a, b \in \mathbb{C}$ with $b - a \geq 0$. We shall consider a segment $[a, b]$ to be a set $\{a, \dots, b\}$. A multisegment is a multiset of non-empty segments. As our convenience, set $[a, a - 1] = \emptyset$, and a segment can be an empty set and a multisegment can also be an empty set.

Two segments Δ_1 and Δ_2 are said to be linked if $\Delta_1 \cup \Delta_2$ is still a segment, and $\Delta_1 \not\subset \Delta_2$ and $\Delta_2 \not\subset \Delta_1$. For two segments $\Delta_1 = [a_1, b_1], \Delta_2 = [a_2, b_2]$, we write $\Delta_1 < \Delta_2$ if Δ_1 and Δ_2 are linked and $a_1 < a_2$.

For $c \in \mathbb{C}$, define χ_c to be a character on $\mathbb{H}_1 = \mathbb{C}[y]$ such that $\psi_c(y) = c$. For each segment $\Delta = [a, b]$, define $\text{St}(\Delta)$ to be the unique simple quotient of

$$\psi_a \times \psi_{a+1} \times \dots \times \psi_b$$

Moreover, $\text{St}(\Delta)$ is one-dimensional and is the sign representation as a S_n -representation. A standard property on parabolic inductions is that: if Δ_1 and Δ_2 are not linked, then

$$(6) \quad \text{St}(\Delta_1) \times \text{St}(\Delta_2) \cong \text{St}(\Delta_2) \times \text{St}(\Delta_1).$$

For a multisegment $\mathbf{m} = \{\Delta_1, \dots, \Delta_k\}$, we label the segments such that

$$(7) \quad \Delta_1 \not\prec \dots \not\prec \Delta_k.$$

Then, let

$$\lambda(\mathbf{m}) := \text{St}(\Delta_1) \times \dots \times \text{St}(\Delta_k).$$

This is called a **standard module**, and the unique simple quotient of $\lambda(\mathbf{m})$ is denoted by $\text{St}(\mathbf{m})$. It follows from (6) that $\lambda(\mathbf{m})$ is independent of a choice of an order of segments in \mathbf{m} . In particular, for writing each segment in \mathbf{m} as $\Delta_i = [a_i, b_i]$, we can choose an ordering satisfying (7) by fixing:

$$\text{Re}(a_1) \geq \text{Re}(a_2) \geq \dots \geq \text{Re}(a_n).$$

Standard modules construct all irreducible \mathbb{H}_m -modules in the following sense:

Theorem 2.4. [Ze80]

- (1) For any irreducible \mathbb{H}_m -module π , there is a multisegment \mathbf{m} such that $\pi \cong \mathrm{St}(\mathbf{m})$.
- (2) For two multisegments $\mathbf{m}_1, \mathbf{m}_2$, if $\mathrm{St}(\mathbf{m}_1) \cong \mathrm{St}(\mathbf{m}_2)$, then $\mathbf{m}_1 = \mathbf{m}_2$.

Remark 2.5. One may notice there is an alternate way to define irreducible \mathbb{H}_m -modules in [Ro85] by using a uniqueness of some S_m -types. This supplements a parallel story (i.e. Lefschetz principle) to using K -types in defining irreducible Harish-Chandra modules in Definition 2.1.

3. ARAKAWA-SUZUKI TYPE FUNCTORS

3.1. ‘Standard’ representations. The definition of our Arakawa-Suzuki type functor requires a choice of a ‘standard representation’ V of $\mathrm{GL}_n(\mathbb{C})$. For reasons that become obvious later in this paper (see Section 5.2 below), we make the following choice of the standard representation:

Lemma 3.1. *Let $V \cong \mathbb{C}^n$ be the conjugate standard representation of $\mathrm{GL}_n(\mathbb{C})$ i.e. $g \in \mathrm{GL}_n(\mathbb{C})$ acting on V by the matrix multiplication of \bar{g} . Then $V = J(\rho, e_1 + \rho)$, where ρ is half the sum of positive roots in the root system determined by H and B , and $\{e_i\}_{1 \leq i \leq n}$ is the standard basis of \mathbb{C}^n .*

Proof. Let $\xi : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be the (complexified) derivative of the conjugate standard representation, that is, for all $X, Y \in \mathfrak{g}_0$, $\xi(X + jY) := \bar{X} + i\bar{Y} \in \mathfrak{gl}(V)$. Under the isomorphism $\mathfrak{g}_0 \oplus \mathfrak{g}_0 \cong \mathfrak{g}$ given in (2), one has

$$\begin{aligned}\xi(P, 0) &= \frac{1}{2}\xi(P - jiP) = \frac{1}{2}(\bar{P} - i(-i\bar{P})) = 0 \\ \xi(0, Q) &= \frac{1}{2}\xi(\bar{Q} + ji\bar{Q}) = \frac{1}{2}(Q + i(-iQ)) = Q\end{aligned}$$

So V is trivial (with infinitesimal character ρ) on the first copy of \mathfrak{g}_0 , and is the standard representation (with infinitesimal character $e_1 + \rho$) on the second copy of \mathfrak{g}_0 . \square

3.2. Casimir element. Let $E_{ij} \in \mathfrak{g}_0$ be the $n \times n$ matrix with 1 on the (i, j) -entry and 0 on the other entries. For $0 \leq k < l \leq m$, define the Casimir element:

$$\Omega_{kl} := \sum_{1 \leq i, j \leq n} 1^{\otimes k} \otimes E_{ij} \otimes 1^{\otimes l-k-1} \otimes E_{ji} \otimes 1^{m-l} \in U(\mathfrak{g}_0)^{\otimes(m+1)}.$$

Here 1 is the unit in $U(\mathfrak{g}_0)$.

Note that $\{E_{ji}\}$ is a dual basis for $\{E_{ij}\}$ under the pairing:

$$(E, E') \mapsto \mathrm{tr}(EE'^t)$$

and so for any $g \in \mathrm{GL}_n(\mathbb{C})$, Ω_{kl} is invariant under the adjoint action of $\mathrm{GL}_n(\mathbb{C})$:

$$(8) \quad \mathrm{Ad}(g)(\Omega_{kl}) := \sum_{1 \leq i, j \leq n} 1^{\otimes k} \otimes gE_{ij}g^{-1} \otimes 1^{\otimes(l-k-1)}gE_{ji}g^{-1} \otimes 1^{\otimes(m-l)} = \Omega_{kl}.$$

3.3. Arakawa-Suzuki realization of \mathbb{H}_m .

Theorem 3.2. [AS98] *There is an injective algebra (over \mathbb{C}) homomorphism $\Theta : \mathbb{H}_m \longrightarrow U(\mathfrak{g}_0)^{\otimes(m+1)}$ defined by:*

$$\begin{cases} \Theta(s_i) := -\Omega_{i,i+1}, & 1 \leq i < m; \\ \Theta(y_l) := \sum_{0 \leq x < l} \Omega_{x,l} + \frac{n}{2}(1^{\otimes(m+1)}), & 1 \leq l \leq m. \end{cases}$$

We shall write $\Theta_{\mathbb{H}_m}$ for Θ if we have to specify the underlying graded Hecke algebra.

Recall that V is defined in Lemma 3.1. Let X be in \mathcal{HC}_n . Define the algebra homomorphism $\Lambda : U(\mathfrak{g}_0)^{\otimes(m+1)} \rightarrow \text{End}(X \otimes V^{\otimes m})$ determined by:

$$(9) \quad \Lambda(1^{\otimes k} \otimes E \otimes 1^{\otimes(m-k)})(\mathbf{x} \otimes v_1 \otimes \dots \otimes v_m) = \mathbf{x} \otimes v_1 \otimes \dots \otimes (0, E) \cdot v_k \otimes \dots \otimes v_n,$$

where $(0, E)$ is the corresponding complexified Lie algebra action on V . (When $k = 0$, the copy $(0, E)$ acts on \mathbf{x} .)

Definition 3.3. (c.f. [AS98, CT12]) Let X be in \mathcal{HC}_n , and V be the conjugate standard representation as in Lemma 3.1. We define the Arakawa-Suzuki type functor:

$$\Gamma_{n,m} : \mathcal{HC}_n \longrightarrow \mathcal{H}_m$$

to be the exact covariant functor given by

$$(10) \quad \Gamma_{n,m}(X) := \text{Hom}_K(\text{triv}, X \otimes V^{\otimes m}).$$

where the \mathbb{H}_m -action on $\Gamma_{n,m}(X)$ is given by the map

$$\Lambda \circ \Theta : \mathbb{H}_m \rightarrow \text{End}_{\mathbb{C}}(X \otimes V^{\otimes m}).$$

We remark that it follows from (8) that the action of \mathbb{H}_m in Definition 3.3 is well-defined i.e. $(\Lambda \circ \Theta(h))(\Gamma_{n,m}(X)) \subset \Gamma_{n,m}(X)$ for any $h \in \mathbb{H}_m$.

3.4. Another formulation. Define \bar{V} to be the contragredient of the standard representation of $\text{GL}_n(\mathbb{C})$. Then one can check as in Lemma 3.1 that $\bar{V} = J(-e_n + \rho, \rho)$.

We define $\zeta : \mathfrak{g}_0 \oplus \mathfrak{g}_0 \rightarrow \mathfrak{g}_0 \oplus \mathfrak{g}_0$ by:

$$\zeta(E, E') = (-E', -E).$$

This comes from the anti-involution $g \mapsto \bar{g}^{-t}$ and induces an auto-equivalence of categories on \mathcal{HC}_n , still denoted by ζ . Define $\bar{\zeta} : \mathfrak{h}_0 \oplus \mathfrak{h}_0 \rightarrow \mathfrak{h}_0 \oplus \mathfrak{h}_0$ by $\bar{\zeta}(\lambda_L, \lambda_R) = (-\lambda_R, -\lambda_L)$ so that

$$\zeta(J(\lambda_L, \lambda_R)) = J(\bar{\zeta}(\lambda_L, \lambda_R)).$$

We also note that $\zeta(V) = \bar{V}$.

Define

$$\bar{\Lambda} : U(\mathfrak{g}_0)^{\otimes(m+1)} \rightarrow \text{End}_{\mathbb{C}}(X \otimes \bar{V}^{\otimes m})$$

given by

$$\bar{\Lambda}(1^{\otimes k} \otimes E \otimes 1^{\otimes(m-k)})(\mathbf{x} \otimes v_1 \otimes \dots \otimes v_m) = \mathbf{x} \otimes v_1 \otimes \dots \otimes (E, 0) \cdot v_k \otimes \dots \otimes v_n.$$

and define the dual functor $\bar{\Gamma}_{n,m} : \mathcal{HC}_n \rightarrow \mathcal{H}_m$ given by

$$\bar{\Gamma}_{n,m}(X) = \text{Hom}_K(\text{triv}, X \otimes \bar{V}^{\otimes m})$$

with the action of \mathbb{H}_m given by $\bar{\Lambda} \circ \Theta$. As a result, we have:

$$\bar{\Gamma}_{n,m} \circ \zeta \cong \Gamma_{n,m}.$$

3.5. Other Arakawa-Suzuki type functor involving the Bernstein-Gelfand functor. We discuss another way to define a Arakawa-Suzuki type functor for \mathcal{HC}_n . We first recall a functor due to Bernstein-Gelfand [BG80] connecting to the BGG category \mathcal{O}_n for $\mathfrak{gl}_n(\mathbb{C})$. To facilitate the setup in [BG80], we shall identify, \mathcal{HC}_n with the category \mathcal{HC}_n^b of Harish-Chandra $U(\mathfrak{g}_0)$ -bimodules i.e. the category of $(U(\mathfrak{g}_0), U(\mathfrak{g}_0))$ -modules with a local \mathfrak{k} -finiteness condition (in the sense of [BG80, Section 5.1]). See [BG80, Appendix II] for more details on such identification. We shall now define another functor from some subcategory of \mathcal{HC}_n^b to \mathcal{H}_m .

Let $\chi \in \mathfrak{h}_0^*$ be a dominant character. Then χ determines the central character $\chi^* : Z(\mathfrak{g}_0) \rightarrow \mathbb{C}$ given by $\chi^*(z) := (\chi - \rho)(\text{pr}(z))$, where $\text{pr} : U(\mathfrak{g}_0) \rightarrow U(\mathfrak{h}_0)$ by setting other PBW monomials zero. Let $\mathcal{HC}_n^{R,\chi}$ be the subcategory of \mathcal{HC}_n^b such that all $X \in \mathcal{HC}_n^{R,\chi}$ is annihilated by $\phi^R(\ker(\chi^*))$. For instance, for any $\xi \in \mathfrak{h}_0^*$ with $\xi - \chi$ to be integral, all composition factors of the principal series representation $X(\xi, \chi)$ are in $\mathcal{HC}_n^{R,\chi}$.

Define the functor $T_\chi : \mathcal{HC}_n^{R,\chi} \rightarrow \mathcal{O}_n$ by

$$T_\chi(X) := X \otimes_{U(\mathfrak{g}_0)} M(\chi),$$

where $M(\chi)$ is the Verma module with highest weight $\chi - \rho$. We also denote by $F_{\chi'}$ the Arakawa-Suzuki functor in [AS98]. Then we can obtain a functor $F_{\chi'} \circ T_\chi$ from $\mathcal{HC}_n^{R,\chi}$ to \mathcal{H}_m . It is an interesting problem to compare $F_{\chi'} \circ T_\chi$ with $\bar{\Gamma}_{n,m}$, which goes beyond our study in this article. On the other hand, our functor is defined in \mathcal{HC}_n and is applicable for some wider potential applications (see Section 12.2).

4. COMPUTATIONS ON THE ACTIONS OF \mathbb{H}_m ON INDUCED MODULES

We shall compute some actions of \mathbb{H}_n on $\Gamma_{n,m}(X)$ for some parabolically induced modules, which will be used to prove Theorem 5.7 in the next section.

4.1. Some identifications. We identify some spaces, which will be useful in defining some generators for \mathbb{H}_m under $\Gamma_{n,m}$ (see Section 4.3). For a parabolic subgroup P of $GL_n(\mathbb{C})$ containing B and a Harish-Chandra module τ of P ,

- $\text{Ind}_P^G(\tau) \otimes V^{\otimes m} \cong \text{Ind}_P^G(\tau \otimes V^{\otimes m})$, as $GL_n(\mathbb{C})$ -representations, via the map: for $f \in \text{Ind}_P^G(\tau)$,

$$f \otimes v_1 \otimes \dots \otimes v_m \xrightarrow{\Phi} (g \mapsto f(g) \otimes g \cdot v_1 \otimes \dots \otimes g \cdot v_m).$$

This is a bijection, see for example [Tr06, Theorem 44.1].

- $(\text{Ind}_P^G(\tau))^K \cong \tau^{K_P}$, where $K_P = K \cap P$. The natural map is given by the restriction and the inverse map is given by: for $v \in \tau^{K_P}$, define $f \in \text{Ind}_P^G \tau$ as:

$$f(pk) = p \cdot v.$$

It is straightforward to check that last map is well-defined.

4.2. Computation via differentiations. We use notations in the previous section. For $E \in \mathfrak{g}_0$ and $f \in \text{Ind}_P^G(\tau)$, define:

$$(E.f)(g) = \left. \frac{d}{ds} f(g \cdot \exp(sE)) \right|_{s=0},$$

and $((jE).f)(g) = \sqrt{-1}((E.f)(g))$. (Here \exp is the usual exponential map.)

Again we identify $\mathfrak{g}_0 \oplus \mathfrak{g}_0$ with \mathfrak{g} and so for $(E, E') \in \mathfrak{g}_0 \oplus \mathfrak{g}_0$, we mean:

$$((E, E').f)(g) = (\phi^L(E).f)(g) + (\phi^R(E').f)(g).$$

We shall frequently use these formulas in our computations in Sections 4.4 to 4.7.

4.3. Cyclic subspace. Let τ be a Harish-Chandra module of $\text{GL}_{n_1}(\mathbb{C}) \times \text{GL}_{n_2}(\mathbb{C})$. Let $n = n_1 + n_2$ and $m = m_1 + m_2$. Note that $\tau \otimes V^{\otimes m}$ contains a subspace spanned by vectors of the form:

$$\mathbf{x} \otimes v_1 \otimes \dots \otimes v_{m_1} \otimes v_{m_1+1} \otimes \dots \otimes v_{m_1+m_2}$$

for $\mathbf{x} \in \tau$, $v_1, \dots, v_{m_1} \in \mathbb{C}^{n_1} \subset \mathbb{C}^n \cong V$ and $v_{m_1+1}, \dots, v_{m_1+m_2} \in \mathbb{C}^{n_2} \subset \mathbb{C}^n \cong V$. Here, the first inclusion by sending \mathbb{C}^{n_1} to the first n_1 -coordinates and the second inclusion by sending \mathbb{C}^{n_2} to the last n_2 -coordinates of V .

We shall denote such subspace by $\mathcal{W}'(\tau, n_1, n_2, m_1, m_2)$. Let $P = P_{n_1, n_2}$. Let $\mathcal{W}(\tau, n_1, n_2, m_1, m_2)$ be the subspace of $\text{Ind}_P^G(\tau \otimes V^{\otimes(m_1+m_2)})$ containing all functions satisfying the properties:

- (1) $F(1) \in \mathcal{W}'(\tau, n_1, n_2, m_1, m_2)$; and
- (2) $F(k) = F(1)$ for all $k \in U(n)$.

Note that (1) and (2) above also imply that $F(1) \in (\tau \otimes V^{\otimes m})^{U(n_1) \times U(n_2)}$ since $P \cap K = U(n_1) \times U(n_2)$.

4.4. Basic formulas. We shall keep using the notations in Section 4.3 until Section 4.7. For each element $E \in \mathfrak{g}_0$, and for an integer k , we define

$$\Delta_k(E) = 1^{\otimes k} \otimes (0, E) \otimes 1^{\otimes(m-k)}$$

and for $k < l$,

$$\Delta_{kl}(E) = 1^{\otimes k} \otimes (0, E) \otimes 1^{\otimes(l-k-1)} \otimes (0, E^t) \otimes 1^{\otimes(m-l)}.$$

By abuse of notations, we write: for $F \in \text{Ind}_P^G(\tau \otimes V^{\otimes m})$,

$$\Delta_l(E) \cdot F := \Phi(\Delta_l(E) \cdot \Phi^{-1}(F)), \quad \Delta_{kl}(E) \cdot F := \Phi(\Delta_{kl}(E) \cdot \Phi^{-1}(F)),$$

where Φ is the isomorphism given in Section 4.1 and the $\Delta_l(E)$ action is defined in (9) (via Λ). We similarly define for $\Delta_{kl}(E) \cdot F$.

For $g \in GL_n(\mathbb{C})$ and $F \in \text{Ind}_P^G(\tau \otimes V^{\otimes m})$, we shall write

$$(g \cdot F)(h) = F(hg).$$

For $(E, E') \in \mathfrak{g}_0 \oplus \mathfrak{g}_0$, we write $(E, E') \cdot F$ to be the induced Lie algebra action.

Lemma 4.1. *Let $F \in \text{Ind}_P^G(\tau \otimes V^{\otimes m})$ with $F(1) = \sum_i \mathbf{x}_i \otimes v_{i,1} \otimes \dots \otimes v_{i,m}$. For $r \geq 1$, one has*

$$(\Delta_r(E) \cdot F)(1) = \sum_i \mathbf{x}_i \otimes v_{i,1} \otimes \dots \otimes (0, E) \cdot v_{i,r} \otimes \dots \otimes v_{i,m}.$$

Proof. We consider $f \otimes v_1 \otimes \dots \otimes v_m \in \text{Ind}_P^G(\tau) \otimes V^{\otimes m}$. Let $F' = \Phi(f \otimes v_1 \otimes \dots \otimes v_m)$. Then

$$(\Delta_r(E) \cdot F')(g) = f(g) \otimes g \cdot v_1 \otimes \dots \otimes (g \cdot (0, E) \cdot v_r) \otimes \dots \otimes g \cdot v_m$$

Now, evaluating at $g = 1$, we have:

$$(\Delta_r(E) \cdot F')(1) = f(1) \otimes v_1 \otimes \dots \otimes (0, E) \cdot v_r \otimes \dots \otimes v_m.$$

This gives the equality for F' . Since Φ is an isomorphism (see Section 4.1), we can now extend linearly. \square

Lemma 4.2. $(\Delta_0(E) \cdot F)(1) = ((0, E) \cdot F)(1) - (\Delta_1(E) \cdot F)(1) - \dots - (\Delta_m(E) \cdot F)(1).$

Proof. We pick $f \in \text{Ind}_P^G(\tau)$ and let $F = \Phi(f \otimes v_1 \otimes \dots \otimes v_m)$. Set s to be a variable. We have:

$$\begin{aligned} & (\exp(s(0, E)) \cdot F)(g) \\ &= f(g \cdot \exp(s(0, E))) \otimes g \cdot \exp(s(0, E)) \cdot v_1 \otimes \dots \otimes g \cdot \exp(s(0, E)) \cdot v_m \end{aligned}$$

Taking differentiation and then evaluating at $s = 0$ yields:

$$\begin{aligned} ((0, E) \cdot F)(g) &= ((0, E) \cdot f)(g) \otimes g \cdot v_1 \otimes \dots \otimes g \cdot v_m + f(g) \otimes g \cdot (0, E) \cdot v_1 \otimes \dots \otimes g \cdot v_m + \\ &\dots + f(g) \otimes g \cdot v_1 \otimes \dots \otimes g \cdot (0, E) \cdot v_m \end{aligned}$$

Note that the first term is equal to $\Phi(\Delta_0(E) \cdot (f \otimes v_1 \otimes \dots \otimes v_m))$. Then evaluating at $g = 1$, it is equal to $(\Delta_0(E) \cdot F)(1)$.

Starting from the second term, we need to do evaluations at $g = 1$ to see that the corresponding term is equal to:

$$f(1) \otimes v_1 \otimes \dots \otimes (0, E) \cdot v_r \otimes \dots \otimes v_m,$$

and so is equal to $(\Delta_r(E) \cdot F)(1)$.

Now, we rearrange the second and higher terms to the LHS, we obtain the formula in the lemma for $F = \Phi(f \otimes v_1 \otimes \dots \otimes v_m)$. But, Φ is an isomorphism (see Section 4.1), and so we also have the general case. \square

4.5. Computing some actions on torus part. For simplicity, set $\widetilde{\mathcal{W}} = \mathcal{W}(\tau, n_1, n_2, m_1, m_2)$.

Lemma 4.3. *Let $F \in \widetilde{\mathcal{W}}$. For $y \leq n_1$ and $m_1 + 1 \leq l$ and any k ,*

$$(\Delta_{kl}(E_{yy}) \cdot F)(1) = 0.$$

Proof. This follows from that E_{yy} acts by zero on the l -th copy of $\mathbb{C}^{n_2} \subset V$. □

Lemma 4.4. *Let $F \in \widetilde{\mathcal{W}}$. For $n_1 + 1 \leq y$ and $1 \leq k \leq m_1$ and any l with $k < l$,*

$$(\Delta_{kl}(E_{yy}) \cdot F)(1) = 0.$$

Proof. This follows from E_{yy} acts by zero on the k -th copy of $\mathbb{C}^{n_1} \subset V$. □

4.6. Computing some actions on unipotent upper triangular part. We first consider the action arising from the elements of the form $\begin{pmatrix} 0_{n_1} & * \\ & 0_{n_2} \end{pmatrix}$:

Lemma 4.5. *Let $F \in \widetilde{\mathcal{W}}$. For $y \leq n_1 < z$,*

$$(\Delta_{kl}(E_{yz}) \cdot F)(1) = 0$$

for all $0 \leq k < l$.

Proof. We first assume that $k \geq 1$. We divide into three cases:

- Case 1: $k < l \leq m_1$. This follows from that E_{yz} acts by zero on the k -th copy of V (which takes the form $\mathbb{C}^{n_1} \subset V$).
- Case 2: $m_1 + 1 \leq k < l$. This follows from that $E_{yz}^t = E_{zy}$ acts by zero on the l -th copy of V (which takes the form $\mathbb{C}^{n_2} \subset V$).
- Case 3: $k \leq m_1 < l$. This is similar to Case 2.

We now assume that $k = 0$. This follows from Lemma 4.2, since unipotent elements act as the identity on τ . □

We now consider upper triangular matrices of the form:

$$\begin{pmatrix} * & 0 \\ 0 & 0_{n_2} \end{pmatrix}.$$

Lemma 4.6. *Let $F \in \widetilde{\mathcal{W}}$. For $y < z \leq n_1$, and for any $k < l$ with $m_1 + 1 \leq l$,*

$$(11) \quad (\Delta_{kl}(E_{yz}) \cdot F)(1) = 0.$$

Proof. This follows from that $E_{yz}^t = E_{zy}$ acts by zero on the l -th copy of V . □

We now consider upper triangular matrices of the form:

$$\begin{pmatrix} 0_{n_1} & 0 \\ 0 & * \end{pmatrix}.$$

Lemma 4.7. *Let $F \in \widetilde{\mathcal{W}}$. For $n_1 + 1 \leq y < z$, and either*

- (1) $0 \leq k < l \leq m_1$, or

$$(2) \quad 1 \leq k \leq m_1 < l,$$

$$(12) \quad (\Delta_{kl}(E_{yz}) \cdot F)(1) = 0.$$

Proof. This follows from that E_{yz} acts by zero on the k -th copy of $\mathbb{C}^{n_1} \subset V$ for (2) and on the l -th copy of $\mathbb{C}^{n_1} \subset V$ for (1). \square

4.7. Computing some actions on lower triangular part. We similarly have the following two cases. The first case is to consider elements of the form:

$$\begin{pmatrix} * & \\ & 0_{n_2} \end{pmatrix}.$$

Lemma 4.8. *Let $F \in \widetilde{\mathcal{W}}$. For any $1 \leq z < y \leq n_1$ and any $k < l$ with $m_1 + 1 \leq l$, $(\Delta_{kl}(E_{yz}) \cdot F)(1) = 0$.*

Proof. This follows from that E_{yz} acts by zero on the l -th copy of $\mathbb{C}^{n_2} \subset V$. \square

The another case is in the form:

$$\begin{pmatrix} 0_{n_1} & \\ & * \end{pmatrix}.$$

Lemma 4.9. *Let $F \in \widetilde{\mathcal{W}}$. Suppose we are in one of the following cases:*

$$(1) \quad 0 \leq k < l \leq m_1; \text{ or}$$

$$(2) \quad 1 \leq k \leq m_1 < l \leq m_1 + m_2$$

Then, for $n_1 < y < z$, $(\Delta_{kl}(E_{yz}) \cdot F)(1) = 0$.

Proof. (2) follows from that E_{yz} acts by zero on the k -th copy of $\mathbb{C}^{n_1} \subset V$ and (1) follows from E_{zy} acts by zero on the l -th copy of $\mathbb{C}^{n_1} \subset V$. \square

We consider matrices of the form $\begin{pmatrix} 0_{n_1} & \\ * & 0_{n_2} \end{pmatrix}$.

Lemma 4.10. *Let $F \in \widetilde{\mathcal{W}}$. Let $m_1 < l$ and $z \leq n_1 < y$. Then,*

$$\sum_{k=0}^{l-1} (\Delta_{kl}(E_{yz}) \cdot F)(1) = 0$$

Proof. We first consider $(\Delta_{0l}(E_{yz}) \cdot F)(1)$. We shall compute from using another expression:

$$(0, E_{yz}) = (-E_{zy}, E_{yz}) + (E_{zy}, 0).$$

We define $\Delta_k^L(E)$ and $\Delta_{kl}^L(E)$ analogously as $\Delta_k(E)$ and $\Delta_{kl}(E)$ respectively, but using $(E, 0)$ instead of $(0, E)$. We still have an analogous formula for Lemma 4.2 for the left version, and so

$$(\Delta_0^L(E_{zy}) \cdot F)(1) = ((E_{zy}, 0) \cdot F)(1) - \sum_{1 \leq k \leq m_1 + m_2} (\Delta_k^L(E_{zy}) \cdot F)(1).$$

Since $(E_{zy}, 0)$ acts by 0 on V , the latter terms are all zero. Since E_{zy} is an upper triangular matrices, we have that

$$((E_{zy}, 0) \cdot F)(1) = (E_{zy}, 0) \cdot F(1),$$

The right hand side is the Lie algebra action on $F(1) \in \tau \otimes V^{\otimes(m_1+m_2)}$, and so we also have such term to be zero. In conclusion, we have: $(\Delta_0^L(E_{zy}) \cdot F)(1) = 0$ and hence $(\Delta_{0l}^L(E_{zy}) \cdot F)(1) = 0$.

Now we compute

$$((\Delta_0(E_{yz}) \cdot F)(1) = ((\Delta_0^L(-E_{zy}) + \Delta_0(E_{yz})) \cdot F)(1)$$

as an action for the whole term $(-E_{zy}, E_{yz})$, where the term $\Delta_0^L(-E_{zy})$ acts by zero since E_{zy} is an upper triangular matrix and so $f(\exp(s(E_{zy}), 0))$ is a constant for any $f \in \text{Ind}_P^G(\tau)$. By using a version of Lemma 4.2 again, we have:

$$(\Delta_0(E_{yz}) \cdot F)(1) = ((-E_{zy}, E_{yz}) \cdot F)(1) - \sum_{1 \leq k \leq m_1+m_2} ((-\Delta_k^L(E_{zy}) + \Delta_k(E_{yz})) \cdot F)(1)$$

The first term in the RHS is zero since $F \in \widetilde{\mathcal{W}}$ is K -invariant and $(-E_{zy}, E_{yz})$ is in \mathfrak{k} . Furthermore, $\Delta_k^L(E_{zy}) \cdot F = 0$ since its action on V is trivial. Thus, the above expression reduces to give:

$$(\Delta_0(E_{yz}) \cdot F)(1) = - \sum_{1 \leq k \leq m_1+m_2} (\Delta_k(E_{yz}) \cdot F)(1).$$

But we see the terms are zero for $k \geq m_1 + 1$ by using Lemma 4.1. Thus, we further have:

$$(13) \quad (\Delta_0(E_{yz}) \cdot F)(1) = - \sum_{1 \leq k \leq m_1} (\Delta_k(E_{yz}) \cdot F)(1)$$

Imposing the action of $(0, E_{zy})$ on the l -th copy, we have:

$$(\Delta_{0l}(E_{yz}) \cdot F)(1) = - \sum_{1 \leq k \leq m_1} (\Delta_{kl}(E_{yz}) \cdot F)(1).$$

This gives that:

$$\sum_{0 \leq k \leq m_1} (\Delta_{kl}(E_{yz}) \cdot F)(1) = 0.$$

Again $(\Delta_{kl}(E_{yz}) \cdot F)(1) = 0$ for $l > k \geq m_1 + 1$, we have:

$$\sum_{0 \leq k \leq l-1} ((\Delta_{kl}(E_{yz}) \cdot F)(1) = 0$$

□

Let $e_1, \dots, e_{n_1+n_2}$ be a standard basis for V . Let \langle, \rangle_V be the inner product on V given by: for $a, b \in \mathbb{C}$,

$$\langle ae_i, be_j \rangle_V = a\bar{b} \cdot \delta_{ij}.$$

We define $\delta_l^i : \tau \otimes V^{\otimes m} \rightarrow \tau \otimes V^{\otimes m}$ determined by:

$$\delta_l^i(\mathbf{x} \otimes v_1 \otimes \dots \otimes v_m) = \langle v_l, e_i \rangle_V \cdot \mathbf{x} \otimes v_1 \otimes \dots \otimes e_i \otimes \dots \otimes v_m,$$

where e_i is in the l -th position.

Lemma 4.11. *Let $F \in \widetilde{\mathcal{W}}$. Let $l \leq m_1$ and $z \leq n_1 < y$. Then,*

$$(\Delta_{0l}(E_{yz}) \cdot F)(1) = -\delta_l^z(F(1)).$$

Proof. The computation for (13) in the previous lemma again yields:

$$(\Delta_0(E_{yz}) \cdot F)(1) = - \sum_{1 \leq k \leq m_1} (\Delta_k(E_{yz}) \cdot F)(1)$$

Since E_{zy} acts zero on $\mathbb{C}^{n_1} \subset V$, we have that $(\Delta_l(E_{zy}) \cdot \Delta_k(E_{yz}) \cdot F)(1) = 0$ for $k \neq l$ and hence:

$$(\Delta_{0l}(E_{yz}) \cdot F)(1) = -(\Delta_l(E_{zy}) \cdot \Delta_l(E_{yz}) \cdot F)(1).$$

By using Lemma 4.1 twice and using $E_{zy}E_{yz}v = E_{zz}v$, we then have

$$(\Delta_{0l}(E_{yz}) \cdot F)(1) = -\delta_l^z(F(1)).$$

□

5. PARABOLIC INDUCTION UNDER ARAKAWA-SUZUKI TYPE FUNCTOR

The main goal of this section is to prove Theorem 5.7. We will first illustrate how the dimensions of parabolically induced modules behave under $\Gamma_{n,m}$ in Section 5.2, which only involves some simpler dimension formulas. Working the full version of Theorem 5.7 requires a more substantial analysis on the module structure.

5.1. Height of parameters and representations.

Definition 5.1. Let $(\lambda_L, \lambda_R) \in \mathfrak{h}_0^* \times \mathfrak{h}_0^*$ with $\lambda_L - \lambda_R$ to be integral. Write $\mu = \lambda_L - \lambda_R = (\mu_1, \dots, \mu_n) \in \mathfrak{h}_0^*$. The height of (λ_L, λ_R) is defined as

$$\text{ht}((\lambda_L, \lambda_R)) := \begin{cases} \sum_{i=1}^n \mu_i & \text{if all } \mu_i \geq 0; \\ -\infty & \text{otherwise.} \end{cases}$$

We also define $\text{ht}(X(\lambda_L, \lambda_R)) := \text{ht}(\lambda_L, \lambda_R)$ and $\text{ht}(J(\lambda_L, \lambda_R)) := \text{ht}(\lambda_L, \lambda_R)$.

The height will be more important later in picking right choices of Arakawa-Suzuki type functors in Theorems 6.4 and 9.4.

5.2. Dimension under $\Gamma_{n,m}$. Before working on functoriality of parabolic induction under $\Gamma_{n,m}$, it may be educative to illustrate some simpler nature of parabolic inductions under $\Gamma_{n,m}$.

We consider a principal series $X = X(\lambda_L, \lambda_R)$. By the identifications in Section 4.1, there is a natural isomorphism:

$$\Gamma_{n,m}(X(\lambda_L, \lambda_R)) \cong (\mathbb{C}_{(\lambda_L, \lambda_R)} \otimes V^{\otimes m})^T.$$

Note that $\mathbb{C}_{(\lambda_L, \lambda_R)}|_T = \mathbb{C}_{\lambda_L - \lambda_R} = \mathbb{C}_\mu$, and $V|_K = J(\rho, e_1 + \rho)|_K \cong F_{e_1}^* = F_{-e_n}$ as K -modules, where F_ξ as the irreducible K -module of highest weight ξ , and F_ξ^* as its contragredient representation (see Equation (24) in the Appendix for details on finite dimensional representations of $GL_n(\mathbb{C})$). In particular, $V|_T = \bigoplus_{i=1}^n \mathbb{C}_{-e_i}$ are the T -weights of V .

It then follows from a direct computation that if $\text{ht}(X(\lambda_L, \lambda_R)) \neq -\infty$, then

$$(14) \quad \dim(\Gamma_{n,m}(X(\lambda_L, \lambda_R))) = \begin{cases} \frac{m!}{\mu_1! \mu_2! \dots \mu_n!} & m = \text{ht}(X(\lambda_L, \lambda_R)) \\ 0 & \text{otherwise} \end{cases}$$

If $\text{ht}(X(\lambda_L, \lambda_R)) = -\infty$, then for all m ,

$$\Gamma_{n,m}(X(\lambda_L, \lambda_R)) = 0.$$

Proposition 5.2. *Let X_1 be irreducible in \mathcal{HC}_{n_1} and X_2 be irreducible in \mathcal{HC}_{n_2} . Let $m_1 = \text{ht}(X_1)$ and let $m_2 = \text{ht}(X_2)$. Then*

$$(15) \quad \dim \Gamma_{n_1+n_2, m_1+m_2}(X_1 \times X_2) = \dim \Gamma_{n_1, m_1}(X_1) \times \dim \Gamma_{n_2, m_2}(X_2).$$

Proof. We only sketch the proof. It is a standard fact that X_1 and X_2 can be written as a linear combination of standard representations of $\text{GL}_{n_1}(\mathbb{C})$ and $\text{GL}_{n_2}(\mathbb{C})$ respectively. Thus, one can express the terms $\Gamma_{n_1+n_2, m_1+m_2}(X_1 \times X_2)$, $\Gamma_{n_1, m_1}(X_1)$ and $\Gamma_{n_2, m_2}(X_2)$ from such linear combinations and (14). In order to compute the RHS of (15), we need an additional formula recorded in the following lemma. \square

Lemma 5.3. *Let π_1 and π_2 in \mathcal{H}_{m_1} and \mathcal{H}_{m_2} respectively. If both π_1 and π_2 have finite dimensions, we have:*

$$(16) \quad \dim(\pi_1 \times \pi_2) = \frac{\dim(\pi_1) \cdot \dim(\pi_2) \cdot (m_1 + m_2)!}{m_1! \cdot m_2!}.$$

Proof. This follows from the well-known fact that $t_w \otimes v$ form a basis for $\mathbb{H}_{m_1+m_2} \otimes_{\mathbb{H}_{m_1} \otimes \mathbb{H}_{m_2}} (\pi_1 \boxtimes \pi_2)$, where w runs for all minimal representatives in $S_{m_1} \times S_{m_2} \setminus S_{m_1+m_2}$ and v runs for a fixed basis for $\pi_1 \boxtimes \pi_2$. \square

5.3. Description on subspaces of the functor. For fixed m_1, m_2 , let S^{m_1, m_2} be the set of minimal representatives of $(S_{m_1} \times S_{m_2}) \setminus S_{m_1+m_2}$. Let $e_1, \dots, e_{n_1}, e_{n_1+1}, \dots, e_{n_1+n_2}$ be the standard basis for $\mathbb{C}^{n_1+n_2}$. Let $V_1 = \text{span}\{e_1, \dots, e_{n_1}\}$ and let $V_2 = \text{span}\{e_{n_1+1}, \dots, e_{n_1+n_2}\}$. For $w \in S^{m_1, m_2}$, let \mathcal{X}_w be the subspace of $\tau \otimes V^{\otimes(m_1+m_2)}$ spanned by the vectors

$$\mathbf{x} \otimes v_1 \otimes v_2 \otimes \dots \otimes v_{m_1+m_2}$$

with $v_i \in V_1$ if $w(i) \in \{1, \dots, m_1\}$ and $v_i \in V_2$ if $w(i) \in \{m_1 + 1, \dots, m_1 + m_2\}$, and $\mathbf{x} \in \tau$.

We have the following simple linear algebra lemma, whose proof is elementary and so omitted.

Lemma 5.4. *For each representative $w \in S^{m_1, m_2}$, let p_w be a non-zero element in \mathcal{X}_w . Then those p_w 's form a set of linearly independent vectors.*

Lemma 5.5. *We use the notations in the previous lemma. For each representative $w \in S^{m_1, m_2}$, define a linear isomorphism $\iota_w : \mathcal{X}_1 \rightarrow \mathcal{X}_w$ given by*

$$\mathbf{x} \otimes v_1 \otimes \dots \otimes v_{m_1+m_2} \mapsto \mathbf{x} \otimes v_{w^{-1}(1)} \otimes \dots \otimes v_{w^{-1}(m_1+m_2)}.$$

The map induces a linear isomorphism from $\mathcal{X}_1^{K_1 \times K_2}$ to $\mathcal{X}_w^{K_1 \times K_2}$.

Proof. This follows from the fact that $K_1 \times K_2$ -action commutes with permutations in $S_{m_1+m_2}$. \square

Lemma 5.6. *Retain the notations in the previous two lemmas. For each $w \in S^{m_1, m_2}$, let $\mathcal{W}_w^{m_1, m_2}$ be the subspace of $\text{Ind}_P^G(\tau \otimes V^{\otimes(m_1+m_2)})^K$ satisfying $f(1) \in \mathcal{X}_w$. Then*

$$\bigoplus_{w \in S^{m_1, m_2}} \mathcal{W}_w^{m_1, m_2} \subset \text{Ind}_P^G(\tau \otimes V^{\otimes(m_1+m_2)})^K$$

and for each w , $\dim \mathcal{W}_w^{m_1, m_2} = \dim \mathcal{W}_1^{m_1, m_2}$.

Proof. The direct sum follows from Lemma 5.4 and the dimension follows from Lemma 5.5. \square

5.4. Parabolic induction under $\Gamma_{n, m}$: functoriality. We now study the parabolic induction under $\Gamma_{n, m}$.

Theorem 5.7. *Let Y_1 be in \mathcal{HC}_{n_1} and let Y_2 be in \mathcal{HC}_{n_2} . Then there exists a natural isomorphism:*

$$\Gamma_{n_1+n_2, m}(Y_1 \times Y_2) \cong \bigoplus_{m_1+m_2=m} \Gamma_{n_1, m_1}(Y_1) \times \Gamma_{n_2, m_2}(Y_2).$$

Proof. Let $\tau = Y_1 \boxtimes Y_2$. For fixed m_1, m_2 with $m_1 + m_2 = m$, let $\mathcal{W}_w^{m_1, m_2}$ as in Lemma 5.6. Set

$$\mathcal{P}_{m_1, m_2} = \bigoplus_{w \in S^{m_1, m_2}} \mathcal{W}_w^{m_1, m_2}.$$

Similar to Lemma 5.4, one sees that

$$\Gamma_{n_1+n_2, m}(Y_1 \times Y_2) \cong (\text{Ind}_P^G(\tau \otimes V^{\otimes m}))^K = \bigoplus_{m_1+m_2=m} \mathcal{P}_{m_1, m_2}.$$

Claim 1: Each \mathcal{P}_{m_1, m_2} is invariant under \mathbb{H}_m -action, and furthermore,

$$\mathcal{P}_{m_1, m_2} \cong \Gamma_{n_1, m_1}(Y_1) \times \Gamma_{n_2, m_2}(Y_2).$$

Note that the theorem follows from Claim 1. We consider $\mathbb{H}_{m_1} \otimes \mathbb{H}_{m_2}$ as a subalgebra of $\mathbb{H}_{m_1+m_2}$ in Section 2.3. Our proof of Claim 1 relies on another claim:

Claim 2: $\mathcal{W}_1^{m_1, m_2}$ is invariant under the action of the subalgebra $\mathbb{H}_{m_1} \otimes \mathbb{H}_{m_2}$, and is isomorphic to $\Gamma_{n_1, m_1}(Y_1) \boxtimes \Gamma_{n_2, m_2}(Y_2)$.

Assume Claim 2 holds in the meanwhile. It is well-known that any element in \mathbb{H}_m can be written as $\sum_{w \in S^{m_1, m_2}} w^{-1} h_w$ for $h_w \in \mathbb{H}_{m_1} \otimes \mathbb{H}_{m_2}$. The element w^{-1} sends $\mathcal{W}_1^{m_1, m_2}$ to $\mathcal{W}_w^{m_1, m_2}$ (by a direct computation using Lemma 4.1). This shows that \mathcal{P}^{m_1, m_2} is invariant under the action of \mathbb{H}_m .

Moreover, by Frobenius reciprocity, we obtain a non-zero map ψ from $\mathbb{H}_m \otimes \mathbb{H}_{m_1} \otimes \mathbb{H}_{m_2} (\Gamma_{n_1, m_1}(Y_1) \boxtimes \Gamma_{n_2, m_2}(Y_2))$ to $\Gamma_{n_1+n_2, m_1+m_2}(Y_1 \times Y_2)$. By Lemma 5.4, we have that ψ is injective. But, then by comparing the dimensions in Lemmas 5.6 and 5.3, we have that ψ is also surjective. This shows that we have an isomorphism from \mathcal{P}^{m_1, m_2} to $\Gamma_{n_1, m_1}(Y_1) \times \Gamma_{n_2, m_2}(Y_2)$, proving Claim 1.

It remains to prove Claim 2.

Proof of Claim 2: We first consider the action of y_l for $1 \leq l \leq m_1$ on $\mathcal{W}_1^{m_1, m_2}$. Let $F \in \mathcal{W}_1^{m_1, m_2}$. There is a natural isomorphism:

$$\Phi : \Gamma_{n_1, m_1}(Y_1) \boxtimes \Gamma_{n_2, m_2}(Y_2) \rightarrow \mathcal{W}_1^{m_1, m_2}$$

given by

$$\mathbf{z}_1 \otimes \mathbf{z}_2 \mapsto (1 \mapsto \mathbf{z}_1 \otimes \mathbf{z}_2)$$

We shall show that Φ is a $\mathbb{H}_{m_1} \otimes \mathbb{H}_{m_2}$ -map.

From Definition 3.3, the action of y_l is given by

$$\begin{aligned} \Theta_{\mathbb{H}_m}(y_l) = & \sum_{k=0}^{l-1} \left(\sum_{1 \leq y, z \leq n_1} \Delta_{kl}(E_{yz}) + \sum_{1 \leq y \leq n_1 < z \leq n_1 + n_2} \Delta_{kl}(E_{yz}) \right) \\ & + \sum_{k=0}^{l-1} \left(\sum_{1 \leq z \leq n_1 < y \leq n_1 + n_2} \Delta_{kl}(E_{yz}) + \sum_{n_1 + 1 \leq y, z \leq n_1 + n_2} \Delta_{kl}(E_{yz}) \right) + \frac{n_1 + n_2}{2} \end{aligned}$$

Again $\Theta_{\mathbb{H}_m}(y_l)F$ is determined by $(\Theta_{\mathbb{H}_m}(y_l)F)(1)$ and so it suffices to compute that. We shall compute the four parts separately. Now, by Lemma 4.5, for all $0 \leq k < l$,

$$\left(\sum_{1 \leq y \leq n_1 < z \leq n_1 + n_2} \Delta_{kl}(E_{yz}) \cdot F \right) (1) = 0.$$

By Lemmas 4.4, 4.7 and 4.9, for all $0 \leq k < l$

$$\left(\sum_{n_1 < y, z \leq n_1 + n_2} \Delta_{kl}(E_{yz}) \cdot F \right) (1) = 0$$

By Lemma 4.11 and the fact that $v_k = \sum_{i=1}^{n_1} \langle v_k, e_i \rangle e_i$ (since $v_k \in V_1$),

$$\sum_{k=0}^{l-1} \left(\sum_{1 \leq z \leq n_1 < y \leq n_1 + n_2} \Delta_{kl}(E_{yz}) \cdot F \right) (1) = -n_2 F(1).$$

As a result, $\Theta_{\mathbb{H}_m}(y_l)$ is reduced to considering

$$(17) \quad \left(\sum_{k=0}^{l-1} \sum_{1 \leq y, z \leq n_1} \Delta_{kl}(E_{yz}) + \frac{n_1 - n_2}{2} \right) \cdot F (1).$$

One may then apply similarly the formulas in Lemmas 4.1. Putting the normalized factor $\delta^{1/2}$ in the parabolic induction into consideration, one cancels the effect of $-\frac{n_2}{2}$ in the expression of (17), so that it agrees with

$$\Theta_{\mathbb{H}_{m_1}}(y_l) \cdot \Phi^{-1}(F) = \Theta_{\mathbb{H}_{m_1}}(y_l) \cdot (F(1)).$$

Checking the action for y_l for $m_1 + 1 \leq l \leq m_1 + m_2$ is similar and we shall be slightly brief. Again, we divide the computations into four parts as follows:

- For the matrices coming from $\begin{pmatrix} * \\ * \end{pmatrix}$, the action is zero by Lemmas 4.3, 4.6 and 4.8.
- For the matrices coming from $\begin{pmatrix} * \\ * \end{pmatrix}$, the action is zero by Lemma 4.5.
- For the matrices coming from $\begin{pmatrix} * \\ * \end{pmatrix}$, one uses Lemma 4.10 to get to zero by considering a sum.
- For the matrices E coming from $\begin{pmatrix} * \\ * \end{pmatrix}$, one gets the terms $\Delta_{kl}(E)$ acting by zero for $1 \leq k \leq m_1$ by using Lemmas 4.4, and 4.7 and 4.9.

Then $\Theta_{\mathbb{H}_m}(y_l) \cdot F$ reduces to:

$$\left(\sum_{\substack{k=0, \\ m_1 < k < l}} \sum_{n_1+1 \leq y, z \leq n_1+n_2} \Delta_{kl}(E_{yz}) + \frac{n}{2} \right) \cdot F \quad (1).$$

With the effect of the normalizing factor $-\frac{n_1}{2}$ from parabolic induction, we have that the term agrees with

$$\Theta_{\mathbb{H}_{m_2}}(y_l) \cdot \Phi^{-1}(F) = \Theta_{\mathbb{H}_{m_2}}(y_l) \cdot (F(1))$$

This also shows the isomorphism in Claim 2. \square

5.5. General form. We now have the general form for preserving parabolic inductions.

Corollary 5.8. *Let X_i be Harish-Chandra modules of $\mathrm{GL}_{n_i}(\mathbb{C})$ for $i = 1, \dots, r$. Then*

$$\Gamma_{n,m}(X_1 \times \dots \times X_r) = \bigoplus_{m_1, \dots, m_r} \Gamma_{n,m_1}(X_1) \times \dots \times \Gamma_{n,m_r}(X_r),$$

where m_1, \dots, m_r run for all non-negative integers such that $m_1 + \dots + m_r = m$.

6. STANDARD MODULES

In this section, we study the image of $\Gamma_{n,m}$ for principal series $X(\lambda_L, \lambda_R)$ of $\mathrm{GL}_n(\mathbb{C})$. In fact, Theorem 5.7 reduces the study to a character of $\mathrm{GL}_1(\mathbb{C})$, whose image under $\Gamma_{1,m}$ will be computed in Section 6.2.

6.1. Reduction to a particular $\Gamma_{n,m}$.

Lemma 6.1. *Let $m' = \mathrm{ht}(J(\lambda_L, \lambda_R))$. If $m \neq m'$, then $\Gamma_{n,m}(X(\lambda_L, \lambda_R)) = 0$ and $\Gamma_{n,m}(J(\lambda_L, \lambda_R)) = 0$.*

Proof. This follows from the computations in Section 5.2. \square

With this lemma, we only have to study $\Gamma_{n,m}(X)$ when m is the height of a principal series or an irreducible Harish-Chandra module X in the rest of sections.

6.2. Image of $\Gamma_{1,m}$ on a character.

Lemma 6.2. *Let $\chi_{a,b}$ be a $\mathrm{GL}_1(\mathbb{C})$ -character defined in (4). Suppose $m := a - b \geq 0$. Then*

$$\Gamma_{1,m}(\chi_{a,b}) = \mathrm{St}([b + \frac{1}{2}, a - \frac{1}{2}])$$

Proof. It is clear that $\Gamma_{1,m}(\chi_{a,b})$ is one-dimensional. Now the action follows from a straightforward computation. For example, the element 1 in $\mathfrak{h}^* \cong \mathbb{C}$ acts on $\chi_{a,b}$ according to the formula (1),

$$\frac{1}{2}[(b+a) + (\sqrt{-1})^2(a-b)] = b$$

and hence y_1 in \mathbb{H}_m acts by $b + \frac{1}{2}$. \square

6.3. Image of $\Gamma_{n,m}$ on a standard module. For $(\lambda_L, \lambda_R) \in \mathfrak{h}_0^* \times \mathfrak{h}_0^*$ with $\mathrm{ht}(\lambda_L, \lambda_R) \neq -\infty$, we define a multisegment: write $\lambda_L = (\lambda_{L,1}, \dots, \lambda_{L,n})$ and $\lambda_R = (\lambda_{R,1}, \dots, \lambda_{R,n})$,

$$\mathfrak{m}(\lambda_L, \lambda_R) = \left\{ [\lambda_{R,1} + \frac{1}{2}, \lambda_{L,1} - \frac{1}{2}], \dots, [\lambda_{R,n} + \frac{1}{2}, \lambda_{L,n} - \frac{1}{2}] \right\},$$

where we drop all the empty sets (this happens when $\lambda_{L,i} = \lambda_{R,i}$).

Lemma 6.3. *Consider a principal series $X(\lambda_L, \lambda_R)$ of $\mathrm{GL}_n(\mathbb{C})$. Write $\lambda_L = (\lambda_{L,1}, \dots, \lambda_{L,n})$. If*

$$\mathrm{Re}(\lambda_{L,1}) \geq \dots \geq \mathrm{Re}(\lambda_{L,n}),$$

then $X(\lambda_L, \lambda_R)$ is a standard module.

Proof. Recall that $\chi_{r_1, s_1} \times \chi_{r_2, s_2}$ is reducible if and only if there exists integers $p, q \in \mathbb{Z}$ with $p, q > 0$ such that $(\chi_{r_1, s_1} \chi_{r_2, s_2}^{-1})(z) = z^p \bar{z}^q$ (see [Go70]). From this, if we have $\mathrm{Re}(r_1 + s_1) \geq \mathrm{Re}(r_2 + s_2)$ and $\mathrm{Re}(r_1) \leq \mathrm{Re}(s_1)$, we have that $\chi_{r_1, s_1} \times \chi_{r_2, s_2}$ is irreducible and so, by [Vo81, Proposition 4.1.12],

$$(18) \quad \chi_{r_1, s_1} \times \chi_{r_2, s_2} \cong \chi_{r_2, s_2} \times \chi_{r_1, s_1}.$$

The proof for the lemma is an elementary exercise by repeatedly using (18) and using the definition of standard module in Section 2.2, which we leave for the reader. \square

Theorem 6.4. *Let $\lambda_L = (\lambda_{L,1}, \dots, \lambda_{L,n})$ and $\lambda_R = (\lambda_{R,1}, \dots, \lambda_{R,n})$. Suppose $m := \mathrm{ht}(X(\lambda_L, \lambda_R)) \neq -\infty$. Let $\Delta_i = [\lambda_{R,i} + \frac{1}{2}, \lambda_{L,i} - \frac{1}{2}]$. Then*

$$\Gamma_{n,m}(X(\lambda_L, \lambda_R)) = \mathrm{St}(\Delta_1) \times \dots \times \mathrm{St}(\Delta_n).$$

In particular, $\Gamma_{n,m}$ maps standard modules $X(\lambda_L, \lambda_R)$ to standard modules $\lambda(\mathfrak{m}(\lambda_L, \lambda_R))$.

Proof. This follows from the definition of $X(\lambda_L, \lambda_R)$ in Equation (5), Theorem 5.7 and Lemma 6.2. The second statement follows from the definitions of standard modules and Lemma 6.3. \square

Remark 6.5. Note that $\Gamma_{n,m}$ maps tempered representations (see [Knp86, Page 286] for a definition of tempered representations) to tempered \mathbb{H}_m -modules or zero (see [Ev96, Definition 1.4] for the definition of tempered \mathbb{H}_m -modules). In the next section, we will show that $\Gamma_{n,m}$ maps unitary modules to unitary modules.

7. HERMITIAN FORMS UNDER $\Gamma_{n,m}$

In this section, we prove that $\Gamma_{n,m}$ preserves Hermitianity and unitarity.

7.1. Hermitian form for Harish-Chandra modules.

Definition 7.1. Let X be a Harish-Chandra module of $GL_n(\mathbb{C})$. The Hermitian dual, denoted by X^h , of X is the space of K -finite vectors in the algebraic dual of X with a non-degenerate pairing $\langle \cdot, \cdot \rangle : X \times X^h \rightarrow \mathbb{C}$ determining (\mathfrak{g}, K) -module structure on X^h :

$$\langle x, (E_1, E_2) \cdot f \rangle = -\langle (\overline{E_2}, \overline{E_1}) \cdot x, f \rangle$$

$$\langle x, k \cdot f \rangle = \langle k^{-1} \cdot x, f \rangle$$

for any $E_1, E_2 \in \mathfrak{g}_0$, $k \in K$ and $x \in X$ and $f \in X^h$. We say that X is **Hermitian** if $X \cong X^h$. Moreover, we say that X is **unitary** if $X \cong X^h$ and the induced pairing can be chosen to be positive-definite.

We first see that taking the Hermitian dual preserves the height of an irreducible representation.

Proposition 7.2. $\text{ht}(J(\lambda_L, \lambda_R)) = \text{ht}(J(\lambda_L, \lambda_R)^h)$.

Proof. One has $J(\lambda_L, \lambda_R)^h = J(-\overline{\lambda_R}, -\overline{\lambda_L})$ (c.f. [Ba89, Theorem 2.4]), so that $J(\lambda_L, \lambda_R)$ is Hermitian if and only if $(\lambda_L, \lambda_R) = w(-\overline{\lambda_R}, -\overline{\lambda_L})$ for some $w \in S_n$ by Theorem 2.2(2). Now the proposition follows from computing $\text{ht}(J(\lambda_L, \lambda_R))$ and $\text{ht}(J(-\overline{\lambda_R}, -\overline{\lambda_L}))$ directly. \square

7.2. Hermitian form for modules of graded Hecke algebras. We first need a suitable notion of Hermitian modules for \mathbb{H}_m (see e.g. [BCT12]), translated from p -adic groups:

Definition 7.3. A Hermitian anti-involution $*$: $\mathbb{H}_m \rightarrow \mathbb{H}_m$ is a linear map determined by:

$$s_i^* = s_i, \quad y_i^* = -w_0(y_{m+1-i})w_0^{-1}.$$

Definition 7.4. Let π be an \mathbb{H}_m -module. The Hermitian dual, denoted π^* , of π is the space of linear functionals from π to \mathbb{C} determined by:

$$(h \cdot f)(x) = f(h^* \cdot x).$$

In particular, there exists a non-degenerate pairing $\langle \cdot, \cdot \rangle : \pi \times \pi^* \rightarrow \mathbb{C}$ such that

$$\langle h \cdot x, f \rangle = \langle x, h^* \cdot f \rangle.$$

We say that π is ***-Hermitian** if $\pi \cong \pi^*$. We say that π is ***-unitary** if π is *-Hermitian and the induced pairing on π can be chosen to be positive-definite.

7.3. Preserving Hermitian structure. One may compare the proof with [CT11, Theorem 4.2.2].

Theorem 7.5. *Let X be in \mathcal{HC}_n . Then*

$$\Gamma_{n,m}(X^h) \cong \Gamma_{n,m}(X)^*.$$

Proof. We write \langle, \rangle for the natural pairing between X and X^h as in Definition 7.1. Recall that \langle, \rangle_V is the standard Hermitian pairing on V .

We define a Hermitian form \langle, \rangle' on $X \otimes V^{\otimes m}$:

$$\langle f \otimes e_{i_1} \otimes \dots \otimes e_{i_m}, f' \otimes e_{i'_1} \otimes \dots \otimes e_{i'_m} \rangle' = \langle f, f' \rangle \cdot \langle e_{i_1}, e_{i'_1} \rangle_V \cdot \dots \cdot \langle e_{i_m}, e_{i'_m} \rangle_V.$$

This descends to a pairing on $\Gamma_{n,m}(X) \times \Gamma_{n,m}(X^h)$ by taking the K -invariant.

We have to check that: for $p \in \Gamma_{n,m}(X)$ and $p' \in \Gamma_{n,m}(X^h)$,

$$(19) \quad \langle \Theta(s_{l,l+1}) \cdot p, p' \rangle' = \langle p, \Theta(s_{l,l+1}) \cdot p' \rangle'$$

$$(20) \quad \langle \Theta(y_l) \cdot p, p' \rangle' = \langle p, \Theta(y_l^*) \cdot p' \rangle'.$$

Checking (19) is straightforward.

We now consider (20). We first consider p takes the form:

$$f \otimes e_{i_1} \otimes \dots \otimes e_{i_m}$$

and p' takes the form:

$$f' \otimes e_{i'_1} \otimes \dots \otimes e_{i'_m}$$

(which are not necessarily in $\Gamma_{n,m}(X)$ or $\Gamma_{n,m}(X^h)$).

Then

$$\begin{aligned} \langle \Theta(y_l) \cdot p, p' \rangle' &= \sum_{1 \leq i, j \leq n} \langle (0, E_{ij}) \cdot f, f' \rangle \cdot \langle e_{i_1}, e_{i'_1} \rangle_V \cdot \dots \cdot \langle (0, E_{ji}) \cdot e_{i_l}, e_{i'_l} \rangle_V \cdot \dots \cdot \langle e_{i_m}, e_{i'_m} \rangle_V \\ &\quad + \sum_{1 \leq i, j \leq n} \sum_{1 \leq k < l} \langle f, f' \rangle \cdot \dots \cdot \langle (0, E_{ij}) \cdot e_{i_k}, e_{i'_k} \rangle_V \cdot \dots \cdot \langle (0, E_{ji}) \cdot e_{i_l}, e_{i'_l} \rangle_V \cdot \dots \cdot \langle e_{i_m}, e_{i'_m} \rangle_V \\ &= \sum_{1 \leq i, j \leq n} \langle (-E_{ji}, E_{ij}) \cdot f, f' \rangle \cdot \langle e_{i_1}, e_{i'_1} \rangle_V \cdot \dots \cdot \langle (0, E_{ji}) \cdot e_{i_l}, e_{i'_l} \rangle_V \cdot \dots \cdot \langle e_{i_m}, e_{i'_m} \rangle_V \\ &\quad + \sum_{1 \leq i, j \leq n} \sum_{1 \leq k < l} \langle f, f' \rangle \cdot \dots \cdot \langle (-E_{ji}, E_{ij}) \cdot e_{i_k}, e_{i'_k} \rangle_V \cdot \dots \cdot \langle (0, E_{ji}) \cdot e_{i_l}, e_{i'_l} \rangle_V \cdot \dots \cdot \langle e_{i_m}, e_{i'_m} \rangle_V \\ &\quad + \sum_{1 \leq i, j \leq n} \langle (E_{ji}, 0) \cdot f, f' \rangle \cdot \dots \cdot \langle (0, E_{ji}) \cdot e_{i_l}, e_{i'_l} \rangle_V \cdot \dots \cdot \langle e_{i_m}, e_{i'_m} \rangle_V + \frac{n}{2} \langle p, p' \rangle' \end{aligned}$$

where the second equality follows by using $(E_{ji}, 0)$ acts by zero on V . The first three terms in the last sum will be denoted by (I), (II) and (III) respectively.

We now proceed to compute $\langle p, \Theta(y_l^*) \cdot p' \rangle'$. Note that the action of w_0 simply permutes those $e_{i'_a}$ terms. Now,

$$\begin{aligned}
& \langle p, \Theta(y_l^*) \cdot p' \rangle' \\
&= - \langle p, \Theta(w_0 y_{m+1-l} w_0) \cdot p' \rangle' \\
&= - \sum_{1 \leq i, j \leq n} \langle f, (0, E_{ij}) \cdot f' \rangle \cdot \langle e_{i_1}, e_{i'_1} \rangle_V \cdots \langle e_{i_l}, (0, E_{ji}) \cdot e_{i'_l} \rangle_V \cdots \langle e_{i_m}, e_{i'_m} \rangle_V \\
&\quad - \sum_{1 \leq i, j \leq n} \sum_{l < k \leq m} \langle f, f' \rangle \cdots \langle e_{i_l}, (0, E_{ji}) \cdot e_{i'_l} \rangle_V \cdots \langle e_{i_k}, (0, E_{ij}) \cdot e_{i'_k} \rangle_V \cdots \langle e_{i_m}, e_{i'_m} \rangle_V \\
&\quad - \frac{n}{2} \langle p, p' \rangle \\
&= - \sum_{1 \leq i, j \leq n} \langle f, (0, E_{ij}) \cdot f' \rangle \cdot \langle e_{i_1}, e_{i'_1} \rangle_V \cdots \langle e_{i_l}, (0, E_{ji}) \cdot e_{i'_l} \rangle_V \cdots \langle e_{i_m}, e_{i'_m} \rangle_V \\
&\quad - \sum_{1 \leq i, j \leq n} \sum_{l < k \leq m} \langle f, f' \rangle \cdots \langle e_{i_l}, (0, E_{ji}) \cdot e_{i'_l} \rangle_V \cdots \langle e_{i_k}, (-E_{ji}, E_{ij}) \cdot e_{i'_k} \rangle_V \cdots \langle e_{i_m}, e_{i'_m} \rangle_V \\
&\quad - \frac{n}{2} \langle p, p' \rangle,
\end{aligned}$$

where the last equation again follows by $(E_{ij}, 0)$ acts by zero on V . We shall label the first two terms in the last sum by (IV) and (V) (without the subtractions).

We now consider one more formula:

$$(21) \quad \sum_{1 \leq i, j \leq n} \langle f, f' \rangle \cdot \langle e_{i_1}, e_{i'_1} \rangle_V \cdots \langle (0, E_{ji}) \cdot (-E_{ji}, E_{ij}) \cdot e_{i_l}, e_{i'_l} \rangle_V \cdots \langle e_{i_m}, e_{i'_m} \rangle_V = n \cdot \langle p, p' \rangle,$$

which follows from $(0, E_{ji}) \cdot (-E_{ji}, E_{ij}) \cdot e_{i_l} = e_{i_l}$ if $j = i_l$ and $= 0$ otherwise.

For computing $\langle \Theta(y_l) \cdot p, p' \rangle - \langle p, \Theta(y_l^*) \cdot p' \rangle'$, we express the two terms in the above ways. Note that (III) is cancelled with (IV) by using $\langle (0, E_{ji}) \cdot f, f' \rangle = -\langle f, (0, E_{ji}) \cdot f' \rangle$ and $\langle e_{i_l}, (0, E_{ji}) \cdot e_{i'_l} \rangle_V = \langle (0, E_{ij}) \cdot e_{i_l}, e_{i'_l} \rangle$. Now the terms (I), (II) and (V) with the formula (21) give that

$$(22) \quad \langle \Theta(y_l) \cdot p, p' \rangle' - \langle p, \Theta(y_l^*) \cdot p' \rangle' = \sum_{1 \leq i, j \leq n} \langle \Delta_l((0, E_{ij})) \cdot ((-E_{ji}, E_{ij}) \cdot p), p' \rangle',$$

We emphasize here that $(-E_{ji}, E_{ij}) \cdot p$ means the Lie algebra action of $(-E_{ji}, E_{ij})$ on $p \in X \otimes V^{\otimes m}$.

Now, we consider $p \in \Gamma_{n,m}(X)$ (and $p' \in \Gamma_{n,m}(X^h)$) and the formula (22) extends linearly. However, by using the K -invariant, we have that $(-E_{ji}, E_{ij}) \cdot p = 0$. In other words, $\langle \Theta(y_l) \cdot p, p' \rangle' = \langle p, \Theta(y_l^*) \cdot p' \rangle'$ as desired. \square

Corollary 7.6. $\Gamma_{n,m}$ maps Hermitian Harish-Chandra modules to $*$ -Hermitian \mathbb{H}_m -modules or zero, and maps unitary Harish-Chandra modules to $*$ -unitary \mathbb{H}_m -modules or zero.

Remark 7.7. The Lefschetz principle for the unitary dual is also explicated in [Ta09].

Remark 7.8. The unitary characters of $GL_{n'}(\mathbb{C})$ are the building blocks of unitary representations of $GL_n(\mathbb{C})$ in the sense that all unitary representations are products of those characters. We shall see from Theorem 9.4 that the image of a unitary character of $GL_{n'}(\mathbb{C})$ is a (unitary) Speh module

of \mathbb{H}_m (defined in Section 10.2). It is well-known from [Ta86] that Speh modules play a similar role of building blocks for the unitary dual of \mathbb{H}_m .

Remark 7.9. As seen in [ALTV20], even one is interested in studying the unitary dual problem, it is useful to study other Hermitian forms. The anti-involution on $\mathrm{GL}_n(\mathbb{C})$ is given by:

$$g \mapsto \overline{g}^t$$

i.e. the transpose with the complex conjugation. For Harish-Chandra modules, the anti-involution corresponds to the anti-involution $(E, E') \mapsto (\overline{E}^t, \overline{E'}^t)$. On the other hand, there is an anti-involution on \mathbb{H}_m which is the identity on the generators s_l and y_l in Definition 2.3. It is more straightforward to check that the Hermitian forms associated to the anti-involutions correspond to each other under the functor $\Gamma_{n,m}$ (c.f. [Su98]).

8. BERNSTEIN-ZELEVINSKY DERIVATIVES AND TENSOR PRODUCTS

We introduce the tool of Bernstein-Zelevinsky derivatives and connect to the tensor product problem in this section. Some connections to the original notion of Bernstein-Zelevinsky derivatives for p -adic groups will be discussed in Section 10.

8.1. Generalized BZ derivatives. For each $\tau \in \mathrm{Irr}(S_i)$, define

$$\mathbf{BZ}_\tau : \mathcal{H}_n \rightarrow \mathcal{H}_{n-i}$$

is defined as:

$$\mathbf{BZ}_\tau(\pi) = \mathrm{Hom}_{S_i}(\tau, \pi),$$

where π is regarded as a $\mathbb{C}[S_i]$ -module via the embedding $s_l \mapsto s_l$. The \mathbb{H}_{n-i} -module structure on $\tilde{\pi} = \mathbf{BZ}_\tau(\pi)$ is via the following actions:

$$(y_k \cdot_{\tilde{\pi}} f)(x) := y_{i+k} \cdot_{\pi} f(x), \quad (s_k \cdot_{\tilde{\pi}} f)x := s_{i+k} \cdot_{\pi} f(x)$$

for $x \in \tau$.

When τ is the sign representation of S_i , it is studied from the viewpoint of representations of p -adic groups [CS19, Ch22+]. One nice property is the following:

Proposition 8.1. *Let \mathbf{m} be a multisegment. Write $\mathbf{m} = \{\Delta_1, \dots, \Delta_k\}$. For each $\Delta_i = [a_i, b_i]$, write ${}^-\Delta = [a_i + 1, b_i]$ and ${}^-\mathbf{m} = \{{}^-\Delta_1, \dots, {}^-\Delta_k\}$. Let triv be the trivial representation of S_k . Then*

$$\mathbf{BZ}_{\mathrm{triv}}(\mathrm{St}(\mathbf{m})) \cong \mathrm{St}({}^-\mathbf{m}).$$

Proof. This follows from the highest derivative of an irreducible representation due to Zelevinsky [Ze80, Theorem 8.1], and the translation to \mathbb{H}_m in [CS19, Theorem 6.9]. Note that to translate the version from the sign representation to triv , one also applies the Iwahori-Matsumoto involution (see more discussions in the proof of Theorem 11.1). \square

8.2. Arakawa-Suzuki functor and Bernstein-Zelevinsky derivatives. Let $\text{Irr}(S_i)$ be the set of irreducible representations of S_i . Recall that, for $\tau \in \text{Irr}(S_i)$, $\mathbb{S}_\tau(V) = \text{Hom}_{S_i}(\tau, V^{\otimes i})$, where S_i acts by sign permutation on $V^{\otimes i}$ i.e. $w.(v_1 \otimes \dots \otimes v_i) = (-1)^{l(w)} v_{w(1)} \otimes \dots \otimes v_{w(i)}$. We regard $\mathbb{S}_\tau(V)$ as a natural $GL_n(\mathbb{C})$ -representation, and the classical Schur-Weyl duality asserts that $\mathbb{S}_\tau(V)$ is either irreducible or zero.

Theorem 8.2. *Let $\tau \in \text{Irr}(S_i)$. For $m \geq i$, the following diagram is commutative:*

$$\begin{array}{ccc} \mathcal{HC}_n & \xrightarrow{T_\tau} & \mathcal{HC}_n \\ \downarrow \Gamma_{n,m} & & \downarrow \Gamma_{n,m-i} \\ \mathcal{H}_m & \xrightarrow{\mathbf{BZ}_\tau} & \mathcal{H}_{m-i} \end{array}$$

where T_τ is the functor defined by $T_\tau(X) := X \otimes \mathbb{S}_\tau(V)$.

Proof. Since $\Theta(w)$ acts by a sign permutation on the $V^{\otimes i}$, we have the decomposition $X \otimes V^{\otimes m}$:

$$\bigoplus_{\omega \in \text{Irr}(S_i)} \omega \boxtimes (X \otimes \mathbb{S}_\omega(V) \otimes V^{\otimes(m-i)}).$$

Here we may regard as a natural $\mathbb{C}[S_i] \otimes U(\mathfrak{g}_0) \otimes \mathbb{C}[S_{m-i}]$ representation.

Taking the K -invariant on the decomposition and then applying \mathbf{BZ}_τ , we have:

$$\mathbf{BZ}_\tau \circ \Gamma_{n,m}(X) = (X \otimes \mathbb{S}_\tau(V) \otimes V^{\otimes(m-i)})^K.$$

In other words,

$$\mathbf{BZ}_\tau \circ \Gamma_{n,m}(X) \quad \text{and} \quad \Gamma_{n,m-i} \circ T_\tau(X)$$

share the same underlying space. It remains to see that the equality above holds for \mathbb{H}_{m-i} -module action. It is quite straightforward to check that it is true for elements of the form s_l . For y_l , recall that the action from $\mathbf{BZ}_\tau \circ \Gamma_{n,m}(X)$ is given by

$$\sum_{k=0}^{l-1} \Omega_{kl} + \frac{n}{2}$$

The terms $\sum_{k=0}^i \Omega_{kl}$ can be grouped together. Using the Lie algebra action on $X \otimes \mathbb{S}_\tau(V)$, the action of such term agrees respectively with the action $\Omega_{0,l-i}$ from $\Gamma_{n,m-i} \circ T_\tau(X)$. The actions of the terms $\Omega_{i+1,l}, \dots, \Omega_{l-1,l}$ then agree with the ones $\Omega_{1,l-i}, \dots, \Omega_{l-i-1,l-i}$ from $\Gamma_{n,m-i} \circ T_\tau(X)$. Thus, the action of y_l coincides on $\mathbf{BZ}_\tau \circ \Gamma_{n,m}(X)$ and $\Gamma_{n,m-i} \circ T_\tau(X)$. \square

9. IRREDUCIBILITY UNDER $\Gamma_{n,m}$

We combine the tools of Theorems 6.4, 7.5 and 8.2 to prove that $\Gamma_{n,m}$ preserves the irreducibility in Theorem 9.4.

9.1. Non-zero image for the thickened case.

Definition 9.1. We write $\lambda_L = (\lambda_{L,1}, \dots, \lambda_{L,n})$ and $\lambda_R = (\lambda_{R,1}, \dots, \lambda_{R,n})$. We say that the parameter (λ_L, λ_R) is **thickened** if $\lambda_{L,i} - \lambda_{R,j} \geq 0$ for any $1 \leq i, j \leq n$.

Lemma 9.2. Let $X = J(\lambda_L, \lambda_R)$, where $(\lambda_L, \lambda_R) \in \mathfrak{h}_0^* \times \mathfrak{h}_0^*$ is a thickened parameter. Then $\Gamma_{n,m}(X)$ is non-zero, and has a simple quotient isomorphic to $\text{St}(\mathfrak{m}(\lambda_L, \lambda_R))$.

Proof. It is clear that $(\lambda_L, w\lambda_R)$ is also thickened for any $w \in S_n$. Without loss of generality, we assume that

$$\text{Re}(\lambda_{L,1}) \geq \dots \geq \text{Re}(\lambda_{L,n})$$

so that $\Gamma_{n,m}(X(\lambda_L, w\lambda_R))$ is a standard module.

By using the thickenedness, $m = \text{ht}(X(\lambda_L, w\lambda_R))$ is not equal to $-\infty$ and is a constant for all $w \in S_n$. Then it follows from Theorem 6.4 that $\Gamma_{n,m}(X(\lambda_L, w\lambda_R))$ is non-zero.

We pick w_1, \dots, w_k to be a choice of elements in S_n such that $J(\lambda_L, w_1\lambda_R), \dots, J(\lambda_L, w_k\lambda_R)$ are all irreducible modules with the same infinitesimal character as $J(\lambda_L, \lambda_R)$ without redundancy. By Theorems 2.2(b), 2.4 and 6.4, $\{\Gamma_{n,m}(X(\lambda_L, w_i\lambda_R))\}_i$ is linearly independent in the Grothendieck group of \mathbb{H}_m -modules. Moreover, the multisegments $\{\mathfrak{m}(\lambda_L, w_i\lambda_R)\}_i$ are closed under intersection-union operations on multisegments. In other words, by [Ze80, Theorem 7.1], we have that:

$$\text{span}_{\mathbb{C}} \{\text{St}(\mathfrak{m}(\lambda_L, w_i\lambda_R))\}_i = \text{span}_{\mathbb{C}} \{\lambda(\mathfrak{m}(\lambda_L, w_i\lambda_R))\}_i,$$

where we regard the representations in the Grothendieck group of \mathcal{H}_m over \mathbb{C} .

Combining above, we must then have: $\Gamma_{n,m}(J(\lambda_L, w_i\lambda_R))$ is non-zero for all i . By the exactness of $\Gamma_{n,m}$ and Lemma 6.3, we then have a non-zero surjection from $\Gamma_{n,m}(X(\lambda_L, w_i\lambda_R))$ to $\Gamma_{n,m}(J(\lambda_L, w_i\lambda_R))$. This gives the proposition by Theorem 6.4. \square

9.2. Non-zero image for the general case. We shall use the tool of Bernstein-Zelevinsky derivatives to deduce the general case.

Let $\chi : \text{GL}_n(\mathbb{C}) \rightarrow \mathbb{C}^\times$ be the character $\chi(g) = \overline{\det(g)}^{-1}$. Let triv_k be the trivial representation of S_k .

Proposition 9.3. Suppose $m := \text{ht}(J(\lambda_L, \lambda_R)) \neq -\infty$. Then $\Gamma_{n,m}(J(\lambda_L, \lambda_R))$ is non-zero. Moreover, $\Gamma_{n,m}(J(\lambda_L, \lambda_R))$ has a unique simple quotient isomorphic to $\text{St}(\mathfrak{m}(\lambda_L, \lambda_R))$.

Proof. Let $\lambda'_R = \lambda_R - (k, \dots, k)$ for sufficiently large k such that $(\lambda_L, w\lambda'_R)$ is thickened for any $w \in S_n$. By Lemma 9.2, we have that $\Gamma_{n,m+kn}(J(\lambda_L, \lambda'_R)) \neq 0$ and has the simple composition factor $\text{St}(\mathfrak{m}(\lambda_L, \lambda_R))$. Then, one applies the derivative $\mathbf{BZ}_{\text{triv}_n}$ k -times and the resulting module

$$\mathbf{BZ}_{\text{triv}_n} \circ \dots \circ \mathbf{BZ}_{\text{triv}_n} \circ \Gamma_{n,m+kn}(J(\lambda_L, \lambda'_R))$$

is still non-zero. Note that $\mathbb{S}_{\text{triv}_n}(V) = \wedge^n V = \chi^{-1}$ and $J(\lambda_L, \lambda'_R) \otimes \chi^{\otimes(-k)} = J(\lambda_L, \lambda_R)$. Now the statement then follows by using $\mathbf{BZ}_{\text{triv}_n} \circ \Gamma_{n,m'+n}(X) = \Gamma_{n,m'}(\chi^{-1} \otimes X)$ several times from Theorem 8.2 and Proposition 8.1. \square

9.3. Irreducibility under $\Gamma_{n,m}$.

Theorem 9.4. *Let $X = J(\lambda_L, \lambda_R)$ be an irreducible Harish-Chandra module of $GL_n(\mathbb{C})$ such that $\text{ht}(X) \neq -\infty$. Then $\Gamma_{n,m}(X)$ is irreducible, and moreover, $\Gamma_{n,m}(X) \cong \text{St}(\mathfrak{m}(\lambda_L, \lambda_R))$.*

Proof. We assume that $\text{Re}(\lambda_L + \lambda_R)$ is dominant. By Theorem 7.5, we have:

$$(23) \quad \Gamma_{n,m}(J(-\overline{\lambda_R}, -\overline{\lambda_L})) \cong \Gamma_{n,m}(J(\lambda_L, \lambda_R)^h) \cong \Gamma_{n,m}(J(\lambda_L, \lambda_R))^*$$

By Proposition 9.3, $\Gamma_{n,m}(J(\lambda_L, \lambda_R))$ has a unique simple quotient $\text{St}(\mathfrak{m}(\lambda_L, \lambda_R))$. Then $\Gamma_{n,m}(J(\lambda_L, \lambda_R))^*$ has a unique simple submodule $\text{St}(\mathfrak{m}(\lambda_L, \lambda_R))^* \cong \text{St}(\mathfrak{m}(-\overline{\lambda_R}, -\overline{\lambda_L}))$. By (23), $\Gamma_{n,m}(J(-\overline{\lambda_R}, -\overline{\lambda_L}))$ also has such unique simple submodule.

On the other hand, by Theorem 6.4 and the exactness of $\Gamma_{n,m}$, $\Gamma_{n,m}(J(-\overline{\lambda_R}, -\overline{\lambda_L}))$ has a unique simple quotient isomorphic to $\text{St}(\mathfrak{m}(-\overline{\lambda_R}, -\overline{\lambda_L}))$, and no other composition factor isomorphic to that. Thus, combining with the previous paragraph, we then have that

$$\Gamma_{n,m}(J(-\overline{\lambda_R}, -\overline{\lambda_L})) \cong \text{St}(\mathfrak{m}(-\overline{\lambda_R}, -\overline{\lambda_L})).$$

In particular, it is irreducible. Since (λ_L, λ_R) is arbitrary, we are then done. \square

As a result, we also have the following consequence:

Corollary 9.5. *If $\Gamma_{n,m}(X(\lambda_L, \lambda_R)) \neq 0$, then $\Gamma_{n,m}(J(\lambda_L, \lambda_R)) \neq 0$.*

Remark 9.6. (1) The approach for proving irreducibility in [AS98, CT12] is to compare the geometry to obtain coincidence of character formulas.

(2) The analogous statement of Corollary 9.5 does not hold for the original Arakawa-Suzuki functor [AS98] in general.

(3) In general, for two irreducible Harish Chandra modules X_1 and X_2 of $GL_n(\mathbb{C})$, it is possible that $\Gamma_{n,m}(X_1) \cong \Gamma_{n,m}(X_2)$, but $X_1 \not\cong X_2$. For example, when $n = 1$, one may take $X_1 = \chi_{r,r}$ and $X_2 = \chi_{s,s}$ for $r \neq s$. However, this happens only for those 'degenerate' cases. For example, if both X_1 and X_2 are thickened with that all those inequalities for $\lambda_{L,i} - \lambda_{R,j}$ in Definition 9.1 are all strict, then we have $\Gamma_{n,m}(X_1) \cong \Gamma_{n,m}(X_2)$ if and only if $X_1 \cong X_2$.

9.4. Digression: Matching of $U(n)$ -types and S_m -types. Indeed, the ideas in Section 8.2 above could also be used to match the $U(n)$ -structure of X and the S_m -structure of $\Gamma_{n,m}(X)$. Recall that $K = U(n)$. For a K -representation Y , we denote by Y^* the contragredient K -representation of Y .

Theorem 9.7. *Let X be in \mathcal{HC}_n . We have that:*

$$\text{Hom}_{U(n)}(\mathbb{S}_\tau(V)^*, X|_{U(n)}) \cong \text{Hom}_{S_m}(\tau, \Gamma_{n,m}(X)|_{S_m}).$$

Proof. We have that

$$\text{Hom}_{S_m}(\tau, \Gamma_{n,m}(X)) \cong (X \otimes \mathbb{S}_\tau(V))^K \cong \text{Hom}_K(X, \mathbb{S}_\tau(V)^*),$$

where the first isomorphism is similar to discussions in the proof of Theorem 8.2 and the second one follows from the adjointness between Hom and tensor product. \square

Remark 9.8. The highest weight of V , as a K -representation, is $(0, \dots, 0, -1)$. The irreducible representation of S_m can be parameterized by partitions of m so that the trivial representation corresponds to (m) and the sign representation corresponds to $(1, \dots, 1)$.

For a thickened parameter (λ_L, λ_R) with integral $\lambda_L - \lambda_R$, write $\lambda_L = (\lambda_{L,1}, \dots, \lambda_{L,n})$ and $\lambda_R = (\lambda_{R,1}, \dots, \lambda_{R,n})$. This determines a partition $\alpha = (\lambda_{L,1} - \lambda_{R,1}, \dots, \lambda_{L,n} - \lambda_{R,n})$, whose transpose determines an irreducible representation, denoted τ_α , of S_m .

The classical Schur-Weyl duality gives that $\mathbb{S}_{\tau_\alpha}(V)^*$, as a K -representation, has the highest weight $\lambda_L - \lambda_R$. On the other hand, Theorems 9.4 and 9.7 give

$$\mathrm{Hom}_{U(n)}(\mathbb{S}_{\tau_\alpha}(V)^*, J(\lambda_L, \lambda_R)) \cong \mathrm{Hom}_{S_m}(\tau_\alpha, \mathrm{St}(\mathbf{m}(\lambda_L, \lambda_R))).$$

This may be seen as a theoretical explanation on Remark 2.5.

10. LEFSCHETZ PRINCIPLE FOR DIRAC COHOMOLOGY

In the 1990s, Vogan introduced the notion of *Dirac cohomology* for (\mathfrak{g}, K) -modules. The relationship between the infinitesimal character of a (\mathfrak{g}, K) -module and its Dirac cohomology is studied in [HP02] in full detail. Later on, an analogous study of Dirac cohomology for graded Hecke algebra is carried out in [BCT12]. In this section, we will use $\Gamma_{n,m}$ to relate various results of Dirac cohomologies on representations of $\mathrm{GL}_n(\mathbb{C})$ and \mathbb{H}_m .

10.1. Finite-dimensional representations versus ladder representations. An irreducible \mathbb{H}_m -module $\mathrm{St}(\mathbf{m})$ is said to be a **ladder representation** (in the sense in [LM16]) if the multisegment

$$\mathbf{m} = \{[a_1, b_1], \dots, [a_n, b_n]\}$$

satisfies that $a_1 > a_2 > \dots > a_n$ and $b_1 > b_2 > \dots > b_n$.

One can relate finite dimensional representations of $\mathrm{GL}_n(\mathbb{C})$ (see Appendix A for details) and ladder representations using $\Gamma_{n,m}$. Namely, by Theorem 9.4 and the first paragraph of Appendix A, all ladder representations can be realized as the image of some finite-dimensional representations of $\mathrm{GL}_n(\mathbb{C})$ under $\Gamma_{n,m}$.

By Theorem A.1, all finite-dimensional representations of complex groups with nonzero Dirac cohomology must be of the form $X = J(\lambda, -w_0\lambda)$ for regular integral λ . So we are interested in studying Dirac cohomology of the ladder representations $\Gamma_{n,m}(X) = \mathrm{St}(\mathbf{m}(\lambda, -w_0\lambda))$ with $\mathrm{ht}(X) = m > 0$. In order to invoke the classification of all ladder representations with nonzero Dirac cohomology given in [Ch18, Section 7], we define the *content* of a multisegment $\mathbf{m} = \{[a_1, b_1], \dots, [a_n, b_n]\}$ to be the multiset

$$\mathrm{con}(\mathbf{m}) := \{a_1, a_1 + 1, \dots, b_1, \dots, a_n, a_n + 1, \dots, b_n\}$$

(if $a_i = b_i$, we only count it once in the content). And we say \mathbf{m} is **twisted-elliptic** if there exists a multisegment of the form $\mathbf{m}^{temp} = \{[-c_1, c_1], \dots, [-c_m, c_m]\}$ such that $\mathrm{con}(\mathbf{m}) = \mathrm{con}(\mathbf{m}^{temp})$ (c.f. [Ch18, Corollary 7.3]).

For example, $\mathbf{m} = \{[3, 4], [0, 1], [-1, 0], [-4, -3]\}$ has content $\{-4, -3, -1, 0, 0, 1, 3, 4\}$, and is not twisted-elliptic. However, $\mathbf{m}'' = \{[0, 7], [-3, 4], [-4, 3], [-7, 0]\}$ is twisted-elliptic, with

$$\text{con}(\mathbf{m}'') = \text{con}(\{[-7, 7], [-4, 4], [-3, 3], [0, 0]\}).$$

Lemma 10.1. *Let $X = J(\lambda, -w_0\lambda)$ be such that λ is regular and integral. Then the multisegment $\mathbf{m}(\lambda, -w_0\lambda)$ is twisted-elliptic if and only if the following holds for $\lambda = (\lambda_1 > \lambda_2 > \dots > \lambda_n)$:*

If $0 > \lambda_{i+1} > \dots > \lambda_n$, then there exists $a_1 < \dots < a_{n-i}$ such that $\lambda_{a_j} = -\lambda_{n+1-j}$.

Proof. This can be checked directly by the definition of twisted-elliptic multisegments. For instance, for the *if* part of the lemma, let λ^- be obtained by removing all λ_{a_j} and λ_{n+1-j} 's from λ . Then it can be checked that

$$\text{con}(\mathbf{m}(\lambda, -w_0\lambda)) = \text{con}(\mathbf{m}(\lambda^-, -w_0\lambda^-)),$$

so that one is reduced to the case when there are no negative coordinates of λ . In such a case, one can easily prove that the lemma holds. The *only if* part can be proved similarly. \square

Remark 10.2. For **all** finite dimensional representations $X = J(\lambda, -w_0\lambda)$ with nonzero Dirac cohomology, one can always ‘thicken’ X by, for instance, tensoring with a unitary character (see Lemma 10.5 below), so that the thickened module $X'' = J(\lambda'', -w_0\lambda'')$ with the i -th coordinate $\lambda''_i \geq 0$ for all i , satisfying the hypothesis of the above Lemma.

Corollary 10.3. *Let $X = J(\lambda, -w_0\lambda)$ be a finite dimensional representation of $GL_n(\mathbb{C})$ such that $\text{ht}(X) = m > 0$. Then $\Gamma_{n,m}(X)$ has nonzero Dirac cohomology if and only if the multisegment $\mathbf{m}(\lambda, -w_0\lambda)$ is twisted-elliptic.*

Furthermore, if λ is such that $2(\lambda - \rho)$ has nonnegative coordinates, then the W -type of $\Gamma_{n,m}(X)$ contributing to its Dirac cohomology is $\tau_{2(\lambda - \rho)}$.

Proof. The first statement is immediate from [Ch18, Theorem 7.8]. For the second statement, note that the proof of Theorem A.1 implies that $F_{2(\lambda - \rho)}$ (appearing with multiplicity one) contributes to the Dirac cohomology of X . By the Schur-Weyl duality (Theorem 9.7), $\tau_{2(\lambda - \rho)}$ appears in $\Gamma_{n,m}(X)$ with multiplicity one. Then the result follows from checking that the multiplicity

$$\dim \text{Hom}_{\widetilde{W}}(\tau_{\lambda(\mathbf{m}^{temp})}, \tau_{2(\lambda - \rho)} \otimes S)$$

is nonzero (c.f. [Ch18, Lemma 7.20]). \square

Example 10.4. *Let $X = J((7, 3, -3), (3, -3, -7))$ be a finite dimensional $GL_3(\mathbb{C})$ -module. Then $\mathbf{m}((7, 3, -3), (3, -3, -7))$ is twisted-elliptic by Lemma 10.1, with*

$$\text{con}(\mathbf{m}((7, 3, -3), (3, -3, -7))) = \text{con}(\mathbf{m}((7), (-7))) = \left\{ -\frac{13}{2}, -\frac{11}{2}, \dots, \frac{11}{2}, \frac{13}{2} \right\}.$$

By the first paragraph of Corollary 10.3, $\Gamma_{3,14}(X) = \text{St}(\mathfrak{m}((7, 3, -3), (3, -3, -7)))$ has nonzero Dirac cohomology - indeed, the character formula of X is given by (see (25) in the Appendix):

$$\begin{aligned} \sum_{s \in S_3} \det(s) X((7, 3, -3), s(3, -3, -7)) &= X((7, 3, -3), (3, -3, -7)) - X((7, 3, -3), (-3, 3, 7)) \\ &\quad - X((7, 3, -3), (3, -7, -3)) + X((7, 3, -3), (-3, -7, 3)) \\ &\quad + X((7, 3, -3), (-7, 3, -3)) - X((7, 3, -3), (-7, -3, 3)). \end{aligned}$$

Consequently, one can obtain the character formula of $\Gamma_{3,14}(X)$ by applying Theorem 6.4 on the above terms. Indeed, the last summand of the above formula maps to zero by $\Gamma_{3,14}$, while the second last summand yields a tempered module $\text{St}([- \frac{13}{2}, \frac{13}{2}])$. Note that it is the only tempered summand in the character formula of $\Gamma_{3,14}(X)$. As a result, [Ch18, Theorem 6.4] implies that the Dirac index (and hence the Dirac cohomology) of $\Gamma_{3,14}(X)$ is nonzero.

Since X does not satisfy the second hypothesis of Corollary 10.3, one has to invoke the algorithm in [Ch18, Section 7.6] to conclude that the W -type in $\Gamma_{3,14}(X)$ contributing to Dirac cohomology is $\tau_{(12,1,1)}$, occurring with multiplicity one. On the other hand, the decomposition of $X|_K \cong F_{(6,3,-2)} \otimes F_{(6,3,-2)}$ into K -types contains $F_{(12,1,1)}$ with multiplicity one. This matches with the Schur-Weyl duality given in Theorem 9.7.

Finally, if one considers the ‘thickened’ module $X'' := J((10, 6, 0), (0, -6, -10)) = X \otimes (\frac{\det}{|\det|})^6$, then X'' satisfies both hypotheses of Corollary 10.3, and the W -type contributing to the Dirac cohomology of $\Gamma_{3,32}(X'')$ is $\tau_{2(\lambda-\rho)} = \tau_{(18,12,2)}$.

10.2. Dirac series. An interesting part of the unitary dual is called the *Dirac series*, that is, unitary representations with nonzero Dirac cohomology. As we see in the previous section, the Arakawa-Suzuki functor $\Gamma_{n,m}$ maps irreducible, unitary representations of $\text{GL}_n(\mathbb{C})$ to irreducible, unitary representations \mathbb{H}_m . As a special case, one may apply Theorem 9.4 and get:

Lemma 10.5. *Let*

$$\text{Sp}_{n,d} := J\left(\left(\frac{(n-1)+d}{2}, \frac{(n-3)+d}{2}, \dots, \frac{-(n-1)+d}{2}\right), \left(\frac{(n-1)-d}{2}, \frac{(n-3)-d}{2}, \dots, \frac{-(n-1)-d}{2}\right)\right)$$

be the Harish-Chandra module of the unitary character $g \mapsto \det^d(g)/|\det^d(g)|$ in $\text{GL}_n(\mathbb{C})$. Suppose $d > 0$, so that $m := \text{ht}(\text{Sp}_{n,d}) = nd > 0$. Then $\Gamma_{n,m}(\text{Sp}_{n,d}) = a(n, d)$, where

$$a(n, d) := \text{St}\left(\left\{\left[\frac{n-d}{2}, \frac{n+d}{2} - 1\right], \left[\frac{n-d}{2} - 1, \frac{n+d}{2} - 2\right], \dots, \left[-\frac{n+d}{2} + 1, -\frac{n-d}{2}\right]\right\}\right)$$

is the **Speh module** (here we use the notations in [BC14, Section 3.2]).

In Type A , the classification of Dirac series for $\text{GL}_n(\mathbb{C})$ and \mathbb{H}_m are given in [BP11], [DW20] and [BC14] respectively. In particular, one has:

Corollary 10.6. *Let X be in the Dirac series of $\text{GL}_n(\mathbb{C})$ such that $\text{ht}(X) = m > 0$. Then $\Gamma_{n,m}(X) \neq 0$ is also in the Dirac series of \mathbb{H}_m .*

Proof. By the description of Dirac series given in the beginning of [DW20, Section 2], π is in the Dirac series if and only if

$$X \cong (\mathrm{Sp}_{n_1, d_1} \times \cdots \times \mathrm{Sp}_{n_l, d_l}) \times (\mathrm{Sp}_{n'_1, d'_1} \times \cdots \times \mathrm{Sp}_{n'_t, d'_t}),$$

where $n_i + d_i - 1 \equiv 0 \pmod{2}$ for all $1 \leq i \leq l$, $n'_j + d'_j - 1 \equiv 1 \pmod{2}$ for all $1 \leq j \leq t$, and

$$\begin{aligned} n_1 + d_1 - 1 &\geq -n_1 + d_1 + 1 > n_2 + d_2 - 1 \geq -n_2 + d_2 - 1 > \cdots > n_l + d_l - 1 \geq -n_l + d_l + 1; \\ n'_1 + d'_1 - 1 &\geq -n'_1 + d'_1 + 1 > n'_2 + d'_2 - 1 \geq -n'_2 + d'_2 - 1 > \cdots > n'_t + d'_t - 1 \geq -n'_t + d'_t + 1. \end{aligned}$$

On the other hand, Lemma 10.5 implies that

$$\Gamma_{n,m}(X) = (a(n_1, d_1) \times \cdots \times a(n_l, d_l)) \times (a(n'_1, d'_1) \times \cdots \times a(n'_t, d'_t)).$$

Since $n + d - 1 \geq |n - d| + 1 \geq -n + d + 1$ whenever $n, d > 0$, it is easy to see that the inequalities above satisfy Equation (3.6.2) of [BC14], and hence $\Gamma_{n,m}(X)$ has nonzero Dirac cohomology. \square

It would be beneficial to give a theoretical account of the results in this section without invoking [BC14], [Ch18] or [DW20]. It is of interest to see how the Dirac operator ‘transforms’ upon applying the Arakawa-Suzuki functor. As seen in [BCT12], the generators y_l in Definition 2.3 play the role of differential operators for real groups. Hence, a more general question is to see how the work of [Hu93] and [BKZ09] – which roughly speaking uses differential operators to study reducibility for generalized Verma modules – can be transferred to the \mathbb{H}_m -modules.

11. APPLICATIONS FROM BERNSTEIN-ZELEVINSKY DERIVATIVES

11.1. Tensor products on $GL_n(\mathbb{C})$. In this section, we study \mathbf{BZ}_τ for $\tau = \mathrm{triv}$, the trivial representation of S_i and sgn , the sign representation. We first recall the following multiplicity-free result:

Theorem 11.1. [CS19, Ch21, Ch22+] *Let $\psi \in \mathrm{Irr}(\mathbb{H}_m)$. Then, for any $\psi' \in \mathrm{Irr}(\mathbb{H}_{m-i})$,*

$$\dim \mathrm{Hom}_{\mathbb{H}_{m-i}}(\mathbf{BZ}_{\mathrm{triv}}(\psi), \psi') \leq 1,$$

$$\dim \mathrm{Hom}_{\mathbb{H}_{m-i}}(\mathbf{BZ}_{\mathrm{sgn}}(\psi), \psi') \leq 1.$$

Proof. The original version is stated for the sign representation, see [Ch21, Proposition 2.5] and [Ch22+]. But one can then transfer to the trivial representation by using the Iwahori-Matsumoto dual. More precisely, the Iwahori-Matsumoto involution $IM_k : \mathbb{H}_k \rightarrow \mathbb{H}_k$ is given as:

$$w \mapsto (-1)^{l(w)} w \quad \text{for } w \in S_k, \quad y_j \mapsto -y_j \quad \text{for all } j.$$

Then one checks that $IM_{n-i} \circ \mathbf{BZ}_{\mathrm{sgn}} \circ IM_n = \mathbf{BZ}_{\mathrm{triv}}$, where sgn and triv are the sign and trivial representations of S_i respectively. Since those IM defines a categorical equivalence, the result then follows from the sign case in [CS19, Ch22+]. The multiplicity freeness comes from the Bernstein-Zelevinsky theory and the multiplicity one of branching law in [AGRS10, Ch23]. \square

Using the functor $\Gamma_{n,m}$, we deduce that:

Corollary 11.2. *Let X be an irreducible Harish-Chandra module of $\mathrm{GL}_n(\mathbb{C})$. Let $S^i V$ and $\wedge^i V$ be the i -th symmetric tensor representation and the i -th exterior tensor representation of V respectively. Then, for any irreducible Harish-Chandra module X' of $\mathrm{GL}_n(\mathbb{C})$,*

$$\dim \mathrm{Hom}_{\mathrm{GL}_n(\mathbb{C})}(X \otimes \wedge^i V, X') \leq 1$$

$$\dim \mathrm{Hom}_{\mathrm{GL}_n(\mathbb{C})}(X \otimes S^i V, X') \leq 1.$$

Proof. We only explain the statement for $\wedge^i V$ and the one for $S^i V$ is very similar. We shall consider k to be sufficiently large such that $\chi^{\otimes k} \otimes \pi'$ is thickened. Let $\tilde{X}' = \chi^{\otimes k} \otimes \pi'$ and let $\tilde{X} = \chi^{\otimes k} \otimes X$. Then, it suffices to show that

$$\dim \mathrm{Hom}_{\mathrm{GL}_n(\mathbb{C})}(\tilde{X} \otimes \wedge^i V, \tilde{X}') \leq 1.$$

By the exactness of $\Gamma_{n,m}$, we have:

$$\dim \mathrm{Hom}_{\mathrm{GL}_n(\mathbb{C})}(\tilde{X} \otimes \wedge^i V, \tilde{X}') \leq \dim \mathrm{Hom}_{\mathbb{H}_{m-i}}(\Gamma_{n,m-i}(\tilde{X} \otimes \wedge^i V), \Gamma_{n,m-i}(\tilde{X}')).$$

Recall that $\mathbb{S}_{\mathrm{triv}}(V) = \wedge^i V$ by the Schur-Weyl duality. But now, by Theorems 8.2 and 11.1, the latter one has dimension at most one, as desired. \square

We remark that [Ch22+d] also determines (up to applying the Iwahori-Matsumoto involution) when the inequalities in Theorem 11.1 are equalities. This in particular gives a list of possible simple quotients for $X \otimes \wedge^i V$ and $X \otimes S^i V$. One may expect that the list should exhaust those simple quotients.

Incorporating the improved multiplicity result in [Ch23, Theorem 1], we also have:

Corollary 11.3. *Let X be a standard module in \mathcal{HC}_n . Then for any irreducible Harish-Chandra module X' of $\mathrm{GL}_n(\mathbb{C})$,*

$$\dim \mathrm{Hom}_{\mathrm{GL}_n(\mathbb{C})}(X \otimes S^i V, X') \leq 1.$$

Remark 11.4. While Corollaries 11.2 and 11.3 are stated for our choice of V , one can apply the Hermitian involution or the involution from $g \mapsto g^{-t}$ to get to another standard representations as well as their contragredients.

Remark 11.5. We explain an instance how the tensor product in Corollary 11.2 comes into the play of branching laws and this suggests some kind of Lefschetz principle in that direction. Let $\mathcal{S}(\mathbb{C}^n)$ and $\mathcal{S}(\mathbb{C}^n - 0)$ be the spaces of Schwartz functions on \mathbb{C}^n and $\mathbb{C}^n - 0$ respectively. The branching law for the equal rank Fourier-Jacobi model is to study simple $\mathrm{GL}_n(\mathbb{C})$ -quotients of

$$\pi \hat{\otimes} \mathcal{S}(\mathbb{C}^n)$$

for some irreducible representation π of $\mathrm{GL}_n(\mathbb{C})$. Using Borel's lemma, we have a natural short exact sequence for $\mathrm{GL}_n(\mathbb{C})$ -representations,

$$0 \rightarrow \mathcal{S}(\mathbb{C}^n - 0) \rightarrow \mathcal{S}(\mathbb{C}^n) \rightarrow \mathbb{C}[[z_1, \dots, z_n, \bar{z}_1, \dots, \bar{z}_n]] \rightarrow 0$$

and so $0 \rightarrow \mathcal{S}(\mathbb{C}^n - 0) \hat{\otimes} \pi \rightarrow \mathcal{S}(\mathbb{C}^n) \hat{\otimes} \pi \rightarrow \mathbb{C}[[z_1, \dots, z_n, \bar{z}_1, \dots, \bar{z}_n]] \hat{\otimes} \pi \rightarrow 0$. Here $\hat{\otimes}$ is the completed projective tensor product. In particular, the latter tensor product admits a filtration of the form $\pi \otimes S^i V \otimes S^j V'$ (which one may interpret as an analogue of a layer of Bernstein-Zelevinsky filtration in [Ch22+d]) and so we expect [Ch22+, Ch22+d] will be useful in determining those simple quotients). Here V and V' are the conjugate standard and standard representations of $\mathrm{GL}_n(\mathbb{C})$ respectively.

12. APPLICATIONS ON PARABOLIC INDUCTIONS

12.1. Reducibility of parabolic inductions. Recall that χ is defined in Section 9.2. Determining the irreducibility of parabolic inductions for Harish-Chandra modules can be transferred to the same problem for the graded Hecke algebra side in the following Corollary 12.1 and the precise way to do so is indicated in its proof.

Corollary 12.1. *The irreducibility of parabolic induction for the Harish-Chandra category \mathcal{HC}_n can be detected from \mathcal{H}_m (for some m) by using $\Gamma_{n,m}$.*

Proof. Let X and Y be two irreducible Harish-Chandra modules of $\mathrm{GL}_{n_1}(\mathbb{C})$ and $\mathrm{GL}_{n_2}(\mathbb{C})$ respectively. We can choose sufficiently large k such that for any composition factor Z in $X \times Y$, $\chi^{\otimes k} \otimes Z$ is thickened. Since $\chi^{\otimes k} \otimes$ defines an equivalence of categories, it is clear that $\chi^{\otimes k} \otimes (X \times Y)$ is irreducible if and only if $X \times Y$ is irreducible. Now, let m be the height of Z for some (and so all) composition factor Z in $\chi^{\otimes k} \otimes (X \times Y)$. Since $\Gamma_{n_1+n_2,m}$ sends any simple composition factor of $\chi^{\otimes k} \otimes (X \times Y)$ to a non-zero irreducible module (Theorem 9.4), we then have that $\chi^{\otimes k} \otimes (X \times Y)$ is irreducible if and only if $\Gamma_{n_1+n_2,m}(\chi^{\otimes k} \otimes (X \times Y)) \cong \Gamma_{n_1,m_1}(\chi^{\otimes k} \otimes X) \times \Gamma_{n_2,m_2}(\chi^{\otimes k} \otimes Y)$ is irreducible. Here $m_1 = \mathrm{ht}(\chi^{\otimes k} \otimes X)$ and $m_2 = \mathrm{ht}(\chi^{\otimes k} \otimes Y)$, and the last isomorphism follows from Theorem 5.7 and Lemma 6.1. \square

For example, Corollary 12.1 gives another perspective on the classical result that the product of two irreducible unitary representations is still irreducible [Sa89, Theorem 3.1] (c.f. [Vo86, Theorem 6.18(b)] and [Bu03]).

Recall in Section 10 that, for a finite-dimensional representation X of $\mathrm{GL}_n(\mathbb{C})$, $X \cong J(\lambda_L, \lambda_R)$ for some regular dominant λ_L and λ_R in \mathfrak{h}_0^* . In particular, $\Gamma_{n,m}(X)$ is a ladder representation if it is non-zero.

Corollary 12.2. *Let X_1 and X_2 be irreducible representations in \mathcal{HC}_{n_1} and \mathcal{HC}_{n_2} respectively. Suppose X_1 or X_2 is finite-dimensional. Then $X_1 \times X_2$ has unique simple quotient and unique simple submodule.*

Proof. Again, we find a sufficiently large k such that $\chi^{\otimes k} \otimes Z$ is thickened for any simple composition factor Z in $X_1 \times X_2$. We have that $\chi^{\otimes k} \otimes (X_1 \times X_2) \cong (\chi^{\otimes k} \otimes X_1) \times (\chi^{\otimes k} \otimes X_2)$, and so by Theorem 5.7,

$$\Gamma_{n,m}(\chi^{\otimes k} \otimes (X_1 \times X_2)) \cong \Gamma_{n,m_1}(\chi^{\otimes k} \otimes X_1) \times \Gamma_{n,m_2}(\chi^{\otimes k} \otimes X_2),$$

where m_1 and m_2 are the heights of $\chi^{\otimes k} \otimes X_1$ and $\chi^{\otimes k} \otimes X_2$ respectively.

Now $\Gamma_{n,m_2}(\chi^{\otimes k} \otimes X_2)$ is a ladder representation, and so the product has a unique simple quotient [LM16] (with the translation via [Bo76] and [Lu89]). Since $\Gamma_{n,m}$ is exact and sends all the composition factors in $\chi^{\otimes k} \otimes (X_1 \times X_2)$ to a non-zero module, we then must have $\chi^{\otimes k} \otimes (X_1 \times X_2)$ has a unique simple quotient and so does $X_1 \times X_2$. The statement for submodule is the same. \square

We also have the following variation of Corollary 12.2:

Corollary 12.3. *Let X be an irreducible Harish-Chandra representation in \mathcal{HC}_n . Let Y be an irreducible finite-dimensional representation in $\mathcal{HC}_{n'}$ with $n' < n$. Then there exists at most one irreducible Harish-Chandra module Z of $\mathrm{GL}_{n-n'}(\mathbb{C})$ such that X is the unique simple submodule of $Y \times Z$. The statement also holds if one replaces $Y \times Z$ by $Z \times Y$.*

Proof. The analogous statement for \mathbb{H}_m -modules holds for the product between a ladder representation and an arbitrary irreducible module. The argument for passing from Harish-Chandra modules to \mathbb{H}_m -modules is similar to the one in Corollary 12.2. Here we have also used Remark 9.6(3) that after some thickening for irreducible X_1 and X_2 , $\Gamma_{n,m}(X_1) \cong \Gamma_{n,m}(X_2) \neq 0$ implies $X_1 \cong X_2$. \square

When Y is a character, there exists algorithms (e.g. [LM16]) to compute such Y in Corollary 12.3.

Remark 12.4. Here we explain how one can deduce some new explicit quotient branching laws from results of parabolic inductions. Let X be an irreducible Casselman-Wallach representation of $\mathrm{GL}_{n+1}(\mathbb{C})$. Given a character μ of $\mathrm{GL}_1(\mathbb{C})$, one can then find, using results of p -adic group side (c.f. Corollary 12.3), the Langlands parameter of an irreducible Harish-Chandra module Y of $\mathrm{GL}_n(\mathbb{C})$ (if such Y exists) such that

$$\mathrm{Hom}_{\mathrm{GL}_{n+1}(\mathbb{C})}(X, Y \times \mu) \neq 0.$$

Via Frobenius reciprocity, one then has:

$$\mathrm{Hom}_{P_{n,1}}(X, \delta^{1/2}(Y \boxtimes \mu)) \neq 0,$$

Then restricting to $\mathrm{GL}_n(\mathbb{C})$ (viewed as a subgroup of $P_{n,1}$ via the embedding $g \mapsto \begin{pmatrix} g & \\ & 1 \end{pmatrix}$), we obtain a branching law:

$$\mathrm{Hom}_{\mathrm{GL}_n(\mathbb{C})}(X, \delta^{1/2}Y) \neq 0.$$

For example, let $X = J((2, 2, 1), (1, 1, 0))$. Then $J((2, 1), (1, 0)) \times J((2, 1))$ has X as a simple submodule and so $\delta^{1/2}Y = J((\frac{5}{2}, \frac{3}{2}), (\frac{3}{2}, \frac{1}{2}))$ is a simple quotient of $X|_{\mathrm{GL}_2(\mathbb{C})}$.

Remark 12.5. There are other results of parabolic inductions (or in the form of Jacquet functors) in e.g. [LM16, Ch22+b, Ch22+c, Ch22+d] and references therein. The reader is invited to translate those results and find applications.

12.2. Remarks on higher structure. As seen in [Ch22+b], it is useful to determine some higher structure for branching problems. However, for such applications, we need to show that $\Gamma_{n,m}$ preserves some higher structure. The related application is to transfer the results in [Ch22+b] to the Harish-Chandra category as well as determining the equalities in Corollary 11.2. This question for the Arakawa-Suzuki functor [AS98] is studied in [Fu18], which shows a fully-faithful embedding from the deformed BGG category to the completion of module category of \mathbb{H}_m . We leave our case for future navigation.

APPENDIX A. DIRAC COHOMOLOGY OF FINITE DIMENSIONAL REPRESENTATIONS

In this section, we study the Dirac cohomology of finite dimensional representations for all complex connected reductive Lie groups, which includes $GL_n(\mathbb{C})$ as a special case. We shall use the notations in [BP11, Section 2], which in particular coincide with notations for the $GL_n(\mathbb{C})$ case. Firstly, an irreducible Harish-Chandra module $J(\lambda_L, \lambda_R)$ is finite-dimensional if and only if $\lambda_L, \lambda_R \in \mathfrak{h}_0^*$ are regular and integral. Moreover, its K -spectrum is equal to:

$$(24) \quad J(\lambda_L, \lambda_R)|_K = F_{\lambda_L - \rho} \otimes F_{\lambda_R - \rho}^* = F_{\lambda_L - \rho} \otimes F_{-w_0(\lambda_R - \rho)}$$

In particular, the PRV component of the above tensor product has extremal weight $(\lambda_L - \rho) + w_0(-w_0(\lambda_R - \rho)) = \lambda_L - \lambda_R$, which is equal to that of the lowest K -type of $J(\lambda_L, \lambda_R)$.

The main result of [HP02] (reformulated in Equation (2.2) of [BP11] for complex groups) implies that all finite dimensional representations of complex groups with nonzero Dirac cohomology must be of the form $X = J(\lambda, -w_0\lambda)$, where λ is regular integral and $w_0 \in W$ is the longest element in the Weyl group.

Now consider the *Dirac operator*

$$D : X \otimes S \longrightarrow X \otimes S,$$

where S is the spinor module, which is isomorphic to $2^{\lfloor n/2 \rfloor} F_\rho$ as a \tilde{K} -module by [BP11, Lemma 2.2]. By Weyl's unitarity trick, there is a inner product \langle, \rangle_X on π such that $\langle Ev, v' \rangle_X = -\langle v, Ev' \rangle_X$ for all $E \in \mathfrak{k}_0 + j\mathfrak{s}_0$ and $v, v' \in X$ (note that this defines a positive definite Hermitian form with respect to the anti-involution given in Remark 7.9). Consequently, as in Remark 3.2.4 of [HP06], one can define an inner product on $X \otimes S$ such that D is *skew*-Hermitian with respect to this inner product.

As a result, the Dirac cohomology of π is equal to $\ker(D) = \ker(D^2)$, and the question of finding Dirac cohomology of π reduces to studying $D^2 : X \otimes S \rightarrow X \otimes S$, where $X \otimes S \cong 2^{\lfloor n/2 \rfloor} F_{\lambda - \rho} \otimes F_{\lambda - \rho}$ as \tilde{K} -modules.

By the formula of D^2 given in [HP06, Proposition 3.1.6] for instance, D^2 acts on the γ -isotypic component of $X \otimes S$ by the scalar $-||2\lambda||^2 + ||\gamma + \rho||^2$. Then it is easy to check that the only γ -isotypic component which D^2 acts by zero is when $\gamma = (\lambda - \rho) + (\lambda - \rho) + \rho$, which implies the following:

Theorem A.1. *Let G be any complex connected reductive Lie group. Then all finite dimensional (\mathfrak{g}, K) -modules with nonzero Dirac cohomology must be of the form $X = J(\lambda, -w_0\lambda)$ for some regular integral λ . In particular, its Dirac cohomology is equal to $2^{\lfloor n/2 \rfloor} F_{2\lambda-\rho}$.*

Finally, we remark that the *twisted Dirac index* of $X = J(\lambda, -w_0\lambda)$ can also be obtained easily. Namely, the character formula of X is given by:

$$(25) \quad \sum_{w \in W} \det(w) X(\lambda, -w w_0 \lambda).$$

By Theorem 5.1 of [BPT19], only the last summand $w = w_0$ in the above formula contributes to the twisted Dirac index of X , and is equal to $\pm 2^{\lfloor n/2 \rfloor} F_{2\lambda-\rho}$ (the \pm sign depends on the choice of ϵ in the theorem). This gives another evidence on the validity of Theorem A.1.

REFERENCES

- [ALT20] J. Adams, M. van Leeuwen, P. Trapa, D. Vogan, Unitary representations of real reductive groups, *Astérisque*, **417**, 2020.
- [AGRS10] A. Aizenbud, D. Gourevitch, S. Rallis and G. Schiffmann, Multiplicity one theorems, *Ann. of Math.* (2) **172** (2010), no. 2, 1407-1434.
- [AS98] T. Arakawa, T. Suzuki, Duality between $\mathfrak{sl}_n(\mathbb{C})$ and the degenerate affine Hecke algebra, *J. Algebra* **209** (1998), no. 1, 288-304.
- [Ba89] D. Barbasch, The unitary dual for complex classical Lie groups, *Invent. Math.* **96** (1989), no. 1, 103-176.
- [Ba10] D. Barbasch, The unitary spherical spectrum for split classical groups, *J. Inst. Math. Jussieu* **9** (2010), no. 2, 265-356.
- [BC14] D. Barbasch, D. Ciubotaru, Unitary Hecke algebra modules with nonzero Dirac cohomology, *Progr. Math.*, **257** (2014), Birkhäuser/Springer, 1-20
- [BC15] D. Barbasch, D. Ciubotaru, Ladder representations of $GL(n, \mathbb{Q}_p)$. In: Nevins, M., Trapa, P. (eds) *Representations of Reductive Groups. Progress in Mathematics* **312** (2015) Birkhäuser, Cham.
- [BCT12] D. Barbasch, D. Ciubotaru, P. Trapa, Dirac cohomology for graded affine Hecke algebras, *Acta Math.* **202** (2012), no. 2, 197-227.
- [BP11] D. Barbasch, P. Pandžić, Dirac cohomology and unipotent representations of complex groups, *Noncommutative geometry and global analysis, Contemp. Math.* **546**, Amer. Math. Soc., Providence, RI, 2011, 1-22.
- [BPT19] D. Barbasch, P. Pandžić, P. Trapa, Dirac index and twisted characters, *Trans. Amer. Math. Soc.* **371**, Number 3 (2019) 1701-1733.
- [BKZ09] L. Barchini, A. C. Kable, R. Zierau, Conformally invariant systems of differential operators, *Adv. Math.* **221** Issue 3 (2009), 788-811. doi.org/10.1016/j.aim.2009.01.006.
- [Bu03] Baruch, E.M., Proof of Kirillov's conjecture, *Ann. of Math.* **158** (2003), 207-252.
- [BG80] J. Bernstein and S. Gelfand, Tensor products of finite and infinite dimensional representations of semisimple Lie algebras, *Compositio Math.* **41** (1980), 245-285.
- [Bo76] A. Borel, Admissible representations of a semi-simple group over a local field with vectors fixed under an Iwahori subgroup, *Invent Math* **35** (1976), 233-259. https://doi.org/10.1007/BF01390139
- [Ca22] K. Calvert, Compact Schur-Weyl duality and the Type B/C VW-algebra, *Adv. Math.* **407** (2022), 108575. https://doi.org/10.1016/j.aim.2022.
- [Ch18] K.Y. Chan, A vanishing theorem for Dirac cohomology of standard modules. *Adv. Math.* **325** (2018), 274-311.
- [Ch21] K.Y. Chan, Homological branching law for $(GL_{n+1}(F), GL_n(F))$: projectivity and indecomposability, *Invent. math.* **225** (2021), 299-345. https://doi.org/10.1007/s00222-021-01033-5
- [Ch23] K.Y. Chan, Ext-multiplicity theorem for standard representations of (GL_{n+1}, GL_n) , *Math. Z.* **303**, 45 (2023). https://doi.org/10.1007/s00209-022-03198-y
- [Ch22+] K.Y. Chan, Construction of simple quotients of Bernstein-Zelevinsky derivatives and highest derivative multisegments I: reduction to combinatorics, preprint (2022).
- [Ch22+b] K.Y. Chan, On the product functor on the inner forms of general linear group over a non-Archimedean local field, arXiv:2206.10928

- [Ch22+c] K.Y. Chan, On commutations of derivatives and integrals of \square -irreducible representations for p -adic GL , arXiv:2210.17249 (2022)
- [Ch22+d] K.Y. Chan, Quotient branching law for p -adic (GL_{n+1}, GL_n) I: generalized Gan-Gross-Prasad relevant pairs, arXiv:2212.05919 (2022)
- [CS19] K.Y. Chan and G. Savin, Bernstein-Zelevinsky derivatives: a Hecke algebra approach, International Mathematics Research Notices, **3** (2019), 731-760. <https://doi.org/10.1093/imrn/rnx138>
- [CT11] D. Ciubotaru and P. Trapa, Functors for unitary representations of real classical groups and affine Hecke algebras, Adv. Math. **227** no. 4 (2011), 1585-1611.
- [CT12] D. Ciubotaru and P. Trapa, Duality for $GL(n, \mathbb{R})$, $GL(n, \mathbb{Q}_p)$, and the degenerate affine Hecke algebra for $gl(n)$, Amer. J. Math. **134** (2012), 1-30.
- [DW20] C.-P. Dong and K.D. Wong, Scattered representations of $SL(n, \mathbb{C})$, Pacific J. Math. **309** (2020), no. 2, 289-312.
- [Du75] M. Duflo, Représentations irréductibles des groupes semi-simples complexes, Lecture Notes in Math. **497** (1975), 26-88.
- [EFM09] P. Etingof, R. Freund, X. Ma, A Lie-theoretic construction of representations of the degenerate affine and double affine Hecke algebras of type BC_n , Representation Theory **13** (2009), 33-49.
- [Ev96] S. Evens, The Langlands classification for graded Hecke algebras, Proc. Amer. Math. Soc., **124** (4) (1996), 1285-1290.
- [Fu18] R. Fujita, Tilting modules of affine quasi-hereditary algebras, Advances in Mathematics, **324** (2018) 241-266. doi.org/10.1016/j.aim.2017.11.013.
- [Go70] R. Godement, Notes on Jacquet-Langlands' theory, preprint. Princeton (1970)
- [HP02] J.-S. Huang and P. Pandžić, Dirac cohomology, unitary representations and a proof of a conjecture of Vogan, J. Amer. Math. Soc. **15**(1) (2002), 185-202.
- [HP06] J.-S. Huang, P. Pandžić, Dirac Operators in Representation Theory, Mathematics: Theory and Applications, Birkhäuser, 2006.
- [Hu93] J.-S. Huang, Intertwining differential operators and reducibility of generalized Verma modules. Math. Ann. **297** (1993), 309-324. <https://doi.org/10.1007/BF01459504>
- [Knp86] A. Knap, Representation Theory of Real Semisimple Groups: an Overview Based on Examples, Princeton University Press, Princeton, New Jersey, 1986.
- [LM16] E. Lapid, A. Mínguez, On parabolic induction on inner forms of the general linear group over a non-Archimedean local field, Sel. Math. New Ser. **22** (2016), 2347-2400.
- [Lu89] G. Lusztig, Affine Hecke algebras and their graded version, J. Amer. Math. Soc. **2** (1989), 599-685.
- [Od07] H. Oda, Generalization of Harish-Chandra's basic theorem for Riemannian symmetric spaces of non-compact type, Adv. Math. **208** (2) (2007) 549-596.
- [PRV67] K.R. Parthasarathy, R. Ranga Rao, and V. S. Varadarajan. Representations of Complex Semi-Simple Lie Groups and Lie Algebras, Annals of Mathematics **85**, no. 3 (1967): 383-429. doi.org/10.2307/1970351.
- [Ro85] J.D. Rogawski, On modules over the Hecke algebra of a p -adic group, Invent. Math. **79**, 443-465 (1985). <https://doi.org/10.1007/BF01388516>
- [Ro86] J.D. Rogawski, Representations of $GL(n)$ over a p -adic field with an Iwahori-fixed vector. Israel J. Math. **54** (1986), 242-256. <https://doi.org/10.1007/BF02764944>
- [Sa89] S. Sahi, On Kirillov's conjecture for archimedean fields, Compos. Math. **72** (1989), 67-86.
- [Su98] T. Suzuki, Rogawski's conjecture on the Jantzen filtration for the degenerate affine Hecke algebra of type A, Represent. Theory **2** (1998), 393-409
- [St67] E. M. Stein, Analysis in Matrix Spaces and Some New Representations of $SL(N, \mathbb{C})$, Annals of Mathematics, Second Series, **86**, No. 3 (1967), 461-490.
- [Ta86] M. Tadić, On the classification of irreducible unitary representations of $GL(n)$ and the conjectures of Bernstein and Zelevinsky, Ann. Sci. École Norm. Sup., **19** (1986), 335-382.
- [Ta09] M. Tadić, $GL(n, \mathbb{C})^\wedge$ and $GL(n, \mathbb{R})^\wedge$, Automorphic forms and L-functions II. Local aspects (2009), 285-313.
- [Tr06] F. Trèves, Topological vector spaces, distributions and kernels, Mineola, N.Y.: Dover Publications.
- [Vo81] D. Vogan, Representations of real reductive groups, Progr. Math., vol. 15, Birkhäuser Boston, Boston, MA, 1981.
- [Vo86] D. Vogan, The unitary dual of $GL(n)$ over an archimedean field. Invent Math **83**, 449-505 (1986). <https://doi.org/10.1007/BF01394418>
- [Ze80] A. Zelevinsky, Induced representations of reductive p -adic groups II, Ann. Sci. Ecole Norm. Sup. **13** (1980), 154-210.

- [Ze85] A. Zelevinsky, Two remarks on graded nilpotent classes, *Uspehi Mat. Nauk*, 40 (1985), no. 1 (241), 199-200.
[Zh74] D. P. Zhelobenko, *Harmonic analysis on complex semisimple Lie groups*, Mir, Moscow, 1974.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF HONG KONG
Email address: `kychan1@hku.hk`

SCHOOL OF SCIENCE AND ENGINEERING, THE CHINESE UNIVERSITY OF HONG KONG, SHENZHEN, GUANGDONG
518172, CHINA
Email address: `kayue.wong@gmail.com`