

Chapter 1

Personal History with MEF and Some Related Topics

Helen Au-Yang* and Jacques H.H. Perk

*Department of Physics, Oklahoma State University,
145 Physical Sciences, Stillwater, OK, 74078-3072, USA
perk@okstate.edu*

We present our personal histories with Michael Fisher. We describe how each one of us first came to Cornell University. We also discuss our many subsequent interactions and successful collaborations with him on various physics projects.

1. From Shanghai to Cornell University

Born under the name Ou Yang Yee Sun in Shanghai and after having suffered through much of the great famine of 1959–1962, Helen got official permission to join her parents in Hong Kong. It was quite a shock, not knowing Cantonese and English, as she only spoke Shanghainese, Mandarin and some Russian that she had learned in high school. Fortunately, the written Chinese is universal and Helen found a poster announcing a competition for two fellowships at Chu Hai College, Kowloon. Even though Helen had not yet finished high school, she won one of them and was admitted under the new name Helen Au-Yang. After graduating in 1965 she was admitted to San Diego State University.

In 1968 Helen entered graduate school at the State University of New York in Stony Brook. Here she generalized the work of McCoy and Wu¹ on the Ising model on the half plane with boundary magnetic field to the case with boundary coupling different from the coupling in the bulk. Thus she provided exact results² related to the series, Monte Carlo and mean field results of Binder and Hohenberg^{3,4} on 2D and 3D Ising and Heisenberg models with a boundary.

Helen continued with the main topic of her PhD thesis:⁵ Layered Ising models with period n , deriving exact results for the specific heat⁶ and for the pair correlation in a row parallel to the layering.⁷ That last calculation led to a block Toeplitz determinant with a 2×2 matrix generating function $a(\xi)$. After reviewing the then known theorems for such determinants, especially for those for which $a(\xi)$ can be properly factorized, Helen gave $n = 2$ results for the $T < T_c$ spontaneous

*Helen's part of the story is told by her coauthor and husband of 40 years, as, at the time of writing, she was not (yet) sufficiently recovered from a stroke.

magnetization and for the leading long-distance behavior of the pair correlation for $T < T_c$ and for $T > T_c$.

In 1973 Helen graduated with a PhD in Theoretical Physics, with her thesis signed by Professors B. M. McCoy, C. N. Yang, T. T. Wu and others.^a She became a postdoc with Michael Fisher at Cornell University.

2. Helen at Cornell University

With Helen in his group, Michael Fisher continued the work on his project “Bounded and inhomogeneous Ising models” started with Arthur Ferdinand^{9,10} many years earlier. In 1969 they had published finite-size effects on the specific heat of an $m \times n$ Ising lattice, asymptotically for large n with $\xi = m/n$ fixed.¹⁰

Helen and Michael first studied the specific-heat scaling function for an infinitely long Ising strip of width n in the limit $n \rightarrow \infty$.¹¹ After this, they studied the finite-size effects of regularly spaced point defects,^{12–14} also incorporating some earlier calculations of Ferdinand.

Michael was very happy to have Helen around, as he liked things to be done exactly, if they could be done exactly. One time a student solved a cubic equation numerically by computer. Michael told Helen, “Helen, you show him how that is done exactly!”

Michael wanted to keep Helen for a longer time by creating a special research position for her. He asked C. N. Yang to take her back as a postdoc for a year, while he created such a new position at Cornell. At Stony Brook Helen calculated Ising multispin correlations along the diagonal¹⁵ to check the operator reduction formulae of Kadanoff. She also calculated a new scaling form for the four-spin correlation along the diagonal.¹⁶

Back at Cornell University, Helen got involved in a generalization of earlier work,¹¹ namely the study of an $n \times \infty$ Ising strip with field h_1 on the first infinite layer and field h_n on the last (n th) layer. One motivation for this work was to investigate the validity of the *ad hoc* local free energy functional of Fisher and de Gennes.¹⁷ In spite of serious limitations of that theory, their most striking predictions were confirmed and finite-size scaling forms were derived in the limit $n \rightarrow \infty$.^{18,19} A more detailed summary of this research is given in section 4.3 of Fisher’s “Simple Ising Models Still Thrive!” lecture.²⁰

At the same time period much numerical work was done to study inhomogeneous differential approximants for power series of one or more variables²¹ Nowadays, with Maple and Mathematica available, this is a lot easier, but then it was not. At that time using Fortran, one had to prepare a stack of punchcards for each example with up to 80 characters per card and hand that in for processing overnight, carefully keeping the stacks of cards in the right order. This work was very labor intensive,

^aThe publication of the thesis⁵ and the papers^{6,7} got delayed, as her advisor Barry McCoy was involved with the massive project on Painlevé III scaling functions in 2D Ising correlations.⁸

as not only existing high- and low-temperature series coefficients and activity (high-field) expansions of various well-known spin models were examined, but also many test examples derived from mathematical functions with a given singularity and various background noise contributions.

One case was studied in detail: Universality tests at Heisenberg bicritical points.²² The experimentally determined amplitude ratio $Q_{\text{fit}} \approx 1.6 \pm 0.35$ found in MnF_2 was much smaller than theory seemed to predict, $Q_{\text{th}} \approx 2.39$. Noise contributions in the theory could bring that value about 10% down, but that is not enough. Fisher et al. then note²² that shifts in the critical lines compatible with experimental uncertainties could raise Q_{fit} enough to bring theory and experiment in agreement. In other words, Q_{fit} should have been reported with significantly larger error bars.

During a visit back to Stony Brook, Helen calculated the scaling form for the four-spin correlation function of two parallel nonaligned pairs in the planar Ising model.²³

Some time in 1980, Helen's oldest sister and her family were released from labor camp in Western China to settle in Hongkong with the rest of the family there. Helen then abruptly quit her position at Cornell in order to help tutor her two nieces into the Hongkong school system, using the earlier experience that she had had moving from Shanghai to Hongkong in 1962. In 1982 Helen wanted to return to the USA, and Barry McCoy and Joel Lebowitz organized a special invitation to the December 1982 Rutgers Stat. Mech. Meeting.

3. Jacques' visits to Cornell and Rutgers

In 1978 I (Jacques) received the Royal Dutch Shell Oil Travel Prize awarded at Leiden University for my thesis research. This award allowed me to visit research groups at several universities and research institutes in the USA. Thus I came to also visit Professors Fisher and Widom at Cornell University, giving a seminar on (in)stability of critical behavior,^{24,25} as that seemed more suitable than my thesis work on time-dependent correlations in XY chains. Students and postdocs had beforehand told me that Fisher had "destroyed" the speakers in the seminar the previous several weeks, but that did not happen this time. I had seen Fisher in action at earlier conferences and I was prepared. I got a few questions about how things were proved that I could answer.

Noteworthy was the party at Fisher's home. It started with a box of Chinese metal wire puzzles put in front of me. As I had about the same collection, I knew these well. I quickly took a few apart and put them together again, upon which Fisher removed the box, causing a sigh of relief among the others. Fisher then got his Spanish guitar out. He was quite accomplished and entertaining. This and the special treats Mrs. Fisher had prepared were the high points of the party.

At the December 1979 Stat. Mech. meeting at Rutgers, I asked Leo Kadanoff

if he knew that there are continuously varying critical exponents if one varies the coupling constants within one line in the bulk of the square-lattice Ising model.^{26–28} With Michael Fisher and others standing by, Kadanoff said that that is impossible, upon which I said to expect a preprint soon. Fisher and Kadanoff followed that up with some further remarks on the phenomenon.^{29,30}

At this point, we note that, like many others, both of us were very surprised that Fisher and Kadanoff did not share the 1982 Nobel Prize in Physics with Kenneth Wilson.

4. Helen and Jacques married

It all started with Barry McCoy asking Jacques to pick up Helen, who had visited Michael Fisher at Cornell, at the Greyhound bus terminal in Manhattan on New Year's eve of 1982 and to bring her to the McCoy home. Jacques and Helen did not know that Barry and Martha McCoy conspired and had a detailed plan designed to get both of us married. All the scheming worked and six days later we were engaged to be married, with the wedding taking place on January 22, beginning a very happy marriage lasting more than 40 years so far.

Soon after Helen asked, “Jacques, have you ever done anything with your quadratic difference equations for 2D Ising correlations?” This led to the first of many joint papers following, namely a letter with fairly complete results to calculate critical Ising pair correlations³¹ and even a result for the monomer-monomer correlation in the square dimer problem. Many years later, students in the statistical mechanics class of a colleague told us that this letter was cited in Pathria's textbook.³²

Having seen several papers on chiral clock models written at Cornell University during her time there, Helen was led to extend the search for integrable Potts models to chiral ones, discovering the genus 10 curve condition for solving the star-triangle equations for the 3-state non-selfdual chiral Potts model.³³ The full parametrization of the N -state case was found during a visit to Canberra³⁴ and the history of these discoveries is described in a Topical Review celebrating Baxter's 75th birthday and implicitly Helen's 70th.³⁵

5. Later collaborations with Michael Fisher

Michael Fisher had been very apprehensive about our marriage at the beginning, but he noted that Helen was very happy, seeing her at several of Lebowitz's Stat. Mech. meetings at Rutgers.

Then, at the end of 2008, Fisher called me (Jacques) that he wanted a calculation done and that Helen was the only one he trusted to do it correctly. I told him that he was free to ask Helen and let her decide what to do. Fisher then emailed a letter in his typical style, which is reproduced in the appendix. Fisher wanted some exact calculations done on the specific heat to qualitatively understand some features seen



Fig. 1. Helen and Michael at December 2013 Stat. Mech. Meeting at Rutgers University.

in the experiments of Gasparini's group in Buffalo.^{36,37}

There was some delay, but in the end two papers were published on 2D Ising models with alternating layers with weaker and stronger interactions,^{38,39} the second with the details left out in the first one. Fisher was very happy that I helped with some of the LaTeX coding and cleaning up the figures.

Later, Helen thought of a better way to compare with an array of 3D cubes of Helium covered with a 2D film. Correspondingly she studied 2D Ising layers connected by layers of 1D Ising strings, with the strings 1, 2, or 3 units apart. First she did a 60 page calculation of the free energy in her usual fine print handwriting using Hamm's method,⁴⁰ followed by a 20 page calculation, reproducing the results using another method.¹³ When she showed me the results, we could guess the general answer for Ising strings N units apart and next prove it.^{41,42} We also obtained exact results for the spontaneous magnetization.^{41,43}

Finally, Fisher contacted me, as he had to do something about the 3D Ising deceptions of Zhang Zhidong. He asked me if I had any follow-up on my original comment. I sent him what I had, including my last comment.⁴⁴ Fisher got also very upset that his friend of long ago, Norman March, was fooled by the deceptions.

Together we wrote another comment,⁴⁵ in which we also introduced the statistical mechanics community to the new bootstrap results for 3D Ising critical exponents. Fisher was particularly happy, when I pointed out that the result for the correction to scaling exponent $\Delta = \omega\nu = 0.5231(12)$ is very close to the $\frac{1}{2}$ proposed by Andrea Liu and him⁴⁶

In conclusion, it has been a great pleasure to work with Michael Fisher. In some sense Michael was the conscience of the statistical mechanics community, keeping people honest with probing questions during and after their talks and often making good suggestions. He will be dearly missed.

References

1. McCoy, B. M. and Wu, T. T., Theory of Toeplitz determinants and the spin correlations of the two-dimensional Ising model. IV, *Phys. Rev.* **162**, 2, pp. 436–475 (1967).
2. Au-Yang, H., Thermodynamics of an anisotropic boundary of a two-dimensional Ising model, *J. Math. Phys.* **14**, 7, pp. 937–946 (1973).
3. Binder, K. and Hohenberg, P. C., Phase transitions and static spin correlations in Ising models with free surfaces, *Phys. Rev. B* **6**, 9, pp. 3461–3487 (1972).
4. Binder, K. and Hohenberg, P. C., Surface effects on magnetic phase transitions, *Phys. Rev. B* **9**, 5, pp. 2194–2214 (1974).
5. Au-Yang, H., A Study on Layered Ising Model, Ph. D. Thesis, State University of New York at Stony Brook, 1974, pp. i–v, 1–94.
6. Au-Yang, H. and McCoy, B. M., Theory of layered Ising models: Thermodynamics, *Phys. Rev. B* **10**, 3, pp. 886–891 (1974).
7. Au-Yang, H. and McCoy, B., Theory of layered Ising models. II. Spin correlation functions parallel to the layering, *Phys. Rev. B* **10**, 9, pp. 3885–3905 (1974).
8. Wu, T. T., McCoy, B. M., Tracy, C. A. and Barouch, E., Spin-spin correlation functions for the two dimensional Ising model: Exact theory in the scaling region, *Phys. Rev. B* **13**, 1, pp. 315–374 (1976).
9. Fisher, M. E. and Ferdinand, A. E., Interfacial, boundary, and size effects at critical points, *Phys. Rev. Lett.* **19**, 4, pp. 169–172 (1967).
10. Ferdinand, A. E. and Fisher, M. E., Bounded and inhomogeneous Ising models. I. Specific-heat anomaly of a finite lattice, *Phys. Rev.* **185**, 2, pp. 832–846 (1969).
11. Au-Yang, H. and Fisher, M. E., Bounded and inhomogeneous Ising models. II. Specific-heat scaling function for a strip, *Phys. Rev. B* **11**, 5, pp. 3469–3487 (1975).
12. Fisher, M. E. and Au-Yang, H., Regularly spaced defects in Ising models, *J. Phys. C* **8**, 20, pp. L418–L421 (1975).
13. Au-Yang, H., Fisher, M. E. and Ferdinand, A. E., Bounded and inhomogeneous Ising models. III. Regularly spaced point defects, *Phys. Rev. B* **13**, 3, pp. 1238–1265 (1976).
14. Au-Yang, H., Bounded and inhomogeneous Ising models. IV. Specific-heat amplitude for regular defects, *Phys. Rev. B* **13**, 3, pp. 1266–1271 (1976).
15. Au-Yang, H., Multispin correlation functions for Ising models, *Phys. Rev. B* **15**, 5, pp. 2704–2707 (1977).
16. Au-Yang, H., Four-spin correlation function along the diagonal, *Phys. Rev. B* **16**, 11, pp. 5016–5031 (1977).
17. Fisher, M. E. and de Gennes, P.-G., Wall phenomena in a critical binary mixture, *C. R. Acad. Sc. B* **287**, 8, pp. 207–209 (1978).

18. Fisher, M. E. and Au-Yang, H., Critical wall perturbations and a local free energy functional, *Physica A* **101**, 1, pp. 255–264 (1980).
19. Au-Yang, H. and Fisher, M. E., Wall effects in critical systems: Scaling in Ising model strips, *Phys. Rev. B* **21**, 9, pp. 3956–3970 (1980).
20. Fisher, M. E., Simple Ising models still thrive! A review of some recent progress, *Physica A* **106**, 1–2, pp. 28–47 (1981).^b
21. Fisher, M. E. and Au-Yang, H., Inhomogeneous differential approximants for power series, *J. Phys. A* **12**, 10, pp. 1677–1692 (1979), **13**, 4, p. 1517, (1980) (Corrigenda).
22. Fisher, M. E., Chen, J.-H. and Au-Yang, H., Universality tests at Heisenberg bicritical points, *J. Phys. C* **13**, 18, pp. L459–L464 (1980).
23. Au-Yang, H., Four-spin correlation function of two parallel nonaligned pairs in the planar Ising model, *Phys. Rev. B* **23**, 9, pp. 4749–4754 (1981).
24. Capel, H. W., Perk, J. H. H. and Den Ouden, L. W. J., Critical-exponent renormalization and first-order transitions, *Phys. Lett. A* **66**, 6, pp. 437–439 (1978).
25. Capel, H. W., Den Ouden, L. W. J. and Perk, J. H. H., Stability of critical behaviour, critical-exponent renormalization and first-order transitions, *Physica A* **95**, 3, pp. 371–416 (1979).
26. Bariev, R. Z., Effect of linear defects on the local magnetization of a plane Ising lattice, *Zh. Eksp. Teor. Fiz.* **77**, 3, pp. 1217–1229 (1979) [English translation: *Sov. Phys. JETP* **50**, 3, pp. 613–619 (1979)].
27. McCoy, B. M. and Perk, J. H. H., Two-spin correlation functions of an Ising model with continuous exponents, *Phys. Rev. Lett.* **44**, 13, pp. 840–844 (1980).
28. McCoy, B. M. and Perk, J. H. H., *Continuous exponents of spin correlation functions of inhomogeneous layered Ising models*, *Lecture Notes in Mathematics*, Vol. 925, Springer, Berlin, pp. 12–27 (1982).
29. Section 5 of Simple Ising Models Still Thrive!²⁰
30. Kadanoff, L. P., Many-point correlation functions in a modified Ising model, *Phys. Rev. B* **24**, 9, pp. 5382–5383 (1981).
31. Au-Yang, H. and Perk, J. H. H., Ising correlations at the critical temperature, *Phys. Lett. A* **104**, 3, pp. 131–134 (1984).
32. Pathria R. K. and Beale, P. D., *Statistical Mechanics*, Third Edition, Elsevier, Oxford UK, 2011, Section 13.4.
33. Au-Yang, H., McCoy, B. M., Perk, J. H. H., Tang, S. and Yan, M.-L., Commuting transfer matrices in the chiral Potts models: Solutions of the star-triangle equations with genus greater than 1, *Phys. Lett. A* **123**, 5, pp. 219–223 (1987).
34. Baxter, R. J., Perk, J. H. H. and Au-Yang, H., New solutions of the star-triangle relations for the chiral Potts model, *Phys. Lett. A* **128**, 3–4, pp. 138–142 (1988).
35. Perk, J. H. H., The early history of the integrable chiral Potts model and the odd-even problem, *J. Phys. A: Math. Theor.* **49**, 15, (2016) 153001 (20 pp.).
36. Gasparini, F. M., Kimball, M. O., Mooney, K. P. and Diaz-Avila, M., Finite-size scaling of ⁴He at the superfluid transition, *Rev. Mod. Phys.* **80**, 3, pp. 1009–1059 (2008).
37. Perron, J. K. and Gasparini, F. M., Critical point coupling and proximity effects in ⁴He at the superfluid transition, *Phys. Rev. Lett.* **109**, 3, (2012) 035302 (4 pp.).
38. Au-Yang, H. and Fisher, M. E., Criticality in alternating layered Ising models: I. Effects of connectivity and proximity, *Phys. Rev. E* **88**, 3, (2013) 032147 (12 pp.).
39. Au-Yang, H., Criticality in alternating layered Ising models: II. Exact scaling theory,

^bNote that most of Helen’s papers up to that time are cited in this work. Helen is specially mentioned in the Acknowledgements.

- Phys. Rev. E* **88**, 3, (2013) 032148 (8 pp.).
40. Hamm, J. R., Regularly spaced blocks of impurities in the Ising model: Critical temperature and specific heat, *Phys. Rev. B* **15**, 11, pp. 5391–5411 (1977).
 41. Au-Yang, H. and Perk, J. H. H., Ising models with holes: Crossover behavior, arXiv:1806.00873.
 42. Au-Yang, H. and Perk, J. H. H., Specific heat of Ising models with holes: Mathematical details using dimer approaches, arXiv:1808.07525.
 43. Au-Yang, H. and Perk, J. H. H., Spin-spin correlations in central rows of Ising models with holes, arXiv:1907.04748.
 44. Perk, J. H. H., Comment on ‘Mathematical structure of the three-dimensional (3D) Ising model’, *Chinese Physics B* **22**, 8, (2013) 080508 (5 pp.).
 45. Fisher, M. E. and Perk, J. H. H., Comments concerning the Ising model and two letters by N.H. March, *Phys. Lett. A* **380**, 13, pp. 1339–1340 (2016).
 46. Liu, A. J. and Fisher, M. E., On the corrections to scaling in three-dimensional Ising models, *J. Stat. Phys.* **58**, 3/4, pp. 431–442 (1990).

Appendix




Fig. 2. Page 1 of Fisher's January 2009 email to Helen.

From M. Fisher 1 Jan 2009 2.

Project "Theory of Proximity & Weak Coupling Effects for Strongly Ordered Systems".

§0 To be studied via Exact Calculations for 2D Ising chequered and layered systems, and the Development (and testing) of appropriate phenomenological & scaling theories.

§R Some Relevant References.

[F] MEF J. Phys Soc Japan 26 (Suppl.) (1969) 87-93
 {Conf. Review} See Fig. 3 and text of sec. 7 pp. 90-91.

[AYMcC] H. Au-Yang & B.M. McCoy, PR B 10 (1974) 881-
 {General Layered Ising Models}

[155] [FAY] MEF & HAY J. Phys. C 8 (1975) L418-421.
 {Regular Point defects: results, scaling theory, etc.}

[156] [AYF²] HAY, MEF and A.E. Ferdinand PR B 13 (1975) 1238-65.
 "Bordered & Inhomogeneous Ising Models III" = Calc. for Point Defects.

[156A] Note Also HAY PR B 13 (1975) 1266-71. Specific Max Amplitudes

As Regards Expts on Finite Size Scaling in Helium II
 see Gasparini Refs [G1] [G2] etc on attached list.
 NOTE [G4] is a Rev. Mod. Phys. (2008) Review. to [G*]

As Regards Josephson Effects in He II see [J1] and by Packard [J2].
 " Superconductors see [S1, S2] & [S3]
 " Coupling in Bose Condensed Arrays [B1, B2]

Below: §1 Minimal Layered Ising Model; §2 Alternate Layered Ising Model; §3 Chequered Ising Model; §4 Helium Expts. Refs. *See pp. 6, 7 below.

Fig. 3. Page 2 of Fisher's January 2009 email to Helen.




Fig. 4. Page 3 of Fisher's January 2009 email to Helen.

4.

§2 First Extensions: For 2D to 1D Crossover

Exptly, strips with visible rounding are too small to observe spec. bands: So one needs many of them. But for Helium the long strips must be filled! This can be achieved by a thin coupling layer [G4, G*] That can be modeled via the

• Alternating Layered Ising Model

* The needed basic analysis is covered by [AYMcC]

The anticipated results are then just as sketched in Fig. 1 [F] and the questions are as in §1. i.e. Effectively 2 transitions & 4 distortions

§3 2nd Extension • Chequered Ising Model

to model [2D] to [0D] crossover - Basic Finite-Size Expt. But with couplings - see [G*] and "Lack of scaling in an array of boxes" [G3] *

This generalizes [FAY] & [AYF²]: Basic $(m+n') \times (n+n')$ cell.

• Simplest model just has just J & $J' = J'' = J'''$, but likely that having J' , J'' & J''' distinct is no harder.

Leading Questions are as for §1 & §2

How do the weak coupling parameters J', \dots, J'' and m', n' affect T_c & coupling?

• THIS specifies the crucial models •

Fig. 5. Page 4 of Fisher's January 2009 email to Helen.




Fig. 6. Page 5 of Fisher's January 2009 email to Helen.




Fig. 7. Page 6 of Fisher's January 2009 email to Helen.

References (contd.)

- [18] Fisher, M. E. The theory of critical point singularities. In Green, M. (ed.) *Critical Phenomenon Proc. 51st "Enrico Fermi" Summer School, Varenna, Italy* (Academic Press, NY, 1971).
- [19] Costa, R. M., Ferreira, L. M., Vieira, V. N., Pureur, P. & Schaf, J. Coherence transition in granular $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$, $\text{YBa}_2\text{Cu}_{2.95}\text{Zn}_{0.05}\text{O}_{7-\delta}$, and $\text{YBa}_{1.75}\text{Sr}_{0.25}\text{Cu}_3\text{O}_{7-\delta}$ superconductors. *European Physical Journal B* **58**, 107–113 (2007).
- [G7] [20] Mehta, S. & Gasparini, F. M. Specific heat and scaling of ^4He confined in a planar geometry. *Phys. Rev. Lett.* **78**, 2596–2599 (1997).
- [G8] [21] Kimball, M. O. & Gasparini, F. M. Superfluid fraction of ^3He - ^4He mixtures confined at $0.0483\text{ }\mu\text{m}$ between silicon wafers. *Phys. Rev. Lett.* **86**, 1558–1561 (2001).
- [22] Greywall, D. S. & Ahlers, G. Second-sound velocity, scaling, and universality in He-II under pressure near the superfluid transition. *Phys. Rev. Lett.* **28**, 1251–1254 (1972).
- [23] Kosterlitz, J. M. & Thouless, D. J. Ordering, metastability and phase transitions in two-dimensional systems. *J. Phys. C* **6**, 1181–1203 (1973).
- [24] Nelson, D. R. & Kosterlitz, J. M. Universal jump in the superfluid density of two-dimensional superfluids. *Phys. Rev. Lett.* **39**, 1201–1205 (1977).
- [25] Mamaladze, Y. G. & Cheishvili, O. D. Flow of a superfluid liquid in porous media. *Soviet Physics JETP* **23**, 112–117 (1967).
- [G9] [26] Kimball, M. O., Mooney, K. P. & Gasparini, F. M. Three-dimensional critical behavior with 2D, 1D, and 0D dimensionality crossover: Surface and edge specific heats. *Phys. Rev. Lett.* **92**, 115301 (2004).
- [G10] [27] Kimball, M. O. *Helium Confined to a Restricted Geometry – A Study of Lower Dimensionality Crossover and Universality*. Ph.D. thesis, University at Buffalo, The State University of New York (2005).

List Extracted from UNDER REVIEW article:

- [G*] Superfluid transition in granular ^4He : Coupling and proximity effects

UNPUBLISHED

Justin K. Perron,* Mark O. Kimball,† Kevin P. Mooney,‡ and Francis M. Gasparini§

Department of Physics, University at Buffalo,
the State University of New York, Buffalo, NY 14260

*Electronic address: jkperron@buffalo.edu

†Electronic address: mok2@enthalpy.physics.buffalo.edu

‡Electronic address: mooney.kp@gmail.com

§Electronic address: fmg@buffalo.edu

Fig. 8. Page 7 of Fisher's January 2009 email to Helen.