

Topological Phases with Average Symmetries: the Decohered, the Disordered, and the Intrinsic

Ruochen Ma,^{1,2,3,*} Jian-Hao Zhang,^{4,*} Zhen Bi,⁴ Meng Cheng,⁵ and Chong Wang¹

¹*Perimeter Institute for Theoretical Physics, Waterloo, Ontario, Canada N2L 2Y5*

²*Department of Physics and Astronomy, University of Waterloo, Waterloo, ON N2L 3G1, Canada*

³*Kadanoff Center for Theoretical Physics, University of Chicago, Chicago, Illinois 60637, USA*

⁴*Department of Physics, The Pennsylvania State University, University Park, Pennsylvania 16802, USA*

⁵*Department of Physics, Yale University, New Haven, Connecticut 06511-8499, USA*

Global symmetries greatly enrich the landscape of topological quantum phases, playing an essential role from topological insulators to fractional quantum Hall effect. Topological phases in mixed quantum states, originating from *decoherence* in open quantum systems or *disorders* in imperfect crystalline solids, have recently garnered significant interest. Unlike pure states, mixed quantum states can exhibit *average symmetries* – symmetries that keep the total ensemble invariant but not on each individual state. In this work, we present a systematic classification and characterization of average symmetry-protected topological (ASPT) phases applicable to generic symmetry groups, encompassing both average and exact symmetries, for bosonic and fermionic systems. Moreover, we formulate the theory of average symmetry-enriched topological (ASET) orders in disordered bosonic systems. Our systematic approach helps clarify nuanced issues in previous literature and uncovers compelling new physics. Notably, we discover that (1) the definition and classification of ASPT phases in decohered and disordered systems exhibit subtle differences; (2) despite these differences, ASPT phases in both settings can be classified and characterized under a unified framework of defect decoration and spectral sequence; (3) this systematic classification uncovers a plethora of ASPT phases that are *intrinsically mixed*, implying they can exclusively manifest in decohered or disordered systems where part of the symmetry is average; (4) similarly for ASET, we find intrinsically disordered phases exhibiting exotic anyon behaviors – the ground states of such phases necessarily contain localized anyons, with gapless (yet still localized) excitation spectral.

CONTENTS

		E. An example with Lieb-Schultz-Mattis anomaly	19
I. Introduction	1	V. Summary and outlook	20
II. Average SPT: generalities	3	Acknowledgments	21
A. Decohered ASPT	3	A. On the definition of disordered ASPT	21
1. Strong-to-weak spontaneous symmetry breaking	4	B. A brief review of spectral sequence	22
2. Symmetrically invertible states	5	C. Constructions of (2+1)d fermionic SPTs and ASPTs	22
3. Example: cluster chain and edge state	6	References	24
B. Disordered ASPT	7		
1. Interplay with localization physics	8		
C. Mathematical framework: Spectral sequence	9		
III. Intrinsic ASPT: examples	10		
A. Fixed-point model for bosonic ASPT	10		
B. Berry-free ASPT and gapless SPT	12		
C. Fermionic intrinsic ASPT	13		
IV. Average symmetry-enriched topological orders	15		
A. General structures	15		
B. Example: $\mathbb{Z}_2 \times \mathbb{Z}_2$ toric code with \mathbb{Z}_2^A symmetry	17		
C. Intrinsically disordered ASET with \mathbb{Z}_2^A symmetry	17		
D. ASET with 't Hooft anomaly: an example with $\mathbb{Z}_2 \times \mathbb{Z}_2^A$	18		

I. INTRODUCTION

Topological quantum phases are often enriched by global symmetries [1]. The most familiar example is the fractional quantum Hall effect (FQHE). Independent of any global symmetry, the FQHE ground states are characterized by intrinsic *topological orders* that support anyon excitations. Global symmetries such as U(1) charge conservation, however, can further enrich the topological orders, for example by assigning fractional charges to the anyon excitations. FQHE enriched by U(1) charge conservation is a simple example of *symmetry-enriched topological orders* (SET) [2]. Even quantum states with trivial topological orders can be enriched

* These authors contributed equally.

by global symmetries. Familiar examples include topological insulators [3, 4] and the Haldane spin-1 chain [5]. These states do not carry exotic excitations such as anyons, yet they cannot be adiabatically deformed to unentangled product states (like atomic insulators) without breaking relevant symmetries. Such states are known as *symmetry-protected topological* (SPT) states [6–12].

Traditionally the concepts of SPT and SET phases are defined for a pure quantum state $|\Psi\rangle$ – typically the ground state of some local Hamiltonian H . Our understanding of SPT and SET phases in pure states has reached a quite mature stage over the past few decades. For mixed quantum states, however, much less is known. The two relevant physical contexts to study topological phases in mixed states are systems with *decoherence* and *disorders*. In decohered systems (such as noisy intermediate-scale quantum (NISQ) simulators [13–16]), mixed states arise naturally from the coupling between the system and the environment. In a disordered system (such as an imperfect crystalline solid), the Hamiltonian is drawn from a statistically random ensemble $\{H_I\}$, where I labels disorder realizations with a certain probability P_I . Given the ubiquity of decoherence and disorder in realistic systems, it is both natural and important to try to extend our understanding of topological phases, including SPT and SET, to mixed states.

In contrast to pure states, there are two distinct types of symmetry for mixed states. An *exact symmetry* (or *strong symmetry*) is a symmetry for each individual pure state $|\Psi_I\rangle$ in the mixed state ensemble. In contrast, an *average symmetry* (or *weak symmetry*) is not necessarily a symmetry for each individual pure state, but is a symmetry for the ensemble density matrix $\rho = \sum_I P_I |\Psi_I\rangle\langle\Psi_I|$. Namely, for an average symmetry operator g , we have

$$g\rho g^{-1} = \rho. \quad (1)$$

For example, in a solid with quenched disorder, where the electrons experience a random potential, the crystalline symmetries are (at best) average symmetries, while the $U(1)$ electric charge symmetry remains an exact symmetry.

The central task of this work is to systematically examine how exact and average symmetries enrich topological phases in mixed states, in both decohered and disordered systems. Historically, it was found that some free-fermion SPT phases remain nontrivial [17–19], in the sense that the boundary states remain delocalized when part of the protecting symmetries become average due to disorders. For bosonic systems, it was found [20] that, at least in $(1+1)d$, SPT phases protected by exact (strong) symmetries are stable against decoherence [20]. More recently, it was realized [21] that the general notion of SPT phases can be extended to systems with both exact and average symmetries. These “average SPT” (ASPT) phases are characterized by nontrivial topological response, or equivalently decorated defects, involving both the exact and average symmetries. Many examples of ASPT were

studied in Refs. [21–23]. Closely related notions, such as average symmetry defect and anomaly, have also been studied recently in the field theory context [24].

In this work, we systematically classify and characterize topological phases, both SPT and SET, with average and exact symmetries. This allowed us to clarify many subtle issues from previous literature, and discover new physics that are intrinsically associated with mixed states.

First, in Sec. II we discuss the general framework to classify ASPT, in both decohered and disordered systems. Even though the physical states in both scenarios can be described by density matrices of the form $\rho = \sum_I P_I |\Psi_I\rangle\langle\Psi_I|$, in a disordered system the state is also further endowed with an ensemble of Hamiltonians $\{H_I\}$ with probability $\{P_I\}$. This makes the physics of the two scenarios quite different. In particular, the classification of ASPT phases will be different in the two scenarios. Pleasantly, despite the physical difference between decohered and disordered systems, the ASPT phases in both scenarios can be characterized and classified under the unified mathematical framework of spectral sequence. The physics behind the spectral sequence framework is the decoration of symmetry defects [10, 39], which is familiar from the classification of pure state SPT phases.

For decohered systems, we carefully examine the conditions under which the classification based on decorated symmetry defects applies. As a necessary condition, in order for the topological invariants to be well-defined, we find that not only the usual spontaneous symmetry breaking, defined by long-range order in two-point correlation function of order parameters, should be forbidden, but a more subtle form of spontaneous symmetry breaking from an exact/strong symmetry down to an average/weak symmetry can also invalidate the classification. To circumvent the problem, we introduce a notion of invertibility for SRE mixed states, generalizing a similar concept for ground states.

As an appealing consequence of our classification, we discover a plethora of ASPT phases that are *intrinsically mixed*. These are SPT phases that can only exist with average symmetries, in a decohered or disordered system, and by definition cannot exist as pure state SPT. In fact, if we try to deform an intrinsically mixed ASPT to a pure state, for example by reducing the disorder strength, the state reduces to the so-called “intrinsically gapless SPT”. We discuss many examples of intrinsically mixed ASPT, in both bosonic and fermionic systems, in Sec. III.

In Sec. IV we move on to nontrivial intrinsic topological orders in $(2+1)d$ in the presence of quenched disorders, and discuss how average (and exact) symmetries enrich the structure of topological orders. In clean systems, global symmetries can (1) permute different anyon types and (2) fractionalize on certain anyons (such as the fractional $U(1)$ charges of FQHE quasi-particles). Certain enrichment patterns come with ’t Hooft anomalies – such anomalous SET states can be viewed as the sur-

face states of certain SPT phases in one higher dimension. We show that average symmetries can also permute anyons. Furthermore an average symmetry and an exact symmetry can jointly fractionalize on anyons. However, the fractionalization of average symmetry alone becomes ill-defined on the anyons – unless the fractionalization pattern involves certain ’t Hooft anomalies.

Another interesting feature of disordered systems is that certain obstructions of symmetry enrichment, known as \mathcal{H}^3 obstruction, are lifted once the symmetries involved become average. Such intrinsically disordered ASET comes with additional features. For instance, the system will have localized anyons that lead to a gapless spectrum, yet the system still hosts short-range correlated ground states. We discuss many examples, featuring various nontrivial properties of ASETs mentioned above, in Sec. IV.

We end with a summary and outlook in Sec V. Some technical details are discussed in the Appendices.

II. AVERAGE SPT: GENERALITIES

In this section, we first give an overview of the basic notions of ASPT in two different physical scenarios, with decoherence (Sec. IIA) and disorders (Sec. IIB), in the simplest cases where the total symmetry \tilde{G} is the direct product of the average symmetry G and the exact symmetries A . In Sec. IIC we show that for more general symmetry structures, the classification of ASPT can be described using the Atiyah-Hirzebruch spectral sequence, for both the decohered and disordered scenarios (with different input data and consistency conditions). The goal of this Section is to not only review, but significantly systematize and clarify earlier discussions in Ref. [21].

A. Decohered ASPT

We now review the notion of a decohered ASPT state, which is relevant for open quantum systems.

Consider a mixed state ρ on a lattice system. We define ρ to be short-range entangled (SRE) [21, 25] if it can be prepared from a pure product state $|0\rangle$ via a finite-depth local quantum channel \mathcal{E} :

$$\rho = \mathcal{E}(|0\rangle\langle 0|). \quad (2)$$

Since the finite-depth channel can be equivalently viewed as acting a finite-depth local unitary U on the system together with a bath (ancilla) and then tracing out the bath, the short-range entangled mixed state ρ can be equivalently defined as one that can be purified to a short-range entangled state $|\Psi_0\rangle = U|0\rangle$ in an enlarged Hilbert space. Another defining feature of an SRE mixed state is that all physical operators (local or extended) have only short-ranged correlation functions.

We can now enrich the structure of SRE mixed states by considering global symmetries. Let us first consider

on-site unitary symmetries. A symmetry A is called *exact* if for any $a \in A$, $a\rho = e^{i\theta}\rho$. In other words, each individual state in the mixed ensemble is an eigenstate of A with the same eigenvalue. In contrast, a symmetry G is called *average* if for any $g \in G$, $g\rho g^\dagger = \rho$. This means that the density matrix ρ can be diagonalized in a G -symmetric basis, but possibly with different eigenvalues. Below we will consider the simplest situation in which the total symmetry $\tilde{G} = G \times A$, and defer the more general discussion to Sec. IIC.

Recall that SPT phases of pure states are classified as equivalence classes under symmetric adiabatic evolutions or symmetric finite-depth local unitaries. For mixed states, the natural analogues of finite-depth unitaries are finite-depth quantum channels. We can always view the finite-depth local channel as a finite-depth local unitary acting on an enlarged Hilbert space (system $\mathcal{H} \oplus$ bath \mathcal{H}_B):

$$\mathcal{E}(\rho) = \text{tr}_B [U(\rho \otimes |0_B\rangle\langle 0_B|)U^\dagger], \quad (3)$$

where $|0_B\rangle$ is a trivial symmetric product state in the ancillary (bath) Hilbert space. The exact and average symmetries manifest in different manners in the quantum channels [20, 21]: for average symmetries, we demand U to be composed of symmetric local gates; for exact symmetries, we further demand that states in the bath Hilbert space \mathcal{H}_B to transform trivially under the symmetries – in other words, the system and the bath do not exchange exact symmetry charges. In earlier literature on mixed states, the average and exact symmetries are also called “weak” and “strong” symmetries, respectively [20, 26–29].

We then define different *decohered* ASPT phases as equivalence classes of mixed states under symmetric finite-depth local channels. More precisely, two ASPT density matrices ρ_1 and ρ_2 are equivalent if there exist symmetric finite-depth local quantum channels \mathcal{E}_{12} and \mathcal{E}_{21} such that $\rho_2 = \mathcal{E}_{12}(\rho_1)$ and $\rho_1 = \mathcal{E}_{21}(\rho_2)$ (recall that channels, unlike unitaries, are in general not invertible). Notice that the ancillary Hilbert space \mathcal{H}_B will always transform nontrivially under time-reversal (because of the complex conjugation) and space symmetries. Therefore for the purpose of SPT classification, anti-unitary symmetries, and lattice symmetries should always be considered as average symmetries.

For bosonic systems, the decohered ASPT phases in $D < 4$ space dimensions with symmetry $\tilde{G} = A \times G$ (A being exact and G being average) are classified [21] by

$$\bigoplus_{p=0}^D \mathcal{H}^p(G, \mathcal{H}^{D+1-p}(A, U(1))). \quad (4)$$

The above formula has a decorated domain wall interpretation [10] as follows. The G -symmetric state can be obtained from a trivial spontaneous G -breaking state by proliferating domain walls (or more general defects) associated with G . On each proliferated codimension- p defect

we can decorate an SPT of the exact symmetry A in $D-p$ space dimensions, labeled by an element in the group cohomology [7] $\mathcal{H}^{D+1-p}(A, \text{U}(1))$. The decoration pattern is labeled by an element in $\mathcal{H}^p(G, \mathcal{H}^{D+1-p}(A, \text{U}(1)))$.

Recall that for pure states, with both A and G being exact symmetries, the group-cohomology classification of SPT is given by the Künneth formula

$$\mathcal{H}^{D+1}(A \times G, \text{U}(1)) = \bigoplus_{p=0}^{D+1} \mathcal{H}^p(G, \mathcal{H}^{D+1-p}(A, \text{U}(1))). \quad (5)$$

The last term in the sum $\mathcal{H}^{D+1}(G, \text{U}(1))$ is absent in the mixed state classification in Eq. (4). This is because the term $\mathcal{H}^{D+1}(G, \text{U}(1))$, classifying SPT protected purely by the average symmetry G , describes the nontrivial phase factors in the quantum superposition of different domain-wall configurations, which are irrelevant in the mixed state context. In other words, in the mixed state, the domain walls proliferate in terms of classical probability, and superposition phase factors are not well-defined (equivalently, the phase factors from the bra and ket in ρ cancel out). To be more concrete, a representative wavefunction of a clean SPT phase has the following form

$$|\Psi\rangle = \sum_{\mathcal{D}} \sqrt{p_{\mathcal{D}}} e^{i\theta_{\mathcal{D}}} |\Psi_{\mathcal{D}}\rangle |a_{\mathcal{D}}\rangle \quad (6)$$

where $|a_{\mathcal{D}}\rangle$ in the “ancilla” space describing the quantum state of defect network of G , $|\Psi_{\mathcal{D}}\rangle$ is the decorated A -symmetric invertible phase, and $e^{i\theta_{\mathcal{D}}}$ is the phase factor that encodes the information in the $\mathcal{H}^{D+1}(G, \text{U}(1))$ term. The clean SPT wavefunction is basically composed of the quantum superposition of different defect networks with decorated A -symmetric invertible phases. Once the G degrees of freedom are decohered, the relative phases of different quantum states with different configurations of defect networks are no longer relevant, and a mixed ensemble describes the decohered ASPT phase with the following density matrix

$$\rho = \sum_{\mathcal{D}} p_{\mathcal{D}} |\Psi_{\mathcal{D}}\rangle \langle \Psi_{\mathcal{D}}|. \quad (7)$$

We now discuss subtle issues missed in previous literature. As already mentioned above, the group-cohomology classification is established via decorated domain walls, which requires the condition of symmetry localization: global symmetry action applied to a subregion of the system is equivalent to an operator acting on the boundary of the region. It is not entirely clear that our definition of SRE states implies this condition. In fact, we will reexamine the notion of spontaneous symmetry breaking (SSB) in Sec. II A 1 and show that additional conditions must be supplied in order to apply the decorated domain wall classification. We will then discuss the notion of *invertibility* in the mixed state context in Sec. II A 2, followed by some simple examples in Sec. II A 3.

1. Strong-to-weak spontaneous symmetry breaking

For pure states, the notion of SPT requires the symmetry to be not spontaneously broken, in the sense that two-point correlation functions of charged operators vanish exponentially $\langle O(x)O^\dagger(y) \rangle \sim e^{-|x-y|/\xi}$. Otherwise the topological invariants (like those in Eq. (4)) become ill-defined. For mixed states, the two-point function detects spontaneous breaking of weak symmetries – examples include usual SSB in thermal states. For strong symmetry, a non-vanishing two-point function $\langle O(x)O^\dagger(y) \rangle$ of some charged operator $O(x)$ means that the strong symmetry is completely broken – it does not even survive as a weak symmetry. It is then natural to ask: is it possible for a strong symmetry to be spontaneously broken to a weak symmetry?

Recall that a weak symmetry is *explicitly* broken if the one-point function $\text{tr}(\rho O(x)) \neq 0$ for some charged operator $O(x)$. Similarly, a strong symmetry is *explicitly* broken to a weak symmetry if $\text{tr}(O(x)\rho O^\dagger(x)\rho) \neq 0$, as this would require ρ to contain states with different total charge. To define spontaneous (as opposed to explicit) symmetry breaking, we generalize from one-point to two-point functions. So we will define a state to have strong-to-weak SSB if the ordinary 2-point correlator is exponentially vanishing, while the Rényi-2 correlator is non-vanishing:

$$\frac{\text{tr}(O(x)O^\dagger(y)\rho O^\dagger(x)O(y)\rho)}{\text{tr}\rho^2} \sim O(1), \quad (8)$$

for some charged operator $O(x)$, $O(y)$ with $|x-y| \rightarrow \infty$. As an immediate sanity check, we notice that such strong-to-weak SSB is only possible for mixed state since for pure state the Rényi-2 correlator is simply the square of the ordinary correlator.

For a simple example of strong-to-weak SSB, we consider a $1d$ qubit chain $\rho = I + \prod_i X_i$ (normalization omitted) with a strong \mathbb{Z}_2 symmetry generated by $X \equiv \prod_i X_i$ (X_i being the Pauli- x operator on site i). The state is simply the maximally mixed state within the sector of \mathbb{Z}_2 charge $+1$. Now consider the \mathbb{Z}_2 charged operator $O(x) = Z_x$, one can readily verify that Eq. (8) indeed holds. In fact, if we write ρ in Z -basis, it is a mixture (convex sum) of cat (GHZ) states:

$$\rho = \sum_s (|s\rangle + X|s\rangle)(\langle s| + \langle s|X), \quad (9)$$

where s is a bit string in the Z -basis. In this basis, the Rényi-2 Z -correlator becomes very similar to the Edward-Anderson correlator used for spin-glass orders $\overline{\langle Z_x Z_y \rangle_s^2}$, where the overline means averaging over the bit strings.

Another viewpoint is to recall that for pure states, a cat state with SSB $|+\rangle \equiv |\uparrow\uparrow\uparrow \dots\rangle + |\downarrow\downarrow\downarrow \dots\rangle$ is locally indistinguishable from $|-\rangle \equiv |\uparrow\uparrow\uparrow \dots\rangle - |\downarrow\downarrow\downarrow \dots\rangle$, and any superposition of the two cat states will explicitly break the symmetry. Similarly, the two mixed states

$\rho_+ = I + X$ and $\rho_- = I - X$ are locally indistinguishable. In fact, if we trace out only one spin, both states will reduce to a maximally mixed state (the identity matrix) on the remaining qubits. So even though ρ_{\pm} are orthogonal in a very strong sense (each individual state in ρ_+ is orthogonal with each state in ρ_- as they carry different \mathbb{Z}_2 charge), from local properties, we cannot distinguish ρ_{\pm} from any convex superposition $P_+\rho_+ + P_-\rho_-$. The latter, however, explicitly breaks the strong \mathbb{Z}_2 symmetry down to a weak symmetry. Now recall that for a typical thermal ensemble, a state with fixed symmetry charge (also known as the canonical ensemble) and a state without (also known as the grand canonical ensemble) are locally indistinguishable. This leads to the conclusion that a typical thermal ensemble spontaneously breaks any strong symmetry to a weak symmetry.

Just as usual SSB is avoided when considering pure state SPT, we should also forbid strong-to-weak SSB when considering mixed state SPT. Otherwise the topological invariants such as those in Eq. (4) may become ill-defined.

2. Symmetrically invertible states

For pure states, a crucial property of SPT states is *invertibility*. Namely, even though a nontrivial SPT state $|\Psi\rangle \in \mathcal{H}$ cannot be deformed to a trivial product state using a finite-depth symmetric unitary circuit, we can find an “inverse” state $|\tilde{\Psi}\rangle$ from an auxiliary Hilbert space $\tilde{\mathcal{H}}$, such that $|\Psi\rangle \otimes |\tilde{\Psi}\rangle$ can be deformed to a trivial product state in $\mathcal{H} \otimes \tilde{\mathcal{H}}$, using a symmetric finite-depth unitary circuit. The invertibility condition provides a foundation for classifying SPT phases using invertible topological quantum field theory [30, 31]. In fact, it is often more satisfactory to demand symmetric invertibility (instead of simply SRE), as it allows us to naturally incorporate phases like the integer quantum Hall effect.

Another consequence of symmetric invertibility for pure states crucial for our purpose is the “decorated domain wall” picture. To be more concrete, let us consider a state $|\Psi\rangle$ and an on-site unitary symmetry transformation g . Denote the restriction of g to a subregion Γ by g_{Γ} . For a symmetric (i.e. no SSB) SRE pure state, we expect that

$$g_{\Gamma} |\Psi\rangle = V_{\partial\Gamma} |\Psi\rangle, \quad (10)$$

where $V_{\partial\Gamma}$ is a unitary operator supported on the boundary $\partial\Gamma$ of Γ . In 1D, $V_{\partial\Gamma}$ further decomposes into local unitaries supported at the two ends of Γ and this is equivalent to the existence of string order parameters. In general, if one assumes that $V_{\partial\Gamma}$ is a finite-depth unitary circuit, then the group-cohomology classification follows.

We now briefly overview how Eq. (10) follows from the assumption of symmetric invertibility. That is, there exists a state $|\tilde{\Psi}\rangle$ from an ancilla system and a symmetric finite-depth unitary U such that $U |\Psi\rangle \otimes |\tilde{\Psi}\rangle = |0\rangle$, where

$|0\rangle$ is a symmetric product state. Note that g is extended to act on-site on the ancilla Hilbert space as \tilde{g} . U should be *locally* symmetric, meaning that each gate in U commutes with $g \otimes \tilde{g}$. We will denote $\mathbf{g} = g \otimes \tilde{g}$. Then for the symmetry action \mathbf{g}_{Γ} restricted to a subregion Γ

$$\mathbf{g}_{\Gamma} |\Psi\rangle \otimes |\tilde{\Psi}\rangle = \mathbf{g}_{\Gamma} U^{-1} |0\rangle = \mathbf{g}_{\Gamma} U^{-1} \mathbf{g}_{\Gamma}^{-1} U |\Psi\rangle \otimes |\tilde{\Psi}\rangle. \quad (11)$$

Since U consists of gates commuting with \mathbf{g} , we must have

$$\mathbf{g}_{\Gamma} U^{-1} \mathbf{g}_{\Gamma}^{-1} U = V_{\partial\Gamma}. \quad (12)$$

Here $V_{\partial\Gamma}$ is a unitary supported on the boundary $\partial\Gamma$ of Γ . Since the on left hand side $\mathbf{g}_{\Gamma} |\Psi\rangle \otimes |\tilde{\Psi}\rangle$ still factorizes between the system and the ancilla, the same must be true on the right hand side, which implies that $V_{\partial\Gamma}$ factorizes as well and we find Eq. (10).

The notion of symmetrically invertible states can be defined similarly for mixed states: given a state ρ (in Hilbert space \mathcal{H}) and a set of strong (A) and weak (G) symmetries, we call ρ symmetrically invertible if there is a state $\tilde{\rho}$ from an auxiliary Hilbert space $\tilde{\mathcal{H}}$ with appropriate symmetry action of A and G , such that $\rho \otimes \tilde{\rho}$ is two-way connected to a pure product state through some symmetric finite-depth channels. Motivated by the study of pure state invertible phases, in our study of decohered SPT we will focus on states that are symmetrically invertible.

A peculiar feature of mixed states is that it is possible to have a state ρ that is SRE but not symmetrically invertible. In fact, a simple example of such a state is the strong-to-weak SSB state $\rho = I + \prod_i X_i$ (normalization omitted) in a 1D qubit chain with a strong \mathbb{Z}_2 symmetry generated by X (Sec. II A 1). To see that ρ is SRE, we can start from the pure product state $|X_i = +1\rangle$, introduce another copy of the spin chain (with spin operators denoted as Z', X' etc.) in the maximally mixed state $\rho' = I$ (which is SRE). We then measure each ancilla spin in Z basis: if $Z'_i = +1$, we do nothing, and if $Z'_i = -1$, we apply $Z_i Z_{i+1}$ to flip two neighboring spins in the physical system. After tracing out the ancilla spins, we are left with $\rho = I + X$.¹

To see that $\rho = I + X$ is not symmetrically invertible, suppose on the contrary that $\exists \tilde{\rho}$ such that $\mathcal{E} : \rho \otimes \tilde{\rho} \rightarrow |0\rangle\langle 0|$ for some finite-depth channel \mathcal{E} that preserves the strong symmetry $\prod_i X_i \otimes \prod_i \tilde{g}_i$, where \tilde{g}_i is the local \mathbb{Z}_2 symmetry operator acting on the auxiliary Hilbert space $\tilde{\mathcal{H}}$. We can purify this channel to a unitary U acting on a further enlarged Hilbert space (Eq. (3)) $\mathcal{H} \otimes \tilde{\mathcal{H}} \otimes \mathcal{H}'$. Recall by definition the strong symmetry remains $g = \prod_i X_i \otimes \prod_i \tilde{g}_i$ (it acts trivially on \mathcal{H}'), and U is made of local gates commuting with g . Now consider the symmetry action on a large segment of the system A spanning from x to y : $g_A \equiv \prod_{x < i < y} X_i \otimes \tilde{g}_i$. When acting

¹ We thank Yijian Zou for pointing out this construction.

on the final state $|0\rangle\langle 0|$, we have $\langle g_A \rangle_0 = 1$. Therefore $\langle U^\dagger g_A U \rangle = 1$ on the original state $\rho \otimes \tilde{\rho} \otimes |0\rangle\langle 0|$. Since U is a finite-depth circuit made of gates commuting with g , we must have $U^\dagger g_A U = O_x O_y g_A$ where O_x (O_y) is some operator localized around x (y). One can directly verify that $\langle O_x O_y g_A \rangle = 0$ for the initial state, for any choice of local $O_{x,y}$, which leads to a contradiction. Therefore, the density matrix ρ is not symmetrically invertible. We have thus shown that the absence of any form of SSB is a necessary condition for symmetric invertibility for SRE mixed states.

From the above argument, we also conclude that if a strongly \mathbb{Z}_2 -symmetric state is symmetrically invertible, then *there must exist some kind of string order parameter* $\text{tr} \rho o_x o_y \prod_{x < i < y} X_i \sim O(1)$ (now $o(x)$ is a local operator around x that only acts on the physical Hilbert space \mathcal{H}). The situation is then analogous to the pure state case as discussed under Eq. (10). We will see more concrete examples in Sec. II A 3. Similar analysis can be extended to higher dimensions with general strong symmetries, and one can conclude that for symmetrically invertible states, strong symmetry defects must be decorated with some boundary operator o_∂ , which can be further classified based on symmetry and topology. This offers a complementary point of view on the group-cohomology classification in Eq. (4).

Another virtue of focusing on symmetrically invertible states (instead of simply SRE) is that it mitigates the complication due to the non-invertibility of quantum channels. To make this point more manifest, let us assume a slightly stronger form of symmetric invertibility: we demand $\rho \otimes \rho^*$ to be two-way connected to a pure product state through symmetric finite-depth channels. In other words, we demand the complex conjugation ρ^* to be the inverse of ρ . This is motivated by the anticipation that nontrivial topological structure from quantum mechanics should manifest as some nontrivial phase factors – this intuition is true at least for invertible pure states, and may even be true for general symmetrically invertible mixed states (we are not aware of counter examples). The strong form of symmetric invertibility leads to the following appealing result: if two strongly symmetrically invertible states are one-way connected, namely $\exists \mathcal{E}_{FD, \text{symm}}^{12} : \rho_1 \rightarrow \rho_2$, then the two states are automatically two-way connected, namely $\exists \mathcal{E}_{FD, \text{symm}}^{21} : \rho_2 \rightarrow \rho_1$. The “inverse” channel can be constructed in a few simple steps: $\rho_2 \rightarrow \rho_2 \otimes \rho_1 \otimes \rho_1^* \rightarrow \rho_2 \otimes \rho_1 \otimes \rho_2^* \rightarrow \rho_1$, where we have used $(\mathcal{E}_{FD, \text{symm}}^{12})^* : \rho_1^* \rightarrow \rho_2^*$ in the second arrow, and strong symmetric invertibility $\rho_2 \otimes \rho_2^* \rightarrow |0\rangle\langle 0|$ in the last arrow.

3. Example: cluster chain and edge state

As an illustrative example, let us consider one-dimensional qubit chain with $\mathbb{Z}_2 \times \mathbb{Z}_2^{\text{ave}}$ symmetry, where the exact \mathbb{Z}_2 acts on even-sites as $\prod_{i=2n} X_i$ (i labeling the lattice sites), and the average $\mathbb{Z}_2^{\text{ave}}$ acts on odd-sites as

$\prod_{i=2n+1} X_i$. ASPT phases with this symmetry are classified by $\mathcal{H}^1(\mathbb{Z}_2, \mathcal{H}^1(\mathbb{Z}_2, \text{U}(1))) = \mathbb{Z}_2$, with one nontrivial phase. A representative density matrix of the nontrivial phase, on a closed chain with $2N$ sites, is

$$\rho_{\text{cluster}} = \frac{1}{2^N} \sum_{z_{2n+1}=\pm 1} |\Psi_{\{z_{2n+1}\}}\rangle \langle \Psi_{\{z_{2n+1}\}}|$$

$$|\Psi_{\{z_{2n+1}\}}\rangle = \bigotimes_{i=2n+1} |Z_i = z_i\rangle \bigotimes_{j=2n} |X_j = z_{j-1} z_{j+1}\rangle \quad (13)$$

Essentially, we have a classical ensemble of $\mathbb{Z}_2^{\text{ave}}$ domain-wall configurations, and at each domain wall, a nontrivial exact \mathbb{Z}_2 charge is decorated. It is not difficult to check that ρ_{cluster} is symmetrically invertible, by explicitly constructing symmetric finite-depth channels connecting $\rho_{\text{cluster}} \otimes \rho_{\text{cluster}}$ to a pure product state.

We can characterize the ASPT state using a string order parameter. One can show the following string order parameter is nontrivial for the density matrix defined in Eq. (13),

$$\langle Z_{2n-1} X_{2n} X_{2n+2} \dots X_{2m} Z_{2m+1} \rangle \xrightarrow{|m-n| \gg 1} O(1). \quad (14)$$

On the contrast, it was shown in Ref. [21] that states prepared from a pure product state with symmetric finite-depth channels cannot have the above string order parameter being $O(1)$. Furthermore, if we deform the average cluster state using a finite-depth symmetric channel, as long as we keep the state symmetrically invertible (see Sec. II A 2), the $O(1)$ string order parameter will survive – the only possible change is that the end point operator may evolve from Z_i to some other local operator charged under the weak \mathbb{Z}_2 symmetry. Therefore, the density matrix in Eq. 13 represents a nontrivial ASPT. A similarly robust order parameter is strange correlators defined in Refs. [22, 23].

We now show that the string order parameter Eq. (14) implies nontrivial edge correlations, similar to the clean SPT. Consider an open chain, say from $i = 1$ to $i = L$. Far away from the two boundaries the system should be indistinguishable from the closed cluster chain. This means that the string order parameter Eq. (14) should be $\sim O(1)$ as long as the two ends are not too close to the boundary. But if the system has the exact (strong) \mathbb{Z}_2 symmetry, $\prod_{i=2k} X_i \rho = \pm \rho$. So the string order parameter can be equivalently expressed as

$$\langle X_2 \dots X_{2n-2} Z_{2n-1} \cdot Z_{2m+1} X_{2m+2} \dots X_{2\lfloor \frac{L}{2} \rfloor} \rangle \xrightarrow{|m-n| \gg 1} O(1). \quad (15)$$

Furthermore, average symmetry requires that

$$\langle X_2 \dots X_{2n-2} Z_{2n-1} \rangle = \langle Z_{2m+1} X_{2m+2} \dots X_{2\lfloor \frac{L}{2} \rfloor} \rangle = 0. \quad (16)$$

Therefore a nontrivial edge correlation is enforced by the $\mathbb{Z}_2 \times \mathbb{Z}_2^{\text{ave}}$ and the bulk topology.²

² In Ref. [21] the same state was analyzed, and it was concluded

B. Disordered ASPT

We now review the notion of disordered ASPT, which is relevant for zero-temperature systems with disordered Hamiltonians.

We consider an ensemble of disordered Hamiltonians. For concreteness, the Hamiltonian takes the form

$$H_I = H_0 + \sum_i (v_i^I \mathcal{O}_i + h.c.), \quad (17)$$

where v_i^I is a quenched disorder potential drawn from a classical probability distribution $P[v^I]$ (I labeling a particular realization and i labeling a lattice site), \mathcal{O} is a local operator, and H_0 is the non-random part of the Hamiltonian. We require the disorder to be at most short-range correlated, namely $\overline{v_i^* v_j}$ (averaged over the classical probability $P[v^I]$) should decay exponentially with $|i - j|$.

We now consider the ensemble of ground states $\{|\Omega_I\rangle\}$ of the Hamiltonians $\{H_I\}$. We call the ensemble short-range entangled (SRE) if each $|\Omega_I\rangle$ is short-range entangled with a finite correlation length ξ_I that is upper-bounded in the entire ensemble³. Superficially, the ensemble gives a density matrix $\rho = \sum_I P_I |\Psi_I\rangle\langle\Psi_I|$ and the situation appears similar to the decohered open system. However, a crucial difference for the disordered Hamiltonian system is that the states $\{|\Psi_I\rangle\}$ form a preferred basis for the ensemble. For example, two states $|\Psi_I\rangle$ and $|\Psi_{I'}\rangle$ may have equal probability of realization, which means $|\Psi_I\rangle + |\Psi_{I'}\rangle$ is an equally good eigenstate of the density matrix ρ . But in the disordered setting the latter state has no physical meaning. This makes the disordered systems physically quite different from the decohered systems. However, as we will see later in Sec. II C, there is a unified mathematical framework for the classification of SPT phases for the two different settings.

Following the decohered case, we can now define exact and average symmetries. A symmetry A is exact if it commutes with H_I for any disorder realization I . An important difference with the decohered systems is that in disordered systems, time-reversal symmetry can be exact. A symmetry G is average if any element $g \in G$ takes a realization H_I to a different realization $H_{I'} = g H_I g^{-1}$ with $P[v^{I'}] = P[v^I]$. In other words, the disorder potential v may transform nontrivially under G , but the probability $P[v]$ is symmetric under G transforms. We call the

ensemble of states $\{|\Psi_I\rangle\}$ symmetric if both the exact and average symmetries are not spontaneously broken. To align with the discussion for decohered systems, we also demand that the entire ensemble of states $\{|\Psi_I\rangle\}$ to transform identically under the exact symmetry A . This can be viewed as a “canonical ensemble” for a disordered system – the condition is imposed for convenience and is not strictly required⁴.

We define two SRE ensembles (call them $\{H_I, |\Omega_I\rangle\}$ and $\{H'_I, |\Omega'_I\rangle\}$) to be in the same ASPT phase if $\{H_I\}$ can be continuously deformed to $\{H'_I\}$ while keeping all the conditions listed above throughout the deformation: (a) the disorder potentials remain short-range correlated; (b) the symmetries (both exact and average) are not broken explicitly or spontaneously; and (c) the ground states remain short-range entangled. We note that the conditions imposed here are slightly simpler than those originally discussed in Ref. [21], and in Appendix A we show that they are largely equivalent.

The disordered bosonic ASPT, as defined above, with symmetry $\tilde{G} = A \times G$ (A being exact and G being average) are classified (see Ref. [21] and Appendix B) by

$$\bigoplus_{p=0}^{D-1} \mathcal{H}^p(G, h_I^{D+1-p}(A)), \quad (18)$$

where $h_I^q(A)$ is the classification of invertible phases in q spacetime dimension with symmetry A (for bosonic systems at $D < 3$ it is simply the group-cohomology $\mathcal{H}^q(A, \text{U}(1))$).

Similar to Eq. (4), the classification from Eq. (18) also has a decorated domain wall interpretation: on each G -domain wall we can decorate an invertible state with symmetry A . Compared to the clean case (with exact \tilde{G}), the $p = D$ and $p = D + 1$ terms are missing. Similar to the decohered case, the $p = D + 1$ term is absent because the G -domain walls proliferate classically (probabilistically) without a superposition phase factor.

The absence of the $p = D$ term in Eq. (18), which describes decorating a zero-dimensional defect with an A charge, is more interesting. In the definition of SRE ensembles, we only demanded each ground state wavefunction $|\Omega_I\rangle$ to be short-range entangled, and made no requirement on the energy spectrum – we do not demand H_I to be gapped. This is appropriate for disordered systems – for example, even a fully localized Anderson insulator, with un-entangled product-state wavefunction, can be gapless. Once we forgo the requirement on the energy gap, zero-dimensional states with

that there was no nontrivial edge correlation, and the only feature from the open boundary was that the exact \mathbb{Z}_2 charge would fluctuate within the ensemble. What was not appreciated in Ref. [21] was that the fluctuating \mathbb{Z}_2 charge breaks the \mathbb{Z}_2 from exact to average symmetry, and since $\mathbb{Z}_2^{\text{ave}} \times \mathbb{Z}_2^{\text{ave}}$ does not have a nontrivial topological phase, the edge correlation disappears once the symmetry is lowered.

³ It may be possible to impose only a soft bound on ξ_I to allow rare-region effects. For simplicity, we will not study such rare-region effects in this work.

⁴ This condition of fixed total charge under A is necessary for decohered mixed states, otherwise there is no difference between exact and average symmetries since one can always simultaneously diagonalize the density matrix and the average symmetry operator. For disordered systems, however, the fixed-charge condition is optional, since there is the preferred basis given by the Hamiltonians and we are not allowed to freely re-diagonalize the density matrix.

different symmetry charges can now be deformed to each other by continuously tuning the Hamiltonian (a zero-dimensional state is always SRE by definition). This means that decorating zero-dimensional defects will not produce nontrivial phases. Nevertheless, certain patterns of zero-dimensional decoration will come with nontrivial consequences, with an intriguing connection to the physics of localization – we discuss this aspect in detail in Sec. II B 1.

The notion of 't Hooft anomaly can also be generalized naturally to the boundary of disordered ASPT states [21]. Specifically, if the 't Hooft anomaly (labeled by the bulk ASPT phase) is nontrivial, then a symmetric boundary must be long-range entangled with probability 1 in the thermodynamic limit. More precisely, for any finite $\xi_0 > 0$, the probability of a boundary state with correlation length $\xi \leq \xi_0$ vanishes as the system size $L \rightarrow \infty$.

1. Interplay with localization physics

We now return to zero-dimensional decorations in the context of disordered ASPT. As discussed in Sec. II B, decorating with $(0+1)d$ SPT (namely charges of the exact symmetry) does not lead to nontrivial phases. This is because a $(0+1)d$ SPT is nontrivial only if we demand an energy gap, but in disordered systems, we do not require each individual disorder realization to be gapped – we only demand the ground state to be short-range entangled (SRE). In disordered systems, Anderson localization provides a natural mechanism to have a gapless but SRE ground state.

Let us illustrate with a familiar example. Consider a lattice-free fermion system with U(1) charge conservation and lattice translation \mathbb{Z}^d symmetries, with the simplest tight-binding Hamiltonian

$$H = -t \sum_{\langle i,j \rangle} (c_j^\dagger c_i + \text{h.c.}) + \mu \sum_i c_i^\dagger c_i. \quad (19)$$

For $\mu > 2|t|$ or $\mu < -2|t|$, the system is an atomic insulator with U(1) charge per site $q = 0$ or $q = 1$, respectively. The system is metallic for intermediate μ , with a long-range entangled ground state. In fact, any symmetric state interpolating between the two different atomic insulators must be long-range entangled, as guaranteed by the Lieb-Schultz-Mattis theorem [32–34].

Now we add disorders that break the exact translation symmetry to an average symmetry:

$$H_{\text{disorder}} = - \sum_j \epsilon_j c_j^\dagger c_j. \quad (20)$$

As is well known, for sufficiently strong disorder the metallic states (with an intermediate chemical potential μ) become Anderson localized insulators. The ground states of the Anderson insulators are SRE – they are essentially product states of fermions sitting at random locations. The probability P_i for a site i to be occupied can

smoothly change from 0 to 1. This gives a smooth interpolation between the two atomic insulators with charge filling $q = 0$ and $q = 1$.

Another feature of the above localized intermediate state is that it is generally gapless, with localized excitations at arbitrarily small excitation energy in the thermodynamic limit. In order to be more precise, let us fix the total U(1) charge of the system for the entire disordered ensemble – for example, at density ν we can fix $Q = \lfloor \nu L \rfloor$ where L is the system size and $\lfloor \dots \rfloor$ is the integer part. Then the system must be gapless as long as the Hamiltonians are bounded, i.e. the distribution of ϵ_j is bounded (possibly with some rare tails). This is because for a large enough system, we can always find an excitation, e.g. moving a particle from an occupied site to an unoccupied site, that costs arbitrarily small energy. This forced gaplessness can be viewed as the remnant of the Lieb-Schultz-Mattis constraint for fractional charge filling.

In the above example, the lattice sites should be viewed as defects of the translation symmetries [35]. To generalize the above observations to general ASPT with $(0+1)d$ decorations, all we have to do is to replace the lattice sites with the average G -defects and to replace U(1) charge with general Abelian representations of the exact symmetries. The only subtlety is that we need Anderson localization for generic interacting systems – in other words, we need many-body localization (MBL) [36]. Crucially, we only need MBL for low-energy states, with vanishing energy density. Although not rigorously proven for the most general setting, it seems reasonable to assume that such low-energy MBL can be achieved in any dimension without fine-tuning [37]. With the localization assumption in mind, we conclude that

Two disordered ASPT states, with different exact symmetry charge decorations on $(0+1)d$ average symmetry defects, can be smoothly deformed to each other. If the disorder strength is bounded, the intermediate states must have localized excitations with excitation energy vanishing in the thermodynamic limit.

We emphasize that the story of localized states is only relevant for disordered systems, and does not affect the discussion on decohered topological phases. The reason is that even though each individual state in the ensemble of an Anderson insulator is a trivial product state, the disordered ensemble viewed as a density matrix $\rho = \sum_I P_I |\Psi_I\rangle\langle\Psi_I|$ spontaneously breaks the strong symmetry down to a weak symmetry, making it not *symmetrically invertible*, as defined in Sec. II A 2. Indeed, the example of $\rho = I + \prod_i X_i$ discussed in Sec. II A 2 is exactly a \mathbb{Z}_2 version of the Anderson insulator ensemble. At a concrete level, for disordered systems, an SSB from strong to weak symmetry is measured by the Edward-Anderson correlator $\langle Z_i Z_j \rangle^2$. Importantly, for the disor-

dered systems, the calculation should be done in the basis of Hamiltonian ground state $\{|\Psi_I\rangle\}$ – any other basis involving superposition of different $|\Psi_I\rangle$'s will not bear physical meaning. For decohered systems, however, any basis decomposition (purification) of the density matrix is equally legitimate. As explained in Sec. II A 1, we can choose a convenient basis to interpret the density matrix $\rho = I + \prod_i X_i$ as an ensemble of spin-glass orders, indicating a nonzero Edward-Anderson correlator.

C. Mathematical framework: Spectral sequence

In general, for given average symmetry G and exact symmetry A , the total symmetry \tilde{G} does not have to be the direct product $A \times G$. Instead, we only require A to be a normal subgroup of the full symmetry group. In the fermionic case, A contains the fermion parity \mathbb{Z}_2^f as a subgroup. G and A fit into the following short exact sequence:

$$1 \rightarrow A \rightarrow \tilde{G} \rightarrow G \rightarrow 1. \quad (21)$$

For a $D+1$ dimensional SPT, the classification via generalized cohomology theory can be understood by decorations on G domain walls/defects. Mathematically the consistency conditions for domain wall decorations are organized into an (Atiyah-Hirzebruch) spectral sequence [38, 39] (see Appendix B for a brief review), whose E_2 page is given by

$$\bigoplus_{p+q=D+1} E_2^{p,q} = \bigoplus_{p+q=D+1} \mathcal{H}^p(G, h^q(A)). \quad (22)$$

Here $h^q(A)$ is the classification of invertible phases in q spacetime dimension with symmetry A that can be decorated on G -defects. Physically, the term $\mathcal{H}^p(G, h^q(A))$ means decorating invertible states on codimension- p G defects. The decoration pattern must be such that on any G defect there is no 't Hooft anomaly, so it is possible to decorate an invertible state. For example, the condition that any codimension- $(p+1)$ defect is anomaly-free leads to the cocycle condition for the decoration on codimension- p defects, thus the group cohomology.

The exact form of $h^q(A)$ will depend on the physical context. In particular:

- For the standard (pure state, clean) SPT, $h^q(A)$ is the classification of all invertible phases in q spacetime dimension with symmetry A . Note $h^q(A)$ itself can be computed from the classification of invertible phases (without any symmetry) using the same kind of spectral sequence.
- For decohered ASPT, $h^q(A)$ is almost the classification of invertible phases in q spacetime dimension with symmetry A , except (1) $h^0(A) = 0$ for reasons explained in Sec. II A, and (2) $h^q(A)$ does not contain invertible phases that do not require the A

symmetry at all (for example the chiral E_8 state in $(2+1)d$) – this is because such invertible states can be easily trivialized or prepared by a finite-depth quantum channel [21]. Hence the classification of decohered ASPT phases in bosonic systems will be reduced to the Lyndon-Hochschild-Serre (LHS) spectral sequence that replaces $h^q(A)$ by $\mathcal{H}^q[A, U(1)]$ for $q \geq 1$.

- For disordered ASPT, $h^q(A)$ is almost the classification of invertible phases in q spacetime dimension with symmetry A , except $h^0(A) = 0$ and $h^1(A) = 0$ for reasons explained in Sec. II B.

This is not the end of the story. Importantly, not every decoration pattern given by the E_2 page can actually be realized as an ASPT state. A legitimate decoration pattern must satisfy certain consistency conditions. Mathematically, the obstructions to the consistency is given by the “differentials”:

$$d_r : E_2^{p,q} \rightarrow E_2^{p+r, q-r+1}. \quad (23)$$

d_r maps to a decorated domain wall configuration in $(D+2)$ dimension. For bosonic systems with $\tilde{G} = A \times G$ these differentials automatically vanish, so we obtain Eq. (4) and (18) as the classifications. For fermion systems or boson systems with nontrivial group extension, the differentials may not vanish and represent obstructions for certain decoration patterns. Note that explicit expressions for the differentials are available for both bosonic and fermionic systems up to three spatial dimensions [39, 40].

The physical meanings of these differentials correspond to the three consistency conditions of constructing an SPT, as

1. $r \leq D - p$: The decorated G -defects can be gapped without breaking A -symmetry.
2. $r = D - p + 1$: A -charge is preserved during a continuous deformation of the G -defect network. Since $h^1(A) = 0$ for disordered ASPT phases, this obstruction automatically vanishes for disordered ASPT.

Physically, for clean SPTs, a nontrivial d_{D-p+1} obstruction implies that A charge will change (i.e. not be conserved) when we change the G -defect configurations through some local operation. In the disordered setting, however, we change the G -defect configurations by drawing a different disorder realization from the ensemble. Then the change of A -charge no longer requires actual charge non-conservation – all we need is a localized mode, or more precisely a local conserved charge operator Q_A^{local} that changes its ground state eigenvalue as the disorder realization (which determines the G -defect configurations) changes locally. The sample-to-sample fluctuation of the local charge makes the situation similar to the localized states that interpolate between different $(0+1)d$ decorations discussed earlier in Sec. II B 1. Following the logic in

Sec. II B 1, we also conclude that the localization-enabled ASPT with $(0+1)d$ decorations has to be gapless in the thermodynamic limit, as long as the disorder strength is bounded. Given the finite density of localized states at zero energy, such localization-enabled states will also be dubbed “compressible”.

3. $r = D - p + 2$: There is no Berry phase accumulated after a closed path of continuous F -move deformations due to the single-valued property of the SPT wavefunction. Since $h^0(A) = 0$ for both decohered and disordered ASPT, this obstruction automatically vanishes in these two contexts. Physically, the G -defects only proliferate probabilistically to form ASPT phases, so there is no need to assign consistent Berry phases.

At the level of E_2 page, it might appear that there are fewer nontrivial ASPT phases than standard clean SPT phases (absence of $E_2^{p,1}$ for disordered ASPT and absence of $E_2^{p,0}$ for both disordered and decohered ASPT). However, this also means that there are fewer potential obstructions for ASPT phases since each obstruction corresponds to some nontrivial topological phase in one dimension higher. This opens the possibility of ASPT phases that are *intrinsically* disordered or decohered, in the sense that they cannot be viewed as a clean SPT perturbed by disorder or decoherence. Such “intrinsic ASPT” will be one of the main focuses of this work. We dub the disordered ASPT phases enabled by vanishing d_{D-p+1} obstructions *localization-enabled* ASPT or *compressible* ASPT, and ASPT phases (both disordered and decohered) enabled by vanishing d_{D-p+2} obstructions *Berry-free* ASPT.

The differential (23) is also called the *trivialization* map for SPT phases labeled by elements in $E_2^{p+r, q-r+1}$ with decorated domain wall configurations in one higher dimension. The images of the d_r map give trivial SPT phases. The physical meaning of the trivialization map is that the images of d_r are the states with anomalous SPT states [41] on the boundary which is SRE, and the corresponding bulk states should be topologically trivial. We emphasize that the decoherence/disorder does not affect the trivialization of the ASPT phases: the anomalous SPT states on the boundary in the clean systems might be trivial product states in the presence of decoherence or disorder, which are also SRE and manifest the topologically trivial bulk states.

III. INTRINSIC ASPT: EXAMPLES

In this Section, we discuss the “Berry-free” and “compressible” intrinsic ASPT phases in more detail. The Berry-free intrinsic ASPT phases are enabled by the vanishing of the Berry phase obstruction d_{D-p+2} and the compressible intrinsic ASPT phases are enabled by the vanishing of the charge-decoration obstruction d_{D-p+1} , as we discussed in Sec. II C. The Berry-free ASPT can appear in both decohered and disordered settings – the

only difference is that if a phase comes from a $(0+1)d$ decoration it will be trivial in the disordered setting. For Berry-free ASPT states we will mostly not distinguish the two settings in this Section. The compressible ASPT can only appear in disordered settings.

A. Fixed-point model for bosonic ASPT

We will describe a class of “fixed-point” lattice models for Berry-free ASPT phases. The model is a generalization of the group-cohomology model for bosonic SPT phases [6].

Again we denote by G the average symmetry, A the exact symmetry, and the group extension \tilde{G} as defined in Eq. (21). We denote elements of \tilde{G} by $x = (g, a)$, where $g \in G$ and $a \in A$. In $(D+1)$ -dimension, the input to the model is a (homogeneous) $(D+1)$ -cochain $\nu(x_0, \dots, x_{D+1})$, that satisfies the obstructed cocycle condition:

$$(d\nu)(x_0, \dots, x_{D+2}) = O_{D+2}(g_0, \dots, g_{D+2}). \quad (24)$$

Here d is the coboundary operator, g_i is the G -grading of x_i . O_{D+2} is a $(D+2)$ -cocycle in $\mathcal{H}^{D+2}[G, \text{U}(1)]$. To construct a clean SPT state, we will need $O_{D+2} = 1$. For the ASPT construction, this is not necessary.

We will illustrate the construction in $(2+1)d$ in the following, but the same construction works in any dimension. We will work with a triangular lattice. The system consists of a \tilde{G} spin on each site, with an orthonormal basis labeled by group elements, i.e. $\{|x\rangle\}_{x \in \tilde{G}}$. A natural \tilde{G} symmetry action is given by the left multiplication:

$$U_y |x\rangle = |yx\rangle, y \in \tilde{G}. \quad (25)$$

In addition, the lattice has to be equipped with a branching structure, which is essentially an ordering of all sites. For each triangle face Δ of the lattice, denote by i, j, k the three vertices whose ordering satisfies $i < j < k$. We also denote by $s(\Delta) = \pm 1$ the orientation, i.e. whether i, j, k is clockwise or counter-clockwise.

Firstly, we review the standard group cohomology construction when $O_4 = 1$. Define the \tilde{G} -invariant state on the site j :

$$|0_j^{\tilde{G}}\rangle = \frac{1}{\sqrt{|\tilde{G}|}} \sum_{x \in \tilde{G}} |x_j\rangle, \quad (26)$$

and then the trivial paramagnetic state for the whole system:

$$|\Psi_0^{\tilde{G}}\rangle = \bigotimes_j |0_j^{\tilde{G}}\rangle. \quad (27)$$

$|\Psi_0^{\tilde{G}}\rangle$ is the ground state of the following local Hamiltonian:

$$H_{\text{trivial}} = - \sum_j |0_j^{\tilde{G}}\rangle \langle 0_j^{\tilde{G}}|. \quad (28)$$




FIG. 1. Quantum circuit as the entangler of the ASPT density matrix ρ_{ASPT} , from a trivial density matrix ρ_0 . $O(1)$ depicts the finite-depth nature of $V(\{g\})$.

Let us now define the following finite-depth local unitary circuit

$$V = \sum_{\{x\}} \prod_{\Delta_{ijk}} \nu^{s(\Delta_{ijk})}(1, x_i, x_j, x_k) |\{x\}\rangle \langle\{x\}|. \quad (29)$$

V can be viewed as a composition of unitary gates each acting on a triangle. Since all the gates are diagonal in the $|\{x\}\rangle$ basis, they commute with each other and as a circuit V has depth 1. When $O_4 = 1$, one can show that the local gates do not preserve \tilde{G} individually, but the unitary V as a whole does.

The SPT state is given by

$$|\Psi_{\text{SPT}}\rangle = V |\Psi_0^{\tilde{G}}\rangle. \quad (30)$$

The commuting-projector parent Hamiltonian for this state is

$$H = V H_{\text{trivial}} V^\dagger = - \sum_j B_j, \quad (31)$$

$$B_j = V |0_j^{\tilde{G}}\rangle \langle 0_j^{\tilde{G}}| V^\dagger.$$

Here B_j is an operator that acts on the hexagon centered at j .

Now we consider what goes wrong if the cocycle ν is obstructed by a nontrivial $[O_4]$. We find that the state $|\Psi_{\text{SPT}}\rangle$ is no longer symmetric under \tilde{G} . More precisely, it is no longer invariant under the symmetry transformations U_x when $x \in \tilde{G}$ has a nontrivial G -grading: $U_x |\Psi_{\text{SPT}}\rangle \neq e^{i\varphi} |\Psi_{\text{SPT}}\rangle$. However, for $a \in A$, the state is still invariant: $U_a |\Psi_{\text{SPT}}\rangle = |\Psi_{\text{SPT}}\rangle$.

Based on this observation, we now show how to construct the intrinsic ASPT phase. It is now more convenient to think of the \tilde{G} spin as a G spin and an A spin, and treat the G spins as quenched disorder configurations. The configuration of G spins will be collectively denoted as $\{g\}$. In addition, for $h \in G$ under U_h the classical G spin configuration $\{g\}$ transforms to $\{hg\}$.

Instead of $|0_j^{\tilde{G}}\rangle$, we define a trivial paramagnet for the

A spins:

$$|0^A\rangle = \bigotimes_j \left(\frac{1}{\sqrt{|A|}} \sum_{a \in A} |a_j\rangle \right), \quad (32)$$

and then a trivial G spin ensemble:

$$\rho_0 = \frac{1}{|G|^{N_v}} \sum_{\{g\}} |0^A\rangle \langle 0^A| \otimes |\{g\}\rangle \langle\{g\}|, \quad (33)$$

where N_v is the number of vertices of the lattice. This state can be prepared by measuring G spins in the $|0^{\tilde{G}}\rangle$ state (but no post selection). It is easy to verify that the density matrix ρ_0 does not have an exact G symmetry, but still invariant under the average G symmetry.

The follow-up finite-depth unitary circuit $V(\{g\})$ can still be defined in the same way, but now it is viewed as an operator that acts on the A spins conditioned on the G spins. More explicitly:

$$V(\{g\}) = \sum_{\{a\}} \prod_{\Delta_{ijk}} \nu^{s(\Delta_{ijk})}(1, x_i, x_j, x_k) |\{a\}\rangle \langle\{a\}|. \quad (34)$$

We denote it as $V(\{g\})$ to emphasize the G spin dependence. Importantly, even though ν is not a 3-cocycle of the group \tilde{G} , by definition it is a 3-cocycle of A and therefore $V(\{g\})$ is a (globally) A -symmetric finite-depth circuit. On the other hand, under $h \in G$ the unitary transforms as

$$U_h V(\{g\}) U_h^\dagger = e^{i\phi(h;\{g\})} V(\{hg\}). \quad (35)$$

Fixing $\{g\}$, now we can define an A -SPT state:

$$|\Psi_{\text{ASPT}}(\{g\})\rangle = V(\{g\}) |0^A\rangle. \quad (36)$$

It then follows that

$$U_h |\Psi_{\text{ASPT}}(\{g\})\rangle = e^{i\phi(h;\{g\})} |\Psi_{\text{ASPT}}(\{hg\})\rangle. \quad (37)$$

Here $e^{i\phi(h;\{g\})}$ is a phase factor that can be expressed as a product over O_4 , but the exact expression is not important to us – the factors from the bra and ket will cancel out in the density matrix. The parent Hamiltonian of a state $|\Psi(\{g\})\rangle$ in the ensemble is given by

$$H(\{g\}) = -V(\{g\}) \left(\sum_j |0_j^A\rangle \langle 0_j^A| \right) V^\dagger(\{g\}). \quad (38)$$

The collection of states $|\Psi(\{g\})\rangle$ for all the G spins forms a statistical ensemble. More formally, we can write it as the following density matrix:

$$\rho_{\text{ASPT}} = \sum_{\{g\}} p(\{g\}) |\Psi_{\text{ASPT}}(\{g\})\rangle \langle \Psi_{\text{ASPT}}(\{g\})|. \quad (39)$$

Here $p(\{g\})$ is a probability distribution of G spins. In the simplest case, we can simply set p to be a constant independent of $\{g\}$. The density matrix evidently has G average symmetry, but the A symmetry remains exact.

The finite-depth quantum circuit as the entangler of an ASPT density matrix is illustrated in Fig. 1.

Lastly, we discuss two concrete examples. The first example is in $(1+1)d$, with $G = \mathbb{Z}_2$, $A = \mathbb{Z}_2$ and $\tilde{G} = \mathbb{Z}_4$. We will go through the construction of this example in Sec. III B.

The next example is in $(2+1)d$, with $G = \text{SO}(5)$, $A = \mathbb{Z}_2$ and $\tilde{G} = \text{Spin}(5)$. Each state in the ensemble is a Levin-Gu SPT state [42] protected by the exact \mathbb{Z}_2 symmetry. To see why G must be an average symmetry, note that otherwise, we can gauge A to find a double semion topological order with four anyons $\{1, s, s', b\}$. It is enriched by the $\text{SO}(5)$ symmetry, where b transforms as the spinor representation of $\text{SO}(5)$ (required by the group extension). Because $b = s \times s'$, one of s or s' must also transform as a spinor representation, and we will assume it is s . Then s' transforms linearly under $\text{SO}(5)$. In other words, we have effectively a semion topological order $\{1, s\}$ with s being a $\text{SO}(5)$ spinor. This SET is known to have a nontrivial $\text{SO}(5)$ 't Hooft anomaly [43]. Therefore, the original SPT state can not exist with G being an exact symmetry. However, since the only obstruction is the 't Hooft anomaly of G , once G becomes an average symmetry the obstruction no longer matters.

B. Berry-free ASPT and gapless SPT

We now discuss the relationship between the “Berry-free” intrinsic ASPT phases and the recently discussed intrinsically gapless SPT (igSPT) phases [44–46]. Firstly, let us review the physics of igSPTs in $(1+1)d$ [44]. They can be constructed using a “slab”, where the top boundary is an “anomalous” gapped \tilde{G} SPT state, obstructed only by a differential mapped into $[\omega] \in \mathcal{H}^3(G, \text{U}(1))$. The bottom boundary instead has a gapless theory, e.g. a conformal field theory (CFT), where the G symmetry acts faithfully in the low-energy theory with a 't Hooft anomaly given by $[\omega^{-1}]$, and the A symmetry does not act. Together the whole slab is free of any anomaly and can be realized with a non-anomalous \tilde{G} symmetry.

Starting from an igSPT state, we consider turning on a random G symmetry-breaking perturbation. Without loss of generality, we assume that the random perturbation is relevant, so it drives the gapless theory into a disordered SRE ensemble. The result is expected to be an intrinsically ASPT state. In the other direction, any symmetry-preserving clean limit of a Berry-free intrinsic ASPT state must be an igSPT state.

Let us consider an example, with $A = \mathbb{Z}_2$, $G = \mathbb{Z}_2$ and the extension is $\tilde{G} = \mathbb{Z}_4$. For the clean system, there is no nontrivial gapped SPT phase because $\mathcal{H}^2(\mathbb{Z}_4, \text{U}(1)) = \mathbb{Z}_1$. If we look closer, there is a nontrivial $E_2^{1,1}$ term in the LHS spectral sequence, which is however obstructed by a nontrivial d_2 differential into $\mathcal{H}^3(\mathbb{Z}_2, \text{U}(1))$. The same anomaly is realized by a $(1+1)d$ free boson CFT, and together we can construct an igSPT state. We now describe a solvable lattice model for this state. A similar

model was studied in [47].

The Hilbert space of the model consists of Ising spins σ on the sites and τ on links. The symmetries are defined as

$$U_g = \prod_j \sigma_j^x e^{i\frac{\pi}{4} \sum_j (1 - \tau_{j+1/2}^x)}, \quad U_a = \prod_j \tau_{j+1/2}^x. \quad (40)$$

Here g/a is the generator of G/A . The unitaries are on-site, and satisfy $U_a^2 = 1$, $U_g^2 = U_a$. So this is a non-anomalous \mathbb{Z}_4 symmetry.

We define a projector:

$$P = \prod_j P_j, \quad P_j = \frac{1 + \sigma_j^z \tau_{j+1/2}^x \sigma_{j+1}^z}{2}. \quad (41)$$

Physically, in the subspace $P = 1$ an Ising domain wall $\sigma_j^z \sigma_{j+1}^z = -1$ is decorated by a charge $\tau_{j+1/2}^x = -1$. So $P = 1$ enforces domain wall decoration corresponding to the nontrivial element in $\mathcal{H}^1(G, \mathcal{H}^1(A, \text{U}(1)))$. In this subspace, U_g takes the following form:

$$U_g = \prod_j \sigma_j^x e^{i\frac{\pi}{4} \sum_j (1 - \sigma_j^z \sigma_{j+1}^z)}, \quad (42)$$

which takes the form of the anomalous \mathbb{Z}_2 symmetry of the Levin-Gu edge model [42, 48]. It is also easy to see that U_a becomes the identity in this low-energy subspace, at least in the bulk of the spin chain. Define

$$\tilde{\sigma}_j^x = \tau_{j-1/2}^x \sigma_j^x \tau_{j+1/2}^x, \quad \tilde{\sigma}_j^z = \sigma_j^z. \quad (43)$$

$\tilde{\sigma}_j^x$ and $\tilde{\sigma}_j^z$ generate the entire algebra of operators that commute with P . They satisfy the usual commutation relations of Pauli operators.

Now we consider the following Hamiltonian:

$$H_{\text{LG}} = - \sum_j \tilde{\sigma}_j^x (1 - \tilde{\sigma}_{j-1}^z \tilde{\sigma}_{j+1}^z) P. \quad (44)$$

Notice that since the Hamiltonian is written in terms of the $\tilde{\sigma}^{x,z}$ operator it commutes with P . In the low-energy space, Eq. (44) is identical to the edge Hamiltonian of the Levin-Gu model [42]. In addition, the Hamiltonian conserves the number of Ising domain walls, and thus also conserves the \mathbb{Z}_4 charge. The low-energy effective theory of this model is a $c = 1$ free boson, or a Luttinger liquid, with an anomalous U_g symmetry transformation. Thus the model realizes an igSPT phase.

To obtain an ASPT phase, we can proceed in two ways. First, we add some random Ising disorder $-\sum_j h_j \sigma_j^z$ with $h_j = \pm 1$ to break $G = \mathbb{Z}_2$ symmetry. It can be shown that this disorder is a relevant perturbation to the Luttinger liquid. In the strong disorder limit, we can ignore H_{LG} because it does not commute with the random disorder we have added. The Hamiltonian $H_{\mathcal{D}}$ and ground-state wavefunction $|\Psi_{\mathcal{D}}\rangle$ for a specific disorder realization $\{h_j\}$ is

$$H_{\mathcal{D}} = \sum_j P_j + h_j \sigma_j^z \quad (45)$$

$$|\Psi_{\mathcal{D}}\rangle = \bigotimes_j |\sigma_j^z = h_j\rangle \otimes |\tau_{j+1/2}^x = h_j h_{j+1}\rangle.$$

Thus we obtain a disorder ensemble $\{|\Psi_D\rangle\}$.

Alternatively, we construct the following mixed state:

$$\rho = \sum_{\mathcal{D}} p_{\mathcal{D}} |\Psi_{\mathcal{D}}\rangle \langle \Psi_{\mathcal{D}}|, \quad (46)$$

where $p_{\mathcal{D}}$ is the classical probability distribution of the disorder realizations. We now show explicitly that starting from the igSPT (pure) state, one can apply a finite-depth quantum channel to obtain ρ . Denote the density matrix of the igSPT state by ρ_{igSPT} . First we apply the following quantum channel \mathcal{E}^z :

$$\mathcal{E}^z = \mathcal{E}_1^z \circ \mathcal{E}_2^z \circ \dots, \quad \mathcal{E}_j^z[\rho] = \frac{\rho + \sigma_j^z \rho \sigma_j^z}{2}. \quad (47)$$

Notice that this quantum channel only preserves U_g on average, but preserves U_a exactly. After this step $\mathcal{E}^z[\rho_{\text{igSPT}}]$ already takes the form given in Eq. (46), but the probability distribution $p_{\mathcal{D}}$ is long-ranged. In fact, the correlation function of σ^z is the same as that in the pure state.

Then we apply another quantum channel \mathcal{E}^x :

$$\mathcal{E}^x = \mathcal{E}_1^x \circ \mathcal{E}_2^x \circ \dots, \quad \mathcal{E}_j^x[\rho] = \frac{\rho + \tilde{\sigma}_j^x \rho \tilde{\sigma}_j^x}{2}. \quad (48)$$

This channel preserves both U_g and U_a exactly. It is straightforward to check that $(\mathcal{E}^z \circ \mathcal{E}^x)[\rho_{\text{igSPT}}]$ gives a decohered SPT state with $p_{\mathcal{D}} \propto 1$. Notice that this does not imply that the igSPT and the decohered ASPT are in the same phase, as there is no finite-depth local quantum channel that takes ρ (which has only short-ranged correlation functions) to the igSPT (which has power-law correlation functions).

C. Fermionic intrinsic ASPT

We now turn to the fermionic case. For simplicity, we will assume that the “bosonic” symmetry group G becomes average, while the fermion parity conservation \mathbb{Z}_2^f remains an exact symmetry. In the decohered case, it means that the bath is bosonic, so the coupling between the system and the environment preserves fermion parity of the system. The total symmetry group \tilde{G} is a central extension of G by \mathbb{Z}_2^f . We further assume that G is a finite group to simplify the discussion.

The relevant groups of fermionic invertible phases that can be decorated on G -defects, up to $(3+1)d$, are

$$h^0 = \text{U}(1), h^1 = \mathbb{Z}_2, h^2 = \mathbb{Z}_2, h^3 = \mathbb{Z}, h^4 = 0, \quad (49)$$

where $h^1 = \mathbb{Z}_2$ is generated by a complex fermion, $h^2 = \mathbb{Z}_2$ is generated by the Majorana chain, and $h^3 = \mathbb{Z}$ is generated by the $p + ip$ superconductor.

Following the prescription in Sec. II C, for disordered phases we set $h^0 = h^1 = 0$. Thus for disordered ASPT states, the relevant groups are

$$h^0 = 0, h^1 = 0, h^2 = \mathbb{Z}_2, h^3 = \mathbb{Z}, h^4 = 0. \quad (50)$$




FIG. 2. $(3+1)d$ intrinsically decohered fermionic ASPT state from decorating a Majorana chain on the junction of \mathbb{Z}_4 (red) and \mathbb{Z}_2 (blue) domain walls.

For decohered ASPT, the groups become

$$h^0 = 0, h^1 = \mathbb{Z}_2, h^2 = \mathbb{Z}_2, h^3 = \mathbb{Z}_{16}, h^4 = 0. \quad (51)$$

The change of h^3 is because 16 copies of $p + ip$ superconductors are adiabatically equivalent to a E_8 state, which becomes trivialized for decohered phases (assuming the bath to be bosonic).

We now study the classifications in more details. Suppose the spatial dimension is $D \leq 3$. Let us consider the following two terms on the E_2 page: $\mathcal{H}^{D-1}(G, h^2)$ and $\mathcal{H}^D(G, h^1)$, corresponding to decorations of 1-dimensional G defect junctions by Majorana chains, and 0-dimensional G junctions by complex fermions. The potential differentials are

$$d_2 : \mathcal{H}^{D-1}(G, \mathbb{Z}_2) \rightarrow \mathcal{H}^{D+1}(G, h^1), \quad (52)$$

$$d_2 : \mathcal{H}^D(G, \mathbb{Z}_2) \rightarrow \mathcal{H}^{D+2}(G, h^0), \quad (53)$$

and

$$d_3 : \mathcal{H}^{D-1}(G, \mathbb{Z}_2) \rightarrow \mathcal{H}^{D+2}(G, h^0). \quad (54)$$

Explicit expressions for the differentials can be found in [40, 49].

First consider disordered ASPT phases, where $h^0 = h^1 = 0$ and $h^2 = \mathbb{Z}_2$. For the $\mathcal{H}^{D-1}(G, \mathbb{Z}_2)$ part, both d_2 and d_3 automatically vanish. Thus we conclude that any element of $\mathcal{H}^{D-1}(G, \mathbb{Z}_2)$ gives a disordered ASPT phase. Let us list a few examples of intrinsically disordered ASPT phases:

1. $D = 1, G = \mathbb{Z}_2, \tilde{G} = \mathbb{Z}_4^f$. This example is a Majorana chain with \mathbb{Z}_4^f symmetry (i.e. a charge-4e superconductor in the clean limit), which has a d_2 obstruction in the clean case. Below we will describe a concrete model realization of this state in a 1D Kitaev chain with random pairing.
2. $D = 2, G = \mathbb{Z}_2^T, \tilde{G} = \mathbb{Z}_2^T \times \mathbb{Z}_2^f$. Here \mathbb{Z}_2^T is the time-reversal symmetry. The Majorana decoration is classified by $\mathcal{H}^1(\mathbb{Z}_2^T, \mathbb{Z}_2) = \mathbb{Z}_2$, and the nontrivial class is obstructed by d_2 in the clean case. Note that if $\tilde{G} = \mathbb{Z}_4^{Tf}$, then the d_2 obstruction vanishes and the result is the well-known class DIII topological superconductor in 2D (see Appendix C for more details).

3. $D = 3, G = \mathbb{Z}_2, \tilde{G} = \mathbb{Z}_2 \times \mathbb{Z}_2^f$. The Majorana decoration is classified by $\mathcal{H}^2(\mathbb{Z}_2, \mathbb{Z}_2) = \mathbb{Z}_2$. The nontrivial class is obstructed by d_2 in the clean case.
4. $D = 3, G = \mathbb{Z}_2 \times \mathbb{Z}_4, \tilde{G} = \mathbb{Z}_2 \times \mathbb{Z}_4 \times \mathbb{Z}_2^f$. The Majorana decoration is classified by $\mathcal{H}^2(\mathbb{Z}_2 \times \mathbb{Z}_4, \mathbb{Z}_2) = \mathbb{Z}_2^3$. Interestingly, one of them is only obstructed by d_3 in the clean case. More explicitly, denote group elements of $\mathbb{Z}_2 \times \mathbb{Z}_4$ by (a_1, a_2) , where $a_1 = 0, 1$ and $a_2 = 0, 1, 2, 3$. The nontrivial cocycle in $\mathcal{H}^2(G, \mathbb{Z}_2)$ that describes Majorana chain decoration on the junction of the \mathbb{Z}_4 and \mathbb{Z}_2 domain walls (see Fig. 2) is given by

$$n_2(a, b) = a_1 b_2 \pmod{2}. \quad (55)$$

We explicitly check that the d_2 obstruction vanishes and d_3 is nontrivial.

More examples of obstructed fermionic phases can be found in Ref. [50].

For decohered ASPTs, the differentials of the cases with Majorana-chain decoration have been analyzed in the disordered case, and the only difference is that $h^1 = \mathbb{Z}_2$ for decohered ASPTs, so the d_2 differential needs to vanish to ensure the fermion parity conservation. An example of such intrinsically decohered fermionic ASPT is given by Eq. (55) for $G = \mathbb{Z}_4 \times \mathbb{Z}_2$.

For the \mathcal{H}^D part (complex fermion decoration), the d_2 differential always vanishes. An example of a “Berry-free” ASPT phase with complex fermion decoration is $G = \mathbb{Z}_2, \tilde{G} = \mathbb{Z}_2 \times \mathbb{Z}_2^f$ in $D = 3$. The fermionic SPT phase corresponding to the nontrivial element in $\mathcal{H}^3(G, \mathbb{Z}_2) = \mathbb{Z}_2$ is obstructed by d_2 in the clean case, and can now be realized an intrinsically decohered ASPT phase.

Exactly solvable lattice models of fermionic ASPT phases (without $p + ip$ decorations) can be constructed following [40]. We outline the construction in Appendix B for $D = 2$. Below we describe a simple model realization of Majorana chain with an average \mathbb{Z}_4^f symmetry, an example of localization-enabled compressible ASPT phase.

First let us consider a chain of spinless fermions, with the following Hamiltonian:

$$H = - \sum_j (c_j^\dagger c_{j+1} + \text{h.c.}) + \sum_j \Delta_j (c_j c_{j+1} + \text{h.c.}). \quad (56)$$

The \mathbb{Z}_4^f symmetry is generated by $g : c_j \rightarrow ic_j$. The pairing $\Delta_j \rightarrow -\Delta_j$ under the g symmetry. When Δ_j is uniform and nonzero, this is the well-known Hamiltonian of a Kitaev chain, explicitly breaking the \mathbb{Z}_4^f symmetry. When \mathbb{Z}_4^f is exact, we must have $\Delta = 0$ and the ground state is a gapless metal⁵. When \mathbb{Z}_4^f is an average symme-




FIG. 3. Illustration of a random Kitaev chain with average \mathbb{Z}_4^f symmetry. (a) A typical Majorana bond configuration. (b) A uniform bond configuration. (c) Two nearby domain walls on top of the uniform configuration. The total fermion parity of (b) and (c) differ by (-1) .

try, we can turn on a random pairing term with symmetric probability: $P[\Delta_j] = P[-\Delta_j]$. This random pairing term will localize the metallic ground state, resulting in a random Kitaev chain.

In the Hamiltonian Eq. (56), if the configuration Δ_j contains one sign-changing domain wall, we find it harbors a localized complex fermion zero mode. Thus when the sign of the pairing potential is disordered, we expect that there are low-energy states filling the superconducting gap, which we confirm numerically. However, this zero mode is protected by the time reversal symmetry (i.e. the complex conjugation).⁶ In order to obtain a localized state, we lift the local degeneracy from the zero modes by having complex hopping or the pairing terms that break the time-reversal symmetry.

To better appreciate the nature of the the random Kitaev chain with average \mathbb{Z}_4^f symmetry, it is more illuminating to consider the following “fixed point” model. On each lattice site j we have a complex fermion, which we write as two Majoranas $c_j = \frac{1}{2}(\gamma_{A,j} - i\gamma_{B,j})$. The random Hamiltonian takes the form

$$H = - \sum_j \sum_{r_j=A,B} i g_j^{r_j, r_{j+1}} \gamma_{r_j, j} \gamma_{r_{j+1}, j+1}, \quad (57)$$

where g is a random coupling constant taking value in $\{0, \pm 1\}$ such that every lattice link $(j, j+1)$ is covered by exactly one Majorana bond, which makes each state

⁵ There are other terms that can gap out the metal without breaking \mathbb{Z}_4^f , such as a translation-breaking potential. However, to keep the system in a nontrivial Kitaev chain, we need the amplitude of such potential μ to be smaller than Δ . As we recover the exact \mathbb{Z}_4^f symmetry by taking $\Delta \rightarrow 0$, we need to take $\mu \rightarrow 0$ first.

⁶ In fact, $\Delta > 0$ and $\Delta < 0$ belong to distinct topological phases labeled by $\nu = \pm 1$ where $\nu \in \mathbb{Z}_8$ is the topological invariant for topological superconductors with $T^2 = 1$ time-reversal symmetry (the BDI class), so there must be a protected zero mode at the interface.

in the ensemble a nontrivial Kitaev chain. For example, if $g_j^{AB} = 1$, then we must have $g_j^{AA} = g_j^{BB} = g_j^{BA} = 0$, and $g_{j+1}^{BA} = g_{j+1}^{BB} = g_{j-1}^{AA} = g_{j-1}^{BA} = 0$. An example of such random Majorana bond configuration is shown in Fig. 3 (a). Besides this nearest-neighbor constraints, the random coupling g should be uncorrelated in long distance, generated by a probability functional $P[g]$.

Now we examine the condition on $P[g]$ imposed by the average \mathbb{Z}_4^f symmetry, which is generated by $U = \exp(\frac{\pi}{4} \sum_j \gamma_{j,A} \gamma_{j,B})$. Some simple algebra shows that

$$\begin{aligned} U(i\gamma_j^A \gamma_{j+1}^B)U^\dagger &= -i\gamma_j^B \gamma_{j+1}^A \\ U(i\gamma_j^A \gamma_{j+1}^A)U^\dagger &= i\gamma_j^B \gamma_{j+1}^B. \end{aligned} \quad (58)$$

The minus sign above will be important. Now for $P[g]$, we should have

$$P(g^{AB}) = P(-g^{BA}), \quad P(g^{AA}) = P(g^{BB}). \quad (59)$$

We can now further simplify the model by having $g^{AB}, g^{AA}, g^{BB} \in \{0, 1\}$ and $g^{BA} \in \{0, -1\}$, generated by a \mathbb{Z}_4^f -symmetric probability functional $P[g]$.

Now what do all these mean for the total fermion parity $i^L \prod_{j=1}^L \gamma_{j,A} \gamma_{j,B}$? If we have only AB bonds or BA bonds, the total fermion parity on a ring is fixed (-1 for periodic boundary condition). But when we have domain walls between the two bonding patterns, as required by the average \mathbb{Z}_4^f symmetry, the fermion parity starts to change locally. The most illuminating case is when two domain walls are right next to each other, as shown in Fig. 3 (c). In this configuration all links $(j, j+1)$ with $j < 1$ and $j > 3$ are AB bonds, while the intermediate bonds are given by $g_1^{AA} = 1$, $g_2^{BA} = -1$ and $g_3^{BB} = 1$. A simple calculation shows that, compared with the configuration with no domain wall at all ($g_j^{AB} = 1$ for all j , as in Fig. 3 (b)), this configuration has an additional (-1) fermion parity. This means that even though the domain wall behave like an \mathbb{Z}_2 object (the bonding configurations return to the original ones after passing through two domain walls), “fusing” two nearby domain walls together will change the fermion parity. As a result, in the clean limit the domain walls cannot condense to recover the exact \mathbb{Z}_4^f symmetry. This is nothing but the manifestation of the d_2 obstruction. In general, the fermion parity of the state depends on the configurations of the g ’s and fluctuate randomly within the ensemble. In a disordered system the nontrivial domain-wall fusion does not lead to any obstruction for short-range entanglement, since the domain walls are pinned by the disorders. The above discussion eventually leads to a long-distance picture, which contains localized fermions (that carry nontrivial fermion parity) randomly located at the \mathbb{Z}_4^f -domain walls. If we take the absolute ground state of each Hamiltonian realization, different states in the ensemble will have different total fermion parity. If we take a “canonical ensemble” and fix the total fermion parity for the entire ensemble, then half of the ensemble will be put in excited states (to match the fermion parity), and the excitation spectral

above such states will in general be gapless (assuming a bounded distribution of g_j).

IV. AVERAGE SYMMETRY-ENRICHED TOPOLOGICAL ORDERS

A. General structures

In this section we consider disordered topologically ordered phases with average symmetry in $(2+1)d$. We will assume that the system is bosonic. In general, bosonic topological order in the ground state of a gapped Hamiltonian is described by a mathematical structure called unitary modular tensor category (UMTC) [51], or the anyon theory, denoted by \mathcal{C} . Physically, \mathcal{C} consists of a set of topological charges (i.e. anyon types), as well as consistency data specifying their fusion and braiding. When the anyon theory is nontrivial, the state is said to be long-range entangled (LRE).

To analyze LRE phases with disorder, let us first define the notion of LRE ensembles, which is a straightforward extension of SRE ensembles. To formulate the definition it is convenient to choose a “reference state”, which can be the ground state of a gapped Hamiltonian in the clean system, and identify its topological order described by the anyon theory \mathcal{C} . We require that all states in the ensemble are smoothly connected to the reference state, thus described by the same topological order \mathcal{C} . Such an ensemble is said to be LRE with topological order \mathcal{C} . Notice that this definition is meaningful only in disordered systems, where the disorder realizations naturally provide a preferred basis for the density matrix. For a general mixed state which has no preferred basis, the definition of topological order is much more subtle and is beyond the scope of this work.

Just like SRE ensembles, LRE ensembles can be enriched by exact and average symmetries. To characterize such average SET orders, we begin with a review of the classification of SET phases in clean systems, and then indicate how it should be modified for ensembles.

Let us consider the ground state of a gapped Hamiltonian with a global symmetry \tilde{G} , whose topological order is described by a UMTC \mathcal{C} . \tilde{G} can enrich the topological order in three ways [2, 52, 53]:

1. There is a group homomorphism from \tilde{G} to the group of auto-equivalence maps $\text{Aut}(\mathcal{C})$ of \mathcal{C} ,

$$\rho : \tilde{G} \rightarrow \text{Aut}(\mathcal{C}). \quad (60)$$

Here $\text{Aut}(\mathcal{C})$ consists of all the permutations of anyon types which keep the fusion and braiding properties invariant⁷. Basically, ρ tells us how \tilde{G} permutes anyons.

⁷ Note that more precisely, $\text{Aut}(\mathcal{C})$ is the group of braided tensor

2. The anyons may carry fractionalized quantum numbers under \tilde{G} . In particular, given ρ , there is a possible obstruction to symmetry fractionalization, which is an element in $\mathcal{H}_\rho^3(\tilde{G}, \mathcal{A})$, where \mathcal{A} is the group of Abelian anyons. When the \mathcal{H}^3 class vanishes, distinct symmetry fractionalization classes form a torsor over $\mathcal{H}_\rho^2(\tilde{G}, \mathcal{A})$.
3. Once ρ and the symmetry fractionalization of anyons are known, we then need to specify the fusion and braiding properties of \tilde{G} symmetry defects. In particular, given ρ and the symmetry fractionalization of anyons, the global symmetry may have a 't Hooft anomaly valued in $\mathcal{H}^4(\tilde{G}, \text{U}(1))$. When the \mathcal{H}^4 anomaly class vanishes, distinct equivalence classes form a torsor over $\mathcal{H}^3(\tilde{G}, \text{U}(1))$, up to further identifications [54, 55].

It is again useful to think of the SET phases in terms of fluctuating symmetry defect lines. Each defect line is associated with an anyon permutation action given by ρ . The \mathcal{H}^3 obstruction means that defect fusion may fail to be associative: an F move of defect lines may nucleate an extra Abelian anyon, violating the locality requirement. When the \mathcal{H}^3 class vanishes, the extra Abelian anyon can be “absorbed” into decorations of tri-junctions of defects by Abelian anyons. In-equivalent patterns of decorations are classified by a torsor over $\mathcal{H}_\rho^2(\tilde{G}, \mathcal{A})$. Lastly, once we have well-defined defect fusions, including decorations on tri-junctions, there may be a Berry phase in the space of states with defects, which gives the \mathcal{H}^4 anomaly. From this interpretation, it is clear that with a non-trivial $\mathcal{H}_\rho^3(\tilde{G}, \mathcal{A})$ class the map ρ does not make sense in a pure $(2+1)d$ system⁸. The \mathcal{H}^4 anomaly means that the \tilde{G} symmetry has a 't Hooft anomaly. Examples include the surface of a $(3+1)d$ bosonic SPT state or $(2+1)d$ lattice models that satisfy Lieb-Schultz-Mattis-type theorems [56].

Let us now consider a LRE ensemble with topological order \mathcal{C} . The symmetry group \tilde{G} fits into the short exact sequence (21), with exact symmetry A and average symmetry G . We again expect \tilde{G} to be mapped to $\text{Aut}(\mathcal{C})$ through a group homomorphism ρ , which is an invariant of the disorder ensemble. The only way to change ρ is to go through a phase transition, which violates the adiabatic connectability within a single disorder ensemble. It is then useful to think of each state in the disorder ensembles as a topological order with exact A symmetry (i.e. fluctuating A symmetry defects) and a static G defect network.

auto-equivalences of \mathcal{C} and there can be nontrivial elements which do not permute any anyons. However, such examples are only known to occur for very complicated \mathcal{C} , and for simplicity, we do not consider them.

⁸ In field theory, an alternative interpretation of the \mathcal{H}^3 obstruction is that the 0-form symmetry G and the 1-form symmetry \mathcal{A} form a nontrivial 2-group structure.

Generalizing the discussions in Sec. IIB 1, in the presence of disorder it is possible to localize Abelian anyons (as long they do not carry any zero modes protected by the exact symmetry, see below). As a result, there is no longer any energetic requirement to have fixed Abelian anyons decorated on G defect junctions. This is in parallel with dropping the 0D charge decoration in the classification of disordered ASPT phases. In fact, one can gauge the exact symmetry A in a disordered ASPT state to get a disordered ASET state.

We first consider a simpler problem, where the entire symmetry group \tilde{G} becomes average (i.e. A is trivial). In this case, all Abelian anyons can be localized and both the $[O_3]$ obstruction class and the symmetry fractionalization class (i.e. decorations of defect junctions by Abelian anyons) lose their meaning. The same is true for the $\mathcal{H}^4(\tilde{G}, \text{U}(1))$ anomaly and $\mathcal{H}^3(\tilde{G}, \text{U}(1))$ torsor since the disordered SET is an ensemble of defects. We thus conclude that with a trivial A , disordered ASET phases are completely classified by the maps ρ , including those with nontrivial \mathcal{H}^3 obstructions. An example of such an intrinsically disordered ASET is presented below in Sec. IV C.

Next we consider ASETs with a nontrivial A , with a given $\rho : \tilde{G} \rightarrow \text{Aut}(\mathcal{C})$. The map ρ is associated with an obstruction class $[O_3] \in \mathcal{H}_\rho^3(\tilde{G}, \mathcal{A})$. The group $\mathcal{H}_\rho^3(\tilde{G}, \mathcal{A})$ can also be decomposed using the Leray-Serre spectral sequence, whose E_2 page consists of

$$E_2^{p, 3-p} = \bigoplus_{p=0}^3 \mathcal{H}^p(G, \mathcal{H}^{3-p}(A, \mathcal{A})). \quad (61)$$

Here we suppress the group action subscripts for clarity. It should be understood that A acts on \mathcal{A} through ρ , and G acts on the coefficient group \mathcal{H}^{3-p} via both ρ and the G action on A .

Let us start with the component that only involves G , i.e. $E^{3,0} = \mathcal{H}_\rho^3(G, \mathcal{A})$. Since now we only have an ensemble of G defects, if the additional Abelian anyon from an F move of G defects can be localized then this component of the obstruction class no longer makes sense. However, given the exact A symmetry, localization requires that the Abelian anyons involved in the F move do not transform under A symmetry action, and carry no projective multi-dimensional representations of A (a projective one-dimensional representation, i.e. a fractional charge, is allowed).

The other components in the decomposition with $p < 3$ all involve A defects, and hence even when G becomes an average symmetry they still represent nontrivial obstructions.

Having dealt with the \mathcal{H}^3 obstruction, we move to the $\mathcal{H}_\rho^2(\tilde{G}, \mathcal{A})$ torsor. Again we can decompose it using the LHS spectral sequence:

$$E_2^{p, 2-p} = \mathcal{H}_\rho^2(G, \mathcal{A}) \oplus \mathcal{H}^1(G, \mathcal{H}^1(A, \mathcal{A})) \oplus \mathcal{H}_\rho^2(A, \mathcal{A}). \quad (62)$$

Here we also suppress the group action subscripts in the middle term. Via the same reasoning, with localization of




FIG. 4. Anyon permutation of double toric code model with average symmetry \mathbb{Z}_2^A . \mathcal{D} is the symmetry domain wall of \mathbb{Z}_2^A separating the red and blue regimes (with Ising spin- \uparrow and spin- \downarrow), and the green curve depicts the string operator of a \mathbb{Z}_2 gauge group connecting two anyons m_1/m_2 and e_2m_1/e_1m_2 .

Abelian anyons, $\mathcal{H}_\rho^2(G, \mathcal{A})$ may become trivialized, provided that the Abelian anyons involved in this decoration transform trivially (at most as 1D projective reps) under A . The next two terms are still meaningful for disordered ASETs. Perhaps the most interesting part is the $\mathcal{H}^1(G, \mathcal{H}^1(A, \mathcal{A}))$ torsor. When ρ is the identity, in a clean SET this term means that G and A symmetries do not commute when acting on certain anyons. When the G symmetry becomes average, one can interpret this term as the A charge carried by the anyon changes when it passes through certain G defects. Examples of such fractionalizations will be discussed below in Sec. IV D.

We can similarly discuss what changes need to be made for the \mathcal{H}^3 torsor and \mathcal{H}^4 anomaly, however, this is identical to the ASPT classification and there is no need to repeat it.

B. Example: $\mathbb{Z}_2 \times \mathbb{Z}_2$ toric code with \mathbb{Z}_2^A symmetry

We consider an example from a (2+1)d ASPT phase with $A = \mathbb{Z}_2 \times \mathbb{Z}_2$ and $G = \mathbb{Z}_2$ while the extension is trivial. The ASPT phase corresponds to the nontrivial element in $\mathcal{H}^1(G, \mathcal{H}^2(A, \text{U}(1)))$: G domain walls are decorated by (1+1)d A -SPTs.

If we gauge the exact symmetry A to obtain a double toric code topological order, with anyons labeled as $\{e_1, m_1, e_2, m_2\}$ and their combinations, then the average symmetry G will permute the anyons according to

$$\rho: \begin{cases} e_1 \leftrightarrow e_1 \\ e_2 \leftrightarrow e_2 \end{cases}, \begin{cases} m_1 \leftrightarrow m_1 e_2 \\ m_2 \leftrightarrow m_2 e_1 \end{cases} \quad (63)$$

See Fig. 4. This anyon permutation can be seen by firstly considering moving an m_1 anyon across a \mathbb{Z}_2^A domain wall. For the (1+1)d SPT state decorated on the domain wall, moving m_1 across is equivalent to inserting a \mathbb{Z}_2 symmetry defect, which binds a charge of the other \mathbb{Z}_2 , i.e. e_2 . Thus the conservation of topological charge requires $m_1 \rightarrow m_1 e_2$ when passing the \mathbb{Z}_2^A domain wall. Similarly we can show $m_2 \rightarrow m_2 e_1$.

We can relabel the anyons as $\tilde{e}_1 = e_1 m_2$, $\tilde{e}_2 = e_2 m_1$, $\tilde{m}_1 = m_1$, and $\tilde{m}_2 = m_2$, and the theory is rephrased

as two copies of \mathbb{Z}_2 gauge theories, with \mathbb{Z}_2^A simply exchanges the two copies.

C. Intrinsically disordered ASET with \mathbb{Z}_2^A symmetry

It is instructive to study a lattice model for an intrinsically disordered \mathbb{Z}_2^A ASET phase. The model realizes a $\mathbb{D}_{16} = \mathbb{Z}_8 \rtimes \mathbb{Z}_2$ gauge theory, denoted by $\text{D}(\mathbb{D}_{16})$. The group \mathbb{D}_{16} has two generators a and r that satisfy $a^8 = r^2 = 1, rar = a^{-1}$.

For a topological gauge theory with a finite gauge group H , a large class of anyonic symmetries can be understood as outer automorphisms $\text{Out}(H)$ of the gauge group. Here $\text{Out}(H)$ is the quotient $\text{Aut}(H)/\text{Inn}(H)$, where $\text{Aut}(H)$ is the group of automorphisms of H , and $\text{Inn}(H)$ is the group of inner automorphisms (i.e. conjugation by a group element). Physically, the inner automorphisms correspond to gauge transformations, so they do not lead to faithful symmetry actions. Each element in $\text{Out}(H)$ corresponds uniquely to an element in $\text{Aut}(\text{D}(H))$ that does not mix electric and magnetic charges of the gauge theory.

Adopting the general framework in Sec. IV A, different ASETs with an average symmetry group G are classified by homomorphisms $\rho: G \rightarrow \text{Out}(H)$. Mathematically, ρ is associated with a $\mathcal{H}^3(G, Z(H))$ class where $Z(H)$ is the center of the group H . This obstruction arises naturally when classifying the extension of G by H : such an extension always gives a homomorphism from G to $\text{Out}(H)$, but the converse is not necessarily true. Only when the $\mathcal{H}^3(G, Z(H))$ obstruction vanishes the group extension can be defined. Physically, gauging the G symmetry whose action is given by $\rho: G \rightarrow \text{Out}(H)$ should result in a new gauge theory, whose gauge group is an extension of G by H . It turns out that this obstruction of group extension is equivalent to the \mathcal{H}^3 obstruction of anyonic symmetry induced by ρ [57], with $Z(H)$ identified as the group of Abelian pure flux anyons.

We now specialize to the \mathbb{D}_{16} gauge theory. Write $G = \mathbb{Z}_2 = \{1, g\}$. We assume that the image of g under ρ is the following automorphism:

$$\rho_g: a \rightarrow a^5, r \rightarrow ra. \quad (64)$$

The permutation action on anyons is order 2 because ρ_g^2 is the conjugation by a^{-3} . Ref. [57] showed that this ρ has a nontrivial $\mathcal{H}^3(\mathbb{Z}_2, \mathbb{Z}_2)$ class: $O_3(g, g, g) = [a^4]$. Here $[a^4]$ should be understood as the a^4 gauge flux in the \mathbb{D}_{16} gauge theory, which is a \mathbb{Z}_2 Abelian boson. Intuitively, inserting a g defect loop introduces an additional $[a^4]$ anyon.

In Ref. [57] a generalization of Kitaev's quantum double model [58] is introduced for this anomalous SET state. The model is defined on a quasi-2D lattice, where each link has a 16-dimensional Hilbert space, with an orthonormal basis labeled by elements of \mathbb{D}_{16} . They can be

viewed as lattice gauge fields. Schematically, the Hamiltonian takes the following form:

$$H_{\text{clean}} = - \sum_v A_v - \sum_p (\delta_{F_p,1} + \delta_{F_p,a^4}). \quad (65)$$

Here A_v implements the Gauss's law at each vertex v , and F_p is the gauge flux through a plaquette p . For the complete description of the model we refer the readers to [57].

Note that the second term imposes the condition that through each plaquette the gauge flux is either 1 or a^4 (note $Z(\mathbb{D}_{16}) = \{1, a^4\}$). This is distinct from the standard quantum double construction where the plaquette term enforces F_p to be 1. H_{clean} has an extensive ground state degeneracy since we can have 1 or a^4 flux through each plaquette. However, if the fluxes are fixed, the ground state is indeed a \mathbb{D}_{16} gauge theory (with background fluxes).

The reason for this unusual magnetic term is to incorporate the obstructed \mathbb{Z}_2 symmetry. Interested readers can find more details in [57], here we just note that a \mathbb{Z}_2 symmetry transformation can be defined so that it implements the ρ symmetry in the \mathbb{D}_{16} gauge theory. However, under this \mathbb{Z}_2 symmetry transformation, we have $F_p \rightarrow F_p a^4$, which explains the form of the plaquette term.

Now we lift this extensive number of ground states using a disordered perturbation:

$$H[\sigma_p] = H_{\text{clean}} - \sum_p \varepsilon_p (\delta_{F_p,1} - \delta_{F_p,a^4}). \quad (66)$$

Here ε_p are independent random variables drawn symmetrically from the $[-W, W]$ with $W < 1$. The \mathbb{Z}_2 symmetry remains as an average symmetry, under which $\varepsilon_p \rightarrow -\varepsilon_p$.

In accordance with the general principle, the model supports gapless modes for the flux anyon $[a^4]$ localized at certain plaquettes, in order to accommodate the \mathcal{H}^3 obstruction. We can think of the ground state as a localized state of the $[a^4]$ flux anyons.

D. ASET with 't Hooft anomaly: an example with $\mathbb{Z}_2 \times \mathbb{Z}_2^A$

We consider the surface of a (3+1) d ASPT with $\mathbb{Z}_2 \times \mathbb{Z}_2^A$ symmetry. The generator of the \mathbb{Z}_2 (\mathbb{Z}_2^A) will be denoted by g (g_A). In the bulk ASPT state, on each (2+1) d domain wall of the average \mathbb{Z}_2^A symmetry we decorate a Levin-Gu state [42] of the exact \mathbb{Z}_2 (see Fig. 5), with the following topological action on a 4-manifold X_4 :

$$S = \int_{X_4} a^3 \cup b \quad (67)$$

where a and b are background gauge fields of \mathbb{Z}_2 and \mathbb{Z}_2^A , respectively. Notice that when both symmetries are




FIG. 5. Surface topological order of (3+1) d ASPT with $\mathbb{Z}_2 \times \mathbb{Z}_2^A$ symmetry. The indigo surface depicts the \mathbb{Z}_2^A domain wall decorated by a Levin-Gu state, and the violet surface depicts the surface chiral spin liquid enriched by $\mathbb{Z}_2 \times \mathbb{Z}_2^A$ symmetry.

exact, there are two other nontrivial topological terms. One of them is given by

$$S' = \int_{X_4} a \cup b^3. \quad (68)$$

It corresponds to decoration of exact \mathbb{Z}_2 charges on junctions of the \mathbb{Z}_2^A symmetry. As explained in Sec. IIB 1, this action S' is trivialized when \mathbb{Z}_2^A becomes average in a disordered ensemble. The other action is $S + S'$, which is now identified with S .

On the (2+1) d surface of this state, we have a random network of \mathbb{Z}_2^A domain walls, each decorated with a Levin-Gu edge theory described by a Luttinger liquid $[\varphi^T = (\varphi^1, \varphi^2)]$,

$$\mathcal{L} = \frac{K_{IJ}}{4\pi} (\partial_x \varphi^I) (\partial_t \varphi^J) + \frac{V_{IJ}}{8\pi} (\partial_x \varphi^I) (\partial_x \varphi^J) \quad (69)$$

where the K -matrix $K = \sigma^x$. The nontrivial \mathbb{Z}_2 symmetry action is defined as

$$\varphi^{1,2} \mapsto \varphi^{1,2} + \pi. \quad (70)$$

We now want to gap out these domain wall modes in a \mathbb{Z}_2 symmetric way, which can be achieved by placing a semion topological order/chiral spin liquid (CSL) on the surface. Then on each domain wall, we have not only the Levin-Gu edge state but also two counter-propagating chiral Luttinger liquids as the edge modes of the semion topological orders on both sides of the domain wall. The total (1+1) d domain wall theory is

$$\mathcal{L} = \frac{K'_{IJ}}{4\pi} (\partial_x \varphi^I) (\partial_t \varphi^J) + \frac{V'_{IJ}}{8\pi} (\partial_x \varphi^I) (\partial_x \varphi^J) \quad (71)$$

where $\varphi^T = (\varphi^1, \varphi^2, \varphi^3, \varphi^4)$, and the K -matrix $K' = \sigma^x \oplus 2\sigma^z$. A semion is created by the operator $e^{i\varphi^3}$ on one

η	λ_g	λ_{g_A}	S_{anomaly}
1	± 1	± 1	0
-1	1	1	$a \cup b^3 + a^3 \cup b$
-1	1	-1	$a^3 \cup b$
-1	-1	1	$a \cup b^3$
-1	-1	-1	0

TABLE I. 't Hooft anomalies for the projective $\mathbb{Z}_2 \times \mathbb{Z}_2^A$ symmetry actions in clean semion topological order.

side of the domain wall, or $e^{i\varphi^4}$ on the other side. Originally, without Levin-Gu edge modes, the domain wall can be gapped by adding a Higgs term $\cos(2\varphi^3 - 2\varphi^4)$, which induces coherent tunneling of the semions across the domain walls. Alternatively, with the Levin-Gu edge modes, we can gap out the theory in a \mathbb{Z}_2 symmetric way by the following Higgs terms,

$$\cos(\varphi^1 + \varphi^2 - 2\varphi^4) + \cos(\varphi^1 - \varphi^2 - 2\varphi^3). \quad (72)$$

In order for this term to preserve the \mathbb{Z}_2 symmetry, $e^{2i\varphi^4}$ and $e^{2i\varphi^3}$ should be invariant under \mathbb{Z}_2 .

We now analyze the symmetry fractionalization between \mathbb{Z}_2 and \mathbb{Z}_2^A in the semion surface theory. It is instructive to start with the clean case, when both symmetries are exact (we will continue denote the symmetry group as $\mathbb{Z}_2 \times \mathbb{Z}_2^A$). Denote the local h symmetry action on a semion by U_h for every $h \in \mathbb{Z}_2 \times \mathbb{Z}_2^A$. The group relations in $\mathbb{Z}_2 \times \mathbb{Z}_2^A$ lead to the following invariants:

$$\begin{aligned} \lambda_h &= U_h^2, h \in \mathbb{Z}_2 \times \mathbb{Z}_2^A \\ \eta &= U_g U_{g_A} U_g^{-1} U_{g_A}^{-1}. \end{aligned} \quad (73)$$

All of them take ± 1 value. They are related by the algebraic identity: $\lambda_g \lambda_{g_A} \lambda_{gg_A} = \eta$. So there are three independent \mathbb{Z}_2 invariants for the projective symmetry action, consistent with the $\mathcal{H}^2(\mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_2) = \mathbb{Z}_2^3$ classification. The four $\eta = 1$ classes can all be realized by one-dimensional representations, while the $\eta = -1$ classes must be realized by at least two-dimensional representations. In particular, the one with $\lambda_g = \lambda_{g_A} = \lambda_{gg_A} = -1$ is realized in the semion chiral spin liquid, where $\mathbb{Z}_2 \times \mathbb{Z}_2^A$ is a subgroup of the $\text{SO}(3)$ symmetry. The other three classes with $\eta = -1$ are all anomalous with respect to the (exact) $\mathbb{Z}_2 \times \mathbb{Z}_2^A$ symmetry. Their 't Hooft anomalies were computed in [59] and we recall the results in Table I.

When \mathbb{Z}_2^A becomes average, as discussed in Sec. IV A, the invariants λ_{g_A} and λ_{gg_A} are ambiguous due to the localization of semions. However, λ_g and η should remain well-defined. The former should be fairly clear since \mathbb{Z}_2 is an exact symmetry, so we now explain how to define η . Since $U_g U_{g_A} = \eta U_{g_A} U_g$, under the g_A action the \mathbb{Z}_2 charge, measured by the eigenvalue of U_g , changes by η . Therefore, η can be measured as the change of the \mathbb{Z}_2 charge when when adiabatically moving a semion across a g_A defect (which implements the g_A symmetry on the semion).

η	λ_g	S_{anomaly}
1	± 1	0
-1	1	$a \cup b^3$
-1	-1	0

TABLE II. Average 't Hooft anomalies in disordered semion topological order.

First we determine λ_g . In the Luttinger liquid formulation (69), $\lambda_g = -1$ means that under the \mathbb{Z}_2 symmetry, φ^3 and φ^4 must shift by an odd multiples of $\pi/2$, and therefore $e^{2i\varphi^{3/4}}$ should be odd under \mathbb{Z}_2 . However, we have seen that for the Higgs term to preserve symmetry $e^{2i\varphi^{3/4}}$ have to be \mathbb{Z}_2 even. Thus we must have $\lambda_g = 1$.

From Eq. (70) and (72), we find that at the domain wall φ^4 is identified with $\varphi^3 + \varphi^2$, therefore under the \mathbb{Z}_2 symmetry, $e^{i\varphi^4}$ has opposite \mathbb{Z}_2 charge to that of $e^{i\varphi^3}$. In other words, adiabatically moving a semion across a domain wall changes its global \mathbb{Z}_2 charge by -1 . This is the manifestation of the non-commutativity between \mathbb{Z}_2 and \mathbb{Z}_2^A acting on the semion in the ensemble.

Let us now discuss how the anomaly inflow works in the average SET. We have shown that λ_g and η remain good invariants, and λ_{g_A} and λ_{gg_A} are not well-defined individually, but their ratio is fixed by $\lambda_g \eta$. Interestingly, with $\eta = -1$ fixed, the anomalies in Table I separate into two groups corresponding to $\lambda_g = \pm 1$. The two classes inside each group differs just by the $a \cup b^3$ term, which describes decorations of \mathbb{Z}_2 charges on \mathbb{Z}_2^A defect junctions, and becomes trivial due to the localization. This ties nicely with the observation that λ_{g_A} is now ambiguous. The new bulk-boundary correspondence for the semion theory with average $\mathbb{Z}_2 \times \mathbb{Z}_2^A$ symmetry is now summarized in Table II.

This example illustrates that for a mixed group of exact and average symmetry, "symmetry fractionalization" can be well-defined on anyons. We have also derived the complete average 't Hooft anomaly matching for $\mathbb{Z}_2 \times \mathbb{Z}_2^A$ symmetry in the semion topological order.

E. An example with Lieb-Schultz-Mattis anomaly

We consider a $(2+1)d$ lattice system with average $\mathbb{Z} \times \mathbb{Z}$ translation symmetry, exact spin $\text{SO}(3)$ rotation symmetry and a spin-1/2 moment per lattice unit cell. This system has a 't Hooft anomaly from Lieb-Schultz-Mattis (LSM) constraints [56, 60]

$$S = \int_{X_4} x \cup y \cup w_2^{\text{SO}(3)}, \quad (74)$$

where x, y are background gauge fields of the x, y translation symmetries, and $w_2^{\text{SO}(3)}$ is the second Stiefel-Whitney class of the background $\text{SO}(3)$ gauge field. This anomaly remains nontrivial as the $\mathbb{Z} \times \mathbb{Z}$ translation symmetry becomes average [21].

In the clean limit, one of the most well-known topological orders that match with the LSM anomaly is a \mathbb{Z}_2 topological order. In this \mathbb{Z}_2 topological order, the e particle carries a projective representation of $\text{SO}(3)$, corresponding to the nontrivial symmetry fractionalization class in $\mathcal{H}^2(\text{SO}(3), \mathbb{Z}_2)$ (in common term, it carries spin-1/2); the m particle transforms projectively under $\mathbb{Z} \times \mathbb{Z}$ (the nontrivial class in $\mathcal{H}^2(\mathbb{Z} \times \mathbb{Z}, \mathbb{Z}_2)$), in the sense that $T_x T_y = -T_y T_x$ when acting on m , where $T_{x,y}$ are the generators of the x, y translations. These symmetry fractionalizations are required for the \mathbb{Z}_2 topological order to match the anomaly. While the spin-1/2 moment on the e particle is not affected by disorders, we do need to explain the meaning of “ $T_x T_y = -T_y T_x$ ” on the m particle. As discussed in Sec. IV A, the expression $T_x T_y = -T_y T_x$ on m can be interpreted as having an e particle localized at each unit cell (the intersection of a T_x domain wall and a T_y domain wall). If e particle does not carry any degeneracy, then we can deform the state by randomly distributing localized e particles among all unit cells, and states with different e particle distributions will be smoothly connected. However, since e particle carries spin-1/2 moment, the additional localized e particles must find their way to form an $\text{SO}(3)$ invariant (singlet) state. We do not expect the singlet state to be achievable without developing further long-range entanglement. Therefore the notion of “one e particle per unit cell” is robust. Equivalently, $T_x T_y = -T_y T_x$ on m particle and the \mathbb{Z}_2 topological order indeed matches the LSM anomaly.

V. SUMMARY AND OUTLOOK

In this work we have developed a systematic framework, based on the physical picture of defect decoration and the mathematical tool of spectral sequence, to classify and characterize average symmetry-protected topological (ASPT) phases, in both decohered and disordered systems. We have also studied average symmetry-enriched topological (ASET) phases in disordered systems. Our main results are:

- We emphasized the subtle differences between ASPT phases in decohered and disordered systems, which leads to different classifications of ASPT in the two scenarios. Nevertheless, they can both be classified and characterized under the same framework of spectral sequence (decorated defects).
- We discovered a plethora of ASPT phases that are intrinsically mixed, in the sense that they can only appear in mixed state systems (decohered or disordered) with part of the symmetry being average. In other words, these states cannot be viewed as clean SPT states deformed by decoherence or disorder. Some of these states can, however, be viewed as intrinsically gapless pure SPT deformed by decoherence or disorder. We discussed many examples,

in both bosonic and fermionic systems.

- We developed a systematic theory of ASET phases in disordered $(2+1)d$ bosonic systems. Compared to clean SET phases, disordered ASET with average (and exact) symmetries have some distinct features:
 1. While an average symmetry can still permute different anyons, its fractional representation on the anyons cannot be robustly defined, unless the fractionalization pattern involves some ’t Hooft anomaly.
 2. An average symmetry and an exact symmetry can jointly have fractional representation on the anyons.
 3. The \mathcal{H}^3 obstructions of symmetry-enrichment patterns are lifted when the relevant symmetries become average. This leads to intrinsically disordered ASET phases without clean limits. The ground states of such ASET phases contain localized anyons, which leads to gapless (yet still localized) excitation spectra.

We end with some open questions:

- *Boundary physics*: In this work, we mostly focused on the bulk classification and characterization of ASPT phases. In [21], it was shown that the boundary of an ASPT state has an average ’t Hooft anomaly. We have discussed a few examples of bulk-boundary correspondence, including $(1+1)d$ $\mathbb{Z}_2 \times \mathbb{Z}_2^{\text{ave}}$ ASPT phase and the surface ASET of a $(3+1)d$ ASPT phase. It will be useful to understand more systematically how average ’t Hooft anomalies constrain the behavior of the boundary states, especially for the intrinsically mixed/disordered ASPTs with no clean limit.
- *Fermionic ASET*: In this work we studied bosonic ASETs in $(2+1)d$ disordered systems. A natural question is to extend the theory to fermionic systems. In the clean limit, classifications of fermionic SET phases have been developed recently in [61–63]. While certain basic elements of the theory are parallel to the bosonic case, there are important new ingredients and subtleties, especially when ’t Hooft anomalies are concerned.
- *Topological order in mixed states*: We have limited ourselves to topological order in disorder ensembles, where the notion of SRE ensembles can be naturally extended to LRE ensembles. Defining the notion of topological order for a general mixed state is an important question. In fact, a large family of examples can be obtained by “classically” gauging average symmetry and “quantum-mechanically” gauging exact symmetry in ASPT phases.

- *Phase transitions*: quantum phase transitions of intrinsically mixed ASPT or ASET will necessarily involve decoherence or disorders in important manners. This makes the study of such (necessarily non-unitary) quantum phase transitions both challenging and exciting.
- *Physical realizations*: an important task is to realize some ASPT or ASET phases, especially the intrinsically mixed ones, in experimental platforms such as NISQ simulators or disordered solid-state systems. On this front, simple preparation protocols such as those outlined in Sec. III are particularly promising.

ACKNOWLEDGMENTS

We thank Vladimir Calvera, Tyler Ellison, Dominic Else, Zheng-Cheng Gu, David Long, Sagar Vijay, Chenjie Wang, Cenke Xu, Yichen Xu, Carolyn Zhang, and Yijian Zou for the helpful discussions. We are especially grateful to Yijian Zou for pointing out a mistake in the previous version of the manuscript. R.M. acknowledges support from the Natural Sciences and Engineering Research Council of Canada (NSERC) through Discovery Grants. Research at Perimeter Institute (R.M. and C.W.) is supported in part by the Government of Canada through the Department of Innovation, Science and Industry Canada and by the Province of Ontario through the Ministry of Colleges and Universities. J.H.Z. and Z.B. are supported by a startup fund at Penn State University. M.C. acknowledges support from NSF under award number DMR-1846109. J.H.Z., Z.B., and C.W. thank the hospitality of the Kavli Institute for Theoretical Physics, which is supported in part by the National Science Foundation under Grant No. NSF PHY-1748958.

Appendix A: On the definition of disordered ASPT

The definition of disordered average SPT originally given in Ref. [21] was slightly more complicated than that given in Sec. IIB. In particular, in Ref. [21] a short-range entangled ensemble requires any two ground states to be adiabatically connected, namely for any disorder realization I, I' , it is required that $|\Psi_I\rangle = U_{II'}|\Psi_{I'}\rangle$ for some finite-depth local unitary $U_{II'}$. Instead, in Sec. IIB we only demand that any ground state in the ensemble $|\Psi_I\rangle$ to be short-range entangled with correlation length $\xi_I < \xi_m$ for some finite maximal correlation length in the ensemble ξ_m . We now discuss to what extent our new definition is equivalent to the old one.

Consider a ground state in the ensemble $|\Psi_I\rangle$ in a large but finite system without boundary. As a short-range entangled state, it belongs to an invertible phase labeled by $\omega_I^{D+1} \in h^{D+1}(A)$, where $h^{D+1}(A)$ classifies invertible phases with exact symmetry A in D space dimen-

sions. On a finite system, we could effectively compactify the space by viewing certain dimensions as points – for example, the entire system can be viewed as a point if we zoom out far enough. When the system is compactified this way to a $(D - p)$ dimensional space, it should be labeled as a $(D - p)$ invertible state labeled by $\omega_I^{D-p+1} \in h^{D-p+1}(A)$ (it should also depend on the compactification cycle but we will omit this information in the notation). In general ω_I^{D-p+1} is not directly decided by the bulk state ω_I^{D+1} . In particular, when $p = D$ the system reduces to a point and the only nontrivial information left is the charge under A symmetry.

Now the condition in Ref. [21] is the statement that ω_I^{D-p+1} is identical for any disorder realization I , for all $0 \leq p \leq D$. We now show that the new condition in Sec. IIB automatically implies the old condition for $0 \leq p \leq D - 1$: if for some $0 \leq p \leq D - 1$, $\omega_I \neq \omega_{I'}$ for two disorder realizations H_I and $H_{I'}$, then we can choose a third disorder realization in the ensemble I'' , which has $H_{I''} = H_I$ in some region R , and $H_{I''} = H_{I'}$ in the complement \bar{R} . Notice that the spatial independence of the disorder potential is crucial for this construction. The difference in ω in R and \bar{R} , for suitably chosen R , implies a long-range entangled boundary state on ∂R , which leads to a correlation length $\sim |\partial R|$. This leads to unbounded correlation length, and therefore violates the new condition in Sec. IIB.

The above argument does not apply for $p = D$, since two regions with different A charges do not need to have nontrivial boundary state. However, our definition of exact symmetry in Sec. IIB requires that the exact charge to be identical for each disorder realization, so the $p = D$ condition is automatically satisfied. There is still one subtle difference between the old and new definition. In Ref. [21] we consider the ensemble of ground states $\{|\Psi_I^{(0)}\rangle\}$ of the Hamiltonians $\{H_I\}$, and if the states have different total A charges, the ensemble is not considered symmetric SRE. In this work, however, we consider ground states $\{|\Psi_I\rangle\}$ of the Hamiltonians $\{H_I\}$, subject to the condition that the A_I charge is some I -independent fixed value Q . This is analogous to canonical ensemble (instead of grand canonical ensemble) in statistical mechanics. In such “canonical ensemble”, an individual state $|\Psi_I\rangle$ may not be the absolute ground state of H_I . In this case $|\Psi_I\rangle$ has a finite excitation energy, namely the excitation energy density should vanish. If the excitation is localized, in the sense that $|\Psi_I\rangle$ can be obtained from the absolute ground state $|\Psi_I^{(0)}\rangle$ using a finite-depth local unitary circuit $|\Psi_I\rangle = U_{FD}|\Psi_I^{(0)}\rangle$, then $|\Psi_I\rangle$ is still short-range entangled if the absolute ground state is short-range entangled. As long as all $|\Psi_I\rangle$'s are short-range entangled and adiabatically connected to each other, we can still consider the ensemble as symmetric SRE, and the discussions on topological phases can be proceeded with no difficulty. A consequence of using the “canonical ensemble” is that states constructed by decorating zero-dimensional average sym-

metry defects are no longer considered nontrivial, as discussed in Sec. II B.

Appendix B: A brief review of spectral sequence

This section provides a brief overview of the LHS spectral sequence of group cohomology, which is relevant for classifying bosonic ASPT and ASET phases. We also extend our discussion to the more general Atiyah-Hirzebruch (AH) spectral sequence.

A spectral sequence consists of an assembly of Abelian groups $E_r^{p,q}$ ($0 \leq r, p, q \in \mathbb{Z}$). For a fixed r , the collection of all $E_r^{p,q}$ are called E_r page. The differentials are defined as the group endomorphism of the E_r page as:

$$d_r : E_r^{p,q} \rightarrow E_r^{p+r, q-r+1}, \quad (\text{B1})$$

which should satisfy $d_r^2 = 0$. Therefore, the E_r pages and the differentials d_r form a cochain complex, with the following isomorphism:

$$E_{r+1}^{p,q} \simeq \frac{\text{Ker}(d_r^{p,q})}{\text{Im}(d_r^{p-r, q+r-1})} = H_*(E_r, d_r), \quad (\text{B2})$$

where $H_*(E_r, d_r)$ is the homology group of the cochain complex $\{E_r, d_r\}$.

For the LHS spectral sequence with the symmetry group given by Eq. (21), we denote the set of n -cochains, n -cocycles, and n -coboundaries of a group G_0 with coefficients in M by $\mathcal{C}^n[G_0, M]$, $\mathcal{Z}^n[G_0, M]$, and $\mathcal{B}^n[G_0, M]$, respectively.

The E_0 page of the LHS spectral sequence is defined as a group of cochains $E_0^{p,q} = \mathcal{C}^p(G, \mathcal{C}^q[A, M])$, and the d_0 differential maps a cochain in $E_0^{p,q}$ to a cochain $E_0^{p, q+1}$ (see Fig. 6(a) for illustration). The kernel of $d_0^{p,q}$ is $\mathcal{C}^p(G, \mathcal{Z}^q[A, M])$, while the image of $d_0^{p, q-1}$ is just $\mathcal{C}^p(G, \mathcal{B}^q[A, M])$. Hence the E_1 page is given according to Eq. (B2) by

$$\begin{aligned} E_1^{p,q} &= \frac{\text{Ker}(d_0^{p,q})}{\text{Im}(d_0^{p, q-1})} = \frac{\mathcal{C}^p(G, \mathcal{Z}^q[A, M])}{\mathcal{C}^p(G, \mathcal{B}^q[A, M])} \\ &= \mathcal{C}^p(G, \mathcal{H}^q[A, M]). \end{aligned} \quad (\text{B3})$$

We recognize that E_1 page is a subgroup of E_0 page: $E_1 \subset E_0$. More precisely, if we label the elements of $E_0^{p,q}$ by $w_0^{p,q}$, the elements of $E_1^{p,q}$ are the equivalence classes of elements in $E_0^{p,q}$ that satisfy the condition $d_0^{p,q} w_0^{p,q} = 0$.

Subsequently, for the E_1 page $E_1 = \mathcal{C}^p(G, \mathcal{H}^q[A, M])$, the differential d_1 maps a cochain in $\mathcal{C}^p(G, \mathcal{H}^q[A, M])$ to a cochain in $\mathcal{C}^{p+1}(G, \mathcal{H}^q[A, M])$ [see Fig. 6(b)]. The kernel of $d_1^{p,q}$ is cocycles in $\mathcal{Z}^p(G, \mathcal{H}^q[A, M])$, and the image of $d_1^{p-1, q}$ is coboundries in $\mathcal{B}^p(G, \mathcal{H}^q[A, M])$. Then according to Eq. (B2), the E_2 page is given by

$$\begin{aligned} E_2^{p,q} &= \frac{\text{Ker}(d_1^{p,q})}{\text{Im}(d_1^{p-1, q})} = \frac{\mathcal{Z}^p(G, \mathcal{H}^q[A, M])}{\mathcal{B}^p(G, \mathcal{H}^q[A, M])} \\ &= \mathcal{H}^p(G, \mathcal{H}^q[A, M]). \end{aligned} \quad (\text{B4})$$

And E_2 page is a subset of E_1 page,

$$E_2 \subset E_1 \subset E_0. \quad (\text{B5})$$

More precisely, elements of E_2 page are those elements in E_0 page that satisfy the conditions

$$d_0 w_0 = 0, \quad d_1 w_0 = 0. \quad (\text{B6})$$

Following the above paradigm, we can further define the arbitrary E_r page, satisfying the following condition:

$$E_r \subset E_{r-1} \subset \cdots \subset E_2 \subset E_1 \subset E_0. \quad (\text{B7})$$

The elements in E_r page should satisfy the following r conditions,

$$d_q w_0 = 0, \quad 0 \leq q \leq r-1. \quad (\text{B8})$$

In particular, if there is a large enough integer r such that the condition $d_r w_0 = 0$ is satisfied over the entire E_r page, then the E_{r+1} page is essentially identical to the E_r page: $E_{r+1} = E_r$, and all higher pages are the same. It is then said that the spectral sequence stabilizes at the E_r page. For the LHS spectral sequence, the E_∞ page is isomorphic to the group cohomology $\mathcal{H}^{p+q}[\tilde{G}, M]$ as a set. In this paper, we set $M = U(1)$ for the classification of ASPT phases.

The generalization to the AH spectral sequence is straightforward: We can construct the E_r pages of the AH spectral sequence from LHS spectral sequence by substituting the term $\mathcal{H}^q[A, U(1)]$ characterizing the classification of A -symmetric bosonic group-cohomology SPT phases in q -dimensional spacetime by the generalized cohomology group $h^q(A)$ characterizing the classification of A -symmetric invertible topological phases. For example, the E_2 page is defined as


$$E_2^{p,q} = \mathcal{H}^p[G, h^q(A)], \quad p+q = d+1. \quad (\text{B9})$$

Likewise, we can also define the differentials as Eq. (B1), and the AH spectral sequence will converge to the generalized cohomology group $h^{d+1}(\tilde{G})$, i.e., E_∞ is isomorphic to $h^{d+1}(\tilde{G})$.

Below we will see how the AH spectral sequence emerges from a construction of fixed-point wavefunctions of $(2+1)d$ fermionic SPT phases in Appendix C.

Appendix C: Constructions of $(2+1)d$ fermionic SPTs and ASPTs

We review the construction of fixed-point wavefunctions for fermionic SPT (FSPT) phases in $(2+1)d$ interacting fermionic systems following [40], and describe how the construction can be modified for fermionic ASPT phases. This is also a good example to illustrate the AH spectral sequence for decorated domain wall construction. Below G_b denotes the physical symmetry group, assumed to be finite for simplicity.

FIG. 6. LHS spectral sequence and differentials in E_0 , E_1 , and E_2 pages.


The fixed-point wavefunction is constructed by proliferating decorated domain wall: the quantum superposition of all possible G_b symmetry-breaking pattern $|\{g_i\}\rangle$ with fermionic decoration, as

$$|\Psi_{\text{FSPT}}\rangle = \sum_{\{g_i\}} \Psi(\{g_i\}) |\{g_i\}\rangle \quad (\text{C1})$$

The basis state $|\{g_i\}\rangle$ is a state decorated by Majorana chains (the nontrivial element in $h^2(\mathbb{Z}_2^f) = \mathbb{Z}_2$) on the G_b symmetry domain walls, and complex fermions (the nontrivial element in $h^1(\mathbb{Z}_2^f) = \mathbb{Z}_2$) on the junctions of the G_b symmetry domain walls. Therefore, on the triangle lattice, there are three layers of degrees of freedom:

1. $|G_b|$ level bosonic state $|g_i\rangle$ ($g_i \in G_b$) on each vertex;
2. $|G_b|$ species of complex fermions c_{ijk}^σ ($\sigma \in G_b$) at the center of each triangle $\langle ijk \rangle$;
3. $|G_b|$ species of complex fermions (split to pairs of Majorana fermions) $a_{ijk}^\sigma = (\gamma_{ij,A}^\sigma + i\gamma_{ij,B}^\sigma)/2$ on two sides of the link $\langle ij \rangle$.

All degrees of freedom on a triangle are summarized as follows



The G_b symmetry transformations of the degrees of freedom are ($g, g_i, \sigma \in G_b$)

$$\begin{aligned} U(g)|g_i\rangle &= |gg_i\rangle \\ U(g)c_{ijk}^\sigma U^\dagger(g) &= (-1)^{\omega_2(g,\sigma)} c_{ijk}^{g\sigma} \\ U(g)\gamma_{ij,A}^\sigma U^\dagger(g) &= (-1)^{\omega_2(g,\sigma)} \gamma_{ij,A}^{g\sigma} \\ U(g)\gamma_{ij,B}^\sigma U^\dagger(g) &= (-1)^{\omega_2(g,\sigma)+s_1(g)} \gamma_{ij,B}^{g\sigma} \end{aligned} \quad (\text{C3})$$

where $\omega_2(g, \sigma)$ is the factor system of the group extension (21) characterizing the group structure of \tilde{G} , and $s_1(g)$ is defined as

$$s_1(g) = \begin{cases} 0 & \text{if } g \text{ is unitary} \\ 1 & \text{if } g \text{ is antiunitary} \end{cases}. \quad (\text{C4})$$

We first consider the Majorana chain decoration on the domain walls of G_b . The configuration of the decorated Majorana chain is specified by $n_1(g_i, g_j) \in \mathbb{Z}_2$ on the link $\langle ij \rangle$. If the Majorana chain does not go across the link $\langle i, j \rangle$, we say $n_1(g_i, g_j) = 0$, and the Majorana fermions $\gamma_{ij,A}^\sigma$ and $\gamma_{ij,B}^\sigma$ on two sides of the link are paired up as $-i\gamma_{ij,A}^\sigma \gamma_{ij,B}^\sigma = 1$ for $\forall \sigma \in G_b$. If the Majorana chain goes across the link $\langle i, j \rangle$, Majorana fermions $\gamma_{ij,A}^{g_i}$ (and $\gamma_{ij,B}^{g_i}$) will form Majorana entanglement pairs with other Majoranas inside the triangle, and all other Majorana fermions $\gamma_{ij,A}^\sigma$ and $\gamma_{ij,B}^\sigma$ with $\sigma \neq g_i$ are still paired up as $-i\gamma_{ij,A}^\sigma \gamma_{ij,B}^\sigma = 1$. The quantity $n_1(g_0, g_1)n_1(g_1, g_2) = 0, 1$ indicates if $\gamma_{12,A}^{g_1}$ and $\gamma_{01,A}^{g_0}$ are paired up or not [see Eq. (C2)]. The arrows in Eq. (C2) specify the direction of the Majorana pair.

The total number of decorated Majorana chains going through the edges of a specific triangle $\langle 012 \rangle$ is

$$dn_1(g_0, g_1, g_2) = n_1(g_0, g_1) + n_1(g_0, g_2) + n_1(g_1, g_2). \quad (\text{C5})$$

Towards a gapped SPT wavefunction, the above quantity should be even to avoid dangling Majorana fermion, i.e.,

$$dn_1 = 0 \pmod{2}. \quad (\text{C6})$$




FIG. 7. F -move of complex fermion decoration. The Majorana chain decorations are omitted.

Therefore n_1 is an element of $\mathcal{Z}^1[G, h^2(\mathbb{Z}_2^f)]$, the $E_2^{1,2}$ term on the E_2 page of the AH spectral sequence.

We see that the symmetry action may change the directions of the Majorana entanglement pairs, and thereby the fermion parity of a specific triangle. One can find that the fermion parity inside a specific triangle under the symmetry action $U(g_0)$ changes by

$$\Delta P_f^\gamma = (-1)^{(\omega_2 \cup n_1 + s_1 \cup n_1 \cup n_1)(g_0, g_0^{-1}g_1, g_1^{-1}g_2)}. \quad (\text{C7})$$

In order to get a fermionic SPT wavefunction, the wavefunction (C1) should have a definite fermion parity, i.e., states $|\{g\}\rangle$ with different bosonic configurations $\{g\}$ should have the same fermion parity. We take a specific triangulated state $|\{g_0\}\rangle$ and consider a global symmetry transformation $U(g)$, say $U(g)|\{g_0\}\rangle = |\{gg_0\}\rangle$, the fermion parity difference between $|\{g_0\}\rangle$ and $|\{gg_0\}\rangle$ would be the product of Eq. (C7) over the whole triangulation, say

$$\Delta P_f = \prod_{\Delta} (-1)^{(\omega_2 \cup n_1 + s_1 \cup n_1 \cup n_1)(g_1, g_2, g_3)}, \quad (\text{C8})$$

where $g_1, g_2, g_3 \in \{g\}$. Towards a well-defined fixed-point wavefunction of FSPT, we should enforce $\Delta P_f = 1$, and obtain that the following expression should be a 3-coboundary with \mathbb{Z}_2 coefficient,

$$\omega_2 \cup n_1 + s_1 \cup n_1 \cup n_1. \quad (\text{C9})$$

It can be shown that this 3-coboundary is actually dn_2 [cf. Eq. (C10)], where n_2 is the topological index of complex fermion decoration [40].

Lastly, we need to check the pentagon identity is satisfied for the F -moves (see Fig. 7), to make sure that there is no nontrivial Berry phase in the space of decorated domain wall states. As shown in Ref. [40], this condition requires a certain $U(1)$ 4-cocycle $O_4[n_2]$ to be cohomologically trivial. The expression for $O_4[n_2]$ can be

found in [40]. Note that n_1 does not directly contribute to the obstruction function.

To summarize, the input data to the construction is a triple (n_1, n_2, ω_3) , where $n_1 \in \mathcal{C}^1(G_b, \mathbb{Z}_2)$, $n_2 \in \mathcal{C}^2(G_b, \mathbb{Z}_2)$ and $\omega_3 \in \mathcal{C}^3(G_b, U(1))$, satisfying the following consistency conditions:

$$\begin{aligned} dn_1 &= 0, \\ dn_2 &= \omega_2 \cup n_1 + s_1 \cup n_1 \cup n_1, \\ d\omega_3 &= O_4[n_2] \end{aligned} \quad (\text{C10})$$

This is precisely the structure expected from the AH spectral sequence, where n_1, n_2 and ω_3 correspond to the E_2 page, and we can identify the differentials. For example:

$$d_2 n_1 \equiv \omega_2 \cup n_1 + s_1 \cup n_1 \cup n_1, \quad (\text{C11})$$

$$d_2 n_2 \equiv n_2 \cup n_2. \quad (\text{C12})$$

We now modify the construction to get disordered ASPT states. First, the complex fermions c_{ijk}^σ is decoupled from the domain wall configurations, and forms an Anderson insulator. Concretely, we can use the following Hamiltonian for the c_{ijk} fermions:

$$H_{\text{AI}}[\{\varepsilon\}] = - \sum_f \sum_\sigma \varepsilon_{f\sigma} (c_f^\sigma)^\dagger c_f^\sigma. \quad (\text{C13})$$

Here f enumerates all the triangles. $\varepsilon_{f\sigma}$ is a random variable drawn symmetrically from $[-W, W]$ where $W > 0$ (the probability distribution should be independent of σ).

The G_b spins $\{g\}$ are now treated as quenched disorders, assumed to be short-range correlated. The Majorana decorations on domain walls proceed as before, so we include Hamiltonian terms that enforce the decoration pattern, denoted by $H_D[\{g\}]$. The full Hamiltonian reads

$$H[\{g\}, \{\varepsilon\}] = H_D[\{g\}] + H_{\text{AI}}[\{\varepsilon\}]. \quad (\text{C14})$$

The ensemble consists of ground state of $H[\{g\}, \{\varepsilon\}]$ in the total fermion parity even sector.

For decohered SPTs, it is still necessary to impose the fermion parity conservation Eq. (C7). Then the “fixed-point” density matrix is given by

$$\rho = \sum_{\{g\}} p(\{g\}) |\{g\}\rangle \langle \{g\}|. \quad (\text{C15})$$

Here $|\{g\}\rangle$ denote the decorated domain wall states, and $p(\{g\})$ is the probability distribution of domain walls.

[1] Xiao-Gang Wen, *Quantum Field Theory of Many-body Systems: From the Origin of Sound to an Origin of Light and Electrons.*, Oxford Graduate Texts (OUP Premium,

2004).

[2] M. Barkeshli, P. Bonderson, M. Cheng, and Z. Wang, “Symmetry fractionalization, defects, and gauging of

- topological phases,” *Phys. Rev. B* **100**, 115147 (2019), [arXiv:1410.4540 \[cond-mat.str-el\]](#).
- [3] M. Z. Hasan and C. L. Kane, “Colloquium: Topological insulators,” *Rev. Mod. Phys.* **82**, 3045–3067 (2010).
- [4] X.-L. Qi and S.-C. Zhang, “Topological insulators and superconductors,” *Rev. Mod. Phys.* **83**, 1057–1110 (2011).
- [5] F. Duncan M. Haldane, “Nonlinear field theory of large-spin heisenberg antiferromagnets: semiclassically quantized solitons of the one-dimensional easy-axis néel state,” *Physical review letters* **50**, 1153 (1983).
- [6] Xie Chen, Zheng-Cheng Gu, Zheng-Xin Liu, and Xiao-Gang Wen, “Symmetry protected topological orders and the group cohomology of their symmetry group,” *Phys. Rev. B* **87**, 155114 (2013), [arXiv:1106.4772 \[cond-mat.str-el\]](#).
- [7] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, “Symmetry-protected topological orders in interacting bosonic systems,” *Science* **338**, 1604–1606 (2012).
- [8] A. Vishwanath and T. Senthil, “Physics of Three-Dimensional Bosonic Topological Insulators: Surface-Deconfined Criticality and Quantized Magnetoelectric Effect,” *Phys. Rev. X* **3**, 011016 (2013).
- [9] C. Wang and T. Senthil, “Interacting fermionic topological insulators/superconductors in three dimensions,” *Phys. Rev. B* **89**, 195124 (2014).
- [10] Xie Chen, Yuan-Ming Lu, and Ashvin Vishwanath, “Symmetry-protected topological phases from decorated domain walls,” *Nature Communications* **5** (2014), [10.1038/ncomms4507](#).
- [11] T. Senthil, “Symmetry-Protected Topological Phases of Quantum Matter,” *Annu. Rev. Condens. Matter Phys.* **6**, 299–324 (2015).
- [12] Zhen Bi, Alex Rasmussen, Kevin Slagle, and Cenke Xu, “Classification and description of bosonic symmetry protected topological phases with semiclassical nonlinear sigma models,” *Physical Review B* **91**, 134404 (2015).
- [13] John Preskill, “Quantum Computing in the NISQ era and beyond,” *Quantum* **2**, 79 (2018).
- [14] Hannes Bernien, Sylvain Schwartz, Alexander Keesling, Harry Levine, Ahmed Omran, Hannes Pichler, Soonwon Choi, Alexander S. Zibrov, Manuel Endres, Markus Greiner, Vladan Vuletić, and Mikhail D. Lukin, “Probing many-body dynamics on a 51-atom quantum simulator,” *Nature* **551**, 579–584 (2017).
- [15] Mohsin Iqbal, Nathanan Tantivasadakarn, Thomas M. Gatterman, Justin A. Gerber, Kevin Gilmore, Dan Gresh, Aaron Hankin, Nathan Hewitt, Chandler V. Horst, Mitchell Matheny, Tanner Mengle, Brian Neyenhuis, Ashvin Vishwanath, Michael Foss-Feig, Ruben Verresen, and Henrik Dreyer, “Topological Order from Measurements and Feed-Forward on a Trapped Ion Quantum Computer,” (2023), [arXiv:2302.01917 \[quant-ph\]](#).
- [16] Mohsin Iqbal, Nathanan Tantivasadakarn, Ruben Verresen, Sara L. Campbell, Joan M. Dreiling, Caroline Figgatt, John P. Gaebler, Jacob Johansen, Michael Mills, Steven A. Moses, Juan M. Pino, Anthony Ransford, Mary Rowe, Peter Siegfried, Russell P. Stutz, Michael Foss-Feig, Ashvin Vishwanath, and Henrik Dreyer, “Creation of Non-Abelian Topological Order and Anyons on a Trapped-Ion Processor,” (2023), [arXiv:2305.03766 \[quant-ph\]](#).
- [17] Zohar Ringel, Yaacov E. Kraus, and Ady Stern, “Strong side of weak topological insulators,” *Phys. Rev. B* **86**, 045102 (2012), [arXiv:1105.4351 \[cond-mat.mtrl-sci\]](#).
- [18] Roger S. K. Mong, Jens H. Bardarson, and Joel E. Moore, “Quantum Transport and Two-Parameter Scaling at the Surface of a Weak Topological Insulator,” *Phys. Rev. Lett.* **108**, 076804 (2012), [arXiv:1109.3201 \[cond-mat.mes-hall\]](#).
- [19] I. C. Fulga, B. van Heck, J. M. Edge, and A. R. Akhmerov, “Statistical Topological Insulators,” *arXiv e-prints*, [arXiv:1212.6191](#) (2012), [arXiv:1212.6191 \[cond-mat.mes-hall\]](#).
- [20] Caroline de Groot, Alex Turzillo, and Norbert Schuch, “Symmetry Protected Topological Order in Open Quantum Systems,” *Quantum* **6**, 856 (2022).
- [21] Ruochen Ma and Chong Wang, “Average Symmetry-Protected Topological Phases,” (2022), [arXiv:2209.02723 \[cond-mat.str-el\]](#).
- [22] Jong Yeon Lee, Yi-Zhuang You, and Cenke Xu, “Symmetry protected topological phases under decoherence,” *arXiv e-prints*, [arXiv:2210.16323](#) (2022), [arXiv:2210.16323 \[cond-mat.str-el\]](#).
- [23] Jian-Hao Zhang, Yang Qi, and Zhen Bi, “Strange Correlation Function for Average Symmetry-Protected Topological Phases,” *arXiv e-prints*, [arXiv:2210.17485](#) (2022), [arXiv:2210.17485 \[cond-mat.str-el\]](#).
- [24] Andrea Antinucci, Giovanni Galati, Giovanni Rizi, and Marco Serone, “Symmetries and topological operators, on average,” *arXiv e-prints*, [arXiv:2305.08911](#) (2023), [arXiv:2305.08911 \[hep-th\]](#).
- [25] Matthew B. Hastings, “Topological Order at Nonzero Temperature,” *Physical Review Letters* **107**, 210501 (2011).
- [26] Victor V. Albert, “Lindbladians with multiple steady states: theory and applications,” *arXiv e-prints*, [arXiv:1802.00010](#) (2018), [arXiv:1802.00010 \[quant-ph\]](#).
- [27] Berislav Buča and Tomaž Prosen, “A note on symmetry reductions of the Lindblad equation: transport in constrained open spin chains,” *New Journal of Physics* **14**, 073007 (2012), [arXiv:1203.0943 \[quant-ph\]](#).
- [28] Victor V. Albert and Liang Jiang, “Symmetries and conserved quantities in lindblad master equations,” *Phys. Rev. A* **89**, 022118 (2014).
- [29] Simon Lieu, Ron Belyansky, Jeremy T. Young, Rex Lundgren, Victor V. Albert, and Alexey V. Gorshkov, “Symmetry breaking and error correction in open quantum systems,” *Phys. Rev. Lett.* **125**, 240405 (2020).
- [30] D. S. Freed, “Short-range entanglement and invertible field theories,” [arXiv:1406.7278 \[cond-mat.str-el\]](#).
- [31] Daniel S. Freed and Michael J. Hopkins, “Reflection positivity and invertible topological phases,” *arXiv e-prints* (2016), [arXiv:1604.06527](#).
- [32] Elliott Lieb, Theodore Schultz, and Daniel Mattis, “Two soluble models of an antiferromagnetic chain,” *Annals of Physics* **16**, 407–466 (1961).
- [33] Masaki Oshikawa, “Commensurability, Excitation Gap, and Topology in Quantum Many-Particle Systems on a Periodic Lattice,” *Phys. Rev. Lett.* **84**, 1535–1538 (2000), [arXiv:cond-mat/9911137 \[cond-mat.str-el\]](#).
- [34] M. B. Hastings, “Lieb-Schultz-Mattis in higher dimensions,” *Phys. Rev. B* **69**, 104431 (2004), [arXiv:cond-mat/0305505 \[cond-mat.str-el\]](#).
- [35] Ryan Thorngren and Dominic V. Else, “Gauging Spatial Symmetries and the Classification of Topological Crystalline Phases,” *Phys. Rev. X* **8**, 011040 (2018).
- [36] Dmitry A. Abanin, Ehud Altman, Immanuel Bloch, and Maksym Serbyn, “Colloquium: Many-body localization,

- thermalization, and entanglement,” *Reviews of Modern Physics* **91**, 021001 (2019), [arXiv:1804.11065 \[cond-mat.dis-nn\]](#).
- [37] Alexander Elgart and Abel Klein, “Localization in the random XXZ quantum spin chain,” [arXiv e-prints](#), [arXiv:2210.14873](#) (2022), [arXiv:2210.14873 \[math-ph\]](#).
- [38] Davide Gaiotto and Theo Johnson-Freyd, “Symmetry Protected Topological phases and Generalized Cohomology,” *JHEP* **05**, 007 (2019), [arXiv:1712.07950 \[hep-th\]](#).
- [39] Qing-Rui Wang, Shang-Qiang Ning, and Meng Cheng, “Domain wall decorations, anomalies and spectral sequences in bosonic topological phases,” (2021), [arXiv:2104.13233 \[cond-mat.str-el\]](#).
- [40] Q.-R. Wang and Z.-C. Gu, “Construction and classification of symmetry-protected topological phases in interacting fermion systems,” *Phys. Rev. X* **10**, 031055 (2020), [arXiv:1811.00536 \[cond-mat.str-el\]](#).
- [41] Q.-R. Wang, Y. Qi, and Z.-C. Gu, “Anomalous symmetry protected topological states in interacting fermion systems,” *Phys. Rev. Lett.* **123**, 207003 (2019).
- [42] M. Levin and Z.-C. Gu, “Braiding statistics approach to symmetry-protected topological phases,” *Phys. Rev. B* **86**, 115109 (2012).
- [43] Chong Wang, Adam Nahum, Max A. Metlitski, Cenke Xu, and T. Senthil, “Deconfined Quantum Critical Points: Symmetries and Dualities,” *Physical Review X* **7**, 031051 (2017), [arXiv:1703.02426 \[cond-mat.str-el\]](#).
- [44] Ryan Thorngren, Ashvin Vishwanath, and Ruben Verresen, “Intrinsically gapless topological phases,” *Physical Review B* **104**, 075132 (2021).
- [45] Ruochen Ma, Liuju Zou, and Chong Wang, “Edge physics at the deconfined transition between a quantum spin Hall insulator and a superconductor,” *SciPost Physics* **12**, 196 (2022), [arXiv:2110.08280 \[cond-mat.str-el\]](#).
- [46] Rui Wen and Andrew C. Potter, “Bulk-boundary correspondence for intrinsically-gapless SPTs from group cohomology,” (2022), [arXiv:2208.09001 \[cond-mat.str-el\]](#).
- [47] Linhao Li, Masaki Oshikawa, and Yunqin Zheng, “Symmetry Protected Topological Criticality: Decorated Defect Construction, Signatures and Stability,” (2022), [arXiv:2204.03131 \[cond-mat.str-el\]](#).
- [48] X. Chen, Z.-X. Liu, and X.-G. Wen, “Two-dimensional symmetry-protected topological orders and their protected gapless edge excitations,” *Phys. Rev. B* **84**, 235141 (2011).
- [49] Q.-R. Wang and Z.-C. Gu, “Towards a complete classification of symmetry-protected topological phases for interacting fermions in three dimensions and a general group supercohomology theory,” *Phys. Rev. X* **8**, 011055 (2018).
- [50] Naren Manjunath, Vladimir Calvera, and Maissam Barkeshli, “Nonperturbative constraints from symmetry and chirality on majorana zero modes and defect quantum numbers in (2+1) dimensions,” *Phys. Rev. B* **107**, 165126 (2023).
- [51] Alexei Kitaev, “Anyons in an exactly solved model and beyond,” *Annals Phys.* **321**, 2–111 (2006), [arXiv:cond-mat/0506438](#).
- [52] Nicolas Tarantino, Netanel H Lindner, and Lukasz Fidkowski, “Symmetry fractionalization and twist defects,” *New J. Phys.* **18**, 035006 (2016).
- [53] Jeffrey C.Y. Teo, Taylor L. Hughes, and Eduardo Fradkin, “Theory of twist liquids: Gauging an anyonic symmetry,” *Annals of Physics* **360**, 349–445 (2015).
- [54] David Aasen, Parsa Bonderson, and Christina Knapp, “Torsorial actions on G-crossed braided tensor categories,” (2021), [arXiv:2107.10270 \[math.QA\]](#).
- [55] Meng Cheng and Dominic J. Williamson, “Relative anomaly in (1+1)*d* rational conformal field theory,” *Physical Review Research* **2**, 043044 (2020).
- [56] Meng Cheng, Michael Zaletel, Maissam Barkeshli, Ashvin Vishwanath, and Parsa Bonderson, “Translational symmetry and microscopic constraints on symmetry-enriched topological phases: A view from the surface,” *Phys. Rev. X* **6**, 041068 (2016).
- [57] Lukasz Fidkowski and Ashvin Vishwanath, “Realizing anomalous anyonic symmetries at the surfaces of three-dimensional gauge theories,” *Physical Review B* **96**, 045131 (2017).
- [58] A. Yu. Kitaev, “Fault-tolerant quantum computation by anyons,” *Annals of Physics* **303**, 2–30 (2003).
- [59] X. Chen, F. J. Burnell, A. Vishwanath, and L. Fidkowski, “Anomalous symmetry fractionalization and surface topological order,” *Phys. Rev. X* **5**, 041013 (2015).
- [60] Max A. Metlitski and Ryan Thorngren, “Intrinsic and emergent anomalies at deconfined critical points,” *Phys. Rev. B* **98**, 085140 (2018), [arXiv:1707.07686 \[cond-mat.str-el\]](#).
- [61] Daniel Bulmash and Maissam Barkeshli, “Anomaly cascade in (2+1)-dimensional fermionic topological phases,” *Phys. Rev. B* **105**, 155126 (2022).
- [62] Daniel Bulmash and Maissam Barkeshli, “Fermionic symmetry fractionalization in (2+1) dimensions,” *Phys. Rev. B* **105**, 125114 (2022).
- [63] David Aasen, Parsa Bonderson, and Christina Knapp, “Characterization and Classification of Fermionic Symmetry Enriched Topological Phases,” (2021), [arXiv:2109.10911 \[cond-mat.str-el\]](#).