

Absence of one-loop effects on large scales from small scales in non-slow-roll dynamics

Jacopo Fumagalli

*Departament de Física Quàntica i Astrofísica i Institut de Ciències del Cosmos,
Universitat de Barcelona,*

Martí i Franquès 1, 08028 Barcelona, Spain

E-mail: jfumagalli@fqa.ub.edu

ABSTRACT: We show that one-loop corrections to the large-scale power spectrum from small-scale modes in non-slow-roll dynamics are always negligible, namely they are volume suppressed by the ratio of the short to long distance scales. One-loop contributions proportional to the long wavelength tree-level power spectrum, and not sharing this suppression, appear only when considering a subset of vertexes, but they cancel exactly when all relevant interactions are taken into account. We prove the previous statement in two different ways, i.e. by using two equivalent forms of the interaction Hamiltonian. Contributions from boundary terms to equal time correlators are included when necessary.

Contents

1	Introduction	2
2	Different methods to compute the relevant one-loop diagrams	3
2.1	Equivalent ways to express the third-order action	3
2.2	Equivalent Hamiltonians to compute one-loop diagrams	4
3	One-loop corrections in non-slow-roll dynamics	5
3.1	Conventions and approximations	5
3.2	Method 1	7
3.2.1	Contribution from bulk interactions	7
3.2.2	Contributions from boundary terms	9
3.3	Method 2	10
3.3.1	[c-c] contribution	11
3.3.2	[d-c] contribution	12
4	Conclusions	13

1 Introduction

Primordial black holes [1–6] are arguably the *Occam’s Razor* principle applied to Dark Matter [7, 8]. Tracing back their origin to an inflationary epoch requires dynamics beyond the vanilla slow-roll (SR) scenario which are able to enhance the amplitude of the primordial scalar fluctuations at short scales by several orders of magnitude compared to their value at CMB scales. A standard way to achieve the desired magnification is via the introduction of a non-slow-roll phase along the inflationary dynamics, such as, for instance, ultra slow-roll (USR) [9–16] or constant-roll [17–19]. How these small-scale enhanced scalar fluctuations influence large scales, for instance, the one probed by the CMB, has recently been the object of an intense debate [20–28]. Computing one-loop contributions to the equal time correlators in this context leads to the appearance of corrections sharing the following form

$$\mathcal{P}_\zeta^{1\text{-loop}}(p) = c \mathcal{P}_\zeta^{\text{tree}}(p) \int d\ln k \mathcal{P}_\zeta^{\text{tree}}(k) + O\left(\frac{p^3}{k^3}\right), \quad p \ll k, \quad (1)$$

where $\mathcal{P}_\zeta^{\text{tree}/1\text{-loop}}$ are the tree-level and one-loop dimensionless primordial power spectrum, p and k the momenta corresponding to the long and short scale modes respectively, and the integral domain is over the range of momenta spanned by the enhanced fluctuations. Corrections as the one in Eq. (1) has been first found in [20], with $c \sim 10$ for a sharp transition from SR to USR and back to SR. Later, it has also been shown [21, 23–26] that important suppression factors multiplying these contributions are present when considering realistic scenarios in which the transition from the ultra slow-roll to the slow-roll phase is smooth.

Taken at face value, Eq. (1) is telling us something, at the very least, counter-intuitive: the relative 1-loop corrections to the tree-level power spectrum from small scales to large scales is independent of how widely separated the two scales are.

Let us trace the source of this type of corrections. Non-slow-roll scenarios, i.e. the ones under investigations here, are characterized by a sudden increase in absolute value of the so-called second slow-roll parameter η . That promptly calls attention to a particular subset of interaction terms present in the Hamiltonian of the scalar fluctuations. For instance, in [20], that was the cubic interaction proportional to the time derivative of η . It is by considering one-loop diagrams built out of these selected terms alone that corrections as the one in (1) arise.

In this work, we note that once additional diagrams—which might seem irrelevant at first sight—are included, corrections as the one highlighted in Eq. (1) precisely cancel. In short, we find the following exact result:

$$\text{In Eq. (1),} \quad c = 0. \quad (2)$$

We prove the previous statement in two different ways. The first is by considering the same interaction Hamiltonian taken into account in [20] but now with the addition of total time derivative terms (boundary terms) which are nevertheless present once the third order action is written in this form. Boundary terms may contribute in general to equal time correlators [29–34]. The intuition that boundary terms might be relevant in this context

comes from the fact that the same terms do play a crucial role in computing contact diagrams such as the bispectrum [30, 31].

The second method consists of rewriting the action of the curvature perturbations in a form in which boundary terms are inconsequential and the relevant bulk interactions are proportional to η and not its first time derivative. By doing the computation in this way, namely with a different but equivalent set of interaction terms, we confirm the exact cancellation leading to our main result in Eq. (2).

2 Different methods to compute the relevant one-loop diagrams

2.1 Equivalent ways to express the third-order action

We consider standard single-field inflation and work in the gauge where the field fluctuations are set to zero, i.e. $\delta\phi = 0$. The spatial part of the metric reads $g_{ij} = a^2\delta_{ij}a^2e^{2\zeta}$, where a is the scale factor of a flat FLRW background and ζ is the comoving curvature perturbation, the only degree of freedom needed here to describe the dynamics of the scalar fluctuations. The standard second-order action for ζ , obtained after integrating the momentum and Hamiltonian constraints, is given by

$$\mathcal{L}^{(2)} = \frac{1}{2}(M_{\text{Pl}}^2 a^3 2\epsilon) \left[\dot{\zeta}^2 - \frac{(\partial_i \zeta)^2}{a^2} \right], \quad (3)$$

where $\dot{} \equiv d/dt$ means derivative with respect to the cosmic time, $\epsilon \equiv -\dot{H}/H^2$ is the first slow-roll parameter, and $H \equiv \dot{a}/a$ the Hubble rate. We consider scenarios where the second slow-roll parameter, defined as

$$\eta \equiv \frac{\dot{\epsilon}}{\epsilon H}, \quad (4)$$

features a transient growth, i.e. the system experiences a non-slow phase where $|\eta| > 1$ in between two SR phases where $\epsilon, |\eta| \ll 1$. Ultra slow-roll, characterized by a phase with $\eta = -6$, is a particular example of this type. A phase of large η selects a subset of interaction terms from the cubic action, i.e. it is natural to neglect terms with no instance of η in this context. These latter give sub-leading contributions suppressed by the first slow-roll parameter which remains always small. We first consider the third-order action for ζ as derived in [35] once it is written to make explicit the size of the cubic interactions that start at second order in the slow-roll parameters. From there, we consider only terms with instances of the second slow-roll parameter η , i.e.

$$\mathcal{L}^{(3)} = M_{\text{Pl}}^2 \frac{a^3 \epsilon}{2} \dot{\eta} \zeta^2 \dot{\zeta} + M_{\text{Pl}}^2 \frac{d}{dt} \left[-\frac{a^3 \epsilon \eta}{2} \zeta^2 \dot{\zeta} + \dots \right] + f(\zeta) \frac{\delta \mathcal{L}^{(2)}}{\delta \zeta}. \quad (5)$$

To the original cubic action present in [35] we have also included total derivative terms (boundary terms) which were not explicitly written there.¹ The total derivative terms not specified in Eq. (5) are of two types: either they do not contain time derivative of the field ζ and so they do not contribute at any order to the correlation functions, or they give

¹See e.g. [30] for the full list of boundary terms corresponding to the third-order action in [35].

contributions which are suppressed on super-Hubble scale —see for instance [30, 31]. The third group of terms in Eq. (5) labels terms proportional to the linear equation of motion (eom), i.e. the one satisfied by the free field, and we defined

$$-\frac{1}{2M_{\text{Pl}}^2} \frac{\delta \mathcal{L}^{(2)}}{\delta \zeta} \equiv \frac{d}{dt} (a^3 \epsilon \dot{\zeta}) - a^3 \epsilon \frac{\partial^2 \zeta}{a^2}. \quad (6)$$

While terms proportional to the eom does not contribute to Feynman diagram at any order in perturbation theory, there is no a priori reason why the boundary term highlighted above should not contribute to equal time correlators.

Now, let us consider only the terms explicitly written in (5) and do a single integration by parts to remove the time derivative on η present on the first term:

$$\mathcal{L}^{(3)} = -M_{\text{Pl}}^2 \eta \frac{d}{dt} \left[\frac{a^3 \epsilon}{2} \zeta^2 \dot{\zeta} \right] + M_{\text{Pl}}^2 \frac{d}{dt} \left[\frac{a^3 \epsilon \eta}{2} \zeta^2 \dot{\zeta} \right] - M_{\text{Pl}}^2 \frac{d}{dt} \left[\frac{a^3 \epsilon \eta}{2} \zeta^2 \dot{\zeta} \right]. \quad (7)$$

The total time derivative cancels and what is left can be rewritten as

$$\mathcal{L}^{(3)} = -M_{\text{Pl}}^2 \frac{1}{2} a \epsilon \eta \zeta^2 \partial^2 \zeta - M_{\text{Pl}}^2 a^3 \epsilon \eta \dot{\zeta}^2 \zeta + \frac{1}{4} \eta \zeta^2 \frac{\delta \mathcal{L}^{(2)}}{\delta \zeta}, \quad (8)$$

where we have singled out a term proportional to the linear equation of motion.²

To compute correlators in transient non-slow roll scenarios, which are the focus of this work, one can equivalently consider either the terms explicitly reported in the action (5) or the first two terms in the action (8). We stress that in both cases, including all terms highlighted above turned out to be crucial.

2.2 Equivalent Hamiltonians to compute one-loop diagrams

To compute correlators we switch to conformal time ($dt/a = d\tau$), i.e. $\mathcal{S} = \int d^3x dt \mathcal{L} = \int d^3x d\tau (a\mathcal{L})$ and use the interaction Hamiltonian defined as

$$\mathcal{H}_I^{(3)} = - \int d^3x (a\mathcal{L}^{(3)}). \quad (9)$$

We wish to compute 1PI one-loop diagrams following from the insertion of two cubic Hamiltonians. In general, equal time correlators are computed with the *in* – *in* formalism [36–38]

$$\langle \zeta^n(t) \rangle = \langle 0 | \left[\bar{T} \left(e^{i \int_{-\infty(1+i\varepsilon)}^{\tau} d\tau' \mathcal{H}_I(\tau')} \right) \right] \zeta_I^n(\tau) \left[T \left(e^{-i \int_{-\infty(1-i\varepsilon)}^{\tau} d\tau'' \mathcal{H}_I(\tau'')} \right) \right] | 0 \rangle \rangle, \quad (10)$$

where T and \bar{T} are the time and anti-time ordering operators, the subscript I (that we neglect from now on) labels fields in the interaction picture, i.e. fields evolving with the linear equation of motions, and the ε prescription is there to project the adiabatic vacuum of the interacting theory into the vacuum of the free theory $|0\rangle$ in the infinite past. The perturbative expansion of Eq. (10) can be recast, in a compact form, in terms of a series of

²As a side note, this term actually cancels exactly a term proportional to the eom already present in the action (5).

nested commutators –see, for instance, [39]. Although this rewriting loses the information on the UV regularization, the two formulations are equivalent as long as one is interested, as it is the case here, on the effects of interaction term which becomes relevant from a given preferred time.³ For our purposes, we find more convenient to work in terms of the nested commutator form. This latter, for the one-loop contribution we are interested in, leads to

$$\langle \hat{\zeta}_{\mathbf{p}}(\tau) \hat{\zeta}_{\mathbf{p}'}(\tau) \rangle_{1\text{-loop}} = - \int_{\tau_{\text{in}}}^{\tau} d\tau_1 \int_{\tau_{\text{in}}}^{\tau_1} d\tau_2 \langle [\mathcal{H}^{(3)}(\tau_2), [\mathcal{H}^{(3)}(\tau_1), \hat{\zeta}_{\mathbf{p}}(\tau) \hat{\zeta}_{\mathbf{p}'}(\tau)]] \rangle, \quad (11)$$

where τ_{in} labels a generic initial time selected by the interaction terms considered, and $\tau \rightarrow 0$ is the time towards the end of inflation where correlators are evaluated.

Following our discussion in Sec. 2.1, to compute one-loop correction to the two-point correlator $\langle \zeta_{\mathbf{p}} \zeta_{\mathbf{p}'} \rangle$ one can use different although equivalent forms of the interaction Hamiltonian. We proceed following the two equivalent methods summarized below and we obtain in both cases the same results.⁴

- **Method 1**

Insert in Eq. (11) two interaction Hamiltonian as derived from the action (5), i.e.

$$\mathcal{H}^{(3)} = M_{\text{Pl}}^2 \int d^3x \left(-\frac{a^2 \epsilon}{2} \eta' \zeta^2 \zeta' + \frac{d}{d\tau} \left[\frac{a^2 \epsilon \eta}{2} \zeta^2 \zeta' \right] \right) \quad (12)$$

- **Method 2**

Insert in Eq. (11) two interaction Hamiltonian as derived from the action (8), i.e.

$$\mathcal{H}^{(3)} = M_{\text{Pl}}^2 \int d^3x \left(a^2 \epsilon \eta (\zeta')^2 \zeta + \frac{1}{2} a^2 \epsilon \eta \zeta^2 \partial^2 \zeta \right). \quad (13)$$

3 One-loop corrections in non-slow-roll dynamics

3.1 Conventions and approximations

Let us list the few ingredients and approximations needed for our purposes. In momentum space⁵, canonical quantization of the free fields leads to

$$\hat{\zeta}_{\mathbf{p}}(\tau) = \zeta_p(\tau) \hat{a}(\mathbf{p}) + \zeta_p^*(\tau) \hat{a}^\dagger(-\mathbf{p}), \quad (14)$$

with annihilation/creation operators satisfying standard commutation relations

$$[\hat{a}(\mathbf{p}), \hat{a}^\dagger(\mathbf{p}')] = (2\pi)^3 \delta(\mathbf{p} - \mathbf{p}'), \quad (15)$$

³UV divergences that one would usually find using the nested commutator form are not present in our computations simply because time integrals are cut at the time the interactions start to become relevant. By using the nested commutator form we will then re-find the same results present in the literature when considering the same interaction terms.

⁴There is at least a third method one may envisage. A standard way to remove boundary terms is by doing a field redefinition on ζ , then compute correlation functions of the new variable and at the end relate the result to the correlation functions of the original ζ [35]. We plan to investigate this path, in this non-slow-roll context, in future works. Here we concentrate directly on the variable of interest, i.e. ζ .

⁵We use the Fourier transform convention $f(\mathbf{x}) = (2\pi)^{-3} \int d\mathbf{k} e^{-i\mathbf{k} \cdot \mathbf{x}} f(\mathbf{k})$.

and $\zeta_p(\tau)$ labelling the mode functions of the free fields, solutions of the linear equations of motion. From the two previous relations, we write explicitly the following equal time commutators:

$$\left[\hat{\zeta}_{\mathbf{k}}(\tau), \hat{\zeta}_{\mathbf{k}'}(\tau)\right] \equiv (2\pi)^3 \delta(\mathbf{k} + \mathbf{k}') 2i \operatorname{Im}(|\zeta_{\mathbf{k}}|^2) = 0, \quad (16)$$

and

$$\left[\hat{\zeta}_{\mathbf{k}}(\tau), \hat{\zeta}'_{\mathbf{k}'}(\tau)\right] \equiv (2\pi)^3 \delta(\mathbf{k} + \mathbf{k}') 2i \operatorname{Im}(\zeta_{\mathbf{k}} \zeta_{\mathbf{k}'}^*) = (2\pi)^3 \delta(\mathbf{k} + \mathbf{k}') \frac{i}{2a^2 M_{\text{Pl}}^2 \epsilon}. \quad (17)$$

Fields commute at equal time and last equality comes from the Wronskian normalization imposed over the mode functions to have standard commutation relations as in Eq. (15).

We consider a non-slow-roll phase in between two standard slow-roll evolutions. A simple way to model this scenario, while working directly at the level of the effective field theory of the perturbations, is by using the following top-hat profile for η :

$$\eta = \tilde{\eta} + \tilde{\tilde{\eta}}, \quad \text{with} \quad \tilde{\eta} = -\Delta\eta \theta(\tau - \tau_e), \quad \text{and} \quad \tilde{\tilde{\eta}} = \Delta\eta \theta(\tau - \tau_s), \quad (18)$$

τ_s and τ_e label respectively the start and the end of the non-slow-roll phase. We stress that our conclusions are independent of the sharpness of the transition, and θ , here labelling the Heaviside function, can also be replaced by smoother alternatives. Within our definitions, what it is usually called ultra slow-roll corresponds to $\Delta\eta = -6$.

η' peaks at τ_s and τ_e . For simplicity and to capture leading effects when considering interaction terms proportional to η' in the correlators, we only consider the peak at τ_e . At τ_s all mode functions are still following their slow-roll, not enhanced, evolution and we thus disregard these contributions. Thus, we use the following approximation

$$\int^{\tau_1} d\tau_2 \eta'(\tau_2) f(\tau_2) \simeq \int \tilde{\eta}'(\tau_2) f(\tau_e) = (\tilde{\eta}(\tau_1) - \tilde{\eta}(-\infty)) f(\tau_e) = [\tilde{\eta}(\tau_1) + |\Delta\eta|] f(\tau_e). \quad (19)$$

with $f(\tau)$ a continuous function at τ_e .⁶

For simplicity, in this study we always work under the approximation $\mathbf{p} \ll \mathbf{k}$ where \mathbf{k} label modes exiting the horizon around the times the ultra slow-roll dynamics takes place while \mathbf{p} labels momenta associated to the long wave modes. Since ζ_p becomes constant well before reaching the USR region, we can set $\zeta_p(\tau \rightarrow 0) \simeq \zeta_p(\tau_i)$, $\forall \tau_i \in [\tau_s, \tau_e]$. We, thus, use

$$[\hat{\zeta}_{\mathbf{k}}(\tau_i), \hat{\zeta}_{\mathbf{p}}(\tau)] \simeq [\hat{\zeta}_{\mathbf{k}}(\tau_i), \hat{\zeta}_{\mathbf{p}}(\tau_i)] \simeq 0, \quad \forall \tau_i \in [\tau_s, \tau_e]. \quad (20)$$

and

$$[\hat{\zeta}_{\mathbf{k}}(\tau_i), \hat{\zeta}'_{\mathbf{p}}(\tau)] \simeq [\hat{\zeta}_{\mathbf{k}}(\tau_i), \hat{\zeta}'_{\mathbf{p}}(\tau_i)] \simeq (2\pi)^3 \delta(\mathbf{p} + \mathbf{k}) \frac{i}{2a^2 M_{\text{Pl}}^2 \epsilon}, \quad \forall \tau_i \in [\tau_s, \tau_e]. \quad (21)$$

To our taste, in the context under investigation here, using the nested commutator form in Eq. (11), together with the compact notation defined in the next section –see Eq. (30)– allow us to considerably simplify computations. As it will be shown, after unfolding the nested commutators, one can use directly the commutation relations between fields and

⁶Although re-written according to our taste, this discussion leads exactly to the same approximation present, for instance, in [20].

fields derivative (16)-(17) and (20)-(21). These small set of relations is all we need to proceed with our computation.

We are ultimately interested in computing the dimensionless power spectrum \mathcal{P}_ζ defined as

$$\langle \hat{\zeta}_{\mathbf{p}}(\tau) \hat{\zeta}_{\mathbf{p}'}(\tau) \rangle = \delta(\mathbf{p} + \mathbf{p}') (2\pi)^3 \frac{2\pi^2}{p^3} \mathcal{P}_\zeta(p, \tau). \quad (22)$$

\mathcal{P}_ζ is computed perturbatively, i.e.

$$\mathcal{P}_\zeta = \mathcal{P}_\zeta^{\text{tree}} + \mathcal{P}_\zeta^{1\text{-loop}} + \dots, \quad (23)$$

where the tree level power spectrum $\mathcal{P}_\zeta^{\text{tree}}$, obtained by taking the correlators of linear fields, is given by⁷

$$\mathcal{P}_\zeta^{\text{tree}} = \frac{p^3}{2\pi^2} |\zeta_p|^2, \quad (24)$$

while the one-loop corrections $\mathcal{P}_\zeta^{1\text{-loop}}$, computed in the next sections, are given by considering, in Eq. (22), the correlator in Eq. (11).

3.2 Method 1

Let us compute (11) with the interaction Hamiltonian (12) divided in the two sub-pieces

$$\begin{aligned} \mathcal{H}^{(3)} &= \mathcal{H}_a^{(3)} + \mathcal{H}_b^{(3)} \\ \mathcal{H}_a^{(3)} &\equiv M_{\text{Pl}}^2 \int d^3x \left(-\frac{a^2 \epsilon}{2} \eta' \zeta^2 \zeta' \right), \quad \mathcal{H}_b^{(3)} \equiv M_{\text{Pl}}^2 \int d^3x \left(\frac{d}{d\tau} \left[\frac{a^2 \epsilon \eta}{2} \zeta^2 \zeta' \right] \right). \end{aligned} \quad (25)$$

3.2.1 Contribution from bulk interactions

We start by considering only instances of $\mathcal{H}_a^{(3)}$. That allows us to recover the result first found in [20] by means of the nested commutator form. From (11) we can write

$$\begin{aligned} \langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[a, a]} &\equiv - \int^\tau d\tau_1 \int^{\tau_1} d\tau_2 \langle [\mathcal{H}_a^{(3)}(\tau_2), [\mathcal{H}_a^{(3)}(\tau_1), \hat{\zeta}_{\mathbf{p}}(\tau) \hat{\zeta}_{\mathbf{p}'}(\tau)]] \rangle \\ &\equiv \int^\tau d\tau_1 \eta'(\tau_1) \int^{\tau_1} d\tau_2 \eta'(\tau_2) \langle \hat{f}_a(\tau_2, \tau_1) \rangle, \end{aligned} \quad (26)$$

where

$$\hat{f}_a(\tau_2, \tau_1) \equiv -\frac{M_{\text{Pl}}^4}{4} a^2(\tau_1) \epsilon(\tau_1) a^2(\tau_2) \epsilon(\tau_2) \int d\mathbf{K} \left[\hat{\zeta}_{\mathbf{k}_1} \hat{\zeta}_{\mathbf{k}_2} \hat{\zeta}'_{\mathbf{k}_3} |_{\tau_2}, \left[\hat{\zeta}_{\mathbf{k}_4} \hat{\zeta}_{\mathbf{k}_5} \hat{\zeta}'_{\mathbf{k}_6} |_{\tau_1}, \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \right] \right] \quad (27)$$

and Fourier transforming has left us with

$$\int d\mathbf{K} \equiv \prod_{i=1}^3 \left[\int \frac{d\mathbf{k}_{1,i}}{(2\pi)^3} \right] (2\pi)^3 \delta \left(\sum_{i=1}^3 \mathbf{k}_{1,i} \right) \prod_{i=1}^3 \left[\int \frac{d\mathbf{k}_{2,i}}{(2\pi)^3} \right] (2\pi)^3 \delta \left(\sum_{i=1}^3 \mathbf{k}_{2,i} \right). \quad (28)$$

⁷Standard Wick contraction between two linear modes gives $\overline{\hat{\zeta}_{\mathbf{p}}(\tau) \hat{\zeta}_{\mathbf{p}'}(\tau)} = \delta(\mathbf{p} + \mathbf{p}') (2\pi)^3 |\zeta_p|^2$.

Before unfolding the nested commutator let us apply multiple times the approximation in Eq. (19) to Eq. (26):

$$\begin{aligned}\langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_a &\simeq \int^\tau d\tau_1 \eta'(\tau_1) \int^{\tau_1} d\tau_2 \tilde{\eta}'(\tau_2) \langle \hat{f}_a(\tau_e, \tau_1) \rangle = \int^\tau d\tau_1 \eta'(\tau_1) [\tilde{\eta}(\tau_1) + |\Delta\eta|] \langle \hat{f}_a(\tau_e, \tau_1) \rangle \\ &\simeq \int^\tau \left(\frac{(\tilde{\eta}\tilde{\eta})'}{2} + \tilde{\eta}'|\Delta\eta| \right) \langle \hat{f}_a(\tau_e, \tau_e) \rangle = \frac{|\Delta\eta|^2}{2} \langle \hat{f}_a(\tau_e, \tau_e) \rangle.\end{aligned}\quad (29)$$

Let us now turn our attention to the nested commutator inside $\langle \hat{f}_a(\tau_e, \tau_e) \rangle$. To simplify notation and focus on the commuting structure, we label with the same index operators coming from the same vertex, i.e.

$$\left[\hat{\zeta}_{\mathbf{k}_{1,1}} \hat{\zeta}_{\mathbf{k}_{1,2}} \hat{\zeta}'_{\mathbf{k}_{1,3}} |_{\tau_2}, \left[\hat{\zeta}_{\mathbf{k}_{2,1}} \hat{\zeta}_{\mathbf{k}_{2,2}} \hat{\zeta}'_{\mathbf{k}_{2,3}} |_{\tau_1}, \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} |_{\tau} \right] \right] \equiv \left[\hat{\zeta}_2^2 \hat{\zeta}_2', \left[\hat{\zeta}_1^2 \hat{\zeta}_1', \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \right] \right]. \quad (30)$$

Note that, after doing all contractions and integrating over the Dirac deltas, the overall momentum conservation allows us to simply replace k and $|\mathbf{p} - \mathbf{k}|$ to the momenta of the mode functions corresponding to internal contractions (contractions of $\hat{\zeta}$'s corresponding to different vertexes) and p to the momenta of the mode functions corresponding to contractions with external legs, all that by leaving behind an overall integration $\int d\mathbf{k}$. From the approximation (29) and using notation (30), what we wish to compute takes the form

$$\langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[a,a]} = -|\Delta\eta|^2 \frac{M_{\text{Pl}}^4 a^4 \epsilon^2 |_{\tau_e}}{8} \int d\mathbf{K} \langle \left[\hat{\zeta}_2^2 \hat{\zeta}_2', \left[\hat{\zeta}_1^2 \hat{\zeta}_1', \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \right] \right] \rangle \Big|_{\tau_1=\tau_2=\tau_e}. \quad (31)$$

We now unfold the nested commutator. As already mentioned, fields operators (with no derivative) commutes at equal time. That simplifies considerably the computation which is then almost trivialized in the few lines below:

$$\begin{aligned}\left[\hat{\zeta}_2^2 \hat{\zeta}_2', \left[\hat{\zeta}_1^2 \hat{\zeta}_1', \hat{\zeta}_{\mathbf{p}}^2 \right] \right] &= 2 \left[\hat{\zeta}_1', \hat{\zeta}_{\mathbf{p}} \right] \cdot \left[\hat{\zeta}_2^2 \hat{\zeta}_2', \hat{\zeta}_1^2 \hat{\zeta}_{\mathbf{p}} \right] \\ &= 2 \left[\hat{\zeta}_1', \hat{\zeta}_{\mathbf{p}} \right] \cdot \hat{\zeta}_2^2 \left(\left[\hat{\zeta}_2', \hat{\zeta}_1^2 \right] \hat{\zeta}_{\mathbf{p}} + \hat{\zeta}_1^2 \left[\hat{\zeta}_2', \hat{\zeta}_{\mathbf{p}} \right] \right) \\ &= 2 \left[\hat{\zeta}_1', \hat{\zeta}_{\mathbf{p}} \right] \cdot \left(2 \left[\hat{\zeta}_2', \hat{\zeta}_1 \right] \hat{\zeta}_2^2 \hat{\zeta}_1 \hat{\zeta}_{\mathbf{p}} + \left[\hat{\zeta}_2', \hat{\zeta}_{\mathbf{p}} \right] \hat{\zeta}_2^2 \hat{\zeta}_1^2 \right).\end{aligned}\quad (32)$$

We replace each building block commutator with (17) which cancels the prefactor $M_{\text{Pl}}^4 a^4 \epsilon^2$ in (31), and we then Wick contract operators left outside the commutators. That promptly leads to

$$\langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[a,a]} = \delta(\mathbf{p} + \mathbf{p}') \frac{|\Delta\eta|^2}{4} \int d\mathbf{k} \left(|\zeta_{\mathbf{p}}|^2 |\zeta_{\mathbf{k}}|^2 + \frac{1}{2} |\zeta_{\mathbf{k}}|^2 |\zeta_{|\mathbf{p}-\mathbf{k}|}|^2 \right). \quad (33)$$

The second term is a convolution of the modes running in the loop. Once going to the dimensionless power spectrum (22), that correction is volume suppressed, i.e. scale as the ratio of the short to the long scales cube:

$$\int d\ln k k^3 |\zeta'_k|^2 |\zeta_{|\mathbf{p}-\mathbf{k}|}|^2 \propto \frac{1}{k^3} \implies \mathcal{P}_{\zeta}^{1-\text{loop}}(p) \propto \left(\frac{p}{k} \right)^3 \simeq 0. \quad (34)$$

We mention this explicitly only once here and neglect this type of contributions in the following sections.⁸

⁸One can check that, by using the approximation (20), we had already neglected volume suppressed terms proportional to convolution of first derivatives of the mode functions with momenta k and $|\mathbf{p} - \mathbf{k}|$.

We use the symbol \cong to identify quantities which are *equal up to volume suppressed terms*.

Considering the first and dominant term in Eq. (33) and using the definition (22) we get, with obvious notation,

$$\mathcal{P}_{\zeta, [a, a]}^{1\text{-loop}}(p) \cong \frac{|\Delta\eta|^2}{4} \mathcal{P}_{\zeta}^{\text{tree}}(p) \int \frac{d\mathbf{k}}{(2\pi)^3} |\zeta_{\mathbf{k}}(\tau_e)|^2. \quad (35)$$

Last expression has the form of the first term in Eq. (1) with $c = |\Delta\eta|^2/4$.

3.2.2 Contributions from boundary terms

Let us now compute contributions from boundary terms. By inserting two instances of \mathcal{H}_b defined in Eq. (25) into Eq. (11) we get

$$\begin{aligned} \langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[b, b]} &\equiv - \int^\tau d\tau_1 \int^{\tau_1} d\tau_2 \langle [\mathcal{H}_b^{(3)}(\tau_2), [\mathcal{H}_b^{(3)}(\tau_1), \hat{\zeta}_{\mathbf{p}}(\tau) \hat{\zeta}_{\mathbf{p}'}(\tau)]] \rangle \\ &\equiv \int^\tau d\tau_1 \int^{\tau_1} d\tau_2 \langle \hat{f}_b(\tau_2, \tau_1) \rangle \end{aligned} \quad (36)$$

with

$$\hat{f}_b(\tau_2, \tau_1) \equiv -\frac{M_{\text{Pl}}^4}{4} \int d\mathbf{K} \left[\left(a^2 \epsilon \eta \hat{\zeta}_{\mathbf{k}_{2,1}} \hat{\zeta}_{\mathbf{k}_{2,2}} \hat{\zeta}'_{\mathbf{k}_{2,3}} \right)' \Big|_{\tau_2}, \left[\left(a^2 \epsilon \eta \hat{\zeta}_{\mathbf{k}_{1,1}} \hat{\zeta}_{\mathbf{k}_{1,2}} \hat{\zeta}'_{\mathbf{k}_{1,3}} \right)' \Big|_{\tau_1}, \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \right] \right], \quad (37)$$

where $d\mathbf{K}$ was defined in Eq. (28). Integration over τ_2 can be performed straightforwardly. Then we expand the derivative in the first nested commutator and obtain

$$\langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[b, b]} = -\frac{M_{\text{Pl}}^4}{4} \int d\mathbf{K} \int^\tau d\tau_1 \langle \left[a^2 \epsilon \eta \hat{\zeta}^2 \hat{\zeta}', \left[\eta (a^2 \epsilon \hat{\zeta}')' + \eta' a^2 \epsilon \hat{\zeta}^2 \hat{\zeta}' + 2a^2 \epsilon (\hat{\zeta}')^2 \hat{\zeta}, \hat{\zeta}_{\mathbf{p}}^2 \right] \right] \rangle, \quad (38)$$

where all operators in the nested commutator are evaluated at the same time τ_1 . By using the linear equations of motion to rewrite the first term in the nested commutator in Eq. (38), i.e. $(a^2 \epsilon \hat{\zeta}')' = a^2 \epsilon \partial^2 \hat{\zeta}$, one notices that this term gives zero contribution since fields commute at equal time. Applying, as in the previous section, the same approximations (19) to the second term in the nested commutator in Eq. (38) we get

$$-\frac{M_{\text{Pl}}^4}{4} \int d\mathbf{K} \int^\tau \frac{(\eta\eta)'}{2} a^4 \epsilon^2 \langle \left[\hat{\zeta}_2 \hat{\zeta}_2 \hat{\zeta}', \left[\hat{\zeta}_1 \hat{\zeta}_1 \hat{\zeta}', \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \right] \right] \rangle \simeq -\frac{|\Delta\eta|^2}{2} \langle \hat{f}_a(\tau_e, \tau_e) \rangle = -\langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[a, a]}, \quad (39)$$

where \hat{f}_a was defined in Eq. (27). By adding the third term in the nested commutator in Eq. (38) we can rewrite this all contribution as

$$\langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[b, b]} = -\langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[a, a]} - M_{\text{Pl}}^4 \frac{|\Delta\eta|^2}{2} \int d\mathbf{K} \int_{\tau_s}^{\tau_e} d\tau_1 a^4 \epsilon^2 \langle \left[\hat{\zeta}_2 \hat{\zeta}_2 \hat{\zeta}', \left[\hat{\zeta}_1' \hat{\zeta}_1' \hat{\zeta}_1, \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \right] \right] \rangle. \quad (40)$$

By proceeding analogously one can show that

$$\langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[a, b]} \cong -\langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[b, a]} \cong -\langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[a, a]}, \quad (41)$$

where, with the same notation as in Eq. (36), $\langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[a,b]}$ refers to Eq. (11) with the insertions $\mathcal{H}_a^{(3)}(\tau_2)$ and $\mathcal{H}_b^{(3)}(\tau_1)$, while $\langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[b,a]}$ refers to Eq. (11) with the insertions $\mathcal{H}_b^{(3)}(\tau_2)$ and $\mathcal{H}_a^{(3)}(\tau_1)$. From Eq. (41) we thus note that the total contribution from boundary terms is just given by $\langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[b,b]}$.

From (40) we proceed as in Eq. (32) (also symmetrizing not commuting operators⁹). After tedious manipulations we get

$$\begin{aligned} [\hat{\zeta}'_1 \hat{\zeta}'_1 \hat{\zeta}'_1, \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'}] &= 4 [\hat{\zeta}'_1, \hat{\zeta}_{\mathbf{p}'}] \left(2 \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_2 \hat{\zeta}'_1 \hat{\zeta}'_2 [\hat{\zeta}_2, \hat{\zeta}'_1] + \hat{\zeta}_2^2 \hat{\zeta}'_1 \hat{\zeta}'_1 [\hat{\zeta}'_2, \hat{\zeta}_{\mathbf{p}}] + \hat{\zeta}_2^2 \hat{\zeta}_{\mathbf{p}} \hat{\zeta}'_1 [\hat{\zeta}'_2, \hat{\zeta}'_1] \right. \\ &\quad \left. + 2 \hat{\zeta}'_2 \hat{\zeta}_1 \hat{\zeta}_{\mathbf{p}} \hat{\zeta}'_2 [\hat{\zeta}_2, \hat{\zeta}'_1] + \hat{\zeta}_1 \hat{\zeta}'_1 \hat{\zeta}_2^2 [\hat{\zeta}'_2, \hat{\zeta}_{\mathbf{p}}] + \hat{\zeta}_{\mathbf{p}} \hat{\zeta}'_1 \hat{\zeta}_2^2 [\hat{\zeta}'_2, \hat{\zeta}'_1] \right). \end{aligned} \quad (42)$$

All commutators are evaluated at equal time so that we can use Eq. (17) to cancel the $4a^2\epsilon^2$ prefactor. Wick contracting operators outside commutators, and using the power spectrum definition in Eq. (22) lead to (still neglecting terms proportional to ζ'_p)

$$\begin{aligned} \mathcal{P}_{\zeta, [b,b]}^{1-\text{loop}} + \mathcal{P}_{\zeta, [a,a]}^{1-\text{loop}} &\cong - \frac{|\Delta\eta|^2}{2} \frac{p^3}{2\pi^2} \int \frac{d\mathbf{k}}{(2\pi)^3} \int d\tau_1 (|\zeta_p|^2 (|\zeta_k|^2)' - |\zeta_p|^2 (|\zeta_k|^2)' - |\zeta_k|^2 (|\zeta_{\mathbf{p}-\mathbf{k}}|^2)') \\ &\cong 0. \end{aligned} \quad (43)$$

Note that terms proportional to the long-wave power spectrum, namely the first two terms, exactly cancel. The last term just gives a volume suppressed contribution, i.e. scales as $(p/k)^3$.

3.3 Method 2

We now carry out an equivalent derivations which does not require the introduction of boundary terms. In a sense, this is a more orthodox way of proceeding. We compute (11) with the interaction Hamiltonian (13). As before, it is useful to divide the Hamiltonian in two sub-pieces:

$$\begin{aligned} \mathcal{H}^{(3)} &= \mathcal{H}_c^{(3)} + \mathcal{H}_d^{(3)} \\ \mathcal{H}_c^{(3)} &= M_{\text{Pl}}^2 \int d^3x a^2 \epsilon \eta (\zeta')^2 \zeta, \quad \mathcal{H}_d^{(3)} = \frac{1}{2} M_{\text{Pl}}^2 \int d^3x a^2 \epsilon \eta \zeta^2 \partial^2 \zeta. \end{aligned} \quad (44)$$

From (11) we have two non-trivial contributions:

$$\langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[c,c]} \equiv - \int^\tau d\tau_1 \int^{\tau_1} d\tau_2 \langle [\mathcal{H}_c^{(3)}(\tau_2), [\mathcal{H}_c^{(3)}(\tau_1), \hat{\zeta}_{\mathbf{p}}(\tau) \hat{\zeta}_{\mathbf{p}'}(\tau)]] \rangle, \quad (45)$$

and

$$\langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[d,c]} \equiv - \int^\tau d\tau_1 \int^{\tau_1} d\tau_2 \langle [\mathcal{H}_d^{(3)}(\tau_2), [\mathcal{H}_c^{(3)}(\tau_1), \hat{\zeta}_{\mathbf{p}}(\tau) \hat{\zeta}_{\mathbf{p}'}(\tau)]] \rangle. \quad (46)$$

Terms where the Hamiltonian in the first nested commutator is equal to $\mathcal{H}_d^{(3)}$ give negligible (volume suppressed) contributions since ζ_p is frozen during the USR phase and fields operators commute at equal time, see (20).

⁹Operators which do not commute at equal time have always to be thought as in their symmetrized version, i.e. $(\hat{\zeta}_1)^2 \hat{\zeta}_1 = 1/2((\hat{\zeta}_1')^2 \hat{\zeta}_1 + \hat{\zeta}_1 (\hat{\zeta}_1')^2)$, we did not write that explicitly at each step just to leave the notation lighter, but we performed the full computations with the symmetrized operators.

3.3.1 [c-c] contribution

Let us consider two instances of \mathcal{H}_c , as defined in (44), into Eq. (11):

$$\begin{aligned} \langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[c, c]} &= -M_{\text{Pl}}^4 |\Delta\eta|^2 \int_{\tau_s}^{\tau_e} d\tau_1 (a^2 \epsilon)|_{\tau_1} \int_{\tau_s}^{\tau_1} d\tau_2 (a^2 \epsilon)|_{\tau_2} \\ &\quad \times \int d\mathbf{K} \langle [(\hat{\zeta}'_{\mathbf{k}_1} \hat{\zeta}'_{\mathbf{k}_2} \hat{\zeta}_{\mathbf{k}_3})|_{\tau_2}, [(\hat{\zeta}'_{\mathbf{k}_4} \hat{\zeta}'_{\mathbf{k}_5} \hat{\zeta}_{\mathbf{k}_6})|_{\tau_1}, \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'}] \rangle. \end{aligned} \quad (47)$$

We start by unfolding the first nested commutator. Using the same compact notation defined in Eq. (30), that becomes

$$\begin{aligned} [\hat{\zeta}'_1 \hat{\zeta}'_1 \hat{\zeta}_1, \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'}] &= [\hat{\zeta}'_1, \hat{\zeta}_{\mathbf{p}'}] (\hat{\zeta}'_1 \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_1 + \hat{\zeta}_{\mathbf{p}} \hat{\zeta}'_1 \hat{\zeta}_1 + \hat{\zeta}_1 \hat{\zeta}'_1 \hat{\zeta}_{\mathbf{p}} + \hat{\zeta}_1 \hat{\zeta}_{\mathbf{p}} \hat{\zeta}'_1) \\ &= [\hat{\zeta}'_1, \hat{\zeta}_{\mathbf{p}'}] (2\hat{\zeta}'_1 \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_1 + [\hat{\zeta}_{\mathbf{p}}, \hat{\zeta}'_1] \hat{\zeta}_1 + 2\hat{\zeta}_1 \hat{\zeta}'_1 \hat{\zeta}_{\mathbf{p}} + \hat{\zeta}_1 [\hat{\zeta}_{\mathbf{p}}, \hat{\zeta}'_1]) \\ &= [\hat{\zeta}'_1, \hat{\zeta}_{\mathbf{p}'}] (4\hat{\zeta}'_1 \hat{\zeta}_1 \hat{\zeta}_{\mathbf{p}} + 2[\hat{\zeta}_1, \hat{\zeta}'_1] \hat{\zeta}_{\mathbf{p}} + 2\hat{\zeta}_1 [\hat{\zeta}_{\mathbf{p}}, \hat{\zeta}'_1]). \end{aligned} \quad (48)$$

In the second line we have added and subtracted terms equal to the first and third one in parenthesis on the first line. On the third line we added and subtracted a term equal to the first in parenthesis on the second line and used the fact that $[\hat{\zeta}_{\mathbf{p}}, \hat{\zeta}'_1] \simeq 0$. Terms proportional to commutators in the parenthesis of the final expression just lead to tadpole diagrams, namely they force contractions between two legs from the same vertexes once embedded in the full expression (47). As before we focus on 1PI diagrams and so we disregard these terms. Inserting Eq. (48) in the full nested commutator in (47), and after similar manipulations as the one just discussed, we get

$$[\hat{\zeta}'_2 \hat{\zeta}'_2 \hat{\zeta}_2, [\hat{\zeta}'_1 \hat{\zeta}'_1 \hat{\zeta}_1, \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'}]] = 8[\hat{\zeta}'_1, \hat{\zeta}_{\mathbf{p}'}] \left([\hat{\zeta}'_2, \hat{\zeta}'_1] \hat{\zeta}'_2 \hat{\zeta}_2 \hat{\zeta}_1 \hat{\zeta}_{\mathbf{p}} + [\hat{\zeta}'_2, \hat{\zeta}'_1] \hat{\zeta}'_1 \hat{\zeta}'_2 \hat{\zeta}_2 \hat{\zeta}_{\mathbf{p}} + [\hat{\zeta}'_2, \hat{\zeta}_{\mathbf{p}}] \hat{\zeta}'_1 \hat{\zeta}_1 \hat{\zeta}'_2 \hat{\zeta}_2 \right). \quad (49)$$

Last term in the previous equation just lead to a volume suppressed contribution, see (34). Let us then consider only the first two terms in (49). After the usual Wick contractions we have that the one-loop correction to the power spectrum from these terms can be written as:¹⁰

$$\mathcal{P}_{\zeta, [c, c]}^{1\text{-loop}}(p) \cong -8M_{\text{Pl}}^2 |\Delta\eta|^2 \mathcal{P}_{\zeta}^{\text{tree}}(p) \int_{\tau_s}^{\tau_e} d\tau_1 \int_{\tau_s}^{\tau_1} d\tau_2 a^2(\tau_2) \epsilon(\tau_2) \int \frac{d\mathbf{k}}{(2\pi)^3} \text{Im}(\mathcal{X}_{\mathbf{k}} \mathcal{Y}_{|\mathbf{p}-\mathbf{k}|}), \quad (50)$$

where

$$\mathcal{X}_{\mathbf{k}} = \zeta'_{\mathbf{k}}(\tau_2) \zeta_{\mathbf{k}}'^*(\tau_1), \quad \text{and} \quad \mathcal{Y}_{\mathbf{k}} = \zeta'_{\mathbf{k}}(\tau_2) \zeta_{\mathbf{k}}^*(\tau_1). \quad (51)$$

As before, we replace commutators such as $[\hat{\zeta}'_1, \hat{\zeta}_{\mathbf{p}}]$ with the Wronskian conditions in Eq. (21). From (50) we single out terms which depend only on τ_2 and we carry out integration by parts using the linear equation of motion for ζ , i.e. $(a^2 \epsilon \zeta')' = a^2 \epsilon \partial^2 \zeta$. As mentioned, we do not need to use the explicit form of the mode functions. Proceeding in this way, the integrals over τ_2 can be rewritten as

$$\int_{\tau_s}^{\tau_1} d\tau_2 a(\tau_2)^2 \epsilon(\tau_2) \zeta'_{\mathbf{k}} \zeta'_{|\mathbf{p}-\mathbf{k}|} = (a^2 \epsilon \zeta'_{\mathbf{k}} \zeta'_{|\mathbf{p}-\mathbf{k}|})|_{\tau_1} + \int_{\tau_s}^{\tau_1} d\tau_2 a^2 \epsilon k^2 \zeta_{\mathbf{k}} \zeta_{|\mathbf{p}-\mathbf{k}|} \quad (52)$$

¹⁰Eq. (50) coincides with the leading order result found in [24] (see Appendix A, Eq. (97)), with the exception that we found a crucial time switch $\tau_1 \leftrightarrow \tau_2$.

We symmetrize the integrand over the two momenta k and $|\mathbf{p} - \mathbf{k}|$, then we insert the previous expression and its complex conjugate in (50). Surface terms cancel and after regrouping what is left, we finally obtain

$$\begin{aligned} \mathcal{P}_{\zeta, [c, c]}^{1-\text{loop}}(p) \cong & -4M_{\text{Pl}}^2 |\Delta\eta|^2 \mathcal{P}_{\zeta}^{\text{tree}}(p) \int_{\tau_s}^{\tau_e} d\tau_1 \int_{\tau_s}^{\tau_1} d\tau_2 a^2(\tau_2) \epsilon(\tau_2) \int \frac{d\mathbf{k}}{(2\pi)^3} (k^2 + |\mathbf{p} - \mathbf{k}|^2) \\ & \times \text{Im} \left(\zeta_k^*(\tau_1) \zeta_{|\mathbf{p}-\mathbf{k}|}^{\prime*}(\tau_1) \zeta_k(\tau_2) \zeta_{|\mathbf{p}-\mathbf{k}|}(\tau_2) \right). \end{aligned} \quad (53)$$

3.3.2 [d-c] contribution

$$\begin{aligned} \langle \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'} \rangle_{[d, c]} = & -M_{\text{Pl}}^4 \frac{|\Delta\eta|^2}{2} \int_{\tau_s}^{\tau_e} d\tau_1 (a^2 \epsilon)|_{\tau_1} \int_{\tau_s}^{\tau_1} d\tau_2 (a^2 \epsilon)|_{\tau_2} \\ & \times \int d\mathbf{K} \cdot \mathcal{A} \langle [(\hat{\zeta}_{\mathbf{k}_1} \hat{\zeta}_{\mathbf{k}_2} \hat{\zeta}_{\mathbf{k}_3})|_{\tau_2}, [(\hat{\zeta}_{\mathbf{k}_4}' \hat{\zeta}_{\mathbf{k}_5}' \hat{\zeta}_{\mathbf{k}_6}')|_{\tau_1}, \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'}] \rangle. \end{aligned} \quad (54)$$

where

$$\mathcal{A} = -\frac{1}{3} \sum_{i=1}^3 k_{2,i}^2. \quad (55)$$

The first nested commutator is the same as in the previous section, we thus use Eq. (48) and neglect tadpoles. Using the same compact notation as defined in (30) and the same approximation of the previous sections we have

$$\begin{aligned} [\hat{\zeta}_2 \hat{\zeta}_2 \hat{\zeta}_2, [\hat{\zeta}_1' \hat{\zeta}_1' \hat{\zeta}_1', \hat{\zeta}_{\mathbf{p}} \hat{\zeta}_{\mathbf{p}'}]] &= 4[\hat{\zeta}_1', \hat{\zeta}_{\mathbf{p}'}] \cdot [\hat{\zeta}_2 \hat{\zeta}_2 \hat{\zeta}_2, \hat{\zeta}_1' \hat{\zeta}_1'] \hat{\zeta}_{\mathbf{p}} \\ &= 4[\hat{\zeta}_1', \hat{\zeta}_{\mathbf{p}'}] \cdot [\hat{\zeta}_2, \hat{\zeta}_1] (2\hat{\zeta}_2 \hat{\zeta}_2 \hat{\zeta}_1' \hat{\zeta}_{\mathbf{p}} + \hat{\zeta}_2 \hat{\zeta}_{\mathbf{p}} [\hat{\zeta}_1', \hat{\zeta}_2] + \hat{\zeta}_1' \hat{\zeta}_2 \hat{\zeta}_2 \hat{\zeta}_{\mathbf{p}}) \\ &\quad + 4[\hat{\zeta}_1', \hat{\zeta}_{\mathbf{p}'}] \cdot [\hat{\zeta}_2, \hat{\zeta}_1] (2\hat{\zeta}_2 \hat{\zeta}_2 \hat{\zeta}_1 \hat{\zeta}_{\mathbf{p}} + \hat{\zeta}_2 \hat{\zeta}_{\mathbf{p}} [\hat{\zeta}_1, \hat{\zeta}_2] + \hat{\zeta}_1 \hat{\zeta}_2 \hat{\zeta}_2 \hat{\zeta}_{\mathbf{p}}). \end{aligned} \quad (56)$$

As before, once replaced by (21), the commutator $[\hat{\zeta}_1', \hat{\zeta}_{\mathbf{p}'}]$ annihilates the prefactor with the τ_1 dependence. Inserting (56) in Eq. (54), and going back to the dimensionless power spectrum, we obtain

$$\mathcal{P}_{\zeta, [d, c]}^{1-\text{loop}}(p) \cong -M_{\text{Pl}}^2 \frac{|\Delta\eta|^2}{2} \mathcal{P}_{\zeta}^{\text{tree}}(p) \int_{\tau_s}^{\tau_e} d\tau_1 \int_{\tau_s}^{\tau_1} (a^2 \epsilon)|_{\tau_2} d\tau_2 \int \frac{d\mathbf{k}}{(2\pi)^3} \cdot \mathcal{A} \cdot 3 \cdot 8 \text{Im}(\mathcal{Z}_k \mathcal{W}_{\mathbf{q}}) \quad (57)$$

where

$$\mathcal{Z}_k = \zeta_k(\tau_2) \zeta_k^{\prime*}(\tau_1), \quad \text{and} \quad \mathcal{W}_k = \zeta_k(\tau_2) \zeta_k^*(\tau_1). \quad (58)$$

After integrating over the various momenta the prefactor in Eq. (55) becomes¹¹

$$\mathcal{A} = -\frac{1}{3} (k^2 + |\mathbf{p} - \mathbf{k}|^2 + p^2) \simeq -\frac{1}{3} (k^2 + |\mathbf{p} - \mathbf{k}|^2), \quad (59)$$

it is then easy to rewrite the final expression for this contribution as

$$\begin{aligned} \mathcal{P}_{\zeta, [d, c]}^{1-\text{loop}}(p) \cong & 4M_{\text{Pl}}^2 |\Delta\eta|^2 \mathcal{P}_{\zeta}^{\text{tree}}(p) \int_{\tau_s}^{\tau_e} d\tau_1 \int_{\tau_s}^{\tau_1} d\tau_2 a^2(\tau_2) \epsilon(\tau_2) \int \frac{d\mathbf{k}}{(2\pi)^3} (k^2 + |\mathbf{p} - \mathbf{k}|^2) \\ & \times \text{Im} \left(\zeta_k^*(\tau_1) \zeta_{|\mathbf{p}-\mathbf{k}|}^{\prime*}(\tau_1) \zeta_k(\tau_2) \zeta_{|\mathbf{p}-\mathbf{k}|}(\tau_2) \right). \end{aligned} \quad (60)$$

¹¹We would have found terms proportional to p^2 also in the previous section by including terms proportional to ζ_p' (which we always neglect) and doing integration by parts as in Eq. (52).

By summing Eq. (53) with Eq. (60), one finds the desired results:

$$\mathcal{P}_{\zeta, [c, c]}^{1\text{-loop}}(p) + \mathcal{P}_{\zeta, [d, c]}^{1\text{-loop}}(p) \cong 0, \quad (61)$$

where, as before, the symbol \cong identifies quantities up to volume suppressed terms.

4 Conclusions

We have shown that one-loop corrections on large scales from short modes which are enhanced due to a period of non-slow-roll evolution are volume suppressed by the ratio of the two scales involved in the problem. Corrections proportional to the tree-level power spectrum of the long wavelength modes appear when considering part of the computation, but exactly cancel when including all relevant diagrams. We prove our main result, i.e. Eq. (2), by using two equivalent forms of the cubic interaction Hamiltonian for the comoving curvature perturbation. From both methods we learn something. Our first approach shows that, in general, boundary terms cannot be neglected in this context. Our second method confirms the explicit cancellation of non-volume suppressed terms without the need of doing any explicit time integrals. This indicates that our result is quite general, i.e. it goes far beyond the specific setup of a sharp transition during an ultra-slow-roll phase, which was the original motivation behind this computation.

There are several possible extensions that one may envisage for future studies. Let us just list a few. By not taking the long-scale short-scale approximation, one can generalize the computations above and derive one-loop effects on all wavelengths as induced by enhanced short modes. A physical effect will likely appear around momenta corresponding to the short scales and a proper renormalization procedure should be implemented there. Further, we focus our attention on 1PI diagrams and neglect tadpoles. The latter should also be computed and properly regularized in this context. Another possible extension of our work would be to check explicitly that the same cancellation manifests itself when going at the next order in the perturbative Hamiltonian (see [24] for a first attempt on that). Moreover, as already mentioned, it would be interesting to prove the cancellation of the would be leading terms by implementing the field redefinition procedure in [35] in this frameworks. We leave all this for future investigations.

We conclude with a speculative thought. Each time in physics there is an exact cancellation, something more deep lies underneath. Here we have only shown the result of a computation, but understanding the deep reasons behind that remains an open question.

Acknowledgments

The author sincerely thanks Jaume Garriga for a first discussion on this issue and for numerous subsequent enlightening conversations. The author is also grateful to Cristiano Germani, Sadra Jazayeri and Alexandre Serantes for different and useful inputs and clarifications. The research of J.F. is supported by the State Agency for Research of the Spanish Ministry of Science and Innovation through the “Unit of Excellence María de Maeztu 2020-2023” award to the Institute of Cosmos Sciences (CEX2019-000918-M) and by grants 2021-SGR00872 and PID2019-105614GB-C22.

References

- [1] Y.B. Zel'dovich and I.D. Novikov, *The Hypothesis of Cores Retarded during Expansion and the Hot Cosmological Model*, *Soviet Astron. AJ (Engl. Transl.)*, **10** (1967) 602.
- [2] S. Hawking, *Gravitationally collapsed objects of very low mass*, *Mon. Not. Roy. Astron. Soc.* **152** (1971) 75.
- [3] B.J. Carr and S.W. Hawking, *Black holes in the early Universe*, *Mon. Not. Roy. Astron. Soc.* **168** (1974) 399.
- [4] P. Meszaros, *The behaviour of point masses in an expanding cosmological substratum*, *Astron. Astrophys.* **37** (1974) 225.
- [5] B.J. Carr, *The Primordial black hole mass spectrum*, *Astrophys. J.* **201** (1975) 1.
- [6] G.F. Chapline, *Cosmological effects of primordial black holes*, *Nature* **253** (1975) 251.
- [7] P. Ivanov, P. Naselsky and I. Novikov, *Inflation and primordial black holes as dark matter*, *Phys. Rev. D* **50** (1994) 7173.
- [8] J. Garcia-Bellido, A.D. Linde and D. Wands, *Density perturbations and black hole formation in hybrid inflation*, *Phys. Rev. D* **54** (1996) 6040 [[astro-ph/9605094](#)].
- [9] N. Tsamis and R.P. Woodard, *Improved estimates of cosmological perturbations*, *Phys. Rev. D* **69** (2004) 084005 [[astro-ph/0307463](#)].
- [10] W.H. Kinney, *Horizon crossing and inflation with large eta*, *Phys. Rev. D* **72** (2005) 023515 [[gr-qc/0503017](#)].
- [11] M.H. Namjoo, H. Firouzjahi and M. Sasaki, *Violation of non-Gaussianity consistency relation in a single field inflationary model*, *EPL* **101** (2013) 39001 [[1210.3692](#)].
- [12] J. Garcia-Bellido and E. Ruiz Morales, *Primordial black holes from single field models of inflation*, *Phys. Dark Univ.* **18** (2017) 47 [[1702.03901](#)].
- [13] C. Germani and T. Prokopec, *On primordial black holes from an inflection point*, *Phys. Dark Univ.* **18** (2017) 6 [[1706.04226](#)].
- [14] H. Motohashi and W. Hu, *Primordial Black Holes and Slow-Roll Violation*, *Phys. Rev. D* **96** (2017) 063503 [[1706.06784](#)].
- [15] G. Ballesteros and M. Taoso, *Primordial black hole dark matter from single field inflation*, *Phys. Rev. D* **97** (2018) 023501 [[1709.05565](#)].
- [16] M.P. Hertzberg and M. Yamada, *Primordial Black Holes from Polynomial Potentials in Single Field Inflation*, *Phys. Rev. D* **97** (2018) 083509 [[1712.09750](#)].
- [17] H. Motohashi, A.A. Starobinsky and J. Yokoyama, *Inflation with a constant rate of roll*, *JCAP* **09** (2015) 018 [[1411.5021](#)].
- [18] S. Inoue and J. Yokoyama, *Curvature perturbation at the local extremum of the inflaton's potential*, *Phys. Lett. B* **524** (2002) 15 [[hep-ph/0104083](#)].
- [19] K. Tzirakis and W.H. Kinney, *Inflation over the hill*, *Phys. Rev. D* **75** (2007) 123510 [[astro-ph/0701432](#)].
- [20] J. Kristiano and J. Yokoyama, *Ruling Out Primordial Black Hole Formation From Single-Field Inflation*, [2211.03395](#).

- [21] A. Riotto, *The Primordial Black Hole Formation from Single-Field Inflation is Not Ruled Out*, [2301.00599](#).
- [22] J. Kristiano and J. Yokoyama, *Response to criticism on "Ruling Out Primordial Black Hole Formation From Single-Field Inflation": A note on bispectrum and one-loop correction in single-field inflation with primordial black hole formation*, [2303.00341](#).
- [23] A. Riotto, *The Primordial Black Hole Formation from Single-Field Inflation is Still Not Ruled Out*, [2303.01727](#).
- [24] H. Firouzjahi, *One-loop Corrections in Power Spectrum in Single Field Inflation*, [2303.12025](#).
- [25] H. Firouzjahi and A. Riotto, *Primordial Black Holes and Loops in Single-Field Inflation*, [2304.07801](#).
- [26] G. Franciolini, A. Iovino, Junior., M. Taoso and A. Urbano, *One loop to rule them all: Perturbativity in the presence of ultra slow-roll dynamics*, [2305.03491](#).
- [27] G. Tasinato, *A large $|\eta|$ approach to single field inflation*, [2305.11568](#).
- [28] S.-L. Cheng, D.-S. Lee and K.-W. Ng, *Primordial perturbations from ultra-slow-roll single-field inflation with quantum loop effects*, [2305.16810](#).
- [29] D. Seery and J.E. Lidsey, *Non-Gaussian Inflationary Perturbations from the dS/CFT Correspondence*, *JCAP* **06** (2006) 001 [[astro-ph/0604209](#)].
- [30] F. Arroja and T. Tanaka, *A note on the role of the boundary terms for the non-Gaussianity in general k-inflation*, *JCAP* **05** (2011) 005 [[1103.1102](#)].
- [31] C. Burrage, R.H. Ribeiro and D. Seery, *Large slow-roll corrections to the bispectrum of noncanonical inflation*, *JCAP* **07** (2011) 032 [[1103.4126](#)].
- [32] G. Rigopoulos, *Gauge invariance and non-Gaussianity in Inflation*, *Phys. Rev. D* **84** (2011) 021301 [[1104.0292](#)].
- [33] N. Agarwal, R.H. Ribeiro and R. Holman, *Why does the effective field theory of inflation work?*, *JCAP* **06** (2014) 016 [[1311.0869](#)].
- [34] S. Garcia-Saenz, L. Pinol and S. Renaux-Petel, *Revisiting non-Gaussianity in multifield inflation with curved field space*, *JHEP* **01** (2020) 073 [[1907.10403](#)].
- [35] J.M. Maldacena, *Non-Gaussian features of primordial fluctuations in single field inflationary models*, *JHEP* **05** (2003) 013 [[astro-ph/0210603](#)].
- [36] J.S. Schwinger, *Brownian motion of a quantum oscillator*, *J. Math. Phys.* **2** (1961) 407.
- [37] R.D. Jordan, *Effective Field Equations for Expectation Values*, *Phys. Rev. D* **33** (1986) 444.
- [38] E. Calzetta and B.L. Hu, *Closed Time Path Functional Formalism in Curved Space-Time: Application to Cosmological Back Reaction Problems*, *Phys. Rev. D* **35** (1987) 495.
- [39] X. Chen, *Primordial Non-Gaussianities from Inflation Models*, *Adv. Astron.* **2010** (2010) 638979 [[1002.1416](#)].