

# A unified quasiparticle approach to the theory of strongly correlated electron liquids

V. A. Khodel,<sup>1</sup> J. W. Clark,<sup>1,2</sup> and M. V. Zverev<sup>3,4</sup>

<sup>1</sup>*McDonnell Center for the Space Sciences & Department of Physics,  
Washington University, St. Louis, MO 63130, USA*

<sup>2</sup>*Centro de Investigação em Matemática e Aplicações,  
University of Madeira, 9020-105 Funchal, Madeira, Portugal*

<sup>3</sup>*National Research Centre Kurchatov Institute, Moscow, 123182, Russia*

<sup>4</sup>*Moscow Institute of Physics and Technology, Dolgoprudny, Moscow District 141700, Russia*

Landau's quasiparticle formalism is generalized to describe a wide class of strongly correlated Fermi systems, in addition to conventional Fermi liquids. This class includes (i) so-called marginal exemplars and (ii) systems that harbor interaction-driven flat bands, in both of which manifestations of non-Fermi-liquid behavior are well documented. Specifically, the advent of such flat bands is attributed to a spontaneous topological rearrangement of the Landau state that supplements the conventional Landau quasiparticle picture with a different set of quasiparticles, the so-called fermion condensate, whose single-particle spectrum is dispersionless. The celebrated Landau-Luttinger theorem is extended to marginal Fermi liquids, in which the density of the augmented quasiparticle system is shown to coincide with the particle density. On the other hand, the total density of a system hosting an interaction-driven flat band turns out to be the sum of the densities of the two quasiparticle subsystems: the Landau-like component and the fermion condensate. We demonstrate that within the framework of the scenario proposed, a long-standing problem faced by theories of  $D$ -wave superconductivity in cuprates, namely a consistent explanation of the so-called Uemura plot, can be naturally resolved.

## 1. Introduction

The BCS paradigm, established more than half a century ago, has successfully explained the phenomenon of superconductivity discovered by Kamerlingh Onnes in 1911. This success rests upon (i) the Cooper scenario for electron pairing in metals<sup>1-3</sup> and (ii) the Landau Fermi-liquid (LFL) theory. LFL theory is a specific quasiparticle formalism designed for the normal state, in which these quasiparticles are presumed to be immortal, with a zero-temperature momentum distribution  $n_L(p) = \theta(p_F - p)$  identical to that of a perfect Fermi gas with Fermi momentum  $p_F$ <sup>4-6</sup>. Electron-phonon interactions, responsible for Cooper pairing in solids, are taken into account within the framework of a nonperturbative Migdal-Eliashberg theory of superconductivity<sup>7,8</sup>.

Subsequently, Larkin and Migdal successfully applied the BCS-LFL approach to the description of superfluid 3D liquid <sup>3</sup>He<sup>9</sup>. Their basic result is this: In conventional Fermi liquids, where, in accord with the Landau postulate, the damping of single-particle excitations is immaterial, the London diamagnetic anomaly is expressed in terms of the carrier density for an arbitrary interaction between particles. This implies that the  $T = 0$  superfluid density coincides with the carrier density, irrespective of the strength of correlations. This result is identical to the content of the Leggett theorem, derived later by him<sup>10</sup>.

The discovery in 1986 of exotic high- $T_c$  superconductivity in the strongly-correlated electron systems of copper oxides by Bednorz and Müller<sup>11,12</sup> dealt an irreparable blow to the BCS-LFL picture, when applied to these strongly correlated electron systems. BCS theory has proven to be inadequate for explanation of a wide range of experimental data on superconductivity in the so-called strong-coupling limit<sup>13</sup>, while LFL theory itself fails to explain the normal-state resistivity  $\rho(T)$  of exotic superconductors, often called strange metals (SMs), in which<sup>14,15</sup>

$$\rho(T) - \rho_0 = A_1 T. \quad (1)$$

This behavior implies that in normal states of exotic superconductors, the damping  $\gamma(T)$  of single-particle excitations varies *linearly* with  $T$ , in striking contrast with the LFL prediction  $\gamma_{\text{LFL}}(T) \propto T^2$ .

A marginal-Fermi-liquid (MFL) phenomenology introduced soon after the Bednorz-Müller discovery was the first model capable of explaining the observed non-Fermi-liquid (NFL) behavior (1). A key feature of this phenomenology is the addition to the LFL self-energy  $\Sigma_{\text{LFL}}$  of a specific *non-analytic* term<sup>16</sup>

$$\Sigma_{\text{MFL}}(\varepsilon, T) \propto \left[ \frac{2\varepsilon}{\pi} \ln \left( \frac{\max(|\varepsilon|, T)}{\varepsilon_c} \right) - i \max(|\varepsilon|, T) \right], \quad (2)$$

triggered in the original model by critical antiferromagnetic fluctuations<sup>17-20</sup>. In accord with Eq. (2), the MFL damping of single-particle excitations does indeed change linearly with energy  $\varepsilon$ , rendering explanation of the NFL

behavior (1) straightforward. However, upon adoption of Eq. (2), the quasiparticle weight  $z$  vanishes, being traditionally determined by the standard Dyson-like formula

$$z^{-1} = 1 - \left( \frac{\partial \text{Re} \Sigma(p_F, \varepsilon)}{\partial \varepsilon} \right)_{\varepsilon=0}. \quad (3)$$

Furthermore, since the first-derivative term appearing here diverges logarithmically, higher terms being even more singular, it becomes inevitable that the Landau quasiparticle formalism and MFL phenomenology are *incompatible*, fostering intense debate about the ultimate nature of basic degrees of freedom in SMs<sup>15</sup>.

In the time that has elapsed since the introduction of MFL phenomenology, it has become clear that MFL predictions are often at variance with experiment. For example, since critical fluctuations soar upward in the vicinity of a quantum critical point (QCP), so does the doping-dependent MFL coefficient  $A_1(x)$ ; yet away from it, the MFL resistivity demonstrates canonical LFL  $T^2$  behavior. In cuprates, however, experiment shows quite different behavior:  $A_1(x)$  spreads over an extended range of doping- $x$  variation, coming to naught at a value  $x_s$  where the SM and LFL states meet each other<sup>14,15,21–24</sup>.

Remarkably, in a number of heavy-fermion metals the linearity of  $\rho(T)$  is accompanied by the most spectacular exhibition of NFL behavior, associated with a gigantic enhancement of the low-temperature normal-state thermal expansion coefficient  $\alpha(T)$ <sup>25–28</sup> beyond its LFL and MFL values, which are proportional to  $T$  and  $T \ln T$  respectively. Substantial contradictions between predictions of the MFL scenario and experiment arise in the SM superconductors, where superconductivity emerges directly as an instability of the SM state<sup>14</sup>. Accordingly, as  $x$  approaches  $x_s$  from the SM side, both the critical temperature  $T_c(x)$  of termination of superconductivity and the coefficient  $A_1(x)$ , the signature of the normal SM state, decline and vanish in unison. Moreover, the MFL phenomenology fails to provide any elucidation of the famous Uemura plot of the Cooper critical temperature  $T_c(T_F)$  against the Fermi temperature  $T_F = p_F^2/2m_e$  (see below).

These profound failures render the MFL scenario for NFL behavior incomplete and vulnerable, suggesting that the theory of NFL phenomena in strongly correlated Fermi systems is hardly exhausted by this model alone. Indeed, an alternative approach to explanation of this behavior was proposed around the same time<sup>29–35</sup>. In a new scenario, NFL behavior is triggered by a specific topological rearrangement of the Landau state, called *fermion condensation* (FC). Within the FC scenario, such a topological transition leads to formation of a fermion condensate, namely an interaction-driven flat band identified as a dispersionless spectrum  $\epsilon(\mathbf{p})$  in a momentum domain  $\Omega$ . Accordingly, the FC density of states  $D(\varepsilon)$  acquires a  $\delta$ -like contribution  $\rho_{\text{FC}}\delta(\varepsilon)$ , in obvious analogy to the singular Bose-liquid term  $\rho_B\delta(\varepsilon)$  that was documented experimentally in scattering of slow neutrons in liquid  $^4\text{He}$ . This correspondence warrants use of the terms fermion condensate and fermion condensation in describing phenomena observed in strongly correlated Fermi systems in which the density of states  $D(\varepsilon)$  diverges in the momentum region  $\Omega$ .

A decade ago, a precursor of such a topological transition (TT) associated with the divergence of the density of states  $D(\varepsilon)$  was observed in the homogeneous two-dimensional electron liquid of MOSFETs<sup>36</sup>. Several years later, the presence of a fermion condensate itself was revealed in ARPES studies performed in graphene intercalated by Gd<sup>37</sup>. In addition, the merging of the spin- and valley-split Landau levels at the chemical potential in a clean strongly interacting 2D electron gas in silicon is confirmed by available experimental data<sup>38</sup>.

Given this background, an implicit question drives our immediate agenda: Is it possible to upgrade the original LFL quasiparticle formalism to equip it for description of a flattening of the single-particle spectrum beyond the TT point, which gives rise to NFL behavior inherent in cuprates that demonstrate exotic superconductivity? As will be seen, the answer to this question is affirmative.

## 2. Key features of the generic quasiparticle formalism

Extant inconsistencies between the LFL formalism and MFL phenomenology have fostered a widespread opinion that in MFLs and other strongly correlated Fermi systems, quasiparticles are simply nonexistent (see e.g.<sup>14,39</sup>). Intense debates and announcements on this subject continue. Their apotheosis, formulated in Ref.<sup>15</sup>, reads: “One thing is clear in this regime. The particle picture breaks down. ... The strange metal raises the distinct possibility that its resolution must abandon the basic building blocks of quantum theory.” However, as will be seen, there is no sound basis for such strong claims. Indeed, as revealed in results from ARPES studies and available experimental data on thermodynamic and kinetic properties, an inherent aspect of the physics of metallic-state resides in the presence of the Fermi surface (FS), inextricably related to the singular, pole-like structure of the single-particle Green function  $G(\mathbf{p}, \varepsilon)$ . Specifically, the FS is comprised of the totality of points where the trajectory  $\varepsilon = \epsilon(\mathbf{p})$  of a zero of the inverse propagator  $G^{-1}(\mathbf{p}, \varepsilon)$  crosses the energy line. This striking feature, present ubiquitously, applies not only to common metals, where LFL theory has no contenders, but also to anisotropic, strange, and “bad” metals. Variations in the structure of the distributions of zeros from one metal to another, even of quite different types, turn out to be small




FIG. 1: Zeros of the inverse single-particle propagator  $G^{-1}(p_F, \varepsilon)$ . The magenta line corresponds to the Landau Fermi liquid case with  $\gamma_{\text{LFL}}(\varepsilon) \propto \varepsilon^2$ , whereas the blue line tracks zeroes of the marginal Fermi-Liquid inverse propagator, with  $\gamma_{\text{MFL}}(\varepsilon) \propto \varepsilon$ . The red dot is related to the fermion condensate.

(see Fig. 1). Thus, as long as there exists a Fermi surface, *quasiparticles* associated with singularities of the Green function  $G(\mathbf{p}, \varepsilon)$ , remain fundamental degrees of freedom.

Any quasiparticle theory is designed to express quantitatively the low- $T$  characteristics of correlated Fermi systems in terms of those of a system of quasiparticles, interacting through phenomenological pair forces. In conventional homogeneous Fermi liquids, the procedure for introducing quasiparticles involves decomposition of the single-particle propagator  $G(p, \varepsilon) = (\varepsilon + \mu - p^2/2m - \Sigma(p, \varepsilon))^{-1}$ , with  $\mu$  being the chemical potential and  $\Sigma$  denoting the self-energy, as an analytic function of energy, into the sum<sup>5,6,9,40,41</sup>

$$G(p, \varepsilon) \equiv zG^q(p, \varepsilon) + G^r(p, \varepsilon). \quad (4)$$

The quasiparticle weight  $z$  is calculated with the aid of the traditional relation (3), while  $G^r$  represents the regular part of  $G$ . The inverse LFL quasiparticle propagator, determined as  $(G^q(p, \varepsilon))^{-1} = \varepsilon - \epsilon(p)$ , coincides with that of a perfect Fermi gas of the same density  $\rho$ . Near the Fermi surface, the quasiparticle spectrum  $\epsilon(p)$ , possessing the single root  $p = p_F$ , is expressed in terms of the self-energy  $\Sigma$  through the LFL formula

$$\epsilon(p) = z \frac{p_F(p - p_F)}{m} \left( 1 + \frac{m}{p_F} \left( \frac{\partial \Sigma(p, 0)}{\partial p} \right)_{p=p_F} \right), \quad (5)$$

where  $m$  stands for the mass of a free particle. This relation is usually written in the Landau form

$$\epsilon(p) = v_F(p - p_F) \quad (6)$$

in terms of the Fermi velocity  $v_F = p_F/m^*$  and the effective mass  $m^*$ .

An additional observation involves the damping  $\gamma_{\text{LFL}}(\varepsilon)$  of single-particle excitations, which in conventional Fermi liquids is proportional to  $\varepsilon^2$ . Since the ratio  $\gamma_{\text{LFL}}(\varepsilon)/\varepsilon$  comes to naught, the role of damping effects turns out to be immaterial, allowing LFL quasiparticles to be treated as immortal.

### 2.1. Introducing quasiparticles in marginal Fermi liquids

The standard LFL procedure summarized above fails for MFLs, since the marginal term  $\Sigma_{\text{MFL}}(\varepsilon)$  cannot be expanded in a Taylor series. However, as we will readily demonstrate, the presence of this term is not an unsurmountable obstacle to extension of the quasiparticle formalism to MFLs. Indeed, the LFL strategy of introducing quasiparticles can be preserved, provided the whole marginal contribution  $\Sigma_{\text{MFL}}(\varepsilon)$  is kept intact, while expanding in a Taylor series involves only the remaining analytic part  $\Sigma_a$  of  $\Sigma_{\text{MFL}}$ .

It is worth noting that the idea of separating off non-analytic parts of the self-energy  $\Sigma$  that defy treatment within the standard Dyson procedure is not new: it dates back to seminal articles by S. Belyaev and L. Gor'kov<sup>3,42</sup> that lay the foundation of theory of Bose liquids and superconductivity (see below). In dealing with MFLs, this strategy yields

$$G^{-1}(p \rightarrow p_F, \varepsilon \rightarrow 0) = z_a^{-1}[\varepsilon - \epsilon(p) - \Sigma_m(\varepsilon)]. \quad (7)$$

Here the quasiparticle energy  $\epsilon(p)$  is still given by Eq. (5), except that  $\Sigma$  is replaced by  $\Sigma_a$ , while a remaining marginal contribution takes the form

$$\Sigma_m(\varepsilon) = \gamma_1 \varepsilon \left[ \frac{2}{\pi} \ln(|\varepsilon|/\varepsilon_c) - i \right], \quad (8)$$

with normalization chosen to match Eq. (2). In Eq. (7) we have also introduced a refined *finite* quasiparticle weight

$$z_a^{-1} = 1 - \left( \frac{\partial \text{Re} \Sigma_a(p_F, \varepsilon)}{\partial \varepsilon} \right)_{\varepsilon=0}. \quad (9)$$

With this upgrade, the revised formula for the inverse quasiparticle propagator reads

$$(G^q(p, \varepsilon))^{-1} \equiv z_a G^{-1}(p \rightarrow p_F, \varepsilon \rightarrow 0) = \varepsilon - \epsilon(p) - \Sigma_m(\varepsilon), \quad (10)$$

revealing that at the FS, the first derivative of this function has a weak logarithmic singularity.

## 2.2. Quasiparticle structure of systems with interaction-driven flat bands

Another option for the onset of NFL behavior, including the linear-in- $T$  variation of the resistivity  $\rho(T)$ , is associated with a spontaneous violation of the topological stability of the original Landau state<sup>29,30,33,35,43–47</sup>. The quintessence of this phenomenon consists in the advent of the fermion condensate, whose spectrum  $\epsilon(\mathbf{p})$  vanishes identically in a compact region  $\Omega$  of momentum space where the condensate propagator takes the form<sup>32</sup>

$$G^q(\mathbf{p}, \varepsilon) = \frac{1 - n_*(\mathbf{p})}{\varepsilon + i\delta} + \frac{n_*(\mathbf{p})}{\varepsilon - i\delta}, \quad \mathbf{p} \in \Omega. \quad (11)$$

The condensate momentum distribution  $0 < n_*(\mathbf{p}) < 1$ , a continuous function in momentum space, is normalized by the total FC density

$$\rho_{\text{FC}} = 2 \sum_{\mathbf{p} \in \Omega} n_*(\mathbf{p}), \quad (12)$$

which in a homogeneous 3D matter becomes

$$\rho_{\text{FC}} = 2 \int_{p_i}^{p_f} n_*(p) \frac{d^3 p}{(2\pi)^3}, \quad (13)$$

where  $p_i$  and  $p_f$  are lower and upper boundaries of condensate occupation.

## 2.3. Mechanism of marginality of systems with interaction-driven flat bands.

In systems hosting flat bands, the mechanism of marginality is known to be associated with the scattering of LFL-like quasiparticles by the fermion condensate<sup>32,34,35</sup>, implying that the situation becomes radically different from the that in the MFL case, where NFL behavior is due to interaction between collective and single-particle degrees of freedom<sup>16</sup>. To discuss this crucial distinction, we borrow a textbook formula for evaluation of the damping rate  $\gamma(\varepsilon)$ . In symbolic notation adopted for brevity, it reads<sup>6,48</sup>

$$\gamma(\varepsilon) = \left( |\Gamma|^2 \text{Im} G_R \text{Im} G_R \text{Im} G_R \right). \quad (14)$$

Here  $\Gamma$  is the scattering amplitude, while  $G_R$  stands for the retarded Green function. Overwhelming contributions to the integral involved come from the vicinity of the FS, which allows us to (i) neglect contributions from regular parts of  $\text{Im} G_R$  and (ii) factor  $|\Gamma|^2$  out of this integral at the FS. The NFL contribution to the damping  $\gamma$  comes from the condensate component of  $\text{Im} G_R(\mathbf{p}, \varepsilon) \propto n_*(\mathbf{p}) \delta(\varepsilon)$ , which, in accord with Eq. (11), differs from zero solely in the FC momentum domain  $\mathbf{p} \in \Omega$ . In what follows we restrict ourselves to realistic small values of the condensate factor  $\eta = \rho_{\text{FC}}/\rho$ , so that all the propagators but one entering Eq. (14) can be replaced by the corresponding LFL-like expressions from Eq. (10). Thereupon the structure of the integrand becomes fully defined, rendering calculations routine to yield<sup>34</sup>

$$\gamma_{\text{FC}}(x, \varepsilon) \propto \eta(x) \varepsilon, \quad (15)$$

which is proportional to the first power of energy, analogously to the MFL result. However, in the scenario for fermion condensation, the coefficient  $A_1(x)$  that determines the damping of single-particle excitations has the form

$$A_1(x) \propto \eta(x). \quad (16)$$

Presence of the condensate factor  $\eta(x)$  in this formula makes a profound difference: the magnitude of  $A_1(x)$  turns out to grow smoothly with variation of the difference  $x_c - x \equiv x_s - x$  (see Fig. 3), while in the extant MFL phenomenology, the NFL factor  $A_1(x)$  is concentrated close to the QCP. We plan to compare the predictions of the both approaches with available experimental data<sup>49–52</sup> elsewhere

#### 2.4. Four topological critical points in overdoped cuprates

We are now in position to comment on the onset of fermion condensation due to violation of the topological stability of the Landau state in strange metals, taking as an illustration the family  $\text{La}_{1-x}\text{Sr}_x\text{CO}_4$  of two-dimensional cuprates, whose crystal lattice is square, as in many other high- $T_c$  superconductors. Since the analysis of problems associated with presence of the pseudogap is beyond the scope of this article, we address hereafter only the overdoped domain of their phase diagrams. At sufficiently large doping values  $x > 0.3$ , the original Landau state applies, and cuprate properties obey LFL theory. However, as the  $x$  value drops, there comes a point where fermion condensation sets in, signaled by the appearance of four new roots of the equation

$$\epsilon(\mathbf{p}_c, x_c) = 0, \quad (17)$$

which, at a critical doping value  $x_c$  in cuprates arise *simultaneously* in momentum space at four *topological critical points*  $\mathbf{p}_c$ . Such a specific degeneracy, due to the presence of a square crystal lattice, is associated with four saddle points, identified by momenta  $\mathbf{p}_1 = (0, \pi)$ ,  $\mathbf{p}_2 = (\pi, 0)$  and  $\mathbf{p}'_1 = (0, -\pi)$ ,  $\mathbf{p}'_2 = (-\pi, 0)$ <sup>30,35</sup>. Evidently, in the MFL phenomenology, where collective degrees of freedom govern the behavior, while the role of single-particle aspects is minimal, such a degeneracy plays no significant role.

In concrete calculations of topological critical points, the following equivalent condition turns out to be more convenient:

$$v(\mathbf{p}_c, x_c) = |\nabla \epsilon(\mathbf{p}, x_c)|_{\mathbf{p}_c} = 0, \quad (18)$$

where  $\nabla = \partial/\partial \mathbf{p}$  and  $v(\mathbf{p})$  is the quasiparticle group velocity. The underlying motivation is that calculations can then be carried out on the disordered side of the transition point, where the function  $v(\mathbf{p})$  is evaluated based on the Pitaevskii equation, which has the standard form<sup>6,40</sup>

$$\mathbf{v}(\mathbf{p}) = \mathbf{v}_0(\mathbf{p}) + 2 \int f(\mathbf{p}, \mathbf{l}) \nabla n(\mathbf{l}) \frac{d^2 \mathbf{l}}{(2\pi)^2}, \quad (19)$$

where  $\mathbf{v}_0(\mathbf{p}) = z\mathcal{T}^\omega(\mathbf{p})/m_e$  is determined by the  $\omega$ -limit of the vertex  $\mathcal{T}^\omega(\mathbf{p}) = \lim \mathcal{T}(\mathbf{p}, \mathbf{k} \rightarrow 0, \omega \rightarrow 0, kv_F/\omega \ll 1)$  and a phenomenological Landau interaction function  $f$  has been introduced. The advantage of applying Eq. (19) in comparison with other approaches lies in the fact that in this equation, the integration over momenta is concentrated near the Fermi line, and it is the structure of the interaction amplitude in that region which determines the behavior of the relevant phenomenological parameters. Results of numerical calculations based on this equation, which will be analyzed in detail elsewhere, demonstrate that the arrangement and configuration of the Fermi line is expressed profoundly in the damping of single-particle excitations and hence in the NFL factor  $A_1(x)$ .

### 3. Upgrading the quasiparticle description for marginal Fermi liquids

The presence in the single-particle propagator of the small term  $\Sigma_m(\varepsilon)$ , whose derivative diverges logarithmically at  $\varepsilon = 0$ , necessitates an upgrade not only of the extant procedure for introducing quasiparticles, but also with respect to incorporation of interaction effects. These tasks are performed along the same lines as in LFL theory, with the aid of two fundamental relations of many-body theory describing the response of Fermi systems to an external long-wavelength longitudinal field  $(\mathbf{pE})(\mathbf{k} \rightarrow 0, \omega \rightarrow 0)$ . It is essential that the form of the relations obtained remains independent of the type of system under consideration. Correspondingly, there are two different but complementary situations to consider. In the first, where  $\mathbf{k} = 0, \omega \neq 0$ , one deals with a so-called diagonal perturbation for which the field effect is *fictitious*, because the imposition of the field results merely in shifting the energy argument in the propagator  $G(\mathbf{p}, \varepsilon)$ . A typical instance of such a perturbation is associated with a variation of the Hamiltonian under a Galilean transformation<sup>6,40,41</sup>. Omitting details that the reader can find in textbooks, we state here the final form of the appropriate relation, taking 3D homogeneous matter as an example:

$$\frac{\partial G^{-1}(\mathbf{p}, \varepsilon)}{\partial \varepsilon} \mathbf{v}_0 = \mathbf{v}_0 + \iint \mathcal{U}(\mathbf{p}, \varepsilon, \mathbf{p}', \varepsilon') \frac{\partial G^r(\mathbf{p}', \varepsilon')}{\partial \varepsilon'} \mathbf{v}'_0 \frac{2d^3 \mathbf{p}' d\varepsilon'}{(2\pi)^4 i}. \quad (20)$$

Here  $\mathbf{v}_0 = \mathbf{p}/m_e$ , while  $\mathcal{U}$  denotes the block of Feynman diagrams for the scattering amplitude irreducible in the particle-hole channel, known to contain no quasiparticle contributions.

In the opposite limit  $k/\omega \rightarrow \infty$ , Eq. (20) is replaced by a completely different relation stemming from gauge invariance,

$$-\nabla G^{-1}(\mathbf{p}, \varepsilon) = \mathbf{v}_0 + \iint \mathcal{U}(\mathbf{p}, \varepsilon, \mathbf{p}', \varepsilon') \nabla G(\mathbf{p}', \varepsilon') \frac{2d^3 \mathbf{p}' d\varepsilon'}{(2\pi)^4 i}. \quad (21)$$

From this point, we employ for brevity a symbolic notation: round brackets imply summation and integration over all intermediate variables, supplemented by appropriate normalization factors. With this convention, Eqs. (20) and (21) take the forms

$$\begin{aligned} \frac{\partial G^{-1}}{\partial \varepsilon} \mathbf{v}_0 &= \mathbf{v}_0 + \left( \mathcal{U} \frac{\partial G^r}{\partial \varepsilon} \mathbf{v}_0 \right), \\ -\nabla G^{-1} &= \mathbf{v}_0 + \left( \mathcal{U} \nabla G \right). \end{aligned} \quad (22)$$

Following Landau's pioneering work<sup>4</sup>, we isolate quasiparticle contributions to Eq. (21) by introducing a universal scattering amplitude  $\Gamma^\omega$  determined by

$$\Gamma^\omega = \mathcal{U} + \left( \Gamma^\omega B \mathcal{U} \right) \equiv \mathcal{U} + \left( \mathcal{U} B \Gamma^\omega \right). \quad (23)$$

Here  $B \equiv \lim_{\omega \rightarrow 0} G(\mathbf{p}, \varepsilon) G(\mathbf{p}, \varepsilon + \omega)$  is, as traditionally, the regular part of the product  $GG$ , devoid of quasiparticle contributions.

To continue, we multiply the first of Eqs. (22) from the left by the product  $\Gamma^\omega B$  and integrate. Aided by Eq. (23), we then find

$$\frac{\partial G^{-1}(\mathbf{p}, \varepsilon)}{\partial \varepsilon} \mathbf{p} = \mathbf{p} + \left( \Gamma^\omega B \mathbf{p} \right). \quad (24)$$

In gaining this result, we have accounted for the obvious cancellation of two regular contributions. The first is associated with  $\partial G^r / \partial \varepsilon$ ; the second, with  $-B \partial G^{-1} / \partial \varepsilon$ .

Here we observe that Eq. (24) splits into two separate relations. The first is a standard equation containing the leading  $\varepsilon$ -independent components of the self-energy and respective components of the interaction function. The second equation, more specific, is designed to generate a logarithmic term  $\propto \ln \varepsilon$  associated with the non-analytic component  $\Sigma_m$ , whose magnitude is proportional to the small condensate density  $\rho_{\text{FC}}$ . If we are not interested in its evaluation, the latter equation can simply be omitted.

In transformation of the second of Eqs. (22), analogous operations are performed to produce

$$-\nabla G^{-1}(\mathbf{p}, \varepsilon \rightarrow 0) = \frac{\partial G^{-1}}{\partial \varepsilon} \mathbf{v}_0 - \left( \Gamma^\omega A \nabla G^{-1} \right), \quad (25)$$

where

$$A = GG - B \equiv z_a^2 G^q G^q, \quad (26)$$

yielding

$$A \nabla G^{-1}(\mathbf{p}, \varepsilon) = -z_a \left( \nabla G^q(\mathbf{p}, \varepsilon) \right). \quad (27)$$

### 3.1. Persistence of the LL theorem

We now turn to the relation between the particle and a quasiparticle numbers, their equality in conventional Fermi liquids being the essence of the celebrated Landau-Luttinger (LL) theorem<sup>6,53,54</sup>. We begin by recasting the standard formula for evaluation of the electron density  $\rho$  in terms of the Fermi system's response to a weak longitudinal electric field, already addressed briefly above,

$$\rho = \frac{N}{V} = -\frac{1}{3} \text{Re} \iint (\mathbf{p} \nabla G(\mathbf{p}, \varepsilon)) \frac{2d^3 \mathbf{p} d\varepsilon}{(2\pi)^4 i}. \quad (28)$$

In symbolic notation, Eq. (28) is decomposed into the sum  $\rho = \rho_A + \rho_B$ , with

$$\begin{aligned}\rho_A &= \text{Re}\left((\mathbf{p}\nabla G^{-1})A\right), \\ \rho_B &= \text{Re}\left((\mathbf{p}\nabla G^{-1})B\right).\end{aligned}\tag{29}$$

It is advantageous to begin the analysis with treatment of the second of these integrals. Substitution of Eq. (25) then yields  $\rho_B = \rho_B^{(1)} + \rho_B^{(2)}$ , with

$$\rho_B^{(1)} = -\text{Re}\left((\mathbf{p}\mathbf{v}_0)B\frac{\partial G^{-1}}{\partial \varepsilon}\right)\tag{30}$$

and

$$\rho_B^{(2)} = \text{Re}\left(\Gamma^\omega B(\mathbf{p}A\nabla G^{-1})\right).\tag{31}$$

We observe that the term  $(GG\partial G^{-1}/\partial \varepsilon) = -(\partial G/\partial \varepsilon)$  vanishes upon energy integration, in turn triggering the vanishing of  $\rho_B^{(1)}$ . We are then left only with  $\rho_B^{(2)}$ , which, with the aid of Eq. (24), is recast in the form

$$\rho_B^{(2)} = \text{Re}\left(\frac{\partial G^{-1}}{\partial \varepsilon}(\mathbf{p}A\nabla G^{-1})\right) - \text{Re}(\mathbf{p}A\nabla G^{-1}).\tag{32}$$

Its summation with  $\rho_A$  given by the first of Eqs. (29) produces

$$\rho = \text{Re}\left(\frac{\partial G^{-1}}{\partial \varepsilon}(\mathbf{p}A\nabla G^{-1})\right),\tag{33}$$

which, with the aid of Eq. (27), yields the relation

$$\rho = -z_a \text{Re}\left(\frac{\partial G^{-1}}{\partial \varepsilon}(\mathbf{p}\nabla G^q(\mathbf{p}, \varepsilon))\right).\tag{34}$$

At this point, we introduce a new integration variable

$$w = \varepsilon - \gamma_1 \varepsilon \left[ \frac{2}{\pi} \ln(\varepsilon/\varepsilon_c) - i \right]\tag{35}$$

in place of  $\varepsilon$ . We then have  $dw = [\partial G^{-1}(\varepsilon)/\partial \varepsilon]d\varepsilon$  and  $G^q(\mathbf{p}, w) = (w - \epsilon(\mathbf{p}))^{-1}$ , leading to the LFL integral

$$\rho = \iint \text{Im}G^q(\mathbf{p}, w) \frac{2d^3\mathbf{p} dw}{(2\pi)^4} = 2 \int \theta(p_F - p) \frac{d^3\mathbf{p}}{(2\pi)^3} = \frac{p_F^3}{3\pi^2}.\tag{36}$$

Thus, the density of the refined quasiparticles of a marginal Fermi liquid coincides with the particle density *provided* the Fermi momentum  $p_F$  is determined, analogously to the LFL case, as the point where the singularity of the single-particle Green function  $G(p, \varepsilon)$  migrates into the upper-half plane in  $\varepsilon$ .

### 3.2. MFL specific heat calculated in the unified quasiparticle formalism

With the above results in play, it is now instructive to determine whether the refined quasiparticle formalism outlined above is capable of evaluation of the MFL specific heat. To this end, we invoke the generic relation<sup>6</sup>

$$\frac{S(T \rightarrow 0)}{V} = -2 \frac{\partial}{\partial T} \left[ T \sum_{\omega_n} \int \ln \mathcal{G}(\mathbf{p}, \omega_n) \frac{d^3\mathbf{p}}{(2\pi)^3} \right],\tag{37}$$

with  $\omega_n = \pi T(2n+1)$ . In a first step, the temperature-dependent Green function  $\mathcal{G}(\mathbf{p}, \omega)$  is expressed straightforwardly in terms of the retarded Green function  $G_R$ <sup>6</sup>:

$$\frac{S(T \rightarrow 0)}{V} = \frac{1}{T} \int \frac{2d^3\mathbf{p}}{(2\pi)^4 i} \int_{-\infty}^{\infty} \varepsilon \left( -\frac{\partial f(\varepsilon)}{\partial \varepsilon} \right) \ln \left( \frac{G_R(\mathbf{p}, \varepsilon)}{G_R^*(\mathbf{p}, \varepsilon)} \right) d\varepsilon,\tag{38}$$

where  $f(\varepsilon) = [1 + \exp(\varepsilon/T)]^{-1}$ . In what follows, we retain only a logarithmically divergent part of the derivative  $dS(T \rightarrow 0)/dT$ , which comes from the pole part  $G_R^q$  entering the quantity  $\ln(G_R^q(\mathbf{p}, \varepsilon)/(G_R^q)^*(\mathbf{p}, \varepsilon))$ . Simple algebra then leads to  $S = S_+ + S_-$ , with<sup>55</sup>

$$S_{\mp}(T \rightarrow 0) \propto \int_0^\infty \frac{z^2 e^z dz}{(1 + e^z)^2} \int_0^\infty dw \tan^{-1} \left[ \frac{\gamma_1}{\pm w + \gamma_1 \ln(\epsilon_c/T)} \right]. \quad (39)$$

Integrations over  $w$  and  $z$  are then separated to yield the MFL result<sup>16–18</sup>,

$$S(T \rightarrow 0)/T \propto C(T \rightarrow 0)/T \propto \gamma_1 \ln \frac{\epsilon_c}{T}. \quad (40)$$

Thus we conclude that there is no impassable abyss between conventional Fermi liquids and marginal ones. Rather, both may be properly described within the framework of the quasiparticle formalism developed here.

#### 4. Quasiparticle formalism for systems hosting flat bands

As seen above, Fermi systems harboring interaction-driven flat bands can be treated as having two components, the first of which is the fermion condensate and the second a set of LFL-like quasiparticles, a distinctive feature being a linear-in-energy damping of single-particle excitations. Expulsion of regular components of the single-particle propagators from the generic expression for the total density  $\rho$  can be performed along the same lines as for MFLs in the derivation of Eq. (33), with the provision, in accord with Eq. (34), that the momentum integration is to be performed over two different regions in which the quasiparticle propagator  $G^q$  takes different forms. After cumbersome manipulations, we arrive at the formula

$$\rho = \rho_{\text{FC}} - \frac{1}{3} \text{Re} \iint_{p \notin \Omega} (\mathbf{p} \nabla G(\mathbf{p}, \varepsilon)) \frac{2 d^3 \mathbf{p} d\varepsilon}{(2\pi)^4 i}, \quad (41)$$

in which  $\rho_{\text{FC}}$  is given by Eq. (12). Further integrations are performed in analogy to those carried out in the preceding development to yield

$$\rho = \rho_{\text{FC}} + p_i^3/3\pi^2. \quad (42)$$

This key result provides a basis for the relation between particle and quasiparticle densities in systems having a fermion condensate.

##### 4.1. Entropy excess as a signature of systems harboring flat bands

Perhaps the most striking feature exhibited by normal states of systems that harbor a flat band is the presence of a *finite* entropy excess, which is evaluated with the aid of the combinatoric Landau-like formula<sup>32</sup>

$$S_{\text{FC}} = -2 \sum_{\mathbf{p} \in \Omega} [n_*(\mathbf{p}) \ln n_*(\mathbf{p}) + (1 - n_*(\mathbf{p})) \ln(1 - n_*(\mathbf{p}))]. \quad (43)$$

Remarkably, the behavior of this entropy excess – *being proportional to the FC density  $\rho_{\text{FC}}$*  – is in stark contradiction to that of the LFL and MFL entropies, which are proportional to  $T$  and  $T \ln T$  respectively. Analogous unorthodox behavior is exhibited by the coefficient of the thermal expansion  $\alpha(T) = l^{-1}(\partial l/\partial T) \equiv \partial S/\partial P$ , where  $P$  is the pressure. According to Eq. (43), one has

$$\alpha_{\text{FC}}(T \simeq T_c) \propto \rho_{\text{FC}}, \quad (44)$$

where  $T_c$  is the critical temperature for termination of superconductivity. The finite condensate density  $\rho_{\text{FC}}$  varies smoothly upon elevation of  $T$ .

Within the LFL and MFL scenarios, modest values of  $\alpha(T_c)$  are predicted, being proportional to  $T_c$  and  $T_c \ln T_c$  respectively, whereas the magnitude of the ratio  $\alpha_{\text{FC}}(T_c)/\alpha_{\text{LFL}}(T_c)$  turns out to be enormously enhanced, especially in heavy-fermion superconductors that commonly have very low  $T_c$  values. Curiously, it is such a huge enhancement effect that was uncovered 20 years ago in measurements performed on the three-dimensional heavy-fermion superconductor CeCoIn<sub>5</sub>, which has a tiny critical temperature  $T_c \simeq 2.3 \text{ K}$ <sup>25</sup>. These measurements demonstrated that the curve



of  $\alpha(T)$ , leaving the origin in accord with the Nernst decree  $S(T = 0) = 0$ , grows very rapidly upon elevation of  $T$ , landing at  $T \simeq T_c \ll T_D$  on a plateau whose height  $\simeq 4.0 \times 10^{-6} \text{ K}^{-1}$  exceeds the corresponding LFL result  $\alpha_{\text{LFL}}(T \simeq T_c)$  by a factor of order  $10^3$ <sup>25–27</sup>. However, this most astonishing behavior, often accompanied by exotic superconductivity (as addressed below) has still not received due attention in the condensed-matter literature to this day.

In the meantime, analogous enhancement factors have been repeatedly observed, in particular quite recently, in the celebrated heavy-fermion metal  $\text{URu}_2\text{Si}_2$ <sup>28</sup>, which undergoes a hidden second-order phase transition at  $T_c = 17.5 \text{ K}$  whose precise nature remains unknown. In this instance, the measured normal-state value of  $\alpha(T \geq T_c) \simeq 5.0 \times 10^{-6} \text{ K}^{-1}$  even exceeds that found in  $\text{CeCoIn}_5$ . Despite the almost tenfold superiority of the  $T_c$  value inherent in  $\text{URu}_2\text{Si}_2$ , the magnitude of the coefficient  $\alpha(T \simeq T_c)$  still remains almost  $T$ -independent, implying that this metal is not at all a LFL or MFL, especially in view of the fact that the corresponding value of the thermal expansion coefficient is very anisotropic<sup>28</sup>.

Concluding the analysis of this subsection, we notice that within the FC scenario and in consonance with Eqs. (15) and (43), the ratio  $A_1(x)/\alpha(T_c, x)$  must be *doping-independent*:

$$\frac{A_1(x)}{\alpha(T_c, x)} = \text{const.} \quad (45)$$

Experimental verification of this scaling behavior involving thermodynamic and kinetic characteristics would be of particular interest.

## 5. Superconducting state

Adaptation of the foregoing theoretical development to the description of superconducting systems that experience Cooper pairing is performed within the framework of the BCS concept, but it now involves *two different* single-particle Green functions  $G_s$  and  $F$ . At  $T = 0$  these obey the set of equations<sup>3,6</sup>

$$\begin{aligned} G_s(\mathbf{p}, \varepsilon) &= \left[ G^{-1}(\mathbf{p}, \varepsilon) + \Delta_0^2(\mathbf{p}) G(-\mathbf{p}, -\varepsilon) \right]^{-1}, \\ F(\mathbf{p}, \varepsilon) &= G(-\mathbf{p}, -\varepsilon) \Delta_0(\mathbf{p}) G_s(\mathbf{p}, \varepsilon). \end{aligned} \quad (46)$$

Here the superconducting gap  $\Delta_0$  is commonly assumed to be  $p$ - and  $\varepsilon$ -independent, being determined from the Gor'kov equation

$$\Delta_0(\mathbf{p}; x) = - \int \sum_{\mathbf{p}_1} \mathcal{V}(\mathbf{p}, \mathbf{p}_1) F(\mathbf{p}_1, \varepsilon_1; x) \frac{d\varepsilon_1}{2\pi i}, \quad (47)$$

where  $\mathcal{V}$  is a block of Feynman diagrams irreducible in the particle-particle channel. The same relations are valid for the quasiparticle components of  $G_s^q$  and  $F^q$  of the respective Green functions. In common metals without involvement of massive impurity-induced scattering, the set of the quasiparticle Green functions has the standard form

$$\begin{aligned} G_s^q(\mathbf{p}, \varepsilon) &= \frac{u^2(\mathbf{p})}{\varepsilon - E(\mathbf{p}) + i\delta} + \frac{v^2(\mathbf{p})}{\varepsilon + E(\mathbf{p}) - i\delta}, \\ F^q(\mathbf{p}, \varepsilon) &= \frac{\Delta_0(\mathbf{p})}{(\varepsilon - E(\mathbf{p}) + i\delta)(\varepsilon + E(\mathbf{p}) - i\delta)}, \end{aligned} \quad (48)$$

where  $u^2(\mathbf{p}) + v^2(\mathbf{p}) = 1$  and

$$n(\mathbf{p}) = v^2(\mathbf{p}) = (E(\mathbf{p}) - \epsilon(\mathbf{p}))/2E(\mathbf{p}) \quad (49)$$

involves the Bogoliubov quasiparticle energy

$$E(\mathbf{p}) = \sqrt{\epsilon^2(\mathbf{p}) + \Delta_0^2(\mathbf{p})}, \quad (50)$$

with  $\epsilon(\mathbf{p})$  given by the FL relation  $\epsilon(\mathbf{p}, n) = \delta E / \delta n(\mathbf{p})$ . Contributions of pairing terms also dependent on the momentum distribution  $n(\mathbf{p})$  should be taken into account to provide a specific  $\Delta$ -dependence of  $\epsilon(\mathbf{p})$ <sup>31</sup>.

The superconducting state is the vacuum with respect to the Bogolyubov quasiparticles, which have momentum distribution  $n_B(\mathbf{p}, T) = [1 + e^{E(\mathbf{p})/T}]^{-1}$ . Insertion of this distribution into the above formula (43) evidently yields  $S(T = 0) = 0$ , disposing of any contradiction of the Nernst theorem. As a result, the system becomes superconducting

with a critical temperature that substantially exceeds the respective BCS value. Of course, Cooper pairing is not the unique mechanism for ensuring recovery of the Nernst theorem down to zero temperature. For example, antiferromagnetism may replace pairing if the Cooper scenario is ruled out for some reason, as in strong external magnetic fields. We expect to address these pertinent issues in a sequel to this paper.

### 5.1. Equality of particle and quasiparticle numbers in superfluid Fermi liquids

In estimates of the gap value (e.g. in MFL phenomenology), the simple consideration that the number of electrons participating in Cooper pairing should be equal to the total particle number is commonly exploited. But a natural issue then arises. In Cooper systems, the standard Luttinger method for proving the LL theorem becomes inappropriate. However, in due course we will find that in systems that experience Cooper pairing, the equality between particle and quasiparticle numbers still persists.

To facilitate analysis, we first address conventional Fermi liquids such as liquid  $^3\text{He}$ . Instead of Eq. (28), one now has

$$\rho = -\frac{1}{3} \iint p_n \frac{\partial G_s(\mathbf{p}, \varepsilon)}{\partial p_n} \frac{2d^3\mathbf{p} d\varepsilon}{(2\pi)^4 i} = \frac{1}{3} \iint \mathbf{p} K_s(\mathbf{p}, \varepsilon) \nabla G^{-1}(\mathbf{p}, \varepsilon) \frac{2d^3\mathbf{p} d\varepsilon}{(2\pi)^4 i}, \quad (51)$$

where

$$K_s(\mathbf{p}, \varepsilon) = \lim_{\mathbf{k} \rightarrow 0} [G_s(\mathbf{p}, \varepsilon) G_s(\mathbf{p} + \mathbf{k}, \varepsilon) - F(\mathbf{p}, \varepsilon) F(\mathbf{p} + \mathbf{k}, \varepsilon)], \quad (52)$$

and  $p_n$  is a component of the momentum  $\mathbf{p}$  normal to the Fermi surface.

In what follows, we employ results obtained in proof of the persistence of the LL theorem in MFLs, decomposing  $K_s$  into the sum of a quasiparticle term  $K_s^q$  and a regular remainder denoted  $B_s$ .

Insertion of the decomposition  $K_s = K_s^q + B_s$  into Eq. (51) leads to

$$\rho = \rho_K + \rho_B, \quad (53)$$

in which

$$\begin{aligned} \rho_K &= \text{Re} \left( (\mathbf{p} \nabla G^{-1}) K_s^q \right), \\ \rho_B &= \text{Re} \left( (\mathbf{p} \nabla G^{-1}) B_s \right). \end{aligned} \quad (54)$$

Hereafter we shall make use of two key relations, namely

$$\frac{\partial G^{-1}(\mathbf{p}, \varepsilon)}{\partial \varepsilon} \mathbf{p} = \mathbf{p} + (\Gamma^\omega B_s \mathbf{p}), \quad (55)$$

and

$$-\nabla G^{-1}(\mathbf{p}, \varepsilon \rightarrow 0) = \frac{\partial G^{-1}}{\partial \varepsilon} \mathbf{v}_0 - (\Gamma^\omega K_s^q \nabla G^{-1}), \quad (56)$$

their derivations being analogous to those of Eq. (24) and Eq. (25), respectively.

Combining this equation with the latter expression of Eq. (51) then yields  $\rho_B = \rho_s^{(1)} + \rho_s^{(2)}$ , in which the term

$$\rho_s^{(1)} = -\text{Re} \left( (\mathbf{p} \mathbf{v}_0) B_s \frac{\partial G^{-1}}{\partial \varepsilon} \right) = -\left( \partial G_s / \partial \varepsilon \right) \quad (57)$$

vanishes. As a result, we are left with

$$\rho_B = \text{Re} \left( \Gamma^\omega B_s \mathbf{p} K_s^q \nabla G^{-1} \right). \quad (58)$$

These manipulations lead us finally to the result

$$\rho = \frac{1}{3} \iint \frac{\partial G^{-1}(\mathbf{p}, \varepsilon)}{\partial \varepsilon} p_n K_s^q(\mathbf{p}, \varepsilon) \frac{\partial G^{-1}(\mathbf{p}, \varepsilon)}{\partial p_n} \frac{2d^3\mathbf{p} d\varepsilon}{(2\pi)^4 i}. \quad (59)$$

Near the quasiparticle pole, one has  $G_s(\mathbf{p}, \varepsilon) = zG_s^q(\mathbf{p}, \varepsilon)$  and

$$K_s^q(\mathbf{p}, \varepsilon) = z^2[G_s^q(\mathbf{p}, \varepsilon)G_s^q(\mathbf{p}, \varepsilon) - F^q(\mathbf{p}, \varepsilon)F^q(\mathbf{p}, \varepsilon)], \quad (60)$$

such that all the  $z$ -factors cancel to produce

$$\rho = -\frac{2}{3} \iint p_n \frac{\partial G_s^q(\mathbf{p}, \varepsilon)}{\partial p_n} \frac{d^3 \mathbf{p} d\varepsilon}{(2\pi)^4 i} = 2 \int v^2(\mathbf{p}) \frac{d^3 \mathbf{p}}{(2\pi)^3}. \quad (61)$$

Accordingly, we have generalized the LL theorem, demonstrating coincidence between the particle and quasiparticle densities in Cooper superconductors. Extension to those electron systems that harbor interaction-driven flat bands is straightforward.

## 6. Applications to unconventional high- $T_c$ superconductivity

In this section, we address the gross structure of the  $T - x$  cuprate phase diagram, which demonstrates profound non-BCS behavior. In doing so we focus on an overdoped part of the phase diagram, because its underdoped part is subject in full force to an enigmatic pseudogap phenomenon, whose nature is beyond the scope of the present analysis.

### 6.1. Elucidation of non-BCS behavior of the Uemura plot

The famous Uemura plot<sup>56</sup> showing critical temperature  $T_c$  versus the Fermi energy  $T_F = p_F^2/2m_e$  is one of the well-established confirmations of the profound failure of the BCS-FL theory of BM superconductors. In Fig. 2, the experimental data are arranged such that logarithms of critical temperatures  $T_c$  are plotted on the vertical axis and logarithms of the corresponding Fermi energies  $T_F$ , on the horizontal axis. As seen, most of the exotic superconductors have  $T_c(\rho)/T_F$  values around 0.01-0.05, whereas all of the conventional BCS examples lie on the far right in the plot. A more vivid demonstration of the collapse of the mainstream BCS scenario and non-BCS nature of BM superconductivity is hard to find.

To understand how the presence of interaction-driven flat bands restores agreement between experiment and theory, it is helpful to recast the gap equation (47) in the standard simplified form:

$$\Delta_0(\mathbf{p}) = - \sum_{\mathbf{l}} \mathcal{V}(\mathbf{p}, \mathbf{l}) \frac{\Delta_0(\mathbf{l})}{2E(\mathbf{l})}. \quad (62)$$

The quintessence of the FC scenario lies in the fact that in systems hosting interaction-driven flat bands, overwhelming contributions to the r.h.s. of Eq. (62) come from the condensate region  $\mathbf{p} \in \Omega$  where the ratio  $\Delta_0/E(\mathbf{p})$  remains of order unity, irrespective of how small the gap value is. As a result, the condensate contributions to the r.h.s. of Eq. (62) turn out to be of order unity, whereas the respective BCS contributions vanish in the limit  $\Delta \rightarrow 0$  as  $\Delta_0 \ln(\Omega_D/\Delta_0)$ , justifying their neglect and leaving the result<sup>29</sup>

$$\frac{\Delta_0^{\text{FC}}(x)}{T_F} \propto \frac{T_c^{\text{FC}}(x)}{T_F} \simeq 0.5 g_{\text{FC}} I_0 \quad (63)$$

where

$$I_0(x) = \frac{2}{\rho} \sum_{\mathbf{p} \in \Omega} \sqrt{n_*(\mathbf{p})(1 - n_*(\mathbf{p}))} \propto \eta(x), \quad (64)$$

with  $n_*(p)$  denoting the normal-state FC momentum distribution and  $\eta = \rho_{\text{FC}}/\rho$  the dimensionless condensate density. The quantity  $g_{\text{FC}}$  is the FC pairing constant, whose magnitude can be markedly in excess of the BCS constant  $g$  (see below).

At the optimal doping value  $x_o$ , Eq. (63) becomes

$$T_c^{\text{max}}/T_F \simeq 0.5 g_{\text{FC}} \eta_o, \quad (65)$$

where  $T_c^{\text{max}}$  stands for the maximum value of the critical temperature  $T_c(x)$ , while  $\eta_o \equiv \eta(x_o) \simeq 0.1$  is a dimensionless parameter. Bearing in mind the enhancement of the ratio  $g_{\text{FC}}/g$ , we infer that within the framework of the FC scenario, the Uemura plot can be properly explained, in contrast to results obtained in the MFL phenomenology.




FIG. 2: Schematic Uemura plot showing the superconducting transition temperature  $T_c$  versus the Fermi temperature  $T_F$ <sup>56,57</sup>. Different families of unconventional superconductors are located between the two red lines, while conventional superconductors lie far to the right.

### 6.2. Critical behavior of the ratio $T_c(x)/A_1(x)$

It is noteworthy that near critical doping value  $x_c$  where the superconducting dome terminates, two completely different quantities, namely

- (i)  $A_1(x)$ , a signature of the *normal* state that identifies the linear-in- $T$  component of the resistivity  $\rho(T, x)$  (whose NFL behavior has already been explained in the Section 2.2 (cf. Eq. (16)), and
  - (ii) the critical temperature  $T_c(x)$ , a key feature of the *superconducting* state,
- turn out to be proportional to the same quantity, namely the condensate density  $\eta(x)$ , and both *come to naught in unison* at the boundary between the SM and LFL states, due to vanishing of  $\eta(x)$  at that point. Thus we observe that these two “birds,”  $A_1$  and  $T_c$ , can be “killed” with one “stone,” the role of the latter being played by density of the fermion condensate (cf. Ref.<sup>15</sup>).

### 6.3. Fermion condensate and $D$ -wave superconductivity of cuprates

The  $D$ -wave character is another and perhaps more salient feature of BM superconductivity of two-dimensional cuprates. To properly elucidate this phenomenon within the framework of the FC scenario, we need to take into account the fact that the symmetry of the crystal lattice promotes simultaneous generation of four different condensate spots, each associated with its own saddle point<sup>33,35</sup>. Thus the domain of the momentum integration in Eq. (62) is divided into four quite small condensate regions  $\mathbf{p} \in \Omega_k$  where the smooth functions  $\mathcal{V}(\phi; \mathbf{l})$  and  $\Delta(\mathbf{l})$  remain practically unchanged and therefore can be factored out of the integrands to yield<sup>59</sup>

$$\Delta(\varphi) = - \sum_k \mathcal{V}_k(\varphi) \frac{\Delta_k}{|\Delta|}, \quad (66)$$

where  $\Delta_k = \Delta(\mathbf{p}_k)$  and  $\mathcal{V}_k(\varphi) = \mathcal{V}(\varphi; \mathbf{p}_k)$ . The four constants  $\Delta_k$  are readily determined based on their definition and Eq. (66) to arrive at the relation

$$\Delta_k = -\frac{I_0}{4} \sum_l \mathcal{V}_{kl} \frac{\Delta_l}{|\Delta|}, \quad (67)$$

in which  $\mathcal{V}_{11} = \mathcal{V}(0, \pi; 0, \pi)$ ,  $\mathcal{V}_{12} = \mathcal{V}(0, \pi; \pi, 0)$ , etc. Hereafter we also employ notation  $\mathcal{V}^+ = \mathcal{V}_{12} \equiv \mathcal{V}_{1,2'}$ .

The set of equations (67) has four different solutions. The relation of our immediate interest is

$$\Delta_S(x) \simeq -\frac{\rho_{\text{FC}}(x)}{8} [\mathcal{V}_{11} + 2\mathcal{V}^+ + \mathcal{V}_{11'}], \quad (68)$$




FIG. 3: Critical temperature  $T_c$  versus doping  $x$ . As seen, at  $x$  quite lower  $x_c$ , critical temperature  $T_c(x)$  is proportional to  $x_c - x$ , while in the vicinity of the topological critical point,  $T_c(x \simeq x_c) \propto (\mathcal{V}^+)^4$ . This behavior agrees well with the experimental data for  $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$  obtained in Ref.<sup>60</sup>.

which determines the gap  $\Delta_S$  in the single-particle spectrum due to  $S$ -wave pairing, provided the expression in the square brackets has a negative sign; otherwise,  $\Delta_S \equiv 0$ . An analogous statement is valid for the  $D$ -wave gap  $\Delta_D$ , given by

$$\Delta_D(x) \simeq -\frac{\rho_{\text{FC}}(x)}{8} [\mathcal{V}_{11} - 2\mathcal{V}^+ + \mathcal{V}_{11'}]. \quad (69)$$

Importantly, it should be noted that both the quantities  $\Delta_S$  and  $\Delta_D$ , being proportional to the FC density  $\rho_{\text{FC}}$ , have nonzero values only on the ordered, topologically nontrivial side of the Lifshitz transition.

To avoid unjustified complications, we restrict ourselves to a region of the phase diagram where both  $\Delta_D$  and  $\Delta_S$  have positive values. In this case, subtracting Eq. (68) from Eq. (69) yields<sup>58</sup>

$$\Delta_D(x) - \Delta_S(x) = 0.5\rho_{\text{FC}}(x)\mathcal{V}^+. \quad (70)$$

Thus, the outcome of competition between the  $S$ -wave and  $D$ -wave pairing options depends solely on sign of the single matrix element  $\mathcal{V}^+ = \mathcal{V}(0, \pi; \pi, 0)$ , corresponding to the momentum transfer  $q \simeq \pi$  at which the electron-phonon exchange becomes immaterial. Hence, in dealing with two-dimensional electron systems of cuprates the sign of the difference (70) and its magnitude are primarily determined by the repulsive Coulomb part of the matrix element  $\mathcal{V}^+$  (cf. Eq. (65)).

By way of illustration, Fig. 3 presents results of our estimation of doping-dependence of the critical temperature  $T_c(x)$ <sup>59</sup>.

## 7. Conclusion

In the present article, a unified quasiparticle theory appropriate for the quantitative description of the challenging NFL behavior of strongly correlated Fermi systems is presented. The general framework of this theory is in fact the same as in the Landau's conception, but this extended formalism is free of shortcomings of the original LFL picture, which is suitable only for conventional Fermi liquids where damping of single-particle excitations is immaterial.

In a recent survey article<sup>15</sup>, Phillips, Hussey, and Abbamonte have raised several pertinent questions in addressing the challenging problem of strange metallicity. In doing so, the authors have proclaimed a breakdown of the quasiparticle pattern of phenomena in condensed matter in the corresponding experimental domain, advocating the introduction of non-quasiparticle states and/or anomalous dimensions. Pending analysis of the results of quantitative application of our extended quasiparticle theory to relevant experimental data on strange metals, we are nevertheless prompted to contend that in dealing conceptually with the theoretical picture advocated in Ref.<sup>15</sup>, we have a typical scenario envisioned by Occam for the action of his Razor, with the apt response: "When you hear hoofbeats, think horses, not unicorns, not zebras." In the same spirit, working within an updated quasiparticle approach, albeit mundane, we have

developed a self-consistent theory not only of strange metallicity, but indeed for diverse NFL phenomena occurring in strongly correlated electron systems.

To be more specific, we summarize below the points where the Landau quasiparticle formalism has been refined and extended, especially to cope with the advent of topological disorder<sup>61</sup>, the presence of a finite entropy excess  $S_*$  at extremely low  $T \simeq T_c$ , and the nature of the Uemura plot  $T_c(T_F)$ .

(i) We have radically altered the original Dyson-like procedure for introducing quasiparticles, in order to eliminate a fundamental drawback of the Landau formalism that prevents it from yielding a correct description of the observed non-Fermi-liquid behavior.

(ii) We have investigated the issue of equality between the densities of particles and quasiparticles in electron systems that demonstrate non-Fermi-liquid behavior. This involves an extension of the celebrated Landau-Luttinger theorem, its equality proof being applicable to conventional Fermi liquids only. In our treatment, equality has been proven for all the systems under consideration, including their superconducting phases as well, such a relation being hitherto unknown.

(iii) Within the framework of our approach, we have evaluated the non-Fermi-liquid part of the electron heat capacity. The results obtained coincide with those found in the marginal Fermi-liquid phenomenology. This implies that our formalism is in fact applicable to the *quantitative* description of these challenging quantum liquids.

(iv) On the basis of the fermion condensate picture we have explained the linear energy dependence of the quasiparticle damping rate responsible for the linear in temperature normal-state resistivity of strange metals.

(v) Within the framework of the fermion-condensate scenario, we have explained the presence of a huge non-Fermi-liquid entropy excess uncovered in measurements of the linear thermal expansion of strange metals at temperatures comparable with the critical temperatures  $T_c$  for termination of superconductivity<sup>25,28</sup>.

(vi) Within the framework of the same scenario, we have explained the Uemura plot<sup>56</sup>, in which the non-BCS nature of Cooper pairing in strange metals exhibits itself in full force.

(vii) On the same basis, we have properly explained the challenging doping dependence of the critical temperatures  $T_c(x)$  of LSCO compounds measured by A. Ino et al. and reported in Ref.<sup>60</sup>.

As the text of this article was being completed, we were made aware of a recent essay by I. Mazin<sup>62</sup>, in which the relevance of Occam's razor was raised in connection with ideas popular in modern condensed-matter physics. Our views happen to be very similar to those espoused in this essay.

In conclusion, the authors express our gratitude to S. Kivelson, P. Phillips, V. Shaginyan, C. Varma, and G. Volovik for fruitful discussions.

The work was carried out within the state assignment of NRC Kurchatov Institute.

- 
- <sup>1</sup> L. N. Cooper, Phys. Rev. **104**, 1189 (1956).  
<sup>2</sup> J. Bardeen, L. N. Cooper, J. R. Schrieffer, Phys. Rev. **106**, 162 (1957); **108**, 1175 (1957).  
<sup>3</sup> L. P. Gor'kov, Sov. Phys. JETP **7**, 505 (1958).  
<sup>4</sup> L. D. Landau, Sov. Phys. JETP **3**, 920 (1957).  
<sup>5</sup> L. D. Landau, Sov. Phys. JETP **8**, 70 (1959).  
<sup>6</sup> A. A. Abrikosov, L. P. Gor'kov, I. E. Dzyaloshinskii, *Methods of Quantum Field Theory in Statistical Physics* (Pergamon Press, Oxford, 1965).  
<sup>7</sup> A. B. Migdal, Sov. Phys. JETP **7**, 996 (1958).  
<sup>8</sup> G. M. Eliashberg, Sov. Phys. JETP **11**, 696 (1960).  
<sup>9</sup> A. I. Larkin, A. B. Migdal, Sov. Phys. JETP **17**, 1146 (1963).  
<sup>10</sup> A. J. Leggett, J. Stat. Phys. **93**, 927 (1998).  
<sup>11</sup> J. G. Bednorz, K. A. Müller, Z. Phys. B **64**, 189 (1986).  
<sup>12</sup> A. J. Leggett, Lecture Notes on Exotic Superconductivity, University of Tokyo Open Course Ware, June 2011.  
<sup>13</sup> Y. Cao et al., Nature **556**, 43 (2018); **556**, 80 (2018).  
<sup>14</sup> B. Keimer, S. A. Kivelson, M. R. Norman, S. Uchida, J. Zaanen, Nature **518**, 179 (2015).  
<sup>15</sup> P. W. Phillips, N. E. Hussey, P. Abbamonte, Science **377**, 169 (2022).  
<sup>16</sup> E. Abrahams, C. M. Varma, Proc. Natl. Acad. Sci. **97**, 5714 (2000).  
<sup>17</sup> C. M. Varma, P. B. Littlewood, S. Schmitt-Rink, E. Abrahams, A. E. Ruckenstein, Phys. Rev. Lett. **63**, 1996 (1989).  
<sup>18</sup> C. M. Varma, P. B. Littlewood, S. Schmitt-Rink, E. Abrahams, A. E. Ruckenstein, Phys. Rev. Lett. **64**, 497 (1990).  
<sup>19</sup> C. M. Varma, Rep. Prog. Phys. **79**, 082501 (2016).  
<sup>20</sup> C. M. Varma, Rev. Mod. Phys. **92**, 031001 (2020).  
<sup>21</sup> R. A. Cooper et al., Science **323**, 603 (2009).  
<sup>22</sup> H. E. Hussey et al., Phil. Trans. R. Soc. A **369**, 1626 (2011).  
<sup>23</sup> I. Bozović, X. He, J. Wu, A. T. Bollinger, Nature **536**, 309 (2016); Low Temp. Phys. **44**, 674 (2018).  
<sup>24</sup> I. Bozović, X. He, J. Wu, A. T. Bollinger, J. Supercond. Nov. Magn. **31**, 2683 (2018).

- <sup>25</sup> N. Oeschler et al., Phys. Rev. Lett. **91**, 076402 (2003).
- <sup>26</sup> J. G. Donath et al., Phys. Rev. Lett. **100**, 136401 (2008).
- <sup>27</sup> N. Oeschler et al., Physica B **403**, 1254 (2008).
- <sup>28</sup> L. Wang et al., Phys. Rev. Lett. **124**, 257601 (2020).
- <sup>29</sup> V. A. Khodel and V. R. Shaginyan, JETP Lett. **51**, 553 (1990).
- <sup>30</sup> G. E. Volovik, JETP Lett. **53**, 222 (1991).
- <sup>31</sup> P. Nozières, J. Phys. I France **2**, 443 (1992).
- <sup>32</sup> V. A. Khodel, V. V. Khodel and V. R. Shaginyan, Phys. Rep. **249**, 1 (1994).
- <sup>33</sup> G. E. Volovik, JETP Lett. **59**, 830 (1994); arXiv:1606.08318.
- <sup>34</sup> J. Dukel'sky et al., Z. Phys. B **102**, 245 (1997).
- <sup>35</sup> V. Yu. Irkhin, A. A. Katanin, M. I. Katsnelson, Phys. Rev. Lett. **89**, 076401 (2002).
- <sup>36</sup> A. Mokashi et al., Phys. Rev. Lett. **109**, 096405 (2012).
- <sup>37</sup> S. Link et al., Phys. Rev. B **100**, 121407 (R) (2019).
- <sup>38</sup> A. A. Shashkin et al., Phys. Rev. Lett. **112**, 186402 (2014).
- <sup>39</sup> D. V. Else, T. Senthil, Phys. Rev. Lett. **127**, 086601 (2021).
- <sup>40</sup> L. P. Pitaevskii, Sov. JETP **10**, 1267 (1960).
- <sup>41</sup> A. B. Migdal, *Theory of Finite Fermi Systems and Applications to Atomic Nuclei* (Wiley, New York, 1967).
- <sup>42</sup> S. T. Belyaev, Sov. Phys. JETP **7**, 289 (1958).
- <sup>43</sup> V. A. Khodel, J. W. Clark, M. V. Zverev, Phys. Rev. B **78**, 075120 (2008).
- <sup>44</sup> J. W. Clark, M. V. Zverev, V. A. Khodel, Ann. Phys. **327**, 3063 (2012).
- <sup>45</sup> V. A. Khodel, J. W. Clark, M. V. Zverev, Phys. At. Nucl. **74**, 1237 (2011).
- <sup>46</sup> G. E. Volovik, Physics-Uspekhi, **61**, 89 (2018).
- <sup>47</sup> V. A. Khodel, J. Low Temp. Phys. **191**, 14 (2018).
- <sup>48</sup> D. Pines, P. Nozières, *The Theory of Quantum Liquids*, Vol. 1, W. A. Benjamin, Inc., New York Amsterdam, 1966, paragraph 8.
- <sup>49</sup> C. Proust, L. Taillefer, Annu. Rev. Condens. Matter Physics **10**, 409 (2019).
- <sup>50</sup> C. Girod et al., Phys. Rev. B **103**, 214506 (2021).
- <sup>51</sup> A. Ataei et al., Nature Physics **18**, 1420 (2022).
- <sup>52</sup> G. Grissonnanche et al., Nature **595**, 667 (2021).
- <sup>53</sup> J. M. Luttinger, J. C. Ward, Phys. Rev. **118**, 1417 (1960).
- <sup>54</sup> J. M. Luttinger, Phys. Rev. **119**, 1153 (1960).
- <sup>55</sup> V. R. Shaginyan et al., Phys. Rev. B **86**, 085147 (2012).
- <sup>56</sup> Y. J. Uemura, J. Phys. Condens. Matter **16**, S4515 (2004).
- <sup>57</sup> A. P. Drozdov, et.al., Nature **538**, 73 (2015).
- <sup>58</sup> V. A. Khodel et al., arXiv:1505.01966.
- <sup>59</sup> V. A. Khodel, J. W. Clark, M. V. Zverev, JETP Lett. **105**, 267 (2017).
- <sup>60</sup> A. Ino et al., Phys. Rev. B **65**, 094504 (2002).
- <sup>61</sup> V. A. Khodel, J. W. Clark, M. V. Zverev, Phys. Rev. B **102**, 201108(R) (2020).
- <sup>62</sup> I. Mazin, arXiv:2204.08284.