

# On Chaplygin models in $f(G)$ gravity

Fidele Twagirayezu<sup>1</sup>, Abraham Ayirwanda<sup>1</sup>, Albert Munyeshyaka<sup>2</sup>, Solange Mukeshimana<sup>3</sup>, Joseph Ntahompagaze<sup>1</sup>, and Leon Fidele Ruganzu Uwimbabazi<sup>3</sup>

<sup>1</sup>Department of Physics, College of Science and Technology, University of Rwanda, Rwanda

<sup>2</sup>Department of Physics, Mbarara University of Science and Technology, Mbarara, Uganda

<sup>3</sup>Department of Mathematics, College of Science and Technology, University of Rwanda, Rwanda.

Correspondence: ntahompagazej@gmail.com

## Abstract

The current work treats cosmological perturbation in a mixture of standard matter, Chaplygin gas as well as Gauss-bonnet fluids using a 1+3 covariant approach in the context of modified  $f(G)$  gravity. We define the gradient variables to obtain linear perturbation equations. After scalar and redshift transformations, we consider both an original Chaplygin and generalized Chaplygin gas models under Gauss-bonnet gravity. For pedagogical purposes, the consideration of polynomial  $f(G)$  gravity model was used to solve the perturbation equations for short- and long- wavelength modes and investigate the late time evolution. The numerical solutions were obtained. The results show that the energy overdensity perturbations decay with an increase in redshift. The treatment recovers GR results under limiting cases.

*keywords:* Cosmology; Chaplygin gas; Covariant perturbations; Dark energy

*PACS numbers:* 04.50.Kd, 98.80.-k, 95.36.+x, 98.80.Cq; MSC 2020: 83F05, 83D05

## 1 Introduction

Recent studies in modern cosmology have revealed information that the present universe is undergoing a cosmic acceleration [1, 2, 3]. Such an acceleration according to standard model of cosmology is caused by a fluid with a negative pressure [4, 5, 6]. This model is built on the Einstein's theory of relativity known as General Relativity (GR), which was so far a fundamental and a successful theory of gravity for several years [7]. However, GR suffers to explain the recent observed cosmic acceleration at least without invoking the cosmological constant [8, 9, 10]. Recently, it was shown that one can consider Chaplygin gas model, a model with a negative pressure by construction, to employ it under cosmological set up. This can help to study and possibly explain the late cosmic acceleration as well as other cosmological features raised through modern observational cosmology [11]. Several works have treated the Chaplygin gas model in GR [12, 13, 14], in  $f(R)$  gravity [15, 17],  $R$  being Ricci scalar, in  $f(T)$  gravity [18],  $T$  being the torsion scalar, to name a few. It was shown that the Chaplygin gas models can contribute to the understanding of the evolution of the universe through perturbation analysis [11]. There are different types of Chaplygin gas models. For instance, the original and generalized Chaplygin gas model was treated in [19] where the scalar field contribution were discussed. Other exploration of the same model was previously done in [16, 15, 17, 20].

In the present work, a mixture of a Chaplygin gas fluid, standard matter fluid as well as

Gauss-bonnet fluid, as non-interacting fluids, is considered using a  $1+3$  covariant approach. The motivation behind the choice of  $1+3$  covariant approach is that the perturbation equations that describe the dynamical evolution of the universe and predicting the large scale structures formation in late time are easily found using that approach rather than metric perturbation. In addition to that, the advantage to consider the  $1+3$  covariant formalism over the metric formalism relies in the fact that it does not leave physical modes in the evolution of perturbations [21, 22]. The  $1+3$  covariant approach being the way of dividing the space-time into foliated hyper-surfaces and a perpendicular 4-vector-field, it was introduced by Ellis and Bruni in 1989 [23] focusing on GR. Progressively after the recent discovery of cosmic acceleration, it gained much attention to extent that it was employed to different modified gravity theories [24, 29].

The next part of this paper is organized as follows: in Section 2, the presentation of the mathematical framework where one describes the  $1+3$  approach, defines the vector gradients and linear evolution of the variables, is done. The section 3 covers the linear evolutions equations, and presents the scalar perturbation equations in redshift space. In Section 4, we presents the asymptotic description of the system with the consideration of both long-and short- wavelength modes in both dust and radiation epochs respectively and present the numerical results of the perturbation equations by considering both  $GR$  limit and the system of matter chaplygin gas and Gauss-Bonnet fluids. The section 5 is devoted to discussions and conclusions. The adopted spacetime signature is  $(-+++)$  and unless stated otherwise, we have used the convention  $8\pi G = c = 1$ , where  $G$  is the gravitational constant and  $c$  is the speed of light.

## 2 Mathematical Framework

In this section, we present the mathematical aspects to describe the cosmic evolution. In this regard, we first of all define the vector gradient variables of individual fluid and find how it evolves. The action for modified Gauss-Bonnet gravity is given as [25, 26, 30, 27, 31, 28]

$$S = \frac{1}{2} \int d^4x \sqrt{-g} \left[ R + f(G) + 2\mathcal{L}_m \right], \quad (1)$$

with Kappa ( $\kappa$ ) is assumed to equals to 1 and  $R$  is the Ricci scalar and  $G$  is the Gasuu-Bonnet parameter. The modified Einstein equation becomes

$$\begin{aligned} R_{\mu\nu} - \frac{1}{2}g^{\mu\nu}R &= T_{\mu\nu}^m + \frac{1}{2}g^{\mu\nu}f - 2f'RR^{\mu\nu} + 4f'R_\lambda^\mu R^{\nu\lambda} - 2f'R^{\mu\nu\sigma\tau}R_b^{\lambda\sigma\tau} \\ &- 4f'R^{\mu\lambda\sigma\nu}R_{\lambda\sigma} + 2R\nabla^\mu\nabla_\nu f' - 2Rg^{\mu\nu}\nabla^2 f' - 4R^{\nu\lambda}\nabla_\lambda\nabla^\mu f' \\ &- 4R^{\mu\lambda}\nabla_\lambda\nabla^\nu f' + 4R^{\mu\nu}\nabla^2 f' + 4g^{\mu\nu}R^{\lambda\sigma}\nabla_\lambda\nabla_\sigma f' - 4R^{\mu\lambda\nu\sigma}\nabla_\lambda\nabla_\sigma f', \end{aligned} \quad (2)$$

where  $f \equiv f(G)$  and  $f' = \frac{\partial f}{\partial G}$  and  $T_{\mu\nu}^m$  is the energy momentum tensor of the matter fluid (photons, baryons, cold dark matter, and light neutrinos) with  $G$  is given as  $R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\alpha\beta}R^{\mu\nu\alpha\beta}$ ,  $R^{\mu\nu}$  is the Ricci tensor and  $R^{\mu\nu\alpha\beta}$  is the Riemann tensor. The energy-momentum tensor of matter fluid forms is given by  $T_{\mu\nu} = \rho u_\mu u_\nu + p h_{\mu\nu} + q_\mu u_\nu + q_\nu u_\mu + \pi_{\mu\nu}$  where  $\rho$ ,  $p$ ,  $q_\mu$  and  $\pi_{\mu\nu}$  are the energy density, isotropic pressure, heat flux and anisotropic pressure of the fluid respectively. The quantities  $\pi_{\mu\nu}$ ,  $q^\mu$ ,  $\rho$  and  $p$  are reffered to as dynamical quantities. The quantities  $\sigma_{\mu\nu}$ ,  $\varpi_{\mu\nu}$ ,  $\theta$  and  $A_\mu$  are reffered as kinematical quantities. The consideration of standard matter fluids (dust, radiation, etc), Chaplygin gas fluid and Gauss-Bonnet contributions leads us to define the total energy density, isotropic pressures, as

$$\rho_t = \rho_m + \rho_{ch} + \rho_G, \quad p_t = p_m + p_{ch} + p_G, \quad (3)$$

where [11, 30, 31, 32, 33, 34]

$$\rho_G = \frac{1}{2}(f'G - f) - 24f''\dot{G}, \quad (4)$$

$$p_G = \frac{1}{2}(f - f'G) + \frac{G\dot{G}}{3H}f'' + 4H^2\ddot{G}f'' + 4H^2\dot{G}^2f''', \quad (5)$$

$$\rho_{ch} = \left[ A + \frac{C}{a^{3(1+\alpha)}} \right]^{\frac{1}{1+\alpha}}, \quad (6)$$

$$p_{ch} = -\frac{A}{(\rho_{ch})^\alpha}. \quad (7)$$

We assume a spatially flat Friedman-Robert-Walker(FRD) universe,

$$ds^2 = -dt^2 + a^2(dx^2 + dy^2 + dz^2), \quad (8)$$

so that the equation corresponding to the Friedmann equation is modified as

$$3H^2 = \frac{1}{2}(Gf' - f - 24\dot{G}H^3f'') + \left[ A + \frac{C}{a^{3(1+\alpha)}} \right]^{\frac{1}{1+\alpha}} + \rho_m, \quad (9)$$

$$G = 24H^2(\dot{H} + H^2), \quad (10)$$

$$R = 6(\dot{H} + 2H^2), \quad (11)$$

where  $A, C$  are arbitrary constants,  $H = \frac{\dot{a}}{a}$  is the Hubble parameter and  $a$  is the scale factor. The energy density and pressure in the modified Gauss-Bonnet gravity are presented as

$$\rho_t = 3H^2, \quad (12)$$

$$p_t = -(3H^2 + 2\dot{H}). \quad (13)$$

## 2.1 The 1+3 Covariant approach in the context of $f(G)$ gravity

The 1 + 3 covariant decomposition is a framework used in describing the linear evolution of the cosmological perturbations [35, 41]. In this approach, a fundamental observer divides space-time into hyper-surfaces and a perpendicular 4-velocity field vector where 1 + 3 indicates the number of dimensions involved in each slice [36]. That is to mean that manifold geometry of the GR is discribed in four dimensional spacs (ie,. time and space). One of the importance of the 1 + 3 covariant approach is to identity a set of covariant variables which describe the inhomogeneity and anisotropy of the universe [37]. In this context, we define a four-vector coordinate as function of cosmological time ( $x^\mu = x^\mu(\tau)$ ) that labels the comoving distance along a world-line and the corresponding velocity given as :

$$u^\mu = \frac{dx^\mu}{d\tau} \quad (14)$$

The projection tensor,  $h_{\alpha\beta}$  into the three dimensional and orthogonal to  $u^\mu$ , satisfy the following condition:

$$h_{\alpha\beta} = g_{\alpha\beta} + u_\alpha u_\beta \Rightarrow h_\beta^\alpha h_\gamma^\beta = h_\gamma^\alpha, \quad (15)$$

$$h_\alpha^\alpha = 3, h_{\alpha\beta} u^\beta = 0. \quad (16)$$

The covariant derivative of the four-velocity in terms of its kinematic quantities [38] is given by:

$$\tilde{\nabla}_a u_b = \frac{1}{3} h_{ab} \tilde{\theta} + \tilde{\sigma}_{ab} + \tilde{\omega}_{ab} - u_a \tilde{u}_b \quad (17)$$

Where  $\theta$ ,  $\tilde{\sigma}_{ab}$ ,  $\tilde{\omega}_{ab}$ ,  $\tilde{u}_b$ , are: the volume expansion, sheartensor, vorocity tensor and four-acceleration respectively. The Hubble parameter is related to  $\theta$  as  $\theta = 3H$ . Assume the fluids in our consideration are irrotational (ie.,  $\tilde{\omega}_{ab} = 0$ ) and shear-free (i.e  $\tilde{\sigma}_{ab} = 0$ ), the rate of expansion is given by the Raychaudhuri and conservation equations as:

$$\dot{\theta} = -\frac{\theta^2}{3} - \frac{1}{2}(\rho_t + p_t) + \tilde{\nabla}^a \dot{u}_a, \quad (18)$$

$$\dot{\rho} = -\theta(\rho + p), \quad (19)$$

$$\tilde{\nabla}_a p - (\rho_t + p_t) \dot{u}_a = 0. \quad (20)$$

## 2.2 General fluids description

In this part, we assume a non-interacting matter fluids with both chaplygin gas and Gauss-Bonnet fluids in the entire Universe where the growth of the energy overdensity fluctuations contribute to the large scale structure formation. We consider an homogenous and expanding (FRW) cosmological background where it is possible to define the spatial gradient of the Gauge invariant variables as  $D_a^m = \frac{a \tilde{\nabla}_a \rho_m}{\rho_m}$ ,  $Z_a = a \tilde{\nabla}^a \theta$ ,  $D_a^{ch} = \frac{a \tilde{\nabla}_a \rho_{ch}}{\rho_{ch}}$ ,  $D_a^G = \frac{a \tilde{\nabla}_a \rho_G}{\rho_G}$ ,  $\mathcal{G}_a = a \tilde{\nabla}_a G$ ,  $\mathcal{G}_a = a \tilde{\nabla}_a \dot{G}$ . The subscripts  $m$ ,  $G$  and  $ch$  stand for matter, Gauss-Bonnet fluid, Chaplygin gas fluid contributions respectively. All the defined gradient variables will be considered to develop a system of cosmological perturbation equations in the context of chaplygin-gas assisted  $f(G)$  gravity.

## 3 Linear evolution equations

Taking the first derivative with respect to time of the defined gradient variables, we get:

$$\begin{aligned} \dot{D}_m^a = & - \left( 1 + w_m + \frac{(1 + w_m) G \dot{G}}{(1 + w_t) \rho_t \theta^2} \right) Z_a + \frac{(1 + w_m) \rho_m}{(1 + w_t) \rho_t} \omega_m \theta D_m^a + \frac{(1 + w_m)}{(1 + w_t) \rho_t} w_{chp} \rho_{chp} D_{chp}^a \\ & + \frac{(1 + w_m)}{(1 + w_t) \rho_t} \rho_{chp} a \nabla^a w_{chp} + \frac{(1 + w_m)}{(1 + w_t) \rho_t} \left( \frac{1}{2} (1 - f' - G f'') + G \dot{G} f''' + \frac{\dot{G} f''}{G^2} \right) \mathcal{G}_a \\ & + \frac{(1 + w_m)}{(1 + w_t) \rho_t} \frac{G f''}{\theta} G_a \end{aligned} \quad (21)$$

$$\begin{aligned} \dot{\Delta}_{ch} = & - \left[ 1 + \frac{G \dot{G}}{\theta (1 + w_t) \rho_t} \right] (1 + w_{ch}) Z + \frac{\theta (1 + w_{ch})}{(1 + w_t) \rho_t} w_m \rho_m \Delta_m + (1 + \rho_{ch}) \frac{\theta (1 + w_{ch})}{(1 + w_t) \rho_t} w_{ch} \rho_{ch} \Delta_{ch} \\ & - a \theta \tilde{\nabla}_a w_{ch} + \frac{\theta (1 + w_{ch})}{(1 + w_t) \rho_t} \left[ \frac{1}{2} (1 - f' - G f'') + G \dot{G} f''' + \frac{\dot{G} f''}{\theta^2} \right] \mathcal{G}_a + \frac{\theta (1 + w_{ch})}{(1 + w_t) \rho_t} G f'' \mathcal{G}, \end{aligned} \quad (22)$$

$$\begin{aligned} \dot{\mathcal{G}}_a = & \left( \frac{\ddot{G}}{\dot{G}} - \frac{\ddot{G} G f''}{(1 + w_t) \rho_t \theta} \right) \mathcal{G}_a - \frac{\ddot{G} \dot{G}}{(1 + w_t) \rho_t \theta^2} Z_a - \frac{\ddot{G}}{(1 + w_t) \rho_t} w_m \rho_m D_a^m - \frac{\ddot{G} \rho_{ch} w_{ch} D_a^{ch}}{(1 + w_t) \rho_t} \\ & - \frac{\ddot{G} \rho_{ch} a \nabla_a w_{ch}}{(1 + w_t) \rho_t} - \frac{\ddot{G}}{2(1 + w_t) \rho_t} (1 - f' - G f'') \mathcal{G}_a - \frac{\ddot{G} \dot{G}}{(1 + w_t) \rho_t} (G f'' + \frac{f''}{\theta^2}) \mathcal{G}_a \end{aligned} \quad (23)$$

$$\begin{aligned}\dot{\mathcal{G}}_a = & \left(1 - \frac{\dot{G}Gf''}{(1+w_t)\rho_t\theta}\right)\mathcal{G}_a + \frac{\dot{G}^2G}{(1+w_t)\rho_t\theta^2}Z_a - \frac{\dot{G}w_m\rho_mD_a^m}{(1+w_t)\rho_t} - \frac{\dot{G}\rho_{ch}w_{ch}D_a^{ch}}{(1+w_t)\rho_t} - \frac{\dot{G}\rho_{ch}a\nabla_a w_{ch}}{(1+w_t)\rho_t} + \\ & \left[\frac{\dot{G}}{2(1+w_t)\rho_t}(f' + Gf'' - 1) - \frac{\dot{G}^2}{2(1+w_t)\rho_t}(Gf'' + \frac{f''}{\theta^2})\right]\mathcal{G}_a,\end{aligned}\quad (24)$$

$$\begin{aligned}\dot{Z}_a = & -\left[\frac{1}{2}(1+3w_m)\rho_m + \left(\frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t\right)\frac{w_m\rho_m}{(1+w_t)\rho_t}\right]D_a^m \\ & -\left[\frac{1}{2}(1+3w_{chp})\rho_{chp} - \left(\frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t\right)\frac{w_{chp}\rho_{chp}}{(1+w_t)\rho_t}\right]D_a^{chp} \\ & -\left[\frac{3}{2}\rho_{chp} - \left(\frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t\right)\frac{\rho_{chp}}{(1+w_t)\rho_t}\right]a\tilde{\nabla}_a w_{chp} \\ & +\left[1 - 3f'' - 12H^3\dot{G}f^{iv} + 12H\left(H(\ddot{G}f''' + \frac{\ddot{G}}{G}f'' + \dot{G}^2f^{iv}) + 2H^3f''\right)\right. \\ & +\left.\left[\frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t\right]\frac{1}{(1+w_t)\rho_t}\left(\frac{1}{2}(1-f'-Gf'-Gf'') + G\dot{G}f''' + \frac{\dot{G}f''}{G^2}\right)\right]\mathcal{G}_a \\ & +\left(24\dot{G}f''' - 12H^3f''' + \left(\frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t\right)\frac{Gf''}{\theta(1+w_t)\rho_t}\right)\mathcal{G}_a \\ & +\left[8H\left(f''\ddot{G} + f'''\dot{G} + 3Hf' - \frac{3}{2}f'''H\dot{G}\right) - \frac{2}{3}\theta - \left(\frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t\right)\frac{G\dot{G}}{\theta^2(1+w_t)\rho_t}\right]Z_a \\ & -\frac{1}{(1+w_t)\rho_t}\nabla^2\left[w_m\rho_mD_a^m + w_{chp}\rho_{chp}D_a^{chp} + \rho_{chp}a\tilde{\nabla}^a w_{chp}\right] \\ & -\frac{1}{(1+w_t)\rho_t}\nabla^2\left[\left(\frac{1}{2}(1-f'-Gf'') + G\dot{G}f''' + \frac{\dot{G}f''}{G^2}\right)\mathcal{G}_a + \frac{Gf''}{\theta}G_a - \frac{G\dot{G}}{\theta^2}Z_a\right],\end{aligned}\quad (25)$$

$$\begin{aligned}\dot{D}_a^G = & -\left[\frac{4H\theta}{\rho_G}\left(2H\dot{G}f''' + (2\dot{H} - 2H^2)f''\right) - \theta\left(4H^2\ddot{f}' + 4H(2\dot{H} - H^2)\dot{f}'\right)\right. \\ & \left. - H^2\dot{f}'\right)\left(\frac{Gf''}{\theta(1+w_t)\rho_t\rho_G}\right)\right]\mathcal{G}_a + 4H\left[H(\ddot{G}f''' + \frac{f''\ddot{G}}{\dot{G}} + \dot{G}^2f''') + (2\dot{H} - H^2)f''\right. \\ & +\left. \theta\left(4H^2\ddot{f}' + 4H(2\dot{H} - H^2)\dot{f}'\right)\left(\frac{\frac{1}{2} + G\dot{G}f'' + \frac{\dot{G}f'''}{\theta^2}}{(1+w_t)\rho_t\rho_G}\right)\right]\mathcal{G}_a + \frac{1}{3\rho_G}\left[\left(2H(f''\ddot{G} + f'''\dot{G}^2)\right.\right. \\ & -\left.8Hf''\dot{G}\left(\frac{G}{12H^3} + 3H\right) + (2\dot{H} - H^2)f''\dot{G} - 3\left(4H^2\ddot{f}' + 4H(2\dot{H} - H^2)\dot{f}'\right)\right. \\ & +\left.\theta(4H^2\ddot{f}' - 4H(2\dot{H} - H^2)\dot{f}')\left(\frac{G\dot{G}\theta^2}{(1+w_t)\rho_t\rho_G}\right)\right. \\ & +\left.3\theta(4H^2\ddot{f}' + 4H(2\dot{H} - H^2)\dot{f}')\left(\frac{(1-f'-Gf'')G\dot{G}}{\theta^2(1+w_t)\rho_t}\right)\right]Z_a \\ & +\frac{\theta}{\rho_G}\left(4H^2\ddot{f}' + 4H(2\dot{H} - H^2)\dot{f}'\right)D_a^G \\ & +\theta\left(-4H^2\ddot{f}' + 4H(2\dot{H} - H^2)\dot{f}'\right)\frac{w_m\rho_m(1-f'-Gf'')}{(1+w_t\rho_t\rho_G)}D_a^m \\ & +\theta\left(4H^2\ddot{f}' + 4H(2\dot{H} - H^2)\dot{f}'\right)\left(\frac{w_{ch}\rho_{ch}}{(1+w_t)\rho_t\rho_a}\right)a\tilde{\nabla}_a w_{ch}\end{aligned}\quad (26)$$

We have used

$$\dot{u}_a = -\frac{\tilde{\nabla}^a p_t}{\rho_t + p_t} \quad (27)$$

$$a\dot{u}_a = -\frac{1}{(1+w_t)\rho_t} [w_m \rho_m D_m^a + w_{chp} \rho_{chp} D_{chp}^a + a \rho_{chp} \tilde{\nabla}^a w_{chp} + \tilde{\nabla}^a p_G], \quad (28)$$

These linear vector evolution equations (eq. 21-26) can be represented in redshift space after applying scalar and harmonic decomposition methods. It can be shown that the scalar perturbation equations for the matter, Chaplygin gas and Gauss-Bonnet energy densities in redshift space evolve as [42, 43, 34, 41]

$$\begin{aligned} \Delta'_m = & \frac{1}{(1+z)H} \left( 1 + w_m + \frac{(1+w_m)}{(1+w_t)\rho_t} \frac{G(-(1+z)H)G'}{\theta^2} \right) Z \\ & - \frac{1}{(1+z)H} \frac{(1+w_m)}{(1+w_t)} \frac{\rho_m}{\rho_t} \omega_m \theta \Delta_m + \frac{1}{(1+z)H} \alpha \frac{(1+w_m)}{(1+w_t)\rho_t} w_{chp} \rho_{chp} \Delta_{chp} \\ & + \frac{(1+w_m)}{(1+w_t)\rho_t} \frac{Gf''}{\theta} \mathbf{G} - \frac{1}{(1+z)H} \frac{(1+w_m)}{(1+w_t)\rho_t} \left( \frac{1}{2}(1-f'-Gf'')G(1+z)HG'f''' \right. \\ & \left. - \frac{(1+z)HG'f''}{G^2} \right) \mathcal{G} \end{aligned}$$

$$\begin{aligned} Z' = & -\frac{1}{(1+z)H} \left[ -\frac{1}{2}(1+3w_m)\rho_m - \left( \frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t \right) \frac{w_m \rho_m}{(1+w_t)\rho_t} - \frac{w_m \rho_m}{(1+w_t)\rho_t} \frac{k^2}{a^2} \right] \Delta_m + \\ & -\frac{1}{(1+z)H} \left[ -\frac{1}{2}(1+3w_{chp})\rho_{chp} + \left( \frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t \right) \frac{w_{chp} \rho_{chp}}{(1+w_t)\rho_t} \right. \\ & \left. \left[ \frac{3}{2}\rho_{chp} - \left( \frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t \right) \frac{\rho_{chp}}{(1+w_t)\rho_t} \right] (\alpha+1)w_{chp} - \frac{(w_{chp}\rho_{chp} - (\alpha+1)w_{chp})}{(1+w_t)\rho_t} \frac{k^2}{a^2} \right] \Delta_{chp} \\ & -\frac{1}{(1+z)H} \left[ 1 - 3f'' - 12H^3 b_1 f^{iv} + 12H \left( H(b_2 f''' + \frac{b_3}{b_1} f'' + b_1^2 f^{iv}) + 2H^3 f'' \right) \right. \\ & + \left( \frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t \right) \frac{1}{(1+w_t)\rho_t} \left( \frac{1}{2}(1-f'-Gf'-Gf'') + Gb_1 f''' + \frac{b_1 f''}{G^2} \right) \\ & \left. - \frac{1}{(1+w_t)\rho_t} \frac{k^2}{a^2} \left( \frac{1}{2}(1-f'-Gf'') + Gb_1 f''' + \frac{b_1 f''}{G^2} \right) \right] \mathcal{G} \\ & -\frac{1}{(1+z)H} \left( 24b_1 f''' - 12H^3 f''' + \left( \frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t \right) \frac{Gf''}{\theta(1+w_t)\rho_t} - \frac{1}{(1+w_t)\rho_t} \frac{k^2}{a^2} \frac{Gf''}{\theta} \right) \mathbf{G} \\ & -\frac{1}{(1+z)H} \left[ 8H \left( f'' b_2 + f''' b_1^2 + 3Hf' - \frac{3}{2} f''' H b_1 \right) - \frac{2}{3}\theta - \left( \frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t \right) \frac{Gb_1}{\theta^2(1+w_t)\rho_t} \right. \\ & \left. + \frac{k^2}{a^2(1+w_t)\rho_t} \frac{Gb_1}{\theta^2} \right] Z \end{aligned} \quad (29)$$

$$(30)$$

$$\begin{aligned}\mathcal{G}' = & -\frac{1}{(1+z)H} \left(1 - \frac{b_1 G f''}{(1+w_t)\rho_t \theta}\right) \mathbf{G} - \frac{1}{(1+z)H} \frac{b_1^2 G}{(1+w_t)\rho_t \theta^2} Z + \frac{1}{(1+z)H} \frac{b_1 w_m \rho_m \Delta_m}{(1+w_t)\rho_t} \\ & - \frac{1}{(1+z)H} \frac{\alpha b_1 \rho_{ch} w_{ch}}{(1+w_t)\rho_t} \Delta_{ch} + \left( \frac{1}{2(1+z)H} \frac{b_1}{(1+w_t)\rho_t} (f' + G f'' - 1) \right. \\ & \left. + \frac{1}{2(1+z)H} \frac{b_1^2}{(1+w_t)\rho_t} \left( G f'' + \frac{f''}{\theta^2} \right) \right) \mathcal{G}\end{aligned}\quad (31)$$

$$\begin{aligned}\mathbf{G}' = & -\frac{1}{(1+z)H} \left( \frac{b_3}{b_1} - \frac{b_1 G f''}{(1+w_t)\rho_t \theta} \right) \mathbf{G} + \left[ \frac{1}{(1+z)H} \frac{b_2 b_1 G}{(1+w_t)\rho_t \theta^2} \right] Z + \left[ \frac{1}{(1+z)H} \frac{b_2}{(1+w_t)\rho_t} w_m \rho_m \right] \Delta_m \\ & - \left[ \frac{1}{(1+z)H} \frac{\alpha b_2 \rho_{ch} w_{ch}}{(1+w_t)\rho_t} \right] \Delta_{ch} - \frac{1}{(1+z)H} \left[ \frac{b_2}{2(1+w_t)\rho_t} (f' + G f'' - 1) - \frac{b_2 b_1}{(1+w_t)\rho_t} \left( G f'' + \frac{f''}{\theta^2} \right) \right] \mathcal{G}\end{aligned}\quad (32)$$

$$\begin{aligned}\Delta'_G = & -\frac{1}{(1+z)H} \left[ -\frac{4H\theta}{\rho_G} \left( 2H b_1 f''' + (2c_1 - 2H^2) f'' \right) \right. \\ & \left. + \theta \left( 4H^2 f'' b_2 + 4H(2c_1 - H^2) f'' b_1 \right) \left( \frac{G f''}{\theta(1+w_t)\rho_t \rho_G} \right) \right] \mathbf{G} \\ & - \frac{1}{(1+z)H} \left[ 4H \left( H(b_2 f''' + \frac{f'' b_2}{b_2} + b_2^2 f''') + (2c_1 - H^2) f'' \right) \right. \\ & \left. + \theta \left( 4H^2 (f'' b_2 + f''' b_1^2) + 4H(2c_1 - H^2) f'' b_1 \left( \frac{\frac{1}{2} + G b_1 f'' + \frac{b_1 f'''}{\theta^2}}{(1+w_t)\rho_t \rho_G} \right) \right) \right] \mathcal{G} + \\ & - \frac{1}{3((1+z)H)\rho_G} \left[ \left( 2H(f'' b_2 + f''' b_1^2) - 8H f'' b_1 \left( \frac{G}{12H^3} + 3H \right) \right. \right. \\ & \left. \left. + (2c_1 - H^2) f'' b_1 - 3 \left( 4H^2 (f'' b_2 + f''' b_1^2) + 4H(2c_1 - H^2) f'' b_1 \right) \right. \right. \\ & \left. \left. + \theta (4H^2 (f'' b_2 + f''' b_1^2) - 4H(2c_1 - H^2) f'' b_1) \left( \frac{G b_1 \theta^2}{(1+w_t)\rho_t \rho_G} \right) \right. \right. \\ & \left. \left. + 3\theta (4H^2 (f'' b_2 + f''' b_1^2) + 4H(2c_1 - H^2) f'' b_1) \left( \frac{(1-f'-G f'') G b_1}{\theta^2 (1+w_t)\rho_t} \right) \right] Z \\ & - \frac{3}{(1+z)\rho_G} \left( 4H^2 (f'' b_2 + f''' b_1^2) + 4H(2c_1 - H^2) f'' b_1 \right) \Delta_G \\ & - \frac{3}{(1+z)} \left( -4H^2 (f'' b_2 + f''' b_1^2) + 4H(2c_1 - H^2) f'' b_1 \right) \frac{w_m \rho_m (1-f'-G f'')}{(1+w_t \rho_t \rho_G)} \Delta_m \\ & + \frac{3}{(1+z)} \left( 4H^2 (f'' b_2 + f''' b_1^2) + 4H(2c_1 - H^2) f'' b_1 \right) \left( \frac{w_{ch} \rho_{ch}}{(1+w_t)\rho_t \rho_G} \right) w_{ch} (\alpha + 1) \Delta_{chp} \quad (33) \\ \Delta'_{ch} = & \frac{1}{(1+z)H} \left[ 1 + \frac{G b_1}{\theta(1+w_t)\rho_t} \right] (1+w_{ch}) Z - \frac{1}{(1+z)H} \frac{\theta(1+w_{ch})}{(1+w_t)\rho_t} w_m \rho_m \Delta_m \\ & - \frac{3}{(1+z)} \left[ (1+\rho_{ch}) \frac{(1+w_{ch})}{(1+w_t)\rho_t} w_{ch} \rho_{ch} - (1+\alpha) w_{ch} \right] \Delta_{ch} \\ & - \frac{\theta(1+w_{ch})}{(1+z)(1+w_t)H\rho_t} \left[ \frac{1}{2} (1-f'-G f'') + G b_1 f'' + \frac{b_1 f'''}{\theta^2} \right] \mathcal{G} \\ & - \frac{\theta(1+w_{ch}) G f''}{(1+z)(1+w_t)H\rho_t} \mathbf{G},\end{aligned}\quad (34)$$

with  $k = \frac{2\pi a}{\lambda}$ ,  $k$  being the wave number and  $\lambda$ , the wavelength of perturbations. The parameters  $b_1$ ,  $b_2$ ,  $b_3$  and  $c_1$  are presented in Appendix. According to eq. 7, the vector term  $a \tilde{\nabla}_a w_{ch}$  presents

its scalar part as  $a\tilde{\nabla}_a w_{ch} = -w_{ch}(1 + \alpha)\Delta_{ch}$ . In GR limits, with normal form of matter, one can obtain a closed system of first-order perturbation equations which is easier to find the analytical solutions. However, the linear perturbation equations are coupled system of first-order equations for the density fluctuations of matter, chaplygin and Gauss-Bonnet fluids which are more complicated to find the analytical solutions. We therefore considered short wavelength( $\frac{k^2}{a^2 H^2} \gg 1$ ) and long wavelength( $\frac{k^2}{a^2 H^2} \ll 1$ ) limits of the perturbation to analyse the large scale structure implications of the numerical results in the redshift space.

## 4 Asymptotic Analysis

In this section, we analyse the evolution of the perturbation equations in both long-wavelength and short-wavelength regimes together with the consideration that the universe is dominated by dust.

### 4.1 Matter density fluctuations in GR limits

In this part, we analyse the behavior of energy overdensity fluctuations for dust fluid in GR limits for both long and short wavelength modes for the case  $f(G) = G$  and no contribution from the chaplygin gas fluid. We also define the normalised energy density contrast for energy density fluid as

$$\delta(z) = \frac{\Delta_m^k(z)}{\Delta(z_0)},$$

where  $\Delta(z_0)$  is the matter energy density at the initial redshift, hereafter  $z_0 = 1100$ . If we assume that the Universe is dominated mainly by dust fluid, the equation of state parameter becomes  $w = 0$ . Consequently eq. 29 through to eq. 34 become

$$\Delta'_m = \frac{1}{(1+z)H}Z, \quad (35)$$

$$Z' = \frac{\rho_d}{2(1+z)H}, \quad (36)$$

$$\mathcal{G}' = 0, \mathcal{G}' = 0, \Delta'_G = 0, \Delta'_{ch} = 0, \quad (37)$$

which admit the numerical solutions presented in fig. 1 for dust dominated universe. The matter energy overdensity fluctuations  $\delta(z)$  decay with increase in redshift. Throughout all the plots, we rescaled the  $\delta(z)$  to make it readable.

### 4.2 Energy overdensity fluctuations in the context of $f(G)$ gravity approach

For pedagogical purpose, we consider a polynomial  $f(G)$  model presents in [34] given by

$$f(G) = \frac{\alpha_1}{G^2} + \alpha_2 G^{\frac{1}{2}}, \quad (38)$$

We choose this  $f(G)$  model for a quantitative analysis of the derived perturbation equations eq. 29 through to eq. 34, it is a viable model which is compatible with cosmological observations and it is a viable model accounts for the late-time acceleration of the universe without the need for dark energy. The Hubble parameter in redshift space for a dust dominated universe ( $w = 0$ ) is






Figure 1: Plot of matter energy overdensity perturbations versus redshift of Eq. 35 using eq. 36-37 for GR limit ( $f(G) = G$ ) for a dust( $w = 0$ ) dominated Universe for  $\Delta_{in} = 10^{-5}$ ,  $Z_{in} = 10^{-5}$ .

presented as  $H(z) = \frac{2m}{3}(1+z)^{\frac{3}{2m}}$ , for GR limits, we set  $m = 1$ . The Gauss-Bonnet term is given by  $G = 24H^2 \left( H^2 + \dot{H} \right)$  which in redshift space is presented as  $G = \frac{64}{9}m^3 \left( \frac{2m}{3} - 1 \right) (1+z)^{\frac{6}{m}}$ . We assume that the dynamics of the universe is driven by the power-law scale factor of the form  $a(t) = t^{\frac{2m}{3(1+w)}}$  and we consider  $H = \frac{\dot{a}}{a}$ . We analyse the perturbation equations in both short- and long- wavelength modes.

#### 4.2.1 Short-Wavelength mode

In this subsection, we first consider the short-wavelength regime where  $\frac{k^2}{a^2} \gg 1$ , with  $k = \frac{2\pi a}{\lambda}$  and  $\lambda$  is the wavelength of the mode. Applying this approximation and considering the dust dominated universe, it means  $w_m = 0$ , eq. 29 through to eq. 34 are represented as

$$\begin{aligned} \Delta'_m = & \frac{1}{(1+z)H} \left( \left( 1 - \frac{1}{(1+w_t)\rho_t} \right) \frac{(1+z)HGG'}{\theta^2} \right) Z \\ & + \frac{1}{(1+z)H} \alpha \frac{1}{(1+w_t)\rho_t} w_{chp} \rho_{chp} \Delta_{chp} \\ & + \frac{1}{(1+z)H} \frac{1}{(1+w_t)\rho_t} \frac{Gf''}{\theta} \mathbf{G} - \frac{1}{(1+z)H} \frac{1}{(1+w_t)\rho_t} \left( \frac{1}{2} (1 - f' - Gf'') G (1+z) H G' f''' \right. \\ & \left. - \frac{(1+z)H G' f''}{G^2} \right) \mathcal{G} \end{aligned} \quad (39)$$

$$\begin{aligned}
Z' = & \frac{1}{(1+z)H} \left( \frac{1}{2} \rho_d + \frac{1}{3} \theta^2 \right) \Delta_m \\
& - \frac{1}{(1+z)H} \left[ -\frac{1}{2} (1+3w_{chp}) \rho_{chp} + \left( \frac{1}{3} \theta^2 + \frac{1}{2} (1+3w_t) \rho_t \right) \frac{w_{chp} \rho_{chp}}{(1+w_t) \rho_t} \right. \\
& \left. \left[ \frac{3}{2} \rho_{chp} - \left( \frac{1}{3} \theta^2 + \frac{1}{2} (1+3w_t) \rho_t \right) \frac{\rho_{chp}}{(1+w_t) \rho_t} \right] (\alpha+1) w_{chp} - \frac{(w_{chp} \rho_{chp} - (\alpha+1) w_{chp})}{(1+w_t) \rho_t} \frac{k^2}{a^2} \right] \Delta_{chp} \\
& - \frac{1}{(1+z)H} \left[ 1 - 3f'' - 12H^3 b_1 f^{iv} + 12H \left( H(b_2 f''' + \frac{b_3}{b_1} f'' + b_1^2 f^{iv}) + 2H^3 f'' \right) \right. \\
& + \left( \frac{1}{3} \theta^2 + \frac{1}{2} (1+3w_t) \rho_t \right) \frac{1}{(1+w_t) \rho_t} \left( \frac{1}{2} (1-f' - Gf' - Gf'') + Gb_1 f''' + \frac{b_1 f''}{G^2} \right) \\
& \left. - \frac{1}{(1+w_t) \rho_t} \frac{k^2}{a^2} \left( \frac{1}{2} (1-f' - Gf'') + Gb_1 f''' + \frac{b_1 f''}{G^2} \right) \right] \mathcal{G} \\
& - \frac{1}{(1+z)H} \left( 24b_1 f''' - 12H^3 f''' + \left( \frac{1}{3} \theta^2 + \frac{1}{2} (1+3w_t) \rho_t \right) \frac{Gf''}{\theta(1+w_t) \rho_t} - \frac{1}{(1+w_t) \rho_t} \frac{k^2}{a^2} \frac{Gf''}{\theta} \right) \mathbf{G} \\
& - \frac{1}{(1+z)H} \left[ 8H \left( f'' b_2 + f''' b_1^2 + 3Hf' - \frac{3}{2} f''' H b_1 \right) - \frac{2}{3} \theta - \left( \frac{1}{3} \theta^2 + \frac{1}{2} (1+3w_t) \rho_t \right) \frac{Gb_1}{\theta^2 (1+w_t) \rho_t} \right. \\
& \left. + \frac{k^2}{a^2 (1+w_t) \rho_t} \frac{Gb_1}{\theta^2} \right] Z
\end{aligned} \tag{40}$$

$$(41)$$

$$\begin{aligned}
\mathcal{G}' = & -\frac{1}{(1+z)H} \left( 1 - \frac{b_1 G f''}{(1+w_t) \rho_t \theta} \right) \mathbf{G} - \frac{1}{(1+z)H} \frac{b_1^2 G}{(1+w_t) \rho_t \theta^2} Z \\
& - \frac{1}{(1+z)H} \frac{\alpha b_1 \rho_{ch} w_{ch}}{(1+w_t) \rho_t} \Delta_{ch} + \left( \frac{1}{2(1+z)H} \frac{b_1}{(1+w_t) \rho_t} (f' + Gf'' - 1) \right. \\
& \left. + \frac{1}{2(1+z)H} \frac{b_1^2}{(1+w_t) \rho_t} \left( Gf'' + \frac{f''}{\theta^2} \right) \right) \mathcal{G}
\end{aligned} \tag{42}$$

$$\begin{aligned}
\mathbf{G}' = & -\frac{1}{(1+z)H} \left( \frac{b_3}{b_1} - \frac{b_1 G f''}{(1+w_t) \rho_t \theta} \right) \mathbf{G} + \left[ \frac{1}{(1+z)H} \frac{b_2 b_1 G}{(1+w_t) \rho_t \theta^2} \right] Z - \left[ \frac{1}{(1+z)H} \frac{\alpha b_2 \rho_{ch} w_{ch}}{(1+w_t) \rho_t} \right] \Delta_{ch} \\
& - \frac{1}{(1+z)H} \left[ \frac{b_2}{2(1+w_t) \rho_t} (f' + Gf'' - 1) - \frac{b_2 b_1}{(1+w_t) \rho_t} \left( Gf'' + \frac{f''}{\theta^2} \right) \right] \mathcal{G}
\end{aligned} \tag{43}$$

$$\begin{aligned}
\Delta'_G = & -\frac{1}{(1+z)H} \left[ -\frac{4H\theta}{\rho_G} \left[ 2Hb_1f''' + (2c_1 - 2H^2)f'' \right] \right. \\
& + \theta \left( 4H^2f''b_2 + 4H(2c_1 - H^2)f''b_1 \right) \left( \frac{Gf''}{\theta(1+w_t)\rho_t\rho_G} \right) \Big] G \\
& - \frac{1}{(1+z)H} \left[ 4H \left[ H(b_2f''' + \frac{f''b_2}{b_2} + b_2^2f''') + (2c_1 - H^2)f'' \right] \right. \\
& + \theta \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \left( \frac{\frac{1}{2} + Gb_1f'' + \frac{b_1f'''}{\theta^2}}{(1+w_t)\rho_t\rho_G} \right) \right) \Big] \mathcal{G} + \\
& - \frac{1}{3((1+z)H)\rho_G} \left[ \left( 2H(f''b_2 + f'''b_1^2) - 8Hf''b_1 \left( \frac{G}{12H^3} + 3H \right) \right. \right. \\
& + (2c_1 - H^2)f''b_1 - 3 \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \right) \\
& + \theta(4H^2(f''b_2 + f'''b_1^2) - 4H(2c_1 - H^2)f''b_1) \left( \frac{Gb_1\theta^2}{(1+w_t)\rho_t\rho_G} \right) \\
& \left. \left. + 3\theta(4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1) \left( \frac{(1-f'-Gf'')Gb_1}{\theta^2(1+w_t)\rho_t} \right) \right] Z \right. \\
& - \frac{3}{(1+z)\rho_G} \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \right) \Delta_G \\
& + \frac{3}{(1+z)} \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \right) \left( \frac{w_{ch}\rho_{ch}}{(1+w_t)\rho_t\rho_G} \right) w_{ch}(\alpha+1)\Delta_{chp} \quad (44) \\
\Delta'_{ch} = & \frac{1}{(1+z)H} \left[ 1 + \frac{Gb_1}{\theta(1+w_t)\rho_t} \right] (1+w_{ch})Z \\
& - \frac{3}{(1+z)} \left[ (1+\rho_{ch}) \frac{(1+w_{ch})}{(1+w_t)\rho_t} w_{ch}\rho_{ch} - (1+\alpha)w_{ch} \right] \Delta_{ch} - \\
& \frac{\theta(1+w_{ch})}{(1+z)(1+w_t)H\rho_t} \left[ \frac{1}{2}(1-f'-Gf'') + Gb_1f'' + \frac{b_1f'''}{\theta^2} \right] \mathcal{G} - \frac{\theta(1+w_{ch})Gf''}{(1+z)(1+w_t)H\rho_t} G \quad (45)
\end{aligned}$$

The energy density for the chaplygin gas fluid  $\Delta_{ch}$  and the energy density for Gauss-Bonnet fluid  $\Delta_G$  do not couple with the matter energy density  $\Delta_m$ . Numerical solutions of eq. 39 through to eq. 45 are presented in fig. 2 for different values of parameter  $m$ . For simplicity we set  $f(G)$  to  $f$  and considered the initial conditions  $\mathcal{G}_{in} = 10^{-5}$ ,  $\mathbf{G}_{in} = 10^{-5}$  and  $\Delta_{in} = 10^{-5}$  to find numerical solutions. The equations above are presented for the generalized Chaplygin gas model. During numerical results computation, where there is  $w_t$  and  $\rho_t$  terms, we replaced these terms by relations of eq. 3-7 and eq. 12-13 throughout all equations. The original Chaplygin gas model can be obtained by setting  $\alpha = 1$  in eq. 39 through to eq. 45 and its evolution equations are presented as

$$\begin{aligned}
\Delta'_m = & \frac{1}{(1+z)H} \left( \left( 1 - \frac{1}{(1+w_t)\rho_t} \right) \frac{(1+z)HGG'}{\theta^2} \right) Z + \frac{1}{(1+z)H} \frac{1}{(1+w_t)\rho_t} w_{chp}\rho_{chp}\Delta_{chp} \\
& + \frac{1}{(1+z)H} \frac{1}{(1+w_t)\rho_t} \frac{Gf''}{\theta} G - \frac{1}{(1+z)H} \frac{1}{(1+w_t)\rho_t} \left( \frac{1}{2}(1-f'-Gf'')G(1+z)HG'f'' \right. \\
& \left. - \frac{(1+z)HG'f''}{G^2} \right) \mathcal{G} \quad (46)
\end{aligned}$$




Figure 2: Plot of energy density perturbations versus redshift of Eq. 39-45 for different values of  $m$  in the dust dominated Universe for a short wavelength mode  $k = 100$  and  $\alpha = 0.2$ .

$$\begin{aligned}
Z' = & \frac{1}{(1+z)H} \left( \frac{1}{2}\rho_d + \frac{1}{3}\theta^2 \right) \Delta_m \\
& - \frac{1}{(1+z)H} \left[ -\frac{1}{2}(1+3w_{chp})\rho_{chp} + \left( \frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t \right) \frac{w_{chp}\rho_{chp}}{(1+w_t)\rho_t} \right. \\
& 2 \left[ \frac{3}{2}\rho_{chp} - \left( \frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t \right) \frac{\rho_{chp}}{(1+w_t)\rho_t} \right] w_{chp} - \frac{(w_{chp}\rho_{chp} - 2w_{chp})}{(1+w_t)\rho_t} \frac{k^2}{a^2} \left. \right] \Delta_{ch} \\
& - \frac{1}{(1+z)H} \left[ 1 - 3f'' - 12H^3b_1f^{iv} + 12H \left( H(b_2f''' + \frac{b_3}{b_1}f'' + b_1^2f^{iv}) + 2H^3f'' \right) \right. \\
& + \left( \frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t \right) \frac{1}{(1+w_t)\rho_t} \left( \frac{1}{2}(1-f' - Gf' - Gf'') + Gb_1f''' + \frac{b_1f''}{G^2} \right) \\
& - \frac{1}{(1+w_t)\rho_t} \frac{k^2}{a^2} \left( \frac{1}{2}(1-f' - Gf'') + Gb_1f''' + \frac{b_1f''}{G^2} \right) \left. \right] \mathcal{G} \\
& - \frac{1}{(1+z)H} \left( 24b_1f''' - 12H^3f''' + \left( \frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t \right) \frac{Gf''}{\theta(1+w_t)\rho_t} - \frac{1}{(1+w_t)\rho_t} \frac{k^2}{a^2} \frac{Gf''}{\theta} \right) \mathbf{G} \\
& - \frac{1}{(1+z)H} \left[ 8H \left( f''b_2 + f'''b_1^2 + 3Hf' - \frac{3}{2}f'''Hb_1 \right) - \frac{2}{3}\theta - \left( \frac{1}{3}\theta^2 + \frac{1}{2}(1+3w_t)\rho_t \right) \frac{Gb_1}{\theta^2(1+w_t)\rho_t} \right. \\
& + \left. \frac{k^2}{a^2(1+w_t)\rho_t} \frac{Gb_1}{\theta^2} \right] Z \\
\end{aligned} \tag{47}$$

$$\begin{aligned}
\mathcal{G}' = & -\frac{1}{(1+z)H} \left( 1 - \frac{b_1Gf''}{(1+w_t)\rho_t\theta} \right) \mathbf{G} - \frac{1}{(1+z)H} \frac{b_1^2G}{(1+w_t)\rho_t\theta^2} Z \\
& - \frac{1}{(1+z)H} \frac{b_1\rho_{ch}w_{ch}}{(1+w_t)\rho_t} \Delta_{ch} + \left( \frac{1}{2(1+z)H} \frac{b_1}{(1+w_t)\rho_t} (f' + Gf'' - 1) \right. \\
& + \left. \frac{1}{2(1+z)H} \frac{b_1^2}{(1+w_t)\rho_t} \left( Gf'' + \frac{f''}{\theta^2} \right) \right) \mathcal{G} \\
\end{aligned} \tag{48}$$

$$\begin{aligned}
\mathbf{G}' = & -\frac{1}{(1+z)H} \left( \frac{b_3}{b_1} - \frac{b_1Gf''}{(1+w_t)\rho_t\theta} \right) \mathbf{G} + \left[ \frac{1}{(1+z)H} \frac{b_2b_1G}{(1+w_t)\rho_t\theta^2} \right] Z - \left[ \frac{1}{(1+z)H} \frac{b_2\rho_{ch}w_{ch}}{(1+w_t)\rho_t} \right] \Delta_{ch} \\
& - \frac{1}{(1+z)H} \left[ \frac{b_2}{2(1+w_t)\rho_t} (f' + Gf'' - 1) - \frac{12b_2b_1}{(1+w_t)\rho_t} (Gf'' + \frac{f''}{\theta^2}) \right] \mathcal{G} \\
\end{aligned} \tag{49}$$

$$\begin{aligned}
\Delta'_G = & -\frac{1}{(1+z)H} \left[ -\frac{4H\theta}{\rho_G} \left[ 2Hb_1f''' + (2c_1 - 2H^2)f'' \right] \right. \\
& + \theta \left( 4H^2f''b_2 + 4H(2c_1 - H^2)f''b_1 \right) \left( \frac{Gf''}{\theta(1+w_t)\rho_t\rho_G} \right) \Big] \mathbf{G} \\
& - \frac{1}{(1+z)H} \left[ 4H \left[ H(b_2f''' + \frac{f''b_2}{b_2} + b_2^2f''') + (2c_1 - H^2)f'' \right] \right. \\
& + \theta \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \left( \frac{\frac{1}{2} + Gb_1f'' + \frac{b_1f'''}{\theta^2}}{(1+w_t)\rho_t\rho_G} \right) \right) \Big] \mathcal{G} + \\
& - \frac{1}{3((1+z)H)\rho_G} \left[ \left( 2H(f''b_2 + f'''b_1^2) - 8Hf''b_1 \left( \frac{G}{12H^3} + 3H \right) \right. \right. \\
& + (2c_1 - H^2)f''b_1 - 3 \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \right) \\
& + \theta(4H^2(f''b_2 + f'''b_1^2) - 4H(2c_1 - H^2)f''b_1) \left( \frac{Gb_1\theta^2}{(1+w_t)\rho_t\rho_G} \right) \Big) \\
& + 3\theta(4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1) \left( \frac{(1-f'-Gf'')Gb_1}{\theta^2(1+w_t)\rho_t} \right) \Big] \mathbf{Z} \\
& - \frac{3}{(1+z)\rho_G} \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \right) \Delta_G \\
& + \frac{6}{(1+z)} \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \right) \left( \frac{w_{ch}\rho_{ch}}{(1+w_t)\rho_t\rho_G} \right) w_{ch}\Delta_{chp} \quad (50) \\
\Delta'_{ch} = & \frac{1}{(1+z)H} \left[ 1 + \frac{Gb_1}{\theta(1+w_t)\rho_t} \right] (1+w_{ch})\mathbf{Z} \\
& - \frac{3}{(1+z)} \left[ (1+\rho_{ch}) \frac{(1+w_{ch})}{(1+w_t)\rho_t} w_{ch}\rho_{ch} - (1+\alpha)w_{ch} \right] \Delta_{ch} \\
& - \frac{\theta(1+w_{ch})}{(1+z)(1+w_t)H\rho_t} \left[ \frac{1}{2}(1-f'-Gf'') + Gb_1f'' + \frac{b_1f'''}{\theta^2} \right] \mathcal{G} \\
& - \frac{\theta(1+w_{ch})Gf''}{(1+z)(1+w_t)H\rho_t} \mathbf{G} \quad (51)
\end{aligned}$$

which admit numerical solutions presented in fig. 3. We considered the initial conditions  $\mathcal{G}_{in} = 10^{-5}$ ,  $\mathbf{G}_{in} = 10^{-5}$  and  $\Delta_{in} = 10^{-5}$  to find numerical solutions of Eq. 46 through to Eq. 51. From the plot, the energy overdensity decay with increase in redshift.

#### 4.2.2 Long-Wavelength mode

In this subsection, we consider the long-wavelength regime where  $\frac{k^2}{a^2} \ll 1$ , this means that all  $k$ 's terms become negligible. Applying this approximation, and consider the dust dominated




Figure 3: Plot of energy density perturbations versus redshift of Eq. 46-51 for different values of  $m$  in the dust dominated Universe for short wavelength mode  $k = 1000$  in a generalized Chaplygin model  $\alpha = 1$ .

Universe, eq.29 through to eq. 34 are represented as

$$\begin{aligned}
\Delta'_m = & \frac{1}{(1+z)H} \left( \left( 1 - \frac{1}{(1+w_t)\rho_t} \right) \frac{(1+z)HGG'}{\theta^2} \right) Z \\
& + \frac{1}{(1+z)H} \alpha \frac{1}{(1+w_t)\rho_t} w_{chp} \rho_{chp} \Delta_{chp} \\
& + \frac{1}{(1+z)H} \frac{1}{(1+w_t)\rho_t} \frac{Gf''}{\theta} G - \frac{1}{(1+z)H} \frac{1}{(1+w_t)\rho_t} \left( \frac{1}{2} (1 - f' - Gf'') G (1+z) H G' f''' \right. \\
& \left. - \frac{(1+z)H G' f''}{G^2} \right) \mathcal{G}
\end{aligned} \tag{52}$$

$$\begin{aligned}
Z' &= \frac{1}{(1+z)H} \left( \frac{1}{2} \rho_d + \frac{1}{3} \theta^2 \right) \Delta_m \\
&- \frac{1}{(1+z)H} \left[ -\frac{1}{2} (1+3w_{chp}) \rho_{chp} + \left( \frac{1}{3} \theta^2 + \frac{1}{2} (1+3w_t) \rho_t \right) \frac{w_{chp} \rho_{chp}}{(1+w_t) \rho_t} \right. \\
&\quad \left. \left[ \frac{3}{2} \rho_{chp} - \left( \frac{1}{3} \theta^2 + \frac{1}{2} (1+3w_t) \rho_t \right) \frac{\rho_{chp}}{(1+w_t) \rho_t} \right] (\alpha+1) w_{chp} \right] \Delta_{chp} \\
&- \frac{1}{(1+z)H} \left[ 1 - 3f'' - 12H^3 b_1 f^{iv} + 12H \left( H(b_2 f''' + \frac{b_3}{b_1} f'' + b_1^2 f^{iv}) + 2H^3 f'' \right) \right. \\
&\quad \left. + \left( \frac{1}{3} \theta^2 + \frac{1}{2} (1+3w_t) \rho_t \right) \frac{1}{(1+w_t) \rho_t} \left( \frac{1}{2} (1-f' - Gf' - Gf'') + Gb_1 f''' + \frac{b_1 f''}{G^2} \right) \right] \mathcal{G} \\
&- \frac{1}{(1+z)H} \left( 24b_1 f''' - 12H^3 f''' + \left( \frac{1}{3} \theta^2 + \frac{1}{2} (1+3w_t) \rho_t \right) \frac{Gf''}{\theta(1+w_t) \rho_t} \right) \mathbf{G} \\
&- \frac{1}{(1+z)H} \left[ 8H \left( f'' b_2 + f''' b_1^2 + 3Hf' - \frac{3}{2} f''' H b_1 \right) - \frac{2}{3} \theta \right. \\
&\quad \left. - \left( \frac{1}{3} \theta^2 + \frac{1}{2} (1+3w_t) \rho_t \right) \frac{G b_1}{\theta^2 (1+w_t) \rho_t} \right] Z
\end{aligned} \tag{53}$$

$$\begin{aligned}
\mathcal{G}' &= -\frac{1}{(1+z)H} \left( 1 - \frac{b_1 G f''}{(1+w_t) \rho_t \theta} \right) \mathbf{G} - \frac{1}{(1+z)H} \frac{b_1^2 G}{(1+w_t) \rho_t \theta^2} Z \\
&- \frac{1}{(1+z)H} \frac{\alpha b_1 \rho_{ch} w_{ch}}{(1+w_t) \rho_t} \Delta_{ch} + \left( \frac{1}{2(1+z)H} \frac{b_1}{(1+w_t) \rho_t} (f' + Gf'' - 1) \right. \\
&\quad \left. + \frac{1}{2(1+z)H} \frac{b_1^2}{(1+w_t) \rho_t} \left( Gf'' + \frac{f''}{\theta^2} \right) \right) \mathcal{G}
\end{aligned} \tag{54}$$

$$\begin{aligned}
\mathbf{G}' &= -\frac{1}{(1+z)H} \left( \frac{b_3}{b_1} - \frac{b_1 G f''}{(1+w_t) \rho_t \theta} \right) \mathbf{G} + \left[ \frac{1}{(1+z)H} \frac{b_2 b_1 G}{(1+w_t) \rho_t \theta^2} \right] Z - \left[ \frac{1}{(1+z)H} \frac{\alpha b_2 \rho_{ch} w_{ch}}{(1+w_t) \rho_t} \right] \Delta_{ch} \\
&- \frac{1}{(1+z)H} \left[ \frac{b_2}{2(1+w_t) \rho_t} (f' + Gf'' - 1) - \frac{b_2 b_1}{(1+w_t) \rho_t} \left( Gf'' + \frac{f''}{\theta^2} \right) \right] \mathcal{G}
\end{aligned} \tag{55}$$

$$\begin{aligned}
\Delta'_G = & -\frac{1}{(1+z)H} \left[ -\frac{4H\theta}{\rho_G} \left[ 2Hb_1f''' + (2c_1 - 2H^2)f'' \right] \right. \\
& + \theta \left( 4H^2f''b_2 + 4H(2c_1 - H^2)f''b_1 \right) \left( \frac{Gf''}{\theta(1+w_t)\rho_t\rho_G} \right) \Big] G \\
& - \frac{1}{(1+z)H} \left[ 4H \left[ H(b_2f''' + \frac{f''b_2}{b_2} + b_2^2f''') + (2c_1 - H^2)f'' \right] \right. \\
& + \theta \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \left( \frac{\frac{1}{2} + Gb_1f'' + \frac{b_1f'''}{\theta^2}}{(1+w_t)\rho_t\rho_G} \right) \right) \Big] \mathcal{G} + \\
& - \frac{1}{3((1+z)H)\rho_G} \left[ \left( 2H(f''b_2 + f'''b_1^2) - 8Hf''b_1 \left( \frac{G}{12H^3} + 3H \right) \right. \right. \\
& + (2c_1 - H^2)f''b_1 - 3 \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \right) \\
& + \theta(4H^2(f''b_2 + f'''b_1^2) - 4H(2c_1 - H^2)f''b_1) \left( \frac{Gb_1\theta^2}{(1+w_t)\rho_t\rho_G} \right) \\
& \left. \left. + 3\theta(4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1) \left( \frac{(1-f'-Gf'')Gb_1}{\theta^2(1+w_t)\rho_t} \right) \right] Z \right. \\
& - \frac{3}{(1+z)\rho_G} \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \right) \Delta_G \\
& \left. + \frac{3}{(1+z)} \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \right) \left( \frac{w_{ch}\rho_{ch}}{(1+w_t)\rho_t\rho_G} \right) w_{ch}(\alpha+1)\Delta_{chp} \right) \quad (56)
\end{aligned}$$

$$\begin{aligned}
\Delta'_{ch} = & \frac{1}{(1+z)H} \left[ 1 + \frac{Gb_1}{\theta(1+w_t)\rho_t} \right] (1+w_{ch})Z \\
& - \frac{3}{(1+z)} \left[ (1+\rho_{ch}) \frac{(1+w_{ch})}{(1+w_t)\rho_t} w_{ch}\rho_{ch} - (1+\alpha)w_{ch} \right] \Delta_{ch} - \\
& \frac{\theta(1+w_{ch})}{(1+z)(1+w_t)H\rho_t} \left[ \frac{1}{2}(1-f'-Gf'') + Gb_1f'' + \frac{b_1f'''}{\theta^2} \right] \mathcal{G} - \frac{\theta(1+w_{ch})Gf''}{(1+z)(1+w_t)H\rho_t} G \quad (57)
\end{aligned}$$






Figure 4: Plot of energy density perturbations versus redshift of Eq. 52-57 for different values of  $m$  in the dust dominated Universe for long wavelength mode  $k = 0$  and  $\alpha = 0.2$ .




Figure 5: Plot of energy density perturbations versus redshift of Eq. 52-57 for different values of  $m$  in the dust dominated Universe for long wavelength mode  $k = 0.001$  and  $\alpha = 0.2$ .

For simplicity we set  $f(G)$  to  $f$  and considered the initial conditions  $\mathcal{G}_{in} = 10^{-5}$ ,  $\mathbf{G}_{in} = 10^{-5}$  and  $\Delta_{in} = 10^{-5}$  to find numerical solutions of Eq. 52 through to Eq. 57 which are presented in Fig. 4 through to Fig. 5. From the plots, the energy overdensity decay with increase in redshift.

The Eq. 52 through to Eq. 57 are presented for the generalized Chaplygin gas model. The original Chaplygin gas model can be obtained by setting  $\alpha = 1$  in eq. 52 through to eq. 57 and its evolution equations are presented as

$$\begin{aligned}\Delta'_m &= \frac{1}{(1+z)H} \left( \left( 1 - \frac{1}{(1+w_t)\rho_t} \right) \frac{(1+z)HGG'}{\theta^2} \right) Z \\ &+ \frac{1}{(1+z)H} \frac{1}{(1+w_t)\rho_t} w_{chp} \rho_{chp} \Delta_{chp} + \frac{1}{(1+z)H} \frac{1}{(1+w_t)\rho_t} \frac{Gf''}{\theta} \mathbf{G} \\ &- \frac{1}{(1+z)H} \frac{1}{(1+w_t)\rho_t} \left( \frac{1}{2} (1 - f' - Gf'') G (1+z) H G' f''' - \frac{(1+z)H G' f''}{G^2} \right) \mathcal{G},\end{aligned}\tag{58}$$

$$\begin{aligned}Z' &= \frac{1}{(1+z)H} \left( \frac{1}{2} \rho_d + \frac{1}{3} \theta^2 \right) \Delta_m \\ &- \frac{1}{(1+z)H} \left[ -\frac{1}{2} (1 + 3w_{chp}) \rho_{chp} + \left( \frac{1}{3} \theta^2 + \frac{1}{2} (1 + 3w_t) \rho_t \right) \frac{w_{chp} \rho_{chp}}{(1+w_t)\rho_t} \right. \\ &2 \left[ \frac{3}{2} \rho_{chp} - \left( \frac{1}{3} \theta^2 + \frac{1}{2} (1 + 3w_t) \rho_t \right) \frac{\rho_{chp}}{(1+w_t)\rho_t} \right] w_{chp} \left. \right] \Delta_{chp} \\ &- \frac{1}{(1+z)H} \left[ 1 - 3f'' - 12H^3 b_1 f^{iv} + 12H \left( H(b_2 f''' + \frac{b_3}{b_1} f'' + b_1^2 f^{iv}) + 2H^3 f'' \right) \right. \\ &+ \left. \left( \frac{1}{3} \theta^2 + \frac{1}{2} (1 + 3w_t) \rho_t \right) \frac{1}{(1+w_t)\rho_t} \left( \frac{1}{2} (1 - f' - Gf' - Gf'') + Gb_1 f''' + \frac{b_1 f''}{G^2} \right) \right] \mathcal{G} \\ &- \frac{1}{(1+z)H} \left( 24b_1 f''' - 12H^3 f''' + \left( \frac{1}{3} \theta^2 + \frac{1}{2} (1 + 3w_t) \rho_t \right) \frac{Gf''}{\theta(1+w_t)\rho_t} \right) \mathbf{G} \\ &- \frac{1}{(1+z)H} \left[ 8H \left( f'' b_2 + f''' b_1^2 + 3Hf' - \frac{3}{2} f''' H b_1 \right) - \frac{2}{3} \theta \right. \\ &- \left. \left( \frac{1}{3} \theta^2 + \frac{1}{2} (1 + 3w_t) \rho_t \right) \frac{G b_1}{\theta^2 (1+w_t)\rho_t} \right] Z\end{aligned}\tag{59}$$

$$\begin{aligned}\mathcal{G}' &= -\frac{1}{(1+z)H} \left( 1 - \frac{b_1 G f''}{(1+w_t)\rho_t \theta} \right) \mathbf{G} - \frac{1}{(1+z)H} \frac{b_1^2 G}{(1+w_t)\rho_t \theta^2} Z \\ &- \frac{1}{(1+z)H} \frac{b_1 \rho_{ch} w_{ch}}{(1+w_t)\rho_t} \Delta_{ch} + \left( \frac{1}{2(1+z)H} \frac{b_1}{(1+w_t)\rho_t} (f' + Gf'' - 1) \right. \\ &+ \left. \frac{1}{2(1+z)H} \frac{b_1^2}{(1+w_t)\rho_t} \left( Gf'' + \frac{f''}{\theta^2} \right) \right) \mathcal{G}\end{aligned}\tag{60}$$

$$\begin{aligned}\mathbf{G}' &= -\frac{1}{(1+z)H} \left( \frac{b_3}{b_1} - \frac{b_1 G f''}{(1+w_t)\rho_t \theta} \right) \mathbf{G} + \left[ \frac{1}{(1+z)H} \frac{b_2 b_1 G}{(1+w_t)\rho_t \theta^2} \right] Z - \left[ \frac{1}{(1+z)H} \frac{b_2 \rho_{ch} w_{ch}}{(1+w_t)\rho_t} \right] \Delta_{ch} \\ &- \frac{1}{(1+z)H} \left[ \frac{b_2}{2(1+w_t)\rho_t} (f' + Gf'' - 1) - \frac{b_2 b_1}{(1+w_t)\rho_t} \left( Gf'' + \frac{f''}{\theta^2} \right) \right] \mathcal{G}\end{aligned}\tag{61}$$

$$\begin{aligned}
\Delta'_G = & -\frac{1}{(1+z)H} \left[ -\frac{4H\theta}{\rho_G} \left[ 2Hb_1f''' + (2c_1 - 2H^2)f'' \right] \right. \\
& + \theta \left( 4H^2f''b_2 + 4H(2c_1 - H^2)f''b_1 \right) \left( \frac{Gf''}{\theta(1+w_t)\rho_t\rho_G} \right) \Big] \mathbf{G} \\
& - \frac{1}{(1+z)H} \left[ 4H \left[ H(b_2f''' + \frac{f''b_2}{b_2} + b_2^2f''') + (2c_1 - H^2)f'' \right] \right. \\
& + \theta \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \left( \frac{\frac{1}{2} + Gb_1f'' + \frac{b_1f'''}{\theta^2}}{(1+w_t)\rho_t\rho_G} \right) \right) \Big] \mathbf{G} + \\
& - \frac{1}{3((1+z)H)\rho_G} \left[ \left( 2H(f''b_2 + f'''b_1^2) - 8Hf''b_1 \left( \frac{G}{12H^3} + 3H \right) \right. \right. \\
& + (2c_1 - H^2)f''b_1 - 3 \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \right) \\
& + \theta(4H^2(f''b_2 + f'''b_1^2) - 4H(2c_1 - H^2)f''b_1) \left( \frac{Gb_1\theta^2}{(1+w_t)\rho_t\rho_G} \right) \Big] \\
& + 3\theta(4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1) \left( \frac{(1-f'-Gf'')Gb_1}{\theta^2(1+w_t)\rho_t} \right) \Big] \mathbf{Z} \\
& - \frac{3}{(1+z)\rho_G} \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \right) \Delta_G \\
& + \frac{6}{(1+z)} \left( 4H^2(f''b_2 + f'''b_1^2) + 4H(2c_1 - H^2)f''b_1 \right) \left( \frac{w_{ch}\rho_{ch}}{(1+w_t)\rho_t\rho_G} \right) w_{ch}\Delta_{ch} \tag{62}
\end{aligned}$$

$$\begin{aligned}
\Delta'_{ch} = & \frac{1}{(1+z)H} \left[ 1 + \frac{Gb_1}{\theta(1+w_t)\rho_t} \right] (1+w_{ch})\mathbf{Z} - \frac{3}{(1+z)} \left[ (1+\rho_{ch}) \frac{(1+w_{ch})}{(1+w_t)\rho_t} w_{ch}\rho_{ch} - 2w_{ch} \right] \Delta_{ch} \\
& - \frac{\theta(1+w_{ch})}{(1+z)(1+w_t)H\rho_t} \left[ \frac{1}{2}(1-f'-Gf'') + Gb_1f'' + \frac{b_1f'''}{\theta^2} \right] \mathbf{G} - \frac{\theta(1+w_{ch})Gf''}{(1+z)(1+w_t)H\rho_t} \mathbf{G} \tag{63}
\end{aligned}$$

which admit the numerical solutions presented in fig. 6. We considered the initial conditions  $\mathcal{G}_{in} = 10^{-5}$ ,  $\mathbf{G}_{in} = 10^{-5}$  and  $\Delta_{in} = 10^{-5}$  to find numerical solutions of Eq. 58 through to Eq. 63. The energy overdensity fluctuations decay with increase in redshift.

## 5 Discussion and Conclusions

The current treatment is focused on a mixture of matter fluid, chaplygin gas and Gauss-Bonnet fluid as non-interacting entities to study cosmological perturbations in  $f(G)$  gravity using 1+3 covariant Gauge invariant formalism. After presenting the covariant form of field equations and the kinematic quantities in the presence of Gauss-Bonnet invariant, we define matter fluids, chaplygin gas fluids and Gauss-Bonnet fluids to derive first order evolution equations for the defined gradient variables. Since our main interest lies in the structure formation of the Universe, we use scalar decomposition method to decompose the evolution equations into their scalar parts believed to be responsible for the spherical clustering of large scale structure. These scalar perturbation equations are considered as input to study the energy overdensity fluctuations in a dust-Chaplygin gas-Gauss-Bonnet system by applying harmonic decomposition method. After we transform the linear perturbation equations into redshift space for further analysis. We apply both the generalized and original chaplygin gas together with the  $f(G)$  model to find numerical results and analyse how the energy density fluctuations evolve with redshift. We considered the short wavelength limits of the perturbation equations in the




Figure 6: Plot of energy density perturbations versus redshift of Eq. 58-63 for different values of  $m$  in the dust dominated Universe for long wavelength mode  $k = 0.00001$  in a generalized Chaplygin model  $\alpha = 1$ .

Universe dominated by a mixture of dust-chaplygin gas-Gauss-Bonnet fluids system and we depict that the energy overdensity perturbations decay with increase in redshift. From all plots, we observe that the decay of matter density contrast is fast decaying with redshift compared to the matter-Chaplygin-Gauss-Bonnet fluids system, since the universe is believed to become less structured in the accelerated phase of the universe evolution. We have considered different initial conditions such as  $\mathcal{G}_{in} = 10^{-5}$ ,  $\mathbf{G}_{in} = 10^{-5}$  and  $\Delta_{in} = 10^{-5}$  to find numerical solutions of different perturbation equations. The use of initial conditions constrained the amplitude of matter energy densities but does not alter the behaviour of the curves, but the values of the parameter  $m$ ,  $k$  and other considered parameters changes the behavior of the curves. From all the plots, there is no oscillatory behaviours observed.

#### Some of the specific highlights of the present work include:

- We have derived the linear perturbation equations eq. 29-34 in the context of  $f(G)$  gravity, believed to contribute to the large scale structure formation.
- The matter energy density fluctuations couples with the energy density fluctuations of the chaplygin gas and that of Gauss-Bonnet fluids which then decouple for the GR limits.
- In the analysis stage, we considered short wavelength and long wavelength modes for the energy overdensity fluctuations in a Universe dominated by a mixture of a dust-chaplygin gas-Gauss-Bonnet fluids by combining chaplygin gas and the  $f(G)$  models which can mimics the  $\Lambda$ CDM for the  $f(G) = G$  case.
- We solved the perturbation equations using numerical method and the numerical results

of the energy overdensity fluctuations are presented in Fig. 1 GR limits and Fig. 2-6 for the combination of chaplygin gas and  $f(G)$  model in the short and long wavelength modes. From these figures the energy overdensity decay with redshift for the considered range of parameter  $m$ .

- From the plots, the numerical results of the perturbation equations are sensible to the parameter  $m$ . As  $m$  changes, we notice an change in the amplitudes of the energy overdensity fluctuations which reduces to the GR limits for  $m = 1$  but the results do not depend much on the parameter  $\alpha$ .
- Under the consideration of both original and generalized Chaplygin gas models, it was observed that there is no significant difference in the behavior of the results.

In general, the combination of chaplygin gas models and  $f(G)$  model offers an alternative for large scale structure formation scenarios even though there is a variation in the growth of amplitudes of the energy overdensity fluctuations which can be beneficial for constraining the parameters of the models by referring to the observational data which provide different cosmological scenarios that are consistent with the  $\Lambda$ CDM limits. We look forward to undertake this aspect of the task to different chaplygin gas models as well as  $f(G)$  models in the 1 + 3 covariant cosmological perturbations framework for the future work.

## Appendix

### Useful Linearised Differential Identities

For all scalars  $f$ , vectors  $V_a$  and tensors that vanish in the background,  $S_{ab} = S_{(ab)}$ , the following linearised identities hold:

$$\left(\tilde{\nabla}_{\langle a}\tilde{\nabla}_{b\rangle}f\right)' = \tilde{\nabla}_{\langle a}\tilde{\nabla}_{b\rangle}\dot{f} - \frac{2}{3}\Theta\tilde{\nabla}_{\langle a}\tilde{\nabla}_{b\rangle}f + \dot{f}\tilde{\nabla}_{\langle a}A_{b\rangle} , \quad (64)$$

$$\varepsilon^{abc}\tilde{\nabla}_b\tilde{\nabla}_cf = 0 , \quad (65)$$

$$\varepsilon_{cda}\tilde{\nabla}^c\tilde{\nabla}_{\langle b}\tilde{\nabla}^d\rangle f = \varepsilon_{cda}\tilde{\nabla}^c\tilde{\nabla}_{(b}\tilde{\nabla}^d)f = \varepsilon_{cda}\tilde{\nabla}^c\tilde{\nabla}_b\tilde{\nabla}^df = 0 , \quad (66)$$

$$\tilde{\nabla}^2\left(\tilde{\nabla}_af\right) = \tilde{\nabla}_a\left(\tilde{\nabla}^2f\right) + \frac{1}{3}\tilde{R}\tilde{\nabla}_af , \quad (67)$$

$$\left(\tilde{\nabla}_af\right)' = \tilde{\nabla}_a\dot{f} - \frac{1}{3}\Theta\tilde{\nabla}_af + \dot{f}A_a , \quad (68)$$

$$\left(\tilde{\nabla}_aS_{b\dots}\right)' = \tilde{\nabla}_a\dot{S}_{b\dots} - \frac{1}{3}\Theta\tilde{\nabla}_aS_{b\dots} , \quad (69)$$

$$\left(\tilde{\nabla}^2f\right)' = \tilde{\nabla}^2\dot{f} - \frac{2}{3}\Theta\tilde{\nabla}^2f + \dot{f}\tilde{\nabla}^aA_a , \quad (70)$$

$$\tilde{\nabla}_{[a}\tilde{\nabla}_{b]}V_c = -\frac{1}{6}\tilde{R}V_{[a}h_{b]c} , \quad (71)$$

$$\tilde{\nabla}_{[a}\tilde{\nabla}_{b]}S^{cd} = -\frac{1}{3}\tilde{R}S_{[a}{}^{(c}h_{b]}{}^{d)} , \quad (72)$$

$$\tilde{\nabla}^a\left(\varepsilon_{abc}\tilde{\nabla}^bV^c\right) = 0 , \quad (73)$$

$$\tilde{\nabla}_b\left(\varepsilon^{cd\langle a}\tilde{\nabla}_cS_d^{b\rangle}\right) = \frac{1}{2}\varepsilon^{abc}\tilde{\nabla}_b\left(\tilde{\nabla}_dS_c^d\right) , \quad (74)$$

$$\text{curlcurl}V_a = \tilde{\nabla}_a\left(\tilde{\nabla}^bV_b\right) - \tilde{\nabla}^2V_a + \frac{1}{3}\tilde{R}V_a , \quad (75)$$

## Used parameters in redshif space

$$H = \frac{2m}{3}(1+z)^{\frac{3}{2m}} \quad (76)$$

$$\dot{H} = c_1 = -\frac{2m}{3}(1+z)^{\frac{3}{m}} \quad (77)$$

$$\dot{G} = b_1 = \frac{256}{9}m^3\left(1 - \frac{2m}{3}\right)(1+z)^{\frac{15}{2m}} \quad (78)$$

$$\ddot{G} = b_2 = 1280m^3\left(\frac{2m}{3} - 1\right)(1+z)^{\frac{9}{m}} \quad (79)$$

$$\ddot{\ddot{G}} = b_3 = \frac{2560}{3}m^3\left(1 - \frac{2m}{3}\right)(1+z)^{\frac{21}{2m}} \quad (80)$$

$$\dot{f} = -(1+z)Hf'. \quad (81)$$

## Acknowledgments

We thank the anonymous reviewer(s) for the constructive comments towards the significant improvement of this manuscript. The authors acknowledge the financial support from UR-SIDA Grant No. DVC-AAR506/2022. AM acknowledges the support from International Science Program (ISP) to Rwanda Astrophysics, Space and Climate Science Research Group (RASC-SRG), University of Rwanda grant No: RWA01 and to East African Astrophysics Research Network(EAARN) grant No: Afro:05. AM also acknowledges the hospitality of the department of Physics of the University of Rwanda, where part of this work was conceptualise and completed.

## References

- [1] Carlos AP Bengaly. Evidence for cosmic acceleration with next-generation surveys: A model-independent approach. *Monthly Notices of the Royal Astronomical Society: Letters*, 499(1):L6–L10, 2020.
- [2] Adam G Riess, Alexei V Filippenko, Peter Challis, Alejandro Clocchiatti, Alan Diercks, Peter M Garnavich, Ron L Gilliland, Craig J Hogan, Saurabh Jha, Robert P Kirshner, et al. Observational evidence from supernovae for an accelerating universe and a cosmological constant. *The astronomical journal*, 116(3):1009, 1998.
- [3] Saul Perlmutter, Goldhaber Aldering, Gerson Goldhaber, RA Knop, Peter Nugent, Patricia G Castro, Susana Deustua, Sebastien Fabbro, Ariel Goobar, Donald E Groom, et al. Measurements of  $\omega$  and  $\lambda$  from 42 high-redshift supernovae. *The Astrophysical Journal*, 517(2):565, 1999.
- [4] Edmund J Copeland and Shinji Tsujikawa. Dynamics of dark energy. *International Journal of Modern Physics D*, 15(11):1753–1935, 2006.
- [5] Bamba Kazuharu, Capozziello Salvatore, Nojiri Shin’ichi and Odintsov Sergei D Dark energy cosmology: the equivalent description via different theoretical models and cosmography test. *Astrophysics and Space Science*, 342:155–228, 2012.
- [6] Yuting Wang, David Wands, Gong-Bo Zhao, and Lixin Xu. Post-planck constraints on interacting vacuum energy. *PHYSICAL REVIEW D Phys Rev D*, 90:023502, 2014.

- [7] Ram Gopal Vishwakarma. Einstein and beyond: a critical perspective on general relativity. *Universe*, 2(2):11, 2016.
- [8] HA Borges, S Carneiro, JC Fabris, and C Pigozzo. Evolution of density perturbations in decaying vacuum cosmology. *Physical Review D*, 77(4):043513, 2008.
- [9] Max Tegmark, Michael A Strauss, Michael R Blanton, Kevork Abazajian, Scott Dodelson, Havard Sandvik, Xiaomin Wang, David H Weinberg, Idit Zehavi, Neta A Bahcall, et al. Cosmological parameters from sdss and wmap. *Physical review D*, 69(10):103501, 2004.
- [10] Suresh Kumar. Remedy of some cosmological tensions via effective phantom-like behavior of interacting vacuum energy. *Physics of the Dark Universe*, 33:100862, 2021.
- [11] Renier Hough, Shambel Sahlu, Heba Sami, Maye Elmardi, Anna-Mia Swart, and Amare Abebe. Confronting the chaplygin gas with data: background and perturbed cosmic dynamics. *arXiv preprint arXiv:2112.11695*, 2021.
- [12] H Saadat and B Pourhassan. Viscous varying generalized chaplygin gas with cosmological constant and space curvature. *International Journal of Theoretical Physics*, 52:3712–3720, 2013.
- [13] Abha Dev, JS Alcaniz, and Deepak Jain. Cosmological consequences of a chaplygin gas dark energy. *Physical Review D*, 67(2):023515, 2003.
- [14] Ujjal Debnath, Asit Banerjee, and Subenoy Chakraborty. Role of modified chaplygin gas in accelerated universe. *Classical and Quantum Gravity*, 21(23):5609, 2004.
- [15] Maye Elmardi, Amare Abebe, and Abiy Tekola. Chaplygin-gas solutions of  $f(R)$  gravity. *International Journal of Geometric Methods in Modern Physics*, 13(10):1650120, 2016.
- [16] Nojiri Shin’ichi and Odintsov Sergei D Inhomogeneous equation of state of the universe: Phantom era, future singularity, and crossing the phantom barrier. *Physical Review D*, 72(2):023003, 2005.
- [17] Heba Sami, Joseph Ntahompagaze, and Amare Abebe. Inflationary  $f(R)$  cosmologies. *Universe*, 3(4):73, 2017.
- [18] Shambel Sahlu, Joseph Ntahompagaze, Maye Elmardi, and Amare Abebe. The chaplygin gas as a model for modified teleparallel gravity? *The European Physical Journal C*, 79(9):1–31, 2019.
- [19] Vittorio Gorini, Alexander Kamenshchik, Ugo Moschella, and Vincent Pasquier. The chaplygin gas as a model for dark energy. In *The Tenth Marcel Grossmann Meeting: On Recent Developments in Theoretical and Experimental General Relativity, Gravitation and Relativistic Field Theories (In 3 Volumes)*, pages 840–859. World Scientific, 2005.
- [20] Vittorio Gorini, Alexander Kamenshchik, and Ugo Moschella. Can the chaplygin gas be a plausible model for dark energy? *Physical Review D*, 67(6):063509, 2003.
- [21] George FR Ellis and Marco Bruni. Covariant and gauge-invariant approach to cosmological density fluctuations. *Physical Review D*, 40(6):1804, 1989.
- [22] P Dunsby, M Bruni, and GFR Ellis. Cosmological perturbations and the physical meaning of gauge-invariant variables. *Astrophys. J*, 395:34, 1992.
- [23] George FR Ellis and Henk Van Elst. Cosmological models: Cargese lectures 1998. *Theoretical and Observational Cosmology*, pages 1–116, 1999.

- [24] Amare Abebe, Mohamed Abdelwahab, De la Cruz-Dombriz, Peter KS Dunsby, et al. Covariant gauge-invariant perturbations in multifluid  $f(R)$  gravity. *Classical and Quantum Gravity*, 29, 2012.
- [25] Nojiri Shin'ichi and Odintsov Sergei D Modified Gauss–Bonnet theory as gravitational alternative for dark energy. *Physics Letters B*, 631(1-2),1–6,2005.
- [26] Cognola Guido, Elizalde Emilio, Nojiri Shin'ichi, Odintsov Sergei D and Zerbini Sergio Dark energy in modified Gauss-Bonnet gravity: Late-time acceleration and the hierarchy problem. *Physical Review D*,73(8):084007, 2006.
- [27] Nojiri Shin'ichi and Odintsov Sergei D Unified cosmic history in modified gravity: from  $F(R)$  theory to Lorentz non-invariant models. *Physics Reports*, 505, (2-4):59–144,2011.
- [28] Nojiri Sh, Odintsov SD, Oikonomou VK3683913 Modified gravity theories on a nutshell: inflation, bounce and late-time evolution. *Physics Reports*,692:1–104, 2017.
- [29] Amare Abebe, Alvaro de la Cruz-Dombriz, and Peter KS Dunsby. Large scale structure constraints for a class of  $f(R)$  theories of gravity. *Physical review D*, 88(4):044050, 2013.
- [30] Baojiu Li, John D Barrow, and David F Mota. Cosmology of modified Gauss-Bonnet gravity. *Physical Review D*, 76(4):044027, 2007.
- [31] Nadiezhda Montelongo Garcia, Tiberiu Harko, Francisco SN Lobo, and José P Mimoso. Energy conditions in modified Gauss-Bonnet gravity. *Physical Review D*, 83(10):104032, 2011.
- [32] Alexander Kamenshchik, Ugo Moschella, and Vincent Pasquier. An alternative to quintessence. *Physics Letters B*, 511(2-4):265–268, 2001.
- [33] Albert Munyeshyaka, Joseph Ntahompagaze, and Tom Mutabazi. Cosmological perturbations in  $f(G)$  gravity. *International Journal of Modern Physics D*, 30(07):2150053, 2021.
- [34] SA Venikoudis, KV Fasoulakos, and FP Fronimos. Late-time cosmology of scalar field assisted  $f(G)$  gravity. *International Journal of Modern Physics D*, 31(05):2250038, 2022.
- [35] Shambel Sahlu, Joseph Ntahompagaze, Amare Abebe, Álvaro de la Cruz-Dombriz, and David F Mota. Scalar perturbations in  $f(T)$  gravity using the 1+ 3 covariant approach. *The European Physical Journal C*, 80(5):422, 2020.
- [36] Chris Clarkson. Covariant approach for perturbations of rotationally symmetric spacetimes. *Physical Review D*, 76(10):104034, 2007.
- [37] Joseph Ntahompagaze, Amare Abebe, and Manasse Mbonye. A study of perturbations in scalar–tensor theory using 1+ 3 covariant approach. *International Journal of Modern Physics D*, 27(03):1850033, 2018.
- [38] Joseph Ntahompagaze, Shambel Sahlu, Amare Abebe, and Manasse R Mbonye. On multi-fluid perturbations in scalar–tensor cosmology. *International Journal of Modern Physics D*, 29(16):2050120, 2020.
- [39] Sante Carloni, Peter KS Dunsby, and Claudio Rubano. Gauge invariant perturbations of scalar-tensor cosmologies: The vacuum case. *Physical Review D*, 74(12):123513, 2006.
- [40] Albert Munyeshyaka, Abraham Ayirwanda, Fidele Twagirayezu, Beatrice Murorunkwere, and Joseph Ntahompagaze. Multifluid cosmology in  $f(G)$  gravity. *International Journal of Geometric Methods in Modern Physics*, page 2350031, 2022.



- [41] Albert Munyeshyaka, Joseph Ntahompagaze, Tom Mutabazi and Manasse R Mbonye. On 1+3 covariant perturbations of the quasi-Newtonian space-time in modified Gauss-Bonnet gravity. *arXiv preprint arXiv:2305.01331*, 2023.
  - [42] Albert Munyeshyaka, Joseph Ntahompagaze, Tom Mutabazi, Manasse R Mbonye, Abraham Ayirwanda, Fidele Twagirayezu, and Amare Abebe. Perturbations in the interacting vacuum. *International Journal of Geometric Methods in Modern Physics*, page 2350047, 2022.
  - [43] Shambel Sahlu, Heba Sami, Anna-Mia Swart, Thato Tsabone, Maye Elmardi, and Amare Abebe. Perturbations in a chaplygin gas cosmology. *arXiv preprint arXiv:2010.05173*, 2020.
-