

Logics with probabilistic team semantics and the Boolean negation

Miika Hannula¹[0000–0002–9637–6664], Minna Hirvonen¹[0000–0002–2701–9620],
Juha Kontinen¹[0000–0003–0115–5154], Yasir Mahmood²[0000–0002–5651–5391],
Arne Meier³[0000–0002–8061–5376], and Jonni Virtema⁴[0000–0002–1582–3718]

¹ University of Helsinki, Department of Mathematics and Statistics, FI

² Paderborn University, DICE group, Department of Computer Science, DE

³ Leibniz Universität Hannover, Institut für Theoretische Informatik, Hannover, DE

⁴ University of Sheffield, Department of Computer Science, Sheffield, UK

{miika.hannula,minna.hirvonen,juha.kontinen}@helsinki.fi,

yasir.mahmood@uni-paderborn.de, meier@thi.uni-hannover.de,

j.t.virtema@sheffield.ac.uk

Abstract. We study the expressivity and the complexity of various logics in probabilistic team semantics with the Boolean negation. In particular, we study the extension of probabilistic independence logic with the Boolean negation, and a recently introduced logic FOPT. We give a comprehensive picture of the relative expressivity of these logics together with the most studied logics in probabilistic team semantics setting, as well as relating their expressivity to a numerical variant of second-order logic. In addition, we introduce novel entropy atoms and show that the extension of first-order logic by entropy atoms subsumes probabilistic independence logic. Finally, we obtain some results on the complexity of model checking, validity, and satisfiability of our logics.

Keywords: Probabilistic Team Semantics · Model Checking · Satisfiability · Validity · Computational Complexity · Expressivity of Logics

1 Introduction

Probabilistic team semantics is a novel framework for the logical analysis of probabilistic and quantitative dependencies. Team semantics, as a semantic framework for logics involving qualitative dependencies and independencies, was introduced by Hodges [17] and popularised by Väänänen [25] via his dependence logic. Team semantics defines truth in reference to collections of assignments, called *teams*, and is particularly suitable for the formal analysis of properties, such as the functional dependence between variables, that arise only in the presence of multiple assignments. The idea of generalising team semantics to the probabilistic setting can be traced back to the works of Galliani [6] and Hyttinen et al. [18], however the beginning of a more systematic study of the topic dates back to works of Durand et al. [4].

In *probabilistic team semantics* the basic semantic units are probability distributions (i.e., *probabilistic teams*). This shift from set-based to distribution-based

semantics allows probabilistic notions of dependency, such as conditional probabilistic independence, to be embedded in the framework⁵. The expressivity and complexity of non-probabilistic team-based logics can be related to fragments of (existential) second-order logic and have been studied extensively (see, e.g., [7,5,10]). Team-based logics, by definition, are usually not closed under Boolean negation, so adding it can greatly increase the complexity and expressivity of these logics [19,15]. Some expressivity and complexity results have also been obtained for logics in probabilistic team semantics (see below). However, richer semantic and computational frameworks are sometimes needed to characterise these logics.

Metafinite Model Theory, introduced by Grädel and Gurevich [9], generalises the approach of *Finite Model Theory* by shifting to two-sorted structures, which extend finite structures by another (often infinite) numerical domain and weight functions bridging the two sorts. A particularly important subclass of metafinite structures are the so-called \mathbb{R} -structures, which extend finite structures with the real arithmetic on the second sort. *Blum-Shub-Smale machines* (BSS machines for short) [1] are essentially register machines with registers that can store arbitrary real numbers and compute rational functions over reals in a single time step. Interestingly, Boolean languages which are decidable by a non-deterministic polynomial-time BSS machine coincide with those languages which are PTIME-reducible to the true existential sentences of real arithmetic (i.e., the complexity class $\exists\mathbb{R}$) [2,24].

Recent works have established fascinating connections between second-order logics over \mathbb{R} -structures, complexity classes using the BSS-model of computation, and logics using probabilistic team semantics. In [13], Hannula et al. establish that the expressivity and complexity of probabilistic independence logic coincide with a particular fragment of existential second-order logic over \mathbb{R} -structures and NP on BSS-machines. In [16], Hannula and Virtama focus on probabilistic inclusion logic, which is shown to be tractable (when restricted to Boolean inputs), and relate it to linear programming.

In this paper, we focus on the expressivity and model checking complexity of probabilistic team-based logics that have access to Boolean negation. We also study the connections between probabilistic independence logic and a logic called $\text{FOPT}(\leq_c^\delta)$, which is defined via a computationally simpler probabilistic semantics [12]. The logic $\text{FOPT}(\leq_c^\delta)$ is the probabilistic variant of a certain team-based logic that can define exactly those dependencies that are first-order definable [20]. We also introduce novel entropy atoms and relate the extension of first-order logic with these atoms to probabilistic independence logic.

See Figure 1 for our expressivity results and Table 1 for our complexity results.

⁵ In [21] Li recently introduced *first-order theory of random variables with probabilistic independence (FOTPI)* whose variables are interpreted by discrete distributions over the unit interval. The paper shows that true arithmetic is interpretable in FOTPI whereas probabilistic independence logic is by our results far less complex.




Fig. 1. Landscape of relevant logics as well as relation to some complexity classes. Note that for the complexity classes, finite ordered structures are required. Double arrows indicate strict inclusions.

Logic	MC for sentences	SAT	VAL
$\text{FOPT}(\leq_c^\delta)$	PSPACE (Cor. 20)	RE [12, Thm. 5.2]	coRE [12, Thm. 5.2]
$\text{FO}(\perp_c)$	$\in \text{EXSPACE}$ and NEXPTIME-hard (Thm. 24)	RE (Thm. 26)	coRE (Thm. 26)
$\text{FO}(\sim)$	AEXPTIME[poly] [22, Prop. 5.16, Lem. 5.21]	RE [22, Thm. 5.6]	coRE [22, Thm. 5.6]
$\text{FO}(\approx)$	$\in \text{EXPTIME}$, PSPACE-hard (Thm. 22)	RE (Thm. 26)	coRE (Thm. 26)
$\text{FO}(\sim, \perp_c) \in 3\text{-EXSPACE}$, AEXPTIME[poly]-hard (Thm. 25)		RE (Thm. 26)	coRE (Thm. 26)

Table 1. Overview of our results. Unless otherwise noted, the results are completeness results. Satisfiability and Validity are considered for finite structures.

2 Preliminaries

We assume the reader is familiar with the basics in complexity theory [23]. In this work, we will encounter complexity classes PSPACE, EXPTIME, NEXPTIME, EXPSPACE and the class AEXPTIME[poly] together with the notion of completeness under the usual polynomial time many to one reductions. A bit more formally for the latter complexity class which is more uncommon than the others, AEXPTIME[poly] consists of all languages that can be decided by alternating Turing machines within an exponential runtime of $O(2^{n^{O(1)}})$ and polynomially many alternations between universal and existential states. There exists problems in propositional team logic with generalized dependence atoms that are complete for this class [14]. It is also known that truth evaluation of alternating dependency quantified boolean formulae (ADQBF) is complete for this class [14].

2.1 Probabilistic team semantics

We denote first-order variables by x, y, z and tuples of first-order variables by $\mathbf{x}, \mathbf{y}, \mathbf{z}$. For the length of the tuple \mathbf{x} , we write $|\mathbf{x}|$. The set of variables that appear in the tuple \mathbf{x} is denoted by $\text{Var}(\mathbf{x})$. A vocabulary τ is a finite set of relation, function, and constant symbols, denoted by R, f , and c , respectively. Each relation symbol R and function symbol f has a prescribed arity, denoted by $\text{Ar}(R)$ and $\text{Ar}(f)$.

Let τ be a finite relational vocabulary such that $\{=\} \subseteq \tau$. For a finite τ -structure \mathcal{A} and a finite set of variables D , an *assignment* of \mathcal{A} for D is a function $s: D \rightarrow A$. A *team* X of \mathcal{A} over D is a finite set of assignments $s: D \rightarrow A$.

A *probabilistic team* \mathbb{X} is a function $\mathbb{X}: X \rightarrow \mathbb{R}_{\geq 0}$, where $\mathbb{R}_{\geq 0}$ is the set of non-negative real numbers. The value $\mathbb{X}(s)$ is called the *weight* of assignment s . Since zero-weights are allowed, we may, when useful, assume that X is maximal, i.e., it contains all assignments $s: D \rightarrow A$. The *support* of \mathbb{X} is defined as $\text{supp}(\mathbb{X}) := \{s \in X \mid \mathbb{X}(s) \neq 0\}$. A team \mathbb{X} is *nonempty* if $\text{supp}(\mathbb{X}) \neq \emptyset$.

These teams are called probabilistic because we usually consider teams that are probability distributions, i.e., functions $\mathbb{X}: X \rightarrow \mathbb{R}_{\geq 0}$ for which $\sum_{s \in X} \mathbb{X}(s) = 1$.⁶ In this setting, the weight of an assignment can be thought of as the probability that the values of the variables are as in the assignment. If \mathbb{X} is a probability distribution, we also write $\mathbb{X}: X \rightarrow [0, 1]$.

For a set of variables V , the restriction of the assignment s to V is denoted by $s \upharpoonright V$. The *restriction of a team* X to V is $X \upharpoonright V = \{s \upharpoonright V \mid s \in X\}$, and the *restriction of a probabilistic team* \mathbb{X} to V is $\mathbb{X} \upharpoonright V: X \upharpoonright V \rightarrow \mathbb{R}_{\geq 0}$ where

$$(\mathbb{X} \upharpoonright V)(s) = \sum_{\substack{s' \upharpoonright V = s, \\ s' \in X}} \mathbb{X}(s').$$

⁶ In some sources, the term probabilistic team only refers to teams that are distributions, and the functions $\mathbb{X}: X \rightarrow \mathbb{R}_{\geq 0}$ that are not distributions are called *weighted teams*.

If ϕ is a first-order formula, then \mathbb{X}_ϕ is the restriction of the team \mathbb{X} to those assignments in X that satisfy the formula ϕ . The weight $|\mathbb{X}_\phi|$ is defined analogously as the sum of the weights of the assignments in X that satisfy ϕ , e.g.,

$$|\mathbb{X}_{\mathbf{x}=\mathbf{a}}| = \sum_{\substack{s \in X, \\ s(\mathbf{x})=\mathbf{a}}} \mathbb{X}(s).$$

For a variable x and $a \in A$, we denote by $s(a/x)$, the modified assignment $s(a/x): D \cup \{x\} \rightarrow A$ such that $s(a/x)(y) = a$ if $y = x$, and $s(a/x)(y) = s(y)$ otherwise. For a set $B \subseteq A$, the modified team $X(B/x)$ is defined as the set $X(B/x) := \{s(a/x) \mid a \in B, s \in X\}$.

Let $\mathbb{X}: X \rightarrow \mathbb{R}_{\geq 0}$ be any probabilistic team. Then the probabilistic team $\mathbb{X}(B/x)$ is a function $\mathbb{X}(B/x): X(B/x) \rightarrow \mathbb{R}_{\geq 0}$ defined as

$$\mathbb{X}(B/x)(s(a/x)) = \sum_{\substack{t \in X, \\ t(a/x)=s(a/x)}} \mathbb{X}(t) \cdot \frac{1}{|B|}.$$

If x is a fresh variable, the summation can be dropped and the right-hand side of the equation becomes $\mathbb{X}(s) \cdot \frac{1}{|B|}$. For singletons $B = \{a\}$, we write $X(a/x)$ and $\mathbb{X}(a/x)$ instead of $X(\{a\}/x)$ and $\mathbb{X}(\{a\}/x)$.

Let then $\mathbb{X}: X \rightarrow [0, 1]$ be a distribution. Denote by p_B the set of all probability distributions $d: B \rightarrow [0, 1]$, and let F be a function $F: X \rightarrow p_B$. Then the probabilistic team $\mathbb{X}(F/x)$ is a function $\mathbb{X}(F/x): X(B/x) \rightarrow [0, 1]$ defined as

$$\mathbb{X}(F/x)(s(a/x)) = \sum_{\substack{t \in X, \\ t(a/x)=s(a/x)}} \mathbb{X}(t) \cdot f(t)(a)$$

for all $a \in B$ and $s \in X$. If x is a fresh variable, the summation can again be dropped and the right-hand side of the equation becomes $\mathbb{X}(s) \cdot f(s)(a)$.

Let $\mathbb{X}: X \rightarrow [0, 1]$ and $\mathbb{Y}: Y \rightarrow [0, 1]$ be probabilistic teams with common variable and value domains, and let $k \in [0, 1]$. The k -scaled union of \mathbb{X} and \mathbb{Y} , denoted by $\mathbb{X} \sqcup_k \mathbb{Y}$, is the probabilistic team $\mathbb{X} \sqcup_k \mathbb{Y}: Y \rightarrow [0, 1]$ defined as

$$\mathbb{X} \sqcup_k \mathbb{Y}(s) := \begin{cases} k \cdot \mathbb{X}(s) + (1 - k) \cdot \mathbb{Y}(s) & \text{if } s \in X \cap Y, \\ k \cdot \mathbb{X}(s) & \text{if } s \in X \setminus Y, \\ (1 - k) \cdot \mathbb{Y}(s) & \text{if } s \in Y \setminus X. \end{cases}$$

3 Probabilistic independence logic with Boolean negation

In this section, we define probabilistic independence logic with Boolean negation, denoted by $\text{FO}(\perp_c, \sim)$. The logic extends first-order logic with *probabilistic independence atom* $\mathbf{y} \perp_{\mathbf{x}} \mathbf{z}$ which states that the tuples \mathbf{y} and \mathbf{z} are independent given the tuple \mathbf{x} . The syntax for the logic $\text{FO}(\perp_c, \sim)$ over a vocabulary τ is as follows:

$$\phi ::= R(\mathbf{x}) \mid \neg R(\mathbf{x}) \mid \mathbf{y} \perp_{\mathbf{x}} \mathbf{z} \mid \sim \phi \mid (\phi \wedge \phi) \mid (\phi \vee \phi) \mid \exists x \phi \mid \forall x \phi,$$

where x is a first-order variable, \mathbf{x} , \mathbf{y} , and \mathbf{z} are tuples of first-order variables, and $R \in \tau$.

Let ψ be a first-order formula. We denote by ψ^\neg the formula which is obtained from $\neg\psi$ by pushing the negation in front of atomic formulas. We also use the shorthand notations $\psi \rightarrow \phi := (\psi^\neg \vee (\psi \wedge \phi))$ and $\psi \leftrightarrow \phi := \psi \rightarrow \phi \wedge \phi \rightarrow \psi$.

Let $\mathbb{X}: X \rightarrow [0, 1]$ be a probability distribution. The semantics for the logic is defined as follows:

$$\begin{aligned} \mathcal{A} \models_{\mathbb{X}} R(\mathbf{x}) &\text{ iff } \mathcal{A} \models_s R(\mathbf{x}) \text{ for all } s \in \text{supp}(\mathbb{X}). \\ \mathcal{A} \models_{\mathbb{X}} \neg R(\mathbf{x}) &\text{ iff } \mathcal{A} \models_s \neg R(\mathbf{x}) \text{ for all } s \in \text{supp}(\mathbb{X}). \\ \mathcal{A} \models_{\mathbb{X}} \mathbf{y} \perp_{\mathbf{x}} \mathbf{z} &\text{ iff } |\mathbb{X}_{\mathbf{xy}=s(\mathbf{xy})}| \cdot |\mathbb{X}_{\mathbf{xz}=s(\mathbf{xz})}| = |\mathbb{X}_{\mathbf{xyz}=s(\mathbf{xyz})}| \cdot |\mathbb{X}_{\mathbf{x}=s(\mathbf{x})}| \text{ for all } s: \text{Var}(\mathbf{xyz}) \rightarrow A. \\ \mathcal{A} \models_{\mathbb{X}} \sim \phi &\text{ iff } \mathcal{A} \not\models_{\mathbb{X}} \phi. \\ \mathcal{A} \models_{\mathbb{X}} \phi \wedge \psi &\text{ iff } \mathcal{A} \models_{\mathbb{X}} \phi \text{ and } \mathcal{A} \models_{\mathbb{X}} \psi. \\ \mathcal{A} \models_{\mathbb{X}} \phi \vee \psi &\text{ iff } \mathcal{A} \models_{\mathbb{Y}} \phi \text{ and } \mathcal{A} \models_{\mathbb{Z}} \psi \text{ for some } \mathbb{Y}, \mathbb{Z}, k \text{ such that } \mathbb{Y} \sqcup_k \mathbb{Z} = \mathbb{X}. \\ \mathcal{A} \models_{\mathbb{X}} \exists x \phi &\text{ iff } \mathcal{A} \models_{\mathbb{X}(F/x)} \phi \text{ for some } F: X \rightarrow p_A. \\ \mathcal{A} \models_{\mathbb{X}} \forall x \phi &\text{ iff } \mathcal{A} \models_{\mathbb{X}(A/x)} \phi. \end{aligned}$$

The satisfaction relation \models_s above refers to the Tarski semantics of first-order logic. For a sentence ϕ , we write $\mathcal{A} \models \phi$ if $\mathcal{A} \models_{\mathbb{X}_\emptyset} \phi$, where \mathbb{X}_\emptyset is the distribution that maps the empty assignment to 1.

The logic also has the following useful property called *locality*. Denote by $\text{Fr}(\phi)$ the set of the free variables of a formula ϕ .

Proposition 1 (Locality, [4, Prop. 12]). *Let ϕ be any $\text{FO}(\perp_c, \sim)[\tau]$ -formula. Then for any set of variables V , any τ -structure \mathcal{A} , and any probabilistic team $\mathbb{X}: X \rightarrow [0, 1]$ such that $\text{Fr}(\phi) \subseteq V \subseteq D$,*

$$\mathcal{A} \models_{\mathbb{X}} \phi \iff \mathcal{A} \models_{\mathbb{X}|_V} \phi.$$

In addition to probabilistic conditional independence atoms, we may also consider other atoms. If \mathbf{x} and \mathbf{y} are tuples of variables, then $=(\mathbf{x}, \mathbf{y})$ is a *dependence atom*. If \mathbf{x} and \mathbf{y} are also of the same length, $\mathbf{x} \approx \mathbf{y}$ is a *marginal identity atom*. The semantics for these atoms are defined as follows:

$$\begin{aligned} \mathcal{A} \models_{\mathbb{X}} =(\mathbf{x}, \mathbf{y}) &\text{ iff for all } s, s' \in \text{supp}(\mathbb{X}), s(\mathbf{x}) = s'(\mathbf{x}) \text{ implies } s(\mathbf{y}) = s'(\mathbf{y}), \\ \mathcal{A} \models_{\mathbb{X}} \mathbf{x} \approx \mathbf{y} &\text{ iff } |\mathbb{X}_{\mathbf{x}=\mathbf{a}}| = |\mathbb{X}_{\mathbf{y}=\mathbf{a}}| \text{ for all } \mathbf{a} \in A^{|\mathbf{x}|}. \end{aligned}$$

The next two propositions follow from the fact that dependence atoms and marginal identity atoms can be expressed with probabilistic independence atoms.

Proposition 2 ([3, Prop. 24]). $\text{FO}(=(\dots)) \leq \text{FO}(\perp_c)$.

Proposition 3 ([11, Thm. 10]). $\text{FO}(\approx) \leq \text{FO}(\perp_c)$.

On the other hand, omitting the Boolean negation strictly decreases the expressivity:

Proposition 4. $\text{FO}(\perp_c) < \text{FO}(\perp_c, \sim)$.

Proof. By Theorems 4.1 and 6.5 of [13], over a fixed universe size, any open formula of $\text{FO}(\perp_c)$ defines a closed subset of \mathbb{R}^n for a suitable n depending on the size of the universe and the number of free variables. Now, clearly, this cannot be true for all of the formulas of $\text{FO}(\perp_c, \sim)$ as it contains the Boolean negation, e.g., the formula $\sim x \perp_y z$. \square

4 Metafinite logics

In this section, we consider logics over \mathbb{R} -structures. These structures extend finite relational structures with real numbers \mathbb{R} as a second domain and add functions that map tuples from the finite domain to \mathbb{R} .

Definition 5 (\mathbb{R} -structures). Let τ and σ be finite vocabularies such that τ is relational and σ is functional. An \mathbb{R} -structure of vocabulary $\tau \cup \sigma$ is a tuple $\mathcal{A} = (A, \mathbb{R}, F)$ where the reduct of \mathcal{A} to τ is a finite relational structure, and F is a set that contains functions $f^{\mathcal{A}}: A^{\text{Ar}(f)} \rightarrow \mathbb{R}$ for each function symbol $f \in \sigma$. Additionally,

- (i) for any $S \subseteq \mathbb{R}$, if each $f^{\mathcal{A}}$ is a function from $A^{\text{Ar}(f)}$ to S , \mathcal{A} is called an S -structure,
- (ii) if each $f^{\mathcal{A}}$ is a distribution, \mathcal{A} is called a $d[0, 1]$ -structure.

Next, we will define certain metafinite logics which are variants of functional second-order logic with numerical terms. The numerical σ -terms i are defined as follows:

$$i ::= f(\mathbf{x}) \mid i \times i \mid i + i \mid \text{SUM}_{\mathbf{y}} i \mid \log i,$$

where $f \in \sigma$ and \mathbf{x} and \mathbf{y} are first-order variables such that $|\mathbf{x}| = \text{Ar}(f)$. The interpretations of the numerical terms are defined in the obvious way.

Suppose that $\{=\} \subseteq \tau$, and let $O \subseteq \{+, \times, \text{SUM}, \log\}$. The syntax for the logic $\text{SO}_{\mathbb{R}}(O)$ is defined as follows:

$$\phi ::= i = j \mid \neg i = j \mid R(\mathbf{x}) \mid \neg R(\mathbf{x}) \mid (\phi \wedge \psi) \mid (\phi \vee \psi) \mid \exists x \phi \mid \forall x \phi \mid \exists f \psi \mid \forall f \psi,$$

where i and j are numerical σ -terms constructed using operations from O , $R \in \tau$, x, y , and \mathbf{x} are first-order variables, f is a function variable, and ψ is a $\tau \cup \sigma \cup \{f\}$ -formula of $\text{SO}_{\mathbb{R}}(O)$.

The semantics of $\text{SO}_{\mathbb{R}}(O)$ is defined via \mathbb{R} -structures and assignments analogous to first-order logic, except for the interpretations of function variables f , which range over functions $A^{\text{Ar}(f)} \rightarrow \mathbb{R}$. For any $S \subseteq \mathbb{R}$, we define $\text{SO}_S(O)$ as the variant of $\text{SO}_{\mathbb{R}}(O)$, where the quantification of function variables ranges over $A^{\text{Ar}(f)} \rightarrow S$. If the quantification of function variables is restricted to distributions, the resulting logic is denoted by $\text{SO}_{d[0,1]}(O)$. The existential fragment, in which universal quantification over function variables is not allowed, is denoted by $\text{ESO}_{\mathbb{R}}(O)$.

Proposition 6. $\text{SO}_{\mathbb{R}}(\text{SUM}, \times) \equiv \text{SO}_{\mathbb{R}}(+, \times)$.

Proof. First, note that since the constants 0 and 1 are definable in both logics, we may use them when needed. To show that $\text{SO}_{\mathbb{R}}(\text{SUM}, \times) \leq \text{SO}_{\mathbb{R}}(+, \times)$, it suffices to show that any numerical identity $f(\mathbf{x}) = \text{SUM}_{\mathbf{y}} g(\mathbf{x}, \mathbf{y})$ can also be expressed in $\text{SO}_{\mathbb{R}}(+, \times)$. Suppose that $|\mathbf{y}| = n$. Since the domain of \mathcal{A} is finite, we may assume that it is linearly ordered: a linear order \leq_{fin} can be defined with an existentially quantified binary function variable f such that the formulas $f(x, y) = 1$ and $f(x, y) = 0$ correspond to $x \leq_{\text{fin}} y$ and $x \not\leq_{\text{fin}} y$, respectively. Then, without loss of generality, we may assume that we have an n -ary successor function S defined by the lexicographic order induced by the linear order. Thus, we can existentially quantify a function variable h such that

$$\forall \mathbf{z} (h(\mathbf{x}, \mathbf{min}) = g(\mathbf{x}, \mathbf{min}) \wedge h(\mathbf{x}, S(\mathbf{z})) = h(\mathbf{x}, \mathbf{z}) + g(\mathbf{x}, S(\mathbf{z}))).$$

Then $f(\mathbf{x}) = h(\mathbf{x}, \mathbf{max})$ is as wanted.

To show that $\text{SO}_{\mathbb{R}}(+, \times) \leq \text{SO}_{\mathbb{R}}(\text{SUM}, \times)$, we show that any numerical identity $f(\mathbf{xy}) = i(\mathbf{x}) + j(\mathbf{y})$ can be expressed in $\text{SO}_{\mathbb{R}}(\text{SUM}, \times)$. We can existentially quantify a function variable g such that

$$\begin{aligned} g(\mathbf{xy}, \mathbf{min}) &= i(\mathbf{x}) \wedge g(\mathbf{xy}, \mathbf{max}) = j(\mathbf{y}) \\ &\wedge \forall z ((\neg z = \mathbf{min} \wedge \neg z = \mathbf{max}) \rightarrow g(\mathbf{uv}, z) = 0). \end{aligned}$$

Then $f(\mathbf{xy}) = \text{SUM}_z g(\mathbf{xy}, z)$ is as wanted. Note that since no universal quantification over function variables was used, the proposition also holds for existential fragments, i.e., $\text{ESO}_{\mathbb{R}}(\text{SUM}, \times) \equiv \text{ESO}_{\mathbb{R}}(+, \times)$. \square

Proposition 7. $\text{SO}_{d[0,1]}(\text{SUM}, \times) \equiv \text{SO}_{\mathbb{R}}(+, \times)$.

Proof. Since 1 is definable in $\text{SO}_{\mathbb{R}}(\text{SUM}, \times)$ and the formula $\text{SUM}_{\mathbf{x}} f(\mathbf{x}) = 1$ states that f is a probability distribution, we have that $\text{SO}_{d[0,1]}(\text{SUM}, \times) \leq \text{SO}_{\mathbb{R}}(\text{SUM}, \times) \equiv \text{SO}_{\mathbb{R}}(+, \times)$.

Next, we show that

$$\text{SO}_{\mathbb{R}}(+, \times) \leq \text{SO}_{\mathbb{R}_{\geq 0}}(+, \times) \leq \text{SO}_{[0,1]}(+, \times) \leq \text{SO}_{d[0,1]}(\text{SUM}, \times).$$

To show that $\text{SO}_{\mathbb{R}}(+, \times) \leq \text{SO}_{\mathbb{R}_{\geq 0}}(+, \times)$, let $\phi \in \text{SO}_{\mathbb{R}}(+, \times)$. Note that any function $f: A^{\text{Ar}(f)} \rightarrow \mathbb{R}$ can be expressed as $f_+ - f_-$, where f_+ and f_- are functions $A^{\text{Ar}(f)} \rightarrow \mathbb{R}_{\geq 0}$ such that $f_+(\mathbf{x}) = f(\mathbf{x}) \cdot \chi_{\mathbb{R}_{\geq 0}}(f(\mathbf{x}))$ and $f_-(\mathbf{x}) = f(\mathbf{x}) \cdot \chi_{\mathbb{R} \setminus \mathbb{R}_{\geq 0}}(f(\mathbf{x}))$, where $\chi_S: \mathbb{R} \rightarrow \{0, 1\}$ is the characteristic function of $S \subseteq \mathbb{R}$. Since numerical terms $i(\mathbf{x}) - j(\mathbf{x})$ can clearly be expressed in $\text{SO}_{\mathbb{R}}(+, \times)$, it suffices to modify ϕ as follows: for all quantified function variables f , replace each appearance of term $f(\mathbf{x})$ with $f_+(\mathbf{x}) - f_-(\mathbf{x})$ and instead of f , quantify two function variables f_+ and f_- .

To show that $\text{SO}_{\mathbb{R}_{\geq 0}}(+, \times) \leq \text{SO}_{[0,1]}(+, \times)$, let $\phi \in \text{SO}_{\mathbb{R}_{\geq 0}}(+, \times)$. Note that any positive real number can be written as a ratio $x/(1-x)$, where $x \in [0, 1]$. Since numerical terms of the form $i(\mathbf{x})/(1-i(\mathbf{x}))$ can clearly be expressed in $\text{SO}_{d[0,1]}(+, \times)$, it suffices to modify ϕ as follows: for all quantified function variables f , replace each appearance of term $f(\mathbf{x})$ with $f^*(\mathbf{x})/(1-f^*(\mathbf{x}))$ and instead of f , quantify a function variable f^* such that $f^*(\mathbf{x}) \neq 1$ for all \mathbf{x} .

Lastly, to show that $\text{SO}_{[0,1]}(+, \times) \leq \text{SO}_{d[0,1]}(\text{SUM}, \times)$, it suffices to see that for any $\phi \in \text{SO}_{[0,1]}(+, \times)$, we can compress each function term into a fraction of size $1/n^k$, where n is the size of the finite domain and k the maximal arity of any function variable appearing in ϕ . We omit the proof, since it is essentially the same as the one for Lemma 6.4 in [13]. \square

5 Equi-expressivity of $\text{FO}(\perp\!\!\!\perp_c, \sim)$ and $\text{SO}_{\mathbb{R}}(+, \times)$

In this section, we show that the expressivity of probabilistic independence logic with the Boolean negation coincides with full second-order logic over \mathbb{R} -structures.

Theorem 8. $\text{FO}(\perp\!\!\!\perp_c, \sim) \equiv \text{SO}_{\mathbb{R}}(+, \times)$.

We first show that $\text{FO}(\perp\!\!\!\perp_c, \sim) \leq \text{SO}_{\mathbb{R}}(+, \times)$. Note that by Proposition 7, we have $\text{SO}_{d[0,1]}(\text{SUM}, \times) \equiv \text{SO}_{\mathbb{R}}(+, \times)$, so it suffices to show that $\text{FO}(\perp\!\!\!\perp_c, \sim) \leq \text{SO}_{d[0,1]}(\text{SUM}, \times)$. We may assume that every independence atom is in the form $\mathbf{y} \perp\!\!\!\perp_{\mathbf{x}} \mathbf{z}$ or $\mathbf{y} \perp\!\!\!\perp_{\mathbf{x}} \mathbf{y}$ where \mathbf{x}, \mathbf{y} , and \mathbf{z} are pairwise disjoint tuples. [4, Lemma 25]

Theorem 9. *Let formula $\phi(\mathbf{v}) \in \text{FO}(\perp\!\!\!\perp_c, \sim)$ be such that its free-variables are from $\mathbf{v} = (v_1, \dots, v_k)$. Then there is a formula $\psi_{\phi}(f) \in \text{SO}_{d[0,1]}(\text{SUM}, \times)$ with exactly one free function variable such that for all structures \mathcal{A} and all probabilistic teams $\mathbb{X}: X \rightarrow [0, 1]$, $\mathcal{A} \models_{\mathbb{X}} \phi(\mathbf{v})$ if and only if $(\mathcal{A}, f_{\mathbb{X}}) \models \psi_{\phi}(f)$, where $f_{\mathbb{X}}: A^k \rightarrow [0, 1]$ is a probability distribution such that $f_{\mathbb{X}}(s(\mathbf{v})) = \mathbb{X}(s)$ for all $s \in X$.*

Proof. Define the formula $\psi_{\phi}(f)$ as follows:

1. If $\phi(\mathbf{v}) = R(v_{i_1}, \dots, v_{i_l})$, where $1 \leq i_1, \dots, i_l \leq k$, then $\psi_{\phi}(f) := \forall \mathbf{v} (f(\mathbf{v}) = 0 \vee R(v_{i_1}, \dots, v_{i_l}))$.
2. If $\phi(\mathbf{v}) = \neg R(v_{i_1}, \dots, v_{i_l})$, where $1 \leq i_1, \dots, i_l \leq k$, then $\psi_{\phi}(f) := \forall \mathbf{v} (f(\mathbf{v}) = 0 \vee \neg R(v_{i_1}, \dots, v_{i_l}))$.
3. If $\phi(\mathbf{v}) = \mathbf{v}_1 \perp\!\!\!\perp_{\mathbf{v}_0} \mathbf{v}_2$, where $\mathbf{v}_0, \mathbf{v}_1, \mathbf{v}_2$ are disjoint, then

$$\psi_{\phi}(f) := \forall \mathbf{v}_0 \mathbf{v}_1 \mathbf{v}_2 (\text{SUM}_{\mathbf{v} \setminus (\mathbf{v}_0 \mathbf{v}_1)} f(\mathbf{v}) \times \text{SUM}_{\mathbf{v} \setminus (\mathbf{v}_0 \mathbf{v}_2)} f(\mathbf{v}) = \text{SUM}_{\mathbf{v} \setminus (\mathbf{v}_0 \mathbf{v}_1)} f(\mathbf{v}) \times \text{SUM}_{\mathbf{v} \setminus \mathbf{v}_0} f(\mathbf{v})).$$

4. If $\phi(\mathbf{v}) = \mathbf{v}_1 \perp\!\!\!\perp_{\mathbf{v}_0} \mathbf{v}_1$, where $\mathbf{v}_0, \mathbf{v}_1$ are disjoint, then

$$\psi_{\phi}(f) := \forall \mathbf{v}_0 \mathbf{v}_1 (\text{SUM}_{\mathbf{v} \setminus (\mathbf{v}_0 \mathbf{v}_1)} f(\mathbf{v}) = 0 \vee \text{SUM}_{\mathbf{v} \setminus (\mathbf{v}_0 \mathbf{v}_1)} f(\mathbf{v}) = \text{SUM}_{\mathbf{v} \setminus \mathbf{v}_0} f(\mathbf{v})).$$

5. If $\phi(\mathbf{v}) = \sim \phi_0(\mathbf{v})$, then $\psi_{\phi}(f) := \psi_{\phi_0}^{-}(f)$, where $\psi_{\phi_0}^{-}$ is obtained from $\neg \psi_{\phi_0}$ by pushing the negation in front of atomic formulas.
6. If $\phi(\mathbf{v}) = \phi_0(\mathbf{v}) \wedge \phi_1(\mathbf{v})$, then $\psi_{\phi}(f) := \psi_{\phi_0}(f) \wedge \psi_{\phi_1}(f)$.
7. If $\phi(\mathbf{v}) = \phi_0(\mathbf{v}) \vee \phi_1(\mathbf{v})$, then

$$\begin{aligned} \psi_{\phi}(f) &:= \psi_{\phi_0}(f) \vee \psi_{\phi_1}(f) \\ &\vee (\exists g_0 g_1 g_2 g_3 (\forall \mathbf{v} \forall x (x = l \vee x = r \vee (g_0(x) = 0 \wedge g_3(\mathbf{v}, x) = 0)) \\ &\wedge \forall \mathbf{v} (g_3(\mathbf{v}, l) = g_1(\mathbf{v}) \times g_0(l) \wedge g_3(\mathbf{v}, r) = g_2(\mathbf{v}) \times g_0(r)) \\ &\wedge \forall \mathbf{v} (\text{SUM}_x g_3(\mathbf{v}, x) = f(\mathbf{v})) \wedge \psi_{\phi_0}(g_1) \wedge \psi_{\phi_1}(g_2))). \end{aligned}$$

8. If $\phi(\mathbf{v}) = \exists x \phi_0(\mathbf{v}, x)$, then $\psi_\phi(f) := \exists g(\forall \mathbf{v}(\text{SUM}_x g(\mathbf{v}, x) = f(\mathbf{v})) \wedge \psi_{\phi_0}(g))$.
9. If $\phi(\mathbf{v}) = \exists x \phi_0(\mathbf{v}, x)$, then

$$\psi_\phi(f) := \exists g(\forall \mathbf{v}(\forall x \forall y(g(\mathbf{v}, x) = g(\mathbf{v}, y)) \wedge \text{SUM}_x g(\mathbf{v}, x) = f(\mathbf{v})) \wedge \psi_{\phi_0}(g)).$$

Since the the above is essentially same as the translation in [4, Theorem 14], but extended with the Boolean negation (for which the claim follows directly from the semantical clauses), it is easy to show that $\psi_\phi(f)$ satisfies the claim. \square

We now show that $\text{SO}_{\mathbb{R}}(+, \times) \leq \text{FO}(\perp_c, \sim, \approx)$. By Propositions 3 and 7, $\text{FO}(\perp_c, \sim, \approx) \equiv \text{FO}(\perp_c, \sim)$ and $\text{SO}_{\mathbb{R}}(+, \times) \equiv \text{SO}_{d[0,1]}(\text{SUM}, \times)$, so it suffices to show that $\text{SO}_{d[0,1]}(\text{SUM}, \times) \leq \text{FO}(\perp_c, \sim, \approx)$.

Note that even though we consider $\text{SO}_{d[0,1]}(\text{SUM}, \times)$, where only distributions can be quantified, it may still happen that the interpretation of a numerical term does not belong to the unit interval. This may happen if we have a term of the form $\text{SUM}_{\mathbf{x}} i(\mathbf{y})$ where \mathbf{x} contains a variable that does not appear in \mathbf{y} . Fortunately, for any formula containing such terms, there is an equivalent formula without them [16, Lemma 19]. Thus, it suffices to consider formulas without such terms.

To prove that $\text{SO}_{d[0,1]}(\text{SUM}, \times) \leq \text{FO}(\perp_c, \sim, \approx)$, we construct a useful normal form for $\text{SO}_{d[0,1]}(\text{SUM}, \times)$ -sentences. The following lemma is based on similar lemmas from [4, Lemma, 16] and [16, Lemma, 20].

Lemma 10. *Every formula $\phi \in \text{SO}_{d[0,1]}(\text{SUM}, \times)$ can be written in the form $\phi^* := Q_1 f_1 \dots Q_n f_n \forall \mathbf{x} \theta$, where $Q \in \{\exists, \forall\}$, θ is quantifier-free and such that all the numerical identity atoms are in the form $f_i(\mathbf{u}\mathbf{v}) = f_j(\mathbf{u}) \times f_k(\mathbf{v})$ or $f_i(\mathbf{u}) = \text{SUM}_{\mathbf{v}} f_j(\mathbf{u}\mathbf{v})$ for distinct f_i, f_j, f_k such that at most one of them is not quantified.*

Proof. We begin by defining a formula θ_i for each numerical term $i(\mathbf{x})$ using fresh function symbols f_i .

1. If $i(\mathbf{u}) = g(\mathbf{u})$ where g is a function symbol, then θ_i is defined as $f_i(\mathbf{u}) = \text{SUM}_{\emptyset} g(\mathbf{u})$.
2. If $i(\mathbf{u}\mathbf{v}) = j(\mathbf{u}) \times k(\mathbf{v})$, then θ_i is defined as $\theta_j \wedge \theta_k \wedge f_i(\mathbf{u}\mathbf{v}) = f_j(\mathbf{u}) \times f_k(\mathbf{v})$.
3. If $i(\mathbf{u}) = \text{SUM}_{\mathbf{v}} j(\mathbf{u}\mathbf{v})$, then θ_i is defined as $\theta_j \wedge f_i(\mathbf{u}) = \text{SUM}_{\mathbf{v}} f_j(\mathbf{u}\mathbf{v})$.

Then the formula ϕ^* is defined as follows:

1. If $\phi = i(\mathbf{u}) = j(\mathbf{v})$, then $\phi^* := \exists \mathbf{f}(f_i(\mathbf{u}) = f_j(\mathbf{v}) \wedge \theta_i \wedge \theta_j)$ where \mathbf{f} consists of the function symbols f_k for each subterm k of i or j . The negated case $\phi = \neg i(\mathbf{u}) = j(\mathbf{v})$ is analogous; just add negation in front of $f_i(\mathbf{u}) = f_j(\mathbf{v})$.
2. If ϕ is an atom or a negated atom (of the first sort), then $\phi^* := \phi$.
3. If $\phi = \psi_0 \circ \psi_1$, where $\circ \in \{\wedge, \vee\}$ and $\psi_i^* = Q_1^i f_1^i \dots Q_{m_i}^i f_{m_i}^i \forall \mathbf{x}_i \theta_i$ for $i = 0, 1$, then $\phi^* := Q_1^0 f_1^0 \dots Q_{m_0}^0 f_{m_0}^0 Q_1^1 f_1^1 \dots Q_{m_1}^1 f_{m_1}^1 \forall \mathbf{x}_0 \mathbf{x}_1 (\theta_0 \circ \theta_1)$.
4. If $\phi = \exists y \psi$, where $\psi^* = Q_1 f_1 \dots Q_m f_m \forall \mathbf{x} \theta$, then

$$\phi^* := \exists g Q_1 f_1 \dots Q_m f_m \forall \mathbf{x} \forall y (g(y) = 0 \vee \theta).$$

5. Let $\phi = \forall y \psi$, where $\psi^* = Q_1 f_1 \dots Q_m f_m \forall \mathbf{x} \theta$. Let f_{m+1}, \dots, f_n list all of the free function variables in ϕ . Then define

$$\begin{aligned} \phi^* := & Q_1 f_1^* \dots Q_m f_m^* \exists f_{m+1}^* \dots \exists f_n^* \exists \mathbf{f}_{id} \exists d \forall y y' \forall \mathbf{x} (d(y) = d(y') \wedge \\ & \bigwedge_{i=m+1}^n (f_i^*(y, \mathbf{x}) = f_i^*(y', \mathbf{x}) \wedge \text{SUM}_{\mathbf{y}} f_i^*(y, \mathbf{x}) = f_i(\mathbf{x})) \wedge \\ & f_1^*(y, \mathbf{x}) = f_1^*(y', \mathbf{x}) \circ_1 (f_2^*(y, \mathbf{x}) = f_2^*(y', \mathbf{x}) \circ_2 \dots \\ & \circ_{m-1} (f_m^*(y, \mathbf{x}) = f_m^*(y', \mathbf{x}) \circ_m \theta^*) \dots), \end{aligned}$$

where each f_i^* , for $1 \leq i \leq n$, is such that $\text{Ar}(f_i^*) = \text{Ar}(f_i) + 1$, \mathbf{f}_{id} introduces a new function symbol for each multiplication in θ ,

$$\circ_i := \begin{cases} \wedge & \text{if } Q_i = \exists, \\ \rightarrow & \text{if } Q_i = \forall, \end{cases}$$

and the formula θ^* is obtained from θ by replacing all second sort identities α of the form $f_i(\mathbf{uv}) = f_j(\mathbf{u}) \times f_k(\mathbf{v})$ with

$$f_\alpha(y, \mathbf{uv}) = d(y) \times f_i^*(y, \mathbf{uv}) \wedge f_\alpha(y, \mathbf{uv}) = f_j^*(y, \mathbf{u}) \times f_k^*(y, \mathbf{v})$$

and $f_i(\mathbf{u}) = \text{SUM}_{\mathbf{v}} f_j(\mathbf{uv})$ with $f_i^*(y, \mathbf{u}) = \text{SUM}_{\mathbf{v}} f_j^*(y, \mathbf{uv})$.

6. If $\phi = Q f \psi$, where $Q \in \{\exists, \forall\}$ and $\psi^* = Q_1 f_1 \dots Q_m f_m \forall \mathbf{x} \theta$, then $\phi^* := Q f \psi^*$.

It is straightforward to check that ϕ^* is as wanted. In (5), instead of quantifying for each y a distribution f_y , we quantify a single distribution f^* such that $f^*(y, \mathbf{x}) = \frac{1}{|A|} \cdot f_y(\mathbf{x})$, where A is the domain of our structure. \square

Lemma 11. *We use the abbreviations $\forall^* x \phi$ and $\phi \rightarrow^* \psi$ for the $\text{FO}(\perp\!\!\!\perp_c, \sim, \approx)$ -formulas $\sim \exists x \sim \phi$ and $\sim(\phi \wedge \sim \psi)$, respectively. Let $\phi_\exists := \exists \mathbf{y}(\mathbf{x} \perp\!\!\!\perp \mathbf{y} \wedge \psi(\mathbf{x}, \mathbf{y}))$ and $\phi_\forall := \forall^* \mathbf{y}(\mathbf{x} \perp\!\!\!\perp \mathbf{y} \rightarrow^* \psi(\mathbf{x}, \mathbf{y}))$ be $\text{FO}(\perp\!\!\!\perp_c, \sim)$ -formulas with free variables form $\mathbf{x} = (x_1, \dots, x_n)$. Then for any structure \mathcal{A} and probabilistic team \mathbb{X} over $\{x_1, \dots, x_n\}$,*

- (i) $\mathcal{A} \models_{\mathbb{X}} \phi_\exists$ iff $\mathcal{A} \models_{\mathbb{X}(d/\mathbf{y})} \psi$ for some distribution $d: A^{|\mathbf{y}|} \rightarrow [0, 1]$,
- (ii) $\mathcal{A} \models_{\mathbb{X}} \phi_\forall$ iff $\mathcal{A} \models_{\mathbb{X}(d/\mathbf{y})} \psi$ for all distributions $d: A^{|\mathbf{y}|} \rightarrow [0, 1]$.

Proof. Let $\mathbb{Y} := \mathbb{X}(\mathbf{F}/\mathbf{y})$ for some sequence of functions $\mathbf{F} = (F_1, \dots, F_{|\mathbf{y}|})$ such that $F_i: X(A/y_1) \dots (A/y_i) \rightarrow p_A$. Now

$$\mathcal{A} \models_{\mathbb{Y}} \mathbf{x} \perp\!\!\!\perp \mathbf{y} \iff |\mathbb{Y}_{\mathbf{xy}=s(\mathbf{x})\mathbf{a}}| = |\mathbb{Y}_{\mathbf{x}=s(\mathbf{x})}| \cdot |\mathbb{Y}_{\mathbf{y}=\mathbf{a}}| \text{ for all } s \in X, \mathbf{a} \in A^{|\mathbf{y}|}.$$

Since the variables \mathbf{y} are fresh, the right-hand side becomes $\mathbb{X}(s) \cdot F_1(s)(a_1) \dots \cdot F_{|\mathbf{y}|}(s)(a_{|\mathbf{y}|}) = \mathbb{X}(s) \cdot |\mathbb{Y}_{\mathbf{y}=\mathbf{a}}|$ for all $s \in X, \mathbf{a} \in A^{|\mathbf{y}|}$, i.e., $\mathbb{X}(\mathbf{F}/\mathbf{y}) = \mathbb{X}(d/\mathbf{y})$ for some distribution $d: A^{|\mathbf{y}|} \rightarrow [0, 1]$. It is now straightforward to check that the two claims hold. \square

Theorem 12. Let $\phi(p) \in \text{SO}_{d[0,1]}(\text{SUM}, \times)$ be a formula in the form $\phi^* := Q_1 f_1 \dots Q_n f_n \forall \mathbf{x} \theta$, where $Q \in \{\exists, \forall\}$, θ is quantifier-free and such that all the numerical identity atoms are in the form $f_i(\mathbf{u}\mathbf{v}) = f_j(\mathbf{u}) \times f_k(\mathbf{v})$ or $f_i(\mathbf{u}) = \text{SUM}_{\mathbf{v}} f_j(\mathbf{u}\mathbf{v})$ for distinct f_i, f_j, f_k from $\{f_1, \dots, f_n, p\}$. Then there is a formula $\Phi \in \text{FO}(\perp_c, \sim, \approx)$ such that for all structures \mathcal{A} and probabilistic teams $\mathbb{X} := p^{\mathcal{A}}$,

$$\mathcal{A} \models_{\mathbb{X}} \Phi \text{ if and only if } (\mathcal{A}, p) \models \phi.$$

Proof. Define

$$\begin{aligned} \Phi := & \forall \mathbf{x} Q_1^* \mathbf{y}_1 (\mathbf{x} \perp \mathbf{y}_1 \circ_1 Q_2^* \mathbf{y}_2 (\mathbf{x} \mathbf{y}_1 \perp \mathbf{y}_2 \circ_2 Q_3^* \mathbf{y}_3 (\mathbf{x} \mathbf{y}_1 \mathbf{y}_2 \perp \mathbf{y}_3 \circ_3 \dots \\ & Q_n^* \mathbf{y}_n (\mathbf{x} \mathbf{y}_1 \dots \mathbf{y}_{n-1} \perp \mathbf{y}_n \circ_n \Theta) \dots)), \end{aligned}$$

where $Q_i^* = \exists$ and $\circ_i = \wedge$, whenever $Q_i = \exists$ and $Q_i^* = \forall^*$ and $\circ_i = \rightarrow^*$, whenever $Q_i = \forall$.

By Lemma 11, it suffices to show that for all distributions f_1, \dots, f_n , subsets $M \subseteq A^{|\mathbf{x}|}$, and probabilistic teams $\mathbb{Y} := \mathbb{X}(M/\mathbf{x})(f_1/\mathbf{y}_1) \dots (f_n/\mathbf{y}_n)$, we have

$$\mathcal{A} \models_{\mathbb{Y}} \Theta \iff (\mathcal{A}, p, f_1, \dots, f_n) \models \theta(\mathbf{a}) \text{ for all } \mathbf{a} \in M.$$

The claim is shown by induction on the structure of the formula Θ .

1. If θ is an atom or a negated atom (of the first sort), then clearly we may let $\Theta := \theta$.
2. Let $\theta = f_i(\mathbf{x}_i) = f_j(\mathbf{x}_j) \times f_k(\mathbf{x}_k)$. Then define

$$\Theta := \exists \alpha \beta ((\alpha = 0 \leftrightarrow \mathbf{x}_i = \mathbf{y}_i) \wedge (\beta = 0 \leftrightarrow \mathbf{x}_j \mathbf{x}_k = \mathbf{y}_j \mathbf{y}_k) \wedge \mathbf{x} \alpha \approx \mathbf{x} \beta).$$

Assume first that $(\mathcal{A}, p, f_1, \dots, f_n) \models \theta(\mathbf{a})$ for a given $\mathbf{a} \in M$. Then $f_i(\mathbf{a}_i) = f_j(\mathbf{a}_j) \times f_k(\mathbf{a}_k)$. Define functions $F_\alpha, F_\beta: Y \rightarrow \{0, 1\}$ such that $F_\alpha(s) = 0$ iff $s(\mathbf{x}_i) = s(\mathbf{y}_i)$, and $F_\beta(s) = 0$ iff $s(\mathbf{x}_j \mathbf{x}_k) = s(\mathbf{y}_j \mathbf{y}_k)$. Let $\mathbb{Z} := \mathbb{Y}(F_\alpha/\alpha)(F_\beta/\beta)$. It suffices to show that $\mathcal{A} \models_{\mathbb{Z}} \mathbf{x} \alpha \approx \mathbf{x} \beta$. Now, by the definition of \mathbb{Z} , we have $|\mathbb{Z}_{\mathbf{x}\alpha=\mathbf{a}0}| = |\mathbb{Z}_{\mathbf{x}\mathbf{y}_i=\mathbf{a}\mathbf{a}_i}| = |\mathbb{Y}_{\mathbf{x}=\mathbf{a}}| \cdot f_i(\mathbf{a}_i)$ and $|\mathbb{Z}_{\mathbf{x}\beta=\mathbf{a}0}| = |\mathbb{Z}_{\mathbf{x}\mathbf{y}_j \mathbf{y}_k=\mathbf{a}\mathbf{a}_j \mathbf{a}_k}| = |\mathbb{Y}_{\mathbf{x}=\mathbf{a}}| \cdot f_j(\mathbf{a}_j) \cdot f_k(\mathbf{a}_k)$. Since $f_i(\mathbf{a}_i) = f_j(\mathbf{a}_j) \times f_k(\mathbf{a}_k)$, we obtain $|\mathbb{Z}_{\mathbf{x}\alpha=\mathbf{a}0}| = |\mathbb{Z}_{\mathbf{x}\beta=\mathbf{a}0}|$ and $|\mathbb{Z}_{\mathbf{x}\alpha=\mathbf{a}1}| = |\mathbb{Y}_{\mathbf{x}=\mathbf{a}}| \cdot (1 - f_i(\mathbf{a}_i)) = |\mathbb{Z}_{\mathbf{x}\beta=\mathbf{a}1}|$. Hence, $\mathcal{A} \models_{\mathbb{Y}} \Theta$.

Assume then that $\mathcal{A} \models_{\mathbb{Y}} \Theta$, and define \mathbb{Z} as the extension of \mathbb{Y} such that $\mathbb{Z}_{\alpha=0} = \mathbb{Z}_{\mathbf{x}_i=\mathbf{y}_i}$ and $\mathbb{Z}_{\beta=0} = \mathbb{Z}_{\mathbf{x}_j \mathbf{x}_k=\mathbf{y}_j \mathbf{y}_k}$. Then $|\mathbb{Y}_{\mathbf{x}=\mathbf{a}}| \cdot f_i(\mathbf{a}_i) = |\mathbb{Z}_{\mathbf{x}\mathbf{y}_i=\mathbf{a}\mathbf{a}_i}| = |\mathbb{Z}_{\mathbf{x}\mathbf{x}_i=\mathbf{a}\mathbf{y}_i}| = |\mathbb{Z}_{\mathbf{x}\alpha=\mathbf{a}0}| = |\mathbb{Z}_{\mathbf{x}\beta=\mathbf{a}0}| = |\mathbb{Z}_{\mathbf{x}\mathbf{x}_j \mathbf{x}_k=\mathbf{a}\mathbf{y}_j \mathbf{y}_k}| = |\mathbb{Z}_{\mathbf{x}\mathbf{y}_j \mathbf{y}_k=\mathbf{a}\mathbf{a}_j \mathbf{a}_k}| = |\mathbb{Y}_{\mathbf{x}=\mathbf{a}}| \cdot f_j(\mathbf{a}_j) \cdot f_k(\mathbf{a}_k)$ for all $\mathbf{a} \in M$. Hence, $(\mathcal{A}, p, f_1, \dots, f_n) \models \theta(\mathbf{a})$ for all $\mathbf{a} \in M$.

The negated case $\neg f_i(\mathbf{x}_i) = f_j(\mathbf{x}_j) \times f_k(\mathbf{x}_k)$ is analogous; just add \sim in front of the existential quantification.

3. Let $\theta = f_i(\mathbf{x}_i) = \text{SUM}_{\mathbf{x}_k} f_j(\mathbf{x}_k \mathbf{x}_j)$. Then define

$$\Theta := \exists \alpha \beta ((\alpha = 0 \leftrightarrow \mathbf{x}_i = \mathbf{y}_i) \wedge (\beta = 0 \leftrightarrow \mathbf{x}_j = \mathbf{y}_j) \wedge \mathbf{x} \alpha \approx \mathbf{x} \beta).$$

The negated case $\neg f_i(\mathbf{x}_i) = \text{SUM}_{\mathbf{x}_k} f_j(\mathbf{x}_k \mathbf{x}_j)$ is analogous; just add \sim in front of the existential quantification. The proof is similar to the previous one, so it is omitted.

4. If $\theta = \theta_0 \wedge \theta_1$, then $\Theta = \Theta_0 \wedge \Theta_1$. The claim directly follows from semantics of conjunction.
5. Let $\theta = \theta_0 \vee \theta_1$. Then define

$$\Theta := \exists z (z \perp_{\mathbf{x}} z \wedge ((\Theta_0 \wedge z = 0) \vee (\Theta_1 \wedge \neg z = 0))).$$

Assume first that $(\mathcal{A}, p, f_1, \dots, f_n) \models \theta(\mathbf{a})$ for all $\mathbf{a} \in M$. Then there are M_0, M_1 such that $M_0 \cup M_1 = M$, $M_0 \cap M_1 = \emptyset$, and $(\mathcal{A}, p, f_1, \dots, f_n) \models \theta_i(\mathbf{a})$ for all $\mathbf{a} \in M_i$. Define $F: Y \rightarrow p_A$ such that $F(s) = c_i$ when $s(\mathbf{x}) \in M_i$, where c_i is the distribution defined as

$$c_i(a) := \begin{cases} 1 & \text{if } a = i, \\ 0 & \text{otherwise.} \end{cases}$$

Let $\mathbb{Z}_i := \mathbb{X}(M_i/\mathbf{x})(f_1/\mathbf{y}_1) \dots (f_n/\mathbf{y}_n)(c_i/z)$ and $k = |M_0|/|M|$. Now $\mathbb{Z} = \mathbb{Y}(F/z) = \mathbb{Z}_0 \sqcup_k \mathbb{Z}_1$, and we have $\mathcal{A} \models_{\mathbb{Z}} z \perp_{\mathbf{x}} z$, $\mathcal{A} \models_{\mathbb{Z}_0} \Theta_0 \wedge z = 0$, and $\mathcal{A} \models_{\mathbb{Z}_1} \Theta_1 \wedge \neg z = 0$. By locality, this implies that $\mathcal{A} \models_Y \Theta$.

Assume then that $\mathcal{A} \not\models_Y \Theta$. Let $F: Y \rightarrow p_A$ be such that $\mathcal{A} \models_{\mathbb{Z}} z \perp_{\mathbf{x}} z \wedge ((\Theta_0 \wedge z = 0) \vee (\Theta_1 \wedge \neg z = 0))$ for $\mathbb{Z} = \mathbb{Y}(F/z)$. Let then $k\mathbb{Z}'_0 = \mathbb{Z}_{z=0}$ and $(1-k)\mathbb{Z}'_1 = \mathbb{Z}_{z=1}$ for $k = |\mathbb{Z}_{z=0}|$. Now, we also have $\mathcal{A} \models_{\mathbb{Z}'_i} \Theta_i$ for $i = 0, 1$. Since $\mathcal{A} \models_{\mathbb{Z}} z \perp_{\mathbf{x}} z$, we have either $\mathbb{Z}_{\mathbf{x}=\mathbf{a}} = \mathbb{Z}_{\mathbf{x}z=\mathbf{a}0}$ or $\mathbb{Z}_{\mathbf{x}=\mathbf{a}} = \mathbb{Z}_{\mathbf{x}z=\mathbf{a}1}$ for all $\mathbf{a} \in M$. We get that $\mathbb{Z}_{z=0} = \mathbb{Z}_{\mathbf{x} \in M_0}$ for some $M_0 \subseteq M$. Thus, $\mathbb{Z}'_0 = \frac{|M|}{|M_0|}(\mathbb{X}(M/\mathbf{x})(f_1/\mathbf{y}_1) \dots (f_n/\mathbf{y}_n))_{\mathbf{x} \in M_0} = \mathbb{X}(M_0/\mathbf{x})(f_1/\mathbf{y}_1) \dots (f_n/\mathbf{y}_n)$. Hence, $(\mathcal{A}, p, f_1, \dots, f_n) \models \theta_0(\mathbf{a})$ for all $\mathbf{a} \in M_0$. We obtain $(\mathcal{A}, p, f_1, \dots, f_n) \models \theta_1(\mathbf{a})$ for all $\mathbf{a} \in M \setminus M_0$ by an analogous argument. As a result, we get that $(\mathcal{A}, p, f_1, \dots, f_n) \models \theta(\mathbf{a})$ for all $\mathbf{a} \in M$.

□

6 Probabilistic logics and entropy atoms

In this section we consider extending probabilistic team semantics with novel entropy atoms. For a discrete random variable X , with possible outcomes x_1, \dots, x_n occurring with probabilities $P(x_1), \dots, P(x_n)$, the Shannon entropy of X is given as:

$$H(X) := - \sum_{i=1}^n P(x_i) \log P(x_i),$$

The base of the logarithm does not play a role in this definition (usually it is assumed to be 2). For a set of discrete random variables, the entropy is defined in terms of the vector-valued random variable it defines. Given three sets of discrete random variables X, Y, Z , it is known that X is conditionally independent of Y given Z (written $X \perp_{\mathbf{x}} Y \mid Z$) if and only if the conditional mutual information $I(X; Y \mid Z)$ vanishes. Similarly, functional dependence of Y from X holds if and only if the conditional entropy $H(Y \mid X)$ of Y given X vanishes. Writing UV for the union of two sets U and V , we note that $I(X; Y \mid Z)$ and

$H(Y|X)$ can respectively be expressed as $H(ZX) + H(ZY) - H(Z) - H(ZXY)$ and $H(XY) - H(X)$. Thus many familiar dependency concepts over random variables translate into linear equations over Shannon entropies. In what follows, we shortly consider similar information-theoretic approach to dependence and independence in probabilistic team semantics.

Let $\mathbb{X}: X \rightarrow [0, 1]$ be a probabilistic team over a finite structure \mathcal{A} with universe A . Let \mathbf{x} be a k -ary sequence of variables from the domain of \mathbb{X} . Let $P_{\mathbf{x}}$ be the vector-valued random variable, where $P_{\mathbf{x}}(\mathbf{a})$ is the probability that \mathbf{x} takes value \mathbf{a} in the probabilistic team \mathbb{X} . The *Shannon entropy* of \mathbf{x} in \mathbb{X} is defined as follows:

$$H_{\mathbb{X}}(\mathbf{x}) := - \sum_{\mathbf{a} \in A^k} P_{\mathbf{x}}(\mathbf{a}) \log P_{\mathbf{x}}(\mathbf{a}). \quad (1)$$

Using this definition we now define the concept of an entropy atom.

Definition 13 (Entropy atom). *Let \mathbf{x} and \mathbf{y} be two sequences of variables from the domain of \mathbb{X} . These sequences may be of different lengths. The entropy atom is an expression of the form $H(\mathbf{x}) = H(\mathbf{y})$, and it is given the following semantics:*

$$\mathcal{A} \models_{\mathbb{X}} H(\mathbf{x}) = H(\mathbf{y}) \iff H_{\mathbb{X}}(\mathbf{x}) = H_{\mathbb{X}}(\mathbf{y}).$$

We then define *entropy logic* $\text{FO}(H)$ as the logic obtained by extending first-order logic with entropy atoms. The entropy atom is relatively powerful compared to our earlier atoms, since, as we will show next, it encapsulates many familiar dependency notions such as dependence and conditional independence.

Theorem 14. *The following equivalences hold over probabilistic teams of finite structures with two distinct constants 0 and 1:*

1. $=(\mathbf{x}, \mathbf{y}) \equiv H(\mathbf{x}) = H(\mathbf{xy})$.
2. $\mathbf{x} \perp\!\!\!\perp \mathbf{y} \equiv \phi$, where ϕ is defined as

$$\begin{aligned} \forall z \exists \mathbf{u} \mathbf{v} \Big(& [z = 0 \rightarrow (=(\mathbf{u}, \mathbf{x}) \wedge =(\mathbf{x}, \mathbf{u}) \wedge =(\mathbf{v}, \mathbf{xy}) \wedge =(\mathbf{xy}, \mathbf{v}))] \wedge \\ & [z = 1 \rightarrow (=(\mathbf{u}, \mathbf{y}) \wedge =(\mathbf{y}, \mathbf{u}) \wedge \mathbf{v} = \mathbf{0})] \wedge \\ & [(z = 0 \vee z = 1) \rightarrow H(\mathbf{uz}) = H(\mathbf{vz})] \Big), \end{aligned}$$

where $|\mathbf{u}| = \max\{|\mathbf{x}|, |\mathbf{y}|\}$ and $|\mathbf{v}| = |\mathbf{xy}|$.

Proof. The translation of the dependence atom simply expresses that the conditional entropy of \vec{y} given \vec{x} vanishes, which expresses that \vec{y} depends functionally on \vec{x} .

Consider the translation of the independence atom. Observe that ϕ essentially restricts attention to that subteam \mathbb{Y} in which the universally quantified variable z is either 0 or 1. There, the weight distribution of \mathbf{uz} is obtained by vertically stacking together halved weight distributions of \mathbf{x} and \mathbf{y} . Similarly, \mathbf{vz} corresponds to halving and vertical stacking of \mathbf{xy} and a dummy constant

distribution $\mathbf{0}$. Consider now the effect of halving the weights of the entropy function given in (1):

$$\begin{aligned}
H\left(\frac{1}{2}X\right) &= -\sum_{i=1}^n \frac{1}{2}P(x_i) \log \frac{1}{2}P(x_i) \\
&= -\frac{1}{2} \sum_{i=1}^n P(x_i) (\log \frac{1}{2} + \log P(x_i)) \\
&= -\frac{1}{2} \sum_{i=1}^n P(x_i) \log \frac{1}{2} - \frac{1}{2} \sum_{i=1}^n P(x_i) \log P(x_i) \\
&= \frac{1}{2} + \frac{1}{2}H(X).
\end{aligned}$$

Let us turn back to our subteam \mathbb{Y} , obtained by quantification and split disjunction from some initial team \mathbb{X} . This subteam has to satisfy $H(\mathbf{u}z) = H(\mathbf{v}z)$. What this amounts to, is the following

$$\begin{aligned}
H_{\mathbb{Y}}(\mathbf{u}z) = H_{\mathbb{Y}}(\mathbf{v}z) &\iff H_{\mathbb{X}}\left(\frac{1}{2}\mathbf{x}\right) + H_{\mathbb{X}}\left(\frac{1}{2}\mathbf{y}\right) = H_{\mathbb{X}}\left(\frac{1}{2}\mathbf{x}y\right) + H_{\mathbb{X}}\left(\frac{1}{2}\mathbf{0}\right) \\
&\iff 1 + \frac{1}{2}H_{\mathbb{X}}(\mathbf{x}) + \frac{1}{2}H_{\mathbb{X}}(\mathbf{y}) = \frac{1}{2} + \frac{1}{2}H_{\mathbb{X}}(\mathbf{x}y) + \frac{1}{2}H_{\mathbb{X}}(\mathbf{0}) \\
&\iff H_{\mathbb{X}}(\mathbf{x}) + H_{\mathbb{X}}(\mathbf{y}) = H_{\mathbb{X}}(\mathbf{x}y).
\end{aligned}$$

Thus, the translation captures the entropy condition of the independence atom. \square

Since conditional independence can be expressed with marginal independence, i.e., $\text{FO}(\perp\!\!\!\perp_c) \equiv \text{FO}(\perp\!\!\!\perp)$ [11, Theorem 11], we obtain the following corollary:

Corollary 15. $\text{FO}(\perp\!\!\!\perp_c) \leq \text{FO}(\text{H})$.

It is easy to see at this point that entropy logic and its extension with negation are subsumed by second-order logic over the reals with exponentiation.

Theorem 16. $\text{FO}(\text{H}) \leq \text{ESO}_{\mathbb{R}}(+, \times, \log)$ and $\text{FO}(\text{H}, \sim) \leq \text{SO}_{\mathbb{R}}(+, \times, \log)$.

Proof. The translation is similar to the one in Theorem 9, so it suffices to notice that the entropy atom $H(\mathbf{x}) = H(\mathbf{y})$ can be expressed as

$$\text{SUM}_{\mathbf{z}} f(\mathbf{x}, \mathbf{z}) \log f(\mathbf{x}, \mathbf{z}) = \text{SUM}_{\mathbf{z}'} f(\mathbf{y}, \mathbf{z}') \log f(\mathbf{y}, \mathbf{z}').$$

Since SUM can be expressed in $\text{ESO}_{\mathbb{R}}(+, \times, \log)$ and $\text{SO}_{\mathbb{R}}(+, \times, \log)$, we are done. \square

7 Logic for first-order probabilistic dependencies

Here, we define the logic $\text{FOPT}(\leq_c^\delta)$, which was introduced in [12].⁷ Let δ be a quantifier- and disjunction-free first-order formula, i.e., $\delta ::= \lambda \mid \neg\delta \mid (\delta \wedge \delta)$ for a first-order atomic formula λ of the vocabulary τ . Let x be a first-order variable. The syntax for the logic $\text{FOPT}(\leq_c^\delta)$ over a vocabulary τ is defined as follows:

$$\phi ::= \delta \mid (\delta \mid \delta) \leq (\delta \mid \delta) \mid \sim \phi \mid (\phi \wedge \phi) \mid (\phi \vee \phi) \mid \exists^1 x \phi \mid \forall^1 x \phi.$$

Let $\mathbb{X}: X \rightarrow \mathbb{R}_{\geq 0}$ be any probabilistic team, not necessarily a probability distribution. The semantics for the logic is defined as follows:

$$\begin{aligned} \mathcal{A} \models_{\mathbb{X}} \delta &\text{ iff } \mathcal{A} \models_s \delta \text{ for all } s \in \text{supp}(\mathbb{X}). \\ \mathcal{A} \models_{\mathbb{X}} (\delta_0 \mid \delta_1) \leq (\delta_2 \mid \delta_3) &\text{ iff } |\mathbb{X}_{\delta_0 \wedge \delta_1}| \cdot |\mathbb{X}_{\delta_3}| \leq |\mathbb{X}_{\delta_2 \wedge \delta_3}| \cdot |\mathbb{X}_{\delta_1}|. \\ \mathcal{A} \models_{\mathbb{X}} \sim \phi &\text{ iff } \mathcal{A} \not\models_{\mathbb{X}} \phi \text{ or } \mathbb{X} \text{ is empty.} \\ \mathcal{A} \models_{\mathbb{X}} \phi \wedge \psi &\text{ iff } \mathcal{A} \models_{\mathbb{X}} \phi \text{ and } \mathcal{A} \models_{\mathbb{X}} \psi. \\ \mathcal{A} \models_{\mathbb{X}} \phi \vee \psi &\text{ iff } \mathcal{A} \models_{\mathbb{X}} \phi \text{ or } \mathcal{A} \models_{\mathbb{X}} \psi. \\ \mathcal{A} \models_{\mathbb{X}} \exists^1 x \phi &\text{ iff } \mathcal{A} \models_{\mathbb{X}(a/x)} \phi \text{ for some } a \in A. \\ \mathcal{A} \models_{\mathbb{X}} \forall^1 x \phi &\text{ iff } \mathcal{A} \models_{\mathbb{X}(a/x)} \phi \text{ for all } a \in A. \end{aligned}$$

Next, we present some useful properties of $\text{FOPT}(\leq_c^\delta)$.

Proposition 17 (Locality, [12, Prop. 3.2]). *Let ϕ be any $\text{FOPT}(\leq_c^\delta)[\tau]$ -formula. Then for any set of variables V , any τ -structure \mathcal{A} , and any probabilistic team $\mathbb{X}: X \rightarrow \mathbb{R}_{\geq 0}$ such that $\text{Fr}(\phi) \subseteq V \subseteq D$,*

$$\mathcal{A} \models_{\mathbb{X}} \phi \iff \mathcal{A} \models_{\mathbb{X}|_V} \phi.$$

Over singleton traces the expressivity of $\text{FOPT}(\leq_c^\delta)$ coincides with that of FO. For $\phi \in \text{FOPT}(\leq_c^\delta)$, let ϕ^* denote the FO-formula obtained by replacing the symbols \sim , \vee , \exists^1 , and \forall^1 by \neg , \vee , \exists , and \forall , respectively, and expressions of the form $(\delta_0 \mid \delta_1) \leq (\delta_2 \mid \delta_3)$ by the formula $\neg\delta_0 \vee \neg\delta_1 \vee \delta_2 \vee \neg\delta_3$.

Proposition 18 (Singleton equivalence). *Let ϕ be a $\text{FOPT}(\leq_c^\delta)[\tau]$ -formula, \mathcal{A} a τ -structure, and \mathbb{X} a probabilistic team of \mathcal{A} with support $\{s\}$. Then $\mathcal{A} \models_{\mathbb{X}} \phi$ iff $\mathcal{A} \models_s \phi^*$.*

Proof. The proof proceeds by induction on the structure of formulas. The cases for literals and Boolean connectives are trivial. The cases for quantifiers are immediate once one notices that interpreting the quantifiers \exists^1 and \forall^1 maintain singleton supportness. We show the case for \leq . Let $\|\delta\|_{\mathcal{A},s} = 1$ if $\mathcal{A} \models_s \delta$, and $\|\delta\|_{\mathcal{A},s} = 0$ otherwise. Then

$$\begin{aligned} \mathcal{A} \models_{\mathbb{X}} (\delta_0 \mid \delta_1) \leq (\delta_2 \mid \delta_3) &\iff |\mathbb{X}_{\delta_0 \wedge \delta_1}| \cdot |\mathbb{X}_{\delta_3}| \leq |\mathbb{X}_{\delta_2 \wedge \delta_3}| \cdot |\mathbb{X}_{\delta_1}| \\ &\iff \|\delta_0 \wedge \delta_1\|_{\mathcal{A},s} \cdot \|\delta_3\|_{\mathcal{A},s} \leq \|\delta_2 \wedge \delta_3\|_{\mathcal{A},s} \cdot \|\delta_1\|_{\mathcal{A},s} \\ &\iff \mathcal{A} \models_s \neg\delta_0 \vee \neg\delta_1 \vee \delta_2 \vee \neg\delta_3. \end{aligned}$$

⁷ In [12], two sublogics of $\text{FOPT}(\leq_c^\delta)$, called $\text{FOPT}(\leq^\delta)$ and $\text{FOPT}(\leq^\delta, \perp_c^\delta)$, were also considered. Note that the results of this section also hold for these sublogics.

The first equivalence follows from the semantics of \leq and the second follows from the induction hypotheses after observing that the support of \mathbb{X} is $\{s\}$. The last equivalence follows via a simple arithmetic observation. \square

The following theorem follows directly from Propositions 17 and 18.

Theorem 19. *For sentences we have that $\text{FOPT}(\leq_c^\delta) \equiv \text{FO}$.*

The above result immediately yields the following corollary.

Corollary 20. *For sentences $\text{MC}(\text{FOPT}(\leq_c^\delta))$ is PSPACE-complete.*

Proof. This follows directly from the linear translation of $\text{FOPT}(\leq_c^\delta)$ -sentences into equivalent FO -sentences of Theorem 19 and the well-known fact that the model-checking problem of FO is PSPACE-complete. \square

Theorem 21. $\text{FOPT}(\leq_c^\delta) \leq \text{FO}(\perp_c, \sim)$ and $\text{FOPT}(\leq_c^\delta)$ is non-comparable to $\text{FO}(\perp_c)$ for open formulas.

Proof. We begin the proof of the first claim by showing that $\text{FOPT}(\leq_c^\delta) \leq \text{ESO}_{\mathbb{R}}(\text{SUM}, +, \times)$. Note that we may use numerical terms of the form $i \leq j$ in $\text{ESO}_{\mathbb{R}}(\text{SUM}, +, \times)$, because they can be expressed by the formula $\exists f \exists g (g \times g = f \wedge i + f = j)$.

Let formula $\phi(\mathbf{v}) \in \text{FOPT}(\leq_c^\delta)$ be such that its free-variables are from $\mathbf{v} = (v_1, \dots, v_k)$. Then there is a formula $\psi_\phi(f) \in \text{ESO}_{\mathbb{R}}(\text{SUM}, +, \times)$ with exactly one free function variable such that for all structures \mathcal{A} and all probabilistic teams $\mathbb{X}: X \rightarrow \mathbb{R}_{\geq 0}$, $\mathcal{A} \models_{\mathbb{X}} \phi(\mathbf{v})$ if and only if $(\mathcal{A}, f_{\mathbb{X}}) \models \psi_\phi(f)$, where $f_{\mathbb{X}}: A^k \rightarrow \mathbb{R}_{\geq 0}$ is a function such that $f_{\mathbb{X}}(s(\mathbf{v})) = \mathbb{X}(s)$ for all $s \in X$.

We may assume that the formula is in the form $\phi = Q_1^1 x_1 \dots Q_n^1 x_n \theta(\mathbf{v}, \mathbf{x})$, where $Q_i \in \{\exists, \forall\}$ and θ is quantifier-free. We begin by defining inductively a formula $\theta^*(f, \mathbf{x})$ for the subformula $\theta(\mathbf{v}, \mathbf{x})$. Note that in the following χ_δ refers to the characteristic function of δ , i.e., $\chi_\delta: A^{k+n} \rightarrow \{0, 1\}$ such that $\chi_\delta(\mathbf{a}) = 1$ if and only if $\mathcal{A} \models \delta(\mathbf{a})$. For simplicity, we only write $\theta^*(f, \mathbf{x})$ despite the fact that θ^* may contain free function variables χ_δ in addition to the variables f, \mathbf{x} .

1. If $\theta(\mathbf{v}, \mathbf{x}) = \delta(\mathbf{v}, \mathbf{x})$, then $\theta^*(f, \mathbf{x}) := \forall \mathbf{v} (f(\mathbf{v}) = 0 \vee \chi_\delta(\mathbf{v}, \mathbf{x}) = 1)$.
2. If $\theta(\mathbf{v}, \mathbf{x}) = (\delta_0 \mid \delta_1) \leq (\delta_2 \mid \delta_3)(\mathbf{v}, \mathbf{x})$, then

$$\begin{aligned} \theta^*(f, \mathbf{x}) &:= \text{SUM}_{\mathbf{v}}(f(\mathbf{v}) \times \chi_{\delta_0 \wedge \delta_1}(\mathbf{v}, \mathbf{x})) \times \text{SUM}_{\mathbf{v}}(f(\mathbf{v}) \times \chi_{\delta_3}(\mathbf{v}, \mathbf{x})) \\ &\leq \text{SUM}_{\mathbf{v}}(f(\mathbf{v}) \times \chi_{\delta_2 \wedge \delta_3}(\mathbf{v}, \mathbf{x})) \times \text{SUM}_{\mathbf{v}}(f(\mathbf{v}) \times \chi_{\delta_1}(\mathbf{v}, \mathbf{x})). \end{aligned}$$

3. If $\theta(\mathbf{v}, \mathbf{x}) = \sim \theta_0(\mathbf{v}, \mathbf{x})$, then $\theta^*(f, \mathbf{x}) := \theta_0^{*\neg}(f, \mathbf{x}) \vee \forall \mathbf{v} f(\mathbf{v}) = 0$, where $\theta_0^{*\neg}$ is obtained from $\neg \theta_0^*$ by pushing the negation in front of atomic formulas.
4. If $\theta(\mathbf{v}, \mathbf{x}) = (\theta_0 \circ \theta_1)(\mathbf{v}, \mathbf{x})$, where $\circ \in \{\wedge, \vee\}$, then $\theta^*(f, \mathbf{x}) := (\theta_0^* \star \theta_1^*)(\mathbf{x})$, where $\star \in \{\wedge, \vee\}$, respectively.

For each δ , we define a formula $\xi_\delta \in \text{ESO}_{\mathbb{R}}(\text{SUM}, +, \times)$, which says that χ_δ is the characteristic function of δ . Let $\mathbf{y} = (y_1, \dots, y_{k+n})$ and define ξ_δ as follows:

1. If $\delta(\mathbf{y}) = R(y_{i_1}, \dots, y_{i_l})$, where $1 \leq i_1, \dots, i_l \leq k+n$, then $\xi_\delta := \forall \mathbf{y}((\chi_\delta(\mathbf{y}) = 1 \leftrightarrow R(y_{i_1}, \dots, y_{i_l})) \wedge (\chi_\delta(\mathbf{y}) = 0 \leftrightarrow \neg R(y_{i_1}, \dots, y_{i_l})))$.
2. If $\delta(\mathbf{y}) = \neg \delta_0(\mathbf{y})$, then $\xi_\delta := \forall \mathbf{y}(\chi_{\delta_0}(\mathbf{y}) + \chi_{\neg \delta_0}(\mathbf{y}) = 1)$.
3. If $\delta(\mathbf{y}) = (\delta_0 \wedge \delta_1)(\mathbf{y})$, then $\xi_\delta := \forall \mathbf{y}(\chi_{\delta_0 \wedge \delta_1}(\mathbf{y}) = \chi_{\delta_0}(\mathbf{y}) \times \chi_{\delta_1}(\mathbf{y}))$

Let $\delta_1, \dots, \delta_m$ be a list such that each δ_i , $1 \leq i \leq m$, is a subformula of some formula δ that appears in a function symbol χ_δ of the formula $\theta^*(f, \mathbf{x})$. Now, we can define

$$\psi_\phi(f) := \bigvee_{1 \leq i \leq m} \chi_{\delta_i} \left(Q_1 x_1 \dots Q_k x_k \theta^*(f, \mathbf{x}) \wedge \bigwedge_{1 \leq i \leq m} \xi_{\delta_i}(\chi_{\delta_1}, \dots, \chi_{\delta_m}) \right).$$

This shows that $\text{FOPT}(\leq_c^\delta) \leq \text{ESO}_{\mathbb{R}}(\text{SUM}, +, \times)$. The first claim now follows, since $\text{ESO}_{\mathbb{R}}(\text{SUM}, +, \times) \leq \text{SO}_{\mathbb{R}}(+, \times) \equiv \text{FO}(\perp_c, \sim)$.

We will prove the second claim now. In the proof of Proposition 4, it was noted that the formula $\sim x \perp_y z$ cannot be expressed in $\text{FO}(\perp_c)$. This is not the case for $\text{FOPT}(\leq_c^\delta)$ as it contains the Boolean negation, and thus the formula $\sim x \perp_y z$ can be expressed in $\text{FOPT}(\leq_c^\delta)$ by the results of Section 4.2 in [12].

On the other hand, we have $\text{FO}(=(\dots)) \leq \text{FO}(\perp_c)$ (Prop. 2). Since on the level of sentences, $\text{FO}(=(\dots))$ is equivalent to existential second-order logic [25], there is a sentence $\phi \in \text{FO}(\perp_c)$ such that for all $\mathbb{X}: X \rightarrow [0, 1]$, $\mathcal{A} \models_{\mathbb{X}} \phi$ iff a undirected graph $\mathcal{A} = (V, E)$ is 2-colourable. Since over singleton traces the expressivity of $\text{FOPT}(\leq_c^\delta)$ coincides with FO, the sentence ϕ cannot be expressed in $\text{FOPT}(\leq_c^\delta)$, as 2-colourability cannot be expressed in FO. □

8 Complexity of satisfiability, validity and model checking

We now define satisfiability and validity in the context of probabilistic team semantics. Let $\phi \in \text{FO}(\perp_c, \sim, \approx)$. The formula ϕ is *satisfiable in a structure* \mathcal{A} if $\mathcal{A} \models_{\mathbb{X}} \phi$ for some probabilistic team \mathbb{X} , and ϕ is *valid in a structure* \mathcal{A} if $\mathcal{A} \models_{\mathbb{X}} \phi$ for all probabilistic teams \mathbb{X} over $\text{Fr}(\phi)$. The formula ϕ is *satisfiable* if there is a structure \mathcal{A} such that ϕ is satisfiable in \mathcal{A} , and ϕ is *valid* if ϕ is valid in \mathcal{A} for all structures \mathcal{A} .

For a logic L , the satisfiability problem $\text{SAT}(L)$ and the validity problem $\text{VAL}(L)$ are defined as follows: given a formula $\phi \in L$, decide whether ϕ is satisfiable (or valid, respectively). For the model checking problem $\text{MC}(L)$, we consider the following variant: given a *sentence* $\phi \in L$ and a structure \mathcal{A} , decide whether $\mathcal{A} \models \phi$.

Theorem 22. $\text{MC}(\text{FO}(\approx))$ is in EXPTIME and PSPACE-hard.

Proof. First note that $\text{FO}(\approx)$ is clearly a conservative extension of FO, as it is easy to check that probabilistic semantics and Tarski semantics agree on first-order formulas over singleton traces. The hardness now follows from this and the fact that model checking problem for FO is PSPACE-complete.

For upper bound, notice first that any $\text{FO}(\approx)$ -formula ϕ can be reduced to an almost conjunctive formula ψ^* of $\text{ESO}_R(+, \leq, \text{SUM})$ [16, Lem, 17]. Then the desired bounds follow due to the reduction from Proposition 3 in [16]. The mentioned reduction yields families of systems of linear inequalities \mathcal{S} from a structure \mathcal{A} and assignment s such that a system $S \in \mathcal{S}$ has a solution if and only if $\mathcal{A} \models_s \phi$. For a $\text{FO}(\approx)$ -formula ϕ , this transition requires exponential time and this yields membership in EXPTIME. \square

We now prove the following lemma, which will be used to prove the upper-bounds in the next three theorems.

Lemma 23. *Let \mathcal{A} be a finite structure and $\phi \in \text{FO}(\perp_c, \sim)$. Then there is a first-order sentence $\psi_{\phi, \mathcal{A}}$ over vocabulary $\{+, \times, \leq, 0, 1\}$ such that ϕ is satisfiable in \mathcal{A} if and only if $(\mathbb{R}, +, \times, \leq, 0, 1) \models \psi_{\phi, \mathcal{A}}$.*

Proof. Let ϕ be such that its free variables are from $\mathbf{v} = (v_1, \dots, v_k)$. By locality (Prop. 1), we may restrict to the teams over the variables $\{v_1, \dots, v_k\}$. Define a fresh first-order variable $s_{\mathbf{v}=\mathbf{a}}$ for each $\mathbf{a} \in A^k$. The idea is that the variable $s_{\mathbf{v}=\mathbf{a}}$ represents the weight of the assignment s for which $s(\mathbf{v}) = \mathbf{a}$. For notational simplicity, assume that $A = \{1, \dots, n\}$. Thus, we can write $\mathbf{s} = (s_{\mathbf{v}=\mathbf{1}}, \dots, s_{\mathbf{v}=\mathbf{n}})$ for the tuple that contains the variables for all the possible assignments over \mathbf{v} . Define then

$$\psi_{\phi, \mathcal{A}} := \exists s_{\mathbf{v}=\mathbf{1}} \dots s_{\mathbf{v}=\mathbf{n}} \left(\bigwedge_{\mathbf{a}} 0 \leq s_{\mathbf{v}=\mathbf{a}} \wedge \neg 0 = \sum_{\mathbf{a}} s_{\mathbf{v}=\mathbf{a}} \wedge \phi^*(\mathbf{s}) \right),$$

where $\phi^*(\mathbf{s})$ is constructed as follows:

- If $\phi(\mathbf{v}) = R(v_{i_1}, \dots, v_{i_l})$ or $\phi(\mathbf{v}) = \neg R(v_{i_1}, \dots, v_{i_l})$ where $1 \leq i_1, \dots, i_l \leq k$, then $\phi^*(\mathbf{s}) := \bigwedge_{s \models \phi} s = 0$.
- If $\phi(\mathbf{v}) = \mathbf{v}_1 \perp_{\mathbf{v}_0} \mathbf{v}_2$ for some \mathbf{v}_3 such that $\mathbf{v} = \mathbf{v}_0 \mathbf{v}_1 \mathbf{v}_2 \mathbf{v}_3$, then

$$\begin{aligned} \phi^*(\mathbf{s}) := \bigwedge_{\mathbf{a}_0 \mathbf{a}_1 \mathbf{a}_2} \left(\sum_{\mathbf{b}_2 \mathbf{b}_3} s_{\mathbf{v}=\mathbf{a}_0 \mathbf{a}_1 \mathbf{b}_2 \mathbf{b}_3} \times \sum_{\mathbf{b}_1 \mathbf{b}_3} s_{\mathbf{v}=\mathbf{a}_0 \mathbf{b}_1 \mathbf{a}_2 \mathbf{b}_3} = \right. \\ \left. \sum_{\mathbf{b}_3} s_{\mathbf{v}=\mathbf{a}_0 \mathbf{a}_1 \mathbf{a}_2 \mathbf{b}_3} \times \sum_{\mathbf{b}_1 \mathbf{b}_2 \mathbf{b}_3} s_{\mathbf{v}=\mathbf{a}_0 \mathbf{b}_1 \mathbf{b}_2 \mathbf{b}_3} \right), \end{aligned}$$

- If $\phi(\mathbf{v}) = \sim \theta_0(\mathbf{v})$ or $\phi(\mathbf{v}) = \theta_0(\mathbf{v}) \wedge \theta_1(\mathbf{v})$, then $\phi^*(\mathbf{s}) := \neg \theta_0^*(\mathbf{s})$ or $\phi^*(\mathbf{s}) := \theta_0^*(\mathbf{s}) \wedge \theta_1^*(\mathbf{s})$, respectively.
- If $\phi(\mathbf{v}) = \theta_0(\mathbf{v}) \vee \theta_1(\mathbf{v})$, then

$$\begin{aligned} \phi^*(\mathbf{s}) := \exists t_{\mathbf{v}=\mathbf{1}} r_{\mathbf{v}=\mathbf{1}} \dots t_{\mathbf{v}=\mathbf{n}} r_{\mathbf{v}=\mathbf{n}} \left(\bigwedge_{\mathbf{a}} (0 \leq t_{\mathbf{v}=\mathbf{a}} \wedge 0 \leq r_{\mathbf{v}=\mathbf{a}} \wedge \right. \\ \left. s_{\mathbf{v}=\mathbf{a}} = t_{\mathbf{v}=\mathbf{a}} + r_{\mathbf{v}=\mathbf{a}}) \wedge \theta_0^*(\mathbf{t}) \wedge \theta_1^*(\mathbf{r}) \right). \end{aligned}$$

- If $\phi(\mathbf{v}) = \exists x \theta_0(\mathbf{v}, x)$, then

$$\phi^*(\mathbf{s}) := \exists t_{\mathbf{v}x=11} \dots t_{\mathbf{v}x=nn} \left(\bigwedge_{ab} (0 \leq t_{\mathbf{v}x=ab} \wedge s_{\mathbf{v}=\mathbf{a}} = \sum_{c=1}^n t_{\mathbf{v}x=\mathbf{a}c}) \wedge \theta_0^*(\mathbf{t}) \right).$$

- If $\phi(\mathbf{v}) = \forall x \theta_0(\mathbf{v}, x)$, then

$$\phi^*(\mathbf{s}) := \exists t_{\mathbf{v}x=11} \dots t_{\mathbf{v}x=nn} \left(\bigwedge_{ab} (0 \leq t_{\mathbf{v}x=ab} \wedge s_{\mathbf{v}=\mathbf{a}} = \sum_{c=1}^n t_{\mathbf{v}x=\mathbf{a}c} \wedge \bigwedge_{cd} t_{\mathbf{v}x=\mathbf{a}c} = t_{\mathbf{v}x=\mathbf{a}d}) \wedge \theta_0^*(\mathbf{t}) \right).$$

□

Theorem 24. $\text{MC}(\text{FO}(\perp_c))$ is in EXPSPACE and NEXPTIME-hard.

Proof. For the lower bound, we use the fact that dependence atoms can be expressed by using probabilistic independence atoms. Let \mathcal{A} be a structure and \mathbb{X} be a probabilistic team over \mathcal{A} . Then $\mathcal{A} \models_{\mathbb{X}} (\mathbf{x}, \mathbf{y}) \ll_{\mathbf{x}} \mathbf{y}$ [11, Prop. 3]. The NEXPTIME-hardness follows since the model checking problem for $\text{FO}(= (\dots))$ is NEXPTIME-complete [8, Thm. 5.2].

The upper-bound follows from the fact that when restricted to $\text{FO}(\perp_c)$, the exponential translation in Lemma 23 is an existential sentence, and the existential theory of the reals is in PSPACE. □

Theorem 25. $\text{MC}(\text{FO}(\sim, \perp_c))$ is in 3-EXPSPACE and AEXPTIME[poly]-hard.

Proof. We first prove the lower bound through a reduction from the satisfiability problem for propositional team-based logic, that is, $\text{SAT}(\text{PL}(\sim))$. Given a $\text{PL}(\sim)$ -formula ϕ , the problem asks whether there is a team T such that $T \models \phi$? Let ϕ be a $\text{PL}(\sim)$ -formula over propositional variables p_1, \dots, p_n . For $i \leq n$, let x_i denote a variable corresponding to the proposition p_i . Let $\mathcal{A} = \{0, 1\}$ be the structure over empty vocabulary. Then, ϕ is satisfiable iff $\exists p_1 \dots \exists p_n \phi$ is satisfiable iff $\mathcal{A} \models_{\{\emptyset\}} \exists x_1 \dots \exists x_n \phi'$, where ϕ' is a $\text{FO}(\sim)$ -formula obtained from ϕ by simply replacing each proposition p_i by the variable x_i . This gives AEXPTIME-hardness of $\text{MC}(\text{FO}(\sim))$ (and consequently, of $\text{MC}(\text{FO}(\sim, \perp_c))$) since the satisfiability for $\text{PL}(\sim)$ is AEXPTIME-complete [15].

The upper-bound follows from the exponential translation from $\text{FO}(\sim, \perp_c)$ to real arithmetic in Lemma 23 and the fact that the full theory of the reals is in 2-EXPSPACE. □

Theorem 26. $\text{SAT}(\text{FO}(\perp_c, \sim))$ is RE- and $\text{VAL}(\text{FO}(\perp_c, \sim))$ is coRE-complete.

Proof. It suffices to prove the claim for $\text{SAT}(\text{FO}(\perp_c, \sim))$, since the claim for $\text{VAL}(\text{FO}(\perp_c, \sim))$ follows from the fact that $\text{FO}(\perp_c, \sim)$ has the Boolean negation.

For the lower bound, note that $\text{FO}(\perp_c, \sim)$ is a conservative extension of FO , and hence the claim follows from the r.e.-hardness of $\text{SAT}(\text{FO})$ over the finite.

For the upper-bound, we use Lemma 23. Let ϕ be a satisfiable formula of $\text{FO}(\perp_c, \sim)$. We can verify that $\phi \in \text{SAT}(\text{FO}(\perp_c, \sim))$ by going through all finite structures until we come across a structure in which ϕ is satisfiable. Hence, it suffices to show that for any finite structure \mathcal{A} , it is decidable to check whether ϕ is satisfiable in \mathcal{A} . For this, construct a sentence $\psi_{\mathcal{A}, \phi}$ as in Lemma 23. Then $\psi_{\mathcal{A}, \phi}$ is such that ϕ is satisfiable in \mathcal{A} iff $(\mathbb{R}, +, \times, \leq, 0, 1) \models \psi_{\mathcal{A}, \phi}$. Since real arithmetic is decidable, we now have that $\text{SAT}(\text{FO}(\perp_c, \sim))$ is RE-complete. \square

Corollary 27. *$\text{SAT}(\text{FO}(\approx))$ and $\text{SAT}(\text{FO}(\perp_c))$ are RE- and $\text{VAL}(\text{FO}(\approx))$ and $\text{VAL}(\text{FO}(\perp_c))$ are coRE-complete.*

Proof. The lower bound follows from the fact that $\text{FO}(\approx)$ and $\text{FO}(\perp_c)$ are both conservative extensions of FO. We obtain the upper bound from the previous theorem, since $\text{FO}(\perp_c, \sim)$ includes both $\text{FO}(\approx)$ and $\text{FO}(\perp_c)$. \square

9 Conclusion

We have studied the expressivity and complexity of various logics in probabilistic team semantics with the Boolean negation. Our results give a quite comprehensive picture of the relative expressivity of these logics and their relations to numerical variants of (existential) second-order logic. An interesting question for further study is to determine the exact complexities of the decision problems studied in Section 8. Furthermore, dependence atoms based on various notions of entropy deserve further study, as do the connections of probabilistic team semantics to the field of information theory.

Acknowledgements The first author is supported by the ERC grant 101020762. The second author is supported by Academy of Finland grant 345634. The third author is supported by Academy of Finland grants 338259 and 345634. The fifth author appreciates funding by the German Research Foundation (DFG), project ME 4279/3-1. The sixth author is partially funded by the German Research Foundation (DFG), project VI 1045/1-1.

References

1. Blum, L., Shub, M., Smale, S.: On a theory of computation over the real numbers; NP completeness, recursive functions and universal machines (extended abstract) pp. 387–397 (1988)
2. Bürgisser, P., Cucker, F.: Counting complexity classes for numeric computations II: algebraic and semialgebraic sets. *J. Complex.* **22**(2), 147–191 (2006)
3. Durand, A., Hannula, M., Kontinen, J., Meier, A., Virtema, J.: Approximation and dependence via multiteam semantics. *Ann. Math. Artif. Intell.* **83**(3-4), 297–320 (2018)
4. Durand, A., Hannula, M., Kontinen, J., Meier, A., Virtema, J.: Probabilistic team semantics. In: *FoIKS. Lecture Notes in Computer Science*, vol. 10833, pp. 186–206. Springer (2018)

5. Durand, A., Kontinen, J., de Rugy-Altherre, N., Väänänen, J.: Tractability frontier of data complexity in team semantics. *ACM Trans. Comput. Log.* **23**(1), 3:1–3:21 (2022)
6. Galliani, P.: Game Values and Equilibria for Undetermined Sentences of Dependence Logic (2008), MSc Thesis. ILLC Publications, MoL–2008–08
7. Galliani, P., Hella, L.: Inclusion logic and fixed point logic. In: *CSL. LIPIcs*, vol. 23, pp. 281–295. Schloss Dagstuhl - Leibniz-Zentrum für Informatik (2013)
8. Grädel, E.: Model-checking games for logics of imperfect information. *Theor. Comput. Sci.* **493**, 2–14 (2013)
9. Grädel, E., Gurevich, Y.: Metafinite model theory. *Inf. Comput.* **140**(1), 26–81 (1998)
10. Hannula, M., Hella, L.: Complexity thresholds in inclusion logic. *Inf. Comput.* **287**, 104759 (2022)
11. Hannula, M., Hirvonen, Å., Kontinen, J., Kulikov, V., Virtema, J.: Facets of distribution identities in probabilistic team semantics. In: *JELIA. Lecture Notes in Computer Science*, vol. 11468, pp. 304–320. Springer (2019)
12. Hannula, M., Hirvonen, M., Kontinen, J.: On elementary logics for quantitative dependencies. *Ann. Pure Appl. Log.* **173**(10), 103104 (2022)
13. Hannula, M., Kontinen, J., den Bussche, J.V., Virtema, J.: Descriptive complexity of real computation and probabilistic independence logic. In: *LICS*. pp. 550–563. ACM (2020)
14. Hannula, M., Kontinen, J., Lück, M., Virtema, J.: On quantified propositional logics and the exponential time hierarchy. In: *GandALF. EPTCS*, vol. 226, pp. 198–212 (2016)
15. Hannula, M., Kontinen, J., Virtema, J., Vollmer, H.: Complexity of propositional logics in team semantic. *ACM Trans. Comput. Log.* **19**(1), 2:1–2:14 (2018)
16. Hannula, M., Virtema, J.: Tractability frontiers in probabilistic team semantics and existential second-order logic over the reals. *Ann. Pure Appl. Log.* **173**(10), 103108 (2022)
17. Hodges, W.: Compositional semantics for a language of imperfect information. *Log. J. IGPL* **5**(4), 539–563 (1997)
18. Hyttinen, T., Paolini, G., Väänänen, J.: A logic for arguing about probabilities in measure teams. *Arch. Math. Log.* **56**(5-6), 475–489 (2017)
19. Kontinen, J., Nurmi, V.: Team logic and second-order logic. *Fundam. Informaticae* **106**(2-4), 259–272 (2011)
20. Kontinen, J., Yang, F.: Complete logics for elementary team properties. *The Journal of Symbolic Logic* pp. 1–41 (2022). <https://doi.org/10.1017/jsl.2022.80>
21. Li, C.T.: First-order theory of probabilistic independence and single-letter characterizations of capacity regions. In: *ISIT*. pp. 1536–1541. IEEE (2022)
22. Lück, M.: Team logic: axioms, expressiveness, complexity. Ph.D. thesis, University of Hanover, Hannover, Germany (2020), <https://www.repo.uni-hannover.de/handle/123456789/9430>
23. Papadimitriou, C.H.: Computational complexity. Academic Internet Publ. (2007)
24. Schaefer, M., Stefankovic, D.: Fixed points, nash equilibria, and the existential theory of the reals. *Theory Comput. Syst.* **60**(2), 172–193 (2017)
25. Väänänen, J.A.: Dependence Logic - A New Approach to Independence Friendly Logic, London Mathematical Society student texts, vol. 70. Cambridge University Press (2007)