

Symmetry and stability of orientationally ordered collective motions of self-propelled, semiflexible filaments

Madhuvanthi Guruprasad Athani and Daniel A. Beller*

Department of Physics and Astronomy, Johns Hopkins University, Baltimore, Maryland 21218, USA

Ordered, collective motions commonly arise spontaneously in systems of many interacting, active units, ranging from cellular tissues and bacterial colonies to self-propelled colloids and animal flocks. Active phases are especially rich when the active units are sufficiently anisotropic to produce liquid crystalline order and thus active nematic phenomena, with important biophysical examples provided by cytoskeletal filaments including microtubules and actin. Gliding assay experiments have provided a testbed to study the collective motions of these cytoskeletal filaments and unlocked diverse collective active phases, including states with long-range orientational order. However, it is not well-understood how such long-range order emerges from the interplay of passive and active aligning mechanisms. We use Brownian dynamics simulations to study the collective motions of semiflexible filaments that self-propel in quasi-two dimensions, in order to gain insights into the aligning mechanisms at work in these gliding assay systems. We find that, without aligning torques in the microscopic model, long-range orientational order can only be achieved when the filaments are able to overlap. The symmetry (nematic or polar) of the long-range order that first emerges is shown to depend on the energy cost of filament overlap and on filament flexibility. However, our model also predicts that a long-range-ordered active nematic state is merely transient, whereas long-range polar order is the only active dynamical steady state in systems with finite filament rigidity.

I. INTRODUCTION

Active matter systems are composed of a large number of interacting ‘agents’ that individually convert internal energy into motion. Flocks of birds, herds of animals, and growing or swarming bacterial colonies are some examples of active matter across various length scales [1–4]. The formation of spontaneously ordered collective motion is a common phenomenon exhibited by active systems. Clustering and flocking ordered states seen in many active systems can be modeled using active Brownian particles, in cases where the active agents are spherically symmetric [5–7]. But active matter also comprises anisotropic systems with agents whose aspect ratios differ strongly from unity, such as swimming rod-shaped *bacillus subtilis* [8] or *E. coli* bacterial swarms [9, 10], and cytoskeletal filaments such as actin and microtubules [11–13]. The anisotropy of these individual active particles is responsible for the orientational order and the distinct behaviors of active nematics [14, 15].

Collective phases of anisotropic cytoskeletal biofilaments have been experimentally observed in gliding assay systems, which were originally used to study molecular motors such as kinesin, dynein, and myosin, which walk along cytoskeletal filaments such as actin and microtubules [16–18]. In a conventional *in vitro* gliding assay setup, molecular motors are uniformly dispersed on a glass slide where they are held fixed with their tail end, which attaches to molecular cargo *in vivo*, here attached to the surface of the glass slide. Cytoskeletal filaments glide on these motors, which push the filaments forward in a defined direction, making the system microscopically

polar. When the densities of motors and filaments are sufficiently high in a gliding assay, diverse active collective phases spontaneously emerge, including polar flocks, nematic and polar lanes, asters, and “spools” of filaments [19–22]. One such active state observed is the active nematic state with long-range orientational order. Understanding how this long-range ordered (LRO) nematic state emerges and evolves from microscopically polar, flexible constituents is the main subject of this paper.

The effects of shape on interacting colloidal particles were first studied in thermal systems of passive, hard, rod-like colloidal particle systems to reveal that higher aspect ratio and higher packing fraction favor long-range nematic order [23]. These trends hold also in active systems of self-propelled hard rods; additionally, increasing the self-propulsion strength favors nematic alignment [24–27].

In gliding assays, however, the hard-rod approximation becomes unsuitable due to large deformations of the active filaments. The effects of polymer flexibility on nematic ordering are notable already in passive, lyotropic liquid crystals, where lower rigidity disfavors order and increases the isotropic-nematic phase transition concentration, compared to hard-rod colloids [28]. Rigidity is likewise an important parameter in determining the active phases of self-propelled, semiflexible filaments [22, 29–33].

Past simulation studies of active semiflexible filaments that have reported LRO states [22, 29, 32] employed an attractive or aligning interaction between nearby active filaments in their microscopic model. These assumptions were justified by the use of depletants, which cause filaments to bundle in parallel or antiparallel orientations, in conventional gliding assay experiments where LRO states have been observed [19, 20, 22, 32]. However, recent experiments with microtubules on lipid substrates have re-

* d.a.beller@jhu.edu

ported LRO states without the need for depletants [34]. This experimental evidence motivates us to study how LRO active nematic states emerge and evolve without imposed aligning or attractive interactions in the microscopic model.

Another important feature of experimental gliding assays is that filaments frequently cross over one another, with one filament partially and temporarily delaminating from the substrate [19, 22, 32–34]. These crossovers are forbidden by models that assume strictly two-dimensional motion of impenetrable filaments. As we show in this work, filament crossovers are essential to LRO states in the absence of aligning or attractive interactions at the filament-pair level.

In this paper, we conduct a detailed study of the active phase behaviors of self-propelled, semiflexible filaments without imposed attractive or aligning interactions between the active agents. We determine the conditions under which these filaments develop collective motions with long-range orientational order, by varying parameters including the filaments' flexibility and area fraction.

We find that a long-range orientationally ordered phase does not arise when the filaments are strictly constrained to two dimensions through volume exclusion. Instead, we observe LRO states only in *quasi*-two dimensional systems, in which we replace the volume-excluding steric repulsion with a finite energy cost for filament overlap [31, 35], in order to account for crossovers.

In our simulations, the symmetry of the long-range orientational order that first emerges may be polar or nematic (apolar) depending on the flexibility of the filaments. Surprisingly, we find the LRO active nematic state to be merely transient, with all LRO systems eventually becoming polar. Our model predicts that the time taken by the system to reach the LRO active polar steady state monotonically increases with filament stiffness. These findings suggest that experimental observations of LRO active nematic states in microtubule gliding assays may be better interpreted as transient behavior, observable due to the finite lifetime of experiments and relatively high rigidity of the filaments.

II. MODEL

We use Brownian dynamics to simulate active (self-propelled), semiflexible filaments in two dimensions. Each filament is modeled as a bead-spring chain with a Hookean restoring force between adjacent beads and a bending stiffness $\tilde{\kappa}$. Each bead self-propels with a constant active force in the chain's local tangent direction, resulting in an active velocity v_0 in the absence of other forces. The local tangent direction is calculated at each time step from bead positions and does not have its own relaxation time in this model.

Steric repulsion between beads of different chains is modeled using a modified Weeks-Chandler-Andersen potential [30, 36]:

$$U_{\text{WCA}}(r_{ij}) = \begin{cases} 4\varepsilon \left[\left(\frac{\sigma}{r_{ij} + r_{\text{shift}}} \right)^{12} - \left(\frac{\sigma}{r_{ij} + r_{\text{shift}}} \right)^6 \right] + \varepsilon, & r_{ij} \leq 2^{1/6}\sigma \\ 0, & r_{ij} > 2^{1/6}\sigma \end{cases} \quad (1)$$

Here, r_{ij} is the distance between the centers of two beads, and r_{shift} is a parameter introduced to make the U_{WCA} potential finite when $r_{ij} = 0$; this ensures that chains can interpenetrate. The overlap of chains in our explicitly two-dimensional model mimics the crossover events of a quasi-two-dimensional microtubule gliding assay experimental system. In addition to acting between beads of different chains, U_{WCA} also acts between bead pairs on the same chain separated by more than three bead positions, penalizing self-intersections of the semiflexible chains.

We scale simulation lengths in units of $\tilde{x} = 2^{1/6}\sigma$, the range of the WCA repulsive potential. In these units, the bead diameter is $r_0 = 0.5\tilde{x}$. The chain length $\ell = 30r_0$ is the same for all chains and is kept fixed across all simulation results presented here; polydispersity will be investigated in future work. Time t is scaled with respect to \tilde{x}/v_0 , although in our plots we present time in units of ℓ/v_0 . The Brownian (over-damped) dynamics updates bead positions through

$$\mathbf{x}(t + dt) = \mathbf{x}(t) + \mathbf{v}dt + \sqrt{2Ddt}\boldsymbol{\xi}(t) \quad (2)$$

$$\mathbf{v} = \mathbf{F}_{\text{total}}/\gamma \quad (3)$$

The third term on the right-hand side of Eq. 2 is the stochastic noise term representing random fluctuations in bead positions, with each vector component drawn from a zero-mean Gaussian white noise distribution. We use a second-order stochastic Runge-Kutta (SRK-2) method [37] to update the bead positions, with an adaptive time step. All simulations are conducted in a square box of side length L_x with periodic boundary conditions in both the x and y directions. Full details of the bead-spring-chain model can be found in the Supplemental Material [38].

III. RESULTS

A. Dependence of active state on chain penetrability

We begin by examining the strictly two-dimensional scenario in which crossovers are prohibited, with $r_{\text{shift}} = 0$. This choice makes the repulsion potential $U_{\text{WCA}}(r_{ij})$ diverge when two chains overlap ($r_{ij} = 0$), thus forbidding chains to interpenetrate when they collide. We vary bending rigidity $\tilde{\kappa}$, which in passive systems would favor




FIG. 1. **A.** Polar flocks observed when $r_{\text{shift}} = 0$ and $\epsilon = 1$ in U_{WCA} . The simulation box size $L_x = 250$, area fraction $\phi = 0.5$, and number of chains $N = 2016$. The color wheel legend, here and in subsequent figures, indicates the local tangent direction of the filaments. **B.** Plot of the WCA potential for bead interactions in the cases of steric repulsion (blue) and penetrable filaments (orange).

nematic order as it is increased [28]. However, in this active system, we find by varying $\tilde{\kappa}$ across three orders of magnitude that the system does not exhibit long-range order. Instead, polar flocks are seen for all values of $\tilde{\kappa}$, as shown in Fig. 1A and in Supplemental Video 1 [38].

These results indicate that long-range orientational order for mutually impenetrable active semiflexible chains cannot be achieved without an imposed alignment or attractive potential. Active polar flocks similar to those observed here have been reported in various microtubule and actin filament gliding assays both in experiments using depletants and in simulations where the filaments were strongly confined to two dimensions [12, 27, 31, 32, 39]. Similar polar flocks have also been reported in experiments on rod-shaped *E. coli* bacterial colonies when they were strongly confined to two dimensions [35].

We now introduce chain penetrability by modifying the r_{ij} -dependence of the U_{WCA} interaction potential as shown in Fig. 1B, with r_{shift} fixed at 0.1, so that the potential is finite at $r_{ij} = 0$. We vary ϵ to change the potential barrier for chain interpenetration, which affects the frequency of crossovers (Supplemental Video 2 [38]).

At the lowest ϵ values tested, such as $\epsilon = 10^{-12}$ as shown in Fig. 2A, we obtain an isotropic state, as chains simply cross over each other essentially unimpeded. When $\epsilon = 10^{-10}$ we observe polar density waves (Fig. 2B) traveling parallel to the chain self-propulsion direction, as previously seen in simulations and experiments involving actin filaments [22] (where the term “polar clusters” is applied to this state). A state with no positional order but long-range orientational order (Fig. 2C), our main interest in this work, is obtained when $\epsilon = 10^{-8}$. We therefore maintain U_{WCA} parameters of $\epsilon = 10^{-8}, r_{\text{shift}} = 0.1$ in the following sections of this paper.

At even higher values of ϵ , we obtain density inhomogeneities in the form of flocks. When $\epsilon = 10^{-6}$, a jammed flocking state appears (Fig. 2D), with characteristics distinct from the polar flocks seen in Fig. 1A. The jammed

flocks seen here are formed by counter-moving chains that are dynamically arrested relative to the center of mass of the flock, which then moves as a whole. (In contrast, all the chains in a particular polar flock seen in Fig. 1A are oriented nearly in the forward direction.) The jammed flocking state of (Fig. 2D) can be interpreted as a form of motility-induced phase separation [6], with counter-moving filaments impeding each others’ motions as they overlap.

The density inhomogeneities seen in Fig. 2E and F are dynamic and they reconfigure into different lanes and flocks multiple times in the course of the simulation. While the polar lane in Fig. 2F spans the system, we observe upon increasing the system size a state like the finite flock of Fig. 2E. We therefore interpret the system-spanning lane of Fig. 2F as a finite-size effect.

B. Long-range-ordered states

With ϵ fixed at 10^{-8} henceforth, we now investigate active states with long-range orientational order (LRO) in greater detail. First, we study the effect of changing bending rigidity $\tilde{\kappa}$ on the form of the orientational order. In Fig. 3, we see snapshots of the initially formed long-range orientational order (time $t_* = 200.58[\ell/v_0]$) for a range of $\tilde{\kappa}$ values. To quantify the orientational order we compute two different order parameters: nematic order $S(t) = \left| \langle \exp(i2\theta_j^t) \rangle_j \right|$ and polar order $P(t) = \left| \langle \exp(i\theta_j^t) \rangle_j \right|$, where θ is the average orientation of all the beads in a chain with respect to the positive x -axis, and subscripts j indicate a system average over all chains. We label a state as orientationally ordered when $S(t) \geq 0.7$. An orientationally ordered state is labeled a LRO polar state if $P(t) \geq 0.7$, and a LRO nematic state otherwise.

In Fig. 3, we see that more rigid chains (greater $\tilde{\kappa}$) form a LRO nematic state ($S(t_*) > 0.7$ and $P(t_*) < 0.7$) whereas more flexible chains achieve a LRO polar state ($P(t_*) > 0.7$). The same trend is revealed when nematic




FIG. 2. Changing the energy cost ε for chains to crossover in U_{WCA} we obtain the following states: **A**. Isotropic state with no positional or orientational order. **B**. Polar wave state where chains move in the same direction as the wave. **C**. Long-range orientational ordered state. The orange line in Fig. 1B corresponds to the potential used here. **D**. Jammed flock state. **E-F**. Dynamic polar flocks and lanes. All the simulations have box size $L_x = 100$, $\phi = 0.5$, $\ell = 15.5$, $\tilde{\kappa} = 30$ and $r_{\text{shift}} = 0.1$. The color wheel legend indicates the orientation of each filament, averaged over the tangents at each bead.

and polar order parameters, averaged over 20-run ensembles, are plotted against $\tilde{\kappa}$ (Fig. 4A): whereas $\langle S(t_*) \rangle$ increases modestly with increasing $\tilde{\kappa}$, $\langle P(t_*) \rangle$ decreases dramatically. Biopolymers such as microtubules have a higher bending rigidity (persistence length ≈ 1 mm) compared to actin filaments (persistence length ≈ 10 -20 μm) [40]. While LRO polar states have not been observed in experiment, our results are consistent with the known tendency of actin gliding assays to exhibit short-range polar order whereas nematic order (short- or long-range) is more common in microtubules [12, 19, 22, 32, 34, 39].

The modest increase in nematic order with increasing rigidity can be attributed to the fact that more rigid chains have greater restoring forces against bending deformations, and are therefore less likely to be significantly deflected from their direction of motion due to collisions or random fluctuations. Box plots in Fig. 4B and C show the median and spread of nematic and polar order parameter values at time t_* , when all the simulations have reached long-range orientational order. It is interesting to note that, at larger $\tilde{\kappa}$ values that gave apparently nematic states, the polar order parameter values (Fig. 4C) are significantly above zero and widely spread. This indicates that LRO nematic order coexists with LRO polar order, and raises the question of whether the LRO nematic state is, in fact, the dynamical steady state.

C. Nature of the active steady state

In order to determine whether LRO nematic order with weak LRO polar order is a dynamically stable active state, we now examine the evolution of orientationally ordered systems over times much longer than t_* . Examining the time-evolution represented by the snapshots in Fig. 5A and Supplemental Video 3 [38], we can see that the LRO active nematic state discussed in the previous section is merely transient, and the system reaches an active global polar steady state at longer times. To quantify this evolution of orientational order, we plot in

Fig. 5B the polar and nematic order parameters against time for each simulation in a 20-run ensemble, with bending rigidity $\tilde{\kappa} = 70$. The black dashed line indicates the time t_* when all the simulations have achieved long-range nematic orientational order. However, the nematic (apolar) state is transient, as indicated by $P(t)$ rising to saturation near 1 at much later times. The active steady state is therefore a LRO active polar state. At intermediate times after t_* , the polar order appears to plateau for a significant time interval, though the plateau value of $P(t)$ exhibits wide variation between simulation runs.

To understand the effect of changing bending rigidity on the steady-state long-range order, we plot in Fig. 6 the ensemble-averaged nematic (Fig. 6A) and polar (Fig. 6B, C) order parameters against time for multiple values of $\tilde{\kappa}$. From Fig. 6A we see that nematic order saturates at around the same time t_* for all $\tilde{\kappa}$ values. Fig. 6B shows that, while polar order eventually saturates for all $\tilde{\kappa}$ values, the time of saturation depends strongly on $\tilde{\kappa}$. Systems with larger $\tilde{\kappa}$ tend to take much longer times to reach their LRO active polar steady state. To examine the $\tilde{\kappa}$ -dependence of the $P(t)$ saturation time, we plot in Fig. 6C the same data as in Fig. 6B but with a linear scale on the time axis. Fitting a saturating exponential function $f(t) = r - ae^{-t/\tau}$ to the mean polar order parameter of the ensemble yields typical time scales τ for saturation of polar order at $P(t) \rightarrow r$. As shown in the inset to Fig. 6C, we find that τ monotonically increases with increasing $\tilde{\kappa}$, quantifying the slower polar ordering exhibited by more rigid filaments. The saturation value r varies between simulation runs, and its ensemble average increases slightly with increasing $\tilde{\kappa}$, indicating better polar ordering in more rigid filaments.

We next investigate the effect of changing the area fraction ϕ of filaments on the LRO state (Supplemental Material Fig. S3, S4 [38]). We see that the lowest area fraction tested ($\phi = 0.1$) takes the longest to reach the long-range orientational order (Supplemental Material Fig. S4A [38]). When we simulate the systems for much longer times compared to t_* , and fit the polar order parameter to a saturating exponential as we did in Fig. 6C, we find that the time scale τ required to reach a




FIG. 3. States with long-range orientational order obtained for different bending rigidities $\tilde{\kappa}$. All the simulations have $L_x = 100$, $\phi = 0.3$, $\ell = 15.5$, $\varepsilon = 10^{-8}$ and $r_{\text{shift}} = 0.1$. $S(t_*)$ and $P(t_*)$ are the nematic and polar order parameters respectively of the simulation runs pictured here, at time t_* roughly corresponding to the saturation time for nematic order.




FIG. 4. **A.** Mean nematic order parameter and polar order parameter plotted against bending rigidity at time t_* when all the simulations have reached long-range orientational order. **B.** and **C.** Median and the spread of the nematic (**B.**) and polar (**C.**) order values in 20-run ensembles at time t_* .




FIG. 5. **A.** Snapshots of the time-evolution of a system with $\tilde{\kappa} = 30$ and $\phi = 0.3$. At long times the order is polar, not nematic. **B.** Polar (cyan) and nematic (grey) order parameters for an ensemble of simulations ($n = 20$) with $\tilde{\kappa} = 70$. Magenta and blue curves represent the ensemble-mean polar and nematic order parameters, respectively, at each time. All systems reach long-range orientational order around the same time t_* , marked by the black dashed line.




FIG. 6. Ensemble-averaged ($n = 20$) nematic order parameter and polar order parameter. Error bars (appearing as widths of curves at larger time values) indicate the standard error of the mean. **A.** Global nematic order parameter $S(t)$ saturates at approximately the same time t_* for all $\tilde{\kappa}$, values indicating long-range orientational order. **B.** Global polar order parameter $P(t)$ saturates over time for all $\tilde{\kappa}$, but at later times for larger $\tilde{\kappa}$. **C.** A saturating exponential function $f(t) = r - ae^{-t/\tau}$ (dashed curves) is fit to the ensemble-averaged $P(t)$ data (solid curves). Insets show r (top) and τ (bottom) increasing with increasing $\tilde{\kappa}$.

LRO polar steady state monotonically decreases with increasing ϕ (Supplemental Material Fig. S4C [38]). This effect can be attributed to an increase in the rate of filament interactions for higher ϕ values, giving the system more opportunities for reorientation and hence reaching the steady state sooner.

Changing the system size L_x does not seem to have any significant effect on the typical time scales needed to reach long-range orientational order (t_*) or to reach the LRO active polar steady state (τ) (Supplemental Material Fig. S5 [38]). The plateau values of the polar order seen in Fig. 6B are also seen not to depend significantly on ϕ and L_x . We additionally note that incorporating anisotropic friction into the Brownian dynamics model [31] does not qualitatively change our results, as seen in the Supplemental Material Fig. S2 [38].

IV. DISCUSSION AND CONCLUSIONS

In this study, our objective was to investigate ordered states in the collective motions of self-propelled, semi-flexible filaments. With a microscopic model lacking attractive or aligning mechanisms between filament pairs, we find that crossovers are essential to the emergence of LRO states. In contrast, strictly two-dimensional confinement permits only locally ordered polar flocks. This finding is consistent with a recent experimental study of *E. coli* bacterial colonies on a surface, which are seen to form LRO nematic states when crossovers are allowed, and polar flocks when crossovers are suppressed [35].

When we allow filaments to interpenetrate in order to model crossovers, the symmetry of the LRO state that initially forms is nematic for more rigid filaments and

polar for more flexible ones. However, we find that the LRO nematic state, despite its experimental importance, is only a transient state in our simulations. All LRO states eventually evolve into LRO polar states, which we find to be the unique dynamical steady state. The typical time required for emergence of the polar steady state increases monotonically with filament rigidity.

Our results suggest that experimental realizations of LRO active nematic states in microtubule gliding assays are long-lived transient states rather than dynamical steady states. Due to factors such as ATP consumption, gliding assay experiments necessarily have finite lifetimes. For relatively rigid filaments such as microtubules, our model predicts that the timescale τ for emergence of polar order (and thus, the persistence of an apparently nematic state) may be much longer than the time $\sim t_*$ at which nematic order emerges from an isotropic initial condition. Comparing the typical length and active velocities of microtubules in gliding assay experiments [34] to our simulation, we estimate that microtubule bending rigidity corresponds to $\tilde{\kappa} \approx 250$, which is certainly in the high- $\tilde{\kappa}$ regime according to our results.

Our results also suggest that in the limit of $\tilde{\kappa} \rightarrow \infty$ (active rigid rods) the timescale for polar ordering may diverge, making the active nematic state dynamically stable. Long-lived, long-range nematic order in *E. coli* colonies might be interpreted as an effectively stable state in a system of effectively rigid rods [35].

It remains an open question how strongly our findings depend on the use of periodic boundary conditions and reduced system size compared to typical experiments. Surprisingly, however, we find that changing the system size does not have any significant effect on the typical timescale for emergence of the polar steady state. This

suggests that our simulations may be capturing the true physics of the bulk system. We also find that our choice of isotropic friction, at the level of individual beads, is not important to our findings, as introducing anisotropic friction does not qualitatively change our finding that the LRO polar order is the steady state (Supplemental Material Fig. S2 [38]).

A number of interesting questions are raised by the plateaus in polar order $P(t)$ that characterize the transient LRO nematic state in systems of more rigid filaments, before the late-time emergence of the LRO polar steady state (Fig. 5B and 6B). These plateaus are clear in $P(t)$ plots from individual simulation runs, (Fig. 5B) even though the ensemble-averaged $\langle P(t) \rangle$ appears to be monotonically increasing. We have confirmed that the P -value of this plateau has no significant dependence on the system size or on the area fraction of filaments (Supplemental Material Fig. S4, S5 [38]). The reason for this plateau in the polar order parameter's evolution, and the dependence of the plateau value on $\tilde{\kappa}$, remain open questions.

The variability that we observe in plateau P -values suggests that transient states may have wide variability in their orientational order, in an ensemble of active systems with identical parameters. As nematic order S saturates at earlier times, the polar order P that happens to develop simultaneously is apparently remembered over the long lifetime of the transient LRO nematic state. Increasing the strength of noise in the simulations may help to erase memory of the details of the nematic ordering process and thus reduce differences in order between transient states of different runs, resulting in a narrower distribution of P values.

We can understand the exponential saturation and the stability of the LRO polar state by making a comparison to the ordered state of a mean-field Ising model. Once the system attains nematic order, the conversion to a polar state is directly analogous to the ordering in an Ising model ferromagnet: the orientations of any pair of filaments are parallel or antiparallel, with the parallel configuration favored. Rather than interacting only with a few neighbors on a lattice, here the “spins” interact (through crossovers or collisions) with many other spins over the typical timescale of reorientation, suggesting that a mean-field approach may be appropriate. Mean-field ϕ^4 scalar field theory predicts that a perturbation

to the ordered (i.e. polar) state decays exponentially in time. More details can be found in Supplemental Material Section V [38]. This result suggests that mean-field approaches well-developed in the statistical physics of thermal systems may hold predictive power for gliding assays and other active systems in which the active agents can be described as “weakly interacting”.

In this work, we have only focused on the long-range orientationally ordered states with no density inhomogeneities and established that the LRO nematic state is only a transient state. But gliding assays also spontaneously form a variety of ordered states with non-uniform densities [21, 22, 31, 32, 34]. Our results highlight the potential importance of long-lived transients in active states generally, so it is of interest to investigate whether distinct active behaviors may be found in the time-evolution of other, inhomogeneous active states.

A full understanding of LRO active states, including the role of filament bending rigidity, requires detailed study of binary collisions [22, 32, 34], which is the subject of ongoing work. This understanding will complement our study in this work of the effect of area fraction ϕ . We find that increasing ϕ reduces the time taken required for emergence of the LRO polar steady state. This effect can be attributed to an increase in the rate of collisions with increasing ϕ , giving the system more opportunities to reorient filaments. Other realistic complications of interest, including polydispersity, chirality, and growth, make the active states of self-propelled, semiflexible filaments a subject of rich phenomenology even in the simplest ordered phases.

ACKNOWLEDGMENTS

This material is based upon work supported by the National Science Foundation under Grant No. DMR-2225543 and by the Hellman Fellows Fund. A portion of this work was carried out at the Advanced Research Computing at Hopkins (ARCH) core facility (rockfish.jhu.edu), which is supported by the National Science Foundation (NSF) grant number OAC 1920103. We are grateful to Kinjal Dasbiswas, Ajay Gopinathan, Linda Hirst, Fereshteh Memarian, Patrick Noerr, Nathan Prouse, Niranjana Sarpangala, and Fabian Jan Schwarzen-dahl for helpful discussions.

-
- [1] T. Vicsek and A. Zafeiris, Collective motion, *Physics Reports* **517**, 71 (2012).
 - [2] A. Cavagna, A. Cimorelli, I. Giardina, G. Parisi, R. Santagati, F. Stefanini, and M. Viale, Scale-free correlations in starling flocks, *Proceedings of the National Academy of Sciences* **107**, 11865 (2010).
 - [3] M. Ballerini, N. Cabibbo, R. Candelier, A. Cavagna, E. Cisbani, I. Giardina, V. Lecomte, A. Orlandi, G. Parisi, A. Procaccini, M. Viale, and V. Zdravkovic,

Interaction ruling animal collective behavior depends on topological rather than metric distance: Evidence from a field study, *Proceedings of the National Academy of Sciences* **105**, 1232 (2008).

- [4] J. L. Silverberg, M. Bierbaum, J. P. Sethna, and I. Cohen, Collective motion of humans in mosh and circle pits at heavy metal concerts, *Physical Review Letters* **110**, 228701 (2013).
- [5] T. Vicsek, A. Czirók, E. Ben-Jacob, I. Cohen, and

- O. Shochet, Novel type of phase transition in a system of self-driven particles, *Physical Review Letters* **75**, 1226 (1995).
- [6] M. E. Cates and J. Tailleur, Motility-induced phase separation, *Annual Review of Condensed Matter Physics* **6**, 219 (2015).
- [7] P. Digregorio, D. Levis, A. Suma, L. F. Cugliandolo, G. Gonnella, and I. Pagonabarraga, Full phase diagram of active brownian disks: From melting to motility-induced phase separation, *Physical Review Letters* **121**, 098003 (2018).
- [8] H. P. Zhang, A. Be'er, E.-L. Florin, and H. L. Swinney, Collective motion and density fluctuations in bacterial colonies, *Proceedings of the National Academy of Sciences* **107**, 13626 (2010).
- [9] R. M. Harshey and T. Matsuyama, Dimorphic transition in *Escherichia coli* and *Salmonella typhimurium*: surface-induced differentiation into hyperflagellate swarmer cells., *Proceedings of the National Academy of Sciences* **91**, 8631 (1994).
- [10] M. F. Copeland and D. B. Weibel, Bacterial swarming: a model system for studying dynamic self-assembly, *Soft Matter* **5**, 1174 (2009).
- [11] T. Sanchez, D. T. N. Chen, S. J. DeCamp, M. Heymann, and Z. Dogic, Spontaneous motion in hierarchically assembled active matter, *Nature* **491**, 431 (2012).
- [12] V. Schaller, C. Weber, C. Semmrich, E. Frey, and A. R. Bausch, Polar patterns of driven filaments, *Nature* **467**, 73 (2010).
- [13] T. Butt, T. Mufti, A. Humayun, P. B. Rosenthal, S. Khan, S. Khan, and J. E. Molloy, Myosin motors drive long range alignment of actin filaments2, *Journal of Biological Chemistry* **285**, 4964 (2010).
- [14] A. Doostmohammadi, J. Ignés-Mullol, J. M. Yeomans, and F. Sagués, Active nematics, *Nature Communications* **9**, 3246 (2018).
- [15] M. C. Marchetti, J. F. Joanny, S. Ramaswamy, T. B. Liverpool, J. Prost, M. Rao, and R. A. Simha, Hydrodynamics of soft active matter, *Reviews of Modern Physics* **85**, 1143 (2013).
- [16] R. D. Allen, D. G. Weiss, J. H. Hayden, D. T. Brown, H. Fujiwake, and M. Simpson, Gliding movement of and bidirectional transport along single native microtubules from squid axoplasm: evidence for an active role of microtubules in cytoplasmic transport., *The Journal of Cell Biology* **100**, 1736 (1985).
- [17] A. Hunt, F. Gittes, and J. Howard, The force exerted by a single kinesin molecule against a viscous load, *Biophysical Journal* **67**, 766 (1994).
- [18] R. Phillips, J. Kondev, J. Theriot, H. G. Garcia, and N. Orme, *Physical Biology of the Cell* (Garland Science, 2012).
- [19] D. Inoue, B. Mahmot, A. M. R. Kabir, T. I. Farhana, K. Tokuraku, K. Sada, A. Konagaya, and A. Kakugo, Depletion force induced collective motion of microtubules driven by kinesin, *Nanoscale* **7**, 18054 (2015).
- [20] A. Saito, T. I. Farhana, A. M. R. Kabir, D. Inoue, A. Konagaya, K. Sada, and A. Kakugo, Understanding the emergence of collective motion of microtubules driven by kinesins: role of concentration of microtubules and depletion force, *RSC Adv.* **7**, 13191 (2017).
- [21] Y. Sumino, K. H. Nagai, Y. Shitaka, D. Tanaka, K. Yoshikawa, H. Chaté, and K. Oiwa, Large-scale vortex lattice emerging from collectively moving microtubules, *Nature* **483**, 448 (2012).
- [22] L. Huber, R. Suzuki, T. Krüger, E. Frey, and A. R. Bausch, Emergence of coexisting ordered states in active matter systems, *Science* **361**, 255 (2018).
- [23] L. Onsager, The effects of shape on the interaction of colloidal particles, *Annals of the New York Academy of Sciences* **51**, 627 (1949).
- [24] A. Baskaran and M. C. Marchetti, Hydrodynamics of self-propelled hard rods, *Physical Review E* **77**, 011920 (2008).
- [25] F. Ginelli, F. Peruani, M. Bär, and H. Chaté, Large-scale collective properties of self-propelled rods, *Physical Review Letters* **104**, 184502 (2010).
- [26] H. H. Wensink, J. Dunkel, S. Heidenreich, K. Drescher, R. E. Goldstein, H. Löwen, and J. M. Yeomans, Meso-scale turbulence in living fluids, *Proceedings of the National Academy of Sciences* **109**, 14308 (2012).
- [27] M. Abkenar, K. Marx, T. Auth, and G. Gompper, Collective behavior of penetrable self-propelled rods in two dimensions, *Physical Review E* **88**, 062314 (2013).
- [28] A. Khokhlov and A. Semenov, Liquid-crystalline ordering in the solution of long persistent chains, *Physica A: Statistical Mechanics and its Applications* **108**, 546 (1981).
- [29] K. Kim, N. Yoshinaga, S. Bhattacharyya, H. Nakazawa, M. Umetsu, and W. Teizer, Large-scale chirality in an active layer of microtubules and kinesin motor proteins, *Soft Matter* **14**, 3221 (2018).
- [30] K. R. Prathyusha, S. Henkes, and R. Sknepnek, Dynamically generated patterns in dense suspensions of active filaments, *Physical Review E* **97**, 022606 (2018).
- [31] J. M. Moore, T. N. Thompson, M. A. Glaser, and M. D. Betterton, Collective motion of driven semiflexible filaments tuned by soft repulsion and stiffness, *Soft Matter* **16**, 9436 (2020).
- [32] S. Tanida, K. Furuta, K. Nishikawa, T. Hiraiwa, H. Kojima, K. Oiwa, and M. Sano, Gliding filament system giving both global orientational order and clusters in collective motion, *Physical Review E* **101**, 032607 (2020).
- [33] H. Zhou, W. Jung, T. I. Farhana, K. Fujimoto, T. Kim, and R. Yokokawa, Durability of aligned microtubules dependent on persistence length determines phase transition and pattern formation in collective motion, *ACS Nano* **16**, 14765 (2022).
- [34] F. L. Memarian, J. D. Lopes, F. J. Schwarzendahl, M. G. Athani, N. Sarpangala, A. Gopinathan, D. A. Beller, K. Dasbiswas, and L. S. Hirst, Active nematic order and dynamic lane formation of microtubules driven by membrane-bound diffusing motors, *Proceedings of the National Academy of Sciences* **118**, e2117107118 (2021).
- [35] D. Ghosh and X. Cheng, To cross or not to cross: Collective swimming of *Escherichia coli* under two-dimensional confinement, *Physical Review Research* **4**, 023105 (2022).
- [36] J. D. Weeks, D. Chandler, and H. C. Andersen, Role of repulsive forces in determining the equilibrium structure of simple liquids, *The Journal of Chemical Physics* **54**, 5237 (1971).
- [37] A. C. Brañka and D. M. Heyes, Algorithms for brownian dynamics computer simulations: Multivariable case, *Physical Review E* **60**, 2381 (1999).
- [38] See Supplemental Material for details.
- [39] S. Weitz, A. Deutsch, and F. Peruani, Self-propelled rods exhibit a phase-separated state characterized by the presence of active stresses and the ejection of polar clusters, *Physical Review E* **92**, 012322 (2015).

- [40] F. Gittes, B. Mickey, J. Nettleton, and J. Howard, Flexural rigidity of microtubules and actin filaments measured from thermal fluctuations in shape., *Journal of Cell Biology* **120**, 923 (1993).

Supplemental Material: Symmetry and stability of orientationally ordered collective motions of self-propelled, semiflexible filaments

Madhuvanthi Guruprasad Athani¹ and Daniel A. Beller^{1,*}

¹*Department of Physics and Astronomy, Johns Hopkins University, Baltimore, Maryland 21218, USA*

I. MODEL DETAILS

Each semiflexible filament is modeled as a bead-spring chain composed of circular disks (beads) connected by Hookean springs with spring stiffness k . If $\mathbf{r} = r\hat{\mathbf{r}}$ is the separation vector between the centers of any two connected beads and r_0 is the equilibrium spring distance, then the spring force is given by:

$$\mathbf{F}_{\text{stretch}} = -k(r - r_0)\hat{\mathbf{r}} \quad (\text{S.1})$$

A discrete approximation to the bending energy for a semiflexible chain is $\frac{1}{2}\kappa\frac{\Sigma_i(\Delta\theta_i)^2}{\Delta s}$ where $\Delta\theta_i = \theta_i - \theta_{i-1}$ is the angle change between points separated by a small distance Δs along the contour and $\kappa = \frac{1}{2}l_p k_B T$ is the bending stiffness where l_p is the persistence length of the chain [1]. We incorporate this bending energy as a potential acting between every set of three consecutive beads ($\Delta s \approx 2r_0$) that penalizes the bending of the chain.

$$U_{\text{bend}} = \frac{1}{2}\tilde{\kappa}(\pi - \theta)^2 \quad (\text{S.2})$$

$\tilde{\kappa} = \frac{\kappa}{\Delta s}$ has the units of energy and will be referred to as the bending stiffness hereafter and θ is the bond angle between two adjacent linear springs (Fig. S1).




FIG. S1. Schematic illustration of forces in the active bead-spring chain model. (Left) Illustration of active self-propulsion and WCA repulsion between pairs of chains. (Right) A zoomed-in view of the region enclosed by the dashed lines at left, to illustrate the stretching and bending spring forces.

The total force excluding viscous drag is given by

$$\mathbf{F}_{\text{total}} = \mathbf{F}_{\text{stretch}} - \nabla_{\theta} U_{\text{bend}} + \mathbf{F}_{\text{active}} - \nabla U_{\text{WCA}} \quad (\text{S.3})$$

where U_{WCA} is given by Eq. 1 (main text). The Brownian (over-damped) dynamics [2] updates bead positions through

$$\mathbf{x}(t + dt) = \mathbf{x}(t) + \mathbf{v}dt + \sqrt{2Ddt}\boldsymbol{\xi}(t) \quad (\text{S.4})$$

$$\mathbf{v} = \mathbf{F}_{\text{total}}/\gamma \quad (\text{S.5})$$

Here \mathbf{v} is the velocity of the bead, γ is the drag coefficient, and $\boldsymbol{\xi}(t)$ is the stochastic noise term representing random fluctuations in bead positions, drawn from a zero-mean Gaussian white noise distribution.

* d.a.beller@jhu.edu

II. ANISOTROPIC FRICTION

The model described previously assumes isotropic friction on the beads. But as these beads form long filaments with a high aspect ratio, we examine here whether our results are significantly changed by friction anisotropy. For this purpose, we replace Eq. S.5 with an anisotropic generalization for a particular bead α ,

$$v_i^\alpha = M_{ij}^\alpha \cdot F_{\text{total},i}^\alpha, \quad (\text{S.6})$$

where M_{ij}^α is the anisotropic mobility tensor of the α th bead [2]:

$$M_{ij}^\alpha = \frac{1}{\gamma_\parallel} t_i^\alpha \otimes t_j^\alpha + \frac{1}{\gamma_\perp} (\delta_{ij} - t_i^\alpha \otimes t_j^\alpha). \quad (\text{S.7})$$

Here, γ_\perp and γ_\parallel are the perpendicular and parallel drag coefficients. The symbol \otimes represents the outer product, and \mathbf{t}^α is the tangent unit vector to bead α , found by

$$\mathbf{t}^\alpha = \frac{\mathbf{r}^\alpha + \mathbf{r}^{\alpha+1}}{|\mathbf{r}^\alpha + \mathbf{r}^{\alpha+1}|} \quad (\text{S.8})$$

where \mathbf{r}^α is the separation vector between the centers of $(\alpha - 1)$ th and α th connected beads, and $1 < \alpha < N$ if there are N beads in a chain. For a bead that is located at the ends, $\mathbf{t}^1 = \mathbf{r}^2$ and $\mathbf{t}^N = \mathbf{r}^N$. The diffusion coefficient is also anisotropic and for a wormlike chain, it is given by $D_{ij}^\alpha = k_B T M_{ij}^\alpha$ [2]. This can be used in Equation S.4 to compute the positions of beads at the next timestep.




FIG. S2. Nematic and polar order parameters are plotted for an ensemble of simulations with $\tilde{\kappa} = 30$, $L_x = 100$, $\phi = 0.3$. **A.** Isotropic friction. **B.** Anisotropic friction with an anisotropy $\gamma_\perp = 2\gamma_\parallel$. Black dashed line represents time t_* as described in the main text.

Using anisotropic friction of $\gamma_\perp = 2\gamma_\parallel$ in a 20-run ensemble of simulations with $\tilde{\kappa} = 30$ (Fig. S2), we see no qualitative difference compared to simulations with isotropic friction as both Fig. S2A and B reach an LRO polar steady state after being in a transient nematic state at intermediate times. From $\langle S(t) \rangle$ values of both the cases, we can see that the case with isotropic friction reaches the LRO nematic state at a slightly earlier time compared to the case with anisotropic friction. However, the case with anisotropic friction is seen to reach the LRO polar steady state sooner. We also find that anisotropic friction reduces differences in polar order between transient states of different runs during intermediate times, resulting in a narrower distribution of P -values. There is also a slight increase in $\langle P(t) \rangle$ at these intermediate times compared to an ensemble with isotropic friction.

III. EFFECT OF AREA FRACTION

We investigated the effect of changing the area fraction of filaments on LRO state of the system and its steady state. Fig. S3 shows snapshots of simulations with different area fractions after they have reached orientational order.

All the simulations shown here have filaments with $\tilde{\kappa} = 10$ and U_{WCA} parameters $\epsilon = 10^{-8}$ and $r_{\text{shift}} = 0.1$ (Fig. 2C (main text)). From these snapshots, we see that the symmetry of the LRO state reached for all the area fractions is polar which is consistent with the polar order seen for $\tilde{\kappa} = 10$ in (Fig. 3 (main text)) where $\phi = 0.3$. So increasing the area fraction to $\phi = 0.5$ and $\phi = 0.7$ does not change the type of initial orientational order reached in the system.




FIG. S3. Changing area fraction ϕ of filaments. All the snapshots are taken at a time when the system has reached orientational order, which is polar for all cases shown here. The filaments have $\tilde{\kappa} = 10$ and the system size $L_x = 250$.

To understand the steady state behavior of the system, we looked at the order of the system after a long time compared to t_* . The global nematic order parameter in Fig. S4A reveals that all the simulations reach an orientational order but the system with the lowest area fraction ($\phi = 0.1$) takes the longest time to reach the global orientational order. This is presumably due to filament reorientation interactions being less frequent in the lower-density system. The LRO polar state is still the steady state of the system as seen from Fig. S4B. The system takes lesser time to get to the LRO polar state when the area fraction is higher. Fitting a saturating exponential function $f(t) = r - ae^{-t/\tau}$ to the mean polar order parameter, we see that τ decreases with increasing area fraction (Fig. S4C lower inset) whereas saturation parameter r is not seen to follow any trend with change in area fraction (Fig. S4C upper inset).




FIG. S4. Dependence of orientational order time-evolution upon area fraction ϕ . Orientational order parameters are averaged over 20-run ensembles with $\tilde{\kappa} = 30$ and with various ϕ values. **A.** Nematic order time-evolution. **B.** Polar order time-evolution, with logarithmic scale on the time axis. **C.** Polar order time-evolution (solid curves), with linear scale on the time axis. Dashed curves are fits of $f(t) = r - ae^{-t/\tau}$ to the data. Insets show that r does not depend significantly on ϕ (upper inset), while τ decreases with increasing ϕ (lower inset). Time is plotted in units of ℓ/v_0 .

IV. EFFECT OF SIMULATION BOX SIZE

We examine the effect of changing the simulation box size L_x on the LRO states of the system. We fix $\tilde{\kappa} = 30$ throughout this ensemble of 20 runs for four different L_x values. From Fig. S5A, we see that the initial LRO state is reached at roughly the same time for all system sizes, though there is some trend of larger systems taking slightly more time to reach the LRO state. From Fig. S5B we can see that the time to reach the LRO polar steady state does

not depend significantly on the system size. Interestingly, we can also see that the plateau value of the polar order in Fig. S5B does not show any monotonic trend with increasing system size.




FIG. S5. Nematic (left) and polar (right) order parameters are plotted for a 20-run ensemble with $\tilde{\kappa} = 30$ and different simulation box sizes L_x .

V. STABILITY OF THE LRO POLAR ORDER: MEAN FIELD THEORY

We can understand the exponential saturation to a polar steady state and its stability by comparing our system to the mean field theory of an Ising model. We propose that the late-time polar ordering process of an LRO state is analogous to the relaxation to thermal equilibrium of an Ising-model ferromagnet. Let φ be the mean order parameter, which is the net magnetization in the Ising model and the polar order parameter P in our LRO active system. For the Ising model ferromagnet, Model A relaxation for this non-conserved order parameter then gives

$$\frac{\partial \varphi}{\partial t} = -\Gamma \frac{\delta F}{\delta \varphi} \quad (\text{S.9})$$

where F is the free energy and Γ is a constant coefficient. The corresponding free energy density f has the following form in φ^4 scalar field theory:

$$f(\phi, T) = a(T - T_c)\varphi^2 + u\varphi^4 \quad (\text{S.10})$$

The first derivative with respect to the order parameter can be written

$$\frac{\partial f}{\partial \varphi} = 2a(T - T_c)\varphi + 4u\varphi^3 = 2a(T - T_c)\varphi \left(1 + \frac{4u\varphi^2}{2a(T - T_c)} \right). \quad (\text{S.11})$$

In the uniform ordered state ($T_c > T$), equilibrium is given by

$$\frac{\partial f}{\partial \varphi} = 0 \implies \varphi_0^2 = 2a(T_c - T)/4u. \quad (\text{S.12})$$

From Eq. S.9, S.11 and S.12,

$$\frac{\partial \varphi}{\partial t} = 2\Gamma a(T_c - T)\varphi \left(1 - \frac{\varphi^2}{\varphi_0^2} \right) \quad (\text{S.13})$$

Consider a small perturbation $\delta\varphi$ to the ordered state φ_0 .

$$\varphi = \varphi_0 + \delta\varphi \implies \varphi^2 \approx \varphi_0^2 \left(1 + \frac{2\delta\varphi}{\varphi_0} \right) \implies 1 - \frac{\varphi^2}{\varphi_0^2} \approx -\frac{2\delta\varphi}{\varphi_0}. \quad (\text{S.14})$$

Substituting Eq. S.14 into Eq. S.13

$$\frac{\partial(\delta\varphi)}{\partial t} = \frac{\partial(\varphi - \varphi_0)}{\partial t} = -4\Gamma a(T_c - T)\delta\varphi + O(\delta\varphi^2). \quad (\text{S.15})$$

Therefore, to lowest order in $\delta\varphi$, the perturbation from equilibrium decays exponentially in time,

$$\frac{\partial(\delta\varphi)}{\partial t} \propto -\delta\varphi. \quad (\text{S.16})$$

It is important to note that our active system is far from equilibrium and thus not governed solely by a free energy. Nonetheless, we conjecture that its close resemblance to the ordering process of an Ising-model ferromagnet rationalizes our observation of an exponential approach of polar order $P(t)$ to its saturation value at late times.

-
- [1] C. P. Broedersz and F. C. MacKintosh, *Reviews of Modern Physics* **86**, 995 (2014).
 - [2] J. M. Moore, T. N. Thompson, M. A. Glaser, and M. D. Betterton, *Soft Matter* **16**, 9436 (2020).