# Strong tractability for multivariate integration in a subspace of the Wiener algebra<sup>\*</sup>

Takashi Goda<sup>†</sup>

June 5, 2023

#### Abstract

Building upon recent work by the author, we prove that multivariate integration in the following subspace of the Wiener algebra over  $[0, 1)^d$  is strongly polynomially tractable:

$$F_d := \left\{ f \in C([0,1)^d) \, \middle| \, \|f\| := \sum_{\boldsymbol{k} \in \mathbb{Z}^d} |\hat{f}(\boldsymbol{k})| \max\left( \operatorname{width}(\operatorname{supp}(\boldsymbol{k})), \min_{j \in \operatorname{supp}(\boldsymbol{k})} \log |k_j| \right) < \infty \right\},$$

with  $\hat{f}(\mathbf{k})$  being the **k**-th Fourier coefficient of f, supp $(\mathbf{k}) := \{j \in \{1, \ldots, d\} \mid k_j \neq 0\}$ , and width  $: 2^{\{1,\ldots,d\}} \rightarrow \{1,\ldots,d\}$  being defined by

width(u) := 
$$\max_{j \in u} j - \min_{j \in u} j + 1$$
,

for non-empty subset  $u \subseteq \{1, \ldots, d\}$  and width $(\emptyset) := 1$ . Strong polynomial tractability is achieved by an explicit quasi-Monte Carlo rule using a multiset union of Korobov's *p*-sets. We also show that, if we replace width(supp( $\mathbf{k}$ )) with 1 for all  $\mathbf{k} \in \mathbb{Z}^d$  in the above definition of norm, multivariate integration is polynomially tractable but not strongly polynomially tractable.

**Keywords:** multivariate integration, polynomial tractability, Wiener algebra, quasi-Monte Carlo, exponential sum

AMS subject classifications: 41A55, 41A58, 42B05, 65D30, 65D32

#### **1** Introduction and main results

This paper concerns numerical integration for multivariate functions defined over the *d*-dimensional unit cube. For a Riemann integrable function  $f : [0, 1)^d \to \mathbb{R}$ , we approximate its integral

$$I_d(f) = \int_{[0,1)^d} f(\boldsymbol{x}) \, \mathrm{d}\boldsymbol{x}$$

by

$$Q_{d,n}(f) = \sum_{h=0}^{n-1} w_h f(\boldsymbol{x}_h)$$

<sup>\*</sup>The work of the author is supported by JSPS KAKENHI Grant Number 23K03210.

<sup>&</sup>lt;sup>†</sup>School of Engineering, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656, Japan (goda@frcer.t.u-tokyo.ac.jp)

with sets of *n* sampling points  $\{x_0, \ldots, x_{n-1}\} \subset [0, 1)^d$  and associated weights  $\{w_0, \ldots, w_{n-1}\}$ . Quasi-Monte Carlo (QMC) rule denotes a special case of  $Q_{d,n}$  where all the weights  $w_h$  are equal to 1/n. The worst-case error of an algorithm  $Q_{d,N}$  in a Banach space F with norm  $\|\cdot\|$  is defined by

$$e^{\mathrm{wor}}(F, Q_{d,n}) := \sup_{f \in F, \|f\| \le 1} |I_d(f) - Q_{d,n}(f)|.$$

In the field of information-based complexity [11, 12, 16], we are interested in how the *information* complexity  $n(\varepsilon, d, F)$  grows in the reciprocal of the error tolerance  $\varepsilon \in (0, 1)$  and the dimension d. Here, the information complexity is defined as the minimum number of function values, among all possible  $Q_{d,n}$ , needed to make the worst-case error in F no greater than  $\varepsilon$ , that is,

$$n(\varepsilon, d, F) := \min\{n \in \mathbb{N} \mid \exists Q_{d,n} : e^{\mathrm{wor}}(F, Q_{d,n}) \le \varepsilon\}.$$

In a recent work by the author [6], it has been proven that the information complexity for the following unweighted subspace of the Wiener algebra grows only polynomially both in  $\varepsilon^{-1}$  and d:

$$F_d^1 := \left\{ f \in C([0,1)^d) \, \middle| \, \|f\| := \sum_{\boldsymbol{k} \in \mathbb{Z}^d} |\hat{f}(\boldsymbol{k})| \max\left(1, \min_{j \in \text{supp}(\boldsymbol{k})} \log |k_j|\right) < \infty \right\},$$

with  $\hat{f}(\mathbf{k})$  being the **k**-th Fourier coefficient of f, i.e.,

$$\hat{f}(\boldsymbol{k}) = \int_{[0,1)^d} f(\boldsymbol{x}) \exp(-2\pi i \boldsymbol{k} \cdot \boldsymbol{x}) \, \mathrm{d}\boldsymbol{x},$$

and  $\operatorname{supp}(\mathbf{k}) := \{j \in \{1, \ldots, d\} \mid k_j \neq 0\}$ . More precisely, it has been shown that an upper bound  $n(\varepsilon, d, F_d^1) \leq C_1 \varepsilon^{-3} d^3$  holds for a positive constant  $C_1$ , concluding that the problem of multivariate integration in  $F_d^1$  is *polynomially tractable*. We refer to [9, 10] for more recent progress on this line of research. In this context, an *unweighted* function space F refers to a space where all variables and groups of variables play an equal role. Therefore, for any permutation matrix  $\pi$  and  $f \in F$ , it holds that  $f \circ \pi \in F$  and  $||f \circ \pi|| = ||f||$ . The result presented in [6] builds upon the work of Dick [1], who established polynomial tractability for multivariate integration in the intersection of the Wiener algebra and an unweighted space of Hölder continuous functions.

As a continuation of [6], we prove the following result in this paper:

**Theorem 1.** Let  $F_d^2$  be a subspace of the Wiener algebra defined by

$$F_d^2 := \left\{ f \in C([0,1)^d) \, \middle| \, \|f\| := \sum_{\boldsymbol{k} \in \mathbb{Z}^d} |\hat{f}(\boldsymbol{k})| \max\left( \operatorname{width}(\operatorname{supp}(\boldsymbol{k})), \min_{j \in \operatorname{supp}(\boldsymbol{k})} \log |k_j| \right) < \infty \right\},$$

where width :  $2^{\{1,\ldots,d\}} \rightarrow \{1,\ldots,d\}$  is defined by

width(u) := 
$$\max_{j \in u} j - \min_{j \in u} j + 1$$
,

for non-empty subset  $u \subseteq \{1, \ldots, d\}$ , and width $(\emptyset) = 1$ . Then, there exists a positive constant  $C_2$  such that, for any  $d \in \mathbb{N}$  and  $\varepsilon \in (0, 1)$ , we have

$$n(\varepsilon, d, F_d^2) \le C_2 \varepsilon^{-3} / (\log \varepsilon^{-1})$$

In comparison to the result of [6] for  $F_d^1$ , by replacing 1 (the first argument in taking the maximum for each  $\mathbf{k}$ ) with width(supp( $\mathbf{k}$ )) in the definition of norms, the polynomial dependence of the information complexity on the dimension d does not show up anymore, meaning that the problem of multivariate integration in  $F_d^2$  is strongly polynomially tractable. This result is strengthened by the following theorem on the former space  $F_d^1$ .

**Theorem 2.** For any linear algorithm  $Q_{d,n}$  using n function values, we have  $e^{\text{wor}}(F_d^1, Q_{d,n}) \ge d/(2n^2)$  for any  $d \in \mathbb{N}$  and n > 2d.

Note that there is a significant gap between the lower bound on the worst-case error obtained above and the upper bound of order  $dn^{-1/3}$  shown in [6]. Nevertheless, this result implies that a dependence of the information complexity on the dimension d cannot be eliminated for  $F_d^1$ . Therefore, the problem of multivariate integration in  $F_d^1$  is polynomially tractable but *not* strongly polynomially tractable. As a future research direction, it would be interesting to study whether an intermediate space between  $F_d^1$  and  $F_d^2$  still exhibits strong polynomial tractability for multivariate integration. As we have  $1 \leq |\operatorname{supp}(\mathbf{k})| \leq \operatorname{width}(\operatorname{supp}(\mathbf{k}))$  for all  $\mathbf{k} \in \mathbb{Z}^d$  when defining  $|\operatorname{supp}(\mathbf{0})| = 1$ , one of the most natural spaces we can consider is an unweighted space

$$F_d^3 := \left\{ f \in C([0,1)^d) \, \middle| \, \|f\| := \sum_{\boldsymbol{k} \in \mathbb{Z}^d} |\hat{f}(\boldsymbol{k})| \max\left(|\operatorname{supp}(\boldsymbol{k})|, \min_{j \in \operatorname{supp}(\boldsymbol{k})} \log |k_j|\right) < \infty \right\}.$$

Note that, although the space  $F_d^2$  is weighted, it remains invariant under the reversion of the variables, i.e., if  $f \in F_d^2$ , then we have  $g \in F_d^2$  and ||f|| = ||g|| where  $g(x_1, \ldots, x_d) = f(x_d, \ldots, x_1)$ . This is in contrast to many existing results on strong polynomial tractability for multivariate integration in the worst-case setting, where weight parameters are introduced to model the relative importance of each group of variables, and variables are typically assumed ordered in decreasing importance order. See [2, 5, 11, 12, 15] among many others. In fact, it seems not possible to characterize the space  $F_d^2$  in such a way. The author believes that further tractability studies in subspaces of the Wiener algebra will offer new insights into the field of information-based complexity, particularly regarding (strong) polynomial tractability in (un)weighted spaces.

#### 2 Proof of Theorem 1

This section is devoted to proving Theorem 1 by providing an explicit QMC rule that attains the desired worst-case error bound. The QMC rule considered here is exactly the same as the one discussed in [6]. For an integer  $m \ge 2$ , let

$$\mathbb{P}_m := \{ \lceil m/2 \rceil$$

It is known that there exist constants  $c_{\mathbb{P}}$  and  $C_{\mathbb{P}}$  with  $0 < c_{\mathbb{P}} < \min(1, C_{\mathbb{P}})$  such that

$$c_{\mathbb{P}}\frac{m}{\log m} \le |\mathbb{P}_m| \le C_{\mathbb{P}}\frac{m}{\log m},\tag{1}$$

for all  $m \ge 2$ , see [13, Corollaries 1–3]. Now, given an integer  $m \ge 2$ , we define two different point sets as multiset unions:

$$P_{d,m}^1 = \bigcup_{p \in \mathbb{P}_m} S_{d,p}$$
 and  $P_{d,m}^2 = \bigcup_{p \in \mathbb{P}_m} T_{d,p}$ 

where  $S_{d,p} = \{ \boldsymbol{x}_h^{(p)} \mid 0 \le h < p^2 \}$  and  $T_{d,p} = \{ \boldsymbol{x}_{h,\ell}^{(p)} \mid 0 \le h, \ell are sets with <math>p^2$  points known as *Korobov's p-sets* [1, 5, 7, 8]. These point sets are defined as follows:

$$\boldsymbol{x}_{h}^{(p)} = \left(\left\{\frac{h}{p^{2}}\right\}, \left\{\frac{h^{2}}{p^{2}}\right\}, \dots, \left\{\frac{h^{d}}{p^{2}}\right\}\right),$$

and

$$\boldsymbol{x}_{h,\ell}^{(p)} = \left(\left\{\frac{h\ell}{p}\right\}, \left\{\frac{h\ell^2}{p}\right\}, \dots, \left\{\frac{h\ell^d}{p}\right\}\right),$$

respectively, where we write  $\{x\} = x - \lfloor x \rfloor$  to denote the fractional part of a non-negative real number x. It is important to note that taking the multiset unions of Korobov's p-sets with different primes p is crucial in our error analysis. Trivially we have

$$|P_{d,m}^1| = |P_{d,m}^2| = \sum_{p \in \mathbb{P}_m} p^2.$$

The following result on the exponential sums refines the known results from [7, Lemmas 4.5 & 4.6] as well as [5, Lemmas 4.4 & 4.5].

**Lemma 3.** Let  $d \in \mathbb{N}$  and p be a prime with  $p \geq d$ . For any  $\mathbf{k} \in \mathbb{Z}^d \setminus \{\mathbf{0}\}$  such that there exists at least one index  $j^* \in \{1, \ldots, d\}$  where  $k_{j^*}$  is not divisible by p, i.e.,  $p \nmid \mathbf{k}$ , the following bounds hold:

$$\left. \frac{1}{p^2} \sum_{h=0}^{p^2-1} \exp\left(2\pi i \boldsymbol{k} \cdot \boldsymbol{x}_h^{(p)}\right) \right| \le \frac{\operatorname{width}(\operatorname{supp}(\boldsymbol{k}))}{p},$$

and

$$\frac{1}{p^2} \sum_{h,\ell=0}^{p-1} \exp\left(2\pi i \boldsymbol{k} \cdot \boldsymbol{x}_{h,\ell}^{(p)}\right) \leq \frac{\operatorname{width}(\operatorname{supp}(\boldsymbol{k}))}{p}.$$

*Proof.* Let us consider the first bound. As we have  $\{0, \ldots, p^2 - 1\} = \{h_0 + h_1p \mid 0 \le h_0, h_1 < p\}$  and, for each pair of  $h_0, h_1 \in \{0, \ldots, p - 1\}$ , it holds that

$$\exp\left(2\pi i\boldsymbol{k}\cdot\boldsymbol{x}_{h_0+h_1p}^{(p)}\right) = \exp\left(\frac{2\pi i}{p^2}\sum_{j\in\operatorname{supp}(\boldsymbol{k})}k_j(h_0+h_1p)^j\right)$$
$$= \exp\left(\frac{2\pi i}{p^2}\sum_{j\in\operatorname{supp}(\boldsymbol{k})}k_j\sum_{a=0}^j \binom{j}{a}h_0^a(h_1p)^{j-a}\right)$$
$$= \exp\left(\frac{2\pi i}{p^2}\sum_{j\in\operatorname{supp}(\boldsymbol{k})}k_j(h_0^j+jh_0^{j-1}h_1p)\right),$$

we obtain

$$\begin{aligned} \left| \frac{1}{p^2} \sum_{h=0}^{p^2 - 1} \exp\left(2\pi i \mathbf{k} \cdot \mathbf{x}_h^{(p)}\right) \right| &= \left| \frac{1}{p^2} \sum_{h_0, h_1 = 0}^{p-1} \exp\left(\frac{2\pi i}{p^2} \sum_{j \in \text{supp}(\mathbf{k})} k_j (h_0^j + j h_0^{j-1} h_1 p)\right) \right| \\ &= \left| \frac{1}{p} \sum_{h_0 = 0}^{p-1} \exp\left(\frac{2\pi i}{p^2} \sum_{j \in \text{supp}(\mathbf{k})} k_j h_0^j\right) \frac{1}{p} \sum_{h_1 = 0}^{p-1} \exp\left(\frac{2\pi i h_1}{p} \sum_{j \in \text{supp}(\mathbf{k})} k_j j h_0^{j-1}\right) \right| \\ &\leq \frac{1}{p} \sum_{h_0 = 0}^{p-1} \left| \frac{1}{p} \sum_{h_1 = 0}^{p-1} \exp\left(\frac{2\pi i h_1}{p} \sum_{j \in \text{supp}(\mathbf{k})} k_j j h_0^{j-1}\right) \right| \\ &= \frac{1}{p} \sum_{\sum_{j \in \text{supp}(\mathbf{k})} \sum_{k_j j h_0^{j-1} \equiv 0 \pmod{p}}^{p-1} (\text{mod } p) \end{aligned}$$

where the last equality follows from the well-known character property for the trigonometric functions [3, Lemma 4.3]. Here, by denoting  $j_{\min} = \min_{j \in \text{supp}(\mathbf{k})} j$  and  $j_{\max} = \max_{j \in \text{supp}(\mathbf{k})} j$ , we have

$$\sum_{\substack{j \in \text{supp}(\mathbf{k}) \\ j \in \text{supp}(\mathbf{k})}} k_j j h_0^{j-1} = \sum_{\substack{j=j_{\min} \\ j \in \text{supp}(\mathbf{k})}}^{j_{\max}} k_j j h_0^{j-1} = h_0^{j_{\min}-1} \sum_{\substack{j=j_{\min} \\ j \in \text{supp}(\mathbf{k})}}^{j_{\max}} k_j j h_0^{j-j_{\min}}.$$

As the last sum over j is a polynomial in  $h_0$  with degree  $j_{\text{max}} - j_{\text{min}}$ , the number of solutions of the congruence  $\sum_{j \in \text{supp}(\mathbf{k})} k_j j h_0^{j-1} \equiv 0 \pmod{p}$  is at most  $j_{\text{max}} - j_{\text{min}} + 1 = \text{width}(\text{supp}(\mathbf{k}))$ . Thus the result follows. Since the second bound can be proven in the same manner, we omit the details.

Note that, if  $k_j$  is divisible by p for all j, i.e.,  $p \mid \mathbf{k}$ , then we only have a trivial bound on the exponential sum, which is 1. Using this refined result, we obtain the following bounds on the exponential sums for our point sets  $P_{d,m}^1$  and  $P_{d,m}^2$ .

**Corollary 4.** Let  $d \in \mathbb{N}$  and  $m \geq 2$  with  $\min_{p \in \mathbb{P}_m} p \geq d$ . For any  $\mathbf{k} \in \mathbb{Z}^d \setminus \{\mathbf{0}\}$ , it holds that

$$\left|\frac{1}{|P_{d,m}^{1}|}\sum_{p\in\mathbb{P}_{m}}\sum_{h=0}^{p^{2}-1}\exp\left(2\pi i\boldsymbol{k}\cdot\boldsymbol{x}_{h}^{(p)}\right)\right| \leq \frac{1}{m}\left(4\operatorname{width}(\operatorname{supp}(\boldsymbol{k})) + \frac{8}{c_{\mathbb{P}}}\min_{j\in\operatorname{supp}(\boldsymbol{k})}\log|k_{j}|\right)$$

and

$$\left|\frac{1}{|P_{d,m}^2|}\sum_{p\in\mathbb{P}_m}\sum_{h,\ell=0}^{p-1}\exp\left(2\pi i\boldsymbol{k}\cdot\boldsymbol{x}_{h,\ell}^{(p)}\right)\right| \leq \frac{1}{m}\left(4\operatorname{width}(\operatorname{supp}(\boldsymbol{k})) + \frac{8}{c_{\mathbb{P}}}\min_{j\in\operatorname{supp}(\boldsymbol{k})}\log|k_j|\right)$$

*Proof.* The following proof for the first bound is similar to that of [6, Corollary 2.3], and the second bound can be proven in a similar way, so we omit the details. Using Lemma 3, we have

$$\begin{aligned} \left| \frac{1}{|P_{d,m}^{1}|} \sum_{p \in \mathbb{P}_{m}} \sum_{h=0}^{p^{2}-1} \exp\left(2\pi i \boldsymbol{k} \cdot \boldsymbol{x}_{h}^{(p)}\right) \right| &\leq \frac{1}{|P_{d,m}^{1}|} \sum_{p \in \mathbb{P}_{m}} p \operatorname{width}(\operatorname{supp}(\boldsymbol{k})) + \frac{1}{|P_{d,m}^{1}|} \sum_{\substack{p \in \mathbb{P}_{m}\\p \nmid \boldsymbol{k}}} p^{2} \\ &\leq \frac{1}{|P_{d,m}^{1}|} \sum_{\substack{p \in \mathbb{P}_{m}\\p \nmid \boldsymbol{k}}} p \operatorname{width}(\operatorname{supp}(\boldsymbol{k})) + \frac{m^{2}}{|P_{d,m}^{1}|} \sum_{\substack{p \in \mathbb{P}_{m}\\p \mid \boldsymbol{k}}} 1 \\ &\leq \frac{m|\mathbb{P}_{m}|}{(m/2)^{2}|\mathbb{P}_{m}|} \operatorname{width}(\operatorname{supp}(\boldsymbol{k})) + \frac{m^{2}}{(m/2)^{2}|\mathbb{P}_{m}|} \sum_{\substack{p \in \mathbb{P}_{m}\\p \mid \boldsymbol{k}}} 1 \\ &\leq \frac{4}{m} \operatorname{width}(\operatorname{supp}(\boldsymbol{k})) + \frac{4\log m}{c_{\mathbb{P}}m} \sum_{\substack{p \in \mathbb{P}_{m}\\p \mid \boldsymbol{k}}} 1, \end{aligned}$$

where the last inequality follows from (1). To give a bound on the last sum over  $p \in \mathbb{P}_m$  which divides k, we use the fact that, for any integers  $k, n \in \mathbb{N}$ , k has at most  $\log_n k$  prime divisors larger than or equal to n. With  $\mathbb{I}(\cdot)$  denoting the indicator function, for any index  $j^* \in \text{supp}(k)$ , we get

$$\sum_{\substack{p \in \mathbb{P}_m \\ p \mid \boldsymbol{k}}} 1 = \sum_{p \in \mathbb{P}_m} \prod_{j \in \text{supp}(\boldsymbol{k})} \mathbb{I}(p \mid k_j) \le \sum_{p \in \mathbb{P}_m} \mathbb{I}(p \mid k_{j^*}) \le \log_{\lceil m/2 \rceil + 1} |k_{j^*}| \le \frac{2 \log |k_{j^*}|}{\log m}.$$

Since this inequality applies to any index  $j^* \in \text{supp}(\mathbf{k})$ , it holds that

$$\sum_{\substack{p \in \mathbb{P}_m \\ p \mid \mathbf{k}}} 1 \le \frac{2}{\log m} \min_{j \in \text{supp}(\mathbf{k})} \log |k_j|.$$

This completes the proof.

Now we are ready to prove Theorem 1.

Proof of Theorem 1. Since any function  $f \in F_d^2$  has an absolutely convergent Fourier series, by letting  $Q_{d,n}$  being the QMC rule using  $P_{d,m}^1$  (or  $P_{d,m}^2$ ) for some  $m \ge 2$  with  $\min_{p \in \mathbb{P}_m} p \ge d$ , it follows from Corollary 4 that, with n equal to  $\sum_{p \in \mathbb{P}_m} p^2$ ,

$$\begin{aligned} |I_d(f) - Q_{d,n}(f)| &= \left| I_d(f) - \frac{1}{|P_{d,m}^1|} \sum_{p \in \mathbb{P}_m} \sum_{h=0}^{p^2 - 1} f(\boldsymbol{x}_h^{(p)}) \right| \\ &= \left| \hat{f}(\mathbf{0}) - \frac{1}{|P_{d,m}^1|} \sum_{p \in \mathbb{P}_m} \sum_{h=0}^{p^2 - 1} \sum_{\boldsymbol{k} \in \mathbb{Z}^d} \hat{f}(\boldsymbol{k}) \exp\left(2\pi i \boldsymbol{k} \cdot \boldsymbol{x}_h^{(p)}\right) \right| \\ &= \left| \sum_{\boldsymbol{k} \in \mathbb{Z}^d \setminus \{\mathbf{0}\}} \hat{f}(\boldsymbol{k}) \frac{1}{|P_{d,m}^1|} \sum_{p \in \mathbb{P}_m} \sum_{h=0}^{p^2 - 1} \exp\left(2\pi i \boldsymbol{k} \cdot \boldsymbol{x}_h^{(p)}\right) \right| \\ &\leq \sum_{\boldsymbol{k} \in \mathbb{Z}^d \setminus \{\mathbf{0}\}} |\hat{f}(\boldsymbol{k})| \left| \frac{1}{|P_{d,m}^1|} \sum_{p \in \mathbb{P}_m} \sum_{h=0}^{p^2 - 1} \exp\left(2\pi i \boldsymbol{k} \cdot \boldsymbol{x}_h^{(p)}\right) \right| \\ &\leq \frac{1}{m} \sum_{\boldsymbol{k} \in \mathbb{Z}^d \setminus \{\mathbf{0}\}} |\hat{f}(\boldsymbol{k})| \left(4 \operatorname{width}(\operatorname{supp}(\boldsymbol{k})) + \frac{8}{c_{\mathbb{P}}} \min_{j \in \operatorname{supp}(\boldsymbol{k})} \log |k_j|\right) \\ &\leq \frac{16}{c_{\mathbb{P}}m} \sum_{\boldsymbol{k} \in \mathbb{Z}^d \setminus \{\mathbf{0}\}} |\hat{f}(\boldsymbol{k})| \max\left(\operatorname{width}(\operatorname{supp}(\boldsymbol{k})), \min_{j \in \operatorname{supp}(\boldsymbol{k})} \log |k_j|\right) \leq \frac{16}{c_{\mathbb{P}}m} \|f\|. \end{aligned}$$

This leads to an upper bound on the worst-case error as

$$e^{\operatorname{wor}}(F_d^2, Q_{d,n}) \le \frac{16}{c_{\mathbb{P}}m}.$$

Therefore, in order to make  $e^{\text{wor}}(F_d^2, Q_{d,n})$  less than or equal to  $\varepsilon \in (0, 1)$ , it suffices to choose  $m = \lceil 16c_{\mathbb{P}}^{-1}\varepsilon^{-1} \rceil$  and we have

$$n(\varepsilon, d, F_d^2) \le \sum_{p \in \mathbb{P}_{\lceil 16c_{\mathbb{P}}^{-1}\varepsilon^{-1}\rceil}} p^2 \le C_{\mathbb{P}} \frac{\lceil 16c_{\mathbb{P}}^{-1}\varepsilon^{-1}\rceil}{\log\lceil 16c_{\mathbb{P}}^{-1}\varepsilon^{-1}\rceil} \times \left(\lceil 16c_{\mathbb{P}}^{-1}\varepsilon^{-1}\rceil\right)^2,$$

from which the result follows immediately.

## 3 Proof of Theorem 2

Proof of Theorem 2. We adopt a similar approach as in the proofs of [14, Theorem 1] and [4, Theorem 1]. Consider an arbitrary linear algorithm  $Q_{d,n}(f) = \sum_{h=0}^{n-1} w_h f(\boldsymbol{x}_h)$ . For a set  $\mathcal{A} \subset \mathbb{Z}^d$  with enough cardinality  $|\mathcal{A}| > n$ , we define a function  $g : [0, 1)^d \to \mathbb{C}$  by

$$g(\boldsymbol{x}) = \sum_{\boldsymbol{k} \in \mathcal{A}} c_{\boldsymbol{k}} \exp(2\pi i \boldsymbol{k} \cdot \boldsymbol{x})$$

with  $c_{\mathbf{k}} \in \mathbb{C}$ , which satisfies  $g(\mathbf{x}_h) = 0$  for all  $h = 0, \ldots, n-1$ . In fact, there exists a non-zero vector of  $(c_{\mathbf{k}})_{\mathbf{k}\in\mathcal{A}}$ , as the condition that  $g(\mathbf{x}_h) = 0$  for all  $h = 0, \ldots, n-1$  forms n homogeneous linear equations with  $|\mathcal{A}| > n$  unknowns  $c_{\mathbf{k}}$ . Let us normalize these coefficients in such a way that

$$\max_{\boldsymbol{k}\in\mathcal{A}}|c_{\boldsymbol{k}}|=c_{\boldsymbol{\ell}}=1\quad\text{for some }\boldsymbol{\ell}\in\mathcal{A}.$$

With this  $\ell$  and a positive constant C, we define another function  $\tilde{g}: [0,1)^d \to \mathbb{C}$  as follows:

$$\tilde{g}(\boldsymbol{x}) = C \exp(-2\pi i \boldsymbol{\ell} \cdot \boldsymbol{x}) g(\boldsymbol{x}) = C \sum_{\boldsymbol{k} \in \mathcal{A}} c_{\boldsymbol{k}} \exp(2\pi i (\boldsymbol{k} - \boldsymbol{\ell}) \cdot \boldsymbol{x}).$$

Then we construct a real-valued function  $g^*$  defined on  $[0,1)^d$  by taking the average of  $\tilde{g}$  and its complex conjugate:  $g^*(\boldsymbol{x}) = (\tilde{g}(\boldsymbol{x}) + \overline{\tilde{g}(\boldsymbol{x})})/2$ . Regarding the norm of  $g^*$  in  $F_d^1$ , we have

$$\|g^{\star}\| \leq \frac{\|\tilde{g}\| + \|\tilde{\overline{g}}\|}{2} = \|\tilde{g}\| = C \sum_{\boldsymbol{k} \in \mathcal{A}} |c_{\boldsymbol{k}}| \max\left(1, \min_{j \in \operatorname{supp}(\boldsymbol{k})} \log |k_j - \ell_j|\right)$$
$$\leq C \sum_{\boldsymbol{k} \in \mathcal{A}} \max\left(1, \min_{j \in \operatorname{supp}(\boldsymbol{k})} \log |k_j - \ell_j|\right) \leq C \max_{\boldsymbol{\ell} \in \mathcal{A}} \sum_{\boldsymbol{k} \in \mathcal{A}} \max\left(1, \min_{j \in \operatorname{supp}(\boldsymbol{k})} \log |k_j - \ell_j|\right).$$

To ensure  $||g^{\star}|| \leq 1$ , we set

$$C = \left(\max_{\boldsymbol{\ell} \in \mathcal{A}} \sum_{\boldsymbol{k} \in \mathcal{A}} \max\left(1, \min_{j \in \operatorname{supp}(\boldsymbol{k})} \log |k_j - \ell_j|\right)\right)^{-1}.$$

By construction, we have  $g^*(\boldsymbol{x}_h) = 0$  for all  $h = 0, \ldots, n-1$ , which implies  $Q_{n,d}(g^*) = 0$ . On the other hand, the exact integral is given by

$$I_d(g^*) = Cc_{\ell} = C = \left( \max_{\ell \in \mathcal{A}} \sum_{k \in \mathcal{A}} \max\left( 1, \min_{j \in \text{supp}(k)} \log |k_j - \ell_j| \right) \right)^{-1}.$$

Since  $g^{\star} \in F_d^1$  with  $\|g^{\star}\| \leq 1$ , the worst-case error of any linear algorithm  $Q_{d,n}$  is bounded below by

$$e^{\operatorname{wor}}(F_d^1, Q_{d,n}) \ge |I_d(g^*) - Q_{n,d}(g^*)| = \left(\max_{\ell \in \mathcal{A}} \sum_{\boldsymbol{k} \in \mathcal{A}} \max\left(1, \min_{j \in \operatorname{supp}(\boldsymbol{k})} \log |k_j - \ell_j|\right)\right)^{-1}$$

In what follows, let

 $\mathcal{A} = \{\mathbf{0}\} \cup \left\{ \mathbf{k} \in \mathbb{Z}^d \mid d-1 \text{ of } k_j \text{ are all } 0 \text{ and one non-zero } k_j \text{ is from } \{1, \dots, \lceil n/d \rceil \} \right\}.$ 

It is easy to verify that  $|\mathcal{A}| = 1 + d\lceil n/d\rceil > n$ . For this choice of  $\mathcal{A}$ , we can restrict ourselves to  $\ell = (\ell, 0, \dots, 0)$  for some  $\ell \in \{0, \dots, \lceil n/d\rceil\}$ . By utilizing the assumption n > 2d and the well-known inequality  $\log x \leq x - 1$ , we have

$$\sum_{\boldsymbol{k}\in\mathcal{A}} \max\left(1,\min_{j\in\operatorname{supp}(\boldsymbol{k})}\log|k_{j}-\ell_{j}|\right)$$
  
=  $\max\left(1,\log\ell\right) + \sum_{k_{1}=1}^{\lceil n/d\rceil} \max\left(1,\log|k_{1}-\ell|\right) + \sum_{j=2}^{d}\sum_{k_{j}=1}^{\lceil n/d\rceil} \max\left(1,\log k_{j}\right)$   
 $\leq \log\lceil n/d\rceil + d\sum_{k=1}^{\lceil n/d\rceil} \max\left(1,\log k\right)$   
 $\leq \log\lceil n/d\rceil + d\lceil n/d\rceil \log\lceil n/d\rceil$   
 $\leq (\lceil n/d\rceil - 1) \cdot (1 + d\lceil n/d\rceil) \leq \frac{2n^{2}}{d}.$ 

Since the last bound is independent of  $\ell$ , we obtain

$$e^{\operatorname{wor}}(F_d^1, Q_{d,n}) \ge \left(\max_{\ell \in \mathcal{A}} \sum_{\boldsymbol{k} \in \mathcal{A}} \max\left(1, \min_{j \in \operatorname{supp}(\boldsymbol{k})} \log |k_j - \ell_j|\right)\right)^{-1} \ge \frac{d}{2n^2}.$$

This completes the proof.

•

### References

- [1] J. Dick. Numerical integration of Hölder continuous, absolutely convergent Fourier, Fourier cosine, and Walsh series. *Journal of Approximation Theory*, 183:14–30, 2014.
- [2] J. Dick, D. Gomez-Perez, F. Pillichshammer, and A. Winterhof. Digital inversive vectors can achieve polynomial tractability for the weighted star discrepancy and for multivariate integration. *Proceedings of the American Mathematical Society*, 145:3297–3310, 2017.
- [3] J. Dick, A. Hinrichs, and F. Pillichshammer. Proof techniques in quasi-Monte Carlo theory. Journal of Complexity, 31(3):327–371, 2015.
- [4] J. Dick, G. Larcher, F. Pillichshammer, and H. Woźniakowski. Exponential convergence and tractability of multivariate integration for Korobov spaces. *Mathematics of computation*, 80(274):905–930, 2011.
- [5] J. Dick and F. Pillichshammer. The weighted star discrepancy of Korobov's p-sets. Proceedings of the American Mathematical Society, 143:5043–5057, 2015.
- [6] T. Goda. Polynomial tractability for integration in an unweighted function space with absolutely convergent Fourier series. *Proceedings of the American Mathematical Society*, 2023.
- [7] L. K. Hua and Y. Wang. Applications of Number Theory to Numerical Analysis. Springer-Verlag, Berlin, 1981.
- [8] N. M. Korobov. Number-Theoretic Methods in Approximate Analysis. Fizmatgiz, Moscow, 1963.
- D. Krieg. Tractability of sampling recovery on unweighted function classes. arXiv preprint arXiv:2304.14169, 2023.
- [10] D. Krieg and J. Vybiral. New lower bounds for the integration of periodic functions. arXiv preprint arXiv:2302.02639, 2023.
- [11] E. Novak and H. Woźniakowski. Tractability of Multivariate Problems, Volume I: Linear Information. EMS Press, Zürich, 2008.
- [12] E. Novak and H. Woźniakowski. Tractability of Multivariate Problems, Volume II: Standard Information for Functionals. EMS Press, Zürich, 2010.
- [13] J. B. Rosser and L. Schoenfeld. Approximate formulas for some functions of prime numbers. *Illinois Journal of Mathematics*, 6:64–94, 1962.
- [14] I. H. Sloan and H. Woźniakowski. An intractability result for multiple integration. *Mathematics of Computation*, 66:1119–1124, 1997.
- [15] I. H. Sloan and H. Woźniakowski. When are quasi-Monte Carlo algorithms efficient for high dimensional integrals? *Journal of Complexity*, 14(1):1–33, 1998.
- [16] J. F. Traub, G. W. Wasilkowski, and H. Woźniakowski. Information-Based Complexity. Academic Press, New York, 1988.