

Overlap Splines and Meshless Finite Difference Methods

Oleg Davydov

Abstract We consider overlap splines that are defined by connecting the patches of piecewise functions via common values at given finite sets of nodes, without using any partitions of the computational domain. It is shown that some classical finite difference methods may be interpreted as collocation with overlap splines. Moreover, several versions of the meshless finite difference methods, such as the RBF-FD method, are equivalent to the collocation or discrete least squares with appropriately chosen spaces of overlap splines.

1 Introduction

In this paper we consider spaces of piecewise defined multivariate functions (for example piecewise polynomials) without choosing a partition of the domain. Therefore the pieces overlap each other, and the "overlap spline" is a set of patches rather than a well defined function. We connect the patches by requiring that they coincide at a finite number of points. In particular, it is not even expected that any underlying partition exists such that the patches form a continuous function. The absence of a partition eliminates the need for meshing algorithms in the numerical implementation of the overlap splines. Despite the simplicity of this setting, basic questions such as determining the dimension of the spaces of overlap splines do not seem trivial in general. In the univariate case overlap splines are related to the well known discrete splines [18].

In this paper we mainly concentrate on the simplest case of "interpolatory" overlap splines, and their applications to solving operator equations by collocation and discrete least squares methods. Remarkably, it turns out that classical finite difference methods can be interpreted as collocation with overlap splines. However,

Oleg Davydov
University of Giessen, Department of Mathematics, Arndtstrasse 2, 35392 Giessen, Germany, e-mail: oleg.davydov@math.uni-giessen.de

even more interesting is the connection to modern meshless finite difference methods, in particular RBF-FD methods [12], that have grown in popularity in recent years due to their true meshless and isogeometric nature and excellent performance in numerous numerical experiments. This is where the absence of partitions plays a decisive role. We hope that recasting meshless finite difference methods in terms of overlap splines will contribute to developing theoretical justification of these methods. Note that our recent work [8] in fact makes use of an overlap spline construction in its proof of the error bound for a discrete least squares version of RBF-FD. On the other hand, we hope that the overlap spline perspective will help to develop new, improved versions of the meshless finite difference methods.

The paper is organized as follows. In Section 2 we introduce overlap splines, address the dimension question and provide several examples, including discrete splines, multivariate polynomial and kernel based patches. Section 3 briefly discusses the application of overlap splines to the direct approximation of functions, whereas Section 4 is devoted to the solution of operator equations and demonstrates how collocation and discrete least squares with overlap splines lead to several known versions of the meshless finite difference method, including the RBF-FD method and its oversampled variants.

2 Overlap Splines

For a finite set M let $|M|$ denote its cardinality. For a vector $a \in \mathbb{R}^N$ and any set $J \subset \{1, \dots, N\}$ we denote by $a|_J$ the vector $[a_j]_{j \in J}$.

Let G be a set and $X = \{x_j\}_{j=1}^N$ its finite subset. Consider a collection $\mathcal{G} = \{G_i\}_{i=1}^m$ of subsets of G such that $G = \bigcup_{i=1}^m G_i$, and a collection $\mathcal{P} = \{P_i\}_{i=1}^m$ of finite dimensional vector spaces P_i of real-valued functions defined on G_i , $i = 1, \dots, m$. We set $X_i = \{x_j : j \in J_i\} = X \cap G_i$ and $n_i = |X_i| = |J_i|$.

Note that in practice we may prefer to first choose the sets X_i and only thereafter find suitable enclosing sets G_i , or even just show the existence of G_i with certain desired properties. For example, G_i may be only needed in the proofs, whereas the generation of X_i be part of a computational algorithm. A possible approach to the latter when G is a metric space is to choose centers $c_1, \dots, c_m \in G$, and generate X_i by selecting for each $i = 1, \dots, m$ a set of nearest neighbors of c_i in X . This can be done for example by collecting either all neighbors within certain distance from c_i (range search), or a prescribed number of its nearest neighbors (KNN search).

Definition 1 (Overlap splines). Given X , \mathcal{G} and \mathcal{P} , any collection $s = \{p_i\}_{i=1}^m$ of patches $p_i \in P_i$, $i = 1, \dots, m$, is said to be an *overlap spline* if the *connection condition* $p_i|_{X_i \cap X_j} = p_j|_{X_i \cap X_j}$ holds whenever $X_i \cap X_j \neq \emptyset$. The linear space of all overlap splines is denoted $S(X, \mathcal{G}, \mathcal{P})$.

Clearly, $\dim S(X, \mathcal{G}, \mathcal{P}) \leq \sum_{i=1}^m \dim P_i$, and the equality holds if and only if $P_i|_{X_i \cap X_j} = \{0\}$ for all pairs (i, j) such that $i \neq j$ and $X_i \cap X_j \neq \emptyset$. Moreover, it is easy to see that

$$\dim S(X, \mathcal{G}, \mathcal{P}) \geq |X| + \sum_{i=1}^m \dim P_i - \sum_{i=1}^m |X_i|. \quad (1)$$

Indeed, we may represent each $s = \{p_i\}_{i=1}^m \in S(X, \mathcal{G}, \mathcal{P})$ as a point \mathbb{R}^D , where $D = \sum_{i=1}^m \dim P_i$, by expanding each p_i in a basis of P_i , and this defines a linear mapping $S(X, \mathcal{G}, \mathcal{P}) \rightarrow \mathbb{R}^D$. Each point $x_k \in X$ generates at most $m_k - 1$ linearly independent linear equations for the images of s in \mathbb{R}^D resulting from the conditions $p_i(x_k) = p_j(x_k)$ for all $i \neq j$ such that $x_k \in X_i \cap X_j$, where $m_k := |\{i : x_k \in X_i\}|$. These linear equations define the linear subspace of \mathbb{R}^D formed by the images of s , and it follows that the dimension of this subspace is at least $D - \sum_{k=1}^N (m_k - 1)$. Since $\sum_{k=1}^N m_k = \sum_{i=1}^m |X_i|$, the lower bound (1) follows.

We can say more about the dimension of $S(X, \mathcal{G}, \mathcal{P})$ under additional assumptions. A set $Y \subset G$ is said to be *total* for a finite-dimensional vector space P of functions on G if for any $p \in P$ the condition $p|_Y = 0$ implies $p = 0$. This means that $p \in P$ is completely determined by its values at Y . If in addition $|Y| = \dim P$, then Y is an *interpolation set (I-set)* for P , that is the interpolation problem that prescribes arbitrary values of p at the points in Y is uniquely solvable. In other words, Y is an I-set for P if and only if $|Y| = \dim P|_Y = \dim P$.

Definition 2 (Interpolatory overlap splines). A space $S(X, \mathcal{G}, \mathcal{P})$ of overlap splines is called *interpolatory* if each X_i is an I-set for P_i , $i = 1, \dots, m$.

Proposition 1. *Assume that X_i is total for P_i , for all $i = 1, \dots, m$. Then $\dim S(X, \mathcal{G}, \mathcal{P}) \leq |X|$. For interpolatory overlap splines $\dim S(X, \mathcal{G}, \mathcal{P}) = |X|$.*

Proof. Given $s \in S(X, \mathcal{G}, \mathcal{P})$ we define $\hat{u} \in \mathbb{R}^N$ by setting $\hat{u}_j = p_i(x_j)$ for any i such that $x_j \in X_i$. It follows from Definition 1 that $p_i(x_j)$ does not depend on a particular choice of i . The mapping $T : S(X, \mathcal{G}, \mathcal{P}) \rightarrow \mathbb{R}^N$ given by $s \mapsto \hat{u}$ is linear. If $Ts = 0$, then in particular $p_i|_{X_i} = \hat{u}|_{J_i} = 0$ and hence $p_i = 0$ for all i , which shows that $\ker T = \{0\}$. Hence T is an injection, and $\dim S(X, \mathcal{G}, \mathcal{P}) \leq |X|$. This in particular applies to the interpolatory overlap splines. Moreover, in this case $|X_i| = \dim P_i$, $i = 1, \dots, m$, and hence (1) gives the opposite inequality $\dim S(X, \mathcal{G}, \mathcal{P}) \geq |X|$.

The mapping $T : S(X, \mathcal{G}, \mathcal{P}) \rightarrow \mathbb{R}^N$ used in the above proof is well defined for any overlap splines. We will use the notation $s|_X := Ts$, and call $s|_X$ the *restriction* of s to X . Thus, any $s \in S(X, \mathcal{G}, \mathcal{P})$ gives rise to a discretized function $s|_X$ defined only on X . Under the hypotheses of Proposition 1 we have $\ker T = \{0\}$, and hence each $s \in S(X, \mathcal{G}, \mathcal{P})$ is uniquely determined by $s|_X$.

We may always split the task of computing the dimension of $S(X, \mathcal{G}, \mathcal{P})$ into the dimensions of the kernel and the image of T ,

$$\dim S(X, \mathcal{G}, \mathcal{P}) = \dim \ker T + \dim \operatorname{im} T. \quad (2)$$

For the spaces of interpolatory overlap splines the mapping T is a bijection. Indeed, we have shown in the proof of Proposition 1 that $\ker T = \{0\}$ in this case. To show the surjectivity of T , consider any $\hat{u} \in \mathbb{R}^N$, and for each $i = 1, \dots, m$ choose p_i

to be the unique element of P_i such that $p_i(x_j) = \hat{u}_j$ for all $j \in J_i$. Then $s = \{p_i\}_{i=1}^m$ satisfies $Ts = \hat{u}$. This means that the elements $s \in S(X, \mathcal{G}, \mathcal{P})$ can be parameterized by their ‘values’ $s(x_j)$, $j = 1, \dots, N$, uniquely defined by evaluating at x_j any one of the suitable patches p_i such that $x_j \in X_i$.

Any space $S(X, \mathcal{G}, \mathcal{P})$ of interpolatory overlap splines possesses a *Lagrange basis* ℓ_1, \dots, ℓ_N , with $\ell_j = \{\ell_{ji}\}_{i=1}^m$, where each $\ell_{ji} \in P_i$ is uniquely determined by the interpolation conditions $\ell_{ji}(x_k) = \delta_{jk}$, $k \in X_i$, and δ_{jk} is the Kronecker delta. These basis splines are *local* in the sense that $\ell_{ji} = 0$ as soon as $x_j \notin X_i$.

We now discuss several examples of overlap splines. In all special cases considered in this paper G is a closed set in a smooth manifold, mostly $G \subset \mathbb{R}^d$. However we prefer not to restrict the definitions to this situation, because less usual sets G may quite naturally be of interest, e.g. graphs.

Univariate Generalized Splines and Discrete Splines

Let G be an interval $[a, b] \subset \mathbb{R}$. Consider a partition $\Delta = \{a = t_0 < t_1 < \dots < t_m = b\}$ of $[a, b]$, and choose sets $G_i \supset [t_{i-1}, t_i]$, and spaces P_i , $i = 1, \dots, m$. Given a set $X = \{x_j\}_{j=1}^N \subset [a, b]$, any overlap spline $s = \{p_i\}_{i=1}^m \in S(X, \mathcal{G}, \mathcal{P})$ may be used to construct a *generalized spline* $\tilde{s} : [a, b] \rightarrow \mathbb{R}$ in the sense of [18, Chapter 11], where $\tilde{s}(x) := p_i(x)$, with i satisfying $x \in [t_{i-1}, t_i]$ and $x < t_i$ when $i < m$. The linear functionals of the set Γ_{ij} of [18, Definition 11.1] are the point evaluation functionals for all points in $X_i \cap X_j$.

The spline \tilde{s} is a continuous function on $[a, b]$ if each p_i is continuous on $[t_{i-1}, t_i]$ and $t_i \in X$ for all $i = 1, \dots, m-1$. If $X_i \subset [t_{i-1}, t_i]$, $i = 1, \dots, m$, then the generalized spline \tilde{s} may be directly recognized as an overlap spline in $S(X, \tilde{\mathcal{G}}, \mathcal{P})$, with $\tilde{\mathcal{G}} = \{[t_{i-1}, t_i] : i = 1, \dots, m\}$. In particular, we obtain the classical spaces of continuous piecewise polynomials (finite elements) of degree q if X_i is a set of $q+1$ points in $[t_{i-1}, t_i]$, including the endpoints t_{i-1}, t_i , and P_i is the linear space Π_q^1 of univariate polynomials of degree at most q .

However, an overlap spline does not require a partition. According to Definition 1 we only need to choose points $x_j \in [a, b]$, sets $G_i \subset [a, b]$ and spaces P_i . As an example, we take

$$\begin{aligned} x_j &= a + hj, \quad j = 0, \dots, N, \text{ with } h = (b-a)/N, \\ G_i &= [x_{i-1}, x_{i+1}], \quad i = 1, \dots, N-1, \text{ and} \\ P_i &= \Pi_2^1, \text{ the space of quadratic polynomials.} \end{aligned} \tag{3}$$

Since $X_i = X \cap G_i = \{x_{i-1}, x_i, x_{i+1}\}$, $i = 1, \dots, N-1$, are I-sets for Π_2^1 , the space $S(X, \mathcal{G}, \mathcal{P})$ is interpolatory. The Lagrange basis ℓ_0, \dots, ℓ_N for it, with $\ell_j = \{\ell_{ji}\}_{i=1}^{N-1}$, is given by Lagrange polynomials

$$\ell_{ji}(x) = \frac{1}{h^2} \begin{cases} \frac{1}{2}(x - x_{j-2})(x - x_{j-1}), & i = j - 1, \\ -(x - x_{j-1})(x - x_{j+1}), & i = j, \\ \frac{1}{2}(x - x_{j+1})(x - x_{j+2}), & i = j + 1, \\ 0, & \text{otherwise.} \end{cases} \quad (4)$$

If we now choose the following partition, in order to remove the overlaps, $\Delta = \{a = t_0 < t_1 < \dots < t_{N-1} = b - h\}$ with simply $t_j = x_j$ for all j , and define, for each overlap spline $s = \{p_i\}_{i=1}^{N-1} \in S(X, \mathcal{G}, \mathcal{P})$ the continuous function \tilde{s} on $[a, b - h]$ given by

$$\tilde{s}|_{[t_{i-1}, t_i]} = p_i, \quad i = 1, \dots, N - 1,$$

then we arrive at a special case of *discrete splines*, as defined for example in [18, Section 8.5]. Note that the discrepancy that the discrete splines are defined on the interval $[a, b - h]$ instead of $G = [a, b]$ is technical and is related to the fact that discrete splines use forward differences in the definition of the connection conditions, which causes some asymmetry.

Clearly, various results known for discrete splines, such as dimension formulas, local bases (discrete B-splines) or quasi-interpolation methods, see [18] and references therein, can be carried over to appropriate more general types of univariate overlap splines.

Multivariate Polynomial Overlap Splines

Let $G \subset \mathbb{R}^d$, $d \geq 2$, $X = \{x_j\}_{j=1}^N \subset G$ and, for some choice of $\mathcal{G} = \{G_i\}_{i=1}^m$, $G_i \subset G$, let $\mathcal{P}_q := \{\Pi_q^d|_{G_i}\}_{i=1}^m$ for some $q \geq 1$, where Π_q^d denotes the space of d -variate polynomials of total degree less or equal q . Then $S(X, \mathcal{G}, \mathcal{P}_q)$ is a space of multivariate polynomial overlap splines. To generate interpolatory overlap splines, X_i must satisfy $|X_i| = \dim \Pi_q^d|_{X_i} = \dim \Pi_q^d|_{G_i}$.

Given a finite set $X \subset \mathbb{R}^d$, suppose that $\Delta = \{T_1, \dots, T_m\}$ is a *triangulation* of X , where each $T_i \in \Delta$ is a (closed) non-degenerate simplex with vertices in X , and $T_i \cap T_j$ is a common face of T_i, T_j as soon as $i \neq j$. One way to define overlap splines is by choosing $G = \cup_{i=1}^m T_m$ and $\mathcal{G} = \{T_i : i = 1, \dots, m\}$. Then each $X_i = X \cap T_i$ consists of $d + 1 = \dim \Pi_1^d = \dim \Pi_1^d|_{T_i}$ vertices of T_i , and we obtain interpolatory overlap splines $s = \{p_i\}_{i=1}^m \in S(X, \mathcal{G}, \mathcal{P}_1)$ that may be realized as continuous piecewise linear splines \tilde{s} by setting $\tilde{s}|_{T_i} = p_i$, $i = 1, \dots, m$. Thus, in this way we obtain the classical linear Courant elements. Instead of setting $G_i = T_i$, we may choose G_i to be some simply connected sets enclosing T_i , such that $G_i \cap X = X_i$ still holds, leading to essentially the same space of overlap splines, without making use of a partition of G .

Starting with the same X , but choosing different subsets G_i , we obtain completely different spaces of overlap splines. As an illustration, suppose in the bivariate case that a triangulation $\Delta' = \{T'_1, \dots, T'_m\}$ is chosen such that the set of middle points of edges of all triangles of Δ' coincides with X . Then it is easy to see that by choosing G'_i as the triangles T'_i or some sets $G'_i \subset \mathbb{R}^2$ enclosing T'_i , such that $G'_i \cap X = T'_i \cap X$,

$i = 1, \dots, m$, we obtain a space of overlap splines $S(X, \mathcal{G}', \mathcal{P}_1)$ that can be interpreted as the discontinuous Crouzeix-Raviart finite elements, see e.g. [3].

If we start with a triangulation $\Delta = \{T_1, \dots, T_m\}$, and define X as its set of *domain points* of degree q , that is the union of the domain points for all simplices T_i (see the definitions in [14] for $d = 2, 3$, and for example in [1] in general), then by choosing $X_i = X \cap T_i$ and some enclosing sets $G_i \supset T_i$, with $G_i \cap X = X_i$, we get a space of interpolatory overlap splines that may be identified with continuous piecewise polynomials of degree q on the triangulation Δ in the same way as described before for $q = 1$.

Thus, we obtained the classical spline spaces $S_q^0(\Delta)$ on triangulations, where

$$S_q^r(\Delta) := \{f \in C^r(\cup_{i=1}^m T_i) : f|_{T_i} \in \Pi_q^d\}.$$

In order to interpret $S_q^r(\Delta)$ with $r \geq 1$ as overlap splines we would need to generalize the patch connection conditions $p_i|_{X_i \cap X_j} = p_j|_{X_i \cap X_j}$ of Definition 1 by either requiring that the partial derivatives of p_i and p_j up to order r coincide at $X_i \cap X_j$, or that the differences $p_i - p_j$ belong to appropriate polynomial ideals.

In general no triangulation is needed; in order to define overlap splines we just choose $X \subset G$ and overlapping sets G_i . Let us assume that G is the closure of a domain $\Omega \subset \mathbb{R}^d$ (that is, Ω an open and connected set), choose a collection of domains $\Omega_i \subset \mathbb{R}^d$, $i = 1, \dots, m$, that comprise an open cover of G , and set $G_i = \Omega_i \cap G$. For any $X = \{x_j\}_{j=1}^N \subset G$ and any $q \in \mathbb{N}$ we obtain a space of overlap splines $S(X, \mathcal{G}, \mathcal{P}_q)$. We may also prefer to place some further restrictions on the sets X_i , for example requiring that Ω_i are simply connected or convex.

Similar to the case of classical spline spaces (see [14, Section 9.8]), we may speak of a generic position of the set X .

Definition 3 (Generic position of X). We say that $X = \{x_j\}_{j=1}^N$ is in a *generic position* with respect to \mathcal{G} and q if

$$\dim S(X, \mathcal{G}, \mathcal{P}_q) \leq \dim S(\tilde{X}, \mathcal{G}, \mathcal{P}_q)$$

for all sets $\tilde{X} \subset G$ obtained by a sufficiently small perturbation of the coordinates of the points x_j , $j = 1, \dots, N$.

The following statement gives a justification for this definition, as it shows that small perturbations of X do not change the dimension of $S(X, \mathcal{G}, \mathcal{P}_q)$ if X is in a generic position.

Proposition 2. *Given any X, \mathcal{G}, q , we have $\dim S(X, \mathcal{G}, \mathcal{P}_q) \geq \dim S(\tilde{X}, \mathcal{G}, \mathcal{P}_q)$ for all sets $\tilde{X} = \{\tilde{x}_j\}_{j=1}^N \subset G$ obtained by a sufficiently small perturbation of the coordinates of the points x_j , $j = 1, \dots, N$.*

Proof. Assume that the perturbation is so small that $\tilde{J}_i = J_i$, $i = 1, \dots, m$, where J_i, \tilde{J}_i are index sets such that $X_i := X \cap G_i = \{x_j : j \in J_i\}$ and $\tilde{X}_i := \tilde{X} \cap G_i = \{\tilde{x}_j : j \in \tilde{J}_i\}$. Then any $s = \{p_i\}_{i=1}^m \in S(\tilde{X}, \mathcal{G}, \mathcal{P}_q)$ may be uniquely represented by a sequence c of $m \binom{q+d}{d}$ coefficients of m polynomials $p_i \in \Pi_q^d$ satisfying the equation

$A(\tilde{X})c = 0$, where $A(\tilde{X})$ is a matrix whose rows are generated from all connection conditions $p_i|_{\tilde{x}_i \cap \tilde{x}_j} = p_j|_{\tilde{x}_i \cap \tilde{x}_j}$ expressed in terms of the coefficients in c . Hence

$$\dim S(\tilde{X}, \mathcal{G}, \mathcal{P}_q) = \dim \ker A(\tilde{X}) = m \binom{q+d}{d} - \text{rank } A(\tilde{X}).$$

Arguing as in the proof of [14, Theorem 9.32], we choose a nonsingular square submatrix B of $A(X)$ of size $r = \text{rank } A(X)$, and consider the submatrix $B(\tilde{X})$ of $A(\tilde{X})$ corresponding to the same rows and columns. For a sufficiently small perturbation $\det B(\tilde{X}) \neq 0$ since $\det B(\tilde{X})$ is a continuous function of \tilde{X} and $\det B(X) = \det B \neq 0$. Hence $\text{rank } A(\tilde{X}) \geq r$, and the claim follows.

Open Problem (Dimension of the spaces of polynomial overlap splines) In the above setting, determine the dimension of $S(X, \mathcal{G}, \mathcal{P}_q)$, in particular for the case when X is in a generic position. Characterize overlap spline spaces $S(X, \mathcal{G}, \mathcal{P}_q)$ for which the equality holds in (1), that is

$$\dim S(X, \mathcal{G}, \mathcal{P}_q) = |X| + m \binom{q+d}{d} - \sum_{i=1}^m |X_i|. \quad (5)$$

Note that (5) holds in particular for interpolatory overlap splines. Clearly, $S(X, \mathcal{G}, \mathcal{P}_q)$ is interpolatory, with $\dim S(X, \mathcal{G}, \mathcal{P}_q) = |X|$, if and only if each X_i is an I-set for Π_q^d , as in the above examples on triangulations. This means that $|X_i| = \dim \Pi_q^d = \binom{q+d}{d}$ and X_i does not lie on any algebraic hypersurface of degree q .

If we only assume that $|X_i| = \dim \Pi_q^d = \binom{q+d}{d}$ for all $i = 1, \dots, m$, then all X_i may be turned into I-sets for Π_q^d by a single arbitrarily small perturbation of $X = \{x_j\}_{j=1}^N$. Indeed, given such an X , consider the set \mathfrak{X} consisting of all N -tuples $(\tilde{x}_1, \dots, \tilde{x}_N) \in G^N$ such that

- (a) $\tilde{x}_j \neq \tilde{x}_k$ whenever $j \neq k$, and (b) $\tilde{x}_j \in G_k$ if and only if $j \in J_k$,

with J_k defined as in the proof of Proposition 2. For each $i = 1, \dots, m$ denote by \mathfrak{X}_i the subset of \mathfrak{X} that consists of all $(\tilde{x}_1, \dots, \tilde{x}_N) \in \mathfrak{X}$ such that

$$\{\tilde{x}_j : j \in J_i\} \text{ is an I-set for } \Pi_q^d.$$

Then each \mathfrak{X}_i is an open and dense subset of \mathfrak{X} , and hence $\tilde{\mathfrak{X}} := \bigcap_{i=1}^m \mathfrak{X}_i$ is also an open dense subset of \mathfrak{X} . Since $(x_1, \dots, x_N) \in \mathfrak{X}$, by an arbitrary small perturbation of the coordinates of the points x_j , $j = 1, \dots, N$, we may obtain a set $\tilde{X} = \{\tilde{x}_j\}_{j=1}^N$ such that $(\tilde{x}_1, \dots, \tilde{x}_N) \in \tilde{\mathfrak{X}}$ and hence each $\tilde{X}_i = \tilde{X} \cap G_i$ is an I-set for Π_q^d , and thus $S(\tilde{X}, \mathcal{G}, \mathcal{P}_q)$ is interpolatory with $\dim S(\tilde{X}, \mathcal{G}, \mathcal{P}_q) = |\tilde{X}| = |X|$. This implies the following statement.

Proposition 3. *Assume that $|X_i| = \dim \Pi_q^d$ for all $i = 1, \dots, m$. Then X is in a generic position if and only if (5) holds, that is $\dim S(X, \mathcal{G}, \mathcal{P}_q) = |X|$. In particular, X is in a generic position if $S(X, \mathcal{G}, \mathcal{P}_q)$ is interpolatory.*

Proof. If $\dim S(X, \mathcal{G}, \mathcal{P}_q) = |X|$, then X is in a generic position because the inequality $\dim S(\tilde{X}, \mathcal{G}, \mathcal{P}_q) \geq |\tilde{X}| = |X|$ for sufficiently small perturbations follows from (1). Conversely, if X is in a generic position and $\dim S(X, \mathcal{G}, \mathcal{P}_q) > |X|$, then any sufficiently small perturbation \tilde{X} must satisfy $\dim S(\tilde{X}, \mathcal{G}, \mathcal{P}_q) > |X| = |\tilde{X}|$, which contradicts the above argument that we may always find \tilde{X} with interpolatory $S(\tilde{X}, \mathcal{G}, \mathcal{P}_q)$.

Note that the converse of the last claim of Proposition 3 is not true. For example, let $d = 2$, $q = 1$, $m = 1$, let G be any open set, and let X consist of three collinear points in $G_1 = G$. Then $S(X, \mathcal{G}, \mathcal{P}_1)$ is not interpolatory, but it is in a generic position since $\dim S(X, \mathcal{G}, \mathcal{P}_1) = 3 = |X|$.

It follows from [18, Theorem 8.50] that (5) holds for all spaces of univariate discrete splines of [18, Section 8.5] interpreted as overlap splines in the same way as done in the example defined by (3).

However, it is easy to generate an example where (5) fails. Let $d = 2$, $q = 1$, $m = 2$, let Ω_1 and Ω_2 be two connected open sets with nonempty intersection, and $\Omega = \Omega_1 \cup \Omega_2$. We choose X to be a set of N points in $\Omega_1 \cap \Omega_2$. Then $X_1 = X_2 = X$ and the right hand side of (5) is $6 - N$, which is even negative for $N > 6$. It is easy to see that the dimension of $S(X, \mathcal{G}, \mathcal{P}_1)$ is three if the points in X are non-collinear, four if they are collinear and $N \geq 2$, and five if $N = 1$. Thus, (5) is only correct if either $N \leq 2$, or $N = 3$ and the points are non-collinear. Note that X is in a generic position unless $N \geq 3$ and the points are collinear, since otherwise small perturbations of X do not change the dimension of $S(X, \mathcal{G}, \mathcal{P}_1)$.

Kernel-Based Overlap Splines

Recall that a function $K : G \times G \rightarrow R$ is said to be a *positive definite kernel* if the matrix $[K(x_i, x_j)]_{i,j=1}^n$ is positive definite for any finite set $X = \{x_j\}_{j=1}^n \subset G$. Then X is an I-set for the space

$$P_{K,X} := \text{span}\{K(\cdot, x_j) : j = 1, \dots, n\}.$$

Therefore, an interpolatory overlap spline space $S(X, \mathcal{G}, \mathcal{P})$ is obtained for any $X \subset G$ and $\mathcal{G} = \{G_j\}_{j=1}^m$ with usual assumptions if we choose $\mathcal{P} = \mathcal{P}_K := \{P_{K, X \cap G_i}\}_{i=1}^m$.

A kernel K is *conditionally positive definite* with respect to a finite-dimensional space Q of real-valued functions on G if for any finite set $X = \{x_j\}_{j=1}^n \subset G$ the quadratic form $\sum_{i,j=1}^n c_i c_j K(x_i, x_j)$ is positive for all $c \in \mathbb{R}^n \setminus \{0\}$ such that $\sum_{j=1}^n c_j p(x_j) = 0$ for all $p \in Q$. If X is a total set for Q , then X is an I-set for the space

$$P_{K,X,Q} := \left\{ \sum_{j=1}^n c_j K(\cdot, x_j) + \tilde{p} : \tilde{p} \in Q \text{ and } c \in \mathbb{R}^n \text{ with } \sum_{j=1}^n c_j p(x_j) = 0 \forall p \in Q \right\},$$

see for example [23]. Note that $P_{K,X,\{0\}} = P_{K,X}$. For any finite $X \subset G$ and $\mathcal{G} = \{G_j\}_{j=1}^m$ we obtain an interpolatory overlap spline space $S(X, \mathcal{G}, P_{K,Q})$, with $\mathcal{P}_{K,Q} := \{P_{K,X \cap G_i, Q}\}_{i=1}^m$, as soon as each $X_i = X \cap G_i$ is total for Q . If $G \subset \mathbb{R}^d$ and $Q = \Pi_{q-1}^d$ for some $q \geq 1$, then K is said to be *conditionally positive definite of order q* , and we use the notation $P_{K,X,q} := P_{K,X,Q}$ and $\mathcal{P}_{K,q} := \mathcal{P}_{K,Q}$. We also set $P_{K,X,0} := P_{K,X}$ and $\mathcal{P}_{K,0} := \mathcal{P}_K$ to include the positive definite case in the common notation.

We refer to [4, 11, 23] for the theory of (conditionally) positive definite kernels. In particular, for $G \subset \mathbb{R}^d$ several families of kernels are known in the form of a *radial basis function (RBF)* $K(x, y) = \varphi(\|x - y\|_2)$. A typical example of a positive definite RBF is the *Gauss kernel* given by $\varphi(r) = e^{-(\varepsilon r)^2}$, where $\varepsilon > 0$ is the *shape parameter*. The *polyharmonic RBF* given by $\varphi(r) = r^\alpha$ for any $\alpha \in (0, +\infty) \setminus 2\mathbb{N}$ is conditionally positive definite of order $q \geq \lfloor \alpha/2 \rfloor + 1$.

Note that we can also choose different kernels K_i and different spaces Q_i for each $i = 1, \dots, m$. In particular, K_i may be the Gauss kernel with different values of the shape parameter.

3 Approximation of Functions

An overlap spline $s = \{p_i\}_{i=1}^m$, obtained by some computation as an approximation of a function $f : G \rightarrow \mathbb{R}$, may be used directly to estimate function values $f(x)$ or other functionals such as partial derivatives of f at $x \in G$, by evaluating at x suitable patches p_i such that $x \in G_i$. Depending on the application it may be sufficient to just pick one such i and use $p_i(x)$, or we may need to build a (weighted) average of all values $p_i(x)$ with $x \in G_i$, or produce an approximating function $\tilde{s} : G \rightarrow \mathbb{R}$ in a different way. Moreover, the restriction $s|_X$ may already serve as a discrete approximation of f .

In special circumstances, like univariate generalized splines or continuous multivariate polynomial splines on triangulations, the values $p_i(x)$ may be the same for all i with $x \in G_i$, at least for a special choice of G_i , and we obtain a well-defined function \tilde{s} by using these values. However, this is a rare exception if we look for meshless methods, because it typically means that $\mathcal{G} = \{G_i\}_{i=1}^m$ is a partition of G .

We always have at our disposal the *Partition of Unity Method (PUM)* that generates $\tilde{s}(x)$ as a weighted average of $p_i(x)$ as follows. Let $\Gamma = \{\gamma_i\}_{i=1}^m$ be a *partition of unity* associated with \mathcal{G} , such that

$$\gamma_i : G \rightarrow \mathbb{R}, \quad \gamma_i(x) \geq 0, \quad x \in G_i, \quad \gamma_i(x) = 0, \quad x \in G \setminus G_i, \quad i = 1, \dots, m,$$

and

$$\sum_{i=1}^m \gamma_i(x) = 1, \quad x \in G.$$

For each overlap spline $s = \{p_i\}_{i=1}^N \in S(X, \mathcal{G}, \mathcal{P})$ we define a function

$$s_\Gamma(x) = \sum_{\substack{i=1 \\ x \in G_i}}^m \gamma_i(x) p_i(x), \quad x \in G,$$

that can take the role of \tilde{s} . If $S(X, \mathcal{G}, \mathcal{P})$ is interpolatory, then $s_\Gamma|_X = s|_X$.

Note that the patch connections $p_i|_{X_i \cap X_j} = p_j|_{X_i \cap X_j}$ of Definition 1 are not needed in order that s_Γ can serve as an approximation of a given function $f : G \rightarrow \mathbb{R}$. It suffices to find $p_i \in P_i$ that approximate f well on G_i , see e.g. the approximation theory of PUM in [23]. For example, for the approximation of functions from scattered data $f(x_i)$, $i = 1, \dots, N$, with x_i in a smooth manifold G , we may use a kernel-based interpolatory overlap spline $s \in S(X, \mathcal{G}, \mathcal{P})$ and generate s_Γ with desired degree of smoothness by using appropriately smooth partition of unity functions γ_i . However, if we build a *disconnected overlap spline* $s \in S(\emptyset, \mathcal{G}, \mathcal{P})$, with local approximations $p_i \in P_i$ of the data $f|_{X \cap G_i}$ obtained by any suitable method, then s_Γ will also inherit the approximation quality of the patches.

The above *direct approximation* setting, where the input data is given at $x_i \in X$, should be contrasted with the *solution of the operator equations*, which we will consider in the next section. In this case patch connections are crucial, and the discretized solutions $\hat{u} = s|_X$ may in fact be sufficient as the output of the algorithm, without creating a solution \tilde{s} defined on G , or one may leave the latter to a post-processing step with its own algorithms that may in turn rely on PUM or other approaches to scattered data fitting, such as kernel-based and moving least squares methods or spline fitting.

4 Collocation, Least Squares and Finite Difference Methods

We now consider a linear operator equation: Find $u : G \rightarrow \mathbb{R}$, such that

$$Lu(x) = f(x), \quad \forall x \in G, \quad (6)$$

where f is a given function $f : G \rightarrow \mathbb{R}$, and $Lu : G \rightarrow \mathbb{R}$ is well defined for all u in a vector space U of functions defined on G . We assume that the operator L is linear and that (6) has a unique solution $u \in U$ for all $f \in F$, where F is some other vector space of functions on G .

For each $x \in G$, a subset $G_x \subset G$ such that $x \in G_x$ is said to be a *determining set of L at x* if $Lu(x) = Lv(x)$ whenever $u, v \in U$ satisfy $u|_{G_x} = v|_{G_x}$. An overlap spline space $S(X, \mathcal{G}, \mathcal{P})$ is *compatible with L* if $P_i \subset U|_{G_i}$, $i = 1, \dots, m$, and every $G_i \in \mathcal{G}$ is a determining set of L at each $x \in G_i$. Then $Lp(x)$ is well defined for all

$x \in G_i$, $p \in P_i$, and hence $Ls = \{Lp_i\}_{i=1}^m \in S(X, \mathcal{G}, L\mathcal{P})$ is well defined for each $s = \{p_i\}_{i=1}^m \in S(X, \mathcal{G}, \mathcal{P})$, where $L\mathcal{P} := \{LP_i\}_{i=1}^m$, with $LP_i := \{Lp : p \in P_i\}$.

The *Method of Collocation* determines an approximate solution of (6) as an overlap spline s as follows. Choose a space $S(X, \mathcal{G}, \mathcal{P})$ of overlap splines compatible with L , a set of *collocation nodes* $Y = \{y_1, \dots, y_D\} \subset G$, with $D = \dim S(X, \mathcal{G}, \mathcal{P})$, and a mapping $\sigma : \{1, \dots, D\} \rightarrow \{1, \dots, m\}$, such that $y_j \in G_{\sigma(j)}$. Determine $s = \{p_i\}_{i=1}^m \in S(X, \mathcal{G}, \mathcal{P})$ by requiring that

$$Lp_{\sigma(j)}(y_j) = f(y_j), \quad j = 1, \dots, D. \quad (7)$$

If we expand the solution s of (7) in a basis $\{s_1, \dots, s_D\}$ of $S(X, \mathcal{G}, \mathcal{P})$,

$$s = \sum_{k=1}^D c_k s_k, \quad c_k \in \mathbb{R}, \quad (8)$$

where

$$s_k = \{p_{ki}\}_{i=1}^m \in S(X, \mathcal{G}, \mathcal{P}), \quad k = 1, \dots, D,$$

then the unknown coefficients c_k of s satisfy the square system of linear equations

$$\sum_{k=1}^D c_k Lp_{k, \sigma(j)}(y_j) = f(y_j), \quad j = 1, \dots, D. \quad (9)$$

Although in general there is no guarantee that this linear system is non-singular or solvable, we will soon discuss some situations where this has been shown.

The *Method of Discrete Least Squares* differs from the method of collocation in that the number of collocation nodes exceeds the dimension of the space of overlap splines. Moreover, we allow multiple collocation nodes if different patches $p_{\sigma(j)}$ are chosen for the same node y_j . The exact fulfillment of the equations (7) is replaced by the least squares minimization of the residual. More precisely, we choose a sequence $Y = (y_1, \dots, y_M)$ of nodes in G , with $M > D = \dim S(X, \mathcal{G}, \mathcal{P})$, where $y_k = y_j$ is allowed for $k \neq j$, and a mapping $\sigma : \{1, \dots, M\} \rightarrow \{1, \dots, m\}$, such that $y_j \in G_{\sigma(j)}$, and $\sigma(j) \neq \sigma(k)$ whenever $y_j = y_k$ with $j \neq k$. Determine $s = \{p_i\}_{i=1}^m \in S(X, \mathcal{G}, \mathcal{P})$ by solving the minimization problem

$$\min \left\{ \sum_{j=1}^M |Lp_{\sigma(j)}(y_j) - f(y_j)|^2 : \{p_i\}_{i=1}^m \in S(X, \mathcal{G}, \mathcal{P}) \right\}. \quad (10)$$

The algebraic formulation is

$$\min \left\{ \|Ac - b\|_2 : c \in \mathbb{R}^D \right\}, \quad (11)$$

where

$$A = [Lp_{k, \sigma(j)}(y_j)]_{j=1, k=1}^{M, D}, \quad b = [f(y_j)]_{j=1}^M,$$

and $c = [c_k]_{k=1}^D$ is the vector of the coefficients of s in (8). The least squares problem is always solvable. The solution is unique if and only if the system matrix A is of full rank.

As alternative to the formulation (10), also other approaches to the approximate solution of the overdetermined linear system

$$Lp_{\sigma(j)}(y_j) = f(y_j), \quad j = 1, \dots, M, \quad (12)$$

are possible, for example, weighted least squares.

We now discuss several examples emphasizing connections to the Finite Difference Method and its meshless generalizations.

A Boundary Value Problem on an Interval

As a first example we consider the boundary value problem

$$u''(x) = f(x), \quad x \in (a, b), \quad u(a) = u(b) = 0, \quad (13)$$

with unknown $u \in U = C^2[a, b]$. Note that the operator L of (6) takes the form

$$Lu(x) = \begin{cases} u''(x), & x \in (a, b), \\ u(x), & x \in \{a, b\}, \end{cases}$$

and the right hand side is extended to the endpoints of $[a, b]$ by setting $f(a) = f(b) = 0$.

Let us apply the method of collocation using the overlap spline space $S(X, \mathcal{G}, \mathcal{P})$ defined by (3). It is easy to see that this space is compatible with the operator L . The dimension of $S(X, \mathcal{G}, \mathcal{P})$ is $D = N + 1$, and we may look for the overlap spline solution $s = \{p_i\}_{i=1}^{N-1}$ in the form

$$s = \sum_{k=0}^N c_k \ell_k, \quad c_k \in \mathbb{R},$$

using the Lagrange basis (4). As discussed in Section 2, c_k can be interpreted as the values $s(x_k)$ of the overlap spline s because the values of all patches $p_i(x_k)$ such that $x_k \in G_i$ coincide. We choose the collocation nodes $y_j = x_j$, $j = 0, \dots, N$, and corresponding patches $p_{\sigma(j)}$ according to the rule

$$\sigma(j) = j \text{ for } j = 1, \dots, N-1, \quad \sigma(0) = 1, \quad \sigma(N) = N-1.$$

The entries $L\ell_{k, \sigma(j)}(x_j)$ of the system matrix of (9) are easy to compute using the explicit formulas (4), and we arrive at the linear system

$$\begin{aligned} \frac{c_{k-1} - 2c_k + c_{k+1}}{h^2} &= f(x_k), \quad k = 1, \dots, N-1, \\ c_0 &= 0, \quad c_N = 0. \end{aligned} \quad (14)$$

This is in fact the same non-singular linear system that determines the classical finite difference solution of the boundary value problem (13), where c_k are approximations of $u(x_k)$. In particular, the standard theory of finite difference methods applies that provide error bounds for $c_k - u(x_k)$, and thus for $s(x_k) - u(x_k)$, $k = 1, \dots, N-1$.

A second look at the above reveals that the appearance of the second order finite difference

$$\frac{s(x_{k-1}) - 2s(x_k) + s(x_{k+1}))}{h^2},$$

that can be interpreted as the second derivative of the discrete function $s|_X$ at x_k , is not accidental because the finite difference formula

$$p''(x_k) = \frac{p(x_{k-1}) - 2p(x_k) + p(x_{k+1}))}{h^2}, \quad p \in \Pi_2^1,$$

is just a method to compute $p''_k(x_k)$ using $p_k|_{X_k}$ when we set up the collocation system (7) using the Lagrange basis of $S(X, \mathcal{G}, \mathcal{P})$. Therefore, applying the method of collocation with interpolatory overlap splines in terms of their Lagrange basis may always be interpreted as a finite difference method.

Note that collocation with cubic discrete splines has been used in [5] to solve second order boundary value problems.

Dirichlet Problem for the Poisson Equation on the Square

Consider the boundary value problem

$$\Delta u(x) = f(x), \quad x \in \Omega = (0, 1)^2, \quad u|_{\partial\Omega} = 0, \quad (15)$$

where $u \in U = C^2([0, 1]^2)$. We define overlap splines by choosing $G = [0, 1]^2$,

$$\begin{aligned} X &= \{x_{ij}\}_{i,j=0}^N, \quad \text{with } x_{ij} = (ih, jh) \in [0, 1]^2, \quad h = 1/N, \\ \mathcal{G} &= \{G_{ij}\}_{i,j=1}^{N-1}, \quad \text{with } G_{ij} = \{x \in [0, 1]^2 : \|x - x_{ij}\|_2 < h + \varepsilon\}, \end{aligned} \quad (16)$$

where $0 < \varepsilon < (\sqrt{2} - 1)h$. Then $X_{ij} = G_{ij} \cap X = \{x_{ij}, x_{i-1,j}, x_{i+1,j}, x_{i,j-1}, x_{i,j+1}\}$ is the 5-star of the node x_{ij} as used in the standard finite difference discretization of (15).

If we comprise \mathcal{P} of the polynomial spaces $P_{ij} = \Pi_2^2$ then the resulting space $S(X, \mathcal{G}, \mathcal{P}_2)$ is not interpolatory. Its dimension is nevertheless easy to determine by using (2). Indeed, since $\dim \Pi_2^2|_{X_{ij}} = 5 = |X_{ij}|$, we conclude that $\dim \text{im } T = |X| = (N+1)^2$. Further, $\ker T$ is generated by the overlap splines $s = \{p_{ij}\}_{i,j=1}^{N-1}$, with $p_{k\ell}(x_1, x_2) = (x_1 - kh)(x_2 - \ell h)$ for a single pair (k, ℓ) , and $p_{ij} = 0$ whenever $(i, j) \neq (k, \ell)$. There are $(N-1)^2$ overlap splines of this type, hence $\dim \ker T =$

$(N-1)^2$ and $D = \dim S(X, \mathcal{G}, \mathcal{P}_2) = (N+1)^2 + (N-1)^2$, such that (5) holds in this case. However, $\Delta s = \{\Delta p_{ij}\}_{i,j=1}^{N-1} \equiv 0$ for all $s \in \ker T$, which implies that no choice of collocation nodes y_1, \dots, y_D would lead to a regular system matrix for (9).

We now reduce Π_2^2 to a five-dimensional subspace complementary to $\ker T$, by taking

$$\mathcal{P} = \{P_{ij}\}_{i,j=1}^{N-1}, \quad \text{with } P_{ij} = \text{span}\{1, x_1, x_2, x_1^2, x_2^2\}. \quad (17)$$

Then $S(X, \mathcal{G}, \mathcal{P})$ is an interpolatory overlap spline space with dimension $(N+1)^2$. By using its Lagrange basis, choosing collocation nodes $Y = X$, and evaluating Δp_{ij} at $y_{ij} = x_{ij}$, we obtain for the patches p_{ij} of any overlap spline $s = \{p_{ij}\}_{i,j=1}^{N-1} \in S(X, \mathcal{G}, \mathcal{P})$ the identities

$$\Delta p_{ij}(x_{ij}) = \frac{1}{h^2} \left(p_{ij}(x_{i-1,j}) + p_{ij}(x_{i+1,j}) + p_{ij}(x_{i,j-1}) + p_{ij}(x_{i,j+1}) - 4p_{ij}(x_{ij}) \right),$$

and see that the method of collocation in this setting is equivalent to the finite difference method for (15) with the standard five point stencil. Note that P_{ij} of (17) are the spaces of *least interpolation* [2] for the corresponding sets X_{ij} .

As an alternative to reducing the polynomial spaces, a somewhat artificial remedy is to extend each set X_{ij} by one point \tilde{x}_{ij} in G_{ij} not lying in any other $G_{k\ell}$ or on the lines $x_1 = ih$ and $x_2 = jh$. Then $S(\tilde{X}, \mathcal{G}, \mathcal{P}_2)$ with $\tilde{X} = X \cup \{\tilde{x}_{ij}\}_{i,j=1}^{N-1}$ is interpolatory. Since the Lagrange polynomial corresponding to the point \tilde{x}_{ij} is a multiple of $(x_1 - ih)(x_2 - jh)$, its Laplacian vanishes at x_{ij} , and we arrive at the same five point stencil.

Meshless Finite Difference Methods

Overlap spline collocation or discrete least squares for the problem (6) with interpolatory spaces $S(X, \mathcal{G}, \mathcal{P})$ is equivalent to a meshless finite difference method defined as follows.

Definition 4 (Meshless Finite Difference Method). Given a set $X = \{x_i\}_{i=1}^N \subset G$ of *discretization nodes* and a sequence $Y = (y_j)_{j=1}^M \subset G$ of *collocation nodes*, with $M \geq N$, choose for each $j = 1, \dots, M$ a *set of influence* $X_j^{\text{ndf}} \subset X$ and a numerical differentiation formula (ndf)

$$Lv(y_j) \approx \sum_{x_i \in X_j^{\text{ndf}}} w_{ji} v(x_i), \quad v \in U, \quad (18)$$

defined by some weights $w_{ji} \in \mathbb{R}$. Determine the vector $\hat{u} \in \mathbb{R}^N$ from the linear system

$$\sum_{x_i \in X_j^{\text{ndf}}} w_{ji} \hat{u}_i = f(y_j), \quad j = 1, \dots, M, \quad (19)$$

and use \hat{u}_j as approximation of $u(x_j)$, $j = 1, \dots, N$, for the solution u of (6).

If $M = N$ and the system (19) is nonsingular, then \hat{u} is the unique solution of (19). In the case $M > N$ an appropriate method for overdetermined systems has to be applied, usually the method of least squares. We expect that the system matrix of (19) is of full rank, which is normally observed in numerical experiments but only rarely rigorously guaranteed by the design of particular algorithms.

Numerical differentiation formulas (18) are usually obtained by requiring exactness for all elements of appropriately chosen finite-dimensional spaces P_j^{ndf} of functions whose domain of definition contains X_j^{ndf} ,

$$Lp(y_j) = \sum_{x_i \in X_j^{\text{ndf}}} w_{ji} p(x_i), \quad \forall p \in P_j^{\text{ndf}}. \quad (20)$$

this presumes, of course, that (20) is solvable with respect to w_{ji} . If the conditions (20) do not determine the weights w_{ji} uniquely, then a particular solution needs to be chosen.

Several variants of the meshless finite difference method have been proposed, depending on how the collocation nodes are chosen in relation to X , and how the formulas (18) are obtained.

It is easy to see that both the method of collocation and the method of discrete least squares for an interpolatory overlap spline space $S(X, \mathcal{G}, \mathcal{P})$ with the Lagrange basis $\ell_k = \{\ell_{ki}\}_{i=1}^m$, $k = 1, \dots, N$, is equivalent to a version of the meshless finite difference method, where the j th set of influence is the I-set for $P_j^{\text{ndf}} := P_{\sigma(j)}$ given by $X_j^{\text{ndf}} := X_{\sigma(j)} = X \cap G_{\sigma(j)}$, and the weights of (18) are obtained as $w_{jk} = L\ell_{k, \sigma(j)}(y_j)$, or, equivalently, as the unique solution of (20). Then in the case of the method of collocation the systems (9) and (19) are the same, and hence their solution vectors c , respectively \hat{u} , are identical. In the case of the method of discrete least squares the vector c of (11) gives a least squares solution of the overdetermined system (19).

Note that in the case when $Lu(y_j) := u(y_j)$ and $y_j \in X_j^{\text{ndf}}$, the solution of (20) is given by $w_{ji} = \delta_{ji}$, so that the corresponding equation in (19) becomes $\hat{u}_j = f(y_j)$, which is in particular a natural way to handle Dirichlet boundary conditions. It does not matter in this case how we choose X_j^{ndf} .

We now list some particular versions of the interpolatory overlap spline methods that can be interpreted as known meshless finite difference methods. In all of them we assume that G is a closure of a bounded domain in a smooth manifold \mathcal{M} , and (6) is a Dirichlet boundary value problem as long as $G \neq \mathcal{M}$, that is $Lu(x) = u(x)$ for all $x \in \partial G$.

1. The method of collocation for an interpolatory polynomial overlap spline space $S(X, \mathcal{G}, \mathcal{P}_q)$, with $\mathcal{M} = \mathbb{R}^d$, $G = \{G_j\}_{j=1}^N$, where the sets G_j are chosen such that each $X_j = X \cap G_j$, $j = 1, \dots, N$, is an I-set for Π_q^d . In particular, X_j may consist of $\binom{d+q}{d}$ neighbors of x_j , including x_j itself, and the collocation nodes and the mapping σ be given by $y_j = x_j$ and $\sigma(j) = j$, for all $j = 1, \dots, N$. Note that, as explained above, in this case we can also do without choosing X_j

for boundary nodes $x_j \in \partial G$ because the corresponding equations in (9) and (19) are always $c_j = f(x_j)$, respectively, $\hat{u}_j = f(x_j)$. This leads to a meshless finite difference method first considered in [13], where X_j is chosen as the set of $\binom{d+q}{d}$ nearest neighbors of x_j , and subsequently corrected should it fail to be an I-set. More recent approaches, see for example [6, 10, 19, 20] rely on the selection of X_j by algorithms that generate I-sets or subsets of I-sets, such as in classical five point stencil discussed above for the Poisson equation, while trying to optimize the accuracy of the numerical differentiation in (18) or guarantee the solvability of (19).

2. The method of collocation for the polynomial least interpolation overlap spline space $S(X, \mathcal{G}, \mathcal{P}_{\text{least}})$, where $\mathcal{M} = \mathbb{R}^d$, and each P_j is the least interpolation space of polynomials [2] for the set $X_j = X \cap G_j$, $j = 1, \dots, N$. An example of the least interpolation overlap spline space has already been discussed above for the Poisson equation solved by the classical finite difference method with the five point stencil. The meshless finite difference method that results from this setting in the case of $|X| = |\mathcal{G}|$, with X_j being a set of neighbors of $y_j = x_j$ and $\sigma(j) = j$, $j = 1, \dots, N$, has in fact been considered in [9], as a limiting case [17] of the RBF-FD method with the Gauss kernel for its shape parameter going to zero.
3. The method of collocation for the kernel-based overlap spline space $S(X, \mathcal{G}, \mathcal{P}_{K,Q})$, with a positive definite or conditionally positive definite kernel K . The resulting meshless finite difference method, where $|X| = |\mathcal{G}|$, X_j a set of neighbors of $y_j = x_j$ and $\sigma(j) = j$, $j = 1, \dots, N$, was first considered in [21]. It is commonly called the RBF-FD method in the case when K is a radial basis function, see [12] and references therein for more details. A recent overview of parameter choices can be found in [16]. In the case of nontrivial $Q \neq \{0\}$ the sets X_i do not have to be total for Q . It suffices that each X_j , $j = 1, \dots, N$, is such that the exactness equations $Lp(x_j) = \sum_{x_i \in X_j} w_{ji} p(x_i)$, $p \in Q$, are solvable with respect to the weights w_{ji} , see [7]. Then the space $S(X, \mathcal{G}, \mathcal{P}_{K,Q})$ may not be interpolatory, but $\dim P_j|_{X_j} = |X_j|$ holds for all j . Hence either an appropriate reduction of the spaces $P_j = P_{K,X_j,Q}$ or an extension of X_j for all j with $\dim P_j > |X_j|$ will produce an interpolatory space of overlap splines, similar to the five point stencil case discussed in detail above for the solution of the Poisson equation on the square with polynomial overlap splines.
4. The method of discrete least squares for the kernel-based overlap spline space $S(X, \mathcal{G}, \mathcal{P}_{K,Q})$, with $|X| = |\mathcal{G}|$, X_j being a set of neighbors of x_j , $j = 1, \dots, N$. Choose collocation nodes $Y = \{y_i\}_{i=1}^M$ without repetitions, $M > N$, and let $\sigma(i)$ be the index of the node x_j closest to y_i , that is $\|y_i - x_{\sigma(i)}\| = \min_{1 \leq j \leq N} \|y_i - x_j\|$, for some distance in G , for example the Euclidean distance for $G \subset \mathbb{R}^d$. This approach has been considered in [22], which also provides an error analysis for the corresponding meshless finite difference method. In an earlier paper [15] a similar method was applied after transforming kernel-based overlap splines $s \in S(X, \mathcal{G}, \mathcal{P}_{K,Q})$ into functions s_Γ created by a suitable partition of unity Γ .
5. The method of discrete least squares for the kernel-based overlap spline space $S(X, \mathcal{G}, \mathcal{P}_{K,Q})$, where the sequence Y is obtained by combining all sets $X_i = \{x_j : j \in J_i\}$ and $\sigma(k)$ is the index i of the set J_i the node $y_k = x_j$ comes with.

In other words,

$$Y = (x_j, j \in J_1, x_j, j \in J_2, \dots, x_j, j \in J_m),$$

$$(\sigma(1), \dots, \sigma(|Y|)) = (\underbrace{1, \dots, 1}_{|J_1| \text{ times}}, \underbrace{2, \dots, 2}_{|J_2| \text{ times}}, \dots, \underbrace{m, \dots, m}_{|J_m| \text{ times}}).$$

The resulting meshless finite difference method has been considered in [8], with the full rank property and error analysis provided under certain assumptions for elliptic equations (6) on closed manifolds.

References

1. M. Ainsworth, G. Andriamaro, and O. Davydov. Bernstein-Bézier finite elements of arbitrary order and optimal assembly procedures. *SIAM Journal on Scientific Computing*, 33(6):3087–3109, 2011.
2. C. de Boor and A. Ron. The least solution for the polynomial interpolation problem. *Mathematische Zeitschrift*, 210(3):347–378, 1992.
3. S. Brenner and R. Scott. *The Mathematical Theory of Finite Element Methods, 3rd Edition*. Springer Science & Business Media, 2008.
4. M. D. Buhmann. *Radial Basis Functions*. Cambridge University Press, New York, NY, USA, 2003.
5. F. Chen and P. J. Wong. Discrete cubic spline method for second-order boundary value problems. *International Journal of Computer Mathematics*, 91(5):1041–1053, 2014.
6. O. Davydov. Selection of sparse sets of influence for meshless finite difference methods, arXiv:1908.01567, 2019.
7. O. Davydov. Approximation with conditionally positive definite kernels on deficient sets. In M. N. Gregory E. Fasshauer and L. L. Schumaker, editors, *Approximation Theory XVI: Nashville 2019*, pages 27–38. Springer Berlin Heidelberg, 2021.
8. O. Davydov. Error bounds for a least squares meshless finite difference method on closed manifolds. *Advances in Computational Mathematics*, 49:article number 48, 2023.
9. O. Davydov and D. T. Oanh. On the optimal shape parameter for Gaussian radial basis function finite difference approximation of the Poisson equation. *Comput. Math. Appl.*, 62:2143–2161, 2011.
10. O. Davydov and M. Safarpour. A meshless finite difference method for elliptic interface problems based on pivoted QR decomposition. *Applied Numerical Mathematics*, 161:489–509, 2021.
11. G. Fasshauer. *Meshfree Approximation Methods with MATLAB*. World Scientific Publishing Co., Inc., River Edge, NJ, USA, 2007.
12. B. Fornberg and N. Flyer. *A Primer on Radial Basis Functions with Applications to the Geosciences*. Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 2015.
13. P. S. Jensen. Finite difference techniques for variable grids. *Comput. Struct.*, 2(1-2):17–29, 1972.
14. M.-J. Lai and L. L. Schumaker. *Spline functions on triangulations*. Cambridge University Press, 2007.
15. E. Larsson, V. Shcherbakov, and A. Heryudono. A least squares radial basis function partition of unity method for solving PDEs. *SIAM Journal on Scientific Computing*, 39(6):A2538–A2563, 2017.
16. S. Le Borne and W. Leinen. Guidelines for RBF-FD discretization: Numerical experiments on the interplay of a multitude of parameter choices. *Journal of Scientific Computing*, 95(1):8, 2023.

17. R. Schaback. Multivariate interpolation by polynomials and radial basis functions. *Constructive Approximation*, 21(3):293–317, 2005.
18. L. Schumaker. *Spline Functions: Basic Theory*. Cambridge University Press, 2007.
19. B. Seibold. Minimal positive stencils in meshfree finite difference methods for the Poisson equation. *Comput. Methods Appl. Mech. Eng.*, 198(3-4):592–601, 2008.
20. L. Shen, G. Lv, and Z. Shen. A finite point method based on directional differences. *SIAM Journal on Numerical Analysis*, 47(3):2224–2242, 2009.
21. A. I. Tolstykh and D. A. Shirobokov. On using radial basis functions in a ‘finite difference mode’ with applications to elasticity problems. *Computational Mechanics*, 33(1):68–79, 2003.
22. I. Tominec, E. Larsson, and A. Heryudono. A least squares radial basis function finite difference method with improved stability properties. *SIAM Journal on Scientific Computing*, 43(2):A1441–A1471, 2021.
23. H. Wendland. *Scattered Data Approximation*. Cambridge University Press, 2004.