

Lieb-Schultz-Mattis constraint, symmetries, excitation and anomaly of the quantum spin ice models in the planar pyrochlore lattice

Zijian Xiong^{1,2,3}

¹*Department of Physics, Chongqing University, Chongqing, 401331, China*

²*Chongqing Key Laboratory for Strongly Coupled Physics, Chongqing, 401331, China*

³*Department of Applied Physics, University of Tokyo, Tokyo 113-8656, Japan*

We consider the quantum spin ice models in the planar pyrochlore lattice. The models are obtained by perturbing the Ising model with different lattice symmetry preserving quantum fluctuations. We map these models to a compact U(1) lattice gauge theory and discuss its global symmetries which include the 1-form U(1) global symmetry. Utilizing the higher form generalization of the Lieb-Schultz-Mattis theorem and also the monopole effect in the compact lattice gauge theory, we show that these models have "persistent" translation symmetry breaking valence bond solid phase around the Ising limit even the perturbation explicitly breaks all the spin rotation symmetry. We find the excitation in this phase can be described by an abelian Higgs model with charge ± 2 monopoles, this model is recently found to have a mixed 't Hooft anomaly. The one of the consequences of this anomaly is the domain wall in this phase may support deconfined excitations on it, we explicitly construct a solvable domain wall carries such anomaly. The spectrum of the excitations is also studied, we find the spectral weight of the spectrum in the low energy sector along the high symmetry momentum paths is strongly restricted by the 1-form global symmetries. Specially, we show the spectrum directly measures on the domain wall along $q_2 = \pm q_1$ momentum paths.

I. INTRODUCTION

The Lieb-Schultz-Mattis (LSM) theorem [1] and its higher dimensional generalizations [2, 3] are remarkable in quantum many body physics. On the one hand, they provide a non-trivial way for constraining the low energy physics by the microscopic degrees of freedom. On the other hand, the consequence of these theorems rule out the symmetric, trivial (non-degenerate) gapped ground state for spin systems with a certain spatial and internal symmetries. Thus, for a spin system which satisfies the conditions of the theorem, the ground state must either be i) symmetry broken; or ii). gapless; or iii). topological ordered (this can only happen in systems in two or higher dimensions). This actually provides a shortcut for hunting the novel quantum spin liquids [4, 5], specifically, seeking for the systems which fulfill the requirement for the symmetry and show a gapped phase without symmetry breaking.

The original LSM theorem [1–3] relies on the translation symmetry and continuous spin rotation symmetry. Recently, the theorem has been generalized to include space group symmetries [6–9], discrete internal symmetries [7, 10] and even generalized symmetries [11–17]. These generalizations extend the applicability of the LSM theorem. Among the generalizations, a user friendly version of the theorem is the lattice homotopy method [7, 8]. This method states that all the LSM type constraints are completely determined by the spatial distribution of the microscopic degrees of freedom, once the symmetry group (includes the spatial symmetry and the internal symmetry) is given. Thus, one can rule out the symmetric trivial gapped phase in a certain symmetry group even without knowing the concrete model.

In quantum field theory, in some cases, certain global symmetries in a field theory can not be promoted to the gauge symmetries without introducing nonlocal terms. This obstruction for gauging the global symmetries is called 't Hooft

anomaly [18–21], and it also leads to a constraint for the infrared (IR) physics similar to the LSM theorem which precludes the symmetric, trivial gapped vacuum. Indeed, this similarity is not a coincidence. Recent investigation in the symmetry protected topological (SPT) phase and their relation to the 't Hooft anomaly brings new understanding for the LSM theorem [22–30]. It was recognized that the LSM theorem can be understood as a mixed 't Hooft anomaly between the spatial symmetry and the internal symmetry [22, 25–27]. Via the anomaly inflow mechanism, the system triggered LSM constraint can also be viewed as the boundary of some higher dimensional SPT bulk [22]. Since the anomalies relating to the LSM physics can be traced back to the lattice model, these anomalies is sometimes called LSM anomaly or intrinsic anomaly [27]. Besides that, there are also non-LSM anomalies [27, 28].

In addition to the LSM theorems, mixed 't Hooft anomaly also plays an important role in many novel physics of condensed matter, for example, deconfined quantum criticality [20, 31–33], emergibility [28, 34] and classification of SPT phases [23, 24, 35, 36]. Recently, an interesting phenomenon is found that the domain wall in the confined Z_2 valence bond solid (VBS) phase can support deconfined excitations on it. And this nontriviality of the domain wall is guaranteed by mixed 't Hooft anomaly [19, 37].

Furthermore, an interesting observation from the numerical studies of the spin ice models in the planar pyrochlore lattice is that the ground states of these models around the Ising limit have "persistent" translation symmetry breaking VBS phase [38–47]. Here, the "persistent" means VBS remains even the (lattice symmetry preserving) perturbation breaks all the spin rotation symmetry. In other words, no symmetric trivial gapped phase. Motivated by these, in this paper, we study the quantum spin ice models in the planar pyrochlore lattice (see Fig.1 (a)) which are known to have a two fold degenerate VBS phase [38–40, 42, 48] and a faithful lattice gauge theory description [49–52]. The corresponding gauge theory is

equivalent to the quantum link model in high energy physics [48, 53, 54]. This is useful for establishing the low energy field theory from the lattice model. Benefit from this, we can study the ground state by both the LSM theorem in the ultra-violet (UV) level lattice model and the 't Hooft anomaly in the IR theory. On the other hand, although the properties of the ground state of these ice models are well studied, the excitation spectrum is still not fully understood. Another goal of this paper is to figure out the excitation spectrum studied by quantum monte carlo recently [55].

This paper is organized as follows. In Sec. II, we introduce the lattice homotopy method, and apply this method to analyze the LSM physics for the common spin ice models in the planar pyrochlore lattice. Sec. III is devoted to the two dimensional quantum spin ice model and its gauge theory. We introduce the mapping procedure, global symmetries of the gauge theory and show how the LSM theorem emerges in this gauge theory. The continuum field theory dual to this gauge theory and the monopole effect is also studied. In Sec. IV, we study the excitations in the two fold degenerate VBS phase and explicitly construct a solvable domain wall which supports the deconfined excitations on it. And we find the excitations in this phase can be described by an abelian Higgs model with a mixed 't Hooft anomaly which is studied in ref. [19]. We also discuss the spectrum (dynamical structure factor) which is related to our recent work [55]. We find most of the features of the spectrum can be understood from symmetries. Remarkably, we find the spectrum directly measures on the domain wall along some high symmetry paths in the momentum space, and the spectrum is also solvable. In Appendix A, we provide the partition function for the monopoles. In Appendix B, we discuss the deformation of the 't Hooft loop operator. In Appendix C, we discuss the charge conjugation symmetry in the low energy theory.

II. LATTICE HOMOTOPY ANALYSIS

In this work, we are interested in the spin-1/2 model on the planar pyrochlore lattice, see Fig. 1 (a). Before doing any calculation for a concrete model, we would like to know the necessary symmetries to trigger the LSM theorem. This goal can be easily fulfilled by the lattice homotopy method even without writing down the concrete model.

The input data of this method is the symmetry group G , here we assume $G = G_{\text{int}} \times G_s$, where G_{int} represents the internal symmetry group which here is related to the spin, and G_s represents the spatial symmetry group of the lattice. In the spirit of the lattice homotopy, for a lattice system, one can keep the symmetry group $G_{\text{int}} \times G_s$ and deform the lattice, then the degrees of freedom can be moved to the high symmetry positions (i.e. irreducible Wcykoff positions) of G_s . With the help of the three no go theorems about the basic space group symmetries [6, 7]:

- Bieberbach no go theorem: define the fundamental domain as a region which can tile the lattice under the translations and glide symmetries. If the fundamental domain carries the nontrivial projective representation

under G_{int} , then the symmetric trivial gapped phase is forbidden. This theorem is for nonsymmorphic G_s .

- Mirror no go theorem: if there is a translation symmetry along the mirror line, and the projective representation on the translation unit along this mirror line is nontrivial, then the symmetric trivial gapped phase is forbidden.
- Rotation no go theorem: if the projective representation at an even fold rotation center is nontrivial, then the symmetric trivial gapped phase is forbidden.

Then, the lattice homotopy method [7, 8, 28] tells that the existence of the LSM type constraint will be determined by the distribution of the physical degrees of freedom (the representation of G_{int}) on the irreducible Wcykoff positions (given by G_s).

For the first step, we consider the G_s . Since the largest rotation symmetry in the planar pyrochlore lattice is four fold, and there are mirror lines intersecting at $\pi/4$, so the space group of this lattice is the No.11 wallpaper group p4mm [56]. The generators of this group can be chosen as translation symmetry along $\mathbf{e}_1$ and $\mathbf{e}_2$ direction, 4 fold rotation around the plaquette center and mirror line along the $\mathbf{e}_2$. The translation unitcell of this group is shown in Fig. 1 (b), where a,b,c denote the irreducible Wcykoff positions. Putting the physical degrees of freedom (DOF) on the c positions in the translation unitcell, the planar pyrochlore lattice can be generated by the translation unitcell with the primitive translation vectors $\mathbf{e}_1$ and $\mathbf{e}_2$. As the c position is the 2 fold rotation center, the rotation no go theorem will be triggered once the DOF carries nontrivial projective representation of G_{int} and the symmetric trivial gapped phases will be forbidden.

For the second step, we consider the projective representation of some common internal symmetries in the spin model. It is known that the projective representation (PR) for a group G_{int} is classified by the second cohomology group [36] $\mathcal{H}^2[G_{\text{int}}, U(1)]$. We list $\mathcal{H}^2[G_{\text{int}}, U(1)]$ for some common internal symmetry groups [36] in Table I.

- Z_2 symmetry group. $\mathcal{H}^2[Z_2, U(1)] = Z_1$ tells that there is only one trivial PR, thus the trivial phase is allowed.




FIG. 1. (a) Planar pyrochlore lattice, the black dots denote the spins. (b) Translation unitcell of the No.11 wallpaper group p4mm, a,b,c are the Wcykoff Letters. a,b are the 4 fold rotation centers, while c is the 2 fold rotation center, they are denoted by the squares and ellipses, respectively.

TABLE I. Table of the second cohomology group $\mathcal{H}^2[G_{\text{int}}, U(1)]$ for some common symmetry groups G_{int} .

Symmetry group G_{int}	$\mathcal{H}^2[G_{\text{int}}, U(1)]$
Z_2	Z_1
$U(1)$	Z_1
$Z_2 \times Z_2$	Z_2
$U(1) \rtimes Z_2$	Z_2
$SO(3)$	Z_2

A typical model with this symmetry group is the transverse field Ising model [41–43] $H = J \sum_{i,j} S_i^x S_j^x - \Gamma \sum_i S_i^z$, where the generator can be chosen as the π rotation around the spin x axis: $e^{-i\pi S^x}$, where $S^x = \sum_i S_i^x$.

- $U(1)$ symmetry group. There is only one trivial PR for this group, and the trivial phase is allowed. Typical model is the XXZ model under the Zeeman field [44–47].
- $Z_2 \times Z_2$ symmetry group. A representative model is XYZ model $H = J_x \sum_{i,j} S_i^x S_j^x + J_y \sum_{i,j} S_i^y S_j^y + J_z \sum_{i,j} S_i^z S_j^z$. Z_2 symmetries can be selected as the π rotation around the spin x and y axis, respectively. There is a nontrivial PR which is carried by half integer spin. Further, this model has time reversal symmetry which also forms a Z_2 group. There are sufficient symmetries to forbid the trivial phase even without considering the time reversal symmetry.
- $U(1) \rtimes Z_2$ symmetry group. A representative model is XXZ model [38–40, 55] $H = J_{xy} \sum_{i,j} [S_i^x S_j^x + S_i^y S_j^y] + J_z \sum_{i,j} S_i^z S_j^z$. This model has a $U(1)$ symmetry around the spin z axis and Z_2 symmetry (π rotation) around the spin x (or y) axis. The $U(1)$ symmetry forms a group with element $U_\theta^z = e^{-i\theta S^z}$, $S^z = \sum_i S_i^z$. The Z_2 symmetry also forms a group $\{U_\pi^x, (U_\pi^x)^2 = 1\}$. Since $(U_\pi^x)^{-1} U_\theta^z U_\pi^x = U_{-\theta}^z$, and $U_\pi^x U_\theta^z U_\pi^x = U_\pi^x U_\theta^z (U_\pi^x)^{-1} U_\theta^z U_\pi^x = U_\pi^x U_\theta^z U_\pi^x$, we can define a group homomorphism $\varphi : Z_2 \rightarrow \text{Aut}(U(1))$ by $\varphi_h(n) = hnh^{-1}$, for all $h \in Z_2$, $n \in U(1)$. Such that $(h_1, n_1)(h_2, n_2) = (h_1 h_2, \varphi_{h_1}(n_1) n_2)$. Hence the symmetry group is $U(1) \rtimes Z_2$. This group has a nontrivial PR which is known to be carried by half integer spin. There are also enough symmetry to trigger LSM constraint even without considering the time reversal symmetry. In addition, the existence of the nontrivial projective representation for this symmetry group can be easily deduced from the fact that it has a subgroup $Z_2 \times Z_2$ group.
- $SO(3)$ symmetry group. A representative model is the Heisenberg model [57–60]. There is a nontrivial PR which is carried by half integer spin.

From above analysis, we find that the ground states of the spin-1/2 XXZ model, XYZ model and Heisenberg model on the planar pyrochlore lattice are constrained by LSM type theorem which forbids the symmetric trivial gapped phase.

Recently, the 't hooft anomaly in the Heisenberg type model on the planar pyrochlore lattice is studied in ref.[60]. And the LSM physics in the XYZ model on the lattice with the same space group has been studied in ref.[10]. However, in this paper, we are interested in the quantum spin ice models on the planar pyrochlore lattice which are usually obtained from perturbing the classical Ising model by quantum fluctuations. Some common models are transverse field Ising model (TFIM) [41–43], TFIM with Zeeman field [41, 43, 61], TFIM with different nearest neighbor and next nearest neighbor interactions [43], XXZ model [38–40, 51, 55], XXZ model with Zeeman field [44–47]. It is clear that only the XXZ model triggers the LSM theorem from the lattice homotopy method, but this does not mean other models must be trivial.

III. QUANTUM SPIN ICE MODELS

We begin with the Ising model

$$H^{\text{Ising}} = J_z \sum_{\langle i,j \rangle} S_i^z S_j^z, \quad (1)$$

where $J_z > 0$, $\langle i, j \rangle$ denotes the nearest-neighbor bond and the connected diagonal bond, as shown in Fig.1 (a). Due to the corner sharing geometry of the lattice, the Ising model can be rewritten as

$$H^{\text{Ising}} = \frac{J_z}{2} \sum_{\square} (S_{\square}^z)^2 + C, \quad (2)$$

where $S_{\square}^z = \sum_{i \in \square} S_i^z$ is the summation of z component of 4 spins belonging to the same plaquette with diagonal bonds. We dub such plaquette as crossing plaquette, and the plaquette without diagonal bonds as empty plaquette. C is a constant. This Ising model is known to be solvable and equivalent to the classical six vertex model [62]. The ground state of the Ising model eq.(2) satisfies $S_{\square}^z = 0$ for every crossing plaquette. Such local constraint is usually called the ice rule in the literatures [38, 39]. For this lattice, there are six classical configurations fulfill this constraint in every crossing plaquette, thus the classical ground state has extensive degeneracy. And this extensively degenerate ground state manifold is usually dubbed as ice manifold.

Now we introduce the quantum fluctuations. We first consider the XXZ model

$$H = -J_{xy} \sum_{\langle i,j \rangle} (S_i^+ S_j^- + S_i^- S_j^+) + J_z \sum_{\langle i,j \rangle} S_i^z S_j^z, \quad (3)$$

where $J_{xy} > 0$. The choice of the ferromagnetic quantum fluctuation does not matter, but this choice is free from the sign problem and can be simulated by quantum monte carlo [40, 46, 55]. Even for $J_{xy} \ll J_z$, the quantum effect will be triggered, which greatly modifies the classical ground state via the tunneling among different classical configurations in the ice manifold and lifts the degeneracy. This quantum tunneling can be described by an effective Hamiltonian from the degenerate perturbation theory

$$H^{\text{eff}} = -t \sum_{\square} (S_i^+ S_j^- S_k^+ S_l^- + S_i^- S_j^+ S_k^- S_l^+), \quad (4)$$




FIG. 2. PVBS phase in the planar pyrochlore lattice. (a),(b) are two degenerate configurations, the purple and green circles denote the flippable plaquettes.

where i, j, k, l label the spins in an empty plaquette sequentially, $t = 4J_{xy}^2/J_z$. Due to the quantum tunneling effect, the flippable plaquette with alternating up and down spins in the empty plaquette has lower energy than the unflippable one and this effective Hamiltonian can be regarded as some kinetic energy. Moreover, the flippable plaquette repels each other. In other words, all the flippable plaquettes tend to locate in one of the two sublattices. Based on these properties, the quantum ground state is expected to be a plaquette valence bond solid (PVBS) phase with the broken translation symmetry, see Fig.2. And this is confirmed by some related numerical simulations [38–40, 48, 53, 54].

It is worth nothing that the transverse field Ising model [41, 42, 51] $H = J_z \sum_{i,j} S_i^z S_j^z - \Gamma \sum_i S_i^x$ has the same effective Hamiltonian eq.(4) with modified t from the lowest non-vanishing term in the degenerate perturbation theory. The reason for the same effective Hamiltonian can be easily understood from the gauge theory below.

Moreover, there is a local conservation symmetry of $S_{\square}^z$ in the effective Hamiltonian eq.(4), and the symmetry generator is $\exp(i\alpha S_{\square}^z)$. This local symmetry implies the U(1) gauge structure for the low energy physics of the quantum ice [52]. Interestingly, this local U(1) symmetry is emergent in the low energy sector and enforced by the large Ising interaction [63], since both the XXZ model and transverse field Ising model do not have such symmetry.

For XXZ model, when $J_{xy} \gg J_z$, the ice rule no longer holds, and the ground state is a simple ferromagnetic (FM) phase. The transition point separating PVBS and FM phases [46] locates around $J_{xy}/J_z \sim 0.065$. The structure of the phase diagram is consistent with the lattice homotopy analysis, where no place for the symmetric trivial gapped phase. For TFIM, there is a transition point separating the PVBS and paramagnetic phases at $\Gamma/J_z \sim 0.28$ [43]. For TFIM with different nearest neighbor J_1 and next neighbor neighbor J_2 interactions $H = J_1 \sum_{\langle i,j \rangle} S_i^z S_j^z + J_2 \sum_{\langle\langle i,j \rangle\rangle} S_i^z S_j^z - \Gamma \sum_i S_i^x$, PVBS phase lies in the region $1.0013 > J_2/J_1 > 0.9996$ when $\Gamma/J_1 = 0.2$ [43]. It seems that PVBS phase “persists” in these models although they have lower spin symmetry.

A. Mapping to a lattice gauge theory

Since there is local U(1) symmetry in the low energy sector, now we rewrite the theory into a gauge theory [49–52]. It is convenient to express the spin in terms of link variable: $S_i^z = S_{\mu}^z(\mathbf{r})$. Here $\mathbf{r}$ denotes the sites of the square lattice in Fig.1 (a), the spin sits on the link (denote by $\mathbf{r} + \mathbf{e}_{\mu}$, $\mu = 1, 2$) of the square lattice. $\mathbf{e}_{\mu}$ is shown in Fig.1 (a). Then we introduce the rotor representation for the spin

$$\begin{aligned} S_{\mu}^z(\mathbf{r}) &= n_{\mu}(\mathbf{r}) - 1/2, \\ S_{\mu}^{\pm}(\mathbf{r}) &= e^{\pm i\phi_{\mu}(\mathbf{r})}, \end{aligned} \quad (5)$$

where $[\phi_{\mu}(\mathbf{r}), n_{\nu}(\mathbf{r}')] = i\delta_{\mu,\nu}\delta_{\mathbf{r},\mathbf{r}'}$. For spin-1/2, it is clear that $n_{\mu}(\mathbf{r})$ takes 1,0 corresponding to spin up and down, respectively. As $n_{\mu}(\mathbf{r})$ takes integer value, $\phi_{\mu}(\mathbf{r})$ is an angular variable and takes value in $[0, 2\pi)$. It can be checked that this representation keeps the relation $[S^{\pm}, S^z] = \mp S^{\pm}$. In this representation, the ice rule is expressed as

$$\begin{aligned} \sum_{i \in \square} S_i^z &= S_1^z(\mathbf{r}) + S_1^z(\mathbf{r} - \mathbf{e}_1) + S_2^z(\mathbf{r}) + S_1^z(\mathbf{r} - \mathbf{e}_2) = 0, \\ \rightarrow n_1(\mathbf{r}) + n_1(\mathbf{r} - \mathbf{e}_1) + n_2(\mathbf{r}) + n_2(\mathbf{r} - \mathbf{e}_2) &= 2, \end{aligned} \quad (6)$$

Actually, this rotor representation induces a dimer representation [49, 50], $n_{\mu}(\mathbf{r})$ can be regarded as the number of dimer in the link $\mathbf{r} + \mathbf{e}_{\mu}$. In this understanding, the ice physics here is equivalent to a link model [48, 53, 54] defined on the square lattice with the local constraint that every site is required to have two and only two dimers. This dimer scenario will be used in the following section.

As revealed in eq.(6), every site is required to have two and only two dimers, consider a square lattice with $L \times L$ size and take the periodic boundary condition, then we have a global constraint due to the local constraint, namely, $\sum_{\mathbf{r}, \mu} n_{\mu}(\mathbf{r}) = L^2$. This can be understood as, for a square lattice with L^2 sites, there are $2L^2$ spins, the ground state is restricted by the ice rule which implies that only half of the spins pointing up direction ($n=1$) and this leads to the global constraint. Usually, n field can be relaxed to take all the integer values, to become a faithful representation of the original spin-1/2 problem, it is readily to add a Lagrange multiplier term $\frac{1}{2g} \sum_{\mathbf{r}, \mu} [(n_{\mu}(\mathbf{r}) - \frac{1}{2})^2 - \frac{1}{4}]$.

Now, the fields can be defined by

$$\begin{aligned} E_{\mu}(\mathbf{r}) &= e^{i\mathbf{Q}_0 \cdot \mathbf{r}} (n_{\mu}(\mathbf{r}) - \frac{1}{2}), \\ a_{\mu}(\mathbf{r}) &= e^{i\mathbf{Q}_0 \cdot \mathbf{r}} \phi_{\mu}(\mathbf{r}), \end{aligned} \quad (7)$$

where $\mathbf{Q}_0 = (\pi, \pi)$ is a staggering wave vector. And this definition assigns the positive directions for the fields. These fields are canonically conjugated to each other $[a_{\mu}(\mathbf{r}), E_{\nu}(\mathbf{r}')] = i\delta_{\mu,\nu}\delta_{\mathbf{r},\mathbf{r}'}$. E field takes value in $\pm 1/2$ (can be extended to take all the half integer values). a field is an angular variable. Based on the canonical relation, two useful relations can be derived

$$\begin{aligned} [e^{i\omega a_{\mu}(\mathbf{r})}, E_{\mu}(\mathbf{r})] &= -\omega e^{i\omega a_{\mu}(\mathbf{r})}, \\ [a_{\mu}(\mathbf{r}), e^{i\omega E_{\mu}(\mathbf{r})}] &= -\omega e^{i\omega E_{\mu}(\mathbf{r})}, \end{aligned} \quad (8)$$

where ω is an arbitrary function. Consider the eigenstate of $E_\mu(\mathbf{r})$, $E_\mu(\mathbf{r})|\mathcal{E}_\mu(\mathbf{r})\rangle = \mathcal{E}_\mu(\mathbf{r})|\mathcal{E}_\mu(\mathbf{r})\rangle$, then $E_\mu(\mathbf{r})e^{\pm ia_\mu(\mathbf{r})}|\mathcal{E}_\mu(\mathbf{r})\rangle = (\mathcal{E}_\mu(\mathbf{r}) \pm 1)e^{\pm ia_\mu(\mathbf{r})}|\mathcal{E}_\mu(\mathbf{r})\rangle$. Thus, $e^{\pm ia_\mu(\mathbf{r})}$ plays the role of the ladder operator for $E_\mu(\mathbf{r})$.

Define the lattice derivative [64] as $\Delta_\mu^- f(\mathbf{r}) = f(\mathbf{r}) - f(\mathbf{r} - \mathbf{e}_\mu)$, then the ice rule now has a more familiar Gauss's law form

$$\begin{aligned} 0 &= E_1(\mathbf{r}) - E_1(\mathbf{r} - \mathbf{e}_1) + E_2(\mathbf{r}) - E_2(\mathbf{r} - \mathbf{e}_2) \\ &= \sum_\mu \Delta_\mu^- E_\mu(\mathbf{r}), \end{aligned} \quad (9)$$

further, we define the lattice divergence as

$$(\text{div } E)_\mathbf{r} = \sum_\mu \Delta_\mu^- E_\mu(\mathbf{r}) = Q_\mathbf{r}, \quad (10)$$

which measure the effective charge (we will call it charge in the following) at the site $\mathbf{r}$. And, $Q_\mathbf{r}$ takes values in 0, ± 1 , ± 2 for the spin-1/2 model. In contrast to the well studied gauge theory description for the dimer model, there is no static background charge in our problem [49, 50].

The kinetic term eq.(4) is now expressed as

$$\begin{aligned} H^{\text{eff}} &= -2t \sum_{\mathbf{r}} \cos[a_1(\mathbf{r}) + a_2(\mathbf{r} + \mathbf{e}_1) - a_1(\mathbf{r} + \mathbf{e}_2) - a_2(\mathbf{r})] \\ &= -2t \sum_{\mathbf{r}} \cos[\Delta_1^+ a_2(\mathbf{r}) - \Delta_2^+ a_1(\mathbf{r})] \\ &= -2t \sum_{\mathbf{r}} \cos\left[\sum_{\mu,\nu} \epsilon_{\mu\nu} \Delta_\mu^+ a_\nu(\mathbf{r})\right], \end{aligned} \quad (11)$$

here, $\Delta_\mu^+ f(\mathbf{r}) = f(\mathbf{r} + \mathbf{e}_\mu) - f(\mathbf{r}) = \Delta_\mu^- f(\mathbf{r} + \mathbf{e}_\mu)$. And $\epsilon_{\mu\nu}$ is the Levi-Civita symbol, e.g. $\epsilon_{1,2} = -\epsilon_{2,1} = 1$. We can define the lattice curl as

$$(\text{curl } a)_\mathbf{r} = \sum_{\mu,\nu} \epsilon_{\mu\nu} \Delta_\mu^+ a_\nu(\mathbf{r}). \quad (12)$$

With these settings, we have a faithful description for the low energy physics of the spin ice when $g \rightarrow 0$. Now the effective Hamiltonian takes a compact quantum electrodynamics (cQED) form [50, 65]

$$H^{\text{cQED}} = \frac{1}{2g} \sum_{\mathbf{r},\mu} [E_\mu^2(\mathbf{r}) - \frac{1}{4}] - 2t \sum_{\mathbf{r}} \cos[(\text{curl } a)_\mathbf{r}]. \quad (13)$$

It is known that the quantum dimer model in the square lattice takes a very similar form [49, 50] as eq.(13), but the constraints (local and global) are modified.

It can be checked that $[(\text{div } E)_\mathbf{r}, (\text{curl } a)_\mathbf{r}] = 0$, thus, $[(\text{div } E)_\mathbf{r}, H^{\text{cQED}}] = 0$. This implies a gauge symmetry generated by $U = e^{i \sum_{\mathbf{r}} \theta_{\mathbf{r}} Q_\mathbf{r}}$, where $\theta_{\mathbf{r}}$ is an arbitrary function but not operator. So H^{cQED} is gauge invariant. It is obviously that this gauge symmetry is from the local rotation symmetry. Using eq.(8), one can find $a_\mu(\mathbf{r})$ transform as the usual vector potential

$$U a_\mu(\mathbf{r}) U^\dagger = a_\mu(\mathbf{r}) + \theta_{\mathbf{r}} - \theta_{\mathbf{r}+\mathbf{e}_\mu} = a_\mu(\mathbf{r}) - \Delta_\mu^+ \theta_{\mathbf{r}}, \quad (14)$$

under the gauge transformation. Further, one can deduce that the simplest gauge invariant and physical dynamics is

from $\cos[(\text{curl } a)_\mathbf{r}]$, which is equivalent to eq.(4). Hence, the gauge invariant principle forces the XXZ type perturbation, the TFIM type perturbation and other perturbation act as the same way in the low energy at least in the lowest order. Furthermore, the gauge invariant principle also implies that any higher order perturbation terms act on the ice manifold should be loop like operators, otherwise they will create charges and correspond to the higher energy.

B. Symmetries and LSM type constraint for the gauge theory

Besides the gauge symmetry, there are also some global symmetries in the gauge theory eq.(13). Some of them will be useful to restrict the possible ground states.

- The translation symmetries of the square lattice still remain.
- There is a charge conjugation symmetry C which maps $E_\mu(\mathbf{r})$ to $-E_\mu(\mathbf{r})$, $a_\mu(\mathbf{r})$ to $-a_\mu(\mathbf{r})$ and also $Q_\mathbf{r}$ to $-Q_\mathbf{r}$. Actually, this symmetry can be recognized as a Z_2 symmetry U_π^x (the π rotation around spin x axis) in the original spin model eq.(3). This can be seen by $U_\pi^x S^\alpha (U_\pi^x)^{-1} = -S^\alpha$, where $\alpha = y, z$, and, $U_\pi^x S^\pm (U_\pi^x)^{-1} = S^\mp$. Recall the definition of the fields $E_\mu(\mathbf{r}) = e^{i\mathbf{Q}_0 \cdot \mathbf{r}} S_\mu^z(\mathbf{r})$, $S_\mu^\pm(\mathbf{r}) = \exp(\pm i\mathbf{e}_\mu \cdot \mathbf{r}) a_\mu(\mathbf{r})$. Hence, $U_\pi^x : E \rightarrow -E, a \rightarrow -a$. Further discussion on the charge conjugation symmetry is provided in Appendix C.
- Topological symmetry. This symmetry is associated with the conservation of the flux of a , since $\sum_{\mathbf{r}} (\text{curl } a)_\mathbf{r} = 0$ in PBC (closed manifold). The monopole like configuration of a field is still allowed [66–68] in the theory eq.(13). In the continuum theory, this symmetry is clear. If the fluctuation of a field is small, we may expand the cosine term in eq.(13), and let us denote $B_\mathbf{r} = (\text{curl } a)_\mathbf{r}$, then eq.(13) reduces to a familiar Maxwell form $E^2 + cB^2$, c is a coefficient. It is usually written as F^2 in continuum field theory, where $F = da$ is the electromagnetic field tensor. It is known that in (2+1) dimension, one can define a topological number $Q_{\text{top}} = \frac{1}{2\pi} \int_{M_2} da$ in 2-manifold M_2 . And it is associated with a global $U(1)$ topological symmetry $U(1)_{\text{top}}$ [69].
- 1-form $U(1)$ global symmetry [11, 15, 16]. This can be seen as following. Using the canonical relation, we can find

$$[(\text{curl } a)_\mathbf{r}, \sum_{\mathbf{r}' \in \gamma} E_\mu(\mathbf{r}') + E_\mu(\mathbf{r}' + \mathbf{e}_\nu)] = 0, \quad (15)$$

where $\mu \neq \nu$, γ is some "curve" which will be clear below. Hence, under the periodic boundary condition with $L \times L$ system size, we can define some conserved quantities on the noncontractible loops (for example, γ_1

and γ_2 in Fig.3 (a)),

$$\begin{aligned}\tilde{W}_{\gamma_1} &= \sum_{\mathbf{r} \in \gamma_1} E_2(\mathbf{r}) = \sum_{n=0}^{L-1} E_2(\mathbf{r} + n\mathbf{e}_1), \\ \tilde{W}_{\gamma_2} &= \sum_{\mathbf{r} \in \gamma_2} E_1(\mathbf{r}) = \sum_{n=0}^{L-1} E_1(\mathbf{r} + n\mathbf{e}_2),\end{aligned}\quad (16)$$

it can be readily checked that $[\tilde{W}_{\gamma_i}, H^{\text{cQED}}] = 0$, $i = 1, 2$ and $\tilde{W}_{\gamma_i} \leq L/2$. These operators are the so called 't Hooft loop operators [50, 69, 70], and their eigenvalues are usually called winding numbers [71]. From these properties, the winding number is stable against the local fluctuations, thus, it is sometimes called topological number. One can also find another relation $[(\text{curl } a)_{\mathbf{r}}, \sum_{\mathbf{r}' \in \gamma} E_{\mu}(\mathbf{r}') - E_{\nu}(\mathbf{r}' + \mathbf{e}_{\mu})] = 0$, where $\mu \neq \nu$. Using this relation, we can define the 't Hooft loop operator with the staircase like γ .

Now, we define the Wilson line operator $W_{\omega} = e^{i \sum_{\mathbf{r} \in \omega} a_{\mu}(\mathbf{r})}$, ω is an arbitrary "curve" in the lattice, it can be open with two ends, contractible loop or non-contractible loop (such as ω_1 , ω_2 and ω_3 in Fig.3 (b)). Acting on the vacuum (ice manifold), the Wilson operator defined on the open "curve" will create a pair of charges on its ends. Since only the loop operator is gauge invariant, so we only consider the loop operator below. For a noncontractible loop (say, ω_3 in Fig.3 (b)), it can be written explicitly as

$$W_{\omega_3} = e^{i \sum_{\mathbf{r} \in \omega_3} a_{\mu}(\mathbf{r})} = e^{i \sum_{n=0}^{L-1} a_2(\mathbf{r} + n\mathbf{e}_2)}. \quad (17)$$

From eq.(8), it is clear that the Wilson operator W_{ω} does not commute with the 't Hooft operator $\tilde{W}_{\gamma}$ only when the "curve" ω intersects with the "curve" γ odd number of times, otherwise they commute. We consider the case that they intersect once, say, ω_3 and γ_1 in Fig.3 (b). Using eq.(8) again, one can find

$$\begin{aligned}[W_{\omega_3}, \tilde{W}_{\gamma_1}] &= -W_{\omega_3}, \\ [W_{\omega_3}^{\dagger}, \tilde{W}_{\gamma_1}] &= W_{\omega_3}^{\dagger}.\end{aligned}\quad (18)$$

Consider the eigenstate $\tilde{W}_{\gamma_1} |\tilde{w}_{\gamma_1}\rangle = \tilde{w}_{\gamma_1} |\tilde{w}_{\gamma_1}\rangle$, then $\tilde{W}_{\gamma_1} W_{\omega_3} |\tilde{w}_{\gamma_1}\rangle = (\tilde{w}_{\gamma_1} + 1) W_{\omega_3} |\tilde{w}_{\gamma_1}\rangle$. Similarly, $\tilde{W}_{\gamma_1} W_{\omega_3}^{\dagger} |\tilde{w}_{\gamma_1}\rangle = (\tilde{w}_{\gamma_1} - 1) W_{\omega_3}^{\dagger} |\tilde{w}_{\gamma_1}\rangle$. So the Wilson operators W_{ω_3} , $W_{\omega_3}^{\dagger}$ play the role of the ladder operators for the 't Hooft loop operator. Here, we can see that in the ground state, only the nonlocal operator can change the value of the winding number, hence, it is topological (we can also see the topological properties of the 't Hooft loop operators from deforming the loop, see Appendix.B).

Remarkably, eq.(18) also implies

$$e^{i\alpha \tilde{W}_{\gamma_1}} W_{\omega_3} e^{-i\alpha \tilde{W}_{\gamma_1}} = e^{i\alpha} W_{\omega_3}, \quad (19)$$

this satisfies the definition of the 1-form symmetry [16]. In this sense, $e^{i\alpha \tilde{W}_{\gamma_1}}$ is the 1-form $U(1)$ symmetry operator. The object charged under this symmetry is the Wilson loop, and the 1-form charge is the winding number.

Under this symmetry transformation, $a_{\mu}(\mathbf{r})$ transforms as

$$e^{i\alpha \tilde{W}_{\gamma_1}} a_{\mu}(\mathbf{r}) e^{-i\alpha \tilde{W}_{\gamma_1}} = a_{\mu}(\mathbf{r}) + \alpha, \quad (20)$$

if we require that α is a flat 1-form connection ($\text{curl } \alpha = 0$), then $(\text{curl } a)_{\mathbf{r}}$ is invariant under this 1-form symmetry. In this sense, the 1-form symmetry can be regarded as a symmetry: Hamiltonian (function of 1-form gauge field a) which is invariant under shifting a by a flat 1-form connection.

It is worth nothing that this 1-form symmetry remains even if we consider any higher order perturbation terms (strictly speaking, only contractible loops are allowed, since we assume the theory is local) from the spin model. Since the perturbation terms acting on the ice manifold are loop like, and the higher order terms correspond to larger loops which consist of many 4 site loops. From eq.(15), we know that any possible higher order terms in a Hamiltonian restricted by locality will commute with the 't Hooft loop, since they can only be contractible loops. We should also keep in mind that this 1-form symmetry is an emergent symmetry for the ice manifold, it will be ruined when one consider the charge.

- Fractonic symmetry [72]. Since $\tilde{W}_{\gamma_1}$ and $\tilde{W}_{\gamma_2}$ are conserved, so as $\sum_y \tilde{W}_{\gamma_1}$ and $\sum_x \tilde{W}_{\gamma_2}$. We write them as

$$\begin{aligned}P_1 &= \sum_{\mathbf{r}} E_1(\mathbf{r}), \\ P_2 &= \sum_{\mathbf{r}} E_2(\mathbf{r}),\end{aligned}\quad (21)$$

the symbol P actually means electric dipole moment [72]. This can be seen by

$$\begin{aligned}\mathbf{P} &= \sum_{\mathbf{r}} \mathbf{r} (\text{div } E)_{\mathbf{r}} \\ &= \sum_{\mathbf{r}} \mathbf{r} [E_1(\mathbf{r}) - E_1(\mathbf{r} - \mathbf{e}_1) + E_2(\mathbf{r}) - E_2(\mathbf{r} - \mathbf{e}_2)] \\ &= \sum_{\mathbf{r}} \mathbf{r} E_1(\mathbf{r}) - (\mathbf{r} + \mathbf{e}_1) E_1(\mathbf{r}) + \mathbf{r} E_2(\mathbf{r}) - (\mathbf{r} + \mathbf{e}_2) E_2(\mathbf{r}) \\ &= - \sum_{\mathbf{r}} [\mathbf{e}_1 E_1(\mathbf{r}) + \mathbf{e}_2 E_2(\mathbf{r})].\end{aligned}\quad (22)$$

This identification also tells us that $P_1 = P_2 = 0$ since $(\text{div } E)_{\mathbf{r}} = 0$ in the ice manifold. Further, the staggered magnetization $\sum_{\mathbf{r}, \mu} e^{i\mathbf{Q}_0 \cdot \mathbf{r}} S_{\mu}^z(\mathbf{r}) = P_1 + P_2$ is zero, so the ice manifold preserves the charge conjugation symmetry.

With the translation symmetry and this 1-form $U(1)$ global symmetry, now we can consider the adiabatic flux insertion process [2] where the flux corresponds to the 1-form symmetry. Suppose that we consider the translation invariant system with a finite many body gap at time $t=0$ and this gap does not close during the flux insertion process. We can choose a




FIG. 3. (a) Two noncontractible loops γ_1 and γ_2 for 't Hooft loop operators under the periodic boundary condition. (b) Different "curves" for Wilson line operators. ω_1 is an open curve with two ends. ω_2 and ω_3 are contractible and noncontractible loops, respectively. The black dots denote the link variables.

ground state $|\psi_0\rangle$ of $H(t=0)$ as initial state, which has momentum p_0 , $T_1|\psi_0\rangle = e^{ip_0}|\psi_0\rangle$. T_1 is the translation operator along $\mathbf{e}_1$. Assume we can keep the translation invariant during the flux insertion process. After the adiabatic unit flux insertion, the Hamiltonian becomes $H(\phi)$ and the initial state evolves to $|\psi'_0\rangle$ with the same momentum p_0 . It is known that the systems should be equivalent to the one after the unit flux insertion [2], this is usually written as $UH(\phi)U^{-1} = H(t=0)$, where U is a large gauge transformation. And $U|\psi'_0\rangle$ should be one of the ground states under the assumption that the system has a finite many body gap.

Now we consider the commutation relation between the translation along $\mathbf{e}_1$ direction ($T_1^\dagger E_2(\mathbf{r})T_1 = E_2(\mathbf{r}-\mathbf{e}_1)$) and the large gauge transformation $U_{\gamma_1} = \exp[-i\frac{2\pi}{L}\sum_{n=1}^L(x+n)E_2(\mathbf{r}+\mathbf{ne}_1)]$, where $\mathbf{r} = (x, y)$, we find

$$T_1^\dagger U_{\gamma_1} T_1 = U_{\gamma_1} \exp(2\pi i E_2(\mathbf{r})) \exp(-2\pi i \frac{\tilde{W}_{\gamma_1}}{L}). \quad (23)$$

Then,

$$T_1 U_{\gamma_1} |\psi'_0\rangle = \exp(-2\pi i E_2(\mathbf{r})) \exp(2\pi i \frac{\tilde{W}_{\gamma_1}}{L}) e^{ip_0} U |\psi'_0\rangle. \quad (24)$$

Recall that $E_2(\mathbf{r}) = \pm 1/2$ and $\tilde{W}_{\gamma_1} \leq L/2$, thus, if $\tilde{W}_{\gamma_1} \neq L/2$, then $U|\psi'_0\rangle$ has different momentum than p_0 and the ground state is degenerate. Above procedure leads to a LSM type constraint for the ground state, it can be stated as:

In the presence of translation symmetry and the 1-form symmetry, if the filling of the 1-form charge $\tilde{W}_{\gamma_1}/L \neq 1/2$, then the ground state can not be symmetric trivial gapped. It can either be 1). translation invariant and gapped with degeneracy, or 2). translation symmetry breaking or 3). gapless.

We also use the fact that the continuous p-form symmetry in d dimensional space-time can not be spontaneous broken if $p \geq d-2$ [15, 73]. In fact, this 1-form LSM is recently proved in ref.[11] for quantum dimer model. Hence, the low energy theory of the ice models has a mixed 't Hooft anomaly between the 1-form U(1) global symmetry and the lattice translation symmetry [16]. In addition, people recently find that the unbroken 1-form symmetry implies confinement [15, 74].

Thus the ground state is also a confined phase, we will discuss the confinement in next part.

According to the recent numerical studies for the XXZ model and quantum link model [40, 48, 53, 54], the ground state of the gauge theory eq.(13) is a translation symmetry breaking PVBS phase with $\tilde{W}_{\gamma_1} \ll L/2$, this is consistent with above LSM type constraint.

Now we consider a Zeeman field term in the spin model [41, 44–47, 61], $h \sum_i S_i^z = h \sum_i e^{i\mathbf{Q}_0 \cdot \mathbf{r}} E_\mu(\mathbf{r})$. It is obvious that this term is gauge invariant, but the translation symmetries along $\mathbf{e}_1$ and $\mathbf{e}_2$ are explicitly broken. We can still choose the translation symmetry T_d along $\mathbf{e}_1 + \mathbf{e}_2$ direction [11] ($T_d^\dagger E_i(\mathbf{r})T_d = E_i(\mathbf{r}-\mathbf{e}_1-\mathbf{e}_2)$) to repeat the flux insertion argument with the large gauge transformation $U_{\gamma_d} = \exp[-i\frac{2\pi}{L}\sum_{x=1}^L[xE_2(x, x) - xE_1(x, x+1)]]$. One can find

$$T_d^\dagger U_{\gamma_d} T_d = U_{\gamma_d} \exp(-2\pi i \frac{\tilde{W}_{\gamma_d}}{L}), \quad (25)$$

where $\tilde{W}_{\gamma_d} = \sum_{x=1}^L [E_2(x, x) - E_1(x, x+1)]$. Hence, for a suitable filling, the symmetric trivial gapped ground state can still be ruled out.

Compare to the conventional LSM theorem from the spatial symmetry and the 0-form internal symmetry, this 1-form version is not that powerful, since the filling of the 1-form charge is unknown before we find the ground state [11].

Now, we compare this IR 1-form LSM constraint with the previous UV lattice homotopy analysis. Firstly, the XXZ model triggers the LSM theorem, and the translation symmetry breaking PVBS ground state satisfies both LSMs. These results are consistent, but the IR anomaly is non-LSM nature, since we may have the same anomaly in IR even without any UV LSM.

While there is no UV LSM constraint for the other ice models (XXZ model with Zeeman field [44–47], TFIM [42, 43], TFIM with Zeeman field [41, 61] and TFIM with different nearest neighbor and next nearest neighbor interactions [43]). According to the numerical studies, the ground state of these models around the Ising limit is still a translation symmetry breaking PVBS phase even if all the spin rotation symmetries are explicitly broken. It seems that this PVBS phase (detected by translation symmetry breaking, strong plaquette correlation and flippability) is not protected by onsite spin symmetry. Further, the low energy effective models for these lower symmetry ice models are the same (constrained by LSM), and the PVBS phase seems satisfy the IR 1-form LSM. So for these models, there are also non-LSM anomaly in IR [27, 28].

C. Monopole effect in the gauge theory

To complete the analysis for the U(1) gauge theory eq.(13), we study the Monopole effect. The standard procedure is given in ref.[50]. In the first step, we introduce the fields in the dual lattice (see Fig.4)

$$E_\mu(\mathbf{r}) = \sum_{\nu} \epsilon_{\mu\nu} [\Delta_\nu^- h(\mathbf{x}) + B_\nu(\mathbf{x} - \mathbf{e}_\nu)], \quad (26)$$




FIG. 4. E field and a field are the link variables in the black square lattice, while h field is the site variable and B field is link variable in the dual lattice (green dashed lattice). For a given site $\mathbf{r}$ in the square lattice, the corresponding dual site is $\mathbf{x}$.

then the Gauss's law eq.(9) is rewritten as

$$\begin{aligned} (\text{div } E)_{\mathbf{r}} &= \sum_{\mu, \nu} [\epsilon_{\mu\nu} \Delta_{\mu}^{-} \Delta_{\nu}^{-} h(\mathbf{x}) + \epsilon_{\mu\nu} \Delta_{\mu}^{-} B_{\nu}(\mathbf{x} - \mathbf{e}_{\nu})] \\ &= (\text{curl } B)_{\mathbf{x} - \mathbf{e}_1 - \mathbf{e}_2} \\ &= 0, \end{aligned} \quad (27)$$

where we use $\sum_{\mu, \nu} \epsilon_{\mu\nu} \Delta_{\mu}^{-} \Delta_{\nu}^{-} h(\mathbf{x}) \equiv 0$. Once we choose a reference site $h(\mathbf{x}_0) = 0$, then eq.(26) is actually a rule for determining all other $h(\mathbf{x})$ on the dual lattice if a classical E field configuration is given (E field configuration is defined by eq.(7) from a classical spin configuration). Let us denote the site $\mathbf{r} = (x, y)$ with (even, even) or (odd, odd) coordinate as A sublattice, and (even, odd) or (odd, even) as B sublattice. Then the rule for determining $h(\mathbf{x})$ can be stated as: run around the A (B) sublattice site clockwise (anticlockwise), $h(\mathbf{x})$ decreases by $-1/2$ if one crosses a spin up, while crossing a spin down, $h(\mathbf{x})$ increases by $1/2$. This is indeed a height rule, and $h(\mathbf{x})$ is a height field [64, 75]. It is clear that $B_{\nu}(\mathbf{x}) \equiv 0$ once a reference site $h(\mathbf{x}_0) = 0$ is chosen, this is equivalent to choose a gauge for B field [64]. In addition, the average height over 4 plaquette need to be compactified to avoid overcounting [54, 64], we identify the average height h with $h+1$, this can be seen by choosing different reference site.

Now we define the canonical conjugate operator $P(\mathbf{x})$ for the height field $h(\mathbf{x})$: $[P(\mathbf{x}), h(\mathbf{x}')] = i\delta_{\mathbf{x}, \mathbf{x}'}$. Analogously to eq.(8), $e^{\pm i P(\mathbf{x})}$ is the ladder operator for the height field which shifts the value of $h(\mathbf{x})$ by ± 1 . While in the original theory eq.(13), $\cos[(\text{curl } a)_{\mathbf{r}}]$ flips the flippable spin configuration in the plaquette, see Fig.5, which also changes the value of $h(\mathbf{x})$ by ± 1 , thus we identify $\cos[(\text{curl } a)_{\mathbf{r}}] = \cos[P(\mathbf{x})]$. We can now rewrite the theory eq.(13) in terms of the variables living in the dual lattice

$$H^{\text{cQED}} = \frac{1}{2g} \sum_{\mathbf{x}, \nu} [(\Delta_{\nu}^{-} h(\mathbf{x}))^2 - \frac{1}{4}] - 2t \sum_{\mathbf{x}} \cos[P(\mathbf{x})], \quad (28)$$

the theory is invariant under a global shift of the height field $h(\mathbf{x}) \rightarrow h(\mathbf{x}) + c$.

The monopole effect can be studied by path integral. The partition function is $Z = \text{Tr } e^{-\beta H^{\text{cQED}}}$, where β is the imaginary time. Divide the imaginary time into N_{τ} parts, each part is Δ_{τ} : $N_{\tau} \Delta_{\tau} = \beta$. Choosing the eigenbasis $|\{h(\mathbf{x}, \tau_j)\}\rangle$ of the height

field operator, where $\tau_j = j\Delta_{\tau}$. Then the partition function can be written as

$$Z = \lim_{\substack{\Delta_{\tau} \rightarrow 0, \\ N_{\tau} \rightarrow \infty}} \sum_{\{h(\mathbf{x}, \tau_j)\}} \prod_{j=1}^{N_{\tau}} \langle \{h(\mathbf{r}, \tau_j)\} | e^{-\Delta_{\tau} H^{\text{cQED}}} | \{h(\mathbf{r}, \tau_{j+1})\} \rangle, \quad (29)$$

the first term in eq.(28) is diagonal in this basis, thus the corresponding matrix element can be written directly. While second term is off-diagonal, there are several methods to calculate its matrix element [50, 76], all of them give the same form $\exp[-f(t\Delta_{\tau})(\Delta_0^{-} h(\mathbf{x}, \tau_j))^2]$, where $f(t\Delta_{\tau})$ is a function of $t\Delta_{\tau}$, $\Delta_0^{-} h(\mathbf{x}, \tau_j) = h(\mathbf{x}, \tau_j) - h(\mathbf{x}, \tau_{j-1})$. Now, the partition function is

$$Z = \lim_{\substack{\Delta_{\tau} \rightarrow 0, \\ N_{\tau} \rightarrow \infty}} \sum_{\{h(\mathbf{x}, \tau_j)\}} e^{-S_{\text{eff}}[h]}, \quad (30)$$

with the action

$$S_{\text{eff}}[h] = f(t\Delta_{\tau}) \sum_{\mathbf{x}, j} (\Delta_0^{-} h(\mathbf{x}, \tau_j))^2 + \frac{\Delta_{\tau}}{2g} \sum_{\mathbf{x}, \nu, j} [(\Delta_{\nu}^{-} h(\mathbf{x}, \tau_j))^2 - \frac{1}{4}]. \quad (31)$$

In the following, we will denote the "space-time" $(\mathbf{x}, \tau)$ by x , and $x + \mathbf{e}_0 = (\mathbf{x}, \tau + \Delta_{\tau})$. From the height rule, we know that $h(x) = Z/2$, $Z \in \text{integer}$. It is convenient to define a integer value field $H(x) = 2h(x)$. Then we can use Poisson summation to lift the discrete field $H(x)$ into a continuum field $\varphi(x)$ with the price that introducing another discrete field $m(x)$, the partition function now can be written as

$$\begin{aligned} Z &= \lim_{\substack{\Delta_{\tau} \rightarrow 0, \\ N_{\tau} \rightarrow \infty}} \sum_{\{m(x)\}} \int D\varphi e^{i2\pi \sum_x m(x)\phi(x) - S_{\text{eff}}[\varphi]} \\ &= \lim_{\substack{\Delta_{\tau} \rightarrow 0, \\ N_{\tau} \rightarrow \infty}} \sum_{\{m(x)\}} \int D\varphi e^{-S_{\text{eff}}[\varphi, m]} \end{aligned} \quad (32)$$




FIG. 5. The rule for determining h field. (a) Assume the site $\mathbf{r}$ belongs to the A sublattice and suppose the height value is $h(\mathbf{x})$ for the upper right dual site $\mathbf{x}$. Then run around the $\mathbf{r}$ site clockwise, $h(\mathbf{x})$ decreases by $-1/2$ if one crosses a spin up, while crossing a spin down, $h(\mathbf{x})$ increases by $1/2$. (b) is obtained from (a) after flipping the upper right plaquette. It is clear that flip a plaquette will shift the height value by ± 1 .

where

$$S_{\text{eff}}[\varphi, m] = \frac{1}{4} f(t\Delta_\tau) \sum_x (\Delta_0^- \varphi(x))^2 + \frac{\Delta_\tau}{8g} \sum_{x,y} [(\Delta_y^- \varphi(x))^2 - \frac{1}{4}] - i2\pi \sum_x m(x)\varphi(x). \quad (33)$$

It is known that m is the monopole field [50, 67, 77]. Since the theory is a Gaussian action, we can integrate out φ field (see the Appendix).

Now, we consider the consequence of the monopole effect. Under the dilute monopole gas approximation [66, 78, 79], m field can be "integrated out", the remained action has been worked out in ref.[51, 79]

$$S'_{\text{eff}}[\varphi] = \sum_x \frac{1}{4} f(t\Delta_\tau) [(\Delta_0^- \varphi_A)^2 + (\Delta_0^- \varphi_B)^2] + \frac{\Delta_\tau}{8g} (\varphi_A - \varphi_B)^2 - z \cos(2\pi\varphi_A) + z \cos(2\pi\varphi_B). \quad (34)$$

where z is the fugacity for the monopole gas, A,B denote the sublattice of the dual lattice since it is bipartite. The continuum limit of eq.(34) can be found by introducing average height $\chi_1 = (\varphi_A + \varphi_B)/2$, difference in height $\chi_2 = (\varphi_A - \varphi_B)/2$ and expanding the action in terms of the derivatives of these fields. It is clear that χ_2 is massive from eq.(34), integrating it out and rescale the coefficients, a sine-Gordon action is found [51, 79]

$$S_{\text{sg}} = \int d^2\mathbf{r} d\tau \left[\frac{K}{2} (\partial_\mu \chi_1)^2 - \gamma \cos(4\pi\chi_1) \right]. \quad (35)$$

In this dual theory, the global $U(1)$ topological symmetry is usually expressed as the theory which is invariant under the global shift [19, 50, 69] $\chi \rightarrow \chi + c$. It is clear that this topological symmetry is ruined by the cosine term, and it is reduced to a Z_2 symmetry: $Z_{2,\text{top}} : \chi \rightarrow \chi + \frac{1}{2}$. Such topological symmetries is known to be related to the lattice symmetry [32, 79]. In addition, the charge conjugation symmetry is $C : \chi \rightarrow -\chi$.

There are two minima in this field theory which implies the 2-fold degenerate ground states. It is known that the cosine term is marginally relevant [51] which pins the χ_1 field to one of the minima. This is actually the Polyakov confinement mechanism [65]. Eventually, the ground state breaks the $Z_{2,\text{top}}$ symmetry (corresponds to the translation symmetry [79]) and preserves the charge conjugation symmetry. From this point of view, the ice models studied here should have lattice symmetry broken ground states around the Ising limit and the symmetric trivial gapped ground state is ruled out if the perturbation preserves the lattice symmetry. This is consistent with the 1-form LSM theorem. And here the consequence of the monopole effect in the dual theory may imply the filling of the 1-form charge always triggering the 1-form LSM theorem and lead to a translation symmetry breaking ground state. So far, we find that this PVBS phase can be understood from the mixed 't Hooft anomaly (1-form LSM) and also from the monopole effect in the dual theory, both of them lead to a translation symmetry breaking phase.

Further, recognizing $e^{\pm i2\pi\chi}$ as the standard monopole creation/annihilation operators, then, this sine-Gordon theory

tells us that the effective charge of the monopoles is ± 2 [51, 79]. In addition, according to the Polyakov's pioneering work [65], this sine-Gordon theory also implies the charges are always confined [50, 51]. However, this well known charge confinement is not the whole story. It is recently found that such sine-Gordon theory with even monopole charge is an effective description for Z_2 VBS which can be confined in the bulk but supports deconfined excitations on the domain wall [19, 37]. We will study such excitations in the following section.

Usually, the compact gauge field can be divided into a continuous part and a singular part [32, 80, 81], like the spin wave and vortex in the XY model [82]. Above analysis of the monopole tells us that the continuum limit of the theory eq.(13) can be written as

$$-\frac{1}{4e^2} |da|^2 + \mathcal{L}_M, \quad (36)$$

where the first term is the Maxwell term, $\mathcal{L}_M$ is the monopole term.

IV. EXCITATIONS AND ANOMALY

As we mentioned before, eq.(5) actually induces a dimer representation for the ice problem. This can be stated as: put a dimer on the link if there is a spin up on that link, and spin down corresponds to empty. Ice rule then is expressed as: there are two and only two dimers for each site. A dimer configuration for PVBS phase is shown in Fig.6 (a). The kinetic term eq.(4) will flip the dimer configuration in a plaquette between the vertical one and horizontal one $\parallel \leftrightarrow \perp$. This dimer representation is useful when we discuss the excitation out of the ice manifold.

Now we consider the charge excitations. The basis operation in the XXZ model is flipping a pair of spins, which corresponds to move a dimer belonging to a site to the empty position belonging to the same site, as in Fig.6 (b),(c). And such process may create a pair of charges on the different sites belonging to the same sublattice. According to our definition of charge eq.(10), we can check that such a pair of charges carry opposite charge. Moreover, the bond flipping process may also hops the charge in the same sublattice. In contrast, for the transverse field case, the basis operation is flipping a spin, which corresponds to create or remove a dimer. Then this process may creates a pair of charges carry opposite charge locating on the different sublattices, see Fig.6 (d). It is clear that in a single flipping process one can only hops the charge ± 1 excitation, charge ± 2 will split into charges ± 1 . To hop the charges ± 2 , one needs the higher order process, and they require higher energy, thus we will neglect these higher charge excitations.

Based on these observations, we can slightly modify the definition eq.(5) to include the charge

$$S_\mu^\pm(\mathbf{r}) = \exp \left[\pm i e^{i\mathbf{Q}_0 \cdot \mathbf{r}} (a_\mu(\mathbf{r}) + \Theta_{\mathbf{r}} - \Theta_{\mathbf{r}+\mathbf{e}_\mu}) \right], \quad (37)$$

where $\exp(\pm i\Theta)$ describe the charge (matter) field Φ . $\exp(i\Theta_{\mathbf{r}})$ will increase the charge by 1 on site $\mathbf{r}$, and $\exp(-i\Theta_{\mathbf{r}'})$ will




FIG. 6. (a) Dimer representation for the PVBS phase. The red bar denotes the dimer. (b),(c) A pair of charge excitations with opposite charge on the same sublattice can be created by moving the dimer. (d) A pair of charge excitations with opposite charge on the different sublattices can be created by creating or removing the dimer. The blue and gray dots denote the charges on the different sublattices, the filled and empty dots denote the opposite charges.

decrease the charge by 1 on site $\mathbf{r}'$. Since there are two sublattices in our case, this fact leads to two flavors of the field. Further, we introduce a number operator $N_{\mathbf{r}}$ whose canonical conjugated operator is $\Theta_{\mathbf{r}}$. Then, $e^{\pm i\Theta_{\mathbf{r}}}$ are the ladder operators for $N_{\mathbf{r}}$. In the ice manifold, $N_{\mathbf{r}} = 0$ for every site.

Obviously, eq.(37) describes the minimal coupling between the matter field Φ and the gauge field a . And we know that create a charge requires the energy around $J_z/2$, which implies the fields are massive. We may also add interactions for the fields like Bose-Hubbard model. Then a simplest theory to include all these effects is

$$\mathcal{L} = -\frac{1}{4e^2}|da|^2 + \sum_{i=1}^2 [|D_a \Phi_i|^2 + m^2 |\Phi_i|^2 + u |\Phi_i|^4] + w |\Phi_1|^2 |\Phi_2|^2 + \mathcal{L}_M, \quad (38)$$

where D_a is the covariant derivative. This is exactly the same abelian Higgs model which is studied in ref.[19]. The global symmetries of this theory are: topological symmetry $Z_{2,top}$, charge conjugation symmetry C (now also acts on the matter), $U(1)$ flavor symmetry and the Z_2 exchange symmetry $\Phi_1 \leftrightarrow \Phi_2$. It is found that there is a 't Hooft anomaly among these symmetries. The anomaly can be seen by turning on the 1-form background gauge fields [19]: A couples to the topological symmetry, and B couples to the flavor symmetry. Then we need to add a term $A \wedge \frac{da}{2\pi} + A \wedge \frac{dB}{2\pi}$ and replace D_a by D_{a+B} in the theory eq.(38). In the presence of the Z_2 exchange symmetry (now also acts on B , $Z_2 : B \rightarrow -B$), we can not gauge the topological symmetry and flavor symmetry consistently,

hence, anomaly. And we can also see that the charge conjugation symmetry is never broken in the bulk, hence the bulk can carry the anomaly among these 4 symmetries.

An interesting prediction from ref.[19] is that the domain wall interpolating between different vacuums (different Z_2 PVBS) in this theory will also carry an anomaly and there are deconfined excitations on the domain wall. For the domain wall, $Z_{2,top}$ symmetry is restored, while the C symmetry is broken, hence the domain wall carries anomaly among the charge conjugation symmetry, Z_2 exchange symmetry and the flavor symmetry.

As mentioned previously, the flippable plaquette repels each other, and each flippable plaquette can be located on one of the two sublattices. Thus, one can find two simplest domain wall, see Fig.7 (a),(d), we call them staircase type and ladder type, respectively. Actually, the staircase type domain wall (Fig.7 (a),(c)) is "anomalous", the charges move freely on the domain wall by rotating a dimer $\pi/2$ degree (see Fig.7 (b)). And the movement of the charges on the domain wall does not affect the bulk. Now go back to the spin representation, then such a "anomalous" domain wall configuration is just the antiferromagnetic ordered spin chain (staircase like), thus it breaks the Z_2 symmetry around the spin x axis (which corresponds to charge conjugation symmetry) and supports the deconfined excitations. However, in the ladder type domain wall (Fig.7 (d)-(f)), the movement of charges will induce the fluctuation of the gauge field, and they will frustrate each other. Even worse, the movement of charges will affect the bulk. Hence, only the stair case type domain wall is stable. We also note that it is possible to discuss the domain wall from the sine-Gordon theory eq.(35) with suitable boundary condition [83, 84], and it is left for future work.

A. Spectrum

Now we study the spectrum (dynamical structure factor) $S^{zz}(\mathbf{Q}, \omega) = \int \frac{dt}{2\pi} e^{i\omega t} \langle S_{\mathbf{Q}}^z(t) S_{-\mathbf{Q}}^z(0) \rangle$, which is closely related to our recent work [55]. Here $S_{\mathbf{Q}}^z = \sum_{\alpha} S_{\mathbf{Q},\alpha}^z$, and $S_{\mathbf{Q},\alpha}^z = \frac{1}{N} \sum_i e^{-i\mathbf{Q} \cdot \mathbf{r}_{i,\alpha}} S_{i,\alpha}^z$, $\mathbf{r}_{i,\alpha} = \mathbf{R}_i + \rho_{\alpha}$. i labels the unitcell and $\alpha = 1, 2$ labels the sublattice. ρ_{α} is the position of the α spin in a unitcell, which are read as $\rho_1 = \frac{1}{2}e_1$, $\rho_2 = \frac{1}{2}e_2$. The expectation value is calculated in the ground state $|\psi_{spin}\rangle$ of the spin Hamiltonian, and the transferred momentum is $\mathbf{Q} = (q_1, q_2)$.

1. $q_2 = \pm q_1$ paths

Firstly, we study the $q_2 = q_1 = q$ path.

$$\begin{aligned} S_{-\mathbf{Q}}^z &= \frac{1}{N} \sum_{\mathbf{r}, \mu} e^{i\mathbf{Q} \cdot \mathbf{r}} S_{\mu}^z(\mathbf{r}) = \frac{1}{N} \sum_{\mathbf{r}, \mu} e^{i(\pi+q)(x+y)} e^{i\mathbf{Q} \cdot \rho_{\mu}} E_{\mu}(\mathbf{r}) \\ &= \frac{1}{N} \sum_{n=0}^{L-1} \sum_{x=0}^{L-1} e^{i(\pi+q)n} e^{iq/2} (E_1(x, -x+n) + E_2(x, -x+n)), \end{aligned} \quad (39)$$

where $\mathbf{r} = (x, y)$, and we sum over the $E_{\mu}(\mathbf{r})$ on different "curves" ($y = -x+n$, n takes integer), see the "curves" in dif-




FIG. 7. The simplest domainwalls interpolate between different PVBS configurations. (a),(b)(c), staircase type domain wall. This staircase one is actually the antiferromagnetic ordered spin chain which breaks the Z_2 symmetry around the spin x axis and supports the deconfined charge excitations on the domain wall. (d)-(f), ladder type domain wall, the movement of the charge will create the flip-pable dimers then frustrating itself.

ferent colors in Fig.8. Using the canonical relation between a and E , it is not difficult to find

$$[(\text{curl } a)_r, \sum_{x=0}^{L-1} (E_1(x, -x+n) + E_2(x, -x+n))] = 0, \quad (40)$$

thus $\sum_{x=0}^{L-1} (E_1(x, -x+n) + E_2(x, -x+n))$ is a conserved 1-form charge $\tilde{W}_n$. So eq.(39) is actually conserved in the gauge theory eq.(13). Let us denote the ground state of the gauge theory eq.(13) as $|\psi_{cQED}\rangle$. In the low energy (ice manifold), $|\psi_{spin}\rangle$ can be approximated by $|\psi_{cQED}\rangle$, then,

$$\begin{aligned} \langle S_{\mathbf{Q}}^z(t) S_{-\mathbf{Q}}^z(0) \rangle &= \langle \psi_{spin} | S_{\mathbf{Q}}^z(t) S_{-\mathbf{Q}}^z(0) | \psi_{spin} \rangle \\ &\approx \langle \psi_{cQED} | e^{iH^{cQED}t} \tilde{W}(q) e^{-iH^{cQED}t} \tilde{W}(-q) | \psi_{cQED} \rangle \\ &= c, \end{aligned} \quad (41)$$

where $\tilde{W}(-q) = \frac{1}{N} \sum_{n=0}^{L-1} \sum_{x=0}^{L-1} e^{i(\pi+q)n} \tilde{W}_n$, c does not depend on t . So there is no spectral weight in $S^{zz}(\mathbf{Q}, \omega)$ in the low energy sector along $q_2 = q_1$. Similarly, we can also find the same conclusion along $q_2 = -q_1$.




FIG. 8. Decompose the summation over $E_{\mu}(\mathbf{r})$ on the lattice with periodic boundary condition.

Specially, there are two high symmetry points in this path, i.e. $(0,0)$ and (π, π) .

a). For $(0,0)$ momentum, $S_{-\mathbf{Q}}^z = \frac{1}{N} \sum_{\mathbf{r}, \mu} S_{\mu}^z(\mathbf{r})$ is conserved in the XXZ model, so there is no spectral weight. In other models, for example, TFIM, $\sum_{\mathbf{r}, \mu} S_{\mu}^z(\mathbf{r})$ is not conserved in the spin model, so we only expect there is vanishing spectral weight in the low energy sector.

b). For (π, π) momentum, $S_{-\mathbf{Q}}^z = \frac{1}{N} \sum_{\mathbf{r}, \mu} e^{i\pi/2} E_{\mu}(\mathbf{r})$ is related to the fractonic symmetry, we expect that there is no spectral weight in the low energy sector. And the ice models considered in this paper do not have these fractonic symmetries (conserve staggered magnetization) in the lattice model level.

2. domain wall

Actually, the decomposition used in eq.(39) has a clear physical meaning, which is expanding the lattice in the patterns of domain wall. See Fig.8 and Fig.7, one can immediately recognize that the dynamical structure factor $S^{zz}(\mathbf{Q}, \omega)$ along $q_2 = \pm q_1$ directly measures the excitations in the domain wall. Hence, we expect the spectrum along the $q_2 = \pm q_1$ path is just the longitudinal dynamical structure factor of the spin-1/2 chain in the antiferromagnetic phase, which can be solved by the Bethe ansatz [85].

3. $q_i = \pi$ paths

Next, we consider $q_1 = \pi$ path. $\mathbf{Q} = (\pi, q)$,

$$\begin{aligned} S_{-\mathbf{Q}}^z &= \frac{1}{N} \sum_{\mathbf{r}, \mu} e^{i\pi x + iqy} e^{i\mathbf{Q} \cdot \mathbf{r}} S_{\mu}^z(\mathbf{r}) = \frac{1}{N} \sum_{\mathbf{r}, \mu} e^{i(\pi+q)y} e^{i\mathbf{Q} \cdot \mathbf{r}} E_{\mu}(\mathbf{r}) \\ &= \frac{1}{N} \sum_y e^{i(\pi+q)y} (e^{iq/2} \tilde{W}_y + e^{i\pi/2} \sum_x E_1(\mathbf{r})), \end{aligned} \quad (42)$$

which is not a fully conserved term even in the low energy sector. Hence $\langle \psi_{spin} | S_{\mathbf{Q}}^z(t) S_{-\mathbf{Q}}^z(0) | \psi_{spin} \rangle$ will be in general time dependent, and the energy scale is determined by the kinetic term $\propto t$. And we expect the spectrum will be almost flat

with energy scale t . The high symmetry points in this path are $(\pi, 0)$ and (π, π) . Since there is no spectral weight in (π, π) in the low energy, so we naively expect the spectral weight decrease from $(\pi, 0)$ to (π, π) .

4. $q_i = 0$ paths

Lastly, we consider $q_2 = 0$ path. $\mathbf{Q} = (q, 0)$

$$S_{-\mathbf{Q}}^z = \frac{1}{N} \sum_{\mathbf{r}, \mu} e^{iqx} e^{i\mathbf{Q} \cdot \rho_\mu} S_\mu^z(\mathbf{r}) = \frac{1}{N} \sum_{\mathbf{r}, \mu} e^{i(\pi+q)x} e^{i\pi y} e^{i\mathbf{Q} \cdot \rho_\mu} E_\mu(\mathbf{r}), \quad (43)$$

this is not related to the conserved quantity for general q . Combine with above conclusions, we expect the spectrum will be almost flat with energy scale t and increasing spectral weight from $(0, 0)$ to $(\pi, 0)$.

All of the expectations above are consistent with our recent numerical work on the XXZ model [55], especially the one dimensional like spectrum along the $q_2 = \pm q_1$ paths. Here we also prove that there is no spectral weight from the low energy sector along that path, thus it is possible to extract more accurate one dimensional informations (for example, the effective Ising anisotropy and something else) from the improved numerical study.

V. CONCLUSIONS

In this paper, we have discussed the i).constraints for the ground state, ii).excitations and spectrum of different quantum spin ice models in the planar pyrochlore lattice.

For i), we begin with the lattice homotopy method and find the necessary spin rotation symmetries ($Z_2 \times Z_2$, $U(1) \rtimes Z_2$) to trigger the LSM theorem. The models with these symmetries will not have the symmetric trivial gapped phase. But the LSM theorem does not answer why the ground states of the lower symmetry models (XXZ model with Zeeman field, TFIM, TFIM with Zeeman field, TFIM with different nearest neighbor and next nearest neighbor interactions) are still translation symmetry breaking VBS even if the perturbation completely breaks all the spin rotation symmetry. Using the fact the spin ice models have a local conservation of z component of spin, we map the spin ice model to a compact $U(1)$ lattice gauge theory. There are some global symmetries in this gauge theory, the translation symmetry and the 1-form $U(1)$ global symmetry lead to a 1-form LSM theorem which rules out the symmetric trivial gapped phase if the filling of the 1-form charge satisfies some conditions. We find the results from the numerical studies on these ice models are consistent with this 1-form LSM theorem. We also study the dual theory of this gauge theory and find that it is indeed a sine-Gordon theory with charge ± 2 monopoles. The same as the Polyakov's mechanism, the monopole term is marginally relevant and leads to a lattice symmetry breaking phase. Thus, the "persistent" translation symmetry breaking VBS phases in these spin ices models can be understood from both the emergent IR anomaly in the compact $U(1)$ lattice gauge theory

and Polyakov's mechanism in the dual theory description. In short, the Ising model with lattice symmetry preserving perturbations on the planar pyrochlore lattice will have translation symmetry breaking VBS ground states. This is consistent with the existed numerical studies.

For ii), we study the charge excitation by the dimer representation of the planar pyrochlore spin ice. We find the charge excitation can be described by the abelian Higgs model with charge ± 2 monopoles. It is recently found that there is a mixed 't Hooft anomaly in this model. Due to the anomaly, the domain wall in the PVBS phase can support deconfined excitations on it, while the bulk excitations are confined. And we explicitly construct such a domain wall and find it is indeed an antiferromagnetic spin chain which is known to have deconfined excitations. In the last part, we study the spectrum. We find the symmetries in the gauge theory strongly restrict the spectral weight in the low energy sector along the high symmetry momentum paths. Remarkably, we show that there are only domain wall contributions along the $q_2 = \pm q_1$ momentum paths. From such analysis, almost all the features of the spectrum calculated in our recent numerical work [55] can be understood by the symmetry. Finally, we find that the spin ice model on the planar pyrochlore lattice can be realized by superconducting qubits [86] and Rydberg atom arrays [40, 87], we hope the analysis of the excitation and ground state will be helpful for experiments.

ACKNOWLEDGMENTS

We thank Linhao Li, Haruki Watanabe, Yohei Fuji, Ryohei Kobayashi and Takamasa Ando for helpful discussion. This work is supported by the International Postdoctoral Exchange Fellowship Program 2022 by the Office of China Postdoctoral Council: No.PC2022072 and the National Natural Science Foundation of China: No.12147172.

Appendix A: Integrate out φ

We can also integrate out φ in eq.(32) rather than m . This can be done as follow, in eq.(32)

$$\begin{aligned} S_{\text{eff}}[\varphi] &= \frac{1}{4} f(t\Delta_\tau) \sum_x (\Delta_0^- \varphi(x))^2 + \frac{\Delta_\tau}{8g} \sum_{x,v} [(\Delta_v^- \varphi(x))^2 - \frac{1}{4}] \\ &= \frac{1}{4} f(t\Delta_\tau) \sum_x 2\varphi(x) \Delta_0^- \varphi(x) + \frac{\Delta_\tau}{8g} \sum_{x,v} [2\varphi(x) \Delta_v^- \varphi(x) - \frac{1}{4}], \end{aligned} \quad (A1)$$

where we have use the discrete version of the "integration by parts":

$$\begin{aligned} \sum_x (\Delta_\mu^- \varphi)^2 &= \sum_x \varphi^2(x) + \varphi^2(x - \mathbf{e}_\mu) - 2\varphi(x)\varphi(x - \mathbf{e}_\mu) \\ &= 2 \sum_x \varphi(x) \Delta_\mu^- \varphi(x), \end{aligned} \quad (A2)$$

where $\mu = 0, 1, 2$. In the continuous case, we usually use $\int (\partial_\mu \varphi)^2 = \varphi \partial_\mu \varphi - \int \varphi \partial_\mu^2 \varphi$, which corresponds to

$\sum_x (\Delta_\mu^+ \varphi(x)) (\Delta_\mu^- \varphi(x)) = -\sum_x \varphi(x) \Delta_\mu^2 \varphi(x)$. Since $S_{\text{eff}}[\varphi]$ is a Gaussian action, we can integrate out φ ,

$$Z \propto \lim_{\substack{\Delta_\tau \rightarrow 0, \\ N_\tau \rightarrow \infty}} \sum_{\{m(x)\}} [\det A]^{-1/2} \exp^{-\frac{1}{2} \sum_{x,y} (2\pi)^2 m(x) A^{-1}(x,y) m(y)}, \quad (\text{A3})$$

where $A = f(t\Delta_\tau)\Delta_0^- + \frac{\Delta_\tau}{2g} \sum_\nu \Delta_\nu^-$, it is slightly different from ref.[50].

Appendix B: Deformation of the 't Hooft loop operator

The noncontractible loops in the 't Hooft loop operators eq.(16) are parallel to $\mathbf{e}_1$ and $\mathbf{e}_2$, respectively. Actually, we can also choose different loops, some of them are used in the main context. The 't Hooft loop operator is the summation of E fields on the chosen loop which are orthogonal to the loop direction, see Fig.9, E fields are denoted by the red arrows. To commute with the Hamiltonian, every plaquette has two arrows, and one of the arrow is clockwise and another is counterclockwise. If we regard the loop as a surface, then all of the arrows are normal vectors.

We can deform the 't Hooft loop operator by the lattice divergence $(\text{div } E)_r$ (which is zero everywhere in the ground state manifold), for example, in Fig.9, $\tilde{W}_{\gamma_1} + (\text{div } E)_r = \tilde{W}_{\gamma_3}$. With this property, we can find that the eigenvalue of this loop operator (winding number) is unchanged from the deformation. This is consistent with the higher form symmetry [15]: the q-form (q=1) symmetry in d-dimensions (d=2+1 here) is supported in a closed manifold M^{d-q-1} (M^1), and the manifold M^{d-q-1} can be deformed slightly without changing the physical results. And there are two independent winding numbers (which correspond to two inequivalent noncontractible loops on the torus) for each eigenstate.

Appendix C: Charge conjugation symmetry

In the main context, we recognize the Z_2^x symmetry (π rotation around the x axis) as the charge conjugation symmetry for the XXZ type model. Here, we discuss the charge conjugation in the ice manifold. First, we consider the Zeeman field

effect in the Ising model,

$$H^0 = J_z \sum_{\langle i,j \rangle} S_i^z S_j^z - h \sum_i S_i^z, \quad (\text{C1})$$

this can be recasted into

$$H^0 = \frac{J_z}{2} \sum_{\boxtimes} [(S_{\boxtimes}^z) - \frac{h}{2J_z}]^2 + C, \quad (\text{C2})$$

for a certain range of h , the ground state still satisfies the ice rule $S_{\boxtimes}^z = 0$. And the global constraint still works, in other words, $\sum_{\mathbf{r},\mu} S_\mu^z(\mathbf{r}) = \sum_{\mathbf{r},\mu} (n_\mu(\mathbf{r}) - \frac{1}{2}) = \sum_{\mathbf{r},\mu} e^{i\mathbf{Q}_0 \cdot \mathbf{r}} E_\mu(\mathbf{r}) = c$ in the low energy manifold. This implies that this translation preserving UV perturbation plays no role in the low energy theory eq.(13). Further, we can add a term in eq.(13) to impose the Gauss's law explicitly, $\frac{J_z}{2} \sum_{\mathbf{r}} ((\text{div } E)_r)^2$. And the low energy theory still has the symmetry $E \rightarrow -E$ and $a \rightarrow -a$. Here we can see the charge conjugation is an emergent symmetry in the low energy theory, even the UV Hamiltonian breaks the Z_2^x symmetry.




FIG. 9. Deform the 't Hooft loop operator by the lattice divergence (which is denoted by the blue dot in (a)). We can regard the loop γ_1 (γ_3) as a surface, then the E fields (which are represented by the red arrows) in the loop operator are normal vectors.

-
- [1] E. Lieb, T. Schultz, and D. Mattis, *Annals of Physics* **16**, 407 (1961).
- [2] M. Oshikawa, *Phys. Rev. Lett.* **84**, 1535 (2000).
- [3] M. B. Hastings, *Phys. Rev. B* **69**, 104431 (2004).
- [4] L. Balents, *Nature* **464**, 199 (2010).
- [5] L. Savary and L. Balents, *Reports on Progress in Physics* **80**, 016502 (2016).
- [6] H. Watanabe, H. C. Po, A. Vishwanath, and M. Zaletel, *Proceedings of the National Academy of Sciences* **112**, 14551 (2015).
- [7] H. C. Po, H. Watanabe, C.-M. Jian, and M. P. Zaletel, *Phys. Rev. Lett.* **119**, 127202 (2017).
- [8] D. V. Else and R. Thorngren, *Phys. Rev. B* **101**, 224437 (2020).
- [9] Y. Yao and A. Furusaki, *Phys. Rev. B* **106**, 045125 (2022).
- [10] Y. Yao and M. Oshikawa, *Phys. Rev. Lett.* **126**, 217201 (2021).
- [11] R. Kobayashi, K. Shiozaki, Y. Kikuchi, and S. Ryu, *Phys. Rev. B* **99**, 014402 (2019).
- [12] H. He, Y. You, and A. Prem, *Phys. Rev. B* **101**, 165145 (2020).
- [13] O. Dubinkin, J. May-Mann, and T. L. Hughes, *Phys. Rev. B* **103**, 125133 (2021).
- [14] O. M. Aksoy, A. Tiwari, and C. Mudry, *Phys. Rev. B* **104**, 075146 (2021).
- [15] D. Gaiotto, A. Kapustin, N. Seiberg, and B. Willett, *Journal of High Energy Physics* **2015**, 172 (2015).
- [16] J. McGreevy, *Annual Review of Condensed Matter Physics* **14**, null (2023).
- [17] L. Li and Y. Yao, *Phys. Rev. B* **106**, 224420 (2022).
- [18] G. Hooft, Recent developments in gauge theories, 135 (1980).
- [19] Z. Komargodski, T. Sulejmanpasic, and M. Ünsal, *Phys. Rev. B* **97**, 054418 (2018).
- [20] Z. Komargodski, A. Sharon, R. Thorngren, and X. Zhou, *SciPost Phys.* **6**, 003 (2019).
- [21] C. Cordova, D. S. Freed, H. T. Lam, and N. Seiberg, *SciPost Phys.* **8**, 001 (2020).
- [22] M. Cheng, M. Zaletel, M. Barkeshli, A. Vishwanath, and P. Bonderson, *Phys. Rev. X* **6**, 041068 (2016).
- [23] X.-G. Wen, *Phys. Rev. D* **88**, 045013 (2013).
- [24] X.-G. Wen, *Phys. Rev. B* **91**, 205101 (2015).
- [25] G. Y. Cho, C.-T. Hsieh, and S. Ryu, *Phys. Rev. B* **96**, 195105 (2017).
- [26] C.-M. Jian, Z. Bi, and C. Xu, *Phys. Rev. B* **97**, 054412 (2018).
- [27] M. A. Metlitski and R. Thorngren, *Phys. Rev. B* **98**, 085140 (2018).
- [28] W. Ye, M. Guo, Y.-C. He, C. Wang, and L. Zou, *SciPost Phys.* **13**, 066 (2022).
- [29] M. Cheng and N. Seiberg, arXiv preprint arXiv:2211.12543 (2022).
- [30] T. Sulejmanpasic and Y. Tanizaki, *Phys. Rev. B* **97**, 144201 (2018).
- [31] T. Senthil, A. Vishwanath, L. Balents, S. Sachdev, and M. P. A. Fisher, *Science* **303**, 1490 (2004).
- [32] T. Senthil, L. Balents, S. Sachdev, A. Vishwanath, and M. P. A. Fisher, *Phys. Rev. B* **70**, 144407 (2004).
- [33] C. Wang, A. Nahum, M. A. Metlitski, C. Xu, and T. Senthil, *Phys. Rev. X* **7**, 031051 (2017).
- [34] L. Zou, Y.-C. He, and C. Wang, *Phys. Rev. X* **11**, 031043 (2021).
- [35] J. Wang, X.-G. Wen, and E. Witten, *Phys. Rev. X* **8**, 031048 (2018).
- [36] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, *Phys. Rev. B* **87**, 155114 (2013).
- [37] T. Sulejmanpasic, H. Shao, A. W. Sandvik, and M. Ünsal, *Phys. Rev. Lett.* **119**, 091601 (2017).
- [38] N. Shannon, G. Misguich, and K. Penc, *Phys. Rev. B* **69**, 220403 (2004).
- [39] R. Moessner, O. Tchernyshyov, and S. L. Sondhi, *Journal of Statistical Physics* **116**, 755 (2004).
- [40] A. W. Glaetzle, M. Dalmonte, R. Nath, I. Rousochatzakis, R. Moessner, and P. Zoller, *Phys. Rev. X* **4**, 041037 (2014).
- [41] R. Moessner and S. L. Sondhi, *Phys. Rev. B* **63**, 224401 (2001).
- [42] L.-P. Henry and T. Roscilde, *Phys. Rev. Lett.* **113**, 027204 (2014).
- [43] M. Sadrzadeh, R. Haghshenas, and A. Langari, *Phys. Rev. B* **99**, 144414 (2019).
- [44] A. Sen, K. Damle, and T. Senthil, *Phys. Rev. B* **76**, 235107 (2007).
- [45] S. Wessel, *Phys. Rev. B* **78**, 075112 (2008).
- [46] A. Ralko, F. Trouselet, and D. Poilblanc, *Phys. Rev. Lett.* **104**, 127203 (2010).
- [47] S. Wessel, *Phys. Rev. B* **86**, 140501 (2012).
- [48] D. Banerjee, F.-J. Jiang, P. Widmer, and U.-J. Wiese, *Journal of Statistical Mechanics: Theory and Experiment* **2013**, P12010 (2013).
- [49] E. Fradkin and S. Kivelson, *Modern Physics Letters B* **04**, 225 (1990).
- [50] E. Fradkin, *Field theories of condensed matter physics* (Cambridge University Press, 2013).
- [51] A. H. Castro Neto, P. Pujol, and E. Fradkin, *Phys. Rev. B* **74**, 024302 (2006).
- [52] M. Hermele, M. P. A. Fisher, and L. Balents, *Phys. Rev. B* **69**, 064404 (2004).
- [53] F. Tschirsich, S. Montangero, and M. Dalmonte, *SciPost Phys.* **6**, 028 (2019).
- [54] X. Ran, Z. Yan, Y.-C. Wang, J. Rong, Y. Qi, and Z. Y. Meng, arXiv preprint arXiv:2209.10728 (2022).
- [55] Z. Xiong, Y. Xu, and X.-F. Zhang, arXiv preprint arXiv:2111.11445 (2021).
- [56] T. Hahn, U. Shmueli, and J. W. Arthur, *International tables for crystallography*, Vol. A (Springer, 2005).
- [57] P. Sindzingre, J.-B. Fouet, and C. Lhuillier, *Phys. Rev. B* **66**, 174424 (2002).
- [58] J.-S. Bernier, C.-H. Chung, Y. B. Kim, and S. Sachdev, *Phys. Rev. B* **69**, 214427 (2004).
- [59] O. A. Starykh, A. Furusaki, and L. Balents, *Phys. Rev. B* **72**, 094416 (2005).
- [60] S. C. Furuya and Y. Horinouchi, *Phys. Rev. B* **100**, 174435 (2019).
- [61] H. Ishizuka, Y. Motome, N. Furukawa, and S. Suzuki, *Journal of Physics: Conference Series* **320**, 012054 (2011).
- [62] E. H. Lieb, *Phys. Rev. Lett.* **18**, 692 (1967).
- [63] A. Rasmussen, Y.-Z. You, and C. Xu, arXiv preprint arXiv:1601.08235 (2016).
- [64] E. Ardonne, P. Fendley, and E. Fradkin, *Annals of Physics* **310**, 493 (2004).
- [65] A. Polyakov, *Nuclear Physics B* **120**, 429 (1977).
- [66] A. M. Polyakov, *Gauge fields and strings* (Taylor & Francis, 1987).
- [67] M. B. Einhorn and R. Savit, *Phys. Rev. D* **17**, 2583 (1978).
- [68] T. A. DeGrand and D. Toussaint, *Phys. Rev. D* **22**, 2478 (1980).
- [69] D. Tong, Lecture notes, DAMTP Cambridge **10** (2018).
- [70] J. Herzog-Arbeitman, S. Mantilla, and I. Sodemann, *Phys. Rev. B* **99**, 245108 (2019).

- [71] D. S. Rokhsar and S. A. Kivelson, [Phys. Rev. Lett. **61**, 2376 \(1988\)](#).
- [72] D. J. Williamson, Z. Bi, and M. Cheng, [Phys. Rev. B **100**, 125150 \(2019\)](#).
- [73] E. Lake, arXiv preprint arXiv:1802.07747 (2018).
- [74] C. Cordova, T. T. Dumitrescu, K. Intriligator, and S.-H. Shao, arXiv preprint arXiv:2205.09545 (2022).
- [75] C. L. Henley, [Journal of Statistical Physics **89**, 483 \(1997\)](#).
- [76] A. Paramekanti, L. Balents, and M. P. A. Fisher, [Phys. Rev. B **66**, 054526 \(2002\)](#).
- [77] R. Savit, [Phys. Rev. B **17**, 1340 \(1978\)](#).
- [78] J. V. José, L. P. Kadanoff, S. Kirkpatrick, and D. R. Nelson, [Phys. Rev. B **16**, 1217 \(1977\)](#).
- [79] N. Read and S. Sachdev, [Phys. Rev. B **42**, 4568 \(1990\)](#).
- [80] A. Vishwanath, L. Balents, and T. Senthil, [Phys. Rev. B **69**, 224416 \(2004\)](#).
- [81] M. Levin and T. Senthil, [Phys. Rev. B **70**, 220403 \(2004\)](#).
- [82] J. M. Kosterlitz and D. J. Thouless, [Journal of Physics C: Solid State Physics **6**, 1181 \(1973\)](#).
- [83] Y. Fuji, [Phys. Rev. B **93**, 104425 \(2016\)](#).
- [84] Z. Wu and L. Li, arXiv preprint arXiv:2208.10518 (2022).
- [85] I. P. Castillo, arXiv preprint arXiv:2005.10729 (2020).
- [86] A. D. King, C. Nisoli, E. D. Dahl, G. Poulin-Lamarre, and A. Lopez-Bezanilla, [Science \(2021\)](#), [10.1126/science.abe2824](#).
- [87] S. Ebadi, T. T. Wang, H. Levine, A. Keesling, G. Semeghini, A. Omran, D. Bluvstein, R. Samajdar, H. Pichler, W. W. Ho, S. Choi, S. Sachdev, M. Greiner, V. Vuletić, and M. D. Lukin, [Nature **595**, 227 \(2021\)](#).