α -Robust Error Analysis of $L2-1_{\sigma}$ Scheme on Graded Mesh for Time-fractional Nonlocal Diffusion Equation

Pari J. Kundaliya¹

Abstract

In this work, a time-fractional nonlocal diffusion equation is considered. Based on the $L2\text{-}1_{\sigma}$ scheme on a graded mesh in time and the standard finite element method (FEM) in space, the fully-discrete $L2\text{-}1_{\sigma}$ finite element method is investigated for a time-fractional nonlocal diffusion problem. We prove the existence and uniqueness of fully-discrete solution. The α -robust error bounds are derived, i.e. bounds remain valid as $\alpha \to 1^-$, where $\alpha \in (0, 1)$ is the order of a temporal fractional derivative. The numerical experiments are presented to justify the theoretical findings.

Keywords: Nonlocal; weak singularity; $L2-1_{\sigma}$ scheme; graded mesh; α -robust error estimate.

AMS(MOS): 65M60, 65M15, 35R11.

1 Introduction

In recent years, many researchers have focused on standard nonlocal parabolic partial differential equations (PDEs) (see [2, 4, 21, 24, 27]). In this work, we consider the following time-fractional nonlocal diffusion equation:

 ${}_{0}^{c}D_{t}^{\alpha}u(x,t) - a(l(u))\,\Delta u(x,t) = f(x,t) \qquad \text{in} \quad \Omega \times (0,T], \tag{1a}$

$$u(x,t) = 0$$
 on $\partial \Omega \times (0,T]$, (1b)

$$u(x,0) = u_0(x) \qquad \text{in} \quad \Omega, \tag{1c}$$

where $\Omega \subseteq \mathbb{R}^d$ (d = 1 or 2) is a bounded domain with smooth boundary $\partial \Omega$, the function $a : \mathbb{R} \longrightarrow \mathbb{R}$ is a diffusion coefficient, and $l(u) = \int_{\Omega} u(x,t) dx$. The term ${}_0^c D_t^{\alpha} u(x,t)$ is the α^{th} order Caputo fractional derivative of u with $\alpha \in (0,1)$ and it is defined in [9, Definition 3.1] as

$${}_{0}^{c}D_{t}^{\alpha}u(x,t) = \frac{1}{\Gamma(1-\alpha)} \int_{0}^{t} (t-s)^{-\alpha} \frac{\partial u(x,s)}{\partial s} ds, \quad t > 0.$$

¹Department of Mathematics, Institute of Infrastructure, Technology, Research And Management, Ahmedabad, Gujarat, India, (pariben.kundaliya.pm@iitram.ac.in)

The problem considered here is nonlocal due to the presence of the term a(l(u)), which is determined by a global quantity. Nonlocal problems often lead to a number of mathematical and computational difficulties. This type of nonlocal problem appears in population dynamics, where the diffusion coefficient a depends on the entire population in the domain rather than on a local density [5, 24, 27].

The solution of time-fractional PDEs in general exhibits weak singularity near time t = 0 [12, 14, 16, 22, 23, 25]. This makes the existing numerical methods with uniform time mesh often less accurate. In [14], the authors consider a semilinear time-fractional diffusion equation and show that if the solution has singularity near time t = 0, then the L1 scheme on uniform mesh (with step size τ) attains $\mathcal{O}(\tau^{\alpha})$ order of convergence in L^{∞} -norm in time. The graded temporal meshes offer an efficient way of computing a reliable numerical solution near time t = 0 [15,22,25]. The L1 scheme is considered in [5] on graded mesh for solving the time-fractional nonlocal diffusion problem. It is shown that, in the case of weak singularity of solution unear the initial time t = 0, the optimal convergence result with $\mathcal{O}(\tau^{2-\alpha})$ in temporal direction is achieved. It is also known that L1 scheme can give at most $2 - \alpha$ order of convergence in temporal direction. Therefore, in the present work, we investigate a higher order scheme; namely, the $L2-1_{\sigma}$ scheme. This scheme was first proposed by Alikhanov [1] on uniform mesh and latter this scheme was applied in [6, 11, 13, 18] on general meshes. In [5], the error estimates are established, which are α -nonrobust although the numerical scheme performs well as $\alpha \to 1^-$. In present work, the α -robust error bounds are derived in L^2 -norm and H^1 -norm. The concept of α -robust bound was first introduced in [7] and then it was used in [8, 12, 13] for different problems.

Notation: Henceforth, C > 0 denotes a generic constant, which is independent of mesh parameters h and N. We write (\cdot, \cdot) for the inner product and $\|\cdot\|$ for the norm on $L^2(\Omega)$. For $m \in \mathbb{N}$, the notation $H^m(\Omega)$ denotes the standard Sobolev space with the norm $\|\cdot\|_m$ and the space $H^1_0(\Omega)$ is consist of functions from $H^1(\Omega)$ whose trace vanish on the boundary $\partial\Omega$.

Throughout, the following hypotheses are made on the given data a, f and u_0 :

- H1: There exist $m_1, m_2 > 0$ such that $0 < m_1 \le a(x) \le m_2 < \infty, \forall x \in \mathbb{R}$.
- H2: *a* is Lipschitz continuous, i.e., $|a(x_1) a(x_2)| \le L|x_1 x_2|, (x_1, x_2 \in \mathbb{R}).$

H3: $f \in L^2(0,T;L^2(\Omega))$ and $u_0 \in H^1_0(\Omega) \cap H^2(\Omega)$.

In Section 2, we give a fully-discrete scheme for problem (1). We derive a priori bound for a fully-discrete solution in Section 3. The existence and uniqueness of fully-discrete solution are proved in Section 4. Error analysis of our proposed scheme is presented in Section 5. Finally, our theoretical results are confirmed in Section 6.

2 Fully-discrete Scheme

In this section, we discretize the differential equation (1) in both space and time. Firstly, we write the weak formulation of problem (1) and it is given below. For each $t \in (0, T]$, find $u(\cdot, t) \in H_0^1(\Omega)$ such that

$$u(\cdot, 0) = u_0(\cdot), \quad \text{in } \Omega. \tag{2b}$$

Note that the above problem (2) has a unique weak solution under the Hypotheses H1-H3 stated in the introduction [20, Theorems 2.4 and 2.5].

In order to establish a fully-discrete scheme, we assume that τ_h is a quasi uniform partition of Ω into disjoint intervals in \mathbb{R}^1 or triangles in \mathbb{R}^2 and h is the mesh size. Let V_h be the M-dimensional subspace of $H_0^1(\Omega)$, which consists of continuous piecewise linear polynomials.

Let $N \in \mathbb{N}$ and $\tau_N := \left\{ t_n : t_n = T\left(\frac{n}{N}\right)^r, n = 0, \dots, N \right\}$ be a non-uniform partition of [0, T] with time step $\tau_n = t_n - t_{n-1}$ and mesh grading parameter $r \ge 1$. Let $u^n := u(t_n)$ and U^n be the approximate value of u at t_n . For $\sigma = \frac{\alpha}{2}$, we set $t_{n-\sigma} := (1-\sigma) t_n + \sigma t_{n-1}$ and $U^{n,\sigma} := (1-\sigma) U^n + \sigma U^{n-1}$.

The $L2-1_{\sigma}$ approximation to ${}_{0}^{c}D_{t_{n-\sigma}}^{\alpha}u$ on a graded mesh is given below [13, P.5]. For $n = 1, \ldots, N$,

Here $g_{1,1} = \tau_1^{-1} a_{1,1}$ and for $n \ge 2$,

$$g_{n,j} = \begin{cases} \tau_j^{-1} (a_{n,1} - b_{n,1}) & \text{(if } j = 1) \\ \tau_j^{-1} (a_{n,j} + b_{n,j-1} - b_{n,j}) & \text{(if } 2 \le j \le n - 1) \\ \tau_j^{-1} (a_{n,n} + b_{n,n-1}) & \text{(if } j = n), \end{cases}$$
(4)

where

$$a_{n,n} = \frac{1}{\Gamma(1-\alpha)} \int_{t_{n-1}}^{t_{n-\sigma}} (t_{n-\sigma}-\eta)^{-\alpha} d\eta, \text{ for } n \ge 1.$$

$$a_{n,j} = \frac{1}{\Gamma(1-\alpha)} \int_{t_{j-1}}^{t_j} (t_{n-\sigma}-\eta)^{-\alpha} d\eta, \text{ for } 1 \le j \le n-1.$$

$$b_{n,j} = \frac{2}{\Gamma(1-\alpha) (t_{j+1}-t_{j-1})} \int_{t_{j-1}}^{t_j} (t_{n-\sigma}-\eta)^{-\alpha} (\eta-t_{j-\frac{1}{2}}) d\eta, \text{ for } 1 \le j \le n-1.$$

Lemma 2.1. Let $v(t) \in C[0,T] \cap C^3(0,T]$ be a function with $|v^{(q)}(t)| \leq C(1+t^{\alpha-q})$, for q = 0, 1, 2, 3 and for $t \in (0,T]$. Then, for n = 1, ..., N,

$$t_{n-\sigma}^{\alpha} \left\|_{0}^{c} D_{t_{n-\sigma}}^{\alpha} v - D_{N}^{\alpha} v^{n-\sigma} \right\| \leq C N^{-\min\{3-\alpha, r\alpha\}}$$

Proof. The proof follows from [6, Lemmas 1 and 7].

Using the above notations, the *fully-discrete scheme* for (1) is following: For each n = 1, ..., N, find $U^n \in V_h$ such that $\forall v_h \in V_h$,

$$\left(D_N^{\alpha}U^{n-\sigma}, v_h\right) + a\left(l(U^{n,\sigma})\right)\left(\nabla U^{n,\sigma}, \nabla v_h\right) = (f^{n-\sigma}, v_h), \tag{5a}$$

$$U^0 = u_h^0, \tag{5b}$$

where u_h^0 is some approximation of $u_0(x)$.

Now, from the definition of D_N^{α} and as ϕ_i 's are the basis functions for V_h , from (5a), for each $i = 1, \ldots, M$, we have

$$g_{n,n}(U^{n}, \phi_{i}) - \left(g_{n,1}U^{0} + \sum_{k=2}^{n} (g_{n,k} - g_{n,k-1})U^{k-1}, \phi_{i}\right) + a(l(U^{n,\sigma}))(\nabla U^{n,\sigma}, \nabla \phi_{i})$$

= $(f^{n-\sigma}, \phi_{i}).$ (6)

Since $U^n \in V_h$, there exists $\alpha_j^n \in \mathbb{R}$ (for each j = 1, ..., M) such that

$$U^n = \sum_{j=1}^M \alpha_j^n \phi_j$$

Set $\alpha^n = [\alpha_1^n, \alpha_2^n, \dots, \alpha_M^n]$. It can be noticed that (6) is a nonlinear equation in U^n . If we directly use the Newton's method for solving the equation (6), then the Jacobian matrix becomes dense due to the presence of nonlocal term a(l(u)) [5]. Due to this term, the scheme becomes computationally expensive. To overcome this difficulty, we follow the idea of [10]. This technique was first proposed for elliptic nonlocal problems and latter it was analyzed in [3–5] for solving nonlocal parabolic problems. For this, we reformulate (6) as follows: Find $U^n \in V_h$ and $d \in \mathbb{R}$ such that

$$l(U^{n,\sigma}) - d = 0, \tag{7}$$

and for $1 \leq i \leq M$,

$$(U^{n}, \phi_{i}) - \frac{1}{g_{n,n}} \left(g_{n,1}U^{0} + \sum_{k=2}^{n} (g_{n,k} - g_{n,k-1})U^{k-1}, \phi_{i} \right) + \frac{1}{g_{n,n}} a(d) (\nabla U^{n,\sigma}, \nabla \phi_{i}) - \frac{1}{g_{n,n}} \left(f^{n-\sigma}, \phi_{i} \right) = 0.$$
(8)

Now, in order to use the Newton's method, we rewrite (7) and (8) as follows:

$$F_i(U^n, d) = 0, \text{ for } 1 \le i \le M+1,$$
 (9)

where, for each $1 \leq i \leq M$,

$$F_{i}(U^{n},d) = (U^{n},\phi_{i}) - \frac{1}{g_{n,n}} \Big(g_{n,1}U^{0} + \sum_{k=2}^{n} (g_{n,k} - g_{n,k-1})U^{k-1},\phi_{i} \Big) \\ + \frac{1}{g_{n,n}} a(d) (\nabla U^{n,\sigma}, \nabla \phi_{i}) - \frac{1}{g_{n,n}} \left(f^{n-\sigma}, \phi_{i} \right),$$

and

$$F_{M+1}(U^n, d) = l(U^{n,\sigma}) - d.$$

An application of Newton's method in (9) gives the following system of equation:

$$J\begin{bmatrix}\alpha^n\\d\end{bmatrix} = \begin{bmatrix}A & b\\c & -1\end{bmatrix}\begin{bmatrix}\alpha^n\\d\end{bmatrix} = \begin{bmatrix}\bar{F}\\F_{M+1}\end{bmatrix},$$

where J is the Jacobian matrix, $\alpha^n = [\alpha_1^n, \ldots, \alpha_M^n]^T$, $\overline{F} = [F_1, \ldots, F_M]^T$, and the entries of the matrices $A_{M \times M}$, $b_{M \times 1}$ and $c_{1 \times M}$ are given below.

$$A_{ij} = (\phi_j, \phi_i) + \frac{\sigma}{g_{n,n}} a(d) (\nabla \phi_j, \nabla \phi_i),$$

$$b_{i1} = \frac{1}{g_{n,n}} a'(d) (\nabla U^{n,\sigma}, \nabla \phi_i),$$

$$c_{1j} = \sigma \int_{\Omega} \phi_j \, dx.$$

Now, the sparsity of J follows from the sparsity of matrix A. Note that the solutions of (6) and (7)-(8) are equivalent [5, Theorem 3.1].

3 A priori Bound for Fully-discrete Solution

In this section, we provide a priori bound for fully-discrete solution of problem (5). In order to this, we shall need the definition of the Discrete Laplacian and the coercivity property of $L2-1_{\sigma}$ scheme.

Definition 3.1. [26, P.10] The discrete Laplacian is a map $\Delta_h : V_h \to V_h$ defined as

$$(\Delta_h u, v) = -(\nabla u, \nabla v), \quad \forall u, v \in V_h.$$

Lemma 3.2. [6, Corollary 1] For any function $v = v(\cdot, t)$ defined on a graded mesh $\{t_j\}_{j=0}^N$, one has

$$\frac{1}{2}D_N^{\alpha}\|v^{n-\sigma}\|^2 \leq \left(D_N^{\alpha}v^{n-\sigma}, v^{n,\sigma}\right), \quad for \ n = 1, \dots, N.$$

For further analysis, for n = 1, ..., N, we define the coefficients $p_{n-i}^{(n)}$ as follows:

$$p_{n-i}^{(n)} := \begin{cases} \frac{1}{g_{i,i}} \sum_{k=i+1}^{n} \left(g_{k,i+1} - g_{k,i} \right) p_{n-k}^{(n)} & \text{(if } i = 1, \dots, n-1) \\ \frac{1}{g_{n,n}} & \text{(if } i = n). \end{cases}$$
(10)

Lemma 3.3. [13, P.46 (15)] Let $\gamma \in (0, 1)$ be any constant. For n = 1, ..., N, the coefficients defined in (10) satisfy the following inequality.

$$\sum_{i=1}^{n} p_{n-i}^{(n)} i^{r(\gamma-\alpha)} \leq \frac{11\Gamma(1+\gamma-\alpha)}{4\Gamma(1+\gamma)} T^{\alpha} \left(\frac{t_n}{T}\right)^{\gamma} N^{r(\gamma-\alpha)}.$$

Now, we state the discrete fractional Gr \ddot{o} nwall inequality, which will be used in deriving *a priori* bounds and error estimates.

Lemma 3.4. [13, Lemma 4.1] Let $n \ge 1$ and let λ_i be non-negative constants with $0 < \sum_{i=0}^{n} \lambda_i \le \Lambda$, where Λ is some positive constant independent of n. Suppose that the non-negative sequences $\{\xi^n, \eta^n : n \ge 1\}$ are bounded and the non-negative grid function $\{v^n : n \ge 0\}$ satisfies

$$D_N^{\alpha}(v^{n-\sigma})^2 \leq \sum_{i=1}^n \lambda_{n-i} (v^{i,\sigma})^2 + \xi^n v^{n,\sigma} + (\eta^n)^2, \text{ for } n \geq 1.$$

If the non-uniform grid satisfies the criterion $\max_{1 \le n \le N} \tau_n \le \frac{1}{\sqrt[\alpha]{\frac{11}{2}} \Gamma(2-\alpha) \Lambda}$, then

$$v^{n} \leq 2E_{\alpha} \left(\frac{11}{2} \Lambda t_{n}^{\alpha}\right) \left[v^{0} + \max_{1 \leq k \leq n} \sum_{j=1}^{k} p_{k-j}^{(k)}(\xi^{j} + \eta^{j}) + \max_{1 \leq j \leq n} \left\{\eta^{j}\right\}\right].$$

Remark 3.5. The L2-1 $_{\sigma}$ scheme on graded mesh satisfies all the hypotheses, which are required for the proof of Lemmas 3.3 and 3.4 [17, Example 3, P.222].

The next theorem is one of the main results of this article.

Theorem 3.6. The fully-discrete solution U^n of (5) satisfies

$$\|U^n\| \leq C(1+\|U^0\|), \tag{11}$$

$$\|\nabla U^n\| \leq C \left(1 + \|\nabla U^0\|\right), \tag{12}$$

for each n = 1, ..., N.

Proof. First we note that, by substituting $\gamma = \alpha$ in Lemma 3.3, we can get

$$\sum_{j=1}^{k} p_{k-j}^{(k)} \le \frac{11 t_k^{\alpha}}{4 \Gamma(1+\alpha)} \le \frac{11 T^{\alpha}}{4 \Gamma(1+\alpha)}.$$
(13)

Next, by taking $v_h = U^{n,\sigma}$ in (5a), we get

$$\left(D_N^{\alpha}U^{n-\sigma}, U^{n,\sigma}\right) + a\left(l(U^{n,\sigma})\right)\left(\nabla U^{n,\sigma}, \nabla U^{n,\sigma}\right) = (f^{n-\sigma}, U^{n,\sigma}), \ \forall v_h \in V_h.$$
(14)

Applying Hypothesis H1 and the Cauchy-Schwartz inequality in (14), we have

$$\left(D_N^{\alpha}U^{n-\sigma}, U^{n,\sigma}\right) \leq \|f^{n-\sigma}\| \, \|U^{n,\sigma}\|.$$
(15)

Using Lemma 3.2 and the inequality $b_1b_2 \leq \frac{b_1^2}{2} + \frac{b_2^2}{2}$ in (15), we get

$$\frac{1}{2}D_N^{\alpha} \|U^{n-\sigma}\|^2 \le \frac{1}{2}\|f^{n-\sigma}\|^2 + \frac{1}{2}\|U^{n,\sigma}\|^2.$$
(16)

An application of Lemma 3.4 into (16) gives

$$||U^{n}|| \leq 2E_{\alpha} \left(\frac{11}{4}t_{n}^{\alpha}\right) \left(||U^{0}|| + \max_{1 \leq k \leq n} \left(\max_{1 \leq j \leq k} ||f^{j-\sigma}||\right) \sum_{j=1}^{k} p_{k-j}^{(k)} + \max_{1 \leq j \leq n} ||f^{j-\sigma}||\right).$$

Thus, using the estimate (13) in the above inequality, we have

$$\|U^n\| \le C\left(1 + \|U^0\|\right),$$

where $C = 2E_{\alpha}\left(\frac{11}{4}t_n^{\alpha}\right) \max\left\{1, \left(\frac{11T^{\alpha}}{4\Gamma(1+\alpha)} + 1\right) \max_{1\le j\le n} \|f^{j-\sigma}\|\right\}.$

To derive an estimate for $\|\nabla U^n\|$, we rewrite (5a) using the operator Δ_h as follows:

$$\left(D_N^{\alpha}U^{n-\sigma}, v_h\right) - a\left(l(U^{n,\sigma})\right)\left(\Delta_h U^{n,\sigma}, v_h\right) = (f^{n-\sigma}, v_h), \ \forall v_h \in V_h.$$
(17)

Now we take $v_h = -\Delta_h U^{n,\sigma}$ in (17) and then proceed in a similar manner as above to obtain

$$\|\nabla U^n\| \le C\left(1 + \|\nabla U^0\|\right),\,$$

where $C = 2 E_{\alpha} \left(\frac{11}{2} t_n^{\alpha}\right) \max \left\{ 1, \frac{1}{\sqrt{m_1}} \left(\frac{11 T^{\alpha}}{4 \Gamma(1+\alpha)} + 1\right) \max_{1 \le j \le n} \|f^{j-\sigma}\| \right\}$ and the constant m_1 is as per Hypothesis H1 in the introduction. This completes the proof. \Box

4 Existence-Uniqueness of Fully-discrete Solution

For proving the existence and the uniqueness of a fully-discrete solution, we shall need the following proposition, which is a consequence of the Brouwer's fixed point theorem [26, P.237].

Proposition 4.1. Let $S : X \to X$ be a continuous map on a finite dimensional Hilbert space X such that (S(w), w) > 0, $\forall w \in X$ with $||w||_X = \zeta$, $\zeta > 0$. Then there exists $v \in X$ such that S(v) = 0, $||v||_X \leq \zeta$.

We also define

$$\tau := \frac{2m_1 \left(a_{n,n} + b_{n,n-1} \right)}{(1-\sigma) \left\{ LCR_1 (1-2\sigma) \right\}^2},\tag{18}$$

where $R_1 = C(1 + \|\nabla U^0\|).$

In next result, we prove the existence and uniqueness of a fully-discrete solution. The proof is almost similar to the proof of [5, Theorem 4.1]. However, we prove it here for the sake of completeness.

Theorem 4.2. Let U^0, \ldots, U^{n-1} be given. Then for all $1 \le n \le N$, there exists a unique solution $U^n \in V_h$ of (5), provided $\max_{1 \le n \le N} \tau_n \le \tau$.

Proof: First we prove the existence. Rewriting (5a) as follows:

$$(U^{n}, v_{h}) + \frac{1}{g_{n,n}} a \left(l(U^{n,\sigma}) \right) (\nabla U^{n,\sigma}, \nabla v_{h}) - \frac{1}{g_{n,n}} \left(f^{n-\sigma}, v_{h} \right) - \frac{1}{g_{n,n}} \left(g_{n,1} U^{0} + \sum_{k=2}^{n} (g_{n,k} - g_{n,k-1}) U^{k-1}, v_{h} \right) = 0.$$
(19)

Now, we define a map $G: V_h \longrightarrow V_h$ such that

$$(G(X^{n}), v_{h}) = (X^{n}, v_{h}) - \frac{1}{g_{n,n}} \left(g_{n,1}U^{0} + \sum_{k=2}^{n} (g_{n,k} - g_{n,k-1})U^{k-1}, v_{h} \right) + \frac{(1 - \sigma)}{g_{n,n}} a (l(U^{n,\sigma})) (\nabla X^{n}, \nabla v_{h}) - \frac{1}{g_{n,n}} (f^{n-\sigma}, v_{h}) + \frac{\sigma}{g_{n,n}} a (l(U^{n,\sigma})) (\nabla U^{n-1}, \nabla v_{h}).$$

$$(20)$$

Then the map G is continuous. By choosing $v_h = X^n$ in (20), we get

$$(G(X^{n}), X^{n}) \geq \left(\|X^{n}\| - \frac{g_{n,1}}{g_{n,n}} \|U^{0}\| - \frac{1}{g_{n,n}} \sum_{k=2}^{n} (g_{n,k} - g_{n,k-1}) \|U^{k-1}\| - \frac{\sigma m_{1}}{g_{n,n}} \|\nabla U^{n-1}\| - \frac{1}{g_{n,n}} \|f^{n-\sigma}\| \right) \|X^{n}\|.$$

$$(21)$$

Thus, $(G(X^n), X^n) > 0$, if

$$||X^{n}|| > \frac{g_{n,1}}{g_{n,n}} ||U^{0}|| + \frac{1}{g_{n,n}} \sum_{k=2}^{n} (g_{n,k} - g_{n,k-1}) ||U^{k-1}|| + \frac{\sigma m_{1}}{g_{n,n}} ||\nabla U^{n-1}|| + \frac{1}{g_{n,n}} ||f^{n-\sigma}||.$$
(22)

Hence, by Proposition 4.1, we can assure that (5) has a solution.

Next we prove the uniqueness. Suppose, if possible, there exist two solutions of (5), say U_1^n and U_2^n at time $t = t_n$. For simplicity, we denote U_1^n by U_1 , U_2^n by U_2 , $U_1^{n,\sigma}$ by U_1^{σ} and $U_2^{n,\sigma}$ by U_2^{σ} , respectively. Let $U_1 - U_2 = \mathcal{Y}$. From (19), we can get

$$(U_1 - U_2, v_h) + \frac{(1 - \sigma)}{g_{n,n}} a(l(U_1^{\sigma})) (\nabla U_1, \nabla v_h) - \frac{(1 - \sigma)}{g_{n,n}} a(l(U_2^{\sigma})) (\nabla U_2, \nabla v_h) + \frac{\sigma}{g_{n,n}} \left\{ a(l(U_1^{\sigma})) - a(l(U_2^{\sigma})) \right\} (\nabla U^{n-1}, \nabla v_h) = 0.$$
(23)

By subtracting $\frac{(1-\sigma)}{g_{n,n}} a(l(U_1^{\sigma})) (\nabla U_2, \nabla v_h)$ from both sides of (23) and then choosing $v_h = \mathcal{Y}$, we get

$$(\mathcal{Y}, \mathcal{Y}) + \frac{(1-\sigma)}{g_{n,n}} a(l(U_1^{\sigma}))(\nabla \mathcal{Y}, \nabla \mathcal{Y}) = \frac{(1-\sigma)}{g_{n,n}} \Big\{ a(l(U_2^{\sigma})) - a(l(U_1^{\sigma})) \Big\} (\nabla U_2, \nabla \mathcal{Y}) \\ - \frac{\sigma}{g_{n,n}} \Big\{ a(l(U_1^{\sigma})) - a(l(U_2^{\sigma})) \Big\} (\nabla U^{n-1}, \nabla \mathcal{Y}).$$

By using (12), Hypotheses H1-H2, the Cauchy-Schwartz inequality, and the Young's inequality in the above equation, we have

$$\left(1 - \frac{(1-\sigma)\{LCR_1(1-2\sigma)\}^2}{2m_1g_{n,n}}\right) \|\mathcal{Y}\|^2 \le 0.$$

Since $\max_{1 \le n \le N} \tau_n \le \tau$, it follows that $\left(1 - \frac{(1-\sigma)\{LCR_1(1-2\sigma)\}^2}{2m_1 g_{n,n}}\right) > 0$. Hence, $U_1 = U_2$. This completes the proof.

5 Error Estimates

For deriving a priori error estimates in Theorem 5.3, the following definition and lemma will be required.

Definition 5.1. [26, P.8] The Ritz-projection is a map $R_h : H_0^1(\Omega) \to V_h$ such that

$$(\nabla w, \nabla v) = (\nabla R_h w, \nabla v), \quad \forall w \in H_0^1(\Omega) \text{ and } \forall v \in V_h$$

Utilizing (25), one can easily prove that R_h satisfies $\|\nabla R_h w\| \leq C$. Since V_h is the space of linear polynomials, it is well known [26, Lemma 1.1] that

$$\|w - R_h w\| + h \|\nabla (w - R_h w)\| \le Ch^2 \|\Delta w\|, \quad \forall w \in H^2 \cap H^1_0.$$
(24)

Lemma 5.2. [11, P.9 (32)] If $u \in C^2(0,T]$ and $\|\partial_t^q u\| \leq C(1+t^{\alpha-q})$, for q = 0, 1, 2, 3, then, for n = 1, ..., N,

$$t_{n-\sigma}^{\alpha} \| u^{n,\sigma} - u^{n-\sigma} \| \le C N^{-\min\{r\alpha,2\}}.$$

Now, in the following theorem, we state and prove the most significant result of the present work. For this, we have to assume that the solution u satisfies the following conditions.

$${}_{0}^{c}D_{t}^{\alpha}u \in L^{\infty}(0,T; H^{2}(\Omega)), \qquad u \in L^{\infty}(0,T; H_{0}^{1}(\Omega) \cap H^{3}(\Omega)),$$

and $\|\partial_{t}^{q}u\|_{3} \leq C (1+t^{\alpha-q}), \quad \text{for } q = 0, 1, 2, 3.$ (25)

Theorem 5.3. For $1 \leq n \leq N$, let u^n and U^n be the solutions of (1) and (5), respectively. Then

$$\max_{1 \le n \le N} \|u^n - U^n\| \le C \left(h^2 + N^{-\min\{2, r\alpha\}}\right), \tag{26}$$

$$\max_{1 \le n \le N} \|\nabla (u^n - U^n)\| \le C \left(h + N^{-\min\{2, r\alpha\}}\right).$$
(27)

Proof. For the proof, we split the error $(u^n - U^n)$ as follows:

$$u^n - U^n = u^n - R_h u^n + R_h u^n - U^n = \rho^n + \theta^n,$$

where $\rho^n := u^n - R_h u^n$ and $\theta^n := R_h u^n - U^n$. As the estimate for ρ^n can be given by (24), it is sufficient to find estimate for θ^n for proving the theorem.

For any $v_h \in V_h$, the estimate for θ^n is given by

$$\begin{pmatrix} D_N^{\alpha} \theta^{n-\sigma}, v_h \end{pmatrix} + a (l(U^{n,\sigma})) (\nabla \theta^{n,\sigma}, \nabla v_h) \\ = (D_N^{\alpha} R_h u^{n-\sigma} - {}_0^c D_{t_{n-\sigma}}^{\alpha} u, v_h) - a (l(u^{n-\sigma})) (\Delta u^{n,\sigma} - \Delta u^{n-\sigma}, v_h) \\ - \{a (l(U^{n,\sigma})) - a (l(u^{n-\sigma}))\} (\Delta u^{n,\sigma}, v_h).$$

$$(28)$$

Now, we choose $v_h = \theta^{n,\sigma}$ in (28) to get

$$\left(D_N^{\alpha} \theta^{n-\sigma}, \theta^{n,\sigma} \right) + a \left(l(U^{n,\sigma}) \right) \left(\nabla \theta^{n,\sigma}, \nabla \theta^{n,\sigma} \right) = \left(D_N^{\alpha} R_h u^{n-\sigma} - {}^c_0 D_{t_{n-\sigma}}^{\alpha} u, \theta^{n,\sigma} \right) - \left\{ a \left(l(U^{n,\sigma}) \right) - a \left(l(u^{n-\sigma}) \right) \right\} (\Delta u^{n,\sigma}, \theta^{n,\sigma}) - a \left(l(u^{n-\sigma}) \right) (\Delta u^{n,\sigma} - \Delta u^{n-\sigma}, \theta^{n,\sigma}).$$
(29)

An application of the bound of a, the Cauchy-Schwartz inequality and the Triangle inequality in (29) gives

$$\begin{aligned} \left(D_N^{\alpha} \theta^{n-\sigma}, \theta^{n,\sigma} \right) + m_1 \| \nabla \theta^{n,\sigma} \|^2 &\leq \| D_N^{\alpha} R_h u^{n-\sigma} - {}_0^c D_{t_{n-\sigma}}^{\alpha} R_h u \| \| \theta^{n,\sigma} \| \\ &+ \| {}_0^c D_{t_{n-\sigma}}^{\alpha} R_h u - {}_0^c D_{t_{n-\sigma}}^{\alpha} u \| \| \theta^{n,\sigma} \| \\ &+ m_2 \| \Delta u^{n,\sigma} - \Delta u^{n-\sigma} \| \| \theta^{n,\sigma} \| \\ &+ \left| a \left(l(U^{n,\sigma}) \right) - a \left(l(u^{n-\sigma}) \right) \right| \| \Delta u^{n,\sigma} \| \| \theta^{n,\sigma} \|. \end{aligned}$$

Since function a is Lipschitz continuous, we have

$$\begin{aligned} \left| a \left(l(U^{n,\sigma}) \right) - a \left(l(u^{n-\sigma}) \right) \right| &\leq L C \left\| U^{n,\sigma} - u^{n-\sigma} \right\| \\ &\leq L C \left\{ \left\| U^{n,\sigma} - u^{n,\sigma} \right\| + \left\| u^{n,\sigma} - u^{n-\sigma} \right\| \right\} \\ &\leq L C \left\{ \left\| \rho^{n,\sigma} \right\| + \left\| \theta^{n,\sigma} \right\| + \left\| u^{n,\sigma} - u^{n-\sigma} \right\| \right\}. \end{aligned}$$
(31)

Also, it follows from (25) and (24) that

$$\|{}_{0}^{c}D_{t_{n-\sigma}}^{\alpha}R_{h}u - {}_{0}^{c}D_{t_{n-\sigma}}^{\alpha}u\| \leq C h^{2} \|\Delta_{0}^{c}D_{t_{n-\sigma}}^{\alpha}u\| \leq C h^{2}.$$
(32)

From Lemma 3.2, (24), (30), (31) and (32), we have

$$D_{N}^{\alpha} \|\theta^{n-\sigma}\|^{2} \leq C_{1} \|\theta^{n,\sigma}\|^{2} + C_{2} \Big\{ \|D_{N}^{\alpha}R_{h}u^{n-\sigma} - {}_{0}^{c}D_{t_{n-\sigma}}^{\alpha}R_{h}u\| + \|u^{n,\sigma} - u^{n-\sigma}\| \\ + \|\Delta u^{n,\sigma} - \Delta u^{n-\sigma}\| + h^{2} \Big\} \|\theta^{n,\sigma}\|, \quad (33)$$

where C_1 is dependent on C, L and C_2 is dependent on C, L, m_2 .

An application of Lemma 3.4 into (33) leads to

$$\|\theta^{n}\| \leq 2 E_{\alpha} \Big(\frac{11}{2} C_{1} t_{n}^{\alpha} \Big) \Big(\|\theta^{0}\| + C_{2} \max_{1 \leq k \leq n} \sum_{j=1}^{k} p_{k-j}^{(k)} \Big\{ h^{2} + \|u^{j,\sigma} - u^{j-\sigma}\| + \|D_{N}^{\alpha} R_{h} u^{j-\sigma} - {}_{0}^{c} D_{t_{j-\sigma}}^{\alpha} R_{h} u\| + \|\Delta u^{j,\sigma} - \Delta u^{j-\sigma}\| \Big\} \Big).$$
(34)

Now, from Lemmas 2.1 and 5.2, we have

$$\begin{split} &\| D_N^{\alpha} R_h u^{j-\sigma} - {}^c_0 D_{t_{j-\sigma}}^{\alpha} R_h u \| + \| u^{j,\sigma} - u^{j-\sigma} \| + \| \Delta u^{j,\sigma} - \Delta u^{j-\sigma} \| \\ &\leq C t_{j-\sigma}^{-\alpha} \left(N^{-\min\{3-\alpha,r\alpha\}} + N^{-\min\{2,r\alpha\}} \right) \\ &\leq C \left(2N \right)^{r\alpha} N^{-\min\{2,r\alpha\}} j^{r(l_N-\alpha)}, \end{split}$$
(35)

where the fact that $t_{j-\sigma}^{-\alpha} \leq j^{-r\alpha} (2N)^{r\alpha} \leq (2N)^{r\alpha} j^{r(l_N-\alpha)}$ is used with $l_N = \frac{1}{\ln N}$.

Using (35) in (34), we can obtain

$$\|\theta^{n}\| \leq 2 E_{\alpha} \left(\frac{11}{2} C_{1} t_{n}^{\alpha}\right) \left(\|\theta^{0}\| + C_{2} h^{2} \max_{1 \leq k \leq n} \sum_{j=1}^{k} p_{k-j}^{(k)} + C (2N)^{r\alpha} N^{-\min\{2, r\alpha\}} \max_{1 \leq k \leq n} \sum_{j=1}^{k} p_{k-j}^{(k)} j^{r(l_{N}-\alpha)}\right).$$
(36)

Now, using Lemma 3.3 with $\gamma = l_N$, we can have

$$\sum_{j=1}^{k} p_{k-j}^{(k)} j^{r(l_N - \alpha)} \leq \frac{11 \Gamma(1 + l_n - \alpha)}{4 \Gamma(1 + l_N)} T^{\alpha} N^{r(l_N - \alpha)}.$$
(37)

Using (13) and (37) in (36), we get

$$\begin{aligned} \|\theta^{n}\| &\leq 2 E_{\alpha} \Big(\frac{11}{2} C_{1} t_{n}^{\alpha} \Big) \Big(\|\theta^{0}\| + \frac{11 C_{2} T^{\alpha}}{4 \Gamma(1+\alpha)} h^{2} \\ &+ C (2N)^{r\alpha} \frac{11 \Gamma(1+l_{n}-\alpha)}{4 \Gamma(1+l_{N})} T^{\alpha} N^{r(l_{N}-\alpha)} N^{-\min\{2,r\alpha\}} \Big). \end{aligned}$$
(38)

Choosing $U^0 = R_h u_0$, we get $\|\theta^0\| = 0$. Therefore, from (38), we have,

$$\|\theta^n\| \le C\left(h^2 + N^{-\min\{2, r\alpha\}}\right),\tag{39}$$

where $C = 2 E_{\alpha} \left(\frac{11}{2} C_1 t_n^{\alpha} \right) \max \left\{ \frac{11 C_2 T^{\alpha}}{4 \Gamma(1+\alpha)}, 2^{r\alpha} C \frac{11 \Gamma(1+l_n-\alpha)}{4 \Gamma(1+l_N)} N^{rl_N} T^{\alpha} \right\}.$

Finally, an application of the Triangle inequality and the estimates (24) and (39) give (26).

Now, to derive the error estimate in H_0^1 -norm, we use the definition of discrete Laplacian Δ_h and rewrite (28) as follows:

$$\begin{pmatrix} D_N^{\alpha} \theta^{n-\sigma}, v_h \end{pmatrix} - a (l(U^{n,\sigma})) (\Delta_h \theta^{n,\sigma}, v_h) \\ = \left(D_N^{\alpha} R_h u^{n-\sigma} - {}_0^c D_{t_{n-\sigma}}^{\alpha} u, v_h \right) - a (l(u^{n-\sigma})) (\Delta u^{n,\sigma} - \Delta u^{n-\sigma}, v_h) \\ - \left\{ a (l(U^{n,\sigma})) - a (l(u^{n-\sigma})) \right\} (\Delta u^{n,\sigma}, v_h).$$

$$(40)$$

We take $v_h = -\Delta_h \theta^{n,\sigma}$ in (40) and then use the similar arguments as above together with the Poincaré inequality to obtain

$$\|\nabla u^n - \nabla U^n\| \le C \left(h + N^{-\min\{2, r\alpha\}}\right).$$

This completes the proof.

Corollary 5.4. For $1 \leq n \leq N$, let u^n and U^n be the solutions of (1) and (5), respectively. Then

$$\max_{1 \le n \le N} \|u^n - U^n\|_1 \le C \left(h + N^{-\min\{2, r\alpha\}}\right).$$

Proof. This proof follows from the Poincaré inequality and Theorem 5.3. \Box

Remark 5.5. Above analysis can be easily extended for the case when f = f(u) is Lipschitz continuous, except the H_0^1 -norm error analysis. More precisely, in case of f = f(u), we get a term $\|\nabla (f(u^{n-\sigma}) - f(U^{n,\sigma}))\|$ which creates problem in getting H_0^1 -norm error estimate.

6 Numerical Experiments

In this section, for confirmation of our theoretical estimates, we perform two numerical experiments with known exact solution. In both numerical experiments, the time interval is taken to be [0, 1]. In order to achieve optimal order of convergence rate, we choose mesh grading parameter $r = \frac{2}{\alpha}$. We consider the tolerance $\epsilon = 10^{-12}$ for stopping the iterations in Newton's method. Spatial domain Ω is partitioned with (M_s+1) node points in each direction and the time interval is partitioned into N number of sub-intervals. We set $M_s = N$ and calculate error for different values of N to get the rate of convergence in time direction. Similarly, we set $N = M_s$ and calculate error for different values of M_s to conclude the rate of convergence in spatial direction.

Example 6.1. For this example, we consider equation (1) with a spatial domain $\Omega = (0, \pi)$ and $a(w) = 3 + \sin w$. Then we can choose f(x, t) such that the analytical solution of equation (1) is $u(x, t) = (t^3 + t^{\alpha}) \sin x$.

Table 1 shows the order of convergence in the temporal direction in $L^{\infty}(0, T; L^{2}(\Omega))$ norm on graded mesh for $\alpha = 0.4$, $\alpha = 0.6$ and $\alpha = 0.8$. From this table, it can be seen that using graded mesh, we can achieve $O(N^{-2})$ convergence rate in the temporal direction. The order of convergence in the spatial direction in L^{2} -norm and H_{0}^{1} -norm for $\alpha = 0.4$, $\alpha = 0.6$ and $\alpha = 0.8$ are given in Tables 2 and 3, respectively. From these tables, one can observe that the numerical convergence results are in accordance with our theoretical convergence estimates.

N	$\alpha = 0.4$		$\alpha = 0.6$		$\alpha = 0.8$	
	Error	OC	Error	OC	Error	OC
2^{6}	1.22E-03	1.9748	1.04E-03	2.0270	1.01E-03	2.0136
2^{7}	3.11E-04	2.0069	2.54E-04	2.0239	2.49E-04	2.0143
2^{8}	7.75E-05	2.0073	6.26E-05	2.0149	6.17E-05	2.0138
2^{9}	1.93E-05	2.0068	1.55E-05	2.0125	1.53E-05	2.0128
2^{10}	4.80E-06	-	3.84E-06	-	3.79E-06	-

Table 1: Error and order of convergence in $L^{\infty}(0,T; L^{2}(\Omega))$ norm in temporal direction on graded mesh for Example 6.1.

M_s	$\alpha = 0.4$		$\alpha = 0.6$		$\alpha = 0.8$	
	Error	OC	Error	OC	Error	OC
2^{6}	7.97E-04	2.0785	1.04E-03	2.0270	1.01E-03	2.0136
2^{7}	1.89E-04	2.0778	2.54E-04	2.0274	2.49E-04	2.0143
2^{8}	4.47E-05	2.0629	6.24E-05	2.0243	6.17E-05	2.0139
2^{9}	1.07E-05	2.0472	1.53E-05	2.0203	1.53E-05	2.0128
2^{10}	2.59E-06	-	3.78E-06	-	3.79E-06	-

Table 2: Error and order of convergence in L^2 -norm in space for Example 6.1.

M_s	$\alpha = 0.4$		$\alpha = 0.6$		$\alpha = 0.8$	
	Error	OC	Error	OC	Error	OC
2^{6}	3.55E-02	0.9999	3.55E-02	1.0000	3.55E-02	1.0000
2^{7}	1.78E-02	0.9999	1.78E-02	1.0000	1.78E-02	1.0000
2^{8}	8.88E-03	0.9999	8.88E-03	1.0000	8.88E-03	1.0000
2^{9}	4.44E-03	0.9999	4.44E-03	1.0000	4.44E-03	1.0000
2^{10}	2.22E-03	-	2.22E-03	-	2.22E-03	-

Table 3: Error and order of convergence in H_0^1 -norm in space for Example 6.1.

Example 6.2. In this second example, consider equation (1) with the spatial domain $\Omega = (0, 1) \times (0, 1)$ and $a(w) = 3 + \sin w$. Then we can choose f(x, y, t) in such a way that the analytical solution of the equation (1) is $u(x, y, t) = (t^3 + t^{\alpha})(x - x^2)(y - y^2)$.

In Table 4, we have given the order of convergence in the temporal direction in $L^{\infty}(0,T;L^2(\Omega))$ norm on graded mesh for $\alpha = 0.5$, $\alpha = 0.7$ and $\alpha = 0.9$. It can be seen from this table that using graded mesh, we obtain optimal convergence rate in time direction, as predicted by Theorem 5.3. Tables 5 and 6 confirm the theoretical convergence estimates in L^2 -norm & H_0^1 -norm in spatial direction for $\alpha = 0.5$, $\alpha = 0.7$ and $\alpha = 0.9$.

N	$\alpha = 0.5$		$\alpha = 0.7$		$\alpha = 0.9$	
	Error	OC	Error	OC	Error	OC
2^{3}	5.59E-03	1.8107	4.94E-03	1.9108	4.40E-03	1.9553
2^{4}	1.59E-03	1.9193	1.31E-03	1.9680	1.14E-03	1.9872
2^{5}	4.21E-04	1.9644	3.36E-04	1.9875	2.86E-04	1.9960
2^{6}	1.08E-04	1.9836	8.46E-05	1.9947	7.18E-05	1.9987
2^{7}	2.73E-05	-	2.12E-05	-	1.80E-05	-

Table 4: Error and order of convergence in $L^{\infty}(0,T; L^{2}(\Omega))$ norm in time on graded mesh for Example 6.2.

M_s	$\alpha = 0.5$		$\alpha = 0.7$		$\alpha = 0.9$	
	Error	OC	Error	OC	Error	OC
2^{3}	5.59E-03	1.8107	4.94E-03	1.9108	4.40E-03	1.9553
2^{4}	1.59E-03	1.9193	1.31E-03	1.9680	1.14E-03	1.9872
2^{5}	4.21E-04	1.9644	3.36E-04	1.9875	2.86E-04	1.9960
2^{6}	1.08E-04	1.9836	8.46E-05	1.9947	7.18E-05	1.9987
2^{7}	2.73E-05	-	2.12E-05	-	1.80E-05	-

Table 5: Error and order of convergence in L^2 -norm in space for Example 6.2.

M_s	$\alpha = 0.5$		$\alpha = 0.7$		$\alpha = 0.9$	
	Error	OC	Error	OC	Error	OC
2^3	4.67E-02	1.0359	4.59E-02	1.0232	4.53E-02	1.0120
2^{4}	2.28E-02	1.0171	2.26E-02	1.0080	2.25E-02	1.0035
2^5	1.13E-02	1.0055	1.12E-02	1.0022	1.12E-02	1.0009
2^{6}	5.61E-03	1.0015	5.60E-03	1.0006	5.60E-03	1.0002
2^{7}	2.80E-03	-	2.80E-03	-	2.80E-03	-

Table 6: Error and order of convergence in H_0^1 -norm in space for Example 6.2.

7 Conclusions

In this work, we have proposed $L2\text{-}1_{\sigma}$ Galerkin finite element method for solving timefractional PDE with nonlocal diffusion term. Under certain regularity assumptions on the exact solution, we have derived α -robust *a priori* error estimates in L^2 -norm and H^1 -norm. We also have confirmed our theoretical estimates by numerical experiments. The derivation of regularity assumptions (in Section 5) on the exact solution u is under the investigation and it will be reported in our future work.

Declarations:

Conflict of interest- The author declares no competing interests.

References

- A. Alikhanov, A new difference scheme for the time fractional diffusion equation, J. Comput. Phys., 280 (2015), 424-438.
- [2] Rui M. P. Almeida, José C. M. Duque, Jorge Ferreira, and Rui J. Robalo, The Crank-Nicolson-Galerkin finite element method for a nonlocal parabolic equation with moving boundaries, Numer. Methods Partial Differ. Eq., 31 (5) (2015), 1515-1533.
- [3] S. Chaudhary, V. Srivastava, V. V. K. Srinivas Kumar, and B. Srinivasan, *Finite element approximation of nonlocal parabolic problem*, Numer. Methods Partial Differ. Eq., 33 (2017), 786-313.

- [4] S. Chaudhary, Finite element analysis of nonlocal coupled parabolic problem using Newton's method, Comput. Math. Appl., 75(3) (2018), 981-1003.
- [5] S. Chaudhary, and P. J. Kundaliya, L1 scheme on graded mesh for subdiffusion equation with nonlocal diffusion term, Math. Comput. Simul., 195(2022) 119-137.
- [6] H. Chen, and M. Stynes, Error analysis of a second-order method on fitted meshes for a time fractional diffusion problem, J. Sci. Comput., 79(2) (2019), 624-647.
- [7] H. Chen, and M. Stynes, Blow-up of error estimates in time-fractional initialboundary value problems, IMA J. Numer. Anal., 41(2) (2021), 974-997.
- [8] H. Chen, Y. Wang, and H. Fu, α-robust H1-norm error estimate of nonuniform Alikhanov scheme for fractional sub-diffusion equation, Appl. Math. Lett., 125 (2022), 107771.
- [9] K. Diethelm, The analysis of fractional differential equations: An applicationoriented exposition using differential operators of Caputo type, Lecture Notes in Mathematics, Springer, 2010.
- [10] T. Gudi, Finite element method for a nonlocal problem of Kirchhoff type, SIAM J. Numer. Anal., 50(2) (2012), 657-668.
- [11] C. Huang, and M. Stynes, Optimal spatial H¹-norm analysis of a finite element method for a time-fractional diffusion equation, J. Comput. Appl. Math., 367 (2020), 112435.
- [12] C. Huang, and M. Stynes, α-robust error analysis of a mixed finite element method for a time-fractional biharmonic equation, Numer. Algor., 87 (2021), 1749-1766.
- [13] C. Huang, and M. Stynes, A sharp α -robust $L^{\infty}(H^1)$ error bound for a timefractional Allen-Cahn problem discretised by the Alikhanov L2-1_{σ} scheme and a standard FEM, J. Sci. Comput., 91, Article number: 43 (2022).
- [14] B. Jin, B. Li, and Z. Zhou, Numerical analysis of nonlinear subdiffusion equations, SIAM J. Numer. Anal., 56(1) (2018), 1-23.
- [15] N. Kopteva, Error analysis of the L1 method on graded and uniform meshes for a fractional derivative problem in two and three dimensions, Math. Comp., 8(2019), 2135-2155.

- [16] D. Li, H. Qin, and J. Zhang, Sharp pointwise-in-time error estimate of L1 scheme for nonlinear subdiffusion equations, arXiv:2101.04554v1, (2021).
- [17] H. Liao, W. Mclean, and J. Zhang, A discrete Grönwall inequality with applications to numerical schemes for sub-diffusion problems, SIAM J. Numer. Anal., 57(1) (2019), 218-237.
- [18] H. Liao, W. McLean, and J. Zhang, A second-order scheme with nonuniform time steps for a linear reaction-subdiffusion equation, Commun. Comput. Phys., 30(2) (2021), 567-601.
- [19] J. Manimaran, L. Shangerganesh, A. Debbouche, and V. Antonov, Numerical solutions for time-fractional cancer invasion system with nonlocal diffusion, Front. Phys., 7 (2019), 93.
- [20] J. Manimaran, and L. Shangerganesh, Error estimates for Galerkin finite element approximations of time-fractional nonlocal diffusion equation, Int. J. Comput. Math., 98(7) (2020), 1365-1384.
- [21] S. B. Menezes, *Remarks on weak solutions for a nonlocal parabolic problem*, Int. J. Mathematics and Math. Sci., 2006 (2006), 1-10.
- [22] K. Mustapha, and W. McLean, A second-order accurate numerical method for a fractional wave equation, Numer. Math., 105 (2007), 481-510.
- [23] K. Sakamoto, and M. Yamamoto, Initial value/boundary value problems for fractional diffusion-wave equations and applications to some inverse problems, J. Math. Anal. Appl., 382 (2011), 426-447.
- [24] V. Srivastava, S. Chaudhary, V. V. K. Srinivas Kumar, and B. Srinivasan, Fully discrete finite element scheme for nonlocal parabolic problem involving the Dirichlet energy, J. Appl. Math. Comput., 53 (2017), 413-443.
- [25] M. Stynes, E. O'Riordan, and J. Gracia, Error analysis of a finite difference method on graded meshes for a time-fractional diffusion equation, SIAM J. Numer. Anal., 55 (2017), 1057-1079.
- [26] V. Thomée, Galerkin finite element methods for parabolic problems, 2nd revised and expanded ed., Springer, Berlin, 2006.
- [27] S. Zheng, and M. Chipot, Asymptotic behaviour of solutions to nonlinear parabolic equations with nonlocal terms, Asymptot. Anal., 45(3-4)(2005) 301-312.