

Gauge fields through the Big Bang

Martina Adamo*, Flavio Mercati†

Departamento de Física, Universidad de Burgos, 09001 Burgos, Spain.

June 21, 2023

Abstract

Recent studies have demonstrated the possibility to uphold classical determinism within gravitational singularities, showcasing the ability to uniquely extend Einstein's equations across the singularity in certain symmetry-reduced models. This extension can be achieved by allowing the orientation of spatial hypersurfaces to dynamically change. Furthermore, a crucial aspect of the analysis revolves around the formulation of the dynamical equations in terms of physical degrees of freedom, demonstrating their regularity at the singularity. Remarkably, singular behavior is found to be confined solely to the gauge/unphysical degrees of freedom. This paper extends these results to gravity coupled with Abelian and non-Abelian gauge fields in a symmetry-reduced model (homogeneous anisotropic universe). Near the Big Bang, the dynamics of the geometry and the gauge fields is reformulated in a way that shows that determinism is preserved, assuming a change in orientation at the singularity. The gauge fields are demonstrated to maintain their orientation throughout the singularity, indicating that the predicted orientation change of spatial hypersurfaces holds physical significance. This observation suggests that an observer can discern the specific side of the Big Bang they inhabit.

Contents

1	Introduction	2
2	Hamiltonian formulation of Einstein–Maxwell theory	5
3	Homogeneous <i>ansatz</i>	6
4	Solving the diffeomorphism constraints	7
5	Inevitability of collapse	9
6	One-dimensional <i>ansatz</i>	11
7	Generic Einstein–Maxwell system	19
8	Conclusions	21
A	Einstein–Yang–Mills model under one-dimensional <i>ansatz</i>	26

*marti.adamo@gmail.com

†flavio.mercati@gmail.com

1 Introduction

General relativity (GR) predicts the existence of gravitational singularities: regions of the spacetime manifold where certain physical quantities become meaningless in a coordinate-independent way. In these regions, for example, some components of the stress-energy tensor may diverge, as well as some curvature invariants, or the geodesic equation may be singular (*i.e.*, geodesic incompleteness, as predicted by the Penrose–Hawking singularity theorems [1–5]).

Currently, quantum gravity effects are considered the most promising approach to regularize gravitational singularities [6–9], similar to how QED renders the energy of a point-like electric charge finite, thanks to the uncertainty principle [10]. However, spacetime singularities differ significantly from those in electromagnetism. Unlike the latter, spacetime singularities arise directly from the evolution (via Einstein’s equations) of regular initial data. This makes them physical predictions of the theory, while the diverging energy of a point-like charge is a consequence of the idealization of a point particle, which is introduced manually into the initial conditions.

One of the most remarkable implications of gravitational singularities is the apparent breakdown of determinism. In Lorentzian field theories like GR, classical determinism refers to the ability to uniquely predict the values of the physical fields anywhere within a region of spacetime known as the *causal diamond*, provided that the initial values of these fields on some space-like region are given. The presence of a gravitational singularity appears to violate determinism, making it impossible to predict the values of the fields throughout a future causal cone originating from the singularity. The loss of predictability in GR around these regions can be summarized by Hawking’s words: “One does not know what will come out of a singularity” [11].

Nevertheless, recent works [12–15] have revealed the possibility of preserving determinism in certain symmetry-reduced models that exhibit Big Bang or black hole singularities. Ref. [12] proved that, under a homogeneous but not necessarily isotropic *ansatz*, it is possible to reformulate Einstein’s equations in terms of a set of variables that satisfies a theorem of existence and uniqueness at the singularity. This result has been established in [12–15] for the initial singularity of the *Bianchi-IX model*, a homogeneous non-isotropic universe with an S^3 topology [16, 17], filled with *stiff matter*. The presence of this type of matter source is necessary to regularize the eternal chaotic dynamics that would otherwise occur as the singularity is approached. In the absence of stiff matter, the Bianchi-IX singularity exhibits Misner’s *mixmaster* behavior [16, 17]: as the singularity is approached, the spatial volume goes to zero, while the *shape* degrees of freedom (which measure the anisotropy of the spatial metric) oscillate chaotically. This intricate motion persists indefinitely in coordinate time, with the shape variables oscillating an infinite number of times before reaching the singularity. However, the singularity itself is reached within a finite amount of proper time: the behavior is then that of an *essential* singularity (analogous to $\lim_{x \rightarrow 0} \sin(1/x)$). This essential singularity prevents knowledge of the exact values of all the physical degrees of freedom at the singularity. The presence of stiff matter regularizes this behavior, ensuring that the system enters a final phase of *quiescence*, *i.e.*, a non-chaotic anisotropic collapse described by the *Bianchi-I* (or *Kasner*) model [18]. This condition is indispensable for extending the solution of Einstein’s equation through the singularity since each physical degree of freedom must admit a well-defined limit at the Big Bang.

It is possible to identify a set of physical variables that remain well-defined at the

singularity,¹ enabling the formulation of the equations of motion in a manner consistent with the *Picard–Lindelöf theorem* on the existence and uniqueness of solutions. This implies that the newly introduced regular variables continue to evolve uniquely through the singularity. As the singularity resides at the boundary of the configuration space, this suggests the need to extend the configuration space of GR. In [12], this is achieved by allowing changes in the orientation of space. The interpretation of the regular variables beyond the singularity is as follows: they describe the geometry of spatial hypersurfaces with reversed orientations that lie beyond the Big Bang. Consequently, a “second universe” emerges from the singularity with an inverted spatial orientation.

In [14], it was conjectured that the preservation of determinism is not limited to homogeneous models such as Bianchi IX but is a general characteristic of realistic Big Bang and black hole singularities. Firstly, quiescence, which was a key feature of the original result [12], is not exclusive to models with a stiff matter source. As noted in [14], the Starobinski model, which involves an effective action for gravity that includes the lowest-order quantum corrections to the Einstein–Hilbert action, also exhibits this characteristic (in addition to being the most promising candidate for explaining inflation [24]). Therefore, it can be said that pure (semiclassical) gravity alone, without any matter sources, tends towards a quiescent behavior.

Secondly, the assumption of homogeneity, which allowed the treatability of the models examined thus far, does not seem to be a prerequisite for the continuity results to hold. A strong evidence supporting this idea is provided by the Belinsky–Khalatnikov–Lifshitz (BKL) conjecture [25], which states that as one approaches a space-like singularity, the time derivatives in Einstein’s equations dominate over spatial derivatives, implying that the asymptotic dynamics is described by an (infinite) set of decoupled ordinary differential equations, one for each spatial point. These equations are identical to the equations of motion for a Bianchi-IX universe, and in quiescent models, they exhibit the continuation result under discussion. Interestingly, the BKL conjecture is essentially proven for universes with stiff matter sources [26], providing strong indications that inhomogeneities will not change the result regarding continuation through singularities.

Furthermore, the models examined thus far have only included scalar matter fields, which, as we remarked, can be seen as an effective description of quantum gravitational degrees of freedom, in the case of Starobinski’s model [14]. It is commonly understood that “matter does not matter” near a singularity [8, 27, 28]. In the vicinity of an isotropic FLRW solution, a simple scaling argument shows that the contributions to Friedman’s equations arising from Standard Model matter (a^{-3}), radiation (a^{-4}), the cosmological constant (a^0), and spatial curvature (a^{-2}) are all suppressed compared to the contribution of anisotropic shear, which scales as a^{-6} (where a denotes the FLRW scale factor). Here, by anisotropic shear, we refer to what we later refer to as *shape kinetic energy*, which represents the term analogous to kinetic energy associated with the change in anisotropy parameters (visualize a scenario in which we are in close proximity to an initially isotropic spatial metric that is gradually losing its isotropy). It should be noted that the only exception to this behavior is scalar fields, which contribute to the Friedman equations with terms that scale as a^{-6} . This intuition does not hold when we delve deep into the anisotropic regime [29]. If the pressure of matter sources becomes anisotropic, it can interact in a complex manner with the shape degrees of freedom,

¹See [19–23] for a reformulation of the dynamics of GR as a non-hamiltonian system based on these variables.

and it is not possible to demonstrate that matter or radiation decouples in the same way as in the isotropic regime. The motto “matter does not matter” does not hold in an anisotropic universe, therefore the continuability of Einstein’s equations through the Big Bang needs to be proven separately in presence of matter or radiation fields.

In this paper, our focus is on radiation, specifically electromagnetic and Yang–Mills fields. We aim to provide a comprehensive analysis of their dynamics near a Big Bang singularity in a simplified model, namely, under the assumption of homogeneity. Our first objective is to rigorously prove that the radiation degrees of freedom truly decouple from the gravitational ones (in the sense that they disappear from the equations of motion of the latter), while being driven by their own evolution. This will be the first goal of this paper.

The second goal is to study how the gauge degrees of freedom evolve under the influence of the gravitational ones as we progress through the singularity. This question is intriguing because, although the orientation of spatial slices is reversed upon crossing the singularity, it is not evident whether this reversal can be observed, for example by means of parity-breaking tests like beta decay [30]. It remains uncertain whether such tests could determine the side of the singularity we find ourselves on. For this, we need to know what happens to gauge fields and fermions, whether for example their direction is flipped or not. This paper takes the first step towards addressing this question by analyzing the behavior of the gauge fields.

Our results suggest that gauge fields do not reverse their direction across the singularity, although we cannot prove this yet in a fully general context. Our analysis is restricted to homogeneous gauge fields, and furthermore, it applies in full generality only to Abelian gauge groups. In the non-Abelian case, our analysis is limited to a “one-dimensional” *ansatz*, meaning that both the gauge vector potential and its conjugate momentum are assumed to point in the same spatial direction throughout the evolution. The relaxation of these assumptions will be the focus of future investigations.

The paper is structured as follows: Section 2 provides a review of the Hamiltonian formulation of the Einstein–Maxwell system. Sections 3 and 4 focus on the phase-space reduction to the homogeneous case, assuming a spatial topology of a three-sphere and the invariance under translations of both the metric and the gauge fields throughout the evolution. This *ansatz* is compatible with the Hamiltonian evolution and reduces the degrees of freedom to a finite set, whose equations of motion are ordinary differential equations in time.

Section 5 introduces the Misner variables commonly used to discuss homogeneous universes with a three-sphere topology (Bianchi-IX models) and demonstrates the inevitability of the singularity even in the presence of gauge fields. In Section 6, a further simplification is introduced through the one-dimensional *ansatz* for the gauge fields discussed earlier. Under this *ansatz*, the continuation result can be (relatively) easily proven.

Section 7 considers the relaxation of the one-dimensional *ansatz* for Abelian gauge fields and the continuation result is proven. The extension of this result to $SU(2)$ and $SU(3)$ gauge fields under the one-dimensional *ansatz* is detailed in Appendix A. However, the relaxation of this *ansatz* in non-Abelian gauge theories goes beyond the scope of the present paper. Finally, in Section 8, we draw conclusions based on the knowledge gained thus far. In Table 1 we summarize the notations used in the paper.

Greek indices are spacetime indices	$\mu, \nu, \rho, \dots \in \{0, 1, 2, 3\}$
Latin lowercase indices from the end of the alphabet are spatial indices	$i, j, k, \dots \in \{1, 2, 3\}$
Latin lowercase indices from the beginning of the alphabet are internal/dreibein indices	$a, b, c, \dots \in \{1, 2, 3\}$
Latin uppercase indices are internal $SO(N)$ indices	$I, J, K, \dots \in \{1, 2, \dots, N^2 - 1\}$

Table 1: Notations used in the paper.

2 Hamiltonian formulation of Einstein–Maxwell theory

Our goal is to extend the model of [12, 14] to the case of the Einstein–Maxwell system: GR minimally coupled with electromagnetism, whose action is given by

$$S = \int d^4x \sqrt{-h} \left(R - \frac{1}{4} h^{\mu\nu} h^{\rho\sigma} F_{\mu\rho} F_{\nu\sigma} \right), \quad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu. \quad (1)$$

In the Arnowitt–Deser–Misner Hamiltonian formalism [31], the four-dimensional metric $h_{\mu\nu}$ is split into its spatial components g_{ij} , which serve as canonical variables, and four other fields: the *lapse* scalar N and the *shift* vector N^i ,² which are Lagrange multipliers because their time derivatives do not appear in the action. The relations between these quantities and the spacetime metric components are given by $h_{ij} = g_{ij}$, $h_{00} = -N^2 + N_i N^i$, and $h_{0i} = h_{i0} = N_i$.

These quantities have the following physical interpretations: the spatial components g_{ij} represent the three-dimensional metric of equal-time hypersurfaces, the shift generates infinitesimal spatial translations along the hypersurfaces, and the lapse represents the proper time measured by observers moving orthogonally between neighboring hypersurfaces. The time derivatives of g_{ij} are replaced, through a Legendre transform, by the conjugate momenta $\pi^{ij} = (\sqrt{g}/2N) (g^{ik} g^{jl} - g^{ij} g^{kl}) (\dot{g}_{kl} - \mathcal{L}_{\vec{N}} g_{kl})$, where g^{ij} is the inverse matrix of g_{ij} , and $\mathcal{L}_{\vec{N}}$ is the Lie derivative w.r.t. the shift N^i .

The Hamiltonian decomposition of the electromagnetic action is similar to the familiar one in Minkowski spacetime: the canonical variables are the spatial components of the electromagnetic potential A_i , while the time component A_0 (which is the scalar potential) acts as a fifth Lagrange multiplier. The spatial components of the Faraday tensor are given by $F_{ij} = \partial_i A_j - \partial_j A_i$. Finally, the momenta canonically conjugate to A_i are the components of the electric field $E^i = (\sqrt{g}/N) g^{ij} (F_{0j} - N^k F_{kj})$.

The following equal-time Poisson-bracket relations hold for conjugate pairs of variables:

$$\begin{aligned} \{g_{ij}(x), \pi^{kl}(y)\} &= \frac{1}{2} (\delta_i^k \delta_j^l + \delta_i^l \delta_j^k) \delta^{(3)}(x - y), \\ \{A_i(x), E^j(y)\} &= \delta_i^j \delta^{(3)}(x - y), \end{aligned} \quad (2)$$

while all other brackets are zero. The time evolution is given by the total Hamiltonian,

²The spatial metric, shift, and lapse are a two-tensor, a vector and a scalar field, respectively, under diffeomorphisms of the spatial hypersurface.

which for our system is a linear combination of the following constraints:

$$\begin{aligned}\mathcal{H}[N] &= \int d^3x N \left(\frac{1}{\sqrt{g}} \left(\pi^{ij} \pi_{ij} - \frac{1}{2} \pi^2 + \frac{1}{2} g_{ij} E^i E^j \right) + \sqrt{g} \left(\frac{1}{4} g^{ij} g^{kl} F_{ik} F_{jl} - K \right) \right), \\ \mathcal{H}_i[N^i] &= \int d^3x N^i \left(E^j F_{ij} - 2 g_{ij} \nabla_k \pi^{jk} \right), \\ \mathcal{G}[A_0] &= - \int d^3x A_0 \left(\nabla_i E^i \right),\end{aligned}\tag{3}$$

where K is the Ricci scalar and ∇_i is the covariant derivative, both w.r.t. the metric g_{ij} . These constraints are first-class, and close an extension of the so-called *hypersurface deformation algebra* [32, 33] (or rather *algebroid* [34]). The first and the last lines in Eq. (3) represent the Hamiltonian and Gauss constraints, which generate time evolution and electromagnetic gauge transformations respectively. The three constraints \mathcal{H}_i can be expressed (up to boundary terms) as a linear combination of Gauss and diffeomorphism constraints:

$$\mathcal{H}_i[N^i] = \mathcal{D}_i[N^i] - \mathcal{G}[A_i N^i] + (\text{boundary terms}),\tag{4}$$

where:

$$\mathcal{D}_i[N^i] = \int d^3x \left(E^i \mathcal{L}_{\vec{N}} A_i + \pi^{ij} \mathcal{L}_{\vec{N}} g_{ij} \right).\tag{5}$$

3 Homogeneous *ansatz*

We now impose the homogeneous *ansatz*, which has been the starting point of previous works such as [12, 14]. The most general spatially-homogeneous universe with the topology of a three-sphere is described by the Bianchi-IX model [16, 17, 35]. The spatial metric is assumed to have three independent Killing vectors that generate spatial translations (homogeneity). On S^3 , coordinatized by the usual hyperspherical coordinates $\theta \in [0, \pi]$, $\phi \in [0, \pi]$, $\psi \in [0, 2\pi)$, it is possible to construct a basis of vector fields that are invariant under these translations, as well as a dual basis of one-forms:

$$\begin{cases} \chi_1 = -\sin \psi \partial_\theta + \cos \psi \csc \theta \partial_\phi - \cos \psi \cot \theta \partial_\psi \\ \chi_2 = -\cos \psi \partial_\theta - \sin \psi \csc \theta \partial_\phi + \sin \psi \cot \theta \partial_\psi \\ \chi_3 = \partial_\psi \end{cases}, \quad \begin{cases} \sigma^1 = -\sin \psi d\theta + \cos \psi \sin \theta d\phi \\ \sigma^2 = -\cos \psi d\theta - \sin \psi \sin \theta d\phi \\ \sigma^3 = d\psi + \cos \theta d\phi \end{cases},\tag{6}$$

where the duality is given by

$$\sigma_i^a \chi_a^j = \delta_i^j, \quad \sigma_i^a \chi_b^i = \delta_b^a.\tag{7}$$

The most generic homogeneous (but not necessarily isotropic) metric on S^3 can be expressed as a quadratic form (with spatially constant coefficients) in this basis. By imposing the homogeneous *ansatz* also on the conjugate momenta p^{ij} (which is necessary to preserve the homogeneous *ansatz* for the metric under time evolution), as well as on the electromagnetic fields A_i and E^i , we obtain:

$$\begin{aligned}g_{ij}(t, x) &= q_{ab}(t) \sigma_i^a(x) \sigma_j^b(x), \quad \pi^{ij}(t, x) = p^{ab}(t) \chi_a^i(x) \chi_b^j(x) \det \sigma(x), \\ A_i(t, x) &= A_a(t) \sigma_i^a(x), \quad E^i(t, x) = E^a(t) \chi_a^i(x) \det \sigma(x),\end{aligned}\tag{8}$$

$$F_{ij}(t, x) = F_{bc}(t) \sigma_i^b(x) \sigma_j^c(x) = -A_a(t) \delta^{ad} \varepsilon_{dbc} \sigma_i^b(x) \sigma_j^c(x).$$

Notice that the conjugate momenta p^{ij} and E^i require a term $\det \sigma(x) = \sin \theta$ to ensure the correct transformation behavior under diffeomorphisms (that of a tensor density of weight +1). In this basis, all these tensor fields have homogeneous components: q_{ab} , p^{ab} , A_a , E^a , with only a time dependence. Under this *ansatz*, all the constraints (each of which constrains one degree of freedom per spatial point) can be smeared over arbitrary functions and become the following global constraints:

$$\begin{aligned}\mathcal{H}[N] &= n \left(p^{ab} p^{cd} q_{bc} q_{da} - \frac{1}{2} (p^{ab} q_{ab})^2 + q_{ab} q_{cd} \delta^{bc} \delta^{da} - \frac{1}{2} (q_{ab} \delta^{ab})^2 \right. \\ &\quad \left. + \frac{1}{2} q_{ab} E^a E^b + \frac{1}{4} \det q q^{ab} q^{cd} F_{ac} F_{bd} \right), \\ \mathcal{H}_i[N^i] &= n^d \left(E^a F_{da} + 2 p^{ab} q_{ac} \varepsilon_{bdf} \delta^{fc} \right) = \mathcal{D}_i[N^i],\end{aligned}\tag{9}$$

$$\mathcal{G}[\varphi] = 0,$$

where:

$$n = \frac{1}{\sqrt{\det q}} \int d\theta d\phi d\psi \sin \theta N(x), \quad n^a = \int d\theta d\phi d\psi \sin \theta \sigma_i^a(x) N^i(x), \tag{10}$$

are four leftover Lagrange multipliers (the spatial averages of the lapse and shift). Notice that the Gauss constraint is automatically solved by the homogeneous *ansatz*.

The homogeneous *ansatz* is dynamically consistent, *i.e.*, it is preserved by the evolution [16, 17], as can be verified by substituting the *ansatz* into the RHS of the Einstein equations.

4 Solving the diffeomorphism constraints

In order to eliminate the non-physical degrees of freedom, we need to gauge-fix the three diffeomorphism constraints using Dirac's procedure for constrained Hamiltonian systems [36–38]. The diffeomorphism generators are:

$$\begin{cases} \xi_1 = \mathcal{D}_1 = 2 (p^{13} q_{12} - p^{12} q_{13} + p^{23} q_2 - p^2 q_{23} + p^3 q_{23} - p^{23} q_3) + E^3 A_2 - E^2 A_3 \approx 0, \\ \xi_2 = \mathcal{D}_2 = 2 (p^1 q_{13} - p^{13} q_1 + p^{12} q_{23} - p^{23} q_{12} + p^{13} q_3 - p^3 q_{13}) + E^1 A_3 - E^3 A_1 \approx 0, \\ \xi_3 = \mathcal{D}_3 = 2 (p^2 q_{12} - p^{12} q_2 + p^{23} q_{13} - p^{13} q_2 + p^{12} q_1 - p^1 q_{12}) + E^2 A_1 - E^1 A_2 \approx 0. \end{cases}\tag{11}$$

A suitable choice [14, 33] for the gauge-fixing constraints is:

$$\xi_4 = q_{23} \approx 0, \quad \xi_5 = q_{13} \approx 0, \quad \xi_6 = q_{12} \approx 0.\tag{12}$$

These gauge-fixing constraints are second-class w.r.t. ξ_1, ξ_2, ξ_3 everywhere except at the three planes of symmetry $q_1 = q_2$, $q_2 = q_3$ and $q_3 = q_1$. Except for a measure-zero set of solutions that takes entirely place on these planes, all solutions that intersect these planes can be uniquely continued through them by making a different local choice of gauge-fixing. We can solve the diffeomorphism constraints w.r.t. the non-diagonal components of p^{ab} , and this choice is regular everywhere away from the three symmetry

planes:

$$\begin{cases} p^{23} = \frac{E^2 A_3 - E^3 A_2}{2(q_2 - q_3)}, \\ p^{13} = \frac{E^3 A_1 - E^1 A_3}{2(q_3 - q_1)}, \\ p^{12} = \frac{E^1 A_2 - E^2 A_1}{2(q_1 - q_2)}. \end{cases} \quad (13)$$

Using the six second-class constraints ξ_α , $\alpha = 1, \dots, 6$, we can construct the Dirac matrix, which, when evaluated on the constraints hypersurface in the space of solutions of the system (*i.e.*, on-shell), reads:

$$C_{\alpha\beta} = \{\xi_\alpha, \xi_\beta\} \approx \left(\begin{array}{c|c} 0 & M \\ \hline -M & 0 \end{array} \right), \quad (14)$$

where $M = \text{diag}(q_3 - q_2, q_1 - q_3, q_2 - q_1)$. The inverse Dirac matrix is then simply:

$$(C^{-1})^{\alpha\beta} \approx \left(\begin{array}{c|c} 0 & M^{-1} \\ \hline -M^{-1} & 0 \end{array} \right). \quad (15)$$

Therefore, the Dirac bracket:

$$\{f, g\}_* = \{f, g\} - \{f, \xi_\alpha\} (C^{-1})^{\alpha\beta} \{\xi_\beta, g\}, \quad (16)$$

is canonical on the diagonal components of the metric and their momenta, and on the three components of the electromagnetic potential and their momenta:

$$\{q_a, p^b\}_* = \delta_a^b, \quad \{A_a, E^b\}_* = \delta_a^b, \quad (17)$$

and all the other Dirac brackets are zero.

We started with a system described by eighteen (not all physical) degrees of freedom: six components of the symmetric three-dimensional metric q_{ab} , six components of metric momenta p^{ab} , three components of the electromagnetic potential A_a , and three of the electromagnetic momenta E_a (the electric field). After the diffeomorphism gauge-fixing, we fixed six degrees of freedom: the three off-diagonal components of the metric q_{12} , q_{23} , q_{13} (which are set to zero), and the three off-diagonal components of the metric momenta p^{12} , p^{23} , p^{13} (which are now functions of all the other variables). Therefore, we end up with twelve degrees of freedom. Of these, ten are truly physical, in the sense that they are the minimum number of independent variables necessary to uniquely determine a solution. The remaining two are constrained by the Hamiltonian constraint (first equation in (9)) and its gauge-fixing (*i.e.*, the fact that we can freely choose initial conditions among the different points of the solution curve without changing the solution itself).

The equations of motion generated by the Dirac bracket are the canonical equations of motion obtained from the on-shell Hamiltonian:

$$\begin{aligned} \mathcal{H}[N] = n \left[\mathcal{H}_{BIX} + \frac{q_2 q_3 (M_1)^2}{2(q_2 - q_3)^2} + \frac{q_1 q_3 (M_2)^2}{2(q_1 - q_3)^2} + \frac{q_1 q_2 (M_3)^2}{2(q_1 - q_2)^2} \right. \\ \left. + \frac{1}{2} q_1 ((E^1)^2 + (A_1)^2) + \frac{1}{2} q_2 ((E^2)^2 + (A_2)^2) + \frac{1}{2} q_3 ((E^3)^2 + (A_3)^2) \right], \end{aligned} \quad (18)$$

where we called $q_{11} = q_1$, $q_{22} = q_2$, $q_{33} = q_3$, $p^{11} = p^1$, $p^{22} = p^2$, $p^{33} = p^3$, and:

$$M_1 = E^2 A_3 - E^3 A_2, \quad M_2 = E^3 A_1 - E^1 A_3, \quad M_3 = E^1 A_2 - E^2 A_1, \quad (19)$$

and where:

$$\mathcal{H}_{BIX} = (p^1 q_1)^2 + (p^2 q_2)^2 + (p^3 q_3)^2 - \frac{(p^3 q_3 + p^2 q_2 + p^1 q_1)^2}{2} + q_1^2 + q_2^2 + q_3^2 - \frac{(q_1 + q_2 + q_3)^2}{2}, \quad (20)$$

is the Hamiltonian constraint of an empty Bianchi-IX universe [14].

5 Inevitability of collapse

With the following canonical transformation:

$$\begin{cases} q_1 = a_0^2 \exp\left(\frac{x^0 - \sqrt{3}x^1 + x^2}{\sqrt{3}}\right), \\ q_2 = a_0^2 \exp\left(\frac{x^0 + \sqrt{3}x^1 + x^2}{\sqrt{3}}\right), \\ q_3 = a_0^2 \exp\left(\frac{x^0 - 2x^2}{\sqrt{3}}\right), \end{cases} \quad \begin{cases} p^1 = a_0^{-2} \left(\frac{k_2 - \sqrt{3}k_1 + 2k_0}{2\sqrt{3}}\right) \exp\left(-\frac{x^0 - \sqrt{3}x^1 + x^2}{\sqrt{3}}\right), \\ p^2 = a_0^{-2} \left(\frac{k_2 + \sqrt{3}k_1 + 2k_0}{2\sqrt{3}}\right) \exp\left(-\frac{x^0 + \sqrt{3}x^1 + x^2}{\sqrt{3}}\right), \\ p^3 = a_0^{-2} \left(\frac{k_0 - k_2}{\sqrt{3}}\right) \exp\left(-\frac{x^0 - 2x^2}{\sqrt{3}}\right), \end{cases} \quad (21)$$

where a_0 is a dimensional constant (a reference scale), the Hamiltonian constraint takes the form of a diagonal quadratic kinetic term for the metric variables k_a , plus a potential-like term that depends on the metric and electromagnetic variables:

$$\mathcal{H}[N] = n \left[\frac{1}{2} (-k_0^2 + k_1^2 + k_2^2) + \frac{1}{2} U(x, A, E) \right], \quad (22)$$

$$U(x, A, E) = a_0^4 e^{\frac{2x^0}{\sqrt{3}}} C(x^1, x^2) + a_0^2 e^{\frac{x^0}{\sqrt{3}}} V(x^1, x^2, A, E) + W(x^1, x^2, A, E).$$

In the previous equation:

$$\begin{aligned} C(x^1, x^2) = & e^{-2x^1 + \frac{2}{\sqrt{3}}x^2} + e^{2x^1 + \frac{2}{\sqrt{3}}x^2} + e^{-\frac{4}{\sqrt{3}}x^2} \\ & - 2 \left(e^{\frac{2}{\sqrt{3}}x^2} + e^{-x^1 - \frac{1}{\sqrt{3}}x^2} + e^{x^1 - \frac{1}{\sqrt{3}}x^2} \right), \end{aligned} \quad (23)$$

is the Bianchi-IX potential [14, 33], and

$$\begin{aligned} V(x^1, x^2, A, E) = & e^{-x^1 + \frac{x^2}{\sqrt{3}}} ((E^1)^2 + (A_1)^2) \\ & + e^{x^1 + \frac{x^2}{\sqrt{3}}} ((E^2)^2 + (A_2)^2) + e^{-\frac{2x^2}{\sqrt{3}}} ((E^3)^2 + (A_3)^2), \end{aligned} \quad (24)$$

$$W(x^1, x^2, A, E) = \frac{e^{x^1 + \sqrt{3}x^2} (M_1)^2}{(e^{x^1 + \sqrt{3}x^2} - 1)^2} + \frac{e^{x^1 + \sqrt{3}x^2} (M_2)^2}{(e^{x^1} - e^{\sqrt{3}x^2})^2} + \frac{e^{2x^1} (M_3)^2}{(e^{2x^1} - 1)^2},$$

are two new contributions depending on the electromagnetic field.

The variables x^1 and x^2 represent the *shape* degrees of freedom, which quantify the anisotropy of the spatial metric. Their conjugate momenta, k_1 and k_2 , correspond to the rate of change of these anisotropies. The variable x^0 and its conjugate momentum k_0 are related to the volume of the universe v and its conjugate momentum τ , known as the *York time*. These relationships are expressed as follows:

$$v = a_0^3 e^{\frac{\sqrt{3}}{2}x^0}, \quad \tau = \frac{2}{\sqrt{3}} a_0^{-3} e^{-\frac{\sqrt{3}}{2}x^0} k_0. \quad (25)$$

The variable τ , which is named after James York [39], is associated with a specific foliation of spacetime known as *CMC* (*constant-mean extrinsic curvature*). In this foliation, the initial-value problem can be formulated as a system of elliptic equations, whose solution exists and is unique. In a cosmological setting, τ is proportional to (minus) the Hubble parameter [33]. The adoption of this foliation is motivated by the fact that the physical degrees of freedom of GR are spatial-conformal invariants, leading to the proposal of reformulating GR as a three-dimensional conformal field theory known as *Shape Dynamics* [33, 40]. The concepts discussed in the present paper, such as the identification of the shape degrees of freedom as the physical ones, are compatible with the principles of Shape Dynamics, although it does not rely on the shape dynamical interpretation of GR. Therefore, one can view this paper as a result in Shape Dynamics or as an entirely independent result within Hamiltonian GR.

Consider now the equations of motion for x^0 and k_0 :

$$\dot{x}^0 = -n k_0, \quad \dot{k}_0 = -n \left(\frac{1}{\sqrt{3}} a_0^4 e^{\frac{2x^0}{\sqrt{3}}} C + \frac{1}{2\sqrt{3}} a_0^2 e^{\frac{x^0}{\sqrt{3}}} V \right), \quad (26)$$

and assuming, without loss of generality, $n = 1$ and $a_0 = 1$, we can use these equations to calculate the second time derivative of the quantity $\exp(-x^0/\sqrt{3})$, which is a certain power of the volume:

$$\begin{aligned} \frac{d^2}{dt^2} \left(e^{-\frac{x^0}{\sqrt{3}}} \right) &= \frac{d}{dt} \left(e^{-\frac{x^0}{\sqrt{3}}} k_0 \right) = -\frac{1}{3} e^{-\frac{x^0}{\sqrt{3}}} \left(-k_0^2 + e^{\frac{2x^0}{\sqrt{3}}} C \right) - \frac{1}{6} V \\ &\approx \frac{1}{3} e^{-\frac{x^0}{\sqrt{3}}} (k_1^2 + k_2^2 + W) + \frac{1}{6} V, \end{aligned} \quad (27)$$

where we used the Hamiltonian constraint in the last step. The RHS is non-negative, because both V and W are positive-definite. Thus, we have proven that the quantity $\exp(-x^0/\sqrt{3})$ is concave upwards. Consequently, it will monotonically decrease for half of each solution, reaching a single minimum (which may potentially be infinitely far in time, resulting in strictly increasing or decreasing behavior), and then it will monotonically increase for the rest of the solution.

The volume, given by the (square root of the) inverse of this quantity, generally will monotonically increase for half of each solution, will reach a maximum, and subsequently will monotonically decrease to zero. As remarked above, there may also exist degenerate solutions that undergo either monotonic growth or shrinking, reaching maximal expansion only as $t \rightarrow +\infty$ (or $t \rightarrow -\infty$) while exhibiting a single Big Bang singularity as $t \rightarrow -\infty$ (or $t \rightarrow +\infty$). Our focus in this paper is solely on the behavior of the system near one singularity, while the behavior far away from it, where matter fields, cosmological constant terms, and inhomogeneities dominate the dynamics, does not concern us.

The results above represent a cosmological reformulation of the Penrose–Hawking singularity theorems [1–5]. According to these theorems, once a solution begins to collapse, it cannot be halted and will continue to shrink until it reaches a singularity. It is important to note that, although the Big Bang is only reached as $t \rightarrow \pm\infty$, this does not mean that it is in the infinite future (or past). In fact, a finite amount of proper time elapses between any finite value of t and $t \rightarrow \pm\infty$, as proved in [14].

6 One-dimensional *ansatz*

In this section, we consider a simpler but illustrative case in which the electromagnetic field has only one spatial component:

$$A_1 = A_2 = 0, \quad E^1 = E^2 = 0. \quad (28)$$

These conditions are preserved by the equations of motion:

$$\begin{aligned} \{A_1, \mathcal{H}[N]\}_* \Big|_{\substack{A_1=A_2=0 \\ E^1=E^2=0}} &\approx 0, & \{A_2, \mathcal{H}[N]\}_* \Big|_{\substack{A_1=A_2=0 \\ E^1=E^2=0}} &\approx 0, \\ \{E_1, \mathcal{H}[N]\}_* \Big|_{\substack{A_1=A_2=0 \\ E^1=E^2=0}} &\approx 0, & \{E_2, \mathcal{H}[N]\}_* \Big|_{\substack{A_1=A_2=0 \\ E^1=E^2=0}} &\approx 0, \end{aligned} \quad (29)$$

$$\begin{aligned} \{A_1, \mathcal{G}[\varphi]\}_* \Big|_{\substack{A_1=A_2=0 \\ E^1=E^2=0}} &= 0, & \{A_2, \mathcal{G}[\varphi]\}_* \Big|_{\substack{A_1=A_2=0 \\ E^1=E^2=0}} &= 0, \\ \{E_1, \mathcal{G}[\varphi]\}_* \Big|_{\substack{A_1=A_2=0 \\ E^1=E^2=0}} &= 0, & \{E_2, \mathcal{G}[\varphi]\}_* \Big|_{\substack{A_1=A_2=0 \\ E^1=E^2=0}} &= 0. \end{aligned} \quad (30)$$

In our *ansatz*, we arbitrarily chose to keep the third component as the non-zero one, but this choice does not make the model lose any generality. In fact, if we were to choose the first or second component of A_a and E^a as the non-zero one in our *ansatz*, the dynamics would remain identical, with the labels for the first, second and third components permuted accordingly. This can be proven by performing a simultaneous reflection transformation, such as $q_1 \rightarrow q_2$, $q_3 \rightarrow q_1$, $q_2 \rightarrow q_3$ and likewise for p^a . This transformation does not change the Bianchi-IX part of the Hamiltonian constraint: this is due to a discrete symmetry of our system, which remains invariant under the aforementioned reflection transformations.

The Hamiltonian constraint in Eq. (18) now reads:

$$\overline{\mathcal{H}}[N] = \mathcal{H}[N] \Big|_{\substack{A_1=A_2=0 \\ E^1=E^2=0}} = n \left[\mathcal{H}_{BIX} + \frac{1}{2} q_3 ((E^3)^2 + (A_3)^2) \right]. \quad (31)$$

If we now consider the following quantity:

$$H_{HO}^{1D} = (E^3)^2 + (A_3)^2, \quad (32)$$

it is immediate to prove that it is conserved, because it is first-class w.r.t. the Hamiltonian constraint:

$$\{H_{HO}^{1D}, \overline{\mathcal{H}}[N]\}_* \approx 0, \quad (33)$$

therefore, we can assign a constant of motion ε to it, which will remain unchanged along the whole solution. As a result, the geometric degrees of freedom will evolve according to the following effective Hamiltonian constraint:

$$\mathcal{H}_{eff}[N] = n \left[\mathcal{H}_{BIX} + \frac{1}{2} q_3 \varepsilon \right]. \quad (34)$$

We can combine the new term with the potential term present in \mathcal{H}_{BIX} . For any finite value of ε , we can describe the dynamics of the geometrical degrees of freedom as being controlled by an effective potential given by:

$$U_{1D} = q_1^2 + q_2^2 + q_3^2 - \frac{(q_1+q_2+q_3)^2}{2} + \frac{1}{2} q_3 \varepsilon. \quad (35)$$

Now let us demonstrate that the additional term in the potential does not alter the result regarding the continuation through the singularity.

1 Quiescence is unchanged. The structure of the Hamiltonian constraint resembles that of Bianchi IX, although with a deformed potential. The solutions exhibit similar characteristics to those of Bianchi IX: stretches of inertial motion known as *Kasner epochs* when the potential term is negligible, interrupted by brief quasi-elastic bounces referred to as *Taub transitions* that dissipate some of the shape momenta k_1, k_2 [33].

During a Kasner epoch, the dynamics can be well approximated by that of a free particle. In these phases, both the shape degrees of freedom x^i and the scale degree of freedom x^0 evolve linearly w.r.t. parameter time t , so the spatial volume v decreases exponentially. Notice that the proper time s measured by a comoving observer is exponentially related to t (s is proportional to the t -integral of the volume). Therefore, if a Kasner epoch were to extend all the way to the singularity at $t \rightarrow +\infty$, only a finite amount of proper time would have elapsed [14]. Conversely, in a Taub transition, the configuration point bounces against the Bianchi-IX potential, leading to rapid changes in certain shape variables (such as the direction of motion in configuration space), while x^0 undergoes rapid changes in speed (*i.e.*, its conjugate momentum varies rapidly), but not significantly in magnitude. The resulting motion for x^0 is that of a segmented curve, with periods of straight-line motion separated by rapid changes in slope. Proper time remains finite all the way to the singularity. The Big Bang is thus reached within a finite amount of proper time, but the system undergoes an infinite number of Taub transitions. This chaotic behavior prevents certain degrees of freedom from having a well-defined limit at the singularity.

For instance, consider the angular variable of the polar coordinates of the (x^1, x^2) plane. If a Kasner epoch were to extend to the singularity, this variable would settle into a limiting value. However, each Taub transition makes it change again. If we were to plot its value against proper time, near the singularity it would resemble something similar to the function $\sin(1/x)$ as $x \rightarrow 0$, exhibiting an essential singularity. Consequently, it is impossible to determine the specific value this variable takes at the Big Bang. This prevents any attempt to continue these solutions though it [14, 33].

If we introduce a scalar field, we can induce a state of *quiescence*, meaning that the chaotic behavior stops after a finite number of Taub bounces, and the solution settles onto a last Kasner epoch lasting all the way to the Big Bang. However, the additional term in the potential (35) could, in principle, change the conditions for quiescence. This is not the case for potential terms that are polynomial in the metric components q_1, q_2, q_3 , as the one we have under the one-dimensional *ansatz*. This is easily proven by considering the following scenario: let us assume that we begin during a Kasner epoch. The solution takes the following form:

$$x^\alpha(t) = \eta^{\alpha\beta} k_\beta t + x^\alpha(0), \quad k_\alpha(t) = v_\alpha, \quad v_0 = +\sqrt{(v_1)^2 + (v_2)^2}, \quad (36)$$

where $\eta^{\alpha\beta} = \text{diag}(-1, 1, 1)$, $\alpha, \beta = 0, 1, 2$. The plus sign in the dispersion relation for the integration constants v_α has been chosen so that the Big Bang singularity occurs at $t \rightarrow +\infty$. Replacing the solution into the metric components (21), we obtain:

$$\begin{aligned} q_1(t) &= a_0^2 \exp\left(\frac{x^0(t) - \sqrt{3}x^1(t) + x^2(t)}{\sqrt{3}}\right) \propto \exp\left(-\frac{\sqrt{(v_1)^2 + (v_2)^2} + \sqrt{3}v_1 - v_2}{\sqrt{3}}t\right) = e^{-\rho_1 t}, \\ q_2(t) &= a_0^2 \exp\left(\frac{x^0(t) + \sqrt{3}x^1(t) + x^2(t)}{\sqrt{3}}\right) \propto \exp\left(-\frac{\sqrt{(v_1)^2 + (v_2)^2} - \sqrt{3}v_1 - v_2}{\sqrt{3}}t\right) = e^{-\rho_2 t}, \\ q_3(t) &= a_0^2 \exp\left(\frac{x^0(t) - 2x^2(t)}{\sqrt{3}}\right) \propto \exp\left(-\frac{\sqrt{(v_1)^2 + (v_2)^2} + 2v_2}{\sqrt{3}}t\right) = e^{-\rho_3 t}. \end{aligned} \quad (37)$$

In polar coordinates $(v_1, v_2) = |\vec{v}|(\cos \varphi, \sin \varphi)$, the three coefficients ρ_a appearing in the equations above can be expressed as follows:

$$\begin{cases} \rho_1 = \frac{|\vec{v}|}{\sqrt{3}} \left(1 + \sqrt{3} \cos \varphi - \sin \varphi \right) , \\ \rho_2 = \frac{|\vec{v}|}{\sqrt{3}} \left(1 - \sqrt{3} \cos \varphi - \sin \varphi \right) , \\ \rho_3 = \frac{|\vec{v}|}{\sqrt{3}} (1 + 2 \sin \varphi) , \end{cases} \quad (38)$$

and, for any value of φ , one of the coefficients ρ_a is always negative (except for the three special directions along the symmetry axes of the potential, $\varphi = \frac{\pi}{2}, \frac{7\pi}{6}, \frac{11\pi}{6}$, which, however, only concern a measure-zero set of solutions). This can be observed in Fig. 1.




Figure 1: Plot of the three coefficients ρ_a vs. the direction φ .

Now, if we introduce a homogeneous scalar field without mass nor potential, the Hamiltonian constraint (34) changes into:

$$\mathcal{H}_{eff}[N] = n \left[\mathcal{H}_{BIX} + \frac{1}{2} q_3 \varepsilon + \frac{1}{2} k_3^2 \right] , \quad (39)$$

where k_3 is the conjugate momentum to a homogeneous scalar field, which we call x^3 . A Kasner epoch in this case looks exactly the same, with the difference that the dispersion relation appearing in (36) now looks like:

$$v_0 = \sqrt{(v_1)^2 + (v_2)^2 + (v_3)^2} , \quad (40)$$

where the constant of motion v_3 is the (conserved) value of k_3 . Now, Eqs. (37) take the same form, except that the ρ_a coefficients change into:

$$\begin{cases} \rho_1(w) = \frac{|\vec{v}|}{\sqrt{3}} \left(w + \sqrt{3} \cos \varphi - \sin \varphi \right) , \\ \rho_2(w) = \frac{|\vec{v}|}{\sqrt{3}} \left(w - \sqrt{3} \cos \varphi - \sin \varphi \right) , \\ \rho_3(w) = \frac{|\vec{v}|}{\sqrt{3}} (w + 2 \sin \varphi) , \end{cases} \quad w = \sqrt{1 + \frac{(v_3)^2}{(v_1)^2 + (v_2)^2}} . \quad (41)$$

The parameter w takes value 1 when $v_3 = 0$ (no scalar field), and $w > 1$ when $v_3 \neq 0$. Each Taub transition ends in a new Kasner epoch with a lower value of $(v_1)^2 + (v_2)^2$ (see [14, 33] for the proof), so the parameter w progressively grows larger after each bounce. When it reaches values equal to or larger than $w = 2$, all the $\rho_a(w)$ functions become positive everywhere. We reach a situation in which all the terms in any potential that is polynomial in q_a can only decrease with time. The solution settles with increasing

accuracy around a single Kasner epoch all the way to the singularity, without further Taub bounces.

As we mentioned earlier, the effective potential of the one-dimensional model (35) is polynomial in q_a (it includes quadratic terms from the Bianchi-IX part and a linear term in q_3). Therefore, the conditions for quiescence remains completely unchanged.

However, it is important to notice that the polynomiality of the potential is not guaranteed in general. From Eq. (18), we can observe that in the general case where the electromagnetic field has more than one spatial component, there are non-polynomial terms, such as $q_1 q_2 / (q_1 - q_2)$, and so on.

2 Continuing the dynamics through the singularity. In the previous Paragraph, we have proved that the presence of a one-dimensional electromagnetic field does not alter the quiescent behavior as the system approaches the Big Bang. This provides the foundation for extending the continuation result of [12, 14] to GR minimally coupled with electromagnetism under the one-dimensional *ansatz*.

It is important to note that the variables $x^0, x^1, x^2, x^3, k_0, k_1, k_2, k_3$ are not a suitable set for describing the system at the Big Bang. For example, the singularity is located at the boundary of the (x^1, x^2) plane, where $(x^1)^2 + (x^2)^2 \rightarrow \infty$. Therefore, when expressed in terms of $x^0, k_0, \dots, x^3, k_3$, the solutions become degenerate at the Big Bang. The values of certain variables (such as $(x^1)^2 + (x^2)^2$) at the singularity do not depend on the choice of initial values and, in this sense, they are not predictive. However, we can demonstrate that this loss of predictability at the Big Bang is coordinate-dependent. It is possible to find a sufficiently large number of variables that tend to finite *nontrivial* limits at the Big Bang, and at the same time possess the property that specifying their values at any instant, including at the singularity, uniquely determines the solution.

Specifically, we can demonstrate that the equations of motion in these variables form an autonomous set of ordinary differential equations (ODEs) that are regular at the Big Bang. This means that the RHSs of the equations of motion tend to finite limits, as do their first derivatives. At the singularity, these equations satisfy the conditions required by the Picard-Lindelöf theorem of existence and uniqueness of solutions of ODEs [41]. Thus, it is possible to set an initial value problem at the Big Bang that has a unique solution. Consequently, the Big Bang is not necessarily a region where determinism fails, as no information about the dynamical system is lost there.

If the singularity is a region where the existence and uniqueness theorem holds, a unique solution should depart from any of its points, *in two directions*. One direction leads to the interior of the configuration space we used so far. However, it is not clear at this point where the other direction should lead. In fact, the singularity lies at the boundary of the configuration space, and we need to extend this space in order to discuss the fate of the solutions that reach the Big Bang. The aforementioned regular variables enable us to achieve such an extension in a natural manner: the metric and scalar variables x^1, x^2, x^3 are related to the three regular variables β, θ, φ through a *gnomonic map*:

$$\begin{cases} x^1 = |\tan \beta| \sin \theta \cos \varphi, \\ x^2 = |\tan \beta| \sin \theta \sin \varphi, \\ x^3 = |\tan \beta| \cos \theta, \end{cases} \quad (42)$$

where $\beta, \theta \in [0, \pi]$, $\varphi \in [0, 2\pi)$ are hyperspherical coordinates on a three-sphere. These coordinates project the configuration space (x^1, x^2, x^3) onto a hemisphere of a three-sphere. The gnomonic map defines a double cover of an N -dimensional plane by an

N -sphere (see Fig. 2a), in which each hemisphere is mapped to a (x^1, x^2, x^3) hyperplane, extending the original configuration space into two copies of itself. Physically, we interpret each hyperplane as the configuration space of a three-geometry (plus a scalar field) with a different spatial orientation that flips upon crossing the boundary of the two hyperplanes (the equator of the three-sphere) [12, 14]. This implies that a universe approaching the singularity with a certain spatial orientation will collapse at the Big Bang into a degenerate zero-volume one-dimensional geometry, in which two spatial directions are infinitely smaller than the third one.³ Once the Big Bang is reached, the volume can start growing again, but a universe with an opposite spatial orientation will emerge. This entire process can be described using the extended configuration space (namely, the gnomonic three-sphere) where the singularity is projected from the boundary of the plane associated with a fixed spatial orientation onto the equator of the sphere ($\beta = \frac{\pi}{2}$). Therefore, the Big Bang is approached as $\beta \rightarrow \frac{\pi}{2}^\pm$, while the angles θ, φ represent the direction in which the equator is approached in the extended configuration space. Quiescent solutions, which were straight lines in the configuration plane, are projected onto half great circles on the gnomonic sphere. Each half great circle has a unique natural and regular continuation, which corresponds to the other half of the same great circle in the other hemisphere (see Fig. 2b).




Figure 2: Gnomonic map in two dimensions. (a) Gnomonic projection from (two copies of) a plane to a sphere. (b) Straight lines on planes are projected onto half great circles on the gnomonic sphere.

³There is also a measure-zero set of solutions of two-dimensional degenerate geometries, in which one direction is infinitely smaller than the other two [14].

At this point, we have a continuation result for purely Kasner solutions: they are great circles on the gnomonic sphere, which correspond to two (generally distinct) straight lines on the two (x^1, x^2, x^3) hyperplanes associated with the two spatial orientations. To extend this result to Bianchi-IX solutions, where the straight lines only exist in a neighborhood of the singularity, we must identify an additional set of five variables that exhibit a finite nontrivial value at the singularity.

The shape and scalar conjugate momenta k_1, k_2, k_3 can be regularized through the following change of variables:

$$\begin{cases} J = \text{sgn}(\tan \beta) \frac{x^1 k_1 + x^2 k_2 + x^3 k_3}{\sqrt{(x^1)^2 + (x^2)^2 + (x^3)^2}}, \\ L_1 = x^2 k_3 - x^3 k_2, \\ L_3 = x^1 k_2 - x^2 k_1, \end{cases} \quad (43)$$

while the scale variable and its conjugate momentum, x^0 and k_0 , require the following transformation:

$$\begin{cases} \eta = \text{sgn}(\tan \beta) \left(x^0 + k_0 \frac{(x^1)^2 + (x^2)^2 + (x^3)^2}{x^1 k_1 + x^2 k_2 + x^3 k_3} \right), \\ \kappa = |k_0|. \end{cases} \quad (44)$$

The variables we introduced tend to a finite limit as the singularity is approached by a quiescent solution:

$$\begin{cases} J \rightarrow \sin \theta (v_1 \cos \varphi + v_2 \sin \varphi) + v_3 \cos \theta, \\ L_1 \rightarrow \tan \beta (v_3 \sin \theta \sin \varphi - v_2 \cos \theta), \\ L_3 \rightarrow \tan \beta \sin \theta (v_2 \cos \varphi - v_1 \sin \varphi), \\ \eta \rightarrow \text{sgn}(\tan \beta) x^0 + v_0 J^{-1} \tan \beta, \\ \kappa \rightarrow v_0, \end{cases} \quad \text{as } \beta \rightarrow \frac{\pi}{2}^\pm. \quad (45)$$

The solution identified by the initial data $v_0, v_1, v_2, v_3, x^0(0), x^1(0), x^2(0), x^3(0)$ can be matched to a unique solution belonging to the other hemisphere with initial data $-v_0, -v_1, -v_2, -v_3, -x^0(0), -x^1(0), -x^2(0), -x^3(0)$. It is worth noting that the second solution reaches the limit as $t \rightarrow -\infty$, *i.e.*, the Big Bang singularity of the universe with the opposite spatial orientation is reached as $t \rightarrow -\infty$.

We have yet to discuss the electromagnetic degrees of freedom. Under the one-dimensional *ansatz*, there are two electromagnetic variables, A_3 and E^3 . Their equations of motion w.r.t. the effective Hamiltonian (39) (for $n = 1$) are the following:

$$\dot{A}_3 = q_3 E^3, \quad \dot{E}^3 = -q_3 A_3. \quad (46)$$

At the singularity, A_3 and E^3 become conserved as q_3 goes to zero. The electromagnetic variables are unaffected by the orientation flip, meaning that the constant values to which these variables tend are the same regardless of whether the singularity is approached from the left or the right ($\beta \rightarrow \frac{\pi}{2}^+$ or $\beta \rightarrow \frac{\pi}{2}^-$). Thus, these variables are already regular and effectively describe the evolution of electromagnetic degrees of freedom in the entire extended configuration space.

Expressed in terms of the variables $\beta, \theta, \varphi, \eta, J, L_1, L_3, \kappa, A_3, E^3$, and assuming the quiescence conditions are satisfied (*i.e.*, neglecting the potential terms), the Hamiltonian constraint (39) becomes:

$$\mathcal{H}_{Kasner} = \frac{1}{2} \left[\kappa^2 - J^2 - \frac{L_1^2}{\tan^2 \beta \sin^2 \varphi} + \frac{L_3^2 (\cos^2 \varphi - \sin^2 \theta)}{\tan^2 \beta \sin^2 \varphi} - \frac{L_1 L_3}{\tan^2 \beta \tan \theta \tan \varphi \sin \varphi} \right], \quad (47)$$

where we set $n = 1$ for sake of simplicity.

The equations of motion for the variables $\beta, \theta, \varphi, \eta, J, L_1, L_3, \kappa, A_3, E^3$ with respect to the coordinate time t can be obtained by calculating the Dirac brackets using the Hamiltonian constraint (39). However, t is not a suitable choice of independent variable at the singularity as it diverges there. Instead, a natural choice of independent variable is the arc-length on the gnomonic sphere:

$$d\ell = \sqrt{d\beta^2 + \sin^2 \beta (d\theta^2 + \sin^2 \theta d\varphi^2)}, \quad (48)$$

which is automatically monotonic everywhere on a solution and tends to a finite limit at the singularity. The equations of motion give us the relationship between the two independent variables:

$$\frac{d\ell}{dt} = \Lambda^{-1} \cos^2 \beta, \quad \Lambda = \left(J^2 + \sin^{-2} \beta \left((L_3 \sin \theta)^2 + \left(\frac{L_1}{\sin \varphi} + \frac{L_3}{\tan \theta \tan \varphi} \right)^2 \right) \right)^{-\frac{1}{2}}. \quad (49)$$

The equations of motion w.r.t. the arc-length during quiescence read:

$$\begin{aligned} \frac{d\beta}{d\ell} &= \Lambda J, & \frac{dJ}{d\ell} &= \Lambda \Theta \cos \beta \sin^{-3} \beta, \\ \frac{d\theta}{d\ell} &= -\Lambda \left(L_1 + L_3 \frac{\cos \varphi}{\tan \theta} \right) \sin^{-2} \beta \sin^{-1} \varphi, & \frac{dL_1}{d\ell} &= 0, \\ \frac{d\varphi}{d\ell} &= \Lambda L_3 \sin^{-2} \beta \sin^{-2} \theta, & \frac{dL_3}{d\ell} &= 0, \\ \frac{d\eta}{d\ell} &= -\Lambda \Theta \kappa J^{-2} \sin^{-2} \beta, & \frac{d\kappa}{d\ell} &= 0, \\ \frac{dA_3}{d\ell} &= 0, & \frac{dE^3}{d\ell} &= 0, \end{aligned} \quad (50)$$

where:

$$\Theta = \left(\frac{L_3^2}{\tan^2 \theta \tan^2 \varphi} + \frac{L_3^2}{\sin^3 \theta} + \frac{2L_1 L_3}{\tan \theta \tan \varphi \sin \varphi} + \frac{L_1^2}{\sin^2 \varphi} \right) \sin \theta. \quad (51)$$

It is important to notice that this model has ten degrees of freedom, but only eight of them are truly physical, as two are redundant due to the Hamiltonian constraint and its gauge-fixing. To eliminate the two remaining non-physical degrees of freedom, we need to solve the Hamiltonian constraint (47) w.r.t. one of the variables and impose a gauge-fixing condition. A straightforward choice for gauge fixing the Hamiltonian constraint, which also serves as the generator of the dynamics, is to fix a specific instant of time. In our case, the natural choice is $\beta = \frac{\pi}{2}$, which represent the instant of the singularity. The suitability of fixing β as a gauge for (47) can be verified by calculating the Dirac bracket between β and (47) and observing that it is never zero at $\beta = \frac{\pi}{2}$.

At the singularity, the Hamiltonian constraint tends to the simple expression

$$\mathcal{H}_{Kasner} = \frac{1}{2}(\kappa^2 - J^2). \quad (52)$$

Therefore, it can be easily solved w.r.t. either the variable κ or J . Once we compute the equations of motion and their first derivatives, we can impose the condition $\kappa = J$ (keeping in mind that both κ and J are positive-definite at the singularity) if we wish to eliminate this last redundant degree of freedom.

We are now prepared to present the continuation result. The equations of motion (50) are regular at the singularity, meaning that they admit the same left and right limit

as $\beta \rightarrow \frac{\pi}{2}$:

$$\begin{aligned}
\frac{d\beta}{d\ell} &\rightarrow \Lambda_{\pi/2} J, & \frac{dJ}{d\ell} &\rightarrow 0, \\
\frac{d\theta}{d\ell} &\rightarrow -\Lambda_{\pi/2} \left(L_1 + L_3 \frac{\cos \varphi}{\tan \theta} \right) \sin^{-1} \varphi, & \frac{dL_1}{d\ell} &\rightarrow 0, \\
\frac{d\varphi}{d\ell} &\rightarrow \Lambda_{\pi/2} L_3 \sin^{-2} \theta, & \frac{dL_3}{d\ell} &\rightarrow 0, \\
\frac{d\eta}{d\ell} &\rightarrow -\Lambda_{\pi/2} \Theta \kappa J^{-2}, & \frac{d\kappa}{d\ell} &\rightarrow 0, \\
\frac{dA_3}{d\ell} &\rightarrow 0, & \frac{dE^3}{d\ell} &\rightarrow 0,
\end{aligned} \tag{53}$$

where:

$$\Lambda_{\pi/2} = \lim_{\beta \rightarrow \pi/2} \Lambda = \left(J^2 + (L_3 \sin \theta)^2 + \left(\frac{L_1}{\sin \varphi} + \frac{L_3}{\tan \theta \tan \varphi} \right)^2 \right)^{-\frac{1}{2}}. \tag{54}$$

The assumptions of the Picard–Lindelöf theorem require the Lipschitz continuity of the right-hand side of the equations of motion. However, in our case, we can prove an even stronger condition: differentiability. In fact, the first derivatives of the RHSs of (50) w.r.t. all the variables $\beta, \theta, \varphi, \eta, J, L_1, L_3, \kappa, A_3, E^3$ are all regular as $\beta \rightarrow \frac{\pi}{2}^\pm$:

$$\begin{aligned}
\partial_\beta \left(\frac{dJ}{d\ell} \right) &\rightarrow -\Lambda_{\pi/2} \Theta, \\
\partial_\theta \left(\frac{d\beta}{d\ell} \right) &\rightarrow -\Lambda_{\pi/2}^3 J \sin \theta \left(L_3^2 \cos \theta - \frac{L_3^2}{\tan \theta \tan^2 \varphi \sin^3 \theta} - \frac{L_1 L_3}{\sin \varphi \tan \varphi \sin^3 \theta} \right), \\
\partial_\theta \left(\frac{d\theta}{d\ell} \right) &\rightarrow \Lambda_{\pi/2}^3 \frac{L_3}{\sin^2 \theta} \left(\frac{L_1 L_3 \cos \theta \sin^3 \theta}{\sin \varphi} + \frac{J^2 + \frac{1}{2} L_3^2 \sin^2 \theta (3 + \cos(2\theta))}{\tan \varphi} \right), \\
\partial_\theta \left(\frac{d\varphi}{d\ell} \right) &\rightarrow \Lambda_{\pi/2}^3 \frac{L_3}{\sin^2 \theta} \left(\frac{L_3^2}{\tan \theta \tan^2 \varphi \sin^2 \theta} + \frac{L_1 L_3}{\tan \varphi \sin^2 \theta \sin \varphi} - \frac{2J^2}{\tan \theta} \right. \\
&\quad \left. - \frac{2}{\tan \theta} \left(\frac{L_1}{\sin \varphi} + \frac{L_3}{\tan \theta \tan \varphi} \right)^2 - 3 L_3^2 \cos \theta \sin \theta \right), \\
\partial_\theta \left(\frac{d\eta}{d\ell} \right) &\rightarrow \Lambda_{\pi/2}^3 \frac{\kappa}{J^2} \left(-L_3 \Theta \sin \theta \left(\frac{L_3}{\tan \theta \tan^2 \varphi \sin^3 \theta} + \frac{L_1}{\tan \varphi \sin^3 \theta \sin \varphi} - L_3 \cos \theta \right) \right. \\
&\quad \left. + \Lambda_{\pi/2}^{-2} \frac{L_3}{\sin \theta} \left(\frac{L_3}{\tan \theta} \left(\frac{2}{\tan^2 \varphi} + \frac{3}{\sin \theta} \right) + \frac{2L_1}{\tan \varphi \sin \varphi} \right) - \Lambda_{\pi/2}^{-2} \Theta \cos \theta \right), \\
\partial_\varphi \left(\frac{d\beta}{d\ell} \right) &\rightarrow \Lambda_{\pi/2}^3 \frac{J}{\sin \varphi} \left(\frac{L_3}{\tan \theta \tan \varphi} + \frac{L_1}{\sin \varphi} \right) \left(\frac{L_1}{\tan \varphi} + \frac{L_3}{\tan \theta \sin \varphi} \right), \\
\partial_\varphi \left(\frac{d\theta}{d\ell} \right) &\rightarrow \Lambda_{\pi/2}^3 \frac{2J^2 + L_3^2 + L_3^2 \cos(2\theta)}{2 \sin^2 \varphi} \left(L_1 \cos \varphi + \frac{L_3}{\tan \theta} \right), \\
\partial_\varphi \left(\frac{d\varphi}{d\ell} \right) &\rightarrow \Lambda_{\pi/2}^3 \frac{L_3}{\sin^2 \theta \sin \varphi} \left(\frac{L_3}{\tan \theta \tan \varphi} + \frac{L_1}{\sin \varphi} \right) \left(\frac{L_1}{\tan \varphi} + \frac{L_3}{\tan \theta \sin \varphi} \right), \\
\partial_\varphi \left(\frac{d\eta}{d\ell} \right) &\rightarrow \Lambda_{\pi/2}^3 \frac{\kappa \sin^3 \theta}{J^2 \sin^2 \varphi} \left(\frac{2L_1^2}{\tan \varphi} + \frac{2L_3^2}{\tan \varphi \tan^2 \theta} + \frac{L_1 L_3 (3 + \cos(2\theta))}{\tan \theta \sin \varphi} \right) \\
&\quad \left(2L_3^2 - \frac{L_3^2}{\sin^5 \theta} + \frac{2J^2}{\sin^2 \theta} + \left(\frac{L_3}{\tan \theta \tan \varphi \sin \theta} + \frac{L_1}{\sin \varphi \sin \theta} \right)^2 \right), \\
\partial_J \left(\frac{d\beta}{d\ell} \right) &\rightarrow \Lambda_{\pi/2}^3 \left((L_3 \sin \theta)^2 + \left(\frac{L_1}{\sin \varphi} + \frac{L_3}{\tan \theta \tan \varphi} \right)^2 \right), \\
\partial_J \left(\frac{d\theta}{d\ell} \right) &\rightarrow \Lambda_{\pi/2}^3 \frac{J^2}{\sin \varphi} \left(L_1 + \frac{L_3 \cos \varphi}{\tan \theta} \right), \\
\partial_J \left(\frac{d\varphi}{d\ell} \right) &\rightarrow -\Lambda_{\pi/2}^3 \frac{J L_3}{\sin^2 \theta}, \\
\partial_J \left(\frac{d\eta}{d\ell} \right) &\rightarrow \Lambda_{\pi/2}^3 \Theta \frac{\kappa}{J^3} \left(3J^2 + 2 (L_3 \sin \theta)^2 + 2 \left(\frac{L_1}{\sin \varphi} + \frac{L_3}{\tan \theta \tan \varphi} \right)^2 \right),
\end{aligned}$$

(continued on next page)

(55)

(continued from previous page)

$$\begin{aligned}
\partial_{L_1} \left(\frac{d\beta}{d\ell} \right) &\rightarrow -\Lambda_{\pi/2}^3 \frac{J}{\sin \varphi} \left(\frac{L_3}{\tan \theta \tan \varphi} + \frac{L_1}{\sin \varphi} \right), \\
\partial_{L_1} \left(\frac{d\theta}{d\ell} \right) &\rightarrow -\Lambda_{\pi/2}^3 \frac{J^2 + L_3^2 \sin^2 \theta}{\sin \varphi}, \\
\partial_{L_1} \left(\frac{d\varphi}{d\ell} \right) &\rightarrow -\Lambda_{\pi/2}^3 \frac{L_3}{\sin^2 \theta \sin \varphi} \left(\frac{L_3}{\tan \theta \tan \varphi} + \frac{L_1}{\sin \varphi} \right), \\
\partial_{L_1} \left(\frac{d\eta}{d\ell} \right) &\rightarrow -\Lambda_{\pi/2}^3 \frac{\kappa \sin^3 \theta}{J^2 \sin \varphi} \left(\frac{L_3}{\tan \theta \tan \varphi} + \frac{L_1}{\sin \varphi} \right) \\
&\quad \left(2L_3^2 - \frac{L_3^2}{\sin^5 \theta} + \frac{2J^2}{\sin^2 \theta} + \left(\frac{L_3}{\tan \theta \tan \varphi \sin \theta} + \frac{L_1}{\sin \varphi \sin \theta} \right)^2 \right), \\
\partial_{L_3} \left(\frac{d\beta}{d\ell} \right) &\rightarrow -\Lambda_{\pi/2}^3 J \left(L_3 \sin^2 \theta + \frac{L_3}{\tan^2 \theta \tan^2 \varphi} + \frac{L_1}{\tan \theta \tan \varphi \sin \varphi} \right), \\
\partial_{L_3} \left(\frac{d\theta}{d\ell} \right) &\rightarrow \Lambda_{\pi/2}^3 \left(\frac{L_1 L_2 \sin^2 \theta}{\sin \varphi} - \frac{J^2}{\tan \theta \tan \varphi} \right), \\
\partial_{L_3} \left(\frac{d\varphi}{d\ell} \right) &\rightarrow \Lambda_{\pi/2}^3 \left(\frac{J^2}{\sin^2 \theta} + \frac{L_1 L_3}{\tan \theta \tan \varphi \sin \varphi \sin^2 \theta} + \frac{L_1^2}{\sin^2 \theta \sin^2 \varphi} \right), \\
\partial_{L_3} \left(\frac{d\eta}{d\ell} \right) &\rightarrow \Lambda_{\pi/2}^3 \frac{\kappa}{J^2} \left(\left(\frac{L_3 \cos \theta}{\tan \theta \tan^2 \varphi} + \frac{L_1 \cos \theta}{\tan \varphi \sin \varphi} \right) \left(2J^2 + \left(\frac{L_3}{\tan \theta \tan \varphi} + \frac{L_1}{\sin \varphi} \right)^2 \right) \right. \\
&\quad \left. - \frac{L_3^3 \cos^2 \theta \sin \theta}{\tan^2 \varphi} + \frac{L_1^2 L_3 \sin^3 \theta}{\sin^2 \varphi} - L_3^3 \right. \\
&\quad \left. - \frac{L_3}{\sin^2 \theta} \left(2J^2 + \left(\frac{L_3}{\tan \theta \tan \varphi} + \frac{L_1}{\sin \varphi} \right) \left(\frac{L_3}{\tan \theta \tan \varphi} + \frac{2L_1}{\sin \varphi} \right) \right) \right), \\
\partial_{\kappa} \left(\frac{d\theta}{d\ell} \right) &\rightarrow -\Lambda_{\pi/2} \Theta J^{-2},
\end{aligned}$$

and all the other derivatives tend to zero. Notice that imposing the asymptotic solution of the Hamiltonian constraint, $\kappa = J$, does not alter the regularity of the RHSs.

In full generality, when the potential terms cannot be neglected, the Hamiltonian constraint (39) assumes the following form in the new variables:

$$\begin{aligned}
\mathcal{H}_{eff} = \mathcal{H}_{Kasner} &+ \frac{1}{2} e^{\frac{2}{\sqrt{3}} \operatorname{sgn}(\tan \beta) (\eta - \kappa J^{-1} \tan \beta)} C(\beta, \theta, \varphi) \\
&+ \frac{1}{2} e^{\frac{1}{\sqrt{3}} \operatorname{sgn}(\tan \beta) (\eta - \kappa J^{-1} \tan \beta)} e^{-\frac{2}{\sqrt{3}} |\tan \beta| \sin \theta \sin \varphi} \varepsilon,
\end{aligned} \tag{56}$$

where $C(\beta, \theta, \varphi)$ represents the Bianchi-IX potential (23) as a function of β, θ, φ . When the quiescent approximation is relaxed, the equations of motion (50) acquire additional “force” terms arising from the potential. However, these terms are strongly suppressed near the equator/singularity, due to the exponential factors in Eq. (56), which tend to zero as $\beta \rightarrow \frac{\pi}{2}$ like $\exp(-\text{const.} |\tan \beta|)$ (after solving the Hamiltonian, *e.g.*, w.r.t. κ , and substituting the solution back into the equations of motion). In the equations of motion, the suppressing exponentials appear multiplied by powers of $\tan \beta$. Although the positive powers diverge, they do so slower than the exponentials and end up suppressed as well. As a result, the full equations of motion asymptotically tend to the quiescent ones (50). This holds true for the first-derivative expressions (55) as well, once again due to the presence of the suppressing exponentials.

7 Generic Einstein–Maxwell system

In the most generic situation, the electromagnetic field has all three components. The Hamiltonian constraint is given by Eq. (22). In this case as well, we can identify a

conserved quantity (*i.e.*, one that is first-class w.r.t. the Hamiltonian constraint):

$$H_{HO}^{3D} = \sum_{a=1}^3 ((E^a)^2 + (A_a)^2) , \quad \{H_{HO}^{3D}, \mathcal{H}[N]\}_* = 0 . \quad (57)$$

This can be readily proven by observing that $\mathcal{H}[N]$ depends on the electromagnetic variables only through the six terms $(E^a)^2 + (A_a)^2$ and M_a (the latter defined in Eq. (19)), and each of these terms commutes separately with H_{HO}^{3D} . These six terms correspond to the conserved quantities of a three-dimensional Harmonic oscillator, namely, three “energies” and three components of the angular momentum. Hence, we can associate a constant of motion ε_{3D} to H_{HO}^{3D} . It is important to notice that, although this quantity is not explicitly present in (22), it establishes bounds on the possible values of the electromagnetic field and momenta:

$$|A_a| \leq \sqrt{\varepsilon_{3D}} , \quad |E^a| \leq \sqrt{\varepsilon_{3D}} . \quad (58)$$

We now demonstrate that, with the inclusion of a scalar field, as discussed in Paragraph 1 of Section 6, the conditions for quiescence are still satisfied, even without the one-dimensional *ansatz* for the electromagnetic field.

As mentioned before, the relevant Hamiltonian constraint in this scenario is given by Eq. (22), to which we add a scalar field (referred to as x^3) without mass nor potential terms:

$$\mathcal{H}[N] = n \left[\frac{1}{2} (-k_0^2 + k_1^2 + k_2^2 + k_3^2) + \frac{1}{2} U(x, A, E) \right] , \quad (59)$$

where k_3 represents the conjugate momentum to x^3 . Since the electromagnetic variables only appear in the potential term $U(x, A, E)$, the removal of the one-dimensional *ansatz* does not affect the results obtained in Paragraph 1 of Section 6: during a Kasner epoch, the metric components q_a progressively decrease in time, along with any polynomial quantity derived from them. However, in the absence of the one-dimensional *ansatz*, the potential U acquires two additional terms (see Eqs. (23) and (24)), one of which is not even polynomial in q_a . Consequently, they need separate discussion.

The potential U now consists of a combination of three quantities: C , V , and W . The Bianchi-IX potential $C(x^1, x^2)$ depends only on the metric variables and is polynomial in q_a , hence its behavior is analogous to that of the effective potential in the one-dimensional model (35). The potential $V(x^1, x^2, A, E)$ is again polynomial in q_a , but its coefficients are functions of the electromagnetic variables. The presence of the conserved quantity (57) implies that E^a and A_a can only oscillate within finite and fixed values, thus the behavior of V is controlled by that of q_a . Similarly, the behavior of $W(x^1, x^2, A, E)$ is determined by q_a . However, in this case, the dependence of W on q_a is non-polynomial. W depends on the following three functions of q_a :

$$\begin{aligned} \frac{q_1 q_2}{(q_1 - q_2)^2} &= \frac{e^{2|\vec{v}|t \cos \varphi}}{(e^{2|\vec{v}|t \cos \varphi} - 1)^2} , & \frac{q_2 q_3}{(q_2 - q_3)^2} &= \frac{e^{|\vec{v}|t(\cos \varphi + \sqrt{3} \sin \varphi)}}{(e^{|\vec{v}|t(\cos \varphi + \sqrt{3} \sin \varphi)} - 1)^2} , \\ \frac{q_3 q_1}{(q_3 - q_1)^2} &= \frac{e^{|\vec{v}|t(\cos \varphi + \sqrt{3} \sin \varphi)}}{(e^{|\vec{v}|t \cos \varphi} - e^{\sqrt{3}|\vec{v}|t \sin \varphi})^2} , \end{aligned} \quad (60)$$

where q_a has been replaced by the solutions of the equations of motion during a Kasner epoch, as given by Eq. (37), with the velocities expressed in polar coordinates as in Eq. (38). As $t \rightarrow +\infty$, the three quantities in Eq. (60) tend to zero for all values of φ , except $\varphi = \frac{\pi}{6}, \frac{\pi}{2}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{3\pi}{2}, \frac{11\pi}{6}$. These six directions are parallel to the three axes of symmetry of the shape potential $C(x_1, x_2)$ [12, 14]. Along these directions, two of

the metric components q_a are identical, and one of the quantities in Eq. (60) becomes infinite. This singularity only affects a measure-zero set of solutions (those confined along the symmetry axes), and their continuability can be discussed separately.

We have demonstrated that the removal of the one-dimensional *ansatz* does not hinder quiescence: all the potential terms decrease with time, allowing the solution to settle around a single Kasner epoch all the way to the singularity.

Having established that the entire system (*i.e.*, including all six electromagnetic degrees of freedom) exhibits quiescent behavior as it approaches the Big Bang, we now proceed to demonstrate that the continuation result holds as well. To prove this, we follow the same procedure as described in Paragraph 2 of Section 6. In terms of the variables $\beta, \theta, \varphi, \eta, J, L_1, L_3, \kappa, A_1, E^1, A_2, E^2, A_3, E^3$, the dynamics governed by the Hamiltonian constraint (59) is indistinguishable from that generated by (39) when the quiescence conditions are satisfied. Therefore, the equations of motion can be well approximated by Eqs. (50), with the addition of:

$$\frac{dA_1}{d\ell} = 0, \quad \frac{dA_2}{d\ell} = 0, \quad \frac{dE^1}{d\ell} = 0, \quad \frac{dE^2}{d\ell} = 0, \quad (61)$$

whose RHSs are differentiable, similar to the other equations of motion, as we have previously demonstrated.

Due to the presence of additional potential terms, Eq. (56) modifies as:

$$\begin{aligned} \mathcal{H}_{eff} = \mathcal{H}_{Kasner} &+ \frac{1}{2}e^{\frac{2}{\sqrt{3}}\text{sgn}(\tan\beta)(\eta-\kappa J^{-1}\tan\beta)}C(\beta, \theta, \varphi) \\ &+ \frac{1}{2}e^{\frac{1}{\sqrt{3}}\text{sgn}(\tan\beta)(\eta-\kappa J^{-1}\tan\beta)}V(\beta, \theta, \varphi, A, E) + \frac{1}{2}W(\beta, \theta, \varphi, A, E). \end{aligned} \quad (62)$$

The equations of motion acquire additional “force” terms compared to the system under the one-dimensional *ansatz*; however, these terms are highly suppressed near the singularity. All the potential terms go exponentially to zero as $\beta \rightarrow \frac{\pi}{2}$, and the presence of a generic electromagnetic field does not affect this behavior. This is because all components of the electromagnetic field are bounded within fixed and finite values, and thus they do not lead any divergent contribution.

We are able to extend part of these results to non-Abelian gauge fields. However, this is only true under the one-dimensional simplifying *ansatz*, in which the first and the second components of the gauge fields (and momenta) are set to zero. At the moment we cannot prove the continuation result in full generality in the non-Abelian case, therefore we relegated the discussion of the present state of our understanding to Appendix A.

8 Conclusions

In Ref. [12], we conjectured that it is possible to continue Einstein’s *classical* equations through the Big Bang singularity into another universe with an opposite time direction and spatial orientation, which preserves all the information about the state of universe on the other side of the singularity (although it might become irretrievably scrambled in the process due to a chaotic phase of the dynamics). This is intimately related to far-reaching issues such as black hole unitarity and the nature of the Big Bang.

This conjecture was proven in simplified cases, including homogeneous cosmologies [12], inflationary models [13, 14], and the Schwarzschild-scalar system [15]. Our approach is to gradually increase the complexity of the models under consideration, test the validity of the conjecture, and gain insight into the behavior of physical fields across

the singularity. As mentioned in the previous paper [14], the next natural step in this process is to determine if the predicted reversal of orientation at the singularity can be physically measurable. In other words, can the inhabitants of the universe determine, through an experiment, which side of the Big Bang singularity they live in?

To answer this question, three ingredients are necessary. Firstly, we need to understand what happens to the orientation of space defined by the vielbein/frame fields. It has been established that these fields undergo a sign change at the singularity in the original paper [12]. Secondly, we must establish the behavior of vector (gauge) fields and fermions. If all of these fields undergo a “flipping” transition at the singularity, it might cancel out the orientation reversal effect of the vielbeins, making it unobservable. Finally, we need to investigate what happens to experimentally realizable processes, such as beta decays. Ultimately, the crucial factor is whether the parity-breaking vertices of the Standard Model remain unchanged across the singularity when considering their dependence on the spacetime vielbeins.

In the present paper, we conducted a detailed analysis of gauge fields. We first determined that the continuation result remains unchanged in the presence of Abelian gauge fields (in general) and non-Abelian gauge fields (under the simplifying assumption of the one-dimensional ansatz). Additionally, we established that the behavior of the gauge fields near the singularity is straightforward: their values freeze, with zero time derivatives at the exact instant of the singularity, and they evolve through it without flipping their orientation. The next logical step is to analyze fermion fields, which will allow us to determine the fate of the parity-breaking vertices of the Standard Model. Another interesting extension of this work would be to relax the one-dimensional ansatz for non-Abelian gauge fields, although this step has not been feasible thus far. This paper provides compelling evidence that the general case, beyond the one-dimensional ansatz, does not affect the continuation outcome nor the conclusion that gauge fields do not “flip” at the singularity. However, there is still some uncertainty around this matter, and further research is needed for confirmation.

Acknowledgements

This work has been partially supported by Agencia Estatal de Investigación (Spain) under grant PID2019-106802GB-I00/AEI/10.13039/501100011033, by the Regional Government of Castilla y León (Junta de Castilla y León, Spain), and by the Spanish Ministry of Science and Innovation MICIN and the European Union NextGenerationEU (PRTR C17.I1). The authors would also like to acknowledge the contribution of the COST Action CA18108 “Quantum gravity phenomenology in the multi-messenger approach”.

References

- [1] R. Penrose, “Gravitational collapse and space-time singularities,” *Phys. Rev. Lett.* **14** (1965) 57–59.
- [2] S. Hawking, “The Occurrence of singularities in cosmology,” *Proc. Roy. Soc. Lond.* **A294** (1966) 511–521.
- [3] S. Hawking, “The Occurrence of singularities in cosmology. II,” *Proc. Roy. Soc. Lond.* **A295** (1966) 490–493.

- [4] S. Hawking, “The occurrence of singularities in cosmology. III. Causality and singularities,” *Proc. Roy. Soc. Lond.* **A300** (1967) 187–201.
- [5] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 2011.
- [6] C. W. Misner, “Quantum cosmology. I,” *Phys. Rev.* **186** (1969) 1319–1327.
- [7] B. K. Berger, “Singularity avoidance in the semiclassical Gowdy T^3 cosmological model,” *Phys. Lett.* **108B** (1982) 394–398.
- [8] A. Ashtekar, A. Henderson, and D. Sloan, “A Hamiltonian Formulation of the BKL Conjecture,” *Phys. Rev.* **D83** (2011) 084024, [arXiv:1102.3474 \[gr-qc\]](#).
- [9] M. Bojowald, “Quantum cosmology: a review,” *Rep. Prog. Phys.* **78** no. 2, (Jan, 2015) 023901.
- [10] J. S. Schwinger, “Quantum electrodynamics. 2. Vacuum polarization and selfenergy,” *Phys. Rev.* **75** (1948) 651.
- [11] S. W. Hawking, “Breakdown of Predictability in Gravitational Collapse,” *Phys. Rev.* **D14** (1976) 2460–2473.
- [12] T. A. Koslowski, F. Mercati, and D. Sloan, “Through the Big Bang,” *Phys. Lett.* **B778** (2018) 339–343, [arXiv:1607.02460 \[gr-qc\]](#).
- [13] D. Sloan, “Scalar Fields and the FLRW Singularity,” *Class. Quant. Grav.* **36** no. 23, (2019) 235004, [arXiv:1907.08287 \[gr-qc\]](#).
- [14] F. Mercati, “Through the Big Bang in inflationary cosmology,” *JCAP* **2019** no. 10, (Oct, 2019) 025–025.
- [15] F. Mercati and D. Sloan, “Traversing through a black hole singularity,” *Phys. Rev. D* **106** no. 4, (Aug, 2022) .
- [16] C. W. Misner, “Mixmaster Universe,” *Phys. Rev. Lett.* **22** no. 20, (1969) 1071–1074.
- [17] C. Misner, K. Thorne, J. Wheeler, and D. Kaiser, *Gravitation*. Princeton University Press, 2017.
- [18] J. D. Barrow, “Quiescent cosmology,” *Nature* **272** no. 5650, (1978) 211.
- [19] D. Sloan, “Dynamical Similarity,” *Phys. Rev. D* **97** no. 12, (2018) 123541, [arXiv:1803.04472 \[gr-qc\]](#).
- [20] D. Sloan, “New action for cosmology,” *Phys. Rev. D* **103** no. 4, (2021) 043524, [arXiv:2010.07329 \[gr-qc\]](#).
- [21] S. Gryb and D. Sloan, “When scale is surplus,” *Synthese* **199** no. 5-6, (2021) 14769–14820, [arXiv:2103.07384 \[physics.hist-ph\]](#).
- [22] D. Sloan, “Scale Symmetry and Friction,” *Symmetry* **13** no. 9, (2021) 1639, [arXiv:2107.09172 \[gr-qc\]](#).

- [23] D. Sloan, “Herglotz Action for Homogeneous Cosmology,” [arXiv:2206.11322 \[gr-qc\]](#).
- [24] Planck Collaboration, “Planck 2018 results. X. Constraints on inflation,” *A&A* **641** (Sep, 2020) A10.
- [25] V. A. Belinsky, I. M. Khalatnikov, and E. M. Lifshitz, “A General Solution of the Einstein Equations with a Time Singularity,” *Adv. Phys.* **31** (1982) 639–667.
- [26] L. Andersson and A. Rendall, “Quiescent Cosmological Singularities,” *Commun. Math. Phys.* **218** (01, 2000) .
- [27] D. Garfinkle, “Numerical Relativity Beyond Astrophysics,” *Rept. Prog. Phys.* **80** no. 1, (2017) 016901, [arXiv:1606.02999 \[gr-qc\]](#).
- [28] C. Uggla, H. van Elst, J. Wainwright, and G. F. R. Ellis, “Past attractor in inhomogeneous cosmology,” *Phys. Rev. D* **68** (Nov, 2003) 103502.
- [29] D. Sloan, “Anisotropic matter in cosmology: locally rotationally symmetric Bianchi *I* and *VII_o* models,” *Class. Quant. Grav.* **33** no. 10, (2016) 105011, [arXiv:1601.00348 \[gr-qc\]](#).
- [30] P. D. Bolton, F. F. Deppisch, and P. S. B. Dev, “Neutrinoless double beta decay via light neutralinos in R-parity violating supersymmetry,” *JHEP* **2022** no. 3, (Mar, 2022) .
- [31] R. Arnowitt, S. Deser, and C. W. Misner, “Republication of: The dynamics of general relativity,” *Gen. Relativ. Gravit.* **40** no. 9, (Aug, 2008) 1997–2027.
- [32] S. A. Hojman, K. Kuchař, and C. Teitelboim, “Geometrodynamics regained,” *Ann. Phys.* **96** no. 1, (1976) 88–135.
- [33] F. Mercati, *Shape Dynamics: Relativity and Relationalism*. Oxford Univ. Press, 2018. [arXiv:1409.0105](#). (early version on the arXiv).
- [34] C. Blohmann, M. C. B. Fernandes, and A. Weinstein, “Groupoid symmetry and constraints in general relativity,” *Commun. Contemp. Math.* **15** no. 01, (2013) 1250061, [arXiv:1003.2857 \[math.DG\]](#).
- [35] L. Bianchi, “On the Three-Dimensional Spaces Which Admit a Continuous Group of Motions,” *Gen. Relativ. Gravit.* **33** (12, 2001) 2171–2253.
- [36] P. Dirac, *Lectures on Quantum Mechanics*. Belfer Graduate School of Science, monograph series. Dover Publications, 2001.
- [37] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems*. Princeton paperbacks. Princeton University Press, 1992.
- [38] A. Hanson, T. Regge, and C. Teitelboim, *Constrained Hamiltonian Systems*. Accademia nazionale dei Lincei : Contributi del centro linceo interdisciplinare di scienze matematiche e loro applicazioni. Accademia Nazionale dei Lincei, 1976.
- [39] J. W. York, “Role of Conformal Three-Geometry in the Dynamics of Gravitation,” *Phys. Rev. Lett.* **28** (Apr, 1972) 1082–1085.

- [40] H. Gomes, S. Gryb, and T. Koslowski, “Einstein gravity as a 3D conformally invariant theory,” *Class. Quant. Grav.* **28** (2011) 045005, [arXiv:1010.2481 \[gr-qc\]](#).
- [41] G. Teschl, *Ordinary Differential Equations and Dynamical Systems*. Graduate studies in mathematics. American Mathematical Society, 2012.

Appendices

A Einstein–Yang–Mills model under one-dimensional *ansatz*

The results presented in this work for the Einstein–Maxwell system can be extended to the Einstein–Yang–Mills systems with $SU(2)$ and $SU(3)$ structure groups, under the one-dimensional *ansatz*. These non-abelian models are described by the action:

$$S = \int d^4x \sqrt{-h} \left(R - \frac{1}{4} h^{\mu\nu} h^{\rho\sigma} F_{\mu\rho}^I F_{\nu\sigma}^J \delta_{IJ} \right), \quad (63)$$

with Faraday tensor $F_{\mu\nu}^I = \partial_\mu A_\nu^I - \partial_\nu A_\mu^I + c_{JK}^I A_\mu^J A_\nu^K$. The structure constants are given by the three-dimensional Levi-Civita symbol $c_{JK}^I = \delta^{IL} \varepsilon_{LJK}$ for $SU(2)$, and, in the case of $SU(3)$, by a totally-antisymmetric symbol $c_{JK}^I = \delta^{IL} f_{LJK}$, where $f_{123} = 1$, $f_{147} = f_{165} = f_{246} = f_{257} = f_{345} = f_{376} = 1/2$, $f_{458} = f_{678} = \sqrt{3}/2$, and all the others (which are not permutations of these indices) are zero. The scalar product in the internal gauge space is given by the group metric δ_{IJ} , which is also used for raising and lowering internal indices.

1 Homogeneous *ansatz* and global constraints. In the Hamiltonian formalism, after imposing the homogeneous *ansatz*, a generic Einstein–Yang–Mills system undergoes time evolution governed by a Hamiltonian that is a linear combination of the following global constraints:

$$\begin{aligned} \mathcal{H}[N] &= n \left(p^{ab} p^{cd} q_{bc} q_{da} - \frac{1}{2} (p^{ab} q_{ab})^2 + q_{ab} q_{cd} \delta^{bc} \delta^{da} - \frac{1}{2} (q_{ab} \delta^{ab})^2 \right. \\ &\quad \left. + \frac{1}{2} q_{ab} \delta^{IJ} E_I^a E_J^b + \frac{1}{4} \det q q^{ab} q^{cd} \delta_{IJ} F_{ac}^I F_{bd}^J \right), \\ \mathcal{D}_i[N^i] &= n^d \left(E_I^a A_b^I \varepsilon_{adc} \delta^{cb} + 2 p^{ab} q_{ac} \varepsilon_{bdf} \delta^{fc} \right), \end{aligned} \quad (64)$$

$$\mathcal{G}_I[A_0^I] = a_0^I (A_a^J E_K^a c_{JI}^K),$$

where δ^{IJ} is the inverse group metric, n and n^a are defined in Eq. (10), and

$$a_0^I = \int d\theta d\phi d\psi \sin \theta A_0^I(x), \quad (65)$$

are $N^2 - 1$ new Lagrange multipliers, corresponding to the spatial average of the scalar potential A_0^I , where $\dim SU(N) = N^2 - 1$ is the dimension of the gauge group. It should be noted that unlike the Einstein–Maxwell model where the Gauss constraint is automatically satisfied by the homogeneous *ansatz* (as shown in Eqs. (9)), in the case of the Einstein–Yang–Mills system, the Gauss constraint becomes a set of $N^2 - 1$ new proper constraints that need to be solved and gauge-fixed.

2 Gauge-fixing the diffeomorphism constraints. The diffeomorphism constraints in Eqs. (64) share the same functional expression as the abelian ones (the electromagnetic contribution to the diffeomorphism constraints in Eqs. (9) can be rewritten as $E^a A_b \varepsilon_{adc} \delta^{cb}$). Consequently, we can use the same gauge-fixing as in Eqs. (12). By

solving the constraints, the following solutions are obtained:

$$\begin{cases} p^{23} = \frac{E_I^2 A_3^I - E_I^3 A_2^I}{2(q_2 - q_3)}, \\ p^{13} = \frac{E_I^3 A_1^I - E_I^1 A_3^I}{2(q_3 - q_1)}, \\ p^{12} = \frac{E_I^1 A_2^I - E_I^2 A_1^I}{2(q_1 - q_2)}. \end{cases} \quad (66)$$

By applying the same procedure as outlined in Section 4, we derive the following on-shell Hamiltonian constraint:

$$\begin{aligned} \mathcal{H}[N] = n & \left[\mathcal{H}_{BIX} + \frac{q_2 q_3 (M_1)^2}{2(q_2 - q_3)^2} + \frac{q_1 q_3 (M_2)^2}{2(q_1 - q_3)^2} + \frac{q_1 q_2 (M_3)^2}{2(q_1 - q_2)^2} \right. \\ & + \frac{1}{2} \sum_{I=1}^{N^2-1} (q_1 ((E_I^1)^2 + (A_1^I)^2) + q_2 ((E_I^2)^2 + (A_2^I)^2) + q_3 ((E_I^3)^2 + (A_3^I)^2)) \\ & \left. + f(AAA) + g(AAAA) \right], \end{aligned} \quad (67)$$

where in this case, the kinetic term of the gauge fields incorporates the contribution from all the gauge components, and there are also two additional terms (cubic and quartic in the vector potential A_a^I) arising from the interaction of the non-abelian gauge field with itself. The Gauss constraints, which are independent of the metric variables, remain unchanged.

3 One-dimensional ansatz. As we did for the electromagnetic case, we consider a gauge field with a single spatial component:

$$A_1^I = A_2^I = 0, \quad E_I^1 = E_I^2 = 0, \quad \forall I \in \{1, \dots, N^2 - 1\}. \quad (68)$$

This *ansatz* is well-posed for the same reasons discussed at the beginning of Section 6.

The Hamiltonian constraint (Eq. (67)) and Gauss constraints (the last equation in (64)) become:

$$\bar{\mathcal{H}}[N] = \mathcal{H}[N] \Big|_{\substack{A_1^I = A_2^I = 0 \\ E_I^1 = E_I^2 = 0}} = n \left[\mathcal{H}_{BIX} + \frac{1}{2} q_3 \sum_{I=1}^{N^2-1} ((E_I^3)^2 + (A_3^I)^2) \right], \quad (69)$$

$$\mathcal{G}_I[A_0^I] = a_0^I (A_3^J E_K^3 c_{JI}^K).$$

Notice that, under the one-dimensional *ansatz*, the solution of the diffeomorphism constraints becomes $p^{12} = p^{23} = p^{13} = 0$, similar to the one-dimensional abelian case. Additionally, the self-interaction terms $f(AAA)$ and $f(AAAA)$ in the Hamiltonian constraint are also zero. Therefore, the Hamiltonian constraint of a one-dimensional non-abelian system takes the same form as the abelian one under the same *ansatz*, *viz.* Eq. (31). However, the Gauss constraints still need to be solved and gauge-fixed.

4 Gauge-fixing the Gauss constraints. A non-abelian model with a gauge group $SU(N)$ has $N^2 - 1$ non-zero Gauss constraints. However, under the one-dimensional *ansatz*, not all of these constraints are independent. In the case of the groups we are interested in, namely $SU(2)$ and $SU(3)$, it is found that there are only two (out of three) linearly independent Gauss constraint for $SU(2)$, and six (out of eight) linearly independent Gauss constraints for $SU(3)$. This observation is consistent with the number of Casimir operators of these groups: $SU(2)$ has one Casimir, whereas $SU(3)$ has two. The Casimir operators represent the number of free parameters used to label the group representations, while the remaining parameters are determined by the choice of gauge for the independent Gauss constraints.

By arbitrarily selecting $\mathcal{G}_{1,2}$ as the independent gauge generators for $SU(2)$ and $\mathcal{G}_{1,2,4,5,6,7}$ for $SU(3)$, we can find a well-posed gauge-fixing:

$$\begin{aligned} SU(2) : \quad & \begin{cases} \mathcal{G}_1, \mathcal{G}_2 \approx 0, \\ A_3^1, A_3^2 \approx 0, \end{cases} \\ SU(3) : \quad & \begin{cases} \mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_4, \mathcal{G}_5, \mathcal{G}_6, \mathcal{G}_7 \approx 0, \\ A_3^1, A_3^2, A_3^4, A_3^5, A_3^6, A_3^7 \approx 0. \end{cases} \end{aligned} \tag{70}$$

Once the Gauss constraints are solved, the conjugate momenta E_I^3 corresponding to the gauge-fixed components A_3^I must be zero. As a result, the remaining independent gauge components are A_3^3, E_3^3 for $SU(2)$, and $A_3^3, A_3^8, E_3^3, E_8^3$ for $SU(3)$.

5 Effective Hamiltonian constraint. After solving the Gauss constraints, the only remaining constraint is the Hamiltonian one:

$$\begin{aligned} SU(2) : \quad & \overline{\mathcal{H}}[N] = n \left[\mathcal{H}_{BIX} + \frac{1}{2} q_3 ((E_3^3)^2 + (A_3^3)^2) \right], \\ SU(3) : \quad & \overline{\mathcal{H}}[N] = n \left[\mathcal{H}_{BIX} + \frac{1}{2} q_3 ((E_3^3)^2 + (E_8^3)^2 + (A_3^3)^2 + (A_3^8)^2) \right]. \end{aligned} \tag{71}$$

As discussed in Section 6, we can identify a conserved quantity, which is a first-class quantity w.r.t. the Hamiltonian constraint. In the case of $SU(2)$, this conserved quantity corresponds again a one-dimensional harmonic oscillator:

$$H_{HO}^{1D} = (E_3^3)^2 + (A_3^3)^2, \tag{72}$$

(compare this to the abelian case, Eq. (32)), while for $SU(3)$ the conserved quantity is:

$$H_{HO}^{2D} = (E_3^3)^2 + (E_8^3)^2 + (A_3^3)^2 + (A_3^8)^2. \tag{73}$$

H_{HO}^{1D} and H_{HO}^{2D} are constants of motion, and we can set them to a positive constant ε for both gauge groups without loss of generality. This final step allows us to describe the dynamics of both $SU(2)$ and $SU(3)$ gauge systems with the same effective Hamiltonian:

$$\mathcal{H}_{eff}[N] = n \left[\mathcal{H}_{BIX} + \frac{1}{2} q_3 \varepsilon \right]. \tag{74}$$

Since this effective Hamiltonian is identical to that of the one-dimensional abelian case (see Eq. (34)), the continuation result proven in Section 6 also applies to the Einstein–Yang–Mills systems with $SU(2)$ and $SU(3)$ as structure groups under the one-dimensional *ansatz*.