Second order error bounds for POD-ROM methods based on first order divided differences

Bosco García-Archilla^a, Volker John^{b,c}, Julia Novo^{d,*}

^aDepartamento de Matemática Aplicada II, Universidad de Sevilla, Sevilla, Spain ^bWeierstrass Institute for Applied Analysis and Stochastics (WIAS), Mohrenstr. 39, 10117 Berlin, Germany ^cFreie Universität Berlin, Dept. of Mathematics and Computer Science, Arnimallee 6, 14195 Berlin, Germany ^dDepartamento de Matemáticas, Universidad Autónoma de Madrid, Spain

Abstract

This note proves, for simplicity for the heat equation, that using BDF2 as time stepping scheme in POD-ROM methods with snapshots based on difference quotients gives both the optimal second order error bound in time and pointwise estimates.

Keywords: Heat equation; POD-ROM methods; BDF2; Pointwise error estimates

1. Introduction

Most numerical methods using reduced order models based on proper orthogonal decomposition (POD-ROM methods) apply basis functions based on the snapshots (or values at different times) of the full order model (FOM). Recently, it has been shown that adding their first divided differences to the snapshots, or even using only these divided differences to obtain the basis functions, allows for pointwise-in-time error bounds [1, 2, 4, 6]. However, all pointwise-in-time error bounds in the literature are only first order with respect to time.

Although the first divided differences are only first order approximations to the time derivatives of the snapshots, we show in this note that for POD-ROM methods based only on them it is possible to obtain pointwise-in-time second order error bounds if the two step backward differentiation formula (BDF2) is used to integrate the POD-ROM equations. This result is a theoretical support for the observation that second order methods allow for larger step sizes than first order ones without spoiling the error, thus resulting in more efficient POD-ROM simulations.

2. Model problem and proper orthogonal decomposition

Throughout this note, standard notations for Sobolev spaces and their norms will be used. As a model problem problem, we consider the heat equation

 $\begin{array}{rcl} \partial_t u(t, \boldsymbol{x}) - \nu \Delta u(t, \boldsymbol{x}) &=& f(t, \boldsymbol{x}), \quad (t, \boldsymbol{x}) \in (0, T] \times \Omega, \\ u(t, \boldsymbol{x}) &=& 0, \qquad (t, \boldsymbol{x}) \in (0, T] \times \partial \Omega, \\ u(0, \boldsymbol{x}) &=& u^0(\boldsymbol{x}), \quad \boldsymbol{x} \in \Omega, \end{array}$

^{*}Corresponding author

Email addresses: bosco@esi.us.es; research is supported by Spanish MCINYU under grants PGC2018-096265-B-I00 and PID2019-104141GB-I00 (Bosco García-Archilla), john@wias-berlin.de, ORCID 0000-0002-2711-4409 (Volker John), julia.novo@uam.es; research is supported by Spanish MINECO under grants PID2019-104141GB-I00 and VA169P20 (Julia Novo)

in a bounded domain $\Omega \subset \mathbb{R}^d$, $d \in \{2, 3\}$. Let C_p be the constant in the Poincaré inequality

$$\|v\|_0 \le C_p \|\nabla v\|_0, \quad v \in H_0^1(\Omega).$$
 (1)

Let us denote by X_h^l a finite element method based on piece-wise continuous polynomials of degree l that satisfies the homogeneous Dirichlet boundary conditions. The semi-discrete Galerkin approximation, the FOM, consists in finding $u_h : [0,T] \to X_h^l$ such that

$$(\partial_t u_h, v_h) + \nu(\nabla u_h, \nabla v_h) = (f, v_h), \quad \forall \ v_h \in X_h^l.$$

The following error estimation is well-known:

$$\max_{0 \le s \le T} (\|(u - u_h)(s)\|_0 + h\|(u - u_h)(s)\|_1) \le C(u)h^{l+1}.$$
(2)

Fix T > 0 and set $\Delta t = T/M$. Let $t^n = n\Delta t$, $n = 0, \dots, M$, N = M + 1, and define the space

$$\boldsymbol{U} = \operatorname{span}\left\{\sqrt{N}w_{0}, \tau \frac{u_{h}(t^{1}) - u_{h}(t^{0})}{\Delta t}, \tau \frac{u_{h}(t^{2}) - u_{h}(t^{1})}{\Delta t} \dots, \tau \frac{u_{h}(t^{M}) - u_{h}(t^{M-1})}{\Delta t}\right\},$$

where w_0 is either $w_0 = u_h(t^0)$ or $w_0 = \overline{u}_h = \sum_{j=0}^M u_h(t^j)/(M+1)$, and τ is a time scale to make the snapshots dimensionally correct. Denote $U = \operatorname{span}\{y_h^1, y_h^2, \ldots, y_h^N\}$. Let X be either $X = L^2(\Omega)$ or $X = H_0^1(\Omega)$, and denote the correlation matrix by $K = ((k_{i,j})) \in \mathbb{R}^{N \times N}$ with $k_{i,j} = (y_h^i, y_h^j)_X/N, i, j = 1, \ldots, N$, and $(\cdot, \cdot)_X$ being the inner product in X. We denote by $\lambda_1 \geq \lambda_2 \ldots \geq \lambda_d > 0$ the positive eigenvalues of K and by $v_1, \ldots, v_d \in \mathbb{R}^N$ the associated eigenvectors. The orthonormal POD basis functions of U are given by $\varphi_k = (\sum_{j=1}^N v_k^k y_h^j)/(\sqrt{N}\sqrt{\lambda_k})$, where v_k^j is the j-th component of v_k . For any $1 \leq r \leq d$ denote by $U^r = \operatorname{span}\{\varphi_1, \varphi_2, \ldots, \varphi_r\}$, and denote by $P^r : X_h^l \to U^r$ the X-orthogonal projection onto U^r . Then, it holds

$$\frac{1}{N}\sum_{j=1}^{N} \|y_h^j - P^r y_h^j\|_X^2 = \sum_{k=r+1}^d \lambda_k.$$
(3)

The stiffness matrix of the POD basis is given by $S = ((s_{i,j})) \in \mathbb{R}^{d \times d}$, with $s_{i,j} = (\nabla \varphi_i, \nabla \varphi_j)_X$. If $X = L^2(\Omega)$ the following inequality holds for all $v \in U$, see [5, Lemma 2],

$$\|\nabla v\|_0 \le \sqrt{\|S\|_2} \|v\|_0. \tag{4}$$

3. Error analysis

Let us denote by $D^1v^n = (v^n - v^{n-1})/\Delta t$ and by $D^2v^n = ((3/2)v^n - 2v^{n-1} + (1/2)v^{n-2})/\Delta t$, then the POD-ROM method is defined in the following way: Find $u_r^n \in U^r$ such that

$$(Du_r^n, v) + \nu(\nabla u_r^n, \nabla v) = (f^n, v), \quad \forall \ v \in U^r,$$

where $D = D^1$ for n = 1 and $D = D^2$ for $2 \le n \le M$.

Lemma 1. Let T > 0, let X be a Banach space, $z^n = z(t^n) \in X$, then

$$\max_{0 \le k \le M} \|z^k\|_X^2 \le 2\|z^0\|_X^2 + \frac{2T^2}{M} \sum_{n=1}^M \|D^1 z_n\|_X^2,$$
(5)

$$\max_{0 \le k \le M} \|z^k\|_X^2 \le 2\|\overline{z}\|_X^2 + \frac{8T^2}{M} \sum_{n=1}^M \|D^1 z_n\|_X^2, \quad \text{with} \quad \overline{z} = \sum_{j=0}^M z^j / (M+1).$$
(6)

Proof. The proof of (5) can be found in [1, Lemma 3.3]. For proving (6), we observe that

$$z^{k} = z^{0} + \Delta t \sum_{n=1}^{k} D^{1} z^{n}, \quad \overline{z} = z^{0} + \frac{1}{M+1} \left(\Delta t D^{1} z^{1} + \ldots + \Delta t \sum_{n=1}^{M} D^{1} z^{n} \right).$$
(7)

Taking norms yields $\|z^k\|_X \le \|z^0\|_X + \Delta t \sum_{n=1}^M \|D^1 z^n\|_X$ and $\|z^0\|_X \le \|\overline{z}\|_X + \Delta t \sum_{n=1}^M \|D^1 z^n\|_X$, so that

$$\|z^{k}\|_{X} \leq \|\overline{z}\|_{X} + 2\Delta t \sum_{n=1}^{M} \|D^{1}z^{n}\|_{X} \leq \|\overline{z}\|_{X} + 2T^{1/2}(\Delta t)^{1/2} \left(\sum_{n=1}^{M} \|D^{1}z^{n}\|_{X}^{2}\right)^{1/2},$$

hich we reach (6).

from which we reach (6).

In the sequel we define $\tilde{C} = 1$ if $w_0 = u_h(t^0)$ and $\tilde{C} = 4$ if $w_0 = \overline{u}_h$, and $C_X = 1$ if $X = L^2(\Omega)$ and $C_X = C_p^2$ f $X = H_0^1(\Omega)$.

Lemma 2. The following bound holds

$$\max_{0 \le n \le M} \|u_h^n - P^r u_h^n\|_0^2 \le \left(2 + 4\tilde{C}\frac{T^2}{\tau^2}\right) C_X \sum_{k=r+1}^d \lambda_k.$$
(8)

Proof. Taking $z = u_h - P^r u_h$ in (5) or (6), depending on the selection of the first element in U, and applying (3) and $N \leq 2M$, we reach (8).

Lemma 3. Let $\{z^n\}_{n=0}^N \in U^r$ and $\{\tau_1^n\}_{n=1}^N, \{\tau_2^n\}_{n=1}^N \in X_h^l$ satisfying

$$(Dz^n, v) + \nu(\nabla z^n, \nabla v) = (\tau_1^n, v) + \nu(\nabla \tau_2^n, \nabla v), \quad \forall \ v \in \boldsymbol{U}^r,$$
(9)

where $D = D^1$ for n = 1 and $D = D^2$ for $2 \le n \le M$. Then, it holds for $\Delta t < T/4$ and $n \ge 1$

$$\|z^{n}\|_{0}^{2} + 2\nu \sum_{j=1}^{n} \Delta t \|\nabla z^{j}\|_{0}^{2} \leq e^{4} \left(17\|z^{0}\|_{0}^{2} + 28(\Delta t)^{2} \|\tau_{1}^{1}\|_{0}^{2} + 2\Delta tT \sum_{j=2}^{N} \|\tau_{1}^{n}\|_{0}^{2} + 14\nu\Delta t \|\nabla \tau_{2}^{1}\|_{0}^{2} + 2\nu\Delta t \sum_{j=2}^{N} \|\nabla \tau_{2}^{n}\|_{0}^{2} \right).$$

$$(10)$$

Proof. We take $v = \Delta t z^n$ in (9). If n = 1 then $D = D^1$ and Young's inequality yields

$$\frac{1}{2} \|z^1\|_0^2 - \frac{1}{2} \|z^0\|_0^2 + \nu \Delta t \|\nabla z^1\|_0^2 \le \Delta t(\tau_1^1, z^1) + \nu \Delta t(\nabla \tau_2^1, \nabla z^1).$$
(11)

For $n \ge 2$ then $D = D^2$ and one gets

$$\frac{1}{4}\|z^n\|_0^2 + \frac{1}{4}\|\hat{z}^n\|_0^2 - \frac{1}{4}\|z^{n-1}\|_0^2 - \frac{1}{4}\|\hat{z}^{n-1}\|_0^2 + \nu\Delta t\|\nabla z^n\|_0^2 \le \Delta t(\tau_1^n, z^n) + \nu\Delta t(\nabla \tau_2^n, \nabla z^n),$$

where $\hat{z}^n = 2z^n - z^{n-1}$. The Cauchy–Schwarz and Young inequality give

$$\Delta t(\tau_1^n, z^n) + \nu \Delta t(\nabla \tau_2^n, \nabla z^n) \le \frac{\Delta t}{2T} \|z^n\|_0^2 + \frac{T}{2} \|\tau_1^n\|_0^2 + \Delta t \frac{\nu}{2} \|\nabla z^n\|_0^2 + \Delta t \frac{\nu}{2} \|\nabla \tau_2^n\|_0^2.$$
(12)

Multiplying by 4, applying (12), and summing from 2 to n, one gets

$$\|z^{n}\|_{0}^{2} + 2\nu \sum_{j=2}^{n} \Delta t \|\nabla z^{j}\|_{0}^{2} \le \|z^{1}\|_{0}^{2} + \|\hat{z}^{1}\|_{0}^{2} + 2\sum_{j=2}^{n} \frac{\Delta t}{T} \|z^{j}\|_{0}^{2} + 2T \sum_{j=2}^{n} \Delta t \|\tau_{1}^{j}\|_{0}^{2} + 2\nu \sum_{j=2}^{n} \Delta t \|\nabla \tau_{2}^{j}\|_{0}^{2}.$$
(13)

Young's inequality yields $\|\hat{z}^1\|_0^2 \leq 6\|z^1\|_0^2 + 3\|z^0\|^2$, so that $\|z^1\|_0^2 + \|\hat{z}^1\|_0^2 \leq 7\|z^1\|_0^2 + 3\|z^0\|^2$. Using again Young's inequality gives

$$\Delta t(\tau_1^1, z^1) + \nu \Delta t(\nabla \tau_2^1, \nabla z^1) \le \frac{1}{4} \left\| z^1 \right\|_0^2 + (\Delta t)^2 \left\| \tau_1^1 \right\|_0^2 + \Delta t \frac{\nu}{2} \left\| \nabla z^1 \right\|_0^2 + \Delta t \frac{\nu}{2} \left\| \nabla \tau_2^1 \right\|_0^2,$$

so that we obtain from (11)

$$\left\|z^{1}\right\|_{0}^{2} + 2\nu\Delta t \left\|\nabla z^{1}\right\|_{0}^{2} \leq 2\left\|z^{0}\right\|_{0}^{2} + 4(\Delta t)^{2}\left\|\tau_{1}^{1}\right\|_{0}^{2} + 2\Delta t\nu\left\|\nabla \tau_{2}^{1}\right\|_{0}^{2}.$$

Together with (13), it follows that for $n \ge 1$

$$\begin{aligned} \|z^{n}\|_{0}^{2} + 2\nu \sum_{j=1}^{n} \Delta t \|\nabla z^{j}\|_{0}^{2} &\leq 17 \|z^{0}\|_{0}^{2} + 28(\Delta t)^{2} \|\tau_{1}^{1}\|_{0}^{2} + 2\sum_{j=2}^{n} \frac{\Delta t}{T} \|z^{j}\|_{0}^{2} + 2T \sum_{j=2}^{n} \Delta t \|\tau_{1}^{j}\|_{0}^{2} \\ &+ 14\Delta t\nu \|\nabla \tau_{2}^{1}\|_{0}^{2} + 2\nu \sum_{j=1}^{n} \Delta t \|\nabla \tau_{2}^{j}\|_{0}^{2}, \end{aligned}$$

from where (10) follows by applying Gronwall's Lemma [3, Lemma 5.1] for $\Delta t \leq T/4$.

Let $X = L^2(\Omega)$ and let us denote by $e_r^n = u_r^n - P^r u_h^n$ and by $\eta_h^n = P^r u_h^n - u_h^n$. Arguing as in the proof of [2, Theorem 4.6], one gets

$$(De_r^n, v) + \nu(\nabla e_r^n, \nabla v) = (\partial_t u_h^n - Du_h^n, v) - \nu(\nabla \eta_h^n, \nabla v), \quad \forall \ v \in U^r.$$
(14)

Lemma 4. The following bounds hold

$$\left\|\partial_{t}u_{h}^{1} - D^{1}u_{h}^{1}\right\|_{j} \leq \frac{\Delta t}{2} \max_{0 \leq t \leq t_{1}} \left\|\partial_{tt}u_{h}\right\|_{j}, \quad j = 0, 1,$$
(15)

$$\left\|\partial_{t}u_{h}^{n} - D^{2}u_{h}^{n}\right\|_{j} \leq \sqrt{5}(\Delta t)^{3/2} \left(\int_{t_{n-2}}^{t_{n}} \left\|\partial_{ttt}u_{h}(t)\right\|_{j}^{2} dt\right)^{1/2}, \quad n = 2, \dots, N, \ j = 0, 1.$$
(16)

Proof. For $D = D^1$, (15) follows easily from

$$\partial_t u_h^n - D u_h^n = \frac{1}{\Delta t} \int_{t_{n-1}}^{t_n} (\partial_t u_h(t_n) - \partial_t u_h(s)) \, ds = \frac{1}{\Delta t} \int_{t_{n-1}}^{t_n} \left(\int_s^{t_n} \partial_{tt} u_h(t) \, dt \right) \, ds.$$

For $D = D^2$, Taylor series expansion with integral reminder reveals that

$$\partial_t u_h^n - D u_h^n = \frac{1}{\Delta t} \int_{t_{n-2}}^{t_n} \left(2(t - t_{n-1})_+^2 - \frac{1}{2}(t - t_{n-2})^2 \right) \partial_{ttt} u_h \, dt,$$

where $x_{+} = \max(0, x)$, for $x \in \mathbb{R}$. Then, a straightforward calculation shows that

$$\|\partial_t u_h^n - Du_h^n\|_j \le \left(\frac{2}{\sqrt{5}} + \frac{2\sqrt{2}}{\sqrt{5}}\right) (\Delta t)^{3/2} \left(\int_{t_{n-2}}^{t_n} \|\partial_{ttt} u_h(t)\|_j^2 dt\right)^{1/2},$$

and then (16) follows by noticing that $2 + 2\sqrt{2} < 5$.

Lemma 5. Let $X = L^2(\Omega)$. It holds

$$\nu \sum_{j=1}^{n} \Delta t \|\nabla \eta_h^j\|_0^2 \le \nu T \|S\|_2 \left(2 + 4\tilde{C}\frac{T^2}{\tau^2}\right) \sum_{k=r+1}^{d} \lambda_k.$$
(17)

Proof. The proof of (17) follows easily by applying (4) and (8).

Theorem 1 (Bound for $X = L^2(\Omega)$). Let $X = L^2(\Omega)$, then it holds for $\Delta t \leq T/4$

$$\max_{1 \le n \le M} \|u_r^n - u^n\|_0^2 \le 60e^4 \left(\|e_r^0\|_0^2 + (\Delta t)^4 \max_{0 \le s \le \Delta t} \|\partial_{tt} u_h(s)\|_0^2 + T(\Delta t)^4 \int_0^T \|\partial_{ttt} u_h(s)\|_0^2 \, ds \right)
+ 3(1 + 14T\nu e^4 \|S\|_2) \left(2 + 4\tilde{C} \frac{T^2}{\tau^2} \right) \sum_{k=r+1}^d \lambda_k + 3C(u)^2 h^{2(l+1)}.$$
(18)

Proof. From (14) and (10), applying (15), (16) (noting that most integrals over time intervals $[t_{j-1}, t_j]$ appear twice when summing over n), and (17), we obtain

$$\begin{aligned} \|e_r^n\|_0^2 + \nu \sum_{j=1}^n \Delta t \|\nabla e_r^j\|_0^2 &\leq e^4 \left(17\|e_r^0\|_0^2 + 7(\Delta t)^4 \max_{0 \leq s \leq \Delta t} \|\partial_{tt} u_h(s)\|_0^2 \\ + 20T(\Delta t)^4 \int_0^T \|\partial_{ttt} u_h(s)\|_0^2 ds + 14\nu T \|S\|_2 \left(2 + 4\tilde{C}\frac{T^2}{\tau^2} \right) \sum_{k=r+1}^d \lambda_k \right). \end{aligned}$$

To simplify, we replace the factors 17 and 7 by 20. To finish the proof, apply the decomposition $u_r^n - u^n = (u_r^n - P_h u_h^n) + (P_h u_h^n - u_h^n) + (u_h^n - u^n)$, followed by (8) and (2).

Let $X = H_0^1(\Omega)$. Arguing as in the proof of [2, Theorem 4.1] yields

$$(De_r^n, v) + \nu(\nabla e_r^n, \nabla v) = (\partial_t u_h^n - P_r(Du_h^n), v), \quad \forall v \in U^r.$$

Applying Lemma 3 with $z^n = e_r^n$, $\tau_1^n = \partial_t u_h^n - P_r D u_h^n$ and $\tau_2 = 0$ we get

$$\|e_r^n\|_0^2 + 2\nu \sum_{j=1}^n \Delta t \|\nabla e_r^j\|_0^2 \le e^4 \left(17\|e_r^0\|_0^2 + 28(\Delta t)^2 \|\tau_1^1\|_0^2 + 2\Delta tT \sum_{j=2}^N \|\tau_1^n\|_0^2 \right).$$
(19)

Theorem 2 (Bound for $X = H_0^1(\Omega)$). Let $X = H_0^1(\Omega)$, $\Delta t \leq T/4$, and $C_1 = 4e^4 (10 + \Delta t/T) + 2 + 4\tilde{C}$. Then it holds

$$\max_{1 \le n \le M} \|u_r^n - u^n\|_0^2 \le 60e^4 \left(\|e_r^0\|_0^2 + (\Delta t)^4 \max_{0 \le s \le \Delta t} \|\partial_{tt} u_h(s)\|_0^2 + 2(\Delta t)^4 \int_0^T \|\partial_{ttt} u_h(t)\|_0^2 dt \right) \\ + 3C_1 C_p^2 \left(\frac{T}{\tau}\right)^2 \sum_{j=r+1}^d \lambda_k + 3C^2(u)h^{2(l+1)}.$$

$$(20)$$

Proof. The last two terms on the right-hand side of (19) are bounded by the triangle inequality

$$\|\tau_1^n\|_0^2 = \|\partial_t u_h^n - P^r (Du_h^n)\|_0^2 \le 2 \|\partial_t u_h^n - (Du_h^n)\|_0^2 + 2 \|(I - P^r) (Du_h^n)\|_0^2.$$
(21)

For n = 1, the first term is bounded by (15) and the second one by (1) and (3), giving

$$(\Delta t)^2 \|\tau_1^1\|_0^2 \le \frac{(\Delta t)^4}{2} \max_{0 \le s \le \Delta t} \|\partial_{tt} u_h(s)\|_0^2 + \frac{4T}{\tau^2} C_p^2 \Delta t \sum_{k=r+1}^d \lambda_k.$$

For $n \ge 2$, the first term of (21) is estimated by (16). To bound the other term observe that -2 n (a (a) = 1

$$D^{2}u_{h}^{n} = (3/2)D^{1}u^{n} - (1/2)D^{1}u^{n-1},$$

and, consequently,

$$2 \left\| (I - P^r) \left(D^2 u_h^n \right) \right\|_0^2 \le \frac{9}{2} \left\| (I - P^r) \left(D^1 u_h^n \right) \right\|_0^2 + \frac{1}{2} \left\| (I - P^r) \left(D^1 u_h^n \right) \right\|_0^2,$$

so that, by using (1) and (3), one obtains

$$2T\sum_{j=2}^{n} \Delta t \left\| (I - P^{r}) \left(D^{2} u_{h}^{n} \right) \right\|_{0}^{2} \leq 10T\sum_{j=1}^{n} \Delta t \left\| (I - P^{r}) \left(D^{1} u_{h}^{n} \right) \right\|_{0}^{2} \leq 20C_{p}^{2} \left(\frac{T}{\tau} \right)^{2} \sum_{j=r+1}^{d} \lambda_{k}.$$

Collecting the estimates for n = 1 and $n \ge 2$ leads to

$$\begin{aligned} \|e_{r}^{n}\|_{0}^{2} + 2\nu \sum_{j=1}^{n} \Delta t \|\nabla e_{r}^{j}\|_{0}^{2} &\leq e^{4} \left(17\|e_{r}^{0}\|_{0}^{2} + 14(\Delta t)^{4} \max_{0 \leq s \leq \Delta t} \|\partial_{tt}u_{h}(s)\|_{0}^{2} \\ &+ 40(\Delta t)^{4} \int_{0}^{T} \|\partial_{ttt}u_{h}(t)\|_{0}^{2} dt + 4\left(10 + \frac{\Delta t}{T}\right) C_{p}^{2} \frac{T^{2}}{\tau^{2}} \sum_{j=r+1}^{d} \lambda_{k} \right). \end{aligned}$$
Now, the proof is finished in the same way as the proof of Theorem 1.

Now, the proof is finished in the same way as the proof of Theorem 1.

Second order error bounds in time of form (18) and (20) can be derived if the finite differences in U are replaced with the temporal derivatives $\{\partial_t u_h^n\}_{n=0}^M$, with only slight modifications in the analysis. If the set of snapshots is $\{u_h^n\}_{n=0}^M$, then a second order estimate for $\sum_{j=1}^M \Delta t \|u_r^n - u^n\|_0^2$ can be shown along the lines of the presented analysis but neither pointwise estimates nor optimal estimates in the H^1 norm can be obtained.

References

- [1] S. L. Eskew and J. R. Singler. A new approach to proper orthogonal decomposition with difference quotients. Adv. Comput. Math., 49(2):Paper No. 13, 33, 2023.
- [2] B. García-Archilla, V. John, and J. Novo. Pod-roms for incompressible flows including snapshots of the temporal derivative of the full order solution. SIAM J. Numer. Anal., 2023.
- [3] J. G. Heywood and R. Rannacher. Finite-element approximation of the nonstationary Navier-Stokes problem. IV. Error analysis for second-order time discretization. SIAM J. Numer. Anal., 27(2):353-384, 1990.
- [4] B. Koc, S. Rubino, M. Schneier, J. Singler, and T. Iliescu. On optimal pointwise in time error bounds and difference quotients for the proper orthogonal decomposition. SIAM J. Numer. Anal., 59(4):2163–2196, 2021.
- [5] K. Kunisch and S. Volkwein. Galerkin proper orthogonal decomposition methods for parabolic problems. Numer. Math., 90(1):117-148, 2001.
- [6] J. Novo and S. Rubino. Error analysis of proper orthogonal decomposition stabilized methods for incompressible flows. SIAM J. Numer. Anal., 59(1):334-369, 2021.