

Tetrad Fields, Reference Frames, and the Gravitational Energy-Momentum in the Teleparallel Equivalent of General Relativity

J. W. Maluf^{(a)(1)}, F. L. Carneiro^{(b)(2)},
S. C. Ulhoa^{(c)(1)} and J. F. da Rocha-Neto^{(d)(1)}

- (1) Universidade de Brasília, Instituto de Física,
70.910-900 Brasília DF, Brazil
(2) Universidade Federal do Norte do Tocantins,
Centro de Ciências Integradas, 77824-838 Araguaína TO, Brazil

Abstract

We review the concept and definitions of the energy-momentum and angular momentum of the gravitational field in the teleparallel equivalent of general relativity (TEGR). The importance of these definitions is justified by three major reasons. First, the TEGR is a well established and widely accepted formulation of the gravitational field, whose basic field strength is the torsion tensor of the Weitzenböck connection. Second, in the phase space of the TEGR there exists an algebra of the Poincaré group. Not only the definitions of the gravitational energy-momentum and 4-angular momentum satisfy this algebra, but also the first class constraints related to these definitions satisfy the algebra. And third, numerous applications of these definitions lead to physically consistent results. These definitions follow from a well established Hamiltonian formulation, and rely on the idea of localization of the gravitational energy. In this review we revisit the concept of localizability of the gravitational energy, in light of results obtained in recent years. We have studied the behaviour of free particles in the space-time of plane fronted gravitational waves (pp-waves). Free particles are here understood as particles that are not subject to external

forces other than the gravitational acceleration due to pp-waves. Since these particles acquire or loose kinetic energy locally, the transfer of energy from or to the gravitational field must also be localized. We consider this theoretical result an important and definite argument in favour of the localization of the gravitational energy-momentum, and by extension, of the gravitational 4-angular momentum.

- (a) jwmaluf@gmail.com, wadih@unb.br
- (b) fernandolessa45@gmail.com
- (c) sc.ulhoa@gmail.com
- (d) jfrocha@unb.br, jfrn74@yahoo.com

Contents

1	Introduction	1
2	Tetrad fields as reference frames in space-time	8
2.1	Fermi-Walker transported frames	12
2.2	Static frames in the Schwarzschild space-time	15
2.3	Frame in free fall in the Schwarzschild space-time	18
2.4	Static frames in the Kerr space-time	19
3	The gravitational energy-momentum definitions in the TEGR	23
3.1	Main results from the Hamiltonian formulation	25
3.2	Gravitational energy-momentum from the Lagrangian field equations	28
4	Gravitational energy in the Kerr space-time	30
5	Distribution of gravitational energy in the Schwarzschild-de Sitter space-time	35
6	Gravitational energy-momentum in the Bondi-Sachs space-time	39
7	Plane gravitational waves, the kinetic energy of free particles and localization of the gravitational energy	48
8	Final Remarks	55

1 Introduction

The concept of energy is fundamental and indispensable in physics. Energy is a quantity that can *potentially* produce motion of particles or physical bodies. We recall the statement by Galileo, according to which “*My purpose is to set forth a very new science dealing with a very ancient subject. There is, in nature, perhaps nothing older than motion*”, which implies that energy is, likewise, one of the oldest manifestations in nature. Energy is an attribute of basically all particles and fields in the universe, but the energy

of the gravitational field is, surprisingly, an issue whose concept and definition are still in debate and discussion in the physics community. There are various approaches to the definition of the energy of the gravitational field, in part because there are several descriptions of the relativistic gravitational field that are equivalent to each other, and also in part because there are misconceptions about the idea of localization of the gravitational energy. All (or most) the descriptions mentioned above lead (or are equivalent) to Einstein's field equations, and for this reason there is an ongoing debate about which definitions for the energy, momentum and 4-angular momentum of the gravitational field are correct.

The traditional and dominant view of the concept of gravitational energy takes place in the metric formulation of general relativity. According to this view, there does not exist an energy-momentum (or stress-energy) tensor for the gravitational field, because the gravitational field "can always be locally set to zero", and therefore the gravitational energy cannot be localized. The argument is the following.

I. It is always possible to choose a "locally inertial frame" (this is essentially a choice of coordinates) where the gravitational "force" vanishes.

II. This result is accomplished by arguing that the Christoffel symbols vanish on a space-time event or over a timelike geodesic, and thus the metric tensor is locally flat.

III. Finally, it is argued that this vanishing is a manifestation of the principle of equivalence in general relativity. Note that, here, the manifestation of the principle of equivalence is due to (and obtained by) a choice of coordinates, despite the fact that a choice of coordinates on an arbitrary manifold should not have relevant physical (dynamical) consequences.

As a result, many physicists adhere (or feel comfortable) to the use of pseudo-tensors, of which there are numerous definitions in the literature. Pseudo-tensors are used in many analyses of the energy of gravitational waves. The difficulties mentioned above regarding a definition for the stress-energy tensor of the gravitational field gave rise to the concept of quasi-local *mass*, which is related to the energy enclosed by a 2-surface in space. This quantity, in turn, is not associated to any form of energy density, or scalar density on the manifold. The applicability of the definition of quasi-local gravitational mass has its own difficulties, as we will discuss ahead in this

review. It must be emphasized that the vanishing mentioned above of the Christoffel symbols takes place on a space-time event or, more generally, along any geodesic, timelike or spacelike geodesic.

The fact that we may set the Christoffel symbols to vanish on a spacelike geodesic means that this vanishing is a just feature of differential geometry [3]. It is certainly not a consequence of any principle in physics, including the principle of equivalence. In the teleparallel equivalent of general relativity (TEGR), the Christoffel symbols may be constructed, of course, but they do not play any major role in the theory. They may be defined as a secondary, or auxiliary concept. The field strength of the TEGR is the torsion tensor of the Weitzenböck connection, and the torsion tensor cannot be made to vanish by means of a coordinate transformation.

Einstein's formulation of the principle of equivalence has been discussed at length by Norton [1], and readdressed in Ref. [2]. In summary, we just highlight two difficulties presented in Ref. [1], that allows to contradict the statements **I**, **II**, **III** above regarding the impossibility of having a localized definition for the gravitational energy. The first difficulty is that the version of the principle of equivalence that is widely adopted in the literature, as the one displayed above, is due to Pauli [1], and is rather distinct from Einstein's version of the principle. According to Pauli's formulation, “in every infinitely small neighborhood of the space-time, there always exists a coordinate system in which gravitation has no influence either on the motion of particles or any other physical processes”. This conclusion is justified by the vanishing of Christoffel symbols at a point within this neighborhood. But we know that what really vanishes at this point are the first derivatives of the metric tensor, not the second or higher derivatives. Thus, the vanishing of the gravitational field even at the space-time point considered is questionable. We quote here a statement from Ref. [1] (and reproduced in Ref. [2]):

It has rarely been acknowledged that Einstein never endorsed the principle of equivalence which results, here called the “infinitesimal principle of equivalence”. Moreover, his early correspondence (with Pauli) contains a devastating objection to this principle: in infinitesimal regions of the space-time manifold it is impossible to distinguish geodesics from many other curves and therefore impossible to decide whether a point mass is in free fall.

Thus, in infinitesimal regions of the space-time, it is impossible to assert whether gravitation has or has not influence on the motion of free particles.

The second difficulty presented in Ref. [1] is that Einstein considered the principle of equivalence as a consequence of frame transformations, not coordinate transformations. His formulation may be summarized as follows [1].

Consider a reference frame K (a Galilean system) and a reference frame K' , which is uniformly accelerated with respect to K . Then one asks whether an observer in K' must understand his condition as accelerated, or whether there remains a point of view according to which he can interpret his condition as at “rest”. Einstein concludes that by assuming the existence of a homogeneous gravitational field in K' it is possible to consider the latter as at rest. In his words: The assumption that one may treat K' as at rest, in all strictness without any laws of nature not being fulfilled with respect to K' , I call the Principle of Equivalence.

The concepts of frame transformations and coordinate transformations are very much clear in general relativity, and are different from each other. Frame transformations act on tetrad fields, for instance. “*The happiest thought of my life*”, asserted by Einstein and thoroughly described in the book by Schucking and Surowitz [4], is realized by means of a frame transformation, it is an actual free fall, and definitely is not a coordinate transformation. The principle of equivalence is a very beautiful idea, but the eminent physicist Synge [5] claimed that he does not understand it as a principle for general relativity, because either there is or there is not a gravitational field in any arbitrary neighborhood of the space-time, since other relevant tensors such as the Riemann-Christoffel or torsion tensors do not vanish as a consequence of a coordinate transformation. Synge’s argument is logically sound. The existence or removal of the gravitational field cannot depend on a choice of coordinates.

In conclusion, the arguments **I**, **II** and **III** presented above regarding the impossibility of the existence of a localized expression for the gravitational energy are at the same time very weak and insufficient, and place no impediment for the localization of the gravitational energy-momentum.

This review addresses the gravitational energy-momentum and 4-angular momentum in the context of the TEGR. This theory has been investigated by several authors over the years, see Refs. [6]-[23]. It turns out that the TEGR is a suitable formulation for the relativistic gravitational field, that yields interesting results in cosmology, in the framework of modified $f(T)$

theories [24]-[29]. In this review we will adopt the definitions and notations of Ref. [13]. In recent times, we have presented several results in the literature regarding the consistency and plausibility of the definitions of energy-momentum and 4-angular momentum of the gravitational field in the TEGR. All definitions are based on densities which are well behaved under coordinate transformations, and therefore our definitions are localized.

The definitions in consideration here do not rely on abstract mathematical elaborations whose purpose is to yield the Schwarzschild mass, for instance. Rather, they follow from the field equations of the TEGR. The Lagrangian field equations allow the establishment of the gravitational energy-momentum 4-vector P^a , the flux (balance) equations and the fluxes definitions, but the complete structure formed by the gravitational energy-momentum and 4-angular momentum is obtained in the realm of the Hamiltonian formulation. As we will discuss ahead, the 6 extra degrees of freedom of the tetrad field (which has 16 a priori degrees of freedom, compared to the 10 degrees of freedom of the metric tensor) allow the definition of the 3-angular momentum and of the 3-centre of mass momentum of the gravitational field. These latter definitions follow from the Hamiltonian first class constraints of the theory. Thus, all definitions to be considered in this review are consequence of the Hamiltonian field equations of the theory, as we will show.

In recent years, we have investigated the trajectories of idealised free particles in the presence of plane-fronted gravitational waves (pp-waves). The pp-waves are very simple solutions of Einstein's field equations in vacuum, even simpler than the Schwarzschild solution. The idealised particles in consideration are particles with negligible mass, that do not influence the gravitational field around them, and are “free” in the sense that they are subject only to the gravitational field of the pp-waves, i.e., they follow geodesics in space-time. In a series of articles [30, 31, 32, 33] we found that the free particles undergo a velocity memory effect, and also that they gain or loose kinetic energy after the passage of the pp-wave. Since the only physical entity in question is the pp-wave, there is an energy transfer between the particles and the gravitational field of the waves. Evidently, this energy transfer takes place at the position of the particle, and thus the gravitational energy transfer also takes place at the localization of the pointwise particle. It is obvious, therefore, that energy of the gravitational field of the pp-wave must be localized. It does not make sense to relate this energy transfer to closed spatial 2-surfaces of arbitrary radius around the particle. Needless to say, it is even

more absurd to relate the energy transfer to the information storaged on a 2-surface at spatial infinity, if the total gravitational energy in question were an ADM type energy [34], for instance.

We consider this feature of the physical system pp-wave + idealised particle as a strong and definite argument in favour of the localization of the gravitational energy. This argument holds true also in the context of linearised gravitational waves, where the velocity memory effect is also predicted to take place, but which is not considered here.

The plan of this review is the following. In Section 2 we will review the concept of tetrad fields as frames adapted to arbitrary observers in space-time. This interpretation is the primary attribute of tetrad fields. Contrary to many applications in the literature, where tetrad fields are treated as just a collection of four orthonormal 4-vectors in space-time, here the tetrad fields are adapted (attached) to the worldline of an observer. The inevitable conclusion is that tetrad fields provide information of both the observer and the gravitational field. In Section 3 we present both the Lagrangian and Hamiltonian formulations of the TEGR, the definitions of the energy-momentum and 4-angular momentum of the gravitational field, and the Poincaré algebra between these definitions, in the phase space of the theory. To our knowledge, the Hamiltonian formulation of the TEGR is the only framework where the algebra of the Poincaré group is realized by means of the Poisson brackets between the energy-momentum and 4-angular momentum definitions.

We will review only three major applications of the definitions of gravitational energy-momentum. In Section 4 we will readdress the gravitational energy enclosed by the external event horizon of the Kerr black hole. This energy has also been investigated by the quasi-local approach. This is an opportunity to compare our result with the latter and to show that the result obtained in the TEGR is quite remarkable. We will show that energy contained within the external event horizon of the Kerr black hole is extremely close to $2M_{irr}$, where M_{irr} is the irreducible mass of the black hole.

In Section 5 we will consider the Schwarzschild-de Sitter space-time, with the purpose of analysing the distribution of gravitational energy in the vicinity of the cosmological constant Λ . The sharp increase of the gravitational energy density as one approaches the cosmological horizon is a feature that shares similarities with the dark energy problem, in spite of the fact that we are considering a static physical configuration. In Section 6 we readdress the Bondi-Sachs space-time, characterised by the mass aspect M and by two functions, c and d . The two news functions $\partial_0 c$ and $\partial_0 d$ carry information

both at spacelike and null infinities, and therefore they should contribute to the total energy of the space-time. That is indeed what we find in the context of the TEGR. The total energy of the Bondi-Sachs space-time is normally given by the mass function $m(u)$ only, which is an integral of the mass aspect, but here we show that the news functions also contribute, since they are manifestations of the Bondi-Sachs line element at spatial infinity. Section 7 is devoted to the study of geodesics of free particles in the space-time of plane-fronted gravitational waves. Only few examples suffice to demonstrate that the kinetic energy of free particles increases or decreases with the passage of some simple constructions of pp-waves. The results of this section not only support the idea of localization of gravitational energy, but they show that the latter is mandatory. And finally, in Section 8 we present our conclusions.

Notation:

1. Space-time indices μ, ν, \dots and SO(3,1) (Lorentz) indices a, b, \dots run from 0 to 3. Time and space indices are indicated according to $\mu = 0, i$, $a = (0), (i)$.
2. The tetrad fields are represented by $e^a{}_\mu$, and the torsion tensor by $T_{a\mu\nu} = \partial_\mu e_{a\nu} - \partial_\nu e_{a\mu}$. The flat, tangent space Minkowski space-time metric tensor raises and lowers tetrad indices, and is fixed by $\eta_{ab} = e_{a\mu} e_{b\nu} g^{\mu\nu} = (-1, +1, +1, +1)$.
3. The frame components are given by the inverse tetrads $\{e_a{}^\mu\}$, although we may as well refer to $\{e^a{}_\mu\}$ as the frame. The determinant of the tetrad fields is represented by $e = \det(e^a{}_\mu)$

The torsion tensor defined above is often related to the object of anholonomy $\Omega^\lambda{}_{\mu\nu}$ via $\Omega^\lambda{}_{\mu\nu} = e_a{}^\lambda T^a{}_{\mu\nu}$. However, we assume that the space-time geometry is defined by the tetrad fields only, and in this case the only possible nontrivial definition for the torsion tensor is given by $T^a{}_{\mu\nu}$. This torsion tensor is related to the antisymmetric part of the Weitzenböck connection $\Gamma^\lambda_{\mu\nu} = e^{a\lambda} \partial_\mu e_{a\nu}$, which establishes the Weitzenböck space-time. The metric and torsion-free Christoffel symbols are denoted by ${}^0\Gamma^\lambda_{\mu\nu}$, and the associated torsion-free Levi-Civita connection by ${}^0\omega_{\mu ab}$, in the equations below. The parameter c denotes the speed of light, but sometimes it will be omitted by assuming $c = 1$.

2 Tetrad fields as reference frames in space-time

A set of tetrad fields is a collection of four orthonormal, linearly independent vector fields in space-time, $\{e^{(0)}_\mu, e^{(1)}_\mu, e^{(2)}_\mu, e^{(3)}_\mu\}$. They may constitute the local reference frame of an observer that moves along a trajectory C , represented by the worldline $x^\mu(\tau)$ [35, 36, 37]. The parameter τ is the proper time of the observer. The components $e^{(0)}_\mu$ and $e^{(i)}_\mu$ are timelike and spacelike vectors, respectively; e^a_μ transforms as covariant vector fields under coordinate transformations of the space-time manifold, and as contravariant vector fields under SO(3,1) (Lorentz) transformations, i.e., $\tilde{e}^a_\mu = \Lambda^a{}_b e^b_\mu$, where the matrices $\{\Lambda^a{}_b\}$ are representations of the SO(3,1) group and satisfy $\Lambda^a{}_c \Lambda^b{}_d \eta_{ab} = \eta_{cd}$. The metric tensor $g_{\mu\nu}$ is obtained by the standard relation $e^a_\mu e^b_\nu \eta_{ab} = g_{\mu\nu}$.

Tetrad fields e^a_μ allow the projection of vectors and tensors in space-time in the local frame of an observer. Vectors and tensors components are abstract, space-time dependent quantities. In order to measure field quantities with magnitude and direction, an observer must project these quantities on the frame carried by the observer. The projection of a vector $V^\mu(x)$ at a position x^μ , on a particular frame, is simply given by $V^a(x) = e^a_\mu(x) V^\mu(x)$.

Given a worldline C of an observer, represented by $x^\mu(\tau)$, the velocity of the observer along C is denoted by $u^\mu(\tau) = dx^\mu/d\tau$. We identify the observer's velocity with the $a = (0)$ component of $e_a{}^\mu$. Thus, $e_{(0)}{}^\mu = u^\mu(\tau)/c$. The acceleration a^μ of the observer is given by the absolute derivative of u^μ along C ,

$$a^\mu = \frac{Du^\mu}{d\tau} = c \frac{De_{(0)}{}^\mu}{d\tau} = c u^\alpha \nabla_\alpha e_{(0)}{}^\mu, \quad (1)$$

where the covariant derivative is constructed out of the Christoffel symbols ${}^0\Gamma_{\alpha\beta}^\mu$. The last equality is obtained as follows,

$$\begin{aligned} \frac{De_{(0)}{}^\mu}{d\tau} &= \frac{de_{(0)}{}^\mu}{d\tau} + {}^0\Gamma_{\alpha\beta}^\mu \frac{dx^\alpha}{d\tau} e_{(0)}{}^\beta \\ &= \frac{dx^\alpha}{d\tau} \frac{\partial e_{(0)}{}^\mu}{\partial x^\alpha} + {}^0\Gamma_{\alpha\beta}^\mu \frac{dx^\alpha}{d\tau} e_{(0)}{}^\beta \\ &= u^\alpha \nabla_\alpha e_{(0)}{}^\mu. \end{aligned} \quad (2)$$

Thus, $e_a{}^\mu$ allows to obtain the velocity and acceleration of an observer along the worldline. Therefore, for a given set of tetrad fields, $e_{(0)}{}^\mu$ describes a congruence of timelike curves, which determines a field of observers. These observers are characterized by the velocity field $u^\mu = c e_{(0)}{}^\mu$, and are endowed with acceleration a^μ . If $e^a{}_\mu \rightarrow \delta_\mu^a$ in the spacelike limit $r \rightarrow \infty$, then $e^a{}_\mu$ is adapted to static observers at spatial infinity.

The geometrical characterization of the whole set of tetrad fields as an observer's frame may be given by considering the acceleration of the frame along an arbitrary path $x^\mu(\tau)$ of an observer. The acceleration of the whole frame is determined by the absolute derivative of $e_a{}^\mu$ along $x^\mu(\tau)$. Thus, assuming that the observer carries an orthonormal tetrad frame $e_a{}^\mu$, the acceleration of the frame along the path is given by [38, 39]

$$\frac{De_a{}^\mu}{d\tau} = \phi_a{}^b e_b{}^\mu, \quad (3)$$

where ϕ_{ab} is the antisymmetric acceleration tensor. According to Refs. [38, 39], in analogy with the Faraday tensor we may identify $\phi_{ab} \rightarrow (\mathbf{a}/c, \boldsymbol{\Omega})$, where \mathbf{a} is the translational acceleration ($\phi_{(0)(i)} = a_{(i)}/c$) and $\boldsymbol{\Omega}$ is the frequency of rotation of the local spatial frame with respect to a non-rotating, Fermi-Walker transported frame. It follows from Eq. (3) that

$$\phi_a{}^b = e^b{}_\mu \frac{De_a{}^\mu}{d\tau} = e^b{}_\mu u^\lambda \nabla_\lambda e_a{}^\mu. \quad (4)$$

The acceleration vector a^μ defined by Eq. (1) may be projected on a frame. The result is

$$a^b = e^b{}_\mu a^\mu = c e^b{}_\mu u^\alpha \nabla_\alpha e_{(0)}{}^\mu = c \phi_{(0)}{}^b. \quad (5)$$

Thus, a^μ and $\phi_{(0)(i)}$ are not different translational accelerations of the frame.

The expression of a^μ given by Eq. (1) may be rewritten as

$$\begin{aligned} a^\mu/c &= u^\alpha \nabla_\alpha e_{(0)}{}^\mu = u^\alpha \nabla_\alpha u^\mu = \frac{dx^\alpha}{d\tau} \left(\frac{\partial u^\mu}{\partial x^\alpha} + {}^0\Gamma_{\alpha\beta}^\mu u^\beta \right) \\ &= \frac{d^2 x^\mu}{d\tau^2} + {}^0\Gamma_{\alpha\beta}^\mu \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau}, \end{aligned} \quad (6)$$

where ${}^0\Gamma_{\alpha\beta}^\mu$ are the Christoffel symbols. We see that if $u^\mu = c e_{(0)}{}^\mu$ represents a geodesic trajectory, then the frame is in free fall and $a^\mu/c = 0 = \phi_{(0)(i)}$.

Therefore we conclude that non-vanishing values of the latter quantities do represent *inertial* (i.e., non-gravitational) accelerations of the frame. A static frame in an asymptotically flat space-time is an example of a frame that is subject to inertial accelerations.

Since the tetrads are orthonormal vectors, we may rewrite Eq. (4) as $\phi_a^b = -u^\lambda e_a^\mu \nabla_\lambda e^b_\mu$, where $\nabla_\lambda e^b_\mu = \partial_\lambda e^b_\mu - {}^0\Gamma_{\lambda\mu}^\sigma e^b_\sigma$. Now we take into account the identity

$$\partial_\lambda e^b_\mu - {}^0\Gamma_{\lambda\mu}^\sigma e^b_\sigma + {}^0\omega_\lambda^b{}_c e^c_\mu \equiv 0,$$

where ${}^0\omega_\lambda^b{}_c$ is the metric compatible, torsion-free Levi-Civita connection,

$$\begin{aligned} {}^0\omega_{\mu ab} &= -\frac{1}{2}e^c_\mu(\Omega_{abc} - \Omega_{bac} - \Omega_{cab}), \\ \Omega_{abc} &= e_{a\nu}(e_b^\mu \partial_\mu e_c^\nu - e_c^\mu \partial_\mu e_b^\nu), \end{aligned} \quad (7)$$

and express ϕ_a^b as

$$\phi_a^b = c e_{(0)}^\mu({}^0\omega_\mu^b{}_a). \quad (8)$$

At last we consider the *identity* ${}^0\omega_\mu^a{}_b = -K_\mu^a{}_b$, where $K_\mu^a{}_b$ is the contortion tensor defined by

$$K_{\mu ab} = \frac{1}{2}e_a^\lambda e_b^\nu(T_{\lambda\mu\nu} + T_{\nu\lambda\mu} + T_{\mu\lambda\nu}), \quad (9)$$

and $T_{\lambda\mu\nu} = e^a_\lambda T_{a\mu\nu}$ (see Eq. (4) of Ref. [40]; the identity may be obtained by direct calculation). After simple manipulations, we finally obtain

$$\phi_{ab} = \frac{c}{2}[T_{(0)ab} + T_{a(0)b} - T_{b(0)a}] = -\phi_{ba}. \quad (10)$$

The expression above is clearly not invariant under local SO(3,1) transformations, but is invariant under coordinate transformations.

The values of ϕ_{ab} for given tetrad fields may be used to characterize the frame. We interpret ϕ_{ab} as the inertial accelerations along the trajectory $x^\mu(\tau)$. Therefore, given any set of tetrad fields for an arbitrary space-time, its geometrical interpretation may be obtained,

(i) either by suitably identifying $e_{(0)}^\mu = u^\mu/c$, together with the orientation in the three-dimensional space of the components $e_{(1)}^\mu, e_{(2)}^\mu, e_{(3)}^\mu$, considering the symmetry of the physical configuration, or,

(ii) by the values of the acceleration tensor $\phi_{ab} = -\phi_{ba}$, which characterize the inertial state of the frame.

The condition $e_{(0)}^\mu = u^\mu/c$ fixes only the three components $e_{(0)}^1, e_{(0)}^2, e_{(0)}^3$, because the component $e_{(0)}^0$ is determined by the normalization condition $u^\mu u^\nu g_{\mu\nu} = -c^2$. In both cases, the fixation of the frame requires the fixation of 6 components of the tetrad fields.

Two simple, familiar and straightforward applications of Eqs. (1-10) are the following:

(1) The tetrad fields adapted to observers at rest in Minkowski space-time is given by $e^a_\mu(ct, x, y, z) = \delta_\mu^a$. By means of a time-dependent boost in the x direction, say, the tetrad fields read

$$e^a_\mu(ct, x, y, z) = \begin{pmatrix} \gamma & -\beta\gamma & 0 & 0 \\ -\beta\gamma & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad (11)$$

where $\gamma = (1 - \beta^2)^{-1/2}$, $\beta = v/c$ and $v = v(t)$. In the expression above a and μ label rows and columns, respectively. The frame above is adapted to observers whose four-velocity is $u^\mu(t, x, y, z) = c e_{(0)}^\mu = (c\gamma, c\beta\gamma, 0, 0)$. After simple calculations we obtain

$$\begin{aligned} \phi_{(0)(1)} &= c \frac{d}{dx^0} [\beta\gamma] = \frac{d}{dt} \left[\frac{v/c}{\sqrt{1 - v^2/c^2}} \right], \\ \phi_{(0)(2)} &= 0, \\ \phi_{(0)(3)} &= 0, \end{aligned} \quad (12)$$

and $\phi_{(i)(j)} = 0$.

(2) The frame adapted to an observer at rest in Minkowski space-time whose four-velocity is $u^\mu = (c, 0, 0, 0)$, endowed with spatial axes that rotate around the z axis, say, with angular velocity $\omega(t)$, reads

$$e^a_\mu(ct, x, y, z) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \omega(t) & -\sin \omega(t) & 0 \\ 0 & \sin \omega(t) & \cos \omega(t) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (13)$$

After simple calculations we obtain

$$\begin{aligned}\phi_{(2)(3)} &= 0, \\ \phi_{(3)(1)} &= 0, \\ \phi_{(1)(2)} &= -c \frac{d\omega}{dx^0},\end{aligned}\tag{14}$$

and $\phi_{(0)(i)} = 0$. The examples above support the interpretation of ϕ_{ab} as the inertial accelerations of the frame.

2.1 Fermi-Walker transported frames

Fermi-Walker transported frames define a standard of non-rotation for accelerated observers. These are frames for which the frequency of rotation $\phi_{(i)(j)}$ vanishes. The vanishing of $\phi_{(i)(j)}$ is the maximum requirement one can impose on an arbitrary frame, in any space-time, regarding its rotational state. Here we show how to transform an arbitrary frame into a Fermi-Walker transported frame, by means of a local Lorentz transformation. We will adopt the notation, presentation and results of Ref. [5].

The absolute derivative of a vector V^μ along a world-line C is defined as usual by

$$\frac{DV^\mu}{d\tau} = \frac{dV^\mu}{d\tau} + {}^0\Gamma_{\alpha\beta}^\mu V^\alpha \frac{dx^\beta}{d\tau}.\tag{15}$$

We consider four 4-vectors, A^μ , B^μ , C^μ and D^μ that satisfy the following equations and properties:

$$\frac{DA^\mu}{d\tau} = bB^\mu,\tag{16}$$

$$\frac{DB^\mu}{d\tau} = cC^\mu + bA^\mu,\tag{17}$$

$$\frac{DC^\mu}{d\tau} = dD^\mu - cB^\mu,\tag{18}$$

$$\frac{DD^\mu}{d\tau} = -dC^\mu,\tag{19}$$

where $A^\mu A_\mu = -1$, $B^\mu B_\mu = C^\mu C_\mu = D^\mu D_\mu = 1$, and b, c, d are non-negative coefficients. Starting with the vector A^μ , Eq. (16) defines B^μ , Eq. (17)

defines C^μ and Eq. (18) defines D^μ . Equation (19) is verified in view (i.e., it is a consequence) of Eqs. (16-18). It is not difficult to verify that Eqs. (16-19) imply that A^μ, B^μ, C^μ and D^μ form an orthonormal set of vectors, i.e., $A^\mu B_\mu = 0$, etc.

We may identify A^μ with the unit vector tangent to the trajectory C , $A^\mu = dx^\mu/ds$. In this case, A^μ, B^μ, C^μ and D^μ establish the Frenet-Serret frame, and Eqs. (16-19) are called the Frenet-Serret equations. B^μ, C^μ and D^μ are the first, second and third normals to C , and b, c, d are the first, second and third curvatures of C , respectively [5]. Particular values of the coefficients b, c, d yield special curves in space-time. For instance, if $b = c = d = 0$, the curve C is a geodesic; if $b = \text{constant}$ and $c = d = 0$, C represents a hyperbola, and if $b = \text{constant}$, $c = \text{constant}$ and $d = 0$, then C is a helix [5].

Let us consider a timelike trajectory C represented by $x^\mu = x^\mu(\tau)$, in a space-time determined by the metric tensor $g_{\mu\nu}$. The Fermi-Walker transport of a vector F^μ along C is defined by [5]

$$\frac{DF^\mu}{d\tau} = bF_\alpha(A^\mu B^\alpha - A^\alpha B^\mu). \quad (20)$$

Given the value $F^\mu(\tau_0)$ at a certain initial position τ_0 , Eq. (20) allows the determination of F^μ along C . The Fermi-Walker transport of a second rank tensor along C is defined by

$$\frac{DT^{\mu\nu}}{d\tau} = bT_\alpha{}^\nu(A^\mu B^\alpha - A^\alpha B^\mu) + bT^\mu{}_\alpha(A^\nu B^\alpha - A^\alpha B^\nu). \quad (21)$$

It follows from the equation above that

$$\frac{Dg^{\mu\nu}}{d\tau} = 0 = \frac{D\delta_\nu^\mu}{d\tau}. \quad (22)$$

The velocity vector $A^\mu = dx^\mu/d\tau = u^\mu$ naturally undergoes Fermi-Walker transport. Application of Eq. (20) to A^μ yields Eq. (16).

It is easy to see that the scalar product of two vectors is preserved under the Fermi-Walker transport. Let ϕ represent the scalar product of the vectors Σ^μ and Ψ_μ . Along C we have

$$\begin{aligned} \phi(\tau + d\tau) - \phi(\tau) &= \Sigma^\mu(\tau + d\tau)\Psi_\mu(\tau + d\tau) - \Sigma^\mu(\tau)\Psi_\mu(\tau) \\ &= \Psi_\mu(\delta^{FW}\Sigma^\mu) + \Sigma^\mu(\delta^{FW}\Psi_\mu), \end{aligned} \quad (23)$$

where

$$\begin{aligned}\delta^{FW} \Sigma^\mu &= -{}^0\Gamma_{\alpha\beta}^\mu \Sigma^\beta dx^\alpha + b\Sigma_\alpha (A^\mu B^\alpha - A^\alpha B^\mu) d\tau, \\ \delta^{FW} \Psi_\mu &= {}^0\Gamma_{\alpha\mu}^\lambda \Psi_\lambda dx^\alpha + b\Psi_\alpha (A_\mu B^\alpha - A^\alpha B_\mu) d\tau.\end{aligned}\quad (24)$$

Equations (23) and (24) lead to $\phi(\tau + d\tau) - \phi(\tau) = 0$.

In what follows, we identify the velocity vector A^μ on C with the timelike component of the tetrad fields $e_{(0)}^\mu$ according to

$$\frac{1}{c} A^\mu = \frac{1}{c} \frac{dx^\mu}{d\tau} = e_{(0)}^\mu. \quad (25)$$

The important feature of this identification is that the Fermi-Walker transport of $e_{(0)}^\mu$ along C guarantees that $e_{(0)}^\mu$ will always be tangent to C . The spacelike components $e_{(k)}^\mu$ are everywhere orthogonal to $e_{(0)}^\mu$. Along C , the components $e_{(k)}^\mu$ also undergo Fermi-Walker transport. Given that $e_{(k)}^\mu$ and $e_{(0)}^\mu$ are orthogonal, we have from Eq. (20) that

$$\frac{De_{(k)}^\mu}{d\tau} = b e_{(0)}^\mu e_{(k)\lambda}^\lambda B_\lambda. \quad (26)$$

Taking into account Eq. (16), the equation above may be rewritten as

$$\frac{De_{(k)}^\mu}{d\tau} = c e_{(0)}^\mu e_{(k)\lambda}^\lambda \frac{De_{(0)}^\lambda}{d\tau}. \quad (27)$$

In the equation above, c is the speed of light. Equation (27) determines the transport of the orthonormal basis e_a^μ along an arbitrary timelike curve C , such that $e_{(0)}^\mu$ is always tangent to C .

In view of the definition given by Eq. (1), Eq. (27) may be expressed as

$$\frac{De_{(k)}^\mu}{d\tau} = u^\mu e_{(k)\lambda}^\lambda a^\lambda. \quad (28)$$

Making $b = (k)$ in Eq. (5), we find $e_{(k)\lambda}^\lambda = a_{(k)} = \phi_{(0)(k)}$. Thus the Fermi-Walker transport of the frame may be written as

$$\frac{De_{(k)}^\mu}{d\tau} = u^\mu \phi_{(0)(k)}. \quad (29)$$

On the other hand, it is easy to verify that the total acceleration of the frame components $e_{(k)}^\mu$ given by Eq. (3) may be expressed in terms of $\phi_{(0)(k)}$ and $\phi_{(j)(k)}$ according to

$$\frac{D e_{(k)}{}^\mu}{d\tau} = u^\mu \phi_{(0)(k)} + \phi_{(k)}{}^{(j)} e_{(j)}{}^\mu. \quad (30)$$

Therefore if $\phi_{(j)(k)} = 0$, the frame is Fermi-Walker transported and is non-rotating. It turns out that we can impose $\phi_{(j)(k)} = 0$, at least formally.

Suppose that a frame is given such that $\phi_{(j)(k)} \neq 0$. In terms of the torsion tensor components, the quantities $\phi_{(j)(k)}$ are written as

$$\phi_{(i)(j)} = \frac{1}{2} [e_{(i)}{}^\mu e_{(j)}{}^\nu T_{(0)\mu\nu} + e_{(0)}{}^\mu e_{(j)}{}^\nu T_{(i)\mu\nu} - e_{(0)}{}^\mu e_{(i)}{}^\nu T_{(j)\mu\nu}]. \quad (31)$$

Under a local Lorentz transformation of the spatial components, we have

$$\tilde{e}_{(i)}{}^\mu = \Lambda_{(i)}{}^{(k)} e_{(k)}{}^\mu, \quad (32)$$

$$\begin{aligned} \tilde{T}_{(i)\mu\nu} &= \partial_\mu \tilde{e}_{(i)\nu} - \partial_\nu \tilde{e}_{(i)\mu} \\ &= \Lambda_{(i)}{}^{(k)} T_{(k)\mu\nu} + [\partial_\mu \Lambda_{(i)}{}^{(k)}] e_{(k)\nu} - [\partial_\nu \Lambda_{(i)}{}^{(k)}] e_{(k)\mu}. \end{aligned} \quad (33)$$

The coefficients $\{\Lambda_{(i)}{}^{(j)}(x)\}$ of the spatial components of the local Lorentz transformation are fixed by requiring $\tilde{\phi}_{(i)(j)} = 0$.

It is possible to show that for given non-vanishing values of the quantities $\phi_{(j)(k)}$, the condition $\tilde{\phi}_{(i)(j)} = 0$ is obtained provided the coefficients $\{\Lambda_{(i)}{}^{(j)}\}$ of the Lorentz transformation satisfy the equation [37]

$$e_{(0)}{}^\mu \Lambda_{(m)}{}^{(j)} \partial_\mu \Lambda_{(j)(k)} - \phi_{(k)(m)} = 0. \quad (34)$$

Thus, given an arbitrary frame, it is always possible, at least formally, to rotate the frame and obtain a Fermi-Walker transported frame for which $\tilde{\phi}_{(i)(j)} = 0$. We note that the local Lorentz transformation (32) does not affect the timelike component $e_{(0)}{}^\mu$.

2.2 Static frames in the Schwarzschild space-time

In this and in the following subsections we will analyse some distinguished frames. We will show that the values of the acceleration tensor are in agreement with the physical nature and interpretation of the frames, and have well defined physical meaning. In these subsections we will adopt the speed of light $c = 1$.

A frame is static in the Schwarzschild space-time provided it undergoes an inertial acceleration that exactly cancels the gravitational acceleration on the frame. This is an interesting example because it allows an alternative derivation of the well known gravitational acceleration of a free particle in the Schwarzschild space-time.

In spherical coordinates the Schwarzschild space-time is described by the line element

$$ds^2 = -\left(1 - \frac{2m}{r}\right)dt^2 + \left(1 - \frac{2m}{r}\right)^{-1}dr^2 + r^2d\theta^2 + r^2 \sin^2\theta d\phi^2. \quad (35)$$

A field of static observers in this space-time is characterized by the vector field u^μ such that $u^\mu = (u^0, 0, 0, 0)$, i.e., the spatial components of u^μ vanish. Consequently, in the construction of the tetrad fields we require

$$e_{(0)}{}^i = u^i = 0. \quad (36)$$

In view of the orthogonality of the tetrad components, this condition implies $e^{(k)}{}_0 = 0$. A simple form of $e_{a\mu}$ in (t, r, θ, ϕ) coordinates that satisfies this property and yields Eq. (35) is given by

$$e_{a\mu} = \begin{pmatrix} -\beta & 0 & 0 & 0 \\ 0 & \alpha \sin\theta \cos\phi & r \cos\theta \cos\phi & -r \sin\theta \sin\phi \\ 0 & \alpha \sin\theta \sin\phi & r \cos\theta \sin\phi & r \sin\theta \cos\phi \\ 0 & \alpha \cos\theta & -r \sin\theta & 0 \end{pmatrix}, \quad (37)$$

where

$$\begin{aligned} \alpha &= \left(1 - \frac{2m}{r}\right)^{-1/2}, \\ \beta &= \left(1 - \frac{2m}{r}\right)^{1/2}. \end{aligned} \quad (38)$$

Recall that a and μ label lines and rows, respectively. It is possible to show that in the asymptotic limit $r \rightarrow \infty$ the inverse tetrad components in (t, x, y, z) coordinates satisfy

$$\begin{aligned} e_{(1)}{}^\mu(t, x, y, z) &\cong (0, 1, 0, 0), \\ e_{(2)}{}^\mu(t, x, y, z) &\cong (0, 0, 1, 0), \\ e_{(3)}{}^\mu(t, x, y, z) &\cong (0, 0, 0, 1). \end{aligned} \quad (39)$$

Altogether, conditions (36) and (39) fix 6 degrees of the frame.

The evaluation of the acceleration tensor ϕ_{ab} is straightforward. After a number of manipulations we find that Eq. (37) represents a non-rotating frame, i.e.,

$$\phi_{(i)(j)} = 0. \quad (40)$$

The translational acceleration, however, is non-vanishing. From definition (10) we find

$$\phi_{(0)(i)} = T_{(0)(0)(i)} = e_{(0)}{}^\mu e_{(i)}{}^\nu T_{(0)\mu\nu}. \quad (41)$$

For $a = (0)$, the only non-vanishing component of $T_{a\mu\nu}$ is $T_{(0)01} = \partial_1 \beta$. Thus, the equation above yields

$$\phi_{(0)(i)} = g^{00} g^{11} e_{(0)0} e_{(i)1} T_{(0)01}, \quad (42)$$

from what follows

$$\begin{aligned} \phi_{(0)(1)} &= \frac{m}{r^2} \left(1 - \frac{2m}{r}\right)^{-1/2} \sin \theta \cos \phi, \\ \phi_{(0)(2)} &= \frac{m}{r^2} \left(1 - \frac{2m}{r}\right)^{-1/2} \sin \theta \sin \phi, \\ \phi_{(0)(3)} &= \frac{m}{r^2} \left(1 - \frac{2m}{r}\right)^{-1/2} \cos \theta. \end{aligned} \quad (43)$$

We define the acceleration \mathbf{a} ,

$$\mathbf{a} = \phi_{(0)(1)} \hat{\mathbf{x}} + \phi_{(0)(2)} \hat{\mathbf{y}} + \phi_{(0)(3)} \hat{\mathbf{z}}, \quad (44)$$

which may be written as

$$\mathbf{a} = \frac{m}{r^2} \left(1 - \frac{2m}{r}\right)^{-1/2} \hat{\mathbf{r}}, \quad (45)$$

where

$$\hat{\mathbf{r}} = \sin \theta \cos \phi \hat{\mathbf{x}} + \sin \theta \sin \phi \hat{\mathbf{y}} + \cos \theta \hat{\mathbf{z}}.$$

Equation (45) represents the inertial acceleration necessary to maintain the frame in stationary state in space-time. Therefore it must necessarily

cancel the geodesic gravitational acceleration that is exerted on the frame. In fact,

$$a = -\frac{m}{r^2} \left(1 - \frac{2m}{r}\right)^{-1/2}, \quad (46)$$

is precisely the standard geodesic acceleration of a body in free fall in the Schwarzschild space-time, which may be obtained directly from Eq. (1) [41].

We have seen that Eqs. (36) and (39) fix the six degrees of freedom of the tetrad frame (37). Alternatively, the set of conditions (40), (43) also fix the frame. The sets (40), (43) and (36), (39) are essentially equivalent. This equivalence is in agreement with the discussion presented in the paragraph just below Eq. (10), namely, either we establish the frame by fixing the frame components according to the symmetry of the physical configuration, and which amounts to choosing a suitable congruence of timelike worldlines, or by fixing the components of the acceleration tensor (which, in general, is a less intuitive procedure). We remark, finally, that Eq. (41) is invariant under coordinate transformations.

2.3 Frame in free fall in the Schwarzschild space-time

A frame in free fall in the Schwarzschild space-time is radially accelerated towards the center of the black hole, under the action of the gravitational field only.

An observer that is in radial free fall in the Schwarzschild space-time is endowed with the four-velocity [41]

$$u^\alpha = \left[\left(1 - \frac{2m}{r}\right)^{-1}, -\left(\frac{2m}{r}\right)^{1/2}, 0, 0 \right]. \quad (47)$$

The simplest set of tetrad fields that satisfies the condition

$$e_{(0)}{}^\alpha = u^\alpha, \quad (48)$$

is given by [36]

$$e_{a\mu} = \begin{pmatrix} -1 & -\alpha^2 \eta & 0 & 0 \\ \eta \sin \theta \cos \phi & \alpha^2 \sin \theta \cos \phi & r \cos \theta \cos \phi & -r \sin \theta \sin \phi \\ \eta \sin \theta \sin \phi & \alpha^2 \sin \theta \sin \phi & r \cos \theta \sin \phi & r \sin \theta \cos \phi \\ \eta \cos \theta & \alpha^2 \cos \theta & -r \sin \theta & 0 \end{pmatrix}, \quad (49)$$

where

$$\eta = \left(\frac{2m}{r} \right)^{1/2} = (1 - \alpha^{-2})^{1/2}. \quad (50)$$

Since the frame is in free fall, the equation $\phi_{(0)(i)} = 0$ is verified. It is not difficult to show by direct calculations that this set of tetrad fields also satisfies the conditions

$$\phi_{(i)(j)} = \frac{1}{2}[T_{(0)(i)(j)} + T_{(i)(0)(j)} - T_{(j)(0)(i)}] = 0. \quad (51)$$

Equation (48) fixes 3 conditions on the frame, i.e., $e_{(0)}^i = u^i$. Together with the 3 conditions fixed by Eq. (51), we have six conditions on the frame. These conditions completely fix the structure of the tetrad fields, even though Eq. (51) has been verified *a posteriori*. Therefore Eq. (49) describes a non-rotating frame in radial free fall in the Schwarzschild space-time. No inertial acceleration is imparted to the frame.

2.4 Static frames in the Kerr space-time

Another interesting physical configuration is the frame adapted to static observers in the Kerr space-time. In the weak field approximation, a relationship may be established with the gravitoelectromagnetic (GEM) field quantities [36]. In spherical (Boyer-Lindquist) coordinates, the Kerr's space-time is established by the line element

$$ds^2 = -\frac{\psi^2}{\rho^2}dt^2 - \frac{2\chi \sin^2 \theta}{\rho^2}d\phi dt + \frac{\rho^2}{\Delta}dr^2 + \rho^2 d\theta^2 + \frac{\Sigma^2 \sin^2 \theta}{\rho^2}d\phi^2, \quad (52)$$

with the following definitions:

$$\begin{aligned} \Delta &= r^2 + a^2 - 2mr, \\ \rho^2 &= r^2 + a^2 \cos^2 \theta, \\ \Sigma^2 &= (r^2 + a^2)^2 - \Delta a^2 \sin^2 \theta, \\ \psi^2 &= \Delta - a^2 \sin^2 \theta, \\ \chi &= 2amr. \end{aligned} \quad (53)$$

A static frame in Kerr's space-time is defined by the congruence of time-like curves $u^\mu(\tau)$ such that $u^i = 0$, namely, the spatial velocity of the observers is zero with respect to static observers at spacelike infinity. Since we identify $e_{(0)}^i = u^i$, a static reference frame is established by the condition

$$e_{(0)}^i = 0. \quad (54)$$

As in subsection 2.2, the orthogonality of the tetrads imply $e^{(k)}_0 = 0$. This latter equation remains valid even after a local rotation of the frame: $\tilde{e}^{(k)}_0 = \Lambda^{(k)}_{(j)} e^{(j)}_0 = 0$. Therefore, condition (54) determines the static character of the frame, up to an orientation of the spacelike components of the frame in the three-dimensional space (see Eq. (59) below).

A simple form of the tetrad fields that satisfies Eq. (54) (or, equivalently, $e^{(k)}_0 = 0$) reads [36]

$$e_{a\mu} = \begin{pmatrix} -A & 0 & 0 & -B \\ 0 & C \sin \theta \cos \phi & \rho \cos \theta \cos \phi & -D \sin \theta \sin \phi \\ 0 & C \sin \theta \sin \phi & \rho \cos \theta \sin \phi & D \sin \theta \cos \phi \\ 0 & C \cos \theta & -\rho \sin \theta & 0 \end{pmatrix}, \quad (55)$$

with the following definitions,

$$\begin{aligned} A &= \frac{\psi}{\rho}, \\ B &= \frac{\chi \sin^2 \theta}{\rho \psi}, \\ C &= \frac{\rho}{\sqrt{\Delta}}, \\ D &= \frac{\Upsilon}{\rho \psi}. \end{aligned} \quad (56)$$

In the expression of D we have

$$\Upsilon = (\psi^2 \Sigma^2 + \chi^2 \sin^2 \theta)^{1/2}. \quad (57)$$

We are interested in obtaining the acceleration tensor ϕ_{ab} , and for this purpose we need the inverse tetrad fields e_a^μ . They read

$$e_a{}^\mu = \begin{pmatrix} \frac{\rho}{\psi} & \frac{\rho\chi}{\psi\Upsilon} \sin\theta \sin\phi & -\frac{\rho\chi}{\psi\Upsilon} \sin\theta \cos\phi & 0 \\ 0 & \frac{\sqrt{\Delta}}{\rho} \sin\theta \cos\phi & \frac{\sqrt{\Delta}}{\rho} \sin\theta \sin\phi & \frac{\sqrt{\Delta}}{\rho} \cos\theta \\ 0 & \frac{1}{\rho} \cos\theta \cos\phi & \frac{1}{\rho} \cos\theta \sin\phi & -\frac{1}{\rho} \sin\theta \\ 0 & -\frac{\rho\psi}{\Upsilon} \frac{\sin\phi}{\sin\theta} & \frac{\rho\psi}{\Upsilon} \frac{\cos\phi}{\sin\theta} & 0 \end{pmatrix}, \quad (58)$$

where now the index a labels the columns and μ labels the rows.

The frame determined by Eqs. (55) and (56) is valid in the region outside the ergosphere. The function $\psi^2 = \Delta - a^2 \sin^2\theta$ vanishes over the external surface of the ergosphere (defined by $r = r^* = m + \sqrt{m^2 - a^2 \cos^2\theta}$; over this surface we have $g_{00} = 0$). Various components of Eqs. (55) and (56) are not well defined when $r = r^*$. It is well known that it is not possible to maintain static observers inside the ergosphere of the Kerr space-time.

It follows from Eq. (58) that for large values of r we have

$$e_{(3)}{}^\mu(t, r, \theta, \phi) \cong (0, \cos\theta, -(1/r)\sin\theta, 0),$$

or

$$e_{(3)}{}^\mu(t, x, y, z) \cong (0, 0, 0, 1). \quad (59)$$

Therefore we may assert that the frame given by Eq. (55) is characterized by the following properties: **(i)** the frame is static, because Eq. (54) is verified; **(ii)** the $e_{(3)}{}^\mu$ components are oriented along the symmetry axis of the black hole (the z direction). The second condition is ultimately responsible for the simple form of Eq. (55).

The evaluation of ϕ_{ab} requires more algebraic manipulations than the previous cases, but is straightforward. It is useful to define the vectors

$$\begin{aligned} \hat{\mathbf{r}} &= \sin\theta \cos\phi \hat{\mathbf{x}} + \sin\theta \sin\phi \hat{\mathbf{y}} + \cos\theta \hat{\mathbf{z}}, \\ \hat{\theta} &= \cos\theta \cos\phi \hat{\mathbf{x}} + \cos\theta \sin\phi \hat{\mathbf{y}} - \sin\theta \hat{\mathbf{z}}, \end{aligned} \quad (60)$$

which have well defined meaning as unit vectors in the asymptotic limit $r \rightarrow \infty$. The translational acceleration and the frequency of rotation are denoted in Cartesian components as

$$\mathbf{a} = (\phi_{(0)(1)}, \phi_{(0)(2)}, \phi_{(0)(3)}), \quad (61)$$

$$\boldsymbol{\Omega} = (\phi_{(2)(3)}, \phi_{(3)(1)}, \phi_{(1)(2)}). \quad (62)$$

We obtain the following expressions for \mathbf{a} and $\boldsymbol{\Omega}$ [36]:

$$\mathbf{a} = \frac{m}{\psi^2} \left[\frac{\sqrt{\Delta}}{\rho} \left(\frac{2r^2}{\rho^2} - 1 \right) \hat{\mathbf{r}} + \frac{2ra^2}{\rho^3} \sin \theta \cos \theta \hat{\theta} \right], \quad (63)$$

$$\boldsymbol{\Omega} = -\frac{\chi}{\Upsilon\rho} \cos \theta \hat{\mathbf{r}} + \frac{\psi^2 \sqrt{\Delta}}{2\Upsilon\rho} \sin \theta \partial_r \left(\frac{\chi}{\psi^2} \right) \hat{\theta} - \frac{\psi^2}{2\Upsilon\rho} \sin \theta \partial_\theta \left(\frac{\chi}{\psi^2} \right) \hat{\mathbf{r}}. \quad (64)$$

These vectors also characterize the frame determined by Eq. (55). The fixation of \mathbf{a} and $\boldsymbol{\Omega}$ is equivalent to the fixation of six components of the tetrad fields. Equations (63) and (64) represent the inertial accelerations that one must exert on the frame in order to ensure (i) the static character of the frame (Eq. (54)), and that (ii) the $e_{(3)}^\mu$ components of the tetrad fields asymptotically coincide with the symmetry axis of the black hole (Eq. (59)).

The form of \mathbf{a} and $\boldsymbol{\Omega}$ for large values of r is very interesting. It is easy to verify that in the limit $r \rightarrow \infty$ we obtain

$$\mathbf{a} \cong \frac{m}{r^2} \hat{\mathbf{r}}, \quad (65)$$

$$\boldsymbol{\Omega} \cong -\frac{am}{r^3} \left(2 \cos \theta \hat{\mathbf{r}} + \sin \theta \hat{\theta} \right). \quad (66)$$

By identifying $m \leftrightarrow q$ and $4\pi ma \leftrightarrow \bar{m}$, where q is the electric charge and \bar{m} is the magnetic dipole moment, equations (65) and (66) resemble the electric field of a point charge and the magnetic field of a perfect dipole that points in the z direction, respectively. These equations represent a manifestation of gravitoelectromagnetism.

If we remove all inertial accelerations on the frame, an observer located at a position (r, θ, ϕ) will be subject to a radial acceleration $-\mathbf{a}$ and to a frequency of rotation determined by $-\boldsymbol{\Omega} = \boldsymbol{\Omega}_D$, which is the dragging frequency of the frame. Thus, the gravitomagnetic effect is locally equivalent to inertial effects in a frame endowed with angular frequency $-\boldsymbol{\Omega}_D$. This is precisely the gravitational Larmor's theorem, discussed in Ref. [42].

In view of the standard examples discussed above, it is quite clear that the primary and natural attribute of tetrad fields is the establishment of reference frames in space-time. The inevitable conclusion is that tetrad fields carry information of both the inertial state of the observer and of the gravitational field.

3 The gravitational energy-momentum definitions in the TEGR

In the ordinary metric formulation of general relativity, the gravitational energy-momentum was first given by the standard ADM expressions [34], but the complete set of energy, momentum, angular momentum and centre of mass moment was probably put forward in a concise form by Regge and Teitelboim [43]. These notions were presented in the context of the Hamiltonian formulation of general relativity. The idea was to require the variation of the total Hamiltonian to be well defined in an asymptotically flat space-time, where the standard asymptotic space-time translations and 4-rotations are considered as coordinate transformations at spacelike infinity. This requirement leads to the addition of boundary (surface) terms to the primary Hamiltonian, so that the latter has well defined functional derivatives, and therefore one may obtain the field equations in the Hamiltonian framework (Hamilton's equations) by means of a consistent procedure. In this way, one arrives at the total energy, momentum, angular momentum and centre of mass moment of the gravitational field, given by surface terms of the total Hamiltonian. The work of Regge and Teitelboim indicate the relevance of the Hamiltonian formulation for the establishment of these definitions.

In this Section, we will recall the definitions of energy-momentum P^a and of the 4-angular momentum L^{ab} for the gravitational field in the TEGR. These definitions have been discussed several times in the literature. Here, we will display the summary presented in Refs. [13, 44]. In the present review, the TEGR is constructed out of the tetrad fields only, as we pointed out earlier. In the last Section, we will briefly discuss the so called “covariant” formulation, which encompasses the addition of a flat spin connection.

The first relevant consideration is an identity between the scalar curvature of the Riemann-Christoffel tensor, and an invariant combination of quadratic terms of the torsion tensor,

$$eR(e) \equiv -e \left(\frac{1}{4} T^{abc} T_{abc} + \frac{1}{2} T^{abc} T_{bac} - T^a T_a \right) + 2\partial_\mu(eT^\mu), \quad (67)$$

where $T_a = T^b{}_{ba}$ and $T_{abc} = e_b{}^\mu e_c{}^\nu T_{a\mu\nu}$. The Lagrangian density for the gravitational field in the TEGR is given by [10]

$$L(e) = -k e \left(\frac{1}{4} T^{abc} T_{abc} + \frac{1}{2} T^{abc} T_{bac} - T^a T_a \right) - \frac{1}{c} L_M$$

$$\equiv -ke\Sigma^{abc}T_{abc}-\frac{1}{c}L_M, \quad (68)$$

where $k = c^3/(16\pi G)$, L_M represents the Lagrangian density for the matter fields, and Σ^{abc} is defined by

$$\Sigma^{abc}=\frac{1}{4}\left(T^{abc}+T^{bac}-T^{cab}\right)+\frac{1}{2}\left(\eta^{ac}T^b-\eta^{ab}T^c\right). \quad (69)$$

Thus, the Lagrangian density is geometrically equivalent to the scalar curvature density. The variation of $L(e)$ with respect to $e^{a\mu}$ yields the field equations

$$e_{a\lambda}e_{b\mu}\partial_\nu(e\Sigma^{b\lambda\nu})-e(\Sigma^{b\nu}_aT_{b\nu\mu}-\frac{1}{4}e_{a\mu}T_{bcd}\Sigma^{bcd})=\frac{1}{4kc}eT_{a\mu}, \quad (70)$$

where $T_{a\mu}$ is defined by $\delta L_M/\delta e^{a\mu}=eT_{a\mu}$. These field equations are equivalent to Einstein's equations. It is possible to verify by direct calculations that the equations above can be rewritten as

$$\frac{1}{2}[R_{a\mu}(e)-\frac{1}{2}e_{a\mu}R(e)]=\frac{1}{4kc}T_{a\mu}, \quad (71)$$

The left and right hand sides of Eq. (67) are invariant under arbitrary SO(3,1) local transformations. However, the Lagrangian density (68) does not contain the total divergence that appears on the right hand side of (67). Therefore, the Lagrangian density (68) is not invariant under local Lorentz transformations, but the field equations are covariant under such transformations. As discussed in Ref. [10], the variation of the action integral constructed out of (68) is well defined if one considers asymptotically flat space-times, i.e., one does not need the addition of surface terms to obtain well defined variations in the asymptotic limit $r \rightarrow \infty$.

Equations (70) and (71) prove the equivalence between the TEGR and the standard metric formulation of GR. Both theories are equivalent because the field equations are the same. However, in a theory constructed out of tetrad fields, additional field quantities such as third rank tensors do exist (the torsion tensor, for instance), besides new vector and tensor densities, which cannot be defined in the standard metric formulation of gravity. These additional field quantities are covariant under global Lorentz transformations, but not under local transformations. It turns out that these vector and tensor densities yield the definitions of energy, momentum and 4-angular momentum of the gravitational field, which are covariant under

global Lorentz transformations, as expected. This general framework is not different from what happens in the ordinary formulation of arbitrary field theories, where the energy-momentum and 4-angular momentum are frame dependent field quantities that transform under the global Lorentz group. Thus, energy transforms as the zero component of the energy-momentum four-vector. These features must naturally hold also in the context of the theory for the gravitational field. Consider a static black hole represented by a mass parameter m only. For a distant observer, the total energy of this black hole is given by $E = mc^2$. But at very great distances, this black hole is considered as a particle of mass m , and if it moves with constant velocity v , then its total energy as seen by the same distant observer is $E = \gamma mc^2$, where $\gamma = (1 - v^2/c^2)^{-1/2}$. By the same reasoning, the gravitational momentum, angular momentum and the centre of mass moment are also frame dependent field quantities, whose values are different in different frames and for different observers. On physical grounds, energy, momentum, angular momentum and centre of mass moment cannot be local Lorentz *invariant* field quantities, since these quantities depend on the frame, as we know from special relativity, which is the limit of the general theory of relativity when the gravitational field is weak or negligible.

After some rearrangements, Eq. (70) may be written in the form [40]

$$\partial_\nu(e\Sigma^{a\mu\nu}) = \frac{1}{4k}ee^a{}_\nu(t^{\mu\nu} + \frac{1}{c}\mathbf{T}^{\mu\nu}), \quad (72)$$

where

$$t^{\mu\nu} = k(4\Sigma^{bc\mu}T_{bc}{}^\nu - g^{\mu\nu}\Sigma^{bcd}T_{bcd}), \quad (73)$$

is interpreted as the gravitational energy-momentum tensor [13, 40], and $\mathbf{T}^{\mu\nu} = e_a{}^\mu\mathbf{T}^{a\nu}$.

3.1 Main results from the Hamiltonian formulation

In this subsection we will consider the field equations in the absence of matter fields. The Hamiltonian density of the TEGR is constructed as usual in the phase space of the theory. We first note that the Lagrangian density (68) does not depend on the time derivatives of e_{a0} . Therefore, the latter arise as Lagrange multipliers in the Hamiltonian density H . The momenta canonically conjugated to e_{a0} are denoted by Π^{a0} . The latter are primary constraints of the theory: $\Pi^{a0} \approx 0$. The momenta canonically conjugated to e_{ai} are given by $\Pi^{ai} = \delta L/\delta \dot{e}_{ai} = -4k\Sigma^{a0i}$. The Hamiltonian density H is

obtained by rewriting the Lagrangian density in the form $L = \Pi^{ai}\dot{e}_{ai} - H$, in terms of e_{ai} , Π^{ai} and Lagrange multipliers. After the Legendre transform is performed, we obtain the final form of the Hamiltonian density. It reads [45, 46]

$$H(e, \Pi) = e_{a0}C^a + \lambda_{ab}\Gamma^{ab}, \quad (74)$$

where λ_{ab} are Lagrange multipliers. In the equation above we have omitted a surface term. The expression of $C^a = \delta H/\delta e_{a0}$ in terms of the field variables is rather long. It is formally simplified in Eq. (76) below. The constraints $\Gamma^{ab} = -\Gamma^{ba}$ are defined by

$$\Gamma^{ab} = 2\Pi^{[ab]} + 4ke(\Sigma^{a0b} - \Sigma^{b0a}). \quad (75)$$

After solving the field equations, the Lagrange multipliers are identified as $\lambda_{ab} = (1/4)(T_{a0b} - T_{b0a} + e_a{}^0T_{00b} - e_b{}^0T_{00a})$ [45]. The constraints C^a may be written as

$$C^a = -\partial_i\Pi^{ai} - p^a = 0, \quad (76)$$

where p^a is an intricate expression of the field quantities. In the context of the Lagrangian density (68), p^a may be interpreted as the gravitational energy-momentum density (see Eq. (77) below). The quantities C^a and Γ^{ab} are first class constraints. They satisfy an algebra similar to the algebra of the Poincaré group [46].

The integral form of the constraint equations $C^a = 0$ yields the gravitational energy-momentum P^a [40, 47],

$$P^a = -\int_V d^3x \partial_i\Pi^{ai}, \quad (77)$$

where V is an arbitrary volume of the three-dimensional space and $\Pi^{ai} = -4k\Sigma^{a0i}$. By means of Gauss's law, we may transform the volume integral in Eq. (77) into a surface integral, according to

$$P^a = \oint_S dS_i 4k\Sigma^{a0i}, \quad (78)$$

where S is the 2-dimensional spacelike boundary of the volume V .

In similarity to the definition above, the definition of the gravitational 4-angular momentum follows from the integral form of the constraint equations $\Gamma^{ab} = 0$ [48]. It reads

$$L^{ab} = - \int_V d^3x 2\Pi^{[ab]}, \quad (79)$$

where

$$2\Pi^{[ab]} = (\Pi^{ab} - \Pi^{ba}) = -4ke(\Sigma^{a0b} - \Sigma^{b0a}). \quad (80)$$

However, it has been noted [49] that the second term on the right hand side of Eq. (75) can be rewritten as a total divergence, so that the constraints Γ^{ab} become

$$\Gamma^{ab} = 2\Pi^{[ab]} - 2k\partial_i[e(e^{ai}e^{b0} - e^{bi}e^{a0})] = 0. \quad (81)$$

Therefore, the definition of the total 4-angular momentum of the gravitational field L^{ab} may be given by an integral of a total divergence, in similarity to Eq. (77),

$$L^{ab} = \oint_S dS_i 2k[e(e^{ai}e^{b0} - e^{bi}e^{a0})]. \quad (82)$$

It is easy to show [48] that expressions (77) and (79) satisfy the algebra of the Poincaré group in the phase space of the theory,

$$\begin{aligned} \{P^a, P^b\} &= 0, \\ \{P^a, L^{bc}\} &= \eta^{ab}P^c - \eta^{ac}P^b, \\ \{L^{ab}, L^{cd}\} &= \eta^{ad}L^{cb} + \eta^{bd}L^{ac} - \eta^{ac}L^{db} - \eta^{bc}L^{ad}. \end{aligned} \quad (83)$$

The quantities in the left hand side of the expressions above are Poisson brackets evaluated in the phase space determined by the canonically conjugate field variables (e_{ai}, Π^{ai}) . In the evaluation of the Poisson brackets, we have used the following functional derivatives of P^a and L^{ab} ,

$$\begin{aligned} \frac{\delta L^{ab}}{\delta e_{ck}(z)} &= -\eta^{ac}\Pi^{bk}(z) + \eta^{bc}\Pi^{ak}(z), \\ \frac{\delta L^{ab}}{\delta \Pi_{ck}(z)} &= \delta_c^a e^b{}_k(z) - \delta_c^b e^a{}_k(z), \\ \frac{\delta P^a}{\delta e_{ck}(z)} &= 0, \\ \frac{\delta P^a}{\delta \Pi_{ck}(z)} &= - \int d^3x \delta_c^a \frac{\partial}{\partial x^k} \delta^3(x - z). \end{aligned} \quad (84)$$

Therefore, from a physical point of view, the interpretation of the quantities P^a and L^{ab} as gravitational energy-momentum and 4-angular momentum is consistent. It is well known that the algebra of the Poincaré group is intimately connected to energy, momentum and 4-angular momentum. The values of P^a and L^{ab} may be computed by means of Eqs. (78) and (82).

Definitions (77) and (79) are invariant under coordinate transformations of the three-dimensional space, under time reparametrizations, and under global $\text{SO}(3,1)$ transformations. The gravitational energy is the zero component of the energy-momentum four-vector P^a .

3.2 Gravitational energy-momentum from the Lagrangian field equations

The definition of gravitational energy-momentum may also be obtained in the Lagrangian framework, according to the procedure of Ref. [40]. Again, we assume $c = 1 = G$, and consider the field equations in the form given Eqs. (72) and (73). In view of the antisymmetry property $\Sigma^{a\mu\nu} = -\Sigma^{a\nu\mu}$, it follows that

$$\partial_\lambda \left[e e^a{}_\mu (t^{\lambda\mu} + T^{\lambda\mu}) \right] = 0. \quad (85)$$

The equation above yields the continuity (or balance) equation,

$$\frac{d}{dt} \int_V d^3x e e^a{}_\mu (t^{0\mu} + T^{0\mu}) = - \oint_S dS_j \left[e e^a{}_\mu (t^{j\mu} + T^{j\mu}) \right], \quad (86)$$

where S is the boundary of an arbitrary 3-dimensional volume V , as considered above. Therefore we (again) identify $t^{\lambda\mu}$ as the gravitational energy-momentum tensor [40],

$$P^a = \int_V d^3x e e^a{}_\mu (t^{0\mu} + T^{0\mu}), \quad (87)$$

as the *total* energy-momentum contained within the volume V of the 3-dimensional space,

$$\Phi_g^a = \oint_S dS_j (e e^a{}_\mu t^{j\mu}), \quad (88)$$

as the gravitational energy-momentum flux, and

$$\Phi_m^a = \oint_S dS_j (e e^a{}_\mu T^{j\mu}), \quad (89)$$

as the energy-momentum flux of matter [40]. In view of the field equations (72), Eq. (87) may be written as $P^a = -\int_V d^3x \partial_j \Pi^{aj}$, from what follows

$$P^a = - \oint_S dS_j \Pi^{aj}, \quad (90)$$

where $\Pi^{aj} = -4ke \Sigma^{a0j}$, in similarity to Eq. (78).

We establish Eqs. (78) and (82) as the definitions of the gravitational energy-momentum and gravitational 4-angular momentum, respectively. The reason for considering surface integrals is because the gravitational field on the surface of integration S contains all the information on the interior region, and of course the integral can be carried out more easily, even in the presence of singularities, admitting that the space-time has singularities. We cannot transform the volume integral (77) into the surface integral (78) by means of Gauss theorem, if the space-time has singularities, and for this reason Eq. (78) is conceptually more important than Eq. (77). But we recall that exactly this same difficulty occurs in the definition of the ADM gravitational energy-momentum, whose definitions are given by surface integrals and yield satisfactory and acceptable results in the context of asymptotically flat space-times. Therefore, Eq. (78) represents the gravitational energy-momentum within the surface S .

Definition (78) (or (90)) is invariant under coordinate transformations of the three-dimensional space and under time reparametrizations, which are basic requirements of any gravitational energy-momentum definition. Eq. (85) is a true energy-momentum conservation equation, in contrast to the analogous situation in the metric formulation of general relativity, where the covariant derivative of the energy-momentum tensor is not a strict conservation equation. We repeat and emphasize that in the ordinary formulation of arbitrary field theories, energy, momentum, angular momentum and the centre of mass moment are frame dependent field quantities, that transform under the global $SO(3,1)$ group. This property, which takes place in special relativity, is satisfied by definition (78). In particular, the energy E transforms as the zero component of the energy-momentum four-vector. Thus, we conclude that in the context of the TEGR, there is a smooth transition between the weak field limit of general relativity and special relativity. We have a well defined conservation equation for the matter fields, with or without the gravitational field, in arbitrary coordinate systems.

The components of the vector P^a are $(E/c, \mathbf{P})$. If we assume that the tetrad fields satisfy asymptotic boundary conditions,

$$e_{a\mu} \simeq \eta_{a\mu} + \frac{1}{2} h_{a\mu}(1/r), \quad (91)$$

at spatial infinity, i.e., in the limit $r \rightarrow \infty$, then the total gravitational energy $E = cP^{(0)}$ is the ADM energy,

$$E = \frac{c^4}{16\pi G} \int_{S \rightarrow \infty} dS_k (\partial_i h_{ik} - \partial_k h_{ii}) = E_{ADM}. \quad (92)$$

The expression above is just one of the consistent results that follow from the above definitions.

4 Gravitational energy in the Kerr space-time

The Kerr space-time plays a central role in general relativity. We may assume that practically all single, self gravitating systems in nature rotate around their own symmetry or principal inertia axes, and that the Kerr space-time represents the exterior region of these rotating compact objects. However, the Kerr line element represents a black hole and possesses the ergoregion, which is the region between the external event horizon ($r = r_+$, see below) and the ergosurface ($r = r_+^\star$). It is theoretically predicted that it is possible to extract energy from the region between $r = r_+^\star$ and $r = r_+$ by means of the Penrose process [50]. After the complete extraction of this energy, which is considered to be a rotational energy, when the black hole no longer rotates, there remains the irreducible mass of the black hole M_{irr} , which was calculated by Christodoulou and Ruffini [51]. It is also assumed that no form of energy can escape from the external event horizon of the black hole, during the complete Penrose process. For this reason, the evaluation of the gravitational energy contained within the external event horizon of the Kerr black hole is a relevant issue in the investigation of the localization of the gravitational energy. It is rather obvious that this energy cannot be made to vanish by means of coordinate transformations, or by any formulation of the principle of equivalence, not even at a point inside the external event horizon.

In order to calculate the energy contained within the external event horizon of the Kerr black hole, we need tetrad fields that are defined up to $r = r_+$, i.e., we must have observers that can reach the external event horizon, even under the dragging effect of the black hole, since inside the ergosphere it is not

possible to establish static frames. The tetrad fields considered in subsection 2.4 do not satisfy this requirement, since they are defined from $r = \infty$ up to $r = r_+^*$, which defines the external surface of the ergosphere. The set of tetrad fields that are defined from $r = \infty$ up to $r = r_+$ is the one that satisfies Schwinger's time gauge condition [52].

We will make use of the notation in Section 2.4, and in particular of Eqs. (52) and (53), but not of Eq. (54), as we will explain ahead. The Kerr black hole is characterized by the mass parameter m and by the angular momentum per unit mass $a = J/m$. In the Penrose process [50], the initial mass m and angular momentum J of the black hole vary by dm and dJ , respectively, such that $dm - \Omega_H dJ \geq 0$, where Ω_H is the angular velocity of the external event horizon of the black hole,

$$\Omega_H = \frac{a}{2mr_+} = \frac{a}{a^2 + r_+^2}. \quad (93)$$

The radius of the external event horizon is given by $r_+ = m + \sqrt{m^2 - a^2}$. During the process, the variation of the area A of the black hole satisfies $dA \geq 0$ (because the angular momentum parameter a decreases, and consequently r_+ increases). In the final stage of an idealized process, the mass of the black hole becomes the irreducible mass M_{irr} [51], defined by the relation $m^2 = M_{irr}^2 + J^2/(4M_{irr}^2)$, and the Kerr black hole becomes a Schwarzschild black hole. The expression of the irreducible mass is

$$M_{irr} = \frac{1}{2}\sqrt{r_+^2 + a^2}. \quad (94)$$

An analysis of various gravitational energy expressions for the Schwarzschild and Kerr black holes has been made long time ago in Ref. [53]. Taking into account various expressions for the gravitational mass, it was concluded that the mass contained within the event horizon of the Schwarzschild black hole is $2m$, for all considered definitions. The same value is obtained by the quasi-local expressions (as, for instance, the Brown-York expression [54]). Likewise, one expects that the mass contained within the external event horizon at the final stage of the complete Penrose process, when the black hole becomes non-rotating, is given by $2M_{irr}$, and the gravitational energy contained within the external event horizon of the black hole is, therefore, $2M_{irr}c^2$. In similarity to the Schwarzschild space-time, the total gravitational energy of the Kerr space-time finally becomes $M_{irr}c^2$, after the complete Penrose process.

We have already shown that the definition for the gravitational energy in the TEGR yields an expression for the energy contained within the external event horizon of the Kerr black hole that is strikingly close to $2M_{irr}c^2$. In this Section, we will review this result.

We proceed to establish the frame that satisfies Schwinger's time gauge condition. Here we assume $c = 1$, but the notation will be that of subsection 2.4. As we mentioned above, the frame must be defined such that the radial coordinate r runs from r_+ to infinity, i.e., the frame must be defined in the whole region outside the external event horizon, and consequently also inside the ergosphere of the black hole. The external surface of the ergosphere is defined by $r = r_+^* = m + \sqrt{m^2 - a^2 \cos^2 \theta}$. Inside the ergosphere all observers are necessarily dragged in circular motion by the gravitational field. The four-velocity of observers that circulate around the black hole, outside the external horizon, under the action of the gravitational field of the Kerr space-time, is given by

$$u^\mu(t, r, \theta, \phi) = \frac{\rho\Sigma}{(\psi^2\Sigma^2 + \chi^2 \sin^2 \theta)^{1/2}}(1, 0, 0, \frac{\chi}{\Sigma^2}), \quad (95)$$

where all functions are defined by Eqs. (52) and (53). It is possible to show that if we restrict the radial coordinate to $r = r_+$, the $\mu = 3$ component of Eq. (95) becomes

$$\frac{\chi}{\Sigma^2} = \frac{a}{2mr_+} = \frac{a}{a^2 + r_+^2} = \Omega_H,$$

The quantity

$$\omega(r) = -\frac{g_{03}}{g_{33}} = \frac{\chi}{\Sigma^2}, \quad (96)$$

is the dragging angular velocity of inertial frames.

The tetrad fields **(i)** that are adapted to observers whose four-velocities are given by Eq. (95), i.e., for which $e_{(0)}^\mu = u^\mu$, and consequently defined in the region $r > r_+$, **(ii)** whose $e_{(i)}^\mu$ components in Cartesian coordinates are asymptotically oriented along the unit vectors $\hat{\mathbf{x}}$, $\hat{\mathbf{y}}$, $\hat{\mathbf{z}}$, and **(iii)** that is asymptotically flat, is given by

$$e_{a\mu} = \begin{pmatrix} -A & 0 & 0 & 0 \\ B \sin \theta \sin \phi & C \sin \theta \cos \phi & D \cos \theta \cos \phi & -F \sin \theta \sin \phi \\ -B \sin \theta \cos \phi & C \sin \theta \sin \phi & D \cos \theta \sin \phi & F \sin \theta \cos \phi \\ 0 & C \cos \theta & -D \sin \theta & 0 \end{pmatrix}, \quad (97)$$

where

$$\begin{aligned} A &= \frac{(g_{03}g_{03} - g_{00}g_{33})^{1/2}}{(g_{33})^{1/2}}, \\ B &= -\frac{g_{03}}{(g_{33})^{1/2} \sin \theta}, \\ C &= (g_{11})^{1/2}, \\ D &= (g_{22})^{1/2}, \\ F &= \frac{(g_{33})^{1/2}}{\sin \theta}. \end{aligned} \quad (98)$$

These tetrad fields are the unique configuration that satisfies the above conditions, since six conditions are imposed on $e^a{}_\mu$. It satisfies Schwinger's time gauge condition $e_{(i)}{}^0 = 0$. This frame allows observers to reach the vicinity of the external event horizon of the Kerr black hole. Therefore we may evaluate the gravitational energy contained within any surface S determined by the condition $r > r_+$, and in particular in the limit $r \rightarrow r_+$. Expression (97) is precisely the same set of tetrad fields (Eq. (4.9)) considered in Ref. [47].

The energy contained within the external event horizon of the black hole is calculated by means of the $a = (0)$ component of Eq. (78) or (90),

$$P^{(0)} = E = - \oint_S dS_i \Pi^{(0)i} = - \oint_S d\theta d\phi \Pi^{(0)1}(r, \theta, \phi). \quad (99)$$

S is a surface of constant radius determined by the condition $r = r_+$ (recall that we are assuming $c = 1$). After a number of algebraic calculations we obtain [47]

$$E = m \left[\frac{\sqrt{2p}}{4} + \frac{6p - q^2}{4q} \ln \left(\frac{\sqrt{2p} + q}{p} \right) \right]. \quad (100)$$

The quantities p and λ are defined by

$$p = 1 + \sqrt{1 - q^2}, \quad a = q m, \quad 0 \leq q \leq 1.$$

Note that when $q = 0$, we have $E = 2m$. Equation (100) is functionally very different from $2M_{irr} = \sqrt{r_+^2 + a^2}$. However, the two expressions are very similar, as we can verify in Figure 1.




Figure 1: Gravitational energy in the interior of the external event horizon of the Kerr black hole, as function of $q = a/m$.

In Figure 1, we have plotted (i) E/m , where E is given by Eq. (100), and (ii) $2M_{irr}/m$, as functions of $a/m = q$. Both curves are parametrized by q , which varies from 0 to 1. The upper curve represents Eq. (100), and the lower curve represents $2M_{irr}$. The almost coincidence between the two expressions is striking, and is perhaps the major achievement of definition (90). It shows that Eq. (100) is in very close agreement with $2M_{irr}$, as expected. This result supports the idea of localization of the gravitational energy.

It is not the purpose of this review to address or critique alternative approaches that deal with the same investigation of this Section, but we may just compare our result with the one obtained by other means, notably the quasi-local expression for the gravitational energy. A recent proposal for the quasi-local gravitational energy was given by Wang and Yau [56, 57], and applied to the Kerr space-time [58] to address precisely the irreducible mass of the black hole. In the context of the latter references, it was made an

analysis similar to the one above that led to Figure 1. However, due to mathematical reasons and difficulties, the analysis can be carried out only in the range $0 \leq a \leq \sqrt{3}m/2$ for the values of $a = J/m$. A latter attempt to circumvent this problem was made in Ref. [59], by means of an even different approach to quasilocal mass. By inspecting Figure 3 of the latter reference, which is the equivalent to Figure 1 of the present analysis, we see a sharp and unnatural discontinuity of the curve at the value $a = \sqrt{3}m/2$. In addition, the value of the mass within the horizon exceeds $2m$ for values of a close to m . In our opinion, these are difficulties of the whole method based on quasi-local expressions for the gravitational energy, which cannot reproduce $2M_{\text{irr}}$ in the whole range of values of the parameter a . Furthermore, the latter approach to gravitational energy is based only on the metric tensor, and for this reason it should be independent of frames or observers, but not so long ago the need to introduce “quasi-local observers” was put forward in Ref. [60]. This attempt shows that, one way or another, observers and frames are necessary to address the notion of gravitational energy. After all, the observed mass of the black holes depends on the frame, in case the black hole is moving with respect to the frame. An implicit dependence on the frame already takes place at the very initial considerations of the Schwarzschild or Kerr line elements: the parameter m is the total mass of black hole in a frame where the black hole *is at rest*.

5 Distribution of gravitational energy in the Schwarzschild-de Sitter space-time

The main purpose of this section is to review the fact that a cosmological constant, even in a simple and idealised model such as the Schwarzschild-de Sitter space-time, induces an extremely high density of energy in the vicinity of the cosmological horizon, determined by a surface of constant radius $r \approx R$, where $R = \sqrt{3/\Lambda}$, and Λ is a positive cosmological constant. In the pure de Sitter space-time, the cosmological horizon is determined exactly by $r = R$. To our knowledge, the result of this section can only be obtained in the realm of the TEGR.

The line element of the Schwarzschild-de Sitter space-time is given by

$$ds^2 = -\alpha^2 dt^2 + \frac{1}{\alpha^2} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2, \quad (101)$$

where

$$\alpha^2 = 1 - \frac{2m}{r} - \frac{r^2}{R^2}. \quad (102)$$

Here we are considering natural units and making the speed of light $c = 1$. The Schwarzschild-de Sitter space-time has been considered in the TEGR in Ref. [55]. In this Section we make use of the notation and some results of Ref. [44]. We will restrict the considerations to the region between the Schwarzschild and cosmological horizons. The set of tetrad fields adapted to stationary observers in the space-time is given by

$$e_{a\mu} = \begin{pmatrix} -\alpha & 0 & 0 & 0 \\ 0 & \alpha^{-1} \sin \theta \cos \phi & r \cos \theta \cos \phi & -r \sin \theta \sin \phi \\ 0 & \alpha^{-1} \sin \theta \sin \phi & r \cos \theta \sin \phi & r \sin \theta \cos \phi \\ 0 & \alpha^{-1} \cos \theta & -r \sin \theta & 0 \end{pmatrix}. \quad (103)$$

It is not difficult to calculate the radial (inertial, non-gravitational) accelerations that are necessary to maintain the frame static in space-time. They are given by

$$\begin{aligned} \phi_{(0)(1)} &= \frac{d\alpha}{dr} \sin \theta \cos \phi, \\ \phi_{(0)(2)} &= \frac{d\alpha}{dr} \sin \theta \sin \phi, \\ \phi_{(0)(3)} &= \frac{d\alpha}{dr} \cos \theta. \end{aligned} \quad (104)$$

We define the inertial acceleration vector Φ as

$$\Phi(r) = (\phi_{(0)(1)}, \phi_{(0)(2)}, \phi_{(0)(3)}) \equiv \phi(r)\hat{r} = \frac{d\alpha}{dr}\hat{r}, \quad (105)$$

where $\hat{r} = (\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta)$, and

$$\frac{d\alpha}{dr} = \frac{1}{\alpha} \left(\frac{m}{r^2} - \frac{r}{R^2} \right). \quad (106)$$

Close to the Schwarzschild horizon we have $d\alpha/dr > 0$, and close to the cosmological horizon, $d\alpha/dr < 0$, as it should be to maintain the frame

static in space-time [44] (close to the Schwarzschild horizon the gravitational acceleration is attractive, and close to the de Sitter horizon it is repulsive).

In view of the spherical symmetry of the physical configuration, the gravitational energy obtained from Eq. (90) is reduced to

$$P^{(0)} = - \oint_S dS_1 \Pi^{(0)1}, \quad (107)$$

where $dS_1 = d\theta d\phi$. After a number of simple calculations, we find that

$$-\Pi^{(0)1} = 4ke \Sigma^{(0)01} = \frac{1}{4\pi} r \sin \theta \left(1 - \sqrt{1 - \frac{2m}{r_0} - \frac{r_0^2}{R^2}} \right), \quad (108)$$

and thus the energy contained within a surface of constant radius r_0 is given by

$$P^{(0)} = r_0 \left(1 - \sqrt{1 - \frac{2m}{r_0} - \frac{r_0^2}{R^2}} \right). \quad (109)$$

In the following, we will simplify the considerations and make $m = 0$. By also making $r_0 = R$ in Eq. (109), we find [55]

$$P^{(0)} = R = \sqrt{\frac{3}{\Lambda}}, \quad (110)$$

which is the total gravitational energy contained in the interior of the cosmological horizon. Returning to Eq. (107), we note that it can be written as a volume integral,

$$P^{(0)} = - \int_V dr d\theta d\phi \partial_r \Pi^{(0)1}, \quad (111)$$

where V is any spherical volume of radius r_0 such that $r_0 \leq R$. Now we take the radial derivative of $\Pi^{(0)1}$ in the equation above, and integrate in the angular variables. The resulting expression is

$$P^{(0)} = \int_V dr \varepsilon(r), \quad (112)$$

where $\varepsilon(r)$ expresses the gravitational energy density in the pure de Sitter space, i.e., $\varepsilon(r) dr$ is the gravitational energy contained within the shells of radii r and $r + dr$. It reads [55]

$$\varepsilon(r) = 1 + \frac{2\beta^2 - 1}{\sqrt{1 - \beta^2}}, \quad (113)$$

where $\beta^2 = r^2/R^2$. Note that if $\beta \rightarrow 0$, then $\varepsilon(0) \rightarrow 0$ and if $\beta \rightarrow 1$, ε diverges. These features are shown in Figure 2.




Figure 2: The gravitational energy density in the de Sitter space-time as a function of the radial coordinate only, after integration on the angular variables.

The total energy contained inside the surfaces of radii $0.1R$, $0.5R$, $0.9R$ are given by $E_g = 5.01 \times 10^{-4}R$, $E_g = 0.067R$, $E_g = 0.51R$, respectively. Thus, almost half of the gravitational energy is located between $\beta = 0.9$ and $\beta = 1$.

We see that the cosmological constant induces a concentration of gravitational energy in the vicinity of the cosmological horizon. Therefore, the existence of an extremely high density of energy located very far from idealised observers, and which could play the role of the so called dark energy in cosmological models, is explained by the definition of gravitational energy in the TEGR, even in the absence of matter fields.

It is also very interesting to observe that the cosmological constant gives rise to a classical vacuum energy that is not uniform over the 3-dimensional space, contrary to what is assumed in ordinary quantum field theories in flat space-times. The presence of a very tiny cosmological constant may not be perceived on neighbourhoods such as our planetary system, but its existence modifies the vacuum energy density on a cosmological scale. By inserting the physical constants c and G (the gravitational constant) into Eq. (110), we have

$$P^{(0)} = \frac{E}{c} = \frac{c^3}{G} \sqrt{\frac{3}{\Lambda}}, \quad (114)$$

where $\Lambda \approx 10^{-56} \text{ cm}^{-2}$ and $G = 6,67 \times 10^{-8} \frac{\text{cm}^3}{\text{g} \cdot \text{s}^2}$. It follows that

$$E = E_{vacuum} \approx 21 \times 10^{76} \frac{\text{g} \cdot \text{cm}^2}{\text{s}^2}. \quad (115)$$

This is a significant value that should modify some assumptions in ordinary quantum field theories, regarding the uniformity in space of the vacuum energy density, in case the space-time is endowed with a positive cosmological constant.

6 Gravitational energy-momentum in the Bondi-Sachs space-time

The space-time of a radiating gravitational field configuration is described by the Bondi-Sachs line element. The latter is usually presented in the (u, r, θ, ϕ) coordinates, where r is the ordinary radial variable and u is the retarded time, $u = t - r$ (r, θ and ϕ are spherical coordinates). The main quantities in the Bondi-Sachs line element are the functions $M(u, \theta, \phi)$, which is the *mass aspect*, and $c(u, \theta, \phi)$, $d(u, \theta, \phi)$, which are related to the *news functions*.

The Bondi-Sachs line element is primarily a model for isolated astrophysical configurations that loose energy in the form of gravitational radiation. The space-time around these configurations is not strictly asymptotically flat because although the metric tensor components fall off in the expected way as $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}(1/r)$ in the asymptotic limit $r \rightarrow \infty$, the time derivative of $h_{\mu\nu}$ is of order $1/r$ at spacelike infinity in Cartesian coordinates. The first significant investigation of this physical system was made by Bondi and

collaborators [61], who established the coordinates and notation that are currently employed in the analysis of gravitational radiation. The line element obtained by Bondi and collaborators is constructed out of the functions $M(u, \theta)$ and $c(u, \theta)$ only. These functions do not depend on the azimuthal coordinate ϕ because the line element is axially symmetric. The time derivative $\partial_0 c = \partial c / \partial u$ is identified as the first news function. Bondi's line element was later generalised by Sachs [62], who abandoned the axial symmetry and obtained the most general metric tensor that describes gravitational radiation at spacelike and null infinities. In the work by Sachs, there also appears the function $d(u, \theta, \phi)$, that yields the second news function $\partial_0 d$, and all functions (M, c, d) now depend also on the angular variable ϕ .

The two news functions are interpreted as the radiating degrees of freedom of the gravitational field, and this single feature justifies the importance of the metric tensor that describes the Bondi-Sachs space-time. The mathematical expression of the Bondi-Sachs metric tensor is not given in closed mathematical form. It is presented in powers of the radial coordinate r . There are very good review articles that clarify the several aspects of the subject, and also explain the emergence of the Bondi-Sachs energy-momentum vector [63, 64, 65, 66].

The Bondi-Sachs energy-momentum $m_\mu(u)$ is constructed out of the mass aspect $M(u, \theta, \phi)$ only, i.e., it does not depend on the functions $c(u, \theta, \phi)$ and $d(u, \theta, \phi)$ (see, for instance, Eq. (4.4) of Ref. [66]). However, the news functions are expected to yield real, observable manifestations of the gravitational field at spacelike or null infinities, and so they should contribute to surface integrals, both at spacelike and null infinities. For this reason, it is intriguing that the total Bondi-Sachs energy-momentum does not depend on the news function.

The ADM expression [34] of the total gravitational energy-momentum is evaluated at spacelike infinity. However, the authors of Refs. [67, 68, 69] have expressed the ADM energy-momentum in terms of the Bondi-Sachs energy-momentum $m_\mu(u)$, which is linear in the integration of the mass aspect $M(u, \theta, \phi)$ at null infinity. More specifically, the authors of the latter references rewrote the metric tensor of the Bondi-Sachs line element in the ordinary (t, r, θ, ϕ) coordinates, and parametrized the spacelike hypersurfaces by means of the standard time coordinate t . As these authors explain, the prior assumption is that the ADM energy-momentum is the past limit of the Bondi-Sachs energy-momentum, i.e., $m_\mu(-\infty)$. They found, however, that the resulting expression for the total ADM energy-momentum depends on

the integration of the mass aspect $M(u)$ when $u \rightarrow -\infty$, as expected (as well as on the spatial orientations for the 3-momenta), but also depends on integrations of functions that depend on c and d . Although the total ADM energy-momentum is strictly constructed for asymptotically flat space-times, the new and modified expression for the Bondi-Sachs energy-momentum, and its dependence on the functions c and d , is a very interesting and reasonable result.

In this Section we will show that gravitational energy-momentum (78) yields a quite interesting result. The latter expression renders the usual, expected integration of the mass aspect, plus a term that is interpreted as the energy of the gravitational radiation, since it depends only on the functions c and d . This is a result that can only be achieved in the TEGR. We will present the essential functions of the Bondi-Sachs space-time and the set of tetrad fields, but we will omit several long calculations that have already been presented in the literature [70]. We will emphasize the results only.

The Bondi-Sachs line element is given by

$$\begin{aligned} ds^2 = & g_{00} du^2 + g_{22} d\theta^2 + g_{33} d\phi^2 \\ & + 2g_{01} du dr + 2g_{02} du d\theta + 2g_{03} du d\phi + 2g_{23} d\theta d\phi, \end{aligned} \quad (116)$$

where

$$\begin{aligned} g_{00} &= \frac{V}{r} e^{2\beta} - r^2(e^{2\gamma} U^2 \cosh 2\delta + e^{-2\gamma} W^2 \cosh 2\delta + 2UW \sinh 2\delta), \\ g_{01} &= -e^{2\beta}, \\ g_{02} &= -r^2(e^{2\gamma} U \cosh 2\delta + W \sinh 2\delta), \\ g_{03} &= -r^2 \sin \theta (e^{-2\gamma} W \cosh 2\delta + U \sinh 2\delta), \\ g_{22} &= r^2 e^{2\gamma} \cosh 2\delta, \\ g_{33} &= r^2 e^{-2\gamma} \cosh 2\delta \sin^2 \theta, \\ g_{23} &= r^2 \sinh 2\delta \sin \theta. \end{aligned} \quad (117)$$

We adopt the usual convention $(u, r, \theta, \phi) = (x^0, x^1, x^2, x^3)$. The functions β , γ , δ , U and W in the equations above are not exact. They are given only in asymptotic form, in powers of $1/r$. We will dispense with the powers of $1/r$ of the field quantities that do not contribute to the calculations. Thus, the asymptotic form of the functions above are

$$\begin{aligned}
V &\simeq -r + 2M, \\
\beta &\simeq -\frac{c^2 + d^2}{4r^2}, \\
\gamma &\simeq \frac{c}{r}, \\
\delta &\simeq \frac{d}{r}, \\
U &\simeq -\frac{l(u, \theta, \phi)}{r^2}, \\
W &\simeq -\frac{\bar{l}(u, \theta, \phi)}{r^2},
\end{aligned} \tag{118}$$

where

$$\begin{aligned}
l &= \partial_2 c + 2c \cot \theta + \partial_3 d \csc \theta, \\
\bar{l} &= \partial_2 d + 2d \cot \theta - \partial_3 c \csc \theta.
\end{aligned}$$

In the limit $r \rightarrow \infty$, the asymptotic form of the functions above yield

$$\begin{aligned}
g_{00} &\simeq -1 + \frac{2M}{r}, \\
g_{01} &\simeq -1 + \frac{c^2 + d^2}{2r^2}, \\
g_{02} &\simeq l + \frac{1}{r}(2cl + 2d\bar{l} - p), \\
g_{03} &\simeq \bar{l} \sin \theta + \frac{1}{r}(-2c\bar{l} + 2dl - \bar{p}) \sin \theta, \\
g_{22} &\simeq r^2 + 2cr + 2(c^2 + d^2), \\
g_{33} &\simeq [r^2 - 2cr + 2(c^2 + d^2)] \sin^2 \theta, \\
g_{23} &\simeq 2dr \sin \theta + \frac{4d^3}{3r} \sin \theta.
\end{aligned} \tag{119}$$

The functions p and \bar{p} are defined in Refs. [67, 68]. They depend on functions that are not defined above. However, they will not contribute to the final expressions, and for this reason we will not present their definitions here.

The expressions of the contravariant components of the metric tensor are calculated by means of the usual procedure out of Eqs. (117) (and not out of

Eqs. (119)). The inverse components are given by $g^{\mu\nu} = (-1)^{\mu+\nu}(1/g) M_{\mu\nu}$, where $g = -g_{01}^2(g_{22}g_{33} - g_{23}^2)$ is the determinant of the metric tensor, and $M_{\mu\nu}$ is the co-factor of the $\mu\nu$ component (we have taken into account all necessary powers of $1/r$ of the functions given in Eq. (118)). We find

$$\begin{aligned}
 g^{00} &= g^{02} = g^{03} = 0, \\
 g^{01} &\simeq -1 - \frac{c^2 + d^2}{2r^2}, \\
 g^{11} &\simeq 1 - \frac{2M}{r}, \\
 g^{12} &\simeq \frac{l}{r^2}, \\
 g^{13} &\simeq \frac{\bar{l} \sin \theta}{r^2}, \\
 g^{22} &\simeq \frac{1}{r^2}, \\
 g^{33} &\simeq \frac{1}{r^2 \sin^2 \theta}, \\
 g^{23} &\simeq -\frac{2d}{r^3 \sin^2 \theta}. \tag{120}
 \end{aligned}$$

In order to investigate the gravitational radiation in the context of the Bondi-Sachs space-time, it is convenient to choose a set of tetrad fields adapted to static observers in space-time. The worldline of these observers is characterized by the field of 4-velocities u^μ of the observers such that $u^\mu = (u^0, 0, 0, 0)$. Following the procedure discussed in Section 2, we identify the $a = (0)$ component of the frame $e_a{}^\mu$ as $e_{(0)}{}^\mu = u^\mu$. The three other components of the frame, $e_{(i)}{}^\mu$, are orthogonal to $e_{(0)}{}^\mu$, and may be oriented in the three-dimensional space according to the symmetry of the physical system, in case there are privileged directions in the 3-space. It follows from the requirement $e_{(0)}{}^\mu = u^\mu$ that a static observer in space-time must satisfy the conditions $e_{(0)}{}^i(t, x^k) = (0, 0, 0)$. It is easy to verify, by means of a coordinate transformation, that in terms of the retarded time u we also have $e_{(0)}{}^i(u, x^k) = (0, 0, 0)$.

The Bondi-Sachs space-time is not axially symmetric. It is clear from the mathematical structure of the line element that there are no distinguished directions at spacelike infinity. Since we will evaluate surface integrals at spacelike infinity, i.e., we will be interested only in total quantities (we will

integrate over a surface S determined by $r = \text{constant}$, for r finite but sufficiently distant from the source), any set of tetrad fields that satisfy the asymptotic expansion $e_{a\mu} \simeq \eta_{a\mu} + (1/2)h_{a\mu}(1/r)$ in Cartesian coordinates when $r \rightarrow \infty$, and that satisfy the conditions $e_{(0)}^i(u, r, \theta, \phi) = 0$, will serve our purposes. For such a frame, $e_{(1)\mu}^i$, $e_{(2)\mu}^i$ and $e_{(3)\mu}^i$ will define the usual unit frame vectors in the x , y and z directions, respectively, in the limit $r \rightarrow \infty$, provided e_a^μ is constructed in Cartesian coordinates. These conditions establish a consistent reference frame at spacelike infinity.

In spite of the intricacies of the Bondi-Sachs line element, it is possible to establish a simple set of tetrad fields that yields Eqs. (116) and (117), and that satisfy the conditions $e_{(0)}^i = 0$. Note that $e_{(0)}^i = 0$ implies $e^{(i)}_0 = 0$. One set of tetrad fields that satisfy these requirements, and has the asymptotic form $e_{a\mu} \simeq \eta_{a\mu} + (1/2)h_{a\mu}$ at spacelike infinity, is given in (u, r, θ, ϕ) coordinates by [70]

$$\begin{aligned}
e_{(0)\mu} &= (-A, -E, -F, -G), \\
e_{(1)\mu} &= (0, B_1 \sin \theta \cos \phi + B_2 \cos \theta \cos \phi - B_3 \sin \theta \sin \phi, \\
&\quad C_1 r \cos \theta \cos \phi - C_2 \sin \theta \sin \phi, \\
&\quad -D r \sin \theta \sin \phi), \\
e_{(2)\mu} &= (0, B_1 \sin \theta \sin \phi + B_2 \cos \theta \sin \phi + B_3 \sin \theta \cos \phi, \\
&\quad C_1 r \cos \theta \sin \phi - C_2 \sin \theta \cos \phi, \\
&\quad D r \sin \theta \cos \phi), \\
e_{(3)\mu} &= (0, B_1 \cos \theta - B_2 \sin \theta, -C_1 r \cos \theta, 0). \tag{121}
\end{aligned}$$

The quantities $A, B_1, B_2, B_3, C_1, C_2, D, E, F, G$ are determined by requiring that $e_{a\mu}$ yields the metric tensor components (119) according to $e_{a\mu} e_{b\nu} \eta^{ab} = g_{\mu\nu}$. The determination of exact form of these quantities in terms of the metric tensor components (117) is not so simple. These quantities must satisfy the following equations,

$$\begin{aligned}
-A^2 &= g_{00} \\
-AE &= g_{01} \\
-AF &= g_{02} \\
-AG &= g_{03}
\end{aligned}$$

$$\begin{aligned}
-E^2 + B_1^2 + B_2^2 + B_3^2 \sin^2 \theta &= g_{11} = 0 \\
-EF + B_2(C_1 r) + B_3 C_2 \sin^2 \theta &= g_{12} = 0 \\
-EG + B_3(Dr) \sin^2 \theta &= g_{13} = 0 \\
-F^2 + (C_1 r)^2 + C_2^2 \sin^2 \theta &= g_{22} \\
-G^2 + (Dr)^2 \sin^2 \theta &= g_{33} \\
-FG + C_2(Dr) \sin^2 \theta &= g_{23}. \tag{122}
\end{aligned}$$

In view of the fact that we need the components of the torsion tensor only in the asymptotic limit $r \rightarrow \infty$, we restrict the considerations to quantities whose power of $1/r$ are actually needed in the calculations. Thus, we consider terms of power up to $(1/r)^2$ only. Quantities of order $(1/r)^n$, with $n \geq 3$, do not contribute to the final, total expressions, and will be neglected. We find

$$\begin{aligned}
A &\simeq 1 - \frac{M}{r}, \\
E &\simeq 1 + \frac{M}{r}, \\
F &\simeq -l - \frac{1}{r}(2cl + 2d\bar{l} + Ml - p), \\
G &\simeq -\sin \theta \left[\bar{l} + \frac{1}{r}(-2c\bar{l} + 2dl + M\bar{l} - \bar{p}) \right], \\
B_1 &\simeq 1 + \frac{M}{r}, \\
B_2 &\simeq -\frac{l}{r} - \frac{1}{r^2}(2Ml + cl - p), \\
B_3 &\simeq -\frac{1}{\sin \theta} \left[\frac{\bar{l}}{r} + \frac{1}{r^2}(2M\bar{l} - c\bar{l} + 2dl - \bar{p}) \right], \\
C_1 &\simeq 1 + \frac{c}{r} + \frac{1}{r^2} \left[\frac{l^2}{2} + c^2 - d^2 \right], \\
C_2 &\simeq \frac{1}{\sin \theta} \left[2d + \frac{1}{r}(l\bar{l} + 2cd) \right]. \\
D &\simeq 1 - \frac{c}{r} + \frac{1}{r^2} \left(\frac{\bar{l}^2}{2} + c^2 + d^2 \right). \tag{123}
\end{aligned}$$

The expressions above completely fix the set of tetrad fields given by Eq. (121). We note that the $\sin \theta$ in the denominator of B_3 and C_2 in the expressions above does not lead to a divergence of $e_{a\mu}$ on the z axis of the

coordinate system, when $\theta = 0$. In Eq. (121) these quantities are multiplied by $\sin \theta$. It can be shown that both in (u, x, y, z) or in (t, x, y, z) coordinates, the set of tetrad fields given by (121) is everywhere smooth in the three-dimensional space (except possibly at $r = 0$, but the functions (118) are not exactly defined in the neighbourhood of $r = 0$).

As we mentioned earlier, we will skip many long calculations, and present the main results only. Some important steps in the calculations are presented in Ref. [70]. After many calculations and simplifications, we obtain the total gravitational energy by means of Eq. (78). It reads

$$P^{(0)} = 4k \int_0^{2\pi} d\phi \int_0^\pi d\theta \sin \theta \left[M + \partial_0 F \right], \quad (124)$$

where

$$F = -\frac{1}{4} \left(l^2 + \bar{l}^2 \right) + \frac{1}{2} c^2 + d^2. \quad (125)$$

The quantities under integration in the expression above are the only ones that contribute to the surface integral in the limit $r \rightarrow \infty$.

The new term $\partial_0 F$ generalises the standard Bondi-Sachs energy. As expected, the latter depends not only on the integral of the mass aspect $M(u, \theta, \phi)$, but also on the functions $c(u, \theta, \phi)$ and $d(u, \theta, \phi)$.

The surface S of integration in Eq. (78) is usually a spacelike surface of constant radius r . In the context of Eq. (124), for finite values of the ordinary time t , the limit $r \rightarrow \infty$ corresponds to $u \rightarrow -\infty$, and therefore in Eq. (124) we have $P^{(0)}(u \rightarrow -\infty)$. However, this is not a mandatory condition. We may consider very large but finite values of the radial coordinate r ($r \gg M$ and $r \gg \partial_0 F$), as explained in Ref. [70], such that M and $\partial_0 F$ remain the only contributions to the surface integral in Eq. (124). In this case, $P^{(0)}$ depends on arbitrary and finite values of u . The fact is that real observers are never strictly located at spacelike infinity.

Equation (124) is interpreted as the gravitational energy that a static observer measures at very large distances from the source determined by $M(u, \theta, \phi)$. Assuming the speed of light $c = 1$ as well as $G = 1$, the quantity

$$E_{rad} = \frac{1}{4\pi} \int_0^{2\pi} d\phi \int_0^\pi d\theta \sin \theta (\partial_0 F), \quad (126)$$

is interpreted as the energy of the gravitational radiation, i.e., a form of energy detached from the source.

In the study of electrodynamics of moving but isolated sources, one finds that the total electric field is composed of a term that is usually called the *near field*, which is determined by the charges of the source, and by a term called *far field*, which is determined not only by the charge, but by the velocity and acceleration of the charge. This latter component of the electric field falls off as $1/r$ at spacelike infinity, and represents the first order contribution of the electric field at large distances (it contributes to the Poynting vector, together with the corresponding magnetic field component). By making an analogy with expressions (125) and (126), we may say that E_{rad} given by (126) is a form of *far energy*, that eventually may be detected and measured with the increasing sophistication of the precision instruments presently used in the large detectors of gravitational waves.

The gravitational momenta is calculated in a way very similar to the gravitational energy. Again, the algebraic calculations are somewhat long, but pose no special mathematical difficulty. The gravitational momenta $P^{(i)}$ ($i = 1, 2, 3$) may be presented in a simplified form by first defining the standard textbook vectors,

$$\begin{aligned}\hat{r}^i &= (\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta), \\ \hat{\theta}^i &= (\cos \theta \cos \phi, \cos \theta \sin \phi, -\sin \theta), \\ \hat{\phi}^i &= (-\sin \phi, \cos \phi, 0).\end{aligned}\tag{127}$$

The final expressions are given by

$$\begin{aligned}P^{(i)} &= 4k \int_0^{2\pi} d\phi \int_0^\pi d\theta \sin \theta \left[(M + \partial_0 F) \hat{r}^i \right. \\ &\quad \left. + \frac{1}{4} (l \partial_0 M) \hat{\theta}^i + \frac{1}{4} (\bar{l} \partial_0 M) \hat{\phi}^i \right].\end{aligned}\tag{128}$$

Altogether, $P^{(0)}$ and $P^{(i)}$ constitute the Bondi-Sachs energy-momentum in the realm of the TEGR. One important observation is the following. In view of the field equations for the metric tensor (116), it is known that the time derivative $\partial_0 M$ may be written in terms of the time derivatives of the functions c and d [62, 65] according to

$$\partial_0 M = -[(\partial_0 c)^2 + (\partial_0 d)^2] + \frac{1}{2} \partial_0 \left(\partial_2 l + l \cot \theta + \frac{\partial_3 \bar{l}}{\sin \theta} \right).\tag{129}$$

If we evaluate all integrals in P^a in the limit $r \rightarrow \infty$, we are actually taking the limit $u \rightarrow -\infty$, as we discussed above. If, in addition, we assume that the news functions satisfy the initial conditions

$$\partial_0 c(-\infty) = 0, \quad \partial_0 d(-\infty) = 0, \quad (130)$$

then the total energy-momentum $P^a = (P^{(0)}, P^{(i)})$ given by (124) and (128) reduces to the well known expression for the Bondi-Sachs energy-momentum given in Ref. [66], for instance. These initial conditions are physically reasonable. They are related to the fact that the physical configuration is not initially radiating (i.e., the radiation starts at a finite instant of the retarded time u), but they are not mandatory. We asserted above that we can evaluate $(P^{(0)}, P^{(i)})$ over a surface S of integration sufficiently far from the source (so that only terms of order $1/r$ and $1/r^2$ contribute to the calculations), such that the surface S is large but finite. This fact allows us to dispense with the initial conditions given by Eq. (130), since the retarded time u is not strictly $-\infty$, and treat Eq. (126) as, indeed, the energy of the gravitational radiation.

As a final remark, we mention that in Ref. [70] we have simplified the analysis to an axially symmetric space-time, constructed an analytic expression for the news function $c(u, \theta)$ and obtained an expression for the energy of gravitational radiation E_{rad} as a somewhat complicated infinite sum. We believe that this issue could be re-analysed, since the resulting expression for E_{rad} is a sum of finite values, and maybe it could be simplified to an infinite sum where each term can be easily interpreted.

7 Plane gravitational waves, the kinetic energy of free particles and localization of the gravitational energy

As we asserted at the Introduction of this review, in a series of publications [30, 31, 32, 33] we have shown that the action of plane-fronted gravitational waves (pp-waves) on free particles may increase or decrease the kinetic energy of the particle. The literature on pp-waves is really vast, as we have indicated in the above references. It is important to recall that pp-waves are very simple exact solutions of the vacuum Einstein's field equations, and also of the teleparallel field equations (70) and (72). A very good analysis on this

issue is presented in Refs. [71, 72]. As for any simple solutions of field equations or of equations of motion in physics, these waves are idealizations of more general realistic configurations, but no doubt, the latter is expected to display the basic features of the idealized physical configurations. The belief is that pp-waves may exist in nature not exactly described by the simple mathematical functions that we will present ahead, but by a more elaborate and realistic mathematical description.

We recall that free particles or free material systems are here understood as physical systems that are subject to the gravitational field only, and not to any external inertial forces. The consequence of the action of pp-waves on free particles is two fold, at least. First, since free particles gain or loose energy with the passage of the wave, so does the wave loose or gain energy from the particle. This feature allows the wave to travel in space for periods of time such as billions of years, without dissipating. The waves that are currently being observed in the large terrestrial detectors are, supposedly, waves that have travelled for such long periods of time.

The second consequence is that since the free particles are idealized pointwise objects, the energy transfer is localized in space. This fact demonstrates that the gravitational energy is localized. As we asserted at the Introduction, it does not make sense to relate this energy transfer to any spacelike 2-surface of arbitrary radius that envelops the particle. The energy transfer is effectively pointwise, and this is the simplest description of the physical process.

In the following, we will recall the exposition presented in Ref. [32], in the study of the memory effect due to the pp-waves. The space-time before and after the passage of the wave is assumed to be flat or nearly flat, for suitable amplitudes of the waves. The metric tensor that describes the plane wave space-time has a well known and simple structure. We assume that wave propagates along the z direction. Then, the line element reads

$$ds^2 = dx^2 + dy^2 + 2du\,dv + H(x, y, u)du^2, \quad (131)$$

in (u, v, x, y) coordinates. In the flat space-time region, we may identify

$$u = \frac{1}{\sqrt{2}}(z - t), \quad (132)$$

$$v = \frac{1}{\sqrt{2}}(z + t). \quad (133)$$

The coordinates (x, y) parametrize the planes parallel to the wave front. The line element depends essentially on a function $H(x, y, u)$. Here, the retarded time is not given by u , but by $-u$.

The class of exact solutions considered here belong to the well known Kundt family of space-times. They were studied in great detail by Ehlers and Kundt [71, 72]. The dependence of the function H on the variable u is arbitrary, which is a typical feature of solutions of wave equations. In view of this arbitrariness, the function H may be chosen so that it describes a short burst of gravitational wave, in which case the u dependence of H may be given by smooth Gaussians or derivatives of Gaussians. The function H must only satisfy [71]

$$\nabla^2 H = \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) H = 0. \quad (134)$$

We assume that far from an astrophysical source, a realistic gravitational wave may be approximated by an exact plane wave, exactly as we do in the study of electromagnetic waves. The form of the line element (131), in the coordinates (u, v, x, y) , was first presented by Brinkmann [73].

In order to study the geodesic equations, we need first rewrite the line element above in (t, x, y, z) coordinates. It reads

$$ds^2 = \left(\frac{H}{2} - 1 \right) dt^2 + dx^2 + dy^2 + \left(\frac{H}{2} + 1 \right) dz^2 - H dt dz. \quad (135)$$

We assume $c = 1$. The function H must satisfy only Eq. (134). As already asserted, the dependence of H on the retarded time $(-u)$ is arbitrary. The geodesic equations in terms of the t, x, y, z coordinates are

$$2\ddot{t} + \sqrt{2}H\ddot{u} + \sqrt{2}\dot{H}\dot{u} - \frac{1}{\sqrt{2}}\frac{\partial H}{\partial u}\dot{u}^2 = 0, \quad (136)$$

$$2\ddot{x} - \frac{\partial H}{\partial x}\dot{u}^2 = 0, \quad (137)$$

$$2\ddot{y} - \frac{\partial H}{\partial y}\dot{u}^2 = 0, \quad (138)$$

$$2\ddot{z} + \sqrt{2}H\ddot{u} + \sqrt{2}\dot{H}\dot{u} - \frac{1}{\sqrt{2}}\frac{\partial H}{\partial u}\dot{u}^2 = 0, \quad (139)$$

where the dot represents derivative with respect to the parameter s that appears on the left hand side of Eq. (135). Among the various possibilities for the function $H(-u)$, we have [32]

$$H_1 = A_+(u)(x^2 - y^2), \quad (140)$$

$$H_2 = A_\times(u) xy, \quad (141)$$

or by a linear combination of these quantities. The two expressions above of H satisfy Eq. (134), and have been considered in the literature in several investigations. We anticipate that both expressions lead to the same qualitative behaviour for the geodesics and kinetic energy of the particles. We consider the amplitudes $A_+(u)$ and $A_\times(u)$ to be given by regular Gaussians (normalised, not necessarily to 1), that represent short bursts of gravitational waves. These amplitudes are typically (but not exactly) given by

$$A_{(+,\times)} \simeq \frac{1}{\lambda} e^{-(u^2/2\lambda^2)}. \quad (142)$$

In addition to the ansatz above, we have chosen multiplicative constants in the expression of the amplitude in order to yield satisfactory pictures (see below). In fact, the amplitude does not need to be normalized.

The effect of a gravitational wave on a free particle is assumed to be very weak. For this reason, the kinetic energy per unit mass K of the free particles is written as

$$2K = \frac{1}{2}(V_x^2 + V_y^2 + V_z^2), \quad (143)$$

where the velocities are $V_x = dx/du$, $V_y = dy/du$ and $V_z = dz/du$. Thus, in view of Eq. (132), we have

$$K = \frac{1}{2} \left[\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 + \left(\frac{dz}{dt} \right)^2 \right], \quad (144)$$

which is a valid expression for instants of time before and after the passage of the wave, in which case the space-time is flat, or nearly flat.

We define K_f and K_i as the final and initial kinetic energies of the free particles, i.e., the energy of the particles after and before the passage of the wave. We also define $\Delta K = K_f - K_i$, and the normalized quantity ΔK_N ,




Figure 3: ΔK_N as a function of λ , considering H_1

$$\Delta K_N = \frac{K_f - K_i}{K_f + K_i}. \quad (145)$$

The geodesic equations have been solved numerically, as explained in Ref. [32], and the quantities ΔK and ΔK_N have also been evaluated numerically, which resulted in the figures below. The initial conditions for the constructions of these figures were arbitrarily chosen, and are clearly presented in [32].

In Figure 3, we see an interesting variation of the normalised kinetic energy with respect to the parameter λ that appears in Eq. (142). But most important is the behaviour of $K(u)$ with respect to the variable u , which is given in Figures 4 and 5.

In Figure 4, the particle loses energy to the gravitational field, and the opposite takes place in Figure 5. We have analysed innumerable situations where this energy transfer occurs, and that yield figures much different from Figures 4 and 5. The gain or loss of energy depends essentially in the initial conditions for the geodesic equations. However, an explanation to the variation of the kinetic energy of the free particle may be given by the work-energy relation for the free particle in the gravitational field.

In Ref. [33] we have investigated, in great detail, the validity of the classical Torricelli equation, $v_f^2 = v_i^2 + 2a\Delta x$, for a particle that is subject to




Figure 4: $K(u)$, considering H_1 and $\lambda = 7.30247$. Figure 5: $K(u)$, considering H_1 and $\lambda = 13.6043$.

the gravitational field of a pp-wave. For a given arbitrary trajectory between the initial and final positions denoted by i and f , respectively, the Torricelli equation is written as

$$\frac{1}{2}v_f^2 - \frac{1}{2}v_i^2 = \int_i^f \mathbf{a} \cdot d\mathbf{l}, \quad (146)$$

where \mathbf{a} is the acceleration of the particle due to the gravitational field. The space-time at the initial and final positions of the particle is assumed to be flat. The relation above is just a manifestation of the standard work-energy relation $\Delta K = \Delta W$ of classical Newtonian physics, for a particle that is under the action of a force $\mathbf{F} = m\mathbf{a}$. However, in Eq. (146), both quantities ΔK (the left hand side) and ΔW (the right hand side of (146)) are quantities per unit of mass. As we discussed in Ref. [33], the relativistic form of the right hand side of the equation above is calculated according to the expression

$$\Delta W = - \int_i^f \phi_{(0)(i)} e^{(i)}_j dx^j = - \int_i^f a_j dx^j. \quad (147)$$

The procedure for calculating the right hand side of the equation above is the following. First, we establish a set of tetrad fields adapted to static observers in the space-time of a pp-wave given by the metric tensor (135). Then, we calculate the inertial accelerations that are necessary to maintain the frame static in space-time. We identify the gravitational acceleration on the particle (or on any other physical body) as exactly *minus* the inertial accelerations, since the frame must be static. This identification explains the minus sign in Eq. (147), and in this equation $e^{(i)}_j$ are components of the

static co-frame. The acceleration tensor components $\phi_{(0)(i)}$ are calculated out of Eq. (5) or (10).

As for ΔK , we used

$$\begin{aligned}\Delta K &= \frac{1}{4} \left[\dot{x}^2(u) + \dot{y}^2(u) + \dot{z}^2(u) \right]_{u=-\infty}^{u=\infty} \\ &= \frac{1}{2} \left[\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 + \left(\frac{dz}{dt} \right)^2 \right]_{t=-\infty}^{t=\infty}.\end{aligned}\quad (148)$$

The trajectories and initial conditions for the solution of the geodesic equations are presented in detail in Ref. [33]. We reaffirm that in the initial ($t \rightarrow -\infty$) and final ($t \rightarrow +\infty$) positions of the particle, the space-time is flat, in view of the Gaussian form of the wave, and this fact means that the initial and final geodesic trajectories of the particle are straight lines in space-time. By means of numerical and graphical analysis, we have found a striking and astonishing coincidence between ΔK and ΔW , for various trajectories and initial conditions.

A more realistic treatment of this gravitational field configuration would take into account the contribution of the mass of the particle to the total gravitational field. However, such a solution of the field equations does not exist, even in approximate form. For a particle with very small mass, that generates a gravitational field negligible compared to the field of the wave, the geodesic equations considered in Refs. [30, 31, 32, 33] are consistent and realistic trajectories of the particles and lead to a measurable velocity memory effect. An interesting consequence of the configuration pp-wave + free particles is that it is possible, at least in principle, to extract energy from gravitational waves, keeping in mind that the waves may as well carry away energy from material configurations.

The general conclusion is that we may have $\Delta K > 0$ or $\Delta K < 0$ depending **(i)** on the initial and final conditions of the geodesic trajectories, and **(ii)** on the gravitational acceleration on the particle, due to the pp-wave. In any case, there is an actual transfer of energy between the gravitational field of the pp-wave and the particle. This is a possible explanation to the memory effect, in the context of non-linear plane gravitational waves.

One important conclusion of the present review is that the energy transfer between the particle and the wave (i.e., the gravitational field) is pointwise, which demonstrates the localizability of the gravitational energy.

8 Final Remarks

In this review we have revisited the definitions of energy-momentum P^a (Eq. (78)) and 4-angular momentum L^{ab} (Eq. (82)) of the gravitational field in the TEGR, with emphasis on the localization of the gravitational energy and, by extension, on the localization of the gravitational momentum and 4-angular momentum. We argue that the velocity memory effect is a manifestation of the localization of the gravitational energy: the transfer of energy between an otherwise free particle and the gravitational field of a plane-fronted gravitational wave is pointwise, and so the gravitational energy must be localized at the position of the particle. Therefore, the densities that arise in the TEGR and that yield the definitions of P^a and L^{ab} do make physical sense. In addition, the results obtained in the analysis of distinct configurations like the Kerr black hole, the Schwarzschild-de Sitter space-time and the Bondi-Sachs space-time support the validity of P^a , given that all results are consistent from the physical point of view. These space-times have specific features, and the definition of P^a is in agreement with the expected phenomenology regarding each one of these features, which is an indication of the universality of P^a .

In Ref. [74] we investigated the transfer of angular momentum between a free particle and the gravitational field of a gyratonic pp-wave. Exactly as in the case of the energy transfer, the final angular momentum of the particle may be smaller or higher than the initial angular momentum. This variation of angular momentum of the particle depends on the initial conditions of the particle, as well as on the nature of the gyratonic wave.

We have also revisited the concept of tetrad fields, and by means of the analysis of well known and physically motivated constructions, we have shown, once again, that a set of tetrad fields in space-time yields the reference frame of distinguished observers. The important and inevitable conclusion is that tetrad fields describe at the same time the reference frame of a field of observers, and the gravitational field. These two manifestations are inextricably linked. As we explained in Section 2, the set of tetrad fields may be characterized **(i)** either by fixing the timelike component of the frame $e_{(0)}^\mu$ along the trajectory of an observer, and fixing the three other spacelike components according to the symmetry of the physical configuration, or **(ii)** by requiring the six components of the acceleration tensor to satisfy some conditions on the frame in space-time (as, for instance, to maintain the frame static and non-rotating in space-time).

Two sets of tetrad fields that are related by a local Lorentz transformation may be equivalent in the sense that they yield the same metric tensor, but in general they describe quite distinct congruences of observers (i.e., congruence of timelike curves defined by $e_{(0)}^\mu(x)$), and yield different results for the gravitational energy-momentum, which is covariant under global, but not under local Lorentz transformations. We have made clear in Sections 2 and 4 that the sets of tetrad fields given by Eqs. (55) and (97) for the Kerr line element are physically different. Only the latter one has physical relevance for the result in Section 4, namely, to the evaluation of the irreducible mass of the Kerr black hole.

The calculation of the energy contained within the external event horizon of the Kerr black hole leads to a result that is strikingly close to twice the irreducible mass of the black hole, which is the expected result. The final expression obtained in our analysis is a lot more satisfactory than the result achieved by any other approach in the existing literature on this subject. We must choose the set of tetrad fields adapted to the field of observers that are allowed within the ergosphere of the Kerr black hole. This set of tetrad fields is given by Eq. (97). The frame described by the latter equation yields a field of observers that are dragged in rotational motion by the gravitational field of the black hole inside the ergosphere, which is a well known feature of the Kerr space-time. On the other hand, tetrads defined by Eq. (55) are not defined in the interior of the ergosphere, and for this reason they play no role in the evaluation of the irreducible mass.

In order to present our definitions of P^a and L^{ab} , we have readdressed the Lagrangian and Hamiltonian approaches to the TEGR. The Hamiltonian formulation is always important because it leads to the constraint equations and guarantees the mathematical and physical consistency of the theory. The TEGR is a theory with only first class constraints, which means that the hyperbolic evolution equations are well defined. The definitions of energy, momentum and 4-angular momentum of the gravitational field in the TEGR are essentially a natural development of the Hamiltonian formulation. As we wrote before, they are not abstract mathematical elaborations motivated by circumventing a non-existent problem of localization of the gravitational energy.

The TEGR, as exposed in this review, is constructed out of the tetrad fields only. The addition of a (flat or non-flat) spin connection does not change the dynamics of the theory, and just brings unnecessary complications both for the mathematical evolution of the Hamiltonian field equations

and for the measurable quantities of the theory. There is, currently, a debate in the physics community regarding this issue, as we can see in the exposition of Refs. [75, 76, 77, 78], and references therein. We also call attention to the interesting considerations carried out in Ref. [20], regarding the establishment of tetrad fields in space-time. The notion of ideal or preferred frames has always been pursued in the context of tetrad theories of gravity, of which the TEGR is one prominent realisation.

The Lagrangian and Hamiltonian formulations of the TEGR considered in this review are invariant under the global $SO(3,1)$ group. However, the field equations of the theory, Eqs. (70) or (72), as discussed before, are covariant under the local $SO(3,1)$ group. The fact that the field equations are covariant under local Lorentz transformations means that the theory, defined by the field equations, can be formulated in the frame of an arbitrary observer in space-time. This is the essential and crucial meaning of local Lorentz invariance of a theory. Measurable quantities, of course, depend on the frame. Even in the standard formulation of classical electrodynamics, the electric field of a point charge has an expression in a static frame (where the charged particle is at rest), and a different expression in a frame that is moving with respect to the point charge. Assuming that the space-time is flat and that the velocity of the moving frame is constant, both observers at the static and moving frames move along geodesics in space-time. In spite of moving along geodesics, they measure different values for the electric field, and this is just a consequence of the special theory of relativity. It simply does not make sense to allow this feature to take place in classical electrodynamics, and not in context of a theory for the gravitational field. In the framework of the special theory of relativity, physical measurements are frame dependent, and so they should be in the realm of general relativity. After all, one of the attributes of tetrad fields is exactly to project vectors and tensor components (coordinate dependent quantities) into the local frame of an observer.

References

- [1] J. Norton, “What was Einstein’s principle of equivalence?”, *Stud. Hist. Phil. Sci.* **16**, 203 (1985).

- [2] J. W. Maluf, M. V. O. Veiga and J F. da Rocha-Neto, “Regularized expression for the gravitational energy-momentum in teleparallel gravity and the principle of equivalence”, *Gen. Relativ. Gravit.* **39**, 227-240 (2007).
- [3] S. B. Babak and L. P. Grishchuk, *Phys. Rev. D* **61**, 024038 (1999).
- [4] “Einstein’s Apple: homogeneous Einstein’s fields”, E. L. Schucking and E. J. Surowitz (World Scientific, 2015).
- [5] J. L. Synge, “Relativity: the general theory”, (North-Holland 1976).
- [6] K. Hayashi and T. Shirafuji, “New general relativity”, *Phys. Rev. D* **19**, 3524 (1979); **24**, 3312 (1981).
- [7] F. W. Hehl, “Four Lectures on Poincaré Gauge Field Theory”, in “Cosmology and Gravitation: Spin, Torsion, Rotation and Supergravity”, edited by P. G. Bergmann and V. de Sabbata (Plenum Press, New York, 1980).
- [8] Y. M. Cho, “Einstein Lagrangian as the translational Yang-Mills Lagrangian”, *Phys. Rev. D* **14**, 2521 (1976); “Gauge theory of the Poincaré symmetry”, *Phys. Rev.* **14**, 3335 (1976).
- [9] J. M. Nester, “Is there really a problem with the teleparallel theory?” *Class. Quantum Grav.* **5**, 1003 (1988).
- [10] J. W. Maluf, “The Hamiltonian description of the teleparallel equivalent of general relativity”, *J. Math. Phys.* **35**, 335 (1994).
- [11] J. W. Maluf, “Localization of energy in general relativity”, *J. Math. Phys.* **36**, 4242 (1995).
- [12] Yu. N. Obukhov and J. G. Pereira, “Metric-affine approach to teleparallel gravity”, *Phys. Rev. D* **67**, 044016 (2003).
- [13] J. W. Maluf, “The teleparallel equivalent of general relativity”, *Ann. Phys. (Berlin)* **525**, 339 (2013).
- [14] R. Aldrovandi and J. G. Pereira, “Teleparallel Gravity: an Introduction”, Springer (2013).

- [15] L. Combi and G. E. Romero, “Is teleparallel gravity really equivalent to general relativity?”, *Ann. Phys. (Berlin)* **530**(1), 1700175 (2018).
- [16] L. Combi and G. E. Romero, “Gravitational energy and radiation of a charged black hole”, *Class. Quantum Grav.* **34**, 195008 (2017).
- [17] J. B. Formiga, “The energy-momentum tensor of gravitational waves, Wyman spacetime and freely falling observers”, *Ann. Phys. (Berlin)* **530**, 0320 (2018).
- [18] J. B. Formiga, “Conformal teleparallel theories and Weyl geometry”, *Phys. Rev. D* **99**, 064047 (2019).
- [19] J. B. Formiga, “The Gravitational Energy–Momentum Density of Radially Accelerated Observers in Schwarzschild Spacetime”, *Ann. Phys. (Berlin)* **532** 1900507 (2020).
- [20] J. B. Formiga, “On the teleparallel frame problem”, *Mod. Phys. Lett. A* **37**, 33n34, 2250222 (2022); *Braz. J. Phys.* **51**, 1823-1832 (2021) [arXiv:2004.10788].
- [21] J. B. Formiga and V. R. Gonçalves, “The gravitational energy density of the Universe”, *Mod. Phys. Lett. A* **36**, 18, 2150125 (2021).
- [22] V. R. Gonçalves and J. B. Formiga, “The black hole energy and the energy of the other universe”, *Eur. Phys. J. Plus* **136**, 10, 1063 (2021).
- [23] J. B. Formiga and V. R. Gonçalves, “Generalization of the ADM gravitational energy momentum”, *Phys. Rev. D* **106**, 4, 044021 (2022).
- [24] “Modified teleparallel gravity: Inflation without an inflaton”, R. Ferraro and F. Fiorini, *Phys. Rev. D* **75**, 084031 (2007).
- [25] “Born-Infeld gravity in Weitzenböck spacetime”, R. Ferraro and F. Fiorini, *Phys. Rev. D* **78**, 124019 (2008).
- [26] G. Bengochea and R. Ferraro, “Dark torsion as the cosmic speed-up” *Phys. Rev. D* **79**, 124019 (2009).
- [27] Yi-Fu Cai, Salvatore Capozziello, Mariafelicia De Laurentis, Emmanuel N. Saridakis, “ $f(T)$ teleparallel gravity and cosmology”, *Rept. Prog. Phys.* **79**, 106901 (2016).

- [28] Sebastian Bahamonde, Konstantinos F. Dialektopoulos, Celia Escamilla-Rivera, Gabriel Farrugia, Viktor Gakis, Martin Hendry, Manuel Hohmann, Jackson Levi Said, Jurgen Mifsud, Eleonora Di Valentino, “Teleparallel Gravity: From Theory to Cosmology”, *Rept. Prog. Phys.* **86**, 026901 (2023).
- [29] Salvatore Capozziello, Maurizio Capriolo and Maria Transirico, “The gravitational energy-momentum pseudotensor: the cases of $f(R)$ and $f(T)$ gravity”, *Int. J. Geom. Meth. Mod. Phys.* **15**, 1850164 (2018).
- [30] J. W. Maluf, J. F. da Rocha-Neto, S. C. Ulhoa and F. L. Carneiro, “Plane Gravitational Waves, the Kinetic Energy of Free Particles and the Memory Effect”, *Gravitation and Cosmology* **24**, 261-266 (2018).
- [31] J. W. Maluf, J. F. da Rocha-Neto, S. C. Ulhoa and F. L. Carneiro, “Kinetic energy and angular momentum of free particles in the gyratonic pp-waves space-times”, *Class. Quantum Grav.* **35**, 115001 (2018).
- [32] J. W. Maluf, J. F. da Rocha-Neto, S. C. Ulhoa and F. L. Carneiro, “Variations of the Energy of Free Particles in the pp-Wave Spacetimes”, *Universe* **2018**, *4*, 74.
- [33] J. W. Maluf, J. F. da Rocha-Neto, S. C. Ulhoa and F. L. Carneiro, “The work-energy relation for particles on geodesics in the pp-wave space-times”, *JCAP03* (2019)028.
- [34] R. Arnowitt, S. Deser and C. W. Misner, “The Dynamics of General Relativity”, in “Gravitation: an Introduction to Current Research”, edited by L. Witten, page 227, Wiley, New York (1962).
- [35] F. H. Hehl, J. Lemke and E. W. Mielke, “Two Lectures on Fermions and Gravity”, in “Geometry and Theoretical Physics”, edited by J. Debrus and A. C. Hirshfeld (Springer, Berlin Heidelberg, 1991).
- [36] J. W. Maluf, F. F. Faria and S. C. Ulhoa, “On reference frames in space-time and gravitational energy in freely falling frames”, *Class. Quantum Grav.* **24** (2007) 2743.
- [37] J. W. Maluf and F. F. Faria, “On the construction of Fermi-Walker transported frames”, *Ann. Phys. (Berlin)* **17** (2008) 326.

- [38] B. Mashhoon and U. Muench, “Length measurement in accelerated systems”, *Ann. Phys. (Leipzig)* **11**, 532 (2002).
- [39] B. Mashhoon, “Vacuum electrodynamics of accelerated systems: Non-local Maxwell’s equations”, *Ann. Phys. (Leipzig)* **12**, 586 (2003).
- [40] J. W. Maluf, “The gravitational energy-momentum tensor and the gravitational pressure”, *Ann. Phys. (Berlin)* **14**, 723 (2005).
- [41] J. B. Hartle, “Gravity: An Introduction to Einstein’s General Relativity” (Addison-Wesley, San Francisco, 2003), p. 198.
- [42] B. Mashhoon, “On the gravitational analogue of Larmor’s theorem”, *Phys. Lett. A* **173**, 347 (1993).
- [43] T. Rege and C. Teitelboim, “Role of surface integrals in the Hamiltonian formulation of general relativity”, *Ann. Phys. (NY)* **88**, 286 (1974); “Constrained Hamiltonian Systems” (Accademia Nazionale dei Lincei, Roma, 1976).
- [44] J. W. Maluf, “The Teleparallel Equivalent of General Relativity and the Gravitational Centre of Mass”, *Universe* **2016**, 2, 19.
- [45] J. W. Maluf and J. F. da Rocha-Neto, “Hamiltonian formulation of general relativity in the teleparallel geometry”, *Phys. Rev D* **64**, 084014 (2001).
- [46] J. F. da Rocha-Neto, J. W. Maluf and S. C. Ulhoa, “Hamiltonian formulation of unimodular gravity in the teleparallel geometry”, *Phys. Rev. D* **82**, 124035 (2010); **87**, 069909(E) (2013).
- [47] J. W. Maluf, J. F. da Rocha-Neto, T. M. L. Toríbio and K. H. Castello-Branco, “Energy and angular momentum of the gravitational field in the teleparallel geometry”, *Phys. Rev. D* **65**, 124001 (2002).
- [48] J. W. Maluf, S. C. Ulhoa, F. F. Faria and J. F. da Rocha-Neto, “The angular momentum of the gravitational field and the Poincaré group”, *Class. Quantum Grav.* **23**, 6245 (2006).
- [49] J. F. da Rocha-Neto and J. W. Maluf, “The angular momentum of plane-fronted gravitational waves in the teleparallel equivalent of general relativity”, *Gen. Relativ. Gravit.* **46**, 1667 (2014).

- [50] R. Penrose, “Extraction of rotational energy from a black hole”, *Riv. Nuovo Cimento* **1**, 252 (1969).
- [51] D. Christodoulou and R. Ruffini, “Reversible and Irreversible Transformations in Black-Hole Physics”, *Phys. Rev. Lett.* **25**, 1596 (1970).
- [52] J. Schwinger, “Quantized gravitational field”, *Phys. Rev.* **130**, 1253 (1963).
- [53] G. Bergqvist, “Quasilocal mass for event horizons”, *Classical Quantum Grav.* **9**, 1753 (1992).
- [54] J. D. Brown and J. W. York, Jr, “Quasi-local energy and conserved charges derived from the gravitational action”, *Phys. Rev. D* **47**, 1407 (1993).
- [55] J. W. Maluf, “On the distribution of gravitational energy in the de Sitter space”, *J. Math. Phys.* **37**, 6293 (1996).
- [56] Mu-Tao Wang and Shing-Tung Yau, “Isometric embeddings into the Minkowski space and new quasi-local mass”, *Commun. Math. Phys.* **288**, 919-942 (2009).
- [57] Mu-Tao Wang and Shing-Tung Yau, “Quasilocal mass in general relativity”, *Phys. Rev. Lett.* **102**, 021101 (2009).
- [58] Warner A. Miller, Shannon Ray, Mu-Tao Wang, and Shing-Tung Yau, “Wang and Yau’s Quasi-Local Energy for an Extreme Kerr Spacetime”, *Class. Quantum Grav.* **35**, 055007 (2018).
- [59] Maciej Dunajski and Paul Tod, “The Kijowski-Liu-Yau Quasi-local Mass of the Kerr Black Hole Horizon”, *Class. Quantum Grav.* **38**, 235001 (2021).
- [60] Mu-Tao Wang, “Gravitational energy seen by quasilocal observers”, *Class. Quantum Grav.* **28**, 114011 (2011).
- [61] H. Bondi, M. G. J. van der Burg and A. W. K. Metzner, “Gravitational waves in general relativity, VII. Waves from axi-symmetric isolated system”, *Proc. R. Soc. London A* **269**, 21 (1962).

- [62] R. K. Sachs, “Gravitational waves in general relativity VIII. Waves in asymptotically flat space-time Proc. R. Soc. London **A270**, 103 (1962).
- [63] A. Trautman, “Conservation Laws in General Relativity”, in “Gravitation: an Introduction to Current Research”, page 169, edited by L. Witten (Wiley, New York, 1962).
- [64] F. A. E. Pirani, “Gravitational Radiation”, in “Gravitation: an Introduction to Current Research”, page 199, edited by L. Witten (Wiley, New York, 1962).
- [65] R. K. Sachs, “Gravitational Radiation”, in “Relativity, Groups and Topology”, page 522, lectures delivered at the Les Houches Summer School 1963, edited by C. de Witt and B. de Witt (Gordon and Breach, New York, 1963).
- [66] J. N. Goldberg, “Invariant Transformations, Conservation Laws and Energy-Momentum”, in “General Relativity and Gravitation”, page 469, edited by A. Held (Plenum, New York, 1980).
- [67] Xiao Zhang, “On the relation between ADM and Bondi energy-momenta”, *Adv. Theor. Math. Phys.* **10**, 261 (2006) [arXiv:gr-qc/0511036].
- [68] Wen-Ling Huang, Shing Tung Yau and Xiao Zhang, “Positivity of the Bondi mass in Bondi’s radiating spacetimes”, *Rend. Lincei Mat. Appl.* **17**, 335 (2006) [arXiv:math/0604155].
- [69] Wen-Ling Huang and Xiao Zhang, “On the relation between ADM and Bondi energy-momenta - radiative spatial infinity”, Proc. of the 4th International Congress of Chinese Mathematicians, Hangzhou 2007, Vol. III, 367-379 (Higher Education Press, 2008) [arXiv:gr-qc/0511037].
- [70] J. W. Maluf, J. F. da Rocha-Neto and S. C. Ulhoa, “Bondi–Sachs energy-momentum and the energy of gravitational radiation”, *Gen. Relativ. Gravit.* **47**, 82 (2015).
- [71] J. Ehlers and W. Kundt, “Exact Solutions of the Gravitational Field Equations”, in “Gravitation: an Introduction to Current Research”, edited by L. Witten (Wiley, New York, 1962).

- [72] P. Jordan, J. Ehlers and W. Kundt, “Republication of: Exact solutions of the field equations of the general theory of relativity”, *Gen. Relativ. Grav.* **41**, 2191 (2009).
- [73] H. W. Brinkmann, “On Riemann spaces conformal to Euclidean spaces”, *Proc. Natl. Acad. Sci. U.S.* **9**, 1 (1923); *Math. Ann.* **94**, 119 (1925).
- [74] J. W. Maluf, J. F. da Rocha-Neto, S. C. Ulhoa and F. L. Carneiro, “Kinetic energy and angular momentum of free particles in the gyratonic pp-waves space-times”, *Class. Quantum Grav.* **35** 115001 (2018).
- [75] M. Hohmann, “Variational Principles in Teleparallel Gravity”, *Universe* **2021**, 7, 114.
- [76] M. Hohmann, “A geometric view on local Lorentz transformations in teleparallel gravity”, *Int. J. Geom. Methods in Theor. Phys.* **19**, 2240001 (2022).
- [77] D. Blixt, R. Ferraro, A. Golovnev and M. J. Guzmán, “Lorentz gauge invariant variables in torsion theories of gravity”, *Phys. Rev. D* **105**, 084029 (2022).
- [78] A. Golovnev, “The geometrical meaning of the Weitzenböck connection”, [arXiv:2302.13599].